## On the conjecture of non-inner automorphisms of finite *p*-groups

Mandeep Singh<sup>\*</sup> and Mahak Sharma<sup>\*\*</sup>

\*Department of Mathematics, Arya College, Ludhiana - 141 001, India

\*\* Department of Mathematics, Goswami Ganesh Dutta Sanatan Dharma College, Chandigarh - 160019, India

Let p be a prime number. A long standing conjecture asserts that every finite non-abelian p-group has a non-inner automorphism of order p. In this paper, we prove that if G is an odd order finite non-abelian monolithic p-group such that every maximal subgroup of G is non-abelian and  $[Z(M), g] \leq Z(G)$  for every maximal subgroup M of G and  $g \in G \setminus M$ . Then G has a non-inner automorphism of order p leaving the Frattini subgroup  $\Phi(G)$ elementwise fixed.

## 2010 Mathematics Subject Classification: Primary: 20D15, Secondary: 20D45.

**Keywords:** Finite *p*-groups, Non-inner automorphisms.

1 Introduction Let G be a finite non-abelian p-group, where p is a prime number. In 1966, Gaschütz [16] proved that G has a non-inner automorphism of p-power order. In 1973, Berkovich [5] proposed the following conjecture (see for eg. [13, Problem 4.13]).

## Prove that every finite non-abelian p-group admits an automorphism of order p which is not an inner one.

Mathematicians are interested in finding the non-inner automorphism of order p which fixes one of the subgroups Z(G),  $\Phi(G)$ ,  $\Omega_1(Z(G))$  or  $\Omega_1(\Phi(G))$  elementwise. In 1965, Liebeck [12] proved that if G is of class 2, then G has a non-inner automorphism of order p if p > 2, and order 2 or 4 if p = 2 which fixes  $\Phi(G)$  elementwise. This conjecture has been settled for the following classes of p-groups: G is regular [6, 15]; the nilpotency class of G is 2 or 3 [1, 12];  $C_G(Z(\Phi(G))) \neq \Phi(G)$ [6]; G/Z(G) is powerful [2]; G is a p-group, where p > 2 such that (G, Z(G)) is a Camina pair [8]; or Z(G) is non-cyclic and W(G) is non-abelian, where  $W(G)/Z(G) = \Omega_1(Z_2(G)/Z(G))$ [9]. For more details on this conjecture, the readers are advised to see ([3], [7], [11], [17]) and references therein.

It is observed that in most of the cases if the conjecture is false, then the center Z(G) is cyclic. The main motivation behind this paper is the following natural question:

Question 1: Let G be a finite non-abelian p-group with cyclic center Z(G). Under what additional conditions on G does the conjecture hold?

Most recently, in 2024, Komma [11] answered this question by proving that if G is an odd order p-group with cyclic center satisfying  $C_G(G^p\gamma_3(G)) \cap Z_3(G) \leq Z(\Phi(G))$ , then the conjecture is true.

It follows from the main result of Attar [4] that if G is a finite non-abelian p-group such that  $Z_2(G)/Z(G)$  is cyclic. Then G has a non-inner central automorphism of order p which fixes  $\Phi(G)$  elementwise. Therefore the following natural question also arises.

Question 2: Let G be a finite non-abelian p-group such that  $Z_2(G)/Z(G)$  is non-cyclic abelian. Under what additional conditions on G does the conjecture hold?

In this paper, we give an affirmative answer to these questions in case of monolithic *p*-groups. Infact, we prove that if *G* is an odd order finite non-abelian monolithic *p*-group such that every maximal subgroup of *G* is non-abelian and  $[Z(M), g] \leq Z(G)$  for every maximal subgroup *M* of *G* and  $g \in G \setminus M$ , then *G* has a non-inner automorphism of order *p* leaving the Frattini subgroup  $\Phi(G)$  elementwise fixed.

**2** Main results We first recall from [10] that a group G is said to be monolithic if and only if there exists a non-trivial normal subgroup of G which is contained in every non-trivial normal subgroup of G. Equivalently, a p-group G is said to be monolithic if  $Z(G) \simeq C_p$ . For  $x \in G$ ,  $\{[x,G]\}$  denotes the set  $\{[x,g] \mid g \in G\}$ . It can be checked that for a finite non-abelian p-group G if  $[Z(M),g] \subseteq Z(G)$  for every non-abelian maximal subgroup M of G and  $g \in G \setminus M$ , then  $[Z(M),g] \leq Z(G)$ . All other unexplained notations are standard.

We will use the following well known lemmas.

**Lemma 2.1.** Let  $n \in N$ ,  $x \in Z_2(G)$  and  $y \in G$ . Then

- (i)  $(xy)^n = x^n y^n [y, x]^{\frac{n(n-1)}{2}}$ .
- (*ii*)  $[x^n, y] = [x, y]^n = [x, y^n].$

**Lemma 2.2.** Let G be a finite p-group, M be a maximal subgroup of G and  $g \in G \setminus M$ . Let  $u \in Z(M)$  such that  $(gu)^p = g^p$ . Then the map  $\alpha$  given by  $g \to gu$  and  $m \to m$ , for all  $m \in M$ , can be extended to an automorphism of G of order p that fixes M elementwise.

**Theorem 2.3.** Let G be a finite non-abelian monolithic p-group (p > 2) such that every maximal subgroup of G is non-abelian and  $[Z(M), g] \leq Z(G)$  for every maximal subgroup M of G and  $g \in G \setminus M$ . Then G has a non-inner automorphism of order p which fixes  $\Phi(G)$  elementwise.

Notice that we cannot drop the condition that every maximal subgroup M in the hypothesis is non-abelian. Because if G has an abelian maximal subgroup M, then  $C_G(Z(\Phi(G))) \neq \Phi(G)$ . Suppose, if possible,  $C_G(Z(\Phi(G))) = \Phi(G)$ . Then  $C_G(\Phi(G)) = Z(\Phi(G))$ . Therefore

$$M \le C_G(M) \le C_G(\Phi(G)) = Z(\Phi(G)) = \Phi(G).$$

Thus  $M = \Phi(G)$ , which is a contradiction. It follows from [6] that G has a non-inner automorphism of order p which fixes  $\Phi(G)$  elementwise.

*Proof.* Since G is a finite non-abelian monolithic p-group, |Z(G)| = p. Suppose if possible that G has no non-inner automorphism of order p leaving the Frattini subgroup  $\Phi(G)$  elementwise fixed. It follows from [4] that  $Z_2(G)/Z(G)$  is not cyclic. Also by [6],  $C_G(Z(\Phi(G))) = \Phi(G)$ .

We claim that  $Z_2(G)$  is abelian. It follows from [14, 5.2.22] that

$$\exp(Z_2(G)/Z(G)) \le \exp(Z(G)) = p.$$

Since  $Z_2(G)/Z(G) = Z(G/Z(G))$  is abelian,  $Z_2(G)/Z(G)$  is elementary abelian. Therefore  $g^p \in Z(G)$  for all  $g \in Z_2(G)$  which implies that  $[x, g^p] = 1$  for all  $x \in G$ . Now for  $g \in Z_2(G)$  and  $x \in G$ , it follows from Lemma 2.1 that  $[x^p, g] = [x, g]^p = [x, g^p] = 1$ . Also by Grun's Lemma  $[Z_2(G), G'] = 1$ . Thus

$$[Z_2(G), \Phi(G)] = [Z_2(G), G^p][Z_2(G), G'] = 1.$$

Therefore  $Z_2(G) \leq C_G(\Phi(G)) = Z(\Phi(G))$ . Thus  $Z_2(G)$  is abelian.

Next we prove that  $Z(M) \leq Z_2(G)$  for every maximal subgroup M of G. Suppose, if possible that Z(M) is not contained in  $Z_2(G)$  for some maximal subgroup M of G. Let  $m \in Z(M) \setminus Z(G)$  such that  $m \notin Z_2(G)$ . Then there exists some  $x \in G \setminus M$  such that  $[m, x] \notin Z(G)$ , which is a contradiction to the hypothesis.

Next we prove that  $Z_2(G)$  contains an element of order p which does not belong to Z(G). Since d(Z(G)) = 1, it follows from [2, corollary 2.3] that  $d(Z_2(G)/Z(G)) = d(G)$ . Observe that  $Z_2(G)$  can not be cyclic. Because if  $d(Z_2(G)) = 1$ , then  $d(Z_2(G)/Z(G)) = 1$ , and therefore

$$d(G) = d(Z_2(G)/Z(G)) = 1,$$

which is a contradiction. As  $Z_2(G)$  is abelian, it implies that  $d(\Omega_1(Z_2(G))) = d(Z_2(G)) \ge 2$ . Since |Z(G)| = p, therefore  $Z_2(G)$  contains an element of order p which does not belong to Z(G).

Consider  $u \in \Omega_1(Z_2(G)) \setminus Z(G)$ . Define a map  $\sigma : G \to Z(G)$  by  $\sigma(x) = [x, u]$  for all  $x \in G$ . The map  $\sigma$  is surjective with ker $\sigma = C_G(u)$ . Therefore the subgroup  $M = C_G(u)$  is a maximal subgroup of G. Let  $g \in G \setminus M$ , it follows from Lemma 2.1 that

$$(gu)^p = g^p u^p [u,g]^{\frac{p(p-1)}{2}} = g^p.$$

By Lemma 2.2, the map  $\alpha : G \to G$  defined by  $\alpha(g) = gu$  and  $\alpha(m) = m$  for all  $m \in M$  can be extended to an automorphims of order p. By hypothesis  $\alpha$  must be an inner automorphism. Let  $\alpha = \beta_t$ , the inner automorphim induced by t. Since  $\alpha$  is the identity on M,  $t \in C_G(M)$ . If  $t \notin M$ , then  $G = M\langle t \rangle$ . It implies that  $t \in Z(G) \leq \Phi(G) \leq M$ , which is a contradiction. It follows that  $t \in Z(M) \leq Z_2(G)$ . Thus  $u = g^{-1}\alpha(g) = [g, t] \in Z(G)$ , which contradicts the choice of u. Hence G has a non-inner automorphism of order p which fixes  $\Phi(G)$  elementwise.  $\Box$ 

Attar [4] proved that if G is a finite non-abelian p-group such that  $C_G(Z(\Phi(G))) = \Phi(G)$ and  $\frac{Z_2(G) \cap Z(\Phi(G))}{Z(G)}$  is not elementary abelian of rank rs, where r = d(G) and s = d(Z(G)), then G has a non-inner central automorphism of order p which fixes  $\Phi(G)$  elementwise. The following corollary, which is an immediate consequence of Theorem 2.3, confirms that the conjecture holds if G is a finite non-abelian monolithic p-group of odd order such that  $C_G(Z(\Phi(G))) = \Phi(G)$ and  $\frac{Z_2(G) \cap Z(\Phi(G))}{Z(G)}$  is elementary abelian of rank r, where r = d(G).

**Corollary 2.4.** Let G be a finite non-abelian monolithic p-group (p > 2) such that  $C_G(Z(\Phi(G))) = \Phi(G)$  and  $[Z(M), g] \leq Z(G)$  for every maximal subgroup M of G and  $g \in G \setminus M$ . Then G has a non-inner automorphism of order p which fixes  $\Phi(G)$  elementwise.

*Proof.* Since  $C_G(Z(\Phi(G))) = \Phi(G)$ , every maximal subgroup of G is non-abelian. It follows from the proof of Theorem 2.3 that  $Z_2(G) \leq Z(\Phi(G))$ . Thus  $\frac{Z_2(G) \cap Z(\Phi(G))}{Z(G)} = \frac{Z_2(G)}{Z(G)}$  is elementary abelian and  $d(\frac{Z_2(G) \cap Z(\Phi(G))}{Z(G)}) = d(G)$ . Rest of the proof is similar.

We conclude this paper with an example of a group that satisfies the hypothesis of Theorem 2.3.

Consider

$$G = \langle f_1, f_2, f_3, f_4, f_5, f_6, f_7 \rangle,$$

with the relations:  $[f_2, f_1] = f_3, [f_1, f_7] = [f_2, f_7] = [f_3, f_6] = [f_4, f_5] = 1, f_1^3 = f_4, f_2^3 = f_3, f_3^3 = f_5, f_4^3 = f_6, f_5^3 = f_7, f_6^3 = f_7^3 = 1$ . This group is the group number 194 in the GAP Library of groups of order  $3^7$ . The nilpotency class of G is 4. In this group  $Z(G) = \langle f_7 \rangle$  and therefore |Z(G)| = 3. Thus G is a monolithic 3-group. This group has four maximal subgroups  $M_1, M_2, M_3$  and  $M_4$ , where

$$M_{1} = \langle f_{1}, f_{3}, f_{4}, f_{5}, f_{6}, f_{7} \rangle, \ Z(M_{1}) = \langle f_{6}, f_{7} \rangle,$$
$$M_{2} = \langle f_{2}, f_{3}, f_{4}, f_{5}, f_{6}, f_{7} \rangle, \ Z(M_{2}) = \langle f_{5}, f_{7} \rangle,$$
$$M_{3} = \langle f_{1}f_{2}^{2}, f_{3}, f_{4}, f_{5}, f_{6}, f_{7} \rangle, \ Z(M_{3}) = \langle f_{5}^{2}f_{6}f_{7}, f_{7}^{2} \rangle$$
$$M_{4} = \langle f_{1}f_{2}, f_{3}, f_{4}, f_{5}, f_{6}, f_{7} \rangle, \ Z(M_{4}) = \langle f_{5}f_{6}f_{7}^{2}, f_{7} \rangle.$$

Therefore  $\Phi(G) = \langle f_3, f_4, f_5, f_6, f_7 \rangle$ . Observe that since  $\Phi(G)$  is non-abelian, it implies that  $M_1, M_2, M_3$  and  $M_4$  are also non-abelian.

One can check that  $[Z(M), g] \leq Z(G)$  for all  $M \in \{M_1, M_2, M_3, M_4\}$  and  $g \in G \setminus M$ . This group has 4374 automorphisms. One of these automorphism is  $\alpha$ , where

 $\alpha(f_i) = f_i \text{ for all } i \in \{1, 3, 4, 5, 6, 7\} \text{ and } \alpha(f_2) = f_2 f_6.$ 

Note that  $\alpha$  is a non-inner automorphism of order 3 which fixes  $\Phi(G)$  element-wise.

## References

- A. Abdollahi, Finite p-groups of class 2 have a noninner automorphism of order p, J. Algebra, **312** (2007), 876-879.
- [2] A. Abdollahi, Powerful p-groups have noninner automorphism of order p and some cohomology, J. Algebra 323 (2010), 779-789.
- [3] A. Abdollahi and S. M. Ghoraishi, On Noninner 2-automorphisms of finite 2-groups, Bull. Aust. Math. Soc., 90 (2014), 227–231.
- M. S. Attar, On a conjecture about automorphisms of finite p-groups, Arch. Math., 93 (2009), 399–403.
- [5] Y. Berkovich, Groups of Prime Power Order, vol. 1, Walter de Gruyter, Berlin, (2008).
- [6] M. Deaconescu and G. Silberberg, Noninner automorphism of order p of finite p-groups, J. Algebra, 250 (2002), 283-287.
- [7] S. Fouladi and R. Orfi, Noninner automorphisms of order p in finite p-groups of coclass 2, when p > 2, Bull. Aust. Math. Soc., **90** (2014), 232-236.
- [8] S. M. Ghoraishi, A note on automorphisms of finite p-groups, Bull. Aust. Math. Soc., 87 (2013), 24-26.
- R. Garg and M. Singh, Finite p-groups with non-cyclic center have a non-inner automorphism of order p, Arch. Math., 117 (2) (2021), 129–132.
- [10] P. Hall, On the finiteness of certain soluble groups, Proc. London Math. Soc., 3(9) (1959), 595-622.
- P. Komma, Non-inner automorphisms of order p in finite p-groups of coclass 4 and 5, Int. J. Group Theory, 13(1) (2024), 97–114.
- [12] H. Liebeck, Outer automorphisms in nilpotent p-groups of class 2, J. London Math. Soc., 40 (1965), 268-275.
- [13] V. D. Mazurov and E. I. Khukhro, Unsolved problems in group theory, the kourovka notebook, Vol. 16, Russian Academy of Sciences, Siberian Branch, Sobolev Institute of Mathematics, Novosibirsk (2006).
- [14] D. J. S. Robinson, A course in the Theory of Groups, New York Inc.: Springer-Verlag, (1996).
- [15] P. Schmid, A cohomological property of regular p-groups, Math. Z., 175 (1980), 1-3.
- [16] W. Gaschütz, Nichtabelsche p-Gruppen besitzen äußere p-Automorphismen, J. Algebra, 4 1966, 1-2.
- [17] M. Ruscitti, L. Legarreta and M. K. Yadav, Non-inner automorphisms of order p in finite p-groups of coclass 3, Monatsh Math., 183(4) (2016), 679-697.