

Tilting mutations as generalized Kauer moves for (skew) Brauer graph algebras with multiplicity

VALENTINE SOTO

Abstract. Generalized Kauer moves are local moves of multiple edges in a Brauer graph that yield derived equivalences between Brauer graph algebras of multiplicity identically 1 [Sot24]. Moreover, these derived equivalences are given by a tilting mutation. The goal of this paper is to generalize this result first for Brauer graph algebras with arbitrary multiplicity and second for a generalization of Brauer graph algebras called skew Brauer graph algebras. In these contexts, we prove that the generalized Kauer moves induce derived equivalences via tilting mutations. We also show that skew Brauer graph algebras of multiplicity identically 1 can be seen as the trivial extension of skew gentle algebras.

Contents

Introduction	1
Notations and conventions	4
1. Skew group algebras	4
1.1. Trivial extension	5
1.2. Morphism space of G -invariant objects	6
1.3. Silting mutation	8
2. Generalized Kauer moves for Brauer graphs with multiplicity	10
2.1. Brauer graph algebras with multiplicity	11
2.2. Covering of Brauer graph algebras with multiplicity	13
2.3. Generalized Kauer moves	16
2.4. Compatibility with silting mutations	21
3. Generalized Kauer moves for skew Brauer graph algebras	25
3.1. Skew Brauer graph algebras	25
3.2. Covering of skew Brauer graphs	28
3.3. Generalized Kauer moves	32
3.4. Compatibility with silting mutations	35
References	38

Introduction

Tilting theory plays an important role in the representation theory of algebras as it yields derived equivalences [Hap87; Ric89]. Given a tilting object, one can construct others under some

assumptions via the notion of mutation [BGP73; APR79; RS91; Oku98]. This operation consists in replacing a direct summand of the given object using an approximation. Unfortunately, tilting mutation does not always exist. This problem has led to the introduction of silting mutations in triangulated categories which generalizes these tilting mutations [AI12]. The goal of this paper is to describe the silting mutation for a certain class of algebras called skew Brauer graph algebras, which generalizes the class of Brauer graph algebras.

Brauer graph algebras were first introduced in modular representation theory of finite groups by Donovan and Freislich [DF78]. They are finite dimensional algebras that are constructed thanks to the combinatorial data of a Brauer graph. A Brauer graph is a finite graph equipped with a multiplicity function on each vertex and with a cyclic ordering of the edges around each vertex. Moreover, Brauer graph algebras are symmetric special biserial algebras [Sch15]. Hence, the representation theory of Brauer graph algebras is well-known [DF78; WW85; ES92; Gre74; Rog98]. We refer to the survey [Sch18] for a detailed list of results on Brauer graph algebras. In the recent years, the study of derived equivalences of Brauer graph algebras has drawn some attention. Antipov and Zvonareva proved that this class of algebras is closed under derived equivalences [AZ22]. Moreover, Opper and Zvonareva described a complete numerical derived invariant for Brauer graph algebras [OZ22] (see also [Zvo24] for a survey on derived equivalences of Brauer graph algebras).

Silting mutation in Brauer graph algebras can be understood in terms of generalized Kauer moves which are particular moves of edges in the corresponding Brauer graph. These moves were first described by Kauer in the case of one edge [Kau98]. Aihara proved that these moves of one edge give rise to an irreducible tilting mutation in the case of Brauer tree algebras (i.e. Brauer graph algebras whose underlying graph is a tree). These moves were then generalized in [Sot24] to describe all silting mutations in Brauer graph algebras of multiplicity identically one. Since Brauer graph algebras are symmetric [Sch15], the silting mutation in these algebras are in fact tilting. Hence, these generalized Kauer moves yield derived equivalences of Brauer graph algebras. The first main result of this paper is the generalization of these results for Brauer graph algebras with arbitrary multiplicity.

Theorem (Theorem 2.20). *Let $\Gamma = (\Gamma_0, \Gamma_1)$ be a Brauer graph and B be its corresponding Brauer graph algebra. We assume that the least common multiple \bar{m} of the multiplicities in Γ is invertible in the field k . Then, for any subset of edges E in Γ , the following holds.*

- (1) *The Brauer graph algebra B' obtained by a generalized Kauer move of E in Γ is derived equivalent to B .*
- (2) *The silting mutation of B over the projective B -module corresponding to the edges in $\Gamma_1 \setminus E$ is tilting and its endomorphism algebra is isomorphic to B' .*

The idea of the proof is to construct a $\mathbb{Z}/\bar{m}\mathbb{Z}$ -covering $\tilde{\Gamma}$ of Γ which is a Brauer graph of multiplicity identically one using [Asa19]. In this case, one can recover B from the Brauer graph algebra \tilde{B} associated to $\tilde{\Gamma}$ thanks to the construction of skew group algebras as defined in [RR85] (cf Proposition 2.11). Moreover, there is a commutativity between constructing this covering and applying the generalized Kauer moves. This can be summarized with the following commutative diagram

$$\begin{array}{ccc}
\tilde{\Gamma} & \xrightarrow[\text{move of } \tilde{H}']{\text{generalized Kauer}} & \mu_{\tilde{H}'}^+(\tilde{\Gamma}) \\
\downarrow \mathbb{Z}/\bar{m}\mathbb{Z}\text{-covering} & & \downarrow \mathbb{Z}/\bar{m}\mathbb{Z}\text{-covering} \\
\Gamma & \xrightarrow[\text{move of } H']{\text{generalized Kauer}} & \mu_{H'}^+(\Gamma)
\end{array}$$

where \tilde{H}' is the set of half-edges in $\tilde{\Gamma}$ consisting of all the lifts of the elements in H' .

In a second part of this paper, we generalize the previous result for skew Brauer graph algebras, which generalize the class of Brauer graph algebras. Their construction relies on the combinatorial data of a skew Brauer graph which is a Brauer graph with possible “degenerate” edges, meaning that the corresponding half-edges are fixed by the pairing. The second main result of this paper describes the combinatorics of silting mutation for skew Brauer graph algebras.

Theorem (Theorem 3.19). *Let $\Gamma = (\Gamma_0, \Gamma_1)$ be a skew Brauer graph and B be its corresponding skew Brauer graph algebra. We assume that the least common multiple \bar{m} of the multiplicities in Γ and 2 are invertible in the field k . Then, for any subset of edges E in Γ , the following holds.*

- (1) *The Brauer graph algebra B' obtained by a generalized Kauer move of E in Γ is derived equivalent to B .*
- (2) *The silting mutation of B over the projective B -module corresponding to the edges in $\Gamma_1 \setminus E$ is tilting and its endomorphism algebra is isomorphic to B' .*

The proof for skew Brauer graph algebras is analogous to the one for Brauer graph algebras with arbitrary multiplicity. We first construct a $\mathbb{Z}/2\mathbb{Z}$ -covering $\tilde{\Gamma}$ of Γ which is a standard Brauer graph (with multiplicity). In this case, one can recover B from the Brauer graph algebra \tilde{B} associated to $\tilde{\Gamma}$ thanks to the construction of skew group algebras as defined in [RR85] (cf Proposition 3.9). Again, there is a commutativity between constructing this covering and applying the generalized Kauer moves. This can be summarized with the following commutative diagram

$$\begin{array}{ccc}
\tilde{\Gamma} & \xrightarrow[\text{move of } \tilde{H}']{\text{generalized Kauer}} & \mu_{\tilde{H}'}^+(\tilde{\Gamma}) \\
\downarrow \mathbb{Z}/2\mathbb{Z}\text{-covering} & & \downarrow \mathbb{Z}/2\mathbb{Z}\text{-covering} \\
\Gamma & \xrightarrow[\text{move of } H']{\text{generalized Kauer}} & \mu_{H'}^+(\Gamma)
\end{array}$$

where \tilde{H}' is the set of half-edges in $\tilde{\Gamma}$ consisting of all the lifts of the elements in H' . Using this covering, we also show that skew Brauer graph algebras of multiplicity identically 1 can be understood as trivial extension of skew gentle algebras (cf Theorem 3.11). This generalizes Theorem 3.13 in [Sch18] which interprets Brauer graph algebra of multiplicity identically 1 as trivial extension of gentle algebras.

Note that a notion of skew Brauer graph algebra is also studied in an ongoing project by Elsenner, Guazzelli and Valdivieso. Some of their results were presented in a conference in Oslo.

Organisation of the paper. In the first section, we recall the definition and some properties of skew group algebras. In particular, we study the trivial extension, the silting mutation and the morphism spaces of G -invariant objects in skew group algebras. These are tools needed in the next two sections.

In the second section, we prove our first main result for Brauer graph algebras with arbitrary multiplicity. For this, we first recall the construction of the covering by a Brauer graph of multiplicity identically 1 in [Asa19]. Then, we explain how one can recover B from the Brauer graph algebra associated to its covering thanks to the construction of skew group algebras. Finally, we show a commutativity between the covering and the generalized Kauer moves.

The third section follows the same outline than the second for skew Brauer graph algebras. We also explain how skew Brauer graph algebras of multiplicity identically 1 can be seen as trivial extensions of skew gentle algebras.

Acknowledgments. This paper is part of a PhD thesis supervised by Claire Amiot and supported by the CNRS. The author deeply thanks her advisor for taking the time to read this paper and for the helpful advice and comments on it. Part of this work was done while the author was in Sherbrooke University. The mobility was co funded by the Mitacs Globalink Research Award and the IDEX. She also thanks Thomas Brüstle for interesting discussions and Sherbrooke University for warm welcome.

Notations and conventions

In this paper, all algebras are supposed to be finite dimensional over an algebraically closed field k . For any algebra Λ , we denote by $\text{mod}(\Lambda)$ the category of finitely generated right Λ -modules, $\text{D}^b(\Lambda)$ the bounded derived category associated to $\text{mod}(\Lambda)$ and $\text{per}(\Lambda)$ the full subcategory of $\text{D}^b(\Lambda)$ consisting of perfect complexes. Moreover, arrows in a quiver will be composed from right to left : for arrows α, β , we write $\beta\alpha$ for the path from the start of α to the target of β . Similarly, morphisms will be composed from right to left.

1. Skew group algebras

In this section, we collect some properties of skew group algebras. First, we recall the definition of a skew group algebra from [RR85]. Throughout this section, G denotes a finite abelian group whose order $|G|$ is invertible in the field k . By a G -action on an algebra, we mean a left action by automorphisms.

Definition 1.1. Let Λ be a k -algebra with a G -action. The *skew group algebra* ΛG associated to Λ is the algebra defined by $\Lambda G = \Lambda \otimes_k kG$ as a k -vector space and whose multiplication is induced by the G -action on Λ i.e.

$$(\lambda \otimes g) \cdot (\mu \otimes h) = \lambda(g \cdot \mu) \otimes gh$$

for all $\lambda, \mu \in \Lambda, g, h \in G$ and we extend this relation by linearity and by distributivity.

Denoting by \widehat{G} the dual of G , note that ΛG has a natural \widehat{G} -action which is given by

$$\chi \cdot (\lambda \otimes g) = \chi(g)\lambda \otimes g$$

for all $\chi \in \widehat{G}, \lambda \in \Lambda, g \in G$ and where the relation is extended by linearity.

Moreover, given two algebras Λ and Λ' with a G -action and M a Λ - Λ' -bimodule, one can define $MG := M \otimes_k kG$. It has a natural structure of ΛG - $\Lambda' G$ -bimodule defined by

$$(\lambda \otimes g) \cdot (m \otimes h) = \lambda(g \cdot m) \otimes gh \quad \text{and} \quad (m \otimes h) \cdot (\lambda' \otimes g') = m(h \cdot \lambda') \otimes hg'$$

for all $\lambda \in \Lambda$, $\lambda' \in \Lambda'$, $g, h, g' \in G$ and we extend these relations by linearity and by distributivity.

1.1. Trivial extension

We first study the trivial extension of a skew group algebra. Let us recall the definition of the trivial extension of an algebra. We denote by $D = \text{Hom}_k(-, k)$ the standard duality on $\text{mod}(\Lambda)$.

Definition 1.2. Let Λ be a k -algebra. The *trivial extension* $\text{Triv}(\Lambda)$ of Λ by the Λ - Λ -bimodule $D\Lambda$ is the algebra defined by $\text{Triv}(\Lambda) = \Lambda \oplus D\Lambda$ as a k -vector space and whose multiplication is induced by the Λ - Λ -bimodule structure of $D\Lambda$ i.e.

$$(a, \phi) \cdot (b, \psi) = (ab, a\psi + \phi b)$$

for all $a, b \in \Lambda$ and $\phi, \psi \in D\Lambda$.

Let Λ be a k -algebra with a G -action. Then, G acts on $D\Lambda$ as follows

$$g \cdot \phi : (b \mapsto \phi(g^{-1} \cdot b))$$

for all $g \in G$ and $\phi \in D\Lambda$. In particular, $(D\Lambda)G = D\Lambda \otimes_k \Lambda G$ has a natural structure of ΛG - ΛG -bimodule.

Lemma 1.3. For any k -algebra Λ with a G -action, there is an isomorphism of ΛG - ΛG -bimodules

$$(D\Lambda)G \simeq D(\Lambda G)$$

Proof. The isomorphism of ΛG - ΛG -bimodules is given by

$$\begin{aligned} D(\Lambda G) &\longrightarrow (D\Lambda)G \\ \phi &\longmapsto \sum_{h \in G} \phi_h \otimes h \\ \psi_g &\longleftarrow \psi \otimes g \end{aligned}$$

where $\phi_h : \lambda \mapsto \phi((h^{-1} \cdot \lambda) \otimes h^{-1}) \in D\Lambda$ and $\psi_g : (\lambda \otimes h) \mapsto \psi(\delta_{h^{-1}}^g(g \cdot \lambda)) \in D(\Lambda G)$ with δ being the Kronecker symbol. \square

Hence, the G -action on Λ induces a G -action on $\text{Triv}(\Lambda)$ which is given by

$$g \cdot (a, \phi) = (g \cdot a, g \cdot \phi)$$

for all $g \in G$ and $(a, \phi) \in \text{Triv}(\Lambda)$. In particular, one can construct the skew group algebra associated to $\text{Triv}(\Lambda)$.

Proposition 1.4. For any k -algebra Λ with a G -action, there is an isomorphism of algebras

$$\text{Triv}(\Lambda G) \simeq \text{Triv}(\Lambda)G$$

Proof. Let us define the following linear map

$$\begin{aligned}\Phi : \operatorname{Triv}(\Lambda G) &\longrightarrow \operatorname{Triv}(\Lambda)G \\ (a \otimes g, \phi) &\longmapsto (a, 0) \otimes g + \sum_{h \in G} (0, \phi_h) \otimes h\end{aligned}$$

where ϕ_h is the morphism in $D\Lambda$ defined by

$$\phi_h : b \longmapsto \phi(h^{-1} \cdot b \otimes h^{-1})$$

We denote by $A := \Phi(a \otimes g, \phi)\Phi(a' \otimes g', \phi')$. We have,

$$\begin{aligned}A &= \left[(a, 0) \otimes g + \sum_{h \in G} (0, \phi_h) \otimes h \right] \left[(a', 0) \otimes g' + \sum_{h \in G} (0, \phi'_h) \otimes h \right] \\ &= (a(g \cdot a'), 0) \otimes gg' + \sum_{h \in G} (0, \phi_h \cdot (h \cdot a')) \otimes hg' + \sum_{h \in G} (0, a \cdot (g \cdot \phi'_h)) \otimes gh \\ &= (a(g \cdot a'), 0) \otimes gg' + \sum_{h \in G} (0, (\phi \cdot (a' \otimes g'))_h) \otimes h + \sum_{h \in G} (0, ((a \otimes g) \cdot \phi')_h) \otimes h \\ &= \Phi((a \otimes g, \phi)(a' \otimes g', \phi'))\end{aligned}$$

for all $(a \otimes g, \phi), (a' \otimes g', \phi') \in \operatorname{Triv}(\Lambda G)$. Hence, Φ is a morphism of algebras. Moreover, one can easily check that Φ is also invertible. Its inverse is given by

$$\begin{aligned}\operatorname{Triv}(\Lambda)G &\longrightarrow \operatorname{Triv}(\Lambda G) \\ (b, \psi) \otimes g &\longmapsto (b \otimes g, \psi_g)\end{aligned}$$

where ψ_g is the morphism in $D(\Lambda G)$ given by

$$\psi_g : (a \otimes h) \longmapsto \psi(\delta_{h^{-1}}^g(g \cdot a))$$

where δ denotes the Kronecker symbol. This concludes the proof. \square

1.2. Morphism space of G -invariant objects

In this subsection, we are interested in the morphism space between G -invariant objects in the bounded derived category. The case of the endomorphism space of a G -invariant object has already been examined in [AB22, Theorem 2.10].

A G -action on a finite dimensional k -algebra Λ induces a right G -action on its bounded derived category $\operatorname{D}^b(\Lambda)$ in the sense of [Ela15, Definition 3.1]. This action is given by the autoequivalences $- \otimes_{\Lambda}^{\mathbf{L}} \Lambda_g$ for all $g \in G$, where Λ_g is the Λ -module whose underlying group is Λ and whose action is twisted by g . In the following, X^g denotes $X \otimes_{\Lambda}^{\mathbf{L}} \Lambda_g$ for all $X \in \operatorname{D}^b(\Lambda)$ and $f^g : X^g \rightarrow Y^g$ denotes $f \otimes_{\Lambda}^{\mathbf{L}} \Lambda_g$ for all morphism $f : X \rightarrow Y$ in $\operatorname{D}^b(\Lambda)$. Moreover,

$$\operatorname{D}^b(\Lambda) \begin{array}{c} \xrightarrow{- \otimes_{\Lambda}^{\mathbf{L}} \Lambda G} \\ \xleftarrow{\operatorname{Res}} \end{array} \operatorname{D}^b(\Lambda G)$$

form an adjoint pair of triangulated functors in both directions and the unit of the adjunction splits [LeM20, Lemma 2.3.1]. In particular, we obtain a functorial isomorphism for all $X \in \mathbf{D}^b(\Lambda)$

$$\mathrm{Res}(X \otimes_{\Lambda}^{\mathbf{L}} \Lambda G) \simeq \bigoplus_{g \in G} X^g \quad (1)$$

Definition 1.5. An object $X \in \mathbf{D}^b(\Lambda)$ is said to be *G-invariant* if there exist isomorphisms $\iota_g : X^{g^{-1}} \rightarrow X$ for all $g \in G$ satisfying that $\iota_{gh} = \iota_g(\iota_h)^{g^{-1}}$ for all $g, h \in G$.

Example 1.6. (1) It is easy to check that Λ is a G -invariant object in $\mathbf{D}^b(\Lambda)$ where the isomorphisms $\iota_g : \Lambda^{g^{-1}} \simeq \Lambda_{g^{-1}} \rightarrow \Lambda$ are given by the action of $g \in G$. Moreover, if e is an idempotent of Λ stable under the action of G , the same isomorphisms give us that $e\Lambda$ is also a G -invariant object.

(2) Let $f : X \rightarrow Y$ be a morphism in $\mathbf{D}^b(\Lambda)$ between two G -invariant objects. We denote by $\iota_g^X : X^{g^{-1}} \rightarrow X$ and $\iota_g^Y : Y^{g^{-1}} \rightarrow Y$ the isomorphisms corresponding to X and Y respectively. Let us assume that for all $g \in G$ the following diagram commutes

$$\begin{array}{ccc} X^{g^{-1}} & \xrightarrow{f^{g^{-1}}} & Y^{g^{-1}} \\ \downarrow \iota_g^X & & \downarrow \iota_g^Y \\ X & \xrightarrow{f} & Y \end{array}$$

Then, the cone of f denoted $\mathrm{Cone}(f)$ is also G -invariant. Indeed, one can check that the isomorphisms $\iota_g : \mathrm{Cone}(f)^{g^{-1}} \rightarrow \mathrm{Cone}(f)$ defined by

$$\iota_g = \begin{pmatrix} \iota_g^X[1] & 0 \\ 0 & \iota_g^Y \end{pmatrix}$$

for all $g \in G$ satisfy the condition of Definition 1.5.

We now adapt the results about the endomorphism algebra of a G -invariant object from [AB22] to the morphism space of G -invariant objects. The proof of the following statements follows the same outline than the proofs of Lemma 2.5 and Proposition 2.8 in [AB22].

Lemma 1.7. *Let X and Y be two G -invariant objects in $\mathbf{D}^b(\Lambda)$. Then, G acts on the morphism space $\mathrm{Hom}_{\mathbf{D}^b(\Lambda)}(X, Y)$ as follows*

$$g \cdot f = \iota_g^Y \circ f^{g^{-1}} \circ (\iota_g^X)^{-1}$$

for all $g \in G$ and $f \in \mathrm{Hom}_{\mathbf{D}^b(\Lambda)}(X, Y)$, where $\iota_g^X : X^{g^{-1}} \rightarrow X$ and $\iota_g^Y : Y^{g^{-1}} \rightarrow Y$ are the isomorphisms corresponding to X and Y respectively.

Given two G -invariant objects X and Y in $\mathbf{D}^b(\Lambda)$, $\mathrm{Hom}_{\mathbf{D}^b(\Lambda)}(X, Y)G = \mathrm{Hom}_{\mathbf{D}^b(\Lambda)}(X, Y) \otimes_k kG$ has a natural structure of $(\mathrm{End}_{\mathbf{D}^b(\Lambda)}(Y)G)$ - $(\mathrm{End}_{\mathbf{D}^b(\Lambda)}(X)G)$ -bimodule (cf after Definition 1.1). Moreover, thanks to Theorem 2.10 in [AB22], there is an isomorphism of algebras

$$\mathrm{End}_{\mathrm{D}^b(\Lambda)}(Z)G \simeq \mathrm{End}_{\mathrm{D}^b(\Lambda G)}(Z \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G)$$

for all G -invariant object Z in $\mathrm{D}^b(\Lambda)$. Hence, $\mathrm{Hom}_{\mathrm{D}^b(\Lambda G)}(X \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G, Y \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G)$ has also a natural structure of $(\mathrm{End}_{\mathrm{D}^b(\Lambda)}(Y)G)$ - $(\mathrm{End}_{\mathrm{D}^b(\Lambda)}(X)G)$ -bimodule.

Proposition 1.8. *Let X and Y be two G -invariant objects in $\mathrm{D}^b(\Lambda)$ and $\iota_g^X : X^{g^{-1}} \rightarrow X$ and $\iota_g^Y : Y^{g^{-1}} \rightarrow Y$ be the isomorphisms corresponding to X and Y respectively. Then, there is an isomorphism of $(\mathrm{End}_{\mathrm{D}^b(\Lambda)}(Y)G)$ - $(\mathrm{End}_{\mathrm{D}^b(\Lambda)}(X)G)$ -bimodules given by*

$$\begin{aligned} \mathrm{Hom}_{\mathrm{D}^b(\Lambda)}(X, Y)G &\longrightarrow \mathrm{Hom}_{\mathrm{D}^b(\Lambda G)}(X \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G, Y \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G) \\ f \otimes g &\longmapsto ((f \circ \iota_g^X) \otimes 1_{\Lambda G}) \circ L_g^X \end{aligned}$$

where $L_g^X : X \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G \rightarrow X^{g^{-1}} \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G$ is the isomorphism induced by the multiplication on the left by $1 \otimes g$ in ΛG .

1.3. Silting mutation

Let us now study silting mutations for skew group algebras. We first recall the definition and some properties of silting mutations given in [AI12]. Throughout this section, Λ denotes a finite dimensional k -algebra. For any object M in $\mathrm{per}(\Lambda)$, $\mathrm{add}(M)$ denotes the full subcategory of $\mathrm{per}(\Lambda)$ consisting of direct sums of direct summands of M and $\mathrm{thick}(M)$ denotes the smallest triangulated subcategory of $\mathrm{per}(\Lambda)$ containing M that is closed under direct summands.

Definition 1.9. Let M be an object in $\mathrm{per}(\Lambda)$.

- M is said to be *silting* if $\mathrm{Hom}_{\mathrm{per}(\Lambda)}(M, M[> 0]) = 0$ and $\mathrm{thick}(M) = \mathrm{per}(\Lambda)$.
- M is said to be *tilting* if M is silting and $\mathrm{Hom}_{\mathrm{per}(\Lambda)}(M, M[< 0]) = 0$.

Definition 1.10. Let M be an object in $\mathrm{per}(\Lambda)$.

- We say that a morphism $f : X \rightarrow Y$ in $\mathrm{per}(\Lambda)$ is *left minimal* if the only morphisms $h \in \mathrm{End}_{\mathrm{per}(\Lambda)}(Y)$ satisfying $hf = f$ are isomorphisms.
- We say that a morphism $f : X \rightarrow Y$ in $\mathrm{per}(\Lambda)$ is a *left $\mathrm{add}(M)$ -approximation* of X if $Y \in \mathrm{add}(M)$ and $\mathrm{Hom}_{\mathrm{per}(\Lambda)}(f, M)$ is surjective.
- We say that a morphism $f : X \rightarrow Y$ in $\mathrm{per}(\Lambda)$ is a *left minimal $\mathrm{add}(M)$ -approximation* of X if f is left minimal and a left $\mathrm{add}(M)$ -approximation of X . It is unique up to isomorphisms.
- Let M_0 be a direct summand of M and $f : M/M_0 \rightarrow M'_0$ be the left minimal $\mathrm{add}(M_0)$ -approximation of M/M_0 . The *left mutation* of M over M_0 is the object

$$\mu^+(M; M_0) := M_0 \oplus \mathrm{Cone}(f) \in \mathrm{per}(\Lambda)$$

where $\mathrm{Cone}(f)$ denotes the cone of f .

Theorem 1.11 (Aihara-Iyama [AI12, Theorem 2.32]).

- Any left mutation of a silting object of $\text{per}(\Lambda)$ is a silting object of $\text{per}(\Lambda)$.
- For any M_0 direct summand of a tilting object M in $\text{per}(\Lambda)$, the left mutation $\mu^+(M; M_0)$ is tilting if and only if $\text{Hom}_{\text{per}(\Lambda)}(M_0, f)$ is injective where f is the left minimal $\text{add}(M_0)$ -approximation of M/M_0 .

Remark 1.12. We can define dually a right approximation and a right mutation. Moreover, we have a dual version of Theorem 1.11 for right mutations.

We want now to prove the following result on silting mutations for skew group algebras.

Proposition 1.13. *Let Λ be a finite dimensional k -algebra with a G -action. Let M be a silting object in $\text{per}(\Lambda)$ and M_0 be a direct summand of M . We denote by $f : M/M_0 \rightarrow M'_0$ the left minimal $\text{add}(M_0)$ -approximation of M/M_0 . If M_0 , M/M_0 and M'_0 are G -invariant, then there is an isomorphism in $\text{per}(\Lambda G)$*

$$\mu^+(M \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G; M_0 \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G) \simeq \mu^+(M; M_0) \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G$$

Remark 1.14. Let us assume that the hypotheses of the previous proposition hold. Thanks to Theorem 2.10 of [AB22], we conclude that $\mu^+(M \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G; M_0 \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G)$ is tilting in $\text{per}(\Lambda G)$ if $\mu^+(M; M_0)$ is tilting in $\text{per}(\Lambda)$ and G -invariant.

In order to prove the previous proposition, we need to study how the functor $- \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G$ behaves with left minimal morphisms.

Lemma 1.15. *Let Λ be a finite dimensional k -algebra with a G -action and $f : X \rightarrow Y$ be a left minimal morphism in $\text{per}(\Lambda)$. If X and Y are G -invariant, then $f \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G$ is a left minimal morphism in $\text{per}(\Lambda G)$.*

Proof. Let $h : Y \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G \rightarrow Y \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G$ be an endomorphism in $\text{per}(\Lambda G)$ such that $h \circ (f \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G) = f \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G$. We want to prove that h is an isomorphism. In the following, we denote by g_0, \dots, g_{n-1} the elements of the group G where $g_0 = 1_G$. Thanks to Proposition 2.8 in [AB22], h can be written as the sum of $h_{g_i} \otimes g_i$ for $i = 0, \dots, n-1$ in $\text{End}_{\text{per}(\Lambda)}(Y)G$, for some endomorphisms $h_{g_i} : Y \rightarrow Y$ in $\text{per}(\Lambda)$. Similarly, using Proposition 1.8, $f \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G$ can be written as $f \otimes g_0$ in $\text{Hom}_{\text{per}(\Lambda)}(X, Y)G$ and the equality $h \circ (f \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G) = f \overset{\mathbf{L}}{\otimes}_{\Lambda} \Lambda G$ gives us

$$f \otimes g_0 = \left(\sum_{i=0}^{n-1} h_{g_i} \otimes g_i \right) (f \otimes g_0) = \sum_{i=0}^{n-1} h_{g_i} \circ (g_i \cdot f) \otimes g_i$$

By identification, we deduce that $h_{g_0} f = f$ and $h_{g_i} \circ (g_i \cdot f) = 0$ for $i = 1, \dots, n-1$. Let us prove that h admits a right inverse i.e. that there exist $t_{g_0}, \dots, t_{g_{n-1}} \in \text{End}_{\text{per}(\Lambda)}(Y)$ such that

$$\text{id}_{\Lambda} \otimes g_0 = \left(\sum_{i=0}^{n-1} h_{g_i} \otimes g_i \right) \left(\sum_{j=0}^{n-1} t_{g_j} \otimes g_j \right) = \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} h_{g_i g_j^{-1}} \circ (g_i g_j^{-1} \cdot t_{g_j}) \otimes g_i \quad (2)$$

Thanks to the relations obtained by identification in (2), one can check by a descending induction that for all $i = 1, \dots, n-1$, t_{g_i} can be written as the sum of $h'_{g_l} \circ (g_i g_l^{-1} \cdot t_{g_l})$ for $l = 0, \dots, i-1$ for some $h'_{g_l} : Y \rightarrow Y$ in $\text{per}(\Lambda)$ satisfying that $h'_{g_l} \circ (g_i g_l^{-1} \cdot f) = 0$. In particular, by identification in (2), we obtain

$$\text{id}_\Lambda = h_{g_0} t_{g_0} + \sum_{j=1}^{n-1} h_{g_j^{-1}} \circ (g_j^{-1} \cdot t_{g_j}) = h_{g_0} t_{g_0} + h''_{g_0} t_{g_0}$$

for some endomorphism $h''_{g_0} : Y \rightarrow Y$ in $\text{per}(\Lambda)$ satisfying that $h''_{g_0} f = 0$, thanks to the previous induction. Since $(h_{g_0} + h''_{g_0}) \circ f = f$ and f is left minimal, $h_{g_0} + h''_{g_0}$ is invertible and we can set t_{g_0} to be its inverse. Hence, one can use the previous induction to define the t_{g_i} for $i = 1, \dots, n-1$ so that (2) holds. Similarly, one can check that h admits a left inverse which ends the proof. \square

Proof of Proposition 1.13. We assume that M_0 , M/M_0 and M'_0 are G -invariant. Since the functor $- \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G : \text{per}(\Lambda) \rightarrow \text{per}(\Lambda G)$ is triangulated, note that we have

$$(M/M_0) \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G = (M \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G) / (M_0 \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G)$$

Thanks to Lemma 1.15, it remains to prove that $f \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G$ is a left $\text{add}(M_0 \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G)$ -approximation of $(M/M_0) \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G$. Applying the functor $- \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G$ on the triangle induced by f in $\text{per}(\Lambda)$, we obtain the following triangle in $\text{per}(\Lambda G)$

$$(M/M_0) \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G \xrightarrow{f \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G} M'_0 \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G \longrightarrow \text{Cone}(f) \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G \longrightarrow (M/M_0) \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G[1]$$

where $\text{Cone}(f)$ denotes the cone of f . Hence, applying $\text{Hom}_{\text{per}(\Lambda G)}(-, M_0 \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G)$ to the previous triangle, it suffices to prove that

$$H := \text{Hom}_{\text{per}(\Lambda G)}(\text{Cone}(f) \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G, M_0 \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G[1]) = 0$$

For this, note that we have the following isomorphisms of vector spaces

$$\begin{aligned} H &\simeq \text{Hom}_{\text{per}(\Lambda)}(\text{Cone}(f), \text{Res}(M_0 \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda G[1])) \\ &\simeq \text{Hom}_{\text{per}(\Lambda)}(\text{Cone}(f), \oplus_{g \in G} (M_0)^g[1]) \\ &\simeq \text{Hom}_{\text{per}(\Lambda)}(\text{Cone}(f), \oplus_{g \in G} M_0[1]) \end{aligned}$$

where the second isomorphism comes from (1) and the third arises from the fact that M_0 is G -invariant. Since the left mutation of a silting object is silting by Theorem 1.11, we obtain that $H = 0$ which concludes the proof. \square

2. Generalized Kauer moves for Brauer graphs with multiplicity

In this section, we extend the notion of generalized Kauer moves defined for Brauer graphs of multiplicity one in [Sot24] to the case of Brauer graphs with arbitrary multiplicity. We will show that these generalized Kauer moves can be understood in terms of silting mutations. This generalizes Theorem 3.10 in [Sot24] for arbitrary multiplicity.

2.1. Brauer graph algebras with multiplicity

Let us recall the notion of Brauer graph algebras with multiplicity. These are finite dimensional algebras defined thanks to a combinatorial data called a Brauer graph. As in [Sot24], we will use the definition of a Brauer graph using combinatorial maps (see for instance [Laz14]).

Definition 2.1. A *Brauer graph* is the data $\Gamma = (H, \iota, \sigma, m)$ where

- H is the set of half-edges;
- ι is a permutation of H without fixed points satisfying $\iota^2 = \text{id}_H$: it is called the *pairing*;
- σ is a permutation of H called the *orientation*;
- $m : H \rightarrow \mathbb{Z}_{>0}$ is a map that is constant on a σ -orbit : it is called the *multiplicity*.

To the data $\Gamma = (H, \iota, \sigma, m)$, one can naturally define a graph whose vertex set is $\Gamma_0 = H/\sigma$, edge set is $\Gamma_1 = H/\iota$ and source map is the natural projection $s : H \rightarrow H/\sigma$. Moreover, each cycle in the decomposition of σ gives rise to a cyclic ordering of the edges around a vertex. Since $m : H \rightarrow \mathbb{Z}_{>0}$ is constant on the σ -orbits, it induces a map $\tilde{m} : H/\sigma \rightarrow \mathbb{Z}_{>0}$. Hence, we recover the usual definition of a Brauer graph with multiplicity as defined in [Sch18] for instance.


Remark 2.2. In order to simplify the definition of a Brauer graph algebra, we do not take into consideration the following Brauer graph



which corresponds to $H = \{1^+, 1^-\}$, $\iota = (1^+ 1^-)$, $\sigma = \text{id}_H$ and m being identically 1. More precisely, we assume that such graph does not appear in any connected component of a given Brauer graph. Moreover, the previous graph is not really interesting for our purpose since the generalized Kauer move of the edge 1 is trivial.

From now on, we identify $\Gamma = (H, \iota, \sigma, m)$ with its corresponding graph that we have constructed above. By convention, the orientation of a Brauer graph will correspond to the local embedding of each vertex into the counterclockwise oriented plane.

Example 2.3. Let $\Gamma = (H, \iota, \sigma, m)$ be the Brauer graph where the set of half-edges is $H = \{1^+, 1^-, 2^+, 2^-, 3^+, 3^-, 4^+, 4^-\}$, the involution ι is $(1^+ 1^-)(2^+ 2^-)(3^+ 3^-)(4^+ 4^-)$, the permutation σ is $(1^- 4^- 3^- 2^-)(2^+ 3^+)$ and the multiplicity $m : H \rightarrow \mathbb{Z}_{>0}$ is given by $m(1^-) = m(2^-) = m(3^-) = m(4^-) = m(4^+) = 1$ and $m(2^+) = m(3^+) = m(1^+) = 2$. With our previous convention, this Brauer graph can be represented as follows




where the multiplicity of an half-edge is given by the value in the vertex it is incident to. From now on, we will only write on the graph the multiplicities that are strictly greater than one.

Given a Brauer graph $\Gamma = (H, \iota, \sigma, m)$, one can construct a quiver $Q_\Gamma = (Q_0, Q_1)$ as in [Sch18], which is reinterpreted with our definition as follows

- The vertex set Q_0 is the edge set H/ι of Γ . In the following, we denote by $[h]$ the edge in H/ι associated to $h \in H$.
- The arrow set Q_1 is induced by the orientation σ and the multiplicity m of Γ . More precisely, for any half-edge $h \in H$ that is not fixed by σ , there is an arrow α_h from $[h]$ to $[\sigma h]$. Moreover, for any half-edge $h \in H$ that is fixed by σ , there is an arrow α_h from $[h]$ to itself if and only if $m(h) > 1$.

Note that any half-edge $h \in H$ inducing an arrow α_h from $[h]$ to $[\sigma h]$ in Q_Γ gives rise to an oriented cycle C_h in Q_Γ that begins and ends at $[h]$. We call such an oriented cycle a *special h -cycle* and it is of the form $C_h = \alpha_{\sigma^{n-1}h} \dots \alpha_h$, where n is the size of the σ -orbit of h .

Example 2.4. Let us consider $\Gamma = (H, \iota, \sigma, m)$ the Brauer graph defined in Example 2.3. Then, its associated quiver Q_Γ is given by



In this case, the special h -cycles are given by $C_{1+} = \alpha_{1+}$, $C_{1-} = \alpha_{2-}\alpha_{3-}\alpha_{4-}\alpha_{1-}$, $C_{2+} = \alpha_{3+}\alpha_{2+}$, $C_{2-} = \alpha_{3-}\alpha_{4-}\alpha_{1-}\alpha_{2-}$, $C_{3+} = \alpha_{2+}\alpha_{3+}$, $C_{3-} = \alpha_{4-}\alpha_{1-}\alpha_{2-}\alpha_{3-}$ and $C_{4-} = \alpha_{1-}\alpha_{2-}\alpha_{3-}\alpha_{4-}$. Note that there is no special 4^+ -cycle since 4^+ is fixed by σ and is of multiplicity one so it does not induce an arrow in Q_Γ .

Definition 2.5. Let $\Gamma = (H, \iota, \sigma, m)$ be a Brauer graph and Q_Γ be its associated quiver. The *Brauer graph algebra* B_Γ of Γ is the path algebra kQ_Γ/I_Γ where the ideal of relations I_Γ is generated by

$$(I) \quad (C_h)^{m(h)} - (C_{\iota h})^{m(\iota h)}$$

for any half-edge $h \in H$ such that h and ιh both induce an arrow in Q_Γ .

$$(II) \quad \alpha_h(C_h)^{m(h)}$$

for any half-edge $h \in H$ such that h induces an arrow in Q_Γ .

$$(III) \quad \alpha_{\iota\sigma h}\alpha_h$$

for any half-edge $h \in H$ such that h and $\iota\sigma h$ both induce an arrow in Q_Γ .

In general, the relations that generates the ideal of relations I_Γ are not minimal as we can see in the following example.

Example 2.6. Let us consider $\Gamma = (H, \iota, \sigma, m)$ the Brauer graph defined in Example 2.3. Then, its associated Brauer graph algebra B_Γ is defined as follows : its quiver is described in Example 2.4 and its ideal of relations is generated by

- (I) $\alpha_{1+}^2 - \alpha_{2-}\alpha_{3-}\alpha_{4-}\alpha_{1-}, (\alpha_{3+}\alpha_{2+})^2 - \alpha_{3-}\alpha_{4-}\alpha_{1-}\alpha_{2-}, (\alpha_{2+}\alpha_{3+})^2 - \alpha_{4-}\alpha_{1-}\alpha_{2-}\alpha_{3-};$
- (II) $\alpha_{1+}^3, \alpha_{1-}\alpha_{2-}\alpha_{3-}\alpha_{4-}\alpha_{1-}, \alpha_{2-}\alpha_{3-}\alpha_{4-}\alpha_{1-}\alpha_{2-}, \alpha_{3-}\alpha_{4-}\alpha_{1-}\alpha_{2-}\alpha_{3-}, \alpha_{4-}\alpha_{1-}\alpha_{2-}\alpha_{3-}\alpha_{4-}, \alpha_{2+}(\alpha_{3+}\alpha_{2+})^2, \alpha_{3+}(\alpha_{2+}\alpha_{3+})^2;$
- (III) $\alpha_{1+}\alpha_{2-}, \alpha_{1-}\alpha_{1+}, \alpha_{2+}\alpha_{3-}, \alpha_{2-}\alpha_{3+}, \alpha_{3-}\alpha_{2+}, \alpha_{3+}\alpha_{4-}.$

In particular, the relations (II) can be obtained from the relations (I) and (III).

2.2. Covering of Brauer graph algebras with multiplicity

In this subsection, we recall how Brauer graph algebras with multiplicity can be covered by a Brauer graph algebra with multiplicity identically one [Asa19]. We begin with defining the notion of morphisms of Brauer graphs.

Definition 2.7. Let $\Gamma = (H, \iota, \sigma, m)$ and $\Gamma' = (H', \iota', \sigma', m')$ be Brauer graphs. A *morphism of Brauer graphs* $p : \Gamma \rightarrow \Gamma'$ is a map $p : H \rightarrow H'$ that commutes with the pairings and the orientations i.e. $p \circ \iota = \iota' \circ p$ and $p \circ \sigma = \sigma' \circ p$.

To define this covering, we will need to introduce a particular grading on the Brauer graph with multiplicity called an *admissible* grading.

Definition 2.8. Let $\Gamma = (H, \iota, \sigma, m)$ be a Brauer graph and \bar{m} be the least common multiple of the $m(h)$ for $h \in H$.

- We say that a $\mathbb{Z}/\bar{m}\mathbb{Z}$ -grading $d : H \rightarrow \mathbb{Z}/\bar{m}\mathbb{Z}$ is *admissible* if

$$\sum_{h \in H, s(h)=v} d(h) = \frac{\bar{m}}{\tilde{m}(v)}$$

for all $v \in H/\sigma$, where $s : H \rightarrow H/\sigma$ is the source map of Γ and $\tilde{m} : H/\sigma \rightarrow \mathbb{Z}_{>0}$ is the map induced by m on H/σ (cf after Definition 2.1).

- We say that (Γ, d) is a *$\mathbb{Z}/\bar{m}\mathbb{Z}$ -graded Brauer graph* if $d : H \rightarrow \mathbb{Z}/\bar{m}\mathbb{Z}$ is an admissible $\mathbb{Z}/\bar{m}\mathbb{Z}$ -grading.

Let $\Gamma = (H, \iota, \sigma, m)$ be a Brauer graph. Denoting by \bar{m} the least common multiple of the $m(h)$ for $h \in H$, we equip Γ with an admissible $\mathbb{Z}/\bar{m}\mathbb{Z}$ -grading $d : H \rightarrow \mathbb{Z}/\bar{m}\mathbb{Z}$. One can construct a new Brauer graph $\Gamma_d = (H_d, \iota_d, \sigma_d)$ of multiplicity identically one as follows [Asa19]


- $H_d = H \times (\mathbb{Z}/\bar{m}\mathbb{Z})$: an element of H_d will be denoted h_i for $h \in H$ and $i \in \mathbb{Z}/\bar{m}\mathbb{Z}$;
- For all $h_i \in H_d$, $\iota_d(h_i) = (\iota h)_i$;

- For all $h_i \in H_d$, $\sigma_d(h_i) = (\sigma h)_{i+d(h)}$.


It is clear that the projection $p_\Gamma : H_d \rightarrow H$ defines a morphism of Brauer graphs from Γ_d to Γ . From now on, Γ_d will be called the *Galois covering* of Γ . This terminology is motivated by the following result.

Proposition 2.9 (Asashiba [Asa19, Proposition 1.17 and Theorem 2.11]). *Let (Q_Γ, I_Γ) and (Q_d, I_d) the bound quivers corresponding to Γ and Γ_d respectively. Then the morphism of bound quivers $F : (Q_d, I_d) \rightarrow (Q_\Gamma, I_\Gamma)$ arising from the morphism of Brauer graphs $p : \Gamma_d \rightarrow \Gamma$ defined previously is a Galois covering with group $Z/\overline{m}\mathbb{Z}$.*

Example 2.10. Let $\Gamma = (H, \iota, \sigma, m)$ be the Brauer graph defined in Example 2.3. We equip Γ with an admissible $\mathbb{Z}/2\mathbb{Z}$ -grading $d : H \rightarrow \mathbb{Z}/2\mathbb{Z}$ defined by $d(1^+) = d(3^+) = 1$ and $d(1^-) = d(2^-) = d(2^+) = d(3^-) = d(4^-) = d(4^+) = 0$ which can be represented on the graph as follows



Then the Galois covering $\Gamma_d = (H_d, \iota_d, \sigma_d)$ of the Brauer graph Γ equipped with the admissible $\mathbb{Z}/2\mathbb{Z}$ -grading $d : H \rightarrow \mathbb{Z}/2\mathbb{Z}$ is given by



In what follows, we will prove that the Brauer graph algebra B associated to Γ can be recovered from its Galois covering with the cyclic group $G = (\mathbb{C}_{\overline{m}}, \cdot)$ of order \overline{m} thanks to the construction of the quiver and relations of a skew group algebra for a cyclic group [RR85, Section 2.3]. From now on, we assume that \overline{m} is invertible in k and we fix g a generator of the cyclic group G . Denoting by $e_{[h_i]}$ the idempotent in B_d corresponding to the edge $[h_i] \in H_d/\iota_d$, there is a natural action of G on $B_d = kQ_d/I_d$ given by

- $g \cdot e_{[h_i]} = e_{[h_{i+1}]}$ for all $[h_i] \in H_d/\iota_d$;
- $g \cdot \alpha_{h_i} = \alpha_{h_{i+1}}$ where α_{h_i} is the arrow in Q_d induced by the half-edge $h_i \in H_d$.

Similarly, the dual \widehat{G} of G is also a cyclic group. Let χ be a generator of \widehat{G} . Denoting by $e_{[h]}$ the idempotent in B corresponding to the edge $[h] \in H/\iota$, there is a natural action of \widehat{G} on $B = kQ/I$ given by

- $\chi \cdot e_{[h]} = e_{[h]}$ for all $[h] \in H/\iota$;
- $\chi \cdot \alpha_h = \chi(g)^{-d(h)} \alpha_h$ where α_h is the arrow in Q induced by the half-edge $h \in H$;

Note that the G -action on B_d arises from a G -action on Q_d which preserves the ideal of relations I_d whereas the \widehat{G} -action on B does not come from an action on its quiver.

Proposition 2.11. *Let f be the idempotent in $B_d G$ given by the sum of the $e_{[h_0]} \otimes 1_G$ for all $[h] \in H/\iota$. Then, the algebra $fB_d Gf$ has a natural \widehat{G} -action and the map*

$$\begin{aligned} \phi: \quad B &\longrightarrow fB_d Gf \\ e_{[h]} &\longmapsto e_{[h_0]} \otimes 1_G \\ \alpha_h &\longmapsto \beta_h := \alpha_{h_{-d(h)}} \otimes g^{-d(h)} \end{aligned}$$

is an isomorphism of algebras commuting with the \widehat{G} -actions.

Proof. Since there is a G -action on B_d , one can naturally construct a \widehat{G} -action on $B_d G$ as explained after Definition 1.1. Moreover, if χ denotes a generator of the cyclic group \widehat{G} , note that $\chi \cdot f = f$. Hence, the \widehat{G} -action on $B_d G$ extends onto a \widehat{G} -action on $fB_d Gf$. Furthermore, using the construction in [RR85], the quiver and relations of $fB_d Gf$ are given as follows

- The vertices of the quiver of $fB_d Gf$ are in bijection with the idempotents $e_{[h_0]} \otimes 1_G$ for all $[h] \in H/\iota$. Moreover, the arrows are given by the $\beta_h := \alpha_{h_{-d(h)}} \otimes g^{-d(h)}$ for all $h \in H$.
- The relations of $fB_d Gf$ are determined thanks to a representative in the G -orbit of the relations of B_d . Hence, its ideal of relations is generated by

$$(I') \quad (\beta_{\sigma^{n-1}h} \dots \beta_h)^{m(h)} - (\beta_{\sigma^{n'-1}\iota h} \dots \beta_{\iota h})^{m(\iota h)}$$

for all $h \in H$ such that $h_{-d(h)}$ and $(\iota h)_{-d(\iota h)}$ both induce an arrow in Q_d , where n and n' are the size of the σ -orbit of h and ιh respectively.

$$(II') \quad \beta_h (\beta_{\sigma^{n-1}h} \dots \beta_h)^{m(h)}$$

for all $h \in H$ such that $h_{-d(h)}$ induces an arrow in Q_d , where n is the size of the σ -orbit of h .

$$(III') \quad \beta_{\iota\sigma h} \beta_h$$

for all $h \in H$ such that $h_{-d(h)}$ and $(\iota\sigma h)_{-d(\iota\sigma h)}$ both induce an arrow in Q_d .

Let us detail how the relations (I') are obtained. For any generator R of I_d , the ideal of relations of B_d , there exists a unique relation R' in the G -orbit of R which is of the form $R' : g^t(e_{[h_0]}) \rightarrow e_{[h'_0]}$ for some $0 \leq t < \overline{m}$. In this case, the ideal of relations of fB_dGf is generated by elements of the form $f(R' \otimes g^t)f$. We recall that the type (I) relation generating I_d is of the form

$$R = \alpha_{\sigma_d^{n_d-1}h_i} \dots \alpha_{h_i} - \alpha_{\sigma_d^{n'_d-1}\iota_d h_i} \dots \alpha_{\iota_d h_i}$$

for all $h_i \in H_d$ such that h_i and $\iota_d h_i$ both induce an arrow in Q_d , where n_d and n'_d denotes the size of the σ_d -orbit of h_i and $\iota_d h_i$ respectively. By construction of Γ_d , $n_d = nm(h)$ and $n'_d = n'm(\iota h)$ where n and n' denotes the size of the σ -orbit of h and ιh respectively. Moreover, since $d : H \rightarrow \mathbb{Z}/\overline{m}\mathbb{Z}$ is admissible, the unique representative R' in the G -orbit of R of the form $R' : g^t(e_{[h_0]}) \rightarrow e_{[h'_0]}$ for some $0 \leq t < \overline{m}$ is

$$R' = \alpha_{\sigma_d^{nm(h)-1}h_0} \dots \alpha_{h_0} - \alpha_{\sigma_d^{n'm(\iota h)-1}\iota_d h_0} \dots \alpha_{\iota_d h_0}$$

where $t = 0$ in this case. Using the definition of σ_d , note that

$$\beta_{\sigma^{n-1}h} \dots \beta_h = \alpha_{\sigma_d^{n-1}h_{-d(\sigma^{n-1}h)}} \dots \alpha_{h_{-\sum_{i=0}^{n-1} d(\sigma^i h)}} \otimes g^{-\sum_{i=0}^{n-1} d(\sigma^i h)}$$

Hence, one can check that $f(R' \otimes 1_G)f$ is indeed given by the relations (I'). The relations (II') and (III') are obtained similarly from the type (II) and type (III) relation in B_d respectively.

It is clear that ϕ defines an isomorphism between the quivers of B and fB_dGf which clearly extends on an isomorphism of algebras between the path algebras of these quivers. Moreover, since h_i induces an arrow in Q_d for $i \in \mathbb{Z}/\overline{m}\mathbb{Z}$ if and only if h induces an arrow in Q_Γ , it is clear that the ideals of relations of B and fB_dGf coincide via the previous isomorphism of path algebras. Thus, ϕ defines an isomorphism of algebras between B and fB_dGf . Moreover, one can easily check that this isomorphism commutes with the \widehat{G} -action on these two algebras. \square

2.3. Generalized Kauer moves

Our goal is to define generalized Kauer moves for Brauer graph with arbitrary multiplicity so that they can be understood in terms of silting mutations as for the multiplicity one case [Sot24, Theorem 3.10]. The idea of the proof is to use the Galois covering defined in the previous subsection which can be constructed whenever the Brauer graph is equipped with an admissible grading. Hence, we need to define a graded version of the generalized Kauer moves for Brauer graph with multiplicity. As in [Sot24], we begin with defining these for successive half-edges called *sectors*.

Definition 2.12. Let $\Gamma = (H, \iota, \sigma, m)$ be a Brauer graph and $H' \subset H$ stable under ι .

- We say that $(h, r) \in H \times \mathbb{Z}_{\geq 0}$ is a *sector* in Γ of elements in H' if $r + 1$ is the smallest integer $r' \geq 0$ such that $\sigma^{r'}h \notin H'$.
- We say that $(h, r) \in H \times \mathbb{Z}_{\geq 0}$ is a *maximal sector* in Γ of elements in H' if it is a sector satisfying that $\sigma^{-1}h \notin H'$.

We denote respectively by $\text{sect}(H', \sigma)$ and $\text{Sect}(H', \sigma)$ the set of sectors and maximal sectors in Γ of elements in H' .

From now on, if $m : H \rightarrow \mathbb{Z}_{>0}$ is the multiplicity of a Brauer graph $\Gamma = (H, \iota, \sigma, m)$, we denote by \bar{m} the least common multiple of the $m(h)$ for $h \in H$.

Definition 2.13. Let $(\Gamma, d) = (H, \iota, \sigma, m, d)$ be a $\mathbb{Z}/\bar{m}\mathbb{Z}$ -graded Brauer graph and H' be a subset of H stable under ι . The $\mathbb{Z}/\bar{m}\mathbb{Z}$ -graded *generalized Kauer move* of a sector $(h, r) \in \text{sect}(H', \sigma)$ in Γ gives rise to a $\mathbb{Z}/\bar{m}\mathbb{Z}$ -Brauer graph $\mu_{(h,r)}^+(\Gamma, d) = (H, \iota, \sigma_{(h,r)}, m_{(h,r)}, d_{(h,r)})$ where

$$\begin{aligned} \sigma_{(h,r)} &= (h \sigma^{r+1}h)\sigma(\sigma^r h \iota \sigma^{r+1}h) & \text{and} & & m_{(h,r)} : H &\rightarrow \mathbb{Z}_{>0} \\ & & & & \sigma^i h &\mapsto m(\iota \sigma^{r+1}h) \quad \text{for } i = 0, \dots, r \\ & & & & h' &\mapsto m(h') \quad \text{for } h' \neq \sigma^i h \end{aligned}$$

and where the grading $d_{(h,r)} : H \rightarrow \mathbb{Z}/\bar{m}\mathbb{Z}$ is defined by

$$\begin{aligned} d_{(h,r)} : \iota \sigma^{r+1}h &\mapsto -\sum_{i=0}^r d(\sigma^i h) \\ \sigma^r h &\mapsto \begin{cases} d(\iota \sigma^{r+1}h) + d(\sigma^r h) & \text{if } \iota \sigma^{r+1}h \neq \sigma^{-1}h \\ \sum_{i=-1}^r d(\sigma^i h) + d(\sigma^r h) & \text{else} \end{cases} \\ \sigma^{-1}h &\mapsto \begin{cases} \sum_{i=-1}^r d(\sigma^i h) & \text{if } \iota \sigma^{r+1}h \neq \sigma^{-1}h \\ -\sum_{i=0}^r d(\sigma^i h) & \text{else} \end{cases} \\ h' &\mapsto d(h') \quad \text{for } h' \neq \iota \sigma^{r+1}h, \sigma^r h, \sigma^{-1}h \end{aligned}$$

The underlying Brauer graph $\mu_{(h,r)}^+(\Gamma) = (H, \iota, \sigma_{(h,r)}, m_{(h,r)})$ can be obtained from Γ as follows.





Figure 1: Generalized Kauer move of a sector (h, r)

Remark 2.14. In the special case where $\iota \sigma^{r+1}h = \sigma^{-1}h$, then the $\mathbb{Z}/\bar{m}\mathbb{Z}$ -graded generalized Kauer move of the sector (h, r) does not change the underlying Brauer graph which is given as follows



$$\Gamma = \mu_{(h,r)}^+(\Gamma)$$

However, the degree of the $\sigma^i h$ will a priori change in the process. In particular, the Galois covering of Γ and $\mu_{(h,r)}^+(\Gamma)$ could have different orientation, even in this case.

By construction of the generalized Kauer move described in the previous picture, it is clear that $m_{(h,r)}$ is constant on the $\sigma_{(h,r)}$ -orbits. Moreover, $d_{(h,r)} : H \rightarrow \mathbb{Z}/\overline{m}\mathbb{Z}$ is admissible. Indeed, one can construct a bijection $\phi : H/\sigma \rightarrow H/\sigma_{(h,r)}$ satisfying

$$\begin{aligned} s_{(h,r)}(\sigma^i h) &= \phi(s(\iota\sigma^{r+1}h)) & \text{for } i = 0, \dots, r \\ s_{(h,r)}(h') &= \phi(s(h')) & \text{for } h' \neq \sigma^i h, i = 0, \dots, r \end{aligned}$$

where $s : H \rightarrow H/\sigma$ and $s_{(h,r)} : H \rightarrow H/\sigma_{(h,r)}$ are the source map of Γ and $\mu_{(h,r)}^+(\Gamma)$ respectively. Then, one can easily check that, for all vertex $v \in H/\sigma_{(h,r)}$ in $\mu_{(h,r)}^+(\Gamma)$, we have

$$\sum_{h' \in H, s_{(h,r)}(h')=v} d_{(h,r)}(h') = \sum_{h' \in H, s(h')=\phi^{-1}(v)} d(h') = \frac{\overline{m}}{\tilde{m}(\phi^{-1}(v))}$$

where $\tilde{m} : H/\sigma \rightarrow \mathbb{Z}_{>0}$ is induced by m (cf after Definition 2.1). Note that $\overline{m_{(h,r)}} = \overline{m}$. Moreover, one can easily check that $\tilde{m}(\phi^{-1}(v)) = \tilde{m}_{(h,r)}(v)$ for all $v \in H/\sigma_{(h,r)}$ where $\tilde{m}_{(h,r)} : H/\sigma_{(h,r)} \rightarrow \mathbb{Z}_{>0}$ is induced by $m_{(h,r)}$. Hence, $d_{(h,r)} : H \rightarrow \mathbb{Z}/\overline{m}\mathbb{Z}$ is indeed admissible.

Remark 2.15. By definition, if Γ is a Brauer graph of multiplicity identically one then $\mu_{(h,r)}^+(\Gamma)$ is also a Brauer graph of multiplicity identically one. Hence, this definition generalizes Definition 2.2 in [Sot24].

Thanks to Lemma 2.4 of [Sot24], we can consider successive $\mathbb{Z}/\overline{m}\mathbb{Z}$ -graded generalized Kauer moves. We denote by

$$\mu_{(h_1,r_1)(h_2,r_2)}^+(\Gamma, d) = (H, \iota, \sigma_{(h_1,r_1)(h_2,r_2)}, m_{(h_1,r_1)(h_2,r_2)}, d_{(h_1,r_1)(h_2,r_2)})$$

the $\mathbb{Z}/\overline{m}\mathbb{Z}$ -graded Brauer graph defined by $\mu_{(h_1,r_1)}^+(\mu_{(h_2,r_2)}^+(\Gamma, d))$.

Proposition 2.16. *Let $\Gamma = (H, \iota, \sigma, m)$ be a Brauer graph and H' be a subset of H stable under ι . Let $(h_1, r_1), (h_2, r_2)$ be two distinct maximal sectors in Γ of elements in H' . For any admissible $\mathbb{Z}/\overline{m}\mathbb{Z}$ -grading $d : H \rightarrow \mathbb{Z}/\overline{m}\mathbb{Z}$ of Γ , we have*

$$\mu_{(h_1,r_1)(h_2,r_2)}^+(\Gamma, d) = \mu_{(h_2,r_2)(h_1,r_1)}^+(\Gamma, d)$$

Proof. Thanks to Proposition 2.5 in [Sot24], it only remains to prove that $m_{(h_1, r_1)(h_2, r_2)} = m_{(h_2, r_2)(h_1, r_1)}$. It is clear that this equality holds for h' distinct from the $\sigma^i h_1$ for $i = 0, \dots, r_1$ and from the $\sigma^i h_2$ for $i = 0, \dots, r_2$. Since h_1 and h_2 have a symmetric role, it suffices to check the equality for the $\sigma^i h_1$, $i = 0, \dots, r_1$. On the one hand, we have

$$m_{(h_1, r_1)(h_2, r_2)}(\sigma^i h_1) = m_{(h_1, r_1)(h_2, r_2)}(\sigma_{(h_2, r_2)}^i h_1) = m_{(h_2, r_2)}(\iota \sigma_{(h_2, r_2)}^{r_1+1} h_1) = m(\iota \sigma^{r_1+1} h_1)$$


On the other hand, since $\sigma^i h_1$ is distinct from $\sigma_{(h_1, r_1)}^j h_2 = \sigma^j h_2$ for $j = 0, \dots, r_2$, we have

$$m_{(h_2, r_2)(h_1, r_1)}(\sigma^i h_1) = m_{(h_1, r_1)}(\sigma^i h_1) = m(\iota \sigma^{r_1+1} h_1)$$

□

Definition 2.17. Let $(\Gamma, d) = (H, \iota, \sigma, m, d)$ be a $\mathbb{Z}/\overline{m}\mathbb{Z}$ -Brauer graph and H' be a subset of H stable under ι . The $\mathbb{Z}/\overline{m}\mathbb{Z}$ -graded generalized Kauer move of H' is the succession of the $\mathbb{Z}/\overline{m}\mathbb{Z}$ -graded generalized Kauer move of all the maximal sectors $(h, r) \in \text{Sect}(H', \sigma)$ in Γ . This gives rise to a $\mathbb{Z}/\overline{m}\mathbb{Z}$ -graded Brauer graph that will be denoted by $\mu_{H'}^+(\Gamma, d) = (H, \iota, \sigma_{H'}, m_{H'}, d_{H'})$.

Example 2.18. Let us consider $\Gamma = (H, \iota, \sigma, m, d)$ the $\mathbb{Z}/2\mathbb{Z}$ -graded Brauer graph defined in Example 2.10. Let $H' = \{1^+, 1^-, 2^+, 2^-\}$ be a subset of H stable under ι . Then, the $\mathbb{Z}/2\mathbb{Z}$ -graded Brauer graph $\mu_{H'}^+(\Gamma, d) = (H, \iota, \sigma_{H'}, m_{H'}, d_{H'})$ obtained from the $\mathbb{Z}/2\mathbb{Z}$ -graded generalized Kauer move of H' is given by



where the orientation $\sigma_{H'}$ is $(1^- 4^+ 2^-)(2^+ 4^- 3^-) = (2^- 4^-)(2^+ 3^+)\sigma(1^- 4^+)(2^+ 3^-)$, the multiplicity $m_{H'} : H \rightarrow \mathbb{Z}_{>0}$ is given by $m_{H'}(1^+) = m_{H'}(3^+) = 2$ and $m_{H'}(1^-) = m_{H'}(2^-) = m_{H'}(4^+) = m_{H'}(2^+) = m_{H'}(4^-) = m_{H'}(3^-) = 1$ and the admissible $\mathbb{Z}/2\mathbb{Z}$ -grading $d_{H'} : H \rightarrow \mathbb{Z}/2\mathbb{Z}$ is given by $d_{H'}(1^+) = d_{H'}(3^+) = 1$ and $d_{H'}(1^-) = d_{H'}(2^-) = d_{H'}(4^+) = d_{H'}(2^+) = d_{H'}(4^-) = d_{H'}(3^-) = 0$.

Considering a $\mathbb{Z}/\overline{m}\mathbb{Z}$ -graded generalized Kauer move of a $\mathbb{Z}/\overline{m}\mathbb{Z}$ -graded Brauer graph (Γ, d) , one can construct a Galois covering with group $\mathbb{Z}/\overline{m}\mathbb{Z}$ for $\mu_{H'}^+(\Gamma, d)$ as defined before Proposition 2.9. The following proposition shows a commutativity between constructing this Galois covering and applying a generalized Kauer move.

Proposition 2.19. Let $(\Gamma, d) = (H, \iota, \sigma, m, d)$ be a $\mathbb{Z}/\overline{m}\mathbb{Z}$ -graded Brauer graph and H' be a subset of H stable under ι . We denote by $\Gamma_d = (H_d, \iota_d, \sigma_d)$ the Galois covering of Γ constructed before Proposition 2.9 and $H'_d = H' \times \mathbb{Z}/\overline{m}\mathbb{Z} \subset H_d$. Then, the Galois covering of $\mu_{H'}^+(\Gamma, d)$ is the Brauer graph $\mu_{H'_d}^+(\Gamma_d)$.

The previous proposition can be summarized in the following commutative diagram

$$\begin{array}{ccc}
\Gamma_d & \xrightarrow[\text{move of } H'_d]{\text{generalized Kauer}} & \mu_{H'_d}^+(\Gamma_d) \\
p_\Gamma \downarrow & & \downarrow p_{\mu_{H'}^+(\Gamma)} \\
(\Gamma, d) & \xrightarrow[\text{Kauer move of } H']{\mathbb{Z}/\overline{m}\mathbb{Z}\text{-graded generalized}} & \mu_{H'}^+(\Gamma, d)
\end{array}$$

where $p_\Gamma : \Gamma_d \rightarrow \Gamma$ and $p_{\mu_{H'}^+(\Gamma)} : \mu_{H'_d}^+(\Gamma_d) \rightarrow \mu_{H'}^+(\Gamma)$ are the morphisms of Brauer graphs induced by the natural projection and $\mu_{H'}^+(\Gamma)$ denotes the underlying Brauer graph of $\mu_{H'}^+(\Gamma, d)$.

Proof. It is easy to see that H'_d is a subset of H_d stable under ι_d and that $(h, r) \in \text{Sect}(H', \sigma)$ is a maximal sector in Γ if and only if $(h_i, r) \in \text{Sect}(H'_d, \sigma_d)$ is a maximal sector in Γ_d for all $i \in \mathbb{Z}/\overline{m}\mathbb{Z}$. Thanks to Proposition 2.16, it suffices to prove that the Galois covering of $\mu_{(h,r)}^+(\Gamma, d)$ is given by

$$\mu_{p_\Gamma^{-1}(h,r)}^+(\Gamma_d) = (H_d, \iota_d, (\sigma_d)_{p_\Gamma^{-1}(h,r)})$$

for any maximal sector $(h, r) \in \text{Sect}(H', \sigma)$, where $p_\Gamma^{-1}(h, r)$ is the product of the maximal sectors (h_i, r) for $i \in \mathbb{Z}/\overline{m}\mathbb{Z}$. We denote by

$$\mu_{(h,r)}^+(\Gamma)_{d_{(h,r)}} = (H_{d_{(h,r)}}, \iota_{d_{(h,r)}}, (\sigma_{(h,r)})_{d_{(h,r)}})$$

the Galois covering of $\mu_{(h,r)}^+(\Gamma, d)$. Since $\overline{m_{(h,r)}} = \overline{m}$, the Brauer graphs $\mu_{p_\Gamma^{-1}(h,r)}^+(\Gamma_d)$ and $\mu_{(h,r)}^+(\Gamma)_{d_{(h,r)}}$ have the same set of half-edges and the same pairings. It remains to prove the equality of the orientations. By definition, we have

$$(\sigma_d)_{p_\Gamma^{-1}(h,r)} = \left(\prod_{i \in \mathbb{Z}/\overline{m}\mathbb{Z}} (h_i \quad \sigma_d^{r+1} h_i) \right) \sigma_d \left(\prod_{i \in \mathbb{Z}/\overline{m}\mathbb{Z}} (\sigma_d^r h_i \quad \iota_d \sigma_d^{r+1} h_i) \right)$$

$$(\sigma_{(h,r)})_{d_{(h,r)}} = [(h \quad \sigma^{r+1} h) \sigma (\sigma^r h \quad \iota \sigma^{r+1} h)]_{d_{(h,r)}}$$

It is clear that for any half-edge distinct from the $\sigma_d^{-1} h_i$, $\iota_d \sigma_d^{r+1} h_i$ and $\sigma_d^r h_i$, $i \in \mathbb{Z}/\overline{m}\mathbb{Z}$, the equality holds. We only detail the computations for $\sigma_d^r h_i$, the other cases being similar. Then,

$$\begin{aligned}
(\sigma_d)_{p_\Gamma^{-1}(h,r)}(\sigma_d^r h_i) &= \begin{cases} \sigma_d^{r+1} h_j & \text{if } \sigma_d \iota_d \sigma_d^{r+1} h_i = h_j \text{ for some } j \in \mathbb{Z}/\overline{m}\mathbb{Z} \\ \sigma_d \iota_d \sigma_d^{r+1} h_i & \text{else} \end{cases} \\
&= \begin{cases} (\sigma^{r+1} h)_{i+2 \sum_{k=0}^r d(\sigma^k h) + d(\iota \sigma^{r+1} h)} & \text{if } \sigma \iota \sigma^{r+1} h = h \\ (\sigma \iota \sigma^{r+1} h)_{i+\sum_{k=0}^r d(\sigma^k h) + d(\iota \sigma^{r+1} h)} & \text{else} \end{cases}
\end{aligned}$$

since the first condition is equivalent to $\sigma \iota \sigma^{r+1} h = h$ and in this case $j = i + \sum_{k=0}^r d(\sigma^k h) + d(\iota \sigma^{r+1} h)$. On the other hand, we have

$$\begin{aligned}
(\sigma_{(h,r)} d_{(h,r)} (\sigma_d^r h_i)) &= (\sigma_{(h,r)} d_{(h,r)} ((\sigma^r h)_{i+\sum_{k=0}^{r-1} d(\sigma^k h)})) \\
&= (\sigma_{(h,r)} (\sigma^r h))_{i+\sum_{k=0}^{r-1} d(\sigma^k h)+d_{(h,r)}(\sigma^r h)} \\
&= \begin{cases} (\sigma^{r+1} h)_{i+\sum_{k=0}^{r-1} d(\sigma^k h)+\sum_{i=-1}^r d(\sigma^k h)+d(\sigma^r h)} & \text{if } \sigma \iota \sigma^{r+1} h = h \\ (\sigma \iota \sigma^{r+1} h)_{i+\sum_{k=0}^{r-1} d(\sigma^k h)+d(\iota \sigma^{r+1} h)+d(\sigma^r h)} & \text{else} \end{cases} \\
&= (\sigma_d)_{p_{\Gamma}^{-1}(h,r)} (\sigma_d^r h_i)
\end{aligned}$$

□

2.4. Compatibility with silting mutations

The goal of this part is to prove the following theorem which is an analogous version of Theorem 3.10 in [Sot24] for the case of Brauer graphs with arbitrary multiplicity.

Theorem 2.20. *Let $\Gamma = (H, \iota, \sigma, m)$ be a Brauer graph and H' be a subset of H stable under ι . We assume that \bar{m} is invertible in k . Denoting by B and B' the Brauer graph algebras associated respectively to Γ and $\mu_{H'}^+(\Gamma)$, there is an equivalence of triangulated categories*

$$\text{per}(B') \xrightarrow{-\mathbb{L}_{B'} \mu^+(B; e_{H''} B)} \text{per}(B)$$

where $e_{H''} B$ is the projective B -module corresponding to the edges in $(H \setminus H')/\iota$.

To prove this theorem, we will need the following result.

Proposition 2.21. *Let us consider the setting of Proposition 2.19. We assume that \bar{m} is invertible in k . We denote by B_d and B'_d the Brauer graph algebras associated to Γ_d and $\mu_{H'_d}^+(\Gamma_d)$ respectively. Moreover, let G be the cyclic group $(\mathbb{C}_{\bar{m}}, \cdot)$ of order \bar{m} . Then, there is an equivalence of triangulated categories*

$$\text{per}(B'_d G) \xrightarrow{-\mathbb{L}_{B'_d G} \mu^+(B_d G; e_{H''_d} B_d G)} \text{per}(B_d G)$$

where $e_{H''_d} B_d$ denotes the projective B_d -module corresponding to the edges in $(H_d \setminus H'_d)/\iota_d$.

Proof. By Theorem 3.10 in [Sot24], we know that there is a triangle equivalence

$$\text{per}(B'_d) \xrightarrow{-\mathbb{L}_{B'_d} T_d} \text{per}(B_d)$$

where $T_d := \mu^+(B_d; e_{H''_d} B_d)$. Let us prove that T_d is G -invariant. In this case, we obtain the following equivalence of triangulated categories thanks to Theorem 2.10 in [AB22]

$$\text{per}(B'_d G) \xrightarrow{-\mathbb{L}_{B'_d G} (T_d \otimes_{B_d} B_d G)} \text{per}(B_d G)$$

For all $h_i \in H'_d$, we denote by

$$\alpha(h_i, H'_d) : e_{[h_i]}B_d \longrightarrow e_{[\sigma_d^{r(h_i)+1}h_i]}B_d$$

the morphism induced by the path in B_d from h_i to $\sigma_d^{r(h_i)+1}h_i$ where $r(h_i) + 1 = \min\{r \geq 0 \mid \sigma_d^r h_i \notin H'_d\}$. By an abuse of notation, we set $e_{[\sigma_d^{r(h_i)+1}h_i]}B_d = 0$ if the σ_d -orbit of h_i is contained in H'_d . Then, the left minimal $\text{add}(e_{H'_d}B_d)$ -approximation of $e_{[h_i]}B_d$ is given by

$$\alpha_{[h_i]} : e_{[h_i]}B_d \xrightarrow{\begin{pmatrix} \alpha(h_i, H'_d) \\ \alpha(\iota_d h_i, H'_d) \end{pmatrix}} e_{[\sigma_d^{r(h_i)+1}h_i]}B_d \oplus e_{[\sigma_d^{r(\iota_d h_i)+1}\iota_d h_i]}B_d$$

The two paths in B_d inducing this approximation can be represented in Γ_d as follows




Figure 2: Left minimal $\text{add}(e_{H'_d} B_d)$ -approximation of $e_{[h_i]}B_d$

By definition of the left mutation, we have

$$\mu^+(B_d; e_{H'_d}B_d) = e_{H'_d}B_d \oplus \bigoplus_{[h_i] \in H'_d/\iota_d} \text{Cone}(\alpha_{[h_i]}) = e_{H'_d}B_d \oplus \bigoplus_{[h] \in H'/\iota} \text{Cone}(\bigoplus_{i \in \mathbb{Z}/\overline{m}\mathbb{Z}} \alpha_{[h_i]})$$

Since G acts bijectively on the set of edges $(H_d \setminus H'_d)/\iota_d$, the object $e_{H'_d}B_d$ is G -invariant by Example 1.6 (1). Moreover, for all $[h] \in H'/\iota$,

$$\bigoplus_{i \in \mathbb{Z}/\overline{m}\mathbb{Z}} \alpha_{[h_i]} : X_{[h]} := \bigoplus_{i \in \mathbb{Z}/\overline{m}\mathbb{Z}} e_{[h_i]}B_d \longrightarrow Y_{[h]} := \bigoplus_{i \in \mathbb{Z}/\overline{m}\mathbb{Z}} (e_{[\sigma_d^{r(h_i)+1}h_i]}B_d \oplus e_{[\sigma_d^{r(\iota_d h_i)+1}\iota_d h_i]}B_d)$$

is a morphism in $D^b(B_d)$ between two G -invariant objects. Indeed, this follows again from Example 1.6 (1) and the isomorphisms $\iota_g^{X_{[h]}}$ and $\iota_g^{Y_{[h]}}$ (cf Definition 1.5) are respectively given by the action of g on $X_{[h]}$ and $Y_{[h]}$. Moreover, using the definition of $\alpha_{[h_i]}$, it is not hard to check that the following diagram commutes for all $g \in G$

$$\begin{array}{ccc} X_{[h]}^{g^{-1}} & \xrightarrow{\left(\bigoplus_{i \in \mathbb{Z}/\overline{m}\mathbb{Z}} \alpha_{[h_i]}\right)^{g^{-1}}} & Y_{[h]}^{g^{-1}} \\ \downarrow \iota_g^{X_{[h]}} & & \downarrow \iota_g^{Y_{[h]}} \\ X_{[h]} & \xrightarrow{\bigoplus_{i \in \mathbb{Z}/\overline{m}\mathbb{Z}} \alpha_{[h_i]}} & Y_{[h]} \end{array}$$

By Example 1.6 (2), we deduce that $\text{Cone}(\bigoplus_{i \in \mathbb{Z}/\overline{m}\mathbb{Z}} \alpha_{[h_i]})$ is G -invariant for all $[h] \in H'/\iota$. Hence, we conclude that $\mu^+(B_d; e_{H'_d} B_d)$ is indeed G -invariant.

To complete the proof, it remains to show that

$$T_d G := \mu^+(B_d G; e_{H'_d} B_d G)$$


is isomorphic to $T_d \overset{\mathbf{L}}{\otimes}_{B_d} B_d G$ in $\text{per}(B_d G)$. For this, it suffices to check that the assumptions of Proposition 1.13 hold. We have seen that $e_{H'_d} B_d$ is G -invariant. One can check with a similar argument that $e_{H'_d} B_d$, the projective B_d -module corresponding to the edges in H'_d/ι_d , is also G -invariant. Moreover, the left minimal $\text{add}(e_{H'_d} B_d)$ -approximation of $e_{H'_d} B_d$ is of the form

$$\bigoplus_{[h] \in H'/\iota} \bigoplus_{i \in \mathbb{Z}/\overline{m}\mathbb{Z}} \alpha_{[h_i]} : e_{H'_d} B_d = \bigoplus_{[h] \in H'/\iota} X_{[h]} \longrightarrow \bigoplus_{[h] \in H'/\iota} Y_{[h]}$$

We have seen previously that $Y_{[h]}$ is G -invariant for all $[h] \in H'/\iota$. Hence $\bigoplus_{[h] \in H'/\iota} Y_{[h]}$ is also G -invariant and this concludes the proof. \square

Proof of Theorem 2.20. Let us define an admissible $\mathbb{Z}/\overline{m}\mathbb{Z}$ -grading $d : H \rightarrow \mathbb{Z}/\overline{m}\mathbb{Z}$ on Γ as follows : for every vertex $v \in H/\sigma$,

- If there exist half-edges in $H \setminus H'$ and in H' that are incident with v , then we choose any maximal sector $(h, r) \in \text{Sect}(H', \sigma)$ around v and we set $d(\sigma^{-1}h) = \overline{m}/m(\sigma^{-1}h)$ and $d(h') = 0$ for all $h' \neq \sigma^{-1}h$ incident with v .



The blue half-edge represents the only half-edge in the cyclic ordering around v that has a non trivial degree for d .

- Else, we choose any $h \in H$ incident with v and we set $d(h) = \overline{m}/m(h)$ and $d(h') = 0$ for all $h' \neq h$ incident with v .

Let $\Gamma_d = (H_d, \iota_d, \sigma_d)$ be the Galois covering of Γ constructed before Proposition 2.9 and $H'_d = H' \times \mathbb{Z}/\overline{m}\mathbb{Z} \subset H_d$. We denote by B_d and B'_d the Brauer graph algebras associated to Γ_d and $\mu_{H'_d}^+(\Gamma_d)$ respectively. Moreover, G denotes the cyclic group $(\mathbb{C}_{\overline{m}}, \cdot)$ of order \overline{m} . We have the following commutative diagram

$$\begin{array}{ccc} \text{per}(B'_d G) & \xrightarrow{-\overset{\mathbf{L}}{\otimes}_{B'_d G} T_d G} & \text{per}(B_d G) \\ \downarrow -\overset{\mathbf{L}}{\otimes}_{B'_d G} B'_d G f & & \downarrow -\overset{\mathbf{L}}{\otimes}_{B_d G} B_d G f \\ \text{per}(B') & \overset{-\overset{\mathbf{L}}{\otimes}_{B'} T}{\dashrightarrow} & \text{per}(B) \end{array}$$

where f is the idempotent given by the sum of the $e_{[h_0]} \otimes 1_G$ for all $[h] \in H/\iota$ and $T_d G = \mu^+(B_d G; e_{H'_d} B_d G)$. The vertical arrows arise from Proposition 2.11 since $\mu_{H'_d}^+(\Gamma_d)$ is the Galois covering of $\mu_{H'}^+(\Gamma, d)$ by Proposition 2.19. The top arrow is given by Proposition 2.21. This leads to a derived equivalence between B' and B and the associated tilting object T is given by

$$T = (f B'_d G \overset{\mathbf{L}}{\otimes}_{B'_d G} T_d G) \overset{\mathbf{L}}{\otimes}_{B_d G} B_d G f$$

It remains to prove that $T \simeq \mu^+(B; e_{H''} B)$. We saw in the proof of Proposition 2.11 that there is an isomorphism in $\text{per}(B_d G)$

$$T_d G \simeq T_d \overset{\mathbf{L}}{\otimes}_{B_d} B_d G$$

where $T_d = \mu^+(B_d; e_{H'_d} B_d)$. Hence, denoting by $f_{H'_d}$ the idempotent given by the sum of the $e_{[h_0]} \otimes 1_G$ for all $[h] \in (H \setminus H')/\iota$, we have

$$f B'_d G \overset{\mathbf{L}}{\otimes}_{B'_d G} T_d G \simeq f_{H'_d} B_d G \oplus \bigoplus_{[h_0] \in H'_d/\iota} \text{Cone}(\alpha_{[h_0]} \overset{\mathbf{L}}{\otimes}_{B_d} B_d G)$$

where $\alpha_{[h_0]}$ is the left minimal $\text{add}(e_{H'_d} B_d)$ -approximation of $e_{[h_0]} B_d$. By construction of d , $\alpha_{[h_0]} \overset{\mathbf{L}}{\otimes}_{B_d} B_d G$ is of the form

$$\alpha_{[h_0]} \overset{\mathbf{L}}{\otimes}_{B_d} B_d G : e_{[h_0]} B_d G \longrightarrow e_{[(\sigma^{r(h_0)+1} h)_0]} B_d G \oplus e_{[(\sigma^{r(\iota_d h_0)+1} \iota h)_0]} B_d G$$

Thus, $\alpha_{[h_0]} \overset{\mathbf{L}}{\otimes}_{B_d} B_d G$ is the left minimal $\text{add}(f_{H'_d} B_d G)$ -approximation of $e_{[h_0]} B_d G$. Hence,

$$f B'_d G \overset{\mathbf{L}}{\otimes}_{B'_d G} T_d G \simeq \mu^+(f B_d G; f_{H'_d} B_d G)$$

Moreover, since $-\overset{\mathbf{L}}{\otimes}_{B_d G} B_d G f : \text{per}(B_d G) \rightarrow \text{per}(B_\Gamma)$ is an equivalence of triangulated categories, we conclude that

$$T \simeq \mu^+(f B_d G; f_{H'_d} B_d G) \overset{\mathbf{L}}{\otimes}_{B_d G} B_d G f \simeq \mu^+(f B_d G f; f_{H'_d} B_d G f)$$

which is isomorphic to $\mu^+(B; e_{H''} B)$ by Proposition 2.11. \square

Remark 2.22. Using the classification of Brauer graph algebras up to derived equivalence given in [OZ22], we can directly prove that the algebras B and B' are derived equivalent. Indeed, we have to check the following conditions

- (i) Γ and $\mu_{H'}^+(\Gamma)$ have the same number of edges and vertices;
- (ii) Γ and $\mu_{H'}^+(\Gamma)$ have the same number of faces and their multi-sets of the perimeters of the faces coincide;
- (iii) The multi-sets of the multiplicities of the vertices of Γ and $\mu_{H'}^+(\Gamma)$ coincide;
- (iv) Either both or none of Γ and $\mu_{H'}^+(\Gamma)$ are bipartite.

Thanks to Proposition 1.17 in [Sot24], it only remains to prove the third point. It is clear by the definition of $m_{(h,r)}$ that the multi-sets of the multiplicities of the vertices of Γ and $\mu_{(h,r)}^+(\Gamma)$ coincide for all (h, r) maximal sector in Γ of elements in H' . Since $\mu_{H'}^+(\Gamma)$ is obtained from the succession of generalized Kauer moves of all maximal sectors of elements in H' , we deduce that the third point holds. Hence, the algebras B and B' are derived equivalent.

3. Generalized Kauer moves for skew Brauer graph algebras

In this section, we define the notion of skew Brauer graph algebras which is a generalization of Brauer graph algebras. Our aim is to define a notion of generalized Kauer moves for skew Brauer graph algebras so that it corresponds to a silting mutation. This will again be a generalization of Theorem 3.10 in [Sot24]. Throughout this section, the characteristic of the field k is supposed to be different than 2.

3.1. Skew Brauer graph algebras

In this part, we define the notion of skew Brauer graph which is a Brauer graph where the pairing may have some fixed points. From this combinatorial data, we construct a new algebra called a skew Brauer graph algebra. This construction is analogous to the definition of a Brauer graph algebra.

Definition 3.1. A *skew Brauer graph* is the data $\Gamma = (H, \iota, \sigma, m)$ where

- H is the set of half-edges;
- ι is a permutation of H with possible fixed points satisfying $\iota^2 = \text{id}_H$;
- σ is a permutation of H called the *orientation*;
- $m : H \rightarrow \mathbb{Z}_{>0}$ is a map that is constant on a σ -orbit : it is called the *multiplicity*.


To the data $\Gamma = (H, \iota, \sigma, m)$, one can naturally define a graph. For this, let us denote H_\circ the set of half-edges that are not fixed by ι and H_\times the set of half-edges that are fixed by ι . Hence, the vertex set of this graph is given by $\Gamma_0 \simeq H/\sigma \cup H_\times$ and the edge set is given by $\Gamma_1 = H/\iota$. Moreover, each cycle in the decomposition of σ gives rise to a cyclic ordering of the half-edges around a vertex in H/σ . Since $m : H \rightarrow \mathbb{Z}_{>0}$ is constant on a σ -orbit, it induces a map $\tilde{m} : H/\sigma \rightarrow \mathbb{Z}_{>0}$. We denote by \circ the vertices in H/σ and by \times the vertices in H_\times . By construction of the graph, each \times -vertex has a unique edge incident to it. One can understand an edge incident to a \times -vertex as a “degenerate” edge. Hence, one can see a skew Brauer graph as a Brauer graph with possible “degenerate” edges. From now on, we identify $\Gamma = (H, \iota, \sigma, m)$ with the graph that we have constructed above. By convention, the orientation of a skew Brauer graph corresponds to the local embedding of each \circ -vertex into the counterclockwise oriented plane.

Remark 3.2. For a similar reason than in Remark 2.2, we do not take into consideration the two following skew Brauer graph algebras



where the first one is given in Remark 2.2 and the other is given by $H = \{2\}$, $\iota = \text{id}_H = \sigma$ and m being identically 1. More precisely, we assume that such graphs do not appear in any connected component of a given skew Brauer graph. In particular, since we have excluded such graphs, there is no half-edge that is fixed simultaneously by ι and σ .

Example 3.3. Let $\Gamma = (H, \iota, \sigma, m)$ be a skew Brauer graph where the set of half-edges H is $\{1^+, 1^-, 2, 3, 4^+, 4^-, 5^+, 5^-\}$, the involution ι is $(1^+ 1^-)(4^+ 4^-)(5^+ 5^-)$, the permutation σ is $(1^- 3 2)(1^+ 4^+ 5^+)$ and the multiplicity $m : H \rightarrow \mathbb{Z}_{>0}$ is given by $m(4^-) = 3$, $m(3) = m(2) = m(1^-) = 2$ and $m(4^+) = m(1^+) = m(5^+) = m(5^-) = 1$. With our previous convention, this skew Brauer graph can be represented as follows




where the multiplicity of a half-edge is given by the value in the \circ -vertex it is incident to. From now on, we will only write on the graph the multiplicities that are strictly greater than one.

Given a skew Brauer graph $\Gamma = (H, \iota, \sigma, m)$, one can construct a quiver $Q_\Gamma = (Q_0, Q_1)$ as follows

- The vertex set Q_0 of Q_Γ is given by $H_\circ/\iota \cup (H_\times/\iota \times \mathbb{Z}/2\mathbb{Z})$. We denote by $[h]_i$ a vertex of Q_Γ where $i = \emptyset$ if $[h] \in H_\circ/\iota$ and $i = 0, 1$ if $[h] \in H_\times/\iota$.
- The arrow set Q_1 of Q_Γ is induced by the orientation σ and the multiplicity m . More precisely, for any half-edge $h \in H$ that is not fixed by σ there are arrows ${}^j\alpha_h^i$ from $[h]_i$ to $[\sigma h]_j$ for all $i, j = \emptyset, 0, 1$. Moreover, for any half-edge h that is fixed by σ there is an arrow α_h from $[h]$ to itself if and only if $m(h) > 1$.

Note that any half-edge $h \in H$ inducing an arrow in Q_Γ gives rise to oriented cycles in Q_Γ that begins and ends at $[h]_i$ for $i = \emptyset, 0, 1$. We call such an oriented cycle a *special h_i -cycle*. By definition, there are $2^{|H_\times \cap (\text{Orb}_\sigma(h) \setminus \{h\})|}$ such special cycles where $\text{Orb}_\sigma(h)$ denotes the σ -orbit of h . We denote by \mathcal{C}_{h_i} the set of all special h_i -cycle. Moreover, an element of \mathcal{C}_{h_i} is of the form ${}^i\alpha_{\sigma^{n-1}h} {}^{i_{n-1}} \dots {}^{i_1}\alpha_h^i$ for some $i_1, \dots, i_{n-1} = \emptyset, 0, 1$ where n is the size of the σ -orbit of h .

Example 3.4. Let $\Gamma = (H, \iota, \sigma, m)$ be the skew Brauer graph defined in Example 3.3. Its corresponding quiver is given by



In this case, the sets of special h_i -cycles are given by $\mathcal{C}_{1-} = \{\alpha_2^{i_2} \alpha_3^{i_1} \alpha_{1-}^{i_1} \mid i_1, i_2 = 0, 1\}$, $\mathcal{C}_{2_i} = \{\alpha_3^{i_1} \alpha_{1-}^{i_1} \alpha_2^i \mid i_1 = 0, 1\}$ for $i = 0, 1$, $\mathcal{C}_{3_i} = \{\alpha_{1-} \alpha_2^{i_1} \alpha_3^i \mid i_1 = 0, 1\}$ for $i = 0, 1$, $\mathcal{C}_{1+} = \{\alpha_5 + \alpha_4 + \alpha_{1+}\}$, $\mathcal{C}_{4+} = \{\alpha_{1+} + \alpha_5 + \alpha_{4+}\}$, $\mathcal{C}_{4-} = \{\alpha_{4-}\}$ and $\mathcal{C}_{5+} = \{\alpha_{4+} + \alpha_{1+} + \alpha_{5+}\}$. Note that there is no special 5^- -cycle since 5^- is fixed by σ and is of multiplicity 1 so it does not induce an arrow in Q_Γ .

Definition 3.5. Let $\Gamma = (H, \iota, \sigma, m)$ be a skew Brauer graph and Q_Γ be its associated quiver. The *skew Brauer graph algebra* B_Γ of Γ is the path algebra kQ_Γ/I_Γ where the set of relations I_Γ is generated by

$$(I) \quad (2^{n_\times(h)} C_h)^{m(h)} = (2^{n_\times(\iota h)} C_{\iota h})^{m(\iota h)}$$

for all special cycles $C_h \in \mathcal{C}_h$ and $C_{\iota h} \in \mathcal{C}_{\iota h}$ where $n_\times(h)$ denotes the number of half-edges in the σ -orbit of h that are in H_\times .

$$(II) \quad {}^{i_1} \alpha_h^i (C_{h_i})^{m(h)}$$

for all special cycles $C_{h_i} = {}^i \alpha_{\sigma^{n-1} h}^{i_{n-1}} \dots {}^{i_1} \alpha_h^i \in \mathcal{C}_{h_i}$.

$$(III) \quad {}^{i'} \alpha_{\iota \sigma h} \alpha_h^i$$

for all $h \in H$ such that h and $\iota \sigma h$ both induce an arrow with $\sigma h \in H_\circ$ and $i, i' = \emptyset, 0, 1$.

$$(IV) \quad {}^{i+1} \alpha_{\sigma^{n-1} h}^{i_{n-1}} \dots {}^{i_1} \alpha_h^i (C_{h_i})^{m(h)-1}$$

for all special cycles $C_{h_i} = {}^i \alpha_{\sigma^{n-1} h}^{i_{n-1}} \dots {}^{i_1} \alpha_h^i \in \mathcal{C}_{h_i}$ with $h \in H_\times$.

$$(V) \quad {}^{i'} \alpha_{\sigma h}^0 \alpha_h^i - {}^{i'} \alpha_{\sigma h}^1 \alpha_h^i$$

for all $h \in H$ not fixed by σ such that $\sigma h \in H_\times$ and $i, i' = \emptyset, 0, 1$.

In particular, for $H_\times = \emptyset$, the definitions of skew Brauer graph and skew Brauer graph algebra coincide with the definitions of Brauer graph and Brauer graph algebra. Thanks to the relation (V), note that all the special h_i -cycles are equal in B_Γ . Moreover, these relations are not minimal as we can see in the following example.

Example 3.6. Let $\Gamma = (H, \iota, \sigma, m)$ the skew Brauer graph defined in Example 3.3. The skew Brauer graph algebra associated to Γ is defined as follows : its quiver is described in Example 3.4 and its ideal of relations is generated by

$$(I) \quad \alpha_{1+} + \alpha_5 + \alpha_{4+} - \alpha_{4-}^3 \quad \text{and} \quad 16 (\alpha_2^{i_2} \alpha_3^{i_1} \alpha_{1-}^{i_1})^2 - \alpha_5 + \alpha_4 + \alpha_{1+} \quad \text{for all } i_1, i_2 = 0, 1;$$

$$(II) \quad \alpha_{1+} + \alpha_5 + \alpha_4 + \alpha_{1+}, \alpha_{4+} + \alpha_{1+} + \alpha_5 + \alpha_{4+}, \alpha_5 + \alpha_4 + \alpha_{1+} + \alpha_5, \alpha_{4-}^4 \quad \text{and} \quad {}^{i_1} \alpha_{1-} (\alpha_2^{i_2} \alpha_3^{i_1} \alpha_{1-}^{i_1})^2, \\ {}^{i_1} \alpha_3^i ({}^i \alpha_{1-} \alpha_2^{i_1} \alpha_3^i)^2, \alpha_2^i ({}^i \alpha_3^{i_1} \alpha_{1-} \alpha_2^i)^2 \quad \text{for all } i, i_1, i_2 = 0, 1;$$

$$(III) \quad \alpha_{4-} \alpha_{1+}, \alpha_{4+} \alpha_{4-} \quad \text{and} \quad \alpha_{1+} \alpha_2^i, {}^i \alpha_{1-} \alpha_5 \quad \text{for all } i = 0, 1;$$

$$(IV) \quad {}^{i+1} \alpha_{1-} \alpha_2^{i_1} \alpha_3^i \alpha_{1-} \alpha_2^{i_1} \alpha_3^i, {}^{i+1} \alpha_3^{i_1} \alpha_{1-} \alpha_2^i \alpha_3^{i_1} \alpha_{1-} \alpha_2^i \quad \text{for all } i_1 = 0, 1;$$

$$(V) \quad {}^i \alpha_3^0 \alpha_{1-} - {}^i \alpha_3^1 \alpha_{1-}, \alpha_2^0 \alpha_3^i - \alpha_2^1 \alpha_3^i \quad \text{for all } i = 0, 1.$$

Note that some of the relations in (II) follows from the relations in (I) and (III).

3.2. Covering of skew Brauer graphs

In this section, we explain how skew Brauer graph can be covered by a Brauer graph. This construction is analogous to the Galois covering of a Brauer graph with multiplicity (cf after Definition 2.8). In particular, we need to introduce a particular $\mathbb{Z}/2\mathbb{Z}$ -grading on the skew Brauer graph.

Definition 3.7. Let $\Gamma = (H, \iota, \sigma, m)$ be a skew Brauer graph.

- We say that a $\mathbb{Z}/2\mathbb{Z}$ -grading $d : H \rightarrow \mathbb{Z}/2\mathbb{Z}$ is *0-homogeneous* if

$$\sum_{h \in H, s(h)=v} d(h) = 0$$

for all $v \in H/\sigma$, where $s : H \rightarrow H/\sigma$ is the source map for \circ -vertex in Γ .


- We say that (Γ, d) is a $\mathbb{Z}/2\mathbb{Z}$ -graded skew Brauer graph if $d : H \rightarrow \mathbb{Z}/2\mathbb{Z}$ is 0-homogeneous.

Let $(\Gamma, d) = (H, \iota, \sigma, m, d)$ be a $\mathbb{Z}/2\mathbb{Z}$ -graded skew Brauer graph. One can construct a Brauer graph $\Gamma_d = (H_d, \iota_d, \sigma_d, m_d)$ as follows


- $H_d = H \times (\mathbb{Z}/2\mathbb{Z})$: an element of H_d will be denoted h_i for $h \in H$ and $i \in \mathbb{Z}/2\mathbb{Z}$;
- For all $h_i \in H_d$, $\iota_d(h_i) = \begin{cases} h_{i+1} & \text{if } h \in H_\times \\ (\iota h)_i & \text{if } h \in H_\circ \end{cases}$;
- For all $h_i \in H_d$, $\sigma_d(h_i) = (\sigma h)_{i+d(h)}$;
- For all $h_i \in H_d$, $m_d(h_i) = m(h)$.

One can easily check that ι_d has no fixed points and that m_d is constant on σ_d -orbits. Hence, Γ_d is indeed a Brauer graph. Moreover, it is clear that the projection $p_\Gamma : H_d \rightarrow H$ defined a morphism of skew Brauer graphs from Γ_d to Γ i.e. it commutes with the involutions and the orientations of the skew Brauer graphs (cf Definition 2.7). The Brauer graph Γ_d will be called the *covering* of Γ .

Example 3.8. Let us consider the skew Brauer graph $\Gamma = (H, \iota, \sigma, m)$ defined in Example 3.3. We equip Γ with a 0-homogeneous $\mathbb{Z}/2\mathbb{Z}$ -grading $d : H \rightarrow \mathbb{Z}/2\mathbb{Z}$ defined by $d(h) = 0$ for all $h \in H$ which can be represented on the graph as follows



Then, the covering $\Gamma_d = (H_d, \iota_d, \sigma_d, m_d)$ of the $\mathbb{Z}/2\mathbb{Z}$ -graded skew Brauer graph (Γ, d) is given by the following Brauer graph



In what follows, we will prove that the skew Brauer graph algebra B associated to Γ can be recovered from the Brauer graph algebra B_d associated to the covering Γ_d thanks to the construction of the quiver and relations of a skew group algebra for a cyclic group [RR85, Section 2.3]. We first need to define an action of the cyclic group $G := (\mathbb{C}_2, \cdot)$ of order 2 on B_d whose generator will be denoted by g . Denoting by $e_{[h_i]}$ the idempotent in B_d corresponding to the edge $[h_i] \in H_d/\iota_d$, there is a natural action of G on $B_d = kQ_d/I_d$ given by

- $g \cdot e_{[h_i]} = e_{[h_{i+1}]}$ for all $[h_i] \in H_d/\iota_d$;
- $g \cdot \alpha_{h_i} = \alpha_{h_{i+1}}$ where α_{h_i} is the arrow in Q_d induced by the half-edge $h_i \in H_d$.

Similarly, the dual \widehat{G} of G is also a cyclic group and we denote by χ its generator. Denoting by $e_{[h]_i}$ for $i = \emptyset, 0, 1$ the idempotent(s) in B arising from the edge $[h] \in H/\iota$, there is a natural action of \widehat{G} on $B = kQ/I$ given by

- $\chi \cdot e_{[h]} = e_{[h]}$ for all $[h] \in H_o/\iota$ and $\chi \cdot e_{[h]_i} = e_{[h]_{i+1}}$ for all $[h] \in H_\times/\iota$;
- $\chi \cdot {}^j\alpha_h^i = (-1)^{-d(h)} {}^{j+1}\alpha_h^{i+1}$ where ${}^j\alpha_h^i$ is an arrow in Q induced by the half-edge $h \in H$. By an abuse of notation, if $i = \emptyset$ then $i + 1 = \emptyset$.

Note that the G -action on B_d arises from a G -action on Q_d which preserves the ideal of relations I_d whereas the \widehat{G} -action on B does not come from an action on its quiver.

Proposition 3.9. *Let f be the idempotent in $B_d G$ given by the sum of the $e_{[h_0]} \otimes 1_G$ for all $[h] \in H/\iota$. Then, the algebra $f B_d G f$ has a natural \widehat{G} -action and the map*

$$\begin{aligned} \phi : \quad B &\longrightarrow f B_d G f \\ e_{[h]} &\longmapsto f_{[h]} := e_{[h_0]} \otimes 1_G \\ e_{[h]_i} &\longmapsto f_{[h]_i} := e_{[h_0]} \otimes \frac{1_G + (-1)^i g}{2} \quad \text{for } i = 0, 1 \\ {}^j\alpha_h^i &\longmapsto {}^j\beta_h^i \quad \text{for } i, j = \emptyset, 0, 1 \end{aligned}$$

is an isomorphism of algebras which commutes with the \widehat{G} -actions, where ${}^j\beta_h^i : f_{[h]_i} \rightarrow f_{[h]_j}$ are the arrows in $f B_d G f$.

Proof. Since $\chi \cdot f = f$, the natural \widehat{G} -action on $B_d G$ extends onto a \widehat{G} -action on $f B_d G f$. As for the proof of Proposition 2.11, the isomorphism follows again from the construction of the quiver and relations of $f B_d G f$ given in [RR85].

- The vertices of the quiver of fB_dGf are in bijection with the idempotents $f_{[h]_i}$ defined in the proposition for $i = \emptyset, 0, 1$. By definition, its arrows are of the form ${}^j\beta_h^i : f_{[h]_i} \rightarrow f_{[h]_j}$ for $i, j = \emptyset, 0, 1$.
- The relations of fB_dGf are determined thanks to a representative in the G -orbits of the relations of B_d . Hence, its ideal of relation is generated by

(I')

$$\sum_{\substack{i_k^l = \emptyset, 0, 1 \\ (k, l) \in \{1, \dots, n\} \times \{1, \dots, m(h)\} \setminus \{(n, m(h))\}}} {}^{i+1}\beta_{\sigma^{n-1}h} {}^{i_n^{m(h)}} \dots {}^{i_1^{m(h)}}\beta_h {}^{i_n^{m(h)-1}} \dots {}^{i_1^1}\beta_{\sigma^{n-1}h} {}^{i_{n-1}^1} \dots {}^{i_1^1}\beta_h^i$$

for all $i = 0, 1$ and $h \in H_\times$ inducing an arrow in Q , where n is the size of the σ -orbit of h .

(II')

$$\left(\sum_{i_1, \dots, i_{n-1} = \emptyset, 0, 1} \beta_{\sigma^{n-1}h} {}^{i_{n-1}} \dots {}^{i_1}\beta_h \right)^{m(h)} - \left(\sum_{i_1, \dots, i_{n'-1} = \emptyset, 0, 1} \beta_{\sigma^{n'-1}ih} {}^{i_{n-1}} \dots {}^{i_1}\beta_{ih} \right)^{m(ih)}$$

for all $h \in H_\circ$ such that h and ih both induce an arrow in Q , where n and n' are the size of the σ -orbit of h and ih respectively.

(III')

$$\sum_{\substack{i_k^l = \emptyset, 0, 1 \\ k=1, \dots, n \\ l=1, \dots, m(h)}} {}^j\beta_h {}^{i_n^{m(h)}} {}^{i_n^{m(h)}}\beta_{\sigma^{n-1}h} {}^{i_{n-1}^{m(h)}} \dots {}^{i_1^{m(h)}}\beta_h {}^{i_n^{m(h)-1}} \dots {}^{i_1^1}\beta_{\sigma^{n-1}h} {}^{i_{n-1}^1} \dots {}^{i_1^1}\beta_h^i$$

for all $i, j = \emptyset, 0, 1$ and $h \in H$ inducing an arrow in Q , where n is the size of the σ -orbit of h .

(IV')

$${}^j\beta_{\sigma h} {}^0\beta_h^i - {}^j\beta_{\sigma h} {}^1\beta_h^i$$

for all $i, j = \emptyset, 0, 1$ and $h \in H$ inducing an arrow in Q such that $\sigma h \in H_\times$.

(V')

$${}^j\beta_{\iota\sigma h} \beta_h^i$$

for all $i, j = \emptyset, 0, 1$ and $h \in H$ such that h and $\iota\sigma h$ induce an arrow in Q and $\sigma h \in H_\circ$.

The relations are obtained the same way as in the proof of Proposition 2.11. In this case, there are sums appearing in the relations since we have

$$\alpha_{h-d(h)} \otimes g^{-d(h)} = \sum_{i, j = \emptyset, 0, 1} {}^i\beta_h^j$$

Note that the relations (I') and (II') arise from the type (I) relation in B_d , the relation (III') arise from the type (II) relation in B_d and the relations (IV') and (V') arise from the type (III) relation in B_d . In particular, the relation (IV') means that all special h_i -cycle are equal in fB_dGf . Hence, these five relations may be reformulated to obtain a correspondence with the relations of B under the morphism ϕ . \square

Thanks to this proposition, one can understand skew Brauer graph algebras of multiplicity identically one as the trivial extension of a skew gentle algebra. In particular, this generalizes Theorem 3.13 in [Sch18] which interprets Brauer graph algebras of multiplicity identically one as the trivial extension of gentle algebras. We recall that a skew gentle algebra is a basic algebra that is Morita equivalent to the skew group algebra of a gentle algebra equipped with a $\mathbb{Z}/2\mathbb{Z}$ -action. (cf for instance Section 3.2 in [AB22]).

Definition 3.10. Let $B = kQ/I$ be a skew Brauer graph algebra associated to $\Gamma = (H, \iota, \sigma, m)$ and Δ be a subset of H consisting of representatives of each σ -orbits. An *admissible cut* of B is the set of all arrows in Q induced by the half-edges in Δ . By an abuse of notation, we also denote by Δ the admissible cut.

Theorem 3.11. Let $B = kQ/I$ be a skew Brauer graph algebra of multiplicity identically one associated to $\Gamma = (H, \iota, \sigma)$ and equipped with an admissible cut Δ . Then, the following holds.

- (1) The algebra $B_\Delta := kQ/\langle I \cup \Delta \rangle$ is a skew gentle algebra, where $\langle I \cup \Delta \rangle$ denotes the ideal of kQ generated by $I \cup \Delta$.
- (2) The trivial extension of B_Δ is isomorphic to B .

Proof. Let us consider $d : H \rightarrow \mathbb{Z}/2\mathbb{Z}$ a 0-homogeneous $\mathbb{Z}/2\mathbb{Z}$ -grading on Γ . Hence, one can construct $\Gamma_d = (H_d, \iota_d, \sigma_d)$ the covering of Γ which is a Brauer graph of multiplicity identically one by construction. We denote by B_d the Brauer graph algebra associated to Γ_d and by $p_\Gamma : \Gamma_d \rightarrow \Gamma$ the morphism of skew Brauer graphs given by the projection (cf before Example 3.8). One can easily check that $p_\Gamma^{-1}(\Delta)$ is an admissible cut for Γ_d that is stable under the G -action on Γ_d defined before Proposition 3.9, where $G = (\mathbb{C}_2, \cdot)$ denotes the cyclic group of order 2. By Theorem 3.13 in [Sch18], $(B_d)_{p_\Gamma^{-1}(\Delta)}$ is a gentle algebra whose trivial extension is isomorphic to B_d . Moreover, $(B_d)_{p_\Gamma^{-1}(\Delta)}$ has a natural G -action which can be extended as a G -action onto its trivial extension as explained before Proposition 1.4. One can check that the isomorphism of algebras $\text{Triv}((B_d)_{p_\Gamma^{-1}(\Delta)}) \simeq B_d$ commute with the G -actions. Thanks to Proposition 1.4, we deduce the following isomorphisms of algebras

$$B_d G \simeq \text{Triv}((B_d)_{p_\Gamma^{-1}(\Delta)}) G \simeq \text{Triv}((B_d)_{p_\Gamma^{-1}(\Delta)}) G$$

By Proposition 3.9, we have the following isomorphisms of algebras

$$B \simeq f B_d G f \simeq f \text{Triv}((B_d)_{p_\Gamma^{-1}(\Delta)}) G f$$

Moreover, denoting by A the algebra $(B_d)_{p_\Gamma^{-1}(\Delta)} G$, one can check that the following morphism also gives an isomorphism of algebras

$$\begin{aligned} f \text{Triv}(A) f &\xrightarrow{\sim} \text{Triv}(f A f) \\ f \cdot (a, \phi) \cdot f &\mapsto (f a f, (f \phi f)|_{f A f}) \\ f \cdot (b, \psi') \cdot f &\longleftarrow (f b f, \psi) \end{aligned}$$

where $\psi' : A \rightarrow k$ is defined by $\psi'(c) = \psi(f c f)$ for all $c \in A$. Furthermore, one can easily check that the isomorphism of algebras given in Proposition 3.9 restricts to an isomorphism of algebras between $B_\Delta = B/\langle \Delta \rangle$ and $f A f = f(B_d/\langle p_\Gamma^{-1}(\Delta) \rangle) G f$. This concludes the proof. \square

3.3. Generalized Kauer moves

Our goal is to define generalized Kauer moves for skew Brauer graph algebras so that they can be interpreted in terms of silting mutations. The idea of the proof is analogous to the one used in Section 2. We will use the covering defined in the previous subsection which can be constructed whenever the skew Brauer graph is equipped with a 0-homogeneous $\mathbb{Z}/2\mathbb{Z}$ -grading. Hence, we need to define a graded version of these generalized Kauer moves. We begin by defining them for sectors (cf Definition 2.12).

Definition 3.12. Let $(\Gamma, d) = (H, \iota, \sigma, m, d)$ be a $\mathbb{Z}/2\mathbb{Z}$ -graded skew Brauer graph and $H' \subset H$ be stable under ι . The $\mathbb{Z}/2\mathbb{Z}$ -graded generalized Kauer move of a sector $(h, r) \in \text{sect}(H', \sigma)$ in Γ gives rise to a $\mathbb{Z}/2\mathbb{Z}$ -graded skew Brauer graph $\mu_{(h,r)}^+(\Gamma, d) = (H, \iota, \sigma_{(h,r)}, m_{(h,r)}, d_{(h,r)})$ where

$$\begin{aligned} \sigma_{(h,r)} &= (h \sigma^{r+1} h) \sigma (\sigma^r h \iota \sigma^{r+1} h) & \text{and} & & m_{(h,r)} : & H & \rightarrow & \mathbb{Z}_{>0} \\ & & & & \sigma^i h & \mapsto & m(\iota \sigma^{r+1} h) & \text{for } i = 0, \dots, r \\ & & & & h' & \mapsto & m(h') & \text{for } h' \neq \sigma^i h \end{aligned}$$

and where the $\mathbb{Z}/2\mathbb{Z}$ -grading $d_{(h,r)} : H \rightarrow \mathbb{Z}/2\mathbb{Z}$ is defined by

$$\begin{aligned} d_{(h,r)} : \iota \sigma^{r+1} h & \mapsto \begin{cases} - \sum_{i=0}^r d(\sigma^i h) & \text{if } \sigma^{r+1} h \in H_{\circ} \\ - \sum_{i=0}^r d(\sigma^i h) - 1 & \text{if } \sigma^{r+1} h \in H_{\times} \end{cases} \\ \sigma^r h & \mapsto \begin{cases} d(\iota \sigma^{r+1} h) + d(\sigma^r h) & \text{if } \iota \sigma^{r+1} h \neq \sigma^{-1} h \text{ and } \sigma^{r+1} h \in H_{\circ} \\ d(\iota \sigma^{r+1} h) + d(\sigma^r h) + 1 & \text{if } \iota \sigma^{r+1} h \neq \sigma^{-1} h \text{ and } \sigma^{r+1} h \in H_{\times} \\ \sum_{i=-1}^r d(\sigma^i h) + d(\sigma^r h) & \text{if } \iota \sigma^{r+1} h = \sigma^{-1} h \text{ and } \sigma^{r+1} h \in H_{\circ} \\ \sum_{i=-1}^r d(\sigma^i h) + d(\sigma^r h) + 1 & \text{if } \iota \sigma^{r+1} h = \sigma^{-1} h \text{ and } \sigma^{r+1} h \in H_{\times} \end{cases} \\ \sigma^{-1} h & \mapsto \begin{cases} \sum_{i=-1}^r d(\sigma^i h) & \text{if } \iota \sigma^{r+1} h \neq \sigma^{-1} h \\ d_{(h,r)}(\iota \sigma^{r+1} h) & \text{if } \iota \sigma^{r+1} h = \sigma^{-1} h \end{cases} \\ h' & \mapsto d(h') \quad \text{for } h' \neq \iota \sigma^{r+1} h, \sigma^r h, \sigma^{-1} h \end{aligned}$$

The underlying Brauer graph $\mu_{(h,r)}^+(\Gamma) = (H, \iota, \sigma_{(h,r)}, m_{(h,r)})$ can be obtained from Γ as in Figure 1 and in Remark 2.14 when $\sigma^{r+1} h \in H_{\circ}$ and as follows when $\sigma^{r+1} h \in H_{\times}$





Figure 3: Generalized Kauer move of a sector (h, r) when $\sigma^{r+1}h$ is in H_\times

Remark 3.13. In the special case where $\iota\sigma^{r+1}h = \sigma^{-1}h$, then the $\mathbb{Z}/2\mathbb{Z}$ -graded generalized Kauer move of the sector (h, r) does not change the underlying Brauer graph which is given in Remark 2.14 when $\sigma^{r+1}h \in H_\circ$ and as follows when $\sigma^{r+1}h \in H_\times$



Similarly to Remark 2.14, the degree of the $\sigma^j h$ will a priori change in the process. In particular, the covering of Γ and $\mu_{(h,r)}^+(\Gamma)$ could be different even in this case.

By construction of the generalized Kauer move described in Figure 1 and in the previous figure, it is clear that $m_{(h,r)}$ is constant on the $\sigma_{(h,r)}$ -orbits. Moreover, $d_{(h,r)} : H \rightarrow \mathbb{Z}/2\mathbb{Z}$ is 0-homogeneous. Indeed, one can construct a bijection $\phi : H/\sigma \rightarrow H/\sigma_{(h,r)}$ satisfying

$$\begin{aligned} s_{(h,r)}(\sigma^i h) &= \phi(s(\iota\sigma^{r+1}h)) & \text{for } i = 0, \dots, r \\ s_{(h,r)}(h') &= \phi(s(h')) & \text{for } h' \neq \sigma^i h, i = 0 \dots, r \end{aligned}$$

where $s : H \rightarrow H/\sigma$ and $s_{(h,r)} : H \rightarrow H/\sigma_{(h,r)}$ are the source map for \circ -vertices of Γ and $\mu_{(h,r)}^+(\Gamma)$ respectively. Then, one can easily check that for all vertex $v \in H/\sigma_{(h,r)}$ in $\mu_{(h,r)}^+(\Gamma)$

$$\sum_{h' \in H, s_{(h,r)}(h')=v} d_{(h,r)}(h') = \sum_{h' \in H, s(h')=\phi^{-1}(v)} d(h')$$

Remark 3.14. By definition, if $H_\times = \emptyset$ then Γ is a Brauer graph and by construction $\mu_{(h,r)}^+(\Gamma)$ is also a Brauer graph. Hence, this definition generalizes Definition 2.13.

Thanks to Lemma 2.4 in [Sot24], we can consider successive $\mathbb{Z}/2\mathbb{Z}$ -graded generalized Kauer moves. In the following, we denote by

$$\mu_{(h_1, r_1)(h_2, r_2)}^+(\Gamma, d) = (H, \iota, \sigma_{(h_1, r_1)(h_2, r_2)}, m_{(h_1, r_1)(h_2, r_2)}, d_{(h_1, r_1)(h_2, r_2)})$$


the $\mathbb{Z}/2\mathbb{Z}$ -graded skew Brauer graph defined by $\mu_{(h_1, r_1)}^+(\mu_{(h_2, r_2)}^+(\Gamma, d))$. Using similar computations as in the proof of Proposition 2.5 in [Sot24], we obtain the following result.

Proposition 3.15. *Let $\Gamma = (H, \iota, \sigma, m)$ be a skew Brauer graph and H' be a subset of H stable under ι . Let $(h_1, r_1), (h_2, r_2)$ be two distinct maximal sectors in Γ of elements in H' . For any 0-homogeneous $\mathbb{Z}/2\mathbb{Z}$ -grading $d : H \rightarrow \mathbb{Z}/2\mathbb{Z}$ of Γ , we have*

$$\mu_{(h_1, r_1)(h_2, r_2)}^+(\Gamma, d) = \mu_{(h_2, r_2)(h_1, r_1)}^+(\Gamma, d)$$

Definition 3.16. Let $(\Gamma, d) = (H, \iota, \sigma, m, d)$ be a $\mathbb{Z}/2\mathbb{Z}$ -graded skew Brauer graph and H' be a subset of H stable under ι . The $\mathbb{Z}/2\mathbb{Z}$ -graded generalized Kauer move of H' is the succession of the $\mathbb{Z}/2\mathbb{Z}$ -graded generalized Kauer move of all the maximal sectors $(h, r) \in \text{Sect}(H', \sigma)$ in Γ . This gives rise to a $\mathbb{Z}/2\mathbb{Z}$ -graded skew Brauer graph that will be denoted by $\mu_{H'}^+(\Gamma, d) = (H, \iota, \sigma_{H'}, m_{H'}, d_{H'})$.

Example 3.17. Let us consider $\Gamma = (H, \iota, \sigma, m, d)$ the $\mathbb{Z}/2\mathbb{Z}$ -graded skew Brauer graph defined in Example 3.8. Let $H' = \{1^+, 1^-, 4^+, 4^-\}$ be a subset of H stable under ι . Then, the $\mathbb{Z}/2\mathbb{Z}$ -graded skew Brauer graph $\mu_{H'}^+(\Gamma, d) = (H, \iota, \sigma_{H'}, m_{H'}, d_{H'})$ obtained from a $\mathbb{Z}/2\mathbb{Z}$ -graded generalized Kauer move is given by



where the orientation $\sigma_{H'}$ is $(5^- 1^+ 4^+)(1^- 2 3) = (1^- 3)(1^+ 5^+)\sigma(4^+ 5^-)(1^- 3)$, the multiplicity $m_{H'} : H \rightarrow \mathbb{Z}_{>0}$ is given by $m_{H'}(4^-) = 3$, $m_{H'}(1^-) = m_{H'}(2) = m_{H'}(3) = 2$ and $m_{H'}(5^-) = m_{H'}(1^+) = m_{H'}(4^+) = m_{H'}(5^+) = 1$ and the $\mathbb{Z}/2\mathbb{Z}$ -grading $d_{H'} : H \rightarrow \mathbb{Z}/2\mathbb{Z}$ is given by $d_{H'}(3) = -1$, $d_{H'}(1^-) = 1$ and $d_{H'}(4^-) = d_{H'}(4^+) = d_{H'}(1^+) = d_{H'}(5^-) = d_{H'}(5^+) = d_{H'}(2) = 0$.

Considering a $\mathbb{Z}/2\mathbb{Z}$ -graded generalized Kauer move of a $\mathbb{Z}/2\mathbb{Z}$ -graded skew Brauer graph (Γ, d) , one can construct the covering of $\mu_{H'}^+(\Gamma, d)$ as defined before Example 3.8. The following proposition shows a commutativity between constructing this covering and applying a generalized Kauer move.

Proposition 3.18. *Let $(\Gamma, d) = (H, \iota, \sigma, d)$ be a $\mathbb{Z}/2\mathbb{Z}$ -graded skew Brauer graph and H' be a subset of H stable under ι . We denote by $\Gamma_d = (H_d, \iota_d, \sigma_d, m_d)$ the covering of Γ constructed before Example 3.8 and $H'_d = H' \times \mathbb{Z}/2\mathbb{Z} \subset H_d$. Then, the covering of $\mu_{H'}^+(\Gamma, d)$ is the Brauer graph $\mu_{H'_d}^+(\Gamma_d)$.*

The previous proposition can be summarized in the following commutative diagram

$$\begin{array}{ccc}
\Gamma_d & \xrightarrow[\text{move of } H'_d]{\text{generalized Kauer}} & \mu_{H'_d}^+(\Gamma_d) \\
p_\Gamma \downarrow & & \downarrow p_{\mu^+(\Gamma)} \\
(\Gamma, d) & \xrightarrow[\text{Kauer move of } H']{\mathbb{Z}/2\mathbb{Z}\text{-graded generalized}} & \mu_{H'}^+(\Gamma, d)
\end{array}$$

where $p_\Gamma : \Gamma_d \rightarrow \Gamma$ and $p_{\mu_{H'}^+(\Gamma)} : \mu_{H'_d}^+(\Gamma_d) \rightarrow \mu_{H'}^+(\Gamma)$ are the morphisms of skew Brauer graphs induced by the natural projection and $\mu_{H'}^+(\Gamma)$ denotes the underlying skew Brauer graph of $\mu_{H'}^+(\Gamma, d)$.

Proof. One can easily check that H'_d is a subset of H_d stable under ι_d and that $(h, r) \in \text{Sect}(H', \sigma)$ is a maximal sector in Γ if and only if $(h_i, r) \in \text{Sect}(H'_d, \sigma_d)$ is a maximal sector in Γ_d for all $i \in \mathbb{Z}/2\mathbb{Z}$. By Proposition 3.15, it suffices to prove that the covering of $\mu_{(h,r)}^+(\Gamma, d)$ is given by

$$\mu_{p_\Gamma^{-1}(h,r)}^+(\Gamma_d) = (H_d, \iota_d, (\sigma_d)_{p_\Gamma^{-1}(h,r)}, (m_d)_{p_\Gamma^{-1}(h,r)})$$

for any maximal sector $(h, r) \in \text{Sect}(H', \sigma)$, where $p_\Gamma^{-1}(h, r)$ is the product of the maximal sectors (h_i, r) for $i \in \mathbb{Z}/2\mathbb{Z}$. We denote by

$$\mu_{(h,r)}^+(\Gamma)_{d(h,r)} = (H_{d(h,r)}, \iota_{d(h,r)}, (\sigma_{(h,r)})_{d(h,r)}, (m_{(h,r)})_{d(h,r)})$$

the covering of $\mu_{(h,r)}^+(\Gamma, d)$. By definition, these two Brauer graphs have the same set of half-edges. By construction of the covering, it is clear that they have the same pairing since it only depends on H_\circ and H_\times . Moreover, using similar computations as in the proof of Proposition 2.19, one can check that their orientations are equal. It remains to prove the equality of the multiplicities. It is clear that the equality holds for any half-edge different from the $\sigma_d^i h_j$. Moreover, we have

$$\begin{aligned}
(m_d)_{p_\Gamma^{-1}(h,r)}(\sigma_d^i h_j) &= m_d(\iota_d \sigma_d^{r+1} h_j) = m(\iota \sigma^{r+1} h) \\
(m_{(h,r)})_{d(h,r)}(\sigma_d^i h_j) &= m_{(h,r)}(\sigma^i h) = m(\iota \sigma^{r+1} h)
\end{aligned}$$

This concludes the proof of the proposition. \square

3.4. Compatibility with sifting mutations

The goal of this part is to prove the following theorem which is an analogous version of Theorem 3.10 in [Sot24] for skew Brauer graph algebras.

Theorem 3.19. *Let $\Gamma = (H, \iota, \sigma, m)$ be a skew Brauer graph and H' be a subset of H stable under ι . We assume that 2 and \overline{m} , the least common multiple of the $m(h)$ for $h \in H$, are invertible in the field k . Denoting by B and B' the skew Brauer graph algebras associated respectively to Γ and $\mu_{H'}^+(\Gamma)$, there is an equivalence of triangulated categories*

$$\text{per}(B') \xrightarrow{-\overset{\mathbf{L}}{\otimes}_{B'} \mu^+(B; e_{H''} B)} \text{per}(B)$$

where $e_{H''} B$ is the projective B -module corresponding to the edges in $(H \setminus H')/\iota$.

The idea of the proof is similar to the proof of Theorem 2.20. The following result is an analogous version of Proposition 2.21 for skew Brauer graph algebras.

Proposition 3.20. *Let us consider the setting of Proposition 3.18. We assume that 2 and \bar{m} , the least common multiple of the $m(h)$ for $h \in H$, are invertible in the field k . We denote by B_d and B'_d the Brauer graph algebras associated to Γ_d and $\mu_{H'_d}^+(\Gamma_d)$ respectively. Let G be the cyclic group (\mathbb{C}_2, \cdot) of order 2. Then, there is an equivalence of triangulated categories*

$$\text{per}(B'_d G) \xrightarrow{-\overset{\mathbf{L}}{\otimes}_{B'_d G} \mu^+(B_d G; e_{H'_d} B_d G)} \text{per}(B_d G)$$

where $e_{H'_d} B_d$ denotes the projective B_d -module corresponding to the edges in $(H_d \setminus H'_d)/\iota_d$.

Proof. By Theorem 2.20, there is an equivalence of triangulated categories

$$\text{per}(B'_d) \xrightarrow{-\overset{\mathbf{L}}{\otimes}_{B'_d} T_d} \text{per}(B_d)$$

where $T_d := \mu^+(B_d; e_{H'_d} B_d)$. Similarly to the proof of Proposition 2.21, it suffices to prove that T_d is G -invariant and that the assumptions of Proposition 1.13 hold. Let us first check that T_d is indeed G -invariant. Considering the same notations as in the proof of Proposition 2.21, the left minimal $\text{add}(e_{H'_d} B_d)$ -approximation of $e_{H'_d} B_d$ is given by (cf Figure 2.4)

$$\alpha_{[h_i]} : e_{[h_i]} B_d \xrightarrow{\begin{pmatrix} \alpha(h_i, H'_d) \\ \alpha(\iota_d h_i, H'_d) \end{pmatrix}} e_{[\sigma_d^{r(h_i)+1} h_i]} B_d \oplus e_{[\sigma_d^{r(\iota_d h_i)+1} \iota_d h_i]} B_d$$

Moreover, by definition of the left mutation, we have

$$\mu^+(B_d; e_{H'_d} B_d) = e_{H'_d} B_d \oplus \bigoplus_{[h_i] \in H'_d/\iota_d} \text{Cone}(\alpha_{[h_i]}) = e_{H'_d} B_d \oplus \bigoplus_{[h] \in H'/\iota} \text{Cone}(\beta_{[h]})$$

where $\beta_{[h]} : X_{[h]} \rightarrow Y_{[h]}$ is defined as follows

$$\beta_{[h]} = \begin{cases} \bigoplus_{i \in \mathbb{Z}/2\mathbb{Z}} \alpha_{[h_i]} & \text{if } [h] \in H'_o/\iota \\ \alpha_{[h_0]} & \text{if } [h] \in H'_x/\iota \end{cases} \quad \text{with}$$

$$X_{[h]} = \begin{cases} \bigoplus_{i \in \mathbb{Z}/2\mathbb{Z}} e_{[h_i]} B_d & \text{if } [h] \in H'_o/\iota \\ e_{[h_0]} B_d & \text{if } [h] \in H'_x/\iota \end{cases} \quad \text{and}$$

$$Y_{[h]} = \begin{cases} \bigoplus_{i \in \mathbb{Z}/2\mathbb{Z}} (e_{[\sigma_d^{r(h_i)+1} h_i]} B_d \oplus e_{[\sigma_d^{r(\iota_d h_i)+1} \iota_d h_i]} B_d) & \text{if } [h] \in H'_o/\iota \\ e_{[\sigma_d^{r(h_0)+1} h_0]} B_d \oplus e_{[\sigma_d^{r(h_1)+1} h_1]} B_d & \text{if } [h] \in H'_x/\iota \end{cases}$$

where $H'_\circ = H' \cap H_\circ$ and $H'_\times = H' \cap H_\times$. Since G acts bijectively on the set of edges $(H_d \setminus H'_d)/\iota_d$, the object $e_{H'_d} B_d$ is G -invariant by Example 1.6 (1). Moreover, one can easily check that $X_{[h]}$ and $Y_{[h]}$ are G -invariant objects for all $[h] \in H'/\iota$ thanks to Example 1.6 (1). In this case, the isomorphisms $\iota_g^{X_{[h]}}$ and $\iota_g^{Y_{[h]}}$ (cf Definition 1.5) are respectively given by the action of g on $X_{[h]}$ and $Y_{[h]}$. Furthermore, using the definition of the $\alpha_{[h_i]}$, it is not hard to check that the following diagram commutes for all $g \in G$.

$$\begin{array}{ccc}
X_{[h]}^{g^{-1}} & \xrightarrow{\beta_{[h]}^{g^{-1}}} & Y_{[h]}^{g^{-1}} \\
\downarrow \iota_g^{X_{[h]}} & & \downarrow \iota_g^{Y_{[h]}} \\
X_{[h]} & \xrightarrow{\beta_{[h]}} & Y_{[h]}
\end{array}$$

By Example 1.6 (2), we deduce that $\text{Cone}(\beta_{[h]})$ is G -invariant for all $[h] \in H'/\iota$. Hence, we conclude that $\mu^+(B_d; e_{H'_d} B_d)$ is indeed G -invariant. With similar arguments than in the proof of Proposition 2.21, one can check that the assumptions of Proposition 1.13 holds. This concludes the proof. \square

Proof of Theorem 3.19. Let us consider the 0-homogeneous $\mathbb{Z}/2\mathbb{Z}$ -grading $d : H \rightarrow \mathbb{Z}/2\mathbb{Z}$ defined by $d(h) = 0$ for all $h \in H$. Let $\Gamma_d = (H_d, \iota_d, \sigma_d, m_d)$ be the covering of Γ constructed after Definition 3.7 and $H'_d = H' \times \mathbb{Z}/2\mathbb{Z} \subset H_d$. We denote by B_d and B'_d the Brauer graph algebras associated to Γ_d and $\mu_{H'_d}^+(\Gamma_d)$ respectively. Moreover, G denotes the cyclic group (\mathbb{C}_2, \cdot) of order 2. We have the following commutative diagram

$$\begin{array}{ccc}
\text{per}(B'_d G) & \xrightarrow{-\mathbb{L}_{B'_d G} T_d G} & \text{per}(B_d G) \\
\downarrow -\mathbb{L}_{B'_d G} B'_d G f & & \downarrow -\mathbb{L}_{B_d G} B_d G f \\
\text{per}(B') & \xrightarrow{-\mathbb{L}_{B'} T} & \text{per}(B)
\end{array}$$

where f is the idempotent given by the sum of the $e_{[h_0]} \otimes 1_G$ for all $[h] \in H/\iota$ and $T_d G = \mu^+(B_d G; e_{H'_d} B_d G)$. The vertical arrows arise from Proposition 3.9 since $\mu_{H'_d}^+(\Gamma_d)$ is the covering of $\mu_{H'}^+(\Gamma, d)$ by Proposition 3.18. The top arrow is given by Proposition 3.20. This leads to a derived equivalence between B' and B and the associated tilting object T is given by

$$T = (f B'_d G \mathbb{L}_{B'_d G} T_d G) \mathbb{L}_{B_d G} B_d G f$$

It remains to prove that $T \simeq \mu^+(B; e_{H''} B)$. Similarly to the proof of Theorem 2.20, it suffices to show that $\alpha_{[h_0]} \mathbb{L}_{B_d} B_d G$ is isomorphic to the left minimal $\text{add}(f_{H'_d} B_d G)$ -approximation of $e_{[h_0]} B_d G$ where $f_{H'_d}$ denotes the idempotent in $B_d G$ given by the sum of the $e_{[h_0]} \otimes 1_G$ for all $[h] \in (H \setminus H')/\iota$. Thanks to Proposition 3.9, we know that

$$\alpha_{[h_0]} \mathbb{L}_{B_d} B_d G : e_{[h_0]} B_d G \longrightarrow e_{[\sigma_d^{r(h_0)+1} h_0]} B_d G \oplus e_{[\sigma_d^{r(\iota_d h_0)+1} \iota_d h_0]} B_d G$$

is the left minimal $\text{add}(e_{H'_d} B_d G)$ -approximation of $e_{[h_0]} B_d G$. By definition of d , we have

$$\sigma_d^{r(\iota_d h_0)+1} \iota_d h_0 = \begin{cases} (\sigma^{r(\iota_d h_0)+1} \iota h)_0 & \text{if } h \in H_o \\ (\sigma^{r(\iota_d h_0)+1} h)_1 & \text{if } h \in H_\times \end{cases}$$

Moreover, we have the following isomorphism in $\text{per}(B_d G)$

$$e_{[(\sigma^{r(\iota_d h_0)+1} h)_1]} B_d G \xrightarrow{\sim} e_{[(\sigma^{r(\iota_d h_0)+1} h)_0]} B_d G$$

Indeed, if $\sigma^{r(\iota_d h_0)+1} h \in H_\times$, then $[(\sigma^{r(\iota_d h_0)+1} h)_0] = [(\sigma^{r(\iota_d h_0)+1} h)_1]$ and the isomorphism is just the identity. On the other hand, if $\sigma^{r(\iota_d h_0)+1} h \in H_o$, the isomorphism is given by the multiplication on the left by $(1 \otimes g)$ where g denotes the generator of G . Hence, in each case $\alpha_{[h_0]}^{\mathbb{L}} \otimes_{B_d} B_d G$ can be written as

$$\alpha_{[h_0]}^{\mathbb{L}} \otimes_{B_d} B_d G : e_{[h_0]} B_d G \longrightarrow e_{[(\sigma^{r(h_0)+1} h)_0]} B_d G \oplus e_{[(\sigma^{r(\iota_d h_0)+1} \iota h)_0]} B_d G$$

Thus, $\alpha_{[h_0]}^{\mathbb{L}} \otimes_{B_d} B_d G$ is isomorphic to the left minimal $\text{add}(f_{H'_d} B_d G)$ -approximation of $e_{[h_0]} B_d G$. This concludes the proof. \square

References

- [AB22] C. Amiot and T. Brüstle. “Derived equivalences between skew-gentle algebras using orbifolds”. In: *Documenta Mathematica* 27 (2022), pp. 933–982. DOI: 10.4171/DM/889.
- [AI12] T. Aihara and O. Iyama. “Silting mutation in triangulated categories”. In: *Journal of the London Mathematical Society* 85 (2012). DOI: 10.1112/jlms/jdr055.
- [APR79] M. Auslander, M. I. Platzeck, and I. Reiten. “Coxeter Functors Without Diagrams”. In: *Transactions of the American Mathematical Society* 250 (1979), pp. 1–46. ISSN: 00029947. URL: <http://www.jstor.org/stable/1998978>.
- [Asa19] H. Asashiba. “Smash products of group weighted bound quivers and Brauer graphs”. In: *Communications in Algebra* 47.2 (2019), pp. 585–610. DOI: 10.1080/00927872.2018.1487562. URL: <https://doi.org/10.1080/00927872.2018.1487562>.
- [AZ22] M. Antipov and A. Zvonareva. “Brauer graph algebras are closed under derived equivalence”. In: *Mathematische Zeitschrift* 301 (2022), pp. 1–19. DOI: 10.1007/s00209-021-02937-x.
- [BGP73] I. N. Bernstein, I. M. Gel’fand, and V. A. Ponomarev. “Coxeter Functors and Gabriel’s Theorem”. In: *Russian Mathematical Surveys* 28.2 (1973), pp. 17–32. DOI: 10.1070/RM1973v028n02ABEH001526.
- [DF78] P. Donovan and M. Freislich. “The indecomposable modular representations of certain groups with dihedral sylow subgroup”. In: *Mathematische Annalen* 238 (1978), pp. 207–216. DOI: 10.1007/BF01420248.
- [Ela15] A. Elagin. *On equivariant triangulated categories*. 2015. arXiv: 1403.7027 [math.AG].
- [ES92] K. Erdmann and A. Skowroński. “On Auslander-Reiten Components of Blocks and Self-Injective Biserial Algebras”. In: *Transactions of the American Mathematical Society* 330 (1992). DOI: 10.2307/2154159.
- [Gre74] J. A. Green. “Walking around the Brauer tree”. In: *Journal of the Australian Mathematical Society* 17.2 (1974), pp. 197–213. DOI: 10.1017/S1446788700016761.
- [Hap87] D. Happel. “On the derived category of a finite-dimensional algebra.” In: *Commentarii mathematici Helvetici* 62 (1987), pp. 339–389. URL: <http://eudml.org/doc/140089>.
- [Kau98] M. Kauer. “Derived equivalence of graph algebras”. In: *Contemporary Mathematics* 229, Trends in the representation theory of finite-dimensional algebras (1998), pp. 201–213.

- [Laz14] F. Lazarus. *Combinatorial Graphs and Surfaces from the Computational and Topological Viewpoint Followed by some notes on The Isometric Embedding of the square Flat Torus*. Mémoire d’habilitation à diriger des recherches. 2014.
- [LeM20] P. LeMeur. “Crossed products of Calabi-Yau algebras by finite groups”. In: *Journal of Pure and Applied Algebra* 224.10 (Oct. 2020), p. 106394. DOI: 10.1016/j.jpaa.2020.106394. URL: <https://doi.org/10.1016%2Fj.jpaa.2020.106394>.
- [Oku98] T. Okuyama. *Some examples of derived equivalent blocks of finite groups*. unpublished paper. 1998.
- [OZ22] S. Opper and A. Zvonareva. “Derived equivalence classification of Brauer graph algebras”. In: *Advances in Mathematics* 402 (2022), p. 108341. DOI: 10.1016/j.aim.2022.108341.
- [Ric89] J. Rickard. “Derived categories and stable equivalence”. In: *Journal of Pure and Applied Algebra* 61 (1989), pp. 303–317. DOI: 10.1016/0022-4049(89)90081-9.
- [Rog98] K. Roggenkamp. “Biserial algebras and graphs”. In: *Algebras and Modules II*. American Mathematical Society, 1998, pp. 481–496.
- [RR85] I. Reiten and C. Riedtmann. “Skew group algebras in the representation theory of artin algebras”. In: *Journal of Algebra* 92 (1985), pp. 224–282. URL: <https://api.semanticscholar.org/CorpusID:121741608>.
- [RS91] C. Riedtmann and A. H. Schofield. “On a simplicial complex associated with tilting modules”. English. In: *Commentarii Mathematici Helvetici* 66 (1991), pp. 70–78.
- [Sch15] S. Schroll. “Trivial extensions of gentle algebras and Brauer graph algebras”. In: *Journal of Algebra* 444 (2015), pp. 183–200. ISSN: 0021-8693. DOI: <https://doi.org/10.1016/j.jalgebra.2015.07.037>.
- [Sch18] S. Schroll. “Brauer Graph Algebras”. In: *Homological Methods, Representation Theory and Cluster Algebras*. 2018, pp. 177–223. ISBN: 978-3-319-74584-8. DOI: 10.1007/978-3-319-74585-5_6.
- [Sot24] V. Soto. “Generalized Kauer moves and derived equivalences of Brauer graph algebras”. In: *Journal of Algebra* 657 (2024), pp. 514–548. ISSN: 0021-8693. DOI: <https://doi.org/10.1016/j.jalgebra.2024.05.030>. URL: <https://www.sciencedirect.com/science/article/pii/S0021869324002862>.
- [WW85] B. Wald and J. Waschbüsch. “Tame biserial algebras”. In: *Journal of Algebra* 95.2 (1985), pp. 480–500. ISSN: 0021-8693. DOI: [https://doi.org/10.1016/0021-8693\(85\)90119-X](https://doi.org/10.1016/0021-8693(85)90119-X).
- [Zvo24] A. Zvonareva. *Derived equivalences of Brauer graph algebras*. 2024. arXiv: 2405.05602 [math.RT].

UNIVERSITÉ GRENOBLE ALPES, CNRS, INSTITUT FOURIER, 38610 GIÈRES
E-mail address : valentine.soto@univ-grenoble-alpes.fr