

Analytical solutions of higher-dimensional coupled system of nonlinear time-fractional diffusion-convection-wave equations

K.S. Priyendhu¹, P. Prakash^{1*}, M. Lakshmanan²

¹Department of Mathematics,

Amrita School of Engineering, Coimbatore-641112,

Amrita Vishwa Vidyapeetham, INDIA.

²Department of Nonlinear Dynamics,

Bharathidasan University,

Tiruchirappalli-620 024, INDIA.

E-mail ID's: vishnuindia89@gmail.com & p_prakash@cb.amrita.edu (P Prakash)

: priyendhusasikumar@gmail.com (K.S. Priyendhu)

: lakshman.cnld@gmail.com (M. Lakshmanan)

*-Corresponding Author

Abstract

This article develops how to generalize the invariant subspace method for deriving the analytical solutions of the multi-component $(N + 1)$ -dimensional coupled nonlinear time-fractional PDEs (NTFPDEs) in the sense of Caputo fractional-order derivative for the first time. Specifically, we describe how to systematically find different invariant product linear spaces with various dimensions for the considered system. Also, we observe that the obtained invariant product linear spaces help to reduce the multi-component $(N + 1)$ -dimensional coupled NTFPDEs into a system of fractional-order ODEs, which can then be solved using the well-known analytical methods. More precisely, we illustrate the effectiveness and importance of this developed method for obtaining a long list of invariant product linear spaces for the multi-component $(2 + 1)$ -dimensional coupled nonlinear time-fractional diffusion-convection-wave equations. In addition, we have shown how to find different kinds of generalized separable analytical solutions for a multi-component $(2 + 1)$ -dimensional coupled nonlinear time-fractional diffusion-convection-wave equations along with the initial and boundary conditions using invariant product linear spaces obtained. Finally, we provide appropriate graphical representations of some of the derived generalized separable analytical solutions with various fractional-order values.

Keywords: Invariant subspace method; multi-component $(N + 1)$ -dimensional coupled nonlinear PDEs; analytical solutions; fractional diffusion-convection-wave system; initial and boundary value problems; Laplace transformation technique.

1 Introduction

The theory of fractional calculus is the generalization of integer-order (classical) calculus, and it deals with the theoretical development and applications of arbitrary-order (fractional-order)

derivatives as well as arbitrary-order integrals, commonly known as differ-integrals [1–8]. In the literature, there are different approaches of generalization to the differ-integral definitions. The most prominent among them is the one based on Cauchy n -fold integration, and the widely used definitions of such arbitrary-order derivatives and integrals were proposed by Riemann, Liouville, Caputo and many others [6, 7]. Each of the generalizations has its importance in the sense of its advantages in different applications, especially in the fields of science and engineering [1–8]. Recent studies proved that arbitrary-order derivatives are more efficient and powerful tools for formulating the equations that govern physical processes, which can not only be described by their present state but also by their previous states in the history of the process. This power-law long-term memory property is the central essence of arbitrary-order derivatives, which helps them to describe the physical processes more accurately and precisely than the integer-order derivatives [1–9]. It is well-known that the Caputo fractional derivative considers the initial conditions that can be interpreted physically in science and engineering applications. Also, we know that the Caputo fractional derivative is being utilized to describe many real-world phenomena with a nonlocal nature that depend on their past interactions. In the literature, the Caputo fractional-order derivative is called as singular kernel derivative. Additionally, we note that the singular kernel Caputo fractional-order derivative violates some standard properties, which include the chain rule, Leibniz rule, and semi-group property. Consequently, the Caputo fractional-order derivative is said to possess nonstandard properties. These nonstandard properties allow singular kernel Caputo fractional-order derivative to possess long-term power-law memory property, as well as other fascinating properties such as nonlocal behavior, irreversible nature, and fractal-type properties [1, 2]. Consequently, fractional-order differential equations (FDEs) are used to formulate many real-world processes, primarily in the fields of physics and mechanics [1–9]. For example, one of the most prominent applications of the FDEs in the sense of Caputo fractional-order derivative can be seen in the mathematical formulation of the processes that govern anomalous diffusion problems in transport media like heat transfer in solids, thermo-elasticity without energy dissipation, chemical kinetics, biological population dynamics, vorticity in fluids, electro-dynamics, contamination transport in groundwater, collision in radioactive elements, sorption of organic compounds, ion exchange, geophysical-environmental mechanics, and many more [3–10]. So, in this work, we consider the Caputo fractional-order derivative for our investigation.

1.1 Motivation

The non-Fickian diffusion-convection-wave processes are governed by the two relations as follows:

$$(i) \text{ Conservation law of mass: } \quad \frac{\partial u}{\partial t} = -\nabla \cdot \mathbf{Q}, \&$$

$$(ii) \text{ Constitutive equation [10]: } \quad \mathbf{Q} = \begin{cases} \frac{1}{\Gamma(\alpha)} \frac{\partial}{\partial t} \int_0^t (t-y)^{\alpha-1} \mathbf{q} dy, & \text{if } \alpha \in (0, 1], \\ \frac{1}{\Gamma(\alpha-1)} \int_0^t (t-y)^{\alpha-2} \mathbf{q} dy, & \text{if } \alpha \in (1, 2], \end{cases}$$

where \mathbf{Q} is the matter flux such that $\mathbf{q} = -\kappa \nabla u + \mathbf{v}u$. Here κ and \mathbf{v} are the diffusion coefficient and the velocity of the particle with concentration $u = u(x_1, \dots, x_N, t)$, respectively, and $\nabla = \left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_N} \right)$ is the gradient operator. Thus, the nonlocal-time dependence of the matter flux described by a power-law long-tail memory kernel gives the time-fractional diffusion-convection-wave equation (TFDCWE) [10] in the form

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \kappa \Delta u - \mathbf{v} \cdot \nabla u, \alpha \in (0, 2], \quad (1.1)$$

where $\Delta = \sum_{r=1}^N \frac{\partial^2}{\partial x_r^2}$ denotes the Laplacian and $\frac{\partial^\alpha}{\partial t^\alpha}(\cdot)$ is the Caputo fractional-order derivative on the time variable, $t > 0$ of order α , $\alpha \in (0, 2]$, which is defined [6, 7] as

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \begin{cases} \frac{1}{\Gamma(k-\alpha)} \int_0^t (t-y)^{k-(\alpha+1)} \left(\frac{\partial^k u(x_1, \dots, x_N, y)}{\partial y^k} \right) dy, & \text{if } \alpha \in (k-1, k], k = 1, 2, \\ \frac{\partial^k u}{\partial t^k}, & \text{if } k = \alpha \in \{1, 2\}. \end{cases} \quad (1.2)$$

It is interesting to observe that for $\alpha = 1$, the equation (1.1) is equivalent to the classical integer-order diffusion-convection equation. For $v = 0$ (neglecting convective terms) and $\alpha = 2$, the equation (1.1) represents the classical wave equation. Additionally, when $v = 0$ in (1.1), it denotes sub-diffusion [5, 10–12] if $\alpha \in (0, 1)$ and super-diffusion [5, 10–12] if $\alpha \in (1, 2)$.

Motivated by the above studies, we formulate a multi-component $(2+1)$ -dimensional TFD-CWEs using the following governing equations:

$$(i) \text{ Conservation law of mass: } \quad \frac{\partial u_s}{\partial t} = -\nabla \cdot \mathbf{Q}_s,$$

(ii) Constitutive equation:

$$\mathbf{Q}_s = \begin{cases} \frac{1}{\Gamma(\alpha_s)} \frac{\partial}{\partial t} \int_0^t (t-y)^{\alpha_s-1} (-\kappa_s \nabla u_s(x_1, x_2, y) + \mathbf{v}_s u_s(x_1, x_2, y)) dy, & \text{if } \alpha_s \in (0, 1], \\ \frac{1}{\Gamma(\alpha_s-1)} \int_0^t (t-y)^{\alpha_s-2} (-\kappa_s \nabla u_s(x_1, x_2, y) + \mathbf{v}_s u_s(x_1, x_2, y)) dy, & \text{if } \alpha_s \in (1, 2], \end{cases}$$

where \mathbf{Q}_s is the matter flux, κ_s and \mathbf{v}_s are the diffusion coefficient and the velocity of the s -th particle with concentration $u_s = u_s(x_1, x_2, t)$, $s = 1, 2$, respectively, and $\nabla = \left(\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2} \right)$ is the gradient operator. Substituting (ii) in (i), we obtain the following multi-component $(2 + 1)$ -dimensional TFDCWEs:

$$\frac{\partial^{\alpha_s} u_s}{\partial t^{\alpha_s}} = -\kappa_s \Delta u_s(x_1, x_2, t) + \nabla \cdot \mathbf{v}_s u_s(x_1, x_2, t), \alpha_s \in (0, 2], s = 1, 2, \quad (1.3)$$

where $\Delta = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2}$ denotes the Laplacian operator and $\frac{\partial^{\alpha_s}}{\partial t^{\alpha_s}}(\cdot)$ is the Caputo fractional derivative (1.2). Recent studies show that the significant advantage of diffusion models with time-fractional derivatives is that, in contrast to the classical integer-order diffusion models, they can accurately describe the nonlinear time-dependency of anomalous diffusion processes. Multi-component $(2 + 1)$ -dimensional time-fractional diffusion equations, widely used in the literature, describe many complex physical models involving more than one diffusive particle or when both heat conduction and diffusion co-occur. For example, Zhang et al. [13] have studied the coupled heat conduction and moisture diffusion in a hollow cylindrical pipe in hygrothermoelastic environments using the following multi-component $(2+1)$ -dimensional time-fractional PDEs

$$\begin{aligned} \frac{\partial^{\alpha_1} T}{\partial t^{\alpha_1}} - \beta_1 \frac{\partial^{\alpha_1} C}{\partial t^{\alpha_1}} &= \kappa_1 \Delta T, \\ \frac{\partial^{\alpha_2} C}{\partial t^{\alpha_2}} - \beta_2 \frac{\partial^{\alpha_2} T}{\partial t^{\alpha_2}} &= \kappa_2 \Delta C, \alpha_s \in (0, 2], s = 1, 2, \end{aligned} \quad (1.4)$$

where $T = T(x_1, x_2, t)$ represents the temperature, and $C = C(x_1, x_2, t)$ is the concentration of water vapor in a porous media. In the above system (1.4), $\beta_s = \frac{\gamma_s a_s}{\gamma_s a_j + c_s}$, $j \neq s, j, s = 1, 2$, $\kappa_1 = \frac{\lambda_1}{\gamma_2(c_1 + \gamma_1 a_2)}$, and $\kappa_2 = \frac{\lambda_2 c_2}{c_2 + \gamma_2 a_1}$, where the arbitrary constants $\gamma_1, \gamma_2, c_1, c_2, \lambda_1$ and λ_2 represent the amount of heat released, the density of the material, the specific heat capacity at constant pressure, the volume fraction of moisture, heat conduction and diffusion coefficients, respectively, and $a_s, s = 1, 2$, are material constants. We note that under the substitution $T - \beta_1 C = u_1$ and $C - \beta_2 T = u_2$, the above system (1.4) gets reduced into the following multi-component $(2+1)$ -dimensional coupled time-fractional diffusion-wave equations:

$$\begin{aligned} \frac{\partial^{\alpha_1} u_1}{\partial t^{\alpha_1}} &= \kappa_1 \Delta (\tau_0 u_1 + \tau_1 u_2), \\ \frac{\partial^{\alpha_2} u_2}{\partial t^{\alpha_2}} &= \kappa_2 \Delta (\tau_2 u_1 + \tau_0 u_2), \alpha_s \in (0, 2], s = 1, 2, \end{aligned} \quad (1.5)$$

where $\tau_0 = \frac{1}{1 - \beta_1 \beta_2}$, and $\tau_s = \frac{\beta_s}{1 - \beta_1 \beta_2}$, $s = 1, 2$. Another physical application of the above type of system (1.3) was discussed by Yang et al. [14] to describe the anomalous diffusion of

sodium (Na⁺) and potassium (K⁺) ions in a neuronal phenomenon as follows:

$$\frac{\partial^\alpha u_s}{\partial t^\alpha} = \sum_{i=1}^2 \frac{\partial}{\partial x_i} \left[(\tau_1 u_s + \tau_0) \frac{\partial u_1}{\partial x_i} + (\tau_1 u_s + \tau_0) \frac{\partial u_2}{\partial x_i} \right] - I_s, \alpha \in (0, 1], s = 1, 2, \quad (1.6)$$

where $\frac{\partial^\alpha}{\partial t^\alpha}(\cdot)$ is the Caputo fractional derivative (1.2), τ_0, τ_1 are known constants, $u_s = u_s(x_1, x_2, t)$, $s = 1, 2$, represent the concentration of Na⁺ and K⁺ ions with the current density $I_s, s = 1, 2$. They have developed the Jacobi special collocation method for the above system (1.6), also known as the coupled time-fractional Nernst-Planck equations [14], to find an approximate solution of the above system (1.6) along with the reduction $r = \sqrt{x_1^2 + x_2^2}$.

It is well-known that the essential characteristics of a physical process can be analyzed using its long-term asymptotic behavior and qualitative-quantitative properties. These properties are described exactly by the analytical solutions of the corresponding governing FDEs. However, due to the above-mentioned mysterious properties of fractional derivatives, the task of finding analytical (exact) solutions for nonlinear FDEs is difficult and challenging, particularly for nonlinear fractional PDEs. Further, there is still no well-defined analytical method available to find analytical solutions to FDEs in general. Due to the rigorous investigations in recent years, some numerical and analytical methods have been developed to find solutions for different classes of FDEs which include the Adomain decomposition method [15–17], the exp-function method [18, 19], the sub-equation method [20], the generalized differential transform method [21], the hybrid method [22], the separation of variable methods [23–26], the homotopy analysis method [27], the homotopy perturbation method [28], the Lie symmetry analysis method [11, 12, 29–37], and the invariant subspace method [38–61], to name a few.

Among these analytical methods, the invariant subspace method has proven to be an effective and powerful analytic tool for finding generalized separable analytical solutions for many classes of nonlinear fractional PDEs. This method was initially proposed by Galaktionov and Svirshchevskii [62], which is also addressed as the generalized separation of the variables method or the Titov-Galaktionov method [54], for finding generalized separable analytical solutions of nonlinear integer-order evolution PDEs. Later, the potential of this method was further investigated by Ma [63], Zhu and Qu [64], and many others [54, 65–71] to solve many important classes of nonlinear coupled and scalar integer-order PDEs. In 2013, Gazizov and Kasatkin [38] extended the invariant subspace method to solve NTFPDEs. Recently, this method was extended to solve many classes of nonlinear scalar fractional PDEs in (1+1)-dimensions [39–49, 51–54, 56]. Moreover, this method has been successfully extended and applied to find generalized separable analytical solutions for a (1 + 1)-dimensional coupled nonlinear fractional PDEs [31, 46–48, 50].

In 2019, Sangita et al. [55] applied this method to find generalized separable analytical solutions for the $(2 + 1)$ -dimensional coupled NTFPDEs.

1.2 Novelty of our work

We believe that no one has developed the invariant subspace method to find generalized separable analytical solutions for multi-component $(N + 1)$ -dimensional coupled NTFPDEs as far as we know. So, we describe for the first time the generalization of the invariant subspace method for deriving different kinds of analytical solutions for multi-component $(N + 1)$ -dimensional coupled NTFPDEs. The main aim of our work is to systematically develop the invariant subspace method for obtaining different types of invariant product linear spaces and analytical solutions for multi-component coupled NTFPDEs mentioned above. Here, it is noteworthy that the obtained finite-dimensional invariant product linear spaces help to reduce the multi-component $(N + 1)$ -dimensional coupled NTFPDEs into a system of fractional ODEs (FODEs). More precisely, the importance of the developed invariant subspace method has been discussed through finding the analytical solutions in the generalized separable form for the following initial value problems of the multi-component $(2 + 1)$ -dimensional coupled nonlinear TFDCWEs

$$\left(\frac{\partial^{\alpha_1} u_1}{\partial t^{\alpha_1}}, \frac{\partial^{\alpha_2} u_2}{\partial t^{\alpha_2}} \right) = \mathbf{F}(\mathbf{U}) \equiv (F_1(u_1, u_2), F_2(u_1, u_2)), \alpha_1, \alpha_2 \in (0, 2], \quad (1.7)$$

together with initial conditions:

$$\begin{aligned} (i) \quad & u_s(x_1, x_2, 0) = \omega_s(x_1, x_2) \text{ if } \alpha_s \in (0, 1], s = 1, 2, \\ (ii) \quad & u_s(x_1, x_2, 0) = \omega_s(x_1, x_2), \quad \frac{\partial u_s}{\partial t} \Big|_{t=0} = \vartheta_s(x_1, x_2) \text{ if } \alpha_s \in (1, 2], s = 1, 2, \end{aligned} \quad (1.8)$$

where the components $F_s(u_1, u_2)$, $s = 1, 2$, are considered as follows:

$$\begin{aligned} F_s(u_1, u_2) = & \sum_{i=1}^2 \frac{\partial}{\partial x_i} \left(\beta_{si}(u_1, u_2) \frac{\partial u_1}{\partial x_i} + \rho_{si}(u_1, u_2) \frac{\partial u_2}{\partial x_i} \right) \\ & + \sum_{i=1}^2 \left(\zeta_{si}(u_1, u_2) \frac{\partial u_1}{\partial x_i} + v_{si}(u_1, u_2) \frac{\partial u_2}{\partial x_i} \right). \end{aligned} \quad (1.9)$$

Here $\mathbf{U} = (u_1, u_2)$, $u_s = u_s(x_1, x_2, t)$, $x_i \in \mathbb{R}$, $t > 0$, the functions $\beta_{si}(u_1, u_2)$, $\rho_{si}(u_1, u_2)$ denote the diffusion coefficients and $\zeta_{si}(u_1, u_2)$, $v_{si}(u_1, u_2)$, $s, i = 1, 2$, represent the convective terms of the physical system. We observe that the obtained system (1.5) is a particular case of the considered coupled NTFPDEs (1.7). Additionally, we would like to point out that the above system (1.6) with negligible current density (that is, $I_s = 0$) is a particular case of the discussed coupled NTFPDEs (1.7). In the literature, finding the analytical solutions of the coupled diffusion-convection-wave equations has gained considerable attention because of its widespread

application as a mathematical formulation for the dynamics of long-range interactions between more than one element or species in physical processes like convection-dominated transport media problems. Some important classes of coupled diffusion-convection equations include the coupled Burgers equations, the hydrodynamic model for semiconductor device simulation, magneto-hydrodynamics, shallow water equations, and the Navier-stokes equations [72–74]. In [75], LDG (local discontinuous Galerkin) method was used efficiently to study analytical solutions for the integer-order coupled diffusion-convection equations [75] in the form

$$\frac{\partial \mathbf{U}}{\partial t} = \nabla \cdot \mathbf{F}(\mathbf{U}, D\mathbf{U}), \quad (1.10)$$

where $\mathbf{U} = (u_1, u_2, \dots, u_m)$, $u_s = u_s(x_1, x_2, \dots, x_N, t)$, $x_r \in \mathbb{R}$, $t > 0$, $r = 1, 2, \dots, N$, $s = 1, 2, \dots, m$, $N, m \in \mathbb{N}$, ∇ is the gradient and D is the total derivative of \mathbf{U} of order one. Note that for $\alpha = 1$, the above-coupled equations (1.7) belongs to the class of given coupled diffusion-convection equations (1.10) with $m = N = 2$.

1.3 Structure of the article

This article is structured as follows: In section 2, we develop the invariant subspace method to identify different types of invariant product linear spaces and generalized separable analytical solutions for multi-component $(N + 1)$ -dimensional coupled NTFPDEs. In addition, we systematically discuss how to construct N -different types of invariant product linear spaces for the system mentioned above with their invariance criteria. Section 3 briefly describes how to systematically obtain various kinds of invariant product linear spaces to the multi-component $(2 + 1)$ -dimensional coupled nonlinear TFDCWEs (1.7) with cubic power-law nonlinearity using the developed invariant subspace method. Also, we present a comprehensive explanation of how to find generalized separable analytical solutions for the considered coupled equations (1.7) equipped with initial and boundary conditions. Additionally, we show the graphical representations of some of the obtained generalized separable analytical solutions with various α -values. Finally, section 4 discusses a brief review and concluding remarks on the developed method with their obtained results.

2 Multi-component $(N + 1)$ -dimensional coupled NTFPDEs: Invariant subspace method

This section presents the systematic procedure for deriving different kinds of analytical solutions of the multi-component $(N + 1)$ -dimensional coupled NTFPDEs using the invariant subspace

method. Let us consider the generalized multi-component $(N + 1)$ -dimensional coupled NTF-PDEs as follows:

$$\begin{aligned} \frac{\partial^{\alpha_s} u_s}{\partial t^{\alpha_s}} &= A_s(\mathbf{U}) \\ &= A_s \left(x_1, x_2, \dots, x_N, u_1, \dots, u_m, \frac{\partial u_1}{\partial x_1}, \dots, \frac{\partial^{q_s} u_1}{\partial x_1^{i_1} \partial x_2^{i_2} \dots \partial x_N^{i_N}}, \dots, \right. \\ &\quad \left. \frac{\partial u_m}{\partial x_1}, \dots, \frac{\partial^{q_s} u_m}{\partial x_1^{i_1} \partial x_2^{i_2} \dots \partial x_N^{i_N}} \right), \alpha_s > 0, s = 1, 2, \dots, m, \end{aligned} \quad (2.1)$$

where $A_s(\mathbf{U})$ is the q_s -th order sufficiently smooth differential operator of the m -component vector function $\mathbf{U} = (u_1, \dots, u_m)$, $u_s = u_s(x_1, \dots, x_N, t)$, $\sum_{j=1}^N i_j = q_s$, $i_j \in \{0, 1, \dots, q_s\}$, $q_s \in \mathbb{N}$, $j = 1, 2, \dots, N$, and $\frac{\partial^{\alpha_s}}{\partial t^{\alpha_s}}(\cdot)$ denotes the Caputo derivative (1.2) of order $\alpha_s > 0$, $s = 1, 2, \dots, m$.

In this work, we develop the invariant subspace method for the above-coupled NTFPDEs (2.1) to derive possible types of analytical solutions in the generalized separable form based on $w_{s,i}(x_1, x_2, \dots, x_N)$ as $\mathbf{U} = (u_1, u_2, \dots, u_m)$ with s -th component function

$$u_s := u_s(x_1, \dots, x_N, t) = \sum_{i=1}^{k_s} C_{s,i}(t) w_{s,i}(x_1, \dots, x_N), k_s \in \mathbb{N}, s = 1, 2, \dots, m, \quad (2.2)$$

where the unknown functions $C_{s,i}(t)$ and $w_{s,i}(x_1, x_2, \dots, x_N)$ are to be obtained that depend on the considered coupled NTFPDEs (2.1) and its invariant product linear spaces.

Let us first define the product linear space \mathbf{Y}_n as

$$\begin{aligned} \mathbf{Y}_n &= Y_{1,n_1} \times Y_{2,n_2} \times \dots \times Y_{m,n_m} \\ &= \{\mathbf{y} = (y_1, y_2, \dots, y_m) \mid y_s \in Y_{s,n_s}, s = 1, 2, \dots, m\}, \end{aligned} \quad (2.3)$$

along with linear component spaces Y_{s,n_s} of dimension $n_s (= \dim(Y_{s,n_s}) < \infty)$, for each $s = 1, 2, \dots, m$, under the component-wise addition '+', and scalar multiplication '.' as

$$\mathbf{y} + \mathbf{z} = (y_1 + z_1, \dots, y_m + z_m), \text{ and } \mu \cdot \mathbf{y} = (\mu \cdot y_1, \dots, \mu \cdot y_m),$$

respectively, where $\mathbf{y} = (y_1, \dots, y_m)$, $\mathbf{z} = (z_1, \dots, z_m) \in \mathbf{Y}_n$, and $\mu \in \mathbb{R}$. The following proposition explains how to determine the dimension of the above-mentioned product linear space (2.3).

Theorem 2.1. [76] *Let $\mathbf{Y}_n = Y_{1,n_1} \times Y_{2,n_2} \times \dots \times Y_{m,n_m}$ given in (2.3) be the product linear space with the n_s -dimensional linear component spaces Y_{s,n_s} , where $n_s (= \dim(Y_{s,n_s})) < \infty$, $s = 1, 2, \dots, m$. Then, the dimension of the product linear space \mathbf{Y}_n is $n = n_1 + n_2 + \dots + n_m$. That is, $\dim(\mathbf{Y}_n) = \sum_{s=1}^m \dim(Y_{s,n_s})$.*

Next, we discuss the invariant product linear spaces for the given coupled NTFPDEs (2.1) with their invariant criteria. Also, we present the systematic algorithmic study to derive generalized separable analytical solutions of the form (2.2) using the invariant product linear spaces.

2.1 Finding invariant product linear spaces and generalized separable analytical solutions of the coupled NTFPDEs (2.1)

Let us consider the vector differential operator $\hat{\mathbf{A}}[\mathbf{U}] = (A_1(\mathbf{U}), A_2(\mathbf{U}), \dots, A_m(\mathbf{U}))$, where $A_s(\mathbf{U})$ is the nonlinear differential operator given in (2.1) for $s = 1, 2, \dots, m$. Now, we introduce the product linear space $\hat{\mathbf{W}}_k$ as

$$\begin{aligned} \hat{\mathbf{W}}_k &= W_{1,k_1} \times W_{2,k_2} \times W_{3,k_3} \times \dots \times W_{m,k_m} \\ &= \left\{ \mathbf{w}(x_1, \dots, x_N) = (w_1(x_1, \dots, x_N), \dots, w_m(x_1, \dots, x_N)) \mid \right. \\ &\quad \left. w_s(x_1, \dots, x_N) \in W_{s,k_s}, s = 1, 2, \dots, m \right\}, \end{aligned} \quad (2.4)$$

where W_{s,k_s} are k_s -dimensional linear component spaces for $s = 1, 2, \dots, m$. Additionally, from the Proposition 2.1, we note that $\dim(\hat{\mathbf{W}}_k) = k = \sum_{s=1}^m k_s (< \infty)$. Thus, we say that $\hat{\mathbf{W}}_k$ is an invariant finite-dimensional product linear space for the nonlinear vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$ if either one of the following invariant conditions holds:

- (a) For every $\mathbf{U} = (u_1, \dots, u_m) \in \hat{\mathbf{W}}_k$, $\hat{\mathbf{A}}[\mathbf{U}] = (A_1(\mathbf{U}), \dots, A_m(\mathbf{U})) \in \hat{\mathbf{W}}_k$ or $\hat{\mathbf{A}}[\hat{\mathbf{W}}_k] \subseteq \hat{\mathbf{W}}_k$.
- (b) If $\mathbf{U} \in \hat{\mathbf{W}}_k$, then $A_s(\mathbf{U}) \in W_{s,k_s}$, or $A_s(\hat{\mathbf{W}}_k) = A_s(W_{1,k_1} \times W_{2,k_2} \times \dots \times W_{m,k_m}) \subseteq W_{s,k_s}, \forall u_s \in W_{s,k_s}, s = 1, 2, \dots, m$.

Now, let $B_s = \{w_{s,1}(x_1, x_2, \dots, x_N), w_{s,2}(x_1, x_2, \dots, x_N), \dots, w_{s,k_s}(x_1, x_2, \dots, x_N)\}$ be a basis for the linear component spaces W_{s,k_s} , then $W_{s,k_s}, s = 1, 2, \dots, m$, can be written as follows:

$$\begin{aligned} W_{s,k_s} &= \text{Span}(B_s) \\ &= \left\{ w_s(x_1, \dots, x_N) = \sum_{i=1}^{k_s} \mu_{s,i} w_{s,i}(x_1, \dots, x_N) \mid \mu_{s,i} \in \mathbb{R}, i = 1, 2, \dots, k_s \right\}. \end{aligned} \quad (2.5)$$

Thus, using the basis B_s of the linear component spaces $W_{s,k_s}, s = 1, 2, \dots, m$, we define the basis $\hat{\mathbf{B}}$ for the given finite-dimensional product linear space $\hat{\mathbf{W}}_k$ as follows:

$$\begin{aligned} \hat{\mathbf{B}} &= \left\{ (w_{1,1}, 0, \dots, 0), \dots, (w_{1,k_1}, 0, \dots, 0), \dots, (0, \dots, 0, w_{m,1}), \dots, (0, \dots, 0, w_{m,k_m}) \mid \right. \\ &\quad \left. w_{s,i} = w_{s,i}(x_1, x_2, \dots, x_N) \in B_s, i = 1, 2, \dots, k_s, s = 1, 2, \dots, m \right\}. \end{aligned}$$

Hence, the invariant criteria for the invariant product linear space $\hat{\mathbf{W}}_k$ of $\hat{\mathbf{A}}(\mathbf{U})$ can be obtained as follows:

$$\begin{aligned} \hat{\mathbf{A}}(\mathbf{U}) &= (A_1(\mathbf{U}), \dots, A_m(\mathbf{U})) \text{ with} \\ A_s(\mathbf{U}) &= \sum_{i=1}^{k_s} \Lambda_{s,i}(\lambda_{1,1}, \dots, \lambda_{m,k_m}) w_{s,i}(x_1, x_2, \dots, x_N), \end{aligned} \quad (2.6)$$

whenever $\mathbf{U} = (u_1, u_2, \dots, u_m)$ along with $u_s = \sum_{i=1}^{k_s} \lambda_{s,i} w_{s,i}(x_1, x_2, \dots, x_N)$, $\lambda_{s,i} \in \mathbb{R}$, where the coefficients $\Lambda_{s,i}(\cdot)$, $i = 1, 2, \dots, k_s$, $s = 1, 2, \dots, m$, are obtained from the expansion corresponding to the basis $\hat{\mathbf{B}}$. Thus, we can obtain the analytical solution (2.2) for the considered coupled NTFPDEs (2.1) whenever the invariant criteria (2.6) hold.

Now, we state how to reduce the given multi-component $(N + 1)$ -dimensional coupled NTFPDEs (2.1) into a system of time-fractional ODEs under the invariant product linear space (2.4).

Theorem 2.2. *Let $\hat{\mathbf{A}}[\mathbf{U}] = (A_1(\mathbf{U}), A_2(\mathbf{U}), \dots, A_m(\mathbf{U}))$ be the vector differential operator, which admits the invariant finite-dimensional product linear space $\hat{\mathbf{W}}_k = W_{1,k_1} \times W_{2,k_2} \times \dots \times W_{m,k_m}$, where W_{s,k_s} are the k_s -dimensional linear component spaces defined in (2.5) for $s = 1, 2, \dots, m$. Then, there exists an analytical solution for the given multi-component $(N + 1)$ -dimensional coupled NTFPDEs (2.1) in the generalized separable form $\mathbf{U} = (u_1, \dots, u_m)$ with $u_s = u_s(x_1, \dots, x_N, t)$ as*

$$u_s = \sum_{i=1}^{k_s} C_{s,i}(t) w_{s,i}(x_1, x_2, \dots, x_N), \quad (2.7)$$

and the functions $C_{s,i}(t)$ satisfying the following system of time-fractional ODEs

$$\frac{d^{\alpha_s} C_{s,i}(t)}{dt^{\alpha_s}} = \Phi_{s,i} \left(C_{1,1}(t), \dots, C_{1,k_1}(t), \dots, C_{m,1}(t), \dots, C_{m,k_m}(t) \right), \quad (2.8)$$

for all $i = 1, 2, \dots, k_s$, $s = 1, 2, \dots, m$, where $\Phi_{s,i}(\cdot)$ are the functions based on the expansion corresponding to the basis $\hat{\mathbf{B}}$.

Proof. Let us assume that $\hat{\mathbf{W}}_k$ is an invariant finite-dimensional product linear space of the considered nonlinear vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$. Then, we obtain the invariant criteria given in (2.6). From this invariant criteria (2.6), we get $\mathbf{U} = (u_1, u_2, \dots, u_m)$ with u_s as given in (2.7), and hence we obtain

$$\begin{aligned} A_s(\mathbf{U}) &= A_s \left(\sum_{i=1}^{k_1} C_{1,i}(t) w_{1,i}(x_1, \dots, x_N), \dots, \sum_{i=1}^{k_m} C_{m,i}(t) w_{m,i}(x_1, \dots, x_N) \right) \\ &= \sum_{i=1}^{k_s} \Phi_{s,i} \left(C_{1,1}(t), \dots, C_{1,k_1}(t), \dots, C_{m,1}(t), \dots, C_{m,k_m}(t) \right) w_{s,i}(x_1, \dots, x_N), \end{aligned} \quad (2.9)$$

where $\Phi_{s,i}(\cdot)$, $i = 1, 2, \dots, k_s$, $s = 1, 2, \dots, m$, are the functions based on expansion corresponding to the basis $\hat{\mathbf{B}}$.

Now, we consider the given coupled NTFPDEs (2.1) having the form

$$\frac{\partial^{\alpha_s} u_s}{\partial t^{\alpha_s}} = A_s(\mathbf{U}), \quad \alpha_s > 0, \quad s = 1, 2, \dots, m. \quad (2.10)$$

Then, by the virtue of the linear property of the Caputo fractional-order derivative and, for every $\mathbf{U} = (u_1, u_2, \dots, u_m)$ with u_s as given in (2.7), we obtain the left-hand side of the given coupled NTFPDEs (2.10) as

$$\begin{aligned} \frac{\partial^{\alpha_s} u_s}{\partial t^{\alpha_s}} &= \frac{\partial^{\alpha_s}}{\partial t^{\alpha_s}} \left(\sum_{i=1}^{k_s} C_{s,i}(t) w_{s,i}(x_1, x_2, \dots, x_N) \right) \\ &= \sum_{i=1}^{k_s} \left(\frac{\partial^{\alpha_s} C_{s,i}(t)}{\partial t^{\alpha_s}} \right) w_{s,i}(x_1, x_2, \dots, x_N), s = 1, 2, \dots, m, \end{aligned} \quad (2.11)$$

and the right-hand side of the given coupled NTFPDEs (2.10) reduces into the form (2.9). Thus, substituting the obtained equations (2.9) and (2.11) into the given NTFPDEs (2.10), we get

$$\sum_{i=1}^{k_s} \left[\frac{d^{\alpha_s} C_{s,i}(t)}{dt^{\alpha_s}} - \Phi_{s,i}(C_{1,1}(t), \dots, C_{1,k_1}(t), \dots, C_{m,1}(t), \dots, C_{m,k_m}(t)) \right] w_{s,i} = 0,$$

where $w_{s,i} = w_{s,i}(x_1, \dots, x_N)$ for $s = 1, 2, \dots, m$. Since the functions $w_{s,i} \in B_s$, $i = 1, 2, \dots, k_s$ are linearly independent for $s = 1, 2, \dots, m$, we find that the given coupled NTFPDEs (2.10) reduces into the system of time-fractional ODEs as given in (2.8) whenever $\mathbf{U} = (u_1, \dots, u_m)$ with u_s as given in (2.7), which completes the proof. \square

Note that Theorem 2.2 provides a necessary condition for the existence of the generalized separable analytical solution (2.2) for the given coupled NTFPDEs (2.1). Thus, we can conclude that whenever $\widehat{\mathbf{W}}_k$ is an invariant product linear space of the considered nonlinear vector differential operator $\widehat{\mathbf{A}}(\mathbf{U})$, then we can find the generalized separable analytical solution $\mathbf{U} = (u_1, \dots, u_m)$ with $u_s = u_s(x_1, \dots, x_N, t)$ given in (2.2) for the given multi-component $(N + 1)$ -dimensional coupled NTFPDEs (2.1).

It is interesting to observe that we can derive the following N -different types of generalized separable analytical solutions for the multi-component $(N + 1)$ -dimensional coupled NTFPDEs (2.1):

- The most generalized separable analytical solution for the coupled NTFPDEs (2.1) is obtained as $\mathbf{U} = (u_1, u_2, \dots, u_m)$ with components

$$u_s = \sum_{r_1, \dots, r_N} C_{s,r_1, \dots, r_N}(t) \prod_{i=1}^N \xi_{s,r_i}^i(x_i), \quad (2.12)$$

where $r_i = 1, 2, \dots, k_{s,i}$, $i = 1, 2, \dots, N$, $s = 1, 2, \dots, m$.

- Other $(N-1)$ types of generalized separable analytical solutions for the coupled NTFPDEs (2.1) are obtained as $\mathbf{U} = (u_1, u_2, \dots, u_m)$ with components

$$\begin{aligned}
u_s = & C_{s,0}(t) + \sum_{i=1}^N \sum_{r_i=1}^{k_{s,i}-1} C_{s,r_i}^i(t) \xi_{s,r_i}^i(x_i) + \dots \\
& + \sum_{i_1, \dots, i_{N-p}} \left(\sum_{r_{i_1}, \dots, r_{i_{N-p}}} C_{s,r_{i_1}, \dots, r_{i_{N-p}}}^{i_1, \dots, i_{N-p}}(t) \prod_{j=1}^{N-p} \xi_{s,r_{i_j}}^{i_j}(x_{i_j}) \right), \tag{2.13}
\end{aligned}$$

where $r_{i_j} = 1, 2, \dots, k_{s,i_j} - 1, i_j = 1, 2, \dots, N, j = 1, 2, \dots, N - p, i_1 < i_2 < \dots < i_{N-p}, s = 1, 2, \dots, m, p = 1, 2, \dots, N - 1$.

Note that the unknown functions $\xi_{s,r_i}^i(x_i), \xi_{s,r_{i_j}}^{i_j}(x_{i_j}), C_{s,0}(t), C_{s,r_i}^i(t), \dots, C_{s,r_{i_1}, \dots, r_{i_{N-p}}}^{i_1, \dots, i_{N-p}}(t)$, and $C_{s,r_1, \dots, r_N}(t)$ are to be determined using the considered coupled NTFPDEs (2.1) and corresponding invariant finite-dimensional product linear spaces. The details of finding the above analytical solutions (2.12)-(2.13) using the various kinds of invariant product linear spaces that are discussed in the following subsection.

2.2 Determining different types of linear component spaces W_{s,k_s} with their invariant product linear space $\hat{\mathbf{W}}_k$ and analytical solutions for the given multi-component NTFPDEs (2.1)

This subsection presents the details of determining the N -different types of invariant product linear spaces $\hat{\mathbf{W}}_k$ and generalized separable analytical solutions for the given multi-component coupled NTFPDEs (2.1). Let us first explain the details of the systematic procedure to determine the different types of invariant product linear spaces $\hat{\mathbf{W}}_k$ for the vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$. Before proceeding further, we explain what kind of notations are used for linear spaces in the remaining sections, which are discussed in the following for easy understanding.

(I) W_{s,k_s}^p represents the s -th component type- p linear component space with dimension k_s for $s = 1, 2, \dots, m$.

(II) $\hat{\mathbf{W}}_k^p$ represents the type- p product linear vector space with dimension k .

Now, we aim to construct the linear component spaces W_{s,k_s}^p of the type- p product linear space $\hat{\mathbf{W}}_k^p, p = 1, 2, \dots, N$, using the system of $k_{s,i}$ -th order linear homogeneous ODEs:

$$\begin{aligned}
\mathbf{L}_{s,k_s,i}^i[y_{s,i}(x_i)] \equiv & \frac{d^{k_{s,i}} y_{s,i}(x_i)}{dx_i^{k_{s,i}}} + \varphi_{s,k_s,i-1}^i(x_i) \frac{d^{k_{s,i}-1} y_{s,i}(x_i)}{dx_i^{k_{s,i}-1}} + \dots \\
& + \varphi_{s,0}^i(x_i) y_{s,i}(x_i) = 0, \tag{2.14}
\end{aligned}$$

where $k_{s,i} \in \mathbb{N}$, and $\varphi_{s,j}^i(x_i), j = 0, 1, \dots, k_{s,i} - 1$ are continuous functions of x_i . Here, we consider the basis of the solution space (or the fundamental set of solutions) for given linear homogeneous ODEs (2.14) as

$$H_{s,i} = \left\{ \xi_{s,1}^i(x_i), \xi_{s,2}^i(x_i), \dots, \xi_{s,k_{s,i}}^i(x_i) \right\}, \quad i = 1, 2, \dots, N, s = 1, 2, \dots, m. \quad (2.15)$$

Next, we provide how to construct the linear component spaces W_{s,k_s}^p for $p = 1, 2, \dots, N$, using the basis of the solution space $H_{s,i}$ given in (2.15). Additionally, we explain the details of determining N -different types of invariant product linear spaces $\hat{\mathbf{W}}_k^p, p = 1, 2, \dots, N$ and the generalized separable analytical solutions given in (2.12)-(2.13) for the given coupled NTFPDEs (2.1) using $\hat{\mathbf{W}}_k^p$.

2.2.1 Type-1 invariant product linear space $\hat{\mathbf{W}}_k^1$ for the given vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$:

Let us **consider** the linear component spaces W_{s,k_s}^1 of the type-1 product linear space $\hat{\mathbf{W}}_k^1$ as the linear span of the set

$$B_s^1 = \left\{ \prod_{i=1}^N \xi_{s,r_i}^i(x_i) \mid r_i = 1, \dots, k_{s,i}, i = 1, 2, \dots, N \right\}, \quad s = 1, 2, \dots, m, \quad (2.16)$$

where the functions $\xi_{s,r_i}^i(x_i) \in H_{s,i}$ is given in (2.15). Using the above set B_s^1 , the type-1 linear component space W_{s,k_s}^1 can be defined as

$$\begin{aligned} W_{s,k_s}^1 &= \text{Span} (B_s^1) \\ &= \left\{ w_s(x_1, \dots, x_N) = \sum_{r_1, \dots, r_N} \mu_{s,r_1, \dots, r_N} \prod_{i=1}^N \xi_{s,r_i}^i(x_i) \mid \mu_{s,r_1, \dots, r_N} \in \mathbb{R}, \right. \\ &\quad \left. r_i = 1, \dots, k_{s,i}, i = 1, \dots, N \right\}. \end{aligned} \quad (2.17)$$

Thus, the type-1 product linear space $\hat{\mathbf{W}}_k^1$ with linear component spaces $W_{s,k_s}^1, s = 1, 2, \dots, m$, given in (2.17) can be written as follows:

$$\begin{aligned} \hat{\mathbf{W}}_k^1 &= W_{1,k_1}^1 \times W_{2,k_2}^1 \times W_{3,k_3}^1 \times \dots \times W_{m,k_m}^1 \\ &= \left\{ \mathbf{w}(x_1, \dots, x_N) = (w_1(x_1, \dots, x_N), \dots, w_m(x_1, \dots, x_N)) \mid \right. \\ &\quad w_s(x_1, \dots, x_N) = \sum_{r_1, \dots, r_N} \mu_{s,r_1, \dots, r_N} \prod_{i=1}^N \xi_{s,r_i}^i(x_i), \xi_{s,r_i}^i(x_i) \in H_{s,i}, \\ &\quad \left. \mu_{s,r_1, \dots, r_N} \in \mathbb{R}, r_i = 1, \dots, k_{s,i}, i = 1, \dots, N, s = 1, 2, \dots, m \right\}. \end{aligned} \quad (2.18)$$

Next, we show that the set B_s^1 is a basis for the linear component space W_{s,k_s}^1 , for all $s = 1, 2, \dots, m$.

Proposition 2.1. Consider the set B_s^1 given in (2.16) with $\xi_{s,r_i}^i(x_i) \in H_{s,i}$ for $i = 1, \dots, N$, and $s = 1, 2, \dots, m$, where $H_{s,i}$ given in (2.15) is the basis of the solution spaces for the given linear homogeneous ODEs (2.14). Then, the set B_s^1 forms a basis for the linear component space $W_{s,k_s}^1 = \text{Span}(B_s^1)$ with the dimension $k_s = \prod_{i=1}^N k_{s,i}$, for all $s = 1, 2, \dots, m$. Additionally, the k_s -dimensional linear spaces W_{s,k_s}^1 form the solution spaces for system of linear homogeneous ODEs (2.14) for $s = 1, 2, \dots, m$.

Proof. Assume that $W_{s,k_s}^1 = \text{Span}(B_s^1)$, where $B_s^1 = \left\{ \prod_{i=1}^N \xi_{s,r_i}^i(x_i) \mid r_i = 1, \dots, k_{s,i}, i = 1, \dots, N \right\}$, and $\xi_{s,r_i}^i(x_i) \in H_{s,i}$, $i = 1, \dots, N$, $s = 1, 2, \dots, m$. Now, we show that $W_{s,k_s}^1 = \text{Span}(B_s^1)$ is a k_s -dimensional vector space with basis B_s^1 under the vector addition (+)

$$(w_1 + w_2)(x_1, \dots, x_N) = w_1(x_1, \dots, x_N) + w_2(x_1, \dots, x_N),$$

and the scalar multiplication (\cdot) as

$$(\mu_s \cdot w)(x_1, \dots, x_N) = \mu_s \cdot w(x_1, \dots, x_N),$$

for every $\mu_s \in \mathbb{R}$, $w(x_1, \dots, x_N), w_i(x_1, \dots, x_N) \in W_{s,k_s}^1$, $i = 1, 2, s = 1, 2, \dots, m$. It is sufficient to prove that B_s^1 is a linearly independent set. Let us consider the following linear combination of the elements in the set B_s^1 as

$$\sum_{r_1, \dots, r_N} \mu_{s,r_1, \dots, r_N} \prod_{j=1}^N \xi_{s,r_j}^j(x_j) = \sum_{r_N=1}^{k_{s,N}} \left(\sum_{r_1, \dots, r_{N-1}} \mu_{s,r_1, \dots, r_N} \prod_{j=1}^{N-1} \xi_{s,r_j}^j(x_j) \right) \xi_{s,r_N}^N(x_N) = 0,$$

where $\xi_{s,r_i}^i(x_i) \in H_{s,i}$ for $x_i, \mu_{s,r_1, \dots, r_N} \in \mathbb{R}$, $r_i = 1, 2, \dots, k_{s,i}$, $k_{s,i} \in \mathbb{N}$, $i = 1, 2, \dots, N$, and $s = 1, 2, \dots, m$. Since the functions $\xi_{s,r_N}^N(x_N), r_N = 1, 2, \dots, k_{s,N}$, are linearly independent, we obtain

$$\sum_{r_1, \dots, r_{N-1}} \mu_{s,r_1, \dots, r_N} \prod_{j=1}^{N-1} \xi_{s,r_j}^j(x_j) = 0,$$

$r_i = 1, 2, \dots, k_{s,i}, i = 1, 2, \dots, N-1$. Thus, proceeding in a similar manner for $N-1$ successive steps, we get

$$\mu_{s,r_1, \dots, r_N} = 0, \forall r_i = 1, 2, \dots, k_{s,i}, i = 1, 2, \dots, N,$$

which gives that B_s^1 is a linearly independent set. Hence, we conclude that W_{s,k_s}^1 is a k_s -dimensional linear space over \mathbb{R} with basis B_s^1 , where $k_s = \prod_{i=1}^N k_{s,i}$, for each $s = 1, 2, \dots, m$.

Moreover, we observe that $\prod_{i=1}^N \xi_{s,r_i}^i(x_i)$ is a solution for the given system of linear homogeneous ODEs (2.14) for $r_i \in \{1, 2, \dots, k_{s,i}\}, i = 1, 2, \dots, N$. Since the basis set $H_{s,i}$ spans the solution space of the given linear homogeneous ODEs (2.14), we can easily observe that any solution of the system of linear homogeneous ODEs (2.14) has the form $w_s(x_1, x_2, \dots, x_N) = \sum_{r_1, \dots, r_N} \mu_{s,r_1, \dots, r_N} \prod_{j=1}^N \xi_{s,r_j}^j(x_j)$, for any real arbitrary constants $\mu_{s,r_1, \dots, r_N}, r_i = 1, 2, \dots, k_{s,i}, i = 1, 2, \dots, N$, and $s = 1, 2, \dots, m$. Thus, the linear component spaces W_{s,k_s}^1 with basis B_s^1 constitute the solution spaces of the given system of linear homogeneous ODEs (2.14) for $s = 1, 2, \dots, m$, which completes the proof. \square

From Proposition 2.1 and Theorem 2.1, we note that the dimension of type-1 product linear space $\hat{\mathbf{W}}_k^1$ given in (2.18) is $k = \sum_{s=1}^m \prod_{i=1}^N k_{s,i}$. Now, we say that the type-1 product linear space $\hat{\mathbf{W}}_k^1$ given in (2.18) is an invariant product linear space for the given vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$ if $\mathbf{U} \in \hat{\mathbf{W}}_k^1$ implies that $\hat{\mathbf{A}}(\mathbf{U}) \in \hat{\mathbf{W}}_k^1$, or equivalently $A_s(\mathbf{U}) \in W_{s,k_s}^1, s = 1, 2, \dots, m$. Thus, we can describe the invariant criteria for the product linear space $\hat{\mathbf{W}}_k^1$ under the vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$ through the given system of linear homogeneous ODEs (2.14) as follows:

Additionally, from the above-invariant conditions, the type-1 invariant product linear space $\hat{\mathbf{W}}_k^1$ can be written as follows:

$$\hat{\mathbf{W}}_k^1 = \left\{ (w_1, \dots, w_m) \mid \mathbf{L}_{s,k_s,i}^i[w_s] = 0, w_s = w_s(x_1, \dots, x_N) \in W_{s,k_s}^1, \right. \\ \left. s = 1, 2, \dots, m, i = 1, 2, \dots, N \right\}.$$

By Theorem 2.2, we observe that whenever $\hat{\mathbf{W}}_k^1$ given in (2.18) is a type-1 invariant product linear space for the given vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$, then given coupled NTFPDEs (2.1) admits an analytical solution of the form (2.12).

Next, we present the other types of the invariant product linear space $\hat{\mathbf{W}}_k^{p+1}, p = 1, 2, \dots, N-1$. Also, we discuss how to obtain the generalized separable analytical solution given in (2.13) for the given coupled NTFPDEs (2.1) when $p = 1, 2, \dots, N-1$ using the other types of invariant product linear spaces.

2.2.2 Type-2 invariant product linear space $\hat{\mathbf{W}}_k^2$ of the given vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$:

Now, we consider the linear component spaces W_{s,k_s}^2 of the type-2 invariant product linear space $\hat{\mathbf{W}}_k^2$ as $W_{s,k_s}^2 = \text{Span}(B_s^2), s = 1, 2, \dots, m$, where

$$B_s^2 = \left\{ 1, \xi_{s,r_i}^i(x_i), \dots, \xi_{s,r_{i_1}}^{i_1}(x_{i_1}) \xi_{s,r_{i_2}}^{i_2}(x_{i_2}) \cdots \xi_{s,r_{i_{N-1}}}^{i_{N-1}}(x_{i_{N-1}}) \mid \right. \\ \left. r_{i_j}, r_i = 1, 2, \dots, k_{s,i}, i_j, i = 1, 2, \dots, N, j = 1, 2, \dots, N-1 \right\}, \quad (2.19)$$

$\xi_{s,r_i}^i(x_i) \in H_{s,i}$ given in (2.15) for $r_i = 1, 2, \dots, k_{s,i}$, and $\xi_{s,k_{s,i}}^i(x_i) = 1$, for all $i = 1, 2, \dots, N, s = 1, 2, \dots, m$.

Thus, we can define the type-2 product linear space \widehat{W}_k^2 as follows:

$$\begin{aligned}
\widehat{W}_k^2 &= W_{1,k_1}^2 \times W_{2,k_2}^2 \times W_{3,k_3}^2 \times \cdots \times W_{m,k_m}^2 \\
&= \left\{ \mathbf{w}(x_1, \dots, x_N) = (w_1, w_2, \dots, w_m) \mid w_s = \mu_{s,0} + \sum_{i=1}^N \sum_{r_i=1}^{k_{s,i}-1} \mu_{s,r_i}^i \xi_{s,r_i}^i(x_i) + \right. \\
&\quad \cdots + \sum_{i_1, \dots, i_{N-1}} \left(\sum_{r_{i_1}, \dots, r_{i_{N-1}}} \mu_{s,r_{i_1}, \dots, r_{i_{N-1}}}^{i_1, \dots, i_{N-1}} \prod_{j=1}^{N-1} \xi_{s,r_{i_j}}^{i_j}(x_{i_j}) \right), \mu_{s,0}, \mu_{s,r_i}^i, \\
&\quad \left. \dots, \mu_{s,r_{i_1}, \dots, r_{i_{N-1}}}^{i_1, \dots, i_{N-1}} \in \mathbb{R}, w_s = w_s(x_1, \dots, x_N) \in W_{s,k_s}^2, \xi_{s,r_n}^n(x_n) \in H_{s,n}, \right. \\
&\quad \xi_{s,k_{s,n}}^n(x_n) = 1, r_n = 1, 2, \dots, k_{s,n} - 1, n = i, i_j = 1, \dots, N, j = 1, 2, \dots, N - 1, \\
&\quad \left. i_1 < i_2 < \cdots < i_{N-1}, s = 1, 2, \dots, m \right\}. \tag{2.20}
\end{aligned}$$

Now, we prove that the set B_s^2 is a basis for the linear component space W_{s,k_s}^2 , for all $s = 1, 2, \dots, m$, in the following proposition.

Proposition 2.2. *Suppose that the set B_s^2 given in (2.19) spans the linear space W_{s,k_s}^2 , for all $s = 1, 2, \dots, m$, when $\xi_{s,r_i}^i(x_i) \in H_{s,i}$, $r_i = 1, \dots, k_{s,i}$, and $\xi_{s,k_{s,i}}^i(x_i) = 1$, $i = 1, 2, \dots, N$, in the set $H_{s,i}$ given in (2.15). Then, the set B_s^2 forms a basis for the linear space W_{s,k_s}^2 with the dimension $k_s = 1 + \sum_{i=1}^N (k_{s,i} - 1) + \cdots + \sum_{i_1, \dots, i_{N-1}} \prod_{j=1}^{N-1} (k_{s,i_j} - 1)$, for all $i_j = 1, 2, \dots, N$, $j = 1, 2, \dots, N - 1$, $s = 1, 2, \dots, m$. Additionally, the k_s -dimensional linear spaces W_{s,k_s}^2 form the solution spaces for system of linear homogeneous ODEs (2.14) with $\varphi_{s,0}^i(x_i) = 0$, $i = 1, 2, \dots, N$, $s = 1, 2, \dots, m$.*

Proof. Assume that $W_{s,k_s}^2 = \text{Span}(B_s^2)$, $s = 1, 2, \dots, m$. We know that the set B_s^2 is a subset of the linearly independent set B_s^1 , when $\xi_{s,k_{s,i}}^i(x_i) = 1$, $i = 1, 2, \dots, N$, in the set $H_{s,i}$ given in (2.15). Since subset of any linearly independent set is a linearly independent set, we can say that B_s^2 is a basis of the linear space W_{s,k_s}^2 , $s = 1, 2, \dots, m$.

Additionally, following the above similar arguments as given in the proof of Proposition 2.1, we conclude that the linear spaces W_{s,k_s}^2 form the solution spaces for the system of linear homogeneous ODEs (2.14) with $\varphi_{s,0}^i(x_i) = 0$, $i = 1, 2, \dots, N$, and the dimension of W_{s,k_s}^2 is $k_s = 1 + \sum_{i=1}^N (k_{s,i} - 1) + \cdots + \sum_{i_1, \dots, i_{N-1}} \prod_{j=1}^{N-1} (k_{s,i_j} - 1)$, $i_j = 1, 2, \dots, N$, $j = 1, 2, \dots, N - 1$, for all $s = 1, 2, \dots, m$. \square

Based on Propositions 2.1 and 2.2, we find that the dimension of the type-2 product linear space \widehat{W}_k^2 is

$$k = \sum_{s=1}^m \left[1 + \sum_{i=1}^N (k_{s,i} - 1) + \cdots + \sum_{i_1, \dots, i_{N-1}} \prod_{j=1}^{N-1} (k_{s,i_j} - 1) \right],$$

where $i_j = 1, 2, \dots, N, j = 1, 2, \dots, N - 1$. Also, we known that $B_s^2 \subset B_s^1$. Thus, we conclude that the type-2 product linear space $\hat{\mathbf{W}}_k^2$ is a subspace of the type-1 product linear space $\hat{\mathbf{W}}_k^1$ whenever $1 \in W_{s,k_s}^i, i = 1, 2, s = 1, 2, \dots, m$.

Now, we say that the product linear space $\hat{\mathbf{W}}_k^2$ is a type-2 invariant product linear space of the given nonlinear vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$ if $\hat{\mathbf{A}}(\mathbf{U}) \in \hat{\mathbf{W}}_k^2$ whenever $\mathbf{U} \in \hat{\mathbf{W}}_k^2$, or $A_s \left[\hat{\mathbf{W}}_k^2 \right] \subseteq W_{s,k_s}^2, s = 1, 2, \dots, m$. That is, for each $u_s \in W_{s,k_s}^2, s = 1, 2, \dots, m$, we have $A_s(\mathbf{U}) \in W_{s,k_s}^2, s = 1, 2, \dots, m$. For this case, the invariant criteria for the type-2 invariant product linear space $\hat{\mathbf{W}}_k^2$ of the given vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$ read as, for $s = 1, 2, \dots, m$,

$$\begin{aligned} & \frac{\partial^N A_s(\mathbf{U})}{\partial x_1 \partial x_2 \dots \partial x_N} = 0, \mathbf{L}_{s,k_s,i}^i \left[A_s(\mathbf{U}) \right] = 0, \text{ and } \varphi_{s,0}^i(x_i) = 0, i = 1, 2, \dots, N, \\ \text{if } & \frac{\partial^N u_s}{\partial x_1 \partial x_2 \dots \partial x_N} = 0, \mathbf{L}_{s,k_s,i}^i [u_s] = 0, \text{ and } \varphi_{s,0}^i(x_i) = 0, i = 1, 2, \dots, N. \end{aligned}$$

It is interesting to note that based on the above-invariant criteria, the type-2 invariant product linear space $\hat{\mathbf{W}}_k^2$ can be read as

$$\begin{aligned} \hat{\mathbf{W}}_k^2 = \left\{ \mathbf{w}(x_1, \dots, x_N) = (w_1, \dots, w_m) \mid \mathbf{L}_{s,k_s,i}^i [w_s] = 0, \varphi_{s,0}^i(x_i) = 0, \frac{\partial^N w_s}{\partial x_1 \dots \partial x_N} = 0, \right. \\ \left. w_s = w_s(x_1, x_2, \dots, x_N) \in W_{s,k_s}^2, i = 1, 2, \dots, N, s = 1, 2, \dots, m \right\}. \end{aligned}$$

Additionally, we observe that the given coupled NTFPDEs (2.1) admit the following form of the analytical solution

$$\begin{aligned} u_s = & C_{s,0}(t) + \sum_{i=1}^N \sum_{r_i=1}^{k_{s,i}-1} C_{s,r_i}^i(t) \xi_{s,r_i}^i(x_i) + \dots \\ & + \sum_{i_1, \dots, i_{N-1}} \left(\sum_{r_{i_1}, \dots, r_{i_{N-1}}} C_{s,r_{i_1}, \dots, r_{i_{N-1}}}^{i_1, \dots, i_{N-1}}(t) \prod_{j=1}^{N-1} \xi_{s,r_{i_j}}^{i_j}(x_{i_j}) \right), \end{aligned} \quad (2.21)$$

where $r_{i_j} = 1, 2, \dots, k_{s,i_j} - 1, i_j = 1, 2, \dots, N, j = 1, 2, \dots, N - 1, i_1 < \dots < i_{N-1}, s = 1, 2, \dots, m$, if the type-2 product linear space $\hat{\mathbf{W}}_k^2$ is invariant under $\hat{\mathbf{A}}(\mathbf{U})$. The above solution (2.21) is exactly the same as the analytical solution (2.13) when $p = 1$.

Note 2.1. We conclude that the type-2 product linear space $\hat{\mathbf{W}}_k^2$ is a subspace of the type-1 product linear space $\hat{\mathbf{W}}_k^1$, whenever $1 \in W_{s,k_s}^1, i = 1, 2, s = 1, 2, \dots, m$, which does not imply that the nonlinear vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$ admits the type-2 invariant product linear space $\hat{\mathbf{W}}_k^2$, and the type-1 invariant product linear space $\hat{\mathbf{W}}_k^1$, simultaneously. In other words, the nonlinear vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$ may admit the type-2 invariant product linear space $\hat{\mathbf{W}}_k^2$, but the operator $\hat{\mathbf{A}}(\mathbf{U})$ need not admit the type-1 invariant product linear space $\hat{\mathbf{W}}_k^1$, and vice versa.

Next, we present the type- $(p + 1)$ invariant product linear spaces $\widehat{\mathbf{W}}_k^{p+1}$ for $\widehat{\mathbf{A}}(\mathbf{U})$ along with their invariant criteria when $p = 1, 2, \dots, N - 1$. Also, we discuss how to derive the corresponding generalized separable analytical solution for the given multi-component coupled NTFPDEs (2.1) using $\widehat{\mathbf{W}}_k^{p+1}, p = 1, 2, \dots, N - 1$.

2.2.3 Type- $(p + 1)$ invariant product linear space $\widehat{\mathbf{W}}_k^{p+1}, p = 1, 2, \dots, N - 1$ for the given vector differential operator $\widehat{\mathbf{A}}(\mathbf{U})$:

Let us consider the set B_s^{p+1} , for each $s = 1, 2, \dots, m$, as follows:

$$B_s^{p+1} = \left\{ 1, \xi_{s,r_1}^i(x_1), \dots, \xi_{s,r_{i_1}}^{i_1}(x_{i_1}) \xi_{s,r_{i_2}}^{i_2}(x_{i_2}) \cdots \xi_{s,r_{i_{N-p}}}^{i_{N-p}}(x_{i_{N-p}}) \mid \right. \\ \left. r_{i_j}, r_i = 1, 2, \dots, k_{s,i}, i_j, i = 1, 2, \dots, N, j = 1, 2, \dots, N - p \right\}, \quad (2.22)$$

where $\xi_{s,r_i}^i(x_i) \in H_{s,i}$ given in (2.15) for $r_i = 1, 2, \dots, k_{s,i}$, and $\xi_{s,k_{s,i}}^i(x_i) = 1$, for all $i = 1, 2, \dots, N, s = 1, 2, \dots, m$. Now, we define the linear component spaces W_{s,k_s}^{p+1} of the type- $(p + 1)$ invariant product linear space $\widehat{\mathbf{W}}_k^{p+1}$ as $W_{s,k_s}^{p+1} = \text{Span}(B_s^{p+1})$, $s = 1, 2, \dots, m$. Thus, we construct the type- $(p + 1)$ product linear space $\widehat{\mathbf{W}}_k^{p+1}$ as follows:

$$\begin{aligned} \widehat{\mathbf{W}}_k^{p+1} &= W_{1,k_1}^{p+1} \times W_{2,k_2}^{p+1} \times W_{3,k_3}^{p+1} \times \cdots \times W_{m,k_m}^{p+1} \\ &= \left\{ \mathbf{w}(x_1, \dots, x_N) = (w_1, \dots, w_m) \mid w_s = \mu_{s,0} + \sum_{i=1}^N \sum_{r_i=1}^{k_{s,i}-1} \mu_{s,r_i}^i \xi_{s,r_i}^i(x_i) + \right. \\ &\quad \cdots + \sum_{i_1, \dots, i_{N-p}} \left(\sum_{r_{i_1}, \dots, r_{i_{N-p}}} \mu_{s,r_{i_1}, \dots, r_{i_{N-p}}}^{i_1, \dots, i_{N-p}} \prod_{j=1}^{N-p} \xi_{s,r_{i_j}}^{i_j}(x_{i_j}) \right), \mu_{s,0}, \mu_{s,r_i}^i, \\ &\quad \left. \dots, \mu_{s,r_{i_1}, \dots, r_{i_{N-p}}}^{i_1, \dots, i_{N-p}} \in \mathbb{R}, w_s = w_s(x_1, \dots, x_N) \in W_{s,k_s}^{p+1}, \right. \\ &\quad \xi_{s,r_n}^n(x_n) \in H_{s,n}, \xi_{s,k_{s,i}}^i(x_i) = 1, r_n = 1, 2, \dots, k_{s,n} - 1, \\ &\quad \left. n = i, i_j = 1, 2, \dots, N, j = 1, 2, \dots, N - p, i_1 < \cdots < i_{N-p}, s = 1, 2, \dots, m \right\}, \end{aligned} \quad (2.23)$$

for all $p = 1, 2, \dots, N - 1$. Next, we show that the set B_s^{p+1} is a basis for the linear component space W_{s,k_s}^{p+1} for all $s = 1, 2, \dots, m$.

Proposition 2.3. *Suppose that the linear space W_{s,k_s}^{p+1} is spanned by set B_s^{p+1} given in (2.22) for all $s = 1, 2, \dots, m$, when $\xi_{s,r_i}^i(x_i) \in H_{s,i}, r_i = 1, \dots, k_{s,i}$, and $\xi_{s,k_{s,i}}^i(x_i) = 1, i = 1, 2, \dots, N$, in the set $H_{s,i}$ given in (2.15). Then, the set B_s^{p+1} forms a basis for the linear space W_{s,k_s}^{p+1} with the dimension $k_s = 1 + \sum_{i=1}^N (k_{s,i} - 1) + \cdots + \sum_{i_1, \dots, i_{N-p}} \prod_{j=1}^{N-p} (k_{s,i_j} - 1), i_j = 1, 2, \dots, N, j = 1, 2, \dots, N - p$, for all $s = 1, 2, \dots, m$. Additionally, the k_s -dimensional linear spaces W_{s,k_s}^{p+1} form solution spaces for system of linear homogeneous ODEs (2.14) with $\varphi_{s,0}^i(x_i) = 0, i = 1, 2, \dots, N, s = 1, 2, \dots, m$.*

Proof. Similar to the proof of Proposition 2.2. \square

Based on Propositions 2.1 and 2.3, we find that the dimension of the type- $(p+1)$ product linear space $\widehat{\mathbf{W}}_k^{p+1}$ is

$$k = \sum_{s=1}^m \left[1 + \sum_{i=1}^N (k_{s,i} - 1) + \cdots + \sum_{i_1, \dots, i_{N-p}} \prod_{j=1}^{N-p} (k_{s,i_j} - 1) \right],$$

where $i_j = 1, 2, \dots, N, j = 1, 2, \dots, N-p, p = 1, 2, \dots, N-1$. Additionally, we note that $B_s^{p+1} \subset B_s^p$ whenever $1 \in B_s^1$, for $p = 1, 2, \dots, N-1, s = 1, 2, \dots, m$. Thus, we say that the type- $(p+1)$ product linear space $\widehat{\mathbf{W}}_k^{p+1}$ is a subspace of the type- p product linear space $\widehat{\mathbf{W}}_k^p$ whenever $1 \in W_{s,k_s}^p, p = 1, 2, \dots, N-1, s = 1, 2, \dots, m$. That is, when $1 \in W_{s,k_s}^i, i = 1, 2, \dots, N, s = 1, 2, \dots, m$, we have

$$\widehat{\mathbf{W}}_k^N \subset \widehat{\mathbf{W}}_k^{N-1} \subset \cdots \subset \widehat{\mathbf{W}}_k^2 \subset \widehat{\mathbf{W}}_k^1.$$

A finite-dimensional product linear space $\widehat{\mathbf{W}}_k^{p+1}$ is a type- $(p+1)$ invariant product linear space for the given vector differential operator $\widehat{\mathbf{A}}(\mathbf{U})$ if $\widehat{\mathbf{A}}(\mathbf{U}) \in \widehat{\mathbf{W}}_k^{p+1}$ whenever $\mathbf{U} \in \widehat{\mathbf{W}}_k^{p+1}$ or $A_s \left[\widehat{\mathbf{W}}_k^{p+1} \right] \subseteq W_{s,k_s}^{p+1}, s = 1, 2, \dots, m$. That is, for each $u_s \in W_{s,k_s}^{p+1}$ we have $A_s(\mathbf{U}) \in W_{s,k_s}^{p+1}, p = 1, 2, \dots, N-1, s = 1, 2, \dots, m$.

Thus, the invariant criteria for the type- $(p+1)$ invariant product linear space $\widehat{\mathbf{W}}_k^{p+1}$ of the given vector differential operator $\widehat{\mathbf{A}}(\mathbf{U})$ can be read as, for every $s = 1, 2, \dots, m$,

$$\begin{aligned} \frac{\partial^N A_s(\mathbf{U})}{\partial x_{i_1} \partial x_{i_2} \cdots \partial x_{i_{N-p-1}}} &= 0, \mathbf{L}_{s,k_s,j}^j \left[A_s(\mathbf{U}) \right] = 0, \text{ and } \varphi_{s,0}^j(x_j) = 0, \text{ whenever} \\ \frac{\partial^N u_s}{\partial x_{i_1} \partial x_{i_2} \cdots \partial x_{i_{N-p-1}}} &= 0, \mathbf{L}_{s,k_s,j}^j [u_s] = 0, \text{ and } \varphi_{s,0}^j(x_j) = 0, \end{aligned}$$

where $i_r, j = 1, 2, \dots, N, r = 1, 2, \dots, N-(p+1)$. Also, from the above-invariant criteria, the type- $(p+1)$ invariant product linear space $\widehat{\mathbf{W}}_k^{p+1}$ can be written as

$$\begin{aligned} \widehat{\mathbf{W}}_k^{p+1} &= \left\{ \mathbf{w}(x_1, \dots, x_N) = (w_1, w_2, \dots, w_m) \mid \frac{\partial^N w_s}{\partial x_{i_1} \partial x_{i_2} \cdots \partial x_{i_{N-p-1}}} = 0, \right. \\ &\quad \left. \mathbf{L}_{s,k_s,j}^j [w_s] = 0, \varphi_{s,0}^j(x_j) = 0, w_s = w_s(x_1, \dots, x_N) \in W_{s,k_s}^{p+1}, \right. \\ &\quad \left. i_r, j = 1, 2, \dots, N, s = 1, 2, \dots, m, r = 1, 2, \dots, N-(p+1) \right\}. \end{aligned}$$

It is important to observe that the given coupled NTFPDEs (2.1) obtain the analytical solution of the form given in (2.13) whenever the given vector differential operator $\widehat{\mathbf{A}}(\mathbf{U})$ admits $\widehat{\mathbf{W}}_k^{p+1}, p = 1, 2, \dots, N-1$. Additionally, it is noteworthy to mention that the type-2 product linear space $\widehat{\mathbf{W}}_k^2$ can be obtained from the type- $(p+1)$ invariant product linear space $\widehat{\mathbf{W}}_k^{p+1}$ when $p = 1$.

Note 2.2. We observe that the relation is obtained as $\hat{\mathbf{W}}_k^N \subset \hat{\mathbf{W}}_k^{N-1} \subset \dots \subset \hat{\mathbf{W}}_k^2 \subset \hat{\mathbf{W}}_k^1$ from all N -types of product linear spaces $\hat{\mathbf{W}}_k^p$, whenever $1 \in W_{s,k_s}^p, s = 1, 2, \dots, m, p = 1, 2, \dots, N$. However, we can say that, in general, it need not be provided with the product linear spaces $\hat{\mathbf{W}}_k^1, \hat{\mathbf{W}}_k^2, \dots, \hat{\mathbf{W}}_k^N$, are invariant under the nonlinear vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$, simultaneously. Thus, it can be simply said that the nonlinear vector differential operator $\hat{\mathbf{A}}(\mathbf{U})$ may admit the type- p_1 invariant product linear space $\hat{\mathbf{W}}_k^{p_1}$, but the operator $\hat{\mathbf{A}}(\mathbf{U})$ need not admit the type- p_2 invariant product linear space $\hat{\mathbf{W}}_k^{p_2}$ whenever $p_1 \neq p_2, p_1, p_2 = 1, 2, \dots, N$.

Next, we show the efficient application of the invariant subspace method by determining various dimensional type-1 and type-2 invariant product linear spaces and their generalized separable analytical solutions for the multi-component $(2 + 1)$ -dimensional coupled nonlinear TFDCWEs (1.7) in the following section.

3 Invariant subspace method for the multi-component $(2 + 1)$ -dimensional coupled nonlinear TFDCWEs (7)

In this section, we explain how to apply the invariant subspace method to obtain the analytical solutions of the multi-component $(2 + 1)$ -dimensional coupled nonlinear TFDCWEs (1.7). So, let us first present how to find different type-1 and type-2 invariant product linear spaces for the following vector differential operators of the given coupled equations (1.7):

$$\mathbf{F}(\mathbf{U}) = (F_1(u_1, u_2), F_2(u_1, u_2)), \quad (3.1)$$

where

$$\begin{aligned} F_1(u_1, u_2) &= \sum_{i=1}^2 \frac{\partial}{\partial x_i} \left(\beta_{1i}(u_1, u_2) \frac{\partial u_1}{\partial x_i} + \rho_{1i}(u_1, u_2) \frac{\partial u_2}{\partial x_i} \right) \\ &\quad + \sum_{i=1}^2 \left(\zeta_{1i}(u_1, u_2) \frac{\partial u_1}{\partial x_i} + v_{1i}(u_1, u_2) \frac{\partial u_2}{\partial x_i} \right), \\ \text{and } F_2(u_1, u_2) &= \sum_{i=1}^2 \frac{\partial}{\partial x_i} \left(\beta_{2i}(u_1, u_2) \frac{\partial u_1}{\partial x_i} + \rho_{2i}(u_1, u_2) \frac{\partial u_2}{\partial x_i} \right) \\ &\quad + \sum_{i=1}^2 \left(\zeta_{2i}(u_1, u_2) \frac{\partial u_1}{\partial x_i} + v_{2i}(u_1, u_2) \frac{\partial u_2}{\partial x_i} \right). \end{aligned}$$

In general, we cannot find the invariant product linear spaces for arbitrary differential operators $F_s(u_1, u_2), s = 1, 2$. Thus, we consider the multi-component $(2 + 1)$ -dimensional coupled TFDCWEs (1.7) with cubic nonlinearity. So, let us consider the following forms of diffusion

coefficients and convection terms as

$$\begin{aligned}
\beta_{s1}(u_1, u_2) &= k_{s5}u_1^2 + k_{s4}u_2^2 + k_{s3}u_1u_2 + k_{s2}u_2 + k_{s1}u_1 + k_{s0}, \\
\rho_{s1}(u_1, u_2) &= q_{s5}u_1^2 + q_{s4}u_2^2 + q_{s3}u_1u_2 + q_{s2}u_2 + q_{s1}u_1 + q_{s0}, \\
\beta_{s2}(u_1, u_2) &= p_{s5}u_1^2 + p_{s4}u_2^2 + p_{s3}u_1u_2 + p_{s2}u_2 + p_{s1}u_1 + p_{s0}, \\
\rho_{s2}(u_1, u_2) &= c_{s5}u_1^2 + c_{s4}u_2^2 + c_{s3}u_1u_2 + c_{s2}u_2 + c_{s1}u_1 + c_{s0}, \\
\zeta_{s1}(u_1, u_2) &= \eta_{s5}u_1^2 + \eta_{s4}u_2^2 + \eta_{s3}u_1u_2 + \eta_{s2}u_2 + \eta_{s1}u_1 + \eta_{s0}, \\
v_{s1}(u_1, u_2) &= g_{s5}u_1^2 + g_{s4}u_2^2 + g_{s3}u_1u_2 + g_{s2}u_2 + g_{s1}u_1 + g_{s0}, \\
\zeta_{s2}(u_1, u_2) &= d_{s5}u_1^2 + d_{s4}u_2^2 + d_{s3}u_1u_2 + d_{s2}u_2 + d_{s1}u_1 + d_{s0}, \\
v_{s2}(u_1, u_2) &= h_{s5}u_1^2 + h_{s4}u_2^2 + h_{s3}u_1u_2 + h_{s2}u_2 + h_{s1}u_1 + h_{s0},
\end{aligned}$$

where $k_{si}, q_{si}, p_{si}, c_{si}, \eta_{si}, g_{si}, d_{si}, h_{si} \in \mathbb{R}$, for $s = 1, 2$, and $i = 0, 1, 2, \dots, 5$. In the following subsection, we present a detailed description of finding two different types of invariant product linear spaces $\hat{\mathbf{W}}_k^1$ and $\hat{\mathbf{W}}_k^2$ for the cubic nonlinear vector differential operator $\mathbf{F}(\mathbf{U})$.

3.1 Finding the invariant product linear space $\hat{\mathbf{W}}_k$ for the considered operator $\mathbf{F}(\mathbf{U})$

Let us describe how to construct two types of various finite-dimensional invariant product linear spaces of the cubic nonlinear vector differential operator $\mathbf{F}(\mathbf{U})$ given in (3.1). Thus, we consider the system of $k_{s,i}$ -order linear homogeneous ODEs of the form

$$\begin{aligned}
\mathbf{L}_{s,k_{s,i}}^i[y_{s,i}(x_i)] &\equiv \frac{d^{k_{s,i}}y_{s,i}(x_i)}{dx_i^{k_{s,i}}} + \varphi_{s,k_{s,i}-1}^i(x_i) \frac{d^{k_{s,i}-1}y_{s,i}(x_i)}{dx_i^{k_{s,i}-1}} + \dots \\
&+ \varphi_{s,0}^i(x_i)y_{s,i}(x_i) = 0, k_{s,i} \in \mathbb{N}, s, i = 1, 2
\end{aligned} \tag{3.2}$$

with the basis of the solution spaces for $\mathbf{L}_{s,k_{s,i}}^i[y_{s,i}(x_i)] = 0$ as follows:

$$H_{s,i} = \left\{ \xi_{s,1}^i(x_i), \xi_{s,2}^i(x_i), \dots, \xi_{s,k_{s,i}}^i(x_i) \right\}, s, i = 1, 2. \tag{3.3}$$

Note that the given cubic nonlinear vector differential operator $\mathbf{F}(\mathbf{U})$ involves two space variables x_1 and x_2 . Thus, we consider $N = 2$. Hence, we can define two types of invariant product linear spaces, which are named type-1 product linear spaces $\hat{\mathbf{W}}_k^1$ and type-2 product linear spaces $\hat{\mathbf{W}}_k^2$. Now, let us first explain how to find the type-1 product linear spaces $\hat{\mathbf{W}}_k^1$ for $\mathbf{F}(\mathbf{U})$ in the following.

Type-1 invariant product linear spaces $\hat{\mathbf{W}}_k^1$ preserved by the given coupled equations (3.1): From the definition of type-1 invariant product linear spaces given in (2.18), we

can write the type-1 invariant product linear spaces $\hat{\mathbf{W}}_k^1$ for the given coupled equations (3.1) as follows:

$$\begin{aligned}\hat{\mathbf{W}}_k^1 &= W_{1,k_1}^1 \times W_{2,k_2}^1 \\ &= \left\{ w(x_1, x_2) = (w_1(x_1, x_2), w_2(x_1, x_2)) \mid \mathbf{L}_{s,k_s,i}^i [w_s(x_1, x_2)] = 0, s, i = 1, 2 \right\} \\ &= \left\{ (w_1, w_2) \mid w_s := w_s(x_1, x_2) = \sum_{r_1, r_2} \mu_{r_1, r_2}^s \xi_{s, r_1}^1(x_1) \xi_{s, r_2}^2(x_2) \in W_{s, k_s}^1, \right. \\ &\quad \left. \mu_{i_1, i_2}^s \in \mathbb{R}, r_i = 1, \dots, k_{s, i}, s, i = 1, 2 \right\},\end{aligned}\tag{3.4}$$

where $\mathbf{L}_{s, k_s, i}^i(\cdot)$ is given in (3.2) and $\xi_{s, 1}^i(x_i) \in H_{s, i}$ is given in (3.3) for $s, i = 1, 2$. Note that the dimension of $\hat{\mathbf{W}}_k^1$ is $k = \dim(\hat{\mathbf{W}}_k^1) = k_{1,1}k_{1,2} + k_{2,1}k_{2,2}$.

The above product linear space (3.4) is invariant under $\mathbf{F}(\mathbf{U})$ if $\mathbf{F}[\hat{\mathbf{W}}_k^1] \subseteq \hat{\mathbf{W}}_k^1$, which means that, for every $\mathbf{U} \in \hat{\mathbf{W}}_k^1$, $F_s(\hat{\mathbf{W}}_k^1) = F_s(W_{1,k_1}^1 \times W_{2,k_2}^1) \subseteq W_{s, k_s}^1, \forall s = 1, 2$. These invariant conditions can be rewritten in the following equivalent form

$$\mathbf{L}_{s, k_s, i}^i [F_s(u_1, u_2)] = 0 \text{ whenever } \mathbf{L}_{s, k_s, i}^i [u_s] = 0, s, i = 1, 2.\tag{3.5}$$

Observe that if the type-1 product linear space $\hat{\mathbf{W}}_k^1$ is invariant under $\mathbf{F}(\mathbf{U})$, then we expect the analytical solution for the given coupled equations (1.7) as $\mathbf{U} = (u_1, u_2)$ with component functions

$$u_s = \sum_{r_1=1}^{k_{s,1}} \sum_{r_2=1}^{k_{s,2}} C_{s, r_1, r_2}(t) \xi_{s, r_1}^1(x_1) \xi_{s, r_2}^2(x_2), k_{s, i} \in \mathbb{N}, s, i = 1, 2.$$

Now, we present a detailed illustration for finding the type-1 product linear spaces $\hat{\mathbf{W}}_k^1$ for the given coupled equations (3.1) using the given system of ODEs (3.2) with $k_{s, i} = 2$. Thus, for $k_{s, i} = 2$, the given system of ODEs (3.2) is obtained as follows:

$$\begin{aligned}\mathbf{L}_{1, k_{1, i}}^i [y_{1, i}(x_i)] &\equiv \left[\frac{d^2}{dx_i^2} + a_{i1} \frac{d}{dx_i} + a_{i0} \right] y_{1, i}(x_i) = 0, \\ \text{and } \mathbf{L}_{2, k_{2, i}}^i [y_{2, i}(x_i)] &\equiv \left[\frac{d^2}{dx_i^2} + b_{i1} \frac{d}{dx_i} + b_{i0} \right] y_{2, i}(x_i) = 0,\end{aligned}\tag{3.6}$$

where $\varphi_{1, j}^i(x_i) = a_{ij}, \varphi_{2, j}^i(x_i) = b_{ij} \in \mathbb{R}$, and the fundamental set of solutions of $\mathbf{L}_{1, k_{1, i}}^i [y_{1, i}(x_i)] = 0$ is $H_{s, i} = \{ \xi_{s, 1}^i(x_i), \xi_{s, 2}^i(x_i) \}, j = 0, 1, s, i = 1, 2$. Thus, the given invariant conditions (3.5) for the type-1 invariant product linear space $\hat{\mathbf{W}}_k^1$ of $\mathbf{F}(\mathbf{U})$ are reduced as

$$\left(\frac{d^2}{dx_1^2} + a_{11} \frac{d}{dx_1} + a_{10} \right) F_1(u_1, u_2) = 0, \quad \left(\frac{d^2}{dx_2^2} + a_{21} \frac{d}{dx_2} + a_{20} \right) F_1(u_1, u_2) = 0,\tag{3.7}$$

$$\left(\frac{d^2}{dx_1^2} + b_{11} \frac{d}{dx_1} + b_{10} \right) F_2(u_1, u_2) = 0, \quad \& \left(\frac{d^2}{dx_2^2} + b_{21} \frac{d}{dx_2} + b_{20} \right) F_2(u_1, u_2) = 0,\tag{3.8}$$

whenever

$$\left(\frac{d^2}{dx_i^2} + a_{i1} \frac{d}{dx_i} + a_{i0} \right) u_1 = 0, \quad \text{and} \quad \left(\frac{d^2}{dx_i^2} + b_{i1} \frac{d}{dx_i} + b_{i0} \right) u_2 = 0, i = 1, 2.\tag{3.9}$$

Here, we consider the cubic nonlinear vector differential operators $F_s(u_1, u_2)$, $s = 1, 2$, as follows:

$$\begin{aligned}
F_s(u_1, u_2) = & \frac{\partial}{\partial x_1} \left[(k_{s5}u_1^2 + k_{s4}u_2^2 + k_{s3}u_1u_2 + k_{s2}u_2 + k_{s1}u_1 + k_{s0}) \frac{\partial u_1}{\partial x_1} \right. \\
& \left. + (q_{s5}u_1^2 + q_{s4}u_2^2 + q_{s3}u_1u_2 + q_{s2}u_2 + q_{s1}u_1 + q_{s0}) \frac{\partial u_2}{\partial x_1} \right] \\
& + \frac{\partial}{\partial x_2} \left[(p_{s5}u_1^2 + p_{s4}u_2^2 + p_{s3}u_1u_2 + p_{s2}u_2 + p_{s1}u_1 + p_{s0}) \frac{\partial u_1}{\partial x_2} \right. \\
& \left. + (c_{s5}u_1^2 + c_{s4}u_2^2 + c_{s3}u_1u_2 + c_{s2}u_2 + c_{s1}u_1 + c_{s0}) \frac{\partial u_2}{\partial x_2} \right] \\
& + (\eta_{s5}u_1^2 + \eta_{s4}u_2^2 + \eta_{s3}u_1u_2 + \eta_{s2}u_2 + \eta_{s1}u_1 + \eta_{s0}) \frac{\partial u_1}{\partial x_1} \\
& + (g_{s5}u_1^2 + g_{s4}u_2^2 + g_{s3}u_1u_2 + g_{s2}u_2 + g_{s1}u_1 + g_{s0}) \frac{\partial u_2}{\partial x_1} \\
& + (d_{s5}u_1^2 + d_{s4}u_2^2 + d_{s3}u_1u_2 + d_{s2}u_2 + d_{s1}u_1 + d_{s0}) \frac{\partial u_1}{\partial x_2} \\
& + (h_{s5}u_1^2 + h_{s4}u_2^2 + h_{s3}u_1u_2 + h_{s2}u_2 + h_{s1}u_1 + h_{s0}) \frac{\partial u_2}{\partial x_2}.
\end{aligned} \tag{3.10}$$

Substituting the above operators $F_s(u_1, u_2)$, $s = 1, 2$, into the above invariant conditions (3.7)-(3.8), and equating each the coefficients of u_s , $\frac{\partial u_s}{\partial x_i}$, $u_1 \left(\frac{\partial u_s}{\partial x_i} \right)$, $u_2 \left(\frac{\partial u_s}{\partial x_i} \right)$, $\left(\frac{\partial u_s}{\partial x_i} \right)^2$, $s, i = 1, 2$, etc. to zero, we get the subsequent set of equations:

(i) From the first equation of (3.7), we get

$$\begin{aligned}
c_{11} + p_{12} = 0, & a_{11}b_{10}(2k_{12} + q_{11}) + b_{10}b_{11}(k_{12} + 3q_{11}) - b_{10}(\eta_{12} - 2g_{11}) = 0, \\
b_{10}(b_{21}c_{11} - h_{11}) = 0, & b_{10}(a_{21}p_{12} - d_{12}) = 0, d_{12} - a_{21}p_{12} = 0, 2h_{11} - b_{21}c_{11} = 0, \\
p_{1i} = p_{11} = d_{1i} = d_{11} = c_{1i} = c_{12} = h_{1i} = h_{12} = 0, & i = 3, 4, 5, (b_{11} - a_{11})(b_{21}c_{11} \\
- h_{11}) = 0, & (b_{10} - a_{10})(b_{21}c_{10} - h_{10}) = 0, b_{10}((a_{11} - b_{11})b_{11}q_{11} + (3k_{12} + 2q_{11})a_{10} \\
- a_{11}g_{11} + b_{11}g_{11} + a_{20}p_{12}b_{10}q_{11} + b_{20}c_{11}) = 0, & (2b_{11} - a_{11})b_{10}q_{10} + (a_{10} - b_{10})g_{10} \\
- a_{10}b_{11}q_{10} + (b_{11}g_{10} + b_{20}c_{10} - b_{11}^2q_{10})(b_{11} - a_{11}) = 0, & a_{11}(k_{12}(a_{11} + 2b_{11}) + b_{11}q_{11} \\
- \eta_{12}) + b_{11}^2(k_{12} + 3q_{11}) - (\eta_{12} + 2g_{11})b_{11} - b_{10}(3q_{11} + k_{12}) - 2(a_{10}k_{12} + a_{20}p_{12} \\
+ b_{20}c_{11}) = 0, & (a_{11} - b_{11})b_{11}^2q_{11} + a_{10}k_{12}(a_{11} + 3b_{11}) + (2a_{10}q_{11} + a_{20}p_{12} + 2b_{10}q_{11} \\
+ b_{11}g_{11} + b_{20}c_{11})b_{11} - a_{11}(a_{20}p_{12} + b_{10}q_{11} + b_{11}g_{11} + b_{20}c_{11}) - 2a_{10}\eta_{12} - b_{10}g_{11} = 0, \\
b_{10}(a_{11}b_{11}q_{10} - b_{11}^2q_{10} - a_{10}q_{10} - a_{11}g_{10} + b_{10}q_{10} + b_{11}g_{10} + b_{20}c_{10}) - a_{10}b_{20}c_{10} = 0, \\
(b_{11} - a_{11})(a_{21}p_{12} - d_{12}) = 0, & (a_{11} - b_{11})(p_{12} + c_{11}) = 0, (b_{11} - a_{11})(b_{21}c_{10} - h_{10}) = 0.
\end{aligned}$$

(ii) From the second equation of (3.7), we get

$$\begin{aligned}
& b_{21}^2(a_{21} - b_{21})c_{11} + (b_{21} - a_{21})a_{10}k_{12} + a_{20}p_{12}(a_{21} + 3b_{21}) + (2a_{20}c_{11} - a_{21}h_{11} \\
& + b_{10}q_{11} + 2b_{20}c_{11} + b_{21}h_{11}) - a_{21}(b_{10}q_{11} + b_{20}c_{11}) - b_{20}h_{11} - 2a_{20}d_{12} = 0, \\
& q_{1i} = q_{12} = k_{1i} = k_{11} = \eta_{1i} = \eta_{11} = g_{1i} = g_{12} = 0, i = 3, 4, 5, (b_{21} - a_{21})(b_{11}q_{10} \\
& - g_{10}) = 0, b_{20}b_{21}c_{10}(a_{21} - b_{21}) + a_{20}(-b_{10}q_{10} - b_{20}c_{10}) + b_{20}(-a_{21}h_{10} + b_{10}q_{10} \\
& + b_{20}c_{10} + b_{21}h_{10}) = 0, b_{20}b_{21}c_{11}(a_{21} - b_{21}) + b_{20}(a_{10}k_{12} + 3a_{20}p_{12} + 2a_{20}c_{11} - a_{21}h_{11} \\
& + b_{10}q_{11} + b_{20}c_{11} + b_{21}h_{11}) = 0, b_{20}((2p_{12} + c_{11})a_{21} + b_{21}(p_{12} + 3bc_{11}) - d_{12} + 2h_{11}) = 0, \\
& (b_{11}q_{11} - g_{11})b_{20} = 0, (b_{20}c_{10} - b_{21}^2c_{10} + b_{21}h_{10})(b_{21} - a_{21}) + b_{10}q_{10}(2b_{21} - a_{21}) + (a_{20} \\
& - b_{20})h_{10} - a_{20}b_{21}c_{10} = 0, b_{20}(k_{12} + q_{11}) = 0, a_{21}p_{12}(a_{21} + 2b_{21}) + (b_{21} - b_{20} - 2a_{20})p_{12} \\
& + b_{21}(a_{21} + 3b_{21})c_{11} - 2a_{10}k_{12} - d_{12}(b_{21} + a_{21}) - 2b_{10}q_{11} - 3b_{20}c_{11} - 2b_{21}h_{11} = 0, \\
& g_{11} - b_{11}q_{11} = 0, (b_{20} - a_{20})(b_{11}q_{10} + g_{10}) = 0, b_{20}(a_{11}k_{12} - \eta_{12}) = 0, q_{11} + k_{12} = 0, \\
& (a_{21} - b_{21})(k_{12} + q_{11}) = 0, (b_{21} - a_{21})(b_{11}q_{11} - g_{11}) = 0, (\eta_{12} - a_{11}k_{12})(a_{21} - b_{21}) = 0.
\end{aligned}$$

(iii) From the first equation of (3.8), we get

$$\begin{aligned}
& p_{2i} = p_{21} = d_{2i} = d_{21} = c_{2i} = c_{22} = h_{2i} = h_{22} = 0, i = 3, 4, 5, d_{22} - a_{21}p_{22} = 0, \\
& (p_{22} + c_{21})(b_{11} - a_{11}) = 0, (b_{21}c_{21} - h_{21})(a_{11} - b_{11}) = 0, a_{10}(b_{21}c_{21} - h_{21}) = 0, \\
& a_{10}(a_{21}p_{22} - d_{22}) = 0, (a_{11} - b_{11})(a_{21}p_{20} - d_{20}) = 0, (a_{11} - b_{11})(a_{21}p_{22} - d_{22}) = 0, \\
& (a_{21}p_{20} - d_{20})(a_{10} - b_{10}) = 0, a_{10}(a_{11}(3k_{22} + q_{21}) + b_{11}(k_{22} + 2q_{21}) + g_{21} \\
& - 2\eta_{22}) = 0, a_{10}(a_{11}b_{11} + a_{10}k_{20} + a_{11}\eta_{20} - a_{11}^2k_{10}k_{20} + a_{20}p_{20} - b_{10}k_{20} - b_{11}\eta_{20}) \\
& - a_{20}b_{10}p_{20} = 0, (a_{10}a_{11}k_{22} - \eta_{22}a_{10})(b_{11} - a_{11}) + (a_{10}k_{22} + a_{20}p_{22} + b_{10} + b_{20}c_{21})a_{10} \\
& + a_{20}(2k_{22} + 3q_{21}) = 0, (b_{11} - a_{11})(a_{11}^2k_{20} - a_{11}\eta_{20} - a_{20}p_{20}) + (2a_{11} - b_{11})a_{10}k_{20} \\
& + b_{10}(\eta_{20} - a_{11}k_{20}) - a_{10}\eta_{20} = 0, a_{11}^2(3k_{22} + q_{21}) + a_{11}b_{11}(k_{22} + 2q_{21}) + (b_{11}^2 - a_{10} \\
& - 2b_{10})q_{21} - 3a_{10}k_{22} - a_{11}(2\eta_{22} + g_{21}) - 2a_{20}p_{22} - b_{11}g_{21} - 2b_{20}c_{21} = 0, p_{22} + c_{21} = 0.
\end{aligned}$$

(iv) From the second equation of (3.8), we get

$$\begin{aligned}
& (b_{21} - a_{21}p_{22})a_{21}^2p_{22} + (a_{10}k_{22} + b_{10}q_{21})(a_{21} - b_{21}) - b_{20}(2h_{21} + c_{21}) - b_{21} \\
& + p_{22}(2a_{21} - b_{21})a_{20} + a_{21}(a_{21}d_{22} + 2b_{20}p_{22} + 3b_{20}c_{21} - b_{21}d_{22}) - a_{20}d_{22} = 0, \\
& q_{2i} = q_{22} = k_{2i} = k_{21} = \eta_{2i} = \eta_{21} = g_{2i} = g_{22} = 0, i = 3, 4, 5, \eta_{22} - a_{11}k_{22} = 0, \\
& (a_{11}k_{20} - \eta_{20})(a_{2i} - b_{2i}) = 0, i = 0, 1, 2g_{21} - 2b_{11}q_{21} = 0, a_{20}b_{11}q_{21} - a_{20}g_{21} = 0, \\
& (a_{11}k_{22} - \eta_{22})(a_{21} - b_{21}) = 0, (a_{21} - b_{21})(b_{11}q_{21} - g_{21}) = 0, a_{20}a_{21}(3p_{22} + c_{21}) \\
& + a_{20}b_{21}(p_{22} + 2c_{21}) - a_{20}(2d_{22} + h_{21}) = 0, (b_{21} - a_{21})(k_{22} + q_{21}) = 0, a_{20}(k_{22} \\
& + q_{21}) = 0, a_{20}(a_{21}b_{21}p_{20} - a_{21}^2p_{20} + a_{10}k_{20} + a_{20}p_{20} + a_{21}d_{20} - b_{20}p_{20} - b_{21}d_{20}) \\
& - a_{10}b_{20}k_{20} = 0, a_{20}[(b_{21} - a_{21})a_{21}p_{22} + a_{10}k_{22} + a_{20}p_{22} + a_{21}d_{22} + b_{10}q_{21} - b_{21}d_{22} \\
& + 2b_{20}p_{22} + 3b_{20}c_{21}] = 0, (a_{21}^2p_{20} - a_{10}k_{20} - a_{21}d_{20})(b_{21} - a_{21}) + a_{20}p_{20}(2a_{21} - b_{21}) \\
& + b_{20}(d_{20} - a_{21}p_{20}) - a_{20}d_{20} = 0, a_{21}^2(3p_{22} + c_{21}) + a_{21}b_{21}(p_{22} + 2c_{21}) + (b_{21}^2 - a_{20} \\
& - 2b_{20})c_{21} - 2a_{10}k_{22} - 3a_{20}p_{22} - b_{21}h_{21} - 2b_{10}q_{21} - a_{21}(2d_{22} + h_{21}) = 0, q_{21} + k_{22} = 0.
\end{aligned}$$

Solving the above sets of equations (i)-(iv) together, we obtain the eight-dimensional type-1 invariant product linear spaces $\widehat{\mathbf{W}}_8^1$ with their corresponding multi-component cubic nonlinear vector differential

operators $\mathbf{F}(\mathbf{U})$. However, we cannot generally solve them for the above set of equations (i)-(iv). Thus, let us first consider the case $a_{11} = b_{21} = 0$. For this case, the obtained eight-dimensional type-1 invariant product linear spaces $\hat{\mathbf{W}}_8^1$ and their corresponding non-zero parameter restrictions in the differential operators $F_s(u_1, u_2)$, $s = 1, 2$, are listed from Table 1 to Table 5. Additionally, we wish to point out that some parameters of the given component differential operators $F_s(u_1, u_2)$, $s = 1, 2$, (3.10) are zero that are not listed in Table 1-Table 5 due to the large number of parameters and cases. Hence, only the non-zero parameters of $F_s(u_1, u_2)$, $s = 1, 2$, given in (3.10) are listed in Table 1-Table 5, and the parameters, which are zero, are not shown explicitly.

Similarly, we can compute other type-1 invariant product linear spaces $\hat{\mathbf{W}}_8^1$ with their vector differential operators $\mathbf{F}(\mathbf{U})$ for other cases such as $a_{10} = b_{20} = 0$, $a_{20} = b_{21} = 0$, etc. Moreover, in the same way, we can determine the other k -dimensional type-1 invariant product linear spaces $\hat{\mathbf{W}}_k^1$ for the vector differential operators $\mathbf{F}(\mathbf{U})$.

Remark 1. *It is important to mention that using the definition of type-2 invariant product linear spaces given in (2.20), we can consider the type-2 invariant product linear spaces $\hat{\mathbf{W}}_k^2$ for the given coupled equations (3.1) as follows:*

$$\hat{\mathbf{W}}_k^2 = W_{1,k_1}^2 \times W_{2,k_2}^2, \quad (3.11)$$

where the component spaces W_{s,k_s}^2 , $s = 1, 2$, are given by

$$\begin{aligned} W_{s,k_s}^2 &= \left\{ w_s(x_1, x_2) \mid \mathbf{L}_{s,k_s,i}^i [w_s(x_1, x_2)] = 0, \varphi_{s,0}^i(x_i) = 0, \frac{\partial^2 w_s(x_1, x_2)}{\partial x_1 \partial x_2} = 0, i = 1, 2 \right\} \\ &= \left\{ w_s(x_1, x_2) = \mu_{s,0} + \sum_{i,r_i} \mu_{s,r_i}^i \xi_{s,r_i}^i(x_i) \mid \mu_{s,0}, \mu_{s,r_i}^i \in \mathbb{R}, r_i = 1, \dots, k_{s,i} - 1, i = 1, 2 \right\}. \end{aligned}$$

Note that the dimension of $\hat{\mathbf{W}}_k^2$ is $k = \dim(\hat{\mathbf{W}}_k^2) = \sum_{s=1}^2 (k_{s,1} + k_{s,2} - 1)$. The above product linear space $\hat{\mathbf{W}}_k^2$ is invariant under $\mathbf{F}(\mathbf{U})$ if $F_s(\hat{\mathbf{W}}_k^2) = F_s(W_{1,k_1}^2 \times W_{2,k_2}^2) \subseteq W_{s,k_s}^2$, $\forall s = 1, 2$, whenever $\mathbf{U} = (u_1, u_2) \in \hat{\mathbf{W}}_k^2$, which reduce to the following forms of invariant conditions:

$$\begin{aligned} \mathbf{L}_{s,k_s,i}^i [u_s] = 0, \varphi_{s,0}^i(x_i) = 0, \frac{\partial^2 u_s}{\partial x_1 \partial x_2} = 0, \text{ implies that} \\ \mathbf{L}_{s,k_s,i}^i [F_s(u_1, u_2)] = 0, \varphi_{s,0}^i(x_r) = 0, \frac{\partial^2 F_s(u_1, u_2)}{\partial x_1 \partial x_2} = 0, s, i = 1, 2. \end{aligned} \quad (3.12)$$

Observe that for $k_{s,i} = 2$, $\varphi_{1,j}^i(x_i) = a_{ij} \in \mathbb{R}$, and $\varphi_{2,j}^i(x_i) = b_{ij} \in \mathbb{R}$, $j = 0, 1, s, i = 1, 2$, the above invariant conditions (3.12) read as

$$\begin{aligned} \left(\frac{d^2}{dx_r^2} + a_{r1} \frac{d}{dx_r} \right) F_1(u_1, u_2) = 0, \frac{\partial^2 F_1(u_1, u_2)}{\partial x_1 \partial x_2} = 0, \\ \left(\frac{d^2}{dx_r^2} + b_{r1} \frac{d}{dx_r} \right) F_2(u_1, u_2) = 0, \frac{\partial^2 F_2(u_1, u_2)}{\partial x_1 \partial x_2} = 0, \\ \text{whenever } \left(\frac{d^2}{dx_r^2} + a_{r1} \frac{d}{dx_r} \right) u_1 = 0, \frac{\partial^2 u_1}{\partial x_1 \partial x_2} = 0, \\ \text{and } \left(\frac{d^2}{dx_r^2} + b_{r1} \frac{d}{dx_r} \right) u_2 = 0, \frac{\partial^2 u_2}{\partial x_1 \partial x_2} = 0, r = 1, 2, \end{aligned} \quad (3.13)$$

where the differential operators $F_s(u_1, u_2)$, $s = 1, 2$, are considered as given in (3.10). Proceeding in a way similar to finding type-1 invariant product linear spaces earlier, we can obtain the various six-dimensional type-2 invariant product linear spaces $\hat{\mathbf{W}}_6^2$ and their corresponding cubic nonlinear vector

differential operator $\mathbf{F}(\mathbf{U})$ using the given invariant conditions (3.13). Also, we note that the type-2 invariant product linear space $\hat{\mathbf{W}}_6^2$ is a subspace of the type-1 invariant product linear space $\hat{\mathbf{W}}_8^1$ when $a_{r0} = b_{r0} = 0, r = 1, 2$. That means $\hat{\mathbf{W}}_6^2 \subset \hat{\mathbf{W}}_8^1$, if $a_{r0} = b_{r0} = 0, r = 1, 2$. However, the type-2 product linear space $\hat{\mathbf{W}}_6^2$ is preserved under the nonlinear vector differential operator $\mathbf{F}(\mathbf{U})$ need not imply the same operator $\mathbf{F}(\mathbf{U})$ admits the corresponding type-1 invariant product linear space $\hat{\mathbf{W}}_8^1$, in general.

$$\mathbf{L}_{s,k_s,i}^i[u_s] = 0, s = 1, 2, \dots, m, i = 1, 2, \dots, N,$$

$$\text{implies that } \mathbf{L}_{s,k_s,i}^i[A_s(\mathbf{U})] = 0, s = 1, 2, \dots, m, i = 1, 2, \dots, N.$$

Additionally, we can find the other possible type-2 invariant product linear spaces $\hat{\mathbf{W}}_k^2$ and their corresponding cubic nonlinear vector differential operator $\mathbf{F}(\mathbf{U})$ similarly using suitable values of $n_r^s, r, s = 1, 2$ in the given invariant conditions (3.12). Moreover, we observe that the type-2 invariant product linear spaces $\hat{\mathbf{W}}_k^2$ are more helpful in finding generalized separable analytical solutions of the given coupled equations (1.7), especially when it is not possible to construct a type-1 invariant product linear space $\hat{\mathbf{W}}_k^1$ for the given coupled equations (1.7).

3.2 Generalized separable analytical solutions for the given coupled equations (1.7)

This subsection provides a systematic study for deriving the generalized separable analytical solutions for the given coupled equations (1.7) using the type-1 and type-2 invariant product linear spaces discussed in the previous section.

3.2.1 Generalized separable analytical solutions for the initial value problems of the given coupled equations (1.7)

First, we discuss the systematic procedure for finding generalized separable analytical solutions for the initial value problems of the considered coupled equations (1.7) using the obtained type-1 and type-2 invariant product linear spaces.

Example 3.1. Consider the following multi-component (2+1)-dimensional coupled nonlinear TFDCWEs as

$$\begin{aligned} \frac{\partial^{\alpha_s} u_s}{\partial t^{\alpha_s}} = F_s(u_1, u_2) \equiv & \frac{\partial}{\partial x_1} \left[(k_{s0} - q_{s1}u_2) \frac{\partial u_1}{\partial x_1} + (q_{s1}u_1 + q_{s0}) \frac{\partial u_2}{\partial x_1} \right] \\ & + \frac{\partial}{\partial x_2} \left[(p_{s0} - c_{s1}u_2) \frac{\partial u_1}{\partial x_2} + (c_{s1}u_1 + c_{s0}) \frac{\partial u_2}{\partial x_2} \right] + \eta_{s0} \frac{\partial u_1}{\partial x_1} \\ & + g_{s0} \frac{\partial u_2}{\partial x_1} + d_{s0} \frac{\partial u_1}{\partial x_2} + h_{s0} \frac{\partial u_2}{\partial x_2}, \alpha_s \in (0, 2], s = 1, 2, \end{aligned} \quad (3.14)$$

with appropriate initial conditions

$$\begin{aligned} (i) \quad & u_s(x_1, x_2, 0) = \omega_s(x_1, x_2) \text{ if } \alpha_s \in (0, 1], s = 1, 2, \\ (ii) \quad & u_s(x_1, x_2, 0) = \omega_s(x_1, x_2), \frac{\partial u_s}{\partial t} \Big|_{t=0} = \vartheta_s(x_1, x_2) \text{ if } \alpha_s \in (1, 2], s = 1, 2. \end{aligned} \quad (3.15)$$

The above-coupled equations (3.14) admit the invariant product linear space

$$\hat{\mathbf{W}}_8^1 = \text{Span}\{1, x_1, x_2, x_1x_2\} \times \text{Span}\{1, x_1, x_2, x_1x_2\},$$

Table 1: Type-1 product linear spaces $\hat{\mathbf{W}}_k^1$ and their corresponding parameter restrictions of $\mathbf{F}(\mathbf{U})$ with $a_{11} = b_{21} = 0$ and $d_{10}, \eta_{10}, k_{10}, p_{10}, g_{20}, h_{20}, q_{20}, c_{20} \in \mathbb{R}$.

Cases	Some non-zero parameter restrictions of $\mathbf{F}(\mathbf{U})$	Type-1 invariant product linear space $\hat{\mathbf{W}}_8^1 = W_{1,4}^1 \times W_{2,4}^1$
1.	$\begin{cases} p_{12} = -c_{11}, p_{22} = -c_{21}, \\ c_{21}, c_{11}, c_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{\theta}x_1), \cos(\sqrt{\theta}x_1), x_2 \sin(\sqrt{\theta}x_1), \\ x_2 \cos(\sqrt{\theta}x_1)\}, \theta_1 = b_{10} - b_{11}^2 \\ W_{2,4}^1 = \text{Span}\{e^{-\theta_2 x_1}, e^{-\theta_3 x_1}, x_2 e^{-\theta_2 x_1}, x_2 e^{-\theta_3 x_1}\}, \\ \theta_2 = \frac{b_{11} + \sqrt{b_{11}^2 - 4b_{10}}}{2}, \theta_3 = \frac{b_{11} - \sqrt{b_{11}^2 - 4b_{10}}}{2} \end{cases}$
2.	$\begin{cases} p_{12} = -c_{11}, p_{22} = -c_{21}, \\ g_{10}, q_{10}, c_{21}, c_{11} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{e^{b_{11}x_1}, e^{-b_{11}x_1}, x_2 e^{b_{11}x_1}, x_2 e^{-b_{11}x_1}\} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2), e^{-b_{11}x_1} \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{20}}x_2), e^{-b_{11}x_1} \cos(\sqrt{b_{20}}x_2)\} \end{cases}$
3.	$\begin{cases} p_{12} = -c_{11}, p_{22} = -c_{21}, \\ q_{10}, c_{21}, c_{11}, c_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{a_{10}}x_1), \cos(\sqrt{a_{10}}x_1), \\ x_2 \sin(\sqrt{a_{10}}x_1), x_2 \cos(\sqrt{a_{10}}x_1)\} \\ W_{2,4}^1 = \text{Span}\{1, x_1, x_2, x_1 x_2\} \end{cases}$
4.	$\begin{cases} p_{12} = -c_{11}, p_{22} = -c_{21}, \\ c_{11}, c_{21} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{a_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{a_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2) \sin(\sqrt{a_{10}}x_1), \\ \cos(\sqrt{b_{20}}x_2) \cos(\sqrt{a_{10}}x_1)\} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2), x_1 \sin(\sqrt{b_{20}}x_2), \\ x_1 \cos(\sqrt{b_{20}}x_2)\} \end{cases}$
5.	$\begin{cases} p_{12} = -c_{11}, p_{22} = -c_{21}, \\ p_{20}, c_{21}, c_{11}, c_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{a_{10}}x_1), \cos(\sqrt{a_{10}}x_1), x_2 \sin(\sqrt{a_{10}}x_1), \\ x_2 \cos(\sqrt{a_{10}}x_1)\} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{10}}x_1), \cos(\sqrt{b_{10}}x_1), x_2 \sin(\sqrt{b_{10}}x_1), \\ x_2 \cos(\sqrt{b_{10}}x_1)\} \end{cases}$
6.	$\begin{cases} k_{12} = -q_{11}, k_{22} = -q_{21}, \\ q_{11}, q_{21}, q_{10}, c_{10}, k_{20} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{a_{20}}x_2), \cos(\sqrt{a_{20}}x_2), x_1 \sin(\sqrt{a_{20}}x_2), \\ x_1 \cos(\sqrt{a_{20}}x_2)\} \\ W_{2,4}^1 = \text{Span}\{1, x_1, x_2, x_1 x_2\} \end{cases}$
7.	$\begin{cases} k_{12} = -q_{11}, k_{22} = -q_{21}, \\ q_{11}, q_{21}, q_{10}, k_{20} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{a_{20}}x_2), \cos(\sqrt{a_{20}}x_2), x_1 \sin(\sqrt{a_{20}}x_2), \\ x_1 \cos(\sqrt{a_{20}}x_2)\} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2), x_1 \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2), \\ x_1 \cos(\sqrt{b_{20}}x_2)\} \end{cases}$
8.	$\begin{cases} k_{12} = -q_{11}, k_{22} = -q_{21}, \\ q_{11}, q_{21}, q_{10}, p_{20} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{1, x_1, x_2, x_1 x_2\} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2), x_1 \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2), \\ x_1 \cos(\sqrt{b_{20}}x_2)\} \end{cases}$
9.	$\begin{cases} p_{12} = -c_{11}, p_{22} = -c_{21}, \\ q_{10} = -\frac{b_{20}c_{10}}{b_{10}}, \\ c_{21}, c_{11}, c_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{a_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{a_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{a_{10}}x_1) \cos(\sqrt{b_{20}}x_2), \\ \sin(\sqrt{a_{10}}x_1) \cos(\sqrt{b_{20}}x_2)\} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{20}}x_2) \sin(\sqrt{b_{10}}x_1), \cos(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{10}}x_1) \cos(\sqrt{b_{20}}x_2)\} \end{cases}$
10.	$\begin{cases} p_{12} = -c_{11}, p_{22} = -c_{21}, \\ c_{21}, c_{11}, c_{10}, g_{10}, q_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{e^{b_{11}x_1}, e^{-b_{11}x_1}, e^{b_{11}x_1}x_2, e^{-b_{11}x_1}x_2\} \\ W_{2,4}^1 = \text{Span}\{1, e^{-b_{11}x_1}, x_2, x_2 e^{-b_{11}x_1}\} \end{cases}$

Table 2: Type-1 product linear spaces $\hat{\mathbf{W}}_k^1$ and their corresponding parameter restrictions of $\mathbf{F}(\mathbf{U})$ with $a_{11} = b_{21} = 0$ and $d_{10}, \eta_{10}, k_{10}, p_{10}, g_{20}, h_{20}, q_{20}, c_{20} \in \mathbb{R}$.

Cases	Some non-zero parameter restrictions of $\mathbf{F}(\mathbf{U})$	Type-1 invariant product linear space $\hat{\mathbf{W}}_8^1 = W_{1,4}^1 \times W_{2,4}^1$
11.	$\begin{cases} p_{12} = -c_{11}, p_{22} = -c_{21}, \\ q_{10} = \frac{g_{10}}{b_{11}}, c_{21}, c_{11}, \\ c_{10}, g_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{a_{10}}x_1), \cos(\sqrt{a_{10}}x_1), x_2 \sin(\sqrt{a_{10}}x_1), \\ x_2 \cos(\sqrt{a_{10}}x_1)\} \\ W_{2,4}^1 = \text{Span}\{1, x_2, e^{-b_{11}x_1}, x_2 e^{-b_{11}x_1}\} \end{cases}$
12.	$\begin{cases} p_{12} = -c_{11}, p_{22} = -c_{21}, \\ q_{10} = \frac{b_{11}g_{10} + b_{20}c_{10}}{b_{11}^2}, \\ g_{10}, c_{21}, c_{11}, c_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2), x_1 \sin(\sqrt{b_{20}}x_2), \\ x_1 \cos(\sqrt{a_{20}}x_2)\} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2), e^{-b_{11}x_1} \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{20}}x_2), e^{-b_{11}x_1} \cos(\sqrt{b_{20}}x_2)\} \end{cases}$
13.	$\begin{cases} p_{12} = -c_{11}, p_{22} = -c_{21}, \\ q_{10} = \frac{g_{10}}{b_{11}}, g_{10}, c_{21}, c_{11} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2) \sin(\sqrt{a_{10}}x_1), \\ \cos(\sqrt{b_{20}}x_2) \sin(\sqrt{a_{10}}x_1), \sin(\sqrt{b_{20}}x_2) \cos(\sqrt{a_{10}}x_1), \\ \cos(\sqrt{a_{10}}x_1) \cos(\sqrt{b_{20}}x_2)\} \\ W_{2,4}^1 = \text{Span}\{e^{-b_{11}x_1} \sin(\sqrt{b_{20}}x_2), \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{20}}x_2), e^{-b_{11}x_1} \cos(\sqrt{b_{20}}x_2)\} \end{cases}$
14.	$\begin{cases} p_{12} = -c_{11}, p_{22} = -c_{21}, \\ q_{10} = \frac{g_{10}}{b_{11}}, c_{10} = -\frac{b_{10}g_{10}}{b_{11}b_{20}}, \\ g_{10}, c_{21}, c_{11} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2) \sin(\theta x_1), \\ \cos(\sqrt{b_{20}}x_2) \sin(\theta x_1), \sin(\sqrt{b_{20}}x_2) \cos(\theta x_1), \\ \cos(\theta x_1) \cos(\sqrt{b_{20}}x_2)\}, \theta = \sqrt{b_{10} - b_{11}^2} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2) \sin(\sqrt{b_{10}}x_1), \\ \cos(\sqrt{b_{20}}x_2) \sin(\sqrt{b_{10}}x_1), \cos(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{10}}x_1) \cos(\sqrt{b_{20}}x_2)\} \end{cases}$
15.	$\begin{cases} k_{12} = -\frac{c_{11}a_{20}}{a_{10}}, \\ q_{11} = \frac{c_{11}a_{20}}{a_{10}}, \\ p_{12} = -c_{11}, c_{21}, c_{11}, \\ q_{10}, c_{10} \in \mathbb{R}, \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{a_{10}}x_1) \sin(\sqrt{a_{20}}x_2), \\ \cos(\sqrt{a_{10}}x_1) \sin(\sqrt{a_{20}}x_2), \cos(\sqrt{a_{20}}x_2) \sin(\sqrt{a_{10}}x_1), \\ \cos(\sqrt{a_{20}}x_2) \cos(\sqrt{a_{10}}x_1)\} \\ W_{2,4}^1 = \text{Span}\{1, x_1, x_2, x_1x_2\} \end{cases}$
16.	$\begin{cases} p_{12} = -c_{11}, q_{11} = \frac{c_{11}k_{20}}{p_{20}}, \\ k_{12} = -\frac{c_{11}k_{20}}{p_{20}}, \\ c_{10}, p_{20}, q_{10}, k_{20} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{e^{\theta x_1} \sin(\sqrt{a_{20}}x_2), e^{\theta x_1} \cos(\sqrt{a_{20}}x_2), \\ e^{-\theta x_1} \sin(\sqrt{a_{20}}x_2), e^{-\theta x_1} \cos(\sqrt{a_{20}}x_2)\}, \\ \theta = \sqrt{\frac{p_{20}a_{20}}{k_{20}}} \\ W_{2,4}^1 = \text{Span}\{1, x_1, x_2, x_1x_2\} \end{cases}$
17.	$\begin{cases} k_{12} = -\frac{c_{11}b_{20}}{a_{10}}, \\ q_{11} = \frac{c_{11}b_{20}}{a_{10}}, p_{12} = -c_{11}, \\ c_{11}, p_{10}, q_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{a_{10}}x_1), \cos(\sqrt{a_{10}}x_1), x_2 \sin(\sqrt{a_{10}}x_1), \\ x_2 \cos(\sqrt{a_{10}}x_1)\} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2), x_1 \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2), \\ x_1 \cos(\sqrt{b_{20}}x_2)\} \end{cases}$
18.	$\begin{cases} k_{12} = \frac{c_{11}(a_{20} - b_{20})}{a_{10}}, \\ q_{11} = \frac{(b_{20} - a_{20})c_{11}}{a_{10}}, \\ p_{12} = -c_{11}, c_{11}, q_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{a_{10}}x_1) \sin(\sqrt{a_{20}}x_2), \\ \cos(\sqrt{a_{10}}x_1) \sin(\sqrt{a_{20}}x_2), \cos(\sqrt{a_{20}}x_2) \sin(\sqrt{a_{10}}x_1), \\ \cos(\sqrt{a_{20}}x_2) \cos(\sqrt{a_{10}}x_1)\} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2), x_1 \sin(\sqrt{b_{20}}x_2), \\ x_1 \cos(\sqrt{b_{20}}x_2)\} \end{cases}$
19.	$\begin{cases} k_{12} = -\frac{q_{21}c_{11}}{c_{21}}, \\ k_{22} = -q_{21}, p_{12} = -c_{11}, \\ p_{22} = -c_{21}, q_{11} = \frac{q_{21}c_{11}}{c_{21}}, \\ c_{11}, c_{21}, q_{21}, q_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{\theta}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{\theta}x_1) \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2) \sin(\sqrt{\theta}x_1), \\ \cos(\sqrt{b_{20}}x_2) \cos(\sqrt{\theta}x_1)\}, \theta = \frac{(b_{20} - a_{20})c_{21}}{q_{21}} > 0 \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2), x_1 \sin(\sqrt{b_{20}}x_2), \\ x_1 \cos(\sqrt{b_{20}}x_2)\} \end{cases}$

Table 3: Type-1 product linear spaces $\hat{\mathbf{W}}_k^1$ and their corresponding parameter restrictions of $\mathbf{F}(\mathbf{U})$ with $a_{11} = b_{21} = 0$ and $d_{10}, \eta_{10}, k_{10}, p_{10}, g_{20}, h_{20}, q_{20}, c_{20} \in \mathbb{R}$.

Cases	Some non-zero parameter restrictions of $\mathbf{F}(\mathbf{U})$	Type-1 invariant product linear space $\hat{\mathbf{W}}_8^1 = W_{1,4}^1 \times W_{2,4}^1$
20.	$\begin{cases} p_{12} = -c_{11}, p_{22} = -c_{21}, \\ k_{12} = -\frac{c_{11}q_{21}}{c_{21}}, \\ q_{11} = \frac{c_{11}q_{21}}{c_{21}}, k_{22} = -q_{21}, \\ k_{20} = \frac{p_{20}q_{21}}{c_{21}}, c_{11}, c_{10}, \\ p_{20}, c_{21}, q_{10}, q_{21} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{e^{\theta x_1} \sin(\sqrt{a_{20}}x_2), \\ e^{-\theta x_1} \sin(\sqrt{a_{20}}x_2), e^{\theta x_1} \cos(\sqrt{a_{20}}x_2), \\ \cos(\sqrt{a_{20}}x_2)e^{-\theta x_1}\}, \theta = \sqrt{\frac{c_{21}a_{20}}{q_{21}}} \\ W_{2,4}^1 = \text{Span}\{1, x_1, x_2, x_1x_2\} \end{cases}$
21.	$\begin{cases} k_{12} = -\frac{q_{21}c_{11}}{c_{21}}, \\ k_{22} = -q_{21}, p_{12} = -c_{11}, \\ p_{22} = -c_{21}, q_{11} = \frac{q_{21}c_{11}}{c_{21}}, \\ p_{20}, q_{10}, c_{11}, c_{21} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{\theta}x_1), \cos(\sqrt{\theta}x_1), x_2 \sin(\sqrt{\theta}x_1), \\ x_2 \cos(\sqrt{\theta}x_1)\}, \theta = \frac{c_{21}b_{20}}{q_{21}} > 0 \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2), x_1 \sin(\sqrt{b_{20}}x_2), \\ x_1 \cos(\sqrt{b_{20}}x_2)\} \end{cases}$
22.	$\begin{cases} k_{12} = \frac{(b_{20}-a_{20})c_{11}k_{20}}{(a_{20}p_{20})}, \\ q_{11} = \frac{c_{11}(a_{20}-b_{20})k_{20}}{(a_{20}p_{20})}, \\ p_{12} = -c_{11}, p_{20}, k_{20}, \\ q_{10}, c_{11} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{e^{\sqrt{\theta}x_1} \sin(\sqrt{a_{20}}x_2), e^{-\sqrt{\theta}x_1} \sin(\sqrt{a_{20}}x_2), \\ \cos(\sqrt{a_{20}}x_2)e^{\sqrt{\theta}x_1}, \cos(\sqrt{a_{20}}x_2)e^{-\sqrt{\theta}x_1}\}, \\ \theta = \frac{p_{20}a_{20}}{k_{20}} > 0 \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2), x_1 \sin(\sqrt{b_{20}}x_2), \\ x_1 \cos(\sqrt{b_{20}}x_2)\} \end{cases}$
23.	$\begin{cases} k_{22} = -q_{21}, p_{12} = -c_{11}, \\ p_{22} = -c_{21}, q_{11} = \frac{q_{21}c_{11}}{c_{21}} \\ k_{12} = -\frac{q_{21}c_{11}}{c_{21}}, p_{20}, \\ k_{20}, q_{10}, q_{21}, c_{21}, c_{11} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{e^{\sqrt{\theta}(x_1+x_2)}, e^{-\sqrt{\theta}(x_1+x_2)}, e^{\sqrt{\theta}(x_1-x_2)}, \\ e^{-\sqrt{\theta}(x_1-x_2)}\}, \theta = \frac{c_{21}b_{20}p_{20}}{k_{20}c_{21}-p_{20}q_{21}} > 0 \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2), x_1 \sin(\sqrt{b_{20}}x_2), \\ x_1 \cos(\sqrt{b_{20}}x_2)\} \end{cases}$
24.	$\begin{cases} k_{12} = -q_{11}, k_{22} = -q_{21}, \\ p_{22} = -c_{21}, d_{20}, h_{10}, \\ p_{20}, c_{21}, c_{10}, q_{11}, \\ q_{21}, q_{10}, k_{20} \in \mathbb{R} \\ k_{12} = -q_{11}, k_{22} = -q_{21}, \\ p_{12} = -c_{11}, p_{22} = -c_{21}, \\ d_{20}, h_{10}, p_{20}, c_{21}, c_{11}, c_{10}, \\ q_{11}, q_{21}, q_{10}, k_{20} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2) \sin(\sqrt{b_{10}}x_1), \\ \cos(\sqrt{b_{20}}x_2) \cos(\sqrt{b_{10}}x_1)\} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2) \sin(\sqrt{b_{10}}x_1), \\ \cos(\sqrt{b_{20}}x_2) \cos(\sqrt{b_{10}}x_1)\} \end{cases}$
25.	$\begin{cases} k_{12} = -q_{11}, k_{22} = -q_{21}, \\ q_{10} = -\frac{b_{20}c_{10}}{b_{10}}, \\ c_{10}, q_{11}, q_{21}, q_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{b_{10}}x_1) \sin(\sqrt{a_{20}}x_2), \\ \cos(\sqrt{b_{10}}x_1) \sin(\sqrt{a_{20}}x_2), \cos(\sqrt{a_{20}}x_2) \sin(\sqrt{b_{10}}x_1), \\ \cos(\sqrt{a_{20}}x_2) \cos(\sqrt{b_{10}}x_1)\} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2) \sin(\sqrt{b_{10}}x_1), \\ \cos(\sqrt{b_{20}}x_2) \cos(\sqrt{b_{10}}x_1)\} \end{cases}$
26.	$\begin{cases} k_{12} = -\frac{q_{21}c_{11}}{c_{21}}, k_{22} = -q_{21}, \\ q_{11} = \frac{q_{21}c_{11}}{c_{21}}, p_{12} = -c_{11}, \\ q_{10} = -\frac{b_{20}c_{10}}{b_{10}}, p_{22} = -c_{21}, \\ p_{20}, q_{21}, c_{11}, c_{21}, c_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{\theta}x_1), \cos(\sqrt{\theta}x_1), x_2 \sin(\sqrt{\theta}x_1), \\ x_2 \cos(\sqrt{\theta}x_1)\}, \theta = \frac{(b_{10}q_{21}+b_{20}c_{21})}{q_{21}} > 0 \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2) \sin(\sqrt{b_{10}}x_1), \\ \cos(\sqrt{b_{20}}x_2) \cos(\sqrt{b_{10}}x_1)\} \end{cases}$

Table 4: Type-1 product linear spaces $\hat{\mathbf{W}}_k^1$ and their corresponding parameter restrictions of $\mathbf{F}(\mathbf{U})$ with $a_{11} = b_{21} = 0$ and $d_{10}, \eta_{10}, k_{10}, p_{10}, g_{20}, h_{20}, q_{20}, c_{20} \in \mathbb{R}$.

Cases	Some non-zero parameter restrictions of $\mathbf{F}(\mathbf{U})$	Type-1 invariant product linear space $\hat{\mathbf{W}}_8^1 = W_{1,4}^1 \times W_{2,4}^1$
27.	$\begin{cases} k_{12} = \frac{b_{20}c_{11}}{b_{10}-a_{10}}, \\ q_{10} = -\frac{b_{20}c_{10}}{b_{10}}, \\ p_{12} = -c_{11}, p_{20}, c_{11}, \\ c_{10} \in \mathbb{R}, q_{11} = \frac{c_{11}b_{20}}{a_{10}-b_{10}} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{a_{10}}x_1), x_2 \sin(\sqrt{a_{10}}x_1), \\ \cos(\sqrt{a_{10}}x_1), x_2 \cos(\sqrt{a_{10}}x_1)\} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2) \sin(\sqrt{b_{10}}x_1), \\ \cos(\sqrt{b_{20}}x_2) \cos(\sqrt{b_{10}}x_1)\} \end{cases}$
28.	$\begin{cases} k_{12} = -q_{11}, k_{22} = -q_{21}, \\ q_{10} = -\frac{b_{20}c_{10}}{b_{10}}, c_{10}, q_{11}, \\ q_{21}, q_{10}, p_{20}, k_{20} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{b_{10}}x_1)e^{\sqrt{\theta}x_2}, \\ \cos(\sqrt{b_{10}}x_1)e^{\sqrt{\theta}x_2}, e^{-\sqrt{\theta}x_2} \sin(\sqrt{b_{10}}x_1), \\ e^{-\sqrt{\theta}x_2} \cos(\sqrt{b_{10}}x_1)\}, \theta = \frac{k_{20}b_{10}}{p_{20}} > 0 \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2) \sin(\sqrt{b_{10}}x_1), \\ \cos(\sqrt{b_{20}}x_2) \cos(\sqrt{b_{10}}x_1)\} \end{cases}$
29.	$\begin{cases} k_{12} = \frac{c_{11}(a_{20}-b_{20})}{(a_{10}-b_{10})}, \\ p_{12} = -c_{11}, q_{10} = -\frac{b_{20}c_{10}}{b_{10}}, \\ q_{11} = \frac{(b_{20}-a_{20})c_{11}}{(a_{10}-b_{10})}, \\ c_{11}, c_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{a_{10}}x_1) \sin(\sqrt{a_{20}}x_2), \\ \cos(\sqrt{a_{10}}x_1) \sin(\sqrt{a_{20}}x_2), \cos(\sqrt{a_{20}}x_2) \sin(\sqrt{a_{10}}x_1), \\ \cos(\sqrt{a_{20}}x_2) \cos(\sqrt{a_{10}}x_1)\} \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2) \sin(\sqrt{b_{10}}x_1), \\ \cos(\sqrt{b_{20}}x_2) \cos(\sqrt{b_{10}}x_1)\} \end{cases}$
30.	$\begin{cases} k_{12} = -\frac{q_{21}c_{11}}{c_{21}}, k_{22} = -q_{21}, \\ p_{12} = -c_{11}, p_{22} = -c_{21}, \\ q_{10} = -\frac{b_{20}c_{10}}{b_{10}}, \\ q_{11} = \frac{q_{21}c_{11}}{c_{21}}, q_{21}, \\ c_{21}, c_{11}, c_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{\sin(\sqrt{\theta}x_1) \sin(\sqrt{a_{20}}x_2), \\ \cos(\sqrt{\theta}x_1) \sin(\sqrt{a_{20}}x_2), \cos(\sqrt{a_{20}}x_2) \sin(\sqrt{\theta}x_1), \\ \cos(\sqrt{a_{20}}x_2) \cos(\sqrt{\theta}x_1)\}, \theta = \frac{(b_{10}q_{21}-c_{21}(a_{20}-b_{20}))}{q_{21}} > 0 \\ W_{2,4}^1 = \text{Span}\{\sin(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \\ \cos(\sqrt{b_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2) \sin(\sqrt{b_{10}}x_1), \\ \cos(\sqrt{b_{20}}x_2) \cos(\sqrt{b_{10}}x_1)\} \end{cases}$
31.	$\begin{cases} k_{12} = -q_{11}, k_{22} = -q_{21}, \\ p_{12} = \frac{d_{12}p_{20}}{d_{20}}, \\ c_{11} = -\frac{d_{12}p_{20}}{d_{20}}, d_{12}, d_{20}, h_{10}, \\ k_{20}, p_{20}, c_{10}, q_{10}, q_{11}, q_{21} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{1, x_1, e^{-\frac{d_{20}}{p_{20}}x_2}, x_1 e^{-\frac{d_{20}}{p_{20}}x_2}\} \\ W_{2,4}^1 = \text{Span}\{1, x_1, x_2, x_1 x_2\} \end{cases}$
32.	$\begin{cases} k_{12} = -\frac{g_{11}}{b_{11}}, k_{22} = -\frac{g_{21}}{b_{11}}, \\ q_{10} = \frac{g_{10}}{b_{11}}, q_{11} = \frac{g_{11}}{b_{11}}, \\ q_{21} = \frac{g_{21}}{b_{11}}, g_{11}, g_{21}, \\ g_{10}, c_{10}, k_{20} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{e^{-\theta_1 x_2}, x_1 e^{-\theta_1 x_2}, e^{-\theta_2 x_2}, \\ x_1 e^{-\theta_2 x_2}\}, \theta_1 = \frac{a_{21} + \sqrt{a_{21}^2 - 4a_{20}}}{2}, \\ \theta_2 = \frac{a_{21} - \sqrt{a_{21}^2 - 4a_{20}}}{2} \\ W_{2,4}^1 = \text{Span}\{1, e^{-b_{11}x_1}, x_2, x_2 e^{-b_{11}x_1}\} \end{cases}$
33.	$\begin{cases} k_{12} = -q_{11}, k_{22} = -q_{21}, \\ q_{11}, q_{21}, q_{10}, c_{10}, k_{20} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{e^{-\theta_1 x_2}, x_1 e^{-\theta_1 x_2}, e^{-\theta_2 x_2}, \\ x_1 e^{-\theta_2 x_2}\}, \theta_1 = \frac{a_{21} + \sqrt{a_{21}^2 - 4a_{20}}}{2}, \\ \theta_2 = \frac{a_{21} - \sqrt{a_{21}^2 - 4a_{20}}}{2} \\ W_{2,4}^1 = \text{Span}\{1, x_1, x_2, x_1 x_2\} \end{cases}$

Table 5: Type-1 product linear spaces \hat{W}_k^1 and their corresponding parameter restrictions of $\mathbf{F}(\mathbf{U})$ with $a_{11} = b_{21} = 0$ and $d_{10}, \eta_{10}, k_{10}, p_{10}, g_{20}, h_{20}, q_{20}, c_{20} \in \mathbb{R}$.

Cases	Some non-zero parameter restrictions of $\mathbf{F}(\mathbf{U})$	Type-1 invariant product linear space $\hat{W}_8^1 = W_{1,4}^1 \times W_{2,4}^1$
34.	$\begin{cases} k_{i2} = -\frac{g_{i1}}{b_{11}}, p_{i2} = -c_{i1}, \\ q_{i1} = \frac{g_{i1}}{b_{11}}, i = 1, 2, \\ q_{10} = \frac{g_{10}}{b_{11}}, \eta_{20}, g_{10}, g_{21}, g_{11}, \\ p_{20}, k_{20}, c_{10}, c_{11}, c_{21} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{1, x_1, x_2, x_1 x_2\} \\ W_{2,4}^1 = \text{Span}\{1, e^{-b_{11}x_1}, x_2, x_2 e^{-b_{11}x_1}\} \end{cases}$
35.	$\begin{cases} p_{12} = -\frac{d_{12}c_{21}}{d_{22}}, k_{12} = -q_{11}, \\ p_{20} = -\frac{d_{20}c_{21}}{d_{22}}, k_{22} = -q_{21}, \\ c_{11} = \frac{d_{12}c_{21}}{d_{22}}, p_{22} = -c_{21}, \\ d_{12}, d_{22}, d_{20}, h_{10}, k_{20}, c_{10}, \\ c_{21}, q_{10}, q_{11}, q_{21} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{1, x_1, e^{\frac{d_{22}}{c_{21}}x_2}, x_1 e^{\frac{d_{22}}{c_{21}}x_2}\} \\ W_{2,4}^1 = \text{Span}\{1, x_1, x_2, x_1 x_2\} \end{cases}$
36.	$\begin{cases} k_{12} = -\frac{g_{11}}{b_{11}}, k_{22} = -\frac{g_{21}}{b_{11}}, \\ p_{12} = \frac{d_{12}}{a_{21}}, q_{10} = \frac{g_{10}}{b_{11}}, \\ q_{11} = \frac{g_{11}}{b_{11}}, q_{21} = \frac{g_{21}}{b_{11}}, \\ c_{11} = -\frac{d_{12}}{a_{21}}, d_{12}, k_{20}, g_{10}, \\ g_{21}, g_{11}, q_{10}, q_{11}, q_{21}, c_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{1, x_1, e^{-a_{21}x_2}, x_1 e^{-a_{21}x_2}\} \\ W_{1,4}^1 = \text{Span}\{1, e^{-b_{11}x_1}, x_2, e^{-b_{11}x_1}x_2\} \end{cases}$
37.	$\begin{cases} k_{12} = -\frac{g_{11}}{b_{11}}, k_{22} = -\frac{g_{21}}{b_{11}}, \\ p_{12} = \frac{d_{12}p_{20}}{d_{20}}, q_{10} = \frac{g_{10}}{b_{11}}, \\ q_{11} = \frac{g_{11}}{b_{11}}, q_{21} = \frac{g_{21}}{b_{11}}, \\ c_{11} = -\frac{d_{12}p_{20}}{d_{20}}, d_{12}, d_{20}, k_{20}, p_{20}, \\ g_{10}, g_{21}, g_{11}, q_{10}, q_{11}, q_{21}, c_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{1, x_1, e^{-\frac{d_{20}}{p_{20}}x_2}, x_1 e^{-\frac{d_{20}}{p_{20}}x_2}\} \\ W_{2,4}^1 = \text{Span}\{1, e^{-b_{11}x_1}, x_2, e^{-b_{11}x_1}x_2\} \end{cases}$
38.	$\begin{cases} k_{12} = -\frac{g_{11}}{b_{11}}, k_{22} = -\frac{g_{21}}{b_{11}}, \\ p_{12} = -\frac{d_{12}c_{21}}{d_{22}}, p_{20} = -\frac{d_{20}c_{21}}{d_{22}}, \\ q_{10} = \frac{g_{10}}{b_{11}}, q_{11} = \frac{g_{11}}{b_{11}}, \\ p_{22} = -c_{21}, c_{11} = \frac{d_{12}c_{21}}{d_{22}}, \\ d_{22}, d_{12}, d_{20}, g_{10}, g_{21}, g_{11}, k_{20}, \\ q_{21}, c_{10}, c_{21} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{1, x_1, e^{\frac{d_{22}}{c_{21}}x_2}, x_1 e^{\frac{d_{22}}{c_{21}}x_2}\} \\ W_{2,4}^1 = \text{Span}\{1, e^{-b_{11}x_1}, x_2, e^{-b_{11}x_1}x_2\} \end{cases}$
39.	$\begin{cases} k_{12} = -q_{11}, k_{22} = -q_{21}, \\ p_{12} = -c_{11}, p_{22} = -c_{21}, \\ d_{20}, \eta_{20}, k_{20}, p_{20}, g_{10}, \\ h_{10}, q_{10}, q_{11}, q_{21}, c_{21}, c_{11}, c_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{1, x_1, x_2, x_1 x_2\} \\ W_{2,4}^1 = \text{Span}\{1, x_1, x_2, x_1 x_2\} \end{cases}$
40.	$\begin{cases} k_{12} = -q_{11}, k_{22} = -q_{21}, p_{12} = \frac{d_{12}}{a_{21}}, c_{11} = -\frac{d_{12}}{a_{21}}, \\ d_{12}, k_{20}, h_{10}, q_{10}, q_{11}, q_{21}, c_{10} \in \mathbb{R} \end{cases}$	$\begin{cases} W_{1,4}^1 = \text{Span}\{1, x_1, e^{-a_{21}x_2}, x_1 e^{-a_{21}x_2}\} \\ W_{2,4}^1 = \text{Span}\{1, x_1, x_2, x_1 x_2\} \end{cases}$

which is discussed in case 39 of Table 5. Then, there exists a generalized separable analytical solution for the coupled equations (3.14) in the form

$$u_s(x_1, x_2, t) = \delta_{s1}(t) + \delta_{s2}(t)x_1 + \delta_{s3}(t)x_2 + \delta_{s4}(t)x_1x_2, s = 1, 2. \quad (3.16)$$

Now, we want to find the unknown functions $\delta_{si}(t), i = 1, 2, 3, 4$, of the above analytical solution $u_s = u_s(x_1, x_2, t), s = 1, 2$. Thus, we substitute (3.16) into the above-coupled equations (3.14). Hence, we obtain the following system of FODEs:

$$\begin{aligned} \mathbf{D}_t^{\alpha_s} \delta_{s1}(t) &= g_{s0}\delta_{22}(t) + h_{s0}\delta_{23}(t) + \eta_{s0}\delta_{12}(t) + d_{s0}\delta_{13}(t), \\ \mathbf{D}_t^{\alpha_s} \delta_{s2}(t) &= h_{s0}\delta_{24}(t) + d_{s0}\delta_{14}(t), \\ \mathbf{D}_t^{\alpha_s} \delta_{s3}(t) &= g_{s0}\delta_{24}(t) + \eta_{s0}\delta_{14}(t), \text{ and} \\ \mathbf{D}_t^{\alpha_s} \delta_{s4}(t) &= 0, s = 1, 2, \end{aligned} \quad (3.17)$$

where $\mathbf{D}_t^{\alpha_s}(\cdot) = \frac{d^{\alpha_s}}{dt^{\alpha_s}}(\cdot)$ represents the Caputo fractional-order derivative (1.2) of order $\alpha_s \in (0, 2], s = 1, 2$, with respect to t .

Next, we use the Laplace transformation method to solve the above-reduced system of FODEs (3.17). In [6], the Laplace transformation of the Caputo derivative with fractional-order $\alpha > 0$ is obtained as follows:

$$L(\mathbf{D}_t^{\alpha} \delta(t); s) = s^{\alpha} L(\delta(t); s) - \sum_{q=0}^{n-1} s^{\alpha-(q+1)} \frac{d^q \delta(t)}{dt^q} \Big|_{t=0}, \text{ Re}(s) > 0,$$

where $\alpha \in (n-1, n], n \in \mathbb{N}$. Here, we apply the Laplace transformation on the last equation of the obtained system of FODEs (3.17), and then taking the inverse Laplace transformation of the resultant equation, we have

$$\delta_{s4}(t) = \begin{cases} A_{s4}, & \text{if } \alpha_s \in (0, 1], \\ A_{s4} + B_{s4}t, & \text{if } \alpha_s \in (1, 2], \end{cases} \quad (3.18)$$

where $A_{s4}, B_{s4} \in \mathbb{R}, s = 1, 2$.

Now, we substitute the above-obtained function $\delta_{s4}, s = 1, 2$, in the second and third equations of the given system of FODEs (3.17). Thus, using the Laplace and the inverse Laplace transformations, we get

$$\delta_{12}(t) = \begin{cases} A_{12} + \frac{\mu_1 t^{\alpha_1}}{\Gamma(\alpha_1 + 1)}, & \text{if } \alpha_s \in (0, 1], s = 1, 2, \\ A_{12} + B_{12}t + \left(\frac{\mu_1}{\Gamma(\alpha_1 + 1)} + \frac{d_{10}B_{14}t}{\Gamma(\alpha_1 + 2)} \right) t^{\alpha_1}, & \text{if } \alpha_1 \in (1, 2] \ \& \ \alpha_2 \in (0, 1], \\ A_{12} + \left(\frac{\mu_1}{\Gamma(\alpha_1 + 1)} + \frac{h_{10}B_{24}t}{\Gamma(\alpha_1 + 2)} \right) t^{\alpha_1}, & \text{if } \alpha_1 \in (0, 1] \ \& \ \alpha_2 \in (1, 2], \\ A_{12} + B_{12}t + \left(\frac{\mu_1}{\Gamma(\alpha_1 + 1)} + \frac{\nu_1 t}{\Gamma(\alpha_1 + 2)} \right) t^{\alpha_1}, & \text{if } \alpha_s \in (1, 2], s = 1, 2, \end{cases} \quad (3.19)$$

$$\delta_{22}(t) = \begin{cases} A_{22} + \frac{\mu_2 t^{\alpha_2}}{\Gamma(\alpha_2 + 1)}, & \text{if } \alpha_s \in (0, 1], s = 1, 2, \\ A_{22} + \left(\frac{\mu_2}{\Gamma(\alpha_2 + 1)} + \frac{d_{20}B_{14}t}{\Gamma(\alpha_2 + 2)} \right) t^{\alpha_2}, & \text{if } \alpha_1 \in (1, 2] \ \& \ \alpha_2 \in (0, 1], \\ A_{22} + B_{22}t + \left(\frac{\mu_2}{\Gamma(\alpha_2 + 1)} + \frac{h_{20}B_{24}t}{\Gamma(\alpha_2 + 2)} \right) t^{\alpha_2}, & \text{if } \alpha_1 \in (0, 1] \ \& \ \alpha_2 \in (1, 2], \\ A_{22} + B_{22}t + \left(\frac{\mu_2}{\Gamma(\alpha_2 + 1)} + \frac{\nu_2 t}{\Gamma(\alpha_2 + 2)} \right) t^{\alpha_2}, & \text{if } \alpha_s \in (1, 2], s = 1, 2, \end{cases} \quad (3.20)$$

$$\delta_{13}(t) = \begin{cases} A_{13} + \frac{\gamma_1 t^{\alpha_1}}{\Gamma(\alpha_1 + 1)}, & \text{if } \alpha_s \in (0, 1], s = 1, 2, \\ A_{13} + B_{13}t + \left(\frac{\gamma_1}{\Gamma(\alpha_1 + 1)} + \frac{\eta_{10} B_{14} t}{\Gamma(\alpha_1 + 2)} \right) t^{\alpha_1}, & \text{if } \alpha_1 \in (1, 2] \& \alpha_2 \in (0, 1], \\ A_{13} + \left(\frac{\gamma_1}{\Gamma(\alpha_1 + 1)} + \frac{g_{10} B_{24} t}{\Gamma(\alpha_1 + 2)} \right) t^{\alpha_1}, & \text{if } \alpha_1 \in (0, 1] \& \alpha_2 \in (1, 2], \\ A_{13} + B_{13}t + \left(\frac{\gamma_1}{\Gamma(\alpha_1 + 1)} + \frac{\kappa_1 t}{\Gamma(\alpha_1 + 2)} \right) t^{\alpha_1}, & \text{if } \alpha_s \in (1, 2], s = 1, 2, \text{ and} \end{cases} \quad (3.21)$$

$$\delta_{23}(t) = \begin{cases} A_{23} + \frac{\gamma_2 t^{\alpha_2}}{\Gamma(\alpha_2 + 1)}, & \text{if } \alpha_s \in (0, 1], s = 1, 2, \\ A_{23} + \left(\frac{\gamma_2}{\Gamma(\alpha_2 + 1)} + \frac{\eta_{20} B_{14} t}{\Gamma(\alpha_2 + 2)} \right) t^{\alpha_2}, & \text{if } \alpha_1 \in (1, 2] \& \alpha_2 \in (0, 1], \\ A_{23} + B_{23}t + \left(\frac{\gamma_2}{\Gamma(\alpha_2 + 1)} + \frac{g_{20} B_{24} t}{\Gamma(\alpha_2 + 2)} \right) t^{\alpha_2}, & \text{if } \alpha_1 \in (0, 1] \& \alpha_2 \in (1, 2], \\ A_{23} + B_{23}t + \left(\frac{\gamma_2}{\Gamma(\alpha_2 + 1)} + \frac{\kappa_2 t}{\Gamma(\alpha_2 + 2)} \right) t^{\alpha_2}, & \text{if } \alpha_s \in (1, 2], s = 1, 2, \end{cases} \quad (3.22)$$

where $\gamma_s = g_{s0}A_{24} + \eta_{s0}A_{14}$, $\kappa_s = g_{s0}B_{24} + \eta_{s0}B_{14}$, $\mu_s = h_{s0}A_{24} + d_{s0}A_{14}$, $\nu_s = h_{s0}B_{24} + d_{s0}B_{14}$, and $A_{si}, B_{si} \in \mathbb{R}, i = 2, 3, 4, s = 1, 2$.

Finally, we substitute the equations (3.18)-(3.22) in the first equation of the obtained system (3.17), and then using the Laplace and inverse Laplace transformations, we have

$$\delta_{21}(t) = \begin{cases} A_{21} + \left(\frac{l_2}{\Gamma(\alpha_2 + 1)} + \sum_{i=1}^2 \frac{\varrho_{2i} t^{\alpha_i}}{\Gamma(\alpha_2 + \alpha_i + 1)} \right) t^{\alpha_2}, & \text{if } \alpha_s \in (0, 1], s = 1, 2, \\ A_{21} + \left(\frac{l_2}{\Gamma(\alpha_2 + 1)} + \frac{(\eta_{20} B_{12} + d_{20} B_{13})t}{\Gamma(\alpha_2 + 2)} + \frac{(g_{20} d_{20} + h_{20} \eta_{20}) B_{14} t^{\alpha_2 + 1}}{\Gamma(2\alpha_2 + 2)} \right. \\ \left. + \sum_{i=1}^2 \frac{\varrho_{2i} t^{\alpha_i}}{\Gamma(\alpha_2 + \alpha_i + 1)} + \frac{(d_{20} \eta_{10} + d_{10} \eta_{20}) B_{14} t^{\alpha_1 + 1}}{\Gamma(\alpha_1 + \alpha_2 + 2)} \right) t^{\alpha_2}, & \\ \text{if } \alpha_1 \in (1, 2] \& \alpha_2 \in (0, 1], \\ A_{21} + B_{21}t + \left(\frac{l_2}{\Gamma(\alpha_2 + 1)} + \sum_{i=1}^2 \frac{\varrho_{2i} t^{\alpha_i}}{\Gamma(\alpha_2 + \alpha_i + 1)} + \frac{(g_{20} B_{22} + h_{20} B_{23})t}{\Gamma(\alpha_2 + 2)} \right. \\ \left. + \frac{2g_{20} h_{20} B_{24} t^{\alpha_2 + 1}}{\Gamma(2\alpha_2 + 2)} + \frac{B_{24} (\eta_{20} h_{10} + d_{20} g_{10}) t^{\alpha_1 + 1}}{\Gamma(\alpha_1 + \alpha_2 + 2)} \right) t^{\alpha_2}, & \\ \text{if } \alpha_1 \in (0, 1] \& \alpha_2 \in (1, 2], \\ A_{21} + B_{21}t + \frac{l_2 t^{\alpha_2}}{\Gamma(\alpha_2 + 1)} + \frac{\varsigma_2 t^{\alpha_2 + 1}}{\Gamma(\alpha_2 + 2)} + \sum_{i=1}^2 \left(\frac{\varrho_{2i} t^{\alpha_i}}{\Gamma(\alpha_i + \alpha_2 + 1)} \right. \\ \left. + \frac{\lambda_{2i} t^{\alpha_i + 1}}{\Gamma(\alpha_i + \alpha_2 + 2)} \right) t^{\alpha_2}, & \text{if } \alpha_s \in (1, 2], s = 1, 2, \end{cases} \quad (3.23)$$

$$\delta_{11}(t) = \begin{cases} A_{11} + \left(\frac{l_1}{\Gamma(\alpha_1 + 1)} + \sum_{i=1}^2 \frac{\varrho_{1i} t^{\alpha_i}}{\Gamma(\alpha_1 + \alpha_i + 1)} \right) t^{\alpha_1}, & \text{if } \alpha_s \in (0, 1], s = 1, 2, \\ A_{11} + B_{11}t + \left(\frac{l_1}{\Gamma(\alpha_1 + 1)} + \frac{(\eta_{10} B_{12} + d_{10} B_{13})t}{\Gamma(\alpha_1 + 2)} + \sum_{i=1}^2 \frac{\varrho_{1i} t^{\alpha_i}}{\Gamma(\alpha_1 + \alpha_i + 1)} \right. \\ \left. + \frac{2d_{10} \eta_{10} B_{14} t^{\alpha_1 + 1}}{\Gamma(2\alpha_1 + 2)} + \frac{(g_{10} d_{20} + \eta_{20} h_{10}) B_{14} t^{\alpha_2 + 1}}{\Gamma(\alpha_1 + \alpha_2 + 2)} \right) t^{\alpha_1}, & \\ \text{if } \alpha_1 \in (1, 2] \& \alpha_2 \in (0, 1], \\ A_{11} + \left(\frac{B_{24} (g_{20} h_{10} + g_{10} h_{20}) t^{\alpha_2 + 1}}{\Gamma(\alpha_1 + \alpha_2 + 2)} + \frac{B_{24} (\eta_{10} h_{10} + d_{10} g_{10}) t^{\alpha_1 + 1}}{\Gamma(2\alpha_1 + 2)} \right. \\ \left. + \frac{l_1}{\Gamma(\alpha_1 + 1)} + \sum_{i=1}^2 \frac{\varrho_{1i} t^{\alpha_i}}{\Gamma(\alpha_1 + \alpha_i + 1)} + \frac{(h_{10} B_{23} + g_{10} B_{22})t}{\Gamma(\alpha_1 + 2)} \right) t^{\alpha_1}, & \\ \text{if } \alpha_1 \in (0, 1] \& \alpha_2 \in (1, 2], \\ A_{11} + B_{11}t + \frac{l_1 t^{\alpha_1}}{\Gamma(\alpha_2 + 1)} + \frac{\varsigma_1 t^{\alpha_1 + 1}}{\Gamma(\alpha_1 + 2)} + \sum_{i=1}^2 \left(\frac{\varrho_{1i} t^{\alpha_i}}{\Gamma(\alpha_1 + \alpha_i + 1)} \right. \\ \left. + \frac{\lambda_{1i} t^{\alpha_i + 1}}{\Gamma(\alpha_1 + \alpha_i + 2)} \right) t^{\alpha_1}, & \text{if } \alpha_s \in (1, 2], s = 1, 2, \end{cases} \quad (3.24)$$




Figure 1: 2D graphical representations of the analytical solution $u_1(x_1, x_2, t)$ and $u_2(x_1, x_2, t)$ given in (3.25) for various values of $\alpha_s \in (0, 1], s = 1, 2$, $\eta_{20} = 1, h_{20} = 4, x_1 = 20$, and $g_{20} = x_2 = -10$.

where $l_s = g_{s0}A_{22} + h_{s0}A_{23} + d_{s0}A_{13} + \eta_{s0}A_{12}$, $\varrho_{s1} = d_{s0}\gamma_1 + \eta_{s0}\mu_1$, $\varrho_{s2} = g_{s0}\mu_2 + h_{s0}\gamma_2$, $\varsigma_s = g_{s0}B_{22} + h_{s0}B_{23} + d_{s0}B_{13} + \eta_{s0}B_{12}$, $\lambda_{s1} = d_{s0}\kappa_1 + \eta_{s0}\nu_1$, $\lambda_{s2} = g_{s0}\nu_2 + h_{s0}\kappa_2$, and $A_{si}, B_{si} \in \mathbb{R}, s = 1, 2, i = 1, 2, 3, 4$.

Thus, the obtained generalized separable analytical solution of the given system (3.14) is $\mathbf{U} = (u_1, u_2)$ with components $u_s = u_s(x_1, x_2, t), s = 1, 2$, as follows;


$$\begin{aligned} u_1(x_1, x_2, t) &= \delta_{11}(t) + \delta_{12}(t)x_1 + [\delta_{13}(t) + \delta_{14}(t)x_1]x_2, \text{ and} \\ u_2(x_1, x_2, t) &= \delta_{21}(t) + \delta_{22}(t)x_1 + [\delta_{23}(t) + \delta_{24}(t)x_1]x_2, \end{aligned} \quad (3.25)$$

where the functions $\delta_{si}, s = 1, 2, i = 1, 2, 3, 4$, are obtained in (3.18)-(3.24). We observe that the above analytical solution (3.25) of the given coupled equations (3.14) satisfies the initial conditions (3.15) with

$$\begin{aligned} \omega_s(x_1, x_2) &= A_{s1} + A_{s2}x_1 + A_{s3}x_2 + A_{s4}x_1x_2, \text{ and} \\ \vartheta_s(x_1, x_2) &= B_{s1} + B_{s2}x_1 + B_{s3}x_2 + B_{s4}x_1x_2, s = 1, 2. \end{aligned}$$


Here, note that whenever $\alpha_s = 1$ and $\alpha_s = 2$, the obtained solutions (3.25) coincide with integer-order cases of the discussed coupled system (3.14). Also, we wish to point out that the given equations (1.5) are a particular case of the considered coupled equations (3.14) when the parameters are considered as $q_{s1} = c_{s1} = \eta_{s0} = g_{s0} = d_{s0} = h_{s0} = 0, s = 1, 2, \kappa_{10} = \kappa_1\tau_0, q_{10} = \kappa_1\tau_1, \kappa_{20} = \kappa_2\tau_2$, and $q_{20} = \kappa_2\tau_0$. For the above-coupled equations (3.14), we have obtained a generalized separable analytical solution (3.25) systematically using the developed invariant subspace method for the first time. However, the approximate solution of the above system (1.4) can be expressed in terms of Mittag-Leffler functions, which are studied through the numerical inversion of the Laplace transformation technique [13]. We note that the obtained analytical solution (3.25) of the considered coupled equations (3.14) is valid for all $\alpha_s, \alpha_s \in (0, 2], s = 1, 2$. The two and three-dimensional graphical representations of the obtained analytical solution (3.25) of the given coupled equations (3.14) for arbitrary values of α_s , where $\alpha_s \in (0, 2], s = 1, 2$, are shown from Figure 1 to Figure 6. In these Figures 1-6, we considered the constants in the obtained analytical solution (3.25) as $A_{11} = 3, A_{21} = A_{14} = -h_{10} = d_{10} = A_{22} = A_{24} = B_{12} = B_{22} = 2, A_{13} = \eta_{10} = -A_{21} = A_{23} = -d_{20} = -B_{13} = B_{14} = -B_{21} = B_{23} = -B_{24} = 1, g_{10} = 50$, and $B_{11} = 5$.

Note 3.1. From the graphical representations given in Figure 1 to Figure 6, we observe that the behavior of the corresponding physical process is highly dependent on the value of the orders $\alpha_s, \alpha_s \in (0, 2], s = 1, 2$,



blue ($\alpha_1 = 0.2$ & $\alpha_2 = 0.3$), *green* ($\alpha_1 = 0.45$ & $\alpha_2 = 0.6$), *purple* ($\alpha_1 = 0.7$ & $\alpha_2 = 0.8$), *red* ($\alpha_1 = \alpha_2 = 1$)

(a)



blue ($\alpha_1 = 0.2$ & $\alpha_2 = 0.3$), *green* ($\alpha_1 = 0.45$ & $\alpha_2 = 0.6$), *purple* ($\alpha_1 = 0.7$ & $\alpha_2 = 0.8$), *red* ($\alpha_1 = \alpha_2 = 1$)

(b)

Figure 2: 3D graphical representations of the analytical solution $u_1(x_1, x_2, t)$ and $u_2(x_1, x_2, t)$ given in (3.25) for various values of $\alpha_s \in (0, 1]$, $s = 1, 2$, $\eta_{20} = 1$, $h_{20} = 4$, and $g_{20} = x_2 = -10$.




Figure 3: 2D graphical representations of the analytical solution $u_1(x_1, x_2, t)$ and $u_2(x_1, x_2, t)$ given in (3.25) for various values of $\alpha_1 \in (1, 2]$, $\alpha_2 \in (0, 1]$, $\eta_{20} = 1$, $h_{20} = 4$, $x_1 = 20$, and $g_{20} = x_2 = -10$.


of the Caputo fractional derivative that appears in the given coupled equations (3.14). The nature of the analytical solution (3.25) of the considered coupled equations (3.14) varies according to different values of α_s , $\alpha_s \in (0, 2]$, $s = 1, 2$. We observe that some important properties of the analytical solution (3.25) of the considered coupled equations (3.14) are given below.

- (a) From Figures 1-2, for various values of $\alpha_s \in (0, 1]$, $s = 1, 2$, the rate of change of diffusion occurred fast when $\alpha_1 = \alpha_2 = 1$ and slow if $\alpha_1 = 0.2, \alpha_2 = 0.3$. Note that for $\alpha_s < 1, s = 1, 2$, the slow-diffusion (sub-diffusion) is visible for both the components u_1 and u_2 .
- (b) From the plotted Figures 3-4, we note that the analytical solution (3.25) of the considered coupled equations (3.14) follows the wave structure for the component u_1 whereas the diffusion behavior is prominent for the component u_2 , for various values of $\alpha_1 \in (1, 2]$, and $\alpha_2 \in (0, 1]$.
- (c) For the Caputo derivative of order $\alpha_s \in (1, 2]$, $s = 1, 2$, we observe a prominent wave behavior for the components u_1 and u_2 as visible in Figures 5-6.

Remark 2. In [73], to describe unsteady magnetohydrodynamic flows, Wu et al. have studied the integer-order multi-component $(2 + 1)$ -dimensional coupled diffusion-convection equations having the form of (3.14) with parametric restrictions $\alpha_1 = \alpha_2 = 1$, and $q_{s1} = q_{10} = c_{s1} = c_{10} = \eta_{10} = d_{10} = \kappa_{20} = p_{20} = g_{20} = h_{20} = 0, s = 1, 2$. They have obtained an analytical solution of the considered coupled equations in terms of exponential and trigonometric functions in t, x_1 , and x_2 . We note that the given coupled equations (3.14) with $q_{s1} = q_{10} = c_{s1} = c_{10} = \eta_{10} = d_{10} = \kappa_{20} = p_{20} = g_{20} = h_{20} = 0, s = 1, 2$, preserve the invariant product subspace


$$W_8 = \text{Span}\{\sin(a_1x_1)\cos(a_2x_2), \sin(a_1x_1)\sin(a_2x_2), \cos(a_1x_1)\cos(a_2x_2), \cos(a_1x_1)\sin(a_2x_2)\} \times \text{Span}\{\sin(a_1x_1)\cos(a_2x_2), \sin(a_1x_1)\sin(a_2x_2), \cos(a_1x_1)\cos(a_2x_2), \cos(a_1x_1)\sin(a_2x_2)\},$$

where $a_1, a_2 \in \mathbb{R}$. Thus, using the above invariant product linear space W_8 , we can obtain a generalized separable analytical solution of the considered coupled equations (3.14) with the above-mentioned



blue($\alpha_1 = 1.2$ & $\alpha_2 = 0.3$), *green*($\alpha_1 = 1.45$ & $\alpha_2 = 0.5$), *purple*($\alpha_1 = 1.7$ & $\alpha_2 = 0.6$), *red*($\alpha_1 = 2$ & $\alpha_2 = 1$)

(a)



blue($\alpha_1 = 1.2$ & $\alpha_2 = 0.3$), *green*($\alpha_1 = 1.45$ & $\alpha_2 = 0.5$), *purple*($\alpha_1 = 1.7$ & $\alpha_2 = 0.6$), *red*($\alpha_1 = 2$ & $\alpha_2 = 1$)

(b)

Figure 4: 3D graphical representations of the analytical solution $u_1(x_1, x_2, t)$ and $u_2(x_1, x_2, t)$ given in (3.25) for various values of $\alpha_1 \in (1, 2]$, $\alpha_2 \in (0, 1]$, $\eta_{20} = 1$, $h_{20} = 4$, and $g_{20} = x_2 = -10$.




Figure 5: 2D graphical representations of the analytical solution $u_1(x_1, x_2, t)$ and $u_2(x_1, x_2, t)$ given in (3.25) for various values of $\alpha_s \in (1, 2]$, $s = 1, 2$, $\eta_{20} = -1$, $h_{20} = 20$, $g_{20} = -50$, $x_1 = 20$, and $x_2 = -10$.

parametric restrictions as follows:

$$u_s(x_1, x_2, t) = \sin(a_1 x_1) [\delta_{s1}(t) \cos(a_2 x_2) + \delta_{s2}(t) \sin(a_2 x_2)] + \cos(a_1 x_1) [\delta_{s3}(t) \cos(a_2 x_2) + \delta_{s4}(t) \sin(a_2 x_2)], \quad s = 1, 2, \quad (3.26)$$

where the unknown functions $\delta_{sj}(t)$, $s = 1, 2$, $j = 1, 2, 3, 4$, can be determined by following the same procedure of the invariant subspace method as used in Example 3.1. Note that the obtained analytical solution (3.26) is more general than the derived solution in [73].

Example 3.2. Now, we consider a coupled nonlinear TFDCWEs in $(2+1)$ -dimensions of the following form

$$\begin{aligned} \frac{\partial^{\alpha_1} u_1}{\partial t^{\alpha_1}} &= \frac{\partial}{\partial x_1} \left[(k_{10} - q_{11} u_2) \frac{\partial u_1}{\partial x_1} + (q_{10} + q_{11} u_1) \frac{\partial u_2}{\partial x_1} \right] + \frac{\partial}{\partial x_2} \left[\left(\frac{d_{12}}{a_{21}} u_2 + p_{10} \right) \frac{\partial u_1}{\partial x_2} \right. \\ &\quad \left. + \left(c_{10} - \frac{d_{12}}{a_{21}} u_1 \right) \frac{\partial u_2}{\partial x_2} \right] + \eta_{10} \frac{\partial u_1}{\partial x_1} + (d_{12} u_2 + d_{10}) \frac{\partial u_1}{\partial x_2} + h_{10} \frac{\partial u_2}{\partial x_2}, \\ \frac{\partial^{\alpha_2} u_2}{\partial t^{\alpha_2}} &= \frac{\partial}{\partial x_1} \left[(k_{20} - q_{21} u_2) \frac{\partial u_1}{\partial x_1} + (q_{21} u_1 + q_{20}) \frac{\partial u_2}{\partial x_1} \right] + c_{20} \frac{\partial^2 u_2}{\partial x_2^2} + g_{20} \frac{\partial u_2}{\partial x_1} + h_{20} \frac{\partial u_2}{\partial x_2}, \end{aligned} \quad (3.27)$$

$\alpha_s \in (0, 2]$, $s = 1, 2$, with appropriate initial conditions


$$\begin{aligned} (i) \quad &u_s(x_1, x_2, 0) = \omega_s(x_1, x_2) \quad \text{if } \alpha_s \in (0, 1], s = 1, 2, \\ (ii) \quad &u_s(x_1, x_2, 0) = \omega_s(x_1, x_2), \quad \frac{\partial u_s}{\partial t} \Big|_{t=0} = \vartheta_s(x_1, x_2) \quad \text{if } \alpha_s \in (1, 2], s = 1, 2. \end{aligned} \quad (3.28)$$

The exponential-polynomial product linear space

$$\hat{\mathbf{W}}_8^1 = \text{Span} \{1, x_1, e^{-a_{21} x_2}, x_1 e^{-a_{21} x_2}\} \times \text{Span} \{1, x_1, x_2, x_1 x_2\}$$


is an invariant product linear space for the given coupled equations (3.27), which is discussed in case 40 of Table 5. By proceeding with the similar technique employed in Example 3.1, we obtain the generalized separable analytical solution for the considered coupled equations (3.27) as follows:

$$\begin{aligned} u_1(x_1, x_2, t) &= \delta_{11}(t) + \delta_{12}(t) x_1 + [\delta_{13}(t) + \delta_{14}(t) x_1] e^{-a_{21} x_2}, \quad \text{and} \\ u_2(x_1, x_2, t) &= \delta_{21}(t) + \delta_{22}(t) x_1 + [\delta_{23}(t) + \delta_{24}(t) x_1] x_2, \end{aligned} \quad (3.29)$$



blue($\alpha_1 = 1.2$ & $\alpha_2 = 1.1$), *green*($\alpha_1 = 1.4$ & $\alpha_2 = 1.5$), *purple*($\alpha_1 = 1.7$ & $\alpha_2 = 1.6$), *red*($\alpha_1 = \alpha_2 = 2$)

(a)



blue($\alpha_1 = 1.2$ & $\alpha_2 = 1.1$), *green*($\alpha_1 = 1.4$ & $\alpha_2 = 1.5$), *purple*($\alpha_1 = 1.7$ & $\alpha_2 = 1.6$), *red*($\alpha_1 = \alpha_2 = 2$)

(b)

Figure 6: 3D graphical representations of the analytical solution $u_1(x_1, x_2, t)$ and $u_2(x_1, x_2, t)$ given in (3.25) for various values of $\alpha_s \in (1, 2]$, $s = 1, 2$, $\eta_{20} = -1$, $h_{20} = 20$, $g_{20} = -50$, and $x_2 = -10$.

where

$$\begin{aligned}
\delta_{11}(t) &= \begin{cases} A_{11} + \left(\frac{l_1}{\Gamma(\alpha_1 + 1)} + \sum_{i=1}^2 \frac{\varrho_{1i} t^{\alpha_i}}{\Gamma(\alpha_1 + \alpha_i + 1)} \right) t^{\alpha_1}, & \text{if } \alpha_s \in (0, 1], s = 1, 2, \\ A_{11} + B_{11}t + \left(\frac{l_1}{\Gamma(\alpha_1 + 1)} + \frac{\eta_{10} B_{12} t}{\Gamma(\alpha_1 + 2)} + \sum_{i=1}^2 \frac{\varrho_{1i} t^{\alpha_i}}{\Gamma(\alpha_1 + \alpha_i + 1)} \right) t^{\alpha_1}, \\ & \text{if } \alpha_1 \in (1, 2] \ \& \ \alpha_2 \in (0, 1], \\ A_{11} + \left[\frac{l_1}{\Gamma(\alpha_1 + 1)} + \sum_{i=1}^2 \left(\frac{\varrho_{1i}}{\Gamma(\alpha_1 + \alpha_i + 1)} + \frac{\lambda_{1i} t}{\Gamma(\alpha_1 + \alpha_i + 2)} \right) \right. \\ & \left. + \frac{h_{10} B_{23} t}{\Gamma(\alpha_1 + 2)} \right] t^{\alpha_1}, & \text{if } \alpha_1 \in (0, 1] \ \& \ \alpha_2 \in (1, 2], \\ A_{11} + B_{11}t + \left[\frac{l_1}{\Gamma(\alpha_1 + 1)} + \frac{l_2 t}{\Gamma(\alpha_1 + 2)} + \sum_{i=1}^2 \left(\frac{\varrho_{1i}}{\Gamma(\alpha_1 + \alpha_i + 1)} \right. \right. \\ & \left. \left. + \frac{\lambda_{1i} t}{\Gamma(\alpha_1 + \alpha_i + 2)} \right) t^{\alpha_i} \right] t^{\alpha_1}, & \text{if } \alpha_s \in (1, 2], s = 1, 2, \end{cases} \\
\delta_{12}(t) &= \begin{cases} A_{12} + \frac{h_{10} A_{24} t^{\alpha_1}}{\Gamma(\alpha_1 + 1)}, & \text{if } \alpha_s \in (0, 1], s = 1, 2, \\ A_{12} + B_{12}t + \frac{h_{10} A_{24} t^{\alpha_1}}{\Gamma(\alpha_1 + 1)}, & \text{if } \alpha_1 \in (1, 2] \ \& \ \alpha_2 \in (0, 1], \\ A_{12} + \frac{h_{10} A_{24} t^{\alpha_1}}{\Gamma(\alpha_1 + 1)} + \frac{h_{10} B_{24} t^{\alpha_1 + 1}}{\Gamma(\alpha_1 + 2)}, & \text{if } \alpha_1 \in (0, 1] \ \& \ \alpha_2 \in (1, 2], \\ A_{12} + B_{12}t + \frac{h_{10} A_{24} t^{\alpha_1}}{\Gamma(\alpha_1 + 1)} + \frac{h_{10} B_{24} t^{\alpha_1 + 1}}{\Gamma(\alpha_1 + 2)}, & \text{if } \alpha_s \in (1, 2], s = 1, 2, \end{cases} \\
\delta_{13}(t) &= \begin{cases} A_{13} E_{\alpha_1, 1}(\gamma_0 t^{\alpha_1}) + \eta_{10} A_{14} t^{\alpha_1 - 1} E_{\alpha_1, \alpha_1}(\gamma_0 t^{\alpha_1}) * E_{\alpha_1, 1}(\gamma_0 t^{\alpha_1}), \\ & \text{if } \alpha_1 \in (0, 1] \ \& \ \alpha_2 \in (0, 2], \\ A_{13} E_{\alpha_1, 1}(\gamma_0 t^{\alpha_1}) + B_{13} t E_{\alpha_1, 2}(\gamma_0 t^{\alpha_1}) + \eta_{10} t^{\alpha_1 - 1} E_{\alpha_1, \alpha_1}(\gamma_0 t^{\alpha_1}) \\ & * (A_{14} E_{\alpha_1, 1}(\gamma_0 t^{\alpha_1}) + B_{14} t E_{\alpha_1, 2}(\gamma_0 t^{\alpha_1})), & \text{if } \alpha_1 \in (1, 2] \ \& \ \alpha_2 \in (0, 2], \end{cases} \\
\delta_{14}(t) &= \begin{cases} A_{14} E_{\alpha_1, 1}(\gamma_0 t^{\alpha_1}), & \text{if } \alpha_1 \in (0, 1] \ \& \ \alpha_2 \in (0, 2], \\ A_{14} E_{\alpha_1, 1}(\gamma_0 t^{\alpha_1}) + B_{14} t E_{\alpha_1, 2}(\gamma_0 t^{\alpha_1}), & \text{if } \alpha_1 \in (1, 2] \ \& \ \alpha_2 \in (0, 2]. \end{cases} \\
\delta_{21}(t) &= \begin{cases} A_{21} + \left(\frac{\varrho_{21}}{\Gamma(\alpha_2 + 1)} + \frac{\varrho_{22} t^{\alpha_2}}{\Gamma(2\alpha_2 + 1)} \right) t^{\alpha_2}, & \text{if } \alpha_1 \in (0, 2] \ \& \ \alpha_2 \in (0, 1], \\ A_{21} + B_{21}t + \left(\frac{\varrho_{21}}{\Gamma(\alpha_2 + 1)} + \frac{\lambda_{21} t}{\Gamma(\alpha_2 + 2)} + \frac{\varrho_{22}}{\Gamma(2\alpha_2 + 1)} + \frac{\lambda_{22} t}{\Gamma(2\alpha_2 + 2)} \right) t^{\alpha_2}, \\ & \text{if } \alpha_1 \in (0, 2] \ \& \ \alpha_2 \in (1, 2], \end{cases} \\
\delta_{22}(t) &= \begin{cases} A_{22} + \frac{h_{20} A_{24} t^{\alpha_2}}{\Gamma(\alpha_2 + 1)}, & \text{if } \alpha_1 \in (0, 2] \ \& \ \alpha_2 \in (0, 1], \\ A_{22} + B_{22}t + \frac{h_{20} A_{24} t^{\alpha_2}}{\Gamma(\alpha_2 + 1)} + \frac{h_{20} B_{24} t^{\alpha_2 + 1}}{\Gamma(\alpha_2 + 2)}, & \text{if } \alpha_1 \in (0, 2] \ \& \ \alpha_2 \in (1, 2], \end{cases} \\
\delta_{23}(t) &= \begin{cases} A_{23} + \frac{g_{20} A_{24} t^{\alpha_2}}{\Gamma(\alpha_2 + 1)}, & \text{if } \alpha_1 \in (0, 2] \ \& \ \alpha_2 \in (0, 1], \\ A_{23} + B_{23}t + \frac{g_{20} A_{24} t^{\alpha_2}}{\Gamma(\alpha_2 + 1)} + \frac{g_{20} B_{24} t^{\alpha_2 + 1}}{\Gamma(\alpha_2 + 2)}, & \text{if } \alpha_1 \in (0, 2] \ \& \ \alpha_2 \in (1, 2], \end{cases} \\
\text{and } \delta_{24}(t) &= \begin{cases} A_{24}, & \text{if } \alpha_1 \in (0, 2] \ \& \ \alpha_2 \in (0, 1], \\ A_{24} + B_{24}t, & \text{if } \alpha_1 \in (0, 2] \ \& \ \alpha_2 \in (1, 2], \end{cases}
\end{aligned}$$

Here the two-parameter Mittag-Leffler function [6] is defined as $E_{a,b}(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(ak+b)}$, $\Re(a), \Re(b) > 0$,
 $*$ is the convolution of two functions defined as $w_1(t) * w_2(t) = \int_0^t w_1(y)w_2(t-y)dy$,

$$\begin{aligned} l_1 &= \eta_{10}A_{12} + h_{10}A_{23}, & \varrho_{11} &= h_{10}\eta_{10}A_{24}, & \lambda_{11} &= h_{10}\eta_{10}B_{24}, & \gamma_0 &= p_{10}a_{21}^2 - d_{10}a_{21}, \\ l_2 &= \eta_{10}B_{12} + h_{10}B_{23}, & \varrho_{12} &= h_{10}g_{20}A_{24}, & \lambda_{12} &= h_{10}g_{20}B_{24}, & \lambda_{21} &= h_{20}B_{23} + g_{20}B_{22}, \\ \varrho_{21} &= h_{20}A_{23} + g_{20}A_{22}, & \varrho_{22} &= 2h_{20}g_{20}A_{24}, & \lambda_{22} &= 2h_{20}g_{20}B_{24}, & & \& A_{si}, B_{si} \in \mathbb{R}, \end{aligned}$$

for $s = 1, 2, i = 1, 2, 3, 4$. Finally, we observe that the above analytical solution (3.29) for the given coupled equations (3.27) satisfies the initial conditions (3.28) with

$$\begin{aligned} \omega_1(x_1, x_2) &= A_{11} + A_{12}x_1 + (A_{13} + A_{14}x_1)e^{-a_{21}x_2}, \\ \omega_2(x_1, x_2) &= A_{21} + A_{22}x_1 + (A_{23} + A_{24}x_1)x_2, \\ \vartheta_1(x_1, x_2) &= B_{11} + B_{12}x_1 + (B_{13} + B_{14}x_1)e^{-a_{21}x_2}, \\ \& \vartheta_2(x_1, x_2) &= B_{21} + B_{22}x_1 + (B_{23} + B_{24}x_1)x_2. \end{aligned}$$

Furthermore, we note that the analytical solution (3.29) of the given coupled equations (3.27) is valid for all $\alpha_s, \alpha_s \in (0, 2], s = 1, 2$. Also, we observe that when $\alpha_s = 1$ and $\alpha_s = 2$, the obtained solutions (3.29) coincide with integer-order cases of the given coupled system (3.27).

Example 3.3. Consider the multi-component $(2 + 1)$ -dimensional coupled nonlinear TFDCWEs for $\alpha_s \in (0, 2], s = 1, 2$, as in the form

$$\begin{aligned} \frac{\partial^{\alpha_1} u_1}{\partial t^{\alpha_1}} &= k_{10} \frac{\partial^2 u_1}{\partial x_1^2} + \frac{g_{10}}{b_{11}} \frac{\partial^2 u_2}{\partial x_1^2} + \frac{\partial}{\partial x_2} \left[(-c_{11}u_2 + p_{10}) \frac{\partial u_1}{\partial x_2} + (c_{11}u_1 + c_{10}) \frac{\partial u_2}{\partial x_2} \right] \\ &\quad + \eta_{10} \frac{\partial u_1}{\partial x_1} + g_{10} \frac{\partial u_2}{\partial x_1} + d_{10} \frac{\partial u_1}{\partial x_2}, \\ \frac{\partial^{\alpha_2} u_2}{\partial t^{\alpha_2}} &= q_{20} \frac{\partial^2 u_2}{\partial x_1^2} + \frac{\partial}{\partial x_2} \left[(-c_{21}u_2 + p_{20}) \frac{\partial u_1}{\partial x_2} + (c_{21}u_1 + c_{20}) \frac{\partial u_2}{\partial x_2} \right] \\ &\quad + g_{20} \frac{\partial u_2}{\partial x_1} + h_{20} \frac{\partial u_2}{\partial x_2}, \end{aligned} \tag{3.30}$$

with appropriate initial conditions

$$\begin{aligned} (i) \quad &u_s(x_1, x_2, 0) = \omega_s(x_1, x_2) \text{ if } \alpha_s \in (0, 1], s = 1, 2, \\ (ii) \quad &u_s(x_1, x_2, 0) = \omega_s(x_1, x_2), \quad \frac{\partial u_s}{\partial t} \Big|_{t=0} = \vartheta_s(x_1, x_2) \text{ if } \alpha_s \in (1, 2], s = 1, 2. \end{aligned} \tag{3.31}$$

The product linear space

$$\begin{aligned} \hat{\mathbf{W}}_8^1 &= \text{Span} \{ \sin(\sqrt{a_{10}}x_1), x_2 \sin(\sqrt{a_{10}}x_1), \cos(\sqrt{a_{10}}x_1), x_2 \cos(\sqrt{a_{10}}x_1) \} \\ &\quad \times \text{Span} \{ 1, x_2, e^{b_{11}x_1}, x_2 e^{b_{11}x_1} \}, a_{10} > 0, \end{aligned}$$

is invariant for the coupled equations (3.30), which is discussed in case 11 of Table 2. Now, we apply the above procedure here and obtain the generalized separable analytical solution for the given coupled equations (3.30) along with $\eta_{10} = d_{10} = h_{20} = 0$ as follows:

$$\begin{aligned} u_1(x_1, x_2, t) &= (\delta_{11}(t) + \delta_{12}(t)x_2) \sin(\sqrt{a_{10}}x_1) + (\delta_{13}(t) + \delta_{14}(t)x_2) \cos(\sqrt{a_{10}}x_1), \\ u_2(x_1, x_2, t) &= \delta_{21}(t) + \delta_{22}(t)x_2 + \delta_{23}(t)e^{b_{11}x_1} + \delta_{24}(t)x_2 e^{b_{11}x_1}, \end{aligned} \tag{3.32}$$

where

$$\begin{aligned}\delta_{1i}(t) &= \begin{cases} A_{1i}E_{\alpha_1,1}(-k_{10}a_{10}t^{\alpha_1}), & \text{if } \alpha_1 \in (0, 1] \& \alpha_2 \in (0, 2], \\ A_{1i}E_{\alpha_1,1}(-k_{10}a_{10}t^{\alpha_1}) + B_{1i}tE_{\alpha_1,2}(-k_{10}a_{10}t^{\alpha_1}), \\ & \text{if } \alpha_1 \in (1, 2] \& \alpha_2 \in (0, 2], i = 1, 2, 3, 4, \end{cases} \\ \delta_{2i}(t) &= \begin{cases} A_{2i}, & \text{if } \alpha_1 \in (0, 2] \& \alpha_2 \in (0, 1], \\ A_{2i} + B_{2i}t, & \text{if } \alpha_1 \in (0, 2] \& \alpha_2 \in (1, 2], i = 1, 2, \text{ and} \end{cases} \\ \delta_{2i}(t) &= \begin{cases} A_{2i}E_{\alpha_2,1}(\gamma_0 t^{\alpha_2}), & \text{if } \alpha_1 \in (0, 2] \& \alpha_2 \in (0, 1], \\ A_{2i}E_{\alpha_2,1}(\gamma_0 t^{\alpha_2}) + B_{2i}tE_{\alpha_2,2}(\gamma_0 t^{\alpha_2}), & \text{if } \alpha_1 \in (0, 2] \& \alpha_2 \in (1, 2], i = 3, 4. \end{cases}\end{aligned}$$

Here $\gamma_0 = q_{20}b_{11}^2 - g_{20}b_{11}$, and $A_{si}, B_{si} \in \mathbb{R}, s = 1, 2, i = 1, 2, 3, 4$. Additionally, we note that the obtained generalized separable analytical solution (3.32) of the given coupled equations (3.30) satisfies the initial conditions (3.31) with

$$\begin{aligned}\omega_1(x_1, x_2) &= (A_{11} + A_{12}x_2) \sin(\sqrt{a_{10}}x_1) + (A_{13} + A_{14}x_2) \cos(\sqrt{a_{10}}x_1), \\ \vartheta_1(x_1, x_2) &= (B_{11} + B_{12}x_2) \sin(\sqrt{a_{10}}x_1) + (B_{13} + B_{14}x_2) \cos(\sqrt{a_{10}}x_1), \\ \omega_2(x_1, x_2) &= A_{21} + A_{22}x_2 + (A_{23} + A_{24}x_2)e^{b_{11}x_1}, \\ \& \vartheta_2(x_1, x_2) &= B_{21} + B_{22}x_2 + (B_{23} + B_{24}x_2)e^{b_{11}x_1}.\end{aligned}$$

Also, we would like to mention here that the obtained analytical solution (3.32) of the considered coupled equations (3.30) is valid for all $\alpha_s, \alpha_s \in (0, 2], s = 1, 2$. Additionally, we note that when $\alpha_s = 1$ and $\alpha_s = 2$, the obtained solutions (3.32) coincide with integer-order cases of the coupled system (3.30).

3.2.2 Generalized separable analytical solutions for the initial-boundary value problems of the given coupled equations (1.7)

Next, we provide the systematic approach of the developed method to derive analytical solutions for the initial and the Dirichlet boundary value problems of the considered coupled equations (1.7) based on the determined invariant product subspaces.

Example 3.4. Now, we consider the following multi-component (2+1)-dimensional coupled linear TFD-CWEs as

$$\begin{aligned}\frac{\partial^{\alpha_1} u_1}{\partial t^{\alpha_1}} &= k_{10} \frac{\partial^2 u_1}{\partial x_1^2} + q_{10} \frac{\partial^2 u_2}{\partial x_1^2} + p_{10} \frac{\partial^2 u_1}{\partial x_2^2} + q_{10} b_{11} \frac{\partial u_2}{\partial x_1}, \\ \frac{\partial^{\alpha_2} u_2}{\partial t^{\alpha_2}} &= q_{20} \frac{\partial^2 u_2}{\partial x_1^2} + c_{20} \frac{\partial^2 u_2}{\partial x_2^2} + g_{20} \frac{\partial u_2}{\partial x_1}, \alpha_s \in (0, 2], s = 1, 2,\end{aligned}\tag{3.33}$$

with appropriate initial conditions

$$\begin{aligned}(i) \quad & u_s(x_1, x_2, 0) = \omega_s(x_1, x_2) \quad \text{if } \alpha_s \in (0, 1], s = 1, 2, \\ (ii) \quad & u_s(x_1, x_2, 0) = \omega_s(x_1, x_2), \quad \frac{\partial u_s}{\partial t} \Big|_{t=0} = \vartheta_s(x_1, x_2) \quad \text{if } \alpha_s \in (1, 2], s = 1, 2,\end{aligned}\tag{3.34}$$

and the Dirichlet boundary conditions

$$\begin{aligned}
(B-I) \left\{ \begin{array}{l} u_1(0, x_2, t) = \xi_{11}(t) \sin(\sqrt{b_{20}}x_2) + \xi_{12}(t) \cos(\sqrt{b_{20}}x_2), \text{ if } (0, x_2, t) \in \partial\Omega \times [0, \infty), \\ u_1(\sigma_1, x_2, t) = \xi_{13}(t) \sin(\sqrt{b_{20}}x_2) + \xi_{14}(t) \cos(\sqrt{b_{20}}x_2), \\ \text{if } (\sigma_1, x_2, t) \in \partial\Omega \times [0, \infty), \\ u_1(x_1, 0, t) = \xi_{21}(t) \sin(\sqrt{a_{10}}x_1) + \xi_{22}(t) \cos(\sqrt{a_{10}}x_1), \text{ if } (x_1, 0, t) \in \partial\Omega \times [0, \infty), \\ u_1(x_1, \sigma_2, t) = \xi_{23}(t) \sin(\sqrt{a_{10}}x_1) + \xi_{24}(t) \cos(\sqrt{a_{10}}x_1), \\ \text{if } (x_1, \sigma_2, t) \in \partial\Omega \times [0, \infty), \end{array} \right. \quad (3.35) \\
(B-II) \left\{ \begin{array}{l} u_2(0, x_2, t) = f_{11}(t) \sin(\sqrt{b_{20}}x_2) + f_{12}(t) \cos(\sqrt{b_{20}}x_2), \text{ if } (0, x_2, t) \in \partial\Omega \times [0, \infty), \\ u_2(\sigma_1, x_2, t) = f_{13}(t) \sin(\sqrt{b_{20}}x_2) + f_{14}(t) \cos(\sqrt{b_{20}}x_2), \\ \text{if } (\sigma_1, x_2, t) \in \partial\Omega \times [0, \infty), \\ u_2(x_1, 0, t) = f_{21}(t) + f_{22}(t)e^{-b_{11}x_1}, \text{ if } (x_1, 0, t) \in \partial\Omega \times [0, \infty), \\ u_2(x_1, \sigma_2, t) = f_{23}(t) + f_{24}(t)e^{-b_{11}x_1}, \text{ if } (x_1, \sigma_2, t) \in \partial\Omega \times [0, \infty). \end{array} \right.
\end{aligned}$$

The given coupled equations (3.33) are defined on a domain $\Omega \times [0, \infty)$, where $\Omega = \{(x_1, x_2) \in \mathbb{R}^2 \mid 0 \leq x_i \leq \sigma_i, \sigma_i \in \mathbb{R}, i = 1, 2\}$ is a subset of \mathbb{R}^2 whose boundary $\partial\Omega$ is defined by the lines $x_1 = 0, x_2 = 0, x_1 = \sigma_1$, and $x_2 = \sigma_2$. It is easy to observe that the coupled equations (3.33) admit the following invariant product linear space

$$\begin{aligned}
\hat{\mathbf{W}}_8^1 = & \text{Span} \left\{ \sin(\sqrt{a_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{a_{10}}x_1) \sin(\sqrt{b_{20}}x_2), \sin(\sqrt{a_{10}}x_1) \cos(\sqrt{b_{20}}x_2), \right. \\
& \left. \cos(\sqrt{a_{10}}x_1) \cos(\sqrt{b_{20}}x_2) \right\} \\
& \times \text{Span} \left\{ \sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2), e^{-b_{11}x_1} \sin(\sqrt{b_{20}}x_2), e^{-b_{11}x_1} \cos(\sqrt{b_{20}}x_2) \right\},
\end{aligned}$$

where $a_{10}, b_{20} \geq 0$. Applying the similar procedure as explained in Example 3.1, the obtained generalized separable analytical solution for the given coupled equations (3.33) is $\mathbf{U} = (u_1, u_2)$ with $u_s = u_s(x_1, x_2, t), s = 1, 2$, as

$$\begin{aligned}
u_1 = & [\delta_{11}(t) \sin(\sqrt{a_{10}}x_1) + \delta_{13}(t) \cos(\sqrt{a_{10}}x_1)] \sin(\sqrt{b_{20}}x_2) \\
& + [\delta_{12}(t) \sin(\sqrt{a_{10}}x_1) + \delta_{14}(t) \cos(\sqrt{a_{10}}x_1)] \cos(\sqrt{b_{20}}x_2), \quad (3.36) \\
u_2 = & [\delta_{21}(t) + \delta_{23}(t)e^{-b_{11}x_1}] \sin(\sqrt{b_{20}}x_2) + [\delta_{22}(t) + \delta_{24}(t)e^{-b_{11}x_1}] \cos(\sqrt{b_{20}}x_2),
\end{aligned}$$

where

$$\begin{aligned}
\delta_{1i}(t) = & \begin{cases} A_{1i}E_{\alpha_1,1}(\gamma_0 t^{\alpha_1}), \text{ if } \alpha_1 \in (0, 1] \& \alpha_2 \in (0, 2], \\ A_{1i}E_{\alpha_1,1}(\gamma_0 t^{\alpha_1}) + B_{1i}tE_{\alpha_1,2}(\gamma_0 t^{\alpha_1}), \text{ if } \alpha_1 \in (1, 2] \\ \& \alpha_2 \in (0, 2], i = 1, 2, 3, 4, \end{cases} \\
\delta_{2i}(t) = & \begin{cases} A_{2i}E_{\alpha_2,1}(-c_{20}b_{20}t^{\alpha_2}), \text{ if } \alpha_1 \in (0, 2] \& \alpha_2 \in (0, 1], \\ A_{2i}E_{\alpha_2,1}(-c_{10}b_{20}t^{\alpha_2}) + B_{2i}tE_{\alpha_2,2}(-c_{10}b_{20}t^{\alpha_2}), \\ \text{if } \alpha_1 \in (0, 2] \& \alpha_2 \in (1, 2], i = 1, 2, \end{cases} \quad (3.37) \\
\& \delta_{2i}(t) = & \begin{cases} A_{2i}E_{\alpha_2,1}(\gamma_1 t^{\alpha_2}), \text{ if } \alpha_1 \in (0, 2] \& \alpha_2 \in (0, 1], \\ A_{2i}E_{\alpha_2,1}(\gamma_1 t^{\alpha_2}) + B_{2i}tE_{\alpha_2,2}(\gamma_1 t^{\alpha_2}), \text{ if } \alpha_1 \in (0, 2] \& \alpha_2 \in (1, 2], i = 3, 4. \end{cases}
\end{aligned}$$

Here $\gamma_0 = -k_{10}a_{10} - p_{10}b_{20}, \gamma_1 = q_{20}b_{11}^2 - c_{20}b_{20} - g_{20}b_{11}$, and $A_{si}, B_{si} \in \mathbb{R}, s = 1, 2, i = 1, 2, 3, 4$. We also observe that the above analytical solution (3.36) for the given coupled equations (3.33) satisfies the

initial-boundary conditions (3.34)-(3.35) with

$$\begin{aligned}
\omega_1(x_1, x_2) &= (A_{11} \sin(\sqrt{a_{10}}x_1) + A_{13} \cos(\sqrt{a_{10}}x_1)) \sin(\sqrt{b_{20}}x_2) \\
&\quad + (A_{12} \sin(\sqrt{a_{10}}x_1) + A_{14} \cos(\sqrt{a_{10}}x_1)) \cos(\sqrt{b_{20}}x_2), \\
\vartheta_1(x_1, x_2) &= (B_{11} \sin(\sqrt{a_{10}}x_1) + B_{13} \cos(\sqrt{a_{10}}x_1)) \sin(\sqrt{b_{20}}x_2) \\
&\quad + (B_{12} \sin(\sqrt{a_{10}}x_1) + B_{14} \cos(\sqrt{a_{10}}x_1)) \cos(\sqrt{b_{20}}x_2), \\
\omega_2(x_1, x_2) &= (A_{21} + A_{23}e^{-b_{11}x_1}) \sin(\sqrt{b_{20}}x_2) + (A_{22} + A_{24}e^{-b_{11}x_1}) \cos(\sqrt{b_{20}}x_2), \\
\vartheta_2(x_1, x_2) &= (B_{21} + B_{23}e^{-b_{11}x_1}) \sin(\sqrt{b_{20}}x_2) + (B_{22} + B_{24}e^{-b_{11}x_1}) \cos(\sqrt{b_{20}}x_2), \\
\xi_{11}(t) &= \delta_{12}(t), & \xi_{13}(t) &= C_1\delta_{11}(t) + C_2\delta_{12}(t), \\
\xi_{12}(t) &= \delta_{14}(t), & \xi_{14}(t) &= C_1\delta_{13}(t) + C_2\delta_{14}(t), \\
\xi_{21}(t) &= \delta_{13}(t), & \xi_{23}(t) &= C_3\delta_{11}(t) + C_4\delta_{13}(t), \\
\xi_{22}(t) &= \delta_{14}(t), & \xi_{24}(t) &= C_3\delta_{12}(t) + C_4\delta_{14}(t), \\
f_{11}(t) &= \delta_{21}(t) + \delta_{23}(t), & f_{12}(t) &= \delta_{22}(t) + \delta_{24}(t), \\
f_{13}(t) &= \delta_{21}(t) + \delta_{23}(t)e^{-b_{11}\sigma_1}, & f_{14}(t) &= \delta_{22}(t) + \delta_{24}(t)e^{-b_{11}\sigma_1}, \\
f_{21}(t) &= \delta_{22}(t), & f_{23}(t) &= C_3\delta_{21}(t) + C_4\delta_{22}(t), \\
f_{22}(t) &= \delta_{24}(t), \& & f_{24}(t) &= C_3\delta_{23}(t) + C_4\delta_{24}(t),
\end{aligned}$$

where $C_1 = \sin(\sigma_1\sqrt{a_{10}})$, $C_2 = \cos(\sigma_1\sqrt{a_{10}})$, $C_3 = \sin(\sigma_2\sqrt{b_{20}})$, $C_4 = \cos(\sigma_2\sqrt{b_{20}})$, and the functions δ_{si} as given in (3.37) for $s = 1, 2$, and $i = 1, 2, 3, 4$. Additionally, we observe that the obtained analytical solution (3.36) of the given coupled equations (3.33) is valid for all $\alpha_s, \alpha_s \in (0, 2]$, $s = 1, 2$. Also, we note that the derived solutions (3.36) coincide with integer-order cases of the discussed coupled system (3.33) if $\alpha_s = 1$ and $\alpha_s = 2$.

Note 3.2. It is interesting to observe that in the above Example 3.4, the obtained analytical solution (3.36) for the given initial-boundary value problem (3.33)-(3.35) is spatially periodic. From the physical point of view, linear analysis of possible patterns in an experiment is observed to be repeating if the experimental domain is subjected to periodic boundary conditions. While studying the stability of the underlying patterns, these periodic boundary conditions in a bounded domain are crucial in both fractional-order and integer-order PDEs.

Example 3.5. Next, we consider the following multi-component $(2 + 1)$ -dimensional coupled nonlinear TFDCWEs as

$$\begin{aligned}
\frac{\partial^{\alpha_1} u_1}{\partial t^{\alpha_1}} &= \frac{\partial}{\partial x_1} \left[(k_{10} - q_{11}u_2) \frac{\partial u_1}{\partial x_1} + (q_{11}u_1 + q_{10}) \frac{\partial u_2}{\partial x_1} \right] \\
&\quad + p_{10} \frac{\partial^2 u_1}{\partial x_2^2} + \eta_{10} \frac{\partial u_1}{\partial x_1} + d_{10} \frac{\partial u_1}{\partial x_2}, \\
\frac{\partial^{\alpha_2} u_2}{\partial t^{\alpha_2}} &= \frac{\partial}{\partial x_1} \left[(k_{20} - q_{21}u_2) \frac{\partial u_1}{\partial x_1} + (q_{21}u_1 + q_{20}) \frac{\partial u_2}{\partial x_1} \right] \\
&\quad + c_{20} \frac{\partial^2 u_2}{\partial x_2^2} + g_{20} \frac{\partial u_2}{\partial x_1} + h_{20} \frac{\partial u_2}{\partial x_2},
\end{aligned} \tag{3.38}$$

$\alpha_s \in (0, 2]$, $s = 1, 2$, with appropriate initial conditions

$$\begin{aligned}
(i) \quad u_s(x_1, x_2, 0) &= \omega_s(x_1, x_2) \text{ if } \alpha_s \in (0, 1], s = 1, 2, \\
(ii) \quad u_s(x_1, x_2, 0) &= \omega_s(x_1, x_2), \quad \frac{\partial u_s}{\partial t} \Big|_{t=0} = \vartheta_s(x_1, x_2) \text{ if } \alpha_s \in (1, 2], s = 1, 2,
\end{aligned} \tag{3.39}$$

and the Dirichlet boundary conditions

$$(B-I) \begin{cases} u_1(0, x_2, t) = \xi_{11}(t) \sin(\sqrt{a_{20}}x_2) + \xi_{12}(t) \cos(\sqrt{a_{20}}x_2), \text{ if } (0, x_2, t) \in \partial\Omega \times [0, \infty), \\ u_1(\sigma_1, x_2, t) = \xi_{13}(t) \sin(\sqrt{a_{20}}x_2) + \xi_{14}(t) \cos(\sqrt{a_{20}}x_2), \\ \text{if } (\sigma_1, x_2, t) \in \partial\Omega \times [0, \infty), \\ u_1(x_1, 0, t) = \xi_{21}(t) + \xi_{22}(t)x_1, \text{ if } (x_1, 0, t) \in \partial\Omega \times [0, \infty), \\ u_1(x_1, \sigma_2, t) = \xi_{23}(t) + \xi_{24}(t)x_1, \text{ if } (x_1, \sigma_2, t) \in \partial\Omega \times [0, \infty), \end{cases} \quad (3.40)$$

$$(B-II) \begin{cases} u_2(0, x_2, t) = f_{11}(t) \sin(\sqrt{b_{20}}x_2) + f_{12}(t) \cos(\sqrt{b_{20}}x_2), \text{ if } (0, x_2, t) \in \partial\Omega \times [0, \infty), \\ u_2(\sigma_1, x_2, t) = f_{13}(t) \sin(\sqrt{b_{20}}x_2) + f_{14}(t) \cos(\sqrt{b_{20}}x_2), \\ \text{if } (\sigma_1, x_2, t) \in \partial\Omega \times [0, \infty), \\ u_2(x_1, 0, t) = f_{21}(t) + f_{22}(t)x_1, \text{ if } (x_1, 0, t) \in \partial\Omega \times [0, \infty), \\ u_2(x_1, \sigma_2, t) = f_{23}(t) + f_{24}(t)x_1, \text{ if } (x_1, \sigma_2, t) \in \partial\Omega \times [0, \infty). \end{cases} \quad (3.41)$$

The given coupled equations (3.38) are defined on a domain $\Omega \times [0, \infty)$, where $\Omega = \{(x_1, x_2) \in \mathbb{R}^2 | 0 \leq x_i \leq \sigma_i, \sigma_i \in \mathbb{R}, i = 1, 2\}$ is a subset of \mathbb{R}^2 whose boundary $\partial\Omega$ is defined by the lines $x_1 = 0, x_2 = 0, x_1 = \sigma_1$, and $x_2 = \sigma_2$. It is given in case 7 of Table 1 that the coupled equations (3.38) admit the following invariant product linear space

$$\hat{\mathbf{W}}_8^1 = \text{Span}\{\sin(\sqrt{a_{20}}x_2), \cos(\sqrt{a_{20}}x_2), x_1 \sin(\sqrt{a_{20}}x_2), x_1 \cos(\sqrt{a_{20}}x_2)\} \\ \times \text{Span}\{\sin(\sqrt{b_{20}}x_2), \cos(\sqrt{b_{20}}x_2), x_1 \sin(\sqrt{b_{20}}x_2), x_1 \cos(\sqrt{b_{20}}x_2)\},$$

where $a_{20}, b_{20} \geq 0$. Applying a similar procedure as explained in Example 3.1, the obtained generalized separable analytical solution for the given coupled equations (3.38) along with $\eta_{10} = d_{10} = h_{20} = g_{20} = 0$ is $\mathbf{U} = (u_1, u_2)$ with $u_s = u_s(x_1, x_2, t)$, $s = 1, 2$, as

$$u_1(x_1, x_2, t) = (\delta_{11}(t) + \delta_{13}(t)x_1) \sin(\sqrt{a_{20}}x_2) + (\delta_{12}(t) + \delta_{14}(t)x_1) \cos(\sqrt{a_{20}}x_2), \\ u_2(x_1, x_2, t) = (\delta_{21}(t) + \delta_{23}(t)x_1) \sin(\sqrt{b_{20}}x_2) + (\delta_{22}(t) + \delta_{24}(t)x_1) \cos(\sqrt{b_{20}}x_2), \quad (3.42)$$

where

$$\delta_{1i}(t) = \begin{cases} A_{1i}E_{\alpha_1,1}(-p_{10}a_{20}t^{\alpha_1}), \text{ if } \alpha_1 \in (0, 1] \& \alpha_2 \in (0, 2], \\ A_{1i}E_{\alpha_1,1}(-p_{10}a_{20}t^{\alpha_1}) + B_{1i}tE_{\alpha_1,2}(-p_{10}a_{20}t^{\alpha_1}), \\ \text{if } \alpha_1 \in (1, 2] \& \alpha_2 \in (0, 2], i = 1, 2, 3, 4, \end{cases} \\ \text{and } \delta_{2i}(t) = \begin{cases} A_{2i}E_{\alpha_2,1}(-c_{20}b_{20}t^{\alpha_2}), \text{ if } \alpha_1 \in (0, 2] \& \alpha_2 \in (0, 1], \\ A_{2i}E_{\alpha_2,1}(-c_{10}b_{20}t^{\alpha_2}) + B_{2i}tE_{\alpha_2,2}(-c_{10}b_{20}t^{\alpha_2}), \\ \text{if } \alpha_1 \in (0, 2] \& \alpha_2 \in (1, 2], i = 1, 2, 3, 4. \end{cases} \quad (3.43)$$

Here $A_{si}, B_{si} \in \mathbb{R}$, $s = 1, 2, i = 1, 2, 3, 4$. We also observe that the above analytical solution (3.42) of the given coupled equations (3.38) satisfies the initial-boundary conditions (3.39)-(3.41) with

$$\omega_1(x_1, x_2) = (A_{11} + A_{13}x_1) \sin(\sqrt{a_{20}}x_2) + (A_{12} + A_{14}x_1) \cos(\sqrt{a_{20}}x_2), \\ \vartheta_1(x_1, x_2) = (B_{11} + B_{13}x_1) \sin(\sqrt{a_{20}}x_2) + (B_{12} + B_{14}x_1) \cos(\sqrt{a_{20}}x_2), \\ \omega_2(x_1, x_2) = (A_{21} + A_{23}x_1) \sin(\sqrt{b_{20}}x_2) + (A_{22} + A_{24}x_1) \cos(\sqrt{b_{20}}x_2), \\ \vartheta_2(x_1, x_2) = (B_{21} + B_{23}x_1) \sin(\sqrt{b_{20}}x_2) + (B_{22} + B_{24}x_1) \cos(\sqrt{b_{20}}x_2), \\ \xi_{11}(t) = \delta_{11}(t), \quad \xi_{12}(t) = \xi_{21}(t) = \delta_{12}(t), \quad \xi_{22}(t) = \delta_{14}(t), \\ f_{11}(t) = \delta_{21}(t), \quad f_{12}(t) = f_{21}(t) = \delta_{22}(t), \quad f_{22}(t) = \delta_{24}(t), \\ \xi_{23}(t) = C_1\delta_{11}(t) + C_2\delta_{12}(t), \quad \xi_{13}(t) = \delta_{11}(t) + \sigma_1\delta_{13}(t), \quad \xi_{14}(t) = \delta_{12}(t) + \sigma_1\delta_{14}(t), \\ \xi_{24}(t) = C_1\delta_{13}(t) + C_2\delta_{14}(t), \quad f_{13}(t) = \delta_{21}(t) + \sigma_1\delta_{23}(t), \quad f_{14}(t) = \delta_{22}(t) + \sigma_1\delta_{24}(t), \\ f_{23}(t) = C_3\delta_{21}(t) + C_4\delta_{22}(t), \& \quad f_{24}(t) = C_3\delta_{23}(t) + C_4\delta_{24}(t),$$

where $C_1 = \sin(\sigma_2\sqrt{a_{20}})$, $C_2 = \cos(\sigma_2\sqrt{a_{20}})$, $C_3 = \sin(\sigma_2\sqrt{b_{20}})$, $C_4 = \cos(\sigma_2\sqrt{b_{20}})$, and the functions δ_{si} as given in (3.43) for $s = 1, 2$, and $i = 1, 2, 3, 4$.

Additionally, we wish to mention here that when $I_s = \tau_1 = 0$, $s = 1, 2$, the given system (1.6) can be considered as a particular case of the above-discussed coupled equations (3.38) with $\alpha_s = \alpha$, $\alpha \in (0, 1]$, $q_{s1} = h_{20} = d_{10} = g_{20} = \eta_{10} = 0$, and $\kappa_{s0} = q_{s0} = p_{s0} = c_{s0} = \tau_0$, $s = 1, 2$, for which we have obtained the analytical solution in the generalized separable form (3.42) using the developed invariant subspace method. Also, Yang et al. [14] have found an approximate solution for the given coupled equations (1.6) using the Jacobi special collocation method along with the reduction $r = \sqrt{x_1^2 + x_2^2}$. Also, we note that the obtained analytical solution (3.42) for the considered coupled equations (3.38) is valid for all $\alpha_s, \alpha_s \in (0, 2]$, $s = 1, 2$. Additionally, we observe that the obtained solutions (3.42) coincide with integer-order cases of the given coupled system (3.38) when $\alpha_s = 1$ and $\alpha_s = 2$.

4 Discussions and Concluding remarks

4.1 Discussions

Diffusion is one of the most fundamental phenomena that occur naturally in physical, biological, chemical and ecology systems. With the development of science and technology, the need to study more complicated processes, usually involving the long-time interactions of more than one particle, chemical agent or species has become more significant. In particular, the flow of fluids in a porous media, dispersive sedimentation on the surfaces, and catalytic chemical reactors are studied extensively using the class of coupled diffusion-convection equations [72, 77–79]. One of the revolutionary investigations was made by Turing [80] to formulate the various pattern formations in organisms using coupled diffusion systems. Later, the interest in complex processes governed by more than one particle described mathematically using the class of coupled diffusion equations increased significantly [17, 81–83]. The convective flux in such a complex process has an intrinsic effect whenever an external force is present. For example, considering the effect of gravitational force in the porous medium fluids dynamics, the coupled system of diffusion-convection equations plays a vital role in studying the transport of contamination in groundwater [77].

One of the most celebrated coupled diffusion-convection equations is the coupled Burger's equation, which has many applications in the physical context. The mathematical formulation of the $(1 + 1)$ dimensional coupled Burger's equation [72] having the form

$$\frac{\partial u_i}{\partial t} = v_i \frac{\partial}{\partial x} [(1 - k_{i1}u_1 + k_{i2}u_2)u_i] + d_i \frac{\partial^2 u_i}{\partial x^2}, \quad (4.1)$$

which Esipov [72] proposed initially to describe the poly-dispersive sedimentation on rocks under gravitational force using the following governing equations based on Fick's law:

$$(i) \text{ the continuum equation: } \quad \frac{\partial \mathbf{U}}{\partial t} + \frac{\partial \mathbf{q}}{\partial x} = 0, \text{ and}$$

$$(ii) \text{ the constitutive equation: } \quad q_i = V_i(\mathbf{U}) - D_{ij}(\mathbf{U}) \frac{\partial u_j}{\partial x},$$

where $\mathbf{q} = (q_1, q_2)$, q_i is the i^{th} particle flux and $\mathbf{U} = (u_1, u_2)$, $u_i = u_i(x, t)$ is the i^{th} particle concentration $i = 1, 2$. Here, $V_i(\mathbf{U})$ denotes the concentration-dependent velocity, and $D_{ij}(\mathbf{U})$ is the diffusion coefficient for each $i, j = 1, 2$, which were taken in particular as $-v_i(1 - k_{i1}u_1 + k_{i2}u_2)u_i$ and d_i , respectively for $v_i, d_i, k_{i1}, k_{i2} \in \mathbb{R}$, $i = 1, 2$. Due to its broad applicability, the continuous interest in the class of coupled systems of diffusion-convection equations led to the study of analytical solutions for more complex systems

of this kind, even in higher dimensions. For instance, Fletcher [78] studied the analytical solutions for the $(2 + 1)$ -dimensional coupled Burger's equations of the form

$$\begin{aligned}\frac{\partial u_1}{\partial t} &= \frac{1}{Re} \sum_{r=1}^2 \frac{\partial^2 u_1}{\partial x_r^2} - u_1 \frac{\partial u_1}{\partial x_1} - u_2 \frac{\partial u_1}{\partial x_2} \\ \frac{\partial u_2}{\partial t} &= \frac{1}{Re} \sum_{r=1}^2 \frac{\partial^2 u_2}{\partial x_r^2} - u_1 \frac{\partial u_2}{\partial x_1} - u_2 \frac{\partial u_2}{\partial x_2},\end{aligned}\tag{4.2}$$

where Re represents the Reynold's number, and $u_s = u_s(x_1, x_2, t), x_r \in \mathbb{R}, t > 0$. The above-coupled equations (4.2) coincides with the well-known Navier-Stokes equations for incompressible fluids without pressure gradient terms. These are used to study the properties of various hydro-dynamical models and chemical reactors, to name only a few. Using the local discontinuous Galerkin method, an extensive study on different properties of the most generalized coupled diffusion-convection equations in the form of (1.10) was discussed by Cockburn and Shu [75].

However, the anisotropic or heterogeneous media diffusion processes challenged the scientists with their unpredictable nature, commonly called anomalous diffusion [74, 84]. Such anomalous diffusion has the characteristic property of nonlinear mean square displacement in time. More precisely, $\langle (x - \langle x \rangle)^2 \rangle \propto t^\alpha, \alpha > 0$, where x is the displacement and t is the time [74, 84]. Povstenko [10] considered the nonlocal time-dependence related to the matter flux \mathbf{Q} as follows:

$$\mathbf{Q} = \begin{cases} \frac{1}{\Gamma(\alpha)} \frac{\partial}{\partial t} \int_0^t (t-y)^{\alpha-1} (-\kappa \nabla u) dy, & \text{if } \alpha \in (0, 1], \\ \frac{1}{\Gamma(\alpha-1)} \int_0^t (t-y)^{\alpha-2} (-\kappa \nabla u) dy, & \text{if } \alpha \in (1, 2]. \end{cases}$$

The above-equation gives the time-fractional diffusion-wave equation of the form

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \kappa \Delta u, \alpha \in (0, 2],\tag{4.3}$$

where κ is the diffusive constant, $u = u(x_1, \dots, x_N, t), x_r \in \mathbb{R}, t > 0$, and $\frac{\partial^\alpha}{\partial t^\alpha}(\cdot)$ represents the Caputo fractional derivative (1.2). We observed that the above equation (4.3) is used to describe many real-world processes in semiconductors, chemical reactors, biological systems, thermoelastic materials, viscous-elastic materials, hydrothermal medium, pattern formations, and so on [1–5, 8, 10, 74]. Moreover, to investigate the heat conduction phenomena in an infinite dimensional composite medium, Povstenko [9] used the time-fractional multi-component diffusion-wave (heat) equations of the form

$$\frac{\partial^{\alpha_s} u_s}{\partial t^{\alpha_s}} = \kappa_s \frac{\partial^2 u_s}{\partial x^2}, \alpha_s \in (0, 2], s = 1, 2,\tag{4.4}$$

where κ_s is the heat conduction constant, and $u_s = u_s(x, t)$ is the temperature of the s -th surface, $x \in \mathbb{R}, t > 0$, and $\frac{\partial^{\alpha_s}}{\partial t^{\alpha_s}}(\cdot)$ represents α_s -th order of the Caputo time-fractional derivative (1.2). Lenzi [84] used the multi-component equations (4.4) to describe the transport and anomalous diffusion separated by a membrane in heterogeneous systems. In many complex physical phenomena, the convection terms appear naturally along with the anomalous diffusion processes. Moreover, long-time interactions in systems with more than one species can be described using the class of system of time-fractional diffusion equations [9, 81–84]. The class of coupled TFDCWEs, with or without including source/sink terms, have many potential applications in engineering, biology, chemistry, and physics, primarily in the fluid flow problems in porous media. The approximate analytical solution for the $(1 + 1)$ -dimensional coupled nonlinear time-fractional diffusion-convection equations with source term by imposing the Dirichlet boundary conditions were investigated in [79].

In general, finding analytical solutions for coupled nonlinear fractional diffusion equations is a challenging and difficult task. However, analytical methods like the invariant subspace method [46–48, 50, 55] and Lie symmetry method [31, 85–87] were proven to be helpful in finding analytical solutions of coupled and scalar nonlinear diffusion equations and coupled nonlinear diffusion-convection equations in $(1 + 1)$ -dimensions in both integer-order and fractional-order cases. A few analytical studies only exist in the literature for studying analytical solutions of coupled NTFPDEs, particularly in higher dimensions [55]. Additionally, we wish to point out that we can expect the following kinds of analytical solutions $\mathbf{U} = (u_1(x_1, x_2, t), u_2(x_1, x_2, t))$ for the multi-component $(2 + 1)$ -dimensional coupled nonlinear integer-order diffusion-convection equations as

(a) Functional separable analytical solution: $\mathbf{U} = (\Phi_1(z), \Phi_2(z))$,

$$\text{where } z = \sum_{i_2=1}^{n_2} \sum_{i_1=1}^{n_1} \varphi_{i_1, i_2}(t) \prod_{j=1}^2 \psi_{i_j}(x_j).$$

(b) Travelling-wave analytical solution: $\mathbf{U} = (\Psi_1(z), \Psi_2(z))$,

$$\text{where } z = k_1 t + k_2 x_1 + k_3 x_2, k_r \in \mathbb{R}, r = 1, 2, 3. \quad (4.5)$$

(c) Generalized separable analytical solutions:

$$(I) \quad \mathbf{U} = \left(\sum_{i=1}^{n_1} \varphi_i^1(t) \psi_i^1(z), \sum_{i=1}^{n_2} \varphi_i^1(t) \psi_i^1(z) \right), z = k_1 x_1 + k_2 x_2, k_r \in \mathbb{R}, r = 1, 2,$$

$$(II) \quad \mathbf{U} = \left(\sum_{i_2=1}^{n_2} \sum_{i_1=1}^{n_1} \varphi_{i_1, i_2}^1(t) \prod_{j=1}^2 \psi_{i_j}^1(x_j), \sum_{i_2=1}^{m_2} \sum_{i_1=1}^{m_1} \varphi_{i_1, i_2}^2(t) \prod_{j=1}^2 \psi_{i_j}^2(x_j) \right).$$

Here $n_j, m_j \in \mathbb{N}$, the unknown functions $\varphi_{i_1, i_2}(t)$, $\varphi_{i_1, i_2}^s(t)$ and $\psi_{i_j}(x_j)$, $\psi_{i_j}^s(x_j)$, $s, j = 1, 2$, are to be determined, which depend on the considered coupled system. We can obtain solutions of the form (a), (b) and (c) for the multi-component $(2 + 1)$ -dimensional coupled nonlinear integer-order diffusion-convection equations through the functional and generalized separable methods, respectively. However, for fractional-order case in the sense of the Riemann-Liouville and Caputo fractional-order derivatives, the functional separable solutions (a) and travelling-wave analytical solutions (b) are not possible for the the multi-component $(2 + 1)$ -dimensional coupled TFDCWEs (1.7) with singular kernel fractional-order derivatives. Hence, we can expect only generalized separable solutions (c) for the multi-component $(2 + 1)$ -dimensional coupled TFDCWEs (1.7) under singular kernel fractional derivatives. So, in this work, we discussed the generalized separable solutions (c)-(II) for the given multi-component $(2 + 1)$ -dimensional coupled nonlinear TFDCWEs (1.7) under the Caputo fractional derivative through the invariant subspace method. We also note that only a few papers studied the analytical solutions for the coupled NTFPDEs in higher dimensions. For the first time, we developed the theory of invariant subspace method to find the analytical solutions of the multi-component $(N + 1)$ -dimensional coupled NTFPDEs. In this work, we also studied how to find analytical solutions of a more general class of multi-component $(2 + 1)$ -dimensional coupled nonlinear TFDCWEs with suitable initial-boundary conditions. Here, we used the Dirichlet boundary conditions for the multi-component $(2 + 1)$ -dimensional coupled TFDCWEs defined in a bounded domain. Furthermore, we would like to emphasize that the invariant subspace method discussed in this article is applicable to all kinds of fractional derivatives, which are available in the literature, not only the Caputo fractional derivative. Also, we note that some interesting investigations have been made in the existing literature for the invariant subspace method to find generalized separable analytical solutions for fractional PDEs in $(1 + 1)$ -dimension under different fractional derivatives, such as the Riemann-Liouville fractional derivative [38, 47, 48, 55], and the Regularized Prabhakar fractional derivative [56]. In future, we will study the details of applications for the developed invariant subspace

method to obtain generalized separable analytical solutions for higher-dimensional coupled systems of time-fractional PDEs under different types of fractional derivatives.

In the following subsection, we summarize the importance of the developed method and the results obtained from this work.

4.2 Concluding remarks

In the present work, we have systematically developed the generalization of the invariant subspace method for finding the invariant product linear space and generalized separable analytical solutions of multi-component $(N + 1)$ -dimensional coupled NTFPDEs (2.1) for the first time. Explicitly, we have provided a detailed description for constructing the various kinds of type- p invariant product linear spaces for the coupled NTFPDEs given in (2.1) for $p = 1, 2, \dots, N$. Also, we wish to mention here that one can expect N different types of generalized separable analytical solutions for the coupled NTFPDEs (2.1) based on the obtained type- p invariant product linear spaces, $p = 1, 2, \dots, N$. Additionally, we have explained how to derive analytical solutions for the given coupled NTFPDEs (2.1) by reducing it into a system of FODEs using the invariant product linear spaces determined. More precisely, we successfully illustrated the efficacy of the developed method through the well-known multi-component $(2 + 1)$ -dimensional coupled nonlinear TFDCWEs (1.7) and constructed their invariant product linear spaces. Using the type-1 and type-2 invariant product linear spaces determined for coupled equations (1.7), we have explained how to obtain the analytical solutions of the coupled equations (1.7) with the appropriate initial conditions. Finally, we wish to point out that one can expect two types of analytical solutions for the coupled equations (1.7) based on the type-1 and type-2 invariant product linear spaces.

Additionally, we wish to point out that the discussed invariant subspace method is applicable for finding the analytical solutions of scalar and coupled systems of nonlinear PDEs involving both time and space fractional derivatives. So, we can use this method to find analytical solutions for coupled system of time and space fractional diffusion-convection-wave equations under different fractional-order derivatives. In addition, we wish to point out that analytical solutions of scalar and coupled time-space fractional nonlinear PDEs have been derived through the invariant subspace method in [47, 48]. However, they have not developed the systematic algorithmic approach of the invariant subspace method to obtain analytical solutions for scalar and coupled time-space fractional nonlinear PDEs in the literature. So, in the future, we plan to develop systematically the algorithmic approach of the invariant subspace method for finding the analytical solutions of scalar and coupled time-space fractional nonlinear PDEs. We hope that the discussed systematic investigation will help to study the exact descriptions of many mutually connected real-world complex phenomena more accurately. This study will help to prove the effectiveness of the invariant subspace method, which is a powerful systematic and analytical tool for finding the generalized separable analytical solutions for complex coupled nonlinear fractional PDEs. We strongly believe that due to the nonstandard properties of singular kernel fractional-order derivatives, the present development of the invariant subspace method will be very useful in constructing various kinds of generalized separable analytical solutions for coupled and scalar nonlinear fractional PDEs arising in various areas of science and engineering.

Acknowledgments

The first author is thankful for the financial assistance received in the form of International Mathematical Union Breakout Graduate fellowship-2023 (IMU-BGF-2023-06) provided by the IMU, Germany. Another

author (M.L.) receives support from the Department of Science and Technology, India-SERB National Science Chair position (NSC/2020/000029).

Declaration of competing interest

The authors declare that they have no known competing interest.

References

- [1] V.E. Tarasov and J.J. Trujillo, *Ann. Physics* **334** (2013) 1-23.
- [2] V.E. Tarasov, *Fractional Dynamics: Applications of Fractional Calculus to Dynamics of Particles, Fields and Media, Nonlinear Physical Science* (Springer, Heidelberg, Germany, 2011.)
- [3] C. Ionescu, A. Lopes, D. Copot, J.A.T. Machado and J.H.T. Bates, *Commun. Nonlinear Sci. Numer. Simulat.* **51** (2017) 141-159.
- [4] R.L. Bagley and P.J. Torvik, *ASME J. Appl. Mech.* **51** (1984) 294-298.
- [5] V.E. Tarasov, *Internat. J. Modern Phys. B* **27** (2013) 1330005.
- [6] I. Podlubny, *Fractional Differential Equations* (Academic Press, New York, 1999).
- [7] K. Diethelm, *The Analysis of Fractional Differential Equations* (Springer, Berlin, 2010).
- [8] F. Mainardi, *Fractional Calculus: Some Basic Problems in Continuum and Statistical Mechanics*, in *Fractals and Fractional Calculus in Continuum Mechanics*, eds., A. Carpinteri and F. Mainardi (Springer, New York, 1997), pp. 291-348.
- [9] Y.Z. Povstenko, *J. Thermal Stresses*, **36** (2013) 351-363.
- [10] Y. Povstenko, *Fractional Thermoelasticity*, (Springer, Switzerland, 2015).
- [11] T. Bakkyaraj and R. Sahadevan, *Pramana-J. Phys.* **85** (2015) 849-860.
- [12] S.Y. Lukashchuk, *Nonlinear Dyn.* **80** (2015) 791-802.
- [13] X.Y. Zhang, Y. Peng, Y.J. Xie and X.F. Li, *Z. Angew. Math. Phys.* **70** (2019) 2.
- [14] Y. Yang, W. Qiao, J. Wang and S. Zhang, *Comput. Math. Appl.* **78** (2019) 1431-1449.
- [15] V. Daftardar-Gejji and H. Jafari, *J. Math. Anal. Appl.* **301** (2005) 508-518.
- [16] H. Jafari and V. Daftardar-Gejji, *J. Comput. Appl. Math.* **196** (2006) 644-651.
- [17] V. Gafiychuk, B. Datsko, V. Meleshko and D. Blackmore, *Chaos, Solitons and Fractals* **41** (2009) 1095-1104.
- [18] A. Bekir, Ö. Güner and A.C. Cevikel, *Abstr. Appl. Anal.* **2013** (2013) 426462.
- [19] A. Bekir, O. Guner and A.C. Cevikel, *IEEE/CAA J. Autom. Sin.* **4** (2017) 315-321.
- [20] A. Bekir, E. Aksoy and A.C. Cevikel, *Math. Meth. Appl. Sci.* **38** (2015) 3807-3817.
- [21] Z. Odibat and S. Momani, *Appl. Math. Lett.* **21** (2008) 194-199.
- [22] W.X. Ma, M.M. Mousa and M.R. Ali, *Mod. Phys. Lett. B* **34** (2020) 2050049.
- [23] C. Wu and W. Rui, *Commun. Nonlinear Sci. Numer. Simulat.* **63** (2018) 88-100.
- [24] W. Rui and H. Zhang, *Comp. Appl. Math.* **39** (2020) 1-26.
- [25] W. Rui, X. Yang and F. Chen, *Physica A.* **595** (2022) 127068.
- [26] W. Rui, *Nonlinear Dyn.* **109** (2022) 943-961.
- [27] H. Jafari and S. Seifi, *Commun. Nonlinear Sci. Numer. Simulat.* **14** (2009) 2006-2012.
- [28] H. Jafari and S. Momani, *Phys. Lett. A* **370** (2007) 388-396.
- [29] R.K. Gazizov, A.A. Kasatkin and S.Y. Lukashchuk, *Phys. Scr.* **T136** (2009) 014016.
- [30] R. Sahadevan and T. Bakkyaraj, *J. Math. Anal. Appl.* **393** (2012) 341-347.

- [31] R. Sahadevan and P. Prakash, *Chaos, Solitons and Fractals* **104** (2017) 107-120.
- [32] P. Prakash and R. Sahadevan, *Nonlinear Dyn.* **89** (2017) 305-319.
- [33] A.M. Nass, *Appl. Math. Comput.* **347** (2019) 370-380.
- [34] T. Bakkyaraj, *Eur. Phys. J. Plus* **135** (2020) 126.
- [35] R. Sahadevan and P. Prakash, *Int. J. Dynamical Systems and Differential Equations* **9** (2019) 44-64.
- [36] P. Prakash, *Comput. Appl. Math.* **40** (2021) 162.
- [37] K. Sethukumarasamy, P. Vijayaraju and P. Prakash, *J. Nonlinear Math. Phys.* **28** (2021) 219-241.
- [38] R.K. Gazizov and A.A. Kasatkin, *Comput. Math. Appl.* **66** (2013) 576-584.
- [39] P. Artale Harris and R. Garra, *Nonlinear Stud.* **20** (2013) 471-481.
- [40] R. Sahadevan and T. Bakkyaraj, *Fract. Calc. Appl. Anal.* **18** (2015) 146-162.
- [41] R. Sahadevan and P. Prakash, Exact solution of certain time-fractional nonlinear partial differential equations, *Nonlinear Dyn.* 85(2016) 659-673.
- [42] S. Choudhary and V. Daftardar-Gejji, *Fract. Calc. Appl. Anal.* **20** (2017) 477-493.
- [43] P. Artale Harris and R. Garra, *J. Math. Phys.* **58** (2017) 063501.
- [44] P. Artale Harris and R. Garra, *Commun. Appl. Industr. Math.* **6** (2014) e-487.
- [45] W. Rui, *Appl. Math. Comput.* **339** (2018) 158-171.
- [46] R. Sahadevan and P. Prakash, *Commun. Nonlinear Sci. Numer. Simulat.* **42** (2017) 158-177.
- [47] S. Choudhary and V. Daftardar-Gejji, *Int. J. Model. Simul. Sci. Comput.* **10** (2019) 1941010.
- [48] P. Prakash, *Pramana-J. Phys.* **94** (2020) 103.
- [49] W. Feng and S. Zhao, *Mod. Phys. Lett. B* **32** (2018) 1850401.
- [50] P. Prakash, K.S. Priyendhu and M. Meenakshi, *Comp. Appl. Math.* **43** (2024) 30.
- [51] J.G. Liu, X.J. Yang, Y.Y. Feng, *Mod. Phys. Lett. B* **34** (2020) 2050009.
- [52] P. Prakash, S. Choudhary and V. Daftardar-Gejji, *Eur. Phys. J. Plus* **135** (2020) 490(24p).
- [53] R. Garra, Z. Tomovski, *Math. Model. Anal.* **26** (2021) 72-81.
- [54] A.D. Polyanin and A.I. Zhurov, *Separation of Variables and Exact Solutions to Nonlinear PDEs* (CRC Press, Oxon, 2022).
- [55] S. Choudhary, P. Prakash and V. Daftardar-Gejji, *Comp. Appl. Math.* **38** (2019) 126.
- [56] Y.M. Chu, M. Inc, M.S. Hashemi and S. Eshaghi, *Comp. Appl. Math.* **41** (2022) 271.
- [57] A.H. Abdel Kader, M.S. Abdel Latif and D. Baleanu, *Math. Meth. Appl. Sci.* **44** (2021) 4701-4714.
- [58] P. Prakash, K.S. Priyendhu and K.M. Anjitha, *Comp. Appl. Math.* **41** (2022) 30.
- [59] P. Prakash, K.S. Priyendhu and M. Lakshmanan, *Commun. Nonlinear Sci. Numer. Simulat.* **111** (2022) 106436.
- [60] K.S. Priyendhu, P. Prakash and M. Lakshmanan, *Commun. Nonlinear Sci. Numer. Simulat.* **122** (2023) 107245.
- [61] P. Prakash, R. Thomas and T. Bakkyaraj, *Comp. Appl. Math.* **42** (2023) 97.
- [62] V.A. Galaktionov and S.R. Svirshchevskii, *Exact Solutions and Invariant Subspaces of Nonlinear Partial Differential Equations in Mechanics and Physics* (Chapman and Hall/CRC, London, 2007).
- [63] W.X. Ma, *Sci. China Math.* **55** (2012) 1769-1778.
- [64] C. Zhu and C. Qu, *Symmetry* **8** (2016) 128.
- [65] W.X. Ma and Y. Liu, *Commun. Nonlinear Sci. Numer. Simulat.* **17** (2012) 3795-3801.
- [66] W.X. Ma, Y. Zhang, Y. Tang and J. Tu, *Appl. Math. Comput.* **218** (2012) 7174-7183.
- [67] Y. Ye, W.X. Ma, S. Shen and D. Zhang, *J. Nonlinear Math. Phys.* **21** (2014) 132-148.

- [68] C. Qu and C. Zhu, *J. Phys. A: Math. Theor.* **42** (2009) 475201.
- [69] J. Song, S. Shen, Y. Jin and J. Zhang, *Commun. Nonlinear Sci. Numer. Simulat.* **18** (2013) 2984-2992.
- [70] P. Prakash, *Eur. Phys. J. Plus* **134** (2019) 261.
- [71] A.D. Polyanin and A.I. Zhurov, *Appl. Math. Lett.* **37** (2014) 43-48.
- [72] S.E. Esipov, *Phys. Rev. E* **52** (1995) 3711-3718.
- [73] S. Wu, B. Peng and Z.F. Tian, *Appl. Numer. Math.* **146** (2019) 89-122.
- [74] L.R. Evangelista and E.K. Lenzi, *Fractional Diffusion Equations and Anomalous Diffusion* (Cambridge University Press, Cambridge, 2018).
- [75] B. Cockburn and C.W. Shu, *SIAM J. Numer. Anal.* **35** (1998) 2440-2463.
- [76] S. Axler, *Linear Algebra Done Right* (Springer, Heidelberg, 2014).
- [77] F. Rossi, M.A. Budroni, N. Marchettini and J.C. Landeira, *Chaos* **22** (2012) 037109.
- [78] C.A.J. Fletcher, *Int. J. Numer. Methods. Fluids* **3** (1983) 213-216.
- [79] P. Pandey, S. Kumar and S. Das, *Eur. Phys. J. Plus* **134** (2019) 364.
- [80] A.M. Turing, *Trans. R. Soc. Lond. B*, **237** (1952) 37-72.
- [81] B. Datsko, Y. Luchko and V. Gafiychuk, *Int. J. Bifurcat. Chaos* **22** (2012) 1250087.
- [82] B. Datsko and V. Gafiychuk, *Fract. Calc. Appl. Anal.* **21** (2018) 237-253.
- [83] T.A.M. Langlands, B.I. Henry and S.L. Wearne, *Phys. Rev. E* **77** (2008) 021111.
- [84] E.K. Lenzi, H.V. Ribeiro, A.A. Tateishi, R.S. Zola and L.R. Evangelista, *Proc. R. Soc. A* **472** (2016) 20160502.
- [85] R. Cherniha and J.R. King, *J. Math. Anal. Appl.* **308** (2005) 11-35.
- [86] P. S. Bindu, M. Senthilvelan and M. Lakshmanan, *J. Phys. A: Math. Gen.* **34** (2001) L689.
- [87] P. S. Bindu, M. Senthilvelan and M. Lakshmanan, *Int. J. Bifurcat. Chaos* **14** (2004) 1577-1600.