

THE STRUCTURE OF PERIODIC POINT FREE DISTAL HOMEOMORPHISMS ON THE ANNULUS

ENHUI SHI, HUI XU*, AND ZIQI YU

ABSTRACT. Let A be an annulus in the plane \mathbb{R}^2 and $g : A \rightarrow A$ be a boundary components preserving homeomorphism which is distal and has no periodic points. Then there is a continuous decomposition of A into g -invariant circles such that all the restrictions of g on them share a common irrational rotation number and all these circles are linearly ordered by the inclusion relation on the sets of bounded components of their complements in \mathbb{R}^2 .

1. INTRODUCTION

Recurrence is one of the most fundamental notions in the theory of dynamical system. There are various definitions to describe the recurrence behaviors of a point in a system, such as periodic point, almost periodic point, distal point, recurrent point, regularly recurrent point, and so on. There has been a considerable progress in studying the structures of the dynamical systems all points of which possess some kind of recurrence.

Montgomery [28] proved that every pointwise periodic homeomorphism on a connected manifold is periodic. For an infinite compact minimal metric system each point of which is regularly recurrent, Block and Keesling [5] proved that it is topologically conjugate to an adding machine. Shi, Xu, and Yu [35] showed that every pointwise recurrent expansive homeomorphism is topologically conjugate to a subshift of some symbol system, which extends a classical result of Mañé [26] for minimal expansive homeomorphisms. The structure of pointwise recurrent maps having the pseudo orbit tracing property is completely determined by Mai and Ye [25].

There are also many interesting results around the structures of recurrent maps on low-dimensional spaces. Mai [24] showed that a pointwise recurrent graph map is either topologically conjugate to an irrational rotation on the circle or of finite order. Naghmouchi [31] and Blokh [6] characterized the structures of pointwise recurrent maps on uniquely arcwise connected curves. Kolev and Pérouème [20] showed recurrent homeomorphisms

*Corresponding author.

on compact surfaces with negative Euler characteristic are of finite order. Foland [14] proved that any equicontinuous homeomorphism on a closed 2-cell is topologically conjugate either to a reflection of a disk in a diameter or to a rotation of a disk about its center. Ritter [34] further determined the structure of equicontinuous homeomorphisms on the 2-sphere and annulus. Oversteegen and Tymchatyn [32] proved that recurrent homeomorphisms on the plane are periodic.

The notion of distality was introduced by Hilbert for better understanding equicontinuity [12]. The study of minimal distal systems culminates in the beautiful structure theorem of Furstenberg [16], which describes completely the relations between distality and equicontinuity for minimal systems. Considering minimal distal actions on compact manifolds, Rees [33] proved a sharpening of Furstenberg's structure theorem.

The aim of the paper is to study the structure of distal homeomorphisms on annulus without periodic points. One may consult [3, 7, 15, 17] for many interesting related investigations.

We obtain the following theorem.

Theorem 1.1. *Let A be an annulus in the plane \mathbb{R}^2 and $g : A \rightarrow A$ be a boundary components preserving homeomorphism which is distal and has no periodic points. Then there is a continuous decomposition of A into g -invariant circles such that all the restrictions of g on them share a common irrational rotation number and all these circles are linearly ordered by the inclusion relation on the sets of bounded components of their complements in \mathbb{R}^2 .*

The paper is organised as follows. In Section 2, we will introduce some concepts and facts in the theories of dynamical system and topology. Specially, we will give the definitions of solenoid and adding machine from the viewpoint of topological groups and recall some results around distal homeomorphisms. In Section 3, we will show that there exists no adding machine contained in the boundary of an f -invariant open disk under some appropriate assumptions. Based on this result, we show in Section 4 the existence of an f -invariant circle in the boundary just mentioned. In Section 5, we show further that there are sufficiently many f -invariant circles in the annulus. Relying on all these results, we show in Section 6 the existence of the expected decompositions.

2. PRELIMINARIES

In this section, we will recall some notions, notations, and elementary facts in the theories of dynamical system and topology.

2.1. Recurrence, minimal sets, and factors. By a *dynamical system* we mean a pair (X, f) , where X is a metric space and $f : X \rightarrow X$ is a homeomorphism. For $x \in X$, the *orbit* of x is the set $O(x, f) \equiv \{f^i(x) : i \in \mathbb{Z}\}$. If there is some $n > 0$ such that $f^n(x) = x$, then x is called a *periodic point* of f and the minimal such n is called the *period* of x . A periodic point x of period 1 is called a *fixed point*, that is $f(x) = x$. A subset A of \mathbb{Z} is *syndetic* if there is $l > 0$ with $A \cap \{p, p+1, \dots, p+l\} \neq \emptyset$ for any $p \in \mathbb{Z}$. We call x an *almost periodic point* if for any open neighborhood U of x , the set $N(x, U) \equiv \{i : f^i(x) \in U\}$ is syndetic. If there is a sequence of positive integers $n_1 < n_2 < \dots$ such that $f^{n_i}(x) \rightarrow x$, then we call x a *recurrent point*; and if for any open neighborhood U of x , there always exists $n > 0$ such that $f^n(U) \cap U \neq \emptyset$, then we call x a *nonwandering point*. Clearly, an almost periodic point is recurrent and a recurrent point is nonwandering. If each point of X is nonwandering, then we call f *nonwandering*.

A subset S of X is *f-invariant* if $f(S) = S$; we use $f|_S$ to denote the restriction of f to S . If S is an *f-invariant* nonempty closed subset of X and contains no proper *f-invariant* closed subset, then we call S a *minimal set* of f . If X is a minimal set, we call the system (X, f) is *minimal*. It is clear from the definition that S is minimal if and only if for each $x \in S$, $O(x, f)$ is dense in S . By an argument of Zorn's lemma, we have that if X is compact, then there always exists a minimal set of f . We have known that each point of a compact minimal set is almost periodic (see e.g. [1, Chap.1-Theorem 1]).

For any two dynamical systems (X, f) and (Y, g) , if there is a continuous surjection $\phi : X \rightarrow Y$ such that $\phi \circ f = g \circ \phi$, then we say that (Y, g) is a *factor* of (X, f) and (X, f) is an *extension* of (Y, g) ; we call ϕ a *factor map* or a *semiconjugation* between (X, f) and (Y, g) ; if ϕ is a homeomorphism, then we call (X, f) and (Y, g) are *topologically conjugate*. Clearly, if M is a minimal set of f , then $\phi(M)$ is a minimal set of g . It is well known that if \mathbb{S}^1 is the unit circle and $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ is an orientation preserving homeomorphism without periodic points, then (\mathbb{S}^1, f) is semiconjugate to a rigid minimal rotation on \mathbb{S}^1 (see e.g. [36, Theorem 6.18]).

A topological space U is called an *open disk* if it is homeomorphic to the unit open disk in the plane \mathbb{R}^2 . By Riemann mapping theorem, we know that an open subset U of \mathbb{C} is an open disk if and only if it is simply connected.

The following theorem is implied by Brouwer' lemma (see e.g. [13, 23]).

Theorem 2.1. *If U is an open disk and $f : U \rightarrow U$ is an orientation-preserving nonwandering homeomorphism, then f has a fixed point in U .*

2.2. Continuous and semi-continuous decompositions. Let (X, \mathcal{T}) be a topological space. A *partition* of X is a collection \mathcal{D} of nonempty, mutually disjoint subsets of X such that $\cup \mathcal{D} = X$. Define $\pi : X \rightarrow \mathcal{D}$ by letting $\pi(x)$ be the unique $D \in \mathcal{D}$ such that $x \in D$ for each $x \in X$. We endow \mathcal{D} with the largest topology so that π is continuous, that is $\mathcal{U} \subset \mathcal{D}$ is open iff $\cup \mathcal{U} \in \mathcal{T}$. The topological space \mathcal{D} so defined is called the *decomposition* of X . We also call \mathcal{D} the *quotient space* of X by identifying each element of \mathcal{D} into a point and call π the *quotient map*. The partition \mathcal{D} is called *upper semi-continuous* provided that whenever $D \in \mathcal{D}$, $U \in \mathcal{T}$, and $D \subset U$, there exists $V \in \mathcal{T}$ with $D \subset V$ such that if $A \in \mathcal{D}$ and $A \cap V \neq \emptyset$, then $A \subset U$.

Now suppose X is a compact metric space with metric d . Let 2^X be the collection of all nonempty closed subsets of X and let $C(X) = \{A \in 2^X : A \text{ is connected}\}$. The *Hausdorff metric* H_d on 2^X is defined by $H_d(A, B) = \inf\{\varepsilon : A \subset B_d(B, \varepsilon) \text{ and } B \subset B_d(A, \varepsilon)\}$ for each $A, B \in 2^X$. Then 2^X and $C(X)$ are both compact metric spaces with respect to H_d , called the *hyperspaces* of X (see e.g. [30, Theorems 4.13 and 4.17]). Let \mathcal{D} be a partition of X such that each element of \mathcal{D} is closed. The partition \mathcal{D} is called *continuous* if the quotient map $\pi : X \rightarrow \mathcal{D}$, thought of as a map from X into 2^X , is continuous.

If \mathcal{D} is a partition of X , then it induces an equivalence relation $R \subset X \times X$ by defining $(x, y) \in R$ if $\{x, y\} \subset A$ for some $A \in \mathcal{D}$. The relation R is called a *closed relation* if it is a closed subset of $X \times X$.

The following proposition can be seen in [18, Proposition 2.2].

Proposition 2.2. *Let X be a compact metric space. Let \mathcal{D} be a partition of X and R be the equivalence relation induced by \mathcal{D} . If R is closed, then \mathcal{D} is upper semi-continuous.*

Proposition 2.2 together with the compactness of hyperspaces implies the following proposition.

Proposition 2.3. *Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a homeomorphism and $M \subset \mathbb{R}^2$ be a compact minimal set of f . Let \mathcal{M} be the set of all components of M and let $\mathcal{M}' = \{\{x\} : x \in \mathbb{R}^2 \setminus M\}$. Then $\mathcal{M} \cup \mathcal{M}'$ is an upper semi-continuous decomposition of \mathbb{R}^2 .*

A *continuum* is a connected compact metric space. If X is a continuum contained in the plane \mathbb{R}^2 such that $\mathbb{R}^2 \setminus X$ is connected, then we call that X *does not separate the plane*. The following theorem is due to R. L. Moore (see e.g. [22, p.533, Theorem 8] for a slightly more general form).

Theorem 2.4. *The space of a upper semi-continuous decomposition of \mathbb{R}^2 into continua, which do not separate \mathbb{R}^2 , is homeomorphic to \mathbb{R}^2 .*

2.3. Solenoids and adding machines. Let X be a compact metric space with metric d and $f : X \rightarrow X$ be a homeomorphism. We say (X, f) is *equicontinuous* if for each $\varepsilon > 0$, there is a $\delta > 0$ such that $d(f^i(x), f^i(y)) < \varepsilon$ for any $i \in \mathbb{Z}$, whenever $d(x, y) < \delta$. Let K be a compact abelian metric group and $a \in K$. The *rotation* $\rho_a : K \rightarrow K$ is defined by $\rho_a(x) = ax$ for any $x \in K$. Clearly, if $\{a^n : n \in \mathbb{Z}\}$ is dense in K , then (K, ρ_a) is minimal and equicontinuous.

The following theorem shows that minimal rotations on compact abelian metric groups are the only equicontinuous minimal systems (see [36, Theorem 5.18]).

Theorem 2.5 (Halmos-von Neumann). *Let X be a compact metric space and $f : X \rightarrow X$ be a minimal and equicontinuous homeomorphism. Then (X, f) is topologically conjugate to a minimal rotation on a compact abelian metric group.*

For each positive integer i , let K_i be a compact metric group and let $f_i : K_{i+1} \rightarrow K_i$ be a surjective continuous group homomorphism. The *inverse limit* of $\{K_i, f_i\}$ is

$$\lim_{\leftarrow} \{K_i, f_i\} \equiv \{(x_i) \in \prod_{i=1}^{\infty} K_i : x_i = f_i(x_{i+1})\},$$

which is a compact metric group under the multiplication “ \cdot ” defined by $(x_i) \cdot (y_i) = (x_i y_i)$. If each K_i is the unit circle $\{z \in \mathbb{C} : |z| = 1\}$ and $\lim_{\leftarrow} \{K_i, f_i\}$ is not the circle, then we call $\lim_{\leftarrow} \{K_i, f_i\}$ a *solenoid*; if each K_i is a finite cyclic group and $\lim_{\leftarrow} \{K_i, f_i\}$ is not finite, then we call $\lim_{\leftarrow} \{K_i, f_i\}$ an *adding machine*. As topological spaces, a solenoid is a homogeneous indecomposable circle-like continuum and an adding machine is a Cantor set. We also call a minimal rotation on an adding machine an adding machine.

Since every compact metric group is an inverse limit of compact Lie groups (see e.g. [29, Chap. 4.6-4.7]), and the only connected Lie groups of dimension 1 is the circle group and the only Lie groups of dimension 0 are finite groups, the following proposition is clear.

Proposition 2.6. *If K is a connected compact metric group of dimension 1, then it is either a circle or a solenoid; if K is a compact metric group of dimension 0 and has a dense cyclic subgroup, then it is either a finite cyclic group or an adding machine.*

A curve is an 1-dimensional continuum. The following corollaries are immediate from Theorem 2.5 and Proposition 2.6.

Corollary 2.7. *Let X be a curve and $f : X \rightarrow X$ be a minimal equicontinuous homeomorphism. Then (X, f) is topologically conjugate to a minimal rotation either on the circle or on a solenoid.*

Corollary 2.8. *Let X be a compact metric space of dimension 0 and $f : X \rightarrow X$ be a minimal equicontinuous homeomorphism. Then (X, f) is either a periodic orbit or topologically conjugate to an adding machine.*

The following proposition is shown by Bing [4], which is also implied by the main result in [19].

Proposition 2.9. *Solenoids are not planar continua.*

We call x in a system (X, f) *regularly recurrent* if for any open neighborhood U of x , there is a positive integer n such that $f^{kn}(x) \in U$ for each $k = 0, 1, \dots$.

The following proposition is implied by the definition (see [5] for a characterization of adding machine using regular recurrence).

Proposition 2.10. *If (X, f) is an adding machine, then each point x of X is regularly recurrent.*

2.4. Structures of distal homeomorphisms. Let X be a compact metric space with metric d and let $f : X \rightarrow X$ be a homeomorphism. We call that (X, f) is *distal* if for any $x \neq y \in X$, $\inf_{i \in \mathbb{Z}} \{d(f^i(x), f^i(y))\} > 0$. We suggest the readers to consult [1] for the proofs of the following well known facts: (1) If (X, f) is distal, then X is a disjoint union of minimal sets; (2) If (X, f) is minimal and distal and (Y, g) is a factor of (X, f) , then

(Y, g) is also minimal and distal; (3) Let (X, f) be a minimal distal system. Then it has a maximal equicontinuous factor.

Lemma 2.11. [33, §6] *Let (X, f) be a minimal distal system and $\pi : (X, f) \rightarrow (Y, g)$ be a factor map. Then the covering dimension of the fibers $\pi^{-1}(y), y \in Y$, is constant and*

$$\dim(X) = \dim(Y) + \dim \pi^{-1}(y).$$

Lemma 2.12. [8, p.192, Theorem 3·17.13] *Let (X, f) be a minimal distal system and $\pi : (X, f) \rightarrow (Y, g)$ be a factor map. If (Y, g) is equicontinuous and there is some $y \in Y$ with $\dim \pi^{-1}(y) = 0$, then (X, f) is also equicontinuous.*

Clearly, Lemma 2.12 implies a distal minimal system of zero dimension is equicontinuous. In fact, this is also true for non-minimal distal systems of zero dimension (see [2, Corollary 1.9]).

Proposition 2.13. *Let X be a compact connected metric space and $f : X \rightarrow X$ be a minimal distal homeomorphism. If $\dim(X) = 1$, then (X, f) is equicontinuous.*

Proof. Suppose that (X, f) is not equicontinuous. Then the maximal equicontinuous factor of (X, f) is nontrivial. Let $\pi : (X, f) \rightarrow (Y, g)$ be the factor map to its maximal equicontinuous factor. In particular, Y is connected. By Lemma 2.11, we have $\dim(Y) = 1$ and for each $y \in Y$, $\dim \pi^{-1}(y) = 0$. Then it follows from Lemma 2.12 that (X, f) is equicontinuous. This contradiction shows that (X, f) is equicontinuous. \square

3. NONEXISTENCE OF AN ADDING MACHINE IN THE BOUNDARY OF AN OPEN DISK

We call a topological space X is an *arc* (resp. *open arc*) if it is homeomorphic to the closed interval $[0, 1]$ (resp. the open interval $(0, 1)$). We call X a *circle* if it is homeomorphic to the unit circle in the plane, that is X is a simple closed curve.

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be an orientation preserving homeomorphism and $U \subset \mathbb{R}^2$ be a bounded f -invariant open disk. A *cross-cut* of U is an open arc γ in U with $\bar{\gamma}$ being an arc jointing two points of ∂U . A *cross-section* of U is any connected component of $U \setminus \gamma$ for some cross cut γ of U . A *chain* for U is a sequence of sections $\mathcal{C} = (D_i)_{i=1}^{\infty}$ such that $D_1 \supset D_2 \supset \dots$ and $\overline{\partial_U D_i} \cap \overline{\partial_U D_j} = \emptyset$ for all $i \neq j$. Two chains $(D_i)_{i=1}^{\infty}$ and $(D'_i)_{i=1}^{\infty}$ are called *equivalent* if for any $i > 0$ there is $j > i$ such that $D_j \subset D'_i$ and $D'_j \subset D_i$. A chain $(D_i)_{i=1}^{\infty}$ is called a *prime chain* if $\text{diam}(\partial_U D_i) \rightarrow 0$. An equivalence class of prime chains

is called a *prime end* of U . We use $b_\varepsilon(U)$ to denote the set of all prime ends of U . Let $\hat{U} = U \cup b_\varepsilon(U)$.

Now we topologize \hat{U} as follows. For a cross-section D of U and for a prime chain (U_i) representing $p \in b_\varepsilon(U)$, if $U_i \subset D$ for sufficiently large i , then we call p *divides* D . Set $\mathcal{E}(D) = \{p \in b_\varepsilon(U) : p \text{ divides } D\}$. Consider the family \mathcal{B} consisting of all sets of the form $D \cup \mathcal{E}(D)$ for some cross-section D , together with all open subsets of U . Then \mathcal{B} is a topological basis on \hat{U} . We endow \hat{U} with the topology generated by \mathcal{B} .

The following theorem is known as the Carathéodory's prime ends compactification theorem (see [9, 10]).

Theorem 3.1 (Prime ends compactification). *\hat{U} is homeomorphic to the unit closed disk and $b_\varepsilon(U)$ is homeomorphic to the unite circle \mathbb{S}^1 .*

It is well known that the homeomorphism $f|_U$ can be extended to a homeomorphism $\hat{f} : \hat{U} \rightarrow \hat{U}$. We call the rotation number of $\hat{f}|_{\mathbb{S}^1}$ the *prime ends rotation number* of $f|_{\overline{U}}$.

The following theorem is due to Cartwright and Littlewood [11]. One may consult [21] for the proof of the converse direction under more general settings.

Theorem 3.2. *If f is nonwandering and has no periodic point in ∂U , then the prime ends rotation number of $f|_{\overline{U}}$ is irrational.*

Now we use Theorem 3.2 to prove a key result.

Proposition 3.3. *If the prime ends rotation number of $f|_{\overline{U}}$ is irrational, then no minimal set in ∂U is an adding machine.*

Proof. Assume to the contrary that there is a minimal set $K \subset \partial U$, which is an adding machine. Fix $p \in K$. Since the rotation number of $\hat{f}|_{b_\varepsilon(U)}$ is irrational, $\hat{f}|_{b_\varepsilon(U)}$ is semi-conjugate to an irrational rotation on the unit circle. Then we take a cross-cut γ of U such that for each cross-section D of γ , $\mathcal{E}(D)$ contains the closure of a wandering interval (if any) of $\hat{f}|_{b_\varepsilon(U)}$ and such that p is not an endpoint of γ in \overline{U} . Take a sufficiently small $\varepsilon > 0$ such that $V \equiv B(p, \varepsilon) \cap U$ is contained in a cross-section D of γ . Let D' be the cross-section of γ other than D . By Proposition 2.10, there is some positive integer n such that

$$(3.1) \quad f^{kn}(p) \in B(p, \varepsilon)$$




FIGURE 1.




FIGURE 2.

for all $k \geq 0$. Take a sequence (x_i) in V such that $x_i \rightarrow p$. Passing to a subsequence if necessary, we suppose $x_i \rightarrow q \in b_{\mathcal{E}}(U) \subset \hat{U}$. So, there is some $l > 0$ such that

$$(3.2) \quad \hat{f}^{ln}(q) \in \mathcal{E}(D').$$

Then for sufficiently large i , by equations (3.1) and (3.2), and by the continuity, we have both $f^{ln}(x_i) \in V \subset D$ and $f^{ln}(x_i) = \hat{f}^{ln}(x_i) \in D'$ (see Figure 1 and Figure 2). This is a contradiction. \square

4. EXISTENCE OF AN f -INVARIANT CIRCLE IN THE BOUNDARY OF AN OPEN DISK

Proposition 4.1. *Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a nonwandering homeomorphism and let U be an f -invariant open disk. If $f|_{\partial U} : \partial U \rightarrow \partial U$ is distal and f has no periodic points except for an only fixed point $O \in U$, then there is an f -invariant circle C in ∂U such that $(C, f|_C)$ is minimal and O belongs to the bounded component of $\mathbb{R}^2 \setminus C$.*

Proof. Fix a minimal set M in ∂U . Then $\dim(M) \leq 1$. Noting that f is nonwandering and has no periodic point in ∂U , by Theorem 3.2, we have the prime end rotation number of $f|_{\overline{U}}$ is irrational. Then it follows from Corollary 2.8 and Proposition 3.3 that M is not an adding machine. So $\dim(M) = 1$. Thus M has a component K with $\dim(K) = 1$.

Now we discuss into several cases:

Case 1. There is some $n \geq 1$ such that $f^n(K) = K$ and $f^i(K) \cap K = \emptyset$ for $1 \leq i < n$. Clearly, (K, f^n) is minimal. Then, from Proposition 2.13, it is equicontinuous. By Corollary 2.7 and Proposition 2.9, K is a circle.

Subcase 1.1. $n = 1$. Let $C = K$ and let D be the bounded component of $\mathbb{R}^2 \setminus C$. Since f has no periodic points in C , so by Brouwer's fixed point theorem, $O \in D$. Thus C satisfies the requirement.

Subcase 1.2. $n > 1$. Let $C_i = f^i(K)$, $i = 0, \dots, n-1$. Then C_i are pairwise disjoint. Let D_i be the bounded component of $\mathbb{R}^2 \setminus C_i$. If there are $i \neq j$ such that $C_i \subset D_j$, then $f^{i-j}(\overline{D_j}) \subset D_j$. This contradicts the assumption that f is nonwandering. So these D_i are pairwise disjoint. Since each D_i contains a fixed point of f^n by Brouwer's fixed point theorem, this contradicts the assumption that O is the only periodic points of f . So this subcase does not occur.

Case 2. $f^i(K)$, $i \in \mathbb{Z}$, are pairwise disjoint. Write $K_i = f^i(K)$. If K separates the plane, then $\mathbb{R}^2 \setminus K$ has a bounded component, so is each K_i . Similar to the arguments in Subcase 1.2, we have that for any $i \neq j$, K_i is contained in the unbounded component of $\mathbb{R}^2 \setminus K_j$. Thus any bounded component of $\mathbb{R}^2 \setminus K$ is a wandering open set of f . This is a contradiction.

From the above discussions, we get the following claim.

Claim A. Either the conclusion of Proposition 4.1 holds, or M has infinitely many components and any nondegenerate component of M does not separate the plane.

If the conclusion of Proposition 4.1 does not hold, then by Claim A together with Proposition 2.3 and Theorem 2.4, we get a factor $g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ of f by identifying each component of M to a point. Let $\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the factor map. Then $\pi(U)$ is a g -invariant open disk and g is a nonwandering homeomorphism and has no periodic point in $\partial\pi(U)$. So, by Theorem 3.2, the prime ends rotation number of $g|_{\pi(\overline{U})}$ is irrational. Noting that $g|_{\partial\pi(U)}$ is still distal and $\pi(M)$ is totally disconnected and infinite, by Corollary 2.8, we see that $(\pi(M), g)$ is an adding machine contained in the boundary of $\pi(U)$. This contradicts Proposition 3.3.

All together, we complete the proof. \square

5. EXISTENCE OF AN INTERMEDIATE f -INVARIANT CIRCLE

Lemma 5.1. *Let X be a compact metric space and let $f : X \rightarrow X$ be a distal homeomorphism. If K is an f -invariant proper closed subset of X , then there are $\delta > 0$ and a nonempty open subset U of X such that $f^i(U) \cap B(K, \delta) = \emptyset$ for each $i \in \mathbb{Z}$.*

Proof. For each positive integer n , let $V_n = \{x \in X : f^i(x) \in B(K, \frac{1}{n}) \text{ for some } i \in \mathbb{Z}\}$. If the conclusion of Lemma 5.1 does not hold, then V_n is a dense open subset of X for each n . Thus by Baire's Theorem, $G \equiv \bigcap_{n=0}^{\infty} V_n$ is a dense G_δ -set. Take $x \in G \setminus K$. Then $\overline{O(x, f)} \cap K \neq \emptyset$. This contradicts the minimality of $\overline{O(x, f)}$. \square

For any two circles C, C' in the plane \mathbb{R}^2 , write $C \prec C'$ if C is contained in the bounded component of $\mathbb{R}^2 \setminus C'$.

Proposition 5.2. *Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be an orientation preserving nonwandering homeomorphism. Let A be an f -invariant annulus with two boundary circles $C_1 \prec C_2$. Suppose $f|_A$ is distal and f has no periodic points except for an only fixed point O in the bounded component of $\mathbb{R}^2 \setminus C_1$. Then there is an f -invariant circle C with $C_1 \prec C \prec C_2$.*

Proof. By Lemma 5.1, we can take $\delta > 0$ and a nonempty open set U in A such that $f^i(U) \cap B(C_1 \cup C_2, \delta) = \emptyset$ for each $i \in \mathbb{Z}$. Set $W = \bigcup_{i \in \mathbb{Z}} f^i(U)$. Let K be the unbounded component of $\mathbb{R}^2 \setminus W$. Then K is f -invariant and $C_2 \subset \overset{\circ}{K}$. Let V be a component of $\mathbb{R}^2 \setminus K$. Then V is an open disk by a direct application of Jordan separation theorem. Since f is nonwandering, there is some $n \geq 0$ with $f^n(V) = V$. Then, by Theorem 2.1, there is a periodic point of f^n in V , so is for f . Thus by the assumption, we have $O \in V$. This implies $C_1 \subset V$. From the above discussions, we see that V is the only component of $\mathbb{R}^2 \setminus K$, and hence $f(V) = V$. Now applying Proposition 4.1, we get the required circle. \square

6. A DECOMPOSITION OF THE ANNULUS INTO f -INVARIANT CIRCLES

In this section, we will complete the proof of the main Theorem. All assumptions are as in Theorem 1.1. WLOG, we may suppose the annulus $A = \{z \in \mathbb{C} : 1 \leq |z| \leq 2\}$. Then we extend $g : A \rightarrow A$ to $f : \mathbb{C} \rightarrow \mathbb{C}$ by defining

$$f(x) = \begin{cases} \frac{|x|}{2} \cdot g\left(\frac{2x}{|x|}\right), & |x| > 2; \\ g(x), & 1 \leq |x| \leq 2; \\ |x| \cdot g\left(\frac{x}{|x|}\right), & 0 < |x| < 1; \\ 0, & |x| = 0. \end{cases}$$

It is clear from the definition that f is an orientation preserving nonwandering homeomorphism on the plane and has no periodic points except for the only fixed point 0.

Write $C_1 = \{z : |z| = 1\}$ and $C_2 = \{z : |z| = 2\}$. For each circle C in the plane, we use $D(C)$ and $OD(C)$ to denote the bounded component and unbounded component of $\mathbb{R}^2 \setminus C$, respectively. Let \prec be the transitive order defined in Section 5; that is, for circles C and C' in the plane, $C \prec C'$ iff $C \subset D(C')$. Let

$$\mathcal{C} = \{C : C \text{ is an } f\text{-invariant circle and } C_1 \prec C \prec C_2\} \cup \{C_1, C_2\}.$$

Let \mathcal{T} be the family of all chains of \mathcal{C} respect to \prec . Then \mathcal{T} is a partial set with respect to the inclusion relation on the power set of \mathbb{R}^2 . Using Zorn's lemma, there is a maximal chain \mathcal{P} in \mathcal{T} .

Claim A. \mathcal{P} is a partition of A .

Proof of Claim A. Assume to the contrary that there is some $v \in A \setminus \cup \mathcal{P}$. Since $C_1, C_2 \in \mathcal{P}$ by the maximality of \mathcal{P} , we have $v \in \overset{\circ}{A}$. Set $\mathcal{P}_1 = \{C \in \mathcal{P} : v \notin D(C)\}$ and $\mathcal{P}_2 = \{C \in \mathcal{P} : v \in D(C)\}$. Then $C_1 \in \mathcal{P}_1$ and $C_2 \in \mathcal{P}_2$.

Now we discuss into several cases.

Case 1. \mathcal{P}_1 has no maximal element. Let $U = \cup_{C \in \mathcal{P}_1} D(C)$. Then U is an f -invariant open disk. From Proposition 4.1, we have an f -invariant circle C_3 in ∂U with $0 \in D(C_3)$. Clearly, $C_3 \notin \mathcal{P}$ and $C \prec C_3$ for any $C \in \mathcal{P}_1$. If $C_3 \prec C$ for any $C \in \mathcal{P}_2$, then $\{C_3\} \cup \mathcal{P}$ is a chain, which contradicts the maximality of \mathcal{P} . So, there must exist a $C_4 \in \mathcal{P}_2$ such that $C_3 \cap C_4 \neq \emptyset$. Let V be a component of $D(C_4) \setminus \overline{D(C_3)}$. Then V is a component of $\mathbb{R}^2 \setminus (C_3 \cup C_4)$, and hence it is an open disk. Noting that f is nonwandering, we have




FIGURE 3.

$f^n(V) = V$ for some $n \geq 0$. Then by Theorem 2.1, f has a periodic point in V . This is a contradiction. So, Case 1 does not happen.

Case 2. \mathcal{P}_2 has no minimal element. We consider the Riemann sphere $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$. Let $\mathbb{L} = \hat{\mathbb{C}} \setminus \{0\}$. Then \mathbb{L} is a plane. Define $\hat{f} : \mathbb{L} \rightarrow \mathbb{L}$ by letting $\hat{f}(\infty) = \infty$ and $\hat{f}(x) = f(x)$ for any $x \in \mathbb{C} \setminus \{0\}$. Then \hat{f} is an orientation preserving nonwandering homeomorphism on the plane \mathbb{L} and has no periodic points except for the only fixed point ∞ . Similar to the discussions in Case 1, we see that Case 2 does not happen.

Case 3. \mathcal{P}_1 has the maximal element C_5 and \mathcal{P}_2 has the minimal element C_6 . Then $C_5 \prec C_6$. Applying Proposition 5.2, we get an f -invariant circle C_7 such that $C_5 \prec C_7 \prec C_6$. Then $\{C_7\} \cup \mathcal{P}$ is a chain. This contradicts the maximality of \mathcal{P} . Thus Case 3 does not occur.

So $A = \cup \mathcal{P}$; that is \mathcal{P} is a partition of A . □

Claim B. \prec is a dense and complete linear order on \mathcal{P} .

Proof of Claim B. (1) Linearity. For any distinct $C, C' \in \mathcal{P}$, we have $C \cap C' = \emptyset$ and hence either $C \subset D(C')$ or $C' \subset D(C)$. This shows that either $C \prec C'$ or $C' \prec C$. Thus \prec is a linear order.

(2) Density. Let $C, C' \in \mathcal{P}$ be with $C \prec C'$. Then we have $\overline{D(C)} \subsetneq D(C')$. Now for any $x \in D(C') \setminus \overline{D(C)}$, there is some $C'' \in \mathcal{P}$ such that $x \in C''$. It follows from the definition of \prec that $C \prec C'' \prec C'$. This shows that \prec is a dense order.

(3) Completeness. To the contrary, assume that \prec is incomplete. That is there is a Dedekind Gap, which means that there is a partition $\mathcal{P} = \mathcal{L} \cup \mathcal{U}$ such that

- For any $C \in \mathcal{L}$ and $C' \in \mathcal{U}$, $C \prec C'$,
- \mathcal{L} has a maximal element C^* and \mathcal{U} has a minimal element C_* .

Then we have

$$\bigcup_{C \in \mathcal{L}} C = \overline{D(C^*)} \setminus D(C_1) \text{ and } \bigcup_{C \in \mathcal{U}} C = \overline{OD(C_*)} \setminus OD(C_2),$$

both of which are closed in A . But this contradicts the connectedness of A . This shows that \prec is complete. \square

Now Claim B implies that (\mathcal{P}, \prec) endowed with the ordering topology is homeomorphism to a closed interval. WLOG, we may assume that $(\mathcal{P}, \prec) \cong [1, 2]$ and use C_r to denote elements of \mathcal{P} with $r \in [1, 2]$.

Recall that 2^A is the hyperspace of A endowed with the Hausdorff metric.

Claim C. \mathcal{P} is closed in 2^A . Specially, \mathcal{P} is a continuous decomposition.

Proof of Claim C. Let C_{r_n} be a sequence in \mathcal{P} that converges to K in 2^A under Hausdorff metric. Since $(\mathcal{P}, \prec) \cong [1, 2]$, by passing to some subsequence, we may assume that $C_{r_n} \xrightarrow{\prec} C_r$ under the ordering topology for some $r \in [1, 2]$.

Next we will show that $K = C_r$ which implies that \mathcal{P} is closed in 2^A . We may assume as well $C_{r_n} \prec C_r$ for each n . Fix an $x \in K$. Then it follows from the definition of Hausdorff metric that there is a sequence (x_n) with $x_n \in C_{r_n}$ such that $x_n \rightarrow x$. Since $x_n \in C_{r_n} \subset D(C_r)$, we have $x \in \overline{D(C_r)}$. To show $x \in C_r$, we assume that $x \in D(C_r)$. Then there is some $s \in [1, r)$ such that $x \in C_s$. Since $r_n \rightarrow r$, we have $s < r_n \leq r$ and hence $C_s \prec C_{r_n} \preceq C_r$ for any sufficiently large n . But in this case, x_n cannot converge to x ; this is a contradiction. To sum up, we have $x \in C_r$. Hence $K \subset C_r$ as x is arbitrary. On the other hand, for any $y \in C_r$, there is some subsequence (y_{n_i}) with $y_{n_i} \in C_{r_{n_i}}$ such that $y_{n_i} \rightarrow y$. Indeed, take any point $z \in C_1$ and let L be the segment connecting z and y in A . Then $L \cap C_{r_n} \neq \emptyset$ for each n and we choose some $y_n \in L \cap C_{r_n}$. It is clear that y is a limit point of (y_n) . The above arguments show that $K = C_r$ and the closedness of \mathcal{P} in 2^A is followed. \square

For any orientation preserving homeomorphism ϕ on a circle, we use $\rho(\phi)$ to denote the rotation number of ϕ .

Claim D. The rotation numbers of $f|_C$, $C \in \mathcal{P}$, are the same irrational number.

Proof of Claim D. Assume to contrary that there are $C' \neq C'' \in \mathcal{P}$ such that the rotation numbers $\rho(f|_{C'}) \neq \rho(f|_{C''})$. Let \tilde{A} be the annulus in A with boundary $C' \cup C''$. Write $\alpha' = \rho(f|_{C'})$ and $\alpha'' = \rho(f|_{C''})$. Take a homeomorphism $h : \tilde{A} \rightarrow \mathbb{S}^1 \times [0, 1]$ such that $h(C') = \mathbb{S}^1 \times \{0\}$ and $h(C'') = \mathbb{S}^1 \times \{1\}$. Let $\tilde{f} = hf|_A h^{-1}$. Then $\rho(\tilde{f}|_{h(C')}) = \alpha'$ and $\rho(\tilde{f}|_{h(C'')}) = \alpha''$. Let $\mathbb{R} \times [0, 1]$ be the universal covering of $\mathbb{S}^1 \times [0, 1]$ and let $F : \mathbb{R} \times [0, 1] \rightarrow \mathbb{R} \times [0, 1]$ be a lift of \tilde{f} . Noting that \tilde{f} is homotopic to the identity, we have $TF = FT$, where T is the unit translation on $\mathbb{R} \times [0, 1]$ defined by $T(s, t) = (s + 1, t)$. Let $\pi : \mathbb{R} \times [0, 1] \rightarrow \mathbb{R}$ be the projection to the first coordinate, that is $\pi(s, t) = s$. For any map $\psi : \mathbb{R} \times [0, 1] \rightarrow \mathbb{R} \times [0, 1]$, write $\psi_1 = \pi\psi$. Then there are $m', m'' \in \mathbb{Z}$ such that

$$\lim_{n \rightarrow \infty} \frac{F_1^n(x, 0) - x}{n} = \alpha' + m'$$

and

$$\lim_{n \rightarrow \infty} \frac{F_1^n(x, 1) - x}{n} = \alpha'' + m''.$$

WLOG, suppose $\alpha' + m' < \alpha'' + m''$. Take a rational number $\frac{p}{q}$ with

$$\alpha' + m' < \frac{p}{q} < \alpha'' + m''.$$

Then we have

$$\lim_{n \rightarrow \infty} \frac{(T^{-p}F^q)_1^n(x, 0) - x}{n} = q(\alpha' + m') - p < 0$$

and

$$\lim_{n \rightarrow \infty} \frac{(T^{-p}F^q)_1^n(x, 1) - x}{n} = q(\alpha'' + m'') - p > 0.$$

Since f is nonwandering, it follows from [15, Theorem 3.3] that \tilde{f}^q has a fixed point in $\mathbb{S}^1 \times [0, 1]$. That is \tilde{f} has a periodic point, so is f . This is a contradiction. So the rotation numbers of $f|_C$, $C \in \mathcal{P}$, are the same, the irrationality of which clearly follows from the Poincaré's classification theorem for circle homeomorphisms. \square

All together, we complete the proof of Theorem 1.1.

REFERENCES

- [1] J. Auslander, *Minimal flows and their extensions*, in: North-Holland Mathematics Studies, Vol. 153, in: Notas de Matemática [Mathematical Notes], vol. 122, North-Holland Publishing Co., Amsterdam, 1988.
- [2] J. Auslander, E. Glasner and B. Weiss, *On recurrence in zero dimensional flows*, Forum Math. 19 (2007), 107-114.
- [3] F. Béguin, S. Crovisier, F. Le Roux, and A. Patou, *Pseudo-rotations of the closed annulus: variation on a theorem of J Kwapisz*, Nonlinearity 17 (2004) 1427-1453.
- [4] R. H. Bing, *A simple closed curve is the only homogeneous bounded plane continuum that contains an arc*, Canad. J. Math. 12 (1960), 209-230.
- [5] L. Block and J. Keesling, *A characterization of adding machine maps*, Top. Appl. 140 (2004), 151-161.
- [6] A. Blokh, *Pointwise-recurrent maps on uniquely arcwise connected locally arcwise connected spaces*, Proc. Amer. Math. Soc. 143 (2015), 3985-4000.
- [7] F. Botelho, *Rotation sets of maps of the annulus*, Pacific J. Math. 133 (1988), 251-266.
- [8] I. U. Bronstein, *Extensions of minimal transformation groups*, Sitjthoff and Noordhoff, 1979.
- [9] C. Carathéodory, *Über die Begrenzung einfach zusammenhängender Gebiete*, Math. Ann. 73 (1913), 323-370.
- [10] C. Carathéodory, *Über die gegenseitige Beziehung der Ränder bei der konformen Abbildung des Inneren einer Jordanschen Kurve auf einen Kreis*, Math. Ann. 73 (1913), 305-320.
- [11] M. L. Cartwright and J.E. Littlewood, *Some fixed point theorems*, with an appendix by H. D. Ursell, Ann. of Math. 54 (1951), 1-37.
- [12] R. Ellis, *Distal transformation Groups*, Pacific J. Math. 8 (1958) 401-405
- [13] A. Fathi, *An orbit closing proof of Brouwer's lemma on translation arcs*, L'enseignement Mathématique 33 (1987), 315-322.
- [14] N. E. Foland, *A characterization of the almost periodic homeomorphisms on the closed 2-cell*, Proc. Amer. Math. Soc. 16 (1965), 1031-1034.
- [15] J. Franks, *Generalizations of the Poincare-Birkhoff theorem*, Annals of Mathematics, 128 (1988), 139-151.
- [16] H. Furstenberg, *The Structure of distal flows*, Amer. J. Math. 85 (1963), 477-515.
- [17] M. Handel, *The rotation set of homeomorphism of the annulus is closed*, Commun. Math. Phys. 127, 339-349 (1990).
- [18] T. Hauser and T. Jäger, *Monotonicity of maximal equicontinuous factors and an application to toral flows*, Proc. Amer. Math. Soc. 147 (2019), 4539-4554.

- [19] L. C. Hoehn, L. G. Oversteegen, *A complete classification of homogeneous plane continua*, Acta Math. 216 (2016), 177-216.
- [20] B. Kolev and M. Pérouème, *Recurrent surface homeomorphisms*, Math. Proc. Cambridge Philos. Soc. 124 (1998), 161-168.
- [21] A. Koropecski, P. Le Calvez, and M. Nassiri, *Prime ends rotation numbers and periodic points*, Duke Math. J. 164 (2015), 403-472.
- [22] K. Kuratowski, *Topology*, Vol. II, Academic Press, New York, N.Y., 1968.
- [23] P. Le Calvez, *From Brouwer theory to the study of homeomorphisms of surfaces*, European Mathematical Society (EMS), Zürich, 2006, 77-98.
- [24] J. H. Mai, *Pointwise-recurrent graph maps*, Ergodic Theory Dynam. Systems 25 (2005), 629- 637.
- [25] J. H. Mai and X. D. Ye, *The structure of pointwise recurrent maps having the pseudo orbit tracing property*, Nagoya Math. J. 166 (2002), 83-92.
- [26] R. Mañé, *Expansive homeomorphisms and topological dimension*, Trans. Amer. Math. Soc. 252 (1979), 313-319.
- [27] M. Misiurewicz and K. Ziemian, *Rotation sets for maps of tori*, J. London Math. Soc. (2) 40, 490-506 (1989).
- [28] D. Montgomery, *Pointwise periodic homeomorphisms*, Amer. J. Math. 59 (1937), 118-120.
- [29] D. Montgomery and L. Zippin, *Topological transformation groups*, Interscience Publishers, New York-London, 1955.
- [30] S. Nadler, *Continuum theory-An introduction*, Monographs and Textbooks in Pure and Applied Mathematics, 158. Marcel Dekker, Inc., New York, 1992.
- [31] I. Naghmouchi, *Pointwise recurrent dendrite maps*, Ergod. Theory Dyn. Syst. 33 (2013), 1115-1123.
- [32] L. G. Oversteegen and E. D. Tymchatyn, *Recurrent homeomorphisms on \mathbb{R}^2 are periodic*, Proc. Amer. Math. Soc. 110 (1990), 1083-1088.
- [33] M. Rees, *On the structure of minimal distal transformation groups with manifolds as phase spaces*, thesis, University of Warwick, Coventry, England, 1977.
- [34] G. X. Ritter, *A characterization of almost periodic homeomorphisms on the 2-sphere and the annulus*, General Topology Appl. 9 (1978), 185-191.
- [35] E. H. Shi, H. Xu, and Z. Q. Yu, *The structure of pointwise recurrent expansive homeomorphisms*, Ergodic Theory Dynam. Systems 43 (2023), 3448-3459.
- [36] P. Walters, *An introduction to ergodic theory*, GTM 79, Springer-Verlag, 1982.

SCHOOL OF MATHEMATICS AND SCIENCES, SOOCHOW UNIVERSITY, SUZHOU, JIANGSU 215006,
CHINA

Email address: ehshi@suda.edu.cn

DEPARTMENT OF MATHEMATICS, SHANGHAI NORMAL UNIVERSITY, SHANGHAI 200234, CHINA

Email address: huixu@shnu.edu.cn

SCHOOL OF MATHEMATICS AND SCIENCES, SOOCHOW UNIVERSITY, SUZHOU, JIANGSU 215006,
CHINA

Email address: 20204207013@stu.suda.edu.cn