

EXACT SOLUTION OF GENERALIZED TRIPLE ISING CHAINS WITH MULTI-SPIN INTERACTIONS

PAVEL KHRAPOV AND NIKITA VOLKOV

ABSTRACT. In this paper, we obtain the exact physical characteristics of the triple-chain Ising model on a torus with all possible multispin interactions invariant with respect to rotation by the angle $2\pi/3$. The exact value of the partition function in a finite cyclically closed strip of length L , as well as the free energy, internal energy, specific heat, magnetization, susceptibility, and entropy in the thermodynamic limit at $L \rightarrow \infty$ are found by the transfer-matrix method for the model. The spectrum of the transfer-matrix and the structure of its eigenvectors are found. For two special cases — for the model with multispin interactions of even number of spins and for the model with some interactions of two, three, four and six spins, simplified expressions of the mentioned physical characteristics are obtained; in the thermodynamic limit they are expressed through the logarithm of the root of the quadratic equation. For the model with multispin interactions of an even number of spins, a kind of pair correlations in the thermodynamic limit is found, and it is shown that the magnetization at zero magnetic field is equal to zero; the structure of the ground states of the system is found and examples of their projections of seven-dimensional space onto three-dimensional space and examples of configurations corresponding to these ground states are given. The correlation length is shown and its graphs are given. For percolation invariant with respect to rotation about the central axis of rotation through the angle $2\pi/3$, a limit relation for the non-percolation probability is derived. As special cases, we consider the planar triangular model with all possible interactions, including, perhaps, different triple interactions inside neighboring triangles, and the planar model with nearest-neighbor, next nearest-neighbor, and plaquette interactions. For them the main exact physical characteristics have been found. This allowed us to obtain them for the planar gonihedric model as well. Graphs of physical characteristics illustrating the obtained results are constructed.

CONTENTS

1. Introduction	2
2. Model description and main result	6
3. The main special case: a model with multispin interactions of even number of spins	11
3.1. Model description and results	11
3.2. Pair correlations of the model with multispin interactions of even number of spins	12
3.3. Ground states and correlation length of the model with multispin interactions of even number of spins	16

2010 *Mathematics Subject Classification.* 82B20, 82B23.

Key words and phrases. Chain Ising model, transfer-matrix, partition function, free energy, specific heat, internal energy, magnetization, susceptibility, entropy, thermodynamic limit, pair correlations, correlation length, ground states, triangular model, planar gonihedric model.

4. Model with some interactions of two, three, four and six spins	25
5. Exact value of invariant percolation in the triple-chain generalized Ising model with all possible multispin interactions	27
6. Planar model with nearest neighbor, next-nearest neighbor and plaquette interactions	29
6.1. Planar gonihedric model	33
7. Planar triangular model	35
8. Examples of physical characteristics in the thermodynamic limit	39
9. Proofs of theorems	43
9.1. Proof of the theorems 2.1, 2.2	43
9.2. Proof of theorems 3.1, 3.2	45
9.3. Proof of theorems 4.1, 4.2	46
9.4. Proof of theorems 5.1, 5.2	47
10. Conclusion	48
11. Appendix	49
References	51

1. INTRODUCTION

The Ising model, proposed by Lenz and introduced in 1925 by Ising [1] to describe magnetic transitions, is still the subject of intensive research because it allows us to study the characteristics of magnetic systems and a number of different related physical phenomena, such as frustration.

The one-dimensional model was solved by Ising himself, and the exact value of the partition function and free energy for the Ising model with nearest-neighbor interaction without an external magnetic field on a $n \times \infty$ cylinder was first obtained by Onsager [2].

Various numerical and analytical solution methods are applied to investigate such models. Among numerical methods, variations of the Monte Carlo method or the Metropolis [10] algorithm predominate, while the transfer-matrix method, originally proposed in [3], [4], and the combinatorial method [5], are used for the exact analytical solution.

In [13] the exact value of the free energy for the model with one central and three side spins is found, the Hamiltonian of which includes the interactions of the central spin of a layer with the central spins on the nearest neighboring layers, with the side spins of this layer, and also the interactions of the side spins with the nearest side spins of the neighboring layers.

In [6] for the triple-chain Ising model with interactions of nearest and next-nearest neighbors without an external field, the exact solution of the secular transfer-matrix equation is obtained and all its eigenvalues and the form of eigenvectors are obtained; formulas for specific heat, parallel and perpendicular susceptibilities are derived; the behavior of these values as a function of temperature for various choices of the model parameters is considered.

In [7], a triple-chain model with different interactions of nearest neighbors and next-nearest neighbors is studied by the transfer-matrix method, for which all eigenvalues and eigenvectors of the transfer-matrix are found, the structure of ground states is shown, and expressions for pair correlations of spins in the thermodynamic

limit are obtained. These expressions are calculated using the spin-layer matrices, the structure of which is constructed by analogy with the two-dimensional [8] and one-dimensional [9] cases.

Besides triple-chain model, other models in three-dimensional space, such as those with cubic lattices, and various planar and linear models are also being actively investigated.

Thus, for example, in [14] the exact solution of two Ising models with a cyclic-closed lattice $2 \times 2 \times \infty$, the first of which has completely anisotropic interactions, and the second consists of two different types of linear chains including non-intersecting diagonal interactions on the outer faces of the parallelepiped, is obtained using the transfer-matrix method, and the behavior of specific heat is analyzed, and in [15] by the transfer-matrix method, taking into account its symmetry, the eigenvalues and the structure of eigenvectors for the three-dimensional cubic model $n = m = 2$ with nearest-neighbor interactions are obtained, among which the interactions between spins located vertically in one layer, horizontally in one layer, and all interactions between neighboring layers are equal.

In [33] the generalized Ising model in the $2 \times 2 \times \infty$ strip with a Hamiltonian invariant with respect to the central axis of rotation through the angle $\pi/2$, which includes all possible multiplicative interactions of an even number of spins in the unit cube is considered. The exact value of free energy and specific heat in the thermodynamic limit is found.

A separate chapter is devoted to the gonihedric model in the $2 \times 2 \times \infty$ strip, the exact values of the free energy and specific heat in the thermodynamic limit are found for the case of free boundary conditions and an analogue of cyclically closed boundary conditions in both directions perpendicular to the central axis of rotation.

In [17] for a cubic lattice with nearest-neighbor and next-nearest-neighbor interactions, a detailed study of phase diagrams is carried out.

In [32] we obtain formulas for finding the free energy in the thermodynamic limit on the set of exact disordered solutions for the two-dimensional generalized Ising model in an external field with the interaction of nearest neighbors, next-nearest neighbors, all possible triple interactions and the interaction of four spins for the planar model, and for the three-dimensional generalized model in an external field with all possible interactions in the tetrahedron.

In [20] the spectra of stochastic operators are investigated.

The technique of cluster decompositions [21] has been very influential in the study of lattice models.

In [24] cluster properties and bound states of the transfer matrix of the Yang-Mills model with a compact gauge group are shown, and in [27] cluster expansion and spectrum of the transfer matrix of the two-dimensional ising model with strong external field are shown.

In addition to classical ones, alternative methods for solving these models are being developed. For example, in [30] on a quantum computer emulator, in particular, a triple-chain model with a Hamiltonian invariant with respect to rotation by an angle $\frac{2\pi}{3}$ and taking into account pair and plaquette interactions and interaction with the external field is investigated, - a method of searching with the help of a variational quantum algorithm for the free energy and magnetization in the thermodynamic limit for one-, double- and triple-chain Ising models, and with the help

of special parameterization of the state of the qubit system and transfer-matrix decomposition, the free energy and magnetization are calculated for the triple-chain model, and in [12] the study of the cubic lattice model with nearest-neighbor, next-nearest-neighbor, and plaquette interactions by the method of cluster variation is carried out. In [31] the Fourier transform of the elementary transfer-matrix of the generalized two-dimensional Ising model with special boundary conditions with a helical-type shift and a type of Hamiltonian covering the generalized Ising model with a multispin interaction, as well as models equivalent to models on a triangular lattice with a checkerboard-type Hamiltonian, is performed.

In [32] we obtain formulas for finding the free energy in the thermodynamic limit on the set of exact disordered solutions for the two-dimensional generalized Ising model in an external field with the interaction of nearest neighbors, next-nearest neighbors, all possible triple interactions and the interaction of four spins for the planar model, and for the three-dimensional generalized model in an external field with all possible interactions in the tetrahedron.

A meaningful review and results on percolation theory can be found in [23].

In [34] percolation in the Ising model is studied, based on the generalized results of [35]. Percolation in a strip of finite width for an independent field and the Ising model was studied in [36], for continuous models in [22].

In recent years, interest in simple systems of Ising spins has arisen as a result of considering the random surface model in the context of string theory [25],[26]. This model was called gonihedric and its three-dimensional case was studied by approximation methods.

In this paper, for the generalized triple-chain Ising model on the torus with a Hamiltonian with all possible multispin interactions invariant with respect to rotation by an angle $\frac{2\pi}{3}$, we obtain the exact values of the partition function in a finite closed strip of length L , as well as the free energy, internal energy, specific heat, magnetization, susceptibility, and entropy for a closed chain of finite length L and in the thermodynamic limit at $L \rightarrow \infty$.

For a model with multispin interactions of an even number of spins, the form of pair correlations in the thermodynamic limit is found, and it is shown that the magnetization at zero external field is equal to zero; the structure of the ground states of the system is found. A formula for the correlation length is derived. The obtained results are in agreement with the results of the paper [7], in which the interaction Hamiltonian containing nearest-neighbor and next-nearest neighbor interactions is considered.

Limit relations are obtained for percolation invariant with respect to rotation by an angle $\frac{2\pi}{3}$.

As special cases, exact physical characteristics are obtained for the planar triangular model with all possible interactions, including various triple interactions inside neighboring triangles, and for the planar model with nearest-neighbor, next-nearestneighbor and cladding interactions. For specific parameters of interactions this gives exact physical characterizations for the planar gonihedric model.

Next, let us describe the structure of the paper.

In paragraph 2 we describe the model and its physical characteristics, introduce the set of elementary generating carriers of the Hamiltonian of the system and the Hamiltonian itself, formulate two theorems for the general case of the model with all possible multispin interactions in the thermodynamic limit and in the strip of

finite length, which were proved: exact analytical expressions for the physical characteristics in the thermodynamic limit are given and the expression for the partition function in the strip of finite length and all eigenvalues of the transfer-matrix are written out.

The third paragraph presents the first special case — a triple-chain model with multispin interactions of even number of spins, for which two theorems similar to the theorems of the second paragraph are formulated and all eigenvalues of its transfer-matrix are written out.

For this model, exact analytical expressions of pair correlations in the thermodynamic limit are obtained using the spin-layer matrices and the transfer-matrix, for which the transition matrix to the diagonal form is written out. With its help it is shown that the magnetization of this model in the absence of an external field is zero, which coincides with the results of [19]. The general expressions for pair correlations for the model with multispin interactions of an even number of spins coincide with the expressions for pair correlations of the triple-chain model with nearest- and next-nearest neighbor interactions considered in [7].

Besides, for this case we describe the structure of ground states [16], give examples of corresponding model configurations and two example-illustrations explicitly showing the picture of ground states in two partial projections of the seven-dimensional space of ground states onto the three-dimensional one, and show with their help the possibility of special behavior of the correlation length near the boundary points of the regions of ground states.

In paragraph 4 we present the second special case — a model with some interactions of two, three, four and six spins. For it two theorems similar to the theorems of the second paragraph are formulated and all eigenvalues are written out, two of which have multiplicity 1 and two have multiplicity 3.

In the fifth paragraph, we formulate two theorems on the non-percolation probability, invariant with respect to rotation by an angle $\frac{2\pi}{3}$, in a closed strip of finite length L , and in the thermodynamic limit.

In the sixth paragraph we consider a model with nearest neighbor, next-nearest neighbor and plaquette interactions. This is an important special case of the more general model of paragraph 3, and all the results of the third paragraph are true for it. The results for the gonihedric model are formulated separately.

In the seventh paragraph we consider the planar triangular model, the validity of the theorems 2.1 and 2.2 is shown for it.

In the eighth paragraph we consider an example illustrating all physical characteristics given in the main theorem of the second paragraph in the thermodynamic limit for the general case of the model with all possible multispin interactions at two variable parameters corresponding to the next-nearest neighbor interactions and equal to each other, and a variable temperature — three-dimensional plots of these quantities and cross sections of some of them at fixed temperatures are shown.

In paragraph 9 we prove six theorems (1 - 4, 6, 7), formulated in paragraphs 2 — 4: with the help of the theorem on preservation of eigenspaces by commuting matrices [29] we obtained the structures of eigenvectors of transfer-matrices for each case, due to which the problem of finding eigenvalues of initial transfer-matrices of size 8×8 was reduced to the search of roots of characteristic equations of matrices of smaller dimensions — for the general case of the model with all possible multispin interactions, 4 eigenvalues, including the largest one, were found as roots

of the equation of the fourth degree, the solution of which was obtained by the Ferrari method, and the other four as roots of two quadratic trinomials. For the first special case, four eigenvalues, including the largest one, are found as roots of two square trinomials, and the remaining four are found as roots of four linear equations. In the second special case, all eigenvalues are found as roots of two quadratic trinomials — two roots of multiplicity 1, and two roots of multiplicity 3.

In the appendix, we give formulas for the Ferrari method of solving the fourth degree equation, and a scheme for computing the partial derivatives on the external field H of the coefficients of the equation (2.21).

2. MODEL DESCRIPTION AND MAIN RESULT




FIGURE 1. Triple-chain lattice of the Ising model

Consider a three-dimensional triple-chain cyclically closed lattice model of size $3 \times L$ (Fig. 1) with a total number of nodes in the lattice $N = 3L$, where L is the number of layers in the model:

$$\mathcal{L}_L = \{t_i^m; m = 0, 1, \dots, L-1; t_i^L = t_i^0; i = 0, 1, 2\} \quad (2.1)$$

Let us consider that in each node t_i^m there is a particle, the state of which is determined by the value of spin $\sigma_i^m \equiv \sigma_{t_i^m}$, $m = 0, 1, \dots, L-1; i = 0, 1, 2$.

Let $\Omega_P^m = \{t_i^m, t_i^{m+1}, i = 0, 1, 2\}$ be a triangular prism (Fig. 1). Let us introduce

the set of elementary generating carriers of the Hamiltonian:

$$\begin{aligned}
 \Phi_1^m &= \{\{t_0^m\}, \{t_0^{m+1}\}\}, \\
 \Phi_2^m &= \{\{t_0^m, t_1^m\}, \{t_0^m, t_0^{m+1}\}, \{t_0^m, t_1^{m+1}\}, \{t_0^m, t_2^{m+1}\}, \{t_0^{m+1}, t_1^{m+1}\}\}, \\
 \Phi_3^m &= \{\{t_0^m, t_1^m, t_2^m\}, \{t_0^m, t_1^m, t_0^{m+1}\}, \{t_0^m, t_1^m, t_1^{m+1}\}, \{t_0^m, t_1^m, t_2^{m+1}\}, \{t_0^m, t_0^{m+1}, t_2^{m+1}\}, \\
 &\quad \{t_0^m, t_0^{m+1}, t_1^{m+1}\}, \{t_0^m, t_1^{m+1}, t_2^{m+1}\}, \{t_0^{m+1}, t_1^{m+1}, t_2^{m+1}\}\}, \\
 \Phi_4^m &= \{\{t_0^m, t_1^m, t_2^m, t_0^{m+1}\}, \{t_0^m, t_1^m, t_0^{m+1}, t_1^{m+1}\}, \{t_0^m, t_1^m, t_0^{m+1}, t_2^{m+1}\}, \\
 &\quad \{t_0^m, t_1^m, t_1^{m+1}, t_2^{m+1}\}, \{t_0^m, t_0^{m+1}, t_1^{m+1}, t_2^{m+1}\}\}, \\
 \Phi_5^m &= \{\{t_0^m, t_1^m, t_2^m, t_1^{m+1}, t_2^{m+1}\}, \{t_1^m, t_2^m, t_0^{m+1}, t_1^{m+1}, t_2^{m+1}\}\}, \\
 \Phi_6^m &= \{\{t_0^m, t_1^m, t_2^m, t_0^{m+1}, t_1^{m+1}, t_2^{m+1}\}\}.
 \end{aligned} \tag{2.2}$$

Let us introduce an elementary generating Hamiltonian for all interactions:

$$\hat{H}^m = \hat{H}_1^m + \hat{H}_2^m + \hat{H}_3^m + \hat{H}_4^m + \hat{H}_5^m + \hat{H}_6^m, \tag{2.3}$$

for which

$$\begin{aligned}
 \hat{H}_r^m &= - \sum_{\alpha_m \in \Phi_r^m} J_{\alpha_m} \sigma_{\alpha_m}, \\
 r &= 1, 2, \dots, 6
 \end{aligned}$$

— components of the Hamiltonian responsible for the interaction of r spins in the Ω_P^m prism, J_{α_m} are the corresponding coefficients of interspin interactions.

For the function $\varphi(\sigma_0^m, \sigma_1^m, \sigma_2^m, \sigma_0^{m+1}, \sigma_1^{m+1}, \sigma_2^{m+1})$ that depends on σ_t , $t \in \Omega_P^m$, we introduce the "rotation" operator $R_{\frac{2\pi}{3}}^m$ by the angle $\frac{2\pi}{3}$:

$$R_{\frac{2\pi}{3}}^m(\varphi(\sigma_0^m, \sigma_1^m, \sigma_2^m, \sigma_0^{m+1}, \sigma_1^{m+1}, \sigma_2^{m+1})) = \varphi(\sigma_1^m, \sigma_2^m, \sigma_0^m, \sigma_1^{m+1}, \sigma_2^{m+1}, \sigma_0^{m+1}), \tag{2.4}$$

then the Hamiltonian of the model has the form

$$\mathcal{H}_L(\sigma) = \sum_{m=0}^{L-1} H^m, \tag{2.5}$$

where

$$H^m = \hat{H}^m + R_{\frac{2\pi}{3}}^m \hat{H}^m + (R_{\frac{2\pi}{3}}^m)^2 \hat{H}^m, \tag{2.6}$$

that is, all carriers of the Hamiltonian in the prism Ω_P^m are generated by elementary generating carriers (2.2) at the rotation of the model by the angle $\frac{2\pi}{3}$ and the angle $\frac{4\pi}{3}$.

In this formulation of the Hamiltonian for the interaction $J_{\{t_0^m, t_1^m, t_2^m\}}$ and for \hat{H}_6^m one must take into account the $\frac{1}{3}$ multiplier compensating the rotations.

In addition, when going from the prism Ω_P^m to the prism Ω_P^{m+1} , the elementary generating carriers $\{t_0^m\}, \{t_0^m, t_1^m\}, \{t_0^m, t_1^m, t_2^m\}$ in the prism Ω_P^m generate, respectively, elementary generating carriers $\{t_0^{m+1}\}, \{t_0^{m+1}, t_1^{m+1}\}, \{t_0^{m+1}, t_1^{m+1}, t_2^{m+1}\}$, so that the latter can be omitted explicitly in (2.2), but we include them to emphasize the following clearly:

$$\begin{aligned}
 \Phi_4^m &= \{\Omega_P^m \setminus \{t_0^m, t_1^m\}, \Omega_P^m \setminus \{t_0^m, t_0^{m+1}\}, \Omega_P^m \setminus \{t_0^m, t_1^{m+1}\}, \\
 &\quad \Omega_P^m \setminus \{t_0^m, t_2^{m+1}\}, \Omega_P^m \setminus \{t_0^{m+1}, t_1^{m+1}\}\}, \\
 \Phi_5^m &= \{\Omega_P^m \setminus \{t_0^m\}, \Omega_P^m \setminus \{t_0^{m+1}\}\}.
 \end{aligned} \tag{2.7}$$

Note also that this transition must be taken into account when composing the Hamiltonian of the model, indicating in front of its components, containing the interactions $\{t_0^m\}$, $\{t_0^m, t_1^m\}$, $\{t_0^m, t_1^m, t_2^m\}$, $\{t_0^{m+1}\}$, $\{t_0^{m+1}, t_1^{m+1}\}$, $\{t_0^{m+1}, t_1^{m+1}, t_2^{m+1}\}$ and generated by them, the multiplier $\frac{1}{2}$.

The partition function of the model will be written in the form:

$$Z_L = \sum_{\sigma} \exp \left\{ - \frac{\mathcal{H}_L(\sigma)}{k_B T} \right\}, \quad (2.8)$$

where k_B is the Boltzmann constant (for simplification in further calculations we take $k_B = 1$), and the summation is taken over all configurations of spins.

To find the partition function, we introduce a transfer-matrix of size 8×8 . Its nonzero elements are defined as follows:

$$\theta_{k,l} = \theta_{(\sigma_0^m, \sigma_1^m, \sigma_2^m), (\sigma_0^{m+1}, \sigma_1^{m+1}, \sigma_2^{m+1})} = \exp \left\{ - \frac{H^m}{T} \right\}, \quad (2.9)$$

$$k = \frac{1 - \sigma_0^m}{2} + (1 - \sigma_1^m) + 2(1 - \sigma_2^m),$$

$$l = \frac{1 - \sigma_0^{m+1}}{2} + (1 - \sigma_1^{m+1}) + 2(1 - \sigma_2^{m+1}).$$

Then the partition function of the model is expressed as follows:

$$Z_L = \text{tr}(\theta^L). \quad (2.10)$$

The free energy of the system per lattice node is defined in the standard way [9]:

$$f(H, T) = - \frac{T}{N} \ln Z_L(H, T), \quad (2.11)$$

where $N = 3L$ is the number of nodes of the considered lattice.

Internal energy per lattice node is equal to [9]:

$$u(H, T) = -T^2 \frac{\partial}{\partial T} \left(\frac{f(H, T)}{T} \right). \quad (2.12)$$

The specific heat per node is defined as [9]:

$$c(H, T) = \frac{\partial}{\partial T} u(H, T) = -2T \frac{\partial}{\partial T} \left(\frac{f(H, T)}{T} \right) - T^2 \frac{\partial^2}{\partial T^2} \left(\frac{f(H, T)}{T} \right). \quad (2.13)$$

Magnetization and susceptibility, respectively, are equal to [9]:

$$M(H, T) = - \frac{\partial}{\partial H} f(H, T), \quad (2.14)$$

$$\chi(H, T) = \frac{\partial}{\partial H} M(H, T), \quad (2.15)$$

and entropy:

$$S(H, T) = - \frac{\partial}{\partial T} f(H, T). \quad (2.16)$$

To formulate the following theorems, we introduce matrices:

$$\tau^1 = \begin{pmatrix} \tau_{11}^1 & \tau_{12}^1 & \tau_{13}^1 & \tau_{14}^1 \\ \tau_{21}^1 & \tau_{22}^1 & \tau_{23}^1 & \tau_{24}^1 \\ \tau_{31}^1 & \tau_{32}^1 & \tau_{33}^1 & \tau_{34}^1 \\ \tau_{41}^1 & \tau_{42}^1 & \tau_{43}^1 & \tau_{44}^1 \end{pmatrix} = \begin{pmatrix} \theta_{11} & \theta_{12} + \theta_{13} + \theta_{15} & \theta_{14} + \theta_{16} + \theta_{17} & \theta_{18} \\ \theta_{21} & \theta_{22} + \theta_{23} + \theta_{25} & \theta_{24} + \theta_{26} + \theta_{27} & \theta_{28} \\ \theta_{41} & \theta_{42} + \theta_{43} + \theta_{45} & \theta_{44} + \theta_{46} + \theta_{47} & \theta_{48} \\ \theta_{81} & \theta_{82} + \theta_{83} + \theta_{85} & \theta_{84} + \theta_{86} + \theta_{87} & \theta_{88} \end{pmatrix}, \quad (2.17)$$

$$\tau^2 = \begin{pmatrix} \tau_{11}^2 & \tau_{12}^2 \\ \tau_{21}^2 & \tau_{22}^2 \end{pmatrix} = \begin{pmatrix} e^{\frac{2\pi}{3}i}\theta_{52} + e^{-\frac{2\pi}{3}i}\theta_{53} + \theta_{55} & e^{-\frac{2\pi}{3}i}\theta_{54} + e^{\frac{2\pi}{3}i}\theta_{56} + \theta_{57} \\ e^{\frac{2\pi}{3}i}\theta_{72} + e^{-\frac{2\pi}{3}i}\theta_{73} + \theta_{75} & e^{-\frac{2\pi}{3}i}\theta_{74} + e^{\frac{2\pi}{3}i}\theta_{76} + \theta_{77} \end{pmatrix}, \quad (2.18)$$

$$\tau^3 = \begin{pmatrix} \tau_{11}^3 & \tau_{12}^3 \\ \tau_{21}^3 & \tau_{22}^3 \end{pmatrix} = \begin{pmatrix} e^{-\frac{2\pi}{3}i}\theta_{52} + e^{\frac{2\pi}{3}i}\theta_{53} + \theta_{55} & e^{\frac{2\pi}{3}i}\theta_{54} + e^{-\frac{2\pi}{3}i}\theta_{56} + \theta_{57} \\ e^{-\frac{2\pi}{3}i}\theta_{72} + e^{\frac{2\pi}{3}i}\theta_{73} + \theta_{75} & e^{\frac{2\pi}{3}i}\theta_{74} + e^{-\frac{2\pi}{3}i}\theta_{76} + \theta_{77} \end{pmatrix}, \quad (2.19)$$

and also expressions:

$$\begin{aligned} b_0 &= \frac{a}{2} - \sqrt{\frac{a^2}{4} - b + y_1}, \\ c_0 &= \frac{y_1}{2} - \text{sign}\left(\frac{a}{2}y_1 - c\right) \sqrt{\frac{y_1^2}{4} - d}, \\ b_1 &= \frac{a}{2} + \sqrt{\frac{a^2}{4} - b + y_1}, \\ c_1 &= \frac{y_1}{2} + \text{sign}\left(\frac{a}{2}y_1 - c\right) \sqrt{\frac{y_1^2}{4} - d}, \end{aligned} \quad (2.20)$$

where a, b, c, d are the coefficients of the characteristic equation of the fourth degree of the matrix (2.17):

$$\lambda^4 + a\lambda^3 + b\lambda^2 + c\lambda + d = 0, \quad (2.21)$$

and y_1 is one of the roots of the cubic equation obtained for equation (2.21) according to Ferrari's method described in the appendix.

Theorem 2.1. Main theorem

In the thermodynamic limit, the free energy, internal energy, specific heat, magnetization, susceptibility per lattice node and entropy, respectively, are calculated as follows:

$$f(H, T) = -T \frac{\ln(\lambda_{max}(H, T))}{3}, \quad (2.22)$$

$$u(H, T) = T^2 \frac{\partial}{\partial T} \frac{\ln(\lambda_{max}(H, T))}{3}, \quad (2.23)$$

$$c(H, T) = 2T \frac{\partial}{\partial T} \frac{\ln(\lambda_{max}(H, T))}{3} + T^2 \frac{\partial^2}{\partial T^2} \frac{\ln(\lambda_{max}(H, T))}{3}, \quad (2.24)$$

$$M(H, T) = \frac{T}{3} \frac{1}{\lambda_{max}(H, T)} \frac{\partial \lambda_{max}(H, T)}{\partial H}, \quad (2.25)$$

$$\chi(H, T) = \frac{T}{3} \left(\frac{1}{\lambda_{max}(H, T)} \frac{\partial^2 \lambda_{max}(H, T)}{\partial H^2} + \left(\frac{1}{\lambda_{max}(H, T)} \frac{\partial \lambda_{max}(H, T)}{\partial H} \right)^2 \right), \quad (2.26)$$

$$S(H, T) = \frac{1}{3} \left(\ln(\lambda_{max}(H, T)) + \frac{T}{\lambda_{max}(H, T)} \frac{\partial \lambda_{max}(H, T)}{\partial T} \right), \quad (2.27)$$

where the largest eigenvalue $\lambda_{max}(H, T)$ of the transfer-matrix θ has the form:

$$\lambda_{max}(H, T) = \frac{-b_0 + \sqrt{b_0^2 - 4c_0}}{2} \quad (2.28)$$

and its partial derivatives:

$$\frac{\partial \lambda_{max}(H, T)}{\partial T} = -\frac{a'_T \lambda_{max}^3 + b'_T \lambda_{max}^2 + c'_T \lambda_{max} + d'_T}{4\lambda_{max}^3 + 3a\lambda_{max}^2 + 2b\lambda_{max} + c}, \quad (2.29)$$

$$\frac{\partial \lambda_{max}(H, T)}{\partial H} = -\frac{a'_H \lambda_{max}^3 + b'_H \lambda_{max}^2 + c'_H \lambda_{max}}{4\lambda_{max}^3 + 3a\lambda_{max}^2 + 2b\lambda_{max} + c}, \quad (2.30)$$

$$\begin{aligned} \frac{\partial^2 \lambda_{max}(H, T)}{\partial H^2} = & -\frac{1}{4\lambda_{max}^3 + 3a\lambda_{max}^2 + 2b\lambda_{max} + c} \left(12\lambda_{max}^2 \left(\frac{\partial \lambda_{max}(H, T)}{\partial H} \right)^2 + \right. \\ & + a''_{HH} \lambda_{max}^3 + 6a'_H \lambda_{max}^2 \frac{\partial \lambda_{max}(H, T)}{\partial H} + 6a\lambda_{max} \left(\frac{\partial \lambda_{max}(H, T)}{\partial H} \right)^2 + b''_{HH} \lambda_{max}^2 + \\ & \left. + 4b'_H \lambda_{max} \frac{\partial \lambda_{max}(H, T)}{\partial H} + 2b \left(\frac{\partial \lambda_{max}(H, T)}{\partial H} \right)^2 + c''_{HH} \lambda_{max} + 2c'_H \frac{\partial \lambda_{max}(H, T)}{\partial H} \right). \end{aligned} \quad (2.31)$$

Theorem 2.2. *The partition function in a finite closed strip of length L can be written as:*

$$Z_L = \sum_{k=1}^8 \lambda_k^L, \quad (2.32)$$

where λ_k , $k = 1, \dots, 8$ are the roots of the characteristic polynomial of the transfer matrix θ , for which:

$$\lambda_{1,2} = \frac{-b_0 \pm \sqrt{b_0^2 - 4c_0}}{2}, \quad (2.33)$$

$$\lambda_{3,4} = \frac{-b_1 \pm \sqrt{b_1^2 - 4c_1}}{2} \quad (2.34)$$

– are the roots of the characteristic equation (2.21) of the matrix (2.17), found by the Ferrari method, and the eigenvalues:

$$\lambda_{5,6} = \frac{-b_2 \pm \sqrt{b_2^2 - 4c_2}}{2}, \quad (2.35)$$

$$\lambda_{7,8} = \frac{-b_3 \pm \sqrt{b_3^2 - 4c_3}}{2} \quad (2.36)$$

- are respectively the roots of the characteristic square polynomials of the matrices (2.18) and (2.19), for which:

$$\begin{aligned} b_j &= -(\tau_{11}^j + \tau_{22}^j), \\ c_j &= \tau_{11}^j \tau_{22}^j - \tau_{12}^j \tau_{21}^j, \\ & j = 2, 3. \end{aligned} \quad (2.37)$$

3. THE MAIN SPECIAL CASE: A MODEL WITH MULTISPIN INTERACTIONS OF EVEN NUMBER OF SPINS

3.1. Model description and results.

In this case, the components $\hat{H}_1^m, \hat{H}_3^m, \hat{H}_5^m$ of the elementary generating Hamiltonian (2.3) are zero, and the transfer-matrix θ has the center-symmetric form.

To formulate the following theorems, we introduce matrices:

$$\tau^4 = \begin{pmatrix} \tau_{11}^4 & \tau_{12}^4 \\ \tau_{21}^4 & \tau_{22}^4 \end{pmatrix} = \begin{pmatrix} \theta_{11} + \theta_{18} & \theta_{12} + \theta_{13} + \theta_{14} + \theta_{15} + \theta_{16} + \theta_{17} \\ \theta_{21} + \theta_{28} & \theta_{22} + \theta_{23} + \theta_{24} + \theta_{25} + \theta_{26} + \theta_{27} \end{pmatrix}, \quad (3.1)$$

$$\tau^5 = \begin{pmatrix} \tau_{11}^5 & \tau_{12}^5 \\ \tau_{21}^5 & \tau_{22}^5 \end{pmatrix} = \begin{pmatrix} \theta_{11} - \theta_{18} & \theta_{12} + \theta_{13} - \theta_{14} + \theta_{15} - \theta_{16} - \theta_{17} \\ \theta_{21} - \theta_{28} & \theta_{22} + \theta_{23} - \theta_{24} + \theta_{25} - \theta_{26} - \theta_{27} \end{pmatrix}. \quad (3.2)$$

Theorem 3.1. *In the thermodynamic limit, the free energy, internal energy, specific heat, and entropy per lattice node are respectively of the form (2.22), (2.23), (2.24), (2.27), where the largest eigenvalue λ_{max} of the transfer-matrix θ is of the form:*

$$\lambda_{max} = \frac{-b_4 + \sqrt{b_4^2 - 4c_4}}{2} \quad (3.3)$$

Theorem 3.2. *The partition function in a finite closed band of length L can be written in the form (2.32), where $\lambda_k, k = 1, \dots, 8$ are the roots of the characteristic polynomial of the transfer-matrix θ , for which:*

$$\lambda_{1,2} = \frac{-b_4 \pm \sqrt{b_4^2 - 4c_4}}{2}, \quad (3.4)$$

$$\lambda_{3,4} = \frac{-b_5 \pm \sqrt{b_5^2 - 4c_5}}{2} \quad (3.5)$$

— are respectively the roots of the characteristic square polynomials of matrices (3.1) and (3.2), for which:

$$\begin{aligned} b_j &= -(\tau_{11}^j + \tau_{22}^j), \\ c_j &= \tau_{11}^j \tau_{22}^j - \tau_{12}^j \tau_{21}^j, \\ j &= 4, 5, \end{aligned}$$

and the remaining eigenvalues are of the form:

$$\lambda_5 = (\theta_{22} + \theta_{27}) + e^{\frac{2\pi}{3}i}(\theta_{23} + \theta_{26}) + e^{-\frac{2\pi}{3}i}(\theta_{24} + \theta_{25}), \quad (3.6)$$

$$\lambda_6 = (\theta_{22} - \theta_{27}) + e^{\frac{2\pi}{3}i}(\theta_{23} - \theta_{26}) - e^{-\frac{2\pi}{3}i}(\theta_{24} - \theta_{25}), \quad (3.7)$$

$$\lambda_7 = (\theta_{22} + \theta_{27}) + e^{-\frac{2\pi}{3}i}(\theta_{23} + \theta_{26}) + e^{\frac{2\pi}{3}i}(\theta_{24} + \theta_{25}), \quad (3.8)$$

$$\lambda_8 = (\theta_{22} - \theta_{27}) + e^{-\frac{2\pi}{3}i}(\theta_{23} - \theta_{26}) - e^{\frac{2\pi}{3}i}(\theta_{24} - \theta_{25}), \quad (3.9)$$

where θ_{2l} are the elements of the transfer-matrix θ , $l = 2, \dots, 7$.

3.2. Pair correlations of the model with multispin interactions of even number of spins.

The Hamiltonian of this model coincides with (2.5), for which:

$$\begin{aligned}
H^m = & -\frac{J_1}{2} \left(\sigma_0^m \sigma_1^m + \sigma_0^m \sigma_2^m + \sigma_1^m \sigma_2^m + \sigma_0^{m+1} \sigma_1^{m+1} + \sigma_0^{m+1} \sigma_2^{m+1} + \sigma_1^{m+1} \sigma_2^{m+1} \right) - \\
& -J_2 \left(\sigma_0^m \sigma_0^{m+1} + \sigma_1^m \sigma_1^{m+1} + \sigma_2^m \sigma_2^{m+1} \right) - \\
& -J_3 \left(\sigma_0^m \sigma_1^{m+1} + \sigma_1^m \sigma_0^{m+1} + \sigma_2^m \sigma_0^{m+1} + \sigma_0^m \sigma_2^{m+1} + \sigma_1^m \sigma_2^{m+1} + \sigma_2^m \sigma_1^{m+1} \right) - \\
& -J_4 \left(\sigma_0^m \sigma_1^m \sigma_2^m \sigma_0^{m+1} + \sigma_0^m \sigma_1^m \sigma_2^m \sigma_1^{m+1} + \sigma_0^m \sigma_1^m \sigma_2^m \sigma_2^{m+1} + \right. \\
& \left. + \sigma_0^m \sigma_0^{m+1} \sigma_1^{m+1} \sigma_2^{m+1} + \sigma_1^m \sigma_0^{m+1} \sigma_1^{m+1} \sigma_2^{m+1} + \sigma_2^m \sigma_0^{m+1} \sigma_1^{m+1} \sigma_2^{m+1} \right) - \\
& -J_5 \left(\sigma_0^m \sigma_1^m \sigma_0^{m+1} \sigma_1^{m+1} + \sigma_0^m \sigma_2^m \sigma_0^{m+1} \sigma_2^{m+1} + \sigma_1^m \sigma_2^m \sigma_1^{m+1} \sigma_2^{m+1} \right) - \\
& -J_6 \left(\sigma_0^m \sigma_1^m \sigma_0^{m+1} \sigma_2^{m+1} + \sigma_0^m \sigma_2^m \sigma_1^{m+1} \sigma_2^{m+1} + \sigma_1^m \sigma_2^m \sigma_0^{m+1} \sigma_1^{m+1} + \sigma_0^m \sigma_1^m \sigma_1^{m+1} \sigma_2^{m+1} + \right. \\
& \left. + \sigma_0^m \sigma_2^m \sigma_0^{m+1} \sigma_1^{m+1} + \sigma_1^m \sigma_2^m \sigma_0^{m+1} \sigma_2^{m+1} \right) - \\
& -J_7 \sigma_0^m \sigma_1^m \sigma_2^m \sigma_0^{m+1} \sigma_1^{m+1} \sigma_2^{m+1}, \tag{3.10}
\end{aligned}$$

and the transfer-matrix (2.9) has the structure of the form

$$\theta = \begin{pmatrix} a & b & b & c & b & c & c & d \\ b & e & f & g & f & g & h & c \\ b & f & e & g & f & h & g & c \\ c & g & g & e & h & f & f & b \\ b & f & f & h & e & g & g & c \\ c & g & h & f & g & e & f & b \\ c & h & g & f & g & f & e & b \\ d & c & c & b & c & b & b & a \end{pmatrix}, \tag{3.11}$$

for which

$$\begin{aligned}
 a &= \exp \left\{ \frac{3J_1 + 3J_2 + 6J_3 + 6J_4 + 3J_5 + 6J_6 + J_7}{T} \right\}, \\
 b &= \exp \left\{ \frac{J_1 + J_2 + 2J_3 - 2J_4 - J_5 - 2J_6 - J_7}{T} \right\}, \\
 c &= \exp \left\{ \frac{J_1 - J_2 - 2J_3 + 2J_4 - J_5 - 2J_6 + J_7}{T} \right\}, \\
 d &= \exp \left\{ \frac{3J_1 - 3J_2 - 6J_3 - 6J_4 + 3J_5 + 6J_6 - J_7}{T} \right\}, \\
 e &= \exp \left\{ \frac{-J_1 + 3J_2 - 2J_3 - 2J_4 + 3J_5 - 2J_6 + J_7}{T} \right\}, \\
 f &= \exp \left\{ \frac{-J_1 - J_2 + 2J_3 - 2J_4 - J_5 + 2J_6 + J_7}{T} \right\}, \\
 g &= \exp \left\{ \frac{-J_1 + J_2 - 2J_3 + 2J_4 - J_5 + 2J_6 - J_7}{T} \right\}, \\
 h &= \exp \left\{ \frac{-J_1 - 3J_2 + 2J_3 + 2J_4 + 3J_5 - 2J_6 - J_7}{T} \right\}.
 \end{aligned} \tag{3.12}$$

and its eigenvalues (according to 3.4 - 3.9):

$$\begin{aligned}
 \lambda_{1,2} &= \frac{a + d + e + 2f + 2g + h \pm \sqrt{C_1}}{2}, \\
 \lambda_{3,4} &= \frac{a - d + e + 2f - 2g - h \pm \sqrt{C_2}}{2}, \\
 \lambda_{5,6} &= e + h - g - f, \\
 \lambda_{7,8} &= e - h + g - f,
 \end{aligned} \tag{3.13}$$

for which:

$$\begin{aligned}
 C_1 &= (a + d - e - 2f - 2g - h)^2 + 12(b + c)^2, \\
 C_2 &= (-a + d + e + 2f - 2g - h)^2 + 12(b - c)^2.
 \end{aligned} \tag{3.14}$$

Theorem 3.3. *In the thermodynamic limit the correlations of two spins for the model with Hamiltonian (3.10) are written as:*

$$\begin{aligned}
 &G_{(i,0),(j,0)} = G_{(i,1),(j,1)} = G_{(i,2),(j,2)} = \\
 &= \left(A_1 B_2 + \frac{1}{3} B_1 A_2 \right)^2 \left(\frac{\lambda_3}{\lambda_1} \right)^{j-i} + \left(A_1 A_2 - \frac{1}{3} B_1 B_2 \right)^2 \left(\frac{\lambda_4}{\lambda_1} \right)^{j-i} + \frac{8}{9} B_1^2 \left(\frac{\lambda_7}{\lambda_1} \right)^{j-i},
 \end{aligned} \tag{3.15}$$

$$\begin{aligned}
 &G_{(i,0),(j,1)} = G_{(i,0),(j,2)} = G_{(i,1),(j,2)} = \\
 &= \left(A_1 B_2 + \frac{1}{3} B_1 A_2 \right)^2 \left(\frac{\lambda_3}{\lambda_1} \right)^{j-i} + \left(A_1 A_2 - \frac{1}{3} B_1 B_2 \right)^2 \left(\frac{\lambda_4}{\lambda_1} \right)^{j-i} - \\
 &\quad - \frac{4}{9} B_1^2 \left(\frac{\lambda_7}{\lambda_1} \right)^{j-i},
 \end{aligned} \tag{3.16}$$

$$S_2 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}. \quad (3.24)$$

The expression (3.20) can be rewritten as:

$$\langle \sigma_\alpha^i \sigma_\beta^j \rangle = \frac{\text{Tr}(\theta^{i-1} S_\alpha \theta^{j-i} S_\beta \theta^{L-j+1})}{\text{Tr} \theta^L} = \frac{\text{Tr}(\Lambda^{i-1} S'_\alpha \Lambda^{j-i} S'_\beta \Lambda^{L-j+1})}{\text{Tr} \Lambda^L}. \quad (3.25)$$

Similarly, the expression (3.21) is written in the form

$$\langle \sigma_\gamma^k \rangle = \frac{\text{Tr}(S'_\gamma \Lambda^L)}{\text{Tr} \Lambda^L}, \quad (3.26)$$

for which Λ is the diagonal matrix obtained from the transfer-matrix by diagonalization transformation, and

$$S'_l = P^{-1} S_l P, \quad (3.27)$$

where P is the transition matrix of the transfer-matrix to the diagonal form:

$$P = \frac{1}{\sqrt{6}} \begin{pmatrix} \sqrt{3}A_1 & \sqrt{3}B_1 & 0 & 0 & \sqrt{3}B_2 & -\sqrt{3}A_2 & 0 & 0 \\ B_1 & -A_1 & \omega & 1 & A_2 & B_2 & -\omega & -1 \\ B_1 & -A_1 & \omega^2 & \omega^2 & A_2 & B_2 & -\omega^2 & -\omega^2 \\ B_1 & -A_1 & 1 & \omega & -A_2 & -B_2 & 1 & \omega \\ B_1 & -A_1 & 1 & \omega & A_2 & B_2 & -1 & -\omega \\ B_1 & -A_1 & \omega^2 & \omega^2 & -A_2 & -B_2 & \omega^2 & \omega^2 \\ B_1 & -A_1 & \omega & 1 & -A_2 & -B_2 & \omega & 1 \\ \sqrt{3}A_1 & \sqrt{3}B_1 & 0 & 0 & -\sqrt{3}B_2 & \sqrt{3}A_2 & 0 & 0 \end{pmatrix}, \quad (3.28)$$

$$P^{-1} = \frac{1}{\sqrt{6}} \begin{pmatrix} \sqrt{3}A_1 & B_1 & B_1 & B_1 & B_1 & B_1 & B_1 & \sqrt{3}A_1 \\ \sqrt{3}B_1 & -A_1 & -A_1 & -A_1 & -A_1 & -A_1 & -A_1 & \sqrt{3}B_1 \\ 0 & \bar{\omega} & \bar{\omega}^2 & 1 & 1 & \bar{\omega}^2 & \bar{\omega} & 0 \\ 0 & 1 & \bar{\omega}^2 & \bar{\omega} & \bar{\omega} & \bar{\omega}^2 & 1 & 0 \\ \sqrt{3}B_2 & A_2 & A_2 & -A_2 & A_2 & -A_2 & -A_2 & -\sqrt{3}B_2 \\ -\sqrt{3}A_2 & B_2 & B_2 & -B_2 & B_2 & -B_2 & -B_2 & \sqrt{3}A_2 \\ 0 & -\bar{\omega} & -\bar{\omega}^2 & 1 & -1 & \bar{\omega}^2 & \bar{\omega} & 0 \\ 0 & -1 & -\bar{\omega}^2 & \bar{\omega} & -\bar{\omega} & \bar{\omega}^2 & 1 & 0 \end{pmatrix}, \quad (3.29)$$

$$\omega = e^{\frac{2\pi}{3}i} \quad (3.30)$$

From the expressions (3.27) – (3.29) we obtain the matrix S'_0 :

$$\begin{pmatrix} 0 & 0 & 0 & 0 & A_1 B_2 + \frac{B_1 A_2}{3} & \frac{B_1 B_2 - A_1 A_2}{3} & -\frac{2}{3} e^{-\frac{\pi i}{3}} B_1 & \frac{2}{3} B_1 \\ 0 & 0 & 0 & 0 & B_1 B_2 - \frac{A_1 A_2}{3} & -\frac{A_1 B_2 - B_1 A_2}{3} & \frac{2}{3} e^{-\frac{\pi i}{3}} A_1 & -\frac{2}{3} A_1 \\ 0 & 0 & 0 & 0 & \frac{2}{3} e^{\frac{\pi i}{3}} A_2 & \frac{2}{3} e^{\frac{\pi i}{3}} B_2 & -\frac{1}{3} & -\frac{2}{3} e^{\frac{\pi i}{3}} \\ 0 & 0 & 0 & 0 & -\frac{2}{3} A_2 & -\frac{2}{3} B_2 & -\frac{2}{3} e^{-\frac{\pi i}{3}} & -\frac{1}{3} \\ A_1 B_2 + \frac{B_1 A_2}{3} & B_1 B_2 - \frac{A_1 A_2}{3} & \frac{2}{3} e^{-\frac{\pi i}{3}} A_2 & -\frac{2}{3} A_2 & 0 & 0 & 0 & 0 \\ \frac{B_1 B_2 - A_1 A_2}{3} & -A_1 A_2 - B_1 A_2 - \frac{A_1 B_2}{3} & \frac{2}{3} e^{-\frac{\pi i}{3}} B_2 & -\frac{2}{3} B_2 & 0 & 0 & 0 & 0 \\ -\frac{2}{3} e^{\frac{\pi i}{3}} B_1 & \frac{2}{3} e^{\frac{\pi i}{3}} A_1 & -\frac{1}{3} & -\frac{2}{3} e^{\frac{\pi i}{3}} & 0 & 0 & 0 & 0 \\ \frac{2}{3} B_1 & -\frac{2}{3} A_1 & -\frac{2}{3} e^{-\frac{\pi i}{3}} & -\frac{1}{3} & 0 & 0 & 0 & 0 \end{pmatrix}, \quad (3.31)$$

and the matrix S'_1 :

$$\begin{pmatrix} 0 & 0 & 0 & 0 & A_1 B_2 + \frac{B_1 A_2}{3} & \frac{B_1 B_2 - A_1 A_2}{3} & -\frac{2}{3} e^{\frac{\pi i}{3}} B_1 & -\frac{2}{3} e^{\frac{\pi i}{3}} B_1 \\ 0 & 0 & 0 & 0 & B_1 B_2 - \frac{A_1 A_2}{3} & -\frac{A_1 B_2 - B_1 A_2}{3} & \frac{2}{3} e^{\frac{\pi i}{3}} A_1 & \frac{2}{3} e^{\frac{\pi i}{3}} A_1 \\ 0 & 0 & 0 & 0 & \frac{2}{3} e^{-\frac{\pi i}{3}} A_2 & \frac{2}{3} e^{-\frac{\pi i}{3}} B_2 & -\frac{1}{3} & \frac{2}{3} \\ 0 & 0 & 0 & 0 & \frac{2}{3} e^{-\frac{\pi i}{3}} A_2 & \frac{2}{3} e^{-\frac{\pi i}{3}} B_2 & \frac{2}{3} & -\frac{1}{3} \\ A_1 B_2 + \frac{B_1 A_2}{3} & B_1 B_2 - \frac{A_1 A_2}{3} & \frac{2}{3} e^{\frac{\pi i}{3}} A_2 & \frac{2}{3} e^{\frac{\pi i}{3}} A_2 & 0 & 0 & 0 & 0 \\ \frac{B_1 B_2 - A_1 A_2}{3} & -A_1 A_2 - B_1 A_2 - \frac{A_1 B_2}{3} & \frac{2}{3} e^{\frac{\pi i}{3}} B_2 & \frac{2}{3} e^{\frac{\pi i}{3}} B_2 & 0 & 0 & 0 & 0 \\ -\frac{2}{3} e^{-\frac{\pi i}{3}} B_1 & \frac{2}{3} e^{-\frac{\pi i}{3}} A_1 & -\frac{1}{3} & \frac{2}{3} & 0 & 0 & 0 & 0 \\ -\frac{2}{3} e^{-\frac{\pi i}{3}} B_1 & \frac{2}{3} e^{-\frac{\pi i}{3}} A_1 & \frac{2}{3} & -\frac{1}{3} & 0 & 0 & 0 & 0 \end{pmatrix}. \quad (3.32)$$

From (3.25), taking into account (3.31) and (3.32), we obtain the expressions for $\langle \sigma_0^i \sigma_0^j \rangle$ and $\langle \sigma_0^i \sigma_1^j \rangle$, which in the thermodynamic limit at $L \rightarrow \infty$ coincide with (3.15) and (3.16), and $\langle \sigma_1^i \sigma_1^j \rangle = \langle \sigma_2^i \sigma_2^j \rangle = \langle \sigma_0^i \sigma_0^j \rangle$ and $\langle \sigma_0^i \sigma_2^j \rangle = \langle \sigma_1^i \sigma_2^j \rangle = \langle \sigma_0^i \sigma_1^j \rangle$, given the invariance of the system with respect to rotation.

Thus, from (3.26), given (3.31), (3.32), we have:

$$\langle \sigma_0^{k_0} \rangle = \langle \sigma_1^{k_1} \rangle = \langle \sigma_2^{k_2} \rangle = 0 \quad (3.33)$$

Note that it follows from the expression (3.33) that the magnetization of the system is zero in this case, and the correlations of the two spins (3.19) coincide with (3.15) and (3.16).

3.3. Ground states and correlation length of the model with multispin interactions of even number of spins.

It follows from the structure of the transfer-matrix (3.11) that the Hamiltonian (3.10) corresponding to one prism of the Ω_P^m model can take eight different values depending on the configuration at fixed values of the interspin interactions J_i . Fixing the interactions $J_2 = J_4 = J_6 = J_7 = J$ and varying $\frac{J_1}{J}, \frac{J_3}{J}, \frac{J_5}{J}$ from -15 to 15 , we show a picture of the ground states obtained by minimizing the [16] (??) in this section of the seven-dimensional space (Fig. 2), as well as its section at $\frac{J_5}{J} = 15, 0, -5, -15$ (Fig. 3).

The element of the transfer-matrix corresponding to this ground state, the vertices of the bodies bounding the regions of ground states given in the figures, the color designation, and one of the configurations corresponding to this ground state are given in the table (1), and the coordinates of the indicated vertices are given in the table (2).

Note that all model configurations that are in the same class [16] as those given in the table (2) also correspond to ground states.

Moreover, the number of configurations in the ground state to which the elements b, c, f, g of the transfer-matrix correspond is infinite in the thermodynamic limit according to the structure of this transfer-matrix.




FIGURE 2. The ground states of the model in coordinates $\frac{J_1}{J}, \frac{J_3}{J}, \frac{J_5}{J}$ at $J_2 = J_4 = J_6 = J_7 = J$




FIGURE 3. Ground states of the model in coordinates $\frac{J_1}{J}, \frac{J_3}{J}$ at $J_2 = J_4 = J_6 = J_7 = J$ and a) $\frac{J_5}{J} = 15$, b) $\frac{J_5}{J} = 0$, c) $\frac{J_5}{J} = -5$, d) $\frac{J_5}{J} = -15$

Similarly, fixing the interactions $J_3 = J_4 = J_6 = J_7 = J$ and varying $\frac{J_1}{J}, \frac{J_2}{J}, \frac{J_5}{J}$ from -15 to 15 , we show a picture of the ground states in a given section of seven-dimensional space (Fig. 4), as well as its cross-section at $\frac{J_5}{J} = 15, 0, -5, -15$ (Fig. 5).




FIGURE 4. Ground states of the model in coordinates $\frac{J_1}{J}, \frac{J_2}{J}, \frac{J_5}{J}$ at $J_3 = J_4 = J_6 = J_7 = J$




FIGURE 5. Ground states of the model in coordinates $\frac{J_1}{J}, \frac{J_2}{J}$ at $J_3 = J_4 = J_6 = J_7 = J$ and a) $\frac{J_5}{J} = 15$, b) $\frac{J_5}{J} = 0$, c) $\frac{J_5}{J} = -5$, d) $\frac{J_5}{J} = -15$

TABLE 1. Transfer-matrix elements, boundary points, color designations, and possible configurations corresponding to the ground states indicated in the figures.











Transfer-matrix element and boundary points	Color designation	Configuration example
$a,$ $A_1 A_2 A_3 A_4 A_6 A_7 A_8 A_9 A_{10} A_{11}$ $A_{13} A_{14} A_{19} A_{20} A_{21} A_{22} A_{23} A_{24},$ $B_1 B_2 B_3 B_8 B_9 B_{10} B_{11}$ $B_{15} B_{16} B_{20} B_{21} B_{28} B_{29} B_{32}$		
$b,$ $A_2 A_3 A_{17} A_{18} A_{19} A_{20},$ $B_2 B_3 B_4 B_{10} B_{11} B_{12} B_{27} B_{28}$		
$c,$ $A_{17} A_{18} A_{19} A_{20} A_{23} A_{24} A_{25} A_{27},$ $B_{11} B_{12} B_{21} B_{22} B_{25}$ $B_{26} B_{27} B_{28} B_{29} B_{30}$		
$d,$ $A_9 A_{10} A_{22} A_{23} A_{24} A_{25}$ $A_{26} A_{27} A_{29} A_{32} A_{33},$ $B_{19} B_{20} B_{21} B_{22} B_{29} B_{30} B_{31} B_{32}$		
$e,$ $A_{10} A_{11} A_{12} A_{15} A_{21}$ $A_{22} A_{29} A_{30} A_{31} A_{32},$ $B_6 B_7 B_8 B_9 B_{14} B_{15} B_{16} B_{17}$		
$f,$ $A_3 A_4 A_5 A_6 A_{14}$ $A_{15} A_{16} A_{17} A_{20} A_{21},$ $B_{11} B_{12} B_{13} B_{14} B_{15}$ $B_{21} B_{22} B_{23} B_{24} B_{25}$		
$g,$ $A_{15} A_{16} A_{17} A_{20} A_{21} A_{22}$ $A_{23} A_{27} A_{28} A_{29} A_{30},$ $B_4 B_5 B_6 B_9 B_{10} B_{11} B_{12} B_{13} B_{14} B_{15}$		
$h,$ $A_{11} A_{12} A_{13} A_{14} A_{15} A_{21},$ $B_{14} B_{15} B_{16} B_{17} B_{18}$ $B_{19} B_{20} B_{21} B_{22} B_{23}$		

TABLE 2. Coordinates of the boundary points of the regions of the ground states shown in the figures 2, 4.

Point	Coordinates	Point	Coordinates
A_1	$(15, 15, -15)$	B_1	$(15, 15, -15)$
A_2	$(15, \frac{5}{2}, -15)$	B_2	$(15, 4, -15)$
A_3	$(2, 9, -15)$	B_3	$(4, 15, -15)$
A_4	$(-4, 15, -15)$	B_4	$(2, 15, -15)$
A_5	$(-15, 15, -15)$	B_5	$(-15, 15, -15)$
A_6	$(-15, 15, -4)$	B_6	$(-15, 15, -6)$
A_7	$(-15, 15, 15)$	B_7	$(-15, 15, 15)$
A_8	$(15, 15, 15)$	B_8	$(-6, 15, 15)$
A_9	$(15, -\frac{5}{3}, 15)$	B_9	$(-6, 15, -6)$
A_{10}	$(-\frac{2}{3}, -\frac{5}{3}, 15)$	B_{10}	$(2, 15, -14)$
A_{11}	$(-6, 1, 15)$	B_{11}	$(2, 1, -7)$
A_{12}	$(-15, 1, 15)$	B_{12}	$(2, 1, -15)$
A_{13}	$(-15, 10, 15)$	B_{13}	$(-15, 1, -15)$
A_{14}	$(-15, 10, 1)$	B_{14}	$(-15, 1, 1)$
A_{15}	$(-15, 1, 1)$	B_{15}	$(-6, 1, 1)$
A_{16}	$(-15, 1, -15)$	B_{16}	$(-6, 1, 15)$
A_{17}	$(2, 1, -15)$	B_{17}	$(-15, 1, 15)$
A_{18}	$(15, 1, -15)$	B_{18}	$(-15, -15, 15)$
A_{19}	$(15, 1, -\frac{27}{2})$	B_{19}	$(2, -15, 15)$
A_{20}	$(2, 1, -7)$	B_{20}	$(2, -\frac{13}{3}, 15)$
A_{21}	$(-6, 1, 1)$	B_{21}	$(2, -\frac{13}{3}, -\frac{5}{3})$
A_{22}	$(-\frac{2}{3}, -\frac{5}{3}, 1)$	B_{22}	$(2, -15, -7)$
A_{23}	$(2, -\frac{5}{3}, -\frac{5}{3})$	B_{23}	$(-15, -15, -7)$
A_{24}	$(15, -\frac{5}{3}, -\frac{49}{6})$	B_{24}	$(-15, -15, -15)$
A_{25}	$(15, -\frac{17}{2}, -15)$	B_{25}	$(2, -15, -15)$
A_{26}	$(15, -15, -15)$	B_{26}	$(15, -15, -15)$
A_{27}	$(2, -15, -15)$	B_{27}	$(15, 1, -15)$
A_{28}	$(-15, -15, -15)$	B_{28}	$(15, 1, -\frac{27}{2})$
A_{29}	$(-14, -15, 1)$	B_{29}	$(15, -\frac{13}{3}, -\frac{49}{6})$
A_{30}	$(-15, -15, 1)$	B_{30}	$(15, -15, -\frac{27}{2})$
A_{31}	$(-15, -15, 15)$	B_{31}	$(15, -15, 15)$
A_{32}	$(-14, -15, 15)$	B_{32}	$(15, -\frac{13}{3}, 15)$
A_{33}	$(15, -15, 15)$	B_{33}	$(15, 15, 15)$

The boundary points of the ground states are of particular interest, since near them and in them themselves a change in the behavior of the correlation length can be observed at certain temperatures. The correlation length can be written in the form:

$$\xi^{-1} = \ln \left(\min_{\lambda_i \neq \lambda_{max}} \left| \frac{\lambda_{max}}{\lambda_i} \right| \right). \quad (3.34)$$

As an example, we present plots (Figures 6 – 8) of the inverse correlation lengths in the low-temperature region near some boundary points and directly at the boundary points themselves for the regions shown in Figures 2 and 4.




FIGURE 6. Inverse correlation length plot in the low-temperature region at $\frac{J_1}{J} = 2$, $\frac{J_5}{J} = -\frac{5}{3}$, $J_2 = J_4 = J_6 = J_7 = J$ and $\frac{J_3}{J} = -1.5, -1.6, -\frac{5}{3}$




FIGURE 7. Inverse correlation length plot in the low-temperature region at $\frac{J_1}{J} = -\frac{2}{3}$, $\frac{J_5}{J} = 1$, $J_2 = J_4 = J_6 = J_7 = J$ and $\frac{J_3}{J} = -1.5, -1.6, -\frac{5}{3}$




FIGURE 8. Inverse correlation length plot in the low-temperature region at $\frac{J_1}{J} = 2$, $\frac{J_5}{J} = -\frac{5}{3}$, $J_3 = J_4 = J_6 = J_7 = J$ and $\frac{J_2}{J} = -4.2, -4.3, -\frac{13}{3}$

The figures show that, first of all, at the boundary points of the ground state regions, the inverse correlation length can be non-zero when the temperature tends to zero.

In addition, peaks can be observed at and near the boundary points at certain temperature values in the inverse correlation length plots.

4. MODEL WITH SOME INTERACTIONS OF TWO, THREE, FOUR AND SIX SPINS

Let us consider the second case in which only interspin interactions generated by elementary carriers $\{t_0^m, t_0^{m+1}\}$, $\{t_0^m, t_1^m, t_2^m\}$, $\{t_0^{m+1}, t_1^{m+1}, t_2^{m+1}\}$, $\{t_0^m, t_1^m, t_2^{m+1}\}$, $\{t_0^m, t_1^{m+1}, t_2^{m+1}\}$, $\{t_0^m, t_1^m, t_0^{m+1}, t_1^{m+1}\}$, $\{t_0^m, t_1^m, t_2^m, t_0^{m+1}, t_1^{m+1}, t_2^{m+1}\}$ are nonzero.

The component (2.6) of the Hamiltonian (2.5) in this case has the form:

$$\begin{aligned}
 H^m = & - \left[J_1 \left(\sigma_0^m \sigma_0^{m+1} + \sigma_1^m \sigma_1^{m+1} + \sigma_2^m \sigma_2^{m+1} \right) + \right. \\
 & + \frac{1}{2} J_2 \left(\sigma_0^m \sigma_1^m \sigma_2^m + \sigma_0^{m+1} \sigma_1^{m+1} \sigma_2^{m+1} \right) + \\
 & + J_3 \left(\sigma_0^m \sigma_1^m \sigma_2^{m+1} + \sigma_0^m \sigma_2^m \sigma_1^{m+1} + \sigma_1^m \sigma_2^m \sigma_0^{m+1} \right) + \\
 & + J_4 \left(\sigma_0^m \sigma_1^{m+1} \sigma_2^{m+1} + \sigma_1^m \sigma_0^{m+1} \sigma_2^{m+1} + \sigma_2^m \sigma_0^{m+1} \sigma_1^{m+1} \right) + \\
 & + J_5 \left(\sigma_0^m \sigma_1^m \sigma_0^{m+1} \sigma_1^{m+1} + \sigma_0^m \sigma_2^m \sigma_0^{m+1} \sigma_2^{m+1} + \sigma_1^m \sigma_2^m \sigma_1^{m+1} \sigma_2^{m+1} \right) + \\
 & \left. + J_6 \sigma_0^m \sigma_1^m \sigma_2^m \sigma_0^{m+1} \sigma_1^{m+1} \sigma_2^{m+1} \right], \tag{4.1}
 \end{aligned}$$

where $\frac{1}{2}$ before J_2 is a coefficient that accounts for the transition from the n th to the $(n+1)$ th layer of the model.

To simplify further formulations we introduce the notations $p = \exp \left\{ \frac{J_1}{T} \right\}$, $q_1 = \exp \left\{ \frac{J_2}{T} \right\}$, $q_2 = \exp \left\{ \frac{J_3}{T} \right\}$, $q_3 = \exp \left\{ \frac{J_4}{T} \right\}$, $u = \exp \left\{ \frac{J_5}{T} \right\}$, $r = \exp \left\{ \frac{J_6}{T} \right\}$.

To formulate the following theorems, we introduce notation:

$$\begin{aligned}
 h &= p^6 q_1 q_2^4 q_3^4 r^2 u^2, \\
 h_1 &= -p^5 q_2 q_3 r^3 u \left(3q_2^2 q_3^2 (q_1^2 + q_2^2 q_3^2) + p^4 u^4 (1 + q_1^2 q_2^6 q_3^6) \right), \\
 h_2 &= q_1 \left(9p^4 q_2^4 q_3^4 (r^4 - p^4) - 3p^4 u^4 (q_2^8 + q_3^8) + 3p^8 r^4 u^4 (1 + q_2^8 q_3^8) + q_2^4 q_3^4 u^8 (p^{12} r^4 - 1) \right), \\
 h_3 &= p^5 q_2 q_3 r^3 u \left(q_2^2 q_3^2 (q_1^2 + q_2^2 q_3^2) - p^4 u^4 (1 + q_1^2 q_2^6 q_3^6) \right), \\
 h_4 &= q_1 \left(p^4 q_2^4 q_3^4 (r^4 - p^4) + p^4 u^4 (q_2^8 + q_3^8) - p^8 r^4 u^4 (1 + q_2^8 q_3^8) + q_2^4 q_3^4 u^8 (p^{12} r^4 - 1) \right)
 \end{aligned}$$

Theorem 4.1. *In the thermodynamic limit, the free energy, internal energy, specific heat, and entropy per lattice node are respectively (2.22), (2.23), (2.24), (2.27), where the largest eigenvalue λ_{max} of the transfer-matrix θ has the form:*

$$\lambda_{max} = \frac{-h_1 + \sqrt{h_1^2 - 4hh_2}}{2h} \tag{4.2}$$

Theorem 4.2. *The partition function in a finite closed strip of length L can be written in the form (2.32), where $\lambda_k, k = 1, \dots, 8$ are the roots of the characteristic polynomial of the transfer-matrix θ , two of which have multiplicity 3, and expressed in the following form:*

$$\lambda_{1,2} = \frac{-h_1 \pm \sqrt{h_1^2 - 4hh_2}}{2h}, \quad (4.3)$$

$$\lambda_{3,4,5} = \frac{-h_3 + \sqrt{h_3^2 - 4hh_4}}{2h}, \quad (4.4)$$

$$\lambda_{6,7,8} = \frac{-h_3 - \sqrt{h_3^2 - 4hh_4}}{2h}. \quad (4.5)$$

5. EXACT VALUE OF INVARIANT PERCOLATION IN THE TRIPLE-CHAIN
GENERALIZED ISING MODEL WITH ALL POSSIBLE MULTISPIN INTERACTIONS

Spin configurations in the cyclic-closed strip (2.1), in which there exists a side of -1: $\sigma_0^m = \sigma_1^m = -1$, $\sigma_0^m = \sigma_2^m = -1$, $\sigma_1^m = \sigma_2^m = -1$ at some $m \in \{0, 1, \dots, L-1\}$, we will call the percolation configurations.

If we exclude from the set of all model configurations the set of percolation configurations, we obtain the set of non-percolation configurations.

Accordingly, the non-percolation probability is the probability that the random state of the model will be from the set of non-percolation configurations. The non-percolation probability of a model is determined by the ratio of two quantities:

$$P_L = \frac{Z'_L}{Z_L}, \quad (5.1)$$

where Z_L is the partition function (2.8) of the model with Hamiltonian (2.5), Z'_L is the analog of the partition function, in which the summation is performed only on the non-percolation configurations. We will call it the partition function of non-percolation configurations.

The transfer-matrix of non-percolation configurations θ' coincides with the transfer-matrix (2.9) of all configurations of the model, but with zero matrix elements corresponding to the percolation configurations:

$$\theta' = \begin{pmatrix} * & * & * & 0 & * & 0 & 0 & 0 \\ * & * & * & 0 & * & 0 & 0 & 0 \\ * & * & * & 0 & * & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ * & * & * & 0 & * & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad (5.2)$$

To formulate the following theorems, we introduce the matrix:

$$\tau' = \begin{pmatrix} \theta'_{11} & \theta'_{12} + \theta'_{13} + \theta'_{15} \\ \theta'_{21} & \theta'_{22} + \theta'_{23} + \theta'_{25} \end{pmatrix} \quad (5.3)$$

Theorem 5.1. The main percolation theorem

In the thermodynamic limit for the non-percolation probability P_L (5.1) of lattice model with Hamiltonian (2.5) there is an equality:

$$\lim_{L \rightarrow \infty} \frac{\ln P_L}{L} = \ln \frac{\mu_{max}}{\lambda_{max}}, \quad (5.4)$$

where $\mu_{max} = \mu_{max}(H, T)$ is the largest root of the characteristic equation of the matrix τ' (5.3), and $\lambda_{max} = \lambda_{max}(H, T)$ is the largest root (2.28) of the characteristic equation of the matrix τ^1 (2.17).

Theorem 5.2. *In a finite closed strip of length L for the non-percolation probability P_L (5.1) of the lattice model with Hamiltonian (2.5) there is an equality:*

$$P_L = \frac{\sum_{i=1}^4 \lambda_i^L}{\sum_{k=1}^8 \lambda_k^L}, \quad (5.5)$$

where $\lambda'_{1,2}$ are the roots of the characteristic square equation of the matrix (5.3), and λ'_3, λ'_4 , respectively, are equal:

$$\lambda'_3 = \theta'_{52} e^{\frac{2\pi}{3}i} + \theta'_{53} e^{-\frac{2\pi}{3}i} + \theta'_{55}, \quad (5.6)$$

$$\lambda'_4 = \theta'_{52} e^{-\frac{2\pi}{3}i} + \theta'_{53} e^{\frac{2\pi}{3}i} + \theta'_{55}, \quad (5.7)$$

and λ_k , $k = 1, \dots, 8$ coincide with the roots (2.33) – (2.36).

The proofs of the formulated theorems 2.1 – 3.2, 4.1 – 5.2 are given in paragraph 9.

6. PLANAR MODEL WITH NEAREST NEIGHBOR, NEXT-NEAREST NEIGHBOR AND PLAQUETTE INTERACTIONS

The triple-chain generalized Ising model with multispin interactions is the minimal possible model on the torus, which unfolding yields a model on the plane.

Consider a planar model with nearest neighbor, next-nearest neighbor, and plaquette interactions, one layer of which contains three plaquettes.

This model is obtained by unfolding the lattice shown in Figure 1 in the plane with the addition of additional boundary conditions $t_3^m \equiv t_0^m$ and $t_3^{m+1} \equiv t_0^{m+1}$ (Fig. 9):




FIGURE 9. Planar Ising model with nearest-neighbor, next-nearestneighbor and plaquette interactions obtained by unfolding the triple-chain model on the plane

Here J_1 corresponds to nearest neighbor interactions on the same layer, J_2 corresponds to nearest neighbor interactions on adjacent layers, J_3 corresponds to next-nearest neighbor interactions, and J_4 is not explicitly shown in the figure 9 but corresponds to plaquette interactions.

The Hamiltonian of the system coincides with (2.5), where the Hamiltonian of one layer of the model is of the form:

$$\begin{aligned}
 H^m = & -\frac{J_1}{2} \left(\sigma_0^m \sigma_1^m + \sigma_1^m \sigma_2^m + \sigma_2^m \sigma_3^m + \sigma_0^{m+1} \sigma_1^{m+1} + \sigma_1^{m+1} \sigma_2^{m+1} + \sigma_2^{m+1} \sigma_3^{m+1} \right) - \\
 & -J_2 \left(\frac{1}{2} \sigma_0^m \sigma_0^{m+1} + \sigma_1^m \sigma_1^{m+1} + \sigma_2^m \sigma_2^{m+1} + \frac{1}{2} \sigma_3^m \sigma_3^{m+1} \right) - \\
 & -J_3 \left(\sigma_0^m \sigma_1^{m+1} + \sigma_1^m \sigma_2^{m+1} + \sigma_2^m \sigma_3^{m+1} + \sigma_1^m \sigma_0^{m+1} + \sigma_2^m \sigma_1^{m+1} + \sigma_3^m \sigma_2^{m+1} \right) - \\
 & -J_4 \left(\sigma_0^m \sigma_1^m \sigma_0^{m+1} \sigma_1^{m+1} + \sigma_1^m \sigma_2^m \sigma_1^{m+1} \sigma_2^{m+1} + \sigma_2^m \sigma_3^m \sigma_2^{m+1} \sigma_3^{m+1} \right),
 \end{aligned} \tag{6.1}$$

and as a transfer-matrix we can use an eighth-order matrix, the structure of which coincides with the transfer-matrix (3.11), the elements $a - h$ of which are rewritten according to (6.1). Then the eigenvalues of the transfer-matrix of this planar

model coincide with (3.13), and all physical characteristics can be described by the expressions obtained for the triple-chain model with interactions of an even number of spins — the theorems 3.1 and 3.2 are valid for this model. The expressions for pair correlations formulated in Theorem 3.3 are also valid for this model if we assume that all nodes t_0^m, t_1^m, t_2^m are in fact nearest neighbors because of the condition $t_3^m \equiv t_0^m$.

As an example for this planar model, we show plots of free energy, internal energy, specific heat and entropy in the thermodynamic limit in the low-temperature region for four different cases of parameters J_1, \dots, J_4 , one of which concerns the so-called planar gonihedric model for which $J_1 = J_2 = k$, $J_3 = -\frac{k}{2}$, $J_4 = \frac{1-k}{2}$ [11].




FIGURE 10. Plots, respectively, of free energy, internal energy, specific heat and entropy for the cases $J_1 = 0.11$, $J_2 = 0.12$, $J_4 = 0.13$ (left) and $J_1 = -0.11$, $J_2 = -0.12$, $J_4 = -0.13$ (right). $J_3 = r \in [-1, 1]$, $T \in [0.1, 4]$




FIGURE 11. Plots of free energy, internal energy, specific heat and entropy for the case $J_1 = \frac{1}{2}$, $J_2 = 1$, $J_4 = \frac{1}{2}$ and $J_3 = r \in [-3, 3]$, $T \in [0.1, 4]$




FIGURE 12. Plots of free energy, internal energy, specific heat and entropy for the case $J_1 = J_2 = \frac{1}{2}$, $J_4 = \frac{1}{4}$ and $J_3 = r \in [-1, 1]$, $T \in [0.1, 4]$

The last case (Fig. 12) involves an interaction that yields the so-called gonihedric model (when $J_3 = -\frac{1}{4}$).

In [12], while constructing phase diagrams for a three-dimensional cubic lattice at variable next-nearest neighbor interaction (J_3 in our work), a phase transition in the low-temperature region for two different values of J_4 was observed near the value of the parameter J_3 corresponding to the gonihedric model (at $\frac{J_3}{J_1} = -\frac{1}{4}$).

In our case, the special behavior of the plots in Figure 12 in the low-temperature region is observed at $J_3 = -\frac{1}{4}$, which corresponds to the planar gonihedric model ($\frac{J_3}{J_1} = -\frac{1}{2}$). When graphs were plotted on a set of other values of parameters $J_1 = J_2$ and J_4 (the parameters were taken positive), independent of each other, which were not included, a similar picture was observed: a special behavior of the graphs in the low-temperature region occurred at the point where the relation $\frac{J_3}{J_1} = -\frac{1}{2}$ was observed.

6.1. Planar gonihedric model.

A special case of the planar model with nearest neighbor, next-nearest neighbor, and plaquette interactions discussed above is the so-called planar gonihedric model, which was mentioned when graphing in Figure 12.

The Hamiltonian of one layer of the planar gonihedric model coincides with (6.1), for which [11]

$$\begin{aligned} J_1 &= J_2 = k, \\ J_3 &= -\frac{k}{2}, \\ J_4 &= \frac{1-k}{2}, \end{aligned} \tag{6.2}$$

i.e. all interspin interactions J_i in the gonihedric model depend on one parameter k , and the theorems 3.1 – 3.3 are valid for it, as well as for the more general case considered above.

As an example, for the case of the planar gonihedric model we have considered, we plot the free energy, internal energy, specific heat, and entropy at the variables temperature T and coefficient k (Fig. 13).




FIGURE 13. Plots of the physical characteristics of the planar gonihedric model at $k \in [-3, 3], T \in [0.1, 4]$

7. PLANAR TRIANGULAR MODEL

Consider a planar triangular model with different nearest neighbor interactions, triple interactions in the same triangle, and interaction with the external field, which, similar to the previous case of the planar model, is obtained by unfolding the lattice shown in Figure 1 in the plane with the addition of additional boundary conditions $t_3^m \equiv t_0^m$ and $t_3^{m+1} \equiv t_0^{m+1}$ (Fig. 14). For clarity, an additional $m + 2$ th layer of the model is added to the figure 14:




FIGURE 14. Planar triangular Ising model with nearest neighbor, next-nearest neighbor and plaquette interactions obtained by unfolding the triple-chain model on the plane

Here J_1, J_2, J_3 correspond to nearest neighbor interactions within a single triangle, J_4, J_5 are not shown explicitly but correspond to triple interactions within adjacent triangles, and H corresponds to the interaction with the external field.

The Hamiltonian of the system coincides with (2.5), where the Hamiltonian of one layer of the model is of the form:

$$\begin{aligned}
 H^m = & -\frac{J_1}{2} \left(\sigma_0^m \sigma_1^m + \sigma_0^m \sigma_2^m + \sigma_2^m \sigma_3^m + \sigma_0^{m+1} \sigma_1^{m+1} + \sigma_1^{m+1} \sigma_2^{m+1} + \sigma_2^{m+1} \sigma_3^{m+1} \right) - \\
 & -J_2 \left(\frac{1}{2} \sigma_0^m \sigma_0^{m+1} + \sigma_1^m \sigma_1^{m+1} + \sigma_2^m \sigma_2^{m+1} + \frac{1}{2} \sigma_3^m \sigma_3^{m+1} \right) - \\
 & -J_3 \left(\sigma_1^m \sigma_0^{m+1} + \sigma_2^m \sigma_1^{m+1} + \sigma_3^m \sigma_2^{m+1} \right) - \\
 & -J_4 \left(\sigma_0^m \sigma_1^m \sigma_0^{m+1} + \sigma_1^m \sigma_2^m \sigma_1^{m+1} + \sigma_2^m \sigma_3^m \sigma_2^{m+1} \right) - \\
 & -J_5 \left(\sigma_1^m \sigma_0^{m+1} \sigma_1^{m+1} + \sigma_2^m \sigma_1^{m+1} \sigma_2^{m+1} + \sigma_3^m \sigma_2^{m+1} \sigma_3^{m+1} \right) - \\
 & -\frac{H}{2} \left(\frac{1}{2} \sigma_0^m + \sigma_1^m + \sigma_2^m + \frac{1}{2} \sigma_3^m + \frac{1}{2} \sigma_0^{m+1} + \sigma_1^{m+1} + \sigma_2^{m+1} + \frac{1}{2} \sigma_3^{m+1} \right),
 \end{aligned} \tag{7.1}$$

The transfer-matrix of such a model contains 22 different matrix elements, its eigenvalues coincide with (2.33) – (2.36), and the theorems 2.1 and 2.2 are valid for it.

As an example for the planar triangular model, we show plots of free energy, internal energy, specific heat, magnetization, magnetic susceptibility, and entropy in the thermodynamic limit in the low-temperature region for two cases of parameters J_1, \dots, J_5 at a constant external field H — for the case when all $J_i > 0$, and when all $J_i < 0$.




FIGURE 15. Plots, respectively, of free energy, internal energy, specific heat and some of its cross sections, magnetization, some susceptibility cross sections and entropy for the case of $J_1 = 0.1, J_2 = 0.11, J_4 = 0.12, J_5 = 0.13, H = 0.15$. $J_3 = r \in [-3, 3], T \in [0.1, 4]$




FIGURE 16. Plots, respectively, of free energy, internal energy, specific heat and some of its cross sections, magnetization, some susceptibility cross sections and entropy for the case of $J_1 = -0.1, J_2 = -0.11, J_4 = -0.12, J_5 = -0.13, H = 0.15$. $J_3 = r \in [-3, 3], T \in [0.1, 4]$

8. EXAMPLES OF PHYSICAL CHARACTERISTICS IN THE THERMODYNAMIC LIMIT

Let us illustrate the physical characteristics stated in the main theorem 2.1. For this purpose, we will use the analytical expression of the largest eigenvalue λ_{max} (2.28) of the transfer-matrix θ and apply the formulas (2.22) – (2.26) to find the thermodynamic characteristics in the thermodynamic limit in the low-temperature region (Fig. 17 – 21) at the following interactions:

TABLE 3. Values of interspin interactions and elementary carriers generating them

Interaction	Value	Elementary carrier generating interaction
J_1	0.1	$\{t_0^m, t_1^m\}$
J_2	0.11	$\{t_0^m, t_0^{m+1}\}$
J_3	r	$\{t_0^m, t_1^{m+1}\}$
J_4	r	$\{t_0^m, t_2^{m+1}\}$
J_5	0.12	$\{t_0^m, t_1^m, t_2^m\}$
J_6	0.13	$\{t_0^m, t_1^m, t_0^{m+1}\}$
J_7	0.14	$\{t_0^m, t_1^m, t_1^{m+1}\}$
J_8	0.15	$\{t_0^m, t_1^m, t_2^{m+1}\}$
J_9	0.16	$\{t_0^m, t_0^{m+1}, t_2^{m+1}\}$
J_{10}	0.17	$\{t_0^m, t_0^{m+1}, t_1^{m+1}\}$
J_{11}	0.18	$\{t_0^m, t_1^{m+1}, t_2^{m+1}\}$
J_{12}	0.19	$\{t_0^m, t_1^m, t_2^m, t_0^{m+1}\}$
J_{13}	0.2	$\{t_0^m, t_1^m, t_0^{m+1}, t_1^{m+1}\}$
J_{14}	0.21	$\{t_0^m, t_1^m, t_0^{m+1}, t_2^{m+1}\}$
J_{15}	0.22	$\{t_0^m, t_1^m, t_1^{m+1}, t_2^{m+1}\}$
J_{16}	0.23	$\{t_0^m, t_0^{m+1}, t_1^{m+1}, t_2^{m+1}\}$
J_{17}	0.24	$\{t_0^m, t_1^m, t_2^m, t_0^{m+1}, t_1^{m+1}\}$
J_{18}	0.25	$\{t_0^m, t_1^m, t_0^{m+1}, t_1^{m+1}, t_2^{m+1}\}$
J_{19}	0.26	$\{t_0^m, t_1^m, t_2^m, t_0^{m+1}, t_1^{m+1}, t_2^{m+1}\}$
H	0.3	External magnetic field




FIGURE 17. Plots of free energy and internal energy per spin at given interactions $J_1 - J_{19}$ and external field H (table 3), $r \in [-3, 3]$, $T \in [0.1, 4]$




FIGURE 18. Plot of specific heat per spin for given interactions $J_1 - J_{19}$ and external field H (table 3), $r \in [-3, 3]$, $T \in [0.1, 4]$ and its cross sections at $T = 0.1$, $T = 0.2$, $T = 0.3$, $T = 0.4$, $T = 0.5$




FIGURE 19. Plot of magnetization at given interactions $J_1 - J_{19}$ and external field H (table 3), $r \in [-3, 3]$, $T \in [0.1, 4]$

The susceptibility cross sections in the low-temperature region are as follows:




FIGURE 20. Plots of susceptibility at given interactions $J_1 - J_{19}$ and external field H (table 3), $r \in [-0.9, -0.5]$, $T = 0.1$, $T = 0.15$, $T = 0.2$, $T = 0.25$, $T = 0.3$.




FIGURE 21. Entropy graph at given interactions $J_1 - J_{19}$ and external field H (table 3), $r \in [-1, -0.5]$, $T \in [0.05, 4]$.

9. PROOFS OF THEOREMS

9.1. Proof of the theorems 2.1, 2.2.

To prove the corresponding theorems, let us find all eigenvalues of the transfer-matrix θ (2.9).

One of the matrices commuting with θ is the "rotation" matrix D_1 by the angle $\frac{2\pi}{3}$, $D_1 : \{\sigma_0^m \rightarrow \sigma_1^{m+1}, \sigma_1^m \rightarrow \sigma_2^{m+1}, \sigma_2^m \rightarrow \sigma_0^{m+1}, m = 0, \dots, L-1\}$:

$$D_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad (9.1)$$

eigenvalues and basis vectors of corresponding eigenspaces of which are written out in the table 4:

 TABLE 4. Eigenvalues and eigenspaces of matrix D_1

Eigenvalue	The basis vectors of the eigenspace
1	$(1, 0, 0, 0, 0, 0, 0, 0, 0)^\top$ $(0, 1, 1, 0, 1, 0, 0, 0, 0)^\top$ $(0, 0, 0, 1, 0, 1, 1, 0, 0)^\top$ $(0, 0, 0, 0, 0, 0, 0, 0, 1)^\top$
$e^{\frac{2\pi}{3}i}$	$(0, 0, 0, e^{-\frac{2\pi}{3}i}, 0, e^{\frac{2\pi}{3}i}, 1, 0, 0)^\top$ $(0, e^{\frac{2\pi}{3}i}, e^{-\frac{2\pi}{3}i}, 0, 1, 0, 0, 0, 0)^\top$
$e^{-\frac{2\pi}{3}i}$	$(0, 0, 0, e^{\frac{2\pi}{3}i}, 0, e^{-\frac{2\pi}{3}i}, 1, 0, 0)^\top$ $(0, e^{-\frac{2\pi}{3}i}, e^{\frac{2\pi}{3}i}, 0, 1, 0, 0, 0, 0)^\top$

Using the theorem on the conservation of eigenspaces by commuting matrices [29], and considering that any linear combination of vectors of an eigenspace is also an eigenvector of that eigenspace, let us write down three different kinds of eigenvectors of the transfer-matrix θ :

$$\begin{aligned} \vec{y}_1 &= (x_1, x_2, x_2, x_3, x_2, x_3, x_3, x_4)^\top, \\ \vec{y}_2 &= (0, e^{\frac{2\pi}{3}i}x_1, e^{-\frac{2\pi}{3}i}x_1, e^{-\frac{2\pi}{3}i}x_2, x_1, e^{\frac{2\pi}{3}i}x_2, x_2, 0)^\top, \\ \vec{y}_3 &= (0, e^{-\frac{2\pi}{3}i}x_1, e^{\frac{2\pi}{3}i}x_1, e^{\frac{2\pi}{3}i}x_2, x_1, e^{-\frac{2\pi}{3}i}x_2, x_2, 0)^\top. \end{aligned} \quad (9.2)$$

As a result, all 8 eigenvectors of the matrix θ will be represented by these types of vectors, namely: four eigenvectors of the form \vec{y}_1 , two eigenvectors of the form \vec{y}_2 and two eigenvectors of the form \vec{y}_3 .

From the structure of the eigenvectors (9.2) and the structure of the transfer-matrix θ (2.9), four eigenvalues of the matrix θ will coincide with the eigenvalues of the matrix τ^1 (2.17), two will coincide with the eigenvalues of the matrix τ^2 (2.18), and two more will coincide with the eigenvalues of the matrix τ^3 (2.19).

The characteristic equation of the matrix τ^1 (2.17) is an equation of degree four (2.21) whose coefficients a, b, c, d are given in the appendix. In the general case it is solved by the method of Ferrari, also described in the appendix, according to which and taking into account the substitutions (2.20), four of the eight eigenvalues of the transfer-matrix θ have explicit expressions (2.33), (2.34).

The characteristic equations of the matrices τ^2 (2.18), τ^3 (2.19) are quadratic, respectively:

$$\begin{aligned}\lambda^2 + b_2\lambda + c_2 &= 0, \\ \lambda^2 + b_3\lambda + c_3 &= 0,\end{aligned}$$

whose coefficients b_2, c_2, b_3, c_3 respectively coincide with the expressions (2.37) and, consequently, their eigenvalues are expressed as (2.35), (2.36).

The transfer-matrix θ is real, with positive elements. Then, by the Perron-Frobenius theorem, it has a simple largest eigenvalue corresponding to an eigenvector with all positive components: comparing the obtained expressions (2.33) – (2.36), we have that the largest eigenvalue is λ_1 coinciding with (2.28).

The largest eigenvalue $\lambda_{max}(H, T)$ of the transfer-matrix satisfies the equation (2.21). Differentiating and doubly differentiating it by the external field H , we obtain the expressions (note that $d'_H = 0$ according to the form of the matrix (2.17)):

$$\begin{aligned}4\lambda_{max}^3 \frac{\partial \lambda_{max}}{\partial H} + a'_H \lambda_{max}^3 + 3a\lambda_{max}^2 \frac{\partial \lambda_{max}}{\partial H} + b'_H \lambda_{max}^2 + 2b\lambda_{max} \frac{\partial \lambda_{max}}{\partial H} + \\ + c'_H \lambda_{max} + c \frac{\partial \lambda_{max}}{\partial H} = 0, \\ 12\lambda_{max}^2 \left(\frac{\partial \lambda_{max}}{\partial H} \right)^2 + 4\lambda_{max}^3 \frac{\partial^2 \lambda_{max}}{\partial H^2} + a''_{HH} \lambda_{max}^3 + 6a'_H \lambda_{max}^2 \frac{\partial \lambda_{max}}{\partial H} + \\ + 6a\lambda_{max} \left(\frac{\partial \lambda_{max}}{\partial H} \right)^2 + 3a\lambda_{max}^2 \frac{\partial^2 \lambda_{max}}{\partial H^2} + b''_{HH} \lambda_{max}^2 + 4b'_H \lambda_{max} \frac{\partial \lambda_{max}}{\partial H} + \\ + 2b \left(\frac{\partial \lambda_{max}}{\partial H} \right)^2 + 2b\lambda_{max} \frac{\partial^2 \lambda_{max}}{\partial H^2} + c''_{HH} \lambda_{max} + 2c'_H \frac{\partial \lambda_{max}}{\partial H} + c \frac{\partial^2 \lambda_{max}}{\partial H^2} = 0,\end{aligned}$$

from which the expressions (2.30), (2.31) are obtained.

Similarly, differentiating equation (2.21) by temperature T , we obtain the expression:

$$\begin{aligned}4\lambda_{max}^3 \frac{\partial \lambda_{max}}{\partial T} + a'_T \lambda_{max}^3 + 3a\lambda_{max}^2 \frac{\partial \lambda_{max}}{\partial T} + b'_T \lambda_{max}^2 + 2b\lambda_{max} \frac{\partial \lambda_{max}}{\partial T} + \\ + c'_T \lambda_{max} + c \frac{\partial \lambda_{max}}{\partial T} + d'_T = 0,\end{aligned}$$

from which we get (2.29).

Expressions for the partial derivatives of $a'_H, b'_H, c'_H, a''_{HH}, b''_{HH}, c''_{HH}$ are given in the appendix.

9.2. Proof of theorems 3.1, 3.2.

In the first special case, in addition to commuting with the matrix D_1 (9.1), the transfer-matrix θ commutes with the matrix D_2 due to central symmetry:

$$D_2 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad (9.3)$$

eigenvalues and basis vectors of corresponding eigenspaces of which are written out in the table 5:

TABLE 5. Eigenvalues and eigenspaces of the matrix D_2

Eigenvalue	The basis vectors of the eigenspace
1	$(1, 0, 0, 0, 0, 0, 0, 1)^\top$ $(0, 1, 0, 0, 0, 0, 1, 0)^\top$ $(0, 0, 1, 0, 0, 1, 0, 0)^\top$ $(0, 0, 0, 1, 1, 0, 0, 0)^\top$
-1	$(-1, 0, 0, 0, 0, 0, 0, 1)^\top$ $(0, -1, 0, 0, 0, 0, 1, 0)^\top$ $(0, 0, -1, 0, 0, 1, 0, 0)^\top$ $(0, 0, 0, -1, 1, 0, 0, 0)^\top$

Using the theorem on the conservation of eigenspaces by commuting matrices [29], and assuming that all eigenvectors of matrix D_2 are either centrally symmetric or centrally antisymmetric, and that the transfer-matrix is centrally symmetric for this case, let us write down the first two possible kinds of eigenvectors of the transfer-matrix, which will be either centrally symmetric or centrally antisymmetric [28]:

$$\begin{aligned} \vec{y}_4 &= (x_1, x_2, x_2, x_2, x_2, x_2, x_2, x_1)^\top, \\ \vec{y}_5 &= (x_1, x_2, x_2, -x_2, x_2, -x_2, -x_2, -x_1)^\top. \end{aligned} \quad (9.4)$$

Moreover, using the commutation θ with the matrix D_1 , applying to which the condition of central symmetric or central antisymmetric eigenvector, we find linear combinations of vectors from eigenspaces (table 4) that are centrally symmetric or centrally antisymmetric, which are defined uniquely (without taking into account some multiplier x):

$$\begin{aligned} \vec{y}_6 &= (0, e^{\frac{2\pi}{3}ix}, e^{-\frac{2\pi}{3}ix}, x, x, e^{-\frac{2\pi}{3}ix}, e^{\frac{2\pi}{3}ix}, 0)^\top, \\ \vec{y}_7 &= (0, -e^{\frac{2\pi}{3}ix}, -e^{-\frac{2\pi}{3}ix}, x, -x, e^{-\frac{2\pi}{3}ix}, e^{\frac{2\pi}{3}ix}, 0)^\top, \\ \vec{y}_8 &= (0, x, e^{-\frac{2\pi}{3}ix}, e^{\frac{2\pi}{3}ix}, e^{\frac{2\pi}{3}ix}, e^{-\frac{2\pi}{3}ix}, x, 0)^\top, \\ \vec{y}_9 &= (0, -x, -e^{-\frac{2\pi}{3}ix}, e^{\frac{2\pi}{3}ix}, -e^{\frac{2\pi}{3}ix}, e^{-\frac{2\pi}{3}ix}, x, 0)^\top. \end{aligned} \quad (9.5)$$

As a result, all 8 eigenvectors of the matrix θ will be represented by these types of vectors, namely: two eigenvectors of the form \vec{y}_4 , two eigenvectors of the form \vec{y}_5 , and one eigenvector of the form $\vec{y}_6 - \vec{y}_9$ each.

From the structure of the eigenvectors (9.4), (9.5), two eigenvalues of the matrix θ will coincide with the eigenvalues of the matrix τ^4 (3.1), two will coincide with the eigenvalues of the matrix τ^5 (3.2), and four will coincide with the expressions (3.6) – (3.9).

Similar to the previous proof, the expressions for the eigenvalues of matrices τ^4, τ^5 of size 2×2 coincide with the expressions (3.4), (3.5), and the largest eigenvalue of matrix θ is λ_1 , the expression for which coincides with (3.3).

9.3. Proof of theorems 4.1, 4.2.

In the second special case, in addition to commuting with the matrix D_1 (9.1), the transfer-matrix θ commutes with the matrix D_3 :

$$D_3 = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}, \quad (9.6)$$

eigenvalues and basis vectors of corresponding eigenspaces of which are written out in the table 6:

TABLE 6. Eigenvalues and eigenspaces of the matrix D_3

Eigenvalue	The basis vectors of the eigenspace
1	$(0, 0, 0, 0, 1, 0, 0, 1)^\top$ $(0, 0, 0, 0, 0, 1, 1, 0)^\top$ $(1, 0, 0, 1, 0, 0, 0, 0)^\top$ $(0, 1, 1, 0, 0, 0, 0, 0)^\top$
-1	$(0, 0, 0, 0, -1, 0, 0, 1)^\top$ $(0, 0, 0, 0, 0, -1, 1, 0)^\top$ $(-1, 0, 0, 1, 0, 0, 0, 0)^\top$ $(0, -1, 1, 0, 0, 0, 0, 0)^\top$

Similarly to the previous proofs, using the eigenspace conservation theorem for commuting matrices and considering that linear combinations of vectors from eigenspaces are also eigenvectors of this subspace, let us write down one of the possible kinds of eigenvectors of the matrix θ , taking as a basis the eigenspace of the matrix D_3 corresponding to the eigenvalue 1:

$$\vec{y}_{10} = (x_1, x_2, x_2, x_3, x_2, x_3, x_3, x_4)^\top \quad (9.7)$$

Comparing the types of vectors \vec{y}_1 и \vec{y}_{10} , уточним вид собственного вектора трансфер-матрицы θ :

$$\vec{y}_{10_1} = (x_1, x_2, x_2, x_1, x_2, x_1, x_1, x_2)^\top \quad (9.8)$$

From the form of the eigenvector (9.8), we obtain that two eigenvalues of the matrix θ will coincide with the eigenvalues of the matrix τ^6 :

$$\begin{aligned} \tau^6 &= \begin{pmatrix} \theta_{11} + \theta_{14} + \theta_{16} + \theta_{17} & \theta_{12} + \theta_{13} + \theta_{15} + \theta_{18} \\ \theta_{21} + \theta_{24} + \theta_{26} + \theta_{27} & \theta_{22} + \theta_{23} + \theta_{25} + \theta_{28} \end{pmatrix} = \\ &= \begin{pmatrix} \frac{q_1 r (3 + p^4 q_2^4 q_3^4 u^4)}{p u q_2 q_3} & \frac{3 p q_2}{q_3 r u} + \frac{q_3^3 u^3}{p^3 q_2^3 r} \\ \frac{3 p q_2}{q_3 r u} + \frac{q_3^3 u^3}{p^3 q_2^3 r} & \frac{r (3 q_2^4 q_3^4 + p^4 u^4)}{p q_1 q_2^3 q_3^3 u} \end{pmatrix}, \end{aligned} \quad (9.9)$$

expressions for which coincide with (4.3).

To find the remaining six eigenvalues of the matrix θ , we divide its characteristic polynomial by the characteristic polynomial of the matrix τ^6 . As a result, we have a polynomial which is factorized into three identical square trinomials:

$$\begin{aligned} &\left(p^6 q_1 q_2^4 q_3^4 r^2 u^2 \lambda^2 + p^5 q_2 q_3 r^3 u \left[q_2^2 q_3^2 (q_1^2 + q_2^2 q_3^2) - p^4 u^4 (1 + q_1^2 q_2^6 q_3^6) \right] \lambda + \right. \\ &+ q_1 \left[p^4 q_2^4 q_3^4 (r^4 - p^4) + p^4 u^4 (q_2^8 + q_3^8) - p^8 r^4 u^4 (1 + q_2^8 q_3^8) + q_2^4 q_3^4 u^8 (p^{12} r^4 - 1) \right] \Big)^3 = \\ &= \left(h \lambda^2 + h_3 \lambda + h_4 \right)^3, \end{aligned} \quad (9.10)$$

from which we obtain two eigenvalues of the matrix θ , each of which has multiplicity 3 and whose expressions coincide with (4.4), (4.5).

Comparing the expressions (4.3) – (4.5) for the eigenvalues of the matrix θ , we obtain that the largest one is λ_1 , coinciding with (4.2).

9.4. Proof of theorems 5.1, 5.2.

The proof is based on commuting the matrix θ' (5.2) with the matrix D_1 (9.1).

Proceeding from the structure of the transfer-matrix θ' and the type of eigenvectors (9.2) obtained from the eigen subspaces of the matrix D_1 , let us write down three possible types of eigenvectors of the matrix θ' :

$$\begin{aligned} \vec{y}_1' &= (x_1, x_2, x_2, 0, x_2, 0, 0, 0)^\top, \\ \vec{y}_2' &= (0, e^{\frac{2\pi}{3}i} x_1, e^{-\frac{2\pi}{3}i} x_1, 0, x_1, 0, 0, 0)^\top, \\ \vec{y}_3' &= (0, e^{-\frac{2\pi}{3}i} x_1, e^{\frac{2\pi}{3}i} x_1, 0, x_1, 0, 0, 0)^\top. \end{aligned} \quad (9.11)$$

As a result, four of the eight eigenvectors of the matrix θ' will be represented by these types of vectors, namely: two eigenvectors of the form \vec{y}_1' , one eigenvector of the form \vec{y}_2' and one eigenvector of the form \vec{y}_3' .

The remaining four eigenvectors of the matrix θ' are zero.

The structure of the eigenvectors (9.11) shows that two of the eight eigenvalues of the matrix θ' coincide with the eigenvalues of the matrix τ' (5.3). Two more, respectively, with the expressions (5.6), (5.7), and the remaining four eigenvalues are zero.

Moreover, the largest eigenvalue of the transfer-matrix θ' coincides with the

largest eigenvalue of the matrix τ' , and the analog of the partition function Z'_L is equal to the sum of L th powers of the eigenvalues of the transfer-matrix θ' .

Thus, in the thermodynamic limit, the equality (5.4) is true, and in the finite-length strip, the expression for the probability of non-percolation coincides with (5.5).

10. CONCLUSION

In this paper, for the considered triple-chain rotation invariant generalized Ising model with all possible multispin interactions for a closed spin chain, the exact values of the partition function, free energy, internal energy, specific heat, magnetization, susceptibility, and entropy in the thermodynamic limit at $L \rightarrow \infty$, as well as the exact expression for the partition function in a finite band of length L are found by the transfer-matrix method.

The main results of the paper are formulated and proved in the form of two theorems for the general case. In the first theorem, the expressions of physical characteristics of the model in the thermodynamic limit are calculated using the exact expression for the largest eigenvalue of the transfer-matrix θ . In the second theorem, an expression for finding the partition function in a finite strip of length L is obtained using all 8 eigenvalues of the transfer-matrix θ , the expressions for which were obtained explicitly. Exact expressions for all 8 eigenvalues of the constructed transfer-matrix of size 8×8 are found: 4 eigenvalues are expressed through the roots of the fourth degree equation, which is solved by the Ferrari method described in the appendix, and the other four eigenvalues are expressed through the roots of two quadratic trinomials. The structure of the eigenvectors of the transfer-matrix is also shown.

With the use of analytical expressions of the largest eigenvalue of the transfer-matrix, an example illustrating the mentioned physical characteristics in the thermodynamic limit is given: at a variable temperature and two variable spin interactions corresponding to the interaction of next-nearest neighbors and equal to each other, three-dimensional plots of free energy, internal energy, specific heat, magnetization, susceptibility and entropy and cross sections of some of them at fixed temperatures are constructed.

The theorems analogous to the general case are formulated and proved for two special cases: the first of which is a model with multispin interactions of even number of spins, and the second is a model with some interactions of two, three, four and six spins.

For both special cases additional matrices (9.3, 9.6) different from the matrix for the general case (9.1), commuting with the transfer-matrix, were found, which allowed to clarify the type of eigenvectors and simplify the results obtained.

For the main special case, exact analytical expressions for all eigenvalues are given using the transfer-matrix method, four of which, including the largest one, are found as roots of two quadratic equations, and another four as roots of four linear equations. We obtain exact analytic expressions for the partition function in a strip of finite length L , as well as exact expressions for the free energy, internal energy, specific heat, and entropy of the model in the thermodynamic limit at

$L \rightarrow \infty$, which, due to the simple form of the largest eigenvalue of the transfer-matrix, also take a simplified form. In addition, the structure of the eigenvectors of the transfer-matrix is also shown.

An exact analytical expression of pair correlations in the thermodynamic limit was also found for this case, the structure of ground states was shown, for each of which an example of the corresponding configuration of the model was given, and it was also shown that the model has zero magnetization in the absence of an external field. With the help of two special projections of the ground states, the possibility of a special behavior of the correlation length at the boundaries of the ground states was illustrated.

In the second special case, by means of the theorem on the conservation of eigenspaces by commuting matrices, the exact expressions for all eigenvalues of the transfer-matrix are found as two roots of multiplicity one of one square trinomial, and two roots of multiplicity three of the second square trinomial.

Two theorems similar to those for the general case are also formulated and proved for this case, and expressions for the partition function in a finite strip of length L and expressions for the free energy, internal energy, specific heat and entropy in the thermodynamic limit at $L \rightarrow \infty$ are explicitly derived.

We formulate and prove two theorems on the probability of non-percolation, invariant with respect to rotation by an angle $2\pi/3$, in a closed strip of finite length L , and in the thermodynamic limit.

As an important special case of the model with multispin interactions of even number of spins, the model with nearest-neighbor, next-nearest neighbor and plaquette interactions is considered separately.

Results for the flat gonihedric model are formulated separately.

The large parameter selection possibilities allowed us to obtain accurate characterizations of physical quantities for the planar triangular model as well.

11. APPENDIX

Ferrari method for solving the fourth degree equation

To find the roots of the fourth degree equation (2.21), we will use the Ferrari method [38].

Suppose there is an equation of degree four:

$$\lambda^4 + a\lambda^3 + b\lambda^2 + c\lambda + d = 0, \quad (11.1)$$

for which, in the context of our work.

$$\begin{aligned}
a &= -\tau_{11}^1 - \tau_{22}^1 - \tau_{33}^1 - \tau_{44}^1, \\
b &= \tau_{11}^1\tau_{22}^1 + \tau_{11}^1\tau_{33}^1 + \tau_{11}^1\tau_{44}^1 + \tau_{22}^1\tau_{33}^1 + \tau_{22}^1\tau_{44}^1 + \tau_{33}^1\tau_{44}^1 - \\
&\quad - \tau_{24}^1\tau_{42}^1 - \tau_{34}^1\tau_{43}^1 - \tau_{23}^1\tau_{32}^1 - \tau_{12}^1\tau_{21}^1 - \tau_{13}^1\tau_{31}^1 - \tau_{14}^1\tau_{41}^1, \\
c &= -\tau_{11}^1\tau_{22}^1\tau_{33}^1 - \tau_{11}^1\tau_{22}^1\tau_{44}^1 - \tau_{11}^1\tau_{33}^1\tau_{44}^1 + \tau_{11}^1\tau_{24}^1\tau_{42}^1 + \tau_{11}^1\tau_{34}^1\tau_{43}^1 + \tau_{11}^1\tau_{23}^1\tau_{32}^1 - \\
&\quad - \tau_{22}^1\tau_{33}^1\tau_{44}^1 - \tau_{24}^1\tau_{43}^1\tau_{32}^1 - \tau_{23}^1\tau_{34}^1\tau_{42}^1 + \tau_{24}^1\tau_{42}^1\tau_{33}^1 + \tau_{43}^1\tau_{34}^1\tau_{22}^1 + \tau_{32}^1\tau_{23}^1\tau_{44}^1 + \\
&\quad + \tau_{12}^1\tau_{21}^1\tau_{33}^1 + \tau_{12}^1\tau_{21}^1\tau_{44}^1 - \tau_{12}^1\tau_{24}^1\tau_{41}^1 - \tau_{12}^1\tau_{23}^1\tau_{31}^1 - \tau_{13}^1\tau_{32}^1\tau_{21}^1 - \tau_{13}^1\tau_{34}^1\tau_{41}^1 + \\
&\quad + \tau_{13}^1\tau_{31}^1\tau_{22}^1 + \tau_{13}^1\tau_{31}^1\tau_{44}^1 + \tau_{14}^1\tau_{41}^1\tau_{22}^1 + \tau_{14}^1\tau_{41}^1\tau_{33}^1 - \tau_{14}^1\tau_{21}^1\tau_{42}^1 - \tau_{14}^1\tau_{43}^1\tau_{31}^1, \\
d &= \tau_{11}^1\tau_{22}^1\tau_{33}^1\tau_{44}^1 + \tau_{11}^1\tau_{24}^1\tau_{43}^1\tau_{32}^1 + \tau_{11}^1\tau_{23}^1\tau_{34}^1\tau_{42}^1 - \tau_{11}^1\tau_{24}^1\tau_{42}^1\tau_{33}^1 - \tau_{11}^1\tau_{43}^1\tau_{34}^1\tau_{22}^1 - \tau_{11}^1\tau_{32}^1\tau_{23}^1\tau_{44}^1 - \\
&\quad - \tau_{12}^1\tau_{21}^1\tau_{33}^1\tau_{44}^1 - \tau_{12}^1\tau_{24}^1\tau_{43}^1\tau_{31}^1 - \tau_{12}^1\tau_{23}^1\tau_{34}^1\tau_{41}^1 + \tau_{12}^1\tau_{24}^1\tau_{41}^1\tau_{33}^1 + \tau_{12}^1\tau_{21}^1\tau_{34}^1\tau_{43}^1 + \tau_{12}^1\tau_{31}^1\tau_{23}^1\tau_{44}^1 + \\
&\quad + \tau_{13}^1\tau_{21}^1\tau_{32}^1\tau_{44}^1 + \tau_{13}^1\tau_{31}^1\tau_{42}^1\tau_{24}^1 + \tau_{13}^1\tau_{34}^1\tau_{41}^1\tau_{22}^1 - \tau_{13}^1\tau_{24}^1\tau_{32}^1\tau_{41}^1 - \tau_{13}^1\tau_{34}^1\tau_{42}^1\tau_{21}^1 - \tau_{13}^1\tau_{31}^1\tau_{22}^1\tau_{44}^1 - \\
&\quad - \tau_{14}^1\tau_{21}^1\tau_{32}^1\tau_{43}^1 - \tau_{14}^1\tau_{31}^1\tau_{42}^1\tau_{23}^1 - \tau_{14}^1\tau_{41}^1\tau_{22}^1\tau_{33}^1 + \tau_{14}^1\tau_{23}^1\tau_{32}^1\tau_{41}^1 + \tau_{14}^1\tau_{21}^1\tau_{42}^1\tau_{33}^1 + \tau_{14}^1\tau_{31}^1\tau_{43}^1\tau_{22}^1.
\end{aligned} \tag{11.2}$$

By Ferrari method, if y_1 is an arbitrary valid positive solution of an auxiliary equation of degree three:

$$y^3 + Ay^2 + By + C = 0, \tag{11.3}$$

for which

$$\begin{aligned}
A &= -b, \\
B &= ac - 4d, \\
C &= -a^2d + 4bd - c^2,
\end{aligned}$$

then the four roots of the original equation of degree four (11.1) are found as roots of two quadratic equations

$$\lambda^2 + \frac{a}{2}\lambda + \frac{y_1}{2} = \pm \sqrt{\left(\frac{a^2}{4} - b + y_1\right)\lambda^2 + \left(\frac{a}{2}y_1 - c\right)\lambda + \left(\frac{y_1^2}{4} - d\right)}, \tag{11.4}$$

where the root expression in the right side is a complete square trinomial.

Using the Cardano-Vieta formula [37], we write one of the roots of the equation (11.3) in the form:

$$y_1 = \frac{1}{3} \left(-A + 2\sqrt{A^2 - 3B} \sin \left[\frac{1}{3} \left(\arcsin \left[\frac{2A^3 - 9AB + 27C}{2\sqrt{(A^2 - 3B)^3}} \right] + 2\pi \right) \right] \right), \tag{11.5}$$

then the equation (11.4) can be represented as two quadratic equations:

$$\lambda^2 + \frac{a}{2}\lambda + \frac{y_1}{2} = \pm \left(\sqrt{\frac{a^2}{4} - b + y_1}\lambda + \text{sign} \left(\frac{a}{2}y_1 - c \right) \sqrt{\frac{y_1^2}{4} - d} \right), \tag{11.6}$$

solving which, we obtain four roots of the original equation of degree four.

Computation of partial derivatives of the coefficients of equation (2.21) by the external field H.

For clarity, let us write out a matrix consisting of partial derivatives on the outer

field H matrix elements of matrix (2.17):

$$\begin{pmatrix} \frac{3}{T}\tau_{11}^1 & \frac{2}{T}\tau_{12}^1 & \frac{1}{T}\tau_{13}^1 & 0 \\ \frac{2}{T}\tau_{21}^1 & \frac{1}{T}\tau_{22}^1 & 0 & -\frac{1}{T}\tau_{24}^1 \\ \frac{1}{T}\tau_{31}^1 & 0 & -\frac{1}{T}\tau_{33}^1 & -\frac{2}{T}\tau_{34}^1 \\ 0 & -\frac{1}{T}\tau_{42}^1 & -\frac{2}{T}\tau_{43}^1 & -\frac{3}{T}\tau_{44}^1 \end{pmatrix} \quad (11.7)$$

From the expressions for the coefficients (11.2) we obtain expressions for partial derivatives:

$$a'_H = \frac{1}{T} \left(-3\tau_{11}^1 - \tau_{22}^1 + \tau_{33}^1 + 3\tau_{44}^1 \right),$$

$$b'_H = \frac{1}{T} \left(-4\tau_{11}^1\tau_{22}^1 + 2\tau_{11}^1\tau_{33}^1 - 2\tau_{22}^1\tau_{44}^1 - 4\tau_{33}^1\tau_{44}^1 + 2\tau_{24}^1\tau_{42}^1 + 4\tau_{34}^1\tau_{43}^1 - 4\tau_{12}^1\tau_{21}^1 - 2\tau_{13}^1\tau_{31}^1 \right),$$

$$\begin{aligned} c'_H = \frac{1}{T} \left(-3\tau_{11}^1\tau_{22}^1\tau_{33}^1 - \tau_{11}^1\tau_{22}^1\tau_{44}^1 + \tau_{11}^1\tau_{33}^1\tau_{44}^1 + \tau_{11}^1\tau_{24}^1\tau_{42}^1 - \tau_{11}^1\tau_{34}^1\tau_{43}^1 + 3\tau_{11}^1\tau_{23}^1\tau_{32}^1 + \right. \\ \left. + 3\tau_{22}^1\tau_{33}^1\tau_{44}^1 + 3\tau_{24}^1\tau_{43}^1\tau_{32}^1 + 3\tau_{23}^1\tau_{34}^1\tau_{42}^1 - 3\tau_{24}^1\tau_{42}^1\tau_{33}^1 - 3\tau_{43}^1\tau_{34}^1\tau_{22}^1 - 3\tau_{32}^1\tau_{23}^1\tau_{44}^1 + \right. \\ \left. + 3\tau_{12}^1\tau_{21}^1\tau_{33}^1 + \tau_{12}^1\tau_{21}^1\tau_{44}^1 - \tau_{12}^1\tau_{24}^1\tau_{41}^1 - 3\tau_{12}^1\tau_{23}^1\tau_{31}^1 - 3\tau_{13}^1\tau_{32}^1\tau_{21}^1 + \tau_{13}^1\tau_{34}^1\tau_{41}^1 + \right. \\ \left. + 3\tau_{13}^1\tau_{31}^1\tau_{22}^1 - \tau_{13}^1\tau_{31}^1\tau_{44}^1 + \tau_{14}^1\tau_{41}^1\tau_{22}^1 - \tau_{14}^1\tau_{41}^1\tau_{33}^1 - \tau_{14}^1\tau_{21}^1\tau_{42}^1 + \tau_{14}^1\tau_{43}^1\tau_{31}^1 \right), \end{aligned}$$

$$a''_{HH} = \frac{1}{T^2} \left(-9\tau_{11}^1 - \tau_{22}^1 - \tau_{33}^1 - 9\tau_{44}^1 \right),$$

$$b''_{HH} = \frac{1}{T^2} \left(16\tau_{11}^1\tau_{22}^1 + 4\tau_{11}^1\tau_{33}^1 + 4\tau_{22}^1\tau_{44}^1 + 16\tau_{33}^1\tau_{44}^1 - 4\tau_{24}^1\tau_{42}^1 - 16\tau_{34}^1\tau_{43}^1 - 16\tau_{12}^1\tau_{21}^1 - 4\tau_{13}^1\tau_{31}^1 \right),$$

$$\begin{aligned} c''_{HH} = \frac{1}{T^2} \left(-9\tau_{11}^1\tau_{22}^1\tau_{33}^1 - \tau_{11}^1\tau_{22}^1\tau_{44}^1 - \tau_{11}^1\tau_{33}^1\tau_{44}^1 + \tau_{11}^1\tau_{24}^1\tau_{42}^1 + \tau_{11}^1\tau_{34}^1\tau_{43}^1 + 9\tau_{11}^1\tau_{23}^1\tau_{32}^1 - \right. \\ \left. - 9\tau_{22}^1\tau_{33}^1\tau_{44}^1 - 9\tau_{24}^1\tau_{43}^1\tau_{32}^1 - 9\tau_{23}^1\tau_{34}^1\tau_{42}^1 + 9\tau_{24}^1\tau_{42}^1\tau_{33}^1 + 9\tau_{43}^1\tau_{34}^1\tau_{22}^1 + 9\tau_{32}^1\tau_{23}^1\tau_{44}^1 + \right. \\ \left. + 9\tau_{12}^1\tau_{21}^1\tau_{33}^1 + \tau_{12}^1\tau_{21}^1\tau_{44}^1 - \tau_{12}^1\tau_{24}^1\tau_{41}^1 - 9\tau_{12}^1\tau_{23}^1\tau_{31}^1 - 9\tau_{13}^1\tau_{32}^1\tau_{21}^1 - \tau_{13}^1\tau_{34}^1\tau_{41}^1 + \right. \\ \left. + 9\tau_{13}^1\tau_{31}^1\tau_{22}^1 + \tau_{13}^1\tau_{31}^1\tau_{44}^1 + \tau_{14}^1\tau_{41}^1\tau_{22}^1 + \tau_{14}^1\tau_{41}^1\tau_{33}^1 - \tau_{14}^1\tau_{21}^1\tau_{42}^1 - \tau_{14}^1\tau_{43}^1\tau_{31}^1 \right). \end{aligned}$$

REFERENCES

- [1] Ising E., Beitrag zur Theorie des Ferromagnetismus, Zeitschrift für Physik, 31 (1925), 253-258. DOI:10.1007/bf02980577
- [2] Onsager L., Crystal statistics. I. A two-dimensional model with an order-disorder transition, Physical Review, 65(3-4) (1944), 117-149. DOI: 10.1103/PhysRev.65.117
- [3] Kramers H. A., Wannier G. H. Statistics of the two-dimensional ferromagnet. Part I //Physical Review. - 1941. - T. 60. - №. 3. - C. 252. DOI:10.1103/physrev.60.252
- [4] Kramers H. A., Wannier G. H. Statistics of the two-dimensional ferromagnet. Part II //Physical Review. - 1941. - T. 60. - №. 3. - C. 263. DOI:10.1103/physrev.60.263
- [5] Kac M., Ward J. C. A combinatorial solution of the two-dimensional Ising model //Physical Review. - 1952. - T. 88. - №. 6. - C. 1332. DOI: 10.1103/PhysRev.88.1332
- [6] Yurishchev M.A., Statistical Mechanics of a triple Ising chain. Sov. J. Low Temp. Phys., 1984, v. 10, N. 6, p.625-635.
- [7] Yokota, T. (1989). Pair correlations for double-chain and triple-chain Ising models with competing interactions. Physical Review B, 39(16), 12312-12315. DOI:10.1103/physrevb.39.12312

- [8] Kalok, L., and L. C. De Menezes. "Competing interactions in double-chains: Ising and spherical models." *Zeitschrift für Physik B Condensed Matter* 20.2 (1975): 223-229. DOI:10.1007/bf01315695
- [9] Baxter, R.J. *Exactly Solved Models in Statistical Mechanics*; Academic Press: San Diego, California, USA, 2016; ISBN 0-12-083180-5.
- [10] Binder K., Heermann D. W., Binder K. *Monte Carlo simulation in statistical physics*. – Berlin : Springer-Verlag, 1992. – T. 8.
- [11] Bathas G. K. et al. Two-and three-dimensional spin systems with gonihedric action // *Modern Physics Letters A*. – 1995. – T. 10. – №. 35. – C. 2695-2701. DOI:10.1142/S0217732395002829
- [12] Cirillo, E. N. M., Gonnella, G., & Pelizzola, A. (1997). Critical behavior of the three-dimensional Ising model with nearest-neighbor, next-nearest-neighbor, and plaquette interactions. *Physical Review E*, 55(1), R17–R20. DOI:10.1103/physreve.55.r17
- [13] Yurishchev, M. A. (1989). Ising Model Consisting of Central and Three Side Chains. *Physica Status Solidi (b)*, 153(2), 703–710. DOI:10.1002/pssb.2221530229
- [14] Yurishchev, M. A. (2007). Ising models on the $2 \times 2 \times \infty$ lattices. *Journal of Experimental and Theoretical Physics*, 104(3), 461–466. DOI:10.1134/s1063776107030120
- [15] Ratner I. M., Eigenvalues of the Transfer Matrix of the Three-Dimensional Ising Model in the Particular Case $n = m = 2$, *Theoretical and Mathematical Physics*, 199(3) (2019), 909-921. DOI: 10.1134/S0040577919060102
- [16] Kashapov, I. A. Structure of ground states in three-dimensional using model with three-step interaction. *Teoreticheskaya i Matematicheskaya Fizika* 33.1 (1977): 110-118. DOI:10.1007/bf01039015
- [17] Dos Anjos, R. A., Viana, J. R., de Sousa, J. R., & Plascak, J. A. (2007). Three-dimensional Ising model with nearest- and next-nearest-neighbor interactions. *Physical Review E*, 76(2). DOI:10.1103/physreve.76.022103
- [18] Fisher, Michael E., and Robert J. Burford. "Theory of critical-point scattering and correlations. I. The Ising model." *Physical Review* 156.2 (1967): 583. DOI:10.1103/PhysRev.156.583
- [19] Fedro A. J. New method for obtaining exact solutions of the Ising problem for finite arrays of coupled chains // *Physical Review B*. – 1976. – V. 14. – №. 7. – C. 2983. DOI: 10.1103/PhysRevB.14.2983
- [20] Minlos, R. A., and Sinai, Y. G. (1970). Spectra of stochastic operators arising in lattice models of a gas. *Theoretical and Mathematical Physics*, 2(2), 167–176. DOI:10.1007/bf01036789
- [21] Malyshev, V. A., and Minlos, R. A. (2012). *Gibbs random fields: cluster expansions (Vol. 44)*. Springer Science and Business Media.
- [22] Minlos R. A., Khrapov P.V., Percolation in a finite strip for continuous systems, *Vestnik Moskovskogo Universiteta. Seriya 1. Matematika. Mekhanika.*, **1** (1985), 56-60.
- [23] Men'shikov M. V., Molchanov S.A., and Sidorenko A.F., Percolation theory and some applications, *Journal of Soviet Mathematics*, **42(4)** (1988), 1766-1810.
- [24] Minlos, R. A., and Khrapov, P. V. (1984). Cluster properties and bound states of the transfer matrix of the Yang-Mills model with a compact gauge group. 1. *Teor. Mat. Fiz.:(USSR)*, 61(3), 460-465.
- [25] Savvidy G. K., Wegner F. J. Geometrical string and spin systems, *Nuclear Physics B*, **413(3)** (1994), 605-613.
- [26] Johnston D.A., Mueller M., Janke W., Plaquette Ising models, degeneracy and scaling, *The European Physical Journal Special Topics*, **226(4)** (2017), 749-764. DOI: 10.1140/epjst/e2016-60329-4
- [27] Khrapov, P. V. (1984). Cluster expansion and spectrum of the transfer matrix of the two-dimensional ising model with strong external field. *Theoretical and Mathematical Physics*, 60(1), 734–735. DOI:10.1007/bf01018259
- [28] Andrew A. L. Eigenvectors of certain matrices // *Linear Algebra and its Applications*. – 1973. – T. 7. – №. 2. – C. 151-162. DOI: 10.1016/0024-3795(73)90049-9
- [29] Horn R. A., Johnson C. R. *Matrix analysis*. – Cambridge university press, 2012.

- [30] Andreev A. S., Khrapov P. V. Simulating Ising-like models on quantum computer. *Sovremennye informacionnye tehnologii i IT-obrazovanie = Modern Information Technologies and IT-Education*. – 2023. – T. 19. – №. 3.
- [31] Khrapov P.V. Fourier Transform of Transfer Matrices of Plane Ising Models. *Sovremennye informacionnye tehnologii i IT-obrazovanie = Modern Information Technologies and IT-Education*. 2019; 15(2):306-311. DOI: 10.25559/SITITO.15.201902.306-311
- [32] Khrapov P.V. Disorder Solutions for Generalized Ising Model with Multi-spin Interaction. *Sovremennye informacionnye tehnologii i IT-obrazovanie = Modern Information Technologies and IT-Education*. 2019; 15(2):312-319. DOI: 10.25559/SITITO.15.201902.312-319.
- [33] Khrapov, Pavel, and Veronika Nikishina. Exact solution Ising-like $2 \times 2 \times \infty$ models with multispin interactions. arXiv preprint arXiv:2205.15640 (2022). DOI: 10.48550/arXiv.2205.15640
- [34] Coniglio, A., Nappi, C. R., Peruggi, F., Russo, L., Percolation and phase transitions in the Ising model, *Communications in Mathematical Physics*, **51(3)** (1976), 315-323.
- [35] Miyamoto, A remark to Harris's theorem on percolation, *Communications in Mathematical Physics*, **44(2)** (1975), 169-173.
- [36] Khrapov P.V., Percolation in a finite strip, *Vestnik Moskovskogo Universiteta. Seriya 1. Matematika. Mekhanika.*, **4** (1985), 10-13.
- [37] Tabachnikov S.L., Fuks D.B., Mathematical divertissement, MCNMO, (2011). <https://elibrary.ru/item.asp?id=19464850>
- [38] Cardano G., Witmer T. R., Ore O. The rules of algebra: Ars Magna. – Courier Corporation, 2007. – T. 685.

PAVEL KHRAPOV, DEPARTMENT OF MATHEMATICS, BAUMAN MOSCOW STATE TECHNICAL UNIVERSITY, UL. BAUMANSKAYA 2-YA, 5/1, MOSCOW, 105005, MOSCOW, RUSSIAN FEDERATION
Email address: khrapov@bmstu.ru , pvkhrapov@gmail.com

NIKITA VOLKOV, DEPARTMENT OF MATHEMATICS, BAUMAN MOSCOW STATE TECHNICAL UNIVERSITY, UL. BAUMANSKAYA 2-YA, 5/1, MOSCOW, 105005, MOSCOW, RUSSIAN FEDERATION
Email address: volkovns@student.bmstu.ru , nikita.volkov01@mail.ru