# JOINT PARAMETER ESTIMATIONS FOR SPIN GLASSES

WEI-KUO CHEN\*, ARNAB SEN<sup>†</sup>, AND QIANG WU<sup>‡</sup>

Abstract. Spin glass models with quadratic-type Hamiltonians are disordered statistical physics systems with competing ferromagnetic and anti-ferromagnetic spin interactions. The corresponding Gibbs measures belong to the exponential family parametrized by (inverse) temperature  $\beta > 0$  and external field  $h \in \mathbb{R}$ . Given a sample from these Gibbs measures, a statistically fundamental question is to infer the temperature and external field parameters. In 2007, Chatterjee (Ann. Statist. 35 (2007), no. 5, 1931–1946) first proved that in the absence of external field  $h = 0$ , the maximum pseudolikelihood estimator for  $\beta$  is  $\sqrt{N}$ -<br>absence of external field  $h = 0$ , the maximum pseudolikelihood estimator for  $\beta$  is  $\sqrt{N}$ consistent under some mild assumptions on the disorder matrices. It was left open whether the same method can be used to estimate the temperature and external field simultaneously. In this paper, under some easily verifiable conditions, we prove that the bivariate maximum In this paper, under some easily verifiable conditions, we prove that the bivariate maximum<br>pseudolikelihood estimator is indeed jointly  $\sqrt{N}$ -consistent for the temperature and external field parameters. The examples cover the classical Sherrington-Kirkpatrick model and its diluted variants.

#### 1. Introduction and Main Results

Ising model possesses the Gibbs measure that belongs to the bivariate exponential family and takes the form, for  $\sigma \in \{-1, +1\}^N$ ,

$$
P_{\beta,h}(\sigma)=\frac{1}{Z_N(\beta,h)}\exp(\beta\langle \sigma, J\sigma\rangle+h\langle \sigma, \mathbf{1}\rangle),
$$

where  $\beta > 0$  is the (inverse) temperature,  $h \in \mathbb{R}$  is the external field, and  $J \in \mathbb{R}^N \times \mathbb{R}^N$  is a real-valued coupling matrix. In the past decades, estimation problems for the Ising model have received a great deal of attention from various communities due to its broad downstream applications, such as in statistics and computer science etc. One of the major research branches is, if the coupling matrix J is known, about estimating  $\beta$ , h, or both from a given sample of the Gibbs measure. One classical approach to this estimation problem is based on the maximum likelihood method. In the case of mean field Ising model, also known as Curie-Weiss model, where  $J$  is the adjacency matrix of complete graph with proper scaling, Comets and Gidas [\[11\]](#page-28-0) showed that for  $\beta > 0$  and  $h \neq 0$ , the maximum likelihood estimator Comets and Giaas [11] showed that for  $\rho > 0$  and  $n \neq 0$ , the maximum intention destimator (MLE) for  $\beta$  given h or vice versa is always  $\sqrt{N}$ -consistent. However, joint estimation of  $(\beta, h)$  is impossible. In some cases, maximum likelihood estimation for the Ising model can be very challenging since it involves the normalizing constant  $Z_N(\beta, h)$  whose computation, in particular, is nontrivial for models with  $J$  having both positive and negative entries. For example, in the setting of spin glass models, the interaction matrix J consists of random entries; as a result, understanding the limiting behavior of the log-normalizing constant as

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, 127 VINCENT HALL 206 CHURCH ST. SE MINneapolis, MN 55455

 $E\text{-}mail\;addresses: \; \texttt{``wkchen@umn.edu,} \; \texttt{``armab@umn.edu,} \; \texttt{``wuq@umn.edu.}$ 

Date: June 18, 2024.

<sup>2020</sup> Mathematics Subject Classification. Primary: 62F12, 62F10, 82B44.

Key words and phrases. Spin glasses, Sherrington-Kirkpatrick model, Pseudolikelihood estimator.

 $N \to \infty$  is generally a challenging problem and only a few cases are solvable in the literature. Arguably, one of the most intriguing results was the famous Parisi formula [\[17,](#page-28-1) [18\]](#page-28-2) that provided a variational representation for this limit established in the seminal work of Talagand [\[21\]](#page-28-3). Nonetheless, it is still a mystery to understand spin glass models for large but finite N.

Due to the intractability of MLE approach, it is desirable to seek for some other candidate methods that do not involve computing the log-normalizing constants. To this end, the pseudolikelihood approach [\[4,](#page-27-0)[5\]](#page-27-1) was proposed, where the pseudolikelihood function is defined as the product of conditional distributions such that the normalizing constant gets cancelled out. In this case, the maximum pseudolikelihood estimator (MPLE) can indeed alleviate the issue of dealing with a possibly complicated normalizing constant.

The MPLE approach has been studied a lot recently due to its efficiency and simplicity. The first work for the Ising-type model was initiated by Chatterjee [\[8\]](#page-27-2), where the author set  $h = 0$ and considered the single-parameter estimation problem for  $\beta > 0$ . Under the assumption that J has bounded operator norm and  $\liminf_{N\to\infty}N^{-1}\ln Z_N(\beta,h) > 0$ , it was shown that that *J* has bounded operator norm and  $\lim_{N\to\infty} N$   $\lim_{N\to\infty} N_0$ ,  $n > 0$ , it was shown that the MPLE is  $\sqrt{N}$ -consistency estimation for  $\beta$ . In particular, the result includes examples of spin glasses, such as the Sherrington-Kirkpatrick (SK) model [\[20\]](#page-28-4) and the Hopfield model [\[15\]](#page-28-5). Afterwards, the result of  $[8]$  was extended to other generalized Ising models  $[6, 16]$  $[6, 16]$  $[6, 16]$ . All those results have been restricted to the single parameter estimation setting. As we saw in the Curie-Weiss model, joint estimation is generally a much more challenging task. In fact, sometimes MPLE does not be even exist as opposed to the scenario for the single parameter. Incidentally, extending the single parameter estimation results to the joint parameters estimation setting was proposed as one of the open problems by Chatterjee [\[8\]](#page-27-2). An attempt to resolve this question was obtained by Ghosal and Mukherjee [\[14\]](#page-28-7) in the setting of the ferromagnetic Ising model where  $J$  has nonnegative entries and they established a set of sufficient conditions model where *J* has nonnegative entries and they established a set of under which joint estimation for  $(\beta, h)$  is possible with  $\sqrt{N}$ -consistency.

The definition of  $(\beta, n)$  is possible with  $\sqrt{N}$ -consistency.<br>In this paper, we aim to show that MPLE is a  $\sqrt{N}$ -jointly consistent estimator for  $(\beta, h)$ allowing  $J$  to possess positive and negative entries. More precisely, the matrix  $J$  is a weighted adjacency matrix of a graph, where the weights are i.i.d. centered random variable with finite third moment and the graph requires that its averaging degree diverges at least logarithmically fast and is comparable to the maximum degree. This setting covers a collection of spin glass models such as the celebrated SK model and its diluted variants. We show that the validity of our result requires checking three major conditions: the non-flatness of the free energy, the boundedness of the operator norm of  $J$ , and the positivity of the Hessian matrix of the pseudolikelihood function, see Theorem [1.1.](#page-3-0) Among them, the third condition plays a central role and it also appeared in [\[14\]](#page-28-7). Incidentally, the authors therein verified this positivity condition critically relying on the nonnegativity assumption of coupling coefficients which allowed them to use the nonlinear large deviation theory for random graphs [\[3,](#page-27-4) [9,](#page-28-8) [10,](#page-28-9) [13\]](#page-28-10). However, such technique does not seem to work when the coupling matrix  $J$  has both positive and negative entries. Until now, it remained as a main obstacle to extend joint estimation results beyond the non-negative coupling coefficients setting. We are able to check the nonsingularity condition with high probability in the setting of random coupling matrices via the small ball probability arguments  $[19, 22]$  $[19, 22]$  $[19, 22]$ .

1.1. Notation. Let  $\mathcal{G} := ([N], E)$  be a graph with vertex set  $[N] = \{1, 2, \ldots, N\}$  and edge set E. Denote by  $d_i$  and d the degree of vertex  $i \in [N]$  and the average degree of  $\mathcal{G}$ , respectively. In addition,  $d_{\text{max}} := \max_{i \in [N]} d_i$  denotes the maximum degree of  $\mathcal{G}$ . Let  $G = (g_{ij})_{i,j=1}^N$  be a

symmetric matrix and vanishes along the diagonal, *i.e.*,  $g_{ii} = 0$  for  $i \in [N]$ . Set

<span id="page-2-0"></span>
$$
J := \frac{1}{\sqrt{d}} A \circ G,\tag{1}
$$

where  $\circ$  denotes the Hadmard product and A is the adjacency matrix of G, that is,

$$
A_{ij} = \begin{cases} 1, & \text{if } (i,j) \in E, \\ 0, & \text{if } i = j \text{ or } (i,j) \notin E. \end{cases}
$$

If  $U_N$  and  $V_N$  are two non-negative random variables sampled from some probability measure P, we say  $U_N = O_p(V_N)$  if  $(U_N/V_N)_{N \geq 1}$  is a tight family under P, i.e., for each  $\epsilon > 0$ , there exists  $C$  such that

$$
P(U_N/V_N \leqslant C) \geqslant 1 - \epsilon \quad \text{for all } N \geqslant 1.
$$

The notation  $\Omega_p$  is defined similarly.

1.2. **Basic Setting of MPLE.** For any  $\beta > 0, h \in \mathbb{R}$ , the Gibbs measure associated to J is defined as

$$
P_{\beta,h}(\sigma) = \frac{\exp(\beta \langle J\sigma, \sigma \rangle / 2 + h\langle \sigma, \mathbf{1} \rangle)}{Z_N(\beta, h)},\tag{2}
$$

for all  $\sigma \in \Sigma_N := \{-1, +1\}^N$ , where

$$
Z_N(\beta, h) = \sum_{\sigma \in \Sigma_N} \exp(\beta \langle J\sigma, \sigma \rangle / 2 + h\langle \sigma, \mathbf{1} \rangle)
$$

is the normalizing constant, also known as the partition function in statistical physics. For any  $\tau \sim P_{\beta,h}$  and  $f : \Sigma_N \to \mathbb{R}$ , denote by  $\langle f(\tau) \rangle$  the expectation of  $f(\tau)$  with respect to the Gibbs measure. The free energy is defined as

<span id="page-2-1"></span>
$$
F_N(\beta, h) = \frac{1}{N} \log Z_N(\beta, h). \tag{3}
$$

We proceed to formulate the MPLE for  $(\beta, h)$ . In the sequel, we assume that  $\beta_0$ ,  $h_0$  are the ground truth parameters and a sample  $\tau \sim P_{\beta_0,h_0}(\cdot)$ . For  $\sigma \in \{-1,+1\}^N$ , the *pseudolikelihood* function is defined as

$$
L(\beta, h|\sigma) = \log \prod_{i=1}^{N} P_{\beta, h}(\sigma_i|\sigma_{j\neq i})
$$
  
= 
$$
\sum_{i=1}^{N} \sigma_i \left(\beta \sum_{j=1}^{N} J_{ij} \sigma_j + h\right) - \sum_{i=1}^{N} \log \cosh \left(\beta \sum_{j\neq i} J_{ij} \sigma_j + h\right) - \log 2.
$$
 (4)

Taking the partial derivatives of the pseudolikelihood function w.r.t. the parameters  $\beta, h$ , we have the corresponding *score functions*,

$$
S(\beta, h|\sigma) := \frac{\partial L}{\partial \beta} = \sum_{i=1}^{N} m_i(\sigma)(\sigma_i - \tanh(\beta m_i(\sigma) + h)), \tag{5}
$$

<span id="page-2-2"></span>
$$
Q(\beta, h|\sigma) := \frac{\partial L}{\partial h} = \sum_{i=1}^{N} (\sigma_i - \tanh(\beta m_i(\sigma) + h)),
$$
\n(6)

where

$$
m_i(\sigma) := \sum_{j=1}^N J_{ij}\sigma_j = \sum_{j \neq i} J_{ij}\sigma_j.
$$

We further compute the negative Hessian matrix  $H(\beta, h|\sigma)$  for later use as

$$
H(\beta, h|\sigma) := \begin{pmatrix} \sum_{i=1}^{N} m_i(\sigma)^2 \theta_i(\beta, h|\sigma)^2 & \sum_{i=1}^{N} m_i(\sigma) \theta_i(\beta, h|\sigma)^2 \\ \sum_{i=1}^{N} m_i(\sigma) \theta_i(\beta, h|\sigma)^2 & \sum_{i=1}^{N} \theta_i(\beta, h|\sigma)^2 \end{pmatrix},
$$
(7)

where for  $i \in [N], \theta_i(\beta, h|\sigma) := \text{sech}^2(\beta m_i(\sigma) + h)$ . From a straightforward computation, the determinant of the Hessian is

<span id="page-3-1"></span>
$$
|H(\beta, h|\sigma)| = \frac{N^2}{2}\widetilde{T}_N(\beta, h|\sigma),
$$

where

$$
\widetilde{T}_N(\beta, h|\sigma) := \frac{1}{N^2} \sum_{i,j \in [N]} \theta_i(\sigma) \theta_j(\sigma) (m_i(\sigma) - m_j(\sigma))^2.
$$
\n(8)

A quantity accompanied with  $T_N(\beta, h|\sigma)$  is

$$
T_N(\beta, h|\sigma) := \frac{1}{N^2} \sum_{i,j \in [N]} (m_i(\sigma) - m_j(\sigma))^2 = \frac{2}{N} \sum_{i \in [N]} (m_i(\sigma) - \bar{m}(\sigma))^2,
$$
(9)

where

$$
\bar{m}(\sigma) := \frac{1}{N} \sum_{i \in [N]} m_i(\sigma).
$$

For notational brevity, we will often suppress the dependence of  $\theta_i$ ,  $T_N$ , and  $T_N$  on  $\beta$  and  $h$ .

We assume that the coupling matrix  $J$  is given and we have access to a single sample from the ground truth  $P_{\beta_0,h_0}$ . Our ultimate aim is to determine the conditions on the model so that the pseudolikelihood estimator exists and is consistent. The following theorem presents a set of sufficient conditions to attend this purpose.

<span id="page-3-0"></span>**Theorem 1.1** (Existence and consistency of MPLE). Suppose that the coupling matrix  $J \in$  $\mathbb{R}^{N\times N}$  is symmetric and has zeros entries along the diagonal. For fixed  $\beta_0>0$  and  $h_0\in\mathbb{R}$ , let  $\tau \in \{-1,+1\}^N$  be a sample drawn from the Gibbs measure  $P_{\beta_0,h_0}$ . Assume that the following conditions hold.

(i) Non-flatness of the limiting free energy: There exists some  $0 < \beta' < \beta_0$  such that

$$
\liminf_{N \to \infty} (F_N(\beta', h_0) - F_N(0, h_0)) > 0.
$$

(ii) Boundedness of J:

$$
||J|| = O(1),
$$

where  $||J||$  is the operator norm of J.

(iii) Positivity of the Hessian: Under  $P_{\beta_0,h_0}$ ,

$$
\widetilde{T}_N(\tau) = \widetilde{T}_N(\beta_0, h_0 | \tau) = \Omega_p(1).
$$

Then with  $P_{\beta_0,h_0}$ -probability tending to one, the MPLE exists, i.e., the bivariate equation

$$
(S(\beta, h|\tau), Q(\beta, h|\tau)) = (0, 0), \tag{10}
$$

has a unique solution  $(\hat{\beta}(\tau), \hat{h}(\tau)) \in \mathbb{R}^2$ . Moreover,  $(\hat{\beta}(\tau), \hat{h}(\tau))$  is a  $\sqrt{N}$ -consistent estimator of the ground truth  $(\beta_0, h_0)$  in the sense that under  $P_{\beta_0, h_0}$ ,

$$
\|(\hat{\beta}(\tau) - \beta_0, \hat{h}(\tau) - h_0)\|_2 = O_p(N^{-1/2}).
$$

Note that this theorem holds for arbitrary real-valued symmetric matrices J with zeros along the diagonal and they do not need to be of the form [\(1\)](#page-2-0).

Remark 1.2. Under different assumptions, Ghosal-Mukherjee [\[14\]](#page-28-7) obtained similar existence and consistency results as Theorem 1.2, where the matrix  $J$  has nonnegative entries and uniformly bounded row sums. Our proof for the consistency part in Theorem [1.1](#page-3-0) is similar to that of  $[14,$  Theorem 1.2. However, the approach for the existence part in the same theorem does not apply to our setting.

<span id="page-4-0"></span>**Assumption 1.3.** For the underlying graph  $\mathcal{G} = (N, E)$ , we assume the average degree d and the maximum degree  $d_{\text{max}}$  satisfy

 $(i)$   $d \gg \log N$ , (*ii*)  $d_{\text{max}} \leqslant Cd$ ,

where  $C > 0$  is a constant independent of N.

Heuristically, the above condition specifies that the underlying graph can neither be too sparse nor have some vertices with very large degrees. Recall that a random variable  $X$  is  $\kappa^2$ -subgaussian if  $\mathbb{E}X = 0$  and  $\mathbb{E}e^{sX} \leqslant e^{\kappa^2 s^2/2}$  for all  $s \in \mathbb{R}$ . We say that  $G = (g_{ij})_{i,j \in [N]}$  is a symmetric  $\kappa^2$ -subgaussian matrix with vanishing diagonal if its upper triangular entries are independent  $\kappa^2$ -subgaussian. Now we state the first result that verifies the non-flatness of the free energy in temperature.

<span id="page-4-2"></span>**Theorem 1.4.** Suppose that G is a symmetric  $\kappa^2$ -subgaussian matrix with vanishing diagonal and Assumption [1.3](#page-4-0) (i) is in force. Then for any  $\beta > 0$  and  $h \in \mathbb{R}$ , there exist positive constants  $c_0, c, K$  independent of N such that

$$
\mathbb{P}\big(F_N(\beta,h)-F_N(0,h)\leqslant c_0\big)\leqslant K\exp(-cN).
$$

The next result establishes the bounednedness of the operator norm J.

<span id="page-4-3"></span>**Theorem 1.5.** Suppose that G is a symmetric  $\kappa^2$ -subgaussian matrix with vanishing diagonal and Assumption [1.3](#page-4-0) holds. Then there exist constants  $C, K > 0$ , depending on  $\kappa$ , such that

$$
\mathbb{P}(\|J\| \geqslant C) \leqslant K \exp(-d/\log N).
$$

As we shall see below, the proofs for these two theorems are not too difficult, which borrow some standard ideas and results from the treatment of the Ising model as well as the concentration inequalities for sparse random matrices  $[2,12]$  $[2,12]$ . The much more difficult component of this work lies on the verification of the strict positivity of the Hessian stated below.

<span id="page-4-1"></span>**Theorem 1.6.** Suppose that  $G$  is a symmetric random matrix with vanishing diagonal so that its upper triangular entries satisfy

$$
(g_{ij})_{i>j} \ are \ independent \ with \ \mathbb{E} \ g_{ij} = 0, \ \mathbb{E}[g_{ij}^2] = 1, \ and \ \mathbb{E} \ |g_{ij}|^3 \leq B,
$$
 (11)

where  $B > 0$  is some finite constant independent of N. Furthermore, suppose that Assumption [1.3](#page-4-0) (i) is satisfied and  $d \to \infty$  as  $N \to \infty$ . Then there exist positive constants  $c_0, c$ , and K such that

$$
\mathbb{P}\left(\min_{\sigma}\widetilde{T}_N(\sigma)\leqslant c_0\right)\leqslant K\exp(-cN).
$$

For the sample  $\tau$  from ground truth  $P_{\beta_0,h_0}$ , obviously  $\widetilde{T}_N(\tau) \geqslant \min_{\sigma} \widetilde{T}_N(\sigma)$  holds. The above theorem suggests that  $\widetilde{T}_N(\tau) \leq c_0$  holds with exponentially decaying annealed probability.

**Remark 1.7.** If the minimization is over a continuous sphere  $S_N := \{ \sigma \in \mathbb{R}^N : ||\sigma||^2 = N \}$ instead of discrete cube  $\{-1,+1\}^N$ , then the statement of Theorem [1.6](#page-4-1) is not true any more. For example, if J is nonsingular with probability one, which, in particular, holds for the SK model, then we take  $\sigma^* = \gamma J^{-1} \mathbf{1}$  for  $\gamma := N^{-1/2} ||J^{-1} \mathbf{1}||$ , so that  $m(\sigma^*) = \gamma \mathbf{1}$  and consequently,

$$
\widetilde{T}_N(\sigma^*) \leqslant T_N(\sigma^*) = 0.
$$

Therefore, we always have

$$
\min_{\sigma \in S_N} \widetilde{T}_N(\sigma) = 0.
$$

As of now, with high probability, all three conditions in Theorem [1.1](#page-3-0) have been verified under varying assumptions. Putting them all together, we obtain the asserted joint estimation for  $(\beta, h)$ .

<span id="page-5-0"></span>**Theorem 1.8.** For  $\beta_0 > 0$  and  $h_0 \in \mathbb{R}$ , suppose  $\tau \sim P_{\beta_0, h_0}$ . Suppose that

- (1) G is a symmetric  $\kappa^2$ -subgaussian random matrix with vanishing diagonal,
- (2) Assumption [1.3](#page-4-0) holds.

Then with P-probability tending to one, the bivariate MPLE,  $(\hat{\beta}(\tau), \hat{h}(\tau))$ , is jointly  $\sqrt{N}$ consistent.

**Example 1.9.** Letting the upper triangular entries of  $G$  being i.i.d. standard Gaussian, Theorem [1.8](#page-5-0) covers the classical SK model, for which  $\mathcal G$  is a complete graph so that  $d_{\text{max}} =$  $d = N - 1 \gg \log N$  satisfying Assumption [1.3.](#page-4-0) Another example is the diluted SK model, where G is a sparse Erdös-Rényi random graph  $G = G(N, p_N)$  with  $p_N \gg \log N/N$ .

1.3. Proof Sketches. Theorem [1.1](#page-3-0) consists of existence and consistency parts. The latter follows from a standard Delta method argument similar to [\[14,](#page-28-7) Theorem 1.2]. For existence, note that if the pseudolikelihood function diverges to the negative infinity as  $|\beta| + |h| \to \infty$ , then it has a unique maximizer (i.e., the MPLE exists) as it is already strictly concave thanks to the condition (iii). Almost the entirety of the proof of existence part is devoted to showing this divergence. We prove that this is indeed true if  $(\tau_i, m_i(\tau))_{i \in [N]}$  contains all possible signs. We then show that this property holds w.h.p. under the assumptions  $(i)$  and  $(ii)$ . Even though this is intuitively clear since the spins and the fields  $m_i$  should not be in favor of certain choices of signs overwhelmingly, proving it rigorously takes some effort. The key role is played by a general concentration result, see Lemma [2.1](#page-7-0) below, which says that for  $\tau \sim P_{\beta,h}$  and any suitably Lipschitz function f, the linear statistics  $N^{-1} \sum_{i=1}^{N} f(\tau_i, \tau_i m_i(\tau))$ stays around its conditional mean  $N^{-1} \sum_{i=1}^{N} \langle f(\tau_i, \tau_i m_i(\tau)) | \tau_i \rangle \forall i \neq i$ . As an immediate consequence, our lemma implies the concentration for the score functions  $S(\beta, h|\tau)$  and  $Q(\beta, h|\tau)$ (see Corollary [2.2](#page-7-1) below), which was previously established in [\[8\]](#page-27-2) using Stein's method for exchangeable pairs. Our result holds for more general functions and its proof relies on the cavity argument instead.

For Theorem [1.4,](#page-4-2) it suffices to show that  $E F_N(\beta, h)$  is strictly increasing in  $\beta > 0$  since this quantity is concentrated. To this end, we first show that the expected free energy is universal in the sense that  $E F_N(\beta, h)$  is asymptotically the same between Gaussian and subgaussian matrices with proper matching moments. Now for the free energy with Gaussian disorder, one can use the Gaussian integration by parts and then a standard cavity computation to

write

$$
\partial_{\beta} \mathbb{E} F(\beta, h) = \frac{\beta^2}{2} \Big( 1 - \frac{2}{Nd} \sum_{(i,j) \in E} \mathbb{E} \langle \sigma_i^1 \sigma_j^1 \sigma_i^2 \sigma_j^2 \rangle \Big)
$$
  
= 
$$
\frac{\beta^2}{2} \Big( 1 - \frac{2}{Nd} \sum_{(i,j) \in E} \mathbb{E} \langle \tanh(\beta m_i(\sigma) + h) \sigma_j \rangle^2 \Big)
$$

The crucial observation that we will establish below is that with high probability, if  $\sigma \sim$  $P_{\beta,h}(\cdot)$ , then there exists a vertex subset  $U = U(\sigma) \subseteq [N]$  that is of linear size and the fields  $(m_i(\sigma))_{i\in U}$  are uniformly bounded by an absolute constant. Additionally, each  $i \in U$  satisfies  $d_i \geq d/4$ . Using these, we readily see that  $\partial_{\beta} \mathbb{E} F(\beta, h) \geq c$  for some universal constant c as long as  $\beta$  is small enough and yields non-flatness of  $F_N(\beta, h)$ . The proof of Theorem [1.5](#page-4-3) follows straightforwardly by using the asymptotic results [\[2\]](#page-27-5) and the concentration bound [\[12\]](#page-28-13) for the operator norm of sparse subgaussian matrices.

The proof of Theorem [1.6](#page-4-1) is the central ingredient of this work. For simplicity, let us sketch an argument to show that

$$
\mathbb{P}\left(\min_{\sigma} T_N(\sigma) \leqslant c_0\right) \leqslant K \exp(-cN).
$$

We will prove the above by taking a union bound over the spin configuration  $\sigma \in \{-1,1\}^N$ . Notice that

<span id="page-6-0"></span>
$$
T_N(\sigma) = \frac{2}{N} \sum_{i \in [N]} (m_i(\sigma) - \bar{m}(\sigma))^2 = \frac{2}{N} \inf_{\gamma \in \mathbb{R}} ||m(\sigma) - \gamma \mathbf{1}||_2^2, \tag{12}
$$

It is easy to check using Markov inequality that  $||m(\sigma)||_{\infty} \leqslant M := 2^N$  with probability at least  $1 - C(1.5)^{-N}$ . On this event, the infimum over  $\gamma$  in [\(12\)](#page-6-0) can be restricted to the interval  $[-M, M]$ , which can further be approximated by taking minima over an  $\epsilon$ -net  $\mathcal D$  of size  $O(2^N \varepsilon^{-1})$  for small but constant  $\epsilon > 0$ . Therefore, it suffices to show that uniformly for any  $\sigma \in \{-1,1\}^N$  and any  $\gamma \in \mathcal{D}$ ,

<span id="page-6-1"></span>
$$
\mathbb{P}\big(\|m(\sigma) - \gamma \mathbf{1}\|_{2} \leqslant c_{1} \sqrt{N}\big) \leqslant K 10^{-N} \tag{13}
$$

for appropriate positive constant  $c_1$ . The event on the left-hand side can immediately be recognized as a small ball probability event. Had the entries of J been independent, the local fields  $m_i(\sigma)$  were independent. In that case, we can use the standard small ball probability result (e.g., Berry-Esseen theorem) for the linear combinations of independent random variables to infer that for each i, the probability that  $|m_i(\sigma) - \gamma| < c_1$  is at most 1/10 for sufficiently small  $c_1$  and then tensorizes the bound to obtain [\(13\)](#page-6-1). However, since J is symmetric,  $m_i(\sigma)$ are not independent anymore and the tensorization breaks down.

To overcome this obstacle, we borrow an idea from [\[22\]](#page-28-12). We only use the randomness of a principal submatrix of J of linear size while conditioning out the rest of the entries. Let  $T \text{ }\subset N$  be the set of the row (and the column) indices belonging to this principal submatrix, which we denote by  $J_T$ . Dealing with this smaller matrix instead of J has the critical advantage that the entries of  $J_T$  are independent (as long as  $|T| \le N/2$ ). We can then apply the previous small ball probability argument for  $J_T \sigma_T$ . However, successfully implementing this strategy requires that the matrix  $J_T$  is not too sparse in the sense that in the graph G the number of edges between each vertex of T and  $[N] \setminus T$  tends to  $\infty$  as  $N \to \infty$ . We apply Erdös' probabilistic method to guarantee the existence of such a 'good' deterministic set T provided G satisfies Assumption [1.3\(](#page-4-0)i) and  $d \to \infty$  as  $N \to \infty$ . This gives the lower bound for  $\min_{\sigma} T_N(\sigma)$ .

.

#### 8 CHEN, SEN, AND WU

The original term  $\widetilde{T}_N(\sigma)$  has an additional complication since  $\theta_i(\sigma) := \operatorname{sech}^2(\beta m_i(\sigma) + h)$ can be small if  $m_i(\sigma)$  is big. However, it can be shown that with overwhelming probability, there exists a (random) linear subset of indices of T satisfying  $|m_i(\sigma)| \leqslant C$ , which then yields the lower bound for  $\min_{\sigma} T_N(\sigma)$  as it follows from the previous step that the local fields  $(m_i(\sigma))_{i\in\mathcal{I}}$  have sufficient variability.

1.4. Structure of the paper. In the order their statements appear, the proofs for Theorems [1.1,](#page-3-0) [1.4,](#page-4-2) [1.5,](#page-4-3) and [1.6](#page-4-1) are accordingly located in Sections [2,](#page-7-2) [3,](#page-16-0) [4,](#page-19-0) and [5.](#page-21-0)

1.5. Acknowledgements. W-KC is partly supported by NSF grants DMS-1752184 and DMS-2246715 and Simons Foundation grant 1027727. He also thanks the hospitality of National Center for Theoretical Sciences in Taiwan during his visit between May 6-10, 2024, where part of this work was completed. AS is partly supported by Simons Foundation MP-TSM-00002716. QW and AS would like to thank Sumit Mukherjee for helpful discussions.

## 2. Proof of Theorem [1.1](#page-3-0)

<span id="page-7-2"></span>The proof of Theorem [1.1](#page-3-0) contains the existence and consistency of the MPLE. The latter is deferred to Section [2.3.](#page-14-0) For the existence part, as mentioned in the proof sketches, it suffices to show that the pseudolikelihood function diverges to negative infinity at infinity. We verify this property in Section [2.2](#page-13-0) based on a number of lemmas established in the following subsection.

2.1. **Auxiliary Lemmas.** Let  $\beta > 0$  and  $h \in \mathbb{R}$ . Let  $f(x, y)$  be defined on  $\{-1, 1\} \times \mathbb{R}$ . For any  $i \in [N]$  and  $\sigma \in \Sigma_N$ , set

$$
L_i(\sigma) = \langle f(\tau_i, \tau_i m_i(\tau)) | \tau_l = \sigma_l, \forall l \neq i \rangle
$$
  
= 
$$
\frac{e^{\beta m_i(\sigma) + h} f(1, m_i(\sigma)) + e^{-\beta m_i(\sigma) - h} f(-1, -m_i(\sigma))}{e^{\beta m_i(\sigma) + h} + e^{-\beta m_i(\sigma) - h}},
$$

where  $\langle \cdot \rangle$  is the Gibbs expectation with respect to  $\tau \sim P_{\beta,h}$ . Our first lemma establishes the following concentration for the linear statistics of  $(\tau_i, \tau_i m_i(\tau))_{i \in [N]}$ .

<span id="page-7-0"></span>**Lemma 2.1.** Let  $\tau$  be sampled from  $P_{\beta,h}$  for some  $\beta > 0$  and  $h \in \mathbb{R}$ . Assume that  $f(x, y)$ be defined on  $\{-1,1\} \times \mathbb{R}$  and satisfies  $||f||_{\infty} \leq 1$ ,  $||\partial_y f||_{\infty} < \infty$ , and  $||\partial_y^2 f||_{\infty} < \infty$ . For any  $N \geqslant 1$ , we have that

$$
\left\langle \left( \sum_{i=1}^{N} \left( f(\tau_i, \tau_i m_i(\tau)) - L_i(\tau) \right) \right)^2 \right\rangle \le 4N + C_1 N \|J\| + \frac{C_2}{2} N \|J\|^2,
$$
\n(14)

where

<span id="page-7-3"></span>
$$
C_1 := 2\beta + \|\partial_y f\|_{\infty},
$$
  
\n
$$
C_2 := 2\beta + 5\beta^2 + 2\beta \|\partial_y f\|_{\infty} + \|\partial_y^2 f\|_{\infty}.
$$

As an immediate corollary of this lemma with  $f(x, y) = y$  and  $f(x, y) = x$ , we have

<span id="page-7-1"></span>**Corollary 2.2** (Second moment bounds for score functions). Assume that  $||J|| = O(1)$ . Then there exists a constant C such that

$$
\langle (S(\beta,h|\tau))^2 \rangle \leqslant CN, \quad \langle (Q(\beta,h|\tau))^2 \rangle \leqslant CN,
$$

where  $S$  and  $Q$  are defined in  $(5)$  and  $(6)$ .

*Proof of Lemma [2.1.](#page-7-0)* Let distinct  $i, j \in [N]$  be fixed. Set

$$
m_j^{(i)}(\sigma) := \sum_{k \neq i,j} J_{jk} \sigma_k
$$

and  $L_i^{(i)}$  $j^{(i)}(σ)$  is same as  $L_j(σ)$  with  $m_j(σ)$  replaced by  $m_j^{(i)}$  $j^{(i)}(\sigma)$ . Since  $L_i(\tau)$  is the expectation of  $f(\tau_i, \tau_i m_i(\tau))$  conditionally on  $(\tau_i)_{i \neq i}$ , we have

$$
\langle \big(f(\tau_i, \tau_i m_i(\tau)) - L_i(\tau)\big) \big(f(\tau_j, \tau_j m_j(\tau)) - L_j^{(i)}(\tau)\big) \rangle = 0,
$$

which implies

$$
\langle (f(\tau_i, \tau_i m_i(\tau)) - L_i(\tau)) (f(\tau_j, \tau_j m_j(\tau)) - L_j(\tau)) \rangle
$$
  
=  $\langle c_{ij}(\tau) (f(\tau_i, \tau_i m_i(\tau)) - L_j(\tau)) \rangle$ ,

where

$$
c_{ij}(\tau) := L_j^{(i)}(\tau) - L_j(\tau).
$$

Now, using Taylor expansion,

$$
\left| L_j^{(i)}(\tau) - L_j(\tau) + J_{ij}\partial_{J_{ij}} L_j(\tau) \right| \leqslant \frac{J_{ij}^2}{2} \max_{J_{ij} \in \mathbb{R}} \left| \partial_{J_{ij}}^2 L_j(\tau) \right|.
$$

Here,

$$
\partial_{J_{ij}} L_j(\tau) = \frac{e^{\beta m_j(\tau) + h} \partial_y f(1, m_j(\tau)) + e^{-\beta m_j(\sigma) - h} \partial_y f(-1, -m_j(\tau))}{e^{\beta m_j(\tau) + h} + e^{-\beta m_j(\tau) - h}}
$$

$$
+ \beta \frac{e^{\beta m_j(\tau) + h} f(1, m_j(\tau)) - e^{-\beta m_j(\tau) - h} f(-1, -m_j(\tau))}{e^{\beta m_j(\tau) + h} + e^{-\beta m_j(\tau) - h}}
$$

$$
- \beta \frac{e^{\beta m_j(\tau) + h} f(1, m_j(\tau)) + e^{-\beta m_j(\tau) - h} f(-1, -m_j(\tau))}{e^{\beta m_j(\tau) + h} + e^{-\beta m_j(\tau) - h}} \tanh(\beta m_j(\tau) + h)
$$

and

$$
|\partial_{J_{ij}}L_j(\tau)|\leqslant C_1:=2\beta+\|\partial_y f\|_\infty.
$$

Since  $\partial_{J_{ij}}L_j(\tau)$  is independent of  $i,$  it follows that

$$
\left|\sum_{i\neq j} J_{ij}\partial_{J_{ij}} L_j(\tau)\right| = \left|\sum_{i,j=1}^N J_{ij}\partial_{J_{ij}} L_j(\tau)\right| \leqslant C_1 N \|J\|.
$$

On the other hand, since

$$
|\partial_{J_{ij}}^2 L_j(\tau)| \leq C_2 := 2\beta + 5\beta^2 + 2\beta \|\partial_y f\|_{\infty} + \|\partial_y^2 f\|_{\infty},
$$

we have

$$
\left| \sum_{i \neq j} J_{ij}^2 \partial_{J_{ij}}^2 L_j(\tau) \right| = \left| \sum_{i,j=1}^N J_{ij}^2 \partial_{J_{ij}}^2 L_j(\tau) \right|
$$
  
\$\leq C\_2 \sum\_{i,j=1}^N J\_{ij}^2 = C\_2 \mathbf{1}^T (J \circ J) \mathbf{1} \leq C\_2 N || J \circ J || \leq C\_2 N || J ||^2.\$

From these, we arrive at

$$
\sum_{i \neq j} |c_{ij}(\tau)| \leq C_1 N ||J|| + \frac{C_2}{2} N ||J||^2
$$

and thus,

$$
\sum_{i\neq j}\langle \big(f(\tau_i,\tau_i m_i(\tau))-L_i(\tau)\big)\big(f(\tau_j,\tau_j m_j(\tau))-L_j(\tau)\big)\rangle \leqslant C_1N||J||+\frac{C_2}{2}N||J||^2.
$$

This readily yields our assertion.

Let  $0 < \delta < M$  be fixed. Let  $\rho$  be a nonnegative twice differentiable function on R with  $\rho \equiv 1$  on  $(\delta, \infty)$ ,  $\rho \equiv 0$  on  $(-\infty, 0)$ , and  $\|\rho'\|_{\infty}$ ,  $\|\rho''\|_{\infty} < \infty$ . Also, let  $\bar{\rho}$  be a nonnegative twice differentiable function on  $\mathbb R$  with  $\overline{\rho} \equiv 1$  on  $(-\infty, -\delta)$ ,  $\overline{\rho} \equiv 0$  on  $(0, \infty)$ , and  $\|\overline{\rho}'\|_{\infty}, \|\overline{\rho}''\|_{\infty} < \infty$ . Define  $f(x, y) = x \rho(y)$  and  $\bar{f}(x, y) = x \bar{\rho}(y)$ . Set

$$
\Delta(x, z) = f(x, xz) - \frac{e^{\beta z + h} f(1, z) + e^{-\beta z - h} f(-1, -z)}{e^{\beta z + h} + e^{-\beta z - h}},
$$

$$
\bar{\Delta}(x, z) = \bar{f}(x, xz) - \frac{e^{\beta z + h} \bar{f}(1, z) + e^{-\beta z - h} \bar{f}(-1, -z)}{e^{\beta z + h} + e^{-\beta z - h}}.
$$

Our next lemma establishes some useful bounds for the difference on the left-hand side of  $(14)$  associated to f and f defined above.

<span id="page-9-0"></span>Lemma 2.3. We have that

$$
\Delta(1, z) \geqslant 0, \ \forall z \in \mathbb{R} \ and \ \Delta(1, z) \geqslant \frac{1}{1 + e^{2\beta M + 2h}}, \ \forall \delta \leqslant z \leqslant M,
$$
\n
$$
(15)
$$

$$
\Delta(-1, z) \geqslant 0, \ \forall z \in \mathbb{R} \ and \ \Delta(-1, z) \leqslant -\frac{1}{1 + e^{-2\beta\delta - 2h}}, \ \forall \delta \leqslant z \leqslant M,
$$

$$
\bar{\Delta}(1,z) \geqslant 0, \ \forall z \in \mathbb{R} \ and \ \bar{\Delta}(1,z) \geqslant \frac{1}{1 + e^{-2\beta\delta + 2h}}, \ \forall -M \leqslant z \leqslant -\delta,
$$
\n<sup>(17)</sup>

$$
\bar{\Delta}(-1, z) \leq 0, \ \forall z \in \mathbb{R} \ and \ \bar{\Delta}(-1, z) \leq -\frac{1}{1 + e^{2\beta M - 2h}}, \ \forall -M \leq z \leq -\delta. \tag{18}
$$

*Proof.* Computing directly yields that for any  $z \in \mathbb{R}$ ,

$$
\Delta(1, z) = \frac{\rho(z) + \rho(-z)}{1 + e^{2\beta z + 2h}} \ge 0, \quad \Delta(-1, z) = -\frac{\bar{\rho}(z) + \bar{\rho}(-z)}{1 + e^{-2\beta z - 2h}} \le 0,
$$
  

$$
\bar{\Delta}(1, z) = \frac{\bar{\rho}(z) + \bar{\rho}(-z)}{1 + e^{2\beta z + 2h}} \ge 0, \quad \bar{\Delta}(-1, z) = -\frac{\bar{\rho}(z) + \bar{\rho}(-z)}{1 + e^{-2\beta z - 2h}} \le 0.
$$

If  $\delta \leqslant z \leqslant M$ , then

$$
\Delta(1, z) = \frac{1}{1 + e^{2\beta z + 2h}} \ge \frac{1}{1 + e^{2\beta M + 2h}},
$$
  

$$
\Delta(-1, z) = -\frac{1}{1 + e^{-2\beta z - 2h}} \le -\frac{1}{1 + e^{-2\beta \delta - 2h}}.
$$

Also, if  $-M \leqslant z \leqslant -\delta$ ,

$$
\bar{\Delta}(1, z) = \frac{1}{1 + e^{2\beta z + 2h}} \ge \frac{1}{1 + e^{-2\beta \delta + 2h}},
$$
  

$$
\bar{\Delta}(-1, z) = -\frac{1}{1 + e^{-2\beta z - 2h}} \le -\frac{1}{1 + e^{2\beta M - 2h}}
$$

<span id="page-9-4"></span><span id="page-9-3"></span><span id="page-9-2"></span><span id="page-9-1"></span>.

■

For a given  $\sigma \in \{-1, +1\}^N$  and for  $M > \delta > 0$ , let

$$
I_{\delta,M}^+(\sigma) := \{ i : \delta \leq \sigma_i m_i(\sigma) \leq M \},
$$
  

$$
I_{\delta,M}^-(\sigma) := \{ i : -M \leq \sigma_i m_i(\sigma) \leq -\delta \}.
$$

The third lemma shows that if these two sets are linear large and  $\Delta(\sigma_r, \sigma_r m_r(\sigma))$  and  $\bar{\Delta}(\sigma_r, \sigma_r m_r(\sigma))$  for  $1 \leq r \leq N$  are small on average, then there exist four coordinates, on which the pair  $(\sigma_r, m_r(\sigma))$  must take all possible signs.

<span id="page-10-7"></span>**Lemma 2.4.** Assume that there exist some  $0 < \delta < M$  and  $\varepsilon > 0$  such that for some  $N \ge 4$ and  $\sigma \in \{-1, +1\}^N$ , the following inequalities are satisfied

$$
|I_{\delta,M}^+(\sigma)| \geq \varepsilon N, |I_{\delta,M}^-(\sigma)| \geq \varepsilon N,\tag{19}
$$

$$
\left|\frac{1}{N}\sum_{r=1}^{N}\Delta(\sigma_r,\sigma_r m_r(\sigma))\right| \leq \frac{\varepsilon}{2}\min\left(\frac{1}{1+e^{2\beta M+2h}},\frac{1}{1+e^{2\beta M-2h}}\right),\tag{20}
$$

$$
\left|\frac{1}{N}\sum_{r=1}^{N}\bar{\Delta}(\sigma_r,\sigma_r m_r(\sigma))\right| \leq \frac{\varepsilon}{2}\min\left(\frac{1}{1+e^{-2\beta\delta+2h}},\frac{1}{1+e^{-2\beta\delta-2h}}\right),\tag{21}
$$

where  $\Delta$  and  $\bar{\Delta}$  are defined in Lemma [2.3.](#page-9-0) Then there exist distinct i, j, k, l  $\in$  [N] such that

<span id="page-10-6"></span><span id="page-10-3"></span><span id="page-10-1"></span><span id="page-10-0"></span>
$$
\sigma_i = 1, \ \delta \leqslant m_i(\sigma) \leqslant M,\tag{22}
$$

<span id="page-10-4"></span><span id="page-10-2"></span>
$$
\sigma_j = -1, \ -M \leqslant m_j(\sigma) \leqslant -\delta,\tag{23}
$$

$$
\sigma_k = -1, \ \delta \leqslant m_k(\sigma) \leqslant M,\tag{24}
$$

<span id="page-10-5"></span>
$$
\sigma_l = 1, \ -M \le m_l(\sigma) \le -\delta. \tag{25}
$$

Proof. Since  $|I_{\delta,M}^+(\sigma)| \geq \varepsilon N$ , we readily see that either  $\sigma_i = 1, \delta \leq m_i(\sigma) \leq M$  or  $\sigma_j =$  $-1, -M \leqslant m_j(\sigma) \leqslant -\delta$  for some  $i, j$ . If  $\sigma_i = 1$  and  $\delta \leqslant m_i(\sigma) \leqslant M$  for all  $i \in I^+_{\delta,M}(\sigma)$ , then from [\(15\)](#page-9-1) and  $|I_{\delta,M}^+(\sigma)| \geq \varepsilon N$ , we arrive at

$$
\frac{1}{N}\sum_{r=1}^N \Delta(\sigma_r, \sigma_r m_r(\sigma)) \geq \frac{\varepsilon}{1 + e^{2\beta M + 2h}},
$$

contradicting [\(20\)](#page-10-0). If  $\sigma_j = -1$  and  $-M \leqslant m_j(\sigma) \leqslant -\delta$  for all  $j \in I^+_{\delta,M}(\sigma)$ , then from [\(18\)](#page-9-2) and  $|I^+_{\delta,M}(\sigma)| \geqslant \varepsilon N$ ,

$$
\frac{1}{N}\sum_{r=1}^N \bar{\Delta}(\sigma_r, \sigma_r m_r(\sigma)) \leq -\frac{\varepsilon}{1 + e^{2\beta M - 2h}},
$$

contradicting [\(20\)](#page-10-0) again. Hence, we see that there exist distinct i, j such that [\(22\)](#page-10-1) and [\(23\)](#page-10-2) hold. Similarly, the assumption  $|I_{\delta,M}^-(\sigma)| \geq \varepsilon N$  implies that either  $\delta_k = 1, -M \leqslant m_k(\sigma) \leqslant -\delta$ or  $\delta_l = -1, \delta \leqslant m_l(\sigma) \leqslant M$  for some k, l. If  $\sigma_k = 1$  and  $-M \leqslant m_k(\sigma) \leqslant -\delta$  for all  $k \in I_{\delta,M}^-(\sigma)$ , then from [\(17\)](#page-9-3) and  $|I_{\delta,M}^-(\sigma)| \geq \varepsilon N$ 

$$
\frac{1}{N} \sum_{r=1}^{N} \bar{\Delta}(\sigma_r, \sigma_r m_r(\sigma)) \geq \frac{\varepsilon}{1 + e^{-2\beta\delta + 2h}},
$$

contradicting [\(21\)](#page-10-3). Finally, if  $\sigma_l = -1$  and  $\delta \leqslant m_l(\sigma) \leqslant M$  for all  $l \in I_{\delta,M}^-(\sigma)$ , then from [\(16\)](#page-9-4) and  $|I_{\delta,M}(\sigma)| \geq \varepsilon N$ 

$$
\frac{1}{N}\sum_{r=1}^N \Delta(\sigma_r, \sigma_r m_r(\sigma)) \leq -\frac{\varepsilon}{1 + e^{-2\beta\delta - 2h}},
$$

contradicting  $(21)$ . Therefore, there must exist distinct k, l such that  $(24)$  and  $(25)$  hold. This completes our proof.

We have seen that from Lemma [2.1,](#page-7-0) the inequalities  $(20)$  and  $(21)$  hold with high probability as  $N$  tends to infinity. Our next goal will be to validate the assumption  $(19)$ , which relies on the following lemma.

<span id="page-11-0"></span>**Lemma 2.5** (Positivity of the Hamiltonian). Let  $\tau \sim P_{\beta,h}(\cdot)$  for some  $\beta > 0$  and  $h \in \mathbb{R}$ . If

$$
\liminf_{N \to \infty} (F_N(\beta', h) - F_N(0, h)) > 0
$$

for some  $0 < \beta' < \beta$ , then there exists some  $\delta > 0$  such that

$$
\limsup_{N \to \infty} \frac{1}{N} \log P_{\beta,h}\left(\frac{1}{N} \sum_{i=1}^N \tau_i m_i(\tau) < \delta\right) < 0.
$$

*Proof.* For any  $\delta > 0$ , by the Markov inequality,

$$
P_{\beta,h}\left(\frac{1}{N}\sum_{i=1}^{N}\tau_{i}m_{i}(\tau)<\delta\right)\leq e^{\lambda N\delta}\mathbb{E}_{\beta,h}\left(\exp\left(-\lambda\sum_{i=1}^{N}\tau_{i}m_{i}(\tau)\right)\right)
$$

$$
=e^{\lambda N\delta}Z_{N}(\beta-\lambda,h)(Z_{N}(\beta,h))^{-1}
$$

$$
=e^{\lambda N\delta+N(F_{N}(\beta-\lambda,h)-F_{N}(\beta,h))}.
$$

Using the convexity of  $F_N$  in  $\beta$ , we have that for  $\lambda \in (0, \beta)$ ,

$$
F_N(\beta, h) - F_N(\beta - \lambda, h) \geq \frac{\partial}{\partial \beta} F_N(\beta - \lambda, h) \lambda \geq \frac{F_N(\beta - \lambda, h) - F_N(0, h)}{\beta - \lambda} \lambda.
$$

Form the given assumption, letting  $\delta = 2^{-1} \liminf_{N \to \infty} (F_N(\beta', h) - F_N(0, h))$  and  $\lambda = \beta - \beta'$ completes our proof.

Finally, we show that the assumption [\(19\)](#page-10-6) holds with high probability.

<span id="page-11-1"></span>**Lemma 2.6.** Suppose  $\tau \sim P_{\beta,h}(\cdot)$  for  $\beta > 0, h \in \mathbb{R}$  and the condition (i) in Theorem [1.1](#page-3-0) holds, then there exist  $\varepsilon > 0$  and  $0 < \delta < M$  such that w.h.p., the following inequalities hold,

$$
|I_{\delta,M}^+(\tau)|\geqslant \varepsilon N\ and\ |I_{\delta,M}^-(\tau)|\geqslant \varepsilon N.
$$

*Proof.* Let us begin with the following notations. For a given  $\sigma$  and  $0 < \delta < M$ , let

$$
I_M(\sigma) := \{ i \in [N] : |m_i(\sigma)| \leqslant M \}
$$

and

$$
I^+(\sigma) := \{i \in [N] : \sigma_i m_i(\sigma) > 0\}, \ I^-(\sigma) := \{i \in [N] : \sigma_i m_i(\sigma) < 0\},
$$
  

$$
I^+_{\delta}(\sigma) := \{i \in [N] : \sigma_i m_i(\sigma) \geq \delta\}, \ I^-_{\delta}(\sigma) := \{i \in [N] : \sigma_i m_i(\sigma) \leq -\delta\}.
$$

We will abbreviate the above notations by  $I_M$ ,  $I^+, I^-, I^+_\delta, I^-_\delta$  when the configuration is the sample  $\tau$ . By Markov inequality, we readily have

<span id="page-12-2"></span><span id="page-12-0"></span>
$$
|I_M^c| \leq \frac{\sum_{i \in [N]} m_i(\tau)^2}{M^2} \leq \frac{\|J\|^2 N}{M^2}.
$$
 (26)

From [\(26\)](#page-12-0) and the Cauchy-Schwarz inequality,

$$
\sum_{i \in I_M^c} |m_i(\tau)| \leq |I_M^c|^{1/2} \Big(\sum_{i=1}^N |m_i(\tau)|^2\Big)^{1/2} \leq |I_M^c|^{1/2} \|J\| \sqrt{N} \leq \frac{N \|J\|^2}{M}.\tag{27}
$$

Combining [\(26\)](#page-12-0) with Lemma [2.5,](#page-11-0) there exist  $\varepsilon > 0$  and  $0 < \delta < M$  such that w.h.p., we have  $|I^+_{\delta,M}(\tau)| \geqslant \varepsilon N.$ 

Next, we focus on  $I_{\delta,M}^-(\tau)$ . Notice that

$$
S(\beta, h|\tau) = \sum_{i \in [N]} m_i(\tau)(\tau_i - \tanh(\beta m_i(\tau) + h))
$$
  
= 
$$
\sum_{i \in I_M^c} m_i(\tau)(\tau_i - \tanh(\beta m_i(\tau) + h))
$$
 (28)

<span id="page-12-1"></span>
$$
+\sum_{i\in I_M\cap I^+} m_i(\tau)(\tau_i-\tanh(\beta m_i(\tau)+h))\tag{29}
$$

<span id="page-12-4"></span><span id="page-12-3"></span>
$$
+\sum_{i\in I_M\cap I^-} m_i(\tau)(\tau_i-\tanh(\beta m_i(\tau)+h)).\tag{30}
$$

For [\(28\)](#page-12-1), using [\(27\)](#page-12-2), for any  $\varepsilon' > 0$ , one can choose M sufficiently such that

$$
\sum_{i \in I_M} m_i(\tau) (\tau_i - \tanh(\beta m_i(\tau) + h)) \leqslant \varepsilon' N.
$$

Now for  $(29)$ , we have

$$
\sum_{i \in I_M \cap I^+} m_i(\tau)(\tau_i - \tanh(\beta m_i(\tau) + h))
$$
\n
$$
\geqslant \sum_{i \in I_M \cap I^+} m_i(\tau)\tau_i(1 - \tanh(|\beta m_i(\tau)| + |h|))
$$
\n
$$
\geqslant c \Big( \sum_{i \in I^+} m_i(\tau)\tau_i - \sum_{i \in I_M^c} |m_i(\tau)| \Big),
$$

where  $c = 1 - \tanh(|\beta|M + |h|) > 0$ . By Lemma [2.5,](#page-11-0) we have w.h.p.,  $\sum_{i \in I^+} m_i(\tau) \tau_i \geqslant 2\eta N$ for some  $\eta > 0$ . Using [\(27\)](#page-12-2), one can make  $\sum_{i \in I_M} |m_i(\tau)| \leq \eta N$  by choosing M large enough. Therefore,  $(29)$  is bounded from below by  $\eta N$  w.h.p.

Finally, for  $\delta > 0$ , we write [\(30\)](#page-12-4) as

$$
\sum_{i \in I_M \cap I^-} m_i(\tau)(\tau_i - \tanh(\beta m_i(\tau) + h))
$$
  
= 
$$
\sum_{i \in I_{\delta,M}^-(\tau)} m_i(\tau)(\tau_i - \tanh(\beta m_i(\tau) + h))
$$
  
+ 
$$
\sum_{i \in I_M : -\delta \leq \tau_i m_i(\tau) < 0} m_i(\tau)(\tau_i - \tanh(\beta m_i(\tau) + h)).
$$

Since  $|\tau_i - \tanh(\beta m_i(\tau) + h)| \leq 2$ , it follows that

$$
\sum_{i \in I_{\delta,M}^-(\tau)} m_i(\tau) (\tau_i - \tanh(\beta m_i(\tau) + h)) \geq -2M|I_{\delta,M}^-(\tau)|.
$$

Similarly,

$$
\sum_{i\in I_M:-\delta\leqslant\tau_im_i(\tau)<0}m_i(\tau)(\tau_i-\tanh(\beta m_i(\tau)+h))\geqslant-2\delta N.
$$

Thus

$$
\sum_{i\in I_M\cap I^-} m_i(\tau)(\tau_i-\tanh(\beta m_i(\tau)+h))\geqslant -2M|I_{\delta,M}^-(\tau)|-2\delta N.
$$

Putting the bounds for all three terms together with  $\varepsilon' = \eta/4$  and  $\delta = \eta/8$ , we have w.h.p.

$$
S(\beta, h|\tau) \geqslant \frac{\eta}{2}N - 2M|I_{\delta, M}^{-}|.
$$

It now follows from the concentration bound of  $S(\beta, h|\tau)$  established in Corollary [2.2](#page-7-1) that w.h.p.,

$$
|I_{\delta,M}^-(\tau)| \geqslant \frac{\eta}{4M}N.
$$

<span id="page-13-0"></span>2.2. Proof of Theorem [1.1:](#page-3-0) Existence of the MPLE. Before we turn to the proof for the existence part of the MPLE in Theorem [1.1,](#page-3-0) we establish the following lemma first.

<span id="page-13-2"></span>**Lemma 2.7.** Let  $\tau \in \{-1, +1\}^N$  be a sample from the unknown ground truth  $P_{\beta_0, h_0}$ . The maximum pseudolikelihood estimator  $(\hat{\beta}(\tau), \hat{h}(\tau))$  exists if  $\tau$  satisfies the following property: there exist distinct i, j, k,  $l \in [N]$  such that

$$
\tau_i = 1, m_i(\tau) > 0, \ \tau_j = -1, m_j(\tau) < 0, \n\tau_k = -1, m_k(\tau) > 0, \ \tau_l = 1, m_l(\tau) < 0.
$$
\n(31)

 $\setminus$ 

<span id="page-13-1"></span>■

*Proof.* Note that if  $\tau$  satisfies [\(31\)](#page-13-1), then obviously  $\widetilde{T}_N(\tau) > 0$ . From [\(8\)](#page-3-1), the pseudolikelihood function is strictly concave on  $\mathbb{R}^2$ . Thus, to prove the existence of the maximum pseudolikelihood estimator, it suffices to show that

$$
\limsup_{|\beta|+|h|\to\infty} L(\beta,h|\tau) = -\infty.
$$

To this end, noting that

$$
L(\beta, h|\tau) + \log 2 = \sum_{i=1}^{N} \left( \tau_i(\beta m_i(\tau) + h) - \log \cosh(\beta m_i(\tau) + h) \right)
$$
  

$$
\leq \sum_{i=1}^{N} \left( |\beta m_i(\tau) + h| - \log \cosh(\beta m_i(\tau) + h) \right)
$$

and each term in the above sum is nonpositive, it is enough to show that given  $\tau$ , for any  $K > 0$ , there exists some  $C > 0$  such that the following statement holds: for any  $(\beta, h) \in \mathbb{R}^2$ with  $|\beta| + |h| \geq C$ , there exists some  $a = a(\beta, h) \in [N]$  such that

$$
\tau_a(\beta m_a(\tau) + h) - \log \cosh(\beta m_a(\tau) + h) \leqslant -K.
$$

Assume that  $i, j, k$ , and l satisfy [\(31\)](#page-13-1). For each  $(\beta, h) \in \mathbb{R}^2$ , we select a according to the following rules. First, let  $\eta = \min_{a \in \{i,j,k,l\}} |m_a(\tau)| > 0$ .

• If  $\beta, h \neq 0$ , let  $a \in \{i, j, k, l\}$  be such that

$$
sign(m_a(\tau)) = sign(\beta h)
$$
 and  $\tau_a = -sign(h)$ .

Then

$$
\tau_a(\beta m_a(\tau) + h) = -|\beta m_a(\tau)| - |h| \leq -|\beta|\eta - |h|.
$$

- If  $\beta \neq 0, h = 0$ , let  $a \in \{i, j, k, l\}$  be such that  $\tau_a = -\text{sign}(\beta m_a(\tau))$ . Then  $\tau_a(\beta m_a(\tau) + h) = - |\beta| |m_a(\tau)| \leq -|\beta| \eta$ .
- If  $\beta = 0, h \neq 0$ , similarly let  $a \in \{i, j, k, l\}$  with  $\tau_a = -\text{sign}(h)$ . Then

$$
\tau_a(\beta m_a(\tau) + h) = - |h|.
$$

From these and noting that  $\log \cosh(x) \geq 0$  for all  $x \in \mathbb{R}$ , we readily see that for any  $(\beta, h) \in$  $\mathbb{R}^2$ 

$$
\tau_a(\beta m_a(\tau) + h) - \log \cosh(\beta m_a(\tau) + h) \leq \tau_a(\beta m_a(\tau) + h) \leq -\max(|h|, |\beta|\eta).
$$

Now for each  $K > 0$ , our proof is completed by letting  $C = K(1/\eta + 1)$ .

We now turn to the proof for the existence of the MPLE in Theorem [1.1.](#page-3-0) From Lemma [2.7,](#page-13-2) we only need to verify that w.h.p., the sample  $\tau \sim P_{\beta_0, h_0}$  satisfies [\(31\)](#page-13-1). To do this, we will check that w.h.p., the assumptions in Lemma [2.4](#page-10-7) are satisfied. Now, from Lemma [2.6,](#page-11-1) we already proved that there exist  $\varepsilon > 0$  and  $0 < \delta < M$  such that w.h.p.,

$$
|I_{\delta,M}^+(\tau)|,|I_{\delta,M}^-(\tau)|\geq \varepsilon N.
$$

On the other hand, by Lemma [2.1](#page-7-0) with  $f(x, y) = x\rho(y)$  and  $f(x, y) = x\rho(y)$  and with the operator norm bound in the condition (ii) stated in Theorem [1.1,](#page-3-0) we have

$$
\left\langle \left( \sum_{r=1}^{N} \Delta(\tau_r, \tau_r m_r(\tau)) \right)^2 \right\rangle \leq CN, \quad \left\langle \left( \sum_{r=1}^{N} \bar{\Delta}(\tau_r, \tau_r m_r(\tau)) \right)^2 \right\rangle \leq CN
$$

for some constant C. Thus, it follows that the assumptions  $(20)$  and  $(21)$  of Lemma [2.4](#page-10-7) are satisfied w.h.p., validating the conclusion of Lemma [2.4,](#page-10-7) which implies [\(31\)](#page-13-1) and completes our proof.

<span id="page-14-0"></span>2.3. Proof of Theorem [1.1:](#page-3-0) Consistency of MPLE. We begin with a simple lemma:

<span id="page-14-1"></span>**Lemma 2.8.** Fix  $\beta > 0$  and  $h \in \mathbb{R}$ . Let  $\sigma \in \{-1, +1\}^N$ . Suppose that  $||J|| = O(1)$  and  $\widetilde{T}_{N}(\sigma) \geqslant c$ . Then there exists a constant  $M > 0$ , which depends only on c,  $\beta$ , h, and ||J|| such that

$$
\frac{1}{N^2} \sum_{i,j \in I_M(\sigma)} (m_i(\sigma) - m_j(\sigma))^2 \geqslant \frac{c}{2},
$$

where  $I_M(\sigma) = \{i \in [N] : |m_i(\sigma)| \leq M\}.$ 

*Proof.* Let M be sufficiently large such that  $\beta M > |h|$ . We have

$$
\frac{1}{N^2} \sum_{i,j \in I_M(\sigma)} (m_i(\sigma) - m_j(\sigma))^2 \ge \frac{1}{N^2} \sum_{i,j \in I_M(\sigma)} \theta_i(\sigma) \theta_j(\sigma) (m_i(\sigma) - m_j(\sigma))^2
$$
  

$$
\ge c - \frac{1}{N^2} \sum_{\{i,j\} \cap I_M^c(\sigma) \neq \emptyset} \theta_i(\sigma) \theta_j(\sigma) (m_i(\sigma) - m_j(\sigma))^2
$$
  

$$
\ge c - \frac{\text{sech}^2(\beta M - |h|)}{N^2} \sum_{i,j} (m_i(\sigma) - m_j(\sigma))^2.
$$

On the other hand,

$$
\frac{1}{N^2} \sum_{i,j} (m_i(\sigma) - m_j(\sigma))^2 \leq \frac{2}{N} \sum_i m_i(\sigma)^2 \leq 2||J||^2.
$$

Combining these two estimates, the lemma follows that if we choose  $M$  large enough to satisfy  $2 \operatorname{sech}^2(\beta M - |h|) \|J\|$  $2 \leqslant c/2$ .

The proof for the consistency of MPLE in Theorem [1.1](#page-3-0) proceeds as follows. For notational clarity, we will make the dependence of the quantities  $\theta_i$  and  $T_N$  on parameters  $\beta$  and h explicit throughout the proof. Let  $\tau \sim P_{\beta_0,h_0}$ . For  $(\beta,h) \in \mathbb{R}^2$ , let  $\lambda_1(\beta,h|\tau) \geq 0$  be the minimum eigenvalue of the negative Hessian matrix,  $H(\beta, h|\tau)$ . Note that

$$
\lambda_1(\beta, h|\tau) \geq \frac{|H(\beta, h|\tau)|}{\text{Tr}(H(\beta, h|\tau))} = \frac{\frac{1}{2} \sum_{i,j=1}^N \theta_i(\beta, h|\tau) \theta_j(\beta, h|\tau) (m_i(\tau) - m_j(\tau))^2}{\text{Tr}(H(\beta, h|\tau))},\tag{32}
$$

where  $\theta_i(\beta, h|\tau) = \text{sech}^2(\beta m_i(\tau) + h)$ . To bound the right hand side of [\(32\)](#page-15-0) from below, we note

<span id="page-15-0"></span>
$$
\sum_{i,j=1}^{N} \theta_i(\beta, h|\tau) \theta_j(\beta, h|\tau) (m_i(\tau) - m_j(\tau))^2
$$
  
\n
$$
\ge \operatorname{sech}^4(\beta M + |h|) \sum_{i,j \in I_M(\tau)} (m_i(\tau) - m_j(\tau))^2
$$

for any  $M > 0$ . Also, note that since  $\theta_i(\beta, h|\tau) \leq 1$  and  $\sum_{i=1}^N m_i^2(\tau) \leq ||J||^2 N$ ,

$$
\text{Tr}(H(\beta, h|\tau)) = \sum_{i=1}^{N} \theta_i(\beta, h|\tau) (m_i^2(\tau) + 1) \le N(1 + C^2).
$$

Therefore, we obtain

$$
\lambda_1(\beta, h|\tau) \ge \frac{\text{sech}^4(\beta M + |h|)}{2(1 + C^2)N} \sum_{i,j \in I_M(\tau)} (m_i(\tau) - m_j(\tau))^2.
$$
 (33)

This yields the following locally uniform lower bound for  $\lambda_1(\beta, h|\tau)$  that for any  $M > 0$ , there exist some positive  $r, \eta > 0$  such that

$$
\inf_{\|\beta-\beta_0,h-h_0\|\leqslant r} \lambda_1(\beta,h|\tau) \geqslant \frac{\eta}{N} \sum_{i,j\in I_M(\tau)} (m_i(\tau)-m_j(\tau))^2.
$$
 (34)

Next, we construct an interpolation between the pseudolikelihood estimator and the ground truth by letting

<span id="page-15-1"></span>
$$
\beta(t) = t\hat{\beta} + (1-t)\beta_0
$$
 and  $h(t) = t\hat{h} + (1-t)h_0$ 

for  $t \in [0, 1]$ . Set  $f : [0, 1] \rightarrow \mathbb{R}$  as

$$
f(t) = (\hat{\beta} - \beta_0)S(\beta(t), h(t)|\tau) + (\hat{h} - h_0)Q(\beta(t), h(t)|\tau).
$$

Since  $Q(\hat{\beta}, \hat{h}|\tau) = 0$  and  $R(\hat{\beta}, \hat{h}|\tau) = 0$ , it is clear

$$
|f(1) - f(0)| = |(\hat{\beta} - \beta_0)S(\beta_0, h_0|\tau) + (\hat{h} - h_0)Q(\beta_0, h_0|\tau)|.
$$

By Corollary [2.2,](#page-7-1) we have

$$
S(\beta_0, h_0|\tau), Q(\beta_0, h_0|\tau) = O_p(\sqrt{N}).
$$

Applying the Cauchy-Schwarz inequality, we deduce

<span id="page-16-1"></span>
$$
|f(1) - f(0)| = O_p(\sqrt{N}Y_N),
$$
\n(35)

where  $Y_N := ||(\hat{\beta} - \beta_0, \hat{h} - h_0)||_2$ .

Now to bound  $Y_N$ , we need to control  $|f(1) - f(0)|$ . For any  $t \in [0,1]$ , the derivative of f is bounded from below by

$$
f'(t) = (\hat{\beta} - \beta_0, \hat{h} - h_0)H(\beta(t), h(t)|\tau)(\hat{\beta} - \beta_0, \hat{h} - h_0)^T
$$
  
\n
$$
\geq \lambda_1(\beta(t), h(t)|\tau)Y_N^2.
$$

From the fact  $\|(\beta(t) - \beta_0, h(t) - h_0)\|_2 = tY_N$  and the uniform lower bound [\(34\)](#page-15-1), we obtain

$$
f(1) - f(0) = \int_0^1 f'(t)dt \ge \int_0^{\min\left(1, \frac{r}{Y_N}\right)} f'(t)dt
$$
  
 
$$
\ge \min\left(1, \frac{r}{Y_N}\right) \frac{\eta Y_N^2}{N} \sum_{i,j \in I_M(\tau)} (m_i(\tau) - m_j(\tau))^2.
$$
 (36)

From the assumption (iii) of Theorem [1.1,](#page-3-0) it follows that given  $\varepsilon > 0$ , there exists  $c > 0$  such that with  $P_{\beta_0,h_0}$ -probability at least  $1-\varepsilon/2$ , we have  $\widetilde{T}_N(\beta_0,h_0|\tau) \geq c$ . Furthermore, thanks to Lemma  $2.8$ , we can find  $M$  such that on the same event,

$$
\frac{1}{N^2} \sum_{i,j \in I_M(\tau)} (m_i(\tau) - m_j(\tau))^2 \geqslant \frac{c}{2}.
$$

On the other hand,  $(35)$  implies that there exists C such that

<span id="page-16-2"></span>
$$
|f(1) - f(0)| \leqslant C\sqrt{N}Y_N
$$

with  $P_{\beta_0,h_0}$ -probability at least  $1-\varepsilon/2$ . Since both of the conditions hold simultaneously with probability at least  $1 - \varepsilon$ , it now follows from [\(36\)](#page-16-2) that with probability at least  $1 - \varepsilon$ ,

$$
\min(Y_N, r) \leqslant \frac{2C}{\eta c \sqrt{N}}.
$$

Since r is independent of N, it follows that  $Y_N = O_p(q)$ N) as claimed.

3. Proof of Theorem [1.4](#page-4-2)

<span id="page-16-0"></span>To prove the Theorem [1.4,](#page-4-2) we first establish the following universality result in Lemma [3.1,](#page-16-3) which shows that the free energies associated with the subgaussian matrix  $G = (g_{ij})_{1 \leq i,j \leq N}$ is close to that associated to the standard Gaussian disorder. For any  $x = (x_e)_{e \in E}$ , set

$$
F_x(\beta, h) := \frac{1}{N} \ln \sum_{\sigma \in \{-1, +1\}^N} e^{\beta H_x(\sigma) + h \sum_{i=1}^N \sigma_i},
$$

where

$$
H_x(\sigma) := \frac{1}{\sqrt{d}} \sum_{(i,j) \in E} x_{ij} \sigma_i \sigma_j.
$$

Note that  $F_q(\beta, h) = F_N(\beta, h)$ . Let  $z = (z_e)_{e \in E}$  be i.i.d. standard normal independent of g.

<span id="page-16-3"></span>**Lemma 3.1.** There exists a universal constant C independent of  $N, d, \beta, h$  such that

$$
|\mathbb{E} F_g(\beta, h) - \mathbb{E} F_z(\beta, h)| \leqslant \frac{C\beta^3}{\sqrt{d}}.
$$

*Proof.* For  $0 \leq t \leq 1$ , define

$$
\phi(t) = \frac{1}{N} \mathbb{E} \ln e^{\beta H_{N,t}(\sigma)},
$$

where

$$
H_{N,t}(\sigma) := \sqrt{t}H_g(\sigma) + \sqrt{1-t}H_z(\sigma) + h\sum_{i=1}^N \sigma_i.
$$

Then

$$
\phi'(t) = \frac{\beta}{2N} \Big( \frac{\mathbb{E} \langle H_g(\sigma) \rangle_t}{\sqrt{t}} - \frac{\mathbb{E} \langle H_z(\sigma) \rangle_t}{\sqrt{1-t}} \Big) = \frac{\beta}{2N\sqrt{d}} \mathbb{E} \Big[ \sum_{(i,j) \in E} \frac{\langle g_{ij} \sigma_i \sigma_j \rangle_t}{\sqrt{t}} - \sum_{(i,j) \in E} \frac{\langle z_{ij} \sigma_i \sigma_j \rangle_t}{\sqrt{1-t}} \Big].
$$

Here from the Gaussian integration by parts in the Gaussian disorder,

$$
\frac{\mathbb{E}\langle z_{ij}\sigma_i\sigma_j\rangle_t}{\sqrt{1-t}} = \frac{\beta}{\sqrt{d}} \mathbb{E}\langle 1-\sigma_i^1\sigma_j^1\sigma_i^2\sigma_j^2\rangle_t.
$$

On the other hand, from the approximate Gaussian integration by parts in the disorder  $g$ (see, for example,  $[7]$ ), there exists an absolute constant C independent of anything else such that

$$
\Big|\frac{\mathbb{E}\langle g_{ij}\sigma_i\sigma_j\rangle_t}{\sqrt{t}}-\frac{\beta}{\sqrt{d}}\,\mathbb{E}\big\langle1-\sigma_i^1\sigma_j^1\sigma_i^2\sigma_j^2\big\rangle_t\Big|\leqslant \frac{C\beta^2}{d}.
$$

Consequently,

$$
|\phi'(t)|\leqslant \frac{C\beta^3|E|}{2d^{3/2}N}=\frac{C\beta^3}{\sqrt{d}}
$$

and this completes our proof.

We also establish the following concentration result for the free energy  $F_N(\beta, h)$  for later use.

<span id="page-17-0"></span>**Proposition 3.2.** Assume that  $(g_e)_{e \in E}$  is a collection of independent  $\kappa^2$ -subgaussian random variables. For  $t > 0$ , we have

$$
\mathbb{P}\big(|F_N(\beta,h)-\mathbb{E}\, F_N(\beta,h)|\geqslant t\big)\leqslant 2\exp\Big(-\min\Big(\frac{(\ln 2)Nt^2}{2\beta^2\kappa^2},\frac{\sqrt{d\ln 2}Nt}{2\beta\kappa}\Big)\Big).
$$

*Proof.* Denote  $f_N(\beta, h) := NF_N(\beta, h)$ . Let  $(g_i)_{1 \leq i \leq |E|}$  be an enumeration of  $(g_e)_{e \in E}$ . We introduce the following filtration  $\mathcal{F}_k = \sigma(g_k : 1 \leq k \leq |E|)$  for  $1 \leq k \leq |E|$  and  $\mathcal{F}_0 = \emptyset$ . For each  $1 \leq k \leq |E|$ , define the martingale difference

$$
X_k := \mathbb{E}\big[f_N(\beta,h)\big|\mathcal{F}_k\big] - \mathbb{E}\big[f_N(\beta,h)\big|\mathcal{F}_{k-1}\big].
$$

Then  $\mathbb{E}[X_k|\mathcal{F}_{k-1}] = 0$  and

$$
\sum_{k=1}^{|E|} X_k = f_N(\beta, h) - \mathbb{E}\big[f_N(\beta, h)\big].
$$

Fix k. Let  $\tilde{f}_N(\beta, h)$  be the same as  $f_N(\beta, h)$  with  $g_k = 0$  and  $\langle \cdot \rangle_k$  is the corresponding Gibbs expectation. Then

$$
X_k = \mathbb{E}\left[f_N(\beta, h) - \tilde{f}_N(\beta, h)\big|\mathcal{F}_k\right] - \mathbb{E}\left[f_N(\beta, h) - \tilde{f}_N(\beta, h)\big|\mathcal{F}_{k-1}\right]
$$
  
\n
$$
= \mathbb{E}\left[\ln\left\langle\exp\left(\beta g_k \sigma_e/\sqrt{d}\right)\right\rangle_k \big|\mathcal{F}_k\right] - \mathbb{E}\left[\ln\left\langle\exp\left(\beta g_k \sigma_e/\sqrt{d}\right)\right\rangle_k \big|\mathcal{F}_{k-1}\right]
$$
  
\n
$$
\leq \beta d^{-1/2} \mathbb{E}\left[|g_k|\big|\mathcal{F}_k\right] + \beta d^{-1/2} \mathbb{E}\left[|g_k|\big|\mathcal{F}_{k-1}\right]
$$
  
\n
$$
\leq \frac{\beta|g_k|}{\sqrt{d}} + \frac{\beta \mathbb{E}|g_k|}{\sqrt{d}},
$$

where  $\sigma_e = \sigma_i \sigma_j$  and  $e = (i, j)$  is the k-th edge in the enumeration. From the fact that  $g_k$  is  $\kappa^2$ -subgaussian and Jensen's inequality, we have

$$
\mathbb{E}\left[e^{\frac{|X_k|}{A}}\big|\mathcal{F}_{k-1}\right] \leq \mathbb{E}\exp\left(\frac{\beta|g_k|}{A\sqrt{d}} + \frac{\beta \mathbb{E}|g_k|}{A\sqrt{d}}\right)
$$
  

$$
\leq \exp\left(\frac{\beta^2\kappa^2}{2A^2d} + \frac{\beta \mathbb{E}|g_k|}{A\sqrt{d}}\right)
$$
  

$$
\leq \exp\left(\frac{\beta^2\kappa^2}{2A^2d}\right) \mathbb{E}\exp\left(\frac{\beta|g_k|}{A\sqrt{d}}\right) = e^{\frac{\beta^2\kappa^2}{A^2d}} = e^{\ln 2} = 2,
$$

provided

$$
A^2 \equiv \frac{1}{\ln 2} \cdot \frac{\beta^2 \kappa^2}{d}.
$$

Hence, from Bernstein's inequality,

$$
\mathbb{P}\Big(\sum_{k=1}^{|E|} X_k \geq t\Big) \leqslant \exp\Big(-\min\Big(\frac{t^2}{4|E|A^2}, \frac{t}{2A}\Big)\Big).
$$

and thus, using  $|E| = 2^{-1} \sum_{i \in [N]} d_i = 2^{-1} N d$ ,

$$
\mathbb{P}\left(\sum_{k=1}^{|E|} X_k \ge Nt\right) \le \exp\left(-\min\left(\frac{N^2t^2}{4|E|A^2}, \frac{Nt}{2A}\right)\right)
$$

$$
= \exp\left(-\min\left(\frac{(\ln 2)dN^2t^2}{4\beta^2\kappa^2|E|}, \frac{\sqrt{d\ln 2}Nt}{2\beta\kappa}\right)\right)
$$

$$
= \exp\left(-\min\left(\frac{(\ln 2)Nt^2}{2\beta^2\kappa^2}, \frac{\sqrt{d\ln 2}Nt}{2\beta\kappa}\right)\right).
$$

Proof of Theorem [1.4.](#page-4-2) From the Gaussian integration by parts, we have

$$
\partial_{\beta} \mathbb{E} F_z(\beta, h) = \frac{\beta}{N} \mathbb{E} \langle H_z(\sigma) \rangle \n= \frac{\beta^2}{Nd} \Big( |E| - \sum_{(i,j) \in E} \mathbb{E} \langle \sigma_i^1 \sigma_j^1 \sigma_i^2 \sigma_j^2 \rangle \Big) = \frac{\beta^2}{2} \Big( 1 - \frac{2}{Nd} \sum_{(i,j) \in E} \mathbb{E} \langle \sigma_i^1 \sigma_j^1 \sigma_i^2 \sigma_j^2 \rangle \Big),
$$

where by a slight abuse of notation,  $\langle \cdot \rangle$  stands for the Gibbs expectation associated to the free energy  $F_z(\beta, h)$ . Under the given assumption, we could have the following result formally

■

stated and proved in Proposition [5.8.](#page-25-0) It implies that there exist positive  $M$  and  $c$  such that for any  $N \geq 1$  and  $\sigma \sim P_{\beta,h}$ , if we define  $U = U(\sigma) \subseteq [N]$  as

$$
U = \{i : |m_i(\sigma)| \leq M, \ d_i \geq d/4\},\
$$

then w.h.p.,  $|U| \geqslant cn$ .

Notice that

$$
\langle \sigma_i \sigma_j \rangle = \langle \tanh(\beta m_i(\sigma) + h) \sigma_j \rangle.
$$

We have

$$
\langle \sigma_i \sigma_j \rangle = \langle \tanh(\beta m_i(\sigma) + h) \sigma_j; i \in U \rangle + \langle \tanh(\beta m_i(\sigma) + h) \sigma_j; i \in U^c \rangle
$$
  

$$
\leq \tanh(\beta M + h) \langle I(i \in U) \rangle + \langle I(i \in U^c) \rangle
$$
  

$$
= 1 - (1 - \tanh(\beta M + h)) \langle I(i \in U) \rangle.
$$

From the definition of  $U$ , it follows that w.h.p.,

$$
\sum_{(i,j)\in E} I(i\in U) \geqslant \frac{1}{2}\sum_{i\in U}d_i \geqslant \frac{cdN}{8}.
$$

Therefore, we have w.h.p.,

$$
\frac{2}{Nd} \sum_{(i,j)\in E} \langle \sigma_i \sigma_j \rangle \leq \frac{2}{Nd} \Big( |E| - (1 - \tanh(\beta M + |h|)) \Big( \sum_{(i,j)\in E} I(i \in U) \Big) \Big)
$$
  

$$
\leq 1 - (1 - \tanh(\beta M)) \frac{2}{Nd} \cdot \frac{cdN}{8} = 1 - (1 - \tanh(\beta M)) \frac{c}{4},
$$

which implies that there exists a constant  $c' > 0$  such that

<span id="page-19-1"></span>
$$
\partial_{\beta} \mathbb{E} F_z(\beta, h) \geq \beta^2 c' > 0.
$$

Hence,

$$
\partial_{\beta} \mathbb{E} F_g(\beta, h) \geqslant \beta^2 c' - \frac{C\beta^3}{\sqrt{d}}.
$$
\n(37)

Finally, from the concentration of the free energy in the Proposition [3.2](#page-17-0) and the Lemma [3.1,](#page-16-3) we have

$$
\liminf_{N \to \infty} (F_g(\beta, h) - F_g(0, h)) = \liminf_{N \to \infty} (\mathbb{E} F_z(\beta, h) - \mathbb{E} F_z(0, h))
$$

$$
= \liminf_{N \to \infty} \int_0^{\beta} \partial_t \mathbb{E} F_z(t, h) dt
$$

$$
\geq \liminf_{N \to \infty} (\beta - \beta_0) \partial_t \mathbb{E} F_z(\beta_0, h) > 0
$$

for  $\beta_0 < \beta$ , where the last inequality used the fact that  $\partial_t \mathbb{E} F_z(t, h)$  is a nondecreasing function in t as well as  $(37)$ . The asserted exponential tail probability is guaranteed by Proposition [3.2.](#page-17-0) ■

## 4. Proof of Theorem [1.5](#page-4-3)

<span id="page-19-0"></span>We first establish the bound for the expected operator norm of  $J = \frac{1}{\sqrt{2}}$  $\frac{1}{d}A \circ G$ . It relies on the following fundamental result in [\[2,](#page-27-5) Corollary 3.3], which we recall as follows.

**Lemma 4.1** ([\[2,](#page-27-5) Corollary 3.3]). Consider the  $N \times N$  symmetric random matrix X whose entries  $X_{ij} := b_{ij} g_{ij}$  for  $i \geq j$ , where  $b_{ij} \in \mathbb{R}$  and  $g_{ij}$  are independent centered subgaussian. Then

$$
\mathbb{E} \left\| X \right\| \lesssim \max_{i} \sqrt{\sum_{j} b_{ij}^2} + \sqrt{\log N} \cdot \max_{ij} |b_{ij}|.
$$

Appling this result in our setting for the matrix  $J$ , it immediately gives the following lemma.

<span id="page-20-1"></span>**Lemma 4.2.** If the entries of the symmetric matrix  $G$  are independent centered subgaussian random variables, then we have for a universal constant  $C > 0$  such that

$$
\mathbb{E} \|J\| \leqslant C \left( \sqrt{\frac{\log N}{d}} + \sqrt{\frac{d_{\max}}{d}} \right).
$$

*Proof.* The proof is immediate after applying the Corollary 3.3 in [\[2\]](#page-27-5).

Now it can be seen that under the Assumption [1.3,](#page-4-0) we have  $\mathbb{E} ||J|| = O(1)$ . To complete the proof of Theorem [1.5,](#page-4-3) the rest is to prove the following concentration results for  $||J||$ .

<span id="page-20-0"></span>**Lemma 4.3.** If the entries of the symmetric matrix G are independent centered  $\kappa^2$ subgaussian random variables for some  $K \geqslant 1$ , then we have for some universal constant  $C > 0$ ,

$$
\mathbb{P}(|||J|| - \mathbb{E}||J||) \geq t \leq 2 \exp\left(-\frac{Cdt^2}{\kappa^4 \log(Nd)}\right).
$$

The major ingredient for the proof of Lemma [4.3](#page-20-0) is the following concentration result for subgaussian matrices appeared in [\[12\]](#page-28-13). We include this result for reader's convenience.

**Lemma 4.4** ([\[12,](#page-28-13) Proposition A.1]). Let  $X := (X_1, \ldots, X_d)$  be a vector of independent Ksubgaussian random variables for some  $K \geqslant 1$ . Let  $f : \mathbb{R}^d \to \mathbb{R}$  be a convex 1-Lipschitz function. Then for  $s \geq 0$ ,

$$
\mathbb{P}(|f(\boldsymbol{X}) - \mathbb{E}|f(\boldsymbol{X})| \geqslant s) \leqslant 2 \exp\left(-\frac{cs^2}{K^2 \log d}\right),
$$

where  $c > 0$  is some universal constant. Moreover,  $E f(X)$  can be replaced by any median of  $f(\boldsymbol{X}).$ 

*Proof of Lemma [4.3.](#page-20-0)* We will apply the result  $[12,$  Proposition A.1 for subgaussian random matrix to complete the proof. Note that in our setting, the operator norm is obviously a convex function. On the other hand, since for general matrices  $A, B$ ,

$$
|\|A\|-\|B\||\leqslant \|A-B\|_F\,,
$$

where  $\lVert \cdot \rVert_F$  denotes the Frobenius norm. This implies the Lipschitz property of the operator norm. There are in total  $Nd$  many nonzero entries in J, then directly from [\[12,](#page-28-13) Proposition A.1],

$$
\mathbb{P}(|||J|| - \mathbb{E}||J||) \geq t \leq 2 \exp\left(-\frac{Cdt^2}{\kappa^4 \log(Nd)}\right).
$$

Now we put the above two lemmas together to complete the proof of Theorem [1.5.](#page-4-3)

■

*Proof of Theorem [1.5.](#page-4-3)* Using the Assumption [1.3,](#page-4-0) it is easy to see that  $\mathbb{E} \|J\| = O(1)$  from Lemma [4.2.](#page-20-1) On the other hand, since  $d \gg \log d$ , the r.h.s in the concentration bound of Lemma [4.3](#page-20-0) can be bounded as follows,

$$
2\exp\left(-\frac{Cd t^2}{K^2\log(Nd)}\right) \leqslant 2\exp\left(-\frac{Cd t^2}{K^2\log(N)}\right)
$$

Using  $d \gg \log N$  in the Assumption [1.3,](#page-4-0) we have the desired results.

#### 5. Proof of Theorem [1.6](#page-4-1)

<span id="page-21-0"></span>The proof of Theorem [1.6](#page-4-1) uses a linear size set of the vertices of  $\mathcal G$  with all of its vertices having at least cd many neighbors outside the set. We start by showing that such a set always exists.

For a subset of vertices  $Q \subseteq [N]$ , denote by  $d_i^{\text{out}}(Q)$  the number of neighbors of i outside  $Q$ in  $\mathcal{G}$ , i.e.,

$$
d_i^{\text{out}}(Q) = |\{ j \in Q^c : (i, j) \in E \}|.
$$

**Definition 5.1.** For  $\delta > 0, k \ge 1$ , we call a subset of vertices  $T \subseteq [N]$  of  $\mathcal{G}(\delta, k)$ -good if  $|T| \geq \delta N$  and  $d_i^{\text{out}}(T) \geq k$  for all  $i \in T$ .

The following lemma guarantees the existence of a good set T under the assumption of  $d_{\text{max}} \leqslant Cd.$ 

<span id="page-21-1"></span>**Lemma 5.2.** Let G be any graph on  $N \geq 2$  vertices such that  $d_{\text{max}} \leq Cd$ . There exists a subset  $T \subseteq [N]$  that is  $(1/(16C), d/4)$ -good.

To establish this lemma, we need an elementary result that states that given a graph  $\mathcal{G}$ , we can always find a good expander subset  $S \subseteq [N]$  with a large number of outgoing edges, see, e.g. [\[1,](#page-27-7) Theorem 2.2.2].

**Lemma 5.3.** Let  $\mathcal{G} = (N, E)$  be a graph with  $N \geq 2$  vertices. Then there exists  $S \subseteq [N]$  with  $|S| = |N/2|$  and  $|E(S, S^{c})| \geq |E|/2 = N d/4$ , where  $E(S, S^{c}) = \{(i, j) \in E : i \in S, j \in S^{c}\}\$ and d is the total average degree.

*Proof.* Let  $\ell = |N/2|$ . Choose a random set  $U \subseteq [N]$  with  $|U| = \ell$  uniformly at random from the set of all subsets of  $[N]$  of size  $\ell$ . Note that for each  $e \in E$ ,

$$
\mathbb{P}(e \in E(U, U^{c})) = 1 - \frac{\binom{\ell}{2} + \binom{N-\ell}{2}}{\binom{N}{2}} \geq \frac{1}{2}.
$$

Since we can write

$$
|E(U, U^c)| = \sum_{e \in E} 1_{\{e \in E(U, U^c)\}},
$$

it follows from the linearity of expectation that  $\mathbb{E}|E(U, U^c)| \geq |E|/2$ . Therefore, there exists a (deterministic) subset S of size  $\ell$  such that  $|E(S, S^c)| \geq |E|/2$ , as claimed.

*Proof of Lemma [5.2.](#page-21-1)* Fix a subset S as guaranteed by the previous lemma. Let

$$
T=\Big\{i\in S: d_i^{\mathrm{out}}(S)\geqslant \frac{d}{4}\Big\},
$$

.

which contains the vertices in S with at least  $d/4$  many outgoing edges to  $S<sup>c</sup>$ . Now let I be a uniform random variable on S. Write

$$
\frac{1}{|S|} \sum_{i \in S} d_i^{\text{out}}(S) = \mathbb{E} d_I^{\text{out}}(S) \le \max_{i \in S} d_i^{\text{out}}(S) \cdot \mathbb{P}(I \in T) + \frac{d}{4} \mathbb{P}(I \notin T)
$$

$$
\le C d \mathbb{P}(I \in T) + \frac{d}{4} (1 - \mathbb{P}(I \in T)).
$$

Note that since

$$
\frac{1}{|S|} \sum_{i \in S} d_i^{\text{out}}(S) = \frac{E(S, S^c)}{|S|} \ge \frac{Nd/4}{\lfloor N/2 \rfloor} \ge \frac{d}{2},
$$

we have

$$
\frac{1}{2} \le (C - 1/4) \mathbb{P}(I \in T) + \frac{1}{4}.
$$

It follows that

$$
\mathbb{P}(I \in T) \geqslant \frac{1/4}{C - 1/4} \geqslant \frac{1}{4C},
$$

which implies  $|T| \geqslant |S|/(4C) = N/(16C)$ . Furthermore, for each  $i \in T$ , we have  $d_i^{\text{out}}(T) \geqslant$  $d_i^{\text{out}}(S) \geq d/4$ . This completes the proof of Lemma [5.2.](#page-21-1)

From now on, for the rest of Section [5,](#page-21-0) we fix a  $(1/(16C), d/4)$ -good subset T as ensured by Lemma [5.2.](#page-21-1) We employ a union bound over  $\sigma \in \{-1, +1\}^N$  to prove Theorem [1.6;](#page-4-1) the heart of the proof lies the following crucial uniform bound on the  $\ell_2$ -deviation of  $m(\sigma)$  from a constant vector.

<span id="page-22-0"></span>Proposition 5.4. Under the same conditions of Theorem [1.6,](#page-4-1) there exist positive constants  $\delta$  and K such that for any  $N \geqslant 1, \sigma \in \{-1, +1\}^N$ , and  $\gamma \in \mathbb{R}$ , we have

$$
\mathbb{P}\Big(\min_{A \subseteq T: |A| \ge \frac{|T|}{2}} \|m_A(\sigma) - \gamma \mathbf{1}_A\|_2 \le \delta \sqrt{N}\Big) \le \frac{K}{10^N},\tag{38}
$$

,

where  $m_A(\sigma) := (m_i(\sigma))_{i \in A}$  and  $\mathbf{1}_A := (1)_{i \in A}$ .

We first see how this proposition implies that  $T_N(\sigma)$  is bounded away from zero uniformly in  $\sigma$  with high probability, which is a weaker form of Theorem [1.6.](#page-4-1) The proof of Proposition [5.4](#page-22-0) is deferred to Subsection [5.1.](#page-23-0)

Proposition 5.5. Under the same conditions of Theorem [1.6,](#page-4-1) there exist positive constants  $c_1, c, \text{ and } K \text{ such that for any } N \geq 1,$ 

$$
\mathbb{P}\left(\min_{\sigma} T_N(\sigma) \leqslant c_1\right) \leqslant K \exp(-cN).
$$

*Proof.* First, for any  $M > 0$ , if  $\max_{\sigma} |\overline{m}(\sigma)| \leq M$ , we can write

$$
\frac{1}{N} \min_{\sigma} \sum_{i \in [N]} (m_i(\sigma) - \bar{m}(\sigma))^2 = \frac{1}{N} \min_{\sigma} ||m(\sigma) - \bar{m}(\sigma) \mathbf{1}||_2^2
$$

$$
= \frac{1}{N} \min_{\sigma} \inf_{\gamma \in [-M, M]} ||m(\sigma) - \gamma \mathbf{1}||_2^2
$$

which implies that

$$
\mathbb{P}\Big(\frac{1}{N}\min_{\sigma}\inf_{\gamma\in\mathbb{R}}\big\|m(\sigma)-\gamma\mathbf{1}\big\|_{2}^{2}\leqslant c_{1}\Big)\leqslant\mathbb{P}\Big(\max_{\sigma}|\bar{m}(\sigma)|>M\Big)\\qquad \qquad +\mathbb{P}\Big(\frac{1}{N}\min_{\sigma}\inf_{\gamma\in[-M,M]}\|m(\sigma)-\gamma\mathbf{1}\|_{2}^{2}\leqslant c_{1}\Big).
$$

To bound the two probabilities on the right-hand side, first of all, note that

$$
|\bar{m}(\sigma)| = \left|\frac{1}{N}\sum_{i\in[N]} m_i(\sigma)\right| \leq \frac{1}{N}\sum_{i,j\in[N]} |J_{ij}|,
$$

by Markov's inequality, for a large constant  $M = 2^N$ ,

$$
\mathbb{P}\left(\max_{\sigma} |\bar{m}(\sigma)| > M\right) \leq \frac{\sum_{i,j\in[N]} \mathbb{E}|J_{ij}|}{NM} = \frac{d \mathbb{E}|J|}{2^N} \leq \frac{C}{1.5^N}.
$$

As for the second probability, consider an arbitrary partition  $\mathcal{D} = (e_i)_{i \in [2M]}$  of  $[-M, M]$  with  $|e_i - e_{i-1}| < \varepsilon$  for  $i \in [2M]$ . From this,

$$
\frac{1}{N} \min_{\sigma} \inf_{\gamma \in [-M,M]} \|m(\sigma) - \gamma \mathbf{1}\|_2^2 \geq \frac{1}{N} \min_{\sigma} \inf_{\gamma \in \mathcal{D}} \|m(\sigma) - \gamma \mathbf{1}\|_2^2 - \varepsilon^2.
$$

and thus,

$$
\mathbb{P}\Big(\frac{1}{N}\min_{\sigma}\inf_{\gamma\in[-M,M]}\|m(\sigma)-\gamma\mathbf{1}\|\leqslant c_1\Big)\leqslant\mathbb{P}\Big(\frac{1}{N}\min_{\sigma}\inf_{\gamma\in\mathcal{D}}\|m(\sigma)-\gamma\mathbf{1}\|\leqslant c_1+\varepsilon^2\Big).
$$

Applying Proposition [5.4](#page-22-0) with  $\varepsilon^2 + c_1 = \delta^2$ , we have that for any  $\sigma$  and  $\gamma \in \mathcal{D}$ ,

$$
\mathbb{P}(\|m(\sigma)-\gamma\mathbf{1}\|_2\leqslant \delta \sqrt{N})\leqslant \mathbb{P}\Big(\min_{A\subseteq T: |A|\geqslant \frac{|T|}{2}}\|m_A(\sigma)-\gamma\mathbf{1}_A\|_{2,A}\leqslant \delta \sqrt{N}\Big)\leqslant \frac{K}{10^N}.
$$

Finally taking union bound over  $\sigma$  and  $\gamma$  yields

$$
\mathbb{P}\Big(\min_{\sigma}\inf_{\gamma\in\mathcal{D}}\|m(\sigma)-\gamma\mathbf{1}\|\leqslant \delta\sqrt{N}\Big)\leqslant c'\cdot 10^{-N}\cdot 2^{N}\cdot M,
$$

completing our proof.

<span id="page-23-0"></span>5.1. Proof of Proposition [5.4.](#page-22-0) The proof of Proposition [5.4](#page-22-0) relies on certain small-ball probability estimates. To this end, we first recall the following classical small-ball probability result for the linear combination of independent random variables.

<span id="page-23-1"></span>**Lemma 5.6** ([\[19,](#page-28-11) Corollary 2.9]). Let  $\xi_1, \ldots, \xi_n$  be independent centered random variables with variances at least 1 and third moments bounded by B. Then for any  $a \in \mathbb{R}^n$  and  $\varepsilon > 0$ , one has

$$
\sup_{v\in\mathbb{R}}\mathbb{P}\Big(\Big|\sum_{k=1}^n a_k\xi_k-v\Big|\leqslant \varepsilon\Big)\leqslant \sqrt{\frac{2}{\pi}}\frac{\varepsilon}{\|a\|_2}+C_0B\Big(\frac{\|a\|_3}{\|a\|_2}\Big),
$$

where  $C_0$  is some absolute constant.

We also need the following tensorization result.

<span id="page-23-2"></span>**Lemma 5.7** ([\[19,](#page-28-11) Lemma 2.2]). Let  $\zeta_1, \ldots, \zeta_n$  be independent random variables. Assume that there exist constants K and  $\varepsilon_0 > 0$  such that for each i,  $\mathbb{P}(|\zeta_i| < \varepsilon) \leq k \varepsilon$  for all  $\varepsilon \geq \varepsilon_0$ . Then there exists an absolute constant  $C_1$  depending only on K and  $\varepsilon_0$  such that

$$
\mathbb{P}\Bigl(\sum_{i=1}^n \zeta_i^2 < \varepsilon^2 n\Bigr) \leqslant (C_1 K \varepsilon)^n
$$

for all  $\varepsilon \geqslant \varepsilon_0$ .



<span id="page-24-0"></span>FIGURE 1. Block decomposition of matrix J, where  $D_1$  is a  $|T| \times |T|$  matrix with row and column index set  $T$ , while  $D_2$  has row index set  $T$  and column index set  $[N]\backslash T$ . It can be seen that the entries in the block  $D_2$  are independent from each other.

Our goal is to combine Lemma [5.6](#page-23-1) and Lemma [5.7](#page-23-2) to establish some small-ball probability estimate for the vector  $m(\sigma)$ . However, because the matrix J is symmetric, the random variables  $m_1(\sigma), m_2(\sigma), \ldots, m_N(\sigma)$  are not independent. To circumvent this obstacle, we will make use of the fact that the entries in any off-diagonal block sub-matrices of J are independent. In particular, we shall work with the sub-matrix  $D_2$  of J defined on the block  $T \times T^c$ , where T is a  $(1/(16C), d/4)$ -good subset as in Lemma [5.2.](#page-21-1) With this choice of T, we partition the vertex set as  $[N] = T \cup (N \setminus T)$  and accordingly write  $\sigma = (\sigma^{(1)}, \sigma^{(2)}) \in$  $\{-1,+1\}^T \times \{-1,+1\}^{[N]\setminus T}$ . Similarly we represent J (after possible permutation of the N vertices) in the following block form,

<span id="page-24-2"></span>
$$
J = \begin{pmatrix} D_1 & D_2 \\ D_2^T & * \end{pmatrix},\tag{39}
$$

where  $D_1 \in \mathbb{R}^{|T| \times |T|}$  and  $D_2 \in \mathbb{R}^{|T| \times (N-|T|)}$ , also see the visualization in Figure [1.](#page-24-0) Let  $r_i$  be the number of non-zero elements in the *i*-th row of  $D_2$ . Note that  $r_i$  is same as  $d_i^{\text{out}}(T)$  for  $i \in T$ . It follows from the definition of T,

<span id="page-24-1"></span>
$$
\frac{d}{4} \leqslant r_i \leqslant Cd. \tag{40}
$$

Under the above decomposition,

$$
(D_2\sigma^{(2)})_i = \sum_{j \in [N] \setminus T} J_{ij}\sigma_j, \ \forall i \in T,
$$

it can be readily seen that the coordinates in  $D_2\sigma^{(2)}$  are now independently from each other. We now use this property and the bound in  $(40)$  to establish the desired small ball probability estimates and use them to further prove the Proposition [5.4.](#page-22-0)

*Proof of Proposition [5.4.](#page-22-0)* Recall that  $A = (a_{ij})_{i,j}$  is the adjacency matrix of G. First, as we have explained above that  $(D_2 \sigma^{(2)})_{i \in T} = (\langle a_i, g_i \rangle)_{i \in T}$  has independent coordinates and proper moment assumptions, where

$$
\boldsymbol{a}_i := d^{-1/2} (a_{i,|T|+1} \sigma_{|T|+1}, \ldots, a_{i,N} \sigma_N),
$$
  

$$
\boldsymbol{g} := (g_{i,|T|+1}, g_{i,|T|+2}, \ldots, g_{i,N}).
$$

Recall that  $(40)$  implies

$$
r_i = |\{ j \in [N] \setminus T : a_{ij} = 1 \}| \in [d/4, Cd].
$$

In addition, since

$$
\|\boldsymbol{a}_i\|_2^2 = \frac{r_i}{d}
$$
 and  $\|\boldsymbol{a}_i\|_3^3 = \frac{r_i}{d^{3/2}}$ ,

we have

$$
\|\bm{a}_i\|_2 \in [1/2, \sqrt{C}] \text{ and } \left(\frac{\|\bm{a}_i\|_3}{\|\bm{a}_i\|_2}\right)^3 = \frac{1}{\sqrt{r_i}} \leq \frac{2}{\sqrt{d}}.
$$

By Lemma [5.6,](#page-23-1) we have that for any  $i \in T$  and  $\varepsilon > 0$ 

$$
\sup_{v \in \mathbb{R}} \mathbb{P}(|(D_2 \sigma^{(2)})_i - v| \leqslant \varepsilon) \lesssim \varepsilon.
$$

Combining with Lemma [5.7,](#page-23-2) there exists a constant  $\delta > 0$  such that for any  $A \subseteq T$  with  $|A| \geqslant |T|/2 \geqslant N/(32C)$ , we have

$$
\sup_{\boldsymbol{v}\in\mathbb{R}^{|A|}}\mathbb{P}\big(\big\|(D_2\sigma^{(2)})_{i\in A}-\boldsymbol{v}\big\|_2^2\leqslant \delta^2N\big)\lesssim 10^{-N}.
$$

Recall  $m_A(\sigma) = (m_i(\sigma))_{i \in A}$ . Using the previous decomposition of J in [\(39\)](#page-24-2), we observe that

$$
||m_A(\sigma) - \gamma \mathbf{1}||_2^2 \ge ||(D_1 \sigma^{(1)} + D_2 \sigma^{(2)})_{i \in A} - \gamma \mathbf{1}_A||_2^2.
$$

Then

$$
\mathbb{P}(\|m_A(\sigma) - \gamma \mathbf{1}_A\|_2^2 \leq \delta^2 N)
$$
\n
$$
\leq \mathbb{P}(\|(D_1 \sigma^{(1)} + D_2 \sigma^{(2)})_{i \in A} - \gamma \mathbf{1}_A\|_2^2 \leq \delta^2 N)
$$
\n
$$
= \mathbb{E}_{D_1} \Big[ \mathbb{P}(\|(D_1 \sigma^{(1)} + D_2 \sigma^{(2)})_{i \in A} - \gamma \mathbf{1}_A\|_2^2 \leq \delta^2 N |(D_1 \sigma^{(1)})_{i \in A}) \Big]
$$
\n
$$
\leq \mathbb{E}_{D_1} \Big[ \sup_{\mathbf{v} \in \mathbb{R}^{|A|}} \mathbb{P}(\|(D_2 \sigma^{(2)}\|_{i \in A} - \mathbf{v}\|_2^2 \leq \delta^2 N) \Big]
$$
\n
$$
\leq \sup_{\mathbf{v} \in \mathbb{R}^{|A|}} \mathbb{P}(\|(D_2 \sigma^{(2)})_{i \in A} - \mathbf{v}\|_2^2 \leq \delta^2 N) \leq \frac{K}{10^N}
$$

for some constant K. Since A is arbitrary subset of T with  $|A| \geq |T|/2$ , we have

$$
\mathbb{P}\left(\min_{A\subseteq T:|A|\geqslant \frac{|T|}{2}}\|m_A(\sigma)-\gamma\cdot\mathbf{1}_A\|_2\leqslant \delta \sqrt{N}\right)\leqslant \frac{K}{10^N}.
$$

■

5.2. Proof of Theorem [1.6.](#page-4-1) In this section, we will complete the proof of Theorem [1.6.](#page-4-1) First, we prove the following result that states that with high probability, for each  $\sigma$ , the variables  $m_i(\sigma)$  are bounded for at least half of the indices in T.

<span id="page-25-0"></span>**Proposition 5.8.** Assume  $d_{\text{max}} \leq C d$ . There exist constants  $M, K > 0$  such that the following event

$$
\min_{\sigma} |\{i \in T : |m_i(\sigma)| \leq M\}| \geq \frac{|T|}{2}
$$

occurs with probability at least  $1 - K \cdot 10^{-N}$ .

*Proof.* Consider the matrix  $J_T := (D_1, D_2)$ , where  $D_1, D_2$  were defined in [\(39\)](#page-24-2). Note that for each fixed  $\sigma$ ,

$$
m_T(\sigma) = J_T \sigma = D_1 \sigma^{(1)} + D_2 \sigma^{(2)},
$$

Decompose  $D_1$  as

$$
D_1 = D_1^+ + D_1^-,
$$

where

$$
(D_1^+)_{ij} := \begin{cases} (D_1)_{ij}, & \text{if } i < j, \\ 0, & \text{if } i \geq j, \end{cases} \text{ and } (D_1^-)_{ij} := \begin{cases} (D_1)_{ij}, & \text{if } i > j, \\ 0, & \text{if } i \leq j. \end{cases}
$$

Accordingly, we have the decomposition for  $J_T = J_T^+ + J_T^-$  with

$$
J_T^+ := (D_1^+, D_2)
$$
 and  $J_T^- := (D_1^-, 0)$ .

In this way, it can be seen that the entries in  $J^+_{\mathcal{T}}$  $T$  are independent and so are the entries of  $J_T^ T_T^-$ . We let  $m_T^+(\sigma) := J_T^+$  $T^+\sigma$  and  $m_T^-(\sigma) := J_T^ \bar{T}\sigma$ , whose coordinates are now independent. For  $i \in T$ , notice

$$
\mathbb{E} \, m_i^{\pm}(\sigma) = 0 \text{ and } \operatorname{Var}(m_i^{\pm}(\sigma)) = \mathbb{E}[(m_i^{\pm}(\sigma))^2] \leqslant \frac{d_i}{d} \leqslant C.
$$

The last inequality in the variance bound used the assumption  $d_{\text{max}} \leq C d$  and the independence among the entries of  $J_T^+$  $T^+_T, J^-_T$ . Consequently, from Chebyshev's inequality, we have that for  $i \in T$ ,

$$
\mathbb{P}\Big(\big|m_i^{\pm}(\sigma)\big|\geqslant \frac{M}{2}\Big)\leqslant \frac{4C}{M^2}.
$$

Next, set

$$
B_{\sigma}^+ := \left\{ i \in T, \left| \left| m_i^+(\sigma) \right| \leqslant \frac{M}{2} \right\} \text{ and } B_{\sigma}^- := \left\{ i \in T, \left| \left| m_i^-(\sigma) \right| \leqslant \frac{M}{2} \right.\right\}.
$$

Evidently,

 $|B^+_{\sigma}| \sim \text{Binomial}(|T|,p)$  and  $|B^-_{\sigma}| \sim \text{Binomial}(|T|,p)$ ,

where p is some constant with  $p \geq 1-4C/M^2$ . Now, by the binomial concentration, one can choose sufficiently large M such that

$$
P(|B_{\sigma}^+|,|B_{\sigma}^-| \geq \frac{4}{5}|T| \geqslant 1 - \frac{K}{20^N}
$$

for some constant  $K > 0$  independent of N. Let  $B_{\sigma} := B_{\sigma}^+ \cap B_{\sigma}^-$ . Using the identity

$$
|B_\sigma|=|B_\sigma^+|+|B_\sigma^-|-|B_\sigma^+\cup B_\sigma^-|,
$$

with probability at least  $1 - K/20^N$ ,

$$
|B_\sigma| \geqslant \frac{4}{5}\left|T\right| + \frac{4}{5}\left|T\right| - \left|T\right| \geqslant \frac{1}{2}\left|T\right|,
$$

which readily implies that

$$
|m_i(\sigma)| \leq |m_i^+(\sigma)| + |m_i^-(\sigma)| \leq \frac{M}{2} + \frac{M}{2} = M
$$

for  $i \in B_{\sigma}$  and as a consequence,

$$
|\{i \in T : |m_i(\sigma)| \leq M\}| \geq \frac{|T|}{2}.
$$

Finally, applying union bound with resepct to  $\sigma \in \{-1, +1\}^N$  leads to the desired result.

We proceed to establish the proof of Theorem [1.6](#page-4-1) via this proposition.

*Proof of Theorem [1.6.](#page-4-1)* For each fixed  $\sigma$ , let

$$
B_{\sigma} = \{ i \in T : |m_i(\sigma)| \leqslant M \}.
$$

By Proposition [5.8,](#page-25-0) there exist a constant  $M > 0$  sufficiently large such that with probability at least  $1 - K/10^N$ , we have  $\min_{\sigma} |B_{\sigma}| \geq |T|/2 \geq N/(32C)$ . Consequently, for  $i \in B_{\sigma}$ 

$$
\theta_i(\sigma) = \operatorname{sech}(\beta m_i(\sigma) + h) \geq c'' := \operatorname{sech}(\beta M + |h|).
$$

Note that

$$
\widetilde{T}_N(\sigma) \geq \frac{1}{N^2} \sum_{i,j \in B_{\sigma}} (m_i(\sigma) - m_j(\sigma))^2 \theta_i(\sigma) \theta_j(\sigma)
$$
\n
$$
\geq \frac{(c'')^2}{N^2} \sum_{i,j \in B_{\sigma}} (m_i(\sigma) - m_j(\sigma))^2
$$
\n
$$
= \frac{2(c'')^2}{N} \sum_{i \in B_{\sigma}} (m_i(\sigma) - \bar{m}_{B_{\sigma}}(\sigma))^2,
$$

where  $\bar{m}_{B_{\sigma}}(\sigma)$  is the average of  $m_{B_{\sigma}(\sigma)}(\sigma)$ . Therefore, on the event  $|B_{\sigma}| \geq |T|/2$ , we have

$$
\widetilde{T}_N(\sigma) \geq \frac{2(c'')^2}{N} \inf_{\gamma \in [-M,M]} \sum_{i \in B_{\sigma}} (m_i(\sigma) - \gamma)^2
$$
\n
$$
\geq \frac{2(c'')^2}{N} \inf_{\gamma \in [-M,M]} \min_{A \subseteq T : |A| \geq |T|/2} \sum_{i \in A} (m_i(\sigma) - \gamma)^2
$$
\n
$$
\geq \frac{2(c'')^2}{N} \inf_{\gamma \in \mathcal{D}} \min_{A \subseteq T : |A| \geq |T|/2} \left( \|m_A(\sigma) - \gamma \cdot \mathbf{1}\|_2 - \frac{\delta \sqrt{N}}{2} \right)_+^2,\tag{41}
$$

where D is an arbitrary partition of  $[-M, M]$  with mash  $(\delta/2)$  and  $|\mathcal{D}| \leq 10M\delta^{-1}$ . Applying Proposition [5.4](#page-22-0) for a fixed  $\gamma$  and  $\sigma$  in [\(41\)](#page-27-8) and then taking a union bound over  $\gamma \in \mathcal{D}$  and  $\sigma$ , we obtain, for  $c_0 = (c'')^2 \delta^2/2$ ,

$$
\mathbb{P}\left(\min_{\sigma}\widetilde{T}_N(\sigma)\leqslant c_0\right)\leqslant K10^{-N}+|\mathcal{D}|2^N\cdot K10^{-N}.
$$

The proof of Theorem [1.6](#page-4-1) now immediately follows.

#### <span id="page-27-8"></span>**REFERENCES**

- <span id="page-27-7"></span>[1] N. Alon and J. H. Spencer, The probabilistic method, Third, Wiley-Interscience Series in Discrete Mathematics and Optimization, John Wiley & Sons, Inc., Hoboken, NJ, 2008. With an appendix on the life and work of Paul Erdős. MR2437651
- <span id="page-27-5"></span>[2] A. S. Bandeira and R. van Handel, Sharp nonasymptotic bounds on the norm of random matrices with independent entries, Ann. Probab. 44 (2016), no. 4, 2479–2506. MR3531673
- <span id="page-27-4"></span>[3] A. Basak and S. Mukherjee, Universality of the mean-field for the Potts model, Probab. Theory Related Fields 168 (2017), no. 3-4, 557–600. MR3663625
- <span id="page-27-0"></span>[4] J. Besag, Spatial interaction and the statistical analysis of lattice systems, J. Roy. Statist. Soc. Ser. B 36 (1974), 192–236. MR373208
- <span id="page-27-1"></span>[5] J. Besag, Statistical analysis of non-lattice data, Journal of the Royal Statistical Society Series D: The Statistician 24 (1975), no. 3, 179–195.
- <span id="page-27-3"></span>[6] B. B. Bhattacharya and S. Mukherjee, *Inference in Ising models*, Bernoulli 24 (2018), no. 1, 493–525. MR3706767
- <span id="page-27-6"></span>[7] P. Carmona and Y. Hu, Universality in Sherrington–Kirkpatrick's spin glass model 42 (2006), no. 2, 215– 222.
- <span id="page-27-2"></span>[8] S. Chatterjee, Estimation in spin glasses: a first step, Ann. Statist. 35 (2007), no. 5, 1931–1946. MR2363958
- <span id="page-28-8"></span>[9] S. Chatterjee, Large deviations for random graphs, Lecture Notes in Mathematics, vol. 2197, Springer, Cham, 2017. Lecture notes from the 45th Probability Summer School held in Saint-Flour, June 2015, École d'Été de Probabilités de Saint-Flour. [Saint-Flour Probability Summer School]. MR3700183
- <span id="page-28-9"></span><span id="page-28-0"></span>[10] S. Chatterjee and A. Dembo, Nonlinear large deviations, Adv. Math. 299 (2016), 396-450. MR3519474 [11] F. Comets and B. Gidas, Asymptotics of maximum likelihood estimators for the Curie-Weiss model, Ann.
- <span id="page-28-13"></span>Statist. 19 (1991), no. 2, 557–578. MR1105836 [12] N. A. Cook, R. Ducatez, and A. Guionnet, Full large deviation principles for the largest eigenvalue of sub-Gaussian Wigner matrices, 2023.
- <span id="page-28-10"></span>[13] R. Eldan and R. Gross, Decomposition of mean-field Gibbs distributions into product measures, Electron. J. Probab. 23 (2018), Paper No. 35, 24. MR3798245
- <span id="page-28-7"></span>[14] P. Ghosal and S. Mukherjee, Joint estimation of parameters in Ising model, Ann. Statist. 48 (2020), no. 2, 785–810. MR4102676
- <span id="page-28-5"></span>[15] J. J. Hopfield, Neural networks and physical systems with emergent collective computational abilities, Proc. Nat. Acad. Sci. U.S.A. 79 (1982), no. 8, 2554–2558. MR652033
- <span id="page-28-6"></span>[16] S. Mukherjee, J. Son, and B. B. Bhattacharya, Estimation in tensor Ising models, Inf. Inference 11 (2022), no. 4, 1457–1500. MR4526327
- <span id="page-28-1"></span>[17] G. Parisi, Infinite number of order parameters for spin-glasses, Physical Review Letters 43 (1979), no. 23, 1754.
- <span id="page-28-2"></span>[18] G. Parisi, A sequence of approximated solutions to the SK model for spin glasses, Journal of Physics A: Mathematical and General 13 (1980), no. 4, L115.
- <span id="page-28-11"></span>[19] M. Rudelson and R. Vershynin, The Littlewood-Offord problem and invertibility of random matrices, Adv. Math. 218 (2008), no. 2, 600-633. MR2407948
- <span id="page-28-4"></span>[20] D. Sherrington and S. Kirkpatrick, Solvable model of a spin-glass, Physical review letters 35 (1975), no. 26, 1792.
- <span id="page-28-3"></span>[21] M. Talagrand, The Parisi formula, Ann. of Math. (2) 163 (2006), no. 1, 221–263. MR2195134
- <span id="page-28-12"></span>[22] R. Vershynin, Invertibility of symmetric random matrices, Random Structures & Algorithms 44 (2014), no. 2, 135–182.