

Odd-even mass differences of well and rigidly deformed nuclei in the rare earth region: A test of a newly proposed fit of pairing matrix elements

T. V. Nhan Hao* and N. N. Bao Nguyen

*Faculty of Physics, University of Education, Hue University, 34 Le Loi Street, Hue City, Vietnam and
Center for Theoretical and Computational Physics, University of Education,
Hue University, 34 Le Loi Street, Hue City, Vietnam*

D. Quang Tam

*Faculty of Physics, University of Education, Hue University, 34 Le Loi Street, Hue City, Vietnam
Center for Theoretical and Computational Physics, University of Education,
Hue University, 34 Le Loi Street, Hue City, Vietnam and
Faculty of Basic Sciences, University of Medicine and Pharmacy,
Hue University, 34 Le Loi Street, Hue City, Vietnam*

P. Quentin† and L. Bonneau

LP2I, UMR 5797, Université de Bordeaux, CNRS, F-33170, Gradignan, France

Meng-Hock Koh (辜明福)‡

*Department of Physics, Faculty of Science, Universiti Teknologi Malaysia, 81310 Johor Bahru, Johor, Malaysia. and
UTM Centre for Industrial and Applied Mathematics, 81310 Johor Bahru, Johor, Malaysia*

(Dated: June 18, 2024)

We discuss a test of a recently proposed approach to determine average pairing matrix elements within a given interval of single-particle states (sp) around the Fermi level λ as obtained in the so-called uniform gap method (UGM). It takes stock of the crucial role played by the averaged sp level density $\bar{\rho}(e)$. These matrix elements are deduced within the UGM approach, from microscopically calculated $\bar{\rho}(e)$ and gaps obtained from analytical formulae of a semi-classical nature. Two effects generally ignored in similar fits have been taken care of. They are: (a) the correction for a systematic bias in choosing to fit pairing gaps corresponding to equilibrium deformation solutions as discussed by Möller and Nix [Nucl. Phys. A 476, 1 (1992)] and (b) the correction for a systematic spurious enhancement of $\bar{\rho}(e)$ for protons in the vicinity of λ , because of the local Slater approximation used for the treatment of the Coulomb exchange terms in most calculations (see e.g. [Phys. Rev C 84, 014310 (2011)]). This approach has been deemed to be very efficient upon performing Hartree-Fock + BCS (with seniority force and self-consistent blocking when dealing with odd nuclei) calculations of a large sample of well and rigidly deformed even-even rare-earth nuclei. The reproduction of their experimental moments of inertia has been found to be at least of the same quality as what has been obtained in a direct fit of these data [Phys. Rev C 99, 064306 (2019)]. We extend here the test of our approach to the reproduction, in the same region, of three-point odd-even mass differences centered on odd- N or odd- Z nuclei. The agreement with the data is again roughly of the same quality as what has been obtained in a direct fit, as performed in [Phys. Rev C 99, 064306 (2019)].

I. INTRODUCTION

A simple prescription to determine pairing matrix elements from average single-particle (sp) level densities for the ground states of well and rigidly deformed nuclei has been proposed recently in [1] and deemed to be rather successful. These matrix elements V_q (where q stands for the charge state) correspond to their average values around the Fermi energies λ_q . They are to be used in a microscopic approach of the Hartree-Fock-plus-BCS (HF + BCS) type where the pairing correlations are treated within the so-called seniority force approximation.

It takes stock of the strong dependence of the V_q values upon the sp level densities at the Fermi energies $g_q(\lambda_q)$ averaged à la Strutinsky considered as a sound ersatz of a semi-classical approximation. It relies also on the consideration that any fitting of nuclear energies (here any energy related to pairing correlations) in terms of nucleon numbers correspond in an effective way to such a semi-classical approximation. Therefore the input of experimental data is naturally performed through some standard analytical formula expressing the nucleon number dependence of odd-even mass differences defined in different ways noted here as δE (see, e.g., Refs. [2, 3]) taking into account the important corrective contribution of [4].

After the pioneering work of Ref. [5], it has been a current practice to adjust the intensity of pairing correlations to reproduce the data on δE . To a lesser extent, for this purpose, one has also considered moments of in-

* tvnhao@hueuni.edu.vn

† quentin@cenbg.in2p3.fr

‡ kmhock@utm.my

ertia noted as \mathcal{J} , deduced from the first 2^+ excitation energies of well and rigidly deformed even-even heavy nuclei. In a recent paper [6], it has been demonstrated in the region of (or close to) rare-earth deformed nuclei (loosely dubbed as rare-earth nuclei in what follows) that making separate fits on δE and \mathcal{J} yielded similarly good results for the parameters in use to calculate pairing correlations within HF + (seniority) BCS calculations. This was a confirmation that both pieces of data were highly contingent upon a good description of pairing properties and thus asserting their relevance for such fits.

In both cases (with δE or \mathcal{J}), these fitting approaches possibly suffer from accidental local deficiencies of the sp distribution of levels in the vicinity of the Fermi energy. *A priori*, resorting to semi-classical quantities associated to a given microscopic theoretical approach one gets rid of such a practical problem. What remains of course to be assessed is the relevance of the analytical expressions for the (N, Z) dependence of the average δE differences as well as the quality of the averaged sp level density of the canonical basis states.

The method proposed in [1] has been validated by comparing its results for moments of inertia of well and rigidly deformed rare-earth nuclei with those obtained through a specific fit on the \mathcal{J} data as performed in Ref. [6]. A similar quality in the reproduction of such spectroscopic data was obtained.

It is the aim of the present paper to make a similar comparison for the δE data now, between the results obtained within the method developed in [1] and the direct fit of these differences performed in the same nuclear region, in Ref. [6].

In Section II, we will briefly recall the approach developed in [1], whereas Section III will provide some details about the calculations performed here. Our results will be discussed in Section IV ; Section V will be devoted to some conclusions and perspectives.

II. BRIEF SURVEY OF THE APPROACH

The method in use has been presented and discussed in detail in [1]. We will thus restrict here to sketchily describe its main characteristics.

Our starting point is a sp spectrum for a charge state q , obtained *a priori* through any microscopic approach. In what follows it will result from self-consistent HF+BCS calculations within the seniority force simple ansatz for the pairing matrix elements.

We calculate an approximate semi-classically averaged sp level density $\tilde{\rho}(e)$ as a function of the sp energy e through a standard Strutinsky's energy averaging (see Refs. [7, 8]). As discussed in [1] for the considered nuclei located far enough from the neutron drip line, the averaging width is taken as $\gamma = 1.2 \hbar\omega$ where the energy scale as a function of the nucleon number A is the usual expression $\hbar\omega = 41A^{-1/3}$ MeV [9]. Also we take a generalized Laguerre polynomial of degree 2 in the variable

$(e - \tilde{e})/\gamma^2$, with usual notation, to take into account the so-called curvature correction.

For N_q nucleons of the charge state q , we then compute the Fermi energy $\tilde{\lambda}_q$ as

$$N_q = \int_{-\infty}^{\tilde{\lambda}_q} \tilde{\rho}_q(e) de. \quad (1)$$

Within the so-called uniform gap method of Refs. [7] [10], given a suitably average gap $\tilde{\Delta}_q$ (see below), the average pairing matrix element V_q to be used in our HF+BCS approach is given by

$$\frac{1}{V_q} = \int_{\tilde{\lambda}_q - \Omega}^{\tilde{\lambda}_q + \Omega} \frac{\tilde{\rho}_q(e)}{\sqrt{(e - \tilde{\lambda}_q)^2 + \tilde{\Delta}_q^2}} de. \quad (2)$$

As well-known, the value of this matrix element depends on the choice of the energy interval 2Ω centered at the Fermi energy and including the sp states active in the BCS variational determination of occupation probabilities. In this study as in [1] we take $\Omega = 6$ MeV.

The nucleon-number N_q dependence of average gaps $\tilde{\Delta}_q$ could be taken *a priori* from standard formulas (as, e.g., those of Refs. [2], [3]). However Möller and Nix [4] have pointed out that there is a bias in such estimates due to the mere selection of nuclei at equilibrium deformation, corresponding systematically to lower than average quantal sp level densities. As a result of their study, they proposed the following parametrisation of the average gaps

$$\tilde{\Delta}_q = \frac{r B_s}{N_q^{1/3}}, \quad (3)$$

where B_s is set to 1 and $r = 4.8$ MeV.

While we stick to this value for neutrons, we have remarked in [1] that this must not be the case for protons. In most of microscopic approaches of the Hartree-Fock plus BCS, Hartree-Fock-Bogoliubov (HFB) or relativistic mean-field type, one simplifies the treatment of exchange terms of the infinite range Coulomb interaction by having recourse to the local Slater approximation [11]. It has been found long ago [12] and further studied in Refs. [13, 14] that the Slater approximation systematically and significantly overestimates the quantal sp level density near the Fermi energy (especially at the equilibrium deformation).

To ensure a safe use of the Slater approximation in our approach, we must thus quench the parameter r of Eq (3) by a factor R_p given by

$$R_p = 0.0181 E_{cond}^p + 0.781, \quad (4)$$

where E_p^{cond} is the average pairing energy (in MeV). It is estimated using the quantal gap obtained within HF+BCS calculations with some starting ansatz for the proton matrix element V_p such that (see Appendix A of

[1]):

$$E_{cond}^p = \frac{\Delta_p^2}{V_p}. \quad (5)$$

In principle this should imply an iterative process to define new Möller-Nix average gaps and accordingly a new V_q value. However, it has been shown in [1] that choosing some starting V_p within a *reasonable* range of values (as it turns out easily delineated), these iterations do not bring significant modifications of the resulting matrix elements given the rough character of the above corrective process.

III. SOME DETAILS OF THE CALCULATIONS

A. The canonical basis

In our approach, the sp canonical basis states are obtained within self-consistent HF+BCS calculations. Let us recall that throughout this paper the BCS correlations are determined within the seniority force simple approximation. The particle-hole interaction in use is of the standard Skyrme type. Calculations are performed with the SIII parametrisation [15] which has been shown in many instances to provide a rather good description of the spectroscopic properties of well and rigidly deformed nuclei (see, e.g., for a recent account Ref. [16]). The averaged pairing matrix elements V_q are given for each nucleus according to the method summarized in Section II for even-even nuclei. For odd- Z (odd- N respectively) nuclei the retained values of the matrix elements are interpolated between the values found for the neighbouring isotones (isotopes respectively).

The axial and intrinsic parity symmetries have been imposed to the solutions. The nuclei are thus specified by their projection K of their angular momentum on the symmetry axis and their parity π . To solve the Schrödinger equation determining the canonical basis states, we have expanded these states on eigenstates of an axially symmetrical harmonic oscillator Hamiltonian. This basis is defined by a parameter $N_0 = 16$ defining a deformation-dependent selection of relevant basis states corresponding at sphericity to the consideration of 17 shells (see [17]), as well as the inverse-length b and deformation q parameters defined in [17]. The latter two parameters have been optimized for even-even nuclei to give the lowest energy for each equilibrium solution, whereas for odd- A nuclei they have been interpolated from those obtained for even-even nuclei. Integrals involving the local densities are performed using the Gauss-Hermite and Gauss-Laguerre approximate integration methods with 50 points along the symmetry axis and 16 points in the perpendicular direction, respectively.

B. Odd-even mass differences

The data on odd-even mass differences are extracted through a 3-point formula. As discussed in Refs. [18, 19] such differences $\delta_q^{(3)}$ centered on an odd-neutron or odd-proton nucleus are good markers on the degree of pairing correlations. They are, to a large extent, free from single-particle filling effects and are given, e.g., for an isotopic series by

$$\begin{aligned} \delta_n^3(N) &= \frac{(-1)^N}{2} \left[E(N+1, Z) - 2E(N, Z) + E(N-1, Z) \right] \\ &= \frac{(-1)^N}{2} \left[S_n(N, Z) - S_n(N+1, Z) \right] \end{aligned}$$

where N is odd and $S_n(N, Z)$ is the neutron separation energy of a nucleus composed of N neutrons and Z protons whose total energies are denoted as $E(N, Z)$. Similar expressions are easily deduced from the above for odd-proton nuclei.

These energies are compared with those extracted directly from calculated binding energies within the HF+BCS approach. In the case of odd- A nuclei, we have performed selfconsistent blocking calculations (i.e. placing a nucleon in the relevant sp orbit specified by the K^π quantum numbers). The breaking of the time-reversal symmetry implies the presence of new (time-odd) local densities in the expression of the Hamiltonian density resulting in new terms in the corresponding Hartree-Fock potential. As explicitly detailed in Ref. [20], when using the SIII interaction we have considered a restricted choice of time-odd potential fields, yet preserving the Galilean invariance (namely the vector spin field $\mathbf{S}(\mathbf{r})$ and the vector current field $\mathbf{A}(\mathbf{r})$, with usual notation). This choice is dubbed as the *minimal* scheme in Ref. [20].

In the spirit of the Bohr-Mottelson unified model, well suited for these deformed nuclei, we assimilate the nuclear angular momentum and parity quantum numbers I^π to those K^π of the blocked nucleon sp state. This is of course not free for any perturbation of the low-energy nuclear spectra from possible Coriolis coupling which will be ignored here. Particular cases concern solutions where $K = 1/2$ with a decoupling parameter a outside the range $-1 \leq a \leq 4$. They will be specifically discussed in Section IV and shown to be unable to perturb the *natural* rotational band ordering of states.

We have systematically calculated solutions for odd- A nuclei corresponding to the ground state I^π experimental values [21] as well as those where the calculated energies lie below the former or in some cases (see the discussions of Section IV) above generally up to a couple of 100 keV.

C. Choice of sample nuclei

We have chosen odd- A nuclei belonging to the rare-earth region plus Hafnium isotopes (loosely called below

rare-earth nuclei), i.e. some odd-proton isotopes from Europium to Lutetium and odd-neutron isotopes from Samarium to Hafnium. They are chosen to be well and rigidly deformed and, moreover, far enough from a region of transition between deformed and soft nuclei. The first criterion is retained to be able to reduce approximately the collective dynamics to a pure rigid rotation within the Bohr-Mottelson unified model, allowing in particular to limit ourselves to a single BCS state, i.e. ignoring quantal shape fluctuations. The second criterion is retained to avoid wide shape variations (and therefore large sp spectrum changes) between the three isotopes (isotones respectively) entering the calculation of the energy differences $\delta_n^{(3)}$ ($\delta_p^{(3)}$ respectively).

Table I displays the ratio R_{42} of the excitation energies of the first 4^+ and 2^+ states [21] of the 22 even-even nuclei bracketing the odd- A nuclei whose energy differences $\delta_q^{(3)}$ are evaluated in our calculations. It appears that they satisfy reasonably well the first criterion since for all of them one has $R_{42} \geq 3.29$.

On this table also, the intrinsic axial charge quadrupole moments Q_{20}^{int} obtained in our calculations are compared, when available, with the corresponding experimental values deduced either from reduced B(E2) data [22] or those deduced from the spectroscopic moments Q_{20}^{spect} of the first 2^+ state [23], upon using the unified model relation for $I = 2, K = 0$, namely $Q_{20}^{\text{int}} = -3.5 Q_{20}^{\text{spect}}$. One notices a rather good reproduction of the Q_{20}^{int} data with the interaction SIII in use, as shown long ago [24] (somewhat less good however for the $^{156,158}\text{Sm}$ isotopes).

IV. RESULTS

A. Discussion of the ground-state configurations obtained in our calculations for odd- Z nuclei

We compare here the lowest-energy configurations as obtained in our calculations with the experimental values of the angular momentum and parity quantum numbers I^π given in the current version of the NUDAT compilation [21].

As seen on Table II, for 8 nuclei out of 13 (^{161}Tb , ^{163}Tb , ^{167}Ho , ^{169}Ho , ^{169}Tm , ^{171}Tm , ^{177}Lu , ^{179}Lu), our theoretical assignments agree with the data.

We confirm the suggested assignments for 3 nuclei, namely: ^{165}Tb and ^{167}Tb as $3/2^+$ and ^{173}Tm as $1/2^+$. No assignment is proposed in Ref. [21] for the ^{161}Eu nucleus. We suggest a $5/2^-$ configuration noting however that we have obtained a $5/2^+$ solution 179 keV above the $5/2^-$ configuration. In one case (^{159}Eu) the experimental lowest configuration $5/2^+$ is obtained at 145 keV above a $5/2^-$ state. Finally, we remark that the decoupling constant values corresponding to the $1/2^+$ state considered in the three calculated isotopes of Thulium belong to the $[-0.64, -0.60]$ interval ensuring that the band head spin assignment as $1/2$ is correct.

B. Discussion of the ground-state configurations obtained in our calculations for odd- N nuclei

As seen on Table III, for 9 nuclei out of 16 (^{159}Sm , ^{161}Gd , ^{163}Dy , ^{171}Er , ^{173}Yb , ^{175}Yb , ^{177}Yb , ^{179}Hf , ^{181}Hf), our theoretical assignments agree with the data.

We confirm the suggested assignment for one nucleus, namely: ^{157}Sm as $3/2^-$. No assignment is proposed in Ref. [21] for the ^{165}Gd nucleus. We suggest a $7/2^+$ configuration. In ^{167}Dy the suggested ground state configuration $1/2^-$ suggested in Ref. [21] is calculated only 58 keV above a $7/2^+$ state. In one nucleus (^{169}Er) the experimental lowest configuration $1/2^-$ is obtained only 72 keV above a $7/2^+$ state.

Three cases deserve a particular attention. In Ref. [21] two assignments ($5/2^-, 7/2^+$) are suggested for ^{163}Gd . We found the latter ($7/2^+$) as the ground state with the former ($5/2^-$) lying 579 keV higher with a $1/2^-$ configuration located in between at 59 keV. In ^{165}Dy one has found experimentally a $7/2^+$ ground state with a $1/2^-$ isomeric state with an excitation energy of 108 keV. We found both states as the lowest ones but with an inversion, the $1/2^-$ being found 48 keV lower in energy. A similar situation is also present in ^{171}Yb with an experimental $1/2^-$ ground state and a $7/2^+$ isomeric state 95 keV above. Here, too, we found these states as the lowest ones but with an inversion whereby the $1/2$ being found 73 keV lower.

C. Comments on our assignments of spin and parity

In our calculations for both odd- Z and odd- N nuclei, we thus found that we are in agreement in 21 out of 29 cases where an assignment of spin and parity has been reported or suggested in Ref. [21]. We have proposed an assignment for two nuclei. In two instances where an isomeric state has been experimentally found at excitation energies in the 100-150 keV range, we have obtained them, yet with an inversion in their respective ordering, corresponding to an error of 50-70 keV. From all these results implying a sample of 29 odd- A nuclei, yet limiting ourselves to the consideration of intrinsic states within the unified model, we can reasonably hint that our estimates of the low lying bandhead spectra provides a good reproduction of relative energies within not much more than about 150 keV.

Finally we discuss the decoupling constant values corresponding to the $1/2$ states considered in our calculations of these odd- N nuclei. It is found to be equal to -0.61 for the $1/2^-$ state of ^{163}Gd , -0.64 and -0.65 for the $1/2^-$ states of ^{165}Dy and ^{167}Dy , -0.73 for the $1/2^-$ state of ^{169}Er , -0.76 for the $1/2^-$ state of ^{171}Yb and $+0.24$ for the $1/2^-$ state of ^{181}Hf . In all cases, such values warrants that the bandhead spin assignment as $1/2$ is correct.

D. Discussion of the odd-even mass differences obtained in our calculations

Tables II and III display the three-points odd-even mass differences $\delta_q^{(3)}$ for protons and neutrons. The rms differences between experimental (as deduced from the separation energies given in Ref. [21]) and calculated in all cases for the experimental I^π assignments are 165 keV for protons and 141 keV for neutrons. We can compare these values with what has been obtained in Ref. [6] where direct fits of the pairing matrix elements V_n and V_p have been performed on the odd-even mass differences $\delta_n^{(3)}$ and $\delta_p^{(3)}$ for a similar sampling of nuclei. There, the rms deviations were calculated as 182 keV for protons and 78 keV for neutrons. Given the expected accuracy on energies of our approach, we can deem the quality of our current approach to be comparable to those of Ref. [21].

V. CONCLUDING REMARKS

Combining these results with those obtained in the same rare-earth region in Ref.[1], we can conclude that the method proposed in [1] allows a treatment of pairing correlations of the same quality as and simpler than

a lengthy and more or less localized fit process to be performed for each particle-hole interaction. However, it is *a priori* suited to describe pairing correlations only at the equilibrium deformation of well and rigidly deformed nuclei. In practice, all the calculations of the HF+BCS type within the seniority forces approach have assumed that the matrix elements obtained in these particular situations could be widely used in, e.g., computing potential-energy curves or surfaces as functions of some deformation parameters or multipole moments. Using the current approach as a starting point to determine the intensities of pairing residual interactions would free corresponding HF+BCS or HFB calculations from such ambiguities. This improvement is currently under study.

ACKNOWLEDGMENTS

T. V. Nhan Hao, P. Quentin and D. Quang Tam acknowledge the support by the Hue University under the Core Research Program, Grant no. NCM.DHH.2018.09. Another co-author (M.H. Koh) would also like to acknowledge Universiti Teknologi Malaysia for its UTMSHine grant (grant number Q.J130000.2454.09G96).

-
- [1] M.-H. Koh and P. Quentin, Determination of pairing matrix elements from average single particle level densities, arXiv:2405.09013 (submitted for review).
 - [2] A. S. Jensen and P. G. Hansen, Nucl. Phys. A **431**, 393 (1984).
 - [3] D. Madland and J. Nix, Nucl. Phys. A **476**, 1 (1988).
 - [4] P. Möller and J. Nix, Nucl. Phys. A **536**, 20 (1992).
 - [5] A. Bohr, B. R. Mottelson, and D. Pines, Phys. Rev. **110**, 936 (1958).
 - [6] N. M. Nor, N.-A. Rezle, K.-W. Kelvin-Lee, M.-H. Koh, L. Bonneau, and P. Quentin, Phys. Rev. C **99**, 064306 (2019).
 - [7] M. Brack, J. Damgaard, and A. J. *et al.*, Rev. Mod. Phys. **44**, 320 (1972).
 - [8] M. Brack and H. Pauli, Nucl. Phys. A **207**, 401 (1973).
 - [9] S. A. Moszkowski, Handbuch der Physik **XXXIX**, 469 (1957).
 - [10] P. Ring and P. Schuck, The Nuclear Many-body Problem (Springer-Verlag, 1980) p. 240.
 - [11] J. Slater, Phys. Rev. **81**, 385 (1951).
 - [12] C. Titin-Schnaider and P. Quentin, Phys. Lett. B **49**, 397 (1974).
 - [13] J. Skalski, Phys. Rev. C **63**, 024312 (2001).
 - [14] J. L. Bloas, M.-H. Koh, P. Quentin, L. Bonneau, and J. Ithnin, Phys. Rev. C **84**, 0143310 (2011).
 - [15] M. Beiner, H. Flocard, N. V. Giai, and P. Quentin, Nucl. Phys. A **238**, 29 (1975).
 - [16] N. Minkov, L. Bonneau, P. Quentin, J. Bartel, H. Molique, and D. Ivanova, Phys. Rev. C **105**, 044329 (2022).
 - [17] H. Flocard, P. Quentin, A. Kerman, and D. Vautherin, Nucl. Phys. A **203**, 433 (1973).
 - [18] J. Dobaczewski, P. Magierski, W. Nazarewicz, W. Satula, and Z. Szymanski, Phys. Rev. C **63**, 024308 (2001).
 - [19] T. Duguet, P. Bonche, P.-H. Heenen, and J. Meyer, Phys. Rev. C **65**, 014310 (2001).
 - [20] L. Bonneau, N. Minkov, D. D. Duc, P. Quentin, and J. Bartel, Phys. Rev. C **91**, 054307 (2015).
 - [21] NNDC, <https://www.nndc.bnl.gov/nudat3/>.
 - [22] S. Raman, C. N. Jr., and P. Tikkanen, At. Data Nucl. Tables **78**, 1 (2001).
 - [23] NUMOR, https://magneticmoments.info/numor/in_search.php (cut-off date 2019.03.31) and references quoted therein.
 - [24] H. Flocard, P. Quentin, and D. Vautherin, Phys. Lett. B **46**, 304 (1973).

TABLE I. Some static properties of the 22 even-even nuclei considered in this paper. The calculated total binding energies E^{theory} are given in MeV. The experimental axial quadrupole moments of the charge distribution are taken from Refs. [23] ([22] respectively) for spectroscopic first 2^+ states data noted as Q_{20}^{spect} moments (for moments deduced from reduced $B(E2)$ data noted as Q_{20}^{BE2} respectively). They are compared with the corresponding calculated moments Q_{20}^{theory} . All these moments are given in barn.

Nucleus	E^{theory}	R_{42}	Q_{20}^{spect}	Q_{20}^{BE2}	Q_{20}^{theory}
^{156}Sm	-1276.496	3.290	5.85 (7)	—	6.81
^{158}Sm	-1288.582	3.301	6.55 (14)	—	6.99
^{160}Sm	-1299.953	3.292	—	—	7.11
^{160}Gd	-1305.533	3.302	7.28 (14)	7.265 (42)	7.25
^{162}Gd	-1318.251	3.302	—	—	7.40
^{164}Gd	-1330.079	3.295	—	—	7.51
^{166}Gd	-1341.067	3.300	—	—	7.58
^{162}Dy	-1319.955	3.294	—	7.33 (8)	7.38
^{164}Dy	-1333.969	3.301	7.28 (53)	7.503 (33)	7.56
^{166}Dy	-1347.128	3.310	—	—	7.69
^{168}Dy	-1359.411	3.313	—	—	7.76
^{168}Er	-1361.733	3.309	—	7.63 (7)	7.84
^{170}Er	-1375.231	3.310	6.65 (70)	7.65 (7)	7.93
^{172}Er	-1387.702	3.314	—	—	7.72
^{170}Yb	-1374.077	3.293	7.63 (11)	7.63 (9)	7.90
^{172}Yb	-1388.724	3.305	7.77 (14)	7.792 (45)	7.98
^{174}Yb	-1402.460	3.310	7.63 (18)	7.727 (39)	7.77
^{176}Yb	-1415.594	3.310	7.98 (21)	7.30 (13)	7.58
^{178}Yb	-1427.956	3.310	—	—	7.46
^{178}Hf	-1429.291	3.291	7.07 (7)	6.961 (43)	7.22
^{180}Hf	-1442.851	3.307	7.00 (7)	6.85 (9)	7.08
^{182}Hf	-1455.193	3.295	—	—	6.85

TABLE II. Comparison of some spectroscopic properties of the 13 odd- Z nuclei considered in this paper. Along with the calculated total binding energies E^{theory} given in MeV, the intrinsic configuration spins and parities K^π of our solutions are reported. The corresponding experimental ground state values I^π (proposed or suggested - in brackets - when available) for a given nucleus are displayed for the sake of comparison, assuming the validity of the unified model assumption of $I = K$ for the band head states. Both calculated $\delta_{\text{calc}}^{(3)}$ and experimental $\delta_{\text{exp}}^{(3)}$ odd-even mass differences (given in keV) are also reported.

Nucleus	$I^\pi(\text{exp})$	$I^\pi(\text{calc})$	E^{theory}	$\delta_{\text{calc}}^{(3)}$	$\delta_{\text{exp}}^{(3)}$
^{159}Eu	$5/2^+$	$5/2^+$	-1296.280	777	554
		$5/2^-$	-1296.425	633	–
^{161}Eu	–	$5/2^-$	-1308.524	579	–
		$5/2^+$	-1308.345	757	466
^{161}Tb	$3/2^+$	$3/2^+$	-1312.161	583	600
^{163}Tb	$3/2^+$	$3/2^+$	-1325.597	513	529
^{165}Tb	$(3/2^+)$	$3/2^+$	-1338.149	455	550
^{167}Tb	$(3/2^+)$	$3/2^+$	-1349.801	438	579
^{167}Ho	$7/2^-$	$7/2^-$	-1353.783	648	508
^{169}Ho	$7/2^-$	$7/2^-$	-1366.734	588	535
^{169}Tm	$1/2^+$	$1/2^+$	-1367.174	732	602
^{171}Tm	$1/2^+$	$1/2^+$	-1381.392	585	471
^{173}Tm	$(1/2^+)$	$1/2^+$	-1394.585	496	458
^{177}Lu	$7/2^+$	$7/2^+$	-1422.045	397	579
^{179}Lu	$7/2^+$	$7/2^+$	-1435.055	348	669

TABLE III. Same as Table II for the 16 odd- N nuclei considered in this paper. Note that for the ^{165}Dy and ^{171}Yb nuclei, the experimental values of the spin and parity of a low-lying isomeric state have been also displayed.

Nucleus	$I^\pi(\text{exp})$	$I^\pi(\text{calc})$	E^{theory}	$\delta_{\text{calc}}^{(3)}$	$\delta_{\text{exp}}^{(3)}$
^{157}Sm	$(3/2^-)$	$3/2^-$	-1281.764	775	629
^{159}Sm	$5/2^-$	$5/2^-$	-1293.550	718	535
^{161}Gd	$5/2^-$	$5/2^-$	-1311.129	763	605
^{163}Gd	$(5/2^-, 7/2^+)$	$7/2^+$	-1323.398	766	
		$5/2^-$	-1322.819	1346	599
		$1/2^-$	-1323.457	707	
^{165}Gd	–	$7/2^+$	-1334.916	657	507
^{163}Dy	$5/2^-$	$5/2^-$	-1326.097	864	694
^{165}Dy	$7/2^+, 1/2_m^-$	$7/2^+$	-1339.788	760	664
		$1/2^-$	-1339.836	712	–
^{167}Dy	$(1/2^-)$	$1/2^-$	-1352.563	707	661
		$7/2^+$	-1352.620	650	–
^{169}Er	$1/2^-$	$1/2^-$	-1367.745	737	627
		$7/2^+$	-1367.817	665	–
^{171}Er	$5/2^-$	$5/2^-$	-1380.826	640	577
^{171}Yb	$1/2^-, 7/2_m^+$	$1/2^-$	-1380.625	776	703
		$7/2^+$	-1380.698	703	–
^{173}Yb	$5/2^-$	$5/2^-$	-1394.879	713	549
^{175}Yb	$7/2^-$	$7/2^-$	-1408.299	728	522
^{177}Yb	$9/2^+$	$9/2^+$	-1421.236	540	598
^{179}Hf	$9/2^+$	$9/2^+$	-1435.547	524	644
^{181}Hf	$1/2^-$	$1/2^-$	-1448.366	655	512