

BRAID GROUP ACTION AND QUANTUM QUEER SUPERALGEBRA

Jianmin Chen, Zhenhua Li and Hongying Zhu*

ABSTRACT. In this paper, we present explicit actions of braid group on the universal enveloping superalgebra $\mathbf{U}(\mathfrak{q}_n)$ and the quantum queer superalgebra $\mathbf{U}_v(\mathfrak{q}_n)$. Then we provide a new definition of root vectors and some explicit expression for them. With these procedures, we obtain the PBW-type basis containing the product of root vectors.

CONTENTS

1. Introduction	1
2. The braid group	3
3. The queer superalgebra $\mathbf{U}(\mathfrak{q}_n)$ and the action of \mathcal{B} on it	4
4. The root vectors of $\mathbf{U}(\mathfrak{q}_n)$	6
5. The PBW basis of the Kostant \mathbb{Z} -form $\mathbf{U}(\mathfrak{q}_n)_{\mathbb{Z}}$	13
6. The action of \mathcal{B} on the quantum queer superalgebra $\mathbf{U}_v(\mathfrak{q}_n)$	14
7. The root vectors of $\mathbf{U}_v(\mathfrak{q}_n)$	22
8. A PBW-type basis of $\mathbf{U}_v(\mathfrak{q}_n)$	24
References	27

1. INTRODUCTION

The queer Lie superalgebra \mathfrak{q}_n is a super analogue of the general linear Lie algebra \mathfrak{gl}_n and it differs drastically from the basic classical Lie superalgebras. One of the major difference lies in that the Cartan subalgebra of \mathfrak{q}_n is not purely even and there is no invariant bilinear form on \mathfrak{q}_n , which results in very interesting phenomenon on the highest weight space of a finite dimensional irreducible supermodule (see [6, §1.5.4]). For this reason, it is very complicated and challenging task to investigate the structure and representation theory of queer Lie superalgebra \mathfrak{q}_n (see [22]), which has received extensive attention and research.

Quantum groups first arose in the physics literature. In the work of L.D.Faddeev (see [18]), quantum groups had been developed to construct and solve ‘integrable’ quantum systems. In the original form, quantum groups are associative algebras whose defining relations are quantum R-matrix. It was realized independently by V. G. Drinfel’d (see [8], [9]) and M. Jimbo (see [17]) in the mid-1980s that these algebras are Hopf algebras. In many cases, these algebras are deformations of ‘universal enveloping algebras’ of Lie algebras.

Braid groups first appeared, albeit in a disguised form, in an article by Hurwitz published in 1891 and devoted to ramified coverings of surfaces. The notion of a braid was explicitly introduced by Artin [1] in the 1920s to formalize topological objects that model the intertwining of several strings in Euclidean 3-space. Artin pointed out that braids with a fixed number n of strings form a group, called the n -th braid group and denoted

2020 *Mathematics Subject Classification.* 17B37, 17A70, 20G42, 20C08.

Key words and phrases. quantum queer superalgebra, braid group action, root vector, PBW basis.

*Corresponding author

by \mathcal{B} . And it had proved by F. Bohnenblust in [3]. Since then, the braids and the braid groups have been extensively studied by topologists and algebraists, which led to a rich theory. Numerous ramifications now play a role in various parts of mathematics, including knot theory and low dimensional topology. And then the braid group action on quantum group was first formulated as a tool by Lusztig in [19].

The root vectors of an algebra are typically associated with the root system of a Lie algebra. In the theory of root system of a Lie algebra, root vectors are a set of specific vectors labeled by roots, which are elements of a Cartan subalgebra. Specifically, given a semisimple Lie algebra, which includes a Cartan subalgebra, the root system consists of a set of linearly independent elements taking values in this Cartan subalgebra. Each root corresponds to a root vector, and these root vectors constitute the fundamental representation of the Lie algebra.

The Poincaré-Birkhoff-Witt (PBW) theorem was initially formulated and independently proven by Birkhoff [2] and Witt [23] in 1937, which is often used as a springboard for investigating the representation theory of algebras. The classical PBW Theorem associates to any order basis of a Lie algebra is a basis of its universal enveloping algebra. In the case of a quantized universal enveloping algebra, one has no underlying Lie algebra, much less a basis of it, so it is not immediately clear how to obtain an analogue of the PBW theorem for quantum algebra.

The quantized universal enveloping algebra $\mathbf{U}_v(\mathfrak{sl}_n)$ was introduced by Drinfel'd [8]. And the braid group action on its algebraic structures, particularly its introduction in the context of quantum groups, was first formulated by Lusztig in 1988 (see [19]). And then Lusztig [20] gave analogues of 'root vectors' associated with the non-simple roots to construct a basis of $\mathbf{U}_v(\mathfrak{sl}_n)$ of PBW-type. In this work, Lusztig developed the concept of the braid group action and root vectors on quantum algebras, which plays a significant role in the study of quantum groups and their representation theory.

A quantum deformation $\mathbf{U}_v(\mathfrak{q}_n)$ of the universal enveloping algebra $\mathbf{U}(\mathfrak{q}_n)$ was constructed by Olshanski [21] using a modification of the Reshetikhin-Takhtajan-Faddeev method. In this paper, we use the definition of $\mathbf{U}_v(\mathfrak{q}_n)$ given in [12], which is equivalent to the one given by Olshanski. And then we will present a construction of the braid group action on the quantum queer superalgebra $\mathbf{U}_v(\mathfrak{q}_n)$, which will enable the definition of root vectors associated to arbitrary root of \mathfrak{q}_n . And the idea for the definition is already described in [5]: the non-simple roots vectors of \mathfrak{q}_n can be defined by using the action of a finite covering of the Weyl group on \mathfrak{q}_n ; then in the quantum case, the action of the Weyl group is replaced by that of the braid group. Moreover, $\mathbf{U}(\mathfrak{q}_n)$ is the classical limit of $\mathbf{U}_v(\mathfrak{q}_n)$ (see [14]), then we have some relevant results for the classical algebraic structures.

This paper is organized as follows. We recall the definition of the quantized universal enveloping algebra $\mathbf{U}_v(\mathfrak{sl}_n)$ and the braid group action on it in section 2. In section 3, with the basics of the queer Lie superalgebra \mathfrak{q}_n and its universal enveloping superalgebra $\mathbf{U}(\mathfrak{q}_n)$, we provide an action of braid group in $\mathbf{U}(\mathfrak{q}_n)$. After that we introduce the root vector for $\mathbf{U}(\mathfrak{q}_n)$, compute all commutation formulas for these vectors and discuss all the expression of root vectors for $\mathbf{U}(\mathfrak{q}_n)$ in section 4. In section 5, we give the PBW-type basis of the Kostant \mathbb{Z} -form. From section 6 onwards, we investigate the quantum case. In section 6, we recall the equivalent definition of quantum queer superalgebra and present a construction of the braid group action on it. In section 7, we introduce quantum root vectors for $\mathbf{U}_v(\mathfrak{q}_n)$. In the last section, we construct a PBW-type basis for the quantum queer superalgebra.

Throughout this paper, let \mathbb{Q} be the field of rational numbers, and $\mathbb{Q}(v)$ be the field of fractions in one variable v over \mathbb{Q} .

2. THE BRAID GROUP

We recall the definition of the quantized universal enveloping algebra of type A and the braid group associated to it.

Let $A_{n-1} = (a_{i,j})$ be the Cartan matrix of type A with order $n - 1$. More precisely,

$$a_{i,j} = \begin{cases} 2 & \text{if } i = j, \\ -1 & \text{if } |i - j| = 1, \\ 0 & \text{if } |i - j| > 1. \end{cases}$$

It is known that \mathfrak{sl}_n is the simple Lie algebra associated to A_{n-1} .

Definition 2.1. (See [8] and [5, Section 9.1]) The quantum algebra $\mathcal{U}_v(\mathfrak{sl}_n)$ is the associative algebra over $\mathbb{Q}(v)$ generated by E_i, F_i and K_i ($1 \leq i \leq n - 1$) with relations

$$\begin{aligned} K_i K_j &= K_j K_i, & K_i K_i^{-1} &= K_i^{-1} K_i = 1; \\ K_i E_j &= v^{a_{i,j}} E_j K_i, & K_i F_j &= v^{-a_{i,j}} F_j K_i; \\ E_i F_i - F_i E_i &= \delta_{i,j} \frac{K_i - K_i^{-1}}{v - v^{-1}}; \\ E_i E_j &= E_j E_i, & F_i F_j &= F_j F_i \quad \text{for } |i - j| > 1; \\ E_i^2 E_j - (v + v^{-1}) E_i E_j E_i + E_j E_i^2 &= 0, \\ F_i^2 F_j - (v + v^{-1}) F_i F_j F_i + F_j F_i^2 &= 0 \quad \text{for } |i - j| = 1. \end{aligned}$$

The braid group was first introduced by Artin (see [1]). In this paper, we use the definition of braid group given in [19].

Definition 2.2. The braid group $\mathcal{B} = \mathcal{B}(\mathfrak{sl}_n)$ associated to \mathfrak{sl}_n is generated by T_1, \dots, T_{n-1} with relations

$$\begin{aligned} T_i T_j T_i &= T_j T_i T_j \quad \text{if } |i - j| = 1, \\ T_i T_j &= T_j T_i \quad \text{if } |i - j| > 1. \end{aligned} \tag{2.0.1}$$

Notice that the Weyl group of \mathfrak{sl}_n is identified with the symmetric group \mathfrak{S}_n , and the defining relations of \mathcal{B} are the defining relations of \mathfrak{S}_n by replacing s_i with T_i , while the relation $s_i^2 = 1$ being removed.

The following proposition describes the action of \mathcal{B} on $\mathcal{U}_v(\mathfrak{sl}_n)$ in [19].

Proposition 2.3. [19, Proposition 5.1] *For $1 \leq i, j \leq n - 1$, the braid group \mathcal{B} acts by $\mathbb{Q}(v)$ -algebra automorphisms on $\mathcal{U}_v(\mathfrak{sl}_n)$ as*

$$\begin{aligned} T_i(K_i) &= K_i^{-1}, & T_i(E_i) &= -F_i K_i, & T_i(F_i) &= -K_i^{-1} F_i; \\ T_i(K_j) &= K_j K_i \quad \text{for } |i - j| = 1; \\ T_i(E_j) &= -E_i E_j + v^{-1} E_j E_i, & T_i(F_j) &= -F_j F_i + v F_i F_j \quad \text{for } |i - j| = 1; \\ T_i(K_j) &= K_j, & T_i(E_j) &= E_j, & T_i(F_j) &= F_j \quad \text{for } |i - j| > 1. \end{aligned}$$

Assume $w \in \mathfrak{S}_n$ is decomposed into a product of simple reflections $w = s_{i_1} s_{i_2} \dots s_{i_t}$, it is called a reduced expression (or a reduced decomposition) if t is the minimal number of simple reflections which appear in any such expression of w , and t is denoted by $l(w)$, called the length of w .

For the reduced expression $w = s_{i_1} s_{i_2} \dots s_{i_t}$, we denote $T_w := T_{i_1} T_{i_2} \dots T_{i_t}$. Assume that $w = s_{i_1} s_{i_2} \dots s_{i_t}$ and $w = s_{j_1} s_{j_2} \dots s_{j_t}$ are two different reduced expressions, by [7, Theorem 6.45], we have

$$T_{i_1} T_{i_2} \dots T_{i_t} = T_{j_1} T_{j_2} \dots T_{j_t}. \tag{2.0.2}$$

So T_w is well defined. Referring to [5, Proposition 8.1.3], for any $w_1, w_2 \in \mathfrak{S}_n$, we have

$$T_{w_1}T_{w_2} = T_{w_1w_2}, \quad \text{if } l(w_1) + l(w_2) = l(w_1w_2). \quad (2.0.3)$$

3. THE QUEER SUPERALGEBRA $\mathbf{U}(\mathfrak{q}_n)$ AND THE ACTION OF \mathcal{B} ON IT

In this section, we will study the braid group action on queer superalgebra $\mathbf{U}(\mathfrak{q}_n)$.

Following [11], let $I(n|n) = \{1, 2, \dots, n, -1, -2, \dots, -n\}$, and the elementary matrices are accordingly denoted by $E_{i,j}$, with $i, j \in I(n|n)$.

The queer Lie superalgebra \mathfrak{q}_n of order n is the Lie superalgebra consists of the matrices of the form

$$\begin{pmatrix} A & B \\ B & A \end{pmatrix}, \quad (3.0.1)$$

where A, B are $n \times n$ matrices, with rows and columns labeled by $I(n|n)$. The even part $\mathfrak{q}_n^{\bar{0}}$ of \mathfrak{q}_n consists of the matrices of the form (3.0.1) with $B = 0$, while the odd part $\mathfrak{q}_n^{\bar{1}}$ of \mathfrak{q}_n consists of the matrices with $A = 0$.

For any $i, j \in I(n|n)$, let $E_{i,j} \in \mathfrak{q}_n$ be the matrix with 1 at the (i, j) position and 0 elsewhere. It is clear that the set $\{E_{i,j} \mid i, j \in I(n|n)\}$ forms a basis of \mathfrak{q}_n . The parity of $E_{i,j}$ is defined to be

$$p(E_{i,j}) = \begin{cases} 0 & \text{if } ij > 0, \\ 1 & \text{if } ij < 0. \end{cases}$$

Let $\mathfrak{h} = \mathfrak{h}^{\bar{0}} \oplus \mathfrak{h}^{\bar{1}}$ be the standard Cartan subalgebra of \mathfrak{q}_n . Denote

$$h_i = E_{i,i} + E_{-i,-i}, \quad h_{\bar{i}} = E_{-i,i} + E_{i,-i}, \quad i \in \{1, 2, \dots, n\}.$$

It is known that $\mathfrak{h}^{\bar{0}}$ has basis $\{h_1, \dots, h_n\}$, while $\mathfrak{h}^{\bar{1}}$ has basis $\{h_{\bar{1}}, \dots, h_{\bar{n}}\}$.

Let $\{\epsilon_1, \dots, \epsilon_n\}$ be the basis of $(\mathfrak{h}^{\bar{0}})^*$ dual to $\{h_1, \dots, h_n\}$. The dot product on $\{\epsilon_1, \dots, \epsilon_n\}$ is defined as $\epsilon_i \cdot \epsilon_j = \delta_{i,j}$.

For any $1 \leq i \neq j \leq n$, denote

$$\alpha_{i,j} := \epsilon_i - \epsilon_j. \quad (3.0.2)$$

According to [6, Section 1.2.6], the standard Borel subalgebra \mathfrak{b} of \mathfrak{q}_n consists of the matrices of the form (3.0.1) with A, B being upper triangular, and the set of all positive roots corresponding to \mathfrak{b} is

$$\Phi^+ = \{\alpha_{i,j} = \epsilon_i - \epsilon_j \mid 1 \leq i < j \leq n\},$$

and the set of all negative roots is

$$\Phi^- = \{\alpha_{i,j} = \epsilon_i - \epsilon_j \mid 1 \leq i < j \leq n\}.$$

Denote $\alpha_j = \alpha_{j,j+1} = \epsilon_j - \epsilon_{j+1}$ for all $j \in [1, n-1]$, then the set of all simple positive roots could be denoted as

$$\Pi^+ = \{\alpha_j \mid 1 \leq j \leq n-1\}.$$

Denote $\mathbb{Q} := \bigoplus_{j=1}^{n-1} \mathbb{Z}\alpha_j$ to be the set of root lattice.

According to [6, Section 1.2.6], the Weyl group of \mathfrak{q}_n is identified with the symmetric group \mathfrak{S}_n , which is also the Weyl group of \mathfrak{sl}_n . As there is an isomorphism

$$\begin{aligned} \mathfrak{h}^{\bar{0}} &\longrightarrow (\mathfrak{h}^{\bar{0}})^*, \\ h_i &\mapsto \epsilon_i, \quad h_i - h_{i+1} \mapsto \alpha_i, \end{aligned}$$

we get an action of \mathfrak{S}_n on $(\mathfrak{h}^{\bar{0}})^*$ as

$$s_i(\xi) = \xi - \xi(h_i - h_{i+1})\alpha_i \quad (\xi \in (\mathfrak{h}^{\bar{0}})^*). \quad (3.0.3)$$

It is verified that

$$s_i(\alpha_j) = \begin{cases} -\alpha_j & \text{for } i = j, \\ \alpha_i + \alpha_j & \text{for } |i - j| = 1, \\ \alpha_j & \text{otherwise,} \end{cases} \quad (3.0.4)$$

and

$$s_i(\epsilon_j) = \epsilon_{s_i(j)}. \quad (3.0.5)$$

The queer superalgebra $\mathbf{U}(\mathfrak{q}_n)$ is the universal enveloping superalgebra of \mathfrak{q}_n . For any homogeneous $x, y \in \mathbf{U}(\mathfrak{q}_n)$, the Lie superbracket is defined as

$$[x, y] = xy - (-1)^{p(x)p(y)}yx,$$

where for homogeneous z , $p(z)$ is the parity of z .

Following [14, Proposition 1.1], $\mathbf{U}(\mathfrak{q}_n)$ could also be presented as an associative algebra as follows.

Proposition 3.1. *The universal enveloping superalgebra $\mathbf{U}(\mathfrak{q}_n)$ is the associative superalgebra over \mathbb{Q} generated by even generators h_i, e_j, f_j , and odd generators $h_{\bar{i}}, e_{\bar{j}}, f_{\bar{j}}$, with $1 \leq i \leq n$ and $1 \leq j \leq n - 1$, subjected to the following relations:*

$$\begin{aligned} \text{(QS1)} \quad & [h_i, h_j] = 0, \quad [h_i, h_{\bar{j}}] = 0, \quad [h_{\bar{i}}, h_{\bar{j}}] = \delta_{i,j}2h_i; \\ \text{(QS2)} \quad & [h_i, e_j] = (\epsilon_i, \alpha_j)e_j, \quad [h_i, e_{\bar{j}}] = (\epsilon_i, \alpha_j)e_{\bar{j}}; \\ & [h_i, f_j] = -(\epsilon_i, \alpha_j)f_j, \quad [h_i, f_{\bar{j}}] = -(\epsilon_i, \alpha_j)f_{\bar{j}}; \\ \text{(QS3)} \quad & [h_{\bar{i}}, e_j] = (\epsilon_i, \alpha_j)e_{\bar{j}}, \quad [h_{\bar{i}}, f_j] = -(\epsilon_i, \alpha_j)f_{\bar{j}}; \\ & [h_{\bar{i}}, e_{\bar{j}}] = \begin{cases} e_j & \text{if } i = j \text{ or } j + 1, \\ 0 & \text{otherwise,} \end{cases} \quad [h_{\bar{i}}, f_{\bar{j}}] = \begin{cases} f_j & \text{if } i = j \text{ or } j + 1, \\ 0 & \text{otherwise;} \end{cases} \\ \text{(QS4)} \quad & [e_i, f_j] = \delta_{i,j}(h_i - h_{i+1}), \quad [e_{\bar{i}}, f_{\bar{j}}] = \delta_{i,j}(h_i + h_{i+1}), \\ & [e_{\bar{i}}, f_j] = \delta_{i,j}(h_{\bar{i}} - h_{\bar{i}+1}), \quad [e_i, f_{\bar{j}}] = \delta_{i,j}(h_{\bar{i}} - h_{\bar{i}+1}); \\ \text{(QS5)} \quad & [e_i, e_{\bar{j}}] = [e_{\bar{i}}, e_j] = [f_i, f_{\bar{j}}] = [f_{\bar{i}}, f_j] = 0 \quad \text{for } |i - j| \neq 1, \\ & [e_i, e_j] = [f_i, f_j] = 0 \quad \text{for } |i - j| > 1, \\ & [e_i, e_{i+1}] = [e_{\bar{i}}, e_{\bar{i}+1}], \quad [e_i, e_{\bar{i}+1}] = [e_{\bar{i}}, e_{i+1}], \\ & [f_{i+1}, f_i] = [f_{\bar{i}+1}, f_{\bar{i}}], \quad [f_{i+1}, f_{\bar{i}}] = [f_{\bar{i}+1}, f_i]; \\ \text{(QS6)} \quad & [e_i, [e_i, e_j]] = [e_{\bar{i}}, [e_i, e_j]] = 0, \quad [f_i, [f_i, f_j]] = [f_{\bar{i}}, [f_i, f_j]] = 0 \quad \text{for } |i - j| = 1. \end{aligned}$$

Remark 3.2. There is an anti-involution ω of $\mathbf{U}(\mathfrak{q}_n)$ defined on the generators as

$$\begin{aligned} \omega(h_i) &= h_i, & \omega(e_j) &= f_j, & \omega(f_j) &= e_j, \\ \omega(h_{\bar{i}}) &= h_{\bar{i}}, & \omega(e_{\bar{j}}) &= f_{\bar{j}}, & \omega(f_{\bar{j}}) &= e_{\bar{j}}, \end{aligned}$$

with $1 \leq i \leq n$ and $1 \leq j \leq n - 1$.

Considering Proposition 2.3, it is natural to think about the action of \mathcal{B} on $\mathbf{U}(\mathfrak{q}_n)$.

Proposition 3.3. *There is an action of the braid group \mathcal{B} by superalgebra automorphism of $\mathbf{U}(\mathfrak{q}_n)$ on the generators defined as*

$$\begin{aligned} T_i(h_j) &= h_{s_i(j)}, & T_i(h_{\bar{j}}) &= h_{\overline{s_i(j)}}; \\ T_i(e_i) &= -f_i, & T_i(f_i) &= -e_i, & T_i(e_{\bar{i}}) &= -f_{\bar{i}}, & T_i(f_{\bar{i}}) &= -e_{\bar{i}}; \\ T_i(e_j) &= -e_i e_j + e_j e_i, & T_i(f_j) &= -f_j f_i + f_i f_j, \\ T_i(e_{\bar{j}}) &= -e_i e_{\bar{j}} + e_{\bar{j}} e_i, & T_i(f_{\bar{j}}) &= -f_{\bar{j}} f_i + f_i f_{\bar{j}} & \text{for } |i - j| = 1; \\ T_i(e_j) &= e_j, & T_i(f_j) &= f_j, & T_i(e_{\bar{j}}) &= e_{\bar{j}}, & T_i(f_{\bar{j}}) &= f_{\bar{j}} & \text{for } |i - j| > 1. \end{aligned}$$

And the action of T_i^{-1} is given by

$$\begin{aligned} T_i^{-1}(h_j) &= h_{s_i(j)}, & T_i^{-1}(h_{\bar{j}}) &= h_{\overline{s_i(j)}}; \\ T_i^{-1}(e_i) &= -f_i, & T_i^{-1}(f_i) &= -e_i, & T_i^{-1}(e_{\bar{i}}) &= -f_{\bar{i}}, & T_i^{-1}(f_{\bar{i}}) &= -e_{\bar{i}}; \\ T_i^{-1}(e_j) &= -e_j e_i + e_i e_j, & T_i^{-1}(f_j) &= -f_i f_j + f_j f_i, \\ T_i^{-1}(e_{\bar{j}}) &= -e_j e_{\bar{i}} + e_{\bar{i}} e_j, & T_i^{-1}(f_{\bar{j}}) &= -f_{\bar{i}} f_j + f_j f_{\bar{i}} & \text{for } |i - j| = 1; \\ T_i^{-1}(e_j) &= e_j, & T_i^{-1}(f_j) &= f_j, & T_i^{-1}(e_{\bar{j}}) &= e_{\bar{j}}, & T_i^{-1}(f_{\bar{j}}) &= f_{\bar{j}} & \text{for } |i - j| > 1. \end{aligned}$$

Proof. The proof is similar to Theorem 6.3 in Section 6, so we give a sketch here. And by a direct calculation, it is easy to verify that each T_i holds the relations in (QS1)-(QS6). On the other hand, the action of T_i 's on each generator of $\mathbf{U}(\mathfrak{q}_n)$ satisfy the relations in (2.0.1). \square

Remark 3.4. For any $1 \leq i, j \leq n - 1$, $|i - j| = 1$, direct calculation shows that

$$T_i T_j(e_i) = e_j, \quad T_i T_j(e_{\bar{i}}) = e_{\bar{j}}, \quad T_i T_j(f_i) = f_j, \quad T_i T_j(f_{\bar{i}}) = f_{\bar{j}}.$$

Following [16, Proposition 8.20] and with the same discussion, we have the following.

Corollary 3.5. *Assume $w \in \mathfrak{S}_n$ and $\alpha_i, \alpha_k \in \Pi^+$, such that $w(\alpha_i) = \alpha_k$. Then we have*

$$T_w(e_i) = e_k, \quad T_w(e_{\bar{i}}) = e_{\bar{k}}, \quad T_w(f_i) = f_k, \quad T_w(f_{\bar{i}}) = f_{\bar{k}}.$$

4. THE ROOT VECTORS OF $\mathbf{U}(\mathfrak{q}_n)$

In this section, we will study the root vectors of the queer superalgebra $\mathbf{U}(\mathfrak{q}_n)$. Let's start with the relations between the positive roots and the longest element w_0 in the Weyl group \mathfrak{S}_n of \mathfrak{q}_n .

We denote $N = l(w_0)$. For convenience, for any $t \geq 0$ and $\sigma = (i_1, i_2, \dots, i_t) \in \mathbb{N}^t$, we denote $w^\sigma = s_{i_1} s_{i_2} \dots s_{i_t}$. Notice that $w^\sigma = 1$ for the case $t = 0$. When $\sigma = (i_1, i_2, \dots, i_N) \in \mathbb{N}^N$ and $w_0 = w^\sigma = s_{i_1} s_{i_2} \dots s_{i_N}$ is a reduced expression, as \mathfrak{sl}_n and \mathfrak{q}_n have the same simple positive roots, following [5], by the action of the Weyl group on \mathfrak{q}_n , the set of positive roots of \mathfrak{q}_n associated with σ can be described as

$$\Delta_\sigma^+ = \{\beta_{\sigma,1} = \alpha_{i_1}, \quad \beta_{\sigma,2} = s_{i_1}(\alpha_{i_2}), \quad \dots, \quad \beta_{\sigma,N} = s_{i_1} s_{i_2} \dots s_{i_{N-1}}(\alpha_{i_N})\}. \quad (4.0.1)$$

Remark 4.1. According to [7] and [12, Section 2],

$$\sharp \Delta_\sigma^+ = \sharp \Phi^+ = \frac{n(n-1)}{2}, \quad \Delta_\sigma^+ = \Phi^+ = \{\alpha_{i,j} = \epsilon_i - \epsilon_j | 1 \leq i < j \leq n\}, \quad (4.0.2)$$

hence the set Δ_σ^+ does not depend on the chosen of σ and we may denote it as Δ^+ .

Definition 4.2. Fix a reduced decomposition $w_0 = s_{i_1} s_{i_2} \cdots s_{i_N}$ of the longest element in the Weyl group of \mathfrak{q}_n . Define the positive root vectors $\mathbf{e}_k^\sigma, \bar{\mathbf{e}}_k^\sigma$ and negative root vectors $\mathbf{f}_k^\sigma, \bar{\mathbf{f}}_k^\sigma$ of $\mathbf{U}(\mathfrak{q}_n)$ as ($1 \leq k \leq N$)

$$\begin{aligned} \mathbf{e}_k^\sigma &= T_{i_1} T_{i_2} \cdots T_{i_{k-1}}(e_{i_k}), & \bar{\mathbf{e}}_k^\sigma &= T_{i_1} T_{i_2} \cdots T_{i_{k-1}}(e_{\bar{i}_k}), \\ \mathbf{f}_k^\sigma &= T_{i_1} T_{i_2} \cdots T_{i_{k-1}}(f_{i_k}), & \bar{\mathbf{f}}_k^\sigma &= T_{i_1} T_{i_2} \cdots T_{i_{k-1}}(f_{\bar{i}_k}). \end{aligned} \quad (4.0.3)$$

For any $\beta_{\sigma,t} = \alpha_{i,j} \in \Phi^+ \cup \Phi^-$ ($1 \leq i < j \leq n$), we can denote

$$\begin{aligned} \mathbf{e}_{i,j}^\sigma &= \mathbf{e}_t^\sigma, & \bar{\mathbf{e}}_{i,j}^\sigma &= \bar{\mathbf{e}}_t^\sigma, \\ \mathbf{e}_{j,i}^\sigma &= \mathbf{f}_t^\sigma, & \bar{\mathbf{e}}_{j,i}^\sigma &= \bar{\mathbf{f}}_t^\sigma. \end{aligned}$$

And we denote the set of positive and negative root vectors as

$$\begin{aligned} \psi_\sigma^+ &= \{\mathbf{e}_t^\sigma, \bar{\mathbf{e}}_t^\sigma \mid 1 \leq t \leq N\} = \{\mathbf{e}_{i,j}^\sigma, \bar{\mathbf{e}}_{i,j}^\sigma \mid 1 \leq i < j \leq N\}, \\ \psi_\sigma^- &= \{\mathbf{f}_t^\sigma, \bar{\mathbf{f}}_t^\sigma \mid 1 \leq t \leq N\} = \{\mathbf{e}_{j,i}^\sigma, \bar{\mathbf{e}}_{j,i}^\sigma \mid 1 \leq i < j \leq N\}. \end{aligned} \quad (4.0.4)$$

Example 4.3. Fix $n = 3$, there are two choices for reduced decomposition of w_0 .

(1) If we choose $\sigma = (1,2,1)$ and $w_0 = w^\sigma = s_1 s_2 s_1$, the positive roots are

$$\beta_{\sigma,1} = \alpha_1 = \alpha_{1,2}, \quad \beta_{\sigma,2} = s_1(\alpha_2) = \alpha_{1,3}, \quad \beta_{\sigma,3} = s_1 s_2(\alpha_1) = \alpha_{2,3},$$

while the corresponding positive root vectors are

$$\begin{aligned} \mathbf{e}_{1,2}^\sigma &= e_1, & \mathbf{e}_{1,3}^\sigma &= T_1(e_2) = -e_1 e_2 + e_2 e_1, & \mathbf{e}_{2,3}^\sigma &= T_1 T_2(e_1) = e_2, \\ \bar{\mathbf{e}}_{1,2}^\sigma &= e_{\bar{1}}, & \bar{\mathbf{e}}_{1,3}^\sigma &= T_1(e_{\bar{2}}) = -e_1 e_{\bar{2}} + e_{\bar{2}} e_1, & \bar{\mathbf{e}}_{2,3}^\sigma &= T_1 T_2(e_{\bar{1}}) = e_{\bar{2}}. \end{aligned}$$

(2) If we choose $\gamma = (2,1,2)$ and $w_0 = w^\gamma = s_2 s_1 s_2$, then

$$\beta_{\gamma,1} = \alpha_2 = \alpha_{2,3}, \quad \beta_{\gamma,2} = s_2(\alpha_1) = \alpha_{1,3}, \quad \beta_{\gamma,3} = s_2 s_1(\alpha_2) = \alpha_{1,2},$$

and the corresponding positive root vectors are

$$\begin{aligned} \mathbf{e}_{2,3}^\gamma &= e_2, & \mathbf{e}_{1,3}^\gamma &= T_2(e_1) = -e_2 e_1 + e_1 e_2, & \mathbf{e}_{1,2}^\gamma &= T_2 T_1(e_2) = e_1, \\ \bar{\mathbf{e}}_{2,3}^\gamma &= e_{\bar{2}}, & \bar{\mathbf{e}}_{1,3}^\gamma &= T_2(e_{\bar{1}}) = -e_2 e_{\bar{1}} + e_{\bar{1}} e_2, & \bar{\mathbf{e}}_{1,2}^\gamma &= T_2 T_1(e_{\bar{2}}) = e_{\bar{1}}. \end{aligned}$$

In the following of this paper, we fix σ for a reduced decomposition of the longest element \mathfrak{S}_n as follows:

$$\begin{aligned} \sigma &= (1, \quad 2, 1, \quad 3, 2, 1, \quad \cdots, \quad n-1, n-2, \cdots, 2, 1) \in \mathbb{N}^N, \\ w_0 = w^\sigma &= (s_1) \cdot (s_2 s_1) \cdot (s_3 s_2 s_1) \cdots \cdots (s_{n-1} \cdots s_1). \end{aligned} \quad (4.0.5)$$

By a direct computation, we conclude the following lemma.

Lemma 4.4. *Suppose that $1 \leq i < j \leq n$. If $j = i + 1$, then*

$$\beta_{\sigma, \frac{i(i+1)}{2}} = (s_1) \cdot (s_2 s_1) \cdots \cdots (s_i s_{i-1} \cdots s_2)(\alpha_1) = \alpha_{i,i+1};$$

if $j > i + 1$, then

$$\beta_{\sigma, \frac{(j-2)(j-1)}{2} + i - 1} = (s_1) \cdot (s_2 s_1) \cdots \cdots (s_{j-1} s_{j-2} \cdots s_{j-i+1})(\alpha_{j-i}) = \alpha_{i,j}.$$

Proof. By the group action in (3.0.5), we see that

$$\begin{aligned}
& (s_1) \cdot (s_2 s_1) \cdots \cdots (s_i s_{i-1} \cdots s_2)(\alpha_1) \\
&= (s_1) \cdot (s_2 s_1) \cdots \cdots (s_i s_{i-1} \cdots s_2)(\epsilon_1 - \epsilon_2) \\
&= (s_1) \cdot (s_2 s_1) \cdots \cdots (s_i s_{i-1} \cdots s_2)(\epsilon_1) - (s_1) \cdot (s_2 s_1) \cdots \cdots (s_i s_{i-1} \cdots s_2)(\epsilon_2) \\
&= \epsilon_{(s_1) \cdot (s_2 s_1) \cdots \cdots (s_i s_{i-1} \cdots s_2)}(1) - \epsilon_{(s_1) \cdot (s_2 s_1) \cdots \cdots (s_i s_{i-1} \cdots s_2)}(2) \\
&= \epsilon_{(s_1) \cdot (s_2 s_1) \cdots \cdots (s_{i-1} \cdots s_1)}(1) - \epsilon_{(s_1) \cdot (s_2 s_1) \cdots \cdots (s_{i-1} \cdots s_1)}(i+1) \\
&= \epsilon_{(s_1) \cdot (s_2 s_1) \cdots \cdots (s_{i-2} \cdots s_1)}(i) - \epsilon_{i+1} \\
&= \epsilon_i - \epsilon_{i+1} \\
&= \alpha_{i, i+1}.
\end{aligned}$$

Hence, we have proved the first equation. For the second equation, we can prove it directly. For any $1 \leq i, j \leq n$ with $j - i > 1$,

$$\begin{aligned}
& (s_1) \cdot (s_2 s_1) \cdots \cdots (s_{j-1} s_{j-2} \cdots s_{j-i+1})(\alpha_{j-i}) \\
&= (s_1) \cdot (s_2 s_1) \cdots \cdots (s_{j-1} s_{j-2} \cdots s_{j-i+1})(\epsilon_{j-i} - \epsilon_{j-i+1}) \\
&= (s_1) \cdot (s_2 s_1) \cdots \cdots (s_{j-1} s_{j-2} \cdots s_{j-i+1})(\epsilon_{j-i}) \\
&\quad - (s_1) \cdot (s_2 s_1) \cdots \cdots (s_{j-1} s_{j-2} \cdots s_{j-i+1})(\epsilon_{j-i+1}) \\
&= \epsilon_{(s_1) \cdot (s_2 s_1) \cdots \cdots (s_{j-1} s_{j-2} \cdots s_{j-i+1})}(j-i) - \epsilon_{(s_1) \cdot (s_2 s_1) \cdots \cdots (s_{j-1} s_{j-2} \cdots s_{j-i+1})}(j-i+1) \\
&= \epsilon_{(s_1) \cdot (s_2 s_1) \cdots \cdots (s_{j-2} \cdots s_{j-i-1})}(j-i) - \epsilon_j \\
&= \epsilon_{(s_1) \cdot (s_2 s_1) \cdots \cdots (s_{j-3} \cdots s_{j-i-2})}(j-i-1) - \epsilon_j \\
&\quad \vdots \\
&= \epsilon_{(s_1) \cdot (s_2 s_1) \cdots \cdots (s_i \cdots s_1)}(2) - \epsilon_j \\
&= \epsilon_{(s_1) \cdot (s_2 s_1) \cdots \cdots (s_{i-1} \cdots s_1)}(1) - \epsilon_j \\
&= \epsilon_i - \epsilon_j \\
&= \alpha_{i, j}.
\end{aligned}$$

□

Corollary 4.5. *Assume that $1 \leq i < n$, $i + 1 < j < n$. Then we have*

$$\begin{aligned}
e_{i, i+1}^\sigma &= e_i, & \bar{e}_{i, i+1}^\sigma &= e_{\bar{i}}, & e_{i+1, i}^\sigma &= f_i, & \bar{e}_{i+1, i}^\sigma &= \bar{f}_i, \\
e_{i, j}^\sigma &= -e_{i, j-1}^\sigma e_{j-1, j}^\sigma + e_{j-1, j}^\sigma e_{i, j-1}^\sigma = [\cdots [[e_i, -e_{i+1}], -e_{i+2}], \cdots, -e_{j-1}], \\
\bar{e}_{i, j}^\sigma &= -\bar{e}_{i, j-1}^\sigma \bar{e}_{j-1, j}^\sigma + \bar{e}_{j-1, j}^\sigma \bar{e}_{i, j-1}^\sigma = [\cdots [[e_i, -e_{i+1}], -e_{i+2}], \cdots, -e_{\bar{j}-1}], \\
e_{j, i}^\sigma &= -e_{j, j-1}^\sigma e_{j-1, i}^\sigma + e_{j-1, i}^\sigma e_{j, j-1}^\sigma = [-f_{j-1}, [-f_{j-2}, \cdots [-f_{i+1}, f_i]] \cdots], \\
\bar{e}_{j, i}^\sigma &= -\bar{e}_{j, j-1}^\sigma \bar{e}_{j-1, i}^\sigma + e_{j-1, i}^\sigma \bar{e}_{j, j-1}^\sigma = [-\bar{f}_{j-1}, [-f_{j-2}, \cdots [-f_{i+1}, f_i]] \cdots].
\end{aligned}$$

Proof. By Definition 4.2 and Remark 6.6, we see that

$$\begin{aligned}
e_{i, i+1}^\sigma &= (T_1) \cdot (T_2 T_1) \cdots \cdots (T_{i-1} T_{i-2} \cdots T_1) \cdot (T_i T_{i-1} \cdots T_3 T_2)(e_1) \\
&= (T_1) \cdot (T_2 T_1) \cdots \cdots (T_{i-1} T_{i-2} \cdots T_2) \cdot (T_i T_{i-1} \cdots T_3) \cdot T_1 T_2(e_1) \\
&= (T_1) \cdot (T_2 T_1) \cdots \cdots (T_{i-1} T_{i-2} \cdots T_2) \cdot (T_i T_{i-1} \cdots T_3)(e_2) \\
&= (T_1) \cdot (T_2 T_1) \cdots \cdots (T_{i-1} T_{i-2} \cdots T_3) \cdot (T_i T_{i-1} \cdots T_4) \cdot T_2 T_3(e_2) \\
&= (T_1) \cdot (T_2 T_1) \cdots \cdots (T_{i-1} T_{i-2} \cdots T_3) \cdot (T_i T_{i-1} \cdots T_4)(e_3) \\
&\quad \vdots
\end{aligned}$$

$$\begin{aligned}
&= (T_1) \cdot (T_2 T_1) \cdots \cdots (T_{i-2} \cdots T_1)(e_i) \\
&= e_i.
\end{aligned}$$

Then the first equation is proved. More generally, for any $1 \leq i, j \leq n, j > i + 1$, we have

$$\begin{aligned}
\mathbf{e}_{i,j}^\sigma &= (T_1) \cdot (T_2 T_1) \cdots \cdots (T_{i-1} \cdots T_1) \cdot (T_i \cdots T_1) \cdots \cdots (T_{j-1} \cdots T_{j-i+1})(e_{j-i}) \\
&= (T_1) \cdot (T_2 T_1) \cdots \cdots (T_{i-1} \cdots T_2) \cdot [(T_i \cdots T_3) \cdot (T_1 T_2 T_1)] \cdots \cdots (T_{j-1} \cdots T_{j-i+1})(e_{j-i}) \\
&\quad \vdots \\
&= (T_1) \cdot (T_2 T_1) \cdots \cdots (T_{i-1} \cdots T_2) \cdot (T_i \cdots T_1) \cdots \cdots (T_{j-1} \cdots T_{j-i+2}) \cdot T_{j-i} T_{j-i+1}(e_{j-i}) \\
&= (T_1) \cdot (T_2 T_1) \cdots \cdots (T_{i-1} \cdots T_2) \cdot (T_i \cdots T_1) \cdots \cdots (T_{j-1} \cdots T_{j-i+2})(e_{j-i+1}) \\
&= (T_1) \cdot (T_2 T_1) \cdots \cdots (T_{i-1} \cdots T_3)(T_i \cdots T_1) \cdots \cdots (T_{j-1} \cdots T_{j-i+3})(e_{j-i+2}) \\
&\quad \vdots \\
&= (T_1) \cdot (T_2 T_1) \cdots \cdots (T_{i-2} \cdots T_1) \cdot (T_i \cdots T_1) \cdots \cdots (T_{j-2} \cdots T_1)(e_{j-1}) \\
&= T_i T_{i+1} \cdots T_{j-2}(e_{j-1}).
\end{aligned}$$

Applying the last equation by replacing j with $j + 1$, we have

$$\begin{aligned}
\mathbf{e}_{i,j+1}^\sigma &= (T_i T_{i+1} \cdots T_{j-2}) \cdot T_{j-1}(e_j) \\
&= (T_i T_{i+1} \cdots T_{j-2})(-e_{j-1} e_j + e_j e_{j-1}) \\
&= -(T_i T_{i+1} \cdots T_{j-2})(e_{j-1}) \cdot (T_i T_{i+1} \cdots T_{j-2})(e_j) \\
&\quad + (T_i T_{i+1} \cdots T_{j-2})(e_j) \cdot (T_i T_{i+1} \cdots T_{j-2})(e_{j-1}) \\
&= -\mathbf{e}_{i,j}^\sigma e_j + e_j \mathbf{e}_{i,j}^\sigma.
\end{aligned}$$

And thus we have proved the equation for $\mathbf{e}_{i,j}^\sigma$. With a parallel argument, the remaining formulas could be proved. \square

As a natural consequence, we will provide super commutation formulas for these root vectors.

Proposition 4.6. *The following holds for $1 \leq i, j, k, l \leq n$ satisfying $i < j, k < l$ in $U(\mathfrak{q}_n)$:*

(1)

$$\begin{aligned}
[\mathbf{e}_{i,j}^\sigma, \mathbf{e}_{k,l}^\sigma] &= \begin{cases} \mathbf{e}_{k,j}^\sigma & \text{if } i = l \ \& \ j \neq k, \\ -\mathbf{e}_{i,l}^\sigma & \text{if } j = k \ \& \ i \neq l, \\ 0 & \text{otherwise;} \end{cases} \\
[\mathbf{e}_{i,j}^\sigma, \mathbf{e}_{l,k}^\sigma] &= \begin{cases} \mathbf{e}_{i,j}^\sigma & \text{if } i = k \ \& \ j \neq l, \\ -\mathbf{e}_{i,k}^\sigma & \text{if } j = l \ \& \ i \neq k, \\ h_i - h_j & \text{if } i = k \ \& \ j \neq l, \\ 0 & \text{otherwise;} \end{cases} \\
[\mathbf{e}_{j,i}^\sigma, \mathbf{e}_{l,k}^\sigma] &= \begin{cases} -\mathbf{e}_{j,k}^\sigma & \text{if } i = l \ \& \ j \neq k, \\ \mathbf{e}_{i,l}^\sigma & \text{if } j = k \ \& \ i \neq l, \\ 0 & \text{otherwise.} \end{cases}
\end{aligned}$$

(2)

$$\begin{aligned}
[e_{i,j}^\sigma, \bar{e}_{k,l}^\sigma] &= \begin{cases} \bar{e}_{k,j}^\sigma & \text{if } i = l \ \& \ j \neq k, \\ -\bar{e}_{i,l}^\sigma & \text{if } j = k \ \& \ i \neq l, \\ 0 & \text{otherwise ;} \end{cases} \\
[e_{i,j}^\sigma, \bar{e}_{l,k}^\sigma] &= \begin{cases} \bar{e}_{l,j}^\sigma & \text{if } i = k \ \& \ j \neq l, \\ -\bar{e}_{i,k}^\sigma & \text{if } j = l \ \& \ i \neq k, \\ h_{\bar{i}} - h_{\bar{j}} & \text{if } i = k \ \& \ j \neq l, \\ 0 & \text{otherwise ;} \end{cases} \\
[\bar{e}_{i,j}^\sigma, e_{l,k}^\sigma] &= \begin{cases} \bar{e}_{i,j}^\sigma & \text{if } i = k \ \& \ j \neq l, \\ -\bar{e}_{i,k}^\sigma & \text{if } j = l \ \& \ i \neq k, \\ h_{\bar{i}} - h_{\bar{j}} & \text{if } i = k \ \& \ j \neq l, \\ 0 & \text{otherwise ;} \end{cases} \\
[e_{j,i}^\sigma, \bar{e}_{l,k}^\sigma] &= \begin{cases} -\bar{e}_{j,k}^\sigma & \text{if } i = l \ \& \ j \neq k, \\ \bar{e}_{i,l}^\sigma & \text{if } j = k \ \& \ i \neq l, \\ 0 & \text{otherwise .} \end{cases}
\end{aligned}$$

(3)

$$\begin{aligned}
[\bar{e}_{i,j}^\sigma, \bar{e}_{k,l}^\sigma] &= \begin{cases} -e_{k,j}^\sigma & \text{if } i = l \ \& \ j \neq k, \\ -e_{i,l}^\sigma & \text{if } j = k \ \& \ i \neq l, \\ 0 & \text{otherwise ;} \end{cases} \\
[\bar{e}_{i,j}^\sigma, \bar{e}_{l,k}^\sigma] &= \begin{cases} -e_{l,j}^\sigma & \text{if } i = k \ \& \ j \neq l, \\ -e_{i,k}^\sigma & \text{if } j = l \ \& \ i \neq k, \\ h_i + h_j & \text{if } i = k \ \& \ j \neq l, \\ 0 & \text{otherwise ;} \end{cases} \\
[\bar{e}_{j,i}^\sigma, \bar{e}_{l,k}^\sigma] &= \begin{cases} -e_{j,k}^\sigma & \text{if } i = l \ \& \ j \neq k, \\ -e_{i,l}^\sigma & \text{if } j = k \ \& \ i \neq l, \\ 0 & \text{otherwise .} \end{cases}
\end{aligned}$$

(4)

$$\begin{aligned}
[h_k, e_{i,j}^\sigma] &= \begin{cases} e_{i,j}^\sigma & \text{if } k = i, \\ -e_{i,j}^\sigma & \text{if } k = j, \\ 0 & \text{otherwise ;} \end{cases} \\
[h_k, e_{j,i}^\sigma] &= \begin{cases} -e_{j,i}^\sigma & \text{if } k = i, \\ e_{j,i}^\sigma & \text{if } k = j, \\ 0 & \text{otherwise ;} \end{cases} \\
[h_k, \bar{e}_{i,j}^\sigma] &= \begin{cases} \bar{e}_{i,j}^\sigma & \text{if } k = i, \\ -\bar{e}_{i,j}^\sigma & \text{if } k = j, \\ 0 & \text{otherwise ;} \end{cases}
\end{aligned}$$

$$(5) \quad [h_k, \bar{e}_{j,i}^\sigma] = \begin{cases} -\bar{e}_{j,i}^\sigma & \text{if } k = i, \\ \bar{e}_{j,i}^\sigma & \text{if } k = j, \\ 0 & \text{otherwise.} \end{cases}$$

$$\begin{aligned} [h_{\bar{k}}, e_{i,j}^\sigma] &= \begin{cases} \bar{e}_{i,j}^\sigma & \text{if } k = i, \\ -\bar{e}_{i,j}^\sigma & \text{if } k = j, \\ 0 & \text{otherwise;} \end{cases} \\ [h_{\bar{k}}, e_{j,i}^\sigma] &= \begin{cases} -\bar{e}_{j,i}^\sigma & \text{if } k = i, \\ \bar{e}_{j,i}^\sigma & \text{if } k = j, \\ 0 & \text{otherwise;} \end{cases} \\ [h_{\bar{k}}, \bar{e}_{i,j}^\sigma] &= \begin{cases} e_{i,j}^\sigma & \text{if } k = i \text{ or } k = j, \\ 0 & \text{otherwise;} \end{cases} \\ [h_{\bar{k}}, \bar{e}_{j,i}^\sigma] &= \begin{cases} e_{j,i}^\sigma & \text{if } k = i \text{ or } k = j, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Proof. A direct calculation shows that

$$\begin{aligned} [e_{i,j}^\sigma, f_i] &= [[\cdots [[e_i, -e_{i+1}], -e_{i+2}], \cdots, -e_{j-1}], f_i] \\ &= [\cdots [[e_i, -e_{i+1}], f_i], -e_{i+2}], \cdots, -e_{j-1}] \\ &= [f_i, [\cdots [[e_i, -e_{i+1}], -e_{i+2}], \cdots, -e_{j-1}]] - e_{i+1,j}^\sigma \end{aligned}$$

Where the third equality is due to the relation $[e_i, f_i] = h_i - h_{i+1}$ in Proposition 3.1. Similarly, the following holds:

$$[e_{i,j}^\sigma, f_k] = \begin{cases} e_{i+1,j}^\sigma & \text{if } k = i, \\ -e_{i,j-1}^\sigma & \text{if } k = j - 1, \\ 0 & \text{otherwise,} \end{cases}$$

and hence $[e_{i,j}^\sigma, e_{l,i}^\sigma] = e_{l,j}^\sigma$ for $j \neq l$ and $[e_{i,j}^\sigma, e_{j,k}^\sigma] = e_{i,k}$ for $i \neq k$. Thus, by Corollary 4.5, one can deduce that

$$\begin{aligned} [e_{i,j}^\sigma, e_{j,i}^\sigma] &= [e_{i,j}^\sigma, -e_{j,j-1}^\sigma e_{j-1,i}^\sigma + e_{j-1,i}^\sigma e_{j,j-1}^\sigma] \\ &= [e_{i,j-1}^\sigma, e_{j-1,i}^\sigma] + h_{j-1} - h_j \\ &= [e_{i,i+1}^\sigma, e_{i+1,i}^\sigma] + h_{i+1} - h_j \\ &= h_i - h_j. \end{aligned}$$

Hence, the second formula of part (1) holds. Similarly, the other cases can be verified and we skip the detail. \square

It is seen that the root vectors depend on the choice of a reduced expression of w_0 . We then discuss the root vectors of $\mathbf{U}(\mathfrak{q}_n)$ for arbitrary reduced expression of w_0 . Assume $\lambda \in \mathbb{N}^N$ and $w_0 = w^\lambda$ is a reduced expression of w_0 different from (4.0.5). It is known that w^λ could be obtained by applying the following relations

$$\begin{aligned} s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1}, \\ s_i s_j &= s_j s_i \quad \text{for } |i - j| > 1. \end{aligned} \tag{4.0.6}$$

on w^σ in finitely many times. Assume w^ς and w^τ are two reduced expressions of w_0 and w^τ could be obtained by applying one of the equations in (4.0.6), we will find the relations between ψ_ς^+ and ψ_τ^+ .

Case 1: Assume that $w^\varsigma = w' s_i s_j w''$, $w^\tau = w' s_j s_i w''$, with $l(w') = h$ and $|i - j| > 1$.

By the action in (3.0.4), we see that $s_j(\alpha_i) = \alpha_i$ and $s_i(\alpha_j) = \alpha_j$. Therefore, the relationships between the roots in the classical case corresponding to the reduced expressions w^ς and w^τ are as follows:

$$\begin{aligned} \beta_t^\varsigma &= \beta_t^\tau, \quad \text{if } t \neq h+1, h+2; \\ \beta_{h+1}^\varsigma &= w'(\alpha_i) = w' s_j(\alpha_i) = \beta_{h+2}^\tau, \\ \beta_{h+2}^\varsigma &= w' s_i(\alpha_j) = w'(\alpha_j) = \beta_{h+1}^\tau. \end{aligned} \quad (4.0.7)$$

On the other hand, (2.0.2) implies

$$e_t^\varsigma = e_t^\tau, \quad \bar{e}_t^\varsigma = \bar{e}_t^\tau, \quad f_t^\varsigma = f_t^\tau, \quad \bar{f}_t^\varsigma = \bar{f}_t^\tau, \quad \text{for all } t \neq h+1, h+2; \quad (4.0.8)$$

while $T_i(x_j) = x_j$ for $x_j \in \{e_j, e_{\bar{j}}, f_j, f_{\bar{j}}\}$ and $|i - j| > 1$ implies

$$\begin{aligned} e_{h+1}^\varsigma &= T_{w'}(e_i) = T_{w'} T_j(e_i) = e_{h+2}^\tau, & e_{h+2}^\varsigma &= T_{w'} T_i(e_j) = T_{w'}(e_j) = e_{h+1}^\tau, \\ \bar{e}_{h+1}^\varsigma &= T_{w'}(e_{\bar{i}}) = T_{w'} T_j(e_{\bar{i}}) = \bar{e}_{h+2}^\tau, & \bar{e}_{h+2}^\varsigma &= T_{w'} T_i(e_{\bar{j}}) = T_{w'}(e_{\bar{j}}) = \bar{e}_{h+1}^\tau, \\ f_{h+1}^\varsigma &= T_{w'}(f_i) = T_{w'} T_j(f_i) = f_{h+2}^\tau, & f_{h+2}^\varsigma &= T_{w'} T_i(f_j) = T_{w'}(f_j) = f_{h+1}^\tau, \\ \bar{f}_{h+1}^\varsigma &= T_{w'}(f_{\bar{i}}) = T_{w'} T_j(f_{\bar{i}}) = \bar{f}_{h+2}^\tau, & \bar{f}_{h+2}^\varsigma &= T_{w'} T_i(f_{\bar{j}}) = T_{w'}(f_{\bar{j}}) = \bar{f}_{h+1}^\tau. \end{aligned} \quad (4.0.9)$$

Summarizing (4.0.7), (4.0.8), (4.0.9), it follows

$$e_{i,j}^\varsigma = e_{i,j}^\tau, \quad \bar{e}_{i,j}^\varsigma = \bar{e}_{i,j}^\tau, \quad (4.0.10)$$

for all $1 \leq i \neq j \leq n$, hence $\psi_\varsigma^- = \psi_\tau^-$ and $\psi_\varsigma^+ = \psi_\tau^+$.

Case 2: Assume that $w^\varsigma = w' s_i s_j s_i w''$, $w^\tau = w' s_j s_i s_j w''$, with $l(w') = h$ and $|i - j| = 1$. By the action in (3.0.4), we see that $s_i(\alpha_j) = s_j(\alpha_i)$, $s_i s_j(\alpha_i) = \alpha_j$, and $s_j s_i(\alpha_j) = \alpha_i$. Hence, the relationships between the roots in the classical case corresponding to the reduced expressions w^ς and w^τ are as follows:

$$\begin{aligned} \beta_t^\varsigma &= \beta_t^\tau, \quad \text{for all } t \neq h+1, h+2, h+3, \\ \beta_{h+1}^\varsigma &= w'(\alpha_i) = w' s_j s_i(\alpha_j) = \beta_{h+3}^\tau, \\ \beta_{h+2}^\varsigma &= w' s_i(\alpha_j) = w' s_j(\alpha_i) = \beta_{h+2}^\tau, \\ \beta_{h+3}^\varsigma &= w' s_i s_j(\alpha_i) = w'(\alpha_j) = \beta_{h+1}^\tau. \end{aligned} \quad (4.0.11)$$

On the other hand, (2.0.2) implies

$$e_t^\varsigma = e_t^\tau, \quad \bar{e}_t^\varsigma = \bar{e}_t^\tau, \quad f_t^\varsigma = f_t^\tau, \quad \bar{f}_t^\varsigma = \bar{f}_t^\tau, \quad \text{if } t \neq h+1, h+2, h+3, \quad (4.0.12)$$

and Remark 3.4 implies

$$\begin{aligned}
e_{h+1}^s &= T_{w'}(e_i) = T_{w'}T_jT_i(e_j) = e_{h+3}^\tau, \\
\bar{e}_{h+1}^s &= T_{w'}(e_{\bar{i}}) = T_{w'}T_jT_i(e_{\bar{j}}) = \bar{e}_{h+3}^\tau, \\
f_{h+1}^s &= T_{w'}(f_i) = T_{w'}T_jT_i(f_j) = f_{h+3}^\tau, \\
\bar{f}_{h+1}^s &= T_{w'}(f_{\bar{i}}) = T_{w'}T_jT_i(f_{\bar{j}}) = \bar{f}_{h+3}^\tau, \\
e_{h+3}^s &= T_{w'}T_iT_j(e_i) = T_{w'}(e_j) = e_{h+1}^\tau, \\
\bar{e}_{h+3}^s &= T_{w'}T_iT_j(e_{\bar{i}}) = T_{w'}(e_{\bar{j}}) = \bar{e}_{h+1}^\tau, \\
f_{h+3}^s &= T_{w'}T_iT_j(f_i) = T_{w'}(f_j) = f_{h+1}^\tau, \\
\bar{f}_{h+3}^s &= T_{w'}T_iT_j(f_{\bar{i}}) = T_{w'}(f_{\bar{j}}) = \bar{f}_{h+1}^\tau.
\end{aligned} \tag{4.0.13}$$

In the meanwhile, as $T_i(x_j) = -T_j(x_i)$ for $x_j \in \{e_j, e_{\bar{j}}, f_j, f_{\bar{j}}\}$ and $|i - j| = 1$, we have

$$\begin{aligned}
e_{h+2}^s &= T_{w'}T_i(e_j) = -T_{w'}T_j(e_i) = -e_{h+2}^\tau, \\
\bar{e}_{h+2}^s &= T_{w'}T_i(e_{\bar{j}}) = -T_{w'}T_j(e_{\bar{i}}) = -\bar{e}_{h+2}^\tau, \\
f_{h+2}^s &= T_{w'}T_i(f_j) = -T_{w'}T_j(f_i) = -f_{h+2}^\tau, \\
\bar{f}_{h+2}^s &= T_{w'}T_i(f_{\bar{j}}) = -T_{w'}T_j(f_{\bar{i}}) = -\bar{f}_{h+2}^\tau.
\end{aligned} \tag{4.0.14}$$

Summarizing equations (4.0.11), (4.0.12), (4.0.13), (4.0.14), it follows

$$\begin{aligned}
e_{i,j}^s &= -e_{i,j}^\tau, \quad \bar{e}_{i,j}^s = -\bar{e}_{i,j}^\tau, \quad e_{j,i}^s = -e_{j,i}^\tau, \quad \bar{e}_{j,i}^s = -\bar{e}_{j,i}^\tau, \quad \text{if } \alpha_{i,j} = \beta_{h+2}^s = \beta_{h+2}^\tau; \\
e_{i,j}^s &= e_{i,j}^\tau, \quad \bar{e}_{i,j}^s = \bar{e}_{i,j}^\tau, \quad e_{j,i}^s = e_{j,i}^\tau, \quad \bar{e}_{j,i}^s = \bar{e}_{j,i}^\tau, \quad \text{otherwise.}
\end{aligned} \tag{4.0.15}$$

Hence we have proved the following theorem.

Theorem 4.7. *Assume w^γ is a reduced expression of w_0 , $1 \leq i < j \leq n$, then we have*

$$\begin{aligned}
e_{i,j}^\gamma &= r_{i,j}^\gamma e_{i,j}^\sigma, & \bar{e}_{i,j}^\gamma &= r_{i,j}^\gamma \bar{e}_{i,j}^\sigma, \\
e_{j,i}^\gamma &= r_{j,i}^\gamma e_{j,i}^\sigma, & \bar{e}_{j,i}^\gamma &= r_{j,i}^\gamma \bar{e}_{j,i}^\sigma,
\end{aligned}$$

where w^σ is the fixed reduced expression in (4.0.5), and $r_{i,j}^\gamma, r_{j,i}^\gamma \in \{\pm 1\}$.

5. THE PBW BASIS OF THE KOSTANT \mathbb{Z} -FORM $\mathbf{U}(\mathfrak{q}_n)_\mathbb{Z}$

Referring to [11, (3.1)], recall the elements $\mathbf{x}_{i,j}$ and $\bar{\mathbf{x}}_{i,j}$ in $\mathbf{U}(\mathfrak{q}_n)$ for $1 \leq i \neq j \leq n$, and it is trivial that

$$\mathbf{x}_{i,j} = -e_{i,j}^\sigma, \quad \bar{\mathbf{x}}_{i,j} = -\bar{e}_{i,j}^\sigma.$$

For any X , $m \in \mathbb{N}$, and $\mathbf{k} \in \mathbb{N}^n$, set

$$X^{(m)} = \frac{X^m}{m!}, \quad \binom{X}{m} = \frac{X(X-1)\cdots(X-m+1)}{m!}, \quad \binom{\mathbf{h}}{\mathbf{k}} = \prod_{i=1}^n \binom{h_i}{k_i}.$$

By [4, Section 4], the Kostant \mathbb{Z} -form of $\mathbf{U}(\mathfrak{q}_n)$, denoted by $\mathbf{U}(\mathfrak{q}_n)_\mathbb{Z}$, is the \mathbb{Z} -subalgebra generated by

$$\mathcal{G}_\mathbb{Z} = \left\{ \binom{\mathbf{h}}{\mathbf{k}}, h_{\bar{i}}, e_j^{(t)}, e_{\bar{j}}, f_j^{(t)}, f_{\bar{j}} \mid i \in [1, n], j \in [1, n-1], t \in \mathbb{N}, \mathbf{k} \in \mathbb{N}^n \right\}.$$

Let $\mathbf{U}_\mathbb{Z}^0$ be the \mathbb{Z} -subalgebra of $\mathbf{U}(\mathfrak{q}_n)_\mathbb{Z}$ generated by $\binom{\mathbf{h}}{\mathbf{k}}$ and $h_{\bar{i}}$ for $1 \leq i \leq n$, and let $\mathbf{U}_\mathbb{Z}^+$ (resp. $\mathbf{U}_\mathbb{Z}^-$) be the \mathbb{Z} -subalgebra of $\mathbf{U}(\mathfrak{q}_n)_\mathbb{Z}$ generated by e_j and $e_{\bar{j}}$ (resp. $f_j, f_{\bar{j}}$) for $1 \leq j \leq n-1$. Referring to [11, Proposition 3.3], there is a linear space isomorphism

$$\mathbf{U}(\mathfrak{q}_n)_\mathbb{Z} \cong \mathbf{U}_\mathbb{Z}^- \otimes \mathbf{U}_\mathbb{Z}^0 \otimes \mathbf{U}_\mathbb{Z}^+.$$

With the same discussion with [11, Proposition 3.3] and [15, Proposition 6.14], and applying Theorem 4.7, we have

Theorem 5.1. *Assume w^γ is any reduced expression of w_0 , then we have*

- (1) the set $\{\mathbf{e}_{A^+}^\gamma = \prod_{n \geq i \geq 1} \prod_{n \geq j \geq i+1} (\mathbf{e}_{i,j}^\gamma)^{(a_{i,j}^0)} (\bar{\mathbf{e}}_{i,j}^\gamma)^{a_{i,j}^1} \mid A \in M_n(\mathbb{N}|\mathbb{Z}_2)\}$ forms a \mathbb{Z} -basis of $\mathbf{U}_{\mathbb{Z}}^+$;
- (2) the set $\{\mathbf{e}_{A^-}^\gamma = \prod_{n \geq j \geq 2} \prod_{1 \leq i \leq j-1} (\mathbf{e}_{j,i}^\gamma)^{(a_{j,i}^0)} (\bar{\mathbf{e}}_{j,i}^\gamma)^{a_{j,i}^1} \mid A \in M_n(\mathbb{N}|\mathbb{Z}_2)\}$ forms a \mathbb{Z} -basis of $\mathbf{U}_{\mathbb{Z}}^-$;
- (3) the set $\{\mathbf{h}_{A,j} = \prod_{i=1}^n \binom{h_i}{j_i} h_i^{a_{i,i}^1} \mid \mathbf{j} = (j_1, \dots, j_n) \in \mathbb{N}^n, A \in M_n(\mathbb{N}|\mathbb{Z}_2)\}$ forms a \mathbb{Z} -basis of $\mathbf{U}_{\mathbb{Z}}^0$;
- (4) the set $\{\mathbf{b}_A^\gamma = f_A^\gamma \cdot \mathbf{h}_{A,j} \cdot e_A^\gamma \mid \mathbf{j} = (j_1, \dots, j_n) \in \mathbb{N}^n, A \in M_n(\mathbb{N}|\mathbb{Z}_2)\}$ forms a \mathbb{Z} -basis of $\mathbf{U}(\mathfrak{q}_n)_{\mathbb{Z}}$.

6. THE ACTION OF \mathcal{B} ON THE QUANTUM QUEER SUPERALGEBRA $\mathbf{U}_v(\mathfrak{q}_n)$

In [21, Section 4], Olshanski introduced a quantum deformation of the universal enveloping algebra of \mathfrak{q}_n using a modification of the Reshetikhin-Takhtajan-Faddeev method, denoted as $\mathbf{U}_v(\mathfrak{q}_n)$. Du and Wan gives an equivalent definition of $\mathbf{U}_v(\mathfrak{q}_n)$ in [11, Proposition 5.2].

Definition 6.1. [11, Proposition 5.2] The quantum queer superalgebra $\mathbf{U}_v(\mathfrak{q}_n)$ is the (Hopf) superalgebra over $\mathbb{Q}(v)$ generated by even generators $\mathbf{K}_i, \mathbf{K}_i^{-1}, \mathbf{E}_j, \mathbf{F}_j$, and odd generators $\mathbf{K}_{\bar{i}}, \mathbf{E}_{\bar{j}}, \mathbf{F}_{\bar{j}}$, with $1 \leq i \leq n, 1 \leq j \leq n-1$, subjecting to the following relations:

$$(QQ1) \quad \mathbf{K}_i \mathbf{K}_i^{-1} = \mathbf{K}_i^{-1} \mathbf{K}_i = 1, \quad \mathbf{K}_i \mathbf{K}_j = \mathbf{K}_j \mathbf{K}_i, \quad \mathbf{K}_i \mathbf{K}_{\bar{j}} = \mathbf{K}_{\bar{j}} \mathbf{K}_i,$$

$$\mathbf{K}_{\bar{i}} \mathbf{K}_{\bar{j}} + \mathbf{K}_{\bar{j}} \mathbf{K}_{\bar{i}} = 2\delta_{i,j} \frac{\mathbf{K}_i^2 - \mathbf{K}_i^{-2}}{v^2 - v^{-2}};$$

$$(QQ2) \quad \mathbf{K}_i \mathbf{E}_j = v^{(\epsilon_i, \alpha_j)} \mathbf{E}_j \mathbf{K}_i, \quad \mathbf{K}_i \mathbf{E}_{\bar{j}} = v^{(\epsilon_i, \alpha_j)} \mathbf{E}_{\bar{j}} \mathbf{K}_i,$$

$$\mathbf{K}_i \mathbf{F}_j = v^{-(\epsilon_i, \alpha_j)} \mathbf{F}_j \mathbf{K}_i, \quad \mathbf{K}_i \mathbf{F}_{\bar{j}} = v^{-(\epsilon_i, \alpha_j)} \mathbf{F}_{\bar{j}} \mathbf{K}_i;$$

$$(QQ3) \quad \mathbf{K}_{\bar{i}} \mathbf{E}_i - v \mathbf{E}_i \mathbf{K}_{\bar{i}} = \mathbf{E}_{\bar{i}} \mathbf{K}_i^{-1}, \quad v \mathbf{K}_{\bar{i}} \mathbf{E}_{i-1} - \mathbf{E}_{i-1} \mathbf{K}_{\bar{i}} = -\mathbf{K}_i^{-1} \mathbf{E}_{\bar{i}-1},$$

$$\mathbf{K}_{\bar{i}} \mathbf{F}_i - v \mathbf{F}_i \mathbf{K}_{\bar{i}} = -\mathbf{F}_{\bar{i}} \mathbf{K}_i, \quad v \mathbf{K}_{\bar{i}} \mathbf{F}_{i-1} - \mathbf{F}_{i-1} \mathbf{K}_{\bar{i}} = \mathbf{K}_i \mathbf{F}_{\bar{i}-1},$$

$$\mathbf{K}_{\bar{i}} \mathbf{E}_{\bar{i}} + v \mathbf{E}_{\bar{i}} \mathbf{K}_{\bar{i}} = \mathbf{E}_i \mathbf{K}_i^{-1}, \quad v \mathbf{K}_{\bar{i}} \mathbf{E}_{\bar{i}-1} + \mathbf{E}_{\bar{i}-1} \mathbf{K}_{\bar{i}} = \mathbf{K}_i^{-1} \mathbf{E}_{i-1},$$

$$\mathbf{K}_{\bar{i}} \mathbf{F}_{\bar{i}} + v \mathbf{F}_{\bar{i}} \mathbf{K}_{\bar{i}} = \mathbf{F}_i \mathbf{K}_i, \quad v \mathbf{K}_{\bar{i}} \mathbf{F}_{\bar{i}-1} + \mathbf{F}_{\bar{i}-1} \mathbf{K}_{\bar{i}} = \mathbf{K}_i \mathbf{F}_{i-1},$$

$$\mathbf{K}_{\bar{i}} \mathbf{E}_j - \mathbf{E}_j \mathbf{K}_{\bar{i}} = \mathbf{K}_{\bar{i}} \mathbf{F}_j - \mathbf{F}_j \mathbf{K}_{\bar{i}} = \mathbf{K}_{\bar{i}} \mathbf{E}_{\bar{j}} + \mathbf{E}_{\bar{j}} \mathbf{K}_{\bar{i}} = \mathbf{K}_{\bar{i}} \mathbf{F}_{\bar{j}} + \mathbf{F}_{\bar{j}} \mathbf{K}_{\bar{i}} = 0 \text{ for } j \neq i, i-1;$$

$$(QQ4) \quad \mathbf{E}_i \mathbf{F}_j - \mathbf{F}_j \mathbf{E}_i = \delta_{i,j} \frac{\mathbf{K}_i \mathbf{K}_{i+1}^{-1} - \mathbf{K}_i^{-1} \mathbf{K}_{i+1}}{v - v^{-1}},$$

$$\mathbf{E}_{\bar{i}} \mathbf{F}_{\bar{j}} + \mathbf{F}_{\bar{j}} \mathbf{E}_{\bar{i}} = \delta_{i,j} \left(\frac{\mathbf{K}_i \mathbf{K}_{i+1} - \mathbf{K}_i^{-1} \mathbf{K}_{i+1}^{-1}}{v - v^{-1}} + (v - v^{-1}) \mathbf{K}_{\bar{i}} \mathbf{K}_{\bar{i}+1} \right),$$

$$\mathbf{E}_i \mathbf{F}_{\bar{j}} - \mathbf{F}_{\bar{j}} \mathbf{E}_i = \delta_{i,j} (\mathbf{K}_{i+1}^{-1} \mathbf{K}_{\bar{i}} - \mathbf{K}_{\bar{i}+1} \mathbf{K}_i^{-1}), \quad \mathbf{E}_{\bar{i}} \mathbf{F}_j - \mathbf{F}_j \mathbf{E}_{\bar{i}} = \delta_{i,j} (\mathbf{K}_{i+1} \mathbf{K}_{\bar{i}} - \mathbf{K}_{\bar{i}+1} \mathbf{K}_i);$$

$$(QQ5) \quad \mathbf{E}_{\bar{i}}^2 = -\frac{v - v^{-1}}{v + v^{-1}} \mathbf{E}_i^2, \quad \mathbf{F}_{\bar{i}}^2 = \frac{v - v^{-1}}{v + v^{-1}} \mathbf{F}_i^2,$$

$$\mathbf{E}_i \mathbf{E}_{\bar{j}} - \mathbf{E}_{\bar{j}} \mathbf{E}_i = \mathbf{F}_i \mathbf{F}_{\bar{j}} - \mathbf{F}_{\bar{j}} \mathbf{F}_i = 0 \quad \text{for } |i - j| \neq 1,$$

$$\mathbf{E}_i \mathbf{E}_j - \mathbf{E}_j \mathbf{E}_i = \mathbf{F}_i \mathbf{F}_j - \mathbf{F}_j \mathbf{F}_i = \mathbf{E}_{\bar{i}} \mathbf{E}_{\bar{j}} + \mathbf{E}_{\bar{j}} \mathbf{E}_{\bar{i}} = \mathbf{F}_{\bar{i}} \mathbf{F}_{\bar{j}} + \mathbf{F}_{\bar{j}} \mathbf{F}_{\bar{i}} = 0 \quad \text{for } |i - j| > 1,$$

$$\mathbf{E}_i \mathbf{E}_{i+1} - v \mathbf{E}_{i+1} \mathbf{E}_i = \mathbf{E}_{\bar{i}} \mathbf{E}_{\bar{i}+1} + v \mathbf{E}_{\bar{i}+1} \mathbf{E}_{\bar{i}}, \quad \mathbf{E}_i \mathbf{E}_{\bar{i}+1} - v \mathbf{E}_{\bar{i}+1} \mathbf{E}_i = \mathbf{E}_{\bar{i}} \mathbf{E}_{i+1} - v \mathbf{E}_{i+1} \mathbf{E}_{\bar{i}},$$

$$\mathbf{F}_i \mathbf{F}_{i+1} - v \mathbf{F}_{i+1} \mathbf{F}_i = -(\mathbf{F}_{\bar{i}} \mathbf{F}_{\bar{i}+1} + v \mathbf{F}_{\bar{i}+1} \mathbf{F}_{\bar{i}}), \quad \mathbf{F}_i \mathbf{F}_{\bar{i}+1} - v \mathbf{F}_{\bar{i}+1} \mathbf{F}_i = \mathbf{F}_{\bar{i}} \mathbf{F}_{i+1} - v \mathbf{F}_{i+1} \mathbf{F}_{\bar{i}};$$

$$\begin{aligned}
(\text{QQ6}) \quad & E_i^2 E_j - (v + v^{-1}) E_i E_j E_i + E_j E_i^2 = 0, & F_i^2 F_j - (v + v^{-1}) F_i F_j F_i + F_j F_i^2 = 0, \\
& E_i^2 E_{\bar{j}} - (v + v^{-1}) E_i E_{\bar{j}} E_i + E_{\bar{j}} E_i^2 = 0, & F_i^2 F_{\bar{j}} - (v + v^{-1}) F_i F_{\bar{j}} F_i + F_{\bar{j}} F_i^2 = 0, \\
& \text{where } |i - j| = 1.
\end{aligned}$$

Remark 6.2. By observing the relations in (QQ3) and applying the third relation in (QQ4), one provides

$$\begin{aligned}
E_{\bar{j}} &= -v K_{j+1} K_{\bar{j}+1} E_j + K_{j+1} E_j K_{\bar{j}+1}, \\
F_{\bar{j}} &= v K_{j+1}^{-1} K_{\bar{j}+1} F_j - K_{j+1}^{-1} F_j K_{\bar{j}+1}, \\
K_{\bar{j}} &= E_j F_{\bar{j}} K_{j+1} - F_{\bar{j}} E_j K_{j+1} + K_j^{-1} K_{\bar{j}+1} K_{j+1} \\
&= E_j K_{\bar{j}+1} F_j - v^{-1} E_j F_j K_{\bar{j}+1} - v K_{\bar{j}+1} F_j E_j + F_j K_{\bar{j}+1} E_j + K_j^{-1} K_{\bar{j}+1} K_{j+1}.
\end{aligned}$$

Hence by induction on j in descending order from $(n - 1)$ to 1 , one can obtain all the other odd generators in $\mathbf{U}_v(\mathfrak{q}_n)$. In other words, $\mathbf{U}_v(\mathfrak{q}_n)$ could be generated by $\{E_j, F_j, K_i^{\pm 1}, K_{\bar{n}} \mid 1 \leq j \leq n - 1, 1 \leq i \leq n\}$. Moreover, as in [13] and [10], some of the relations (QQ1)–(QQ6) could be omitted.

Following [6], there is an anti-involution Ω over $\mathbf{U}_v(\mathfrak{q}_n)$ with action given by

$$\begin{aligned}
\Omega(v) &= v^{-1}, & \Omega(E_j) &= F_j, & \Omega(F_j) &= E_j, & \Omega(K_i) &= K_i^{-1}, \\
\Omega(E_{\bar{j}}) &= F_{\bar{j}}, & \Omega(F_{\bar{j}}) &= E_{\bar{j}}, & \Omega(K_{\bar{i}}) &= K_{\bar{i}},
\end{aligned} \tag{6.0.1}$$

where $1 \leq i \leq n$ and $1 \leq j \leq n - 1$.

Let $\mathbf{U}_v^{\bar{0}}$ be the subalgebra of $\mathbf{U}_v(\mathfrak{q}_n)$ generated by the even generators $K_i, E_j, F_j (1 \leq i \leq n, 1 \leq j \leq n - 1)$ associated with some defining relations. To obtain the action of \mathcal{B} on $\mathbf{U}_v(\mathfrak{q}_n)$, we consider the action on the even part as the first step.

Denote $\tilde{K}_j = K_j K_{j+1}^{-1}$. Referring to Definition 2.1 and Definition 6.1, it is verified that there is an algebra monomorphism

$$\begin{aligned}
\pi_n : \mathbf{U}_v(\mathfrak{sl}_n) &\rightarrow \mathbf{U}_v^{\bar{0}}, \\
K_j &\mapsto \tilde{K}_j, & E_j &\mapsto E_j, & F_j &\mapsto F_j, & \text{for } j = 1, \dots, n - 1.
\end{aligned}$$

For any $X \in \pi_n(\mathbf{U}_v(\mathfrak{sl}_n))$, we may define the action of \mathcal{B} on it as

$$T_i(X) = \pi_n(T_i(\pi_n^{-1}(X))).$$

Referring to Proposition 2.3, we have

$$\begin{aligned}
T_i(K_i K_{i+1}^{-1}) &= (K_i K_{i+1}^{-1})^{-1} = K_i^{-1} K_{i+1}, \\
T_i(E_i) &= -F_i K_i K_{i+1}^{-1}, & T_i(F_i) &= -K_i^{-1} K_{i+1} E_i, \\
T_i(K_{i+1} K_{i+2}^{-1}) &= (K_{i+1} K_{i+2}^{-1})(K_i K_{i+1}^{-1}) = K_i K_{i+2}^{-1}, \\
T_i(K_{i-1} K_i^{-1}) &= (K_{i-1} K_i^{-1})(K_i K_{i+1}^{-1}) = K_{i-1} K_{i+1}^{-1}, \\
T_i(E_j) &= -E_i E_j + v^{-1} E_j E_i, & T_i(F_j) &= -F_j F_i + v F_i F_j \quad \text{for } |i - j| = 1, \\
T_i(K_j K_{j+1}^{-1}) &= K_j K_{j+1}^{-1}, & T_i(E_j) &= E_j, & T_i(F_j) &= F_j \quad \text{for } |i - j| > 1.
\end{aligned} \tag{6.0.2}$$

Then the action of \mathcal{B} on $\mathbf{U}_v(\mathfrak{sl}_n)$ will induce the action on $\mathbf{U}_v(\mathfrak{q}_n)$.

Theorem 6.3. *The braid group \mathcal{B} acts by $\mathbb{Q}(v)$ -algebra automorphisms on $\mathcal{U}_v(\mathfrak{q}_n)$. The action of \mathcal{B} on the even generators are defined as*

$$\begin{aligned} T_j(\mathbf{K}_i^{\pm 1}) &= \mathbf{K}_{s_j(i)}^{\pm 1} \text{ for } 1 \leq j \leq n-1, \quad 1 \leq i \leq n; \\ T_i(\mathbf{E}_i) &= -\mathbf{F}_i \mathbf{K}_i \mathbf{K}_{i+1}^{-1}, \quad T_i(\mathbf{F}_i) = -\mathbf{K}_i^{-1} \mathbf{K}_{i+1} \mathbf{E}_i; \\ T_i(\mathbf{E}_j) &= -\mathbf{E}_i \mathbf{E}_j + v^{-1} \mathbf{E}_j \mathbf{E}_i, \quad T_i(\mathbf{F}_j) = -\mathbf{F}_j \mathbf{F}_i + v \mathbf{F}_i \mathbf{F}_j \quad \text{for } |i-j|=1; \\ T_i(\mathbf{E}_j) &= \mathbf{E}_j, \quad T_i(\mathbf{F}_j) = \mathbf{F}_j \quad \text{for } |i-j|>1. \end{aligned}$$

The action of \mathcal{B} on the odd generators are defined as

$$\begin{aligned} T_i(\mathbf{K}_{\bar{i}-1}) &= \mathbf{K}_{\bar{i}-1}, \quad T_i(\mathbf{K}_{\bar{i}}) = \mathbf{K}_{\bar{i}+1}, \quad T_i(\mathbf{K}_{\bar{i}+1}) = (v-v^{-1})\mathbf{K}_{\bar{i}+1}\mathbf{F}_i\mathbf{E}_i - (v-v^{-1})\mathbf{F}_i\mathbf{E}_i\mathbf{K}_{\bar{i}+1} + \mathbf{K}_{\bar{i}}, \\ T_i(\mathbf{E}_{\bar{i}}) &= -\mathbf{K}_{\bar{i}+1}\mathbf{F}_i\mathbf{K}_i + v\mathbf{F}_i\mathbf{K}_{\bar{i}+1}\mathbf{K}_i, \quad T_i(\mathbf{F}_{\bar{i}}) = -\mathbf{K}_i^{-1}\mathbf{E}_i\mathbf{K}_{\bar{i}+1} + v^{-1}\mathbf{K}_{\bar{i}+1}\mathbf{K}_i^{-1}\mathbf{E}_i, \\ T_i(\mathbf{E}_{\bar{j}}) &= -\mathbf{E}_i\mathbf{E}_{\bar{j}} + v^{-1}\mathbf{E}_{\bar{j}}\mathbf{E}_i, \quad T_i(\mathbf{F}_{\bar{j}}) = -\mathbf{F}_{\bar{j}}\mathbf{F}_i + v\mathbf{F}_i\mathbf{F}_{\bar{j}} \quad \text{for } |i-j|=1, \\ T_i(\mathbf{K}_{\bar{j}}) &= \mathbf{K}_{\bar{j}}, \quad T_i(\mathbf{E}_{\bar{j}}) = \mathbf{E}_{\bar{j}}, \quad T_i(\mathbf{F}_{\bar{j}}) = \mathbf{F}_{\bar{j}} \quad \text{for } |i-j|>1. \end{aligned}$$

The inverse of T_i is given by

$$\begin{aligned} T_i^{-1}(\mathbf{K}_{i-1}^{\pm 1}) &= \mathbf{K}_{i-1}^{\pm 1}, \quad T_i^{-1}(\mathbf{K}_i^{\pm 1}) = \mathbf{K}_{i+1}^{\pm 1}, \quad T_i^{-1}(\mathbf{K}_{i+1}^{\pm 1}) = \mathbf{K}_i^{\pm 1}, \\ T_i^{-1}(\mathbf{E}_i) &= -\mathbf{K}_{i+1}\mathbf{K}_i^{-1}\mathbf{F}_i, \quad T_i^{-1}(\mathbf{F}_i) = -\mathbf{E}_i\mathbf{K}_i\mathbf{K}_{i+1}^{-1}, \\ T_i^{-1}(\mathbf{E}_j) &= -\mathbf{E}_j\mathbf{E}_i + v^{-1}\mathbf{E}_i\mathbf{E}_j, \quad T_i^{-1}(\mathbf{F}_j) = -\mathbf{F}_i\mathbf{F}_j + v\mathbf{F}_j\mathbf{F}_i \quad \text{for } |i-j|=1, \\ T_i^{-1}(\mathbf{K}_j^{\pm 1}) &= \mathbf{K}_j^{\pm 1}, \quad T_i^{-1}(\mathbf{E}_j) = \mathbf{E}_j, \quad T_i^{-1}(\mathbf{F}_j) = \mathbf{F}_j \quad \text{for } |i-j|>1; \\ T_i^{-1}(\mathbf{K}_{\bar{i}-1}) &= \mathbf{K}_{\bar{i}-1}, \quad T_i^{-1}(\mathbf{K}_{\bar{i}+1}) = \mathbf{K}_{\bar{i}}, \\ T_i^{-1}(\mathbf{K}_{\bar{i}}) &= (v-v^{-1})\mathbf{E}_i\mathbf{F}_i\mathbf{K}_{\bar{i}} - (v-v^{-1})\mathbf{E}_i\mathbf{F}_i\mathbf{K}_{\bar{i}} + \mathbf{K}_{\bar{i}+1}, \\ T_i^{-1}(\mathbf{E}_{\bar{i}}) &= -\mathbf{K}_{i+1}\mathbf{F}_i\mathbf{K}_{\bar{i}} + v\mathbf{K}_{i+1}\mathbf{K}_{\bar{i}}\mathbf{F}_i, \quad T_i^{-1}(\mathbf{F}_{\bar{i}}) = -\mathbf{K}_{\bar{i}}\mathbf{E}_i\mathbf{K}_{i+1}^{-1} + v^{-1}\mathbf{E}_i\mathbf{K}_{\bar{i}}\mathbf{K}_{i+1}^{-1}, \\ T_i^{-1}(\mathbf{E}_{\bar{j}}) &= -\mathbf{E}_j\mathbf{E}_{\bar{i}} + v^{-1}\mathbf{E}_{\bar{i}}\mathbf{E}_j, \quad T_i^{-1}(\mathbf{F}_{\bar{j}}) = -\mathbf{F}_i\mathbf{F}_{\bar{j}} + v\mathbf{F}_{\bar{j}}\mathbf{F}_i \quad \text{for } |i-j|=1, \\ T_i^{-1}(\mathbf{K}_{\bar{j}}) &= \mathbf{K}_{\bar{j}}, \quad T_i^{-1}(\mathbf{E}_{\bar{j}}) = \mathbf{E}_{\bar{j}}, \quad T_i^{-1}(\mathbf{F}_{\bar{j}}) = \mathbf{F}_{\bar{j}} \quad \text{for } |i-j|>1. \end{aligned}$$

We will prove this theorem in several parts.

Proof of the action on the even generators: It is trivial the map π_n holds under the action of T_j . We only need to verify that the action of T_j on \mathbf{K}_i satisfy the equations in (6.0.2) and the definition of braid group \mathcal{B} , for all $1 \leq j \leq n-1$ and $1 \leq i \leq n$.

By $T_i(\mathbf{K}_i) = \mathbf{K}_{i+1}$, $T_{i+1}(\mathbf{K}_i) = \mathbf{K}_i$, $T_i(\mathbf{K}_{i+1}) = \mathbf{K}_i$ and $T_i(\mathbf{K}_{i+2}) = \mathbf{K}_{i+2}$, we have

$$\begin{aligned} T_i(\mathbf{K}_i\mathbf{K}_{i+1}^{-1}) &= T_i(\mathbf{K}_i)T_i(\mathbf{K}_{i+1}^{-1}) = \mathbf{K}_i^{-1}\mathbf{K}_{i+1}, \\ T_i(\mathbf{K}_{i+1}\mathbf{K}_{i+2}^{-1}) &= T_i(\mathbf{K}_{i+1})T_i(\mathbf{K}_{i+2}^{-1}) = \mathbf{K}_i\mathbf{K}_{i+2}^{-1}, \\ T_i(\mathbf{K}_{i-1}\mathbf{K}_i^{-1}) &= T_i(\mathbf{K}_{i-1})T_i(\mathbf{K}_i^{-1}) = \mathbf{K}_{i-1}\mathbf{K}_{i+1}^{-1}, \\ T_i(\mathbf{K}_j\mathbf{K}_{j+1}^{-1}) &= T_i(\mathbf{K}_j)T_i(\mathbf{K}_{j+1}^{-1}) = \mathbf{K}_j\mathbf{K}_{j+1}^{-1} \quad \text{for } |i-j|>1. \end{aligned}$$

And direct calculation leads

$$\begin{aligned} T_i T_{i+1} T_i(\mathbf{K}_{i-1}) &= T_i T_{i+1}(\mathbf{K}_{i-1}) = T_i(\mathbf{K}_{i-1}) = \mathbf{K}_{i-1}, \\ T_{i+1} T_i T_{i+1}(\mathbf{K}_{i-1}) &= T_{i+1} T_i(\mathbf{K}_{i-1}) = T_{i+1}(\mathbf{K}_{i-1}) = \mathbf{K}_{i-1}; \\ T_i T_{i+1} T_i(\mathbf{K}_i) &= T_i T_{i+1}(\mathbf{K}_{i+1}) = T_i(\mathbf{K}_{i+2}) = \mathbf{K}_{i+2}, \\ T_{i+1} T_i T_{i+1}(\mathbf{K}_i) &= T_{i+1} T_i(\mathbf{K}_i) = T_{i+1}(\mathbf{K}_{i+1}) = \mathbf{K}_{i+2}; \\ T_i T_{i+1} T_i(\mathbf{K}_{i+1}) &= T_i T_{i+1}(\mathbf{K}_i) = T_i(\mathbf{K}_i) = \mathbf{K}_{i+1}, \\ T_{i+1} T_i T_{i+1}(\mathbf{K}_{i+1}) &= T_{i+1} T_i(\mathbf{K}_{i+2}) = T_{i+1}(\mathbf{K}_{i+2}) = \mathbf{K}_{i+1}; \end{aligned}$$

$$\begin{aligned} T_i T_{i+1} T_i(\mathbf{K}_{i+2}) &= T_i T_{i+1}(\mathbf{K}_{i+2}) = T_i(\mathbf{K}_{i+1}) = \mathbf{K}_i, \\ T_{i+1} T_i T_{i+1}(\mathbf{K}_{i+2}) &= T_{i+1} T_i(\mathbf{K}_{i+1}) = T_{i+1}(\mathbf{K}_i) = \mathbf{K}_i. \end{aligned}$$

Thus, the composite action $T_i T_{i+1} T_i$ and $T_{i+1} T_i T_{i+1}$ coincide on the generators \mathbf{K}_j . Similarly, we see that the composite action $T_i T_j$ and $T_j T_i$ are equal, where $|i - j| > 1$. Thus, the proposition is completely proven.

Proof of the action on the odd generators: To prove the action on the odd generators, we need the following two lemmas.

Lemma 6.4. *For any $X \in \{\mathbf{K}_{\bar{i}}, \mathbf{E}_{\bar{j}}, \mathbf{F}_{\bar{j}} \mid 1 \leq i \leq n, 1 \leq j \leq n-1\}$, we have*

$$\begin{aligned} T_j T_{j+1} T_j(X) &= T_{j+1} T_j T_{j+1}(X), \\ T_i T_j(X) &= T_j T_i(X) \quad \text{where } |i - j| > 1. \end{aligned}$$

In other word, $T_j (1 \leq j \leq n-1)$ satisfies the relations in (2.0.1).

Proof. For $X = \mathbf{K}_{\bar{j}}$, we see that

$$\begin{aligned} T_j T_{j+1} T_j(\mathbf{K}_{\bar{j-1}}) &= T_j T_{j+1}(\mathbf{K}_{\bar{j-1}}) = \mathbf{K}_{\bar{j-1}}, \\ T_{j+1} T_j T_{j+1}(\mathbf{K}_{\bar{j-1}}) &= T_{j+1} T_j(\mathbf{K}_{\bar{j-1}}) = T_{j+1}(\mathbf{K}_{\bar{j-1}}) = \mathbf{K}_{\bar{j-1}}; \\ T_j T_{j+1} T_j(\mathbf{K}_{\bar{j}}) &= T_j T_{j+1}(\mathbf{K}_{\bar{j+1}}) = T_j(\mathbf{K}_{\bar{j+2}}) = \mathbf{K}_{\bar{j+2}}, \\ T_{j+1} T_j T_{j+1}(\mathbf{K}_{\bar{j}}) &= T_{j+1} T_j(\mathbf{K}_{\bar{j}}) = T_{j+1}(\mathbf{K}_{\bar{j+1}}) = \mathbf{K}_{\bar{j+2}}; \\ T_j T_{j+1} T_j(\mathbf{K}_{\bar{j+1}}) &= T_j T_{j+1}((v - v^{-1})\mathbf{K}_{\bar{j+1}}\mathbf{F}_j\mathbf{E}_j - (v - v^{-1})\mathbf{F}_j\mathbf{E}_j\mathbf{K}_{\bar{j+1}} + \mathbf{K}_{\bar{j}}) \\ &= (v - v^{-1})\mathbf{K}_{\bar{j+2}}\mathbf{F}_{j+1}\mathbf{E}_{j+1} - (v - v^{-1})\mathbf{F}_{j+1}\mathbf{E}_{j+1}\mathbf{K}_{\bar{j+2}} + \mathbf{K}_{\bar{j+1}}, \\ T_{j+1} T_j T_{j+1}(\mathbf{K}_{\bar{j+1}}) &= T_{j+1} T_j(\mathbf{K}_{\bar{j+2}}) = (v - v^{-1})\mathbf{K}_{\bar{j+2}}\mathbf{F}_{j+1}\mathbf{E}_{j+1} - (v - v^{-1})\mathbf{F}_{j+1}\mathbf{E}_{j+1}\mathbf{K}_{\bar{j+2}} + \mathbf{K}_{\bar{j+1}}; \\ T_j T_{j+1} T_j(\mathbf{K}_{\bar{j+2}}) &= T_j T_{j+1}(\mathbf{K}_{\bar{j+2}}) \\ &= T_j((v - v^{-1})\mathbf{K}_{\bar{j+2}}\mathbf{F}_{j+1}\mathbf{E}_{j+1} - (v - v^{-1})\mathbf{F}_{j+1}\mathbf{E}_{j+1}\mathbf{K}_{\bar{j+2}} + \mathbf{K}_{\bar{j+1}}) \\ &= (v - v^{-1})T_j(\mathbf{K}_{\bar{j+2}})T_j(\mathbf{E}_{j+1}) - (v - v^{-1})T_j(\mathbf{F}_{j+1})T_j(\mathbf{E}_{j+1})T_j(\mathbf{K}_{\bar{j+2}}) + T_j(\mathbf{K}_{\bar{j+1}}) \\ &= (v - v^{-1})\mathbf{K}_{\bar{j+2}}(-\mathbf{F}_{j+1}\mathbf{F}_j + v\mathbf{F}_j\mathbf{F}_{j+1})(-\mathbf{E}_j\mathbf{E}_{j+1} + v^{-1}\mathbf{E}_{j+1}\mathbf{E}_j) \\ &\quad - (v - v^{-1})(-\mathbf{F}_{j+1}\mathbf{F}_j + v\mathbf{F}_j\mathbf{F}_{j+1})(-\mathbf{E}_j\mathbf{E}_{j+1} + v^{-1}\mathbf{E}_{j+1}\mathbf{E}_j)\mathbf{K}_{\bar{j+2}} \\ &\quad + (v - v^{-1})\mathbf{K}_{\bar{j+1}}\mathbf{F}_j\mathbf{E}_j - (v - v^{-1})\mathbf{F}_j\mathbf{E}_j\mathbf{K}_{\bar{j+1}} + \mathbf{K}_{\bar{j}} \\ &= (v - v^{-1})\mathbf{K}_{\bar{j+2}}\mathbf{F}_{j+1}\mathbf{F}_j\mathbf{E}_j\mathbf{E}_{j+1} - v^{-1}(v - v^{-1})\mathbf{K}_{\bar{j+2}}\mathbf{F}_{j+1}\mathbf{F}_j\mathbf{E}_{j+1}\mathbf{E}_j \\ &\quad - v(v - v^{-1})\mathbf{K}_{\bar{j+2}}\mathbf{F}_j\mathbf{F}_{j+1}\mathbf{E}_j\mathbf{E}_{j+1} + (v - v^{-1})\mathbf{K}_{\bar{j+2}}\mathbf{F}_j\mathbf{F}_{j+1}\mathbf{E}_{j+1}\mathbf{E}_j \\ &\quad - (v - v^{-1})\mathbf{F}_{j+1}\mathbf{F}_j\mathbf{E}_j\mathbf{E}_{j+1}\mathbf{K}_{\bar{j+2}} + v^{-1}(v - v^{-1})\mathbf{F}_{j+1}\mathbf{F}_j\mathbf{E}_{j+1}\mathbf{E}_j\mathbf{K}_{\bar{j+2}} \\ &\quad + v(v - v^{-1})\mathbf{F}_j\mathbf{F}_{j+1}\mathbf{E}_j\mathbf{E}_{j+1}\mathbf{K}_{\bar{j+2}} - (v - v^{-1})\mathbf{F}_j\mathbf{F}_{j+1}\mathbf{E}_{j+1}\mathbf{E}_j\mathbf{K}_{\bar{j+2}} \\ &\quad + (v - v^{-1})\mathbf{K}_{\bar{j+1}}\mathbf{F}_j\mathbf{E}_j - (v - v^{-1})\mathbf{F}_j\mathbf{E}_j\mathbf{K}_{\bar{j+1}} + \mathbf{K}_{\bar{j}}, \\ T_{j+1} T_j T_{j+1}(\mathbf{K}_{\bar{j+2}}) &= T_{j+1} T_j((v - v^{-1})\mathbf{K}_{\bar{j+2}}\mathbf{F}_{j+1}\mathbf{E}_{j+1} - (v - v^{-1})\mathbf{F}_{j+1}\mathbf{E}_{j+1}\mathbf{K}_{\bar{j+2}} + \mathbf{K}_{\bar{j+1}}) \\ &= (v - v^{-1})T_{j+1}(\mathbf{K}_{\bar{j+2}})T_{j+1}T_{j+1}^{-1}(\mathbf{F}_j)T_{j+1}T_{j+1}^{-1}(\mathbf{E}_j) \\ &\quad - (v - v^{-1})T_{j+1}T_{j+1}^{-1}(\mathbf{F}_j)T_{j+1}T_{j+1}^{-1}(\mathbf{E}_j)T_{j+1}(\mathbf{K}_{\bar{j+2}}) \\ &\quad + T_{j+1}((v - v^{-1})\mathbf{K}_{\bar{j+1}}\mathbf{F}_j\mathbf{E}_j - (v - v^{-1})\mathbf{F}_j\mathbf{E}_j\mathbf{K}_{\bar{j+1}} + \mathbf{K}_{\bar{j}}) \\ &= (v - v^{-1})((v - v^{-1})\mathbf{K}_{\bar{j+2}}\mathbf{F}_{j+1}\mathbf{E}_{j+1} - (v - v^{-1})\mathbf{F}_{j+1}\mathbf{E}_{j+1}\mathbf{K}_{\bar{j+2}} + \mathbf{K}_{\bar{j+1}})\mathbf{F}_j\mathbf{E}_j \\ &\quad - (v - v^{-1})\mathbf{F}_j\mathbf{E}_j((v - v^{-1})\mathbf{K}_{\bar{j+2}}\mathbf{F}_{j+1}\mathbf{E}_{j+1} - (v - v^{-1})\mathbf{F}_{j+1}\mathbf{E}_{j+1}\mathbf{K}_{\bar{j+2}} + \mathbf{K}_{\bar{j+1}}) \\ &\quad + (v - v^{-1})\mathbf{K}_{\bar{j+2}}(-\mathbf{F}_j\mathbf{F}_{j+1} + v\mathbf{F}_{j+1}\mathbf{F}_j)(-\mathbf{E}_{j+1}\mathbf{E}_j + v^{-1}\mathbf{E}_j\mathbf{E}_{j+1}) \end{aligned}$$

$$\begin{aligned}
& - (v - v^{-1})(-F_j F_{j+1} + v F_{j+1} F_j)(-E_{j+1} E_j + v^{-1} E_j E_{j+1}) K_{\overline{j+2}} + K_{\overline{j}} \\
= & (v - v^{-1}) K_{\overline{j+2}} F_{j+1} F_j E_j E_{j+1} - v^{-1} (v - v^{-1}) K_{\overline{j+2}} F_{j+1} F_j E_{j+1} E_j \\
& - v (v - v^{-1}) K_{\overline{j+2}} F_j F_{j+1} E_j E_{j+1} + (v - v^{-1}) K_{\overline{j+2}} F_j F_{j+1} E_{j+1} E_j \\
& - (v - v^{-1}) F_{j+1} F_j E_j E_{j+1} K_{\overline{j+2}} + v^{-1} (v - v^{-1}) F_{j+1} F_j E_{j+1} E_j K_{\overline{j+2}} \\
& + v (v - v^{-1}) F_j F_{j+1} E_j E_{j+1} K_{\overline{j+2}} - (v - v^{-1}) F_j F_{j+1} E_{j+1} E_j K_{\overline{j+2}} \\
& + (v - v^{-1}) K_{\overline{j+1}} F_j E_j - (v - v^{-1}) F_j E_j K_{\overline{j+1}} + K_{\overline{j}}.
\end{aligned}$$

Thus, the compositions $T_j T_{j+1} T_j$ and $T_{j+1} T_j T_{j+1}$ coincide on the generators $K_{\overline{j}}$. Similarly, we also see that

$$T_j T_j (K_{\overline{j}}) = T_j T_j (K_{\overline{j}}) \quad \text{where } |i - j| > 1.$$

For $X = E_{\overline{j}}$ with admissible j , we have

$$\begin{aligned}
T_j T_{j+1} T_j (E_{\overline{j-1}}) &= T_j T_{j+1} (-E_j E_{\overline{j-1}} + v^{-1} E_{\overline{j-1}} E_j) \\
&= -E_{j+1} T_j (E_{\overline{j-1}}) + v^{-1} T_j (E_{\overline{j-1}}) E_{j+1} \\
&= -E_{j+1} (-E_j E_{\overline{j-1}} + v^{-1} E_{\overline{j-1}} E_j) + v^{-1} (-E_j E_{\overline{j-1}} + v^{-1} E_{\overline{j-1}} E_j) E_{j+1} \\
&= E_{j+1} E_j E_{\overline{j-1}} - v^{-1} E_{j+1} E_{\overline{j-1}} E_j - v^{-1} E_j E_{\overline{j-1}} E_{j+1} + v^{-2} E_{\overline{j-1}} E_j E_{j+1}, \\
T_{j+1} T_j T_{j+1} (E_{\overline{j-1}}) &= T_{j+1} T_j (E_{\overline{j-1}}) \\
&= T_{j+1} (-E_j E_{\overline{j-1}} + v^{-1} E_{j-1} E_j) \\
&= -(-E_{j+1} E_j + v^{-1} E_j E_{j+1}) E_{\overline{j-1}} + v^{-1} E_{\overline{j-1}} (-E_{j+1} E_j + v^{-1} E_j E_{j+1}) \\
&= E_{j+1} E_j E_{\overline{j-1}} - v^{-1} E_j E_{j+1} E_{\overline{j-1}} - v^{-1} E_{\overline{j-1}} E_{j+1} E_j + v^{-2} E_{\overline{j-1}} E_j E_{j+1}; \\
T_j T_{j+1} T_j (E_{\overline{j}}) &= T_j T_{j+1} (-K_{\overline{j+1}} F_j K_j + v F_j K_{\overline{j+1}} K_j) = -K_{\overline{j+2}} F_{j+1} K_{j+1} + v F_{j+1} K_{\overline{j+2}} K_{j+1}, \\
T_{j+1} T_j T_{j+1} (E_{\overline{j}}) &= T_{j+1} (E_{\overline{j+1}}) = -K_{\overline{j+2}} F_{j+1} K_{j+1} + v F_{j+1} K_{\overline{j+2}} K_{j+1}; \\
T_j T_{j+1} T_j (E_{\overline{j+1}}) &= T_j T_{j+1} (-E_j E_{\overline{j+1}} + v^{-1} E_{\overline{j+1}} E_j) \\
&= -E_{j+1} T_j T_{j+1} (E_{\overline{j+1}}) + v^{-1} T_j T_{j+1} (E_{\overline{j+1}}) E_{j+1} \\
&= -(v - v^{-1}) K_{\overline{j+2}} F_{j+1} E_{j+1} F_j K_j + (v - v^{-1}) F_{j+1} E_{j+1} K_{\overline{j+2}} F_j K_j \\
&\quad + v (v - v^{-1}) F_j K_{\overline{j+2}} F_{j+1} E_{j+1} K_j - v (v - v^{-1}) F_j F_{j+1} E_{j+1} K_{\overline{j+2}} K_j \\
&\quad - K_{\overline{j+1}} F_j K_j + v F_j K_{\overline{j+1}} K_j, \\
T_{j+1} T_j T_{j+1} (E_{\overline{j+1}}) &= T_{j+1} T_j (-K_{\overline{j+2}} F_{j+1} K_{j+1} + v F_{j+1} K_{\overline{j+2}} K_{j+1}) \\
&= -T_{j+1} (K_{\overline{j+2}}) F_j K_j + v F_j T_{j+1} (K_{\overline{j+2}}) K_j \\
&= -[(v - v^{-1}) K_{\overline{j+2}} F_{j+1} E_{j+1} - (v - v^{-1}) F_{j+1} E_{j+1} K_{\overline{j+2}} + K_{\overline{j+1}}] F_j K_j \\
&\quad + v F_j [(v - v^{-1}) K_{\overline{j+2}} F_{j+1} E_{j+1} - (v - v^{-1}) F_{j+1} E_{j+1} K_{\overline{j+2}} + K_{\overline{j+1}}] K_j \\
&= -(v - v^{-1}) K_{\overline{j+2}} F_{j+1} E_{j+1} F_j K_j + (v - v^{-1}) F_{j+1} E_{j+1} K_{\overline{j+2}} F_j K_j \\
&\quad + v (v - v^{-1}) F_j K_{\overline{j+2}} F_{j+1} E_{j+1} K_j - v (v - v^{-1}) F_j F_{j+1} E_{j+1} K_{\overline{j+2}} K_j \\
&\quad - K_{\overline{j+1}} F_j K_j + v F_j K_{\overline{j+1}} K_j; \\
T_j T_{j+1} T_j (E_{\overline{j+2}}) &= T_j T_{j+1} (E_{\overline{j+2}}) \\
&= T_j (-E_{j+1} E_{\overline{j+2}} + v^{-1} E_{\overline{j+2}} E_{j+1}) \\
&= -(-E_j E_{j+1} + v^{-1} E_{j+1} E_j) E_{\overline{j+2}} + v^{-1} E_{\overline{j+2}} (-E_j E_{j+1} + v^{-1} E_{j+1} E_j) \\
&= E_j E_{j+1} E_{\overline{j+2}} - v^{-1} E_{j+1} E_j E_{\overline{j+2}} - v^{-1} E_{\overline{j+2}} E_j E_{j+1} + v^{-2} E_{\overline{j+2}} E_{j+1} E_j, \\
T_{j+1} T_j T_{j+1} (E_{\overline{j+2}}) &= T_{j+1} T_j (-E_{j+1} E_{j+2} + v^{-1} E_{j+2} E_{j+1})
\end{aligned}$$

$$\begin{aligned}
&= -E_j T_{j+1}(E_{j+2}) + v^{-1} T_{j+1}(E_{j+2}) E_j \\
&= -E_j (-E_{j+1} E_{j+2} + v^{-1} E_{j+2} E_{j+1}) + v^{-1} (-E_{j+1} E_{j+2} + v^{-1} E_{j+2} E_{j+1}) E_j \\
&= E_j E_{j+1} E_{j+2} - v^{-1} E_j E_{j+2} E_{j+1} - v^{-1} E_{j+1} E_{j+2} E_j + v^{-2} E_{j+2} E_{j+1} E_j.
\end{aligned}$$

Thus, the compositions $T_j T_{j+1} T_j$ and $T_{j+1} T_j T_{j+1}$ coincide on the generators $E_{\bar{j}}$. Similarly, we see that

$$T_i T_j(E_{\bar{j}}) = T_j T_i(E_{\bar{j}}) \text{ for } |i - j| > 1.$$

From the previous formulas and applying Ω , we deduce that they also coincide on the generators F_j . \square

Lemma 6.5. *For any $Y \in \mathcal{B}$, Y holds the equations in (QQ1)-(QQ6).*

Proof. According to Proposition 2.3, for any $Y \in \mathcal{B}$, the elements in $\{Y(E_j), Y(F_j), Y(K_i^{\pm 1}) \mid 1 \leq j \leq n-1, 1 \leq i \leq n\}$ satisfy all the generating relations in (QQ1)-(QQ6) composed solely of even generators. Therefore, we need to verify that the elements in $\{Y(E_j), Y(F_j), Y(K_i^{\pm 1}), Y(E_{\bar{j}}), Y(F_{\bar{j}}), Y(K_{\bar{i}}) \mid 1 \leq j \leq n-1, 1 \leq i \leq n\}$ satisfy the remaining relations in (QQ1)-(QQ6).

Due to the group structure of \mathcal{B} and the group action on $\mathbf{U}_v(\mathfrak{q}_n)$, we only need to verify that when $Y = T_i (t \in \{i-1, i, i+1, j-1, j, j+1\})$, the elements in the above set satisfy the remaining generating relations in (QQ1)-(QQ6).

For the relations in (QQ1), when $Y = T_j$, we see that

$$Y(K_i)Y(K_{\bar{j}}) = K_{s_j(i)} K_{\bar{j+1}} = K_{\bar{j+1}} K_{s_j(i)} = Y(K_{\bar{j}})Y(K_i).$$

As for the case $Y = T_i$, we have

$$\begin{aligned}
T_i(K_{\bar{i}})T_i(K_{\bar{i}}) + T_i(K_{\bar{i}})T_i(K_{\bar{i}}) &= 2K_{\bar{i+1}}K_{\bar{i+1}} \\
&= 2\frac{K_{\bar{i+1}}^2 - K_{\bar{i+1}}^{-2}}{v^2 - v^{-2}} \\
&= T_i\left(2\frac{K_i^2 - K_i^{-2}}{v^2 - v^{-2}}\right) \\
&= 2\frac{T_i(K_i)^2 - T_i(K_i)^{-2}}{v^2 - v^{-2}}.
\end{aligned}$$

Similarly, the other cases with equations in (QQ1) can be proved by direct calculation.

For the relations in (QQ2), $Y = T_i, T_{i+1}$ leads

$$\begin{aligned}
T_i(K_i)T_i(E_{\bar{i}}) &= K_{i+1}(-K_{\bar{i+1}}F_i K_i + vF_i K_{\bar{i+1}} K_i) = vT_i(E_{\bar{i}})T_i(K_i), \\
T_{i+1}(K_i)T_{i+1}(F_{\bar{i}}) &= K_i(-F_{\bar{i}}F_{i+1} + vF_{i+1}F_{\bar{i}}) = v^{-1}T_{i+1}(F_{\bar{i}})T_{i+1}(K_i).
\end{aligned}$$

In the same way, the other cases with equations in (QQ2) can be checked.

For the relations in (QQ3), we only give the proof of the second one and the fifth one. The others are analogous to them. If $Y = T_i$, then

$$\begin{aligned}
vT_i(\mathbf{K}_{\bar{i}})T_i(\mathbf{E}_{i-1}) - T_i(\mathbf{E}_{i-1})T_i(\mathbf{K}_{\bar{i}}) &= v\mathbf{K}_{\bar{i+1}}(-\mathbf{E}_i\mathbf{E}_{i-1} + v^{-1}\mathbf{E}_{i-1}\mathbf{E}_i) - (-\mathbf{E}_i\mathbf{E}_{i-1} + v^{-1}\mathbf{E}_{i-1}\mathbf{E}_i)\mathbf{K}_{\bar{i+1}} \\
&= -v\mathbf{K}_{\bar{i+1}}\mathbf{E}_i\mathbf{E}_{i-1} + \mathbf{K}_{\bar{i+1}}\mathbf{E}_{i-1}\mathbf{E}_i + \mathbf{E}_i\mathbf{E}_{i-1}\mathbf{K}_{\bar{i+1}} - v^{-1}\mathbf{E}_{i-1}\mathbf{E}_i\mathbf{K}_{\bar{i+1}} \\
&= (-v\mathbf{K}_{\bar{i+1}}\mathbf{E}_i + \mathbf{E}_i\mathbf{K}_{\bar{i+1}})\mathbf{E}_{i-1} + \mathbf{E}_{i-1}(\mathbf{K}_{\bar{i+1}}\mathbf{E}_i - v^{-1}\mathbf{E}_i\mathbf{K}_{\bar{i+1}}) \\
&= -\mathbf{K}_{\bar{i+1}}^{-1}(-\mathbf{E}_{\bar{i}}\mathbf{E}_{i-1} + v^{-1}\mathbf{E}_{i-1}\mathbf{E}_{\bar{i}}) \\
&= -\mathbf{K}_{\bar{i+1}}^{-1}(-\mathbf{E}_i\mathbf{E}_{\bar{i-1}} + v^{-1}\mathbf{E}_{\bar{i-1}}\mathbf{E}_i) \\
&= -T_i(\mathbf{K}_i^{-1})T_i(\mathbf{E}_{\bar{i-1}}),
\end{aligned}$$

$$\begin{aligned}
T_i(\mathbf{K}_{\bar{i}})T_i(\mathbf{E}_{\bar{i}}) + vT_i(\mathbf{E}_{\bar{i}})T_i(\mathbf{K}_{\bar{i}}) &= \mathbf{K}_{\bar{i+1}}(-\mathbf{K}_{\bar{i+1}}\mathbf{F}_i\mathbf{K}_i + v\mathbf{F}_i\mathbf{K}_{\bar{i+1}}\mathbf{K}_i) + v(-\mathbf{K}_{\bar{i+1}}\mathbf{F}_i\mathbf{K}_i + v\mathbf{F}_i\mathbf{K}_{\bar{i+1}}\mathbf{K}_i)\mathbf{K}_{\bar{i+1}} \\
&= -\mathbf{K}_{\bar{i+1}}^2\mathbf{F}_i\mathbf{K}_i + v^2\mathbf{F}_i\mathbf{K}_{\bar{i+1}}^2\mathbf{K}_i \\
&= (-\mathbf{F}_i\mathbf{K}_i\mathbf{K}_{\bar{i+1}}^{-1})\mathbf{K}_{\bar{i+1}}^{-1} \\
&= T_i(\mathbf{E}_i)T_i(\mathbf{K}_i^{-1}).
\end{aligned}$$

For the second relation in (QQ4), we only give the proof for the case $i = j$. And the other cases with relations in (QQ4) are similar to prove and we omit it. For the case $Y = T_i$, we see that

$$\begin{aligned}
T_i(\mathbf{E}_{\bar{i}}\mathbf{F}_{\bar{i}} + \mathbf{F}_{\bar{i}}\mathbf{E}_{\bar{i}}) &= T_i(\mathbf{E}_{\bar{i}})T_i(\mathbf{F}_{\bar{i}}) + T_i(\mathbf{F}_{\bar{i}})T_i(\mathbf{E}_{\bar{i}}) \\
&= (-\mathbf{K}_{\bar{i+1}}\mathbf{F}_i\mathbf{K}_i + v\mathbf{F}_i\mathbf{K}_{\bar{i+1}}\mathbf{K}_i)(-\mathbf{K}_i^{-1}\mathbf{E}_i\mathbf{K}_{\bar{i+1}} + v^{-1}\mathbf{K}_{\bar{i+1}}\mathbf{K}_i^{-1}\mathbf{E}_i) \\
&\quad + (-\mathbf{K}_i^{-1}\mathbf{E}_i\mathbf{K}_{\bar{i+1}} + v^{-1}\mathbf{K}_{\bar{i+1}}\mathbf{K}_i^{-1}\mathbf{E}_i)(-\mathbf{K}_{\bar{i+1}}\mathbf{F}_i\mathbf{K}_i + v\mathbf{F}_i\mathbf{K}_{\bar{i+1}}\mathbf{K}_i) \\
&= \mathbf{K}_{\bar{i+1}}\mathbf{F}_i\mathbf{E}_i\mathbf{K}_{\bar{i+1}} - v^{-1}\mathbf{K}_{\bar{i+1}}\mathbf{F}_i\mathbf{K}_{\bar{i+1}}\mathbf{E}_i - v\mathbf{F}_i\mathbf{K}_{\bar{i+1}}\mathbf{E}_i\mathbf{K}_{\bar{i+1}} + \mathbf{F}_i\mathbf{K}_{\bar{i+1}}^2\mathbf{E}_i \\
&\quad + \mathbf{E}_i\mathbf{K}_{\bar{i+1}}^2\mathbf{F}_i - v\mathbf{E}_i\mathbf{K}_{\bar{i+1}}\mathbf{F}_i\mathbf{K}_{\bar{i+1}} - v^{-1}\mathbf{K}_{\bar{i+1}}\mathbf{E}_i\mathbf{K}_{\bar{i+1}}\mathbf{F}_i + \mathbf{K}_{\bar{i+1}}\mathbf{E}_i\mathbf{F}_i\mathbf{K}_{\bar{i+1}} \\
&= (v - v^{-1})^2\mathbf{K}_{\bar{i+1}}^2\mathbf{F}_i\mathbf{E}_i - (v - v^{-1})^2\mathbf{K}_{\bar{i+1}}\mathbf{F}_i\mathbf{E}_i\mathbf{K}_{\bar{i+1}} + \mathbf{E}_{\bar{i}}\mathbf{F}_{\bar{i}} + \mathbf{F}_{\bar{i}}\mathbf{E}_{\bar{i}} \\
&= \frac{\mathbf{K}_i\mathbf{K}_{\bar{i+1}} - \mathbf{K}_i^{-1}\mathbf{K}_{\bar{i+1}}^{-1}}{v - v^{-1}} + (v - v^{-1})^2\mathbf{K}_{\bar{i+1}}^2\mathbf{F}_i\mathbf{E}_i \\
&\quad - (v - v^{-1})^2\mathbf{K}_{\bar{i+1}}\mathbf{F}_i\mathbf{E}_i\mathbf{K}_{\bar{i+1}} + (v - v^{-1})\mathbf{K}_{\bar{i+1}}\mathbf{K}_{\bar{i}} \\
&= T_i\left(\frac{\mathbf{K}_i\mathbf{K}_{\bar{i+1}} - \mathbf{K}_i^{-1}\mathbf{K}_{\bar{i+1}}^{-1}}{v - v^{-1}} + (v - v^{-1})\mathbf{K}_{\bar{i}}\mathbf{K}_{\bar{i+1}}\right).
\end{aligned}$$

For (QQ5), we only give the proof of the forth from the bottom and the others are analogous to it. When $Y = T_i$, we have

$$\begin{aligned}
T_i(\mathbf{E}_{\bar{i}})T_i(\mathbf{E}_{\bar{i}}) &= (-\mathbf{K}_{\bar{i+1}}\mathbf{F}_i\mathbf{K}_i + v\mathbf{F}_i\mathbf{K}_{\bar{i+1}}\mathbf{K}_i)(-\mathbf{K}_{\bar{i+1}}\mathbf{F}_i\mathbf{K}_i + v\mathbf{F}_i\mathbf{K}_{\bar{i+1}}\mathbf{K}_i) \\
&= (v - v^{-1})^2\mathbf{F}_i\mathbf{K}_{\bar{i+1}}^2\mathbf{F}_i\mathbf{K}_i^2 + (1 - v^2)\mathbf{F}_i\mathbf{K}_{\bar{i+1}}\mathbf{F}_i\mathbf{K}_{\bar{i+1}}\mathbf{K}_i^2 \\
&\quad + (1 - v^2)\mathbf{F}_i\mathbf{K}_{\bar{i+1}}\mathbf{F}_{\bar{i}}\mathbf{K}_{\bar{i+1}}\mathbf{K}_i^2 + v^2\mathbf{F}_{\bar{i}}^2\mathbf{K}_{\bar{i+1}}^2\mathbf{K}_i^2 \\
&= v^2\frac{v - v^{-1}}{v + v^{-1}}\mathbf{F}_i^2\mathbf{K}_{\bar{i+1}}^2\mathbf{K}_i^2 - v^{-2}\frac{v - v^{-1}}{v + v^{-1}}\mathbf{F}_i^2\mathbf{K}_{\bar{i+1}}^{-2}\mathbf{K}_i^2 + v^2\frac{v - v^{-1}}{v + v^{-1}}\mathbf{F}_i^2\mathbf{K}_{\bar{i+1}}^2\mathbf{K}_i^2 \\
&\quad + v^2(1 - v^2)\mathbf{F}_i^2\mathbf{K}_{\bar{i+1}}^2\mathbf{K}_i^2 - v^2(1 - v^2)\frac{v - v^{-1}}{v + v^{-1}}\mathbf{F}_i^2\mathbf{K}_{\bar{i+1}}^2\mathbf{K}_i^2 \\
&= -v^{-2}\frac{v - v^{-1}}{v + v^{-1}}\mathbf{F}_i^2\mathbf{K}_i^2\mathbf{K}_{\bar{i+1}}^{-2} \\
&= -\frac{v - v^{-1}}{v + v^{-1}}T_i(\mathbf{E}_i)^2,
\end{aligned}$$

$$\begin{aligned}
T_i(\mathbf{E}_i)T_i(\mathbf{E}_{i+1}) - T_i(\mathbf{E}_{i+1})T_i(\mathbf{E}_i) &= (-F_i K_i K_{i+1}^{-1})(-E_i E_{i+1} + v^{-1} E_{i+1} E_i) \\
&\quad - v(-E_i E_{i+1} + v^{-1} E_{i+1} E_i)(-F_i K_i K_{i+1}^{-1}) \\
&= v F_i E_i E_{i+1} K_i K_{i+1}^{-1} - E_{i+1} F_i E_i K_i K_{i+1}^{-1} \\
&\quad - v E_i F_i E_{i+1} K_i K_{i+1}^{-1} + E_{i+1} E_i F_i K_i K_{i+1}^{-1} \\
&= -v \frac{K_i K_{i+1}^{-1} - K_i^{-1} K_{i+1}}{v - v^{-1}} E_{i+1} K_i K_{i+1}^{-1} \\
&\quad + E_{i+1} \frac{K_i K_{i+1}^{-1} - K_i^{-1} K_{i+1}}{v - v^{-1}} K_i K_{i+1}^{-1} \\
&= -v \frac{K_i K_{i+1}^{-1} - K_i^{-1} K_{i+1}}{v - v^{-1}} (K_{i+1}^{-1} E_{i+1} + v E_{i+1} K_{i+1}) K_i \\
&\quad + (K_{i+1}^{-1} E_{i+1} + v E_{i+1} K_{i+1}) \frac{K_i K_{i+1}^{-1} - K_i^{-1} K_{i+1}}{v - v^{-1}} K_i \\
&= (-K_{i+1}^{-1} F_i K_i + v F_i K_{i+1}^{-1} K_i)(-E_i E_{i+1} + v^{-1} E_{i+1} E_i) \\
&\quad + v(-E_i E_{i+1} + v^{-1} E_{i+1} E_i)(-K_{i+1}^{-1} F_i K_i + v F_i K_{i+1}^{-1} K_i) \\
&= T_i(\mathbf{E}_i)T_i(\mathbf{E}_{i+1}) + v T_i(\mathbf{E}_{i+1})T_i(\mathbf{E}_i).
\end{aligned}$$

For the relations in (QQ6), we only detailedly verify the third relation. Then we have

$$\begin{aligned}
&T_i(\mathbf{E}_i)^2 T_i(\mathbf{E}_{i+1}) - (v + v^{-1}) T_i(\mathbf{E}_i) T_i(\mathbf{E}_{i+1}) T_i(\mathbf{E}_i) + T_i(\mathbf{E}_{i+1}) T_i(\mathbf{E}_i)^2 \\
&= (-F_i K_i K_{i+1}^{-1})(-F_i K_i K_{i+1}^{-1})(-E_i E_{i+1} + v^{-1} E_{i+1} E_i) \\
&\quad - (v + v^{-1})(-F_i K_i K_{i+1}^{-1})(-E_i E_{i+1} + v^{-1} E_{i+1} E_i)(-F_i K_i K_{i+1}^{-1}) \\
&\quad + (-E_i E_{i+1} + v^{-1} E_{i+1} E_i)(-F_i K_i K_{i+1}^{-1})(-F_i K_i K_{i+1}^{-1}) \\
&= F_i(E_i F_i - F_i E_i) E_{i+1} - v^{-1} E_{i+1} F_i(E_i F_i - F_i E_i) \\
&\quad - v^{-2}(E_i F_i - F_i E_i) F_i E_{i+1} + v^{-3} E_{i+1}(E_i F_i - F_i E_i) F_i \\
&= F_i \frac{K_i K_{i+1}^{-1} - K_i^{-1} K_{i+1}}{v - v^{-1}} E_{i+1} - v^{-1} E_{i+1} F_i \frac{K_i K_{i+1}^{-1} - K_i^{-1} K_{i+1}}{v - v^{-1}} \\
&\quad - v^{-2} \frac{K_i K_{i+1}^{-1} - K_i^{-1} K_{i+1}}{v - v^{-1}} F_i E_{i+1} + v^{-3} E_{i+1} \frac{K_i K_{i+1}^{-1} - K_i^{-1} K_{i+1}}{v - v^{-1}} F_i \\
&= 0.
\end{aligned}$$

In summary, for each $1 \leq i \leq n-1$, T_i maintain the relation (QQ1)-(QQ6), hence T_i is a superalgebra endomorphism. \square

Remark 6.6. A direct computation shows that

$$T_i T_j(\mathbf{E}_i) = \mathbf{E}_j, \quad T_i T_j(\mathbf{F}_i) = \mathbf{F}_j \quad \text{if } |i - j| = 1,$$

$$T_i T_{i+1}(\mathbf{E}_{\bar{i}}) = \mathbf{E}_{\bar{i+1}}, \quad T_i T_{i+1}(\mathbf{F}_{\bar{i}}) = \mathbf{F}_{\bar{i+1}},$$

$$\begin{aligned}
T_{i+1} T_i(\mathbf{E}_{\bar{i+1}}) &= -E_{i+1} E_i K_{i+2}^{-1} F_{i+1} K_{i+1} + v^{-1} E_i E_{i+1} K_{i+2}^{-1} F_{i+1} K_{i+1} + v E_{i+1} E_i F_{i+1} K_{i+2}^{-1} K_{i+1} \\
&\quad - E_i E_{i+1} F_{i+1} K_{i+2}^{-1} K_{i+1} + v^{-1} K_{i+2}^{-1} F_{i+1} K_{i+1} E_{i+1} E_i - v^{-2} K_{i+2}^{-1} F_{i+1} K_{i+1} E_i E_{i+1} \\
&\quad - F_{i+1} K_{i+2}^{-1} K_{i+1} E_{i+1} E_i + v^{-1} F_{i+1} K_{i+2}^{-1} K_{i+1} E_i E_{i+1}, \\
T_{i+1} T_i(\mathbf{F}_{\bar{i+1}}) &= -K_{i+1}^{-1} E_{i+1} K_{i+2}^{-1} F_i F_{i+1} + v^{-1} K_{i+2}^{-1} K_{i+1}^{-1} E_{i+1} F_i F_{i+1} + v K_{i+1}^{-1} E_{i+1} K_{i+2}^{-1} F_{i+1} F_i \\
&\quad - K_{i+2}^{-1} K_{i+1}^{-1} E_{i+1} F_{i+1} F_i + v F_i F_{i+1} K_{i+1}^{-1} E_{i+1} K_{i+2}^{-1} - v^2 F_{i+1} F_i K_{i+1}^{-1} E_{i+1} K_{i+2}^{-1} \\
&\quad - F_i F_{i+1} K_{i+2}^{-1} K_{i+1}^{-1} E_{i+1} + v F_{i+1} F_i K_{i+2}^{-1} K_{i+1}^{-1} E_{i+1}.
\end{aligned}$$

Proof of the inverse of T_i : Obviously, Remark 6.6 shows that $T_i \in \mathcal{B}$ ($i = 1, \dots, n-1$) are automorphisms. Also, we have

$$\begin{aligned} T_i^{-1}(\mathbf{E}_j) &= T_j(\mathbf{E}_i) = -\mathbf{E}_j\mathbf{E}_i + v^{-1}\mathbf{E}_i\mathbf{E}_j, \\ T_i^{-1}(\mathbf{F}_j) &= T_j(\mathbf{F}_i) = -\mathbf{F}_i\mathbf{F}_j + v\mathbf{F}_j\mathbf{F}_i \quad \text{for } |i-j|=1; \\ T_i^{-1}(\mathbf{E}_{\overline{i+1}}) &= T_{i+1}(\mathbf{E}_{\overline{i}}) = -\mathbf{E}_{i+1}\mathbf{E}_{\overline{i}} + v^{-1}\mathbf{E}_{\overline{i}}\mathbf{E}_{i+1}, \\ T_i^{-1}(\mathbf{F}_{\overline{i+1}}) &= T_{i+1}(\mathbf{F}_{\overline{i}}) = -\mathbf{F}_{\overline{i}}\mathbf{F}_{i+1} + v\mathbf{F}_{i+1}\mathbf{F}_{\overline{i}}. \end{aligned}$$

And by the formulas in Remark 6.2, we see that

$$\begin{aligned} T_i^{-1}(\mathbf{K}_{\overline{i}}) &= T_i^{-1}(T_i(\mathbf{K}_{\overline{i+1}}) - (v - v^{-1})\mathbf{K}_{\overline{i+1}}\mathbf{F}_i\mathbf{E}_i + (v - v^{-1})\mathbf{F}_i\mathbf{E}_i\mathbf{K}_{\overline{i+1}}) \\ &= \mathbf{K}_{\overline{i+1}} - (v - v^{-1})\mathbf{K}_{\overline{i}}(-\mathbf{E}_i\mathbf{K}_i\mathbf{K}_{i+1}^{-1})(-\mathbf{K}_i^{-1}\mathbf{K}_{i+1}\mathbf{F}_i) \\ &\quad + (v - v^{-1})(-\mathbf{E}_i\mathbf{K}_i\mathbf{K}_{i+1}^{-1})(-\mathbf{K}_i^{-1}\mathbf{K}_{i+1}\mathbf{F}_i)\mathbf{K}_{\overline{i}} \\ &= (v - v^{-1})\mathbf{E}_i\mathbf{F}_i\mathbf{K}_{\overline{i}} - (v - v^{-1})\mathbf{K}_{\overline{i}}\mathbf{E}_i\mathbf{F}_i + \mathbf{K}_{\overline{i+1}}, \\ T_i^{-1}(\mathbf{E}_{\overline{i}}) &= T_i^{-1}(\mathbf{K}_{i+1}\mathbf{E}_i\mathbf{K}_{\overline{i+1}} - v\mathbf{K}_{i+1}\mathbf{K}_{\overline{i+1}}\mathbf{E}_i) \\ &= \mathbf{K}_i(-\mathbf{K}_{i+1}\mathbf{K}_i^{-1}\mathbf{F}_i)\mathbf{K}_{\overline{i}} - v\mathbf{K}_i\mathbf{K}_{\overline{i}}(-\mathbf{K}_{i+1}\mathbf{K}_i^{-1}\mathbf{F}_i) \\ &= -\mathbf{K}_{i+1}\mathbf{F}_i\mathbf{K}_{\overline{i}} + v\mathbf{K}_{i+1}\mathbf{K}_{\overline{i}}\mathbf{F}_i, \\ T_{i+1}^{-1}(\mathbf{E}_{\overline{i}}) &= T_{i+1}^{-1}(\mathbf{K}_{\overline{i}}\mathbf{E}_i\mathbf{K}_i - v\mathbf{E}_i\mathbf{K}_{\overline{i}}\mathbf{K}_i) \\ &= \mathbf{K}_{\overline{i}}(-\mathbf{E}_i\mathbf{E}_{i+1} + v^{-1}\mathbf{E}_{i+1}\mathbf{E}_i)\mathbf{K}_i \\ &\quad - v(-\mathbf{E}_i\mathbf{E}_{i+1} + v^{-1}\mathbf{E}_{i+1}\mathbf{E}_i)\mathbf{K}_{\overline{i}}\mathbf{K}_i \\ &= -(\mathbf{K}_{\overline{i}}\mathbf{E}_i - v\mathbf{E}_i\mathbf{K}_{\overline{i}})\mathbf{E}_{i+1}\mathbf{K}_i + v^{-1}\mathbf{E}_{i+1}(\mathbf{K}_{\overline{i}}\mathbf{E}_i - v\mathbf{E}_i\mathbf{K}_{\overline{i}})\mathbf{K}_i \\ &= -(\mathbf{E}_{\overline{i}}\mathbf{K}_i^{-1})\mathbf{E}_{i+1}\mathbf{K}_i + v^{-1}\mathbf{E}_{i+1}(\mathbf{E}_{\overline{i}}\mathbf{K}_i^{-1})\mathbf{K}_i \\ &= -\mathbf{E}_{\overline{i}}\mathbf{E}_{i+1} + v^{-1}\mathbf{E}_{i+1}\mathbf{E}_{\overline{i}}. \end{aligned}$$

Therefore together with the anti-involution Ω given in (6.0.1), Theorem 6.3 can be proved completely.

7. THE ROOT VECTORS OF $\mathbf{U}_v(\mathfrak{q}_n)$

Similar to [5, Proposition 9.1.2], to construct a PBW-type basis of quantum queer superalgebra $\mathbf{U}_v(\mathfrak{q}_n)$, we need to define analogues of root vectors associated with the roots of \mathfrak{q}_n .

Recall the root lattice $\mathbf{Q} = \bigoplus_{j=1}^{n-1} \mathbb{Z}\alpha_j$ and $\alpha_j = \epsilon_j - \epsilon_{j+1}$. Assume $\gamma = \sum_{i=1}^n k_i\epsilon_i \in \mathbf{Q}$, we set $\mathbf{K}_\gamma = \prod_{i=1}^n \mathbf{K}_i^{k_i}$. Then we have the following in $\mathbf{U}_v(\mathfrak{q}_n)$.

Proposition 7.1. *For any $\gamma = \sum_{i=1}^n k_i\epsilon_i \in \mathbf{Q}$, and $w \in \mathfrak{S}_n$, we have $T_w(\mathbf{K}_\gamma) = \mathbf{K}_{w(\gamma)}$.*

Proof. Assume $w = s_{j_1}s_{j_2}\cdots s_{j_r}$ be a reduced expression of w . By Theorem 6.3, for any $h \in [1, n]$, we have $T_j(\mathbf{K}_h) = \mathbf{K}_{s_j(h)}$, and $T_w(\mathbf{K}_h) = T_{j_1}T_{j_2}\cdots T_{j_r}(\mathbf{K}_h) = \mathbf{K}_{w(h)}$. As a consequence,

$$T_w(\mathbf{K}_\gamma) = T_w\left(\prod_{i=1}^n \mathbf{K}_i^{k_i}\right) = \prod_{i=1}^n T_w(\mathbf{K}_i^{k_i}) = \prod_{i=1}^n (T_w(\mathbf{K}_i))^{k_i} = \prod_{i=1}^n \mathbf{K}_{w(i)}^{k_i}.$$

On the other hand, (3.0.5) implies

$$w(\gamma) = w\left(\sum_{i=1}^n k_i \epsilon_i\right) = \sum_{i=1}^n k_i w(\epsilon_i) = \sum_{i=1}^n k_i \epsilon_{w(i)},$$

which means

$$\mathbb{K}_{w(\gamma)} = \prod_{i=1}^n \mathbb{K}_{w(i)}^{k_i} = T_w(\mathbb{K}_\gamma).$$

□

Similar to [5, Definition 8.1.4], we give the following definition.

Definition 7.2. For any expression of $w_0 = w^\gamma = s_{i_1} s_{i_2} \cdots s_{i_N}$ and assume $1 \leq t \leq N$, we define some special elements in $\mathcal{U}_v(\mathfrak{q}_n)$ as

$$\begin{aligned} \mathbb{E}_t^\gamma &= T_{i_1} T_{i_2} \cdots T_{i_{t-1}}(\mathbb{E}_{i_t}), & \mathbb{F}_t^\gamma &= T_{i_1} T_{i_2} \cdots T_{i_{t-1}}(\mathbb{F}_{i_t}), \\ \bar{\mathbb{E}}_t^\gamma &= T_{i_1} T_{i_2} \cdots T_{i_{t-1}}(\bar{\mathbb{E}}_{i_t}), & \bar{\mathbb{F}}_t^\gamma &= T_{i_1} T_{i_2} \cdots T_{i_{t-1}}(\bar{\mathbb{F}}_{i_t}). \end{aligned} \quad (7.0.1)$$

The elements \mathbb{E}_t^γ 's (resp. \mathbb{F}_t^γ 's) are called even positive (resp. negative) root vectors, while the elements $\bar{\mathbb{E}}_t^\gamma$'s (resp. $\bar{\mathbb{F}}_t^\gamma$'s) are called odd positive (resp. negative) root vectors,

We denote the set of positive and negative root vectors by

$$\Psi_\gamma^+ = \{\mathbb{E}_t^\gamma, \bar{\mathbb{E}}_t^\gamma \mid 1 \leq t \leq N\}, \quad \Psi_\gamma^- = \{\mathbb{F}_t^\gamma, \bar{\mathbb{F}}_t^\gamma \mid 1 \leq t \leq N\}.$$

For any $\beta_{\gamma,t} = \alpha_{i,j} \in \Phi^+ \cup \Phi^-$ ($1 \leq i < j \leq n$), we can denote

$$\begin{aligned} \mathbb{E}_{i,j}^\gamma &= \mathbb{E}_t^\gamma, & \bar{\mathbb{E}}_{i,j}^\gamma &= \bar{\mathbb{E}}_t^\gamma, \\ \mathbb{E}_{j,i}^\gamma &= \mathbb{F}_t^\gamma, & \bar{\mathbb{E}}_{j,i}^\gamma &= \bar{\mathbb{F}}_t^\gamma. \end{aligned}$$

Example 7.3. Fix $n = 3$ and we will find the root vectors for $\mathcal{U}_v(\mathfrak{q}_3)$ using two different expressions of the longest element w_0 of \mathfrak{S}_3 .

(1) If we choose $\gamma = (1,2,1)$ and $w_0 = w^\gamma = s_1 s_2 s_1$, the positive roots are

$$\beta_{\gamma,1} = \alpha_1 = \alpha_{1,2}; \quad \beta_{\gamma,2} = s_1(\alpha_2) = \alpha_{1,3}; \quad \beta_{\gamma,3} = s_1 s_2(\alpha_1) = \alpha_{2,3},$$

while the corresponding positive root vectors are

$$\begin{aligned} \mathbb{E}_{1,2}^\gamma &= \mathbb{E}_1, & \mathbb{E}_{1,3}^\gamma &= T_1(\mathbb{E}_2) = -\mathbb{E}_1 \mathbb{E}_2 + v^{-1} \mathbb{E}_2 \mathbb{E}_1, & \mathbb{E}_{2,3}^\gamma &= T_1 T_2(\mathbb{E}_1) = \mathbb{E}_2, \\ \bar{\mathbb{E}}_{1,2}^\gamma &= \bar{\mathbb{E}}_{\bar{1}}, & \bar{\mathbb{E}}_{1,3}^\gamma &= T_1(\bar{\mathbb{E}}_{\bar{2}}) = -\mathbb{E}_1 \bar{\mathbb{E}}_{\bar{2}} + v^{-1} \bar{\mathbb{E}}_{\bar{2}} \mathbb{E}_1, & \bar{\mathbb{E}}_{2,3}^\gamma &= T_1 T_2(\bar{\mathbb{E}}_{\bar{1}}) = \bar{\mathbb{E}}_{\bar{2}}. \end{aligned}$$

(2) If we choose $\gamma = (2,1,2)$ and $w_0 = w^\gamma = s_2 s_1 s_2$, then

$$\beta_{\gamma,1} = \alpha_2 = \alpha_{2,3}; \quad \beta_{\gamma,2} = s_2(\alpha_1) = \alpha_{1,3}; \quad \beta_{\gamma,3} = s_2 s_1(\alpha_2) = \alpha_{1,2},$$

and the corresponding positive root vectors are

$$\begin{aligned} \mathbb{E}_{2,3}^\gamma &= \mathbb{E}_2, & \mathbb{E}_{1,3}^\gamma &= T_2(\mathbb{E}_1) = -\mathbb{E}_2 \mathbb{E}_1 + v^{-1} \mathbb{E}_1 \mathbb{E}_2, & \mathbb{E}_{1,2}^\gamma &= T_2 T_1(\mathbb{E}_2) = \mathbb{E}_1, \\ \bar{\mathbb{E}}_{2,3}^\gamma &= \bar{\mathbb{E}}_{\bar{2}}, & \bar{\mathbb{E}}_{1,3}^\gamma &= T_2(\bar{\mathbb{E}}_{\bar{1}}) = -\mathbb{E}_2 \bar{\mathbb{E}}_{\bar{1}} + v^{-1} \bar{\mathbb{E}}_{\bar{1}} \mathbb{E}_2, \\ \bar{\mathbb{E}}_{1,2}^\gamma &= T_2 T_1(\bar{\mathbb{E}}_{\bar{2}}) = -\mathbb{E}_2 \mathbb{E}_1 \mathbb{K}_{\bar{3}} \bar{\mathbb{F}}_2 \mathbb{K}_2 + v^{-1} \mathbb{E}_1 \mathbb{E}_2 \mathbb{K}_{\bar{3}} \bar{\mathbb{F}}_2 \mathbb{K}_2 + v \mathbb{E}_2 \mathbb{E}_1 \bar{\mathbb{F}}_2 \mathbb{K}_{\bar{3}} \mathbb{K}_2 - \mathbb{E}_1 \mathbb{E}_2 \bar{\mathbb{F}}_2 \mathbb{K}_{\bar{3}} \mathbb{K}_2 \\ &\quad + v^{-1} \mathbb{K}_{\bar{3}} \bar{\mathbb{F}}_2 \mathbb{K}_2 \mathbb{E}_2 \mathbb{E}_1 - v^{-2} \mathbb{K}_{\bar{3}} \bar{\mathbb{F}}_2 \mathbb{K}_2 \mathbb{E}_1 \mathbb{E}_2 - \bar{\mathbb{F}}_2 \mathbb{K}_{\bar{3}} \mathbb{K}_2 \mathbb{E}_2 \mathbb{E}_1 + v^{-1} \bar{\mathbb{F}}_2 \mathbb{K}_{\bar{3}} \mathbb{K}_2 \mathbb{E}_1 \mathbb{E}_2. \end{aligned}$$

With the fixed expression w^σ , similar calculation to Corollary 4.5, then we have the following.

Corollary 7.4. *Assume that $1 \leq i < n$, $i + 1 < j < n$. Then we have*

$$\begin{aligned} E_{i,i+1}^\sigma &= E_i, & \bar{E}_{i,i+1}^\sigma &= E_{\bar{i}}, & E_{i+1,i}^\sigma &= F_i, & \bar{E}_{i+1,i}^\sigma &= F_{\bar{i}}, \\ E_{i,j}^\sigma &= -E_{i,j-1}^\sigma E_{j-1,j}^\sigma + v^{-1} E_{j-1,j}^\sigma E_{i,j-1}^\sigma = [\cdots [[E_i, -E_{i+1}]_{v^{-1}}, -E_{i+2}]_{v^{-1}}, \cdots, -E_{j-1}]_{v^{-1}}, \\ \bar{E}_{i,j}^\sigma &= -E_{i,j-1}^\sigma \bar{E}_{j-1,j}^\sigma + v^{-1} \bar{E}_{j-1,j}^\sigma E_{i,j-1}^\sigma = [\cdots [[E_i, -E_{i+1}]_{v^{-1}}, -E_{i+2}]_{v^{-1}}, \cdots, -E_{j-1}]_{v^{-1}}, \\ E_{j,i}^\sigma &= -E_{j,j-1}^\sigma E_{j-1,i}^\sigma + v E_{j-1,i}^\sigma E_{j,j-1}^\sigma = [-F_{j-1}, [-F_{j-2}, \cdots [-F_{i+1}, F_i]_v]_v \cdots]_v, \\ E_{j,i}^\sigma &= -\bar{E}_{j,j-1}^\sigma E_{j-1,i}^\sigma + v E_{j-1,i}^\sigma \bar{E}_{j,j-1}^\sigma = [-F_{j-1}, [-F_{j-2}, \cdots [-F_{i+1}, F_i]_v]_v \cdots]_v, \end{aligned}$$

where $[a, b]_x = ab - xba$, for any $a, b \in \mathbf{U}_v(\mathfrak{q}_n)$, $x \in \mathbb{Q}(v)$.

Similar to the case for $\mathbf{U}(\mathfrak{q}_n)$ in Section 4, assume w^ς and w^τ are two reduced expressions of w_0 , and w^τ could be obtained by applying one of the equations in (4.0.6), we will find the relations between Ψ_ς^+ and Ψ_τ^+ .

Case 1: Assume that $w^\varsigma = w' s_i s_j w''$, $w^\tau = w' s_j s_i w''$, with $l(w') = h$ and $|i - j| > 1$. In this case, we have

$$\begin{aligned} E_t^\varsigma &= E_t^\tau, & \bar{E}_t^\varsigma &= \bar{E}_t^\tau, & F_t^\varsigma &= F_t^\tau, & \bar{F}_t^\varsigma &= \bar{F}_t^\tau \quad \text{for all } t \neq h+1, h+2; \\ E_{h+1}^\varsigma &= T_{w'}(E_i) = T_{w'} T_j(E_i) = E_{h+2}^\tau, & E_{h+2}^\varsigma &= T_{w'} T_i(E_j) = T_{w'}(E_j) = E_{h+1}^\tau, \\ \bar{E}_{h+1}^\varsigma &= T_{w'}(E_{\bar{i}}) = T_{w'} T_j(E_{\bar{i}}) = \bar{E}_{h+2}^\tau, & \bar{E}_{h+2}^\varsigma &= T_{w'} T_i(E_{\bar{j}}) = T_{w'}(E_{\bar{j}}) = \bar{E}_{h+1}^\tau; \\ F_{h+1}^\varsigma &= T_{w'}(F_i) = T_{w'} T_j(F_i) = F_{h+2}^\tau, & F_{h+2}^\varsigma &= T_{w'} T_i(F_j) = T_{w'}(F_j) = F_{h+1}^\tau, \\ \bar{F}_{h+1}^\varsigma &= T_{w'}(F_{\bar{i}}) = T_{w'} T_j(F_{\bar{i}}) = \bar{F}_{h+2}^\tau, & \bar{F}_{h+2}^\varsigma &= T_{w'} T_i(F_{\bar{j}}) = T_{w'}(F_{\bar{j}}) = \bar{F}_{h+1}^\tau. \end{aligned}$$

Case 2: Assume that $w^\varsigma = w' s_i s_j s_i w''$, $w^\tau = w' s_j s_i s_j w''$, with $l(w') = h$ and $|i - j| = 1$. With the same discussion as Section 4 and referring to Remark 6.6, the relationships between the root vectors are

$$\begin{aligned} E_t^\varsigma &= E_t^\tau, & \bar{E}_t^\varsigma &= \bar{E}_t^\tau, & F_t^\varsigma &= F_t^\tau, & \bar{F}_t^\varsigma &= \bar{F}_t^\tau \quad \text{if } t \neq h+1, h+2, h+3; \\ E_{h+1}^\varsigma &= T_{w'}(E_i) = T_{w'} T_j T_i(E_j) = E_{h+3}^\tau, & \bar{E}_{h+1}^\varsigma &= T_{w'}(E_{\bar{i}}), \\ F_{h+1}^\varsigma &= T_{w'}(F_i) = T_{w'} T_j T_i(F_j) = F_{h+3}^\tau, & \bar{F}_{h+1}^\varsigma &= T_{w'}(F_{\bar{i}}); \\ E_{h+2}^\varsigma &= T_{w'} T_i(E_j) = T_{w'}(-E_i E_j + v^{-1} E_j E_i) = -E_{h+3}^\tau E_{h+1}^\tau + v^{-1} E_{h+1}^\tau E_{h+3}^\tau, \\ \bar{E}_{h+2}^\varsigma &= T_{w'} T_i(E_{\bar{j}}) = T_{w'}(-E_i E_{\bar{j}} + v^{-1} E_{\bar{j}} E_i) = -E_{h+3}^\tau \bar{E}_{h+1}^\tau + v^{-1} \bar{E}_{h+1}^\tau E_{h+3}^\tau, \\ F_{h+2}^\varsigma &= T_{w'} T_i(F_j) = T_{w'}(-F_j F_i + v F_i F_j) = -F_{h+1}^\tau F_{h+3}^\tau + v F_{h+3}^\tau F_{h+1}^\tau, \\ \bar{F}_{h+2}^\varsigma &= T_{w'} T_i(F_{\bar{j}}) = T_{w'}(-F_{\bar{j}} F_i + v F_i F_{\bar{j}}) = -\bar{F}_{h+1}^\tau F_{h+3}^\tau + v F_{h+3}^\tau \bar{F}_{h+1}^\tau; \\ E_{h+3}^\varsigma &= T_{w'} T_i T_j(E_i), & \bar{E}_{h+3}^\varsigma &= T_{w'} T_i T_j(E_{\bar{i}}), \\ F_{h+3}^\varsigma &= T_{w'} T_i T_j(F_i), & \bar{F}_{h+3}^\varsigma &= T_{w'} T_i T_j(F_{\bar{i}}). \end{aligned}$$

if $j = i + 1$, then by Remark 6.6, we have

$$E_{h+3}^\varsigma = E_{h+1}^\tau, \quad \bar{E}_{h+3}^\varsigma = \bar{E}_{h+2}^\tau, \quad F_{h+3}^\varsigma = F_{h+1}^\tau, \quad \bar{F}_{h+3}^\varsigma = \bar{F}_{h+2}^\tau.$$

8. A PBW-TYPE BASIS OF $\mathbf{U}_v(\mathfrak{q}_n)$

In this section, we will construct a PBW-type basis for the quantum queer superalgebra $\mathbf{U}_v(\mathfrak{q}_n)$.

With the fixed reduced expression w^σ of w_0 in (4.0.5), for any $A = (a_{i,j}^{\bar{0}} | a_{i,j}^{\bar{1}}) \in M_n(\mathbb{N}|\mathbb{Z}_2)$, $\mathbf{j} = (j_1, \dots, j_n) \in \mathbb{Z}^n$, denote

$$\begin{aligned} \mathbf{K}_{A,\mathbf{j}} &= \prod_{i=1}^n \mathbf{K}_i^{j_i} \mathbf{K}_i^{a_{i,i}^{\bar{1}}}, \\ \mathbf{E}_A^\sigma &= \prod_{n \geq i \geq 1} \prod_{n \geq j \geq i+1} (\mathbf{E}_{i,j}^\sigma)^{a_{i,j}^{\bar{0}}} (\bar{\mathbf{E}}_{i,j}^\sigma)^{a_{i,j}^{\bar{1}}}, \\ \mathbf{F}_A^\sigma &= \prod_{n \geq j \geq 2} \prod_{1 \leq i \leq j-1} (\mathbf{E}_{j,i}^\sigma)^{a_{j,i}^{\bar{0}}} (\bar{\mathbf{E}}_{j,i}^\sigma)^{a_{j,i}^{\bar{1}}}. \end{aligned}$$

let \mathbf{U}_v^0 be the $\mathbb{Q}(v)$ -subalgebra of $\mathbf{U}_v(\mathfrak{q}_n)$ generated by $\mathbf{K}_i^{\pm 1}$ and \mathbf{K}_i^- for $1 \leq i \leq n$, and let \mathbf{U}_v^+ (respectively. \mathbf{U}_v^-) be the $\mathbb{Q}(v)$ -subalgebra of $\mathbf{U}_v(\mathfrak{q}_n)$ generated by \mathbf{E}_j and $\bar{\mathbf{E}}_j$ (respectively. $\mathbf{F}_j, \bar{\mathbf{F}}_j$) for $1 \leq j \leq n-1$. Referring to [14, Theorem 2.3], there is a linear space isomorphism

$$\mathbf{U}_v(\mathfrak{q}_n) \cong \mathbf{U}_v^- \otimes \mathbf{U}_v^0 \otimes \mathbf{U}_v^+.$$

Recall the root vectors defined in [11](by replacing quantum parameter q to v): for $1 \leq i \leq n-1$, define

$$X_{i,i+1} = \mathbf{E}_i, \quad X_{i+1,i} = \mathbf{F}_i, \quad \bar{X}_{i,i+1} = \bar{\mathbf{E}}_i, \quad \bar{X}_{i+1,i} = \bar{\mathbf{F}}_i;$$

for $|j-i| > 1$, define

$$\begin{aligned} X_{i,j} &:= \begin{cases} X_{i,k} X_{k,j} - v X_{k,j} X_{i,k}, & \text{if } i < j, \\ X_{i,k} X_{k,j} - v^{-1} X_{k,j} X_{i,k}, & \text{if } i > j, \end{cases} \\ \bar{X}_{i,j} &:= \begin{cases} X_{i,k} \bar{X}_{k,j} - v \bar{X}_{k,j} X_{i,k}, & \text{if } i < j, \\ \bar{X}_{i,k} X_{k,j} - v^{-1} X_{k,j} \bar{X}_{i,k}, & \text{if } i > j, \end{cases} \end{aligned} \quad (8.0.1)$$

where k is strictly between i and j . Referring to Corollary 7.4, it is seen that $X_{i,j}, \bar{X}_{i,j}$ differs $\mathbf{E}_{i,j}^\sigma, \bar{\mathbf{E}}_{i,j}^\sigma$ with v and v^{-1} . Hence, similar to [11, Proposition 5.8], we have the following conclusion.

Proposition 8.1. *With the fixed expression w^σ , we have*

- (1) the set $\{\mathbf{E}_A^\sigma = \prod_{n \geq i \geq 1} \prod_{n \geq j \geq i+1} (\mathbf{E}_{i,j}^\sigma)^{a_{i,j}^{\bar{0}}} (\bar{\mathbf{E}}_{i,j}^\sigma)^{a_{i,j}^{\bar{1}}} \mid A \in M_n(\mathbb{N}|\mathbb{Z}_2)\}$ forms a $\mathbb{Q}(v)$ -basis of \mathbf{U}_v^+ ;
- (2) the set $\{\mathbf{F}_A^\sigma = \prod_{n \geq j \geq 2} \prod_{1 \leq i \leq j-1} (\mathbf{E}_{j,i}^\sigma)^{a_{j,i}^{\bar{0}}} (\bar{\mathbf{E}}_{j,i}^\sigma)^{a_{j,i}^{\bar{1}}} \mid A \in M_n(\mathbb{N}|\mathbb{Z}_2)\}$ forms a $\mathbb{Q}(v)$ -basis of \mathbf{U}_v^- ;
- (3) the set $\{\mathbf{K}_{A,\mathbf{j}} = \prod_{i=1}^n \mathbf{K}_i^{j_i} \mathbf{K}_i^{a_{i,i}^{\bar{1}}} \mid \mathbf{j} = (j_1, \dots, j_n) \in \mathbb{Z}^n, A \in M_n(\mathbb{N}|\mathbb{Z}_2)\}$ forms a $\mathbb{Q}(v)$ -basis of \mathbf{U}_v^0 ;
- (4) the set $\{\mathbf{m}_A^\sigma = \mathbf{F}_A^\sigma \cdot \mathbf{K}_{A,\mathbf{j}} \cdot \mathbf{E}_A^\sigma \mid \mathbf{j} = (j_1, \dots, j_n) \in \mathbb{Z}^n, A \in M_n(\mathbb{N}|\mathbb{Z}_2)\}$ forms a $\mathbb{Q}(v)$ -basis of $\mathbf{U}_v(\mathfrak{q}_n)$.

Referring to Lemma 4.4, for $1 \leq i < j \leq n$, we have

$$\begin{aligned} \mathbf{E}_{\frac{i(i+1)}{2}}^\sigma &= \mathbf{E}_{i,i+1}, & \mathbf{F}_{\frac{i(i+1)}{2}}^\sigma &= \mathbf{E}_{i+1,i}, \\ \bar{\mathbf{E}}_{\frac{i(i+1)}{2}}^\sigma &= \bar{\mathbf{E}}_{i,i+1}, & \bar{\mathbf{F}}_{\frac{i(i+1)}{2}}^\sigma &= \bar{\mathbf{E}}_{i+1,i}; \end{aligned} \quad (8.0.2)$$

and when $j > i+1$, we have

$$\begin{aligned} \mathbf{E}_{\frac{(j-2)(j-1)}{2}+i-1}^\sigma &= \mathbf{E}_{i,j}, & \mathbf{F}_{\frac{(j-2)(j-1)}{2}+i-1}^\sigma &= \mathbf{E}_{j,i}, \\ \bar{\mathbf{E}}_{\frac{(j-2)(j-1)}{2}+i-1}^\sigma &= \bar{\mathbf{E}}_{i,j}, & \bar{\mathbf{F}}_{\frac{(j-2)(j-1)}{2}+i-1}^\sigma &= \bar{\mathbf{E}}_{j,i}. \end{aligned} \quad (8.0.3)$$

We define the set for any positive integer m ,

$$(\mathbb{N}|\mathbb{Z}_2)^m = \{\mathbf{k} = (\mathbf{k}^{\bar{0}}|\mathbf{k}^{\bar{1}}) \mid \mathbf{k}^{\bar{0}} = (k_1^{\bar{0}}, k_2^{\bar{0}}, \dots, k_m^{\bar{0}}) \in \mathbb{N}^m, \mathbf{k}^{\bar{1}} = (k_1^{\bar{1}}, k_2^{\bar{1}}, \dots, k_m^{\bar{1}}) \in \mathbb{Z}_2^m\}.$$

And for any $\mathbf{t} \in (\mathbb{N}|\mathbb{Z}_2)^N$, we denote the products of all root vectors of $\mathbf{U}_v(\mathfrak{q}_n)$ by

$$\begin{aligned} \mathbf{E}^{\sigma, \mathbf{t}} &= (\mathbf{E}_N^\sigma)^{t_N^{\bar{0}}} (\bar{\mathbf{E}}_N^\sigma)^{t_N^{\bar{1}}} \cdot (\mathbf{E}_{N-1}^\sigma)^{t_{N-1}^{\bar{0}}} (\bar{\mathbf{E}}_{N-1}^\sigma)^{t_{N-1}^{\bar{1}}} \cdots (\mathbf{E}_1^\sigma)^{t_1^{\bar{0}}} (\bar{\mathbf{E}}_1^\sigma)^{t_1^{\bar{1}}}, \\ \mathbf{F}^{\sigma, \mathbf{t}} &= (\mathbf{E}_{\beta_1}^\sigma)^{t_1^{\bar{0}}} (\bar{\mathbf{F}}_{\beta_1})^{t_1^{\bar{1}}} \cdot (\mathbf{E}_{\beta_2}^\sigma)^{t_2^{\bar{0}}} (\bar{\mathbf{F}}_{\beta_2})^{t_2^{\bar{1}}} \cdots (\mathbf{E}_{\beta_N}^\sigma)^{t_N^{\bar{0}}} (\bar{\mathbf{F}}_{\beta_N})^{t_N^{\bar{1}}}. \end{aligned} \quad (8.0.4)$$

By equations (8.0.2), (8.0.3) and Proposition 8.1, we have the following results:

Theorem 8.2. *With the fixed expression w^σ , we have*

- (1) *the set $\{\mathbf{E}^{\sigma, \mathbf{t}} \mid \mathbf{t} \in (\mathbb{N}|\mathbb{Z}_2)^N\}$ forms a $\mathbb{Q}(v)$ -basis of $\mathbf{U}_v(\mathfrak{q}_n)^+$;*
- (2) *the set $\{\mathbf{F}^{\sigma, \mathbf{t}} \mid \mathbf{t} \in (\mathbb{N}|\mathbb{Z}_2)^N\}$ forms a $\mathbb{Q}(v)$ -basis of $\mathbf{U}_v(\mathfrak{q}_n)^-$;*
- (3) *the set $\{\mathbf{F}^{\sigma, \mathbf{r}} \mathbf{K}_{\mathbf{i}, \mathbf{j}} \mathbf{E}^{\sigma, \mathbf{t}} \mid \mathbf{r}, \mathbf{t} \in (\mathbb{N}|\mathbb{Z}_2)^N, \mathbf{j} = (j_1, \dots, j_n) \in \mathbb{Z}^n, \mathbf{i} = (i_1, \dots, i_n) \in \mathbb{Z}_2^n\}$ forms a $\mathbb{Q}(v)$ -basis of $\mathbf{U}_v(\mathfrak{q}_n)$.*

Open question: For any reduced decomposition w^γ different from w^σ , will Theorem 8.2 still be true?

Acknowledgements. This work was partially supported by the National Natural Science Foundation of China (Grant Nos. 12371040, 12131018 and 12161141001).

REFERENCES

- [1] E. ARTIN, *The theory of braids*, American Scientist, 38 (1946), pp. 112–119.
- [2] G. BIRKHOFF, *Representability of Lie algebras and Lie groups by matrices*, Annals of Mathematics, 38 (1937), p. 526.
- [3] F. BOHNENBLUST, *The algebraical braid group*, Annals of Mathematics, 48 (1947), p. 127.
- [4] J. BRUNDAN AND A. KLESHCHEV, *Modular representations of the supergroup $q(n)$, i* , Journal of Algebra, 260 (2003), pp. 64–98. Special Issue Celebrating the 80th Birthday of Robert Steinberg.
- [5] V. CHARI AND A. PRESSLEY, *A Guide to Quantum Groups*, Cambridge University Press, 10 1995.
- [6] S.-J. CHENG AND W. WANG, *Dualities and representations of Lie superalgebras*, Graduate Studies in Mathematics, 144 (2009).
- [7] B. DENG, J. DU, B. PARSHALL, AND J. WANG, *Finite Dimensional Algebras and Quantum Groups*, vol. 150, American Mathematical Society, Providence, 2008.
- [8] V. DRINFELD, *Hopf algebras and the quantum yang-baxter equation*, Proceedings of the USSR Academy of Sciences, 32 (1985), pp. 254–258.
- [9] V. G. DRINFEL'D, *Quantum groups*, Journal of Soviet Mathematics, 41 (1988), pp. 898–915.
- [10] J. DU, Y. LIN, AND Z. ZHOU, *Quantum queer supergroups via v -differential operators*, Journal of Algebra, (2020).
- [11] J. DU AND J. WAN, *Presenting queer Schur superalgebras*, International Mathematics Research Notices, 2015 (2015), pp. 2210–2272.
- [12] J. DU AND J. WAN, *The queer q -Schur superalgebra*, Journal of the Australian Mathematical Society, 105 (2018), pp. 1–31.
- [13] D. GRANTCHAROV, J. JUNG, S.-J. KANG, M. KASHIWARA, AND M. KIM, *Crystal bases for the quantum queer superalgebra and semistandard decomposition tableaux*, Transactions of the American Mathematical Society, 366 (2011).
- [14] D. GRANTCHAROV, J. H. JUNG, S.-J. KANG, AND M. KIM, *Highest weight modules over quantum queer superalgebra $U_q(\mathfrak{q}(n))$* , Communications in Mathematical Physics, 296 (2009), pp. 827–860.
- [15] H. GU, Z. LI, AND Y. LIN, *The integral Schur-Weyl-Sergeev duality*, Journal of Pure and Applied Algebra, 226 (2022), p. 107044.
- [16] J. C. JANTZEN, *Lectures on quantum groups*, American Mathematical Society, (1995).
- [17] M. JIMBO, *A q -difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation*, Letters in Mathematical Physics, 10 (1985), pp. 63–69.
- [18] L.D.FADDEEV, *Quantum Completely Integrable Models in Field Theory*, Soviet Scientific Reviews, 03 1980.
- [19] G. LUSZTIG, *Quantum deformations of certain simple modules over enveloping algebras*, Advances in Mathematics, 70 (1988), pp. 237–249.
- [20] G. LUSZTIG, *Finite-dimensional hopf algebras arising from quantized universal enveloping algebra*, Journal of the American Mathematical Society, 3 (1990), pp. 257–296.
- [21] G. I. OLSHANSKI, *Quantized universal enveloping superalgebra of type Q and a super-extension of the Hecke algebra*, Letters in Mathematical Physics, 24 (1992).
- [22] A. N. SERGEEV, *The tensor algebra of the identity representation as a module over the lie superalgebra $GL(n, m)$ and $Q(n)$* , Mathematics of The USSR-sbornik, 51 (1985), pp. 419–427.
- [23] E. WITT, *Treue darstellung liescher ringe.*, Journal für die reine und angewandte Mathematik (Crelles Journal), (1937), pp. 152–160.

JIANMIN CHEN, SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN 361005, CHINA

Email address: chenjianmin@xmu.edu.cn

ZHENHUA LI, JIYANG, JINAN 251400, CHINA

Email address: zhen-hua.li@qq.com

HONGYING ZHU, SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN 361005, CHINA

Email address: hongyingz@stu.xmu.edu.cn