

Periodic Orbits in Fermi-Pasta-Ulam-Tsingou Systems

Nachiket Karve, Nathan Rose, and David Campbell
Boston University

(Dated: 18 June 2024)

The FPUT paradox is the phenomenon whereby a one-dimensional chain of oscillators with nonlinear couplings shows non-ergodic behavior. The trajectory of the system in phase space, with a long wavelength initial condition, closely follows that of the Toda model over short times, as both systems seem to relax quickly to a non-thermal, metastable state. Over longer times, resonances in the FPUT spectrum drive the system towards equilibrium, away from the Toda trajectory. Similar resonances are observed in q -breather spectra, suggesting that q -breathers are involved in the route towards thermalization. In this article we investigate such resonances and show that they occur due to exact overlaps of q -breather frequencies of the type $m\Omega_1 = \Omega_k$. The resonances appear as peaks in the energy spectrum. Further, they give rise to new composite periodic orbits, which exist simultaneously with the original q -breathers. We find that such resonances are absent in integrable systems, as a consequence of the (infinite number of) conservation laws associated with integrability.

In 1953, Enrico Fermi, John Pasta, Stanislaw Ulam, and Mary Tsingou (FPUT) studied a one-dimensional chain of oscillators with nonlinear interactions that showed non-ergodic behavior¹. Since their original study, many authors have shown that the FPUT system, when initialized in a long wavelength state, goes through a long phase of metastability before eventually reaching equipartition²⁻⁴. The system is trapped in a non-thermal state during this metastable phase, with the system's energy being exponentially localized around the first mode. Resonant peaks in the energy spectrum are observed to take the system towards equilibrium albeit very slowly. Although previous attempts have been made to explain this behavior^{2,5,6}, they have not fully captured the resonance structure of the FPUT system. Here, we report on exact resonances between periodic orbits in FPUT systems. These result in the formation of resonant peaks of the energy spectra. Moreover, we show that such resonances give rise to composite periodic orbits, which exist exclusively in the nonintegrable FPUT systems and are absent in the integrable Toda lattice and KdV systems.

showed it was equivalent to the Kortewig de Vries (KdV) equation and contained the celebrated “soliton” solutions. Indeed, it was in thw ZK work that the term “solitons” was first coined. The dephasing and subsequent overlapping of these solitons has been used to explain the recurrence phenomenon and predict recurrence times. A similar analysis has been performed in the β -model⁸, whose continuum limit is the modified Kortewig de Vries (mKdV) equation. Alternatively, the Toda lattice is a discrete integrable model close to the discrete α FPUT model⁹. The Toda integrals remain approximately conserved during the time evolution in the FPUT system, which delays thermalization¹⁰. As an integrable system, the Toda lattice also exhibits exact discrete solitary wave solutions, called cnoidal waves¹¹.

A different approach, which we will focus on primarily in this paper, is to consider trajectories of the metastable state as lying close to a special class of exactly periodic orbits of the discrete FPUT systems, known as q -breathers¹². Generally, a breather is a periodic orbit localized in space, which can be continued from the linear limit. A q -breather is likewise a periodic orbit that is localized in mode space. The existence of q breathers at sufficiently small nonlinearity has been proven¹³, and a known numerical technique exists for computing them at finite nonlinearity. A benefit of this approach is that it does not require taking any continuum limits or introducing different models. As the metastable state also features exponential energy localization, it has been argued that the FPUT trajectory is a perturbation of the exactly periodic q -breather orbit¹².

As long as the q -breather is stable, a trajectory at a sufficiently small but finite distance away from it is expected to remain nearby for a long time. Thus, we investigate how the stability of q -breathers changes with the nonlinearity. We find instability regions in both α - and β -FPUT models, corresponding to exact frequency overlaps between different q -breathers. Such resonances result in the creation of new periodic orbits, or composite q -breathers, as we call them here. Composite q -breathers can be thought of as a “superposition” of two or more q -breathers. Note that such periodic orbits are absent in the linear system due to the incommensurable normal mode frequencies. Despite having similar frequency overlaps, these resonances are absent in the discrete Toda model, due to the

I. INTRODUCTION

The Fermi-Pasta-Ulam-Tsingou (FPUT) paradox deals with the extremely slow relaxation dynamics of a chain of anharmonically coupled oscillators prepared far from equilibrium. Suppose the oscillator chain is prepared in a long wavelength mode and the nonlinearity is weak, or equivalently, the energy is small. In that case, one observes a long duration where the system undergoes regular energy recurrences. Furthermore, the energy remains exponentially localized to a small number of modes. This long-lived, non-ergodic behavior is known as the metastable state^{2,3}.

Dynamics of the FPUT systems have often been understood in relation to nearby integrable nonlinear models. While the original FPUT systems correspond to discrete chains of coupled nonlinear oscillators, Zabusky and Kruskal (ZK)⁷ studied the continuum limit of the so-called α -FPUT model and

infinite number of conservation laws in the integrable Toda model, thus hinting at a possible pathway towards equilibrium in nonintegrable systems.

This paper is organized as follows. In Section II we introduce the discrete systems we have examined, namely the α - and β -FPUT models, as well as the Toda lattice. In Section III we discuss the role of solitons in the dynamics of the FPUT models as well as their continuum limits. In Section IV, we explain the numerical procedures for computing q -breathers and their stability and present some sample results. Our main results appear in Section V, where we demonstrate how resonances affect the q -breather solutions in the nonintegrable models and result in orbit bifurcations.

II. THE FPUT MODEL

The Hamiltonian of a one-dimensional chain of N identical oscillators with nonlinear interactions between nearest neighbors has the general form:

$$H = \sum_{n=1}^N \frac{1}{2} p_n^2 + \sum_{n=0}^N V(q_{n+1} - q_n). \quad (1)$$

FPUT originally studied models with cubic and quartic interactions, so that $V_{\alpha\text{-FPUT}}(x) = \frac{1}{2}x^2 + \frac{1}{3}\alpha x^3$ and $V_{\beta\text{-FPUT}}(x) = \frac{1}{2}x^2 + \frac{1}{4}\beta x^4$. The constants α and β capture the strengths of the cubic and quartic nonlinearities. On the other hand, the integrable Toda model has interactions of the form $V_{\text{Toda}}(x) = V_0 \left[e^{x/\sqrt{V_0}} - \frac{x}{\sqrt{V_0}} - 1 \right]$. Note that when V_0 is large, $V(x) \approx \frac{1}{2}x^2 + \frac{1}{6\sqrt{V_0}}x^3 + \mathcal{O}(V_0^{-1})$, which is equivalent to the α -FPUT model with $V_0 = \frac{1}{(2\alpha)^2}$. The Toda model will, therefore, approximate the dynamics of the α -FPUT system at short times^{3,14}.

We will consider here mainly lattices with fixed boundaries ($q_0 = q_{N+1} = 0$) but will switch to periodic boundaries when convenient (see appendix A for more details on systems with periodic boundaries). Different boundary conditions are not expected to introduce qualitative differences in the thermodynamic limit. The equations of motion for a system containing both α and β terms are:

$$\begin{aligned} \ddot{q}_n = & (q_{n+1} + q_{n-1} - 2q_n) + \alpha [(q_{n+1} - q_n)^2 - (q_n - q_{n-1})^2] \\ & + \beta [(q_{n+1} - q_n)^3 - (q_n - q_{n-1})^3]. \end{aligned} \quad (2)$$

The normal mode space is introduced by the following canonical transformations:

$$\begin{pmatrix} Q_k \\ P_k \end{pmatrix} = \sqrt{\frac{2}{N+1}} \sum_{n=1}^N \sin\left(\frac{\pi nk}{N+1}\right) \begin{pmatrix} q_n \\ p_n \end{pmatrix}. \quad (3)$$

The Hamiltonian in the normal space describes N harmonic oscillators with a frequency spectrum of $\omega_k = 2 \sin\left(\frac{k\pi}{2(N+1)}\right)$, interacting via many-body couplings:

$$H = \sum_{k=1}^N \frac{1}{2} (P_k^2 + \omega_k^2 Q_k^2) + \frac{\alpha}{3} \sum_{i,j,k=1}^N A_{i,j,k} Q_i Q_j Q_k$$

$$+ \frac{\beta}{4} \sum_{i,j,k,l=1}^N B_{i,j,k,l} Q_i Q_j Q_k Q_l, \quad (4)$$

where,

$$A_{i,j,k} = \frac{\omega_i \omega_j \omega_k}{\sqrt{2(N+1)}} \sum_{\pm} [\delta_{i\pm j\pm k,0} - \delta_{i\pm j\pm k,2(N+1)}], \quad (5a)$$

$$B_{i,j,k,l} = \frac{\omega_i \omega_j \omega_k \omega_l}{2(N+1)} \sum_{\pm} [\delta_{i\pm j\pm k\pm l,0} - \delta_{i\pm j\pm k\pm l,\pm 2(N+1)}]. \quad (5b)$$

In the absence of nonlinearity, the energy of each normal mode $E_k = \frac{P_k^2 + \omega_k^2 Q_k^2}{2}$ is independently conserved. It is the nonlinearity that allows mixing between different modes. On average, a thermal state has its energy equally distributed among all normal modes. We can, therefore, define a spectral entropy that measures how close a state is to thermalization:

$$S = - \sum_{k=1}^N \varepsilon_k \ln \varepsilon_k, \quad \text{where } \varepsilon_k = \frac{E_k}{\sum_{k'=1}^N E_{k'}}. \quad (6)$$

When all the system's energy is confined to a single mode, the entropy is $S = 0$, whereas the entropy of a thermal state is $S \approx \ln N$.

All FPUT systems can be scaled to have a total energy of $E = 1$ by the transformation $P_k \rightarrow P_k/\sqrt{E}$ and $Q_k \rightarrow Q_k/\sqrt{E}$. The nonlinearities transform as $\alpha \rightarrow \alpha\sqrt{E}$ and $\beta \rightarrow E\beta$ under this scaling. Therefore, the dynamics of an FPUT system can be completely captured by the parameters $E\alpha^2$ and $E\beta$.

Numerical integration of the equations of motion was done using a symplectic fourth order Runge-Kutta integrator¹⁵. The accuracy of this integrator was tested by performing time reversal tests, as described in¹⁶. A feature of studying periodic orbits is that simulations need to be performed over relatively short periods of time, allowing us to use a small timestep of $\Delta t = 10^{-4}$. The KdV and mKdV equations were integrated using a regular fourth-order Runge-Kutta integrator. Various other integrators used for nonlinear equations are discussed in¹⁷.

III. SOLITON DYNAMICS

The existence theorem¹⁸ proves the presence of solitary waves in a wide range of FPUT-like lattices. As will be demonstrated in this section, the dynamics of the FPUT system during the metastable phase is largely dominated by solitons. We will directly compare the evolution of both α - and β -FPUT systems and their continuum limits. Periodic systems will be studied in this section to avoid any complexities introduced by fixed boundaries.

The equations of motion, Eq. (2), for a periodic α -FPUT system ($q_{N+1} = q_0$) composed of particles with mass m and interactions with a spring constant k are:

$$\begin{aligned} m\ddot{q}_n = & k(q_{n+1} + q_{n-1} - 2q_n) \\ & + \alpha [(q_{n+1} - q_n)^2 - (q_n - q_{n-1})^2]. \end{aligned} \quad (7)$$




FIG. 1: Evolution of the KdV (a) and mKdV (b) equations initialized in the state $u(X, 0) = A \cos\left(\frac{2\pi x}{L}\right)$. $A = 1$ and $\delta = 0.022$ in (a), while $A = 7.071$ and $\zeta = 0.0278$ in (b). The state breaks apart into multiple solitons which almost overlap with each other again at later times and hence show recurrences of the initial state.

We take the limit where the lattice spacing a goes to zero at a constant total length of $(N + 1)a = L$. A continuous field $q(x)$ can be defined as one that approximates the lattice: $q(na) = q_n$. Up to fourth order in a , the equations of motion can be expressed in terms of spatial and temporal derivatives of q as:

$$q_{tt} = c^2 q_{xx} + \frac{c^2 a^2}{12} q_{xxxx} + c^2 \varepsilon q_x q_{xx} + \mathcal{O}(a^6), \quad (8)$$

where $c = \sqrt{\frac{k}{m}} a$ describes the speed of propagation on the lattice, and $\varepsilon = \frac{2\alpha a}{k}$. The nonlinear behavior of the lattice can be better understood by making a transformation to the co-moving frame, $X = x - ct$ and $T = \frac{c\varepsilon}{2} t$. Up to first order in ε ,

$$q_{XT} + \frac{a^2}{12\varepsilon} q_{XXXX} + q_X q_{XX} + \mathcal{O}(\varepsilon^2) = 0. \quad (9)$$

Defining $u = q_X$, and $\delta^2 = \frac{a^2}{12\varepsilon}$, we arrive at the KdV equation:

$$u_T + uu_X + \delta^2 u_{XXX} = 0. \quad (10)$$

Similar manipulations can be made to obtain the mKdV equation in the continuum limit of the β -FPUT model:

$$u_T + u^2 u_{XX} + \zeta u_{XXX} = 0, \quad (11)$$

where $\zeta = \frac{1}{36\beta}$.

We perform simulations of these continuum equations where, following⁷ and⁸, we start with a long wavelength initial state $u(X, 0) = A \cos\left(\frac{2\pi x}{L}\right)$. The KdV and mKdV systems quickly break apart into distinct solitons with varying speeds, as shown in Fig. 1. Recurrence is observed when all the soliton trajectories almost overlap again.

Note that in the corresponding FPUT systems, the soliton structure can be seen by studying the evolution of its mechanical momenta since $p(x, t) \approx -m c u(X, T)$. The continuum

initial condition translates to $q_n(t = 0) = \frac{L}{2\pi} A \sin\left(\frac{2\pi n}{N+1}\right)$, and $p_n(t = 0) = -a\sqrt{mk}A \cos\left(\frac{2\pi n}{N+1}\right)$ in the FPUT system, which amounts to initializing the system in the first mode. To view the breakdown of the system into solitons, it is useful to make a transformation to a discrete co-moving frame, defined as:

$$n \rightarrow \left[n - \frac{(N+1)\omega_1}{2\pi} t \right] \pmod{N+1}, \quad t \rightarrow t. \quad (12)$$

The breakdown of the FPUT systems into solitons is almost identical to their continuum counterparts in Fig. 2. After being initialized in the first mode, the system very quickly achieves the metastable state, which is apparent from the time-averaged entropy stabilizing to a non-thermal value. This stage is dominated by highly localized solitons, which persist for a very long time, resulting in the long lifetime of the metastable state. Unlike the continuum models, soliton collisions are expected to lead the system to equipartition eventually.

IV. PERIODIC ORBITS

Although soliton interactions in nonintegrable FPUT-like systems are expected to lead the system towards thermalization, there is a set of measure zero in the phase space containing stable, periodic trajectories, where individual solitons preserve their shape indefinitely. Such periodic orbits, or q -breathers, can be numerically computed through various iterative procedures¹⁹. Here, we describe one such procedure based on the Newton method^{12,19–21}.

A. Numerical Computation

At zero nonlinearity, all trajectories of the system are either periodic or quasi-periodic and lie on N -dimensional tori in the phase space. The Kolmogorov–Arnold–Moser (KAM) theorem²² suggests that some of these tori are continuously deformed, but preserved, as a small amount of nonlinearity is added to the system under some non-resonance conditions. Periodic orbits in the linear system are confined to single normal modes, and as nonlinearity is added, the energy spills over to other nearby modes. Such q -breathers remain exponentially localized around a particular mode.

Consider a q -breather that is continuously connected to the normal mode k_0 in the linear system. We will call k_0 the “seed mode” of this q -breather. The trajectory of this q -breather in the phase space will intersect the half-plane $P_{k_0} = 0$, $Q_{k_0} > 0$ exactly once every time period. Thus, the q -breather is a fixed point of the corresponding Poincaré map, denoted by \mathcal{F} . Under the action of \mathcal{F} , any point \mathbf{y} on the plane is evolved until its trajectory crosses the half-plane.

Fixed points of \mathcal{F} can be numerically computed using the Newton-Raphson method²³. An informed guess $\mathbf{y}^{(0)}$ of the q -breather solution is first made, which is then repeatedly updated by the action of the Newton map \mathcal{N} , defined as:

$$\mathcal{N}[\mathbf{y}] = \mathbf{y} - (\mathbb{1} - \nabla \mathcal{F}[\mathbf{y}])^{-1} (\mathbf{y} - \mathcal{F}[\mathbf{y}]), \quad (13)$$




FIG. 2: (a) Evolution of a 63 particle α -FPUT system with energy $E = 0.032$, and nonlinearity $\alpha = 2.69$ viewed in the discrete co-moving frame. The system quickly breaks apart into solitons and forms the metastable state. Note similarities with the KdV system in Fig. 1. (b) Time-Averaged spectral entropy of the system, scaled by $\ln N$, as a function of time. The metastable state is achieved when the entropy stabilizes to a non-thermal value. The formation of the metastable state coincides with the breakdown of the system into solitons. (c and d) Evolution of a 63 particle β -FPUT model with energy $E = 1.58$ and $\beta = 1$.

where the gradient map $\nabla_{\mathcal{F}}[\mathbf{y}]$ evolves the tangent vectors of \mathbf{y} . After every successive iteration $\mathbf{y}^{(i+1)} = \mathcal{N}[\mathbf{y}^{(i)}]$ the periodicity parameter $\rho = \frac{\|\mathbf{y} - \mathcal{F}[\mathbf{y}]\|}{\|\mathbf{y}\|}$, is computed, where $\|\mathbf{y}\| = \max\{y_k\}$. The tolerance ρ measures how close the current guess is to a periodic solution, and iterations are stopped once the desired accuracy is achieved.

The Newton method is always linearly stable; that is, a finite domain of convergence around a fixed point always exists. When the Poincare map admits multiple fixed points, their domains of convergence can have fractal-like boundaries. It is, therefore, important to make a good initial guess close to the desired solution. To obtain a q -breather with seed mode k_0 in a system with nonlinearity α , a good initial guess might be a q -breather with the same seed mode k_0 but in a system with

nonlinearity $\alpha - \Delta\alpha$. Thus, q -breathers in systems with arbitrary nonlinearities can be obtained by incrementing the nonlinearity in discrete steps, starting from a single normal mode in the linear system. The step size must be small enough such that every breather is in the domain of convergence of the next one.

Additional conditions need to be imposed on \mathcal{F} so that there is a unique fixed point at a given seed mode, energy, and nonlinearity. If we require the q -breather solution to be time-reversal symmetric, then there exists a time when all the mode momenta are simultaneously zero. Further, the mode amplitude Q_{k_0} can be determined given all the other mode amplitudes and the energy of the system. This reduces the number of independent variables in the initial condition of the




FIG. 3: α -FPUT: (a) Energy as a function of the mode number k , averaged over one period of the breather, for a seed mode $k_0 = 1$ q -breather in a 32 particle system with $E\alpha^2 = 4 \times 10^{-3}$. (b) Evolution of its real space momenta. β -FPUT: (c) Energy as a function of the mode number k , averaged over one period of the breather, for a seed mode $k_0 = 1$ q -breather in a 32 particle system with $E\beta = 0.4$. (d) Evolution of its real space momenta.

q -breather to $N - 1$, a huge improvement over the original $2N$.

The gradient map $\nabla \mathcal{F}$ in Eq. 13 can be computed over these $N - 1$ variables by adding perturbations and evolving the linearized equations of motion:

$$\begin{aligned} \delta \ddot{Q}_k = & -\omega_k^2 \delta Q_k - 2\alpha \sum_{i,j=1}^N A_{i,j,k} Q_i \delta Q_j \\ & - 3\beta \sum_{i,j,l=1}^N B_{i,j,k,l} Q_i Q_j \delta Q_l. \end{aligned} \quad (14)$$

By requiring the perturbation to point along a constant energy surface in the phase space, δQ_{k_0} can be determined in terms of all other components.

We use the algorithm described above to successfully compute q -breathers in both α - and β -FPUT systems with a pe-

riodicity tolerance parameter $\rho < 10^{-12}$ (see Fig. 3). As expected, the energy is exponentially localized around the seed mode of the q -breather. In the α -model, its shape is described by $\bar{\epsilon}_k \approx k^2 \left(\frac{\alpha^2 E (N+1)^3 \bar{\epsilon}_1}{\pi^4} \right)^{k-1} \bar{\epsilon}_1$ ²⁴. Note that in the β -model due to its \mathbb{Z}_2 symmetry, only modes of the same parity as the seed mode are excited. The slope of the energy spectrum depends on the nonlinearity of the system, and with increasing nonlinearity, the energy tends to spread out more in the mode space¹².

Studying the evolution of the real-space momenta reveals that q -breathers are composed of solitons (Figs. 3b and 3d). In the α -FPUT model the number of solitons in a q -breather is equal to its seed mode (Fig. 4). The direction of motion depends on the sign of the amplitude of the soliton: a left-




FIG. 4: Evolution of the momenta of seed mode $k_0 = 2$ (a) and $k_0 = 3$ (b) q -breathers in the same system as Fig. 3. The number of solitons in the breather corresponds to the seed mode number.

moving soliton is always positive, whereas a right-moving soliton is negative. On the other hand, the β -model admits both solitons and anti-solitons, and q -breathers always have an equal amount of both. The frequency of a q -breather, Ω_k , which determines the speeds of its constituent solitons, changes monotonically away from the linear mode frequencies, ω_k . For a seed mode $k_0 = 1$ breather in the α -model, perturbation theory predicts²⁴:

$$\Omega_1(E\alpha^2) = \omega_1 + E\alpha^2 \frac{\bar{\epsilon}_1 \omega_1 \omega_2}{8(N+1)(2\omega_1 - \omega_2)} + \dots \quad (15)$$

B. Stability Analysis

The stability of a q -breather can be calculated by linearizing the Poincare map around the corresponding fixed point. The $2N$ eigenvalues $\{\lambda_i\}$ of the Floquet matrix determine the linear stability of the orbit (see appendix B for more details). For a stable q -breather, all eigenvalues must lie on the unit circle. We observe that eigenvalues can only leave the unit circle in pairs, or in other words, due to an eigenvalue collision.

The q -breathers we have found in both α - and β -FPUT model are generally stable for sufficiently small nonlinearity (see Fig. 5). They become unstable around particular higher values of nonlinearity, which we will later show to be due to resonances. On the other hand, q -breathers in the Toda lattice are stable at all nonlinearities.

V. q -BREATHER RESONANCES

Resonant mechanisms in nonintegrable systems are expected to drive them towards equilibrium over a long time. This mechanism has been previously observed in the one-dimensional ϕ^4 model, where kink-antikink collisions can result in bound states²⁵. Wave turbulence theory suggests the presence of six-wave resonances in FPUT systems⁵, leading to a slow diffusion of energy from the first mode to

higher modes. The time to equilibrium is predicted to scale as $(E\alpha^2)^{-4}$ and $(E\beta)^{-8}$ in the α - and β -FPUT systems respectively²⁶. However, numerical experiments suggest that this power law fails in the weakly nonlinear regime²⁷. Instead, energy redistribution is observed to occur due to higher mode resonances. Resonant peaks are seen in the energy spectrum, which lifts it towards equipartition^{2,3}. Similar peaks are found in q -breather spectra, suggesting that q -breather resonances play a role in the thermalization of FPUT systems⁶.

Here, we show that peaks in FPUT q -breather spectra are manifestations of exact resonances between seed mode 1 and higher seed mode q -breathers. These resonances are of the type $m\Omega_1 = \Omega_k$, where Ω_k is the frequency of a higher seed mode q -breather at the same energy and nonlinearity, and m is some integer. We observe such peaks only in nonintegrable systems. Fig. 7 compares α -FPUT and Toda q -breathers at various nonlinearities. Seed mode 1 breathers are numerically computed using the Newton algorithm in a 16-particle α -FPUT system, and their frequencies (Ω_1) are plotted as a function of $E\alpha^2$ in Fig. 7a. We observe that the frequency monotonically increases with the nonlinearity. In the same figure, we plot fractions of the frequencies of certain higher mode q -breathers, Ω_k/m . A resonance between mode 1 and mode k is observed when the two plots cross. q -Breather spectra computed at nonlinearities near such frequency overlaps have peaks at the relevant resonant modes (Fig. 7b). The peak dies down quickly as we move away from the resonance (Fig. 6). Note that the nonlinearities at which we see frequency overlaps exactly correspond to the nonlinearities where the q -breather becomes unstable (Fig. 5). On the other hand, q -breather spectra in the Toda model contain no resonant peaks despite having similar frequency overlaps (Figs. 7c and 7d). This suggests that the resonant mechanism is turned off in integrable systems.

A. Composite q -Breathers

When a resonance of the type $m\Omega_1 = \Omega_k$ occurs in an FPUT system, the period of the seed mode 1 breather is an integer multiple of the period of the higher seed mode breather. This can create a new composite periodic orbit with $k+1$ solitons (Fig. 9b), where k solitons go around the lattice m times during one period. Most of the energy of this new periodic orbit is distributed amongst modes 1 and k (Fig. 9a), unlike orbits continued from single normal modes where the energy is exponentially localized around a single mode. Therefore, this orbit can be thought of as a “superposition” of two q -breathers. Note that such an orbit does not exist in the linear system. Any state in the linear system with more than one mode initialized will necessarily be quasi-periodic.

Numerical computation of the composite orbit can be a bit tricky. Due to frequency overlaps, the Poincare map \mathcal{F} admits two fixed points: one corresponds to the seed mode 1 breather, while the other corresponds to the composite breather. Both solutions can be obtained by dynamically varying the step size of the nonlinearity in the Newton algorithm. This is a brute-force procedure since one determines the step size by trial and




FIG. 5: (a). Magnitude of the largest eigenvalue of the Floquet matrix for computed q -breathers in the 16-particle α -FPUT and Toda models (a) and the 16-particle β -FPUT model (b). Unstable orbits are found in the α -FPUT at particular values of nonlinearity, while all orbits found here for the Toda model are stable. Orbits in the β -FPUT model become and remain unstable above a certain nonlinearity, but remain localized in mode space.




FIG. 6: Energy contained in the 11th mode of a seed mode $k_0 = 1$ q -breather in a 16-particle α -FPUT system, averaged over one period of the breather, as a function of the nonlinearity. The eleventh mode develops a peak when $9\Omega_1 = \Omega_{11}$, which quickly dies down as we move away from the resonance.

error.

Interestingly, it is possible to continue this composite orbit to higher nonlinearities, even when the original breathers stop resonating. The composite breather continues to exist on a separate branch, simultaneously with the original q -breathers. See Fig. 8, where a composite q -breather branch arises in both α - and β -FPUT systems due to a resonance between the first and the eleventh mode. Again, we do not observe a composite breather branch in the Toda model. It begs the question of whether composite periodic orbits exist in the Toda model, or if, like the linear system, they are quasi-periodic.

VI. CONCLUSIONS

In this article, we examined the resonant structure of α - and β -FPUT models. q -Breathers with seed mode $k_0 = 1$ in both systems are seen to develop instabilities at certain non-

linearities. These instabilities correspond to exact frequency overlaps between seed mode $k_0 = 1$ and higher seed mode q -breathers, $m\Omega_1 = \Omega_k$. These manifest as resonant peaks in the q -breather spectrum. Previous works note the similarities between such resonant peaks and those observed in the metastable state⁶. Resonances also result in bifurcations of the q -breather orbit, resulting in new composite q -breathers with multiple prominent modes. This composite breather is seen to exist at higher nonlinearities, even when the original q -breathers stop resonating. Interestingly, such a resonant mechanism is absent in the integrable Toda lattice, indicating that q -breather resonances play a role in pushing the FPUT system towards equilibrium.

Further work needs to be done to understand how breather resonances affect the dynamics of solitons in the system. It will be interesting to study soliton scattering at nonlinearities close to where resonances occur. Interactions between solitons of regular breathers and composite breathers are not well understood either. Furthermore, we speculate the existence of other forms of resonances, such as $m_k\omega_k = m_{k'}\omega_{k'}$. These could also result in new periodic orbits via a period multiplying bifurcation. This would imply the existence of a large number of unique periodic orbits, which only increases with the nonlinearity or energy. How these orbits affect the system's path to equilibrium is an open question.

VII. SUPPLEMENTARY MATERIAL

This section describes the videos included in the supplementary material. Video 1 shows the evolution of the KdV equation from Fig. 1. A timestep of $\Delta t = 10^{-4}$ is used. Video 2 shows the evolution of a seed mode $k_0 = 1$ q -breather in a 32-particle α -FPUT system with $E = 0.1$ and $\alpha = 0.2$. A timestep of $\Delta t = 10^{-3}$ is used for the simulation. In video 3, the energy spectra of seed mode $k_0 = 1$ q -breathers and the energy spectra of composite q -breathers arising from a resonance between modes 1 and 11 in a 16-particle α -FPUT system is plotted with changing $E\alpha^2$. The resonance gives rise




FIG. 7: (a) Frequency of the seed mode 1 q -breather (Ω_1) as a function of the nonlinearity in a 16 particle system, denoted by blue circles. The solid black lines denote fractions of the frequencies of higher mode q -breathers. Exact resonances are observed at $E\alpha^2 \approx 0.029, 0.068, \text{ and } 0.194$. (b) The energy spectrum of the seed mode 1 q -breather at nonlinearities $E\alpha^2 = 0.029, 0.062, \text{ and } 0.176$, which is close to where the exact resonances occur. Peaks are observed at the resonant modes. (c and d) Corresponding q -breathers in the Toda lattice. No resonant peaks are observed despite frequency overlaps.




FIG. 8: Frequencies of composite q -breathers (red triangles) obtained from a resonance between seed mode $k_0 = 1$ (blue circles) and $k_0 = 11$ (black solid line) q -breathers in α -FPUT (a) and β -FPUT (b) systems, both with 16 particles.




FIG. 9: (a) The energy spectrum of a composite q -breather (red triangles) in a 16 particle α -FPUT system with $E\alpha^2 = 0.16$. Modes 1 and 11 contain most of the energy of the breather. The energy spectra of the corresponding seed mode 1 (blue circles) and seed mode 11 (solid black line) q -breathers are shown for comparison. (b) Time evolution of the q -breather.

to a peak at mode 11 and a bifurcation is seen to occur at $E\alpha^2 \approx 0.029$.

Appendix A: Periodic Systems

Here we describe a system with the Hamiltonian (Eq. 1), but with the boundary condition $q_0 = q_{N+1}$. The normal mode space is now complex, and is defined by the canonical transformation:

$$Q_k = \frac{1}{\sqrt{N+1}} \sum_{n=0}^N q_n e^{-i\frac{2\pi kn}{N+1}}. \quad (\text{A1})$$

The condition that the real space variables must be real implies that $Q_k^* = Q_{N+1-k}$, and $P_k^* = P_{N+1-k}$. The normal mode space Hamiltonian for the $\alpha - \beta$ -FPUT system takes the form:

$$H = \sum_{k=1}^N \frac{|P_k|^2 + \omega_k^2 |Q_k|^2}{2} + \frac{\alpha}{3} \sum_{i,j,k=1}^N A_{i,j,k} Q_i Q_j Q_k + \frac{\beta}{4} \sum_{i,j,k,l=1}^N B_{i,j,k,l} Q_i Q_j Q_k Q_l, \quad (\text{A2})$$

with frequencies $\omega_k = 2 \sin\left(\frac{\pi k}{N+1}\right)$, and couplings:

$$A_{i,j,k} = i \frac{\omega_i \omega_j \omega_k}{\sqrt{N+1}} [\delta_{i+j+k, N+1} - \delta_{i+j+k, 2(N+1)}], \quad (\text{A3a})$$

$$B_{i,j,k,l} = -\frac{\omega_i \omega_j \omega_k \omega_l}{N+1} [\delta_{i+j+k+l, N+1} - \delta_{i+j+k+l, 2(N+1)} + \delta_{i+j+k+l, 3(N+1)}]. \quad (\text{A3b})$$

The linear mode energies are given by $E_k = \frac{|P_k|^2 + \omega_k^2 |Q_k|^2}{2}$. Note that $E_{N+1-k} = E_k$.

Appendix B: Floquet Theory

Since q -breathers are periodic in time, one can use Floquet theory²⁸ to study their stability. Floquet theory deals with linear differential equations of the form:

$$\dot{\mathbf{x}}(t) = \mathbf{A}(t)\mathbf{x}(t), \quad (\text{B1})$$

where $\mathbf{A}(t)$ is an $N \times N$ periodic matrix with a period T . This equation can have n linearly independent solutions of the form:

$$\mathbf{x}_i(t) = e^{\mu_i t} \mathbf{p}_i(t), \quad (\text{B2})$$

where $\mathbf{p}_i(t)$ is periodic with a period of T . $\lambda_i = e^{\mu_i T}$ s are called the characteristic multipliers of the system of equations and μ_i s are called the Floquet exponents. Further,

$$\mu_1 + \mu_2 + \dots + \mu_n = \frac{1}{T} \int_0^T dt' \text{tr}(\mathbf{A}(t')). \quad (\text{B3})$$

Note that any solution $\mathbf{x}_i(t)$ gains a phase every period:

$$\mathbf{x}_i(t + nT) = e^{n\mu_i T} \mathbf{x}_i(t). \quad (\text{B4})$$

Thus, the stability of the solution is determined by the Floquet multiplier:

- $|e^{\mu_i T}| < 1$: The solution goes to zero as $t \rightarrow \infty$,
- $|e^{\mu_i T}| = 1$: The solution is stable,
- $|e^{\mu_i T}| > 1$: The solution blows up as $t \rightarrow \infty$.

To analyze the stability of q -breathers in our system we introduce infinitesimal perturbations to the normal modes:




FIG. 10: Floquet multipliers of a stable seed mode $k_0 = 1$ q -breather (a) and a composite q -breather (b) in an 8-particle α -FPUT system.

$Q_k \rightarrow Q_k + \delta Q_k$, $P_k \rightarrow P_k + \delta P_k$. These perturbations obey the following linearized equations of motion:

$$\delta \dot{Q}_k(t) = \delta P_k(t), \quad (\text{B5a})$$

$$\begin{aligned} \delta \dot{P}_k(t) = & -\omega_k^2 \delta Q_k(t) - 2\alpha \sum_{i,j=1}^N A_{i,j,k} Q_i(t) \delta Q_j(t) \\ & - 3\beta \sum_{i,j,l=1}^N B_{i,j,k,l} Q_i(t) Q_j(t) \delta Q_l(t). \end{aligned} \quad (\text{B5b})$$

The above equations are linear and periodic, as required by Eq. B1. The Floquet multipliers of the system can be computed by evolving the perturbations over one period of the breather. Since we are dealing with Hamiltonian systems, there are additional symmetries that are imposed on the Floquet multipliers: for every Floquet multiplier λ_i , both λ_i^* and $1/\lambda_i$ are also Floquet multipliers. Therefore, a q -breather is stable if and only if all the Floquet multipliers lie on the unit circle. Examples of stable and unstable breather Floquet multipliers are given in Fig. 10.

- ¹E. Fermi, J. Pasta, S. Ulam, and M. Tsingou, “Studies of the nonlinear problems,” (1955), 10.2172/4376203.
- ²G. Benettin, A. Carati, L. Galgani, and A. Giorgilli, “The fermi–pasta–ulam problem and the metastability perspective,” in *The Fermi-Pasta-Ulam Problem: A Status Report* (Springer Berlin Heidelberg, Berlin, Heidelberg, 2008) pp. 151–189.
- ³K. A. Reiss and D. K. Campbell, “The metastable state of fermi-pasta-ulam-tsingou models,” *Entropy* **25** (2023).
- ⁴P. Lorenzoni and S. Paleari, “Metastability and dispersive shock waves in the fermi–pasta–ulam system,” *Physica D: Nonlinear Phenomena* **221**, 110–117 (2006).
- ⁵M. Onorato, L. Vozella, D. Proment, and Y. V. Lvov, “Route to thermalization in the fermi-pasta-ulam system,” *Proceedings of the National Academy of Sciences* **112**, 4208–4213 (2015), <https://www.pnas.org/doi/pdf/10.1073/pnas.1404397112>.
- ⁶T. Penati and S. Flach, “Tail resonances of Fermi-Pasta-Ulam q -breathers and their impact on the pathway to equipartition,” *Chaos: An Interdisciplinary Journal of Nonlinear Science* **17** (2007), 10.1063/1.2645141, 023102, https://pubs.aip.org/aip/cha/article-pdf/doi/10.1063/1.2645141/13556322/023102_1_online.pdf.

- ⁷N. J. Zabusky and M. D. Kruskal, “Interaction of “solitons” in a collisionless plasma and the recurrence of initial states,” *Phys. Rev. Lett.* **15**, 240–243 (1965).
- ⁸S. D. Pace, K. A. Reiss, and D. K. Campbell, “The beta Fermi-Pasta-Ulam-Tsingou recurrence problem,” *Chaos: An Interdisciplinary Journal of Nonlinear Science* **29** (2019), 10.1063/1.5122972, 113107, https://pubs.aip.org/aip/cha/article-pdf/doi/10.1063/1.5122972/14622004/113107_1_online.pdf.
- ⁹M. Toda, “Vibration of a chain with nonlinear interaction,” *Journal of the Physical Society of Japan* **22**, 431–436 (1967), <https://doi.org/10.1143/JPSJ.22.431>.
- ¹⁰G. Benettin, H. Christodoulidi, and A. Poino, “The Fermi-Pasta-Ulam Problem and Its Underlying Integrable Dynamics,” *Journal of Statistical Physics* **152**, 195–212 (2013).
- ¹¹B. S. Shastry and A. P. Young, “Dynamics of energy transport in a Toda ring,” *Phys. Rev. B* **82**, 104306 (2010).
- ¹²S. Flach, M. V. Ivanchenko, and O. I. Kanakov, “ q -breathers and the fermi-pasta-ulam problem,” *Phys. Rev. Lett.* **95**, 064102 (2005).
- ¹³R. S. MacKay and S. Aubry, “Proof of existence of breathers for time-reversible or hamiltonian networks of weakly coupled oscillators,” *Nonlinearity* **7**, 1623 (1994).
- ¹⁴A. Hofstrand, “Near-integrable dynamics of the fermi-pasta-ulam-tsingou problem,” *Phys. Rev. E* **109**, 034204 (2024).
- ¹⁵M. P. Calvo and J. M. Sanz-Serna, “The development of variable-step symplectic integrators, with application to the two-body problem,” *SIAM Journal on Scientific Computing* **14**, 936–952 (1993), <https://doi.org/10.1137/0914057>.
- ¹⁶S. D. Pace and D. K. Campbell, “Behavior and breakdown of higher-order Fermi-Pasta-Ulam-Tsingou recurrences,” *Chaos: An Interdisciplinary Journal of Nonlinear Science* **29** (2019), 10.1063/1.5079659, 023132, https://pubs.aip.org/aip/cha/article-pdf/doi/10.1063/1.5079659/14621834/023132_1_online.pdf.
- ¹⁷T. R. Taha and M. I. Ablowitz, “Analytical and numerical aspects of certain nonlinear evolution equations. iii. numerical, korteweg-de vries equation,” *Journal of Computational Physics* **55**, 231–253 (1984).
- ¹⁸G. Friesecke and J. A. D. Wattis, “Existence theorem for solitary waves on lattices,” *Communications in Mathematical Physics* **161**, 391–418 (1994).
- ¹⁹J. L. Marín and S. Aubry, “Breathers in nonlinear lattices: numerical calculation from the anticontinuous limit,” *Nonlinearity* **9**, 1501 (1996).
- ²⁰S. Flach, M. V. Ivanchenko, O. I. Kanakov, and K. G. Mishagin, “Periodic orbits, localization in normal mode space, and the Fermi–Pasta–Ulam problem,” *American Journal of Physics* **76**, 453–459 (2008), https://pubs.aip.org/aapt/ajp/article-pdf/76/4/453/13105345/453_1_online.pdf.
- ²¹S. Flach, “Computational studies of discrete breathers,” in *Energy Localization and Transfer* (World Scientific, 2004) pp. 1–71.
- ²²J. Pöschel, “A lecture on the classical kam theorem,” (2009), [arXiv:0908.2234 \[math.DS\]](https://arxiv.org/abs/0908.2234).
- ²³E. Süli and D. Mayers, *An Introduction to Numerical Analysis*, An Introduction to Numerical Analysis (Cambridge University Press, 2003).
- ²⁴A. Poino, H. Christodoulidi, C. Skokos, and S. Flach, “The two-stage dynamics in the Fermi-Pasta-Ulam problem: From regular to diffusive behavior,” *Chaos: An Interdisciplinary Journal of Nonlinear Science* **21**, 043127 (2011), https://pubs.aip.org/aip/cha/article-pdf/doi/10.1063/1.3658620/14606081/043127_1_online.pdf.
- ²⁵D. K. Campbell, J. F. Schonfeld, and C. A. Wingate, “Resonance structure in kink-antikink interactions in phi4 theory,” *Physica D: Nonlinear Phenomena* **9**, 1–32 (1983).
- ²⁶M. Onorato, Y. V. Lvov, G. Dematteis, and S. Chibbaro, “Wave turbulence and thermalization in one-dimensional chains,” *Physics Reports* **1040**, 1–36 (2023), wave Turbulence and thermalization in one-dimensional chains.
- ²⁷Z. Zhang, J. Kang, and W. Wen, “Behaviors of thermalization for the fermi–pasta–ulam–tsingou system with small number of particles*,” *Chinese Physics B* **30**, 060505 (2021).
- ²⁸B. M. Brown, M. S. P. Eastham, and K. M. Schmidt, “Floquet theory,” in *Periodic Differential Operators* (Springer Basel, Basel, 2013) pp. 1–29.
- ²⁹J. v. Neumann, “Physical applications of the ergodic hypothesis,” *Proceedings of the National Academy of Sciences* **18**, 263–266 (1932), <https://www.pnas.org/doi/pdf/10.1073/pnas.18.3.263>.
- ³⁰J. L. Tuck and M. T. Menzel, “The superperiod of the nonlinear weighted

- string (fpu) problem,” *Advances in Mathematics* **9**, 399–407 (1972).
- ³¹F. M. Izrailev and B. V. Chirikov, “Statistical Properties of a Nonlinear String,” *Soviet Physics Doklady* **11**, 30 (1966).
- ³²F. Fucito, F. Marchesoni, E. Marinari, G. Parisi, L. Peliti, S. Ruffo, and A. Vulpiani, “Approach to equilibrium in a chain of nonlinear oscillators,” *J. Phys. France* **43**, 707–713 (1982).
- ³³J. F., “The fermi-pasta-ulam problem: Paradox turns discovery,” *Physics Reports* **213**, 271–310 (1992).
- ³⁴M. A. Porter, N. J. Zabusky, B. Hu, and D. K. Campbell, “Fermi, pasta, ulam and the birth of experimental mathematics: A numerical experiment that enrico fermi, john pasta, and stanislaw ulam reported 54 years ago continues to inspire discovery,” *American Scientist* **97**, 214–221 (2009).
- ³⁵C. Danieli, D. K. Campbell, and S. Flach, “Intermittent many-body dynamics at equilibrium,” *Physical Review E* **95** (2017), 10.1103/phys-reve.95.060202.
- ³⁶A. Wazwaz, “Chapter 9 the kdv equation,” in *Handbook of Differential Equations: Evolutionary Equations*, Vol. 4, edited by C. M. Dafermos and M. Pokorný (North-Holland, 2008) pp. 485–568.
- ³⁷M. Toda, *Theory of Nonlinear Lattices*, Springer Series in Solid-State Sciences (Springer Berlin Heidelberg, 2012).
- ³⁸R. Hirota, “Exact solution of the korteweg—de vries equation for multiple collisions of solitons,” *Phys. Rev. Lett.* **27**, 1192–1194 (1971).
- ³⁹P. D. Lax, “Integrals of nonlinear equations of evolution and solitary waves,” *Communications on Pure and Applied Mathematics* **21**, 467–490 (1968), <https://onlinelibrary.wiley.com/doi/pdf/10.1002/cpa.3160210503>.
- ⁴⁰D. Hochstrasser, F. G. Mertens, and H. Büttner, “An iterative method for the calculation of narrow solitary excitations on atomic chains,” *Physica D: Nonlinear Phenomena* **35**, 259–266 (1989).
- ⁴¹J. C. Eilbeck and R. Flesch, “Calculation of families of solitary waves on discrete lattices,” *Physics Letters A* **149**, 200–202 (1990).
- ⁴²N. Flytzanis, S. Pnevmatikos, and M. Peyrard, “Discrete lattice solitons: Properties and stability,” *Journal of Physics A: Mathematical and General* **22**, 783 (1989).
- ⁴³A. Vainchtein, “Solitary waves in fpu-type lattices,” *Physica D: Nonlinear Phenomena* **434**, 133252 (2022).