

ITERATED SCHRÖDINGER BRIDGE APPROXIMATION TO WASSERSTEIN GRADIENT FLOWS

MEDHA AGARWAL, ZAID HARCHAOUI, GARRETT MULCAHY, AND SOUMIK PAL

ABSTRACT. We introduce a novel discretization scheme for Wasserstein gradient flows that involves successively computing Schrödinger bridges with the same marginals. This is different from both the forward/geodesic approximation and the backward/Jordan-Kinderlehrer-Otto (JKO) approximations. The proposed scheme has two advantages: one, it avoids the use of the score function, and, two, it is amenable to particle-based approximations using the Sinkhorn algorithm. Our proof hinges upon showing that relative entropy between the Schrödinger bridge with the same marginals at temperature ε and the joint distribution of a stationary Langevin diffusion at times zero and ε is of the order $o(\varepsilon^2)$ with an explicit dependence given by Fisher information. Owing to this inequality, we can show, using a triangular approximation argument, that the interpolated iterated application of the Schrödinger bridge approximation converge to the Wasserstein gradient flow, for a class of gradient flows, including the heat flow. The results also provide a probabilistic and rigorous framework for the convergence of the self-attention mechanisms in transformer networks to the solutions of heat flows, first observed in the inspiring work [SABP22] in machine learning research.

1. INTRODUCTION

Let $\mathcal{P}_2(\mathbb{R}^d)$ denote the space of square-integrable Borel probability measures on \mathbb{R}^d . Throughout this paper we will let $\rho \in \mathcal{P}_2(\mathbb{R}^d)$ refer to both the measure and the its Lebesgue density, whenever it exists. $\mathcal{P}_2(\mathbb{R}^d)$ can be turned into a metric space equipped with the Wasserstein-2 metric [AGS08, Chapter 7]. We will refer to this metric space as simply the Wasserstein space. For any pair of probability measures (ρ_1, ρ_2) in the Wasserstein space, let $\Pi(\rho_1, \rho_2)$ denote the set of couplings, i.e., joint distributions, with marginals (ρ_1, ρ_2) . Further, define the relative entropy, also known as Kullback-Leibler (KL) divergence, between ρ_1 and ρ_2 as

$$(1) \quad H(\rho_1|\rho_2) := \begin{cases} \int \log\left(\frac{d\rho_1}{d\rho_2}\right) d\rho_1 & \text{if } \rho_1 \ll \rho_2 \\ +\infty & \text{otherwise} \end{cases}$$

To understand our results let us start with an example that was first described in [SABP22]. The static Schrödinger bridge ([Léo14]) at temperature ε with both marginals ρ is the solution to the following optimization problem

$$(2) \quad \pi_{\rho,\varepsilon} := \operatorname{argmin}_{\pi \in \Pi(\rho,\rho)} H(\pi|\mu_{\rho,\varepsilon}),$$

Date: June 18, 2024.

2000 *Mathematics Subject Classification.* 49N99, 49Q22, 60J60.

Key words and phrases. Schrödinger bridges, Wasserstein gradient flows, JKO approximation, Fisher information, Sinkformers, Transformers, self-attention.

This research is partially supported by the following grants. All the authors are supported by NSF grant DMS-2134012. Additionally, Pal is supported by NSF grant DMS-2133244; Harchaoui is supported by NSF grant CCF-2019844, DMS-2023166, DMS-2133244. Mulcahy is supported by the National Science Foundation Graduate Research Fellowship Program under Grant No. DGE-2140004. Thanks to PIMS Kantorovich Initiative for facilitating this collaboration supported through a PIMS PRN and the NSF Infrastructure grant DMS 2133244. The authors are listed in alphabetical order.




FIGURE 1. Histograms of $n = 500$ particles at increments of one time unit. Here $(\rho_t, t \in [0, T])$ is the true gradient flow and $(\hat{\rho}_t, t \in [0, T])$ is the piecewise constant interpolation of the analytically known scheme $(\rho_k^\varepsilon, k \in [N_\varepsilon])$.

where $\mu_{\rho, \varepsilon} \in \mathcal{P}(\mathbb{R}^d \times \mathbb{R}^d)$ has density

$$(3) \quad \mu_{\rho, \varepsilon}(x, y) := \rho(x) \frac{1}{(2\pi\varepsilon)^{d/2}} \exp\left(-\frac{1}{2\varepsilon} \|x - y\|^2\right).$$

If $(X, Y) \sim \pi_{\rho, \varepsilon}$, define the **barycentric projection** [PNW22] as a function from \mathbb{R}^d to itself given by

$$(4) \quad \mathcal{B}_{\rho, \varepsilon}(x) = \mathbb{E}_{\pi_{\rho, \varepsilon}}[Y | X = x].$$

Now define a map from $\mathcal{P}_2(\mathbb{R}^d)$ to itself given by the following pushforward

$$(5) \quad \text{SB}_\varepsilon(\rho) := (2\text{Id} - \mathcal{B}_{\rho, \varepsilon})_\# \rho.$$

We call this map $\text{SB}_\varepsilon(\cdot)$ to emphasize the central role of the Schrödinger bridge in its definition. That is, if $(X, Y) \sim \pi_{\rho, \varepsilon}$, then $\text{SB}_\varepsilon(\rho)$ is the law of the random variable $2X - \mathbb{E}_{\pi_{\rho, \varepsilon}}[Y | X]$. If $\varepsilon \approx 0$, then $\pi_{\rho, \varepsilon}$ should closely approximate the quadratic cost optimal transport plan from ρ to itself [Léo12]. That is, one would expect $\mathbb{E}_{\pi_{\rho, \varepsilon}}[Y | X]$ to be approximately close to X itself (see, for instance, [PNW22, Corollary 1]). Therefore, one expects $2X - \mathbb{E}_{\pi_{\rho, \varepsilon}}[Y | X] \approx X$, as well. Hence, the pushforward should not alter ρ by much.

What happens if we iterate this map and rescale time with ε ? That is, fix some $\rho_0 \in \mathcal{P}_2(\mathbb{R}^d)$ and a $T > 0$. Let $N_\varepsilon = \lfloor T\varepsilon^{-1} \rfloor$. With $\rho_0^\varepsilon := \rho_0$, define iteratively

$$\rho_k^\varepsilon := \text{SB}_\varepsilon(\rho_{k-1}^\varepsilon), \quad k \in [N_\varepsilon] := \{1, 2, \dots, N_\varepsilon\}.$$

Define a continuous time piecewise constant interpolant $(\rho^\varepsilon(t) = \rho_{\lfloor t/\varepsilon \rfloor}^\varepsilon, t \in [0, T])$. The question is whether, as $\varepsilon \rightarrow 0+$, this curve on the Wasserstein space has an absolutely continuous limit? If so, can we write down its characterizing continuity equation?

Before we give the answer, consider the simulation in Figure 1. The figure considers a particle system approximation to the above iterative scheme. We push $n = 500$ particles, initially distributed as a mixture of two normal distributions, $\rho_0 := 0.5\mathcal{N}(-2, 1) + 0.5\mathcal{N}(2, 1)$, via the dynamical system for $T = 5$ and step size $\varepsilon = 0.1$. At every step the Schrödinger bridge is approximated by the (matrix) Sinkhorn algorithm [Cut13] applied to the current empirical distribution of the particles and the corresponding barycenter is approximated from this estimate. The pushforward is then applied to the n particles themselves. This gives a time evolving process of empirical distributions that can be thought of as a discrete approximation of our iterative scheme. One can see that the histogram gets smoothed out gradually and ends in a shape that appears to be a Gaussian distribution. In fact, this process of smoothing is reminiscent of the heat flow starting from ρ_0 . In the case of this simple starting distribution, the heat flow $(\rho_t, t \geq 0)$ is known analytically. The curves of the heat flow are superimposed as continuous curves. The broken curves are the exact $\rho^\varepsilon(\cdot)$ which, in this case, can also be computed. The histograms and the two curves all closely match with one another.

One of the results that we prove in this paper shows that, under appropriate assumptions, $(\rho^\varepsilon(t), t \in [0, T])$ indeed converges to the solution of the heat equation $\frac{\partial}{\partial t}\rho_t = \frac{1}{2}\Delta\rho_t$, with initial condition ρ_0 . In a different context and language, this result was first observed in [SABP22, Theorem 1]. However, their proof is incomplete. See the discussion below. The statement of Theorem 4 in this paper lays down sufficient conditions under which this convergence happens.

It is well known from [JKO98, Theorem 5.1] that the heat flow is the Wasserstein gradient flow of (one half) entropy. We prove a generalization of the above result where a similar discrete dynamical system converges, under assumptions, to other Wasserstein gradient flows that satisfy an evolutionary PDE of the type [AGS08, eqn. (11.1.6)]. The general existence and uniqueness of gradient flows (*curves of maximal slopes*) in Wasserstein space are quite subtle [AGS08, Chapter 11]. However, evolutionary PDEs of the type [AGS08, eqn. (11.1.6)] include the classical case of the gradient flow of entropy as the heat equation and that of the Kullback-Leibler divergence (also called relative entropy) as the Fokker-Planck equation [JKO98, Theorem 5.1]. Our iterative approximating schemes always iterate a function of the barycentric projection of the Schrödinger bridge with the *same* marginals. The marginal itself changes with the iteration. Let us give the reader an intuitive idea how such schemes work.

The foundations of all these results can be found in Theorem 1 which shows that the two point joint distribution of a stationary Langevin diffusion is a close approximation of the Schrödinger bridge with equal marginals. More concretely, let $\rho = e^{-g}$ be some element in $\mathcal{P}_2(\mathbb{R}^d)$ for which the Langevin diffusion process with stationary distribution ρ exists and is unique in law. This diffusion process is governed by the stochastic differential equation

$$(6) \quad dX_t = -\frac{1}{2}\nabla g(X_t)dt + dB_t, \quad X_0 \sim \rho,$$

where $(B_t, t \geq 0)$ is the standard d -dimensional Brownian motion. Therefore, for any $\varepsilon > 0$, $\ell_{\rho, \varepsilon} := \text{Law}(X_0, X_\varepsilon)$ is an element in $\Pi(\rho, \rho)$. Then, Theorem 1 shows that, under appropriate conditions on ρ , the symmetrized relative entropy, i.e., the Jensen-Shannon divergence, between $\ell_{\rho, \varepsilon}$ and the Schrödinger bridge $\pi_{\rho, \varepsilon}$ is $o(\varepsilon^2)$. Although this level of precision is sufficient for our purpose, it is shown via explicit computations in (28) that when ρ is Gaussian this symmetrized relative entropy is actually $O(\varepsilon^4)$, an order of magnitude tighter.

In any case, one would expect that, under suitable assumptions, the barycentric projection

$$(7) \quad \mathcal{B}_{\rho, \varepsilon}(x) \approx \mathbb{E}_{\ell_{\rho, \varepsilon}}[Y|X = x] \approx x - \frac{\varepsilon}{2}\nabla g(x) = x + \frac{\varepsilon}{2}\nabla \log \rho(x).$$

The quantity $\nabla \log \rho$ is called the score function. This is a reformulation of [SABP22, Theorem 1] which states that the difference between the two sides converges to zero in $L^2(\rho)$. In other words, one would expect that, under suitable assumptions,

$$(8) \quad \text{SB}_\varepsilon(\rho) := (2\text{Id} - \mathcal{B}_{\rho, \varepsilon})_{\#} \rho \approx \left(\text{Id} - \frac{\varepsilon}{2}\nabla \log \rho(x) \right)_{\#} \rho.$$

We claim this is approximately the solution of the heat equation at time ε , starting from ρ .

To see this, recall that absolutely continuous curves (see [AGS08, Definition 1.1.1]) $(\rho_t, t \geq 0)$ on the Wasserstein space are characterized as weak solutions [AGS08, eqn. (8.1.4)] of continuity equations

$$(9) \quad \frac{\partial}{\partial t}\rho_t + \nabla \cdot (v_t \rho_t) = 0.$$

Here $v_t : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is the time-dependent Borel velocity field of the continuity equation whose properties are given by [AGS08, Theorem 8.3.1]. For any absolutely continuous curve, if v_t

belongs to the tangent space at ρ_t ([AGS08, Definition 8.4.1]), [AGS08, Proposition 8.4.6] states the following approximation result for Lebesgue-a.e. $t \geq 0$

$$(10) \quad \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \mathbb{W}_2(\rho_{t+\varepsilon}, (\text{Id} + \varepsilon v_t)_\# \rho_t) = 0.$$

For example, if ρ is the Wasserstein gradient flow of a function $\mathcal{F} : \mathcal{P}_2(\mathbb{R}^d) \rightarrow (-\infty, \infty]$ satisfying [AGS08, eqn. (11.1.6)], $v_t = -\nabla \frac{\delta \mathcal{F}}{\delta \rho_t}$ where $\frac{\delta \mathcal{F}}{\delta \rho_t}$ is the first variation of \mathcal{F} (see [AGS08, Section 10.4.1]) at ρ_t . For the heat flow, which is the gradient flow of (half) entropy, $\frac{\delta \mathcal{F}}{\delta \rho_t} = \frac{1}{2} \log \rho_t$ and $v_t = -\frac{1}{2} \nabla \log \rho_t$. Comparing (8) with (10) explains the previous claim.

The claimed approximation in (8) is what we call *one-step approximation*. In a somewhat more general setting, this is proved in Theorem 2. Note that a crucial advantage we gain in the Schrödinger bridge scheme that is absent in Langevin diffusion or the continuity equation is that we no longer need to compute the score function $\nabla \log \rho$. Instead it suffices to be able to draw samples from ρ and compute the approximate Schrödinger bridge, say by the Sinkhorn algorithm [Cut13], as done in Figure 1.

To summarize, the flow of ideas in the paper goes as follows. First, we show that the Schrödinger bridge map $\text{SB}_\varepsilon(\rho)$, defined in its full generality in (SB), is an approximation the pushforward of ρ by the tangent vector of the gradient flow (10). This is known to be close to the gradient flow curve itself [AGS08, Proposition 8.4.6]. So one may expect that an iteration of this idea, where an absolutely continuous curve approximated by iterated pushforwards such as (10) can be, under suitable assumptions, approximated by iterated Schrödinger bridge operators $\text{SB}_\varepsilon(\cdot)$. One has to be mindful that the cumulative error remains controlled as we iterate, but in the best of situations this might work. And, in fact, it does for the heat flow, under suitable conditions, as proved in Theorem 4.

Now, a natural question is how general is this method? Can we approximate other gradient flows, say that of Kullback-Leibler with respect to a log-concave density? What about absolutely continuous curves that are not gradient flows? The case of KL is partially settled in this paper in Theorem 5. Let us state our proposed Schrödinger Bridge scheme in somewhat general notation. Consider a suitably nice functional $\mathcal{F} : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R} \cup \{+\infty\}$ and fix an initial measure $\rho_0 \in \mathcal{P}_2(\mathbb{R}^d)$. Suppose the gradient flow of \mathcal{F} can be approximated by a sequence of explicit Euler iterations $(\tilde{\rho}_\varepsilon(k), k \in \mathbb{N})$, where, starting from $\tilde{\rho}_\varepsilon(0) = \rho_0$, iteratively define

$$(11) \quad \tilde{\rho}_\varepsilon(k) = (\text{Id} + \varepsilon \nabla u_{k-1})_\# \tilde{\rho}_\varepsilon(k-1)$$

with $-\nabla u_{k-1}$ being the Wasserstein gradient of \mathcal{F} at $\tilde{\rho}_\varepsilon(k-1)$. Such an approximation scheme will be referred to as an explicit Euler iteration scheme and conditions must be imposed for it to converge to the underlying gradient flow. See Theorem 3 below for sufficient conditions.

Ignoring technical details, our goal is to approximate each such explicit Euler step via a corresponding Schrödinger bridge step. This is accomplished by the introduction of what we call a *surrogate measure*. For simplicity, consider the first step $\tilde{\rho}_\varepsilon(1) = (\text{Id} + \varepsilon \nabla u_0)_\# \rho_0$. Suppose that there is some $\theta \in \mathbb{R} \setminus \{0\}$ such that $\int e^{2\theta u_0} < \infty$, define the **surrogate measure** σ_0 to be one with the density

$$(12) \quad \sigma_0(x) := \exp(2\theta u_0(x) - \Lambda_0(\theta)), \quad \text{where} \quad \Lambda_0(\theta) := \log \int e^{2\theta u_0(y)} dy.$$

As explained in (7), if we consider the Schrödinger bridge at temperature ε with both marginals to be σ_0 , then $\mathcal{B}_{\sigma_0, \varepsilon}(x) \approx x + \frac{\varepsilon}{2} \nabla \log \sigma_0(x) = x + \theta \varepsilon \nabla u_0(x)$. If we now define one step in the SB scheme as

$$(SB) \quad \text{SB}_\varepsilon(\rho_0) = \left(\left(1 - \frac{1}{\theta}\right) \text{Id} + \frac{1}{\theta} \mathcal{B}_{\sigma_0, \varepsilon} \right)_\# \rho_0.$$

Observing that, as $\varepsilon \downarrow 0$, we recover that $\text{SB}_\varepsilon(\rho) \approx (\text{Id} + \varepsilon \nabla u_0)_\# \rho$. Notice that the choice of θ does not play a role except to guarantee that the surrogate σ_0 in (12) is integrable. For the case of the heat flow, $u_0 = -\frac{1}{2} \log \rho_0$. Hence, if we pick $\theta = -1$, $\sigma_0 = \rho_0$ and then the scheme (5) becomes a special case of (SB).

Let us consider the sequence of iterates of (SB), interpolated in continuous time for some $0 < T < \infty$,

$$\rho^\varepsilon((k+1)/\varepsilon) := \text{SB}_\varepsilon(\rho^\varepsilon(k/\varepsilon)), \quad k \in [N_\varepsilon], \quad \text{where } N_\varepsilon := \lfloor T/\varepsilon \rfloor.$$

Theorems 4 and 5 list sufficient conditions under which such a scheme converges uniformly to the gradient flow curve as $\varepsilon \rightarrow 0+$ for specific choices of functional \mathcal{F} .

We should point out that the primary reason why we need the assumptions in Theorems 4 and 5 is to guarantee that the errors go to zero uniformly over a growing number of iterations. Much of the technical work in this paper goes to produce a simple set of sufficient conditions to ensure this. Consequently, our theorems are specialized to the case of gradient flows of geodesically convex regular functions. But the scope of Schrödinger bridge scheme is potentially larger. As shown in Section 5.3 via explicit calculations, the Schrödinger bridge scheme may be used to approximate even time-reversed gradient flows. The corresponding PDEs, such as the backward heat equation/ Fokker-Planck, are ill-posed and care must be taken to implement any approximating scheme. Interestingly, if one replaces (5) by $\text{SB}_\varepsilon(\rho) := (\mathcal{B}_{\rho, \varepsilon})_\# \rho$, then starting from a Gaussian initial distribution, and for all small enough ε , the iterates are shown to converge to the time-reversed gradient flow of one-half entropy.

In Section 3 we also generalize our Schrödinger bridge scheme (SB) by picking the surrogate measure *depending on* ε . This generalization is necessary to generalize the Schrödinger bridge scheme to cover tangent vectors that are not necessarily of the gradient type, but are approximated by a sequence of gradient fields. This generalization also covers the case of the renowned JKO scheme, [JKO98], which can be likened to the implicit Euler discretization scheme of the gradient flow curve. This, however, introduces an additional layer of complexity.

Some of the technical challenges we overcome in this paper involve proving results about Schrödinger bridges at low temperatures when the marginals are not compactly supported. The latter is a common assumption in the literature, see for example [Pal19, PNW22, CCGT22, SABP22]. However, the existence of Langevin diffusion requires an unbounded support. Much of this work is done in Section 2 which may be of independent interest. For example, we extend the expansion of entropic OT cost function from [CT21, Theorem 1.6] for the case of same marginals in (42). In all this analysis an unexpected mathematical object appears repeatedly, the *harmonic characteristic* ([CVR18]) defined below in (14). It appears in the computation of Fisher information of the Langevin and the Schrödinger bridges (see (37)) and in the geodesic approximation of the heat flow (Lemma 7). The role of this object merits a deeper study.

Discussion on [SABP22]. The inspiring work [SABP22] is one of the motivations of our investigations. The authors of [SABP22] propose and analyze a modification to the self-attention mechanism from the Transformer artificial neural network architecture in which the self-attention matrix is enforced to be doubly-stochastic. Since the Sinkhorn algorithm is applied to enforce double stochasticity, the architecture is called the Sinkformer. A major contribution of [SABP22] is the keen observation that self-attention mechanisms can be understood as determining a dynamical system whose particles are the input tokens (possibly after a suitable transformation). See also [GLPR24] for other work in this area. Specifically, [SABP22, Proposition 2,3] observes that, as a regularization parameter vanishes, the associated particle system under doubly-stochastic self-attention evolves according to the heat flow, and this property naturally follows from the doubly-stochastic constraints.

Theorems 1-5 can be understood as probabilistic approaches to [SABP22, Theorem 1], as well as a finer treatment to key steps of the mathematical analysis of the convergence to the heat flow equation. The first statement of [SABP22, Theorem 1] concerns the convergence of the rescaled gradients of entropic potentials for the Schrödinger bridge with the same marginals. Consider the so-called entropic (or, Schrödinger) potentials $(f_\varepsilon, \varepsilon > 0)$ as defined later on in (19). In this language, [SABP22, Theorem 1] states that $\varepsilon^{-1}\nabla f_\varepsilon \rightarrow -\nabla \log \rho$ in $L^2(\rho)$, as $\varepsilon \rightarrow 0+$. We are not able to prove this particular convergence; not least because we work in a slightly different setting, i.e., with measures fully supported on \mathbb{R}^d whereas [SABP22] assumes compact support (although, see the discussion after the proof of Theorem 1 for a similar result). Instead our argument relies on approximating the Schrödinger bridge by the Langevin diffusion which are undefined for measures with compact support. Moreover, in our paper we develop a general machinery that demonstrate the ability of the Schrödinger bridge scheme to approximate gradient flows of other functionals of interest beyond entropy.

2. APPROXIMATION OF SCHRÖDINGER BRIDGE BY LANGEVIN DIFFUSION

2.1. Preliminaries. In this section we fix a measure $\rho \in \mathcal{P}_2(\mathbb{R}^d)$ with density given by $\rho = e^{-g}$ for some function $g : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$. Here, we follow the standard convention in optimal transport of using the same notation to refer to an absolutely continuous measure and its density. We will state assumptions on g later. Let $\mathcal{P}_2^{ac}(\mathbb{R}^d)$ denote the collection of probability measures on \mathbb{R}^d with density and finite second moments. The aim of this section is to prove that when the two marginals are the same, then the Schrödinger bridge and the stationary Langevin diffusion with that marginal are very close in the sense made precise in Theorem 2.

We start by collating standard results about the Langevin diffusion and Schrödinger bridge and introducing the notions from stochastic calculus of which we will make use. The relevant notation is all summarized in Table 5.3.

Let $C^d[0, \infty)$ denote the set of continuous paths from $[0, \infty)$ to \mathbb{R}^d , equipped with the locally uniform metric. Denote the coordinate process by $(\omega_t, t \geq 0)$. Endow this space with a filtration satisfying the *usual conditions* [KS91, Chapter 1, Definition 2.25]. For $x \in \mathbb{R}^d$, let W_x denote the law on $C^d[0, \infty)$ of the standard d -dimensional Brownian motion started from x . All stochastic integrals are Itô integrals.

Let $g \in C^2(\mathbb{R}^d)$ and assume that $(Y_t, t \geq 0)$, the stationary Langevin diffusion with stationary measure equal to $\rho = e^{-g}$, exists. That is, its law on $C^d[0, \infty)$ is a weak solution to (6) with initial condition $Y_0 \sim \rho$. Let Q denote the law of this process on $C^d[0, \infty)$, and let Q_x denote the law of the Langevin diffusion started from x . As in the Introduction, for $\varepsilon > 0$, set $\ell_{\rho, \varepsilon} = \text{Law}(Y_0, Y_\varepsilon)$. Under suitable assumptions on g (see Assumption 1 below), $(Y_t, t \geq 0)$ exists with a.s. infinite explosion time [Roy07, Theorem 2.2.19], and by [Roy07, Lemma 2.2.21] for each time $t > 0$ the laws of Q_x and P_x restricted to $C^d[0, t]$ are mutually absolutely continuous with Radon-Nikodym derivative

$$(13) \quad \frac{dQ_x}{dW_x} \Big|_{C^d[0, t]}(\omega) = \sqrt{\frac{\rho(\omega_t)}{\rho(x)}} \exp \left(- \int_0^t \left(\frac{1}{8} \|\nabla g(\omega_s)\|^2 - \frac{1}{4} \Delta g(\omega_s) \right) ds \right).$$

The expression inside the integral of the $\exp(\cdot)$ is a distinguished quantity in the Langevin diffusion literature (see [LK93]). Denote it by

$$(14) \quad \mathcal{U}(x) := \frac{1}{8} \|\nabla g(x)\|^2 - \frac{1}{4} \Delta g(x).$$

This function is called the harmonic characteristic in [CVR18, eqn. (2)], which provides a heuristic physical interpretation for \mathcal{U} and $\nabla \mathcal{U}$ (the latter of which appears in Theorem 1) as the “mean acceleration” of the Langevin bridge. This quantity also appears in (69) as

the second derivative in time of a quantity of interest, giving another way to see it as mean acceleration.

Let $(p_t(\cdot, \cdot), t > 0)$ and $(q_t(\cdot, \cdot), t > 0)$ denote the transition kernels for the standard d -dimensional Brownian motion and the Langevin diffusion, respectively. By (13)

$$\frac{q_t(x, y)}{p_t(x, y)} = \sqrt{\frac{\rho(y)}{\rho(x)}} \mathbb{E}_{W_x} \left[\exp \left(- \int_0^t \mathcal{U}(\omega_s) ds \right) \middle| \omega_t = y \right].$$

Notice that $\mathbb{E}_{W_x}[\cdot | \omega_t = y]$ is expectation with respect to the law of the Brownian bridge from x to y over $[0, t]$. Define the following function

$$(15) \quad c(x, y, \varepsilon) := - \log \left(\mathbb{E}_{W_x} \left[\exp \left(- \int_0^\varepsilon \mathcal{U}(\omega_s) ds \right) \middle| \omega_\varepsilon = y \right] \right).$$

Recall that $\ell_{\rho, \varepsilon}$ is joint density of (Y_0, Y_ε) under Langevin; it is now equal to

$$(16) \quad \ell_{\rho, \varepsilon}(x, y) = \rho(x) q_\varepsilon(x, y) = \sqrt{\rho(x) \rho(y)} p_\varepsilon(x, y) \exp(-c(x, y, \varepsilon))$$

$$(17) \quad = \sqrt{\rho(x) \rho(y)} \frac{1}{(2\pi\varepsilon)^{d/2}} \exp \left(- \frac{1}{2\varepsilon} \|x - y\|^2 - c(x, y, \varepsilon) \right).$$

Let $(G_t, t \geq 0)$ denote the corresponding Langevin semigroup and L its extended infinitesimal generator (see [RY04, Chapter VII, Section 1] for definitions). That is, we have for $t \geq 0$ and $f : \mathbb{R}^d \rightarrow \mathbb{R}$ bounded and measurable that

$$G_t f(x) := \mathbb{E}_Q[f(\omega_t) | \omega_0 = x].$$

Letting $C^2(\mathbb{R}^d)$ denote the collection of all twice continuously differentiable functions, by [RY04, Chapter VIII, Proposition 3.4] we have for $f \in C^2(\mathbb{R}^d)$ that

$$(18) \quad Lf = \frac{1}{2} \Delta f + \left(-\frac{1}{2} \nabla g \right) \cdot \nabla f.$$

We now define the static Schrödinger bridge with marginals equal to ρ , denoted by $\pi_{\rho, \varepsilon}$. For an excellent and detailed account we refer readers to [Léo14] and references therein; we develop only the notions that we use within our proof. Under mild regularity assumptions on ρ (for instance, finite entropy and a moment condition), it is known that for each $\varepsilon > 0$, $\pi_{\rho, \varepsilon}$ admits an (f, g) -decomposition [Léo14, Theorem 2.8]. As we consider the equal marginal Schrödinger bridge, we can in fact insist that $f = g$. Thus, there exists $f_\varepsilon : \mathbb{R}^d \rightarrow \mathbb{R}$ such that $\pi_{\rho, \varepsilon}$ admits a Lebesgue density of the form

$$(19) \quad \pi_{\rho, \varepsilon}(x, y) = \frac{1}{(2\pi\varepsilon)^{d/2}} \exp \left(\frac{1}{\varepsilon} \left(f_\varepsilon(x) + f_\varepsilon(y) - \frac{1}{2} \|x - y\|^2 \right) \right) \rho(x) \rho(y).$$

We refer to the $(f_\varepsilon, \varepsilon > 0)$ as entropic potentials. Note that, in general, entropic potentials are unique up to a choice of constant; however, our convention of symmetry forces that each f_ε is indeed unique. Moreover, combining (3) and (19), the Schrödinger cost can be written in terms of the entropic potentials as

$$(20) \quad H(\pi_{\rho, \varepsilon} | \mu_{\rho, \varepsilon}) = \mathbb{E}_{\pi_{\rho, \varepsilon}} \left[\log \frac{d\pi_{\rho, \varepsilon}}{d\mu_{\rho, \varepsilon}} \right] = \mathbb{E}_\rho \left[\frac{2}{\varepsilon} f_\varepsilon(X) \right] + \text{Ent}(\rho).$$

As outlined in [Léo14, Proposition 2.3], there is a corresponding dynamic Schrödinger bridge, obtained from $\pi_{\rho, \varepsilon}$ (the “static” Schrödinger bridge) by interpolating with suitably rescaled Brownian bridges. More precisely, let R_ε denote the law of $(\sqrt{\varepsilon} B_t, t \in [0, 1])$ on $C^d[0, 1]$, where

$(B_t, t \geq 0)$ is the reversible d -dimensional Brownian motion. Then, the law of the ε -dynamic Schrödinger bridge on $C^d[0, 1]$ is given by $S_{\rho, \varepsilon}$ where

$$(21) \quad \frac{dS_{\rho, \varepsilon}}{dR_\varepsilon}(\omega) = a^\varepsilon(\omega_0)a^\varepsilon(\omega_1),$$

where $a^\varepsilon(x) := \exp\left(\frac{1}{\varepsilon}f_\varepsilon(x) + \log \rho(x)\right)$. Letting $(P_t, t \geq 0)$ denote the (standard) heat semi-group, we define

$$(22) \quad \psi^\varepsilon(t, x) := \log E_{R_\varepsilon} [a^\varepsilon(\omega_1)|\omega_t = x] = \log E_{R_\varepsilon} [a^\varepsilon(\omega_{1-t})|\omega_0 = x] = \log P_{\varepsilon(1-t)}a^\varepsilon(x),$$

where the second identity follows from the Markov property of R_ε .

Fix $\varepsilon > 0$, set $\rho_s^\varepsilon := (\pi_s)_\# S_{\rho, \varepsilon}$ and call $(\rho_s^\varepsilon, s \in [0, 1])$ the ε -**entropic interpolation** from ρ to itself, explicitly

$$(23) \quad \rho_t^\varepsilon = P_{\varepsilon t} a^\varepsilon P_{\varepsilon(1-t)} a^\varepsilon = \exp(\psi^\varepsilon(t, \cdot) + \psi^\varepsilon(1-t, \cdot)).$$

By [Léo14, Proposition 2.3], we construct a random variable with law ρ_t^ε as follows. Let $(X_\varepsilon, Y_\varepsilon) \sim \pi_{\rho, \varepsilon}$ and $Z \sim N(0, \text{Id})$ be independent to the pair, then

$$(24) \quad X_t^\varepsilon = (1-t)X_\varepsilon + tY_\varepsilon + \sqrt{\varepsilon t(1-t)}Z \sim \rho_t^\varepsilon.$$

Thanks to an analogue of the celebrated Benamou-Brenier formula for entropic cost ([GLR17, Theorem 5.1]), the entropic interpolation is in fact an absolutely continuous curve in the Wasserstein space. From [CT21, Equation (2.14)] the entropic interpolation satisfies the continuity equation (9) with $(\rho_t^\varepsilon, v_t^\varepsilon)_{t \in [0, 1]}$ given by

$$(25) \quad v_t^\varepsilon = \frac{\varepsilon}{2} (\nabla \psi^\varepsilon(t, \cdot) - \nabla \psi^\varepsilon(1-t, \cdot)).$$

Moreover, under mild regularity assumptions on ρ , we obtain the following entropic variant of Benamou-Brenier formulation [GLR17, Corollary 5.8] (with appropriate rescaling)

$$(26) \quad H(\pi_{\rho, \varepsilon} | \mu_{\rho, \varepsilon}) = \inf_{(\rho_t, v_t)} \left(\frac{1}{2\varepsilon} \int_0^1 \|v_t\|_{L^2(\rho_t)}^2 dt + \frac{\varepsilon}{8} \int_0^1 I(\rho_t) dt \right),$$

where $I(\cdot)$ is the Fisher information, defined for $\sigma \in \mathcal{P}_2^{ac}(\mathbb{R}^d)$ by $I(\sigma) = \int \|\nabla \log \sigma\|^2 d\sigma$. The infimum in (26) is taken over all $(\rho_t, v_t)_{t \in [0, 1]}$ satisfying $\rho_0 = \rho_1 = \rho$ and (9). The optimal selection is given in (25). That is, the optimal selection is the entropic interpolation. For an AC curve $(\rho_t, t \in [0, 1])$, we will call the quantity $\int_0^1 I(\rho_t) dt$ the integrated Fisher information.

Lastly, for the same dynamic marginal Schrödinger bridge, we make the following observation about the continuity at $\varepsilon = 0$ of the integrated Fisher information along the entropic interpolation.

Lemma 1. *Let $\rho \in \mathcal{P}_2(\mathbb{R}^d)$ satisfy $-\infty < \text{Ent}(\rho) < +\infty$ and $I(\rho) < +\infty$, then $\lim_{\varepsilon \downarrow 0} \int_0^1 I(\rho_t^\varepsilon) dt = I(\rho)$.*

Proof. By [Léo12, Theorem 3.7(3)], for each $s \in [0, 1]$, ρ_s^ε converges weakly to ρ as $\varepsilon \downarrow 0$. By the lower semicontinuity of Fisher information with respect to weak convergence [Bob22, Proposition 14.2] and Fatou's Lemma,

$$I(\rho) = \int_0^1 I(\rho) dt \leq \int_0^1 \liminf_{\varepsilon \downarrow 0} I(\rho_t^\varepsilon) dt \leq \liminf_{\varepsilon \downarrow 0} \int_0^1 I(\rho_t^\varepsilon) dt.$$

Lastly, by (26) and the suboptimality of the constant curve $\rho_t = \rho, v_t = 0$ for $t \in [0, 1]$

$$\frac{1}{2\varepsilon} \int_0^1 \|v_t^\varepsilon\|_{L^2(\rho_t^\varepsilon)}^2 dt + \frac{\varepsilon}{8} \int_0^1 I(\rho_t^\varepsilon) dt \leq \frac{\varepsilon}{8} I(\rho) \Rightarrow \int_0^1 I(\rho_t^\varepsilon) dt \leq I(\rho).$$

Altogether, it follows

$$0 \leq \limsup_{\varepsilon \downarrow 0} \left(I(\rho) - \int_0^1 I(\rho_t^\varepsilon) dt \right) = I(\rho) - \liminf_{\varepsilon \downarrow 0} \int_0^1 I(\rho_t^\varepsilon) dt \leq I(\rho) - I(\rho) = 0.$$

□

2.2. Langevin Approximation to Schrödinger Bridge. Our main result of the section is the relative entropy bound between $\ell_{\rho,\varepsilon}$ and $\pi_{\rho,\varepsilon}$ given in Theorem 1.

Assumption 1. Let $\rho = e^{-g}$ for $g : \mathbb{R}^d \rightarrow \mathbb{R}$, and recall that $\mathcal{U} = \frac{1}{8}\|\nabla g\|^2 - \frac{1}{4}\Delta g$. Assume that

- (1) ρ is subexponential ([Ver18, Definition 2.7.5]), $g \in C^3(\mathbb{R}^d)$, \mathcal{U} is bounded below, $g \rightarrow \infty$ as $\|x\| \rightarrow \infty$.
- (2) There exists $C > 0$ and $n \geq 1$ such that $\|\nabla \mathcal{U}(x)\|^2 \leq C(1 + \|x\|^n)$.
- (3) $-\infty < \text{Ent}(\rho) < +\infty$ and $I(\rho) < +\infty$.

Note that Assumption 1 (3) ensures the existence $\pi_{\rho,\varepsilon}$ and that (26) holds [GLR17, (Exi),(Reg1),(Reg2)].

These assumptions are, for instance, satisfied by the multivariate Gaussian $N(\mu, \Sigma)$ with $\mu \in \mathbb{R}^d$, $\Sigma \in \mathbb{R}^{d \times d}$ positive definite. This provides an essential class of examples as explicit computations of the Schrödinger bridge between Gaussians are known [JMPC20]. In this instance, $g(x) = \frac{1}{2}(x - \mu)^T \Sigma^{-1}(x - \mu) + \frac{1}{2} \log((2\pi)^d \det \Sigma)$ and thus

$$\mathcal{U}(x) = \frac{1}{8}(x - \mu)^T \Sigma^{-2}(x - \mu) - \frac{1}{4} \text{Tr}(\Sigma^{-1}) \text{ and } \nabla \mathcal{U}(x) = \frac{1}{4} \Sigma^{-2}(x - \mu).$$

Importantly, for $X \sim N(\mu, \Sigma)$ this establishes that $\nabla \mathcal{U}(X) \sim N(0, \frac{1}{16} \Sigma^{-3})$ and thus $\mathbb{E} \|\nabla \mathcal{U}(X)\|^2 = \frac{1}{16} \text{Tr}(\Sigma^{-3})$. More generally, when g is polynomial such that $\int_{\mathbb{R}^d} e^{-g} < +\infty$ these assumptions are also satisfied.

Additionally, we remark that the class of functions g satisfying Assumption 1 is stable under the addition of functions $h \in C^3(\mathbb{R}^d)$ such that h and all of its derivatives up to and including order 3 are bounded (e.g. $h \in C_c^3(\mathbb{R}^d)$). It is also stable under the addition of linear and quadratic functions, so long as the resulting function has a finite integral when exponentiated.

We begin with a preparatory lemma establishing some important integrability properties we make use of in the proof of Theorem 1.

Lemma 2. Let $\rho = e^{-g} \in \mathcal{P}_2(\mathbb{R}^d)$ satisfying Assumption 1.

- (1) For each $\varepsilon > 0$, the densities along the entropic interpolation $(\rho_s^\varepsilon, s \in [0, 1])$ is uniformly subexponential, meaning that for each $T > 0$ there is a constant $K > 0$ such that for $X_s^\varepsilon \sim \rho_s^\varepsilon$ we have $\sup_{\varepsilon \in (0, T)} \sup_{s \in [0, 1]} \|X_s^\varepsilon\|_{\psi_1} \leq K$, where $\|\cdot\|_{\psi_1}$ is the subexponential norm defined in [Ver18, Definition 2.7.5].
- (2) The following integrability conditions hold:
 - (a) The function $\mathcal{U} = \frac{1}{8}\|\nabla g\|^2 - \frac{1}{4}\Delta g$ is in $L^2(\rho_s^\varepsilon)$ for all $s \in [0, 1]$.
 - (b) The function from $[0, 1] \times C^d[0, 1]$ to \mathbb{R} given by $(t, \omega) \mapsto \mathcal{U}(\omega_t)$ is in $L^2(dt \otimes dS_{\rho,\varepsilon})$. Similarly, $(t, \omega) \mapsto \nabla \mathcal{U}(\omega_t)$ is in $L^2(dt \otimes dS_{\rho,\varepsilon})$.
 - (c) For any $T > 0$, $\sup_{\varepsilon \in (0, T)} \int_0^1 \mathbb{E}_{S_{\rho,\varepsilon}} [\|\nabla \mathcal{U}(\omega_s)\|^2] ds < +\infty$.

Proof. Fix $\varepsilon > 0$ and $s \in [0, 1]$. Let (X, Y) be random variables with joint law $\pi_{\rho, \varepsilon}$. Let $Z \sim N(0, \text{Id})$ be independent of (X, Y) . Then X_s^ε as given in (24) has law ρ_s^ε . As $\|\cdot\|_{\psi_1}$ is a norm, observe that for any $T > 0$

$$\sup_{\varepsilon \in (0, T], s \in [0, 1]} \|X_s^\varepsilon\|_{\psi_1} \leq \sup_{\varepsilon \in (0, T], s \in [0, 1]} \left((1-s)\|X\|_{\psi_1} + s\|Y\|_{\psi_1} + \sqrt{\varepsilon s(1-s)}\|Z\|_{\psi_1} \right) < +\infty$$

establishing the uniform upper bound on the subexponential norm.

It then follows that

$$\sup_{\varepsilon \in (0, T], s \in [0, 1]} \mathbb{E}[\|\nabla \mathcal{U}(X_s^\varepsilon)\|^2] \leq \sup_{\varepsilon \in (0, T], s \in [0, 1]} C(1 + \mathbb{E}[\|X_s^\varepsilon\|^n]) < +\infty.$$

This observation also establishes that $(t, \omega) \mapsto \nabla \mathcal{U}(\omega_t)$ is in $L^2(dt \otimes dS_{\rho, \varepsilon})$. By assumption on $\nabla \mathcal{U}$, \mathcal{U} is also of uniform polynomial growth, and thus the same argument establishes the remaining claims. \square

Theorem 1. For $\rho \in \mathcal{P}_2(\mathbb{R}^d)$ satisfying Assumption 1(1,3), for all $\varepsilon > 0$

$$(27) \quad H(\ell_{\rho, \varepsilon} | \pi_{\rho, \varepsilon}) + H(\pi_{\rho, \varepsilon} | \ell_{\rho, \varepsilon}) \leq \frac{1}{2} \varepsilon^2 \left(I(\rho) - \int_0^1 I(\rho_t^\varepsilon) dt \right)^{1/2} \left(\int_0^1 \mathbb{E}_{\rho_t^\varepsilon} \|\nabla \mathcal{U}\|^2 dt \right)^{1/2}.$$

In particular, additionally under Assumption 1(2),

$$\lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon^2} (H(\ell_{\rho, \varepsilon} | \pi_{\rho, \varepsilon}) + H(\pi_{\rho, \varepsilon} | \ell_{\rho, \varepsilon})) = 0.$$

Before proving the theorem, we pause to comment on the sharpness of the inequality presented in (27). As the Schrödinger bridge is explicitly known for Gaussian marginals [JMPC20], consider the case $\rho = N(0, 1)$. Define two matrices

$$\Sigma_1 = \begin{pmatrix} 1 & e^{-\varepsilon/2} \\ e^{-\varepsilon/2} & 1 \end{pmatrix}, \Sigma_2 = \begin{pmatrix} 1 & \frac{1}{2}(\sqrt{\varepsilon^2 + 4} - \varepsilon) \\ \frac{1}{2}(\sqrt{\varepsilon^2 + 4} - \varepsilon) & 1 \end{pmatrix}.$$

It is well known that $\ell_{\rho, \varepsilon} \sim N(0, \Sigma_1)$ is the Ornstein-Uhlenbeck transition density. By [JMPC20], it is known that $\pi_{\rho, \varepsilon} \sim N(0, \Sigma_2)$. One then computes

$$(28) \quad H(\ell_{\rho, \varepsilon} | \pi_{\rho, \varepsilon}) + H(\pi_{\rho, \varepsilon} | \ell_{\rho, \varepsilon}) = \frac{1}{2} \text{Tr}(\Sigma_1^{-1} \Sigma_2) + \frac{1}{2} \text{Tr}(\Sigma_2^{-1} \Sigma_1) - 2 = \frac{1}{1152} \varepsilon^4 + \mathcal{O}(\varepsilon^5).$$

On the other hand, the entropic interpolation between Gaussians is computed in [GLR17], and $I(\rho_t^\varepsilon)$ is minimized at $t = 1/2$. From a computation in Section 5, see (71),

$$I(\rho) - \int_0^1 I(\rho_t^\varepsilon) dt \leq I(\rho) - I(\rho_{1/2}^\varepsilon) = \mathcal{O}(\varepsilon^2).$$

Altogether, the RHS of (27) is then $\mathcal{O}(\varepsilon^3)$. That is, the transition densities for the OU process provide a tighter approximation (an order of magnitude) for same marginal Schrödinger bridge than given by Theorem 1.

Proof of Theorem 1. We argue in the following manner. First, using the densities for $\pi_{\rho, \varepsilon}$ and $\ell_{\rho, \varepsilon}$ given in (16) and (19), respectively, we compute their likelihood ratio and obtain exact expressions for the two relative entropy terms appearing in the statement of the Lemma. The resulting expression is the difference in expectation of the expression $c(x, y, \varepsilon)$ defined in (15) with respect to the Schrödinger bridge and Langevin diffusion. We then extract out the leading order terms from both expectations and show that they cancel. To obtain the stated rate of decay, we then analyze the remaining term with an analytic argument from the continuity equation.

Step 1. The following holds

$$(29) \quad H(\ell_{\rho,\varepsilon}|\pi_{\rho,\varepsilon}) + H(\pi_{\rho,\varepsilon}|\ell_{\rho,\varepsilon}) = \mathbb{E}_{\pi_{\rho,\varepsilon}}[c(X, Y, \varepsilon)] - \mathbb{E}_{\ell_{\rho,\varepsilon}}[c(X, Y, \varepsilon)].$$

To prove (29), from (16) and (19)

$$\frac{d\ell_{\rho,\varepsilon}}{d\pi_{\rho,\varepsilon}} = \frac{1}{\sqrt{\rho(x)\rho(y)}} \exp\left(-\frac{1}{\varepsilon}f_\varepsilon(x) - \frac{1}{\varepsilon}f_\varepsilon(y) - c(x, y, \varepsilon)\right).$$

As both $\pi_{\rho,\varepsilon}, \ell_{\rho,\varepsilon} \in \Pi(\rho, \rho)$, we have that

$$\begin{aligned} H(\ell_{\rho,\varepsilon}|\pi_{\rho,\varepsilon}) &= \mathbb{E}_{\ell_{\rho,\varepsilon}} \left[\log \frac{d\ell_{\rho,\varepsilon}}{d\pi_{\rho,\varepsilon}} \right] = -\frac{2}{\varepsilon} \mathbb{E}_\rho[f_\varepsilon(X)] - \text{Ent}(\rho) - \mathbb{E}_{\ell_{\rho,\varepsilon}}[c(X, Y, \varepsilon)], \\ H(\pi_{\rho,\varepsilon}|\ell_{\rho,\varepsilon}) &= \mathbb{E}_{\pi_{\rho,\varepsilon}} \left[-\log \frac{d\ell_{\rho,\varepsilon}}{d\pi_{\rho,\varepsilon}} \right] = \frac{2}{\varepsilon} \mathbb{E}_\rho[f_\varepsilon(X)] + \text{Ent}(\rho) + \mathbb{E}_{\pi_{\rho,\varepsilon}}[c(X, Y, \varepsilon)]. \end{aligned}$$

Equation (29) follows by adding these expressions together.

Step 2. Recall R_ε as used in (21) and let

$$(30) \quad R(x, y, \varepsilon) := -\log \left(\mathbb{E}_{R_\varepsilon} \left[\exp \left(-\int_0^1 \varepsilon (\mathcal{U}(\omega_s) - \mathcal{U}(\omega_0)) ds \middle| \omega_0 = x, \omega_1 = y \right) \right] \right).$$

Then we claim that

$$(31) \quad H(\ell_{\rho,\varepsilon}|\pi_{\rho,\varepsilon}) + H(\pi_{\rho,\varepsilon}|\ell_{\rho,\varepsilon}) \leq \mathbb{E}_{\pi_{\rho,\varepsilon}}[R(X, Y, \varepsilon)].$$

Recall that W_x is Wiener measure starting from x , and

$$\begin{aligned} c(x, y, \varepsilon) &= -\log \left(\mathbb{E}_{W_x} \left[\exp \left(-\int_0^\varepsilon \mathcal{U}(\omega_s) ds \right) \middle| \omega_\varepsilon = y \right] \right) \\ &= \varepsilon \mathcal{U}(x) - \log \left(\mathbb{E}_{W_x} \left[\exp \left(-\int_0^\varepsilon (\mathcal{U}(\omega_s) - \mathcal{U}(\omega_0)) ds \right) \middle| \omega_\varepsilon = y \right] \right). \end{aligned}$$

For the integral inside the $\exp(\cdot)$, we rescale s from $[0, \varepsilon]$ to $[0, 1]$. By Brownian scaling the law of $(\omega_{\varepsilon s}, s \geq 0)$ under W_x is equal to the law of $(\omega_s, s \geq 0)$ under R_ε with initial condition x , which gives

$$c(x, y, \varepsilon) = \varepsilon \mathcal{U}(x) + R(x, y, \varepsilon).$$

As $\pi_{\rho,\varepsilon} \in \Pi(\rho, \rho)$, thanks to an integration by parts,

$$\begin{aligned} \mathbb{E}_{\pi_{\rho,\varepsilon}}[\mathcal{U}(X)] &= \int_{\mathbb{R}^d} \mathcal{U}(x)\rho(x)dx = \frac{1}{8}I(\rho) - \int_{\mathbb{R}^d} \frac{1}{4}\Delta g(x)e^{-g(x)}dx \\ &= \frac{1}{8}I(\rho) - \int_{\mathbb{R}^d} \frac{1}{4}\|\nabla g(x)\|^2 e^{-g(x)}dx = -\frac{1}{8}I(\rho). \end{aligned}$$

Thus from (29) observe

$$\mathbb{E}_{\pi_{\rho,\varepsilon}}[c(X, Y, \varepsilon)] - \mathbb{E}_{\ell_{\rho,\varepsilon}}[c(X, Y, \varepsilon)] \leq -\frac{\varepsilon}{8}I(\rho) + \mathbb{E}_{\pi_{\rho,\varepsilon}}[R(X, Y, \varepsilon)] - \mathbb{E}_{\ell_{\rho,\varepsilon}}[c(X, Y, \varepsilon)].$$

Thus, to prove (31) it will suffice to show

$$(32) \quad \mathbb{E}_{\ell_{\rho,\varepsilon}}[-c(X, Y, \varepsilon)] \leq \frac{\varepsilon}{8}I(\rho).$$

Observe that

$$\frac{d\ell_{\rho,\varepsilon}}{d\mu_{\rho,\varepsilon}} = \frac{\sqrt{\rho(y)}}{\sqrt{\rho(x)}} \exp(-c(x, y, \varepsilon)).$$

Thus, as $\ell_{\rho,\varepsilon} \in \Pi(\rho, \rho)$ and $-\infty < \text{Ent}(\rho) < +\infty$,

$$(33) \quad H(\ell_{\rho,\varepsilon} | \mu_{\rho,\varepsilon}) = \mathbb{E}_{\ell_{\rho,\varepsilon}} \left[\log \frac{d\ell_{\rho,\varepsilon}}{d\mu_{\rho,\varepsilon}} \right] = \mathbb{E}_{\ell_{\rho,\varepsilon}} [-c(X, Y, \varepsilon)].$$

On $C^d[0, \varepsilon]$ let W_ρ denote the law of Brownian motion with initial distribution ρ . By a well-known formula (see [DGW04, Equation (5.1)] for instance)

$$H(Q_x | W_x) = \frac{1}{2} \mathbb{E}_{Q_x} \left[\int_0^\varepsilon \left\| -\frac{1}{2} \nabla g(\omega_s) \right\|^2 ds \right] = \frac{1}{8} \int_0^\varepsilon \mathbb{E}_{Q_x} [\|\nabla g(\omega_s)\|^2] ds.$$

It then follows from the factorization of relative entropy that

$$\begin{aligned} H(Q | W_\rho) &= \int H(Q_x | W_x) \rho(x) dx = \frac{1}{8} \int \left(\int_0^\varepsilon \mathbb{E}_{Q_x} [\|\nabla g(\omega_s)\|^2] ds \right) \rho(x) dx \\ &= \frac{1}{8} \int_0^\varepsilon \mathbb{E}_Q [\|\nabla g(\omega_s)\|^2] ds = \frac{\varepsilon}{8} I(\rho). \end{aligned}$$

As $\ell_{\rho,\varepsilon} = (\pi_0, \pi_\varepsilon) \# Q$, (32) follows by (33) and the information processing inequality (see [AGS08, Lemma 9.4.5] for instance).

Step 3. Show $\mathbb{E}_{\pi_{\rho,\varepsilon}} [R(X, Y, \varepsilon)]$ is bounded above by the RHS in (27).

To start, apply Jensen's inequality to interchange the $-\log$ and $\mathbb{E}_{R_\varepsilon}[\cdot | \omega_0 = x, \omega_1 = y]$ in the expression for $R(x, y, \varepsilon)$ in (30) to obtain

$$R(x, y, \varepsilon) \leq \varepsilon \mathbb{E}_{R_\varepsilon} \left[\int_0^1 (\mathcal{U}(\omega_s) - \mathcal{U}(\omega_0)) ds \middle| \omega_0 = x, \omega_1 = y \right].$$

Since the dynamic Schrödinger bridge, $S_{\rho,\varepsilon}$ defined in (21), disintegrates as the Brownian bridge once its endpoints are fixed [Léo14, Proposition 2.3], we have that

$$(34) \quad \mathbb{E}_{\pi_{\rho,\varepsilon}} [R(X, Y, \varepsilon)] \leq \varepsilon \mathbb{E}_{\pi_{\rho,\varepsilon}} \left[\mathbb{E}_{R_\varepsilon} \left[\int_0^1 (\mathcal{U}(\omega_s) - \mathcal{U}(\omega_0)) ds \middle| \omega_0 = X, \omega_1 = Y \right] \right]$$

$$(35) \quad = \varepsilon \int_0^1 \mathbb{E}_{S_{\rho,\varepsilon}} [\mathcal{U}(\omega_s) - \mathcal{U}(\omega_0)] ds,$$

where the last equality follows from Fubini's Theorem.

Recall that $(\rho_t^\varepsilon, v_t^\varepsilon)_{t \in [0,1]}$ is a weak solution for the continuity equation (9) if for all $\psi \in C_c^1(\mathbb{R}^d)$ and $t \in [0, 1]$

$$\frac{d}{dt} \mathbb{E}_{\rho_t^\varepsilon} [\psi] = \mathbb{E}_{\rho_t^\varepsilon} [\langle v_t^\varepsilon, \nabla \psi \rangle].$$

Integrating in time over $[0, s]$ for $s \in (0, 1]$ gives

$$\mathbb{E}_{\rho_s^\varepsilon} [\psi] - \mathbb{E}_{\rho_0^\varepsilon} [\psi] = \int_0^s \mathbb{E}_{\rho_u^\varepsilon} [\langle v_u^\varepsilon, \nabla \psi \rangle] du.$$

We now use a density argument to extend the above identity to

$$(36) \quad \mathbb{E}_{S_{\rho,\varepsilon}} [\mathcal{U}(\omega_s) - \mathcal{U}(\omega_0)] = \mathbb{E}_{\rho_s^\varepsilon} [\mathcal{U}] - \mathbb{E}_{\rho_0^\varepsilon} [\mathcal{U}] = \int_0^s \mathbb{E}_{\rho_u^\varepsilon} [\langle v_u^\varepsilon, \nabla \mathcal{U} \rangle] du.$$

For $R > 0$ let $\chi_R : \mathbb{R}^d \rightarrow \mathbb{R}$ be such that $\chi_R \in C_c^\infty(\mathbb{R}^d)$, $0 \leq \chi_R \leq 1$, $\chi_R|_{B(0,R)} \equiv 1$, and $\|\nabla \chi_R\|_\infty \leq 2$. As $\chi_R \mathcal{U} \in C_c^1(\mathbb{R}^d)$,

$$\begin{aligned} \mathbb{E}_{S_{\rho,\varepsilon}} [(\chi_R \mathcal{U})(\omega_s) - (\chi_R \mathcal{U})(\omega_0)] &= \int_0^s \mathbb{E}_{S_{\rho,\varepsilon}} [\langle v_u^\varepsilon(\omega_u), \nabla (\chi_R \mathcal{U})(\omega_u) \rangle] du \\ &= \int_0^s \mathbb{E}_{S_{\rho,\varepsilon}} [\langle v_u^\varepsilon, \nabla \mathcal{U} \rangle \chi_R] ds + \int_0^s \mathbb{E}_{S_{\rho,\varepsilon}} [\langle v_u^\varepsilon, \nabla \chi_R \rangle \mathcal{U}] ds. \end{aligned}$$

By Lemma 2, $\int_0^1 \mathbb{E}_{S_{\rho, \varepsilon}} [\|\nabla \mathcal{U}(\omega_s)\|^2] ds < +\infty$. Moreover, it is shown in (38) that $\int_0^1 \mathbb{E}_{S_{\rho, \varepsilon}} [\|v_s^\varepsilon(\omega_s)\|^2] ds < +\infty$. Thus, by the Dominated Convergence Theorem

$$\lim_{R \uparrow +\infty} \int_0^s \mathbb{E}_{S_{\rho, \varepsilon}} [\langle v_u^\varepsilon, \nabla \mathcal{U} \rangle \chi_R] ds = \int_0^s \mathbb{E}_{S_{\rho, \varepsilon}} [\langle v_u^\varepsilon, \nabla \mathcal{U} \rangle] ds.$$

As $|\chi_R \mathcal{U}(\omega_s) + \chi_R \mathcal{U}(\omega_0)| \leq |\mathcal{U}(\omega_s)| + |\mathcal{U}(\omega_0)|$ and

$$|\langle v_u^\varepsilon, \nabla \chi_R \rangle \mathcal{U}| \leq 2 \|v_u^\varepsilon\| |\mathcal{U}| \leq \|v_u^\varepsilon(\omega_u)\|^2 + |\mathcal{U}(\omega_u)|^2$$

and both upper bounds are integrable in their suitable spaces (again by Lemma 2 and (38)), (36) follows from dominated convergence by sending $R \uparrow +\infty$.

Integrating (36) once more in s over $[0, 1]$, multiplying by ε , and plugging in (25) gives the RHS of (34) as

$$\varepsilon \int_0^1 \mathbb{E}_{S_{\rho, \varepsilon}} [\mathcal{U}(\omega_s) - \mathcal{U}(\omega_0)] ds = \varepsilon^2 \int_0^1 \int_0^s \mathbb{E}_{\rho_u^\varepsilon} \left[\left\langle \frac{1}{\varepsilon} v_u^\varepsilon, \nabla \mathcal{U} \right\rangle \right] du ds.$$

Apply Cauchy-Schwarz on $L^2(C^d[0, 1] \times [0, 1], dS_{\rho, \varepsilon} \otimes du)$ to obtain for each $s \in [0, 1]$

$$\begin{aligned} \left| - \int_0^s \mathbb{E}_{S_{\rho, \varepsilon}} \left[\left\langle \frac{1}{\varepsilon} v_u^\varepsilon(\omega_u), \nabla \mathcal{U}(\omega_u) \right\rangle \right] du \right| &\leq \int_0^1 \mathbb{E}_{S_{\rho, \varepsilon}} \left[\frac{1}{\varepsilon} \|v_u^\varepsilon(\omega_u)\| \|\nabla \mathcal{U}(\omega_u)\| \right] du \\ &\leq \left(\int_0^1 \frac{1}{\varepsilon^2} \|v_u^\varepsilon\|_{L^2(\rho_u^\varepsilon)}^2 du \right)^{1/2} \left(\int_0^1 \mathbb{E}_{\rho_u^\varepsilon} [\|\nabla \mathcal{U}\|^2] du \right)^{1/2} \end{aligned}$$

Thus, altogether we have that

$$(37) \quad \mathbb{E}_{\pi_{\rho, \varepsilon}} [R(X, Y, \varepsilon)] \leq \varepsilon^2 \left(\int_0^1 \frac{1}{\varepsilon^2} \|v_t^\varepsilon\|_{L^2(\rho_t^\varepsilon)}^2 dt \right)^{1/2} \left(\int_0^1 \mathbb{E}_{\rho_t^\varepsilon} [\|\nabla \mathcal{U}\|^2] dt \right)^{1/2}.$$

Recall the entropic Benamou Brenier expression for entropic cost given in (26). By the optimality of (25) and suboptimality of the constant curve $\rho_t = \rho$ for all $t \in [0, 1]$,

$$H(\pi_{\rho, \varepsilon} | \mu_{\rho, \varepsilon}) = \frac{1}{2\varepsilon} \int_0^1 \|v_t^\varepsilon\|_{L^2(\rho_t^\varepsilon)}^2 dt + \frac{\varepsilon}{8} \int_0^1 I(\rho_t^\varepsilon) dt \leq \frac{\varepsilon}{8} I(\rho).$$

Rearranging terms gives

$$(38) \quad \frac{1}{\varepsilon^2} \int_0^1 \|v_t^\varepsilon\|_{L^2(\rho_t^\varepsilon)}^2 dt \leq \frac{1}{4} \left(I(\rho) - \int_0^1 I(\rho_t^\varepsilon) dt \right).$$

With this inequality, (37) becomes the desired

$$\mathbb{E}_{\pi_{\rho, \varepsilon}} [R(X, Y, \varepsilon)] \leq \frac{1}{2} \varepsilon^2 \left(I(\rho) - \int_0^1 I(\rho_t^\varepsilon) dt \right)^{1/2} \left(\int_0^1 \mathbb{E}_{\rho_t^\varepsilon} [\|\nabla \mathcal{U}\|^2] dt \right)^{1/2}.$$

Under Assumption 1 (2), Lemma 2 applies and gives that the rightmost constant has an upper bound once we consider $\varepsilon \in (0, \varepsilon_0)$ for some $\varepsilon_0 > 0$. Thus, the nonnegativity of the LHS of (27) and Lemma 1 give the stated convergence as $\varepsilon \downarrow 0$.

□

2.3. Discussion. Integrated Fisher Information. Under additional assumptions the integrated Fisher information along the entropic interpolation in (27) can be replaced with a potentially more manageable quantity. In certain settings (for instance, when $\rho \in C_c^\infty(\mathbb{R}^d)$ with finite entropy and Fisher information [GLRT20, Lemma 3.2]), there is a conserved quantity along the entropic interpolation called the energy, which we denote $\mathcal{E}_\varepsilon(\rho)$. With $(v_t^\varepsilon, \rho_t^\varepsilon)_{t \in [0,1]}$ as defined in (25), for any $t \in [0, 1]$

$$(39) \quad \mathcal{E}_\varepsilon(\rho) := \frac{1}{2\varepsilon^2} \|v_t^\varepsilon\|_{L^2(\rho_t^\varepsilon)}^2 - \frac{1}{8} I(\rho_t^\varepsilon).$$

Now, evaluate (39) at $t = 1/2$. Then $v_{1/2}^\varepsilon = 0$ and $\mathcal{E}_\varepsilon(\rho) = -\frac{1}{8} I(\rho_{1/2}^\varepsilon)$. On the other hand, integrate (39) over $t \in [0, 1]$, this also gives $\mathcal{E}_\varepsilon(\rho)$ and thus

$$-\frac{1}{8} I(\rho_{1/2}^\varepsilon) = \frac{1}{2\varepsilon^2} \int_0^1 \|v_t^\varepsilon\|_{L^2(\rho_t^\varepsilon)}^2 dt - \frac{1}{8} \int_0^1 I(\rho_t^\varepsilon) dt.$$

This in turn implies

$$\frac{1}{\varepsilon^2} \int_0^1 \|v_t^\varepsilon\|_{L^2(\rho_t^\varepsilon)}^2 dt = \frac{1}{4} \left(\int_0^1 I(\rho_t^\varepsilon) dt - I(\rho_{1/2}^\varepsilon) \right) \leq \frac{1}{4} \left(I(\rho) - I(\rho_{1/2}^\varepsilon) \right).$$

Thus, assuming the conservation of energy holds under Assumption 1, replacing the bound in (38) with the RHS of the above inequality rewrites Theorem 1 as

$$(40) \quad H(\ell_{\rho,\varepsilon}|\pi_{\rho,\varepsilon}) + H(\pi_{\rho,\varepsilon}|\ell_{\rho,\varepsilon}) \leq \frac{1}{2} \varepsilon^2 \left(I(\rho) - I(\rho_{1/2}^\varepsilon) \right)^{1/2} \left(\int_0^1 \mathbb{E}_{\rho_t^\varepsilon} \|\nabla \mathcal{U}\|^2 dt \right)^{1/2}.$$

Alternatively, (40) also holds when ρ is a univariate Gaussian, as explicit computations of the entropic interpolation in this case from [GLR17] demonstrate that $I(\rho_t^\varepsilon)$ is minimized at $t = 1/2$.

Schrödinger Cost Expansion. In the course of proving Theorem 1, we recover a result in alignment with the expansion of Schrödinger cost about $\varepsilon = 0$ developed in [CT21, Theorem 1.6]. Under assumptions on bounded ρ with compact support and finite entropy, [CT21, Theorem 1.6] states

$$(41) \quad H(\pi_{\rho,\varepsilon}|\mu_{\rho,\varepsilon}) = \frac{\varepsilon}{8} I(\rho) + o(\varepsilon).$$

Now, from the Pythagorean Theorem of relative entropy [Csi75, Theorem 2.2]

$$H(\ell_{\rho,\varepsilon}|\mu_{\rho,\varepsilon}) \geq H(\ell_{\rho,\varepsilon}|\pi_{\rho,\varepsilon}) + H(\pi_{\rho,\varepsilon}|\mu_{\rho,\varepsilon}).$$

Recall from Step 2 of Lemma 1 that $H(\ell_{\rho,\varepsilon}|\mu_{\rho,\varepsilon}) \leq \frac{\varepsilon}{8} I(\rho)$, so by Theorem 1 we obtain the following expansion of an upper bound of Schrödinger cost

$$(42) \quad H(\pi_{\rho,\varepsilon}|\mu_{\rho,\varepsilon}) \leq H(\ell_{\rho,\varepsilon}|\mu_{\rho,\varepsilon}) - H(\ell_{\rho,\varepsilon}|\pi_{\rho,\varepsilon}) = \frac{\varepsilon}{8} I(\rho) - o(\varepsilon^2).$$

Thus, in the same-marginal case (and under different regularity assumptions) the second order term in the Schrödinger expansion is zero.

Convergence of Potentials. Revisiting the inequality in (38) and recalling (25), observe that

$$\int_0^1 \|\nabla \psi^\varepsilon(t, \cdot) - \nabla \psi^\varepsilon(1-t, \cdot)\|_{L^2(\rho_t^\varepsilon)}^2 dt \leq I(\rho) - \int_0^1 I(\rho_t^\varepsilon) dt.$$

By Lemma 1, for Lebesgue a.e. $t \in [0, 1]$ it holds that

$$(43) \quad \lim_{\varepsilon \downarrow 0} \|\nabla \psi^\varepsilon(t, \cdot) - \nabla \psi^\varepsilon(1-t, \cdot)\|_{L^2(\rho_t^\varepsilon)}^2 = 0.$$

This fact hints in the direction of a result on the convergence of rescaled gradients of entropic potentials. Suppose that the convergence in (43) holds at $t = 1$ or $t = 0$ (of course, a priori there is no way to guarantee such convergence at a specific t). In this case,

$$\begin{aligned} \nabla\psi^\varepsilon(1, \cdot) - \nabla\psi(0, \cdot) &= \nabla\log a^\varepsilon - \nabla\log P_\varepsilon a^\varepsilon \\ &= \nabla\log a^\varepsilon - \nabla\log \frac{\rho}{a^\varepsilon} = \frac{2}{\varepsilon}\nabla f_\varepsilon + \nabla\log\rho. \end{aligned}$$

As $\rho_0^\varepsilon = \rho_1^\varepsilon = \rho$, (43) would imply that $\lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon}\nabla f_\varepsilon = -\frac{1}{2}\nabla\log\rho$ in $L^2(\rho)$. This would be in accordance with [SABP22, Theorem 1], but now with fully supported densities.

3. ONE STEP ANALYSIS OF SCHRÖDINGER BRIDGE SCHEME

Fix $\rho \in \mathcal{P}_2^{ac}(\mathbb{R}^d)$ and recall the tangent space at ρ , denoted by $\text{Tan}(\rho)$, and defined in [AGS08, Definition 8.4.1] by

$$\text{Tan}(\rho) := \overline{\{\nabla\varphi : \varphi \in C_c^\infty(\mathbb{R}^d)\}}^{L^2(\rho)}.$$

Now, consider a vector field $v = \nabla u \in \text{Tan}(\rho)$, such that the surrogate measure (12) is defined. Our objective in this section is to lay down conditions under which the approximation stated below (SB) holds. More precisely, we show that

$$(44) \quad \lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \mathbb{W}_2(\text{SB}_\varepsilon(\rho), (\text{Id} + \varepsilon v)_\# \rho) = 0.$$

In fact, we will prove the above theorem in a somewhat more general form when the tangent vector field may not even be of a gradient form. The general setting is as follows. Consider a collection of vector fields of gradient type ($v_\varepsilon = \nabla\psi_\varepsilon, \varepsilon > 0$), we wish to tightly approximate the pushforwards $(\text{Id} + \varepsilon v_\varepsilon)_\# \rho$ with $\text{SB}_\varepsilon(\rho)$. As a shorthand, we write

$$(45) \quad S_\varepsilon^1(\rho) := (\text{Id} + \varepsilon v_\varepsilon)_\# \rho.$$

Suppose that for each $\varepsilon > 0$ there exists $\theta_\varepsilon \in \mathbb{R} \setminus \{0\}$ such that the surrogate measure defined in (12) exists. Set

$$(46) \quad \sigma_\varepsilon(x) = \exp(2\theta_\varepsilon\psi_\varepsilon(x) - \Lambda_0(\theta_\varepsilon)), \quad \text{where} \quad \Lambda_0(\theta_\varepsilon) := \log \int e^{2\theta_\varepsilon\psi_\varepsilon(y)} dy.$$

Redefine one step in the SB scheme as

$$(SB) \quad \text{SB}_\varepsilon(\rho) = \left(\left(1 - \theta_\varepsilon^{-1} \right) \text{Id} + \theta_\varepsilon^{-1} \mathcal{B}_{\sigma_\varepsilon, \varepsilon} \right)_\# \rho,$$

where $\mathcal{B}_{\sigma_\varepsilon, \varepsilon}$ is the barycentric projection defined in (4).

In Theorem 2 we show that, under appropriate assumptions,

$$(47) \quad \lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \mathbb{W}_2(\text{SB}_\varepsilon(\rho), S_\varepsilon^1(\rho)) = 0.$$

There are multiple reasons why one might be interested in this general setup. Firstly, we know that, for any $v \in \text{Tan}(\rho)$, there exists a sequence of tangent elements of the type $v_\varepsilon = \nabla\psi_\varepsilon$ such that $\lim_{\varepsilon \rightarrow 0+} v_\varepsilon = v$ in $\mathbf{L}^2(\rho)$. In particular, by an obvious coupling,

$$(48) \quad \limsup_{\varepsilon \rightarrow 0+} \frac{1}{\varepsilon} \mathbb{W}_2((\text{Id} + \varepsilon v_\varepsilon)_\# \rho, (\text{Id} + \varepsilon v)_\# \rho) \leq \lim_{\varepsilon \rightarrow 0+} \|v_\varepsilon - v\|_{\mathbf{L}^2(\rho)} = 0.$$

But, of course, if v is not of the gradient form, one cannot construct a surrogate measure as in (12) to run the Schrödinger bridge scheme. A natural remedy is to show that (47) holds under appropriate assumptions. Then, combined with (48) we recover (44) even when v is not a gradient.

There is one more reason why one might use the above generalized scheme. As explained in (11), we would like to iterate the Schrödinger bridge scheme to track the explicit Euler iterations of a Wasserstein gradient flow. However, the more natural discretization scheme of Wasserstein gradient flows is the implicit Euler or the JKO scheme. See (IE) in Section 4 for the details. In this case, however, the vector field v_ε , which is the Kantorovich potential between two successive steps of the JKO scheme, changes with ε . This necessitates our generalization. Hence, we continue with (46) and the generalized (SB) scheme.

The proof of (47) relies on the close approximation of the Langevin diffusion to the Schrödinger bridge provided in Theorem 1. To harness this approximation, we introduce an intermediary scheme based on the Langevin diffusion, which we denote $\text{LD}_\varepsilon(\cdot)$ and define in the following manner. Let $\varepsilon > 0$ and consider the symmetric Langevin diffusion $(X_t, 0 \leq t \leq \varepsilon)$ with stationary density σ_ε as defined in (46). The drift function for this diffusion is $x \mapsto \frac{1}{2}\nabla \log \sigma_\varepsilon(x) = \theta_\varepsilon \nabla \psi_\varepsilon(x) = \theta_\varepsilon v_\varepsilon(x)$. In analogy with the definition of $\mathcal{B}_{\sigma_\varepsilon, \varepsilon}(\cdot)$ in (4), define $\mathcal{L}_{\sigma_\varepsilon, \varepsilon}(\cdot)$ by

$$(49) \quad \mathcal{L}_{\sigma_\varepsilon, \varepsilon}(x) = \text{E}_{\ell_{\sigma_\varepsilon, \varepsilon}}[Y|X = x].$$

Next, in analogy with the definition of $\text{SB}_\varepsilon(\cdot)$ in (SB), define the Langevin diffusion update

$$(LD) \quad \text{LD}_\varepsilon(\rho) = \left(\left(1 - \theta_\varepsilon^{-1}\right) \text{Id} + \theta_\varepsilon^{-1} \mathcal{L}_{\sigma_\varepsilon, \varepsilon} \right) \# \rho.$$

In analogy with Section 2, for $\varepsilon > 0$ set

$$\mathcal{U}_\varepsilon := \frac{1}{8} \|\nabla \log \sigma_\varepsilon\|^2 + \frac{1}{4} \Delta \log \sigma_\varepsilon.$$

We now make the following assumptions on the collection of densities $(\sigma_\varepsilon, \varepsilon > 0)$ defined in (46).

Assumption 2. For each $\varepsilon > 0$, σ_ε as defined in (46) satisfies Assumption 1. Additionally

- (1) There exists $\sigma_0 \in \mathcal{P}_2^{ac}(\mathbb{R}^d)$ such that $(\sigma_\varepsilon, \varepsilon > 0)$ converges weakly to σ_0 and $I(\sigma_\varepsilon) \rightarrow I(\sigma_0)$ as $\varepsilon \downarrow 0$. Moreover, Assumption 1 holds uniformly in ε in the following manner: there exists $\varepsilon_0 > 0$ and $C, D > 0$, $N \geq 1$ such that for all $\varepsilon \in (0, \varepsilon_0)$, $\|\nabla \mathcal{U}_\varepsilon(x)\|^2 \leq C(1 + \|x\|^N)$ and for $X_\varepsilon \sim \sigma_\varepsilon$, $\|X_\varepsilon\|_{\psi_1} \leq D$.
- (2) The $(\sigma_\varepsilon, \varepsilon \in (0, \varepsilon_0))$ are uniformly semi log-concave, that is, there exists $\lambda \in \mathbb{R}$ such that for all $\varepsilon \in (0, \varepsilon_0)$, $-\nabla^2 \log \sigma_\varepsilon \geq \lambda \text{Id}$. Additionally, $L := \limsup_{\varepsilon \downarrow 0} \text{E}_{\sigma_\varepsilon} \|\nabla^2 \log \sigma_\varepsilon\|_{HS}^2 < +\infty$, where $\|\cdot\|_{HS}$ is the Hilbert-Schmidt norm.
- (3) $\chi := \limsup_{\varepsilon \downarrow 0} \|\rho/\sigma_\varepsilon\|_\infty < +\infty$ and $\theta := \liminf_{\varepsilon \downarrow 0} |\theta_\varepsilon| > 0$.

Remark 1. In all our examples, θ_ε is either always +1 or always -1. Hence, this parameter simply needs to have a positive lower bound (in absolute value) and it does not scale with ε .

While these assumptions may appear stringent, they are indeed satisfied by a wide class of examples, see Section 5. In many of these examples, σ_ε takes the form $\sigma_\varepsilon = \rho e^{V_\varepsilon}$. Thus, condition (3) in Assumption 2 is satisfied if $\liminf_{\varepsilon \downarrow 0} \inf_{x \in \mathbb{R}^d} V_\varepsilon > -\infty$. For instance, when using (SB) to approximate (EE) for the gradient flow of entropy, $\sigma_\varepsilon = \rho$ for all $\varepsilon > 0$ and thus $V_\varepsilon \equiv 0$. Similarly, for approximating the gradient flow of $H(\cdot|N(0, 1))$ from certain starting measures, $V_\varepsilon(x) = \frac{1}{2}x^2 + C$, where C is a normalizing constant, for all $\varepsilon > 0$. See Section 5 for more detailed discussion.

We now provide a quantitative statement about the convergence stated in (47), where we take care to note the dependence on constants appearing in Assumption 2 as this upper bound

must hold uniformly along iterations of this scheme in Section 4. Recall $S_\varepsilon^1(\cdot)$ as defined in (45) and $SB_\varepsilon(\cdot)$ as defined in (SB).

Theorem 2. *Under Assumption 2, there exists a constant $K > 0$ depending on $C, D, N, \lambda, L, \chi, \theta$, such that for all $\varepsilon \in (0, \varepsilon_0)$*

$$\mathbb{W}_2(SB_\varepsilon(\rho), S_\varepsilon^1(\rho)) \leq K\varepsilon \left[\left(I(\sigma_\varepsilon) - \int_0^1 I((\sigma_\varepsilon)_s^\varepsilon) ds \right)^{1/4} + \sqrt{\varepsilon} \right].$$

In particular,

$$\lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \mathbb{W}_2(SB_\varepsilon(\rho), S_\varepsilon^1(\rho)) = 0.$$

Hence, from the discussion following (48) we immediately obtain:

Corollary 1. *Let $v \in \text{Tan}(\rho)$, and let $(\nabla\psi_\varepsilon, \varepsilon > 0) \subset C^\infty(\mathbb{R}^d) \cap \text{Tan}(\rho)$ be such that $\lim_{\varepsilon \rightarrow 0^+} \|\nabla\psi_\varepsilon - v\|_{\mathbf{L}^2(\rho)} = 0$ and the corresponding surrogate measures $(\sigma_\varepsilon > 0)$ satisfy Assumption 2. Then*

$$\lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \mathbb{W}_2((\text{Id} + \varepsilon v)_\# \rho, SB_\varepsilon(\rho)) = 0.$$

Proof of Theorem 2. This result follows from Lemmas 3, 4, and 5 below. Lemmas 3 and 4 together prove an upper bound on $\mathbb{W}_2(SB_\varepsilon(\rho), LD_\varepsilon(\rho))$ that show, in particular,

$$\lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \mathbb{W}_2(SB_\varepsilon(\rho), LD_\varepsilon(\rho)) = 0,$$

and Lemma 5 proves an upper bound on $\mathbb{W}_2(LD_\varepsilon(\rho), S_\varepsilon^1(\rho))$ which implies

$$\lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \mathbb{W}_2(LD_\varepsilon(\rho), S_\varepsilon^1(\rho)) = 0.$$

That is, the three updates $SB_\varepsilon(\rho)$, $LD_\varepsilon(\rho)$, and $S_\varepsilon^1(\rho)$ are within $o(\varepsilon)$ of each other in the Wasserstein-2 metric. Theorem 2 now follows from the triangle inequality and simplifying the choice of the constant K . \square

We begin by converting the relative entropy decay established in Theorem 1 between the Schrödinger bridge and Langevin diffusion into a decay rate between their respective barycentric projections in the Wasserstein-2 metric. The essential tool for this conversion is a Talagrand inequality from [CG14, Theorem 1.8(3)].

Lemma 3. *Let $\varepsilon > 0$. Under Assumption 2, there exists a constant $\alpha > 0$ depending on ε and λ such that*

$$\begin{aligned} \mathbb{W}_2^2((\mathcal{B}_{\sigma_\varepsilon, \varepsilon})_\# \rho, (\mathcal{L}_{\sigma_\varepsilon, \varepsilon})_\# \rho) &\leq \alpha \|\rho/\sigma_\varepsilon\|_\infty H(\pi_{\sigma_\varepsilon, \varepsilon} | \ell_{\sigma_\varepsilon, \varepsilon}) \\ \mathbb{W}_2^2(SB_\varepsilon(\rho), LD_\varepsilon(\rho)) &\leq \alpha \theta_\varepsilon^{-2} \|\rho/\sigma_\varepsilon\|_\infty H(\pi_{\sigma_\varepsilon, \varepsilon} | \ell_{\sigma_\varepsilon, \varepsilon}). \end{aligned}$$

Moreover, with α can be chosen to be increasing in ε .

Recall that $\theta_\varepsilon \in \mathbb{R} \setminus \{0\}$ is a constant chosen for integrability reasons outlined in (46)– it is simply an $O(1)$ term as $\varepsilon \downarrow 0$. See Remark 1.

Proof of Lemma 3. Fix $T > 0$ and $\varepsilon \in (0, T)$. To avoid unnecessary subscripts, set $\sigma := \sigma_\varepsilon$. To start, we produce a coupling of $\ell_{\sigma, \varepsilon}$ and $\pi_{\sigma, \varepsilon}$ by applying a gluing argument (as in [Vil03, Lemma 7.6], for example) to the optimal couplings between corresponding transition densities for the Langevin diffusion and ε -static Schrödinger bridge. From this coupling, we obtain a coupling of the desired barycentric projections of $\ell_{\sigma, \varepsilon}$ and $\pi_{\sigma, \varepsilon}$. We then use the chain rule of relative entropy alongside the Talagrand inequality stated in [CG14, Theorem 1.8(3)] to obtain the inequality stated in the Lemma.

Let $(q_t(\cdot, \cdot), t \geq 0)$ denote the transition densities of the stationary Langevin diffusion associated to σ . By [CG14, Theorem 1.8(3)], there a constant $\alpha > 0$ depending on λ and T such that following Talagrand inequality holds for each $x \in \mathbb{R}^d$

$$\mathbb{W}_2^2(\eta, q_\varepsilon(x, \cdot)) \leq \alpha H(\eta | q_\varepsilon(x, \cdot)) \text{ for all } \eta \in \mathcal{P}(\mathbb{R}^d).$$

Let $(p(x, \cdot), x \in \mathbb{R}^d)$ denote the conditional densities of the ε -static Schrödinger bridge with marginals σ . Construct a triplet of \mathbb{R}^d -valued random variables (X, Y, Z) in the following way. Let $X \sim \rho$, and let the conditional measures $(Y|X = x, Z|X = x)$ have the law of the quadratic cost optimal coupling between $p(x, \cdot)$ and $q_\varepsilon(x, \cdot)$. Denote the law of (X, Y, Z) on $\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d$ by ν . Observe that $(\mathbb{E}_\nu[Y|X], \mathbb{E}_\nu[Z|X])$ is a coupling of $(\mathcal{L}_{\sigma_\varepsilon, \varepsilon})_{\#} \rho$ and $(\mathcal{B}_{\sigma_\varepsilon, \varepsilon})_{\#} \rho$. Moreover, observe that $((1 - \theta_\varepsilon^{-1})X + \theta_\varepsilon^{-1} \mathbb{E}_\nu[Y|X], (1 - \theta_\varepsilon^{-1})X + \theta_\varepsilon^{-1} \mathbb{E}_\nu[Z|X])$ is a coupling of $\text{SB}_\varepsilon(\rho)$ and $\text{LD}_\varepsilon(\rho)$. Thus, to compute an upper bound of $\mathbb{W}_2^2(\text{SB}_\varepsilon(\rho), \text{LD}_\varepsilon(\rho))$, it is sufficient to compute $\mathbb{E} \|\mathbb{E}_\nu[Y|X] - \mathbb{E}_\nu[Z|X]\|^2$.

We first observe for each $x \in \mathbb{R}^d$ that by two applications of the (conditional) Jensen's inequality, the optimality of $(Y|X = x, Z|X = x)$, and the aforementioned Talagrand inequality

$$\|\mathbb{E}_\nu[Y|X = x] - \mathbb{E}_\nu[Z|X = x]\| \leq \mathbb{W}_2(q_\varepsilon(x, \cdot), p(x, \cdot)) \leq \sqrt{\alpha H(p(x, \cdot) | q_\varepsilon(x, \cdot))}.$$

Hence, using the chain rule of relative entropy and the fact that the Langevin diffusion and Schrödinger bridge both start from σ

$$\begin{aligned} \mathbb{W}_2^2((\mathcal{B}_{\sigma_\varepsilon, \varepsilon})_{\#} \rho, (\mathcal{L}_{\sigma_\varepsilon, \varepsilon})_{\#} \rho) &\leq \mathbb{E}_\nu \|\mathbb{E}_\nu[Y|X = x] - \mathbb{E}_\nu[Z|X = x]\|^2 \\ &\leq \int \alpha H(p(x, \cdot) | q_\varepsilon(x, \cdot)) \rho(x) dx = \int \alpha H(p(x, \cdot) | q_\varepsilon(x, \cdot)) \sigma(x) \frac{\rho(x)}{\sigma(x)} dx \\ &\leq \alpha \|\rho / \sigma_\varepsilon\|_\infty H(\pi_{\sigma_\varepsilon, \varepsilon} | \ell_{\sigma_\varepsilon, \varepsilon}). \end{aligned}$$

□

Now, we revisit the bound from Theorem 1 and apply it to Lemma 5.

Lemma 4. *Under Assumption 2, there exists a constant $K > 0$ depending on $N, C,$ and D such that for all $\varepsilon \in (0, \varepsilon_0)$*

$$(50) \quad H(\pi_{\sigma_\varepsilon, \varepsilon} | \ell_{\sigma_\varepsilon, \varepsilon}) + H(\ell_{\sigma_\varepsilon, \varepsilon} | \pi_{\sigma_\varepsilon, \varepsilon}) \leq \frac{\varepsilon^2}{2} K \left(I(\sigma_\varepsilon) - \int_0^1 I((\sigma_\varepsilon)_s^\varepsilon) ds \right)^{1/2}$$

In particular,

$$\lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon^2} (H(\pi_{\sigma_\varepsilon, \varepsilon} | \ell_{\sigma_\varepsilon, \varepsilon}) + H(\ell_{\sigma_\varepsilon, \varepsilon} | \pi_{\sigma_\varepsilon, \varepsilon})) = 0.$$

Proof. Let $\varepsilon \in (0, \varepsilon_0)$ and in analogy with the Section 2 let $((\sigma_\varepsilon)_s^\varepsilon, s \in [0, 1])$ denote the ε -entropic interpolation from σ_ε to itself and set

$$\mathcal{U}_\varepsilon = \frac{1}{8} \|\nabla \log \sigma_\varepsilon\|^2 + \frac{1}{4} \Delta \log \sigma_\varepsilon.$$

By the assumed uniform subexponential bound on the σ_ε and uniform polynomial growth of \mathcal{U}_ε , there exists a constant $K > 0$ as specified in the Lemma statement such that

$$(51) \quad \sup_{\varepsilon \in (0, \varepsilon_0), s \in [0, 1]} \mathbb{E}_{(\sigma_\varepsilon)_s^\varepsilon} [\|\nabla \mathcal{U}_\varepsilon\|^2] \leq K.$$

Theorem 1 and (51) then establish (50) when $\varepsilon \in (0, \varepsilon_0)$. It now remains to show that

$$(52) \quad \lim_{\varepsilon \downarrow 0} \left[I(\sigma_\varepsilon) - \int_0^1 I((\sigma_\varepsilon)_s^\varepsilon) ds \right] = 0.$$

First, we claim that $((\sigma_\varepsilon)_s^\varepsilon, \varepsilon > 0)$ converges weakly to σ_0 as $\varepsilon \downarrow 0$ for all $s \in [0, 1]$. As $(\sigma_\varepsilon, \varepsilon > 0)$ converges weakly to σ_0 , the collection $\{\sigma_0\} \cup (\sigma_\varepsilon, \varepsilon > 0)$ is tight. This implies that $(\pi_{\sigma_\varepsilon, \varepsilon}, \varepsilon > 0)$ is tight. Let π^* denote a weak subsequential limit— we will not denote the subsequence. It is clear that $\pi^* \in \Pi(\sigma_0, \sigma_0)$. In the terminology of [BGN22], the support of each $\pi_{\sigma_\varepsilon, \varepsilon}$ is (c, ε) -cyclically invariant, where $c(x, y) = \frac{1}{2}\|x - y\|^2$. As π^* is a weak subsequential limit of the $(\pi_{\sigma_\varepsilon, \varepsilon}, \varepsilon > 0)$, the exact same argument presented in [BGN22, Lemma 3.1, Lemma 3.2], modified only superficially to allow for the different marginal in each ε , establishes that π^* is c -cyclically monotone. That is, $\pi^* = \pi_{\sigma_0, 0}$ is the quadratic cost optimal transport plan from σ_0 to itself. Moreover, this argument shows that along any subsequence of ε decreasing to 0, there is a further subsequence converging to $\pi_{\sigma_0, 0}$. Hence, the limit holds without passing to a subsequence. Construct a sequence of random variables $((X_\varepsilon, Y_\varepsilon), \varepsilon \geq 0)$ with $\text{Law}(X_\varepsilon, Y_\varepsilon) = \pi_{\sigma_\varepsilon, \varepsilon}$, and let Z be a standard normal random variable independent to the collection. Fix $s \in [0, 1]$, and recall that $\text{Law}(sX_\varepsilon + (1-s)Y_\varepsilon + \sqrt{\varepsilon s(1-s)}Z) = (\sigma_\varepsilon)_s^\varepsilon$. The Continuous Mapping Theorem gives that $sX_\varepsilon + (1-s)Y_\varepsilon + \sqrt{\varepsilon s(1-s)}Z$ converges in distribution to $sX_0 + (1-s)Y_0$ as $\varepsilon \downarrow 0$. As $\text{Law}(sX_0 + (1-s)Y_0) = \sigma_0$, this establishes the weak convergence of $(\sigma_\varepsilon)_s^\varepsilon$ to σ_0 as $\varepsilon \downarrow 0$. By the lower semicontinuity of Fisher information with respect to weak convergence [Bob22, Proposition 14.2] and Fatou's Lemma

$$I(\sigma_0) \leq \int_0^1 \liminf_{\varepsilon \downarrow 0} I((\sigma_\varepsilon)_s^\varepsilon) ds \leq \liminf_{\varepsilon \downarrow 0} \int_0^1 I((\sigma_\varepsilon)_s^\varepsilon) ds.$$

Thus, (52) holds as Assumption 2(1) gives $\lim_{\varepsilon \downarrow 0} I(\sigma_\varepsilon) = I(\sigma_0)$. \square

As the last piece of the proof, we show in the following lemma below that $\text{LD}_\varepsilon(\rho)$ is an $o(\varepsilon)$ approximation of the first step of the Euler iteration $S_\varepsilon^1(\rho)$.

Lemma 5. *For $\rho = e^{-g}$ satisfying Assumption 2,*

$$\mathbb{W}_2^2(S_\varepsilon^1(\rho), \text{LD}_\varepsilon(\rho)) \leq 2\theta_\varepsilon^{-2} \|\rho/\sigma_\varepsilon\|_\infty \varepsilon^3 \mathbb{E}_{\sigma_\varepsilon} \|\nabla^2 \log \sigma_\varepsilon\|_{HS}^2.$$

Proof. The proof follows by producing an intuitive coupling of $\text{GA}_\varepsilon(\rho)$ and $\text{LD}_\varepsilon(\rho)$ and then applying a semigroup property stated in [BGL14, Equation (4.2.3)]. Let $(G_s^{\sigma_\varepsilon}, s \geq 0)$ be the semigroup associated to the Langevin diffusion corresponding to σ_ε , then for all $\varepsilon > 0, s \geq 0$, and $f : \mathbb{R}^d \rightarrow \mathbb{R}$ in the domain of the Dirichlet form

$$(53) \quad \|G_s^{\sigma_\varepsilon} f - f\|_{L^2(\sigma_\varepsilon)}^2 \leq 2s \|\nabla f\|_{L^2(\sigma_\varepsilon)}^2.$$

Let $X \sim \rho$, then by Dynkin's formula

$$\mathcal{L}_{\rho, \varepsilon}(X) = X + \int_0^\varepsilon G_s^{\sigma_\varepsilon} (\theta_\varepsilon \nabla \psi_\varepsilon)(X) ds$$

and thus

$$\begin{aligned} \text{LD}_\varepsilon(\rho) &= \text{Law} \left((1 - \theta_\varepsilon^{-1})X + \theta_\varepsilon^{-1} \left(X + \int_0^\varepsilon G_s^{\sigma_\varepsilon}(\theta_\varepsilon \nabla \psi_\varepsilon)(X) ds \right) \right) \\ &= \text{Law} \left(X + \int_0^\varepsilon G_s^{\sigma_\varepsilon}(\nabla \psi_\varepsilon)(X) ds \right). \end{aligned}$$

Also by definition

$$S_\varepsilon^1(\rho) = \text{Law} (X + \varepsilon \nabla \psi_\varepsilon(X)).$$

Thus, we have the upper bound

$$\begin{aligned} \mathbb{W}_2^2(\text{LD}_\varepsilon(\rho), S_\varepsilon^1(\rho)) &\leq \mathbb{E} \left\| \left(X + \int_0^\varepsilon G_s^{\sigma_\varepsilon}(\nabla \psi_\varepsilon)(X) ds \right) - (X + \varepsilon \nabla \psi_\varepsilon(X)) \right\|^2 \\ &= \mathbb{E} \left\| \int_0^\varepsilon (G_s^{\sigma_\varepsilon}(\nabla \psi_\varepsilon)(X) - \nabla \psi_\varepsilon(X)) ds \right\|^2 \\ &\leq \varepsilon \int_0^\varepsilon \mathbb{E} \|G_s^{\sigma_\varepsilon}(\nabla \psi_\varepsilon)(X) - \nabla \psi_\varepsilon(X)\|^2 ds \\ &= \varepsilon^2 \int_0^1 \|G_{\varepsilon s}^{\sigma_\varepsilon}(\nabla \psi_\varepsilon) - \nabla \psi_\varepsilon\|_{L^2(\rho)}^2 ds, \end{aligned}$$

where the penultimate inequality follows from Jensen's inequality. Then, perform a change of measure and apply (53) to obtain

$$\mathbb{W}_2^2(\text{LD}_\varepsilon(\rho), S_\varepsilon^1(\rho)) \leq 2\|\rho/\sigma_\varepsilon\|_\infty \varepsilon^3 \mathbb{E}_{\sigma_\varepsilon} \|\nabla^2 \psi_\varepsilon\|_{HS}^2.$$

□

4. ITERATED SCHRÖDINGER BRIDGE SCHEME

In this section we present conditions under which suitably interpolated iterations of (SB) converge to the gradient flow of entropy and relative entropy (i.e. solutions to the parabolic heat equation and Fokker-Planck equation). Our results rely on the close approximation of (SB) to explicit Euler iterations that we define later on. We also present a collection of sufficient conditions for which iterations of (SB) converge to the gradient flow of geodesically convex functionals. Roughly put, our assumptions are that (1) the one step approximation given in Theorem 2 holds along iterations of (SB) (which we call consistency) and (2) the error of this approximation does not accumulate too rapidly over a finite time horizon (which is implied by the condition we call contractivity in the limit). Although these conditions seem reasonable (in Section 5 we show examples that satisfy them), it is not immediate how generally they hold.

4.1. Preliminaries. We set up some notation for subdifferential calculus in Wasserstein space. Let $\mathcal{F} : \mathcal{P}_2(\mathbb{R}^d) \rightarrow (-\infty, \infty]$ be a function that is proper, lower-semicontinuous, and λ -geodesically convex functional for some $\lambda \in \mathbb{R}$. Let $D(\mathcal{F}) = \{\rho \in \mathcal{P}_2(\mathbb{R}^d) : \mathcal{F}(\rho) < +\infty\}$. The subdifferential set of \mathcal{F} at $\rho \in D(\mathcal{F})$ is denoted by $\partial\mathcal{F}(\rho)$ (see [AGS08, Definition 10.1.1] for definition of Fréchet subdifferentials). With $D(\partial\mathcal{F}) = \{\rho \in D(\mathcal{F}) : \partial\mathcal{F}(\rho) \neq \emptyset\}$, we make the assumptions that $D(\partial\mathcal{F}) \subset \mathcal{P}_2^{\text{ac}}(\mathbb{R}^d)$, the subset of absolutely continuous probability measures in $\mathcal{P}_2(\mathbb{R}^d)$. Using Brenier's theorem [San15, Theorem 1.22], this implies that there exists a unique optimal transport map from any $\rho \in D(\partial\mathcal{F})$ to any $\nu \in \mathcal{P}_2(\mathbb{R}^d)$. We denote this map by T_ρ^ν . Since \mathcal{F} is λ -geodesically convex, \mathcal{F} is a regular functional [AGS08, Definition 10.1.4] which implies that for all $\rho \in D(\partial\mathcal{F})$, the subdifferential set $\partial\mathcal{F}(\rho)$ contains a unique element with minimum $L^2(\rho)$ norm. This unique minimum selection subdifferential is called the Wasserstein gradient hereon, denoted by $\nabla_{\mathbb{W}}\mathcal{F}(\rho)$, and belongs to the tangent

space $\text{Tan}(\rho)$ [AGS08, Definition 8.4.1]. We further assume that there exists an $\varepsilon_0 > 0$ such that the functional

$$\nu \in \mathcal{P}_2(\mathbb{R}^d) \mapsto \mathcal{F}(\nu) + \frac{1}{2\varepsilon} \mathbb{W}_2^2(\mu, \rho)$$

admits at least one minimum point for any $\varepsilon \in (0, \varepsilon_0)$ and $\rho \in \mathcal{P}_2(\mathbb{R}^d)$.

The two primary classes of examples that we consider later are (i) the (one half) entropy function $\mathcal{F}(\rho) = \frac{1}{2} \text{Ent}(\rho)$ for which the gradient flow is the probabilistic heat flow and (ii) (one half) relative entropy (AKA Kullback-Leibler divergence) with respect to a log-concave probability measure, i.e., $\mathcal{F}(\rho) = \frac{1}{2} H(\cdot | \nu)$, for some log-concave density ν . These are well-known examples for which the above assumptions are satisfied. In Section 5 we will cover examples that are not covered by our assumptions and yet, as we will show by direct computations, our iterated Schrödinger bridge scheme continues to perform well.

4.2. Gradient Flows and their Euler Approximations. Let $(\rho(t), t \in [0, T])$ denote the Wasserstein gradient flow of \mathcal{F} during time interval $[0, T]$. We assume that it exists as an absolutely continuous curve that is the unique solution of the continuity equation

$$\partial_t \rho(t) + \nabla \cdot (v(t) \rho(t)) = 0, \quad \text{where } v(t) = -\nabla_{\mathbb{W}} \mathcal{F}(\rho(t))$$

is assumed to be in $\text{Tan}(\rho(t))$.

Euler iterations of the Wasserstein gradient flow of \mathcal{F} , whether it is explicit or implicit, is a sequence of elements in the Wasserstein space $(\rho_\varepsilon(k), k = 0, 1, 2, \dots)$, starting at some $\rho_\varepsilon(0) = \rho(0)$, given by an iterative map $S_\varepsilon^1(\cdot) : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathcal{P}_2(\mathbb{R}^d)$ of the form (45), in the sense that $\rho_\varepsilon(k) = S_\varepsilon^1(\rho_\varepsilon(k-1)) = S_\varepsilon^k(\rho_\varepsilon(0))$, for $k \in \mathbb{N}$. The continuous-time approximation curve is given by the piecewise constant interpolation

$$(54) \quad \rho_\varepsilon(t) = \rho_\varepsilon(k) \quad t \in [k\varepsilon, (k+1)\varepsilon).$$

Definition 1 (First order approximation). *The sequence $(\rho_\varepsilon(k), k \in \mathbb{N})$, or, equivalently, its continuous time interpolation $(\rho_\varepsilon(t), t \geq 0)$ is called a first-order approximation of a Wasserstein gradient flow $(\rho(t), t \geq 0)$ if, for any $T > 0$,*

$$\lim_{\varepsilon \rightarrow 0} \sup_{t \in [0, T]} \mathbb{W}_2(\rho_\varepsilon(t), \rho(t)) = 0.$$

We now wish to develop sufficient conditions under which $S_\varepsilon^1(\cdot)$ gives a first order approximation. We say that $S_\varepsilon^1(\cdot)$ is **consistent** if for any $T > 0$,

$$(55) \quad \limsup_{\varepsilon \rightarrow 0} \sup_{t \in [0, T-\varepsilon]} \frac{1}{\varepsilon} \mathbb{W}_2(S_\varepsilon^1(\rho(t)), \rho(t+\varepsilon)) = 0,$$

and $S_\varepsilon^1(\cdot)$ is a **contraction in the limit** if

$$(56) \quad \limsup_{\varepsilon \rightarrow 0} \sup_{k \in [N_\varepsilon]} \frac{1}{\varepsilon} \left(\mathbb{W}_2(S_\varepsilon^1(S_\varepsilon^k(\rho(0))), S_\varepsilon^1(\rho(k\varepsilon))) - \mathbb{W}_2(S_\varepsilon^k(\rho(0)), \rho(k\varepsilon)) \right) \leq 0.$$

Remark 2. *Observe that if $S_\varepsilon^1(\cdot)$ is a contraction, i.e. $\mathbb{W}_2(S_\varepsilon^1(\rho), S_\varepsilon^1(\sigma)) \leq \mathbb{W}_2(\rho, \sigma)$, then it is a contraction in the limit as well. Thus, this condition is a more general notion.*

Theorem 3. *Suppose the map $S_\varepsilon^1(\cdot)$ satisfies the two conditions - consistency and contraction in the limit. Then $\rho_\varepsilon(\cdot)$ is a first-order approximation of the gradient flow $(\rho(t), t \in [0, T])$.*

Proof. For any $t \in [k\varepsilon, (k+1)\varepsilon)$, by the triangle inequality and absolute continuity of $\rho(\cdot)$

$$\mathbb{W}_2(\rho_\varepsilon(t), \rho(t)) \leq \mathbb{W}_2(S_\varepsilon^k(\rho(0)), \rho(k\varepsilon)) + \int_{k\varepsilon}^t \|v(s)\|_{L^2(\rho(s))} ds.$$

The second term above converges to 0 with ε because $\|v(t)\|_{L^2(\rho(t))} \in L^1_{\text{loc}}([0, +\infty))$. For the first term, consider the decomposition

$$(57) \quad \mathbb{W}_2 \left(S_\varepsilon^k(\rho(0)), \rho(k\varepsilon) \right) \leq \mathbb{W}_2 \left(S_\varepsilon^k(\rho(0)), S_\varepsilon^1(\rho((k-1)\varepsilon)) \right) + \mathbb{W}_2 \left(S_\varepsilon^1(\rho((k-1)\varepsilon)), \rho(k\varepsilon) \right).$$

Let $\delta > 0$, then by the assumptions of consistency and contraction in the limit, for all $\varepsilon > 0$ small enough

$$\mathbb{W}_2 \left(S_\varepsilon^1(\rho((k-1)\varepsilon)), \rho(k\varepsilon) \right) < \delta\varepsilon,$$

and

$$\mathbb{W}_2 \left(S_\varepsilon^k(\rho(0)), S_\varepsilon^1(\rho((k-1)\varepsilon)) \right) - \mathbb{W}_2 \left(S_\varepsilon^{k-1}(\rho(0)), \rho((k-1)\varepsilon) \right) < \delta\varepsilon,$$

respectively. Thus, (57) gives the following recursion for $\varepsilon \in (0, \varepsilon_0)$ and $k \in [N_\varepsilon]$ with $k \geq 1$

$$\mathbb{W}_2 \left(S_\varepsilon^k(\rho(0)), \rho(k\varepsilon) \right) \leq \mathbb{W}_2 \left(S_\varepsilon^{k-1}(\rho(0)), \rho((k-1)\varepsilon) \right) + 2\delta\varepsilon.$$

Hence, recursively,

$$\limsup_{\varepsilon \rightarrow 0} \sup_{k \in [N_\varepsilon]} \mathbb{W}_2 \left(S_\varepsilon^k(\rho), \rho(k\varepsilon) \right) \leq \mathbb{W}_2(\rho(0), \rho(0)) + \frac{T}{\varepsilon} \cdot (2\delta\varepsilon) = 2T\delta.$$

As $\delta > 0$ was arbitrary, this completes the proof. \square

Explicit Euler Scheme. The explicit Euler scheme in the Wasserstein space can be obtained as an iterative geodesic approximation of the Wasserstein gradient flow minimizing \mathcal{F} for the choice of

$$(EE) \quad S_\varepsilon^1(\rho) = (\text{Id} - \varepsilon \nabla_{\mathbb{W}} \mathcal{F}(\rho))_{\#} \rho.$$

The explicit Euler scheme need not always give a first-order approximation sequence to a Wasserstein gradient flow. However, the assumptions in the statements of Theorems 4 and 5 provides sufficient conditions for which (EE) satisfies consistency and contraction in the limit. Consequently, Theorem 3 gives the uniform convergence of the scheme.

Implicit Euler Scheme. The implicit Euler scheme or, equivalently the JKO scheme [JKO98], in the Wasserstein space can be defined using the following map

$$(IE) \quad S_\varepsilon^1(\rho) = \operatorname{argmin}_{\nu \in \mathcal{P}_2(\mathbb{R}^d)} \left[\mathcal{F}(\nu) + \frac{1}{2\varepsilon} \mathbb{W}_2^2(\nu, \rho) \right].$$

We quickly verify that when $\rho \in \mathcal{P}_2^{ac}(\mathbb{R}^d)$ and $\varepsilon > 0$, there exists $u_\varepsilon : \mathbb{R}^d \rightarrow \mathbb{R}$ such that $S_\varepsilon^1(\rho) = (\text{Id} + \varepsilon \nabla u_\varepsilon)_{\#} \rho$. As $\rho \in \mathcal{P}_2^{ac}(\mathbb{R}^d)$, by Brenier's Theorem there exists a unique gradient of a convex function $\nabla \psi_\varepsilon$ that pushforwards ρ to $S_\varepsilon^1(\rho)$. Setting $u_\varepsilon(x) := \frac{1}{\varepsilon} \left(\psi_\varepsilon(x) - \frac{1}{2} \|x\|^2 \right)$, it holds that $S_\varepsilon^1(\rho) = (\text{Id} + \varepsilon u_\varepsilon)_{\#} \rho$ as desired. The implicit Euler scheme starting from $\rho(0) \in D(\partial \mathcal{F})$ is given by $\rho_\varepsilon(k) = S_\varepsilon^k(\rho(0))$. When \mathcal{F} is geodesically convex, it is known that Theorem 3 holds through [AGS08, Theorem 4.0.7, Theorem 4.0.9, Theorem 4.0.10]. We do not need to verify Definition 1.

In the next section we introduce our iterated Schrödinger bridge scheme that approximates the explicit scheme, under suitable assumptions.

4.3. Uniform Convergence. We now prove two theorems giving conditions under which iterations of (SB) provide a first order approximation to the gradient flow of $\frac{1}{2}\text{Ent}$ and $\frac{1}{2}H(\cdot|\nu)$. The proof of both theorems rely on an appeal to a general result in Lemma 6 that gives sufficient conditions for (SB) iterates to approximate Euler iterates. In principle, Lemma 6 can be applied with $S_\varepsilon^1(\cdot)$ given by implicit or explicit Euler iterations, but our Theorems 4 and 5 both only use explicit Euler iterations.

To simplify the bookkeeping of the constants introduced in Assumptions 2, we define two classes of probability densities. For $C, D, M > 0$, $N \geq 1$, and $\lambda \in \mathbb{R}$ set

$$(58) \quad \Pi_1(C, D, N) := \{\rho \in \mathcal{P}_2^{ac}(\mathbb{R}^d) : \|\rho\|_{\psi_1} \leq D, \nabla \mathcal{U}_\rho \text{ exists, } \|\mathcal{U}_\rho(x)\|^2 \leq C(1 + \|x\|^N)\},$$

and

$$(59)$$

$$\Pi_2(\lambda, M) := \{\rho \in \mathcal{P}_2^{ac}(\mathbb{R}^d) \text{ satisfying Assumption 1, } -\nabla^2 \log \rho \geq \lambda \text{Id, } E_\rho \|\nabla^2 \log \rho\|_{HS}^2 \leq M\}.$$

Lastly, recall the RHS of Theorems 1 and 2 and that $(\rho_s^\varepsilon, s \in [0, 1])$ is the entropic interpolation from ρ to itself. We define

$$(60) \quad \Delta_\varepsilon(\rho) := I(\rho) - \int_0^1 I(\rho_s^\varepsilon) ds.$$

We now state the following theorem for the (probabilistic) heat flow:

$$(61) \quad \partial_t \rho(t) = \frac{1}{2} \Delta_x \rho(t) = \frac{1}{2} \nabla \cdot ((\nabla \log \rho(t)) \rho(t)),$$

starting at $\rho(0)$. Let $S_\varepsilon^1(\cdot)$ denote the explicit Euler iteration map for the heat flow. That is, for $\rho \in \mathcal{P}_2^{ac}(\mathbb{R}^d)$

$$(62) \quad S_\varepsilon^1(\rho) := \left(\text{Id} - \frac{\varepsilon}{2} \nabla \log \rho \right)_\# \rho.$$

Theorem 4. *Assume that $\rho(0)$ is such that*

- (1) $\limsup_{\varepsilon \downarrow 0} \sup_{k \in [N_\varepsilon]} \Delta_\varepsilon(SB_\varepsilon^k(\rho(0))) = 0$
- (2) *There exists $\varepsilon_0 > 0$ and constants $C, D > 0$, and $N \geq 1$ such that for all $\varepsilon \in (0, \varepsilon_0)$, $(S_\varepsilon^k(\rho(0)), k \in \{0\} \cup [N_\varepsilon]) \cup (SB_\varepsilon^k(\rho(0)), k \in [N_\varepsilon]) \subset \Pi_1(C, D, N)$.*
- (3) *There exists $\lambda \in \mathbb{R}$ and $M > 0$ such that for all $\varepsilon \in (0, \varepsilon_0)$ and $k \in [N_\varepsilon]$, $(SB_\varepsilon^k(\rho(0)), k \in [N_\varepsilon]) \subset \Pi_2(\lambda, M)$.*

Then $(SB_\varepsilon^k(\rho_0), k \in [N_\varepsilon])$ is a first order approximation of the (probabilistic) heat flow.

Remark 3. *Theorem 4 lays down sufficient conditions under which [SABP22, Theorem 1] holds. Part of the assumptions are on the Schrödinger bridge iterates which are assumed to be nice enough such that our one step approximations hold uniformly. In Section 5, we show that these assumptions are indeed satisfied when $\rho(0)$ is Gaussian. We are currently unable to verify that these natural conditions must be true once sufficiently stringent conditions (say, strong log concavity and enough smoothness) are assumed on the initial measure $\rho(0)$.*

We now state an analogous theorem for the Wasserstein gradient flow of relative entropy. Fix a reference measure $\nu = e^{-h} \in \mathcal{P}_2(\mathbb{R}^d)$ satisfying assumptions delineated in Theorem 5 below.

Let $S_\varepsilon^1(\cdot)$ denote the explicit Euler iteration for the Fokker-Planck equation

$$(63) \quad \partial_t \rho(t) = \frac{1}{2} \nabla \cdot ((\nabla \log \rho(t) + \nabla h) \rho(t)).$$

That is, for $\rho \in \mathcal{P}_2^{ac}(\mathbb{R}^d)$

$$(64) \quad S_\varepsilon^1(\rho) := \left(\text{Id} - \frac{\varepsilon}{2} \nabla \log \rho - \frac{\varepsilon}{2} \nabla h \right)_{\#} \rho.$$

For $\varepsilon > 0$ and $k \in [N_\varepsilon]$, let $\sigma_\varepsilon^k \in \mathcal{P}_2^{ac}(\mathbb{R}^d)$ denote the surrogate measure obtained in the application of $\text{SB}_\varepsilon^1(\cdot)$ to $\text{SB}_\varepsilon^{k-1}(\rho(0))$, and let $\theta_\varepsilon^k \in \mathbb{R} \setminus \{0\}$ denote the corresponding integrability constant from (46). That is, σ_ε^k is the probability density proportional to $(\rho^{-1}\nu)^{\theta_\varepsilon^k}$, for $\rho = \text{SB}_\varepsilon^{k-1}(\rho(0))$.

Theorem 5. *Let h be convex, and assume that $\rho(0)$ is such that*

- (1) $\limsup_{\varepsilon \downarrow 0} \sup_{k \in [N_\varepsilon]} \Delta_\varepsilon(\sigma_\varepsilon^k) = 0$.
- (2) *There exists $C, D, M > 0$, $N \geq 1$, and $\lambda \in \mathbb{R}$ such that for all $\varepsilon \in (0, \varepsilon_0)$, $(\sigma_\varepsilon^k, k \in [N_\varepsilon]) \subset \Pi_1(C, D, N) \cap \Pi_2(\lambda, M)$, and $\|\text{SB}_\varepsilon^k(\rho(0))/\sigma_\varepsilon^k\|_\infty, (\theta_\varepsilon^k)^{-2} \leq M$ for all $k \in [N_\varepsilon]$.*
- (3) *(Regularity of Fokker-Planck) The collection $\mathcal{F} = (S_\varepsilon^k(\rho(0)), k \in \{0\} \cup [N_\varepsilon]) \cup (\text{SB}_\varepsilon^k(\rho(0)), k \in [N_\varepsilon])$ is such that*

$$\limsup_{\varepsilon \downarrow 0} \sup_{\rho \in \mathcal{F}} \frac{1}{\varepsilon} \mathbb{W}_2(S_\varepsilon^1(\rho), \rho(\varepsilon)) = 0.$$

Then $(\text{SB}_\varepsilon^k(\rho_0), k \in [N_\varepsilon])$ is a first order approximation of the gradient flow of (one-half) KL divergence, i.e. the solution to (63).

Remark 4. *These assumed regularity of the Fokker-Planck is just requiring that the convergence given in [AGS08, Proposition 8.4.6] holds uniformly along the measures in \mathcal{F} . That is, if we start Fokker-Planck from any of the measures ρ in \mathcal{F} , the explicit Euler iteration is a good enough approximation for small times, uniformly in ρ . In Lemma 7 below we show that this statement is true for the heat flow since its corresponding transition kernel is a Gaussian convolution. It is also satisfied by the flow corresponding to the standard Ornstein-Uhlenbeck (OU) process for which the transition kernel is again a Gaussian convolution. In fact, Lemma 7 continues to hold in this case with minor modifications to the proof.*

Observe the difference in the assumptions required for Theorems 4 and 5. The new assumptions in Theorem 5 are regarding the surrogate measures σ_ε^k corresponding to the Schrödinger iterate $(\text{SB}_\varepsilon^k(\rho(0)))$. The reason for this difference is that for the heat flow $\sigma_\varepsilon^k = (\text{SB}_\varepsilon^k(\rho(0)))$ and $\theta_\varepsilon^k = -1$, for every $k \in [N_\varepsilon]$ and any $\varepsilon > 0$. Hence the assumptions simplify.

Before proving Theorems 4 and 5, we present general conditions under which the iterates of (SB), called SB scheme, converge to the gradient flow of a functional \mathcal{F} . For a general case, it is difficult to show that a continuous time interpolation of the SB scheme is a first order approximation of the gradient flow (Definition 1). The following general conditions ensure uniform convergence to the gradient flow via Euler iterations. We then restrict ourselves to the specific cases when \mathcal{F} is entropy and relative entropy.

Assumption 3. *Fix $\mathcal{F} : \mathcal{P}_2(\mathbb{R}^d) \rightarrow (-\infty, +\infty]$ be a geodesically convex functional and fix $T > 0$. With initial measure $\rho(0) \in \mathcal{P}_2^{ac}(\mathbb{R}^d)$, let $(\rho(t), t \geq 0)$ denote the gradient flow of \mathcal{F} . Let $(\rho_\varepsilon(k\varepsilon) = S_\varepsilon^k(\rho_0), k \in [N_\varepsilon])$ be a sequence of Euler iterations that gives rise to a first order approximation of the gradient flow of \mathcal{F} satisfying Definition 1. Moreover, assume that:*

(1) there is **consistency** of $S_\varepsilon^1(\cdot)$ along the $(SB_\varepsilon^k(\rho_0), k \in [N_\varepsilon])$, meaning

$$(65) \quad \limsup_{\varepsilon \downarrow 0} \sup_{k \in [N_\varepsilon]} \frac{1}{\varepsilon} \mathbb{W}_2 \left(SB_\varepsilon^k(\rho_0), S_\varepsilon^1 \left(SB_\varepsilon^{k-1}(\rho_0) \right) \right) = 0.$$

(2) There is **contraction in the limit**, meaning

$$(66) \quad \limsup_{\varepsilon \rightarrow 0} \sup_{k \in [N_\varepsilon]} \frac{1}{\varepsilon} \left(\mathbb{W}_2 \left(S_\varepsilon^k(\rho_0), S_\varepsilon^1 \left(SB_\varepsilon^{k-1}(\rho_0) \right) \right) - \mathbb{W}_2 \left(S_\varepsilon^{k-1}(\rho_0), SB_\varepsilon^{k-1}(\rho_0) \right) \right) \leq 0.$$

These assumptions are reminiscent of (55) and (56). Hence, the proof of the following lemma is very similar to that of Theorem 3.

Lemma 6. *Under Assumption 3,*

$$\limsup_{\varepsilon \downarrow 0} \sup_{k \in [N_\varepsilon]} \mathbb{W}_2(\rho(k\varepsilon), SB_\varepsilon^k(\rho_0)) = 0.$$

Proof. To start, observe that

$$(67) \quad \sup_{k \in [N_\varepsilon]} \mathbb{W}_2(SB_\varepsilon^k(\rho_0), \rho(k\varepsilon)) \leq \sup_{k \in [N_\varepsilon]} \mathbb{W}_2(SB_\varepsilon^k(\rho_0), S_\varepsilon^k(\rho_0)) + \sup_{k \in [N_\varepsilon]} \mathbb{W}_2(S_\varepsilon^k(\rho_0), \rho(k\varepsilon)).$$

By Theorem 3, the rightmost term vanishes as $\varepsilon \downarrow 0$. For the remaining term, for $k \in [N_\varepsilon]$ split it as

$$\mathbb{W}_2(SB_\varepsilon^k(\rho_0), S_\varepsilon^k(\rho_0)) \leq \mathbb{W}_2 \left(SB_\varepsilon^k(\rho_0), S_\varepsilon^1 \left(SB_\varepsilon^{k-1}(\rho_0) \right) \right) + \mathbb{W}_2 \left(S_\varepsilon^1 \left(SB_\varepsilon^{k-1}(\rho_0) \right), S_\varepsilon^k(\rho_0) \right).$$

By Assumption 3(2), one obtains the exact same recursion as in Theorem 3. Hence, the same argument gives that $\limsup_{\varepsilon \downarrow 0} \sup_{k \in [N_\varepsilon]} \mathbb{W}_2(SB_\varepsilon^k(\rho_0), S_\varepsilon^k(\rho_0)) = 0$, completing the Lemma. \square

We now begin the proof of Theorem 4, which follows as an application of Lemma 6 once we verify Assumption 3 holds in this setting. The verification is a lengthy argument that relies on a tight approximation at small time of the explicit Euler step given in (62) and the heat flow, which we present in Lemma below.

Lemma 7. *Let $\rho = e^{-g} \in \Pi_1(C, D, N)$. Let $S_\varepsilon^1(\cdot)$ be as defined in (62), and $(\rho(t), t \geq 0)$ the solution to (61) with $\rho(0) = \rho$. Fix $\varepsilon_0 > 0$, then there is a constant $K > 0$ depending on C, D , and n such that for any $\varepsilon \in (0, \varepsilon_0)$*

$$\mathbb{W}_2(S_\varepsilon^1(\rho), \rho(\varepsilon)) \leq K\varepsilon^2.$$

Proof. We consider two evolving particle systems. Let $(\rho(t, \cdot), t \geq 0)$ denote the (probabilistic) heat flow starting from ρ , and set $g(t, x) = -\log \rho(t, x)$. Let $X, Y : [0, \infty) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ be two particle systems with the same initial configuration $X_0 \sim \rho$ solving the ODE (letting \dot{X}_t denote time derivative)

$$\dot{X}_t = \frac{1}{2} \nabla g(t, X_t) \text{ and } \dot{Y}_t = \frac{1}{2} \nabla g(0, X_0).$$

Observe that $X_\varepsilon \sim \rho(\varepsilon)$ and $Y_\varepsilon \sim S_\varepsilon^1(\rho)$. Now, define $G : [0, +\infty) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ by $G(t, x) = X(t, x) - Y(t, x)$. Observe that $G(0, \cdot) = \dot{G}(0, \cdot) = 0$, giving that

$$(68) \quad G(t) = \int_0^t \int_0^s \ddot{G}(u) du ds.$$

By the chain rule and the fact that $(\rho(t, \cdot), t \geq 0)$ solves the heat equation, observe that

$$\begin{aligned}\ddot{G}(t, X_t) &= \frac{1}{2} \partial_t \nabla_x g(t, X_t) \\ &= \frac{1}{2} \left(\nabla_x^2 g(t, X_t) \dot{X}_t - \frac{1}{2} \nabla_x \left[\frac{\Delta_x \rho(t, X_t)}{\rho(t, X_t)} \right] \right).\end{aligned}$$

In analogy with Section 2, define

$$\mathcal{U}(t, u) = \frac{1}{8} \|\nabla_x g(t, u)\|^2 - \frac{1}{4} \Delta_x g(t, u),$$

and compute that

$$\frac{1}{4} \nabla_x \left[\frac{\Delta \rho(t, u)}{\rho(t, u)} \right] = \nabla_x \left(\frac{1}{8} \|\nabla_x g(t, u)\|^2 + \mathcal{U}(t, u) \right) = \frac{1}{4} \nabla_x^2 g(t, u) \nabla g(t, u) + \nabla_x \mathcal{U}(t, u).$$

Altogether, it now holds that

$$(69) \quad \ddot{G}(t, X_t) = -\nabla_x \mathcal{U}(t, X_t).$$

For a fixed $\varepsilon_0 > 0$, there uniform constant $C' > 0$ and $n' \geq 1$ (potentially slightly modified from C and n) such that for all $t \in [0, \varepsilon_0]$, $\|\nabla_x \mathcal{U}(t, x)\|^2 \leq C'(1 + \|x\|^{n'})$. Let $X_0 \sim \rho$ and $Z \sim N(0, \text{Id})$ be independent, then for $t \geq 0$ the random variable $X_0 + \varepsilon Z$ has law $\rho(t)$. Moreover, as $\|X_0 + tZ\|_{\psi_1} \leq D + t\|Z\|_{\psi_1}$, the $\rho(t)$ have bounded subexponential norm over $t \in [0, \varepsilon_0]$. Hence, there exists a constant K' as described in the Lemma such that for $t \in [0, \varepsilon_0]$, $\mathbb{E}_{\rho(t)} \|\nabla_x \mathcal{U}(t, X)\|^2 \leq K'$. Hence, applying Jensen's inequality twice to (68) gives

$$\mathbb{W}_2^2(S_\varepsilon^1(\rho), \rho(\varepsilon)) \leq \mathbb{E} \|G(\varepsilon, X_0)\|^2 \leq \int_0^\varepsilon \int_0^s \varepsilon s \mathbb{E} [\|\ddot{G}(u, X_u)\|^2] du ds \leq \frac{\varepsilon^4}{3} K',$$

establishing the Lemma. \square

We now prove Theorem 4.

Proof of Theorem 4. It suffices to verify that Assumption 3 holds in this setting.

Step 1: Explicit Euler Iterations are Consistent and Contractive in the Limit.

Since $\rho(0)$ is assumed to be in $\Pi_1(C, D, N)$, by Lemma 7, along the heat flow starting at $\rho(0)$, there is a uniform constant $K > 0$ such that, for any $\varepsilon > 0$ small enough and for all $t \in [0, T - \varepsilon]$, $\mathbb{W}_2(S_\varepsilon^1(\rho(t)), \rho(t + \varepsilon)) \leq K\varepsilon^2$.

Next, we establish contractivity in the limit. For a measure $\mu \in \mathcal{P}(\mathbb{R}^d)$ let $[\mu](t)$ denote the time marginal of the heat flow (61) at time t started from μ . By contractivity of the heat flow with respect to initial data in the Wasserstein-2 metric [AGS08, Theorem 11.1.4],

$$\mathbb{W}_2([S_\varepsilon^k(\rho(0))](\varepsilon), \rho((k+1)\varepsilon)) \leq \mathbb{W}_2(S_\varepsilon^k(\rho(0)), \rho(k\varepsilon)).$$

It then follows from triangle inequality

$$\begin{aligned}\mathbb{W}_2 \left(S_\varepsilon^1 \left(S_\varepsilon^k(\rho(0)) \right), S_\varepsilon^1(\rho(k\varepsilon)) \right) &\leq \mathbb{W}_2(S_\varepsilon^k(\rho(0)), \rho(k\varepsilon)) + \mathbb{W}_2(S_\varepsilon^1(S_\varepsilon^k(\rho(0))), [S_\varepsilon^k(\rho(0))](\varepsilon)) \\ &\quad + \mathbb{W}_2(S_\varepsilon^1(\rho(k\varepsilon)), \rho((k+1)\varepsilon)).\end{aligned}$$

Hence, by assumptions in the theorem statement, for ε small enough there is a universal constant $K > 0$ such that for all $k \in [N_\varepsilon]$

$$\left(\mathbb{W}_2 \left(S_\varepsilon^1 \left(S_\varepsilon^k(\rho(0)) \right), S_\varepsilon^1(\rho(k\varepsilon)) \right) - \mathbb{W}_2 \left(S_\varepsilon^k(\rho(0)), \rho(k\varepsilon) \right) \right) \leq 2K\varepsilon^2,$$

establishing contraction in the limit. This demonstrates that the explicit Euler iterations converge to the heat flow.

Step 2: SB Iterations are Consistent and Contractive in the Limit. To start, fix $\varepsilon > 0$ and $k \geq 1$. To compute $\text{SB}_\varepsilon^k(\rho_0)$ from $\text{SB}_\varepsilon^{k-1}(\rho_0)$, we must obtain the surrogate measure for (SB). Pick $\theta_\varepsilon = -1$, then the surrogate measure is the measure itself, that is, $\sigma_\varepsilon = \text{SB}_\varepsilon^{k-1}(\rho_0)$. This in turn gives $\|\text{SB}_\varepsilon^{k-1}(\rho_0)/\sigma_\varepsilon\|_\infty = 1$. For consistency, by Theorem 2 and the assumptions in the statement of Theorem 4, for small enough $\varepsilon > 0$ there is a universal constant $K > 0$ such that for all $k \in [N_\varepsilon]$ with $k \geq 1$

$$\mathbb{W}_2 \left(\text{SB}_\varepsilon^k(\rho(0)), S_\varepsilon^1 \left(\text{SB}_\varepsilon^{k-1}(\rho(0)) \right) \right) \leq \varepsilon K \left(I(\text{SB}_\varepsilon^{k-1}(\rho(0))) - \int_0^1 I((\text{SB}_\varepsilon^{k-1}(\rho(0)))_s^\varepsilon) ds \right)^{1/4}.$$

By the first assumption in the Theorem statement, consistency holds. Contractivity in the limit holds by the triangle inequality (as in the previous argument) and applying Lemma 7. \square

The proof of Theorem 5 is completely analogous to the proof of Theorem 4 and thus omitted. The only modification is that we replace the role of Lemma 7 with the assumed regularity of Fokker-Planck equation in Theorem 5(3). See also Remark 4.

5. EXAMPLES

In this section we compute explicit examples of iterated Schrödinger bridge approximations to various gradient flows and other absolutely continuous curves on the Wasserstein space. Since Schrödinger bridges are known in closed form for Gaussian marginals ([JMPC20]), our computations are limited to Gaussian models. First, we consider gradient flows of two convex functionals - 1) entropy and 2) KL divergence. Second, while not being gradient flows and aligning with our theoretical analysis in Section 4, we also consider time reversal of these gradient flows to showcase the curiously nice approximation properties of the (SB) scheme for these absolutely continuous curves in the Wasserstein space. These latter examples have recently become important in applications. For example, see [SSDK⁺20] for an application to diffusion based generative models in machine learning.

5.1. Explicit Euler discretization of the gradient flow of Entropy. We show via explicit calculations that (SB) iterations are close to the explicit Euler approximation of the heat flow $(\rho(t), t \in [0, T])$ starting from a Gaussian density. Here $\mathcal{F} = \frac{1}{2}\text{Ent}$ and the continuity equation for the gradient flow of \mathcal{F} is $\partial_t \rho(t) + \nabla \cdot (v(t)\rho(t)) = 0$ where $v(t) = -\frac{1}{2}\nabla \log \rho(t)$. The explicit Euler iterations, $(\rho_\varepsilon(k), k \in [N_\varepsilon])$, where $N_\varepsilon := \lfloor T\varepsilon^{-1} \rfloor$, are given by $\rho_\varepsilon(k+1) = (\text{Id} + \varepsilon v_\varepsilon(k))_{\#} \rho_\varepsilon(k)$ where $v_\varepsilon(k) = -\frac{1}{2}\nabla \log \rho_\varepsilon(k)$.

For $\rho_\varepsilon(0) = \rho = \mathcal{N}(0, \eta^2 I_d)$, $\eta^2 > 0$, we will calculate the (EE) and (SB) iterates and show that the SB step is an $\mathcal{O}(\varepsilon^2)$ approximation of the explicit Euler step. Recall that the form of Schrödinger bridge with equal marginals ρ is explicitly given by [JMPC20, Theorem 1] as

$$\pi_{\rho, \varepsilon} = \mathcal{N} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \eta^2 \begin{pmatrix} I_d & C_{\eta^2}^\varepsilon I_d \\ C_{\eta^2}^\varepsilon I_d & I_d \end{pmatrix} \right) \quad \text{where} \quad C_{\eta^2}^\varepsilon = \frac{1}{\eta^2} \left(\sqrt{\eta^4 + \frac{\varepsilon^2}{4}} - \frac{\varepsilon}{2} \right).$$

From this the barycentric projection can be computed as

$$(70) \quad \mathcal{B}_{\rho, \varepsilon}(x) = \mathbb{E}_{\pi_{\rho, \varepsilon}} [Y | X = x] = C_{\eta^2}^\varepsilon x.$$

For the choice of ρ , $\text{SB}_\varepsilon^1(\rho) = (2\text{Id} - \mathcal{B}_{\rho, \varepsilon})_{\#} \rho = \mathcal{N} \left(0, \left(2 - C_{\eta^2}^\varepsilon \right)^2 \eta^2 I_d \right)$, and the corresponding $\rho_\varepsilon(1) = (\text{Id} - \frac{\varepsilon}{2} \nabla \log \rho)_{\#} \rho = \mathcal{N} \left(0, \left(1 + \frac{\varepsilon}{2\eta^2} \right)^2 \eta^2 I_d \right)$.

Now let's check the sharpness of Theorem 2. By the explicit form of the \mathbb{W}_2 distance between two Gaussians

$$\mathbb{W}_2\left(\rho_\varepsilon(1), \text{SB}_\varepsilon^1(\rho)\right) = d\left|(2 - C_{\eta^2}^\varepsilon)\eta - (1 + \varepsilon\eta^{-2})\eta\right| = d\eta\left|\sqrt{1 + \frac{\varepsilon^2}{4\eta^4}} - 1\right|.$$

Therefore, $\mathbb{W}_2\left(\rho_\varepsilon(1), \text{SB}_\varepsilon^1(\rho)\right) = \mathcal{O}(\varepsilon^2)$, which is an order of magnitude sharper than Theorem 2.

Since both (SB) and (EE) schemes operate via linear pushforwards, then all steps of these schemes are mean-zero Gaussian distributed. For the SB scheme, the surrogate measure at each step is the same as the current measure, i.e. $\sigma_\varepsilon(k) = \text{SB}_\varepsilon^k(\rho)$ for all $k \in [N_\varepsilon]$. Let $\text{SB}_\varepsilon^k(\rho_0) = \mathcal{N}(0, \alpha_k^2 I_d)$ and $\rho_\varepsilon(k) = \mathcal{N}(0, \beta_k^2 I_d)$, then the iterates evolve through the following recursive relationship

$$\alpha_{k+1}^2 = \left(2 - C_{\alpha_k^2}^\varepsilon\right)^2 \alpha_k^2 \quad \text{and} \quad \beta_{k+1}^2 = \left(1 + \frac{\varepsilon}{2\beta_k^2}\right)^2 \beta_k^2.$$

To show that $(\rho_\varepsilon(k), k \in [N_\varepsilon])$ and $(\text{SB}_\varepsilon^k(\rho), k \in [N_\varepsilon])$ are first-order approximations (Definition 1) of the Wasserstein gradient flow $(\rho(t), t \in [0, T])$, we will show that the assumptions of Theorem 4 are satisfied.

First, let us consider assumption (1) of Theorem 4. The formula for the time marginals of the dynamic Schrödinger bridge between $\sigma = \mathcal{N}(0, \eta^2)$ and itself at time $t = 1/2$ is available from [GLR17] and given by

$$(71) \quad (\sigma)_{1/2}^\varepsilon = \mathcal{N}\left(0, \left(\frac{\delta_\varepsilon^2}{4(1 + \delta_\varepsilon)} + 1\right)\eta^2\right) \quad \text{where } \delta_\varepsilon = \frac{1}{2}\left(\varepsilon - 2 + \sqrt{4 + \varepsilon^2}\right).$$

Also, it is known that for $\sigma = \mathcal{N}(0, \eta^2 I_d)$, the Fisher information $I(\sigma) = d\eta^{-2}$. Therefore, for $\sigma_\varepsilon(k) = \text{SB}_\varepsilon^k(\rho) = \mathcal{N}(0, \alpha_k^2)$, we have

$$(72) \quad I(\text{SB}_\varepsilon^k(\rho)) - I((\text{SB}_\varepsilon^k(\rho))_{1/2}^\varepsilon) = \frac{d}{\alpha_k^2} \left(1 - \left(\frac{\delta_\varepsilon^2}{4(1 + \delta_\varepsilon)} + 1\right)^{-1}\right) = \frac{d\delta_\varepsilon^2}{\alpha_k^2 (\delta_\varepsilon^2 + 4(1 + \delta_\varepsilon))}.$$

Since $C_{\nu^2}^\varepsilon \leq 1$ for any $\nu^2 > 0$, $(\alpha_k^2, k \in [N_\varepsilon])$ is monotonically increasing and bounded from below by η^2 . Hence,

$$\sup_{k \in [N_\varepsilon]} \left(I(\text{SB}_\varepsilon^k(\rho)) - I((\text{SB}_\varepsilon^k(\rho))_{1/2}^\varepsilon)\right) \leq I(\rho) - I(\rho_{1/2}^\varepsilon) = \frac{d\delta_\varepsilon^2}{\eta^2 (\delta_\varepsilon^2 + 4(1 + \delta_\varepsilon))}.$$

Since the RHS above converges to zero as $\varepsilon \rightarrow 0+$, we satisfy Assumption (1) of Theorem 4 with rate $\mathcal{O}(\varepsilon^2)$.

Now, we consider assumption (2) of Theorem 4. One can be convinced that for the Gaussian case, if both sequences $(\alpha_k^2, k \in [N_\varepsilon])$ and $(\beta_k^2, k \in [N_\varepsilon])$ are bounded from below and above, then the assumption (2) is satisfied. Trivially, $(\alpha_k^2, k \in [N_\varepsilon])$ and $(\beta_k^2, k \in [N_\varepsilon])$ are bounded from below by η^2 because they are monotonically increasing. Take $\varepsilon \leq \eta^2$. Using the identity $1 - x \leq \sqrt{1 + x^2}$ for all $x \geq 0$, we have that $2 - C_{\alpha_k^2}^\varepsilon \leq 1 + \varepsilon\alpha_k^{-2}$ and therefore $\alpha_{k+1}^2 \leq \alpha_k^2 + 2\varepsilon + \varepsilon^2/\alpha_k^2 \leq \alpha_k^2 + 3\varepsilon$. Consequently, $(\alpha_k^2, k \in [N_\varepsilon])$ is bounded from above by $\eta^2 + 3T$. Now, again using that $\varepsilon \leq \eta^2$, $\beta_{k+1}^2 \leq \beta_k^2 + 5\varepsilon/4$, and therefore $(\beta_k^2, k \in [N_\varepsilon])$ is bounded from above by $\eta^2 + 5T/4$. Step 1 of the proof of Theorem 4 gives uniform convergence of $(\rho_\varepsilon(k), k \in [N_\varepsilon])$ and step 2 gives the uniform convergence of $(\text{SB}_\varepsilon^k(\rho), k \in [N_\varepsilon])$.

5.2. Explicit Euler discretization of the gradient flow of Kullback-Leibler. Now let us consider the Kullback-Leibler (KL) divergence functional $H(\cdot|\nu)$ where $\nu = e^{-g}$ for $\nabla g(x) = x$. This is the Fokker-Planck equation corresponding to the Ornstein-Uhlenbeck process.

Now let us write the (SB) step at $\rho \in \mathcal{P}_2^{ac}(\mathbb{R}^d)$. For this functional, the selection of $\theta \in \mathbb{R} \setminus \{0\}$ in (SB) becomes more delicate. The choice of θ for the surrogate measure $\sigma = \sigma_\varepsilon(0)$ is determined by the integrability of

$$\exp\left(-2\theta\left(\frac{1}{2}\log\rho + \frac{1}{2}g\right)\right) = (\rho e^g)^{-\theta}.$$

If ρ is log-concave, then since g is convex the sign of θ depends on a comparison of the log-concavity of ρ and ν . Suppose ρ is more log-concave than ν , then θ may be taken to be -1 . Consequently, the (SB) step is $\text{SB}_\varepsilon^1(\rho) = (2\text{Id} - \mathcal{B}_{\sigma,\varepsilon})\#\rho$. On the other hand, if ν is more log-concave than the density ρ , then θ may be taken to be 1. In this case, the (SB) step is $\text{SB}_\varepsilon^1(\rho) = (\mathcal{B}_{\sigma,\varepsilon})\#\rho$.

For an initial $\rho = \mathcal{N}(0, \eta^2 I_d)$, $\eta^2 > 0$, we calculate the (SB) and the (EE) step and prove that the (SB) step is an $\mathcal{O}(\varepsilon^2)$ approximation of the (EE) step. If $\eta^2 > 1$, then ρ is less log-concave than e^{-g} , and therefore $\sigma = \rho^{-1}e^{-g} = \mathcal{N}(0, \eta^2/(\eta^2 - 1)I_d)$. This surrogate measure satisfies Assumption 2. By (70), we have that $\text{SB}_\varepsilon^1(\rho) = \mathcal{N}\left(0, (C_{\eta^2/(\eta^2-1)}^\varepsilon)^2 \eta^2 I_d\right)$. Now if $\eta^2 < 1$, then ρ is more log-concave than e^{-g} , and therefore $\sigma = \rho e^g = \mathcal{N}(0, \eta^2/(1 - \eta^2)I_d)$. Note that this choice satisfies Assumption 2 (1-2, 4) but the ratio $\frac{\rho}{\sigma} = \rho^2 e^g$ is unbounded. Nevertheless, we calculate the (SB) update using this surrogate measure and demonstrate one-step convergence, indicating that our assumptions in this paper are stronger than required. By (70) we have $\text{SB}_\varepsilon^1(\rho) = \mathcal{N}(0, (2 - C_{\eta^2/(1-\eta^2)}^\varepsilon)^2 \eta^2 I_d)$. The corresponding (EE) update is $\rho_\varepsilon(1) = (\text{Id} - \frac{\varepsilon}{2}(\nabla \log \rho + \text{Id}))\#\rho = \mathcal{N}\left(0, \left(1 + \frac{\varepsilon(1-\eta^2)}{2\eta^2}\right)^2 \eta^2 I_d\right)$.

Again, we check the sharpness of Theorem 2 by explicitly calculating the \mathbb{W}_2 distance between $\rho_\varepsilon(1)$ and $\text{SB}_\varepsilon^1(\rho)$. For any $\eta^2 > 0$,

$$\mathbb{W}_2\left(\rho_\varepsilon(1), \text{SB}_\varepsilon^1(\rho)\right) = \eta d \left| 2 - C_{\eta^2/(1-\eta^2)}^\varepsilon - 1 - \frac{\varepsilon(1-\eta^2)}{2\eta^2} \right| = \eta d \left| 1 - \sqrt{1 + \frac{\varepsilon^2(1-\eta^2)^2}{4\eta^4}} \right|.$$

Therefore, again $\mathbb{W}_2\left(\rho_\varepsilon(1), \text{SB}_\varepsilon^1(\rho)\right) = \mathcal{O}(\varepsilon^2)$ which is an order of magnitude better than Theorem 2. As mentioned above, here is an example of a fast convergence rate even when the surrogate measure σ does not satisfy Assumption 2(3).

Iterates of the (SB) scheme can be defined beyond the one step in the similar manner if the sign convention remains same throughout the iterative process. The following calculation is done for $\eta^2 < 1$ which is not covered by our theorem. A similar calculation is valid for $\eta^2 > 1$ which is skipped. We claim that if ρ is more log-concave than $\nu = \mathcal{N}(0, I_d)$, then $\text{SB}_\varepsilon^1(\rho)$ is also more log-concave than ν for small enough $\varepsilon > 0$. This is because, using Taylor series expansion around $\varepsilon = 0$, we have

$$\begin{aligned} \left(2 - C_{\eta^2/(1-\eta^2)}^\varepsilon\right)^2 \eta^2 &= \left(2 - \left[\frac{1-\eta^2}{\eta^2} \left(\sqrt{\frac{\eta^4}{(1-\eta^2)^2} + \frac{\varepsilon^2}{4}} - \frac{\varepsilon}{2}\right)\right]\right)^2 \eta^2 \\ &= \eta^2 + (1-\eta^2)\varepsilon - \frac{\varepsilon^2(1-\eta^2)^2}{8\eta^2} + \mathcal{O}(\varepsilon^4). \end{aligned}$$

Therefore, as long as $\varepsilon < \frac{\eta^2}{(1-\eta^2)}$, the SB steps approach monotonically toward the minimizer $\mathcal{N}(0, I_d)$ and the sign convention remains the same. Again, since both (SB) and (EE) schemes evolve via linear pushforwards, all steps are mean-zero Gaussian distributed. Denote $\text{SB}_\varepsilon^k(\rho_0) = \mathcal{N}(0, \alpha_k^2 I_d)$ and $\rho_\varepsilon(k) = \mathcal{N}(0, \beta_k^2 I_d)$ where α_k^2 and β_k^2 evolve via the following recursive relationship

$$\alpha_{k+1}^2 = \left(2 - C_{\alpha_k^2/(1-\alpha_k^2)}^\varepsilon\right)^2 \alpha_k^2 \quad \text{and} \quad \beta_{k+1}^2 = \left(1 + \frac{\varepsilon(1-\beta_k^2)}{\beta_k^2}\right)^2 \beta_k^2.$$

The corresponding surrogate measures $(\sigma_\varepsilon(k), k \in [N_\varepsilon])$ are given by $\sigma_\varepsilon(k) = \mathcal{N}(0, \alpha_k^2/(1-\alpha_k^2)I_d)$.

We again show that the assumptions of Theorem 5 are satisfied to prove that $(\rho_\varepsilon(k), k \in [N_\varepsilon])$ and $(\text{SB}_\varepsilon^k(\rho), k \in [N_\varepsilon])$ are first-order approximation schemes of the Wasserstein gradient flow $(\rho(t), t \in [0, T])$. Using (71), we have that

$$I(\sigma_\varepsilon(k)) - I((\sigma_\varepsilon(k))_{1/2}^\varepsilon) = \left(\frac{1-\alpha_k^2}{\alpha_k^2}\right) \frac{d\delta_\varepsilon^2}{\delta_\varepsilon^2 + 4(1+\delta_\varepsilon)},$$

where $\delta_\varepsilon = \frac{1}{2}(\varepsilon - 2 + \sqrt{4 + \varepsilon^2})$. The sequence $(\alpha_k^2, k \in [N_\varepsilon])$ is monotonically increasing because

$$\alpha_{k+1}^2 = \left(2 - \sqrt{1 + \frac{\varepsilon^2(\alpha_k^2 - 1)^2}{4\alpha_k^4}} + \frac{\varepsilon(1-\alpha_k^2)}{2\alpha_k^2}\right)^2 \alpha_k^2,$$

and the prefactor of α_k^2 can be easily checked to be greater than 1. Moreover, each α_k^2 is bounded from below by its initial value η^2 . Therefore,

$$\sup_{k \in [N_\varepsilon]} \left(I(\sigma_\varepsilon(k)) - I((\sigma_\varepsilon(k))_{1/2}^\varepsilon)\right) \leq I(\sigma(0)) - I((\sigma(0))_{1/2}^\varepsilon) = \left(\frac{1-\eta^2}{\eta^2}\right) \frac{d\delta_\varepsilon^2}{\delta_\varepsilon^2 + 4(1+\delta_\varepsilon)}.$$

Since $\delta_\varepsilon = \mathcal{O}(\varepsilon)$, the RHS converges to zero as $\varepsilon \rightarrow 0+$ at the order $\mathcal{O}(\varepsilon^2)$. Therefore, Assumption (1) of Theorem 5 is satisfied with rate $\mathcal{O}(\varepsilon^2)$.

Now, we show that assumption (2) of Theorem 5 is satisfied. Again, in the Gaussian case, it suffices to show that $(\alpha_k^2/(1-\alpha_k^2), k \in [N_\varepsilon])$ and $(\beta_k^2, k \in [N_\varepsilon])$ is bounded from below and above. Since $C_{\nu^2}^\varepsilon < 1$ for any $\nu^2 > 0$, the sequence $(\alpha_k^2, k \in [N_\varepsilon])$, and equivalently $(\alpha_k^2/(1-\alpha_k^2), k \in [N_\varepsilon])$, is monotonically increasing. Then trivially, $(\alpha_k^2/(1-\alpha_k^2), k \in [N_\varepsilon])$ is lower bounded by $\eta^2/(1-\eta^2)$ and $(\beta_k^2, k \in [N_\varepsilon])$ are lower bounded by η^2 . Take $\varepsilon < \min\left(\frac{\eta^2}{1-\eta^2}, \frac{1}{3}\right)$.

Using the identity $1-x \leq \sqrt{1+x}$ for all $x \geq 0$, we have that $2 - C_{\alpha_k^2/(1-\alpha_k^2)}^\varepsilon \leq 1 + \varepsilon(1-\alpha_k^2)/\alpha_k^2$.

Therefore, $\alpha_{k+1}^2 \leq (1 + \varepsilon(1-\alpha_k^2)/\alpha_k^2)^2 \alpha_k^2$ and $\beta_{k+1}^2 = (1 + \varepsilon(1-\beta_k^2)/\beta_k^2)^2 \beta_k^2$ give recursive inequalities of similar form. Since $\varepsilon \leq \alpha_k^2/(1-\alpha_k^2)$, we have $\alpha_{k+1}^2 \leq \alpha_k^2 + 3\varepsilon(1-\alpha_k^2)$. Recursively, $\alpha_{N_\varepsilon}^2 \leq (1-3\varepsilon)^{N_\varepsilon} \eta^2 + 1 - (1-3\varepsilon)^{N_\varepsilon}$, which is bounded from above. Therefore,

$\left(\frac{\alpha_k^2}{(1-\alpha_k^2)}, k \in [N_\varepsilon]\right)$ is bounded from above by $\frac{\eta^2}{1-\eta^2} + \frac{1-(1-3\varepsilon)^{N_\varepsilon}}{(1-3\varepsilon)^{N_\varepsilon}(1-\eta^2)}$ and similarly, $(\beta_k^2, k \in [N_\varepsilon])$

is bounded from above by $(1-3\varepsilon)^{N_\varepsilon} \eta^2 + (1 - (1-3\varepsilon)^{N_\varepsilon})$, which is bounded as $\varepsilon \downarrow 0$. Consequently, the assumptions of Theorem 5 are satisfied. Step 1 of the proof gives uniform convergence of (EE) scheme and step 2 gives the uniform convergence of the (SB) scheme.

5.3. Time reversal of gradient flows. Let us consider the time reversal of gradient flows of the functionals in the previous two examples. Specifically, if $(\rho(t), t \in [0, T])$ is the gradient flow of a functional $\mathcal{F} : \mathcal{P}_2(\mathbb{R}^d) \rightarrow (-\infty, \infty]$ with velocity field $(v(t), t \in [0, T])$, then let $(\bar{\rho}(t), t \in [0, T])$ denote the time reversed flow defined as $\bar{\rho}(t) := \rho(T-t)$. Naturally, the

velocity field in the reverse process is negative of the velocity in the forward process, i.e. if $(\bar{v}(t), t \in [0, T])$ denotes the velocity field of $(\bar{\rho}(t), t \in [0, T])$, then $\bar{v}(t) = -v(T - t)$.

These absolutely continuous curves are not gradient flows and therefore do not benefit from the theoretical guarantees established in our theorems. Note that the corresponding PDEs, such as the backward heat equation, are well known to be ill-posed ([Olv13, page 129, eqn. 4.29]). However, interestingly, we observe that, for the time reversed gradient flows of entropy and KL divergence functionals with Gaussian marginals, the (SB) scheme approximates these curves with the same convergence rates. For both the examples below, these convergence rates are derived by directly showing that $\text{SB}_\varepsilon^1(\cdot)$ is consistent (55) and a contraction (a stronger version of contraction in limit (56)). Following that, we use Theorem 3 to prove that the (SB) iterates give a first-order approximation of the time reversed flow $(\bar{\rho}(t), t \in [0, T])$. Through the examples, we also highlight a key issue with approximating reversed time gradient flows: to have a valid first-order approximation (Definition 1) of $(\bar{\rho}(t), t \in [0, T])$, the step size ε should be small enough where the bound on ε depends on the properties of the iterates encountered in the future. We include these calculations below for readers' curiosity and their practical importance in score-based generative modeling using diffusions [SSDK⁺20, LYB⁺24].

First, we consider the time reversed gradient flow of $\mathcal{F} = \frac{1}{2}\text{Ent}$. Then, $\bar{v}(t) = \frac{1}{2}\nabla \log \rho(T - t) = \frac{1}{2}\nabla \log \bar{\rho}(t)$. Now we calculate the first (SB) step. For $\rho = \rho(0)$, the surrogate measure is $\sigma = \sigma(0) = e^{\theta \log \bar{\rho}}$ where θ is chosen to ensure integrability of $e^{\theta \log \bar{\rho}}$. Therefore, we may choose $\theta = 1$, which implies $\sigma_\varepsilon(0) = \sigma = \rho$ and the (SB) step is $\text{SB}_\varepsilon^1(\rho) = (\mathcal{B}_{\rho, \varepsilon})_{\#} \rho$. Notice that the sign of θ is flipped compared to Section 5.1. This change ensures that the (SB) for the reverse flow is the opposite direction to the (SB) step of the forward flow. This is interesting as we can approximate both forward and reverse time gradient flow of entropy functional using appropriate direction of (SB) steps, calculated only using the current measure.

Now for an initial $\rho(0) = \mathcal{N}(0, \eta^2 I_d)$, we know that the gradient flow $(\rho(t), t \in [0, T])$ is given by $\rho(t) = \mathcal{N}(0, (\eta^2 + t)I_d)$. Then, we have $\rho = \bar{\rho}(0) = \mathcal{N}(0, (\eta^2 + T)I_d)$ and $\bar{\rho}(\varepsilon) = \mathcal{N}(0, (\eta^2 + T - \varepsilon)I_d)$. Now we show that $\text{SB}_\varepsilon^1(\rho)$ is an $\mathcal{O}(\varepsilon^2)$ close approximation of $\bar{\rho}(\varepsilon)$. For any $\eta^2 > 0$ and $\varepsilon < \eta^2 + T$,

$$\begin{aligned} \mathbb{W}_2(\bar{\rho}(\varepsilon), \text{SB}_\varepsilon^1(\rho)) &= d \left| C_{\eta^2+T}^\varepsilon \sqrt{\eta^2 + T} - \sqrt{\eta^2 + T - \varepsilon} \right| \\ &= d \sqrt{\eta^2 + T} \left| \sqrt{1 + \frac{\varepsilon^2}{4(\eta^2 + T)^2}} - \frac{\varepsilon}{2(\eta^2 + T)} - \sqrt{1 - \frac{\varepsilon}{\eta^2 + T}} \right|. \end{aligned}$$

Denote $x = \varepsilon/(\eta^2 + T)$, then the term containing ε above is $c_\varepsilon = \sqrt{1 + x^2/4} - (x/2) - \sqrt{1 - x}$. Using the identity $\sqrt{a} - \sqrt{b} = (a - b)/(\sqrt{a} + \sqrt{b})$ with $a = 1 + x^2/4$ and $b = \left((x/2) + \sqrt{1 - x} \right)^2$, we can be convinced that constant $c_\varepsilon > 0$. Further, using the identity $\sqrt{1 + a} < 1 + a/2$ for $a > 0$ and $\sqrt{1 - b} > 1 - b/2$ for $0 < b < 1$, we have that $c_\varepsilon < x^2/4 = \frac{\varepsilon^2}{4(\eta^2 + T)^2}$. Therefore, we have the consistency (55) result

$$(73) \quad \mathbb{W}_2(\bar{\rho}(\varepsilon), \text{SB}_\varepsilon^1(\rho)) < \frac{d\varepsilon^2}{4(\eta^2 + T)^{3/2}}.$$

Again, since (SB) scheme evolves via linear pushforwards, the sequence of (SB) steps are $\text{SB}_\varepsilon^k(\rho) = \mathcal{N}(0, \alpha_k^2 I_d)$ where α_k^2 follows the recursive relationship $\alpha_{k+1}^2 = \left(C_{\alpha_k^2}^\varepsilon \right)^2 \alpha_k^2$. Define the piecewise constant interpolation

$$\text{SB}_\varepsilon(t) = \text{SB}_\varepsilon^k(\rho) \quad t \in [k\varepsilon, (k+1)\varepsilon).$$

Unlike the previous two examples, since $(\bar{\rho}(t), t \in [0, T])$ is not a gradient flow, we do not invoke Theorem 6 to prove the uniform convergence of SB scheme. We instead directly show that $(\text{SB}_\varepsilon(t), t \in [0, T])$ is a first-order approximation (Definition 1) of $(\bar{\rho}(t), t \in [0, T])$. We have already shown consistency in (73). Now we show that $\text{SB}_\varepsilon^1(\cdot)$, restricted to Gaussian measures, is Lipschitz with the Lipschitz constant of the order $(1 + \mathcal{O}(\varepsilon))$. Take $\rho_1 = \mathcal{N}(0, \sigma_1^2 I_d)$ and $\rho_2 = \mathcal{N}(0, \sigma_2^2 I_d)$. Then, $\mathbb{W}_2(\text{SB}_\varepsilon^1(\rho_1), \text{SB}_\varepsilon^1(\rho_2))$ can be bounded above by the following string of inequalities

$$\begin{aligned}
&= d \left| \sigma_1 \sqrt{1 + \frac{\varepsilon^2}{4\sigma_1^4}} - \frac{\varepsilon}{2\sigma_1} - \sigma_2 \sqrt{1 + \frac{\varepsilon^2}{4\sigma_2^4}} + \frac{\varepsilon}{2\sigma_2} \right| \\
&\leq \frac{d\varepsilon}{2\sigma_1\sigma_2} |\sigma_1 - \sigma_2| + d \left| \sqrt{\sigma_1^2 + \frac{\varepsilon^2}{4\sigma_1^2}} - \sqrt{\sigma_2^2 + \frac{\varepsilon^2}{4\sigma_2^2}} \right| && \text{(triangle inequality)} \\
&= \frac{d\varepsilon}{2\sigma_1\sigma_2} |\sigma_1 - \sigma_2| + d \frac{|\sigma_1^2 - \sigma_2^2| \left| 1 - \frac{\varepsilon^2}{4\sigma_1^2\sigma_2^2} \right|}{\sqrt{\sigma_1^2 + \frac{\varepsilon^2}{4\sigma_1^2}} + \sqrt{\sigma_2^2 + \frac{\varepsilon^2}{4\sigma_2^2}}} && \left(\sqrt{a} - \sqrt{b} = \frac{a-b}{\sqrt{a} + \sqrt{b}} \right) \\
&\leq \frac{d\varepsilon}{2\sigma_1\sigma_2} |\sigma_1 - \sigma_2| + d \left| 1 - \frac{\varepsilon^2}{4\sigma_1^2\sigma_2^2} \right| |\sigma_1 - \sigma_2| && \left(\sqrt{\sigma_i^2 + \varepsilon^2/4\sigma_i^2} > \sigma_i \text{ for } i \in \{1, 2\} \right).
\end{aligned}$$

For $\varepsilon < 2\sigma_1\sigma_2$, $\left(1 - \frac{\varepsilon^2}{4\sigma_1^2\sigma_2^2}\right) > 0$ and hence we have the contraction

$$(74) \quad \mathbb{W}_2(\text{SB}_\varepsilon^1(\rho_1), \text{SB}_\varepsilon^1(\rho_2)) < \left(1 + \frac{\varepsilon}{2\sigma_1\sigma_2}\right) \mathbb{W}_2(\rho_1, \rho_2)$$

The sequence $(\alpha_k^2, k \in [N_\varepsilon])$ decreases monotonically because

$$\alpha_k^2 = \left(C_{\alpha_{k-1}^2}^\varepsilon\right)^2 \alpha_{k-1}^2 = \left(\sqrt{1 + \frac{\varepsilon^2}{4\alpha_{k-1}^4}} - \frac{\varepsilon}{2\alpha_{k-1}^2}\right)^2 \alpha_{k-1}^2.$$

and $\sqrt{1+x^2} - x < 1$ for all $x > 0$. Now we show that $(\alpha_k^2, k \in [N_\varepsilon])$ is bounded from below because for small enough ε . Using $\sqrt{1+x} \leq 1+x/2$,

$$\alpha_k^2 = \alpha_{k-1}^2 - \varepsilon \sqrt{1 + \frac{\varepsilon^2}{4\alpha_{k-1}^4}} + \frac{\varepsilon^2}{2\alpha_{k-1}^2} \geq \alpha_{k-1}^2 - \varepsilon + \frac{\varepsilon^2}{2\alpha_{k-1}^2} \left(1 - \frac{\varepsilon}{4\alpha_{k-1}^2}\right).$$

If ε is uniformly less than $4\alpha_{k-1}^2$ for all $k \in [N_\varepsilon]$, then $\alpha_k^2 \geq \alpha_{k-1}^2 - \varepsilon$. Consequently, the lowest value is $\alpha_{N_\varepsilon}^2 \geq (\eta^2 + T) - \varepsilon N_\varepsilon = \eta^2$. Therefore, choosing $\varepsilon < \eta^2$, we get that $(\alpha_k^2, k \in [N_\varepsilon])$ is bounded from below by η^2 . This highlights an inherent issue with approximating time-reversed gradient flows: a valid approximation depends on the choice of ε , which in turn depends on the variance of the flow at a future time. As a result, if we run an SB scheme with ε greater than the variance of $\bar{\rho}(T)$, we might encounter the problem of an SB step with a negative variance.

By the triangle inequality

$$\mathbb{W}_2(\bar{\rho}((k+1)\varepsilon), \text{SB}_\varepsilon^{k+1}(\rho)) \leq \mathbb{W}_2(\bar{\rho}((k+1)\varepsilon), \text{SB}_\varepsilon^1(\bar{\rho}(k\varepsilon))) + \mathbb{W}_2(\text{SB}_\varepsilon^1(\bar{\rho}(k\varepsilon)), \text{SB}_\varepsilon^{k+1}(\rho)).$$

Using consistency (73),

$$\mathbb{W}_2(\bar{\rho}((k+1)\varepsilon), \text{SB}_\varepsilon^1(\bar{\rho}(k\varepsilon))) < \frac{d\varepsilon^2}{4(\eta^2 + T - k\varepsilon)^{3/2}},$$

and using Lipschitzness (74),

$$\mathbb{W}_2 \left(\text{SB}_\varepsilon^1(\bar{\rho}(k\varepsilon)), \text{SB}_\varepsilon^{k+1}(\rho) \right) < \left(1 + \frac{\varepsilon}{2(\eta^2 + T - k\varepsilon)\alpha_k} \right) \mathbb{W}_2 \left(\bar{\rho}(k\varepsilon), \text{SB}_\varepsilon^k(\rho) \right).$$

Since for all $k \in [N_\varepsilon]$, $\alpha_k^2 \geq \eta^2$, we have

$$\mathbb{W}_2 \left(\bar{\rho}((k+1)\varepsilon), \text{SB}_\varepsilon^{k+1}(\rho) \right) < \frac{d\varepsilon^2}{4\eta^3} + \left(1 + \frac{\varepsilon}{2\eta^2} \right) \mathbb{W}_2 \left(\bar{\rho}(k\varepsilon), \text{SB}_\varepsilon^k(\rho) \right).$$

Recursively,

$$\mathbb{W}_2 \left(\bar{\rho}(k\varepsilon), \text{SB}_\varepsilon^k(\rho) \right) \leq \frac{d\varepsilon}{2\eta} \left(1 + \frac{\varepsilon}{2\eta^2} \right)^{N_\varepsilon}.$$

Since $N_\varepsilon = \lfloor T\varepsilon^{-1} \rfloor$, therefore, $\lim_{\varepsilon \rightarrow 0} \sup_{k \in [N_\varepsilon]} \mathbb{W}_2 \left(\bar{\rho}(k\varepsilon), \text{SB}_\varepsilon^k(\rho) \right) = 0$, proving the uniform convergence of $\left(\text{SB}_\varepsilon^k(\rho), k \in [N_\varepsilon] \right)$ to the curve $(\bar{\rho}(t), t \in [0, T])$ as $\varepsilon \rightarrow 0+$.

Finally, we consider the case of time reversed case of gradient flow of $\mathcal{F} = \frac{1}{2}H(\cdot|\nu)$ with $\nu = e^{-g} = \mathcal{N}(0, I_d)$. Since this case is very similar to the previous examples on the gradient flow of KL divergence and time reversed gradient flow of entropy, we limit explicit calculations. The velocity field $\bar{v}(t) = \frac{1}{2}(\text{Id} + \nabla \log \bar{\rho}(t))$. For an initial choice $\rho(0) = \mathcal{N}(0, \eta^2)$, the time reversed gradient flow, starting from time $t = T$, is $\bar{\rho}(t) = \mathcal{N} \left(0, \left(\eta^2 e^{-(T-t)} + (1 - e^{-(T-t)}) \right) I_d \right)$.

For brevity of notation, let $\rho = \bar{\rho}(0) = \mathcal{N}(0, \tau^2)$ where $\tau^2 := \eta^2 e^{-T} + (1 - e^{-T})$. Now we will calculate the (SB) step. Again, we choose θ such that the surrogate measure $\sigma = \sigma(0) = (\rho e^g)^\theta$ is integrable. Using the same logic as Section 5.2, if $\eta^2 > 1$ (equivalently $\tau^2 > 1$), then θ may be taken to be -1 giving $\sigma = \rho^{-1} e^{-g} = \mathcal{N}(0, (\tau^2/(\tau^2 - 1))I_d)$ and $\text{SB}_\varepsilon^1(\rho) = (2\text{Id} - \mathcal{B}_{\sigma, \varepsilon})_\# \rho$. Whereas if $\eta^2 < 1$ (equivalently $\tau^2 < 1$), then θ can be taken to be 1 giving $\sigma = \mathcal{N}(0, (\tau^2/(1 - \tau^2))I_d)$ and $\text{SB}_\varepsilon^1(\rho) = (\mathcal{B}_{\sigma, \varepsilon})_\# \rho$. We will present calculations for $\eta^2 < 1$ case. Similar calculations follow for $\eta^2 > 1$ and have been skipped.

Now we prove that the (SB) step is an $\mathcal{O}(\varepsilon^2)$ approximation of $\bar{\rho}(\varepsilon)$. Note that for $\tau^2 < 1$ and $\varepsilon < \log((1 - \tau^2)^{-1})$, $\text{SB}_\varepsilon^1(\rho) = \mathcal{N} \left(0, \left(C_{\tau^2/(1-\tau^2)}^\varepsilon \right)^2 \tau^2 \right)$ and $\bar{\rho}(\varepsilon) = \mathcal{N} \left(0, \tau^2 + (\tau^2 - 1)(e^\varepsilon - 1) \right)$.

Then

$$\mathbb{W}_2 \left(\bar{\rho}(\varepsilon), \text{SB}_\varepsilon^1(\rho) \right) = d\tau \left| \sqrt{1 + \frac{\varepsilon^2(1 - \tau^2)^2}{4\tau^4}} - \frac{\varepsilon(1 - \tau^2)}{2\tau^2} - \sqrt{1 - \frac{1 - \tau^2}{\tau^2}(e^\varepsilon - 1)} \right|.$$

Denote $x = \frac{\varepsilon(1 - \tau^2)}{\tau^2}$, then the term containing ε above is $c_\varepsilon = \sqrt{1 + \frac{x^2}{4}} - \frac{x}{2} - \sqrt{1 - x \left(\frac{e^\varepsilon - 1}{\varepsilon} \right)}$.

It is easy to see that $c_\varepsilon > 0$. Further, using the identity $\sqrt{1 + a} < 1 + a/2$ for $a > 0$ and $\sqrt{1 - b} > 1 - b/2$ for $0 < b < 1$, we have that $c_\varepsilon < \frac{x}{2} \left(\frac{e^\varepsilon - 1}{\varepsilon} \right) + \frac{x^2}{8}$. Therefore, we have the consistency (55) result

$$(75) \quad \mathbb{W}_2 \left(\bar{\rho}(\varepsilon), \text{SB}_\varepsilon^1(\rho) \right) < d\varepsilon^2 \left(\frac{(1 - \tau^2)^2}{8\tau^3} + \frac{C(1 - \tau^2)}{2\tau} \right) \quad \text{for some constant } C.$$

Again since the (SB) scheme evolves via linear pushforwards, $\text{SB}_\varepsilon^k(\rho)$ is a Gaussian measure for all $k \in [N_\varepsilon]$ and let $\text{SB}_\varepsilon^k(\rho) = \mathcal{N}(0, \alpha_k^2 I_d)$. Then α_k^2 evolves via the recursive relationship $\alpha_{k+1}^2 = \left(C_{\alpha_k^2/(1-\alpha_k^2)}^\varepsilon \right)^2 \alpha_k^2$. We prove uniform convergence of the SB scheme in the same way as the time reversed flow of entropy functional. The consistency is show above and now

we show that the (SB) is a contraction for Gaussian measures. Take $\rho_1 = \mathcal{N}(0, \sigma_1^2 I_d)$ and $\rho_2 = \mathcal{N}(0, \sigma_2^2 I_d)$. Then, exactly as in the case of time-reversed gradient flow of entropy,

$$\mathbb{W}_2 \left(\text{SB}_\varepsilon^1(\rho_1), \text{SB}_\varepsilon^1(\rho_2) \right) < \frac{d\varepsilon}{2} \left(1 + \frac{1}{\sigma_1 \sigma_2} \right) |\sigma_1 - \sigma_2| + d \left| 1 - \frac{\varepsilon^2}{4} \left(\frac{1 - \sigma_1^2 \sigma_2^2}{\sigma_1^2 \sigma_2^2} \right) \right| |\sigma_1 - \sigma_2|.$$

For $\varepsilon < \frac{2\sigma_1 \sigma_2}{\sqrt{1 - \sigma_1^2 \sigma_2^2}}$, $\left(1 - \frac{\varepsilon^2}{4} \left(\frac{1 - \sigma_1^2 \sigma_2^2}{\sigma_1^2 \sigma_2^2} \right) \right) > 0$ and hence we have the contraction

$$(76) \quad \mathbb{W}_2 \left(\text{SB}_\varepsilon^1(\rho_1), \text{SB}_\varepsilon^1(\rho_2) \right) < \left(1 + \frac{\varepsilon}{2} \left(1 + \frac{1}{\sigma_1 \sigma_2} \right) \right) \mathbb{W}_2(\rho_1, \rho_2).$$

Because $C_\eta^\varepsilon \leq 1$ for all $\eta^2 > 0$, the sequence $(\alpha_k^2; k \in [N_\varepsilon])$ is monotonically decreasing and bounded from above by its initial value τ^2 . Similar to the reversed time gradient flow of entropy, we get that the variance $(\alpha_k^2; k \in [N_\varepsilon])$ is bounded from below if ε is less than $\frac{4\eta^2}{1 - \eta^2}$, which depends on the variance at time $t = T$. Using the triangular argument from Theorem 3, for any $k \in [N_\varepsilon]$,

$$\mathbb{W}_2 \left(\bar{\rho}(k\varepsilon), \text{SB}_\varepsilon^k(\rho) \right) \leq d\varepsilon \frac{(1 + 4\eta^2 C)}{4\eta(\eta^2 + 1)} \left(1 + \frac{(1 + \eta^{-2})\varepsilon}{2} \right)^{N_\varepsilon}.$$

Therefore, $\lim_{\varepsilon \rightarrow 0} \sup_{k \in [N_\varepsilon]} \mathbb{W}_2 \left(\bar{\rho}(k\varepsilon), \text{SB}_\varepsilon^k(\rho) \right) = 0$, proving the uniform convergence of $(\text{SB}_\varepsilon^k(\rho); k \in [N_\varepsilon])$ to the curve $(\bar{\rho}(t); t \in [0, T])$ as $\varepsilon \rightarrow 0+$.

REFERENCES

- [AGS08] Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. *Gradient flows in metric spaces and in the space of probability measures*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, second edition, 2008.
- [BGL14] Dominique Bakry, Ivan Gentil, and Michel Ledoux. *Analysis and geometry of Markov diffusion operators*, volume 348 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Cham, 2014.
- [BGN22] Espen Bernton, Promit Ghosal, and Marcel Nutz. Entropic optimal transport: geometry and large deviations. *Duke Math. J.*, 171(16):3363–3400, 2022.
- [Bob22] Sergey G. Bobkov. Upper bounds for Fisher information. *Electron. J. Probab.*, 27:Paper No. 115, 44, 2022.
- [CCGT22] Alberto Chiarini, Giovanni Conforti, Giacomo Greco, and Luca Tamanini. Gradient estimates for the schrödinger potentials: convergence to the brenier map and quantitative stability, 2022.
- [CG14] P. Cattiaux and A. Guillin. Semi log-concave Markov diffusions. In *Séminaire de Probabilités XLVI*, volume 2123 of *Lecture Notes in Math.*, pages 231–292. Springer, Cham, 2014.
- [Csi75] I. Csiszár. I -divergence geometry of probability distributions and minimization problems. *Ann. Probability*, 3:146–158, 1975.
- [CT21] Giovanni Conforti and Luca Tamanini. A formula for the time derivative of the entropic cost and applications. *J. Funct. Anal.*, 280(11):Paper No. 108964, 48, 2021.
- [Cut13] Marco Cuturi. Sinkhorn distances: Lightspeed computation of optimal transport. *Advances in neural information processing systems*, 26, 2013.
- [CVR18] Giovanni Conforti and Max Von Renesse. Couplings, gradient estimates and logarithmic Sobolev inequality for Langevin bridges. *Probab. Theory Related Fields*, 172(1-2):493–524, 2018.
- [DGW04] H. Djellout, A. Guillin, and H. Wu. Transportation cost-information inequalities and applications to random dynamical systems and diffusions. *Ann. Probab.*, 32(3B):2702–2732, 2004.
- [GLPR24] Borjan Geshkovski, Cyril Letrouit, Yury Polyanskiy, and Philippe Rigollet. The emergence of clusters in self-attention dynamics. Arxiv preprint 2305.05465, 2024.
- [GLR17] Ivan Gentil, Christian Léonard, and Luigia Ripani. About the analogy between optimal transport and minimal entropy. *Ann. Fac. Sci. Toulouse Math. (6)*, 26(3):569–601, 2017.
- [GLRT20] Ivan Gentil, Christian Léonard, Luigia Ripani, and Luca Tamanini. An entropic interpolation proof of the HWI inequality. *Stochastic Process. Appl.*, 130(2):907–923, 2020.

- [JKO98] Richard Jordan, David Kinderlehrer, and Felix Otto. The variational formulation of the Fokker-Planck equation. *SIAM J. Math. Anal.*, 29(1):1–17, 1998.
- [JMPC20] Hicham Janati, Boris Muzellec, Gabriel Peyré, and Marco Cuturi. Entropic optimal transport between unbalanced Gaussian measures has a closed form. NIPS’20, Red Hook, NY, USA, 2020. Curran Associates Inc.
- [KS91] Ioannis Karatzas and Steven E. Shreve. *Brownian motion and stochastic calculus*, volume 113 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1991.
- [Léo12] Christian Léonard. From the Schrödinger problem to the Monge-Kantorovich problem. *J. Funct. Anal.*, 262(4):1879–1920, 2012.
- [Léo14] Christian Léonard. A survey of the Schrödinger problem and some of its connections with optimal transport. *Discrete Contin. Dyn. Syst.*, 34(4):1533–1574, 2014.
- [LK93] B. C. Levy and A. J. Krener. Dynamics and kinematics of reciprocal diffusions. *Journal of mathematical physics*, 34(5):1846–1875, 1993.
- [LYB⁺24] Sungbin Lim, Eun Bi Yoon, Taehyun Byun, Taewon Kang, Seungwoo Kim, Kyungjae Lee, and Sungjoon Choi. Score-based generative modeling through stochastic evolution equations in Hilbert spaces. *Advances in Neural Information Processing Systems*, 36, 2024.
- [Olv13] P.J. Olver. *Introduction to Partial Differential Equations*. Undergraduate Texts in Mathematics. Springer International Publishing, 2013.
- [Pal19] Soumik Pal. On the difference between entropic cost and the optimal transport cost, 2019. Arxiv 1905.12206. To appear in *The Annals of Applied Probability*.
- [PNW22] Aram-Alexandre Pooladian and Jonathan Niles-Weed. Entropic estimation of optimal transport maps, 2022.
- [Roy07] Gilles Royer. *An initiation to logarithmic Sobolev inequalities*, volume 14 of *SMF/AMS Texts and Monographs*. American Mathematical Society, Providence, RI; Société Mathématique de France, Paris, 2007. Translated from the 1999 French original by Donald Babbitt.
- [RY04] D. Revuz and M. Yor. *Continuous Martingales and Brownian Motion*. Grundlehren der mathematischen Wissenschaften. Springer Berlin Heidelberg, 2004.
- [SABP22] Michael E. Sander, Pierre Ablin, Mathieu Blondel, and Gabriel Peyré. Sinkformers: Transformers with doubly stochastic attention. In Gustau Camps-Valls, Francisco J. R. Ruiz, and Isabel Valera, editors, *Proceedings of The 25th International Conference on Artificial Intelligence and Statistics*, volume 151 of *Proceedings of Machine Learning Research*, pages 3515–3530. PMLR, 28–30 Mar 2022.
- [San15] Filippo Santambrogio. *Optimal transport for applied mathematicians: Calculus of variations, PDEs, and modeling*, volume 87. Springer, 2015.
- [SSDK⁺20] Yang Song, Jascha Sohl-Dickstein, Diederik P Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations*, 2020. ArXiv version 2011.13456.
- [Ver18] Roman Vershynin. *High-dimensional probability*, volume 47 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 2018. An introduction with applications in data science, With a foreword by Sara van de Geer.
- [Vil03] Cédric Villani. *Topics in optimal transportation*, volume 58. American Mathematical Soc., 2003.

NOTATIONS

MEDHA AGARWAL, DEPARTMENT OF STATISTICS, UNIVERSITY OF WASHINGTON, SEATTLE WA 98195, USA,
EMAIL: MEDHAAGA@UW.EDU

ZAID HARCHAOU, DEPARTMENT OF STATISTICS, UNIVERSITY OF WASHINGTON, SEATTLE WA 98195, USA,
EMAIL: ZAID@UW.EDU

GARRETT MULCAHY, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON, SEATTLE WA 98195,
USA, EMAIL: GMULCAHY@UW.EDU

SOUMIK PAL, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON, SEATTLE WA 98195, USA,
EMAIL: SOUMIK@UW.EDU

Symbol	Explanation
$C^d[0, \infty)$	Collection of continuous paths $\omega : [0, \infty) \rightarrow \mathbb{R}^d$
$C^k(\mathbb{R}^d)$	k -times continuously differentiable functions $\mathbb{R}^d \rightarrow \mathbb{R}$
$\mathcal{P}_2(\mathbb{R}^d)$	Space of twice integrable probability measures on \mathbb{R}^d
$\mathcal{P}_2^{ac}(\mathbb{R}^d)$	Space of twice integrable and absolutely continuous probability measures (with respect to Lebesgue measure) on \mathbb{R}^d
$C_c^\infty(\mathbb{R}^d)$	Space of smooth and compactly supported functions in \mathbb{R}^d
$W_2(\mu, \nu)$	Wasserstein-2 distance between measures μ and ν in $\mathcal{P}_2(\mathbb{R}^d)$
T_μ^ν	Optimal transport map from μ to ν (if it exists)
$(\pi_t, t \geq 0)$	Coordinatewise projections on $C^d[0, \infty)$, $\pi_s(\omega) = \omega_s$
$\mu_{\rho, \varepsilon}$	$\mu_{\rho, \varepsilon}(dx, dy) = (2\pi\varepsilon)^{-d/2} \rho(x) \exp(-\ x - y\ ^2/2\varepsilon) dx dy$
$(W_x, x \in \mathbb{R}^d)$	Wiener measure on $C^d[0, \infty)$ with initial condition δ_x
W	Reversible Wiener measure on $C^d[0, \infty)$
R_ε	Law of reversible Brownian motion with diffusion ε
$(P_t, t \geq 0)$	Brownian semigroup
$(p_t(\cdot, \cdot), t > 0)$	Brownian transition densities
Q	Law of stationary Langevin diffusion on $C^d[0, \infty)$
$\ell_{\rho, \varepsilon}$	Joint law of Langevin diffusion at time 0 and ε , i.e. $(\pi_0, \pi_\varepsilon)_\# Q$
$(q_t(\cdot, \cdot), t > 0)$	Transition densities for Langevin diffusion
L	Extended generator for the Langevin diffusion
$(G_t, t \geq 0)$	Langevin semigroup
$\pi_{\rho, \varepsilon}$	ε -static Schrödinger bridge with marginals equal to ρ
$(f_\varepsilon, \varepsilon > 0)$	Entropic potentials associated to $(\pi_{\rho, \varepsilon}, \varepsilon > 0)$
$\mathcal{B}_{\rho, \varepsilon}$	Barycentric projection associated with $\pi_{\rho, \varepsilon}$, $\mathcal{B}_{\rho, \varepsilon}(x) = E_{\pi_{\rho, \varepsilon}}[Y X = x]$
$S_{\rho, \varepsilon}$	Law of ε -dynamic Schrödinger bridge on $C^d[0, 1]$ with initial and terminal distribution ρ
$(\rho_t^\varepsilon, t \in [0, 1])$	Entropic interpolation from ρ to itself, $\rho_t^\varepsilon = (\pi_t)_\# S_{\rho, \varepsilon}$
$\ \cdot\ _{\psi_1}$	Subexponential norm of a random variable, [Ver18, Definition 2.7.5]
$\text{Tan}(\mu)$	Tangent bundle of $\mathcal{P}_2(\mathbb{R}^d)$ at μ
$\text{Ent}(\mu)$	Entropy of $\mu \in \mathcal{P}_2^{ac}(\mathbb{R}^d)$
$H(\mu \nu)$	Kullback-Leibler divergence between μ and ν
$I(\rho)$	Fisher information of ρ , $I(\rho) = E_\rho \ \nabla \log \rho\ ^2$ for $\rho \in \mathcal{P}_2^{ac}(\mathbb{R}^d)$
$\text{SB}_\varepsilon^k(\mu_0)$	k th iterate of Schrödinger bridge scheme starting from μ_0 with stepsize ε
$S_\varepsilon^k(\mu_0)$	k th iterate of pushforward of form $(\text{Id} + \varepsilon v_\varepsilon)_\# \mu_0$, where each $v_\varepsilon : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a vector field specified in context