Embedding formalism for AdS superspaces in five dimensions

Nowar E. Koning and Sergei M. Kuzenko

Department of Physics M013, The University of Western Australia 35 Stirling Highway, Perth W.A. 6009, Australia

Email: nowar.koning@research.uwa.edu.au, sergei.kuzenko@uwa.edu.au

Abstract

The standard geometric description of d-dimensional anti-de Sitter (AdS) space is a quadric in $\mathbb{R}^{d-1,2}$ defined by $(X^0)^2 - (X^1)^2 - \cdots - (X^{d-1})^2 + (X^d)^2 = \ell^2 = \text{const.}$ In this paper we provide a supersymmetric generalisation of this embedding construction in the d = 5 case. Specifically, a bi-supertwistor realisation is given for the \mathcal{N} -extended AdS superspace AdS^{5|8 \mathcal{N}}, with $\mathcal{N} \geq 1$. The proposed formalism offers a simple construction of AdS super-invariants. As an example, we present a model for a massive superparticle in AdS^{5|8 \mathcal{N}} which is manifestly invariant under the AdS isometry supergroup $\mathsf{SU}(2,2|\mathcal{N})$.

Contents

1	Introduction	2
2	\mathbf{AdS}_5	4
	2.1 Algebraic background	. 4
	2.2 Twistor realisation	. 5
	2.3 Bi-twistor realisation	. 6
3	\mathcal{N} -extended AdS superspace in five dimensions	8
	3.1 Algebraic background	. 8
	3.2 Supertwistor realisation	. 9
	3.3 Stabiliser	. 10
	3.4 Bi-supertwistor realisation	. 11
	3.5 $SU(2,2 \mathcal{N})$ -invariant two-point function	. 13
4	Coset representative	14
	4.1 AdS space $(\mathcal{N} = 0)$. 14
	4.2 $\mathcal{N} \neq 0$. 16
5	Superspace geometry	18
	5.1 Geometric structures in $AdS^{5 8N}$. 18
	5.2 Covariant derivatives	. 22
6	AdS superspaces with internal bosonic dimensions	23
	6.1 The $\mathcal{N} = 4$, $m = 4$ case	24
7	Conclusion	27

A Spinors in diverse dimensions 27 A.1 Spinors in 3 + 1 dimensions 27 A.2 Spinors in 4 + 2 dimensions 27 A.3 Spinors in 4 + 1 dimensions 35

1 Introduction

As is well-known, the algebraic underpinning for the AdS/CFT duality is the dual role played by the group $SO_0(d-1,2)$. On the one hand, it is the connected conformal group of the (d-1)-dimensional compactified Minkowski space $\overline{\mathbb{M}}^{d-1} = (S^{d-2} \times S^1)/\mathbb{Z}_2$.¹ On the other hand, it is the connected component of the isometry group of AdS_d , the *d*-dimensional AdS space. The situation in the supersymmetric world is analogous modulo the fact that superconformal groups exist only in six or less space-time dimensions [1]. In particular, for d = 5 the supergroup $SU(2, 2|\mathcal{N})$ is realised either (i) as the superconformal group in four dimensions; or (ii) as the \mathcal{N} -extended AdS supergroup in five dimensions.

From the group-theoretic point of view, an \mathcal{N} -extended AdS superspace in five dimensions is defined as

$$AdS^{5|8\mathcal{N}} = \frac{\mathsf{SU}(2,2|\mathcal{N})}{\mathsf{Spin}(4,1) \times \mathsf{U}(1) \times \mathsf{SU}(\mathcal{N})},\tag{1.1}$$

see, e.g., [2]. In this family of coset spaces, the only conformally flat superspace is

$$AdS^{5|8} = \frac{SU(2,2|1)}{Spin(4,1) \times U(1)} , \qquad (1.2)$$

as was argued in [2].² The $\mathcal{N} = 1$ case is exceptional since there is a unique superconformal algebra in five dimensions [1], the so-called F(4) superalgebra discovered by Kac [5], corresponding to $\mathcal{N} = 1$. This implies that there is only one type of d = 5 superconformal tensor calculus which was developed independently by two groups: Fujita, Kugo, and Ohashi [6–9] and Bergshoeff et al. [10–12]. Finally, there is only one type of d = 5 conformal superspace [13].

¹Strictly speaking, the connected conformal group should be identified with $SO_0(d-1,2)/\mathbb{Z}_2$ if d is odd.

²No explicit construction of a conformally flat $\mathcal{N} = 1$ AdS superspace geometry was given in [2]. Such a construction was derived in [3] by making use of the superspace formulation for the 5D $\mathcal{N} = 1$ Weyl supermultiplet developed in [4].

The simple AdS superspace, eq. (1.2), is a maximally symmetric background of the minimal $\mathcal{N} = 1$ supergravity geometry sketched by Howe [14] (see also [15]) and fully developed in [16,17]. It was shown in [13,18] that AdS^{5|8} is the only maximally supersymmetric solution in the off-shell gauged $\mathcal{N} = 1$ supergravity with the following compensators: (i) the vector multiplet; and (ii) the $\mathcal{O}(2)$ multiplet. Supersymmetric nonlinear σ -models in AdS₅ have interesting off-shell [19] and geometric properties [20].

In the non-supersymmetric case, there exist two different but equivalent realisations of AdS_d : (i) as the coset space O(d-1,2)/O(d-1,1); and (ii) as a hypersurface in $\mathbb{R}^{d-1,2}$

$$-(X^{0})^{2} + (X^{1})^{2} + \dots + (X^{d-1})^{2} - (X^{d})^{2} = -\ell^{2} = \text{const}.$$
 (1.3)

Both realisations have found numerous applications in the literature. Supersymmetric analogues of (1.3) have recently been developed in three and four dimensions, see [21–23]. For the d = 5 case, a bi-twistor realisation for (complexified) AdS₅ also exists, see, e.g., [24, 25]. To the best of our knowledge, a supersymmetric extension of (1.3) in five dimensions is yet to be developed in the literature, although a similar construction exists in the context of superstrings on AdS₅ × S⁵ [26]. In this paper we develop the superembedding formalism³ for AdS^{5|8N}.

Since the work by Ferber [29], supertwistors have found numerous applications in theoretical and mathematical physics. In particular, supertwistor realisations of compactified \mathcal{N} -extended Minkowski superspaces have been developed in four [30,31] and three [32,33] dimensions, and their harmonic/projective extensions have been derived [32–40].⁴ Supertwistor realisations of AdS^(3|p,q) and AdS^{4|4 \mathcal{N}} were provided in [21]. Recently, supertwistor formulations for conformal supergravity theories in diverse dimensions have been proposed [42, 43]. Furthermore, (super)twistor descriptions of (super)particles in AdS spaces have been studied extensively in the literature [44–54].

This paper is organised as follows. In section 2 we begin by introducing the twistor realisation for AdS_5 . Following that, we recall the standard bi-twistor realisation and prove their equivalence. In section 3 we describe a supertwistor realisation for the \mathcal{N} -extended AdS superspace in five dimensions, $AdS^{5|8\mathcal{N}}$. We then introduce a bi-supertwistor realisation for the same superspace and prove their equivalence. Finally we discuss an invariant two-point function on $AdS^{5|8\mathcal{N}}$. In section 4 we develop coset representatives for both the bosonic and supersymmetric cases, and discuss their corresponding coordinate systems. Section 5 is devoted to the study of the geometry of $AdS^{5|8\mathcal{N}}$. Section 6 introduces a family of homogeneous superspaces

³For a pedagogical review of superembeddings see [27, 28].

⁴Recently, there has been an interesting study of the geometric nature of the points at infinity of Minkowski superspace [41].

of the form $\operatorname{AdS}^{5|8\mathcal{N}} \times \mathbb{X}_m^{\mathcal{N}}$, where the internal space $\mathbb{X}_m^{\mathcal{N}}$ is realised in terms of $m \leq \mathcal{N}$ odd supertwistors. The main body of the paper is accompanied by appendix A, which outlines our spinor conventions in diverse dimensions.

2 $\quad \mathbf{AdS}_5$

As is well-known, compactified Minkowski space $\overline{\mathbb{M}}^4 = (S^3 \times S^1)/\mathbb{Z}_2$ can be realised [55–58] as the set of null planes in the twistor space \mathfrak{T} which is a four-dimensional complex vector space, $\mathfrak{T} = \mathbb{C}^4$, endowed with the linear action of the conformal group $\mathsf{SU}(2,2)$,

$$\mathsf{SU}(2,2) = \left\{ g \in \mathsf{SL}(4,\mathbb{C}), \quad g^{\dagger}\Omega g = \Omega, \quad \Omega = \begin{pmatrix} 0 & \mathbb{1}_2 \\ \mathbb{1}_2 & 0 \end{pmatrix} \right\}$$
(2.1)

Twistors can also be used to describe AdS_5 .

2.1 Algebraic background

A twistor $T \in \mathfrak{T}$ is a column vector

$$T = (T_{\hat{\alpha}}) = \begin{pmatrix} f_{\alpha} \\ \bar{g}^{\dot{\alpha}} \end{pmatrix}, \qquad \alpha = 1, 2, \quad \dot{\alpha} = \dot{1}, \dot{2}.$$
(2.2)

The group SU(2,2) naturally acts on \mathfrak{T} by transformations

$$T_{\hat{\alpha}} \longrightarrow g_{\hat{\alpha}}{}^{\hat{\beta}}T_{\hat{\beta}}, \qquad g = \left(g_{\hat{\alpha}}{}^{\hat{\beta}}\right) \in \mathsf{SU}(2,2) .$$
 (2.3)

A unique SU(2,2) invariant inner product on \mathfrak{T} is given by

$$\langle T|S\rangle := T^{\dagger}\Omega S \equiv \bar{T}S , \qquad (2.4)$$

where we have defined

$$\bar{T} = T^{\dagger}\Omega = (\bar{T}^{\hat{\alpha}}), \qquad \bar{T}^{\hat{\alpha}} = (T_{\hat{\beta}})^* \,\Omega^{\hat{\beta}\hat{\alpha}} ,\qquad \Omega = \left(\Omega^{\hat{\alpha}\hat{\beta}}\right). \tag{2.5}$$

We will refer to \overline{T} as the dual of T. It transforms as

$$\bar{T}^{\hat{\alpha}} \longrightarrow \bar{T}^{\hat{\beta}}(g^{-1})_{\hat{\beta}}{}^{\hat{\alpha}}.$$
(2.6)

The space of dual twistors will be denoted $\overline{\mathfrak{T}}$.

In what follows, of special importance for our analysis will be a subgroup of SU(2, 2), denoted USp(2, 2), consisting of those group elements which satisfy the additional condition

$$g^{\mathrm{T}}Cg = C, \qquad C = (\varepsilon^{\hat{\alpha}\hat{\beta}}) = \begin{pmatrix} \varepsilon^{\alpha\beta} & 0\\ 0 & -\varepsilon_{\dot{\alpha}\dot{\beta}} \end{pmatrix},$$
 (2.7)

with $\varepsilon^{\alpha\beta}$ and $\varepsilon_{\dot{\alpha}\dot{\beta}}$ being defined in (A.8). The Lie algebras of SU(2,2) and USp(2,2) consist of elements of the form (A.48) and (A.63), respectively, and are denoted $\mathfrak{su}(2,2)$ and $\mathfrak{usp}(2,2)$.

2.2 Twistor realisation

Now we turn to describing the twistor realisation for AdS_5 . Let us introduce a set of four linearly independent twistors

$$T^{\tilde{\mu}} = (T_{\hat{\alpha}}^{\tilde{\mu}}), \qquad \tilde{\mu} = 1, 2, 3, 4.$$
 (2.8)

We further restrict our attention to those quartets $T^{\tilde{\mu}}$ that satisfy the constraints

$$(T_{\hat{\alpha}}{}^{\tilde{\mu}})^* \Omega^{\hat{\alpha}\hat{\beta}} T_{\hat{\beta}}{}^{\tilde{\nu}} = \Omega^{\tilde{\mu}\tilde{\nu}}, \qquad (\Omega^{\tilde{\mu}\tilde{\nu}}) := \begin{pmatrix} 0 & \mathbb{1}_2\\ \mathbb{1}_2 & 0 \end{pmatrix}, \qquad (2.9a)$$

$$\det\left(T_{\hat{\alpha}}^{\tilde{\mu}}\right) = 1.$$
(2.9b)

The space of all such quartets will be denoted \mathfrak{F} .

On the space \mathfrak{F} , we introduce the equivalence relation

$$T_{\hat{\alpha}}{}^{\tilde{\mu}} \sim T_{\hat{\alpha}}{}^{\tilde{\nu}}\lambda_{\tilde{\nu}}{}^{\tilde{\mu}}, \qquad \lambda \in \mathsf{USp}(2,2).$$
 (2.10)

We emphasise that the constraints (2.9) are invariant under the equivalence transformations (2.10). The group SU(2,2) acts on the space \mathfrak{F} as

$$T_{\hat{\alpha}}{}^{\tilde{\mu}} \to g_{\hat{\alpha}}{}^{\hat{\beta}}T_{\hat{\beta}}{}^{\tilde{\mu}}, \qquad g \in \mathsf{SU}(2,2).$$

$$(2.11)$$

This action is naturally extended to the quotient space \mathfrak{F}/\sim . The latter proves to be a homogeneous space for the group SU(2,2). It turns out that

$$\operatorname{AdS}_5 = \mathfrak{F}/\sim .$$
 (2.12)

To prove (2.12), we introduce the bi-twistor

$$X_{\hat{\alpha}\hat{\beta}} := \ell \mathfrak{C}_{\tilde{\mu}\tilde{\nu}} T_{\hat{\alpha}}{}^{\tilde{\mu}} T_{\hat{\beta}}{}^{\tilde{\nu}} = -X_{\hat{\beta}\hat{\alpha}}, \qquad (2.13)$$

where the matrix \mathfrak{C} is defined as

$$\mathfrak{C} = (\mathfrak{C}^{\tilde{\mu}\tilde{\nu}}) = \begin{pmatrix} \varepsilon & 0\\ 0 & -\varepsilon^{-1} \end{pmatrix}, \qquad \mathfrak{C}^{-1} = (\mathfrak{C}_{\tilde{\mu}\tilde{\nu}}), \qquad \varepsilon = \begin{pmatrix} 0 & 1\\ -1 & 0 \end{pmatrix}.$$
(2.14)

We present the matrix \mathfrak{C} in this form for later convenience. The bi-twistor $X_{\hat{\alpha}\hat{\beta}}$ is invariant under the equivalence relation (2.10), and as such can be used to parametrise the space (2.12).

As shown in appendix A.2, a one-to-one correspondence exists between complex vectors $V^{\underline{a}}$ in 4+2 dimensions and antisymmetric bi-twistors $V_{\hat{\alpha}\hat{\beta}} = -V_{\hat{\beta}\hat{\alpha}}$. It is defined by the rule (A.51),

$$V^{\underline{a}} \rightarrow V_{\hat{\alpha}\hat{\beta}} = V^{\underline{a}}(\Sigma_{\underline{a}})_{\hat{\alpha}\hat{\beta}},$$
 (2.15)

where the matrices $\Sigma_{\underline{a}}$ are given in (A.45). The six-vector corresponding to (2.13),

$$X^{\underline{a}} = \frac{1}{4} (\tilde{\Sigma}^{\underline{a}})^{\hat{\alpha}\hat{\beta}} X_{\hat{\alpha}\hat{\beta}} , \qquad (2.16)$$

is real, because the bi-twistor $X_{\hat{\alpha}\hat{\beta}}$ satisfies the reality condition,

$$\bar{X}^{\hat{\alpha}\hat{\beta}} := \Omega^{\hat{\alpha}\hat{\gamma}} (X^{\dagger})_{\hat{\gamma}\hat{\delta}} \Omega^{\hat{\delta}\hat{\beta}} = \frac{1}{2} \varepsilon^{\hat{\alpha}\hat{\beta}\hat{\gamma}\hat{\delta}} X_{\hat{\gamma}\hat{\delta}} , \qquad (2.17)$$

see (A.55b). In order to prove (2.17) we use the identities

$$T_{\hat{\alpha}}{}^{\tilde{\mu}}T_{\hat{\beta}}{}^{\tilde{\nu}}T_{\hat{\gamma}}{}^{\tilde{\rho}}T_{\hat{\delta}}{}^{\tilde{\sigma}}\varepsilon^{\hat{\alpha}\hat{\beta}\hat{\gamma}\hat{\delta}} = \varepsilon^{\tilde{\mu}\tilde{\nu}\tilde{\rho}\tilde{\sigma}}, \qquad (2.18a)$$

$$\mathfrak{C}^{\tilde{\mu}\tilde{\nu}}\mathfrak{C}^{\tilde{\rho}\tilde{\sigma}} + \mathfrak{C}^{\tilde{\mu}\tilde{\rho}}\mathfrak{C}^{\tilde{\sigma}\tilde{\nu}} + \mathfrak{C}^{\tilde{\mu}\tilde{\sigma}}\mathfrak{C}^{\tilde{\nu}\tilde{\rho}} = -\varepsilon^{\tilde{\mu}\tilde{\nu}\tilde{\rho}\tilde{\sigma}}, \qquad (2.18b)$$

the first of which is equivalent to $\det(T_{\hat{\alpha}}^{\tilde{\mu}}) = 1$. Making use of the completeness relation

$$(\tilde{\Sigma}^{\hat{a}})^{\hat{\alpha}\hat{\beta}}(\tilde{\Sigma}_{\hat{a}})^{\hat{\gamma}\hat{\delta}} = 2\varepsilon^{\hat{\alpha}\hat{\beta}\hat{\gamma}\hat{\delta}}, \qquad (2.19)$$

see (A.50), we obtain

$$X^{\underline{a}}X_{\underline{a}} = -\ell^2 \,. \tag{2.20}$$

2.3 Bi-twistor realisation

A bi-twistor realisation for (complexified) AdS_5 exists in the literature, see [24,25]. In this section we will prove its equivalence with the twistor realisation described above.

Let \mathcal{L} be the set of bi-twistors, $X_{\hat{\alpha}\hat{\beta}} = -X_{\hat{\beta}\hat{\alpha}}$, which obey the following constraints:

$$\bar{X}^{\hat{\alpha}\hat{\beta}} = \Omega^{\hat{\alpha}\hat{\gamma}} (X^{\dagger})_{\hat{\gamma}\hat{\delta}} \Omega^{\hat{\delta}\hat{\beta}} = \frac{1}{2} \varepsilon^{\hat{\alpha}\hat{\beta}\hat{\gamma}\hat{\delta}} X_{\hat{\gamma}\hat{\delta}} , \qquad (2.21a)$$

$$\bar{X}^{\hat{\alpha}\hat{\gamma}}X_{\hat{\gamma}\hat{\beta}} = \ell^2 \delta^{\hat{\alpha}}{}_{\hat{\beta}} \,, \tag{2.21b}$$

$$X_{[\hat{\alpha}\hat{\beta}}X_{\hat{\gamma}\hat{\delta}]} = -\frac{1}{3}\ell^2 \varepsilon_{\hat{\alpha}\hat{\beta}\hat{\gamma}\hat{\delta}} \quad \iff \quad \varepsilon^{\hat{\alpha}\hat{\beta}\hat{\gamma}\hat{\delta}}X_{\hat{\alpha}\hat{\beta}}X_{\hat{\gamma}\hat{\delta}} = -8\ell^2 \,. \tag{2.21c}$$

Our aim is to show that the space \mathcal{L} can be identified with \mathfrak{F}/\sim . First of all, it is not difficult to see that \mathcal{L} provides an equivalent realisation of AdS₅. Making use of the correspondence (A.51), we associate a unique six-vector $X^{\underline{a}}$ with each $X_{\hat{\alpha}\hat{\beta}} \in \mathcal{L}$. The condition (2.21a) means that the six-vector $X^{\underline{a}}$ is real, whereas (2.21b) recreates the AdS condition $X^2 = -\ell^2$. We therefore can identify AdS₅ with \mathcal{L} . It remains to show that every $\hat{X} = (X_{\hat{\alpha}\hat{\beta}}) \in \mathcal{L}$ can be written in the form (2.13), where the twistors $T^{\tilde{\mu}}$ are constrained as in (2.9).

An SU(2,2) transformation acts on $\hat{X} = (X_{\hat{\alpha}\hat{\beta}})$ as

$$\hat{X} \longrightarrow g\hat{X}g^{\mathrm{T}} = X'^{\underline{a}}(\Sigma_{\underline{a}}), \qquad g \in \mathsf{SU}(2,2),$$

$$(2.22)$$

with $X^{\prime \underline{a}}$ given by

$$X^{\underline{a}} = \Lambda^{\underline{a}}{}_{\underline{b}} X^{\underline{b}}, \qquad \Lambda \in \mathsf{SO}_0(4,2).$$
(2.23)

This transformation preserves the conditions (2.21). Now we make use of the fact that AdS_5 is a homogeneous space of $SO_0(4, 2)$, and has a global coset representative s(X), see, e.g., [59]. That is, for any $X^{\underline{a}} \in AdS_5$ it holds that

$$X^{\underline{a}} = s(X)^{\underline{a}}_{\underline{b}} X^{\underline{b}}_{(0)}, \qquad s(X) \in \mathsf{SO}_0(4,2),$$
(2.24)

for some base point $X_{(0)}^{\underline{a}}$. We can choose the base point to be

$$X_{(0)}^{\underline{a}} = (0, 0, 0, 0, 0, -\ell), \qquad (2.25)$$

which corresponds to the bi-twistor

$$X_{(0)} = X^{\underline{a}}_{(0)}(\Sigma_{\underline{a}}) = \ell C^{-1}.$$
(2.26)

Then, any bi-twistor corresponding to a point in AdS_5 can be written as

$$\hat{X} = \ell g C^{-1} g^{\mathrm{T}}, \qquad g \in \mathsf{SU}(2,2).$$
 (2.27)

The group element g, however, is not uniquely defined. It is defined modulo

$$g \sim g\lambda$$
, $\lambda \in \mathsf{USp}(2,2)$. (2.28)

This completes the proof.

3 \mathcal{N} -extended AdS superspace in five dimensions

The supergroup $\mathsf{SU}(2,2|\mathcal{N})$ is the isometry group of five-dimensional \mathcal{N} -extended AdS superspace $\mathrm{AdS}^{5|8\mathcal{N}}$, eq. (1.1). Here we will describe an embedding formalism for $\mathrm{AdS}^{5|8\mathcal{N}}$ constructed in terms of supertwistors of $\mathsf{SU}(2,2|\mathcal{N})$.

3.1 Algebraic background

The supergroup $SU(2, 2|\mathcal{N})$ naturally acts on the space of *even* supertwistors and on the space of *odd* supertwistors.

An arbitrary supertwistor T is a column vector

$$T = (T_A) = \left(\frac{T_{\hat{\alpha}}}{T_i}\right), \qquad i = 1, \dots, \mathcal{N}.$$
 (3.1)

In the case of even supertwistors, $T_{\hat{\alpha}}$ is bosonic and T_i is fermionic. In the case of odd supertwistors, $T_{\hat{\alpha}}$ is fermionic while T_i is bosonic. The even and odd supertwistors are called pure. We introduce the parity function $\epsilon(T)$ defined as: $\epsilon(T) = 0$ if T is even, and $\epsilon(T) = 1$ if T is odd. If we define

$$\epsilon_A = \begin{cases} 0 , & A = \hat{\alpha} \\ 1 , & A = i \end{cases}$$
(3.2)

then the components \boldsymbol{T}_A of a pure supertwistor have the following Grassmann parities

$$\epsilon(\boldsymbol{T}_A) = \epsilon(\boldsymbol{T}) + \epsilon_A \pmod{2} . \tag{3.3}$$

The space of even supertwistors is naturally identified with $\mathbb{C}^{4|\mathcal{N}}$, while the space of odd supertwistors may be identified with $\mathbb{C}^{\mathcal{N}|4}$.

Supertwistor space is equipped with the inner product

$$\langle \boldsymbol{T} | \boldsymbol{S} \rangle = \boldsymbol{T}^{\dagger} \boldsymbol{\Omega} \boldsymbol{S}, \qquad \boldsymbol{\Omega} = \begin{pmatrix} 0 & \mathbb{1}_2 & 0 \\ \mathbb{1}_2 & 0 & 0 \\ 0 & 0 & -\mathbb{1}_{\mathcal{N}} \end{pmatrix} = \begin{pmatrix} \Omega & 0 \\ 0 & -\mathbb{1}_{\mathcal{N}} \end{pmatrix}.$$
 (3.4)

The inner product is invariant under the supergroup $SU(2, 2|\mathcal{N})$ spanned by supermatrices of the form

$$g = (g_A{}^B) \in \mathsf{SL}(4|\mathcal{N}), \qquad g^{\dagger}\Omega g = \Omega.$$
 (3.5)

Associated with a supertwistor T is its dual

$$\bar{\boldsymbol{T}} := \boldsymbol{T}^{\dagger} \boldsymbol{\Omega} = (\bar{\boldsymbol{T}}^{A}) = (\bar{\boldsymbol{T}}^{\hat{\alpha}}, -\bar{\boldsymbol{T}}^{i}), \qquad \bar{\boldsymbol{T}}^{i} := (\boldsymbol{T}_{i})^{*}.$$
(3.6)

 $\mathsf{SU}(2,2|\mathcal{N})$ acts on supertwistors and their duals as

$$\boldsymbol{T}_A \to g_A{}^B \boldsymbol{T}_B, \qquad \bar{\boldsymbol{T}}^A \to \bar{\boldsymbol{T}}^B (g^{-1})_B{}^A.$$
 (3.7)

The superalgebra $\mathfrak{su}(2,2|\mathcal{N})$ is spanned by elements of the form

$$\mathfrak{g} = \begin{pmatrix} \frac{\omega_{\alpha}{}^{\beta} - (\frac{1}{2}\omega^{45} - \frac{\mathrm{i}\mathcal{N}}{\mathcal{N}-4}\tau)\delta_{\alpha}{}^{\beta}}{\frac{\mathrm{i}}{2}(\omega^{a4} - \omega^{a5})(\tilde{\sigma}_{a})^{\dot{\alpha}\beta}} & -\frac{\mathrm{i}}{2}(\omega^{a4} + \omega^{a5})(\sigma_{a})_{\alpha\dot{\beta}}} & 2\psi_{\alpha}{}^{j}}{\frac{\mathrm{i}}{2}(\omega^{a4} - \omega^{a5})(\tilde{\sigma}_{a})^{\dot{\alpha}\beta}} & -\bar{\omega}^{\dot{\alpha}}{}_{\dot{\beta}} + (\frac{1}{2}\omega^{45} + \frac{\mathrm{i}\mathcal{N}}{\mathcal{N}-4}\tau)\delta^{\dot{\alpha}}{}_{\dot{\beta}}} & 2\bar{\epsilon}^{\dot{\alpha}j}}{2\bar{\epsilon}_{i}^{\beta}} & 2\bar{\epsilon}^{\dot{\alpha}j} & \frac{4\mathrm{i}}{\mathcal{N}-4}\tau\delta_{i}{}^{j} + \Lambda_{i}{}^{j}} \end{pmatrix}, \quad (3.8a)$$

with

$$\omega_{\alpha}{}^{\beta} = \frac{1}{2}\omega^{ab}(\sigma_{ab})_{\alpha}{}^{\beta}, \qquad \omega^{\underline{ab}} = -\omega^{\underline{ba}}, \tau \in \mathbb{R}, \qquad \Lambda = (\Lambda_i{}^j) \in \mathfrak{su}(\mathcal{N}).$$
(3.8b)

The elements of $\mathfrak{su}(2,2|\mathcal{N})$ satisfy the conditions

str
$$\mathfrak{g} = 0$$
, $\mathfrak{g}^{\dagger} \Omega + \Omega \mathfrak{g} = 0$. (3.9)

More details on the AdS superalgebra will be elaborated in section 5.

3.2 Supertwistor realisation

In complete analogy with the bosonic construction described in section 2, consider four *even* supertwistors with linearly independent $bodies^5$

$$\boldsymbol{T}^{\tilde{\mu}} = (\boldsymbol{T}_{A}{}^{\tilde{\mu}}), \qquad \boldsymbol{\epsilon}(\boldsymbol{T}_{A}{}^{\tilde{\mu}}) = \boldsymbol{\epsilon}_{A}, \qquad \tilde{\mu} = 1, 2, 3, 4.$$
(3.10)

We restrict our attention to those quartets satisfying the constraint

$$\bar{\boldsymbol{T}}^{\tilde{\mu}A}\boldsymbol{T}_{A}{}^{\tilde{\nu}} = \Omega^{\tilde{\mu}\tilde{\nu}}, \qquad \bar{\boldsymbol{T}}^{\tilde{\mu}A} := (\boldsymbol{T}_{B}{}^{\tilde{\mu}})^{*} \, \Omega^{BA}.$$
(3.11)

The space of such quartets will be denoted $\mathfrak{F}_{\mathcal{N}}$. On $\mathfrak{F}_{\mathcal{N}}$, we introduce the equivalence relation

$$\boldsymbol{T}_{A}{}^{\tilde{\mu}} \sim \boldsymbol{T}_{A}{}^{\tilde{\nu}}\lambda_{\tilde{\nu}}{}^{\tilde{\mu}}, \qquad \lambda \in \mathsf{USp}(2,2),$$

$$(3.12a)$$

$$\boldsymbol{T}_{A}{}^{\tilde{\mu}} \sim e^{i\varphi} \boldsymbol{T}_{A}{}^{\tilde{\mu}}, \qquad \varphi \in \mathbb{R}.$$
 (3.12b)

⁵The terminology "body" and "soul" follows [60].

We emphasise that the condition (3.11) is invariant under the equivalence transformations (3.12). For $\mathcal{N} > 0$, the constraint (2.9b) is substituted with (3.12b).

The supergroup $\mathsf{SU}(2,2|\mathcal{N})$ acts on $\mathfrak{F}_{\mathcal{N}}$ as

$$\boldsymbol{T}_{A}{}^{\tilde{\mu}} \to g_{A}{}^{B}\boldsymbol{T}_{B}{}^{\tilde{\mu}}, \qquad g \in \mathsf{SU}(2,2|\mathcal{N}).$$
 (3.13)

This action is naturally extended to the quotient space $\mathfrak{F}_{\mathcal{N}}/\sim$, which proves to be a homogeneous space of $\mathsf{SU}(2,2|\mathcal{N})$. To show this, we note that the quartets $T_A{}^{\tilde{\mu}}$ can be embedded into a group element

$$g = (\boldsymbol{T}_A{}^{\tilde{\mu}}, \boldsymbol{\Xi}_A{}^{\tilde{i}}) \in \mathsf{SU}(2, 2|\mathcal{N}) , \qquad \tilde{i} = 1, \dots, \mathcal{N} , \qquad (3.14)$$

and $SU(2,2|\mathcal{N})$ acts transitively on itself from the left. It turns out that

$$\mathfrak{F}_{\mathcal{N}}/\sim = \mathrm{AdS}^{5|\mathcal{B}\mathcal{N}}.$$
(3.15)

3.3 Stabiliser

To prove (3.15), it suffices to choose a base point and determine its stabiliser. As a base point, $T^{(0)}$, we choose

$$\boldsymbol{T}^{(0)} = \left(\frac{\mathbb{1}_4}{0}\right) \,. \tag{3.16}$$

This is the simplest quartet satisfying the constraint (3.11). By definition, the stabiliser H of $T^{(0)}$ consists of those elements $h \in SU(2, 2|\mathcal{N})$ that satisfy the condition

$$h\mathbf{T}^{(0)} = \left(\frac{\mathrm{e}^{\mathrm{i}\varphi\lambda}}{0}\right), \qquad \varphi \in \mathbb{R}, \quad \lambda \in \mathsf{USp}(2,2).$$
 (3.17)

That is, it consists of those group elements that map $T^{(0)}$ to an equivalent point with respect to (3.12). These conditions imply that the group elements h take the form

with

$$\varphi \in \mathbb{R}$$
, $\lambda \in \mathsf{USp}(2,2)$, $U \in \mathsf{SU}(\mathcal{N})$. (3.18b)

We therefore have H is isomorphic to

$$\mathsf{USp}(2,2) \times \mathsf{U}(1) \times \mathsf{SU}(\mathcal{N}) \,. \tag{3.19}$$

3.4 Bi-supertwistor realisation

In this subsection we will detail a bi-supertwistor realisation of the AdS superspace described above. Given a point in $\mathfrak{F}_{\mathcal{N}}$, we associate with it the following graded antisymmetric supermatrix

$$\boldsymbol{X}_{AB} := \ell \mathfrak{C}_{\tilde{\mu}\tilde{\nu}} \boldsymbol{T}_{A}{}^{\tilde{\mu}} \boldsymbol{T}_{B}{}^{\tilde{\nu}} = -(-1)^{\epsilon_{A}\epsilon_{B}} \boldsymbol{X}_{BA} \,.$$
(3.20)

These supermatrices are invariant under arbitrary equivalence transformations of the form (3.12a). Because of (3.12b), they are defined modulo the equivalence relation

$$\boldsymbol{X}_{AB} \sim e^{i\varphi} \boldsymbol{X}_{AB}, \qquad \varphi \in \mathbb{R},$$
(3.21)

and hence can be used to parametrise the space $\mathfrak{F}_{\mathcal{N}}/\sim$. Using the dual supertwistors we can introduce the following supermatrix

$$\bar{\boldsymbol{X}} := \boldsymbol{\Omega} \boldsymbol{X}^{\dagger} \boldsymbol{\Omega} \,, \qquad \bar{\boldsymbol{X}}^{AB} = -\ell \mathfrak{C}_{\tilde{\mu}\tilde{\nu}} \bar{\boldsymbol{T}}^{\tilde{\mu}A} \bar{\boldsymbol{T}}^{\tilde{\nu}B} \,. \tag{3.22}$$

The important properties of the supermatrices X and \overline{X} are

$$\boldsymbol{X}_{[AB}\boldsymbol{X}_{CD}\boldsymbol{X}_{E]F} = 0 \implies \boldsymbol{X}_{[AB}\boldsymbol{X}_{CD}\boldsymbol{X}_{EF]} = 0, \qquad (3.23a)$$

$$\bar{\boldsymbol{X}}^{AB}\boldsymbol{X}_{BA} = 4\ell^2\,,\tag{3.23b}$$

$$(-1)^{\epsilon_C} \boldsymbol{X}_{AC} \bar{\boldsymbol{X}}^{CD} \boldsymbol{X}_{DB} = \ell^2 \boldsymbol{X}_{AB}, \qquad (3.23c)$$

where $[...\}$ denotes the graded antisymmetrisation of indices. Property (3.23a) follows from the fact that X_{AB} is constructed from four supertwistors. An important implication of this property is that the body of the bosonic block X_{ij} defined by

$$\boldsymbol{X} = \left(\frac{\boldsymbol{X}_{\hat{\alpha}\hat{\beta}} \| \boldsymbol{X}_{\hat{\alpha}j}}{\boldsymbol{X}_{i\hat{\beta}} \| \boldsymbol{X}_{ij}} \right), \qquad (3.24)$$

vanishes. Indeed, let us consider the case

$$\boldsymbol{X}_{(ij}\boldsymbol{X}_{kl}\boldsymbol{X}_{mn)} = 0. aga{3.25}$$

Since X_{ij} is symmetric, the only non-zero solution to the above is a symmetric, bodiless matrix X_{ij} .

Now, we would like to describe the superspace $\mathrm{AdS}^{5|8\mathcal{N}}$ solely in terms of bi-supertwistors, without any reference to the supertwistor realisation above. Let us consider the space of graded antisymmetric supermatrices X_{AB} ,

$$\boldsymbol{X}_{AB} = -(-1)^{\epsilon_A \epsilon_B} \boldsymbol{X}_{BA}, \qquad (3.26a)$$

$$\epsilon(\boldsymbol{X}_{AB}) = \epsilon_A + \epsilon_B \,. \tag{3.26b}$$

In this space, we consider the surface \mathfrak{L} consisting of those supermatrices which obey the constraints (3.23). We then introduce the quotient space \mathfrak{L}/\sim , where the equivalence relation is given by (3.21). Our goal is to show that \mathfrak{L}/\sim can be identified with $\mathrm{AdS}^{5|8\mathcal{N}}$.

We begin by considering all the elements of ${\mathfrak L}$ with vanishing soul. These elements take the form

$$\boldsymbol{X}_{AB}|_{\Theta=0} := X_{AB} = \left(\begin{array}{c|c} X_{\hat{\alpha}\hat{\beta}} & 0\\ \hline 0 & 0 \end{array}\right), \qquad X_{\hat{\alpha}\hat{\beta}} = -X_{\hat{\beta}\hat{\alpha}} \in \mathbb{C}.$$
(3.27)

The space of such elements will be denoted $\mathfrak{L}|$.

It turns out that $\mathfrak{L}|/\sim$ is a homogeneous space of the subgroup of $\mathsf{SU}(2,2|\mathcal{N})$ spanned by matrices of the form

$$v = \left(\frac{e^{i\varphi\Sigma} \quad 0}{0 \quad |e^{\frac{4i\varphi}{N}}U}\right), \qquad \varphi \in \mathbb{R}, \quad \Sigma \in \mathsf{SU}(2,2), \quad U \in \mathsf{SU}(\mathcal{N}).$$
(3.28)

The proof is analogous to that described in subsection 2.3 and is described below. As a base point we can choose

$$X_{(0)} = \left(\begin{array}{c|c} \ell C^{-1} & 0\\ \hline 0 & 0 \end{array}\right) \,. \tag{3.29}$$

Then, any point in $\mathfrak{L}|/\sim$ can be reached by a group transformation of the form

$$X = v X_{(0)} v^{\mathrm{T}}, \qquad (3.30)$$

with v given by (3.28). As in the bosonic case, the transformation v is not uniquely defined. The stabiliser of the bi-supertwistor $X_{(0)}$ consists precisely of those group elements that take the form (3.18a). We therefore have that $\mathfrak{L}|/\sim$ coincides with the body of the coset superspace $SU(2, 2|\mathcal{N})/(USp(2, 2) \times U(1) \times SU(\mathcal{N}))$.

To prove (3.30), we note that $X_{\hat{\alpha}\hat{\beta}}$ in (3.27) can be written as

$$X_{\hat{\alpha}\hat{\beta}} = X^{\underline{a}}(\Sigma_{\underline{a}})_{\hat{\alpha}\hat{\beta}}.$$
(3.31)

The matrix $\hat{X} = (X_{\hat{\alpha}\hat{\beta}})$ is non-singular. Because of property (3.23c), we have

$$\hat{X}^{\dagger}\Omega\hat{X} = \ell^2\Omega \implies \bar{X}^{\hat{\alpha}\hat{\gamma}}X_{\hat{\gamma}\hat{\beta}} = \ell^2\delta^{\hat{\alpha}}{}_{\hat{\beta}}.$$
(3.32)

This property has two implications:

1. the mutually conjugate six-vectors $\bar{X}^{\underline{a}}$ and $X^{\underline{a}}$ satisfy the following

$$\bar{X}^{\underline{a}}X_{\underline{a}} = -\ell^2 \,, \tag{3.33a}$$

2. $\bar{X}^{\underline{a}}$ and $X^{\underline{a}}$ are linearly dependent, which follows from

$$\bar{X}^{\underline{a}} X^{\underline{b}} (\tilde{\Sigma}_{\underline{a}\underline{b}})^{\hat{\alpha}}{}_{\hat{\beta}} = 0 , \qquad (3.33b)$$

where we have used (3.33a). Property (3.33a) can be seen by taking the trace of (3.32). To prove property (3.33b) it is instructive to write out the above expression explicitly,

$$\bar{X}^{\underline{a}}X^{\underline{b}}(\tilde{\Sigma}_{\underline{a}\underline{b}})^{\hat{\alpha}}{}_{\hat{\beta}} = -\frac{1}{4}\bar{X}^{\underline{a}}X^{\underline{b}}\left(\tilde{\Sigma}_{\underline{a}}\Sigma_{\underline{b}} - \tilde{\Sigma}_{\underline{b}}\Sigma_{\underline{a}}\right)^{\hat{\alpha}}{}_{\hat{\beta}}, \qquad (3.34)$$

and then make use of (A.44). Then we have

$$X^{\underline{a}} = e^{i\varphi}Y^{\underline{a}}, \qquad \varphi \in \mathbb{R}, \quad Y^{\underline{a}} \in AdS_5.$$
 (3.35)

Now we can repeat the argument from the non-supersymmetric case. Specifically, we have

$$X^{\underline{a}} = e^{i\varphi} s(Y)^{\underline{a}} Y^{\underline{b}}_{(0)}, \qquad s(Y) \in \mathsf{SO}_0(4,2), \qquad (3.36)$$

for a base point $Y_{(0)}^{\underline{a}}$. Expression (3.30) follows.

We now note that both the full supergroup, $\mathsf{SU}(2,2|\mathcal{N})$, and a generic bi-supertwistor $\mathbf{X}_{AB} \in \mathfrak{L}/\sim$ have $8\mathcal{N}$ real Grassmann odd degrees of freedom. We conclude that, by replacing v in (3.30) with some $g \in \mathsf{SU}(2,2|\mathcal{N})$, we can reach an arbitrary $\mathbf{X} \in \mathfrak{L}/\sim$ by a transformation acting on $X_{(0)}$. That is, the space \mathfrak{L}/\sim is itself a homogeneous space of $\mathsf{SU}(2,2|\mathcal{N})$, which shows the equivalence of the bi-supertwistor and supertwistor realisations of $\mathrm{AdS}^{5|8\mathcal{N}}$.

3.5 SU(2, 2|N)-invariant two-point function

A powerful application of the formalism described above is the construction of manifestly invariant two-point functions. Indeed, let T and T' be arbitrary points of $\mathfrak{F}_{\mathcal{N}}$. The following two-point function

$$\omega(\boldsymbol{T},\boldsymbol{T}') := -\frac{1}{4} \ell^2 \mathfrak{C}_{\tilde{\mu}\tilde{\nu}} \mathfrak{C}_{\tilde{\rho}\tilde{\sigma}} \langle \boldsymbol{T}^{\tilde{\mu}} | \boldsymbol{T}'^{\tilde{\rho}} \rangle \langle \boldsymbol{T}^{\tilde{\nu}} | \boldsymbol{T}'^{\tilde{\sigma}} \rangle$$
(3.37)

is $SU(2, 2|\mathcal{N})$ -invariant. It is also invariant under the equivalence transformations (3.12), and is therefore defined on the quotient space (3.15). The two-point function (3.37) can be expressed in terms of the bi-supertwistors X and \overline{X} . Let us denote the bi-supertwistors corresponding to T and T' as X and X'. Then the two-point function takes the form

$$\omega(\boldsymbol{X}, \boldsymbol{X}') = -\frac{1}{4} \bar{\boldsymbol{X}}^{AB} \boldsymbol{X}'_{BA}. \qquad (3.38)$$

In the non-supersymmetric case, $\mathcal{N} = 0$, (3.37) and (3.38) coincide with the AdS₅ two-point function $X^{\underline{a}}X'_{\underline{a}}$.

Recently, a similar construction to the above was used to describe a worldline superparticle model in $AdS^{4|4N}$, see [23]. In the present context, such a model would be described by the following action

$$S = -\frac{1}{2} \int \mathrm{d}\tau \, \mathbf{\mathfrak{e}}^{-1} \left\{ -\frac{1}{4} \dot{\bar{\mathbf{X}}}^{AB} \dot{\mathbf{X}}_{BA} + (\mathbf{\mathfrak{e}}m)^2 \right\} \,. \tag{3.39}$$

4 Coset representative

A key role in the study of the geometry of a coset space is played by the coset representative. A global coset representative for a homogeneous space $\mathcal{M} = G/H_{x_0}$, where G is a Lie group and H_{x_0} is the stabiliser (also known as the isotropy group or little group) of some point $x_0 \in \mathcal{M}$, is a bijective map $\mathcal{S} : \mathcal{M} \to G$ such that $\pi \circ \mathcal{S} = \mathrm{id}_{\mathcal{M}}$, where π denotes the canonical projection $\pi : G \to G/H_{x_0}$. For many homogeneous spaces, a global coset representative does not exist. In such cases, local coset representatives $\mathcal{S}_A : U_A \to G$ with the property $\pi \circ \mathcal{S}_A = \mathrm{id}_{U_A}$ can be defined on open charts $\{U_A\}$ that provide an atlas for \mathcal{M} . In the intersection of two charts, U_A and $U_B, U_A \cap U_B \neq \emptyset$, the corresponding coset representatives \mathcal{S}_A and \mathcal{S}_B are related by a little group transformation, $\mathcal{S}_B(x) = \mathcal{S}_A(x)h_{AB}(x)$, with $h_{AB}(x) \in H_{x_0}$.

A global coset representative exists for *d*-dimensional AdS in the context of the standard embedding formalism, see [59]. For the embedding formalism described in this work, suitable for AdS superspace, an exponential coset parametrisation is used. In this case we find local coset representatives and their associated local coordinate systems. These are detailed below.

4.1 AdS space $(\mathcal{N} = 0)$

The Lie algebras $\mathfrak{su}(2,2)$ and $\mathfrak{usp}(2,2)$ are given by expressions (A.48) and (A.63), respectively. Let \mathfrak{K} be the complement of $\mathfrak{usp}(2,2)$ in $\mathfrak{su}(2,2)$. We then have

$$\mathfrak{su}(2,2) = \mathfrak{usp}(2,2) \oplus \mathfrak{K}.$$
(4.1)

The elements of $\mathfrak K$ take the form

$$\mathfrak{t} = \left(\frac{-\frac{1}{2}\omega^{45}\delta_{\alpha}{}^{\beta} \left| -\frac{\mathrm{i}}{2}(\omega^{a5})(\sigma_{a})_{\alpha\dot{\beta}}}{-\frac{\mathrm{i}}{2}(\omega^{a5})(\tilde{\sigma}^{a})^{\dot{\alpha}\beta} \left| \frac{1}{2}\omega^{45}\delta^{\dot{\alpha}}{}_{\dot{\beta}}} \right) \in \mathfrak{K}, \qquad \omega^{\hat{a}5} \in \mathbb{R}.$$

$$(4.2)$$

Then, for a group element $g \in SU(2,2)$ in a neighbourhood of $\mathbb{1}_4$, it holds that

$$g = e^{\mathfrak{t} + \mathfrak{h}}, \qquad \mathfrak{h} \in \mathfrak{usp}(2, 2).$$
 (4.3)

We seek to (locally) factorise g in the following way

$$g = \mathcal{S}(y)h$$
, $\mathcal{S}(y) \in \mathsf{SU}(2,2)$, $h \in \mathsf{USp}(2,2)$, (4.4)

where $\mathcal{S}(y)$ is the coset representative, and y are local coordinates on AdS₅. We can make the following ansatz for $\mathcal{S}(y)$:

$$\mathcal{S}(y) = \left(\begin{array}{c|c} \xi \rho \mathbb{1}_2 & -\mathrm{i}x \\ \hline -\mathrm{i}\tilde{x} & \xi \rho^{-1} \mathbb{1}_2 \end{array}\right) = \mathrm{e}^{\mathfrak{t}}, \qquad (4.5)$$

with

$$x := x^{a}(\sigma_{a}) = x^{\dagger}, \qquad \tilde{x} := x^{a}(\tilde{\sigma}_{a}) = \tilde{x}^{\dagger}, \qquad (4.6a)$$

$$\xi := (1+x^2)^{\frac{1}{2}}, \qquad x^2 = x^a x^b \eta_{ab}, \qquad (4.6b)$$

$$\rho = \bar{\rho} \,. \tag{4.6c}$$

It follows that the local coordinates x^a and ρ are real, and the coordinate chart is specified by

$$x^2 > -1, \qquad \rho \neq 0.$$
 (4.7)

The bi-twistor corresponding to (4.5) is given by

$$X_{\hat{\alpha}\hat{\beta}} = \ell \mathfrak{C}_{\tilde{\mu}\tilde{\nu}} \mathcal{S}(y)_{\hat{\alpha}}{}^{\tilde{\mu}} \mathcal{S}(y)_{\hat{\beta}}{}^{\tilde{\nu}} = \ell \left(\frac{\left(\xi^2 \rho^2 + x^2\right)\varepsilon_{\alpha\beta} \left| -i\xi(\rho + \rho^{-1})x_{\alpha}{}^{\dot{\beta}} \right|}{i\xi(\rho + \rho^{-1})x^{\dot{\alpha}}{}_{\beta} \left| -\left(\xi^2 \rho^{-2} + x^2\right)\varepsilon^{\dot{\alpha}\dot{\beta}} \right|} \right),$$
(4.8a)

where

$$\mathcal{S}(y)_{\hat{\alpha}}{}^{\tilde{\mu}} := \mathcal{S}(y)_{\hat{\alpha}}{}^{\hat{\beta}}T^{(0)}_{\hat{\beta}}{}^{\tilde{\mu}}, \qquad T^{(0)} = \mathbb{1}_4.$$

$$(4.8b)$$

Associated with $X_{\hat{\alpha}\hat{\beta}}$ is the following point in ${\rm AdS}_5$

$$X^{\underline{a}} = \left(\ell\xi(\rho + \rho^{-1})x^{a}, -\frac{1}{2}\ell\xi^{2}(\rho^{2} - \rho^{-2}), -\ell\left(x^{2} + \frac{1}{2}\xi^{2}(\rho^{2} + \rho^{-2})\right)\right).$$
(4.9)

Instead of the coset parametrisation used above, we could choose a coset representative corresponding to Poincaré coordinates in AdS_5 . This coset representative, used in [19, 45], is given by

$$S(y) = \left(\frac{\rho^{\frac{1}{2}} \mathbb{1}_2 \ 0}{-i\rho^{\frac{1}{2}} \tilde{x} \ \rho^{-\frac{1}{2}} \mathbb{1}_2}\right).$$
(4.10)

Its corresponding bi-twistor is

$$X_{\hat{\alpha}\hat{\beta}} = \frac{1}{z} \left(\frac{\varepsilon_{\alpha\beta}}{\mathrm{i}x^{\dot{\alpha}}{}_{\beta}} \left| -((\ell z)^2 + x^2)\varepsilon^{\dot{\alpha}\dot{\beta}} \right| \right), \qquad z := (\ell\rho)^{-1}.$$

$$(4.11)$$

The real coordinates z > 0 and x^a parametrise AdS₅ in the Poincaré patch. They are related to the embedding coordinates $X^{\underline{a}}$ as follows

$$X^{\underline{a}} = \frac{1}{z} \left(x^{a}, -\frac{1}{2} (1 - x^{2} - (\ell z)^{2}), -\frac{1}{2} (1 + x^{2} + (\ell z)^{2}) \right).$$
(4.12)

$4.2 \quad \mathcal{N} \neq 0$

The analysis of the previous section can be extended to the supersymmetric case in similar fashion. Coset representatives for the $\mathcal{N} = 1$ superspaces $\mathrm{AdS}^{5|8} \times S^1$ and $\mathrm{AdS}^{5|8}$ are given in [19,61]. Building on the approach of these references, again denoting the local coordinates by y, we choose the coset representative in the Poincaré patch as

$$\begin{aligned} \mathcal{S}(y) &= g(\mathbf{z}) \cdot g_{S} \cdot g_{D} \end{aligned} \tag{4.13} \\ &= \begin{pmatrix} \frac{1_{2}}{-i\tilde{x}_{+}} & \frac{1}{2} & \frac{2\bar{\theta}}{2\bar{\theta}} \\ \hline 2\theta & 0 & 1_{\mathcal{N}} \end{pmatrix} \begin{pmatrix} \frac{1_{2}}{2} & 2\eta\bar{\eta} & 2\eta \\ \hline 0 & 1_{2} & 0 \\ \hline 0 & 2\bar{\eta} & 1_{\mathcal{N}} \end{pmatrix} \begin{pmatrix} \frac{\rho^{\frac{1}{2}}1_{2}}{2} & 0 & 0 \\ \hline 0 & \rho^{-\frac{1}{2}}1_{2} & 0 \\ \hline 0 & 0 & 1_{\mathcal{N}} \end{pmatrix} \\ &= \begin{pmatrix} \frac{\rho^{\frac{1}{2}}\delta_{\alpha}^{\beta}}{-i\rho^{\frac{1}{2}}x_{+}^{\dot{\alpha}\beta}} & 2\rho^{-\frac{1}{2}}\eta_{\alpha}^{\dot{\alpha}}\bar{\eta}_{k\dot{\beta}} & 2\eta_{\alpha}^{j} \\ \hline -i\rho^{\frac{1}{2}}x_{+}^{\dot{\alpha}\beta} & \rho^{-\frac{1}{2}}(\delta^{\dot{\alpha}}_{\dot{\beta}} - 2ix_{+}^{\dot{\alpha}\gamma}\eta_{\gamma}^{k}\bar{\eta}_{k\dot{\beta}} + 4\bar{\theta}^{\dot{\alpha}k}\bar{\eta}_{k\dot{\beta}}) & 2(\bar{\theta}^{\dot{\alpha}j} - ix_{+}^{\dot{\alpha}\gamma}\eta_{\gamma}^{j}) \\ \hline 2\rho^{\frac{1}{2}}\theta_{i}^{\beta}} & 2\rho^{-\frac{1}{2}}(\bar{\eta}_{i\dot{\beta}} + 2\theta_{i}^{\gamma}\eta_{\gamma}^{k}\bar{\eta}_{k\dot{\beta}}) & \delta_{i}^{j} + 4\theta_{i}^{\gamma}\eta_{\gamma}^{j} \end{pmatrix} \end{aligned}$$

where $x_{\pm}^{a} = x^{a} \pm i\theta_{i}\sigma^{a}\bar{\theta}^{i}$. The coset representative $g(\boldsymbol{z})$ corresponds to that of 4D \mathcal{N} -extended Minkowski superspace.

The action of S(y) on the base point $T^{(0)}$, see (3.16), yields the general form of the quartet $T_A^{\tilde{\mu}}$ in the Poincaré patch

$$\boldsymbol{T}_{A}^{\tilde{\mu}} = \begin{pmatrix} \rho^{\frac{1}{2}} \delta_{\alpha}{}^{\beta} & 2\rho^{-\frac{1}{2}} \eta_{\alpha}{}^{k} \bar{\eta}_{k\dot{\beta}} \\ \hline -\mathrm{i}\rho^{\frac{1}{2}} x_{+}^{\dot{\alpha}\beta} & \rho^{-\frac{1}{2}} \left(\delta^{\dot{\alpha}}{}_{\dot{\beta}} - 2\mathrm{i} x_{+}^{\dot{\alpha}\gamma} \eta_{\gamma}{}^{k} \bar{\eta}_{k\dot{\beta}} + 4\bar{\theta}^{\dot{\alpha}k} \bar{\eta}_{k\dot{\beta}} \right) \\ \hline 2\rho^{\frac{1}{2}} \theta_{i}{}^{\beta} & 2\rho^{-\frac{1}{2}} \left(\bar{\eta}_{i\dot{\beta}} + 2\theta_{i}{}^{\gamma} \eta_{\gamma}{}^{k} \bar{\eta}_{k\dot{\beta}} \right) \end{pmatrix} .$$
(4.14)

Furthermore, the bi-supertwistor X_{AB} is

$$\boldsymbol{X}_{AB} = \begin{pmatrix} \boldsymbol{X}_{\alpha\beta} & \boldsymbol{X}_{\alpha}{}^{\dot{\beta}} & \boldsymbol{X}_{\alpha j} \\ \boldsymbol{X}^{\dot{\alpha}}_{\beta} & \boldsymbol{X}^{\dot{\alpha}\dot{\beta}} & \boldsymbol{X}^{\dot{\alpha}}_{j} \\ \boldsymbol{X}_{i\beta} & \boldsymbol{X}_{i}{}^{\dot{\beta}} & \boldsymbol{X}_{ij} \end{pmatrix}, \qquad (4.15)$$

where the components are given by

$$\boldsymbol{X}_{\alpha\beta} = \frac{1}{z} \left(\epsilon_{\alpha\beta} + 4(\ell z)^2 \eta_{\alpha}{}^k \eta_{\beta}{}^l \bar{\eta}_{k\dot{\gamma}} \bar{\eta}_l{}^{\dot{\gamma}} \right), \tag{4.16a}$$

$$\boldsymbol{X}_{\alpha}{}^{\dot{\beta}} = \frac{1}{z} \left(-\mathrm{i}x_{+\alpha}{}^{\dot{\beta}} + 2(\ell z)^2 \eta_{\alpha}{}^k \bar{\eta}_k{}^{\dot{\gamma}} \left(\delta^{\dot{\beta}}{}_{\dot{\gamma}} - 2\mathrm{i}x_{+}^{\dot{\beta}\gamma} \eta_{\gamma}{}^l \bar{\eta}_{l\dot{\gamma}} + 4\bar{\theta}{}^{\dot{\beta}l} \bar{\eta}_{l\dot{\gamma}} \right) \right), \tag{4.16b}$$

$$\boldsymbol{X}_{\alpha j} = \frac{1}{z} \left(2\theta_{\alpha j} - 4(\ell z)^2 \eta_{\alpha}{}^k \bar{\eta}_{k \dot{\gamma}} \left(\bar{\eta}_j{}^{\dot{\gamma}} + 2\theta_j{}^{\gamma} \eta_{\gamma}{}^l \bar{\eta}_l{}^{\dot{\gamma}} \right) \right), \tag{4.16c}$$

$$\boldsymbol{X}^{\dot{\alpha}\dot{\beta}} = \frac{1}{z} \left(-(\ell z)^2 \left(\delta^{\dot{\alpha}}{}_{\dot{\gamma}} - 2\mathrm{i} x_+^{\dot{\alpha}\gamma} \eta_{\gamma}{}^k \bar{\eta}_{\kappa\dot{\gamma}} + 4\bar{\theta}^{\dot{\alpha}k} \bar{\eta}_{k\dot{\gamma}} \right) \left(\varepsilon^{\dot{\gamma}\dot{\beta}} - 2\mathrm{i} x_+^{\dot{\beta}\sigma} \eta_{\sigma}{}^l \bar{\eta}_l{}^{\dot{\gamma}} + 4\bar{\theta}^{\dot{\beta}l} \bar{\eta}_l{}^{\dot{\gamma}} \right) - x_+^2 \varepsilon^{\dot{\alpha}\dot{\beta}} \right),$$

$$(4.16d)$$

$$\boldsymbol{X}^{\dot{\alpha}}{}_{j} = \frac{1}{z} \Big(-2(\ell z)^{2} \Big(\delta^{\dot{\alpha}}{}_{\dot{\gamma}} - 2\mathrm{i} x^{\dot{\alpha}\gamma}_{+} \eta_{\gamma}{}^{k} \bar{\eta}_{k\dot{\gamma}} + 4\bar{\theta}^{\dot{\alpha}k} \bar{\eta}_{k\dot{\gamma}} \Big) \Big(\bar{\eta}_{j}{}^{\dot{\gamma}} + 2\theta_{j}{}^{\gamma} \eta_{\gamma}{}^{l} \bar{\eta}_{l}{}^{\dot{\gamma}} \Big) - 2\mathrm{i} x^{\dot{\alpha}\gamma}_{+} \theta_{\gamma j} \Big) , \qquad (4.16e)$$

$$\boldsymbol{X}_{ij} = \frac{1}{z} \left(4\theta_i^{\ \alpha} \theta_{\alpha j} - 4(\ell z)^2 \left(\bar{\eta}_{\iota \dot{\alpha}} + 2\theta_i^{\ \gamma} \eta_{\gamma}^{\ k} \bar{\eta}_{k \dot{\alpha}} \right) \left(\bar{\eta}_j^{\ \dot{\alpha}} + 2\theta_j^{\ \delta} \eta_{\delta}^{\ l} \bar{\eta}_{l}^{\ \dot{\alpha}} \right) \right).$$
(4.16f)

The coordinates z and ρ are related as in (4.11).

Using the above formulation we can readily describe the conformal boundary of AdS superspace. To do so, we should switch from \mathbf{X}_{AB} to $z\mathbf{X}_{AB}$ and take the limit $z \to 0$. In this limit, the Grassmann variables η disappear and the resulting bi-supertwistor is given by

$$\boldsymbol{X}_{AB}^{\text{boundary}} = \begin{pmatrix} \frac{\varepsilon_{\alpha\beta} & -ix_{+\alpha}{}^{\dot{\beta}} & 2\theta_{\alpha j} \\ \hline ix_{+}{}^{\dot{\alpha}}{}_{\beta} & -x_{+}^{2}\varepsilon^{\dot{\alpha}\dot{\beta}} & -2ix_{+}{}^{\dot{\alpha}\gamma}\theta_{\gamma j} \\ \hline -2\theta_{i\beta} \left| 2i\theta_{i\gamma}x_{+}{}^{\gamma\dot{\beta}} \right| & 4\theta_{i}{}^{\alpha}\theta_{\alpha j} \end{pmatrix} .$$
(4.17)

This expression coincides with equation (3.17) in [39]. Making use of the results of [39], $\boldsymbol{X}_{AB}^{\text{boundary}}$ can be decomposed into a pair of supertwistors

$$\boldsymbol{T}_{A}{}^{\beta} = \begin{pmatrix} \delta_{\alpha}{}^{\beta} \\ -\mathrm{i}x_{+}^{\dot{\alpha}\beta} \\ \hline 2\theta_{i}{}^{\beta} \end{pmatrix}, \qquad \boldsymbol{T}_{A}{}^{\beta} \sim \boldsymbol{T}_{A}{}^{\gamma}M_{\gamma}{}^{\beta}, \qquad M \in \mathsf{GL}(2\,,\mathbb{C})\,.$$
(4.18)

The local bosonic (x_{+}^{a}) and fermionic (θ_{i}^{α}) coordinates can be interpreted as coordinates in the chiral subspace of a four-dimensional \mathcal{N} -extended Minkowski superspace, $\mathbb{M}^{4|4\mathcal{N}}$. This is supported by the action of the AdS supergroup, $SU(2, 2|\mathcal{N})$, as superconformal transformations on $\mathbb{M}^{4|4\mathcal{N}}$.

An infinitesimal $SU(2, 2|\mathcal{N})$ transformation, $g = \mathbb{1}_{4+\mathcal{N}} + \mathfrak{g}$, acts on $T_A{}^\beta$ as

$$\boldsymbol{T}_{A}{}^{\beta} \to \boldsymbol{T}_{A}{}^{\prime}{}^{\beta} = \left(\delta_{A}{}^{B} + \mathfrak{g}_{A}{}^{B}\right)\boldsymbol{T}_{B}{}^{\gamma}N_{\gamma}{}^{\beta}, \qquad N \in \mathsf{SL}(2,\mathbb{C}),$$

$$(4.19)$$

where \mathfrak{g} is defined in (3.8). One can show that the coordinates x^a_+ and θ^{α}_i transform as

$$\delta \tilde{x}_{+} = \tilde{a} + (\sigma + \bar{\sigma})\tilde{x}_{+} - \bar{\omega}\tilde{x}_{+} - \tilde{x}_{+}\omega + \tilde{x}_{+}b\tilde{x}_{+} + 4i\bar{\epsilon}\theta - 4\tilde{x}_{+}\psi\theta, \qquad (4.20a)$$

$$\delta\theta = \epsilon + \frac{1}{\mathcal{N}} \big((\mathcal{N} - 2)\sigma + 2\bar{\sigma} \big) \theta - \theta\omega + \Lambda\theta + \theta b\tilde{x}_{+} - \mathrm{i}\bar{\psi}\tilde{x}_{+} - 4\theta\psi\theta \,, \tag{4.20b}$$

where the parameters σ , a^a , and b^a are related to those in (3.8) by the rule

$$\sigma = \frac{1}{2} \left(\omega^{45} - \frac{2i\mathcal{N}}{\mathcal{N} - 4} \tau \right), \qquad (4.21a)$$

$$a^{a} = -\frac{1}{2}(\omega^{a4} - \omega^{a5}),$$
 (4.21b)

$$b^{a} = \frac{1}{2} (\omega^{a4} + \omega^{a5}) \,. \tag{4.21c}$$

They correspond to 4D Lorentz transformations $(\omega, \bar{\omega})$, translations (a), special conformal transformations (b), Q-supersymmetry $(\epsilon, \bar{\epsilon})$ and S-supersymmetry $(\psi, \bar{\psi})$ transformations, combined scale and chiral transformations (σ) , and $SU(\mathcal{N})$ transformations (Λ) . The expressions (4.20) coincide with the standard superconformal transformations on Minkowski superspace, see, e.g., [38].

5 Superspace geometry

In this section we will elaborate on details of the supergeometry of the \mathcal{N} -extended AdS superspace introduced above. We introduce local bosonic (x) and fermionic (θ) coordinates on AdS superspace as

$$y^{\hat{M}} = (x^{\hat{m}}, \theta_i{}^{\hat{\mu}}, \bar{\theta}{}^{i}{}_{\hat{\mu}}), \qquad i = 1, \dots \mathcal{N}.$$
(5.1)

5.1 Geometric structures in $AdS^{5|8N}$

We will begin by formalising some of the earlier discussion about the AdS superalgebra. Let us denote the following

$$\mathcal{H} := \mathfrak{usp}(2,2) \oplus \mathfrak{u}(1) \oplus \mathfrak{su}(\mathcal{N}).$$
(5.2)

Denote the complement of \mathcal{H} in $\mathfrak{su}(2,2|\mathcal{N})$ by \mathcal{K} ,

$$\mathfrak{su}(2,2|\mathcal{N}) = \mathcal{K} \oplus \mathcal{H}.$$
(5.3)

The elements $k \in \mathcal{K}$ are given by

$$k = \begin{pmatrix} \frac{-\frac{1}{2}\omega^{45}\delta_{\alpha}{}^{\beta}}{-\frac{i}{2}(\omega^{a5})(\bar{\sigma}_{a})_{\alpha\dot{\beta}}} & 2\eta_{\alpha}{}^{j} \\ \hline \frac{-\frac{i}{2}(\omega^{a5})(\bar{\sigma}_{a})^{\dot{\alpha}\beta}}{2\epsilon_{i}{}^{\beta}} & \frac{1}{2}\omega^{45}\delta^{\dot{\alpha}}{}_{\dot{\beta}}} & 2\bar{\epsilon}^{\dot{\alpha}j} \\ \hline 2\epsilon_{i}{}^{\beta}} & 2\bar{\eta}_{i\dot{\beta}} & 0 \end{pmatrix}, \qquad \omega^{\hat{a}5} \in \mathbb{R}.$$
(5.4)

The elements $m \in \mathcal{H}$ are given by

$$m = \begin{pmatrix} \frac{\frac{1}{2}\omega^{ab}(\sigma_{ab})_{\alpha}{}^{\beta} + \frac{\mathrm{i}\mathcal{N}}{\mathcal{N}-4}\tau\delta_{\alpha}{}^{\beta} & -\frac{\mathrm{i}}{2}(\omega^{a4})(\sigma_{a})_{\alpha\dot{\beta}} & 0\\ \frac{\frac{\mathrm{i}}{2}(\omega^{a4})(\tilde{\sigma}_{a})^{\dot{\alpha}\beta} & \frac{1}{2}\omega^{ab}(\tilde{\sigma}_{ab})^{\dot{\alpha}}{}_{\dot{\beta}} + \frac{\mathrm{i}\mathcal{N}}{\mathcal{N}-4}\tau\delta^{\dot{\alpha}}{}_{\dot{\beta}} & 0\\ \hline 0 & 0 & \frac{4\mathrm{i}}{\mathcal{N}-4}\tau\delta_{i}{}^{j} + \Lambda_{i}{}^{j} \end{pmatrix},$$
(5.5a)

with

$$\omega^{\hat{a}\hat{b}}, \tau \in \mathbb{R}, \qquad \Lambda \in \mathfrak{su}(\mathcal{N}).$$
(5.5b)

Using the coset representative, one can introduce the Maurer-Cartan form, $S^{-1}dS$, which proves to encode all the information about the geometry of our superspace. The Maurer-Cartan form can be decomposed into the vielbein and the connection as

$$\mathcal{S}^{-1}\mathrm{d}\mathcal{S} = \mathbf{E} + \mathbf{\Phi}\,,\tag{5.6a}$$

$$\mathbf{E} = (\mathcal{S}^{-1} \mathrm{d}\mathcal{S})|_{\mathcal{K}}, \qquad (5.6b)$$

$$\Phi = (\mathcal{S}^{-1} \mathrm{d}\mathcal{S})|_{\mathcal{H}}.$$
 (5.6c)

Under a group transformation $g \in SU(2, 2|\mathcal{N})$, the coset representative transforms as

$$\mathcal{S}(y) \to \mathcal{S}(y') = g \,\mathcal{S}(y) \, h^{-1}(g, y) \,, \qquad h \in H \,,$$

$$(5.7)$$

and the vielbein and connection transform as

$$\mathbf{E} \to h \mathbf{E} h^{-1} \,, \tag{5.8a}$$

$$\Phi \to h\Phi h^{-1} - \mathrm{d}h \, h^{-1} \,. \tag{5.8b}$$

Since the vielbein and the connection are elements of the AdS superalgebra, it is useful to introduce a basis for the superalgebra as

$$K_{\hat{A}} := (P_{\hat{a}}, q_{\hat{\alpha}}{}^i, \bar{q}^{\hat{\alpha}}{}_i), \qquad (5.9a)$$

$$H_{\hat{I}} := (M_{\hat{a}\hat{b}}, \mathbb{J}^{i}_{j}, \mathbb{Y}).$$

$$(5.9b)$$

The basis elements $K_{\hat{A}}$ correspond to supertranslations, while $H_{\hat{I}}$ correspond to Lorentz, $SU(\mathcal{N})$, and U(1) transformations. They obey the following graded commutation relations

$$[M_{\hat{a}\hat{b}}, M_{\hat{c}\hat{d}}] = 2\eta_{\hat{d}[\hat{a}}M_{\hat{b}]\hat{c}} - 2\eta_{\hat{c}[\hat{a}}M_{\hat{b}]\hat{d}}, \qquad (5.10a)$$

$$[M_{\hat{a}\hat{b}}, P_{\hat{c}}] = 2\eta_{\hat{c}[\hat{b}}P_{\hat{a}]}, \qquad (5.10b)$$

$$[P_{\hat{a}}, P_{\hat{b}}] = M_{\hat{a}\hat{b}}, \qquad (5.10c)$$

$$[\mathbb{J}_{j}^{i},\mathbb{J}_{l}^{k}] = \delta_{j}^{k}\mathbb{J}_{l}^{i} - \delta_{l}^{i}\mathbb{J}_{j}^{k}, \qquad (5.10d)$$

$$[M_{\hat{a}\hat{b}}, q_{\hat{\alpha}}{}^{i}] = -(\Sigma_{\hat{a}\hat{b}})_{\hat{\alpha}}{}^{\hat{\beta}}q_{\hat{\beta}}{}^{i}, \qquad (5.10e)$$

$$[P_{\hat{a}}, q_{\hat{\alpha}}{}^{i}] = \frac{1}{2} (\gamma_{\hat{a}})_{\hat{\alpha}}{}^{\hat{\beta}} q_{\hat{\beta}}{}^{i}, \qquad (5.10f)$$

$$[P_{\hat{a}}, \bar{q}^{\hat{\alpha}}{}_{i}] = \frac{1}{2} (\gamma_{\hat{a}})^{\hat{\alpha}}{}_{\hat{\beta}} \bar{q}^{\hat{\beta}}{}_{i}, \qquad (5.10g)$$

$$[\mathbb{Y}, q_{\hat{\alpha}}{}^i] = -q_{\hat{\alpha}}{}^i, \qquad (5.10h)$$

$$\{q_{\hat{\alpha}}{}^{i}, \bar{q}^{\hat{\beta}}{}_{j}\} = 2\delta^{i}{}_{j}M^{\hat{a}\hat{b}}(\Sigma_{\hat{a}\hat{b}})_{\hat{\alpha}}{}^{\hat{\beta}} + 2\mathrm{i}\delta^{i}{}_{j}P^{\hat{a}}(\gamma_{\hat{a}})_{\hat{\alpha}}{}^{\hat{\beta}}$$
(5.10i)

$$-4\delta_{\hat{\alpha}}{}^{\beta}\mathbb{J}^{i}{}_{j} - \frac{\mathcal{N}}{\mathcal{N}}{}^{4}\delta^{i}{}_{j}\delta_{\hat{\alpha}}{}^{\beta}\mathbb{Y},$$
$$[\mathbb{J}^{i}{}_{j}, q_{\hat{\alpha}}{}^{k}] = \delta^{k}{}_{j}q_{\hat{\alpha}}{}^{i} - \frac{1}{\mathcal{N}}\delta^{i}{}_{j}q_{\hat{\alpha}}{}^{k}, \qquad (5.10j)$$

with $(\gamma_{\hat{a}})_{\hat{\alpha}}{}^{\hat{\beta}}$ defined as (A.58).

Any element n of the AdS superalgebra, $n \in \mathfrak{su}(2, 2|\mathcal{N})$, can be written as a linear combination of generators

$$n = \frac{1}{2} n^{\hat{a}\hat{b}} M_{\hat{a}\hat{b}} + n_i{}^j \mathbb{J}^i{}_j + i(n_{\mathsf{U}(1)}) \mathbb{Y}$$

+ $n^{\hat{a}} P_{\hat{a}} + i \left(n_i{}^{\hat{\alpha}} q_{\hat{\alpha}}{}^i + \bar{n}^i{}_{\hat{\alpha}} \bar{q}^{\hat{\alpha}}{}_i \right).$ (5.11)

This means that the vielbein can be decomposed as follows

$$\mathbf{E} = \mathbf{E}^{\hat{A}} K_{\hat{A}} = \mathbf{E}^{\hat{a}} P_{\hat{a}} + i \left(\mathbf{E}_{i}{}^{\hat{\alpha}} q_{\hat{\alpha}}{}^{i} + \bar{\mathbf{E}}^{i}{}_{\hat{\alpha}} \bar{q}^{\hat{\alpha}}{}_{i} \right), \qquad (5.12)$$

and the connection can be decomposed as

$$\Phi = \Phi^{\hat{I}} H_{\hat{I}} = \frac{1}{2} \Phi^{\hat{a}\hat{b}} M_{\hat{a}\hat{b}} + \Phi_i{}^j \mathbb{J}^i{}_j + i(\Phi_{\mathsf{U}(1)}) \mathbb{Y}.$$
(5.13)

In the above expressions we have used the definition

$$\mathbf{E}^{\hat{A}} = (\mathbf{E}^{\hat{a}}, \mathbf{E}_{i}^{\hat{\alpha}}, \bar{\mathbf{E}}^{i}_{\hat{\alpha}}).$$
(5.14)

The components of the connection, $\Phi^{\hat{I}}$, can be decomposed with respect to the basis $\mathbf{E}^{\hat{A}}$ to yield the superfields $\Phi_{\hat{A}}{}^{\hat{I}}$:

$$\mathbf{\Phi}^{\hat{I}} = \mathbf{E}^{\hat{A}} \Phi_{\hat{A}}{}^{\hat{I}} \,. \tag{5.15}$$

In accordance with the coset construction, we can introduce the curvature and torsion two forms as

$$\mathbf{R} := \mathrm{d}\boldsymbol{\Phi} - \boldsymbol{\Phi} \wedge \boldsymbol{\Phi}, \qquad \mathbf{T} := \mathrm{d}\mathbf{E} - \mathbf{E} \wedge \boldsymbol{\Phi} - \boldsymbol{\Phi} \wedge \mathbf{E}.$$
(5.16a)

The curvature and torsion can alternatively be expressed as

$$\mathbf{R} = (\mathbf{E} \wedge \mathbf{E})|_{\mathcal{H}}, \qquad \mathbf{T} = (\mathbf{E} \wedge \mathbf{E})|_{\mathcal{K}}.$$
 (5.16b)

Under group transformations, (5.7), they transform covariantly

$$\mathbf{R}' = h\mathbf{R}h^{-1}, \qquad \mathbf{T}' = h\mathbf{T}h^{-1}.$$
(5.17)

The components of the torsion and curvature two forms are defined via the relations

$$\mathbf{T} = \frac{1}{2} \mathbf{E}^{\hat{B}} \wedge \mathbf{E}^{\hat{A}} \left(\mathcal{T}_{\hat{A}\hat{B}}{}^{\hat{c}}P_{\hat{c}} + \mathrm{i} (\mathcal{T}_{\hat{A}\hat{B}k}{}^{\hat{\gamma}}q_{\hat{\gamma}}{}^{k} + \bar{\mathcal{T}}_{\hat{A}\hat{B}}{}^{k}{}_{\hat{\gamma}}\bar{q}^{\hat{\gamma}}{}_{k}) \right),$$
(5.18a)

$$\mathbf{R} = \frac{1}{2} \mathbf{E}^{\hat{B}} \wedge \mathbf{E}^{\hat{A}} \left(\frac{1}{2} \mathcal{R}_{\hat{A}\hat{B}}{}^{\hat{c}\hat{d}} M_{\hat{c}\hat{d}} + \mathcal{R}_{\hat{A}\hat{B}k}{}^{l} \mathbb{J}^{k}{}_{l} + \mathrm{i}(\mathcal{R}_{\mathsf{U}(1)})_{\hat{A}\hat{B}} \mathbb{Y} \right).$$
(5.18b)

Building on the approach utilised in [19, 22, 23], using the definitions (5.18) and the graded commutation relations of the generators, the non-vanishing components of the torsion and curvature can be determined. They are given by

$$\mathcal{T}^{i}{}_{\hat{\alpha}j}{}^{\hat{\beta}\hat{a}} = 2\mathrm{i}\delta^{i}{}_{j}(\gamma^{\hat{a}}){}_{\hat{\alpha}}{}^{\hat{\beta}}, \qquad (5.19\mathrm{a})$$

$$\mathcal{T}^{i}_{\hat{a}\,\hat{\alpha}j}{}^{\hat{\beta}} = -\frac{\mathrm{i}}{2}\delta^{i}{}_{j}(\gamma_{\hat{a}})_{\hat{\alpha}}{}^{\hat{\beta}}, \qquad (5.19\mathrm{b})$$

$$\mathcal{T}_{\hat{a}i}{}^{\hat{\alpha}j}{}_{\hat{\beta}} = -\frac{\mathrm{i}}{2} \delta^{j}{}_{i} (\gamma_{\hat{a}}){}^{\hat{\alpha}}{}_{\hat{\beta}} , \qquad (5.19c)$$

and

$$\mathcal{R}_{\hat{a}\hat{b}}{}^{\hat{c}\hat{d}} = -2\delta_{[\hat{a}}{}^{\hat{c}}\delta_{\hat{b}]}{}^{\hat{d}}, \qquad (5.20a)$$

$$\mathcal{R}_{j}{}^{\hat{\beta}i}{}_{\hat{\alpha}}{}^{\hat{a}\hat{b}} = 4\delta^{i}{}_{j}(\Sigma^{\hat{a}\hat{b}}){}_{\hat{\alpha}}{}^{\hat{\beta}}, \qquad (5.20b)$$

$$\mathcal{R}^{i}{}_{\hat{\alpha}j}{}^{\hat{\beta}}{}_{k}{}^{l} = -4\delta_{\hat{\alpha}}{}^{\hat{\beta}}\delta^{i}{}_{k}\delta^{l}{}_{j}, \qquad (5.20c)$$

$$\left(\mathcal{R}_{\mathsf{U}(1)}\right)^{i}{}_{\hat{\alpha}j}{}^{\hat{\beta}} = \frac{\mathrm{i}(\mathcal{N}-4)}{\mathcal{N}}\delta^{i}{}_{j}\delta_{\hat{\alpha}}{}^{\hat{\beta}}.$$
(5.20d)

5.2 Covariant derivatives

In this subsection we will use the results of the previous subsection to detail the construction of covariant derivatives on AdS superspace and determine their algebra.

Associated with each generator $K_{\hat{A}}$ is a vector field $E_{\hat{A}}$, and each generator $H_{\hat{I}}$ a connection $\Phi^{\hat{I}}$. To determine $E_{\hat{A}}$ it is useful to introduce the vielbein supermatrix $E^{\hat{A}}{}_{\hat{M}}$ used implicitly above:

$$\mathbf{E}^{\hat{A}} = E^{\hat{A}}_{\ \hat{M}} \boldsymbol{\mathcal{E}}^{\hat{M}} , \qquad (5.21a)$$

$$\boldsymbol{\mathcal{E}}^{\hat{M}} = E^{\hat{M}}{}_{\hat{A}} \mathbf{E}^{\hat{A}} \,, \tag{5.21b}$$

with $\boldsymbol{\mathcal{E}}^{\hat{M}}$ given by

$$\boldsymbol{\mathcal{E}}^{\hat{M}} = \left(\mathrm{d}x^{\hat{m}} \,, \mathrm{d}\theta_{i}^{\hat{\mu}} \,, \mathrm{d}\bar{\theta}^{i}_{\hat{\mu}} \right). \tag{5.22}$$

We then have

$$E_{\hat{A}} = E^{\hat{M}}{}_{\hat{A}}\partial_{\hat{M}}, \qquad \partial_{\hat{M}} = (\partial_{\hat{m}}, \frac{\partial}{\partial \theta_{i}{}^{\hat{\mu}}}, \frac{\partial}{\partial \bar{\theta}{}^{i}{}_{\hat{\mu}}}).$$
(5.23)

Now we define the covariant derivatives

$$\mathcal{D}_{\hat{A}} := \left(\mathcal{D}_{\hat{a}}, \mathcal{D}_{\hat{\alpha}}^{i}, \bar{\mathcal{D}}^{\hat{\alpha}}_{i}\right).$$
(5.24)

They take the form

$$\mathcal{D}_{\hat{A}} = E_{\hat{A}} + \frac{1}{2} \Phi_{\hat{A}}{}^{\hat{a}\hat{b}} M_{\hat{a}\hat{b}} + \Phi_{\hat{A}i}{}^{j} \mathbb{J}^{i}{}_{j} + \mathrm{i}(\Phi_{\mathsf{U}(1)})_{\hat{A}} \mathbb{Y} , \qquad (5.25)$$

and their algebra is given by

$$[\mathcal{D}_{\hat{A}}, \mathcal{D}_{\hat{B}}] = -\mathcal{T}_{\hat{A}\hat{B}}{}^{\hat{C}}\mathcal{D}_{\hat{C}} + \frac{1}{2}\mathcal{R}_{\hat{A}\hat{B}}{}^{\hat{c}\hat{d}}M_{\hat{a}\hat{b}} + \mathcal{R}_{\hat{A}\hat{B}\,k}{}^{l}\mathbb{J}^{k}{}_{l} + \mathrm{i}(\mathcal{R}_{\mathsf{U}(1)})_{\hat{A}\hat{B}}\mathbb{Y}.$$
(5.26)

Explicitly, we have

$$\left[\mathcal{D}_{\hat{a}}, \mathcal{D}_{\hat{b}}\right] = -M_{\hat{a}\hat{b}}, \qquad (5.27a)$$

$$[\mathcal{D}_{\hat{a}}, \mathcal{D}_{\hat{\alpha}}^{\ i}] = \frac{\mathrm{i}}{2} (\gamma_{\hat{a}})_{\hat{\alpha}}^{\ \hat{\beta}} \mathcal{D}_{\hat{\beta}}^{\ i}, \qquad (5.27\mathrm{b})$$

$$\left[\mathcal{D}_{\hat{a}}, \bar{\mathcal{D}}^{\hat{\alpha}}_{\ i}\right] = \frac{\mathrm{i}}{2} (\gamma_{\hat{a}})^{\hat{\alpha}}{}_{\hat{\beta}} \bar{\mathcal{D}}^{\hat{\beta}}{}_{\ i} , \qquad (5.27\mathrm{c})$$

$$\{\mathcal{D}_{\hat{\alpha}}{}^{i}, \bar{\mathcal{D}}^{\hat{\beta}}{}_{j}\} = -2\mathrm{i}\delta^{i}{}_{j}(\gamma^{\hat{a}})_{\hat{\alpha}}{}^{\hat{\beta}}\mathcal{D}_{\hat{a}} + 4\delta^{i}{}_{j}M_{\hat{\alpha}}{}^{\hat{\beta}}$$
(5.27d)

 $-4\delta_{\hat{\alpha}}{}^{\hat{\beta}}\mathbb{J}^{i}{}_{j}-\frac{(\mathcal{N}-4)}{\mathcal{N}}\delta^{i}{}_{j}\delta_{\hat{\alpha}}{}^{\hat{\beta}}\mathbb{Y}.$

The above algebra is presented in a different basis to the conventional approaches to 5D supergravity in the literature. We can reconcile them as follows. For $\mathcal{N} = 1$, let us combine the derivatives $\mathcal{D}_{\hat{\alpha}}$ and $\bar{\mathcal{D}}_{\hat{\alpha}}$ into the following $\mathsf{SU}(2)$ doublet

$$\tilde{\mathcal{D}}^{\underline{i}}_{\hat{\alpha}}, \qquad \tilde{\mathcal{D}}^{\underline{1}}_{\hat{\alpha}} = \mathcal{D}_{\hat{\alpha}}, \qquad \tilde{\mathcal{D}}^{\underline{2}}_{\hat{\alpha}} = \bar{\mathcal{D}}_{\hat{\alpha}}, \qquad \underline{i}, \underline{j} = \underline{1}, \underline{2}.$$
(5.28)

The corresponding local coordinates $\theta_{\hat{\alpha}}$ and $\bar{\theta}_{\hat{\alpha}}$ are combined in a similar way and satisfy a pseudo-Majorana condition. The SU(2) indices are raised and lowered by $\varepsilon^{\underline{ij}}$ and $\varepsilon_{\underline{ij}}$, $\varepsilon^{\underline{12}} = \varepsilon_{\underline{21}} = 1$, as $\tilde{\mathcal{D}}_{\hat{\alpha}\underline{i}} = \varepsilon_{\underline{ij}}\tilde{\mathcal{D}}_{\hat{\alpha}}^{\underline{j}}$. We then have the following relations

$$[\mathcal{D}_{\hat{a}}, \tilde{\mathcal{D}}_{\hat{\alpha}}^{\underline{i}}] = \frac{\mathrm{i}}{2} S^{\underline{i}}{\underline{j}} (\gamma_{\hat{a}})_{\hat{\alpha}}{}^{\hat{\beta}} \tilde{\mathcal{D}}_{\hat{\beta}}^{\underline{j}}, \qquad (5.29a)$$

$$\{\tilde{\mathcal{D}}^{\underline{i}}_{\hat{\alpha}}, \tilde{\mathcal{D}}^{\underline{j}}_{\hat{\beta}}\} = -2\mathrm{i}\varepsilon^{\underline{i}\underline{j}}(\gamma^{\hat{a}})_{\hat{\alpha}\hat{\beta}}\mathcal{D}_{\hat{a}} - 4S^{\underline{i}\underline{j}}M_{\hat{\alpha}\hat{\beta}} - 3\varepsilon_{\hat{\alpha}\hat{\beta}}\varepsilon^{\underline{i}\underline{j}}\mathbb{Y}, \qquad (5.29\mathrm{b})$$

with $S^{ij} = S^{ji} = (\sigma_3)^{ij}$. We can see that our algebra above coincides with that of [19], see equations (2.21).⁶

6 AdS superspaces with internal bosonic dimensions

The analysis of the previous sections made use of *even* supertwistors. One can also consider *odd* supertwistors,

$$\Xi = (\Xi_A) = \left(\frac{\Xi_{\hat{\alpha}}}{\Xi_i}\right) , \qquad (6.1)$$

with opposite Grassmann parities

$$\epsilon(\Xi_A) = 1 + \epsilon_A \pmod{2}. \tag{6.2}$$

The superspace $\operatorname{AdS}^{5|8\mathcal{N}}$ can be extended to $\operatorname{AdS}^{5|8\mathcal{N}} \times \mathbb{X}_m^{\mathcal{N}}$, where the internal space $\mathbb{X}_m^{\mathcal{N}}$ is realised in terms of $m \leq \mathcal{N}$ odd supertwistors. In the \mathcal{N} -extended case, there are many choices for the space $\mathbb{X}_m^{\mathcal{N}}$, corresponding to the various ways in which the m odd supertwistors $\mathbf{\Xi}^{\tilde{i}}, \tilde{i} = 1, \ldots, m$ can be constrained. In all cases, the odd supertwistors are required to be orthogonal to the quartet $\mathbf{T}^{\tilde{\mu}}$, which means that they contribute no Grassmann odd degrees of freedom. The remaining constraints on the odd supertwistors should be chosen in such a way that the space $\operatorname{AdS}^{5|8\mathcal{N}} \times \mathbb{X}_m^{\mathcal{N}}$ is a homogeneous space of $\operatorname{SU}(2, 2|\mathcal{N})$. This implies that $\mathbb{X}_m^{\mathcal{N}}$ is a homogeneous space of $\operatorname{SU}(\mathcal{N})$, and can therefore be realised as a coset space

$$\mathbb{X}_{m}^{\mathcal{N}} = \frac{\mathsf{SU}(\mathcal{N})}{\mathcal{G}},\tag{6.3}$$

⁶In [19], $S^{\underline{ij}}$ was denoted J^{ij} .

where \mathcal{G} is a subgroup of $SU(\mathcal{N})$. We will outline a few possibilities below.

Here we consider the superspaces $\mathrm{AdS}^{5|8\mathcal{N}} \times \mathbb{X}_1^{\mathcal{N}}$. Let us accompany the quartet $T^{\tilde{\mu}}$ with a single odd supertwistor, Ξ , subject to the orthonormality conditions

$$\langle \boldsymbol{T}^{\tilde{\mu}} | \boldsymbol{\Xi} \rangle = 0,$$
 (6.4a)

$$\langle \mathbf{\Xi} | \mathbf{\Xi} \rangle = -1 \,. \tag{6.4b}$$

The second condition implies that the body of Ξ_i is non-zero. For $\mathcal{N} = 1$, the internal space is given by S^1 , whereas for $\mathcal{N} > 2$ it can be shown that

$$\mathbb{X}_{1}^{\mathcal{N}} = \frac{\mathsf{SU}(\mathcal{N})}{\mathsf{SU}(\mathcal{N}-1)}, \qquad \mathcal{N} > 2.$$
(6.5)

As a consistency check, we can compare the dimension of the space (6.5) with the degrees of freedom of the supertwistor Ξ . The latter has $2\mathcal{N} - 1$ even degrees of freedom, because of the $2\mathcal{N}$ real components and the constraint (6.4b). This coincides with the $\mathcal{N}^2 - 1 - ((\mathcal{N} - 1)^2 - 1) = 2\mathcal{N} - 1$ dimensions of (6.5).

For $\mathcal{N} = 2$, we can modify the construction as follows. First, we relax the condition (6.4b). We still require that the body of Ξ_i is non-zero. Then, we define the supertwistor Ξ modulo the equivalence relation

$$\Xi \sim d \Xi, \qquad d \in \mathbb{C} - \{0\}. \tag{6.6}$$

The superspace obtained can be seen to be $AdS^{5|16} \times \mathbb{C}P^1$. It is a homogeneous space for SU(2,2|2).

6.1 The $\mathcal{N} = 4$, m = 4 case

As outlined above, for a given \mathcal{N} , the maximal number of odd supertwistors we can consider is \mathcal{N} . With this in mind, for the $\mathcal{N} = 4$ case let us introduce the quartet $\Xi^{\tilde{i}}$, $\tilde{i} = 1, 2, 3, 4$, subject to the orthonormality conditions

$$\langle \boldsymbol{T}^{\tilde{\mu}} | \boldsymbol{\Xi}^i \rangle = 0,$$
 (6.7a)

$$\langle \Xi^{\tilde{i}} | \Xi^{\tilde{j}} \rangle = -\delta^{\tilde{i}\tilde{j}} , \qquad (6.7b)$$

and defined modulo the equivalence relation

$$\boldsymbol{\Xi}^{\tilde{i}} \sim \boldsymbol{\Xi}^{\tilde{j}} \gamma_{\tilde{j}}^{\tilde{i}}, \qquad \gamma \in \mathsf{U}(1) \times \mathsf{USp}(4).$$
(6.8)

Making use of the equivalence relations (3.12b) and (6.8), the U(1) freedom on the even and odd supertwistors can be fixed such that the supermatrix formed by $(\mathbf{T}^{\tilde{\mu}}, \Xi^{\tilde{i}})$ has unit Berezinian. This means that they can be represented as

$$(\boldsymbol{T}^{\tilde{\mu}}, \boldsymbol{\Xi}^{\tilde{i}}) := \mathbb{T} = \left(\frac{\boldsymbol{T}_{\hat{\alpha}}{}^{\tilde{\mu}} \| \boldsymbol{\Xi}_{\hat{\alpha}}{}^{\tilde{i}}}{\boldsymbol{T}_{i}{}^{\tilde{\mu}} \| \boldsymbol{\Xi}_{i}{}^{\tilde{i}}} \right) \in \mathsf{SU}(2, 2|4) \,. \tag{6.9}$$

Furthermore, the equivalence relations that preserve this choice can be written as

$$\mathbb{T} \sim \mathbb{T}\left(\frac{\mathbf{e}^{\mathbf{i}\varphi}\lambda \mid \mathbf{0}}{0 \mid \gamma}\right), \qquad \mathbf{e}^{4\mathbf{i}\varphi} = \det(\gamma).$$
(6.10)

It can be shown that the resulting superspace is

$$\frac{\mathsf{SU}(2,2|4)}{\mathsf{USp}(2,2)\times\mathsf{U}(1)\times\mathsf{USp}(4)}.$$
(6.11)

To prove (6.11), we note that the simplest base point is

$$\mathbb{T}^{(0)} = \left(\begin{array}{c|c} \mathbb{1}_4 & 0\\ \hline 0 & \mathbb{1}_4 \end{array} \right) \,. \tag{6.12}$$

The stabiliser of $\mathbb{T}^{(0)}$ consists of those supergroup elements that map $\mathbb{T}^{(0)}$ to an equivalent point with respect to (6.10). Since the stabiliser of $\mathbb{T}^{(0)}$ is a subgroup of SU(2, 2|4), its elements have unit Berezinian and take the form

$$h = \left(\begin{array}{c|c} \underline{\mathrm{e}^{\mathrm{i}\varphi}\lambda} & 0\\ \hline 0 & \gamma \end{array}\right) \,, \tag{6.13a}$$

with

$$\lambda \in \mathsf{USp}(2,2), \quad \gamma \in \mathsf{U}(1) \times \mathsf{USp}(4), \qquad \mathrm{e}^{4\mathrm{i}\varphi} = \mathrm{det}(\gamma).$$
 (6.13b)

It is therefore isomorphic to

$$\mathsf{USp}(2,2) \times \mathsf{U}(1) \times \mathsf{USp}(4), \qquad (6.14)$$

and we arrive at (6.11). This situation describes the superspace $AdS^{5|32} \times X_4^4$, where the internal space is given by

$$\mathbb{X}_{4}^{4} = \frac{\mathsf{SU}(4)}{\mathsf{U}\mathsf{Sp}(4)} = \frac{\mathsf{SO}(6)}{\mathsf{SO}(5)},$$
 (6.15)

where the latter relation may be established using arguments analogous to those given in subsection 2.3. Thus the space \mathbb{X}_4^4 is equivalent to S^5 .

We can also introduce bi-supertwistors for the space \mathbb{X}_{4}^{4} . Let us introduce the following supermatrix

$$\boldsymbol{\zeta}_{AB} = \boldsymbol{\Xi}_A{}^{\tilde{i}} \boldsymbol{\Xi}_B{}^{\tilde{j}} C_{\tilde{i}\tilde{j}} , \qquad (6.16)$$

where $C_{\tilde{i}\tilde{j}}$ is given by

$$(C_{\tilde{i}\tilde{j}}) = \begin{pmatrix} \varepsilon & 0\\ 0 & \varepsilon \end{pmatrix}.$$
(6.17)

The supermatrix $\boldsymbol{\zeta}_{AB}$ is defined modulo the equivalence relation

$$\boldsymbol{\zeta}_{AB} \sim e^{i\varphi} \boldsymbol{\zeta}_{AB}, \qquad \varphi \in \mathbb{R},$$
(6.18)

which is the remnant of the equivalence transformations (6.8). Because the quartet $\Xi^{\tilde{i}}$ has opposite Grassmann parity to $T^{\tilde{\mu}}$, the supermatrix ζ_{AB} has the following symmetry property

$$\boldsymbol{\zeta}_{AB} = -(-1)^{(1+\epsilon_A)(1+\epsilon_B)} \boldsymbol{\zeta}_{BA} \,. \tag{6.19}$$

It also satisfies the conditions

$$\bar{\boldsymbol{X}}^{AC}\boldsymbol{\zeta}_{CB} = 0, \qquad (6.20a)$$

$$\bar{\boldsymbol{\zeta}}^{AB}\boldsymbol{\zeta}_{BA} = 4\,,\tag{6.20b}$$

where $\bar{\boldsymbol{\zeta}}^{AB}$ is defined in the same way as (3.22).

The bi-supertwistors X_{AB} and ζ_{AB} can be combined in the following way. Let us introduce the supermatrix

$$\mathcal{X} = \left(\begin{array}{c|c} \mathfrak{C}_{\tilde{\mu}\tilde{\nu}} & 0\\ \hline 0 & C_{\tilde{i}\tilde{j}} \end{array} \right) \in \mathsf{SU}(2,2|4) \,. \tag{6.21}$$

Then, we find the following

$$\Upsilon := \mathbb{T} \mathcal{X} \mathbb{T}^{\mathrm{T}} = \ell^{-1} \mathbf{X} + \boldsymbol{\zeta}, \qquad \mathbf{X} = (\mathbf{X}_{AB}), \quad \boldsymbol{\zeta} = (\boldsymbol{\zeta}_{AB}).$$
(6.22)

Due to the U(1) freedom enjoyed by X and ζ , the supermatrix Υ is defined modulo

$$\Upsilon \sim e^{i\varphi}\Upsilon, \qquad \varphi \in \mathbb{R}.$$
 (6.23)

Furthermore, it satisfies the following condition

$$\Upsilon^{\dagger}\Omega\Upsilon = \Omega. \tag{6.24}$$

7 Conclusion

In this paper we presented the supersymmetric generalisation of the embedding formalism for AdS₅. Specifically, we proposed supertwistor and bi-supertwistor realisations for AdS^{5|8 \mathcal{N}}. Making use of the coset construction, we studied the superspace geometry of AdS^{5|8 \mathcal{N}}, and showed that our results are consistent with the known case for the $\mathcal{N} = 1$ superspace AdS^{5|8 \mathcal{N}}. On AdS^{5|8 \mathcal{N}}, we introduced superspace Poincaré coordinates and computed the bi-supertwistor in this coordinate system. We also extended the (bi-)supertwistor realisations to superspaces of the form AdS^{5|8 \mathcal{N}} × $\mathbb{X}_m^{\mathcal{N}}$, where $\mathbb{X}_m^{\mathcal{N}}$ is a homogeneous space of SU(\mathcal{N}). In particular, for the $\mathcal{N} = m = 4$ case, we derived the necessary constraints for the (bi-)supertwistors to describe the AdS₅ × S⁵ superspace (6.11).

Our formalism is useful for constructing superinvariants. Indeed, one of the main results of our paper is the manifestly supersymmetric two-point function, (3.38), which we used to propose a model, (3.39), for a massive superparticle in $AdS^{5|8N}$. More generally, associated with 2n points of $AdS^{5|8N}$ is the following invariant:

$$\bar{\boldsymbol{X}}^{A_1B_1}(1)\boldsymbol{X}_{B_1A_2}(2)(-1)^{\epsilon_{A_2}}\bar{\boldsymbol{X}}^{A_2B_2}(3)\boldsymbol{X}_{B_2A_3}(4)\dots(-1)^{\epsilon_{A_n}}\bar{\boldsymbol{X}}^{A_nB_n}(2n-1)\boldsymbol{X}_{B_nA_1}(2n).$$
(7.1)

Such invariants are expected to appear in correlation functions of quantum fields in AdS_5 .

It would be interesting to recast the type IIB superstring action in $AdS_5 \times S^5$ background [62] in terms of the bi-supertwistors X_{AB} and ζ_{AB} . This will be discussed elsewhere.

Acknowledgements:

The work of SMK is supported in part by the Australian Research Council, project DP230101629. The work of NEK is supported by the Australian Government Research Training Program Scholarship.

A Spinors in diverse dimensions

In this appendix we collect those results of the spinor formalisms in 3 + 1, 4 + 2 and 4 + 1 dimensions, which are used in our work.

A.1 Spinors in 3+1 dimensions

Our two-component spinor conventions coincide with those in the books [63, 64], to which the reader is referred to for a complete discussion. Consider the space $\mathbb{R}^{3,1}$ equipped with metric

$$\eta_{ab} = \operatorname{diag}(-1, +1, +1, +1), \qquad a = 0, 1, 2, 3.$$
(A.1)

The gamma matrices in 3 + 1 dimensions satisfy

$$\{\gamma_a, \gamma_b\} = -2\eta_{ab}\mathbb{1}_4, \qquad (A.2)$$

and can be chosen as

$$\gamma_a = \begin{pmatrix} 0 & \sigma_a \\ \tilde{\sigma}_a & 0 \end{pmatrix}, \tag{A.3}$$

with

$$\sigma_a = (\mathbb{1}_2, \vec{\sigma}) \equiv (\sigma_a)_{\alpha \dot{\alpha}}, \qquad \tilde{\sigma}_a = (\mathbb{1}_2, -\vec{\sigma}) \equiv (\tilde{\sigma}_a)^{\dot{\alpha} \alpha}.$$
(A.4)

The matrix $\gamma_5 = -i\gamma_0\gamma_1\gamma_2\gamma_3$ then takes the form

$$\gamma_5 = \begin{pmatrix} \mathbb{1}_2 & 0\\ 0 & -\mathbb{1}_2 \end{pmatrix} \,. \tag{A.5}$$

The Lorentz generators in the spinor representation are

$$\Sigma_{ab} = -\frac{1}{4} [\gamma_a, \gamma_b] \,. \tag{A.6}$$

They are given by

$$(\Sigma_{ab})_{\hat{\alpha}}^{\ \hat{\beta}} = \begin{pmatrix} (\sigma_{ab})_{\alpha}^{\ \beta} & 0\\ 0 & (\tilde{\sigma}_{ab})^{\dot{\alpha}}_{\ \dot{\beta}} \end{pmatrix}, \qquad (A.7a)$$

$$(\sigma_{ab})_{\alpha}{}^{\beta} = -\frac{1}{4} (\sigma_a \tilde{\sigma}_b - \sigma_b \tilde{\sigma}_a)_{\alpha}{}^{\beta}, \qquad (A.7b)$$

$$(\tilde{\sigma}_{ab})^{\dot{\alpha}}{}_{\dot{\beta}} = -\frac{1}{4} (\tilde{\sigma}_a \sigma_b - \tilde{\sigma}_b \sigma_a)^{\dot{\alpha}}{}_{\dot{\beta}} \,. \tag{A.7c}$$

Two-component spinor indices are raised and lowered using the spinor metrics

$$\varepsilon^{\alpha\beta} = -\varepsilon^{\beta\alpha} , \qquad \varepsilon_{\alpha\beta} = -\varepsilon_{\beta\alpha} , \qquad \varepsilon^{12} = \varepsilon_{21} = 1 ; \qquad (A.8a)$$

$$\varepsilon^{\dot{\alpha}\dot{\beta}} = -\varepsilon^{\dot{\beta}\dot{\alpha}} , \qquad \varepsilon_{\dot{\alpha}\dot{\beta}} = -\varepsilon_{\dot{\beta}\dot{\alpha}} , \qquad \varepsilon^{\dot{12}} = \varepsilon_{2\dot{1}} = 1 , \qquad (A.8b)$$

$$\dot{\epsilon}^{\dot{\alpha}\beta} = -\varepsilon^{\beta\dot{\alpha}} , \qquad \varepsilon_{\dot{\alpha}\dot{\beta}} = -\varepsilon_{\dot{\beta}\dot{\alpha}} , \qquad \varepsilon^{12} = \varepsilon_{\dot{2}\dot{1}} = 1 , \qquad (A.8b)$$

by the rules:

$$\psi^{\alpha} := \varepsilon^{\alpha\beta}\psi_{\beta}, \qquad \psi_{\alpha} = \varepsilon_{\alpha\beta}\psi^{\beta}; \qquad \bar{\phi}^{\dot{\alpha}} := \varepsilon^{\dot{\alpha}\dot{\beta}}\bar{\phi}_{\dot{\beta}}, \qquad \bar{\phi}_{\dot{\alpha}} = \varepsilon_{\dot{\alpha}\dot{\beta}}\bar{\phi}^{\dot{\beta}}. \tag{A.9}$$

The charge conjugation matrix, \mathcal{C} , is defined by

$$\mathcal{C}^{-1}\gamma_a \mathcal{C} = -\gamma_a^{\mathrm{T}} \implies \mathcal{C}^{-1}\gamma_5 \mathcal{C} = \gamma_5^{\mathrm{T}}.$$
 (A.10a)

It has the properties

$$\mathcal{C}^{\dagger} = \mathcal{C}^{\mathrm{T}} = -\mathcal{C} = \mathcal{C}^{-1}, \qquad (A.10b)$$

and can be chosen as

$$\mathcal{C} = \begin{pmatrix} \varepsilon_{\alpha\beta} & 0\\ 0 & \varepsilon^{\dot{\alpha}\dot{\beta}} \end{pmatrix} . \tag{A.10c}$$

A number of useful properties are satisfied by the matrices σ_a and $\tilde{\sigma}_a$. They are:

$$(\sigma_a \tilde{\sigma}_b + \sigma_b \tilde{\sigma}_a)_{\alpha}{}^{\beta} = -2\eta_{ab} \delta_{\alpha}{}^{\beta}, \qquad (A.11a)$$

$$\left(\tilde{\sigma}_a \sigma_b + \tilde{\sigma}_b \sigma_a\right)^{\dot{\alpha}}{}_{\dot{\beta}} = -2\eta_{ab} \delta^{\dot{\alpha}}{}_{\dot{\beta}}, \qquad (A.11b)$$

$$\operatorname{Tr}(\sigma_a \tilde{\sigma}_b) = -2\eta_{ab} \,, \tag{A.11c}$$

$$Tr(\sigma_a \tilde{\sigma}_b) = -2\eta_{ab}, \qquad (A.11c)$$
$$(\sigma^a)_{\alpha \dot{\alpha}} (\tilde{\sigma}_a)^{\dot{\beta}\beta} = -2\delta_{\alpha}{}^{\beta}\delta_{\dot{\alpha}}{}^{\dot{\beta}}. \qquad (A.11d)$$

Associated with a four-vector $x^a \in \mathbb{R}^{3,1}$ are the Hermitian matrices

$$x_{\alpha\dot{\alpha}} = x^a (\sigma_a)_{\alpha\dot{\alpha}}, \quad x^{\dot{\alpha}\alpha} = x^a (\tilde{\sigma}_a)^{\dot{\alpha}\alpha}.$$
 (A.12)

It is sometimes convenient to use the condensed notation

$$x := (x_{\alpha\dot{\alpha}}), \qquad \tilde{x} := (x^{\dot{\alpha}\alpha}). \tag{A.13}$$

A.2Spinors in 4+2 dimensions

In this appendix we will detail the spinor conventions for $SO_0(4,2)$. Consider the space $\mathbb{R}^{4,2}$ parametrised by coordinates $x^{\underline{a}}\,,\ \underline{a}=0,1,2,3,4,5\,,$ and with metric

$$\eta_{\underline{ab}} = \operatorname{diag}(-1, +1, +1, +1, -1). \tag{A.14}$$

The corresponding gamma matrices, $\Gamma_{\underline{a}}$, obey the anti-commutation relations

$$\{\Gamma_{\underline{a}}, \Gamma_{\underline{b}}\} = -2\eta_{\underline{a}\underline{b}}\mathbb{1}_8.$$
(A.15)

We can choose $\Gamma_{\underline{a}}$ in the so-called Weyl repersentation

$$\Gamma_{\underline{a}} = \begin{pmatrix} 0 & \underline{\Sigma}_{\underline{a}} \\ \underline{\tilde{\Sigma}}_{\underline{a}} & 0 \end{pmatrix}, \qquad (A.16)$$

with

$$\underline{\Sigma}_{\underline{a}} = (\mathrm{i}\gamma_a \,, \gamma_5, \mathbb{1}_4) \equiv (\underline{\Sigma}_{\underline{a}})_{\hat{\alpha}\underline{\hat{\beta}}}, \tag{A.17a}$$

$$\underline{\tilde{\Sigma}}_{\underline{a}} = (-\mathrm{i}\gamma_a \,, -\gamma_5 \,, \mathbb{1}_4) \equiv (\underline{\tilde{\Sigma}}_{\underline{a}})^{\underline{\hat{\alpha}}\widehat{\beta}} \,. \tag{A.17b}$$

Here γ_a and γ_5 are the gamma matrices in 3 + 1 dimensions, see appendix A.1.

In accordance with (A.15), the matrices $\underline{\Sigma}$ and $\underline{\tilde{\Sigma}}$ obey the following relations

$$\underline{\underline{\Sigma}}_{\underline{a}}\underline{\underline{\tilde{\Sigma}}}_{\underline{b}} + \underline{\underline{\Sigma}}_{\underline{b}}\underline{\underline{\tilde{\Sigma}}}_{\underline{a}} = -2\eta_{\underline{a}\underline{b}}\mathbb{1}_4, \qquad \underline{\underline{\tilde{\Sigma}}}_{\underline{a}}\underline{\underline{\Sigma}}_{\underline{b}} + \underline{\underline{\tilde{\Sigma}}}_{\underline{b}}\underline{\underline{\Sigma}}_{\underline{a}} = -2\eta_{\underline{a}\underline{b}}\mathbb{1}_4.$$
(A.18)

The Hermitian conjugation properties of $\Gamma_{\underline{a}}$ are

$$\Gamma_{\underline{a}}^{\dagger} = \Gamma_0 \Gamma_5 \Gamma_{\underline{a}} \Gamma_0 \Gamma_5 , \qquad (A.19)$$

which implies the following Hermitian conjugation properties for the matrices $\underline{\Sigma}$ and $\underline{\tilde{\Sigma}}$

$$(\underline{\Sigma}_{\underline{a}})^{\dagger} = \gamma_0 \underline{\tilde{\Sigma}}_{\underline{a}} \gamma_0 , \qquad (\underline{\tilde{\Sigma}}_{\underline{a}})^{\dagger} = \gamma_0 \underline{\Sigma}_{\underline{a}} \gamma_0 .$$
 (A.20)

The Dirac spinor representation is generated by

$$\mathfrak{J}_{\underline{ab}} = -\frac{1}{4} [\Gamma_{\underline{a}}, \Gamma_{\underline{b}}] = \begin{pmatrix} \underline{\Sigma}_{\underline{ab}} & 0\\ 0 & \underline{\tilde{\Sigma}}_{\underline{ab}} \end{pmatrix} .$$
(A.21)

Here we have defined

$$\underline{\Sigma}_{\underline{a}\underline{b}} := -\frac{1}{4} \left(\underline{\Sigma}_{\underline{a}} \underline{\tilde{\Sigma}}_{\underline{b}} - \underline{\Sigma}_{\underline{b}} \underline{\tilde{\Sigma}}_{\underline{a}} \right) \equiv (\underline{\Sigma}_{\underline{a}\underline{b}})_{\hat{\alpha}}{}^{\hat{\beta}}, \qquad (A.22a)$$

$$\underline{\tilde{\Sigma}}_{\underline{a}\underline{b}} := -\frac{1}{4} \left(\underline{\tilde{\Sigma}}_{\underline{a}} \underline{\Sigma}_{\underline{b}} - \underline{\tilde{\Sigma}}_{\underline{b}} \underline{\Sigma}_{\underline{a}} \right) \equiv (\underline{\tilde{\Sigma}}_{\underline{a}\underline{b}})^{\underline{\hat{\alpha}}}_{\underline{\hat{\beta}}}.$$
(A.22b)

Given (A.19), the Hermitian conjugation properties of $\mathfrak{J}_{\underline{ab}}$ are

$$(\mathfrak{J}_{\underline{a}\underline{b}})^{\dagger} = \Gamma_0 \Gamma_5(\mathfrak{J}_{\underline{a}\underline{b}}) \Gamma_0 \Gamma_5 \,. \tag{A.23}$$

For the matrices $\underline{\Sigma}_{\underline{a}\underline{b}}$ and $\underline{\tilde{\Sigma}}_{\underline{a}\underline{b}}$ we have the following

$$(\underline{\Sigma}_{\underline{a}\underline{b}})^{\dagger} = -\gamma_0(\underline{\Sigma}_{\underline{a}\underline{b}})\gamma_0, \qquad (\underline{\tilde{\Sigma}}_{\underline{a}\underline{b}})^{\dagger} = -\gamma_0(\underline{\tilde{\Sigma}}_{\underline{a}\underline{b}})\gamma_0.$$
(A.24)

A Dirac spinor looks like

$$\Psi = \begin{pmatrix} \psi \\ \phi \end{pmatrix}, \qquad \psi = (\psi_{\hat{\alpha}}), \qquad \phi = (\phi^{\underline{\hat{\alpha}}}). \tag{A.25}$$

Its conjugate is defined as

$$\bar{\Psi} := -i\Psi^{\dagger}\Gamma_{0}\Gamma_{5} = \left(\psi^{\dagger}\gamma_{0}, -\phi^{\dagger}\gamma_{0}\right), \quad \psi^{\dagger}\gamma_{0} \equiv (\bar{\psi}^{\hat{\alpha}}), \quad \phi^{\dagger}\gamma_{0} \equiv (\bar{\phi}_{\underline{\hat{\alpha}}}).$$
(A.26)

An infinitesimal $SO_0(4,2)$ transformation acts on Ψ and $\overline{\Psi}$ as

$$\delta \Psi = \frac{1}{2} \omega^{\underline{a}\underline{b}} \mathfrak{J}_{\underline{a}\underline{b}} \Psi , \qquad (A.27a)$$

$$\delta \bar{\Psi} = -\frac{1}{2} \bar{\Psi} \omega^{\underline{a}\underline{b}} \mathfrak{J}_{\underline{a}\underline{b}} \,. \tag{A.27b}$$

This representation is the sum of two irreducible representations, as is clear from (A.21). The twistor (left Weyl spinor) representation is associated with spinors of the form

$$\Psi_{\rm L} = \begin{pmatrix} \psi \\ 0 \end{pmatrix}, \qquad \psi = (\psi_{\hat{\alpha}}), \qquad (A.28)$$

such that

$$\Gamma_7 \Psi_{\rm L} = \Psi_{\rm L} \,, \tag{A.29}$$

where Γ_7 is defined as

$$\Gamma_7 := -i\Gamma_0\Gamma_1\Gamma_2\Gamma_3\Gamma_4\Gamma_5 \equiv \begin{pmatrix} \mathbb{1}_4 & 0\\ 0 & -\mathbb{1}_4 \end{pmatrix}.$$
(A.30)

The Dirac conjugate of $\Psi_{\rm L}$ is given by

$$\bar{\Psi}_{\rm L} = \left(\bar{\psi}, 0\right), \qquad \bar{\psi} := \psi^{\dagger} \gamma_0 = (\bar{\psi}^{\hat{\alpha}}), \qquad (A.31)$$

and transforms according to the dual twistor representation. Following (A.27), the infinitesimal SO(4, 2) transformation laws of Ψ_L and $\bar{\Psi}_L$ are given by

$$\delta\psi_{\hat{\alpha}} = \frac{1}{2}\omega^{\underline{a}\underline{b}}(\underline{\Sigma}_{\underline{a}\underline{b}})_{\hat{\alpha}}{}^{\hat{\beta}}\psi_{\hat{\beta}}, \qquad (A.32a)$$

$$\delta \bar{\psi}^{\hat{\alpha}} = -\frac{1}{2} \bar{\psi}^{\hat{\beta}} \omega^{\underline{a}\underline{b}} (\underline{\Sigma}_{\underline{a}\underline{b}})_{\hat{\beta}}{}^{\hat{\alpha}} . \tag{A.32b}$$

In the main body of this paper, we use the matrices $\Sigma_{\underline{a}}$ and $\tilde{\Sigma}_{\underline{a}}$. These are related to the matrices $\underline{\Sigma}_{\underline{a}}$ and $\underline{\tilde{\Sigma}}_{\underline{a}}$ as described below. We introduce the 8×8 charge conjugation matrix, \mathscr{C} , satisfying

$$\mathscr{C}^{-1}\Gamma_{\underline{a}}\mathscr{C} = -\Gamma_{\underline{a}}^{\mathrm{T}}.$$
(A.33)

It can be chosen as

$$\mathscr{C} = \begin{pmatrix} 0 & \gamma_5 \mathcal{C} \\ -\gamma_5 \mathcal{C} & 0 \end{pmatrix} \equiv \begin{pmatrix} 0 & \mathscr{C}_{\hat{\alpha}}^{\hat{\beta}} \\ \mathscr{C}^{\hat{\alpha}}_{\hat{\beta}} & 0 \end{pmatrix}, \qquad (A.34)$$

with \mathcal{C} as the 3 + 1 dimensional charge conjugation matrix, (A.10). The inverse of \mathscr{C} is

$$\mathscr{C}^{-1} = \begin{pmatrix} 0 & -\mathcal{C}^{-1}\gamma_5 \\ \mathcal{C}^{-1}\gamma_5 & 0 \end{pmatrix} \equiv \begin{pmatrix} 0 & (\mathscr{C}^{-1})^{\hat{\alpha}}_{\hat{\beta}} \\ (\mathscr{C}^{-1})_{\hat{\alpha}}^{\hat{\beta}} & 0 \end{pmatrix}.$$
 (A.35)

We also have

$$\mathscr{C}^{-1}\mathfrak{J}_{\underline{a}\underline{b}}\mathscr{C} = -\mathfrak{J}_{\underline{a}\underline{b}}^{\mathrm{T}}.$$
 (A.36)

Using \mathscr{C} , we can define the charge conjugate spinor

$$\Psi_{\mathscr{C}} := \mathscr{C}\bar{\Psi}^{\mathrm{T}} \,. \tag{A.37}$$

Its infinitesimal transformation rule is

$$\delta\Psi_{\mathscr{C}} = \frac{1}{2}\omega^{\underline{a}\underline{b}}\mathfrak{J}_{\underline{a}\underline{b}}\Psi_{\mathscr{C}},\qquad(A.38)$$

which coincides with (A.27a). Consider the contents of the charge conjugate spinor

$$\Psi_{\mathscr{C}} = \mathscr{C}\bar{\Psi}^{\mathrm{T}}$$

$$= \begin{pmatrix} 0 & \mathscr{C}_{\hat{\alpha}}^{\hat{\beta}} \\ \mathscr{C}^{\hat{\alpha}}_{\hat{\beta}} & 0 \end{pmatrix} \begin{pmatrix} \bar{\psi}^{\hat{\beta}} \\ -\bar{\phi}_{\hat{\beta}} \end{pmatrix}$$

$$= \begin{pmatrix} -\mathscr{C}_{\hat{\alpha}}^{\hat{\beta}}\bar{\phi}_{\hat{\beta}} \\ \mathscr{C}^{\hat{\alpha}}_{\hat{\beta}}\bar{\psi}^{\hat{\beta}} \end{pmatrix}.$$
(A.39)

Since $\Psi_{\mathscr{C}}$ transforms as (A.38), $\bar{\phi}_{\hat{\alpha}} := \mathscr{C}_{\hat{\alpha}}{}^{\hat{\beta}}\bar{\phi}_{\hat{\beta}} = \bar{\phi}_{\hat{\beta}}\mathscr{C}^{\hat{\beta}}_{\hat{\alpha}}$ transforms like a twistor. This means we can convert underlined indices into not-underlined indices.

We have so far considered the twistor representation. The other (right Weyl spinor) representation is equivalent to the dual twistor representation, as

$$\phi^{\hat{\alpha}} := (\mathscr{C}^{-1})^{\hat{\alpha}}{}_{\underline{\hat{\beta}}} \phi^{\underline{\hat{\beta}}} = \phi^{\underline{\hat{\beta}}} (\mathscr{C}^{-1})_{\underline{\hat{\beta}}}^{\hat{\alpha}}$$
(A.40)

transforms like a dual twistor. This also follows from (A.36), which yields

$$-(\underline{\Sigma}_{\underline{a}\underline{b}}^{\mathrm{T}})^{\hat{\alpha}}{}_{\hat{\beta}} = (\mathscr{C}^{-1})^{\hat{\alpha}}{}_{\underline{\hat{\gamma}}}(\underline{\tilde{\Sigma}}_{\underline{a}\underline{b}})^{\underline{\hat{\gamma}}}{}_{\underline{\hat{\delta}}}\mathscr{C}^{\underline{\hat{\delta}}}{}_{\hat{\beta}}.$$
(A.41)

Now, the matrices $\Sigma_{\underline{a}}$ and $\tilde{\Sigma}_{\underline{a}}$ (with twistor indices) are defined as

$$(\Sigma_{\underline{a}})_{\hat{\alpha}\hat{\beta}} := (\underline{\Sigma}_{\underline{a}})_{\hat{\alpha}\hat{\underline{\gamma}}} \mathscr{C}^{\hat{\underline{\gamma}}}_{\hat{\beta}}, \qquad (\tilde{\Sigma}_{\underline{a}})^{\hat{\alpha}\hat{\beta}} := (\mathscr{C}^{-1})^{\hat{\alpha}}_{\hat{\underline{\gamma}}} (\underline{\tilde{\Sigma}}_{\underline{a}})^{\hat{\underline{\gamma}}\hat{\beta}}.$$
(A.42)

They are antisymmetric

$$(\Sigma_{\underline{a}})_{\hat{\alpha}\hat{\beta}} = -(\Sigma_{\underline{a}})_{\hat{\beta}\hat{\alpha}}, \qquad (\tilde{\Sigma}_{\underline{a}})^{\hat{\alpha}\hat{\beta}} = -(\tilde{\Sigma}_{\underline{a}})^{\hat{\beta}\hat{\alpha}}, \qquad (A.43)$$

and satisfy the relations

$$\Sigma_{\underline{a}}\tilde{\Sigma}_{\underline{b}} + \Sigma_{\underline{b}}\tilde{\Sigma}_{\underline{a}} = -2\eta_{\underline{a}\underline{b}}\mathbb{1}_4, \qquad \tilde{\Sigma}_{\underline{a}}\Sigma_{\underline{b}} + \tilde{\Sigma}_{\underline{b}}\Sigma_{\underline{a}} = -2\eta_{\underline{a}\underline{b}}\mathbb{1}_4.$$
(A.44)

It is useful here to give the explicit form of these matrices:

$$\Sigma_{a} = \begin{pmatrix} 0 & \mathrm{i}\sigma_{a}\varepsilon \\ -\mathrm{i}\tilde{\sigma}_{a}\varepsilon^{-1} & 0 \end{pmatrix}, \qquad \tilde{\Sigma}_{a} = \begin{pmatrix} 0 & \mathrm{i}\varepsilon\sigma_{a} \\ -\mathrm{i}\varepsilon^{-1}\tilde{\sigma}_{a} & 0 \end{pmatrix},$$
$$\Sigma_{4} = \begin{pmatrix} -\varepsilon^{-1} & 0 \\ 0 & -\varepsilon \end{pmatrix}, \qquad \tilde{\Sigma}_{4} = \begin{pmatrix} \varepsilon & 0 \\ 0 & \varepsilon^{-1} \end{pmatrix}, \qquad (A.45)$$
$$\Sigma_{5} = \begin{pmatrix} -\varepsilon^{-1} & 0 \\ 0 & \varepsilon \end{pmatrix}, \qquad \tilde{\Sigma}_{5} = \begin{pmatrix} -\varepsilon & 0 \\ 0 & \varepsilon^{-1} \end{pmatrix}.$$

Since the Lorentz generators (A.22a) are invariant under the replacement $\underline{\Sigma}_{\underline{a}} \to \underline{\Sigma}_{\underline{a}}, \underline{\tilde{\Sigma}}_{\underline{a}} \to \underline{\tilde{\Sigma}}_{\underline{a}},$ we can unambiguously write them without an underline. Explicitly, they take the form

$$(\Sigma_{ab})_{\hat{\alpha}}{}^{\hat{\beta}} = \begin{pmatrix} (\sigma_{ab})_{\alpha}{}^{\beta} & 0\\ 0 & (\tilde{\sigma}_{ab})^{\dot{\alpha}}{}_{\dot{\beta}} \end{pmatrix}, \qquad (\Sigma_{a4})_{\hat{\alpha}}{}^{\hat{\beta}} = \begin{pmatrix} 0 & -\frac{i}{2}(\sigma_a)_{\alpha\dot{\beta}}\\ \frac{i}{2}(\tilde{\sigma}_a)^{\dot{\alpha}\beta} & 0 \end{pmatrix}, \qquad (A.46a)$$

$$(\Sigma_{a5})_{\hat{\alpha}}{}^{\hat{\beta}} = \begin{pmatrix} 0 & -\frac{i}{2}(\sigma_a)_{\alpha\dot{\beta}} \\ -\frac{i}{2}(\tilde{\sigma}_a)^{\dot{\alpha}\beta} & 0 \end{pmatrix}, \qquad (\Sigma_{45})_{\hat{\alpha}}{}^{\hat{\beta}} = \begin{pmatrix} -\frac{1}{2}\delta_{\alpha}{}^{\beta} & 0 \\ 0 & \frac{1}{2}\delta^{\dot{\alpha}}_{\dot{\beta}} \end{pmatrix}.$$
(A.46b)

Here we note that we can express a general Lie algebra element as

$$\omega = \frac{1}{2}\omega^{\underline{a}\underline{b}}(\Sigma_{\underline{a}\underline{b}}), \qquad \omega^{\underline{a}\underline{b}} = (\omega^{\underline{a}\underline{b}})^*, \qquad (A.47a)$$

$$\gamma_0 \omega^\dagger + \omega \gamma_0 = 0. \tag{A.47b}$$

We then have the following

$$\frac{1}{2}\omega^{\underline{ab}}(\Sigma_{\underline{ab}})_{\hat{\alpha}}{}^{\hat{\beta}} = \left(\begin{array}{c} \frac{1}{2}\omega^{ab}(\sigma_{ab})_{\alpha}{}^{\beta} - \frac{1}{2}\omega^{45}\delta_{\alpha}{}^{\beta} & -\frac{1}{2}(\omega^{a4} + \omega^{a5})(\sigma_{a})_{\alpha\dot{\beta}} \\ \frac{1}{2}(\omega^{a4} - \omega^{a5})(\tilde{\sigma}_{a})^{\dot{\alpha}\beta} & \frac{1}{2}\omega^{ab}(\tilde{\sigma}_{ab})^{\dot{\alpha}}{}_{\dot{\beta}} + \frac{1}{2}\omega^{45}\delta^{\dot{\alpha}}{}_{\dot{\beta}} \end{array}\right).$$
(A.48)

The matrices (A.42) satisfy a number of additional useful properties. Their Hermitian conjugation properties can be read off using those of $\underline{\Sigma}_{\underline{a}}$ and $\underline{\tilde{\Sigma}}_{\underline{a}}$ as well as the properties of C and γ_5 :

$$(\Sigma_{\underline{a}})^{\dagger} = \gamma_0 \tilde{\Sigma}_{\underline{a}} \gamma_0 \,. \tag{A.49}$$

We have used $C^{\dagger} = C^{-1}$, $\gamma_0^{\mathrm{T}} = \gamma_0$ and (A.10). They also satisfy the following completeness relations

$$\frac{1}{2}\varepsilon_{\hat{\alpha}\hat{\beta}\hat{\gamma}\hat{\delta}}(\tilde{\Sigma}^{\underline{a}})^{\hat{\gamma}\hat{\delta}} = (\Sigma^{\underline{a}})_{\hat{\alpha}\hat{\beta}}, \qquad (A.50a)$$

$$\frac{1}{2}\varepsilon^{\hat{\alpha}\hat{\beta}\hat{\gamma}\hat{\delta}}(\Sigma^{\underline{a}})_{\hat{\gamma}\hat{\delta}} = (\tilde{\Sigma}^{\underline{a}})^{\hat{\alpha}\hat{\beta}}, \qquad (A.50b)$$

$$(\Sigma^{\underline{a}})_{\hat{\alpha}\hat{\beta}}(\Sigma_{\underline{a}})_{\hat{\gamma}\hat{\delta}} = 2\varepsilon_{\hat{\alpha}\hat{\beta}\hat{\gamma}\hat{\delta}}.$$
 (A.50c)

We can now establish the one-to-one correspondence between complex vectors $V^{\underline{a}}$ in $\mathbb{R}^{4,2}$ and antisymmetric bi-twistors $V_{\hat{\alpha}\hat{\beta}} = -V_{\hat{\beta}\hat{\alpha}}$,

$$V_{\hat{\alpha}\hat{\beta}} = V^{\underline{a}}(\Sigma_{\underline{a}})_{\hat{\alpha}\hat{\beta}}, \qquad V^{\underline{a}} = \frac{1}{4}(\tilde{\Sigma}^{\underline{a}})^{\hat{\alpha}\hat{\beta}}V_{\hat{\alpha}\hat{\beta}}, \qquad (A.51a)$$

$$V^{\hat{\alpha}\hat{\beta}} = V^{\underline{a}} (\tilde{\Sigma}_{\underline{a}})^{\hat{\alpha}\hat{\beta}}, \qquad V^{\underline{a}} = \frac{1}{4} (\Sigma^{\underline{a}})_{\hat{\alpha}\hat{\beta}} V^{\hat{\alpha}\hat{\beta}}.$$
(A.51b)

In the above, we have

$$V^{\hat{\alpha}\hat{\beta}} = \frac{1}{2} \varepsilon^{\hat{\alpha}\hat{\beta}\hat{\gamma}\hat{\delta}} V_{\hat{\gamma}\hat{\delta}} , \qquad (A.52)$$

consistent with (A.50). We will now introduce the condensed notation

$$V := (V_{\hat{\alpha}\hat{\beta}}), \qquad \tilde{V} := (V^{\hat{\alpha}\hat{\beta}}). \tag{A.53}$$

We wish to investigate how the bi-twistor associated with $V^{\underline{a}}$ is related to that associated with $\overline{V^{\underline{a}}}$, the complex conjugate of the six-vector $V^{\underline{a}}$. Taking the Hermitian conjugate of V, using (A.49), we find

$$V = V^{\underline{a}}(\Sigma_{\underline{a}}) \to V^{\dagger} = \bar{V}^{\underline{a}}\gamma_0(\tilde{\Sigma}_{\underline{a}})\gamma_0$$

$$= \gamma_0 \tilde{\bar{V}}\gamma_0 .$$
(A.54)

Inserting $V^{\underline{a}} = \overline{V}^{\underline{a}}$ into the above, we find

$$V^{\dagger} = \gamma_0 \tilde{V} \gamma_0 \implies \tilde{V} = \gamma_0 V^{\dagger} \gamma_0 .$$
 (A.55a)

Now, identifying γ_0 with Ω in (2.1) and expressing the reality condition in index notation, we have

$$\Omega^{\hat{\alpha}\hat{\gamma}}(V^{\dagger})_{\hat{\gamma}\hat{\delta}}\Omega^{\hat{\delta}\hat{\beta}} = \frac{1}{2}\varepsilon^{\hat{\alpha}\hat{\beta}\hat{\gamma}\hat{\delta}}V_{\hat{\gamma}\hat{\delta}}.$$
(A.55b)

A.3 Spinors in 4+1 dimensions

Consider the space $\mathbb{R}^{4,1}$ equipped with metric

$$\eta_{\hat{a}\hat{b}} = \text{diag}(-1, +1, +1, +1, +1), \qquad \hat{a} = 0, 1, 2, 3, 4.$$
(A.56)

The corresponding gamma matrices satisfy

$$\{\gamma_{\hat{a}}, \gamma_{\hat{b}}\} = -2\eta_{\hat{a}\hat{b}}\mathbb{1}_4, \qquad (\gamma_{\hat{a}})^{\dagger} = \gamma_0\gamma_{\hat{a}}\gamma_0, \qquad (A.57)$$

and can be chosen as

$$(\gamma_a)_{\hat{\alpha}}{}^{\hat{\beta}} = \begin{pmatrix} 0 & (\sigma_a)_{\alpha\dot{\beta}} \\ (\tilde{\sigma}_a)^{\dot{\alpha}\beta} & 0 \end{pmatrix}, \qquad (\gamma_4)_{\hat{\alpha}}{}^{\hat{\beta}} = \begin{pmatrix} -\mathrm{i}\delta_{\alpha}{}^{\beta} & 0 \\ 0 & \mathrm{i}\delta^{\dot{\alpha}}{}_{\dot{\beta}} \end{pmatrix}.$$
(A.58)

They are related to the Lorentz spinor generators in $\mathbb{R}^{4,2}$, (A.46), as

$$(\Sigma_{\hat{a}5})_{\hat{\alpha}}{}^{\hat{\beta}} = -\frac{\mathrm{i}}{2}(\gamma_{\hat{a}})_{\hat{\alpha}}{}^{\hat{\beta}}.$$
(A.59)

To minimise confusion between the 4+1 and 4+2 cases, we will denote the charge conjugation matrix $\mathbb{R}^{4,1}$ as C. The charge conjugation matrix is defined as

$$C\gamma_{\hat{a}}C^{-1} = (\gamma_{\hat{a}})^{\mathrm{T}}, \qquad C = (\varepsilon^{\hat{\alpha}\hat{\beta}}) = \begin{pmatrix} \varepsilon^{\alpha\beta} & 0\\ 0 & -\varepsilon_{\dot{\alpha}\dot{\beta}} \end{pmatrix}.$$
 (A.60)

The antisymmetric matrices $\varepsilon^{\hat{\alpha}\hat{\beta}}$ and its inverse, $\varepsilon_{\hat{\alpha}\hat{\beta}}$, are used to raise and lower the fourcomponent spinors indices in $\mathbb{R}^{4,1}$.

Now we can define the Lorentz generators for the spinor representation,

$$\Sigma_{\hat{a}\hat{b}} := -\frac{1}{4} [\gamma_{\hat{a}}, \gamma_{\hat{b}}], \qquad (A.61)$$

and choose them in the form

$$(\Sigma_{ab})_{\hat{\alpha}}{}^{\hat{\beta}} = \begin{pmatrix} (\sigma_{ab})_{\alpha}{}^{\beta} & 0\\ 0 & (\tilde{\sigma}_{ab})^{\dot{\alpha}}{}_{\dot{\beta}} \end{pmatrix}, \qquad (\Sigma_{a4})_{\hat{\alpha}}{}^{\hat{\beta}} = \begin{pmatrix} 0 & -\frac{i}{2}(\sigma_a)_{\alpha\dot{\beta}}\\ \frac{i}{2}(\tilde{\sigma}_a)^{\dot{\alpha}\beta} & 0 \end{pmatrix}.$$
(A.62)

We can see that these generators coincide with the 4 + 1 subset of those given in (A.46a). We therefore find an explicit form for a general element of the algebra by 'switching off' the $\omega^{\hat{a}5}$ components of (A.48):

$$\frac{1}{2}\omega^{\hat{a}\hat{b}}(\Sigma_{\hat{a}\hat{b}})_{\hat{\alpha}}{}^{\hat{\beta}} = \left(\frac{\frac{1}{2}\omega^{ab}(\sigma_{ab})_{\alpha}{}^{\beta}}{\frac{1}{2}(\omega^{a4})(\tilde{\sigma}_{a})^{\dot{\alpha}\beta}} \right) \frac{1}{2}\omega^{ab}(\tilde{\sigma}_{ab})^{\dot{\alpha}}{}_{\dot{\beta}}\right) . \tag{A.63}$$

Given (A.57) and (A.60), we have the following properties

$$(\Sigma_{\hat{a}\hat{b}})^{\dagger} = -\gamma_0(\Sigma_{\hat{a}\hat{b}})\gamma_0, \qquad (\Sigma_{\hat{a}\hat{b}})^{\mathrm{T}} = -C(\Sigma_{\hat{a}\hat{b}})C^{-1}.$$
 (A.64)

Writing an element of the Lie algebra $\mathfrak{so}(4,1)$ as (A.63):

$$\omega := \frac{1}{2} \omega^{\hat{a}\hat{b}}(\Sigma_{\hat{a}\hat{b}}), \qquad \omega^{\hat{a}\hat{b}} = (\omega^{\hat{a}\hat{b}})^*, \qquad (A.65)$$

we find the following master equations

$$C\omega + \omega^{\mathrm{T}}C = 0, \qquad (A.66a)$$

$$\gamma_0 \omega + \omega^{\dagger} \gamma_0 = 0. \qquad (A.66b)$$

These correspond to the group equations

$$g^{\mathrm{T}}Cg = C, \qquad (A.67a)$$

$$g^{\dagger}\gamma_0 g = \gamma_0 \,. \tag{A.67b}$$

This group is known as $\mathsf{USp}(2,2)$.

A Dirac spinor in $\mathbb{R}^{4,1}$, $\Psi = (\Psi_{\hat{\alpha}})$, and its Dirac conjugate, $\bar{\Psi} = (\bar{\Psi}^{\hat{\alpha}}) = \Psi^{\dagger} \gamma_0$, look like

$$\Psi_{\hat{\alpha}} = \begin{pmatrix} \psi_{\alpha} \\ \bar{\phi}^{\dot{\alpha}} \end{pmatrix}, \qquad \bar{\Psi}^{\hat{\alpha}} = \begin{pmatrix} \phi^{\alpha} , \bar{\psi}_{\dot{\alpha}} \end{pmatrix}.$$
(A.68)

We can combine $\bar{\Psi}^{\hat{\alpha}} = (\phi^{\alpha}, \bar{\psi}_{\dot{\alpha}})$ and $\Psi^{\hat{\alpha}} = \varepsilon^{\hat{\alpha}\hat{\beta}}\Psi_{\hat{\beta}} = (\psi^{\alpha}, -\bar{\phi}_{\dot{\alpha}})$ into an $\mathsf{SU}(2)$ doublet:

$$\Psi_{\underline{i}}^{\hat{\alpha}} = \left(\Psi_{\underline{i}}^{\alpha}, -\bar{\Psi}_{\dot{\alpha}\underline{i}}\right), \qquad \left(\Psi_{\underline{i}}^{\alpha}\right)^{*} = \bar{\Psi}^{\dot{\alpha}\underline{i}}, \qquad \underline{i} = 1, 2, \qquad (A.69)$$

such that $\Psi_{\underline{1}}^{\alpha} = \phi^{\alpha}$ and $\Psi_{\underline{2}}^{\alpha} = \psi^{\alpha}$. The $\mathsf{SU}(2)$ indices are raised and lowered in the usual way: $\Psi^{\hat{\alpha}\underline{i}} = \varepsilon^{\underline{i}\underline{j}}\Psi_{\underline{j}}^{\hat{\alpha}}$. The spinor $\Psi^{\underline{i}} = (\Psi_{\hat{\alpha}}^{\underline{i}})$ satisfies the pseudo-Majorana condition

$$\bar{\Psi}_{\underline{i}}^{\mathrm{T}} = C\Psi_{\underline{i}} \,. \tag{A.70}$$

References

- [1] W. Nahm, "Supersymmetries and their representations," Nucl. Phys. B 135 (1978), 149.
- [2] I. A. Bandos, E. Ivanov, J. Lukierski and D. Sorokin, "On the superconformal flatness of AdS superspaces," JHEP 06 (2002), 040 [arXiv:hep-th/0205104 [hep-th]].
- [3] S. M. Kuzenko and G. Tartaglino-Mazzucchelli, "Conformally flat supergeometry in five dimensions," JHEP 06 (2008), 097 [arXiv:0804.1219 [hep-th]].

- [4] S. M. Kuzenko and G. Tartaglino-Mazzucchelli, "Super-Weyl invariance in 5D supergravity," JHEP 04 (2008), 032 [arXiv:0802.3953 [hep-th]].
- [5] V. G. Kac, "Lie superalgebras," Adv. Math. 26, 8 (1977).
- [6] T. Kugo and K. Ohashi, "Supergravity tensor calculus in 5D from 6D," Prog. Theor. Phys. 104, 835 (2000) [hep-ph/0006231].
- T. Kugo and K. Ohashi, "Off-shell d = 5 supergravity coupled to matter-Yang-Mills system," Prog. Theor. Phys. 105, 323 (2001) [hep-ph/0010288].
- [8] T. Fujita and K. Ohashi, "Superconformal tensor calculus in five dimensions," Prog. Theor. Phys. 106, 221 (2001) [hep-th/0104130].
- T. Kugo and K. Ohashi, "Gauge and non-gauge tensor multiplets in 5D conformal supergravity," Prog. Theor. Phys. 108, 1143 (2003) [hep-th/0208082].
- [10] E. Bergshoeff, S. Cucu, M. Derix, T. de Wit, R. Halbersma and A. Van Proeyen, "Weyl multiplets of N=2 conformal supergravity in five-dimensions," JHEP 0106, 051 (2001) [hep-th/0104113].
- [11] E. Bergshoeff, S. Cucu, T. de Wit, J. Gheerardyn, R. Halbersma, S. Vandoren and A. Van Proeyen, "Superconformal N = 2, D = 5 matter with and without actions," JHEP **0210**, 045 (2002) [hep-th/0205230].
- [12] E. Bergshoeff, S. Cucu, T. de Wit, J. Gheerardyn, S. Vandoren and A. Van Proeyen, "N = 2 supergravity in five dimensions revisited," Class. Quant. Grav. 21, 3015 (2004) [hep-th/0403045].
- [13] D. Butter, S. M. Kuzenko, J. Novak and G. Tartaglino-Mazzucchelli, "Conformal supergravity in five dimensions: New approach and applications," JHEP 02, 111 (2015) [arXiv:1410.8682 [hep-th]].
- [14] P. S. Howe, "Off-shell N=2 and N=4 supergravity in five-dimensions," in *Quantum Structure of Space and Time*, M. J. Duff and C. J. Isham, Cambridge Univ. Press, 1982, p. 239.
- [15] P. S. Howe and U. Lindström, "The supercurrent in five dimensions," Phys. Lett. B 103, 422 (1981).
- [16] S. M. Kuzenko and G. Tartaglino-Mazzucchelli, "Five-dimensional superfield supergravity," Phys. Lett. B 661, 42 (2008) [arXiv:0710.3440 [hep-th]].
- [17] S. M. Kuzenko and G. Tartaglino-Mazzucchelli, "5D supergravity and projective superspace," JHEP 02, 004 (2008) [arXiv:0712.3102 [hep-th]].
- [18] S. M. Kuzenko, J. Novak and G. Tartaglino-Mazzucchelli, "Symmetries of curved superspace in five dimensions," JHEP 10 (2014), 175 [arXiv:1406.0727 [hep-th]].
- [19] S. M. Kuzenko and G. Tartaglino-Mazzucchelli, "Five-dimensional N = 1 AdS superspace: Geometry, off-shell multiplets and dynamics," Nucl. Phys. B 785 (2007), 34-73 [arXiv:0704.1185 [hep-th]].
- [20] J. Bagger and C. Xiong, "AdS₅ Supersymmetry in N=1 superspace," JHEP 07 (2011), 119 [arXiv:1105.4852 [hep-th]].
- [21] S. M. Kuzenko and G. Tartaglino-Mazzucchelli, "Supertwistor realisations of AdS superspaces," Eur. Phys. J. C 82 (2022) no.2, 146 [arXiv:2108.03907 [hep-th]].
- [22] S. M. Kuzenko and K. Turner, "Embedding formalism for (p, q) AdS superspaces in three dimensions," JHEP 06 (2023), 142 [arXiv:2303.03082 [hep-th]].

- [23] N. E. Koning, S. M. Kuzenko and E. S. N. Raptakis, "Embedding formalism for N-extended AdS superspace in four dimensions," JHEP 11 (2023), 063 [arXiv:2308.04135 [hep-th]].
- [24] A. Sinkovics and E. P. Verlinde, "A Six dimensional view on twistors," Phys. Lett. B 608 (2005), 142-150 [arXiv:hep-th/0410014 [hep-th]].
- [25] T. Adamo, D. Skinner and J. Williams, "Twistor methods for AdS₅," JHEP 08 (2016), 167 [arXiv:1607.03763 [hep-th]].
- [26] R. Roiban and W. Siegel, "Superstrings on AdS(5) x S**5 supertwistor space," JHEP 11 (2000), 024 [arXiv:hep-th/0010104 [hep-th]].
- [27] D. P. Sorokin, "Superbranes and superembeddings," Phys. Rept. 329 (2000), 1-101 [arXiv:hep-th/9906142 [hep-th]].
- [28] I. A. Bandos and D. P. Sorokin, "Superembedding approach to superstrings and super-p-branes," [arXiv:2301.10668 [hep-th]].
- [29] A. Ferber, "Supertwistors and Conformal Supersymmetry," Nucl. Phys. B 132 (1978), 55-64
- [30] Y. I. Manin, "Holomorphic supergeometry and Yang-Mills superfields," J. Sov. Math. 30, 1927 (1985); Gauge Field Theory and Complex Geometry, Springer, Berlin, 1988.
- [31] M. Kotrla and J. Niederle, "Supertwistors and Superspace," Czech. J. Phys. B 35 (1985), 602
- [32] P. S. Howe and M. I. Leeming, "Harmonic superspaces in low dimensions," Class. Quant. Grav. 11 (1994), 2843-2852 [arXiv:hep-th/9408062 [hep-th]].
- [33] S. M. Kuzenko, J. H. Park, G. Tartaglino-Mazzucchelli and R. Unge, "Off-shell superconformal nonlinear sigma-models in three dimensions," JHEP 01 (2011), 146 [arXiv:1011.5727 [hep-th]].
- [34] A. A. Rosly, "Gauge Fields in Superspace and Twistors," Class. Quant. Grav. 2 (1985), 693-699
- [35] J. Lukierski and A. Nowicki, "General Superspaces From Supertwistors," Phys. Lett. B 211 (1988), 276-280
- [36] G. G. Hartwell and P. S. Howe, "(N, p, q) harmonic superspace," Int. J. Mod. Phys. A 10 (1995), 3901-3920 [arXiv:hep-th/9412147 [hep-th]].
- [37] P. S. Howe and G. G. Hartwell, "A Superspace survey," Class. Quant. Grav. 12 (1995), 1823-1880
- [38] S. M. Kuzenko, "On compactified harmonic/projective superspace, 5-D superconformal theories, and all that," Nucl. Phys. B 745 (2006), 176-207 [arXiv:hep-th/0601177 [hep-th]].
- [39] S. M. Kuzenko, "Conformally compactified Minkowski superspaces revisited," JHEP 1210, 135 (2012) [arXiv:1206.3940 [hep-th]].
- [40] E. I. Buchbinder, S. M. Kuzenko and I. B. Samsonov, "Superconformal field theory in three dimensions: Correlation functions of conserved currents," JHEP 06 (2015), 138 [arXiv:1503.04961 [hep-th]].
- [41] N. Boulanger, Y. Herfray and N. Parrini, "Conformal boundaries of Minkowski superspace and their super cuts," JHEP 02 (2024), 177 [arXiv:2312.11222 [hep-th]].
- [42] P. S. Howe and U. Lindström, "Local supertwistors and conformal supergravity in six dimensions," Proc. Roy. Soc. Lond. A 476 (2020) no.2243, 20200683 [arXiv:2008.10302 [hep-th]].

- [43] P. S. Howe and U. Lindström, "Superconformal geometries and local twistors," JHEP 04 (2021), 140 [arXiv:2012.03282 [hep-th]].
- [44] P. Claus, M. Gunaydin, R. Kallosh, J. Rahmfeld and Y. Zunger, "Supertwistors as quarks of SU(2, 2|4)," JHEP 05 (1999), 019 [arXiv:hep-th/9905112 [hep-th]].
- [45] P. Claus, J. Rahmfeld and Y. Zunger, "A Simple particle action from a twistor parametrization of AdS(5)," Phys. Lett. B 466 (1999), 181-189 [arXiv:hep-th/9906118 [hep-th]].
- [46] P. Claus, R. Kallosh and J. Rahmfeld, "BRST quantization of a particle in AdS(5)," Phys. Lett. B 462 (1999), 285-293 [arXiv:hep-th/9906195 [hep-th]].
- [47] I. A. Bandos, J. Lukierski, C. Preitschopf and D. P. Sorokin, "OSp supergroup manifolds, superparticles and supertwistors," Phys. Rev. D 61 (2000), 065009 [arXiv:hep-th/9907113 [hep-th]].
- [48] Y. Zunger, "Twistors and actions on coset manifolds," Phys. Rev. D 62 (2000), 024030 [arXiv:hep-th/0001072 [hep-th]].
- [49] M. Cederwall, "Geometric construction of AdS twistors," Phys. Lett. B 483 (2000), 257-263 [arXiv:hep-th/0002216 [hep-th]].
- [50] M. Cederwall, "AdS twistors for higher spin theory," AIP Conf. Proc. 767 (2005) no.1, 96-105 [arXiv:hep-th/0412222 [hep-th]].
- [51] A. S. Arvanitakis, A. E. Barns-Graham and P. K. Townsend, "Anti-de Sitter Particles and Manifest (Super)Isometries," Phys. Rev. Lett. 118 (2017) no.14, 141601 [arXiv:1608.04380 [hep-th]].
- [52] A. S. Arvanitakis, A. E. Barns-Graham and P. K. Townsend, "Twistor description of spinning particles in AdS," JHEP 01 (2018), 059 [arXiv:1710.09557 [hep-th]].
- [53] D. V. Uvarov, "Supertwistor formulation for massless superparticle in $AdS_5 \times S^5$ superspace," Nucl. Phys. B **936** (2018), 690-713 [arXiv:1807.08318 [hep-th]].
- [54] D. V. Uvarov, "Multitwistor mechanics of massless superparticle on $AdS_5 \times S^5$ superbackground," Nucl. Phys. B **950** (2020), 114830 [arXiv:1907.13613 [hep-th]].
- [55] R. Penrose, "Twistor algebra," J. Math. Phys. 8, 345 (1967); R. Penrose and M. A. H. MacCallum, "Twistor theory: An approach to the quantization of fields and space-time," Phys. Rept. 6, 241 (1972).
- [56] I. E. Segal, Mathematical Cosmology and Extragalactic Astronomy, Academic Press, New York, 1976.
- [57] I. T. Todorov, Conformal Description of Spinning Particles, Springer, Berlin, 1986.
- [58] R. S. Ward and R. O. Wells, Twistor Geometry and Field Theory, Cambridge University Press, Cambridge, 1991.
- [59] S. M. Kuzenko, S. L. Lyakhovich, A. Y. Segal and A. A. Sharapov, "Massive spinning particle on anti-de Sitter space," Int. J. Mod. Phys. A 11 (1996), 3307-3330 [arXiv:hep-th/9509062 [hep-th]].
- [60] B. S. DeWitt, Supermanifolds, Cambridge University Press, Cambridge, 1992.
- [61] S. M. Kuzenko and I. N. McArthur, "Goldstone multiplet for partially broken superconformal symmetry," Phys. Lett. B 522 (2001), 320-326 [arXiv:hep-th/0109183 [hep-th]].

- [62] R. R. Metsaev and A. A. Tseytlin, "Type IIB superstring action in AdS(5) x S**5 background," Nucl. Phys. B 533 (1998), 109-126 [arXiv:hep-th/9805028 [hep-th]].
- [63] J. Wess and J. Bagger, Supersymmetry and Supergravity, Princeton Univ. Press, 1992.
- [64] I. L. Buchbinder and S. M. Kuzenko, Ideas and Methods of Supersymmetry and Supergravity or a Walk Through Superspace Bristol, UK: IOP (1998) 656 p.