# Ajtai's theorem for $T_2^2(R)$ and pebble games with backtracking

Eitetsu Ken $^1$ & Mykyta Narusevych $^2$ 

June 18, 2024

<sup>1</sup>affiliation: Graduate School of Mathematical Sciences, the University of Tokyo supported by: JSPS KAKENHI Grant Number 22KJ1121, Grant-in-Aid for JSPS Fellows, and FoPM program at the University of Tokyo email: yeongcheol-kwon@g.ecc.u-tokyo.ac.jp

<sup>2</sup>affiliation: Department of Mathematics and Physics, Charles University email: mykyta.narusevych@matfyz.cuni.cz supported by the Charles University project PRIMUS/21/SCI/014, Charles University Research Centre program No. UNCE/24/SCI/022 and GA UK project No. 246223

#### Abstract

We introduce a pebble game extended by backtracking options for one of the two players (called Prover) and reduce the provability of the pigeonhole principle for a generic predicate R in the bounded arithmetic  $T_2^2(R)$  to the existence of a particular kind of winning strategy (called oblivious) for Prover in the game.

While the unprovability of the said principle in  $T_2^2(R)$  is an immediate consequence of a celebrated theorem of Ajtai (which deals with a stronger theory  $T_2(R)$ ), up-to-date no methods working for  $T_2^2(R)$  directly (in particular without switching lemma) are known.

Although the full analysis of the introduced pebble game is left open, as a first step towards resolving it, we restrict ourselves to a simplified version of the game. In this case, Prover can use only two pebbles and move in an extremely oblivious way. Besides, a series of backtracks can be made only once during a play. Under these assumptions, we show that no strategy of Prover can be winning.

### 1 Introduction

Ajtai's theorem states the unprovability of the pigeonhole principle in bounded arithmetic and a super-polynomial lower bound for it in constant depth Frege systems (proved in [2], and later strengthened by [19], [22] and [7]). The theorem and its variations ([3], [7], and [8]) are among the strongest lower-bound results in Proof Complexity and independence results for bounded arithmetics. One of the main tools to prove them is Switching Lemma for so-called (shallow) *PHP*-trees. (See [15] or [17] for a clear exposition.) All known frameworks for understanding the proof of Ajtai's theorem, such as forcing with random variables [16] and partially definable forcing [6] in addition to original literature, depend heavily on it.

However, we still do not know any appropriate counterpart of Switching Lemma for resolving long-standing open problems such as:  $I\Delta_0(R)$  (or  $AC^0$ -Frege) v.s. the weak pigeonhole principle or  $V^0(2)$  (or  $AC^0(2)$ -Frege) v.s. the (injective) pigeonhole principle (see [17] for detailed presentations of the problems). Even when we compare the injective pigeonhole principle and the full version of onto pigeonhole principle, Switching Lemma seems not that flexible ([14]). Therefore, it would be very interesting if we could provide another proof of Ajtai's theorem, which is Switching-Lemma-free.

In this article, to make the presentation as simple as possible, we focus on  $T_2^2(R) \not\vdash ontoPHP_n^{n+1}(R)$ , which is an immediate consequence of Ajtai's theorem, and its propositional translation  $LK_{2+\frac{1}{2},O(1)}^*$  v.s. ontoPHP. We present a trial to obtain a Switching-Lemma-free proof of the statement.

Our approach is an extension of pebble-game analysis for resolution ([5]), which is powerful enough to obtain relatively easy independence result:  $T_2^1(R) \not\vdash ontoPHP_n^{n+1}(R)$ . We define a game  $\mathcal{G}_2$ , which is an extended pebble game allowing Spoiler in [5] (in our terminology, **Prover**) to **backtrack** the game record bringing a part of the current partial assignment back to the past. Since **Prover** wins the game easily if there is no restriction for strategies, we introduce **oblivious strategies** of **Prover** and show that if  $T_2^2(R) \vdash ontoPHP_n^{n+1}(R)$ , then some oblivious strategy of **Prover** beats any strategy of the opponent (in our terminology, **Delayer**) in the game  $\mathcal{G}_2$  of appropriate parameters.

Although we leave the complete analysis of  $\mathcal{G}_2$  open, at least we analyze a toy case of  $\mathcal{G}_2$  setting the parameters to easy ones, further restricting oblivious strategies to "automaton-like" strategies, and allowing **Prover** to backtrack just once.

We believe one advantage of our approach compared to tackle the functional version of the problem (that is, we adopt fresh function symbols f and g instead of R to describe the bijective pigeonhole principle) via CPLS, the NP-search problem corresponding to  $T_2^2$  ([18]), is that it is enough for us to consider partial assignments of small size (of order  $\log(n)^{O(1)}$ ). It is at least not immediate for CPLS and 1-Ref(RES) in [18] when we consider dag-like  $R(\log)$  refutations which are not necessarily narrow. Furthermore, our approach differs from those considering  $PLS^{NP}$  and GLS problems discussed in [13],

the  $\Sigma_2^b$ -search problems corresponding to  $T_2^2$ ; roughly speaking, in our game approach, we consider a concrete way to answer each NP-query in  $PLS^{NP}$  computations, at the sake of taking exponentially many steps in general (cf. Corollary 4.7) but following a reasonable restriction (cf. Definition 4.10). On the other hand, the analysis of the game is much more complex than usual pebble games because of the backtracking option.

The article is organized as follows:

§2 presents the convention and the formal system we shall follow and use.

§3 briefly mentions the base case of Ajtai's theorem to locate our work in the course of previous research explicitly:  $T_2^1(R)$  does not prove the pigeonhole principle for relation R

§4 introduces our game notion  $\mathcal{G}_2$ , which is, roughly speaking, a pebbling game allowing **Prover** to backtrack the records of the game. The notion can be regarded as a transposition of Buss's witnessing argument ([9], [11]) to our settings. We connect formal proofs of the pigeonhole principle in  $T_2^2(R)$  and winning strategies of **Prover** for  $\mathcal{G}_2$ .

In §5, we consider a very limited case of  $\mathcal{G}_2$ : **Prover** has only two pebbles, has to use extremely oblivious strategies, and is allowed to backtrack only once, while **Delayer** is of full power. We show that, in this setting, **Delayer** wins  $\mathcal{G}_2$ .

§6 is Acknowledgement.

§7 is an appendix. We sketch the verification of our restriction  $p\Sigma_i$  and  $s\Sigma_i$  of the form of propositional formulae adopted in the paper, present another equivalent definition of  $\mathcal{G}_2$ , and discuss another proof strategy on who wins the toy case of the game  $\mathcal{G}_2$  analyzed in §5.

## 2 Preliminaries

In this article, we consider a variant of  $LK_{2+\frac{1}{2}}^*$ , a propositional counterpart of the relativized bounded arithmetic  $T_2^2(R)$ , whose proofs are of constant heights and consist of cedents of constant cardinalities. See Chapter 8 of [17] for the precise formulation and established translations. Note that it is as powerful as quasipolynomial-sized  $R(\log)$ -proofs when depth-2 formulae are in concern. See Chapter 10 of [17].

To clarify the convention, we explicitly describe our setup.

For each  $i \in \mathbb{N}$ , we adopt the following common notations:  $|i| := \lceil \log_2(i+1) \rceil$ , the length of the binary representation of i, and  $[i] := \{1, \ldots, i\}$  ( $[0] := \emptyset$ ).

For a family  $\mathcal{F}$  of sets and  $s \in \mathbb{N}$ ,  $\mathcal{F}_{\leq s}$  denotes the subfamily of  $\mathcal{F}$  collecting the elements of cardinality  $\leq s$ .

We adopt  $\neg$ , binary  $\lor$ ,  $\land$  and unbounded  $\bigvee$ ,  $\bigwedge$  as propositional connectives, 0 and 1 as propositional constants, and consider only propositional formulae of negation normal form. Given a formula  $\varphi$ ,  $\overline{\varphi}$  denotes the canonical negation normal form of  $\neg \varphi$ . It is called **the complement of**  $\varphi$ .

For a propositional formula  $\varphi$ ,  $|\varphi|$  denotes the size of  $\varphi$ , say, the number of occurrences of variables and connectives in  $\varphi$ . The precise definition does not matter as long as the conventions are polynomially related. See also Chapter 1 of [17].

The distinction between unbounded  $\bigvee$ ,  $\bigwedge$  and binary  $\bigvee$ ,  $\land$  are adopted since the following variation of  $\Sigma_d^{S,t}$  given at the end of section 3.4 of [17] is in our mind: a propositional formula  $\varphi$  is  $p\Sigma_{i+\frac{1}{2}}(z)$   $(i,z\in\mathbb{N})$  if and only if it has the following form:

$$\varphi = \bigvee_{\substack{j_1 \in J_1 \ j_2 \in J_2}} \bigvee_{\substack{j_1, \dots, j_i, \\ \text{exactly } i \text{-times}}} \psi_{j_1, \dots, j_i},$$

where

$$|\varphi| \le z \& \forall k \in [i]. J_k \ne \emptyset$$

and each  $\psi_{\vec{j}}$  is  $\bigvee$ - and  $\bigwedge$ -free and satisfies  $|\psi_{\vec{j}}| \leq \log z$ . Note that if  $\varphi$  is  $p\Sigma_{i+\frac{1}{2}}(z)$ , then such i is unique since we distinguish  $\bigvee$ ,  $\bigwedge$  from  $\bigvee$ ,  $\wedge$ .  $p\Pi_{i+\frac{1}{2}}(z)$  is defined similarly, switching the roles of  $\bigvee$  and  $\bigwedge$ .

We say  $\varphi$  is  $s\Sigma_{i+\frac{1}{2}}(z)$  if it is  $p\Sigma_{i'+\frac{1}{2}}(z)$  for some  $0 \le i' \le i$ , or it is  $p\Pi_{i'+\frac{1}{2}}(z)$  for some  $0 \le i' < i$ . Similarly for  $s\Pi_{i+\frac{1}{2}}(z)$ . We are particularly interested in the case  $z = 2^{|n|^{O(1)}}$  for parameters n and  $i \le 2$ .

We clarify our treatment of trees in our descriptions of sequent calculus and a game.

**Definition 2.1.** Let  $b, h \in \mathbb{N}$ .  $[b]^{\leq h}$  denotes the set of finite sequences on [b] with length  $\leq h$  (including the empty sequence).

For  $v, w \in [b]^{\leq h}$ ,  $v \subseteq w$  means that v is a initial segment of w (or w is an extension of v).

The length of v is denoted by height(v).

Given  $v \in [b]^{\leq h}$  and  $w \in [b]^{\leq k}$ , v \* w denotes the concatenation lying in  $[b]^{\leq h+k}$ .

For  $v = (a_1, \ldots, a_h) \in [b]^h$  and  $k \in \mathbb{N}$ , set

$$v_k := \begin{cases} a_k & (k \le h) \\ -1 & (k > h) \end{cases}.$$

If  $k \leq h$ , then we define  $v_{\leq k} := (v_1, \dots, v_k)$ .

**Remark 2.2.** We often identify [b] as  $[b]^1$  and abuse the notation. For example, we write v \* i for v \* (i).

**Definition 2.3.** A rooted tree of height  $\leq h$  and  $\leq b$ -branching is a subset  $T \subseteq [b]^{\leq h}$  such that:

1.  $T \neq \emptyset$ .

 $2. \ v \subseteq w \in T \Longrightarrow v \in T.$ 

 $\emptyset \in T$  is called **the root of** T.

For  $v, w \in T$ , v < w means the lexicographic order:

$$v < w : \iff \exists k \le h. \ (v_{\le k} = w_{\le k} \& v_{k+1} < w_{k+1})$$

If  $v * i \in T$ , then we say v \* i is a child of v, and v is the parent of v \* i. If  $v * i, v * j \in T$ , then they are said to be siblings.

When  $v, w \in T$  satisfy height(v) = height(w), then we say v is left to w when v < w.

Put

$$height(T) := \max_{v \in T} height(v).$$

**Example 2.4.** If  $v * 1 \in T$ , then v \* 1 must be the leftmost child of v in T.

In this article, we use a one-sided formulation of propositional sequent-calculus, which can be regarded as the non-uniform version of a first-order sequent calculus implementing  $T_2^2(R)$ :

**Definition 2.5.** A cedent is a finite set of propositional formulae. Given a cedent  $\Gamma$ , its **semantic interpretation** is the propositional formula  $\bigvee_{\varphi \in \Gamma} \varphi$ . Here, if  $\Gamma = \emptyset$ , we set  $\bigvee_{\varphi \in \Gamma} \varphi := 0$ . Under a truth assignment,  $\Gamma$  is said to be **true** if and only if its semantic interpretation is true.

We often denote cedents of the form

$$\Gamma_1 \cup \ldots \cup \Gamma_k \cup \{\varphi_1, \ldots, \varphi_m\}$$

by

$$\Gamma_1,\ldots,\Gamma_k,\varphi_1,\ldots,\varphi_m.$$

Given constants c, d, a cedent S and a finite vertex-labeled tree  $\pi = (\mathcal{T}, \mathcal{S})$ ,  $\pi$  is  $\boldsymbol{a}$   $LK_{d+\frac{1}{2},c}^*$ -derivation of S (without redundancy) if and only if the following hold:

- 1.  $height(\mathcal{T}) \leq c$ .
- 2. For each  $v \in \mathcal{T}$ ,  $\mathcal{S}(v)$  is a cedent of cardinality  $\leq c$ .
- 3.  $S(\emptyset) = S$ .
- 4. For each  $v \in \mathcal{T}$ ,  $\mathcal{S}(v)$  is derived from the labels of its children, that is,  $(\mathcal{S}(v*i))_{v*i\in\mathcal{T}}$  by applying one of the following derivation rules:
  - Initial cedent:

$$\Gamma, x, \overline{x}$$
 (where x is a constant or a variable)

• V-Rule:

$$\frac{\Gamma, \varphi_{i_0}}{\Gamma} \quad \text{(where } \bigvee_{i=1}^{I} \varphi_i \in \Gamma, \ 1 \leq i_0 \leq I, \ \varphi_{i_0} \not\in \Gamma)$$

• V-Rule:

$$\frac{\Gamma, \varphi_{i_0}}{\Gamma} \quad \text{(where } \varphi_1 \vee \varphi_2 \in \Gamma, i_0 = 1 \text{ or } i_0 = 2, \, \varphi_{i_0} \not\in \Gamma)$$

•  $\Lambda$ -Rule:

$$\frac{\Gamma, \varphi_1 \qquad \Gamma, \varphi_2 \qquad \cdots \qquad \Gamma, \varphi_I}{\Gamma}$$

where  $\bigwedge_{i=1}^{I} \varphi_i \in \Gamma$ , and  $\varphi_i \not\in \Gamma$  for each  $i \in [I]$ .

• ∧-Rule:

$$\frac{\Gamma, \varphi_1 \qquad \Gamma, \varphi_2}{\Gamma}$$

where  $\varphi_1 \wedge \varphi_2 \in \Gamma$ , and  $\varphi_i \notin \Gamma$  for each i = 1, 2.

• Trivial Cut:

$$\frac{\Gamma, 0}{\Gamma}$$
 (where  $0 \notin \Gamma$ )

•  $p\Sigma_{d+\frac{1}{2}}$ -Induction:

$$\begin{array}{cccc}
\Gamma, \varphi_1 & \Gamma, \overline{\varphi_1}, \varphi_2 & \cdots & \Gamma, \overline{\varphi_{I-1}}, \varphi_I & \Gamma, \overline{\varphi_I} \\
\Gamma & & \Gamma
\end{array}$$

where each  $\varphi_i$  is  $p\Sigma_{d+\frac{1}{2}}(|\pi|)$ , and  $\varphi_i, \overline{\varphi}_i \notin \Gamma$  for  $i \in [I]$ . Here, we define the size  $|\pi|$  of the proof  $\pi$  as

$$|\pi| := \sum_{v \in \mathcal{T}} \sum_{\varphi \in \mathcal{S}(v)} |\varphi|.$$

Note that, when  $I=1, p\Sigma_{d+\frac{1}{2}}$ -Induction is a usual cut-rule for  $p\Sigma_{d+\frac{1}{2}}(|\pi|)$ -formulae.

**Remark 2.6.** In this article, we are particularly interested in d = 1, 2.

**Definition 2.7.** onto  $PHP_n^{n+1}$  is the following cedent<sup>1</sup>:

$$\left\{\bigvee_{p\in[n+1]}\bigwedge_{h\in[n]}\neg r_{ph},\bigvee_{\substack{p\neq p'\in[n+1],\\h\in[n]}}(r_{ph}\wedge r_{p'h}),\bigvee_{\substack{h\in[n]}}\bigwedge_{\substack{p\in[n+1]\\p\in[n+1]}}\neg r_{ph},\bigvee_{\substack{h\neq h'\in[n],\\p\in[n+1]}}(r_{ph}\wedge r_{ph'})\right\}$$

We can also consider a first-order counterpart of the above calculus, and we can regard the above calculus as a propositional translation of it. It is just a reprinting of classical Paris-Wilkie translation [21]. See [15] for a comprehensive presentation of the notion and [11] for a presentation of the closest conventions to ours. See also Appendix 7.1 for our presentation. Using these notions, we have:

**Proposition 2.8** (essentially, Corollary 9.1.4 and the proof of Lemma 9.5.1 in [15]). Suppose  $T_2^d(R) \vdash ontoPHP_n^{n+1}(R)$ , where  $ontoPHP_n^{n+1}(R)$  is a natural formalization of the pigeonhole principle for bijections using (n+1) pigeons and n holes. Then  $ontoPHP_n^{n+1}$  has  $2^{|n|^{O(1)}}$ -sized  $LK_{d+\frac{1}{2},O(1)}^*$ -proofs.

Remark 2.9. Note that our convention  $LK_{d+\frac{1}{2},O(1)}^*$  adopts  $p\Sigma_d$ -Induction rule, which has unbounded arity and makes the translation of  $\Sigma_d^b(R)$ -Induction more straightforward than the argument in the proof of Theorem 9.1.3 in [15]. The resulting proof tree is of constant height, and each cedent is of constant cardinality.

Remark 2.10.  $2^{|n|^C}$ -sized  $LK^*_{d+\frac{1}{2},O(1)}$ -proofs of  $ontoPHP^{n+1}_n$  consist only of  $s\Sigma_{d+\frac{1}{2}}(2^{|n|^C})$ -formulae,  $s\Pi_{d+\frac{1}{2}}(2^{|n|^C})$ -formulae, and subformulae of the formulae in  $ontoPHP^{n+1}_n$ .

# 3 A review of unprovability of the pigeonhole principle over $T_2^1(R)$

In this section, we briefly review the independence result, which can be regarded as the base case of Ajtai's Theorem, in order to locate this article in the course of previous research:

**Theorem 3.1** (essentially by [21]. See for example [6] or [20] for clear expositions).

$$T_2^1(R) \not\vdash ontoPHP_n^{n+1}(R).$$

<sup>&</sup>lt;sup>1</sup>We may have expanded  $\bigvee$  to lower the complexity of the formulae in the cedent, but we are interested in cedents of constant cardinality with respect to the parameter n (recall Definition 2.5 and see Theorem 4.13), so we formulated the cedent  $ontoPHP_n^{n+1}$  as above.

It can be proven in multiple ways: by Paris-Wilkie forcing ([21]. See [6] for a neat presentation), by Riis criterion ([25]), and by converting a  $T_2^1(R)$ -proof of  $ontoPHP_n^{n+1}(R)$  to narrow resolution proofs ([12]) and applying resolution width lower bounds for  $ontoPHP_n^{n+1}$  obtained by pebble-game analysis ([5]).

Our trial can be regarded as a generalization of the last proof strategy to  $T_2^2(R)$ . We sketch our version of the proof of Theorem 3.1 in the rest of this section for completeness and as a showcase of our trial.

Definition 3.2. Let  $\mathcal{M}_n$  be the set of all partial matchings between [n+1] (pigeons) and [n] (holes).

For  $M, M' \in \mathcal{M}_n$ , when  $M \cup M' \notin \mathcal{M}_n$ , in other words, M and M' match a pigeon to different holes or a hole to different pigeons, we say M contradicts M', denoted by  $M \perp M'$ .

We define the following variation of a pebble game.

### **Definition 3.3.** Given $n, C \in \mathbb{N}$ , $\mathcal{G}_1(n, C)$ is the following game:

- 1. Played by two players. We call them **Prover** and **Delayer**. For readability, we use the pronoun "he" for **Prover** and "she" for **Delayer**.
- 2. Possible positions are sequences on  $(\mathcal{M}_n)_{\leq |n|^C}$  with length at most  $2^{|n|^C}$ .
- 3. The initial position is the sequence  $(\emptyset)$  of length 1.
- 4. Now, we describe transitions between positions together with each player's options. suppose the current position is  $(M_0, \ldots, M_l) \in ((\mathcal{M}_n)_{\leq |n|^C})^{l+1}$ . If  $l+1 \geq 2^{|n|^C}$ , then the play ends and **Delayer** wins. If  $l+1 < 2^{|n|^C}$ , then proceed as follows:
  - (a) First, **Prover** plays a subset  $Q \subseteq P_n \dot{\cup} H_n$  of cardinality  $\#Q \leq |n|^C$ , and send it to **Delayer**.
  - (b) **Delayer** plays a matching  $M' \in (\mathcal{M}_n)_{\leq |n|^C}$  such that M' covers Q and is consistent with  $\rho_l$ . If such M' does not exist, the play ends and **Prover** wins. Otherwise, the sequence  $(M_0, \ldots, M_l, M')$  is the next position.

The game is determined since the length of a position increases by 1 as long as the play continues and the game ends when the length reaches  $2^{|n|^C}$ .

**Remark 3.4.** An informal description of  $\mathcal{G}_1(n,C)$  is as follows: **Prover** has  $2|n|^C$  pebbles, and there is a game board having (n+1) pigeons and n holes. He puts some pebbles on some of the pigeons and the holes in a course of the game. Initially, there is no pebbles on the board. In each turn, **Prover** selects at most  $|n|^C$  pigeons and holes and puts one pebble per each. Then **Delayer** answers which holes (resp. pigeons) are matched to the

pebbled pigeons (resp. holes). She must pretend that she knows a bijection between the pigeons and holes, that is, the answers to the pebbles on the board should always be a partial matching between pigeons and holes. If she cannot answer in this way, she loses. If there are more than  $|n|^C$  pebbles on the board, **Prover** must remove some so that the number eventually becomes no more than  $|n|^C$ . Note that we only refer the answers to the pebbles on the current board, and the answers to the removed pebbles become irrelevant to the winning condition anymore.

Moreover, the following is an easy observation:

**Lemma 3.5.** If  $2|n|^C \le n$ , then **Delayer** has a winning strategy for  $\mathcal{G}_1(n, C)$ . Here, **a** winning strategy of **Delayer** for  $\mathcal{G}_1(n, C)$  is a mapping

$$f: (\mathcal{M}_n)_{\leq |n|^C} \times (2^{P_n \dot{\cup} H_n})_{\leq |n|^C} \to (\mathcal{M}_n)_{\leq |n|^C}$$

such that **Delayer** always wins  $\mathcal{G}_1(n, C)$  if she plays f(P, Q) when the current position is P and **Prover** queries Q, regardless **Prover**'s moves.

*Proof.* In each turn, given a position  $P = (M_0, ..., M_l)$  and a query Q, there exists a matching  $\tilde{M}$  such that  $M_l \subseteq \tilde{M}$ ,  $\tilde{M}$  covers Q, and  $\#\tilde{M} \leq 2|n|^C$  since  $2|n|^C \leq n$ . Therefore, it suffices for **Delayer** to answer minimal  $M' \subseteq \tilde{M}$  which covers Q. In this way, **Delayer** can survive arbitrary many turns and win the game.

Corollary 3.6. Given  $C \in \mathbb{N}$ , Delayer has a winning strategy for  $\mathcal{G}_1(n, C)$  for sufficiently large n.

On the other hand, we have the following:

**Theorem 3.7** (essentially by [12]). Suppose  $ontoPHP_n^{n+1}$  has  $LK_{1+\frac{1}{2},O(1)}^*$ -proofs of size  $2^{|n|^{O(1)}}$ . Then there exists a constant C>0 such that, for sufficiently large  $n\in\mathbb{N}$ , **Prover** has a winning strategy for  $\mathcal{G}_1(n,C)$ . Here, **a winning strategy of Prover**  $for \mathcal{G}_1(n,C)$  is a mapping  $f: (\mathcal{M}_n)_{\leq |n|^C} \to (2^{P_n \cup H_n})_{\leq |n|^C}$  such that **Prover** always wins  $\mathcal{G}_1(n,C)$  if he queries f(P) when the current position is P, regardless **Delayer**'s moves.

The proof can be regarded as a "restriction" of the proof of Theorem 4.13, and we only give a sketch here.

**Definition 3.8.** Let  $M \in \mathcal{M}_n$ . Let  $\varphi$  be a  $s\Sigma_0$ -formula. We write  $M \Vdash \varphi$  to denote that a partial assignment  $\rho^M$  induced by M covers the all variables occurring in  $\varphi$  and  $\rho^M \models \varphi$ . Here,  $\rho^M$  is the following assignment:

$$\rho^{M}; r_{ph} \mapsto \begin{cases} 1 & \text{(if } M \text{ matches } p \text{ and } h) \\ 0 & \text{(if } M \perp \{p \mapsto h\}) \\ r_{ph} & \text{(otherwise)} \end{cases}$$

**Definition 3.9.** Let  $\Gamma$  be a cedent consisting only of  $s\Sigma_{2+\frac{1}{2}}$ - or  $s\Pi_{2+\frac{1}{2}}(z)$ -formulae. Let  $\Pi(\Gamma)$  be the set of all  $p\Pi_1(z)$ - or  $p\Pi_2(z)$ -formulae in  $\Gamma$ . Let W be a function defined on  $\Pi(\Gamma)$  and  $M \in \mathcal{M}_n$ . We say (M, W) **falsifies**  $\Gamma$  when the following hold:

- For each  $\varphi \in \Gamma$  which are  $p\Sigma_{0+\frac{1}{2}}(z)$ ,  $M \Vdash \overline{\varphi}$ .
- For each  $\bigwedge_{i=1}^{I} \varphi_i \in \Pi(\Gamma)$ ,  $w := W(\bigwedge_{i=1}^{I} \varphi_i) \in [I]$ . Moreover, if  $\bigwedge_{i=1}^{I} \varphi_i$  is  $p\Pi_{1+\frac{1}{2}}(z)$  (and therefore  $\varphi_i$  is  $p\Sigma_0$ ), then  $M \Vdash \overline{\varphi_w}$ .

We call W a counterexample function for  $\Gamma$ .

Sketch of the proof of Theorem 3.7. By assumption, there exists C > 0 such that, for any sufficiently large n, there exists an  $LK_{1+\frac{1}{2},C}^*$ -derivation  $\pi = (\mathcal{T},\mathcal{S})$  of  $ontoPHP_n^{n+1}$  satisfying  $|\pi| \leq 2^{|n|^C}$ . We extract a winning strategy of **Prover** from  $\pi$ . The main idea is the same as the PLS witnessing for **Prover** given in [11]; in the course of a play, **Prover** always looks at a vertex v in  $\mathcal{T}$  and tries to falsify  $\mathcal{S}(v)$  by  $(M_l, W[P])$ , where we suppose  $P = (M_0, \ldots, M_l)$  is the current position, and W[P] is the counterexample function for  $\mathcal{S}(v)$  obtained in the play so far.

Below, the precise estimation of the number of pebbles we need and some other details to carry out the argument are left as exercise, but we make them explicit. We write " $\Sigma_d$ " (resp. " $\Pi_d$ ") instead of " $\Sigma_{d+\frac{1}{2}}(|\pi|)$ " (resp. " $\Pi_{d+\frac{1}{2}}(|\pi|)$ ") for readability. Initially, **Prover** looks at the root of  $\mathcal{T}$ . The sequent there is  $ontoPHP_n^{n+1}$ , and therefore all the  $s\Pi_1$ -formulae are falsified by  $M_0 = \emptyset$  trivially since there is none. At a position  $P = (M_0, \ldots, M_l)$ , if **Prover** looks at a vertex  $v \in \mathcal{T}$ , then **Prover** queries and transitions depending on the inference rule used to derive  $\mathcal{S}(v)$ :

• Note that S(v) cannot be an Initial cedent:

$$\frac{1}{\Gamma, x, \overline{x}}$$
 (where x is a constant or a variable)

since x and  $\overline{x}$  cannot be falsified simultaneously.

• When  $S(v) = \Gamma$  is derived by  $\vee$ -Rule:

$$\frac{\Gamma, \varphi_{i_0}}{\Gamma} \quad \text{(where } \varphi_1 \vee \varphi_2 \in \Gamma, i_0 = 1 \text{ or } i_0 = 2, \, \varphi_{i_0} \not\in \Gamma)$$

Since  $\varphi_1 \vee \varphi_2$  is  $s\Sigma_2$  or  $p\Pi_1$ , it must be  $s\Sigma_0$ . Therefore, it is already falsified by  $M_l$ . Thus it suffices for **Prover** to query  $\emptyset$  and climb up to the child of v, setting  $M_{l+1} := M_l$  and  $W[P * M_{l+1}] := W[P]$ .

• When  $S(v) = \Gamma$  is derived by Trivial Cut or  $\land$ -rule: similar to the above.

• When  $S(v) = \Gamma$  is derived by  $\bigvee$ -Rule:

$$\frac{\Gamma, \varphi_{i_0}}{\Gamma} \quad \text{(where } \bigvee_{i=1}^{I} \varphi_i \in \Gamma, \ 1 \le i_0 \le I, \ \varphi_{i_0} \not\in \Gamma)$$

If  $\bigvee_{i=1}^{I} \varphi_i$  is one of the formulae in  $ontoPHP_n^{n+1}$ , which is the main difference from the situation of PLS witnessing in [11],  $\varphi_{i_0}$  corresponds to a pigeon or a hole, and **Prover** queries it. The answer must falsify  $\varphi_{i_0}$ , and **Prover** climbs up to the child of v, storing the answer as a counterexample if necessary; we treat the cases for the first two formulae in  $ontoPHP_n^{n+1}$  here:

- If  $\bigvee_{i=1}^{I} \varphi_i = \bigvee_{p \in [n+1]} \bigwedge_{h \in [n]} \neg r_{ph}$ , then **Prover** queries  $i_0 \in [n+1]$ . Receiving **Delayer**'s answer including  $i_0 \mapsto h_0$ , he climbs up to the child of v, setting  $M_{l+1} := M_l \cup \{i_0 \mapsto h_0\}$  and  $W[P * M_{l+1}] := W[P] \dot{\cup} \{\varphi_{i_0} \mapsto h_0\}$ .
- If  $\bigvee_{i=1}^{I} \varphi_i = \bigvee_{\substack{p \neq p' \in [n+1], \\ h \in [n]}} (r_{ph} \wedge r_{p'h})$ ,  $i_0$  is in the form of  $\langle p, p', h \rangle$ . **Prover** queries  $\{p, p'\} \subset [n+1]$ . Receiving **Delayer**'s answer including  $p \mapsto h_0$  and  $p' \mapsto h_1$ , he climbs up to the child of v, setting  $M_{l+1} := M_l \cup \{p \mapsto h_0, p' \mapsto h_1\}$  and  $W[P * M_{l+1}] := W[P]$ .

Otherwise,  $\bigvee_{i=1}^{I} \varphi_i$  is  $p\Sigma_1$ , then  $\varphi_{i_0}$  is  $p\Sigma_0$ , and therefore there exists at most  $|n|^C$ many pigeons and holes, say, a set Q, such that any  $M \in \mathcal{M}_n$  covering Q satisfies  $M \Vdash \varphi_{i_0}$  or  $M \Vdash \overline{\varphi_{i_0}}$ . **Prover** queries Q, and let M be the minimal matching covering Q contained in **Delayer**'s answer. If  $M \Vdash \overline{\varphi_{i_0}}$ , then he climbs up to the child of v, setting  $M_{l+1} := M_l \cup M$  and  $W[P * M_l] := W[P]$ . If  $M \Vdash \varphi_{i_0}$ , then he goes back to the ancestor v' of v where  $\bigvee_{i=1}^{I} \varphi_i$  is eliminated by  $p\Sigma_1$ -Induction, and proceeds to the immediate right siblings, carrying the counterexample  $i_0$  for  $\bigwedge_{i=0}^{I} \overline{\varphi_i}$ ; we consider the position  $P' = (M_0, \dots, M_{l'})$  (l' < l) when **Prover** looked at v'. **Prover** sets  $M_{l+1} := M_{l'} \cup M$  and  $W[P * M_{l+1}] := W[P'] \dot{\cup} \{\bigwedge_{i=0}^{I} \overline{\varphi_i} \mapsto i_0\}$  As for the existence of such ancestor, see the proof of Theorem 4.13.

• When  $S(v) = \Gamma$  is derived by  $\Lambda$ -Rule:

$$\begin{array}{cccc} \Gamma, \varphi_1 & \Gamma, \varphi_2 & \cdots & \Gamma, \varphi_I \\ \hline \Gamma & & \end{array}$$

where  $\bigwedge_{i=1}^{I} \varphi_i \in \Gamma$ , and  $\varphi_i \notin \Gamma$  for each  $i \in [I]$ . We have assumed that the  $s\Pi_1$ formulae in S(v) are already falsified, so **Prover** queries  $Q = \emptyset$  and climbs up to
the child  $v * W(\bigwedge_{i=1}^{I} \varphi_i)$  corresponding to the counterexample, setting  $M_{l+1} := M_l$ and  $W[P * M_{l+1}] := W[P]$ .

• When  $S(v) = \Gamma$  is derived by  $p\Sigma_1$ -Induction:

$$\frac{\Gamma, \varphi_1 \qquad \Gamma, \overline{\varphi_1}, \varphi_2 \qquad \cdots \qquad \Gamma, \overline{\varphi_{I-1}}, \varphi_I \qquad \Gamma, \overline{\varphi_I}}{\Gamma}$$

where each  $\varphi_i$  is  $p\Sigma_1$ , and  $\varphi_i, \overline{\varphi}_i \notin \Gamma$  for  $i \in [I]$ .

Querying  $Q = \emptyset$ , proceed to the leftmost child with  $\Gamma \cup \{\varphi_1\}$ , which does not have a new  $p\Pi_1$  formula, setting  $M_{l+1} := M_l$  and  $W[P * M_{l+1}] := W[P]$ .

We can observe that the vertex v transitions monotonously with respect to the lexicographic ordering on  $\mathcal{T}$ , and therefore the play ends with less than  $2^{|n|^C}$  turns, and it implies **Prover** wins the game.

# 4 A game notion for $T_2^2(R)$

This section aims to give a candidate of an appropriate game notion for  $2^{|n|^{O(1)}}$ -sized  $LK_{2+\frac{1}{2},O(1)}^*$ -proofs.

Towards precise descriptions of them, we define the following notions:

**Definition 4.1.** A tree T is an(n, C)-tree if and only if it satisfies the following:

- 1. T is a tree of height  $\leq C$  and  $\leq 2^{|n|^C}$ -branching,
- 2. If  $v * k \in T$  and  $l \in [k]$ , then  $v * l \in T$ .

We define the game  $\mathcal{G}_2$  as an extended pebble game where **Prover** has another option than putting pebbles and asking queries; he can backtrack the game record, bringing back a certain amount of partial matching he currently has. We manage the record of the game by trees rather than sequences as in Definition 3.3 so that the notion of *obliviousness* in Definition 4.10 and the proof of Theorem 4.13 become easy to describe. The precise definition of  $\mathcal{G}_2$  is as follows (for an informal description of the game, see Remark 4.3):

**Definition 4.2.** Let  $n, C \in \mathbb{N}$ . Suppose we are given an (n, C)-tree T.  $\mathcal{G}_2(n, C, T)$  is the following game:

- 1. Played by two players. We call them in the same way as in Definition 3.3.
- 2. A possible position is a map L such that:
  - (a)  $dom(L) \subseteq T$ .
  - (b) For  $v \in \text{dom}(L)$ , L(v) is of the form  $(M, A_1, \ldots, A_{height(v)})$ , where M is a partial matching between (n+1)-pigeons and n-holes with size  $\leq |n|^C \times height(v)$ , and each  $A_j \in 2^{|n|^C}$ .

- (c) dom(L) is closed downwards under  $\subseteq$ .
- (d)  $v \subseteq w \in \text{dom}(L)$ , if  $(M, \vec{A}) = L(v)$  and  $(M', \vec{A}') = L(w)$ , then they satisfy  $M \subseteq M'$  and  $\vec{A} \subseteq \vec{A}'$ .

For future convenience, we introduce the following notations:

- Set  $c(L) := \max \operatorname{dom}(L)$ .
- Let  $\mathcal{P}$  be the set of all possible positions.
- 3. **The initial position** is  $L_0$ , where dom $(L_0)$  is the rooted tree of height 0, that is, consists only of the root  $\emptyset$ , and  $L_0(\emptyset) = (\emptyset)$ .
- 4. Now, we describe transitions between positions together with each player's options: suppose the current position is L, and  $L(c(L)) = (M, \vec{A})$ .
  - (a) First, **Prover** plays a subset  $Q \subseteq P_n \dot{\cup} H_n$  of cardinality  $\#Q \leq |n|^C$ , and send it to **Delayer**.
  - (b) **Delayer** plays a minimal matching  $M' \in \mathcal{M}_n$  covering Q, and send it back to **Prover**.
  - (c) **Prover** plays a triple  $\langle o, x, B \rangle$ , where  $o \in [3]$ ,  $x \in [2^{|n|^C}]$  if o = 1,  $x \subsetneq c(L)$  if o = 2, 3, and  $B \in [2^{|n|^C}]$ .
  - (d) Depending on o, the next position (or judgment of the winner) is determined as follows:
    - i. Consider the case when o=1. If  $c(L)*x \not\in T$ , then the game ends, and **Prover** loses. Otherwise, extend L to a map on  $\text{dom}(L) \cup \{c(L)*x\}$  by labeling the child c(L)\*x with  $(M \cup M', \vec{A}, B)$ , and let L' be the resulting partial labeling on T'. Note that  $c(L)*x \not\in \text{dom}(L)$  by definition of c(L). If  $M \perp M'$ , then the game ends, and **Prover** wins. Otherwise,  $L' \in \mathcal{P}$ , and it is the next position.
    - ii. If o = 2, proceed as follows. Let  $c(L) = x * k * \sigma$   $(k \in [2^{|n|^C}], \sigma \in [2^{|n|^C}]^{\leq C}$ . Note that  $\sigma$  may be empty). If  $x * (k+1) \notin T$ , the play ends and **Prover** loses.

Consider the case when  $x*(k+1) \in T$ . Note that  $x*(k+1) \notin \text{dom}(L)$  by  $x*k \subseteq \text{dom}(L)$ . Let  $(M'', \vec{\alpha}) = L(x)$ . Extend L by labeling x\*(k+1) with  $(M'' \cup M', \vec{\alpha}, B)$ , and let L' be the resulting map.

If  $M'' \perp M'$  as matchings, then the game ends, and **Prover** wins. Otherwise,  $L' \in \mathcal{P}$ , and it is the next position.

iii. Consider the case when o = 3. Let  $c(L) = x * k * \sigma$ . If  $x * (k - 1) \not\in \text{dom}(L)$ , then the game ends, and **Prover** loses.

Otherwise, let l be the maximal extension of x\*(k-1) within dom(L). If l is a leaf of T, the game ends, and **Prover** loses. Otherwise, let  $(M'', \vec{\alpha}) := L(l)$ . Let

$$T' := (\operatorname{dom}(L) \setminus \{v \in \operatorname{dom}(L) \mid x * k \subseteq v\}) \cup \{l * 1\}$$

Note that  $l*1 \in T$  by assumption on T, and hence  $T' \subseteq T$ . Furthermore,  $l*1 \not\in \text{dom}(L)$  by definition of l. Now, set

$$L' := L \upharpoonright_{(\operatorname{dom}(L) \cap T')} \cup \{l * 1 \mapsto (M'' \cup M', \vec{\alpha}, B)\}$$

If  $M'' \perp M'$  as matchings, the game ends, and **Prover** wins. Otherwise,  $L' \in \mathcal{P}$ , and it is the next position.

### **Remark 4.3.** The intended meaning of each item above is as follows:

- 1. **Delayer** pretends to have a truth assignment falsifying  $ontoPHP_n^{n+1}$ , which is, of course, impossible, and **Prover** wants to disprove it by querying how certain pigeons and holes are mapped to each other.
- 2. Intuitively, L is **Prover**'s partial record of a play of a pebble game. T is a proof-tree  $\mathcal{T}$  of a  $2^{|n|^C}$ -sized  $LK_{2+\frac{1}{2},C}^*$ -derivation  $(\mathcal{T},\mathcal{S})$  of  $ontoPHP_n^{n+1}$ . For  $v \in dom(L) \subseteq T$ ,  $L(v) = (M, \vec{A})$  means M is the state of pebbles played at v, and  $\vec{A}$  is auxiliary information for **Prover** to decide the next move, which may carry at most  $C|n|^C$  bits of information and is insufficient to code the whole play of the game  $\mathcal{G}_2$ .
  - c(L) stands for **the current frontier** of the position L.
- 3. The play starts at the root with no pebbles on the board.
- 4. The play proceeds in turn, starting from **Prover**.
  - (a) **Prover** queries at most  $|n|^C$ -many pigeons and holes Q, and send it to **Delayer**.
  - (b) **Delayer** answers which holes (resp. pigeons) are matched to the pigeons (resp. the holes) queried.
  - (c) Based on **Delayer**'s answer, **Prover** chooses from one of the three options indicated by  $o \in [3]$ . Furthermore, he specifies auxiliary information x and B (described in the following) to determine the next position precisely.
  - (d) i. If o = 1, climb up to the child c(L) \*x of c(L) and add M' to the matching. Prover also remembers B, which can code  $|n|^{O(1)}$  bits of information.

- ii. If o = 2, move backward to  $x \subsetneq c(L)$ , climb up to the immediate right child of x, adding M' to the matching, forgetting the differences of matching and  $\vec{A}$  between x and c(L), but remembering B. Note that we keep a record of the labels of the vertices from x up to c(L).
- iii. If o = 3, move backward to  $x \subsetneq c(L)$ , climb up to the immediate left path from x, adding M' to the matching and remembering B. This amounts to backtracking the game record, bringing back the information M' obtained at the current position. Note that we erase the record of the labels of the vertices between x and c(L), including c(L)'s but not x's.

In any case, if **Prover** cannot play within the tree T, **Prover** loses. If **Delayer**'s answer M' contradicts the matching M labeled at the parent of the subcedent c(L), **Prover** wins.

First, we observe that the game above is determined. Towards it, we introduce the following binary relation:

**Definition 4.4.** Let T, T' be trees. We write  $T \prec T'$  if and only if there exists  $w \in T'$  such that

$$T_{\le w} = T'_{\le w} \& w \notin T.$$

Here, for a vertex v of a tree S,  $S_{\leq v}$  denotes  $\{s \in S \mid s < v\}$ . Note that  $T \prec T'$  implies  $T \neq T'$ . Set  $T \preceq T' :\Leftrightarrow T = T'$  or  $T \prec T'$ .

**Lemma 4.5.**  $\leq$  is a linear order on the set of finite trees.

*Proof.* First,  $\leq$  is antisymmetric. Towards a contradiction, suppose  $T \prec U$  and  $U \prec T$ . Let  $u \in U$  witness  $T \prec U$  and  $t \in T$  witness  $U \prec T$ . If  $u \leq t$ , then it contradicts the two assumptions  $U_{< t} = T_{< t}$  and  $u \notin T$ . Similar for  $t \leq u$ , and it is absurd.

Besides,  $\prec$  is transitive. Suppose  $S \prec T$  and  $T \prec U$ . Let  $t \in T$  witness  $S \prec T$  and  $u \in U$  witness  $T \prec U$ . If  $t \leq u$ , then  $S_{< t} = T_{< t} = U_{< t}$ ,  $t \not\in S$ , and  $t \in U$  because  $T_{< u} = U_{< u}$  and  $u \in U$ . If t > u,  $S_{< u} = T_{< u} = U_{< u}$ , and  $u \not\in T$  together with  $S_{< t} = T_{< t}$  implies  $u \not\in S$ .

Lastly,  $\prec$  satisfies a trichotomy: if  $T \neq U$ , then  $T \prec U$  or  $U \prec T$ . Indeed, by finiteness of T and U, take the maximum  $w \in T \cup U$  such that  $T_{< w} = U_{< w}$ . Note that  $T_{< \emptyset} = U_{< \emptyset}$  trivially. Towards a contradiction, suppose  $w \in T \cap U$ . Then, since  $T \neq U$  and  $T_{< w} = U_{< w}$ , there exists  $v \in T \cup U$  such that w < v. Take the least such v. Then

$$T_{< v} = T_{< w} \cup \{w\} = U_{< w} \cup \{w\} = U_{< v},$$

contradicting the maximality of w. Therefore, w witnesses  $T \prec U$  or  $U \prec T$ .

**Lemma 4.6.** If a position L transitions to L' in a play of  $\mathcal{G}_2(n, C, T)$ , then  $dom(L) \prec dom(L')$ .

*Proof.* Suppose  $\langle o, x, B \rangle$  is **Prover**'s option at L. We split cases:

- 1. if o = 1, then c(L) \* x witnesses  $dom(L) \prec dom(L')$ .
- 2. if o = 2, then x \* (k + 1) witnesses  $dom(L) \prec dom(L')$ , where  $x * k \subseteq c(L)$ .
- 3. if o = 3, then c(L') witnesses  $dom(L) \prec dom(L')$ . Note that c(L') = l \* 1, where l is the maximum leaf of dom(L) extending x \* (k-1). Here,  $x * k \subseteq c(L)$  again.

Corollary 4.7. For any  $n, C \in \mathbb{N}$  and an (n, C)-tree T,  $\mathcal{G}_2(n, C, T)$  ends within  $2^{2^{(C+1)|n|^C}}$ -steps, determining the winner.

*Proof.* When the game ends, it always determines who is the winner. Therefore, it suffices to show that the game ends with at most  $2^{2^{(C+1)|n|^C}}$ -many transitions. Since there are at most  $2^{(2^{|n|^C})^{C+1}}$ -many rooted trees of height  $\leq C$  and  $\leq 2^{|n|^C}$ -branching as subsets of T, the result follows from Lemma 4.6.

Note that, without any restriction, **Prover** easily wins  $\mathcal{G}_2(n, C, T)$  for reasonable parameters:

**Proposition 4.8.** Suppose  $n, C \ge 2$ ,  $2^{|n|^C} > n$ . Let T be the (n, C)-tree as follows:

$$T:=[n+1]^{\leq 1}\cup\{i*1\mid i\in[n+1]\}.$$

Then there exists a winning strategy of Prover for  $\mathcal{G}_2(n, C, T)$ , that is, there is a pair  $(f_1, f_2)$  of functions such that:

- 1.  $f_1: \mathcal{P} \to (2^{P_n \dot{\cup} H_n})_{\leq |n|^C}$ .
- 2.  $f_2: \mathcal{P} \times \mathcal{M}_n \to [3] \times [2^{|n|^C}]^{\leq C} \times [2^{|n|^C}]$ , and  $\langle o, x, B \rangle = f_2(L, M')$  satisfies  $x \in [2^{|n|^C}]$  if o = 1 and  $x \subseteq c(L)$  otherwise.
- 3. If **Prover** keeps playing  $Q = f_1(L)$  at stage 4a in Definition 4.2 and, receiving **Delayer**'s answer M' at stage 4b, plays  $\langle o, x, B \rangle = f_2(L, M')$  at stage 4c, then **Prover** eventually wins  $\mathcal{G}_2(n, C, T)$  regardless **Delayer**'s moves.

*Proof.* Let  $P_n = \{p_1, \dots, p_{n+1}\}$ . Informally, **Prover**'s winning strategy is as follows:

- Taking option o = 1, ramify at the root asking the query  $\{p_1\}$ . The root will have (n+1)-many children.
- One by one, taking option o = 2, ask queries  $\{p_2\}, \ldots, \{p_{n+1}\}$ . There are enough vertices to do it.

• Because of the pigeonhole principle, there will be a < a' such that **Delayer**'s answers  $p_a \mapsto h$  and  $p_{a'} \mapsto h'$  share the same hole: h = h'. Once detecting the pair, repeatedly take option o = 3 and bring back the information  $p_{a'} \mapsto h$  to the vertex storing  $p_a \mapsto h$ .

Now, we give a formal presentation.

For the initial position, set

$$f_1(L_0) := \{p_1\}, \ f_2(L_0, M) := \langle 1, 1, * \rangle,$$

where \* can be any.

Let L be a position with height(c(L)) = 1. If L(c(L)) does not contradict L(l) for any leaf l of dom(L), set

$$f_1(L) := \{p_{a+1}\}$$

where  $a := c(L) \in [n+1]$ .

Let M' be a partial matching of the form  $\{p_{a+1} \mapsto h\}$ . Set

$$f_2(L, M') := \langle 2, \emptyset, * \rangle.$$

Let L be a position with  $height(c(L)) \ge 1$ . If there exists a leaf l of dom(L) such that  $L(l) \perp L(c(L))$ , then let  $a \in [n+1]$  be the maximum index such that  $\{p_a \mapsto h\} \subseteq L(c(L))$  contradicts L(l). Set

$$f_1(L) := \{p_a\}$$

and

$$f_2(L, \{p_a \mapsto h\}) := \langle 3, \emptyset, * \rangle.$$

Regardless values  $f_1(L)$  and  $f_2(L, M')$  for other L and M',  $(f_1, f_2)$  is a winning strategy for **Prover** because of the assumption on the parameters.

**Remark 4.9.** We can see that the part  $\vec{A}$  of the label of positions  $L \in \mathcal{P}$  does not play any role in the proof above. Actually,  $\vec{A}$  is not essential at all for our use of  $\mathcal{G}_2$ . See Appendix 7.2. However, the auxiliary information  $\vec{A}$  is helpful to describe the obliviousness in the Definition 4.10 and in the proof of Theorem 4.13.

Thus, to obtain a nontrivial and possibly useful notion, we must consider a specific limited class of **Prover**'s strategies. The following "obliviousness" is our suggestion (roughly speaking, it restricts **Prover** to refer the information of c(L) and L(c(L)) only):

**Definition 4.10.** Let  $n, C \in \mathbb{N}$ . Suppose T is an (n, C)-tree. **An oblivious strategy** of Prover for  $\mathcal{G}_2(n, C, T)$  is a pair  $(f_1, f_2)$  of functions such that:

• 
$$f_1: \bigcup_{l=0}^C \left( [2^{|n|^C}]^l \times \mathcal{M}_n \times [2^{|n|^C}]^l \right) \to (2^{P_n \dot{\cup} H_n})_{\leq |n|^C}.$$

•  $f_2: \bigcup_{l=0}^C \left( [2^{|n|^C}]^l \times \mathcal{M}_n \times [2^{|n|^C}]^l \times \mathcal{M}_n \right) \to [3] \times [2^{|n|^C}]^{\leq C} \times [2^{|n|^C}], \text{ and } \langle o, x, B \rangle = f_2(v, M, \vec{A}, M') \text{ satisfies } x \in [2^{|n|^C}] \text{ if } o = 1 \text{ and } x \subsetneq v \text{ otherwise.}$ 

An important point of Theorem 4.13 is that, while **Prover** is restricted to oblivious strategies, **Delayer** has no restriction; she can see the whole position L and make a decision:

**Definition 4.11.** A strategy of Delayer for  $\mathcal{G}_2(n,C,T)$  is a function g such that:

•  $g: \mathcal{P} \times (2^{P_n \dot{\cup} H_n})_{\leq |n|^C} \to \mathcal{M}_n$ , and g(L, Q) is a minimal partial matching covering Q for each (L, Q) in the domain.

**Definition 4.12.** Let g be **Delayer**'s strategy for  $\mathcal{G}_2(n, C, T)$ , and  $(f_1, f_2)$  be an oblivious **Prover**'s strategy for  $\mathcal{G}_2(n, C, T)$ .  $(f_1, f_2)$  beats g if and only if **Delayer** wins  $\mathcal{G}_2(n, C, T)$  if the two players play in the following way:

- 1. **Prover** plays  $Q = f_1(v, M, \vec{A})$  at stage 4a in Definition 4.2, where  $v := c(L), (M, \vec{A}) := L(v)$ .
- 2. **Delayer** answers M' = g(L, Q) at stage 4b.
- 3. **Prover** then plays  $\langle o, x, B \rangle = f_2(v, M, \vec{A}, M')$  at stage 4c.

Now, we prove a counterpart of Proposition 3.7 for  $\mathcal{G}_2$ . The proof can be regarded as a transposition of Buss's witnessing argument ([9], [11]).

**Theorem 4.13.** Suppose  $ontoPHP_n^{n+1}$  has  $2^{|n|^{O(1)}}$ -sized  $LK_{2+\frac{1}{2},O(1)}^*$ -proofs. Then, there exists C > 0 such that, for any sufficiently large n, there exists an (n,C)-tree T such that **Prover** has an oblivious strategy  $(f_1, f_2)$  for  $\mathcal{G}_2(n, C, T)$  which beats arbitrary **Delayer**'s strategy g.

*Proof.* Our approach is similar to the sketch given in the last section. The main differences are:

- Since we treat  $LK_{2+\frac{1}{2},O(1)}^*$ -proofs this time, the witness function W[P] will be defined on not only the  $s\Pi_1$ -formulae but also the  $p\Pi_2$ -formulae. It will be represented by  $\vec{A}$  in the following argument.
- $\bullet$  When we consider  $\bigvee$ -Rule:

$$\frac{\Gamma, \varphi_{i_0}}{\Gamma} \quad \text{(where } \bigvee_{i=1}^{I} \varphi_i \in \Gamma, \ 1 \le i_0 \le I, \ \varphi_{i_0} \notin \Gamma)$$

 $\bigvee_{i=1}^{I} \varphi_i$  is either in  $ontoPHP_n^{n+1}$  or  $p\Sigma_1$  or  $p\Sigma_2$ . In the first case, the arguments are the same. In the last case, we argue as if  $\varphi_{i_0}$  was a satisfied  $p\Sigma_0$ -formula in the case when  $\bigvee_{i=1}^{I} \varphi_i$  was  $p\Sigma_1$  in the sketch. If  $\bigvee_{i=1}^{I} \varphi_i$  is  $p\Sigma_1$ , then **Prover** makes a query to decide  $\varphi_{i_0}$ , which is  $p\Sigma_0$ , and if it is falsified, he just climbs up the proof tree, but if it is satisfied, then he backtracks to the point where  $\bigvee$ -Rule is applied to eliminate  $\bigwedge_{i=0}^{k} \overline{\varphi_i}$ . Now he has a counterexample  $i_0$  for it and restarts the play from the point, climbing up the  $\bigvee$ -Rule which he once went through assuming  $\bigwedge_{i=0}^{k} \overline{\varphi_i}$  was true.

Now, we present a precise proof. By assumption, there exists C>0 such that, for any sufficiently large n, there exists an  $LK_{2+\frac{1}{2},C}^*$ -derivation  $\pi=(\mathcal{T},\mathcal{S})$  of  $ontoPHP_n^{n+1}$  satisfying  $|\pi|\leq 2^{|n|^C}$ . By an isomorphism, we may assume that  $\mathcal{T}$  is an (n,C)-tree. Note that each  $\varphi$  appearing in  $\pi$  is  $s\Sigma_2^{2^{|n|^C},|n|^C}$  or  $s\Pi_2^{2^{|n|^C},|n|^C}$  by Remark 2.10. For readability, we omit the superscripts and write  $s\Pi_i$ ,  $s\Sigma_i$ ,  $p\Pi_i$  and so on below.

We extract an oblivious winning strategy of **Prover** for  $\mathcal{G}_2(n, C, \mathcal{T})$  from this  $\pi$ . The strategy sticks to play so that, for each position L,

• Given  $v \in \text{dom}(L)$ , let  $L(v) = (M, A_1, \dots, A_{height(v)})$ . Note that  $height(v) \leq C$  by our formalization of  $LK_{2+\frac{1}{2},C}^*$ . Let  $\varphi \in \mathcal{S}(v)$ .

According to the complexity of  $\varphi$ , the following hold:

- 1. If  $\varphi$  is  $s\Sigma_0$ , then  $M \Vdash \overline{\varphi}$ .
- 2. Consider the case when  $\varphi$  is  $p\Pi_2$ . Let  $\varphi = \bigwedge_{e=1}^E \psi_e$ , where each  $\psi_e$  is  $p\Sigma_1$ . It must be eliminated by  $p\Sigma_2$ -Induction along the path from v to the root in  $\pi$ . Let  $w \subseteq v$  be the vertex such that  $\mathcal{S}(w)$  is derived by  $p\Sigma_2$ -Induction and  $\varphi$  is eliminated there. Then,  $A_{height(w)+1} \leq k$ .
- 3. Consider the case when  $\varphi$  is  $p\Pi_1$ . Let  $\varphi = \bigwedge_{e=1}^E \psi_e$ , where each  $\psi_e$  is  $s\Sigma_0$ .  $\varphi$  must be eliminated along the path from v to the root in  $\pi$ , and the derivation rule must be  $\bigvee$ -Rule.

Let  $u \subseteq v$  be the vertex such that  $S(u) = \Gamma$  is derived by:

$$\frac{\Gamma, \varphi_{i_0}}{\Gamma} \quad \text{(where } \bigvee_{i=1}^{I} \varphi_i \in \Gamma, \ \varphi = \varphi_{i_0}, \ 1 \le i_0 \le I, \ \varphi \not\in \Gamma)$$

Since every formula appearing in  $\pi$  is  $s\Sigma_2$  or  $s\Pi_2$ , we have that  $\bigvee_{i=1}^I \varphi_i$  is  $p\Sigma_2$ , and therefore it must be an element of  $ontoPHP_n^{n+1}$  or eliminated by  $p\Sigma_2$ -Induction along the path from u to the root in  $\pi$ .

In the former case, if

$$\bigvee_{i=1}^{I} \varphi_i = \bigvee_{p \in [n+1]} \bigwedge_{h \in [n]} \neg r_{ph},$$
$$\varphi = \bigwedge_{h \in [n]} \neg r_{i_0h},$$

then  $A := A_{height(u)+1} \in [n]$ . Moreover,  $M \Vdash \overline{\neg r_{i_0 A}}$ . If

$$\bigvee_{i=1}^{I} \varphi_i = \bigvee_{h \in [n]} \bigwedge_{p \in [n+1]} \neg r_{ph},$$
$$\varphi = \bigwedge_{p \in [n+1]} \neg r_{pi_0},$$

then  $A := A_{height(u)+1} \in [n+1]$ . Furthermore,  $M \Vdash \overline{\neg r_{Ai_0}}$ . In the latter case, then,  $A := A_{height(u)+1} \leq I$ , and  $M \Vdash \overline{\psi_A}$ .

Intuitively, according to the strategy, **Prover** always stores a counterexample for each  $p\Pi_1$ -formula  $\varphi \in \mathcal{S}(v)$  and a candidate of a counterexample for each  $p\Pi_2$ -formula, under  $\rho^M$ , in  $\vec{A}$ .

• For each  $w \in \text{dom}(L)$ , if  $S(w) = \Gamma$  is derived by  $p\Sigma_2$ -Induction:

$$\begin{array}{cccc}
\Gamma, \varphi_1 & \Gamma, \overline{\varphi_1}, \varphi_1 & \cdots & \Gamma, \overline{\varphi_{m-1}}, \varphi_m & \Gamma, \overline{\varphi_m}
\end{array}$$

then  $w*(k+1) \in \text{dom}(L)$  implies  $w*k \in \text{dom}(L)$  (note that  $k \in [m]$ ). Furthermore, if  $w*(k+1) \in \text{dom}(L)$ , then, for the maximum leaf  $l \in \text{dom}(L)$  properly extending w\*k,  $S(l) = \Delta$  is derived by  $\bigvee$ -Rule

$$\frac{\Delta, \psi_{i_0}}{\Delta}$$
  $(\bigvee_{i=1}^{I} \psi_i \in \Delta, 1 \le i_0 \le I)$ 

and the principal formula  $\bigvee_{i=1}^{I} \psi_i$  is the  $\varphi_k$  eliminated from  $\mathcal{S}(w * k)$  deriving  $\mathcal{S}(w)$  (therefore each  $\psi_i$  is  $p\Pi_1$ ). Furthermore, if  $L(w * (k+1)) = (M, \vec{A})$ , then  $A_{height(w*(k+1))} = i_0$ .

• For each non-leaf  $w \in \text{dom}(L)$ , if  $S(w) = \Gamma$  is derived by  $\Lambda$ -Rule:

$$\frac{\Gamma, \varphi_1 \qquad \Gamma, \varphi_2 \qquad \cdots \qquad \Gamma, \varphi_I}{\Gamma} \quad \text{(where } \bigwedge_{i=1}^I \varphi_i \in \Gamma, \text{ and } \varphi_i \not\in \Gamma \text{ for each } i \in [I])$$

then w has a unique child in dom(L). If  $\bigwedge_{i=1}^{I} \varphi_i$  is  $p\Pi_1$  or  $p\Pi_2$  in particular, then, letting A be the counterexample (or the candidate of a counterexample)  $A_{height(u)+1}$  for it described in (3) at the first item, the unique child is w \* A.

Now, we describe the strategy explicitly. Suppose we are at a position L satisfying the above conditions. Note that the initial position trivially satisfies them. Let

$$v := c(L), (M, \vec{A}) := L(v).$$

We describe  $f_1(v, M, \vec{A})$  and  $f_2(v, M, \vec{A}, M')$  splitting cases by the rule deriving S(v) in  $\pi$ . Note that, in each case, the conditions above remain satisfied. S(v) is not an Initial cedent since if it was the case, then the literals x and  $\bar{x}$  in S(v) should be falsified by M, which is absurd.

1. The case when  $S(v) = \Gamma$  is derived by  $\vee$ -Rule:

$$\frac{\Gamma, \varphi_{i_0}}{\Gamma} \quad \text{(where } \varphi_1 \vee \varphi_2 \in \Gamma, i_0 = 1 \text{ or } i_0 = 2, \, \varphi_{i_0} \not\in \Gamma)$$

Since  $\varphi_1 \vee \varphi_2$  is  $s\Sigma_2$  or  $s\Pi_2$ , it must be  $s\Sigma_0$ . Set  $f_1(v, M, \vec{A}) := \emptyset$ . **Delayer**'s answer should be  $\emptyset$ , and so set

$$f_2(v, M, \vec{A}, \emptyset) := \langle 1, 1, * \rangle,$$

where \* can be any number in  $[2^{|n|^C}]$ .

- 2. The case when  $S(v) = \Gamma$  is derived by Trivial Cut or  $\land$ -rule: similar as above.
- 3. The case when  $S(v) = \Gamma$  is derived by  $p\Sigma_2$ -Induction:

$$\frac{\Gamma, \varphi_1 \qquad \Gamma, \overline{\varphi_1}, \varphi_2 \qquad \cdots \qquad \Gamma, \overline{\varphi_{I-1}}, \varphi_I \qquad \Gamma, \overline{\varphi_I}}{\Gamma} \quad \text{(where each } \varphi_i \text{ is } p\Sigma_2\text{)}$$

Set  $f_1(v, M, \vec{A}) := \emptyset$ . **Delayer**'s answer should be  $\emptyset$ , and so set

$$f_2(v, M, \vec{A}, \emptyset) := \langle 1, 1, * \rangle,$$

where \* can be any number in  $[2^{|n|^C}]$ .

4. The case when  $S(v) = \Gamma$  is derived by  $\bigvee$ -Rule:

$$\frac{\Gamma, \varphi_{i_0}}{\Gamma} \quad \text{(where } \bigvee_{i=1}^{I} \varphi_i \in \Gamma, \ 1 \le i_0 \le I, \ \varphi_{i_0} \notin \Gamma)$$

We further split cases according to the complexity of  $\bigvee_{i=1}^{I} \varphi_i$ .

(a) First we consider the case when  $\bigvee_{i=1}^{I} \varphi_i \in ontoPHP_n^{n+1}$ . If

$$\bigvee_{i=1}^{I} \varphi_i = \bigvee_{\substack{p \neq p' \in [n+1], \\ h \in [n]}} (r_{ph} \wedge r_{p'h}),$$

let  $i_0 = \langle p, p', h \rangle$ . Then set  $f_1(v, M, \vec{A}) := \{p, p'\}$ . <sup>2</sup> Let M' be **Delayer**'s answer. M' always falsifies  $\varphi_{i_0}$ , so set

$$f_2(v, M, \vec{A}, M') := \langle 1, 1, * \rangle.$$

If

$$\bigvee_{i=1}^{I} \varphi_i = \bigvee_{p \in [n+1]} \bigwedge_{h \in [n]} \neg r_{ph}$$

and  $\varphi_{i_0} = \bigwedge_{h \in [n]} \neg r_{i_0 h}$ , then set

$$f_1(v, M, \vec{A}) := \{i_0\}.$$

Let  $M' = \{i_0 \mapsto h_0\}$  be **Delayer**'s answer. M' together with the witness  $j_0$  falsifies  $\varphi_{i_0}$ , so set

$$f_2(v, M, \vec{A}, M') := \langle 1, 1, h_0 \rangle.$$

If  $\bigvee_{i=1}^{I} \varphi_i$  is one of other elements in  $ontoPHP_n^{n+1}$ , change the roles of pigeons and holes and play similarly as above.

Note that **Prover** does not lose by choosing o = 1 since  $v * 1 \in \mathcal{T}$ .

(b) Else if  $\bigvee_{i=1}^{I} \varphi_i$  is  $p\Sigma_2$  and  $\varphi_{i_0}$  is  $p\Pi_1$ , then, since  $\bigvee_{i=1}^{I} \varphi_i \not\in ontoPHP_n^{n+1}$ , there uniquely exists  $w * e \subseteq v$  such that  $\mathcal{S}(w)$  is derived by  $p\Sigma_2$ -Induction, and  $\bigvee_{i=1}^{I} \varphi_i$  is eliminated there. By definitions,

$$w * (e+1) \in \mathcal{T} \& \bigwedge_{i=1}^{I} \overline{\varphi_i} \in \mathcal{S}(w * (e+1)).$$

Based on this, set

$$f_1(v, M, \vec{A}) := \emptyset, f_2(v, M, \vec{A}, \emptyset) := \langle 2, w, i_0 \rangle.$$

Note that **Prover** does not lose by choosing o = 2 since  $w * (e + 1) \in \mathcal{T}$ .

<sup>&</sup>lt;sup>2</sup>Actually, we may ask just the hole h here, but if we were to consider the case when we deal with the injective pigeonhole principle  $injPHP_n^{n+1}$  instead of  $ontoPHP_n^{n+1}$  and would like to ask only pigeon-queries in the course of the corresponding game, the query  $\{p,p'\}$  is the right one. (note that we do not have the last two formulae of  $ontoPHP_n^{n+1}$  in that case.)

(c) Else if  $\bigvee_{i=1}^{I} \varphi_i$  is  $p\Sigma_1$  and  $\varphi_{i_0}$  is  $s\Sigma_0$ , then, let  $f_1(v, M, \vec{A})$  be the set of pigeons<sup>3</sup> appearing as indices of variables occurring in  $\varphi_{i_0}$ , which is at most cardinality  $\leq |n|^C$ .

Let M' be **Delayer**'s answer. M' decides  $\varphi_{i_0}$ , that is,

$$M' \Vdash \varphi_{i_0} \text{ or } M' \Vdash \overline{\varphi_{i_0}}.$$

If  $M' \Vdash \overline{\varphi_{i_0}}$ , then set

$$f_2(v, M, \vec{A}, M') := \langle 1, 1, * \rangle.$$

If  $M' \Vdash \varphi_{i_0}$ , then, since  $\bigvee_{i=1}^{I} \varphi_i \not\in ontoPHP_n^{n+1}$ , there exists  $w * e \subseteq v$  such that  $S(w) = \Theta$  is derived by  $p\Sigma_2$ -Induction:

$$\Theta, \Phi_1 \qquad \Theta, \overline{\Phi_1}, \Phi_2 \qquad \cdots \qquad \Theta, \overline{\Phi_{E-1}}, \Phi_E \qquad \Theta, \overline{\Phi_E} \qquad \text{(where each } \Phi_e \text{ is } p\Sigma_2\text{)}$$

 $e \geq 2$ , and  $\overline{\Phi_{e-1}} = \bigwedge_{\alpha} \bigvee_{\beta=0}^{B_{\alpha}} \psi_{\alpha\beta}$  has  $\bigvee_{i=1}^{I} \varphi_{i}$  as one of its conjuncts, that is, there is  $\alpha_{0}$  such that  $\bigvee_{i=1}^{I} \varphi_{i} = \bigvee_{\beta=0}^{B_{\alpha_{0}}} \psi_{\alpha_{0}\beta}$ . Furthermore, there exists u with  $w * e \subseteq u \subsetneq v$  such that  $\Lambda = \mathcal{S}(u)$  is derived by  $\Lambda$ -Rule:

$$\frac{\Lambda, \bigvee_{\beta=0}^{B_1} \psi_{1\beta} \qquad \Lambda, \bigvee_{\beta=0}^{B_2} \psi_{2\beta} \qquad \cdots \qquad \Lambda, \bigvee_{\beta=0}^{B_m} \psi_{m\beta}}{\Lambda}$$

where  $\overline{\Phi_{e-1}} = \bigwedge_{\alpha} \bigvee_{\beta=0}^{B_{\alpha}} \psi_{\alpha\beta} \in \Lambda$  and  $u * \alpha_0 \subseteq v$ . We set

$$f_2(v, M, \vec{A}, M') := \langle 3, w, i_0 \rangle.$$

The resulting next position again satisfies the assumptions of the strategy. Indeed, by assumption, the counterexample  $A_{height(w)+1}$  for  $\overline{\Phi_{e-1}}$  stored at w\*e is  $\alpha_0$ . We consider the maximum leaf  $l \in \text{dom } L$  extending w\*(e-1). Note that  $w*(e-1) \in \text{dom}(L)$  by assumption.  $S(l) = \Delta$  is derived by the following V-Rule:

$$\frac{\Delta, \bigwedge_{\beta=0}^{B_{\alpha_0}} \overline{\psi_{\alpha_0 \beta}}}{\Delta} \quad \text{(where } \Phi_{e-1} = \bigvee_{\alpha} \bigwedge_{\beta=0}^{B_{\alpha}} \overline{\psi_{\alpha \beta}} \in \Delta\text{)}$$

The following figure shows the whole situation (the notation @v shows the corresponding vertices):

<sup>&</sup>lt;sup>3</sup>Similarly to before, we may also query holes here.

$$\begin{array}{c} \vdots \\ \hline \Gamma, \varphi_{i_0} \quad (@v*1) \\ \hline \Gamma \quad (@v) \\ \hline \\ \Delta, \bigwedge_{\beta=0}^{B_{\alpha_0}} \overline{\psi_{\alpha_0\beta}} \quad (@l*1) \\ \hline \\ \Delta \quad (@l) \\ \hline \vdots \\ \hline \\ \Theta, \overline{\Phi_{e-2}}, \Phi_{e-1} \quad (@w*(e-1)) \\ \hline \\ \Theta \quad (@w) \\ \hline \end{array} \begin{array}{c} \vdots \\ \hline \\ \Lambda, \bigvee_{\beta=0}^{B_{\alpha_0}} \psi_{\alpha_0\beta} \quad (@u*\alpha_0) \\ \hline \\ \Lambda \quad (@u) \\ \hline \\ \vdots \\ \hline \\ \Theta, \overline{\Phi_{e-1}}, \Phi_{e} \quad (@w*e) \\ \hline \\ \Theta \quad (@w) \\ \hline \end{array}$$

Here, we disregard  $\overline{\Phi_{e-2}}$  when e=2.

Hence, recalling  $\bigwedge_{\beta=0}^{B_{\alpha_0}} \overline{\psi_{\alpha_0\beta}} = \bigwedge_{i=1}^{I} \overline{\varphi_i}$  and  $M' \Vdash \overline{\varphi_{i_0}}$ , the claim follows.

(d) Otherwise,  $\bigvee_{i=1}^{I} \varphi_i$  is  $p\Sigma_0$  and M already falsifies it, so set

$$f_1(v, M, \vec{A}) := \emptyset, f_2(v, M, \vec{A}, \emptyset) := \langle 1, 1, * \rangle.$$

5. The case when  $S(v) = \Gamma$  is derived by  $\Lambda$ -Rule:

$$\frac{\Gamma, \varphi_1 \qquad \Gamma, \varphi_2 \qquad \cdots \qquad \Gamma, \varphi_I}{\Gamma} \quad \text{(where } \bigwedge_{i=1}^I \varphi_i \in \Gamma, \text{ and } \varphi_i \not \in \Gamma \text{ for each } i \in [I])$$

If  $\bigwedge_{i=1}^{I} \varphi_i$  is  $s\Sigma_0$ , it is already falsified, hence there exists k such that  $M \Vdash \overline{\varphi_k}$ . Let k be the least one, and set

$$f_1(v, M, \vec{A}) := \emptyset, f_2(v, M, \vec{A}, \emptyset) := \langle 1, k, * \rangle.$$

If  $\bigwedge_{i=1}^{I} \varphi_i$  is  $p\Pi_1$  or  $p\Pi_2$ , By assumption, a counterexample  $A_e$  for it is stored in  $\vec{A}$ . Set

$$f_1(v, M, \vec{A}) = \emptyset, f_2(v, M, \vec{A}, \emptyset) := \langle 1, A_e, * \rangle,$$

Note that  $M \Vdash \overline{\varphi_{A_e}}$  if  $\bigwedge_{i=1}^I \varphi_i$  is  $p\Pi_1$ .

This completes the description of **Prover**'s strategy. Since **Prover** can continue the play as long as **Delayer** does not make a contradiction, and dom(L) strictly increases in terms of  $\prec$  by Lemma 4.6, if **Delayer** wins, c(L) reaches to one of Initial cedents, which is absurd since an Initial cedent can never be falsified. Hence, **Prover**'s strategy above is a winning one.

### 5 Analysis of simplified $G_2$

As far as we see, analysis of general  $\mathcal{G}_2(n,C,T)$  is rather difficult.

Thus, in this section, we focus on  $\mathcal{G}_2(n, C, T)$  of minimal height C allowing backtracking, namely, C = 2.

Even this case is hard to analyze, so we also assume that all the queries in the play are of size 1. Moreover, we further restrict **Prover**'s oblivious strategy  $(f_1, f_2)$  to the following ones:

- 1. Let  $s \in [2^{|n|^C}]$ .
- 2. There exists a map

$$\mathcal{A} \colon (\mathcal{M}_n)_{\leq 1} \to P_n$$

such that, for  $v \in [2^{|n|^C}]^{\leq 1}$ ,  $f_1(v, M, A) = \{A(M)\}$ .

3. Let  $P_n = \{p_1, \dots, p_{n+1}\}$ . Let  $p_i = \mathcal{A}(\emptyset)$ . Then, for any  $h \in H_n$ ,

$$f_2(\emptyset, L_0(\emptyset), \{p_i \mapsto h\}) = \langle 1, 1, 1 \rangle.$$

4. For  $v \in [2^{|n|^C}]^1$ ,  $M, M' \in \mathcal{M}_n$  of size 1, and  $A \in [s]$ ,

$$f_2(v, M, A, M') = \begin{cases} \langle 2, \emptyset, A + 1 \rangle & \text{ (if } A < s) \\ \langle 3, \emptyset, k \rangle & \text{ (if } A = s, \text{ and } M' \text{ covers } p_k) \end{cases}.$$

5. For  $v \in [2^{|n|^C}]^2$ ,  $M \in \mathcal{M}_n$  of size 2, and  $A_1, A_2 \in [s]$ , if  $A_2 \in [n+1]$  and M matches  $p_{A_2}$  to  $h \in H_n$ , then

$$f_1(v, M, A_1, A_2) = \{p_{A_2}\},\$$

and

$$f_2(v, M, A_1, A_2, \{p_{A_2} \mapsto h\}) = \langle 3, \emptyset, A_2 \rangle.$$

Note that values of  $f_1$  and  $f_2$  for other inputs do not matter to determine the winner. This very limited "toycase" of  $\mathcal{G}_2(n, C, T)$  is still nontrivial since s may be greater than n, which gives advantage to **Prover**. Actually, it turns out that the bound of s above is not essential in this situation.

Below we present the mentioned restriction in a self-contained manner, with notation and terminology consistent with the previous sections.

**Definition 5.1.** Fix an integer  $n \geq 1$ . We define the following **game** (denoted as  $\mathcal{G}_2(n)$ ) played by two players: **Prover** (denoted as **P** and referred to as he) and **Delayer** (denoted as **D** and referred to as she).

**D** claims the existence of an injective function f from the set  $\{0, ..., n\}$  (denoted as  $P_n$ ) into the set  $\{0, ..., n-1\}$  (denoted as  $H_n$ ). We think of  $P_n$  as a set of *pigeons* and  $H_n$  as a set of *holes*.

Through the course of the game **P** asks **D** questions of the form "Where is the pigeon  $p \in P_n$  being mapped to by f" (we denote such question as p itself) and **D** must answer with "p is being mapped to h by f" for some  $h \in H_n$  (we denote such answer as h itself). These answers are then stored by **P** in the form (p, h).

**P**'s goal in the game is to force **D** into giving two contradictory answers. This means **P** wins iff **D** answers h to the question p, while **P**'s list of records contains either (p',h) or (p,h') for  $p' \neq p$  and  $h' \neq h$ .

At each moment during the game,  $\mathbf{P}$  can only store at most 2 records in his list. This means that before asking a new question  $\mathbf{P}$  must make room for an upcoming answer from  $\mathbf{D}$ . We assume  $\mathbf{P}$  always removes the oldest record.

Finally, before the game starts, **P** chooses an integer parameter  $s \ge 1$  which **D** sees, as well. As soon as the total number of the given answers in a game reaches s, the game stops, and the final answer of **D** is compared to all the records that have appeared through the whole game. If there is a single contradiction of the form (p, h), (p', h) or (p, h), (p, h') for  $p' \ne p$  and  $h' \ne h$  and with p being the last question and h being the last answer, **P** wins, otherwise **D** wins.

**Definition 5.2.** A *strategy* for **P** is a pair (s, F) of an integer  $s \ge 1$  and a function F which on input the empty set (the initial game state) or a single record (p, h) (**P**'s list after the removal of an older record) outputs a question p' which **P** should ask next.

We say **P** plays a game according to the strategy (s, F) iff his zeroth move is to claim s as the game's initial parameter, his first move is to ask a question  $F(\emptyset)$  and, finally, after the removal of an older record and being left with a single record (p, h), if the game is not yet finished, **P** asks F((p, h)).

From now on F((p,h)) will be denoted simply as F(p,h).

We say a strategy (s, F) is **winning** for **P** iff, for plays in which **D** never gives two contradictory answers in a row, any last answer of **D** in a game where **P** plays according to (s, F) contradicts at least one prior **D**'s answer.

The ultimate goal is to show that for any  $n \geq 3$ , **P** has no winning strategies. However, before this, we show two simple cases when **P** can win.

**Proposition 5.3.** For  $n \leq 2$  there is a winning strategy for **P** with s being any number > n + 1.

*Proof.* We start with the case n=2.

Let s be 3. **P** starts by asking  $0 \in P_2$ . Then, no matter what the answer is, **P** asks  $1 \in P_2$ . Since **D** cannot give two contradictory answers, after the second step, **P**'s list should contain records (0, h) and (1, h') with  $h \neq h'$ .

**P** removes the older record (0, h) and asks  $2 \in P_2$ . Since s = 3, the game stops after **D** answers that question. Such an answer is of the form h'', but since  $H_n$  is of size 2 and  $h \neq h' \in H_n$ , it follows that h'' = h or h'' = h'. Since we assumed **D** never gives two contradictory answers in a row, it follows h'' = h and so the record (2, h'') contradicts the record (0, h).

For s > 3 it is enough to start as before, but after the 3rd answer, **P** remains to ask  $2 \in P_2$  throughout the rest of the game. Since **D** cannot give two contradictory answers, she will be giving the same answer for the rest of the game and we have seen that the corresponding record (2, h'') necessarily contradicts the first record (0, h).

For the remaining case n = 1, **P** can win even before the game reaches s, since asking first  $0 \in P_1$  and then  $1 \in P_1$  necessary results in a contradiction.

In the definition of the game we have assumed  $P_n = \{0, ..., n\}$  and  $H_n = \{0, ..., n-1\}$ . For the next two propositions, we will consider a more general case with the gap between the sizes of  $P_n$  and  $H_n$  much larger than 1.

**Proposition 5.4.** Let  $P_n$  be  $\{0, ..., 1 + n + n^2 + ... + n^n\}$  and  $H_n$  be  $\{0, ..., n - 1\}$ . Then, for any  $n \ge 1$ , **P** has a winning strategy with s being any parameter  $\ge n + 1$ .

*Proof.* By the same argument as above, we may assume s = n + 1. The general case would follow by repeating the last question until s is reached.

The idea is for  $\mathbf{P}$  to utilize the abundance of possible questions to encode the previous answers of  $\mathbf{D}$ .

Concretely, we start by defining inductively a sequence of sets  $S_n$ . Let  $S_0 = \{0\}$ . For i+1 we define  $S_{i+1} = \{(p,h) \mid p \in S_i, h \in H_n\}$ . Notice that  $|\bigcup_{i \leq n} S_i| = 1 + n + \dots + n^n$ . We identify elements of the set  $P_n \setminus \{1 + n + n^2 + \dots + n^n\}$  with elements of  $\bigcup_{i \leq n} S_i$  via an arbitrary bijection. The remaining single element of  $P_n$  is denoted as \*.

The winning strategy for **P** is as follows. **P** starts with asking  $0 \in S_0$ . Receiving an answer h, he goes on to ask  $(0, h) \in S_1$ .

In the beginning of the (i+1)st step, **P** has record of the form  $(((0,h_1),h_2)...,h_i)$ . If  $h_i$  is among  $h_1,...,h_{i-1}$ , then **P** asks  $(((0,h_1),h_2)...,h_{i-1})$  and **D** must answer  $h_i$  which is contradictory to an answer from some time before. In case  $h_1,...,h_i$  are all different, **P** asks  $(((0,h_1),h_2)...,h_i)$ .

In the sth step, if all  $h_1, ..., h_s$  are different, then **P** asks \*. Otherwise, proceed as in the similar case from above.

Notice that the last question being \* is equivalent to all the holes (elements of  $H_n$ ) being exhausted through the course of the game and so any answer would necessarily contradict an older one. In case **D**'s answer contradicts a previous one already before the sth step, then **P** would notice it since **P**'s record codes all the previous answers given

by  $\mathbf{D}$ . Playing according to the mentioned strategy  $\mathbf{P}$  would then force  $\mathbf{D}$  to give this exact contradictory answer until the game ends.

It is possible to get the gap between the sizes of  $P_n$  and  $H_n$  even smaller.

**Proposition 5.5.** Let  $P_n$  be  $\{0, ..., 2^n - 1\}$  and  $H_n$  be  $\{0, ..., n - 1\}$ . Then, for any  $n \ge 1$  **P**, has a winning strategy with s being any parameter  $\ge n + 1$ .

*Proof.* Fix s = n + 1 as before. We let the set  $P_n$  be labeled by all the subsets of  $H_n$ . The first question is  $\emptyset$ . Having a record of the form  $(S \subset H_n, h)$ , if  $h \in S$ , then **P** asks S. Otherwise, **P** asks  $S \cup \{h\}$ .

At the beginning of the sth step, **P**'s record might be  $(S \subset H_n, h)$  with  $h \in S$ , in which case asking S again results in h, which necessarily contradicts some previous answer. Otherwise, **P**'s record is (S, h) with  $S \cup \{h\} = H_n$ , since s = n + 1. Asking  $H_n$  then results in a record  $(H_n, h')$  with h' being an answer to some previous **P**'s question  $S \subseteq H_n$ .

We leave as an open question whether one can get the gap between the sizes of  $P_n$  and  $H_n$  smaller, even by a constant.

**Question 5.6.** Do there exist integers  $c \ge 2$  and  $n_0, s_0$  such that, for all  $n \ge n_0$  and  $s \ge s_0$ , **P** has a winning strategy for a game played with  $P_n = \{0, ..., 2^n - c\}$  and  $H_n = \{0, ..., n-1\}$ ?

For the rest of the paper, we consider the case with  $P_n = \{0, ..., n\}$  and  $H_n = \{0, ..., n-1\}$  as in the original definition. We already know that  $\mathbf{P}$  can win in a game for  $n \leq 2$ . What we are going to show is that for any  $n \geq 3$ ,  $\mathbf{P}$  cannot win. The proof can be viewed either as a proof-by-contradiction, where we show that no strategy can be a winning one or as a construction of an *anti-strategy* for  $\mathbf{D}$  given a fixed strategy for  $\mathbf{P}$ . Here, anti-strategy is defined as a strategy for  $\mathbf{D}$  which moreover depends on a chosen strategy for  $\mathbf{P}$ .

**Proposition 5.7.** Let  $n \geq 3$  and (s, F) be a strategy for **P**. If  $s \leq n$ , then there is a winning anti-strategy for **D**. Moreover, such anti-strategy does not depend on F.

*Proof.* The anti-strategy is very simple. Being presented with a question p,  $\mathbf{D}$  answers h so that

- 1. if **P** has not asked p before, then h is any hole from  $H_n$  which has not been used by **D** yet;
- 2. if **P** has asked p before, then h is the same as the one **D** answered to p for the first time.

Since  $s \leq n$ , **D** will never run out of holes before the game ends. Notice that through the whole game, all answers of **D** are *compatible* (i.e. are not contradictory) with any previous answers. This shows that the sth answer is not contradictory, as well.

**Definition 5.8.** Any anti-strategy as in the above proof is called a  $canonical\ anti-strategy$  for D.

**Proposition 5.9.** Let  $n \geq 3$  and (s, F) be a strategy for **P**. Assume **D** is utilizing a canonical anti-strategy and case 2. from the proof of 5.7 occurs, i.e. **D** is being asked p which **P** has already asked before in a play where all the answers given so far by **D** are compatible. Then, the chosen anti-strategy is a winning strategy for **D** no matter how large the s is.

*Proof.* Notice that after  $\mathbf{P}$  asks p which he already asked before, the game essentially loops, i.e. the records and their order of appearance are the same as after  $\mathbf{D}$  answered h to p for the first time. Since all  $\mathbf{D}$ 's answers are compatible with all the previous ones, no matter how large s is, the last answer is always a winning one for  $\mathbf{D}$ .

We now need a better way to represent  $\mathbf{P}$ 's strategies. This will help us to construct anti-strategies for  $\mathbf{D}$ .

**Definition 5.10.** Let F be a functional part of  $\mathbf{P}$ 's strategy.

A **graph** associated with F is a directed labeled multigraph  $\mathcal{G}_F = (G_F, E_F, l_F)$  defined as follows.

The set  $G_F$  of nodes equals  $P_n$ , and the labeling function  $l_F$  maps the set  $E_F$  of directed edges to the set  $H_n$ .

The edges  $E_F$  and labels  $l_F$  are defined so that, for any two nodes p, p', there is an edge e going from p to p' labeled by  $h \in H_n$  iff F(p, h) = p'.

Notice that we allow loops, as well.

Two edges e, e' going from p and p', respectively, are *incompatible* (or *inconsistent*) iff either of the following conditions is satisfied

- p = p' and  $l_F(e) \neq l_F(e')$ ;
- $p \neq p'$  and  $l_F(e) = l_F(e')$ .

We say that a node  $p \in G_F$  is an *initial node* iff  $p = F(\emptyset)$ .

A **path** in  $\mathcal{G}_F$  is defined as a directed walk in  $\mathcal{G}_F$  which starts at the initial node. This means the same edges and nodes may appear multiple times in a path.

A path's *length* is defined as the number of edges.

We say that a path is *locally consistent* iff any two subsequent edges are compatible.

A path is called *globally consistent* iff any two edges of the path are consistent.

An edge of a path is called **globally consistent** iff it is consistent with all previous edges of the path.

Notice that two edges e, e' going from p, p', respectively, are consistent iff the set  $\{(p, l_F(e)), (p', l_F(e'))\}$  is a partial one-to-one mapping from  $P_n$  to  $H_n$ .

Figure 1 illustrates a simple example of  $\mathcal{G}_F$  for n=3.




Figure 1 - example of  $\mathcal{G}_F$ 

Fixing a strategy (s, F) of  $\mathbf{P}$  and assuming  $\mathbf{D}$  never gives two contradictory answers in a row, we can identify a play in the game with a locally consistent path in  $\mathcal{G}_F$  of length s. A play results in  $\mathbf{D}$ 's victory iff the last edge of the mentioned path is globally consistent. Thus, there is a winning anti-strategy for  $\mathbf{D}$  against (s, F) iff there is a locally consistent path in  $\mathcal{G}_F$  of length s with the last edge being globally consistent. In particular, there is a winning canonical anti-strategy for  $\mathbf{D}$  against (s, F) iff there is a globally consistent path in  $\mathcal{G}_F$  of length s.

We can now formulate the main theorem of this section in the following way.

**Theorem 5.11.** Let  $n \geq 3$  and (s, F) be a strategy for **P**. Then, there is a locally consistent path in  $\mathcal{G}_F$  of length s with the last edge being globally consistent.

*Proof.* By Proposition 5.7 it is enough to deal with the case  $s \ge n+1$ .

We begin by constructing a globally consistent path in  $\mathcal{G}_F$  of length n. This corresponds to  $\mathbf{D}$  using a canonical anti-strategy during the first n moves.

As was shown in Proposition 5.9, if the path we have constructed so far passes through the same node at least twice, then **D** wins against any s just by using this exact canonical anti-strategy for the rest of the game. Thus, we may assume each node appears exactly once in the path. So, after the path is constructed, all the elements of  $H_n$  viewed as labels of the edges of the path are exhausted along the way.

Without loss of generality, we assume the exact path we get is as shown in Figure 2.





Figure 2 - globally consistent path in  $\mathcal{G}_F$  of length n

Starting from this configuration, our goal is to gradually build up more and more information about  $\mathcal{G}_F$  by subsequently eliminating cases against which we can build winning anti-strategies for  $\mathbf{D}$ .

The final goal is to reduce the whole analysis to the configuration shown in Figure 3.




Let us first show that this reduction is indeed enough to show that **D** wins against any (s, F).

**Lemma 5.12.** Assume  $\mathcal{G}_F$  contains a subgraph as above. Then, **D** can build a locally consistent path of length s with the last edge globally consistent for any  $s \geq n+1$ .

*Proof.* To build such a path we need to analyze four different cases depending on where the edge from  $0 \in P_n$  and labeled by  $0 \in H_n$  go. This is equivalent to analyzing the value a so that F(0,0) = a.

1. a = 0. The desired path is shown in Figure 4.



Giving any  $s \ge n + 1$  and following such a path we are guaranteed that the final question is  $0 \in P_n$  with the answer  $0 \in H_n$ . This is a globally consistent answer.

In the above picture, only one edge is not globally consistent. From now on we will color those edges red to distinguish parts of the path we do not want to end up in and to make the whole analysis easier to visualize.

2. a=1. We have two different paths shown in Figure 5 which **D** should choose depending on the parity of s.




Figure 5 - cover-by-two for case a = 1

We say that these two paths form a **cover-by-two** since any possible  $s \ge n+1$  is covered by one of them in the sense that for each such s either of those paths ends up with an edge being globally consistent.

3. a = 2. Similarly to the previous case, we have two paths shown in Figure 6 forming a cover-by-two.




Figure 6 - cover-by-two for case a=2

4.  $a \ge 3$ . Again, we have two paths shown in Figure 7 forming a cover-by-two.

This exhausts all possible cases.

Goal I: derive that F(2,0) = 1, i.e. that the edge from  $2 \in P_n$  labeled by  $0 \in H_n$  goes to  $1 \in P_n$  (we will omit using this longer expression from now on). By derive we mean to show that any other case results in **D** being able to construct the desired path for any s.

Let F(2,0) = a. We assume  $a \neq 1$ . Notice that the only non-trivial case left is a = 0. All the other choices for a result in a possibility for **D** to construct a single path covering all possible s.




Figure 7 - cover-by-two for case  $a \ge 3$ 




Figure 8

Assume a = 0. Derive F(0, 1) = 1 as is shown in Figure 8.

We mark the edges we can derive immediately by  $\circledast$  from now on. Notice that assuming F(3,1)=1 results in a cover-by-two as is shown in Figure 9.

Let F(3,1) = b. We have just eliminated the possibility of b being equal to 1. There are two non-trivial cases left for b: either b = 0 or b = 2.

Assume b=0. The resulting configuration is shown in Figure 10.

By analyzing all the possible c = F(0,2) we see that any such choice results in **D** being able to construct a single winning path covering all s.

We conclude with b=2. Figure 11 shows this configuration together with an edge which can be immediately derived from it.

Notice that assuming F(2,2) = 0 we derive a cover-by-two shown in Figure 12.


We conclude that F(2,2) = 1. The resulting configuration is shown in Figure 13.

Let d = F(3,0). There are 3 non-trivial cases: d = 0, d = 1, or d = 2. The case d = 0 results in a cover-by-two shown in Figure 14.

The case d=2 results in a cover-by-two, as is shown in Figure 15.

Lastly, if d = 1, we conclude with Figure 16 from which one can easily find a desired path for **D** against any s.

Altogether, this finishes the derivation of the first goal. Thus, we conclude that






Figure 9 - cover-by-two derived from F(3,1) = 1




Figure 10

F(2,0) = 1 as is shown in Figure 17.

**Goal II**: derive that F(3,1) = 2. Notice that, by symmetry, this will imply that F(3,0) = 2, as well.

Let F(3,1) = a. There are 3 non-trivial cases: either a = 0, or a = 1, or a = 2.

Assume a = 0. We conclude with Figure 18.

From this we get a cover-by-two which is shown in Figure 19.

Assume a = 1. This leads to Figure 20, where we first derive that F(1,2) = 0, and then, with this additional information, we derive F(0,0) = 2. (The corresponding edge in the picture below is marked with (\*\*) to emphasize that its derivation comes second after the one marked with (\*).)

This results in a cover-by-two as is shown in Figure 21.

We conclude that the second goal is achieved. We end up with Figure 22. (Again, we mark by \*\*) the additional edge which can be immediately derived from the rest of the picture.)

To finish the whole proof, it is enough to derive that F(2,2) = 1. We assume the only non-trivial case left, namely, that F(2,2) = 0. This is shown in Figure 23.

We conclude with a cover-by-two shown in Figure 24.





Figure 11






Figure 12 - cover-by-two derived from F(2,2) = 0




Figure 13


Hence, F(2,2) = 1 and so we arrive at Figure 25. Applying the Lemma from the beginning, we finish the whole proof.

Below we present an alternative way of proving a result similar to Theorem 5.11. Instead of proving the statement for all  $n \geq 3$  simultaneously, we proceed inductively.

The first drawback is that instead of finding anti-strategies against all s, we will be able to construct them for all s above some threshold  $s_0$  which depends on n.

The second drawback is that we have to deal with the cases n=3 and n=4 separately.

Since we are not aware of any way of dealing with those cases in a principally simpler way than just proving 5.11, what we show is how one deals with the induction step only.







Figure 14 - cover-by-two for case d=0






Figure 15 - cover-by-two for case d=2




Figure 16 - an additional derivation for case d=1




Figure 17

Even though the mentioned proof strategy gives a weaker result than the one already presented, we feel the need to show it to build more intuition around the game  $\mathcal{G}_2$ .




Figure 18




Figure 19 - cover-by-two for case a = 0




Figure 20

We start by introducing a new notion which is used throughout the proof.

**Definition 5.13.** Let  $P_n = \{0, ..., n\}$  and  $H_n = \{0, ..., n-1\}$  as before. We define a **php-tree** for n as a rooted labeled tree  $\mathcal{T} = (V, E, l_V, l_E)$  satisfying the following conditions

• every node  $v \in V$  is labeled by an element  $l_V(v)$  from  $P_n$ ;




Figure 21 - cover-by-two for case a=1




Figure 22





Figure 23






Figure 24 - cover-by-two for case F(2,2)=0

- every edge  $e \in E$  is labeled by an element  $l_E(e)$  from  $H_n$ ;
- the branching number of any node on the kth level is  $\leq n-k$ , where a node v




Figure 25 - configuration for which Lemma 5.12 applies

is said to belong to the kth level iff the distance from v to the root is k and the branching number of a node v on the kth level is defined as the number of edges connecting v to its' children, i.e. nodes on the (k+1)st level;

- along any path starting from the root, any label of a node appears at most once;
- along any path starting from the root, any label of an edge appears at most once.

In Figure 26 we show a simple php-tree for n = 3.




Figure 26 - example of a simple php-tree

**Definition 5.14.** Let  $\mathcal{T}$  be a php-tree. The **depth** of  $\mathcal{T}$  is defined as the maximum level of a node of  $\mathcal{T}$ .

 $\mathcal{T}$  is said to be **complete**, iff its depth is n, and every node on the kth level has branching number exactly n - k.

 $\mathcal{T}$  is said to be **symmetric**, iff the label of a node v and the label of an edge connecting v to its child uniquely determines the label of a child, i.e. for any two nodes v and v' with the same label and any two edges e, e' connecting v to its child node w and v' to w', respectively, if the labels of e, e' are the same, then the labels of w and w' are the same, as well.

Notice that the php-tree as in the above picture is not complete nor symmetric.

Symmetric php-trees naturally represent the structure of all plays against  $\mathbf{P}$ 's strategy with  $\mathbf{D}$  utilizing canonical anti-strategies. This is formalized as follows.

**Proposition 5.15.** Let  $n \geq 3$  and F be a functional part of  $\mathbf{P}$ 's strategy. Then, there is an associated symmetric php-tree  $\mathcal{T}_F$  with the property that  $\mathcal{T}_F$  is complete iff for any  $s \geq n+1$  there is no winning canonical anti-strategy for  $\mathbf{D}$  to play against (s, F).

What this means is that during each play against (s, F) there will be a moment where **D** cannot make her move compatibly with the canonical anti-strategy she has chosen before the game.

*Proof.* We construct the tree as follows.

Label the root as  $F(\emptyset)$ .

Let  $v_i$  be a node that we have already constructed together with its label  $l_V(v_i) \in P_n$ . Let  $(v_0, v_1, ..., v_{i-1}, v_i)$  be the sequence of nodes appearing along the unique path starting from the root  $v_0$  and ending in the node  $v_i$ .

For any  $h \in H_n$  such that answering h to a question  $l_V(v_i)$  results in a valid play according to a canonical anti-strategy for  $\mathbf{D}$  assuming  $\mathbf{D}$  has already answered  $l_E(\{v_{j-1}, v_j\})$  to  $l_V(v_{j-1})$  for all  $0 < j \le i$ , we put a new node  $w_h$  into the tree and connect it to  $v_i$  via an edge  $\{v_i, w_h\}$  iff  $F(l_V(v_i), h)$  equals to  $p \in P_n$  which has not yet appeared as a label of any node  $v_j$  for  $j \le i$ . We label the added  $w_h$  as  $F(l_V(v_i), h)$  and the edge  $\{v_i, w_h\}$  as h.

Notice that the tree we are constructing is, indeed, a php-tree. That the resulting labeled tree is symmetric follows from the fact that the label of a child node w of v depends only on the labels of v and  $\{v, w\}$ , since  $l_V(w) = F(l_V(v), l_E\{v, w\})$ .

Finally, the resulting php-tree is not complete iff there is a valid play against (s, F) with **D** using a canonical anti-strategy so that the records produced are  $(p_0, h_0), ..., (p_i, h_i)$  with  $F(p_i, h_i) = p_j$  for  $j \le i$ . As we have already seen in 5.9 this implies that the chosen anti-strategy is a winning one for **D**.

The tree is complete iff all plays against (s, F) where **D** utilizes every possible canonical anti-strategy eventually exhaust  $H_n$  and so such plays cannot be continued using the rules of a canonical anti-strategy.

**Theorem 5.16.** Let  $n \ge 5$ . Assume **D** has winning anti-strategies for n-1 and n-2 against all strategies (s', F') for n-1 and (s'', F'') for n-2, respectively, where  $s', s'' \ge s_0$  for some particular threshold  $s_0$ . Then, **D** has a winning anti-strategy against any (s, F) for s enjoying  $s \ge \max\{s_0 + 1, 2(n-1) + 2\}$ .

*Proof.* Let  $F(\emptyset)$  be n. There are two principal ways how  $\mathbf{D}$  can utilize winning antistrategies for m < n to come up with a winning anti-strategy against (s, F).

• Commit to the root: **D** chooses an answer h for the first **P**'s question  $n \in P_n$  and then never answers h to any subsequent **P**'s questions. If she is guaranteed

not to encounter the question n anymore, then she can use a winning anti-strategy for  $A'_{n-1} = P_n \setminus \{n\}$  and  $B'_{n-1} = H_n \setminus \{h\}$  against s' = s - 1 and F' which equals F restricted to  $A'_{n-1} \times B'_{n-1}$ .

Formally speaking, the above assumption guarantees only that for any  $(p',h') \in A'_{n-1} \times B'_{n-1}$  it holds that  $F'(p',h') = F(p',h') \in B'_{n-1}$  whenever the record (p',h') can actually appear in a play against (s,F) starting with the record (n,h). However, since we care only about admissible records, we may assume F' is properly defined on all the records from  $A'_{n-1} \times B'_{n-1}$  with values in  $B'_{n-1}$ .

Finally, to properly utilize winning anti-strategy for n-1 we need to relabel  $B'_{n-1}$  by elements of  $H_{n-1} = \{0, ..., n-2\}$  via an arbitrary bijection.

• Forbid holes: D chooses different  $h_1, ..., h_i \in H_n$  and plays without ever using those elements, i.e. she never answers  $h_j$  for  $j \leq i$  to any of the **P**'s questions. If she is guaranteed not to encounter questions  $p_1, ..., p_i \in P_n$  with  $p_j$  all different, then she can use a winning anti-strategy for  $A'_{n-i} = P_n \setminus \{p_1, ..., p_i\}$  and  $B'_{n-i} = H_n \setminus \{h_1, ..., h_i\}$  against s' = s and F' which equals F restricted to  $A'_{n-i}$  and  $B'_{n-i}$ . The same remark as for the "commit to the root" rule applies here, as well. Namely we can assume F' is properly defined and need to relabel  $A'_{n-i}$  by elements of  $P_{n-i} = \{0, ..., n-i\}$  and  $B'_{n-i}$  by elements of  $H_{n-i} = \{0, ..., n-i-1\}$ .

We denote the "commit to the root" rule with h chosen as above as CMT<sub>h</sub> and similarly denote the "forbid holes" rule as FH<sub> $h_1,...,h_i$ </sub>.

One can naturally generalize  $CMT_h$  to  $CMT_{h_1,...,h_i}$  rule so as to be able to fallback to m even smaller than n-1 as in  $FH_{h_1,...,h_i}$  case. However, during the proof, we will only need the case with i=1. Thus, we omit the formal definition of  $CMT_{h_1,...,h_i}$  for i>1.

We first demonstrate the usage of  $FH_{h_1,...,h_i}$  rule to get rid of loops.

We say that a pair (p, h) is a **loop** for F iff F(p, h) = p.

**Lemma 5.17.** Assume there is a loop (p, h) for F. Then, there is a winning anti-strategy for  $\mathbf{D}$  against (s, F) with  $s \ge \max\{s_0, 2(n-2) + 2\}$ .

*Proof.* Let **D** apply FH<sub>h</sub> and play without ever giving two contradictory answers in a row. If she is guaranteed not to encounter p, then there is a winning anti-strategy as was shown above against (s, F) with  $s \ge s_0$ .

Otherwise, by answering **P**'s questions without ever using h, she will eventually be asked p for the first time. Since **D** has not yet used h and has never encountered p, the pair (p,h) is consistent with all the answers given by **D** so far and so **D** answers h. The record produced is (p,h) and so **P** will again ask p, since F(p,h) = p. **D** answers h and continues the loop until the number of rounds reaches s and she wins.

It remains to show that in the above case **D** can win against  $s \ge 2(n-2) + 2$ . Notice that **D** wins against s > s' where s' is the length of the shortest locally consistent path

in  $\mathcal{G}_F$  with the last node of the path being p, p never appearing among the nodes except the last one and labels of all the edges of the path being different from h. We will show that  $s' \leq 2(n-2) + 1$ .

Let  $p_0, p_1, ..., p_{s'}$  be nodes of the path as above and  $h_1, ..., h_{s'}$  be labels of the corresponding edges, i.e.  $h_i$  labels the edge going from  $p_{i-1}$  to  $p_i$  for  $i \leq s'$ . Thus,  $p_0$  is the initial node of  $\mathcal{G}_F$  (i.e.  $F(\emptyset) = p_0$ ) and  $p = p_{s'}$ . Notice that for i < s' it holds that  $p_i \neq p$ . Since we are considering the shortest such path, it follows that for  $p_i \neq p_0$  for all i > 0 (assuming the opposite, i.e. that  $p_i = p_0$  for i > 0, one can get a strictly shorter path  $p_i, ..., p_{s'}$  satisfying all the above assumptions).

To finish the proof of  $s' \leq 2(n-2)+1$  it is enough to show that each  $p' \in P_n \setminus \{p_0, p\}$  appears at most twice in the given path.

For the sake of contradiction, let  $p_{i_1} = p_{i_2} = p_{i_3} \in P_n \setminus \{p_0, p\}$  with  $0 < i_1 < i_2 < i_3 < s'$ . Let  $P_1$  denote  $p_0, p_1, ..., p_{i_1}, P_2$  denote  $p_{i_1}, p_{i_i+1}, ..., p_{i_2}, P_3$  denote  $p_{i_2}, p_{i_2+1}, ..., p_{i_3}$  and  $P_4$  denote  $p_{i_3}, p_{i_3+1}, ..., p_{s'}$ . So the path we have can be written as  $P_1 \circ P_2 \circ P_3 \circ P_4$  with  $\circ$  denoting the usual composition of directed walks in a graph.

Consider  $h_{i_1+1}, h_{i_2+1}$  and  $h_{i_3+1}$  - labels of the edges connecting  $p_{i_j}$  to  $p_{i_j+1}$  in the given path. We now show that all such labels must be distinct.

Assume, for example, that  $h_{i_1+1} = h_{i_2+1}$ . Then the path  $P_1 \circ P_3 \circ P_4$  is locally consistent in  $\mathcal{G}_F$  with the last node of the path being p, p never appearing among the nodes except the last one and labels of all the edges of the path being different from h. Note that  $P_1 \circ P_3 \circ P_4$  is strictly shorter than  $P_1 \circ P_2 \circ P_3 \circ P_4$  contradicting the fact that the latter path is the shortest among the ones satisfying the above condition.

Cases  $h_{i_1+1} = h_{i_3+1}$  and  $h_{i_2+1} = h_{i_3+1}$  are done analogously.

Thus,  $h_{i_1+1}, h_{i_2+1}$  and  $h_{i_3+1}$  are all distinct. In particular, either  $h_{i_2+1}$  or  $h_{i_3+1}$  are different from  $h_{i_1}$ . This means either  $P_1 \circ P_3 \circ P_4$  or  $P_1 \circ P_4$  is a locally consistent path in  $\mathcal{G}_F$  with last node of the path being p, p never appearing among the nodes except the last one and labels of all the edges of the path being different from h. Both such paths are strictly shorter than  $P_1 \circ P_2 \circ P_3 \circ P_4$  and so we get a contradiction.

Thus, we can assume there are no loops for F.

We begin the proof of Theorem 5.16 by generating a symmetric php-tree  $\mathcal{T}_F$  as in 5.15 and we assume  $\mathcal{T}_F$  is complete. Notice that  $\mathcal{T}_F$  does not necessarily capture all the information about F, namely, there may be two different functional parts of  $\mathbf{P}$ 's strategies F, F' so that  $\mathcal{T}_F = \mathcal{T}_{F'}$ .

The above discussion leads to a notion of a loose pair for  $\mathcal{T}_F$ . Let  $p \in P_n$  and  $h \in H_n$ . We say that (p, h) is a **loose pair** for  $\mathcal{T}_F$  iff there is no node v together with an edge  $\{v, w\}$  connecting v to its child w so that  $l_V(v) = p$  and  $l_E(\{v, w\}) = h$ .

**Lemma 5.18.** Assume there are no loose pairs for  $\mathcal{T}_F$ . Then, there is a winning antistrategy for **D** against (s, F) for any  $s \geq s_0 + 1$ .

*Proof.* Recall that  $\mathcal{T}_F$  we are considering is complete and its root is labeled by n. Assuming no pair (p,h) is loose, we derive that, for any possible record (p,h), it holds that  $F(p,h) \neq n$ . Indeed, the fact that (p,h) is not loose guarantees that there exists a node v such that  $l_V(v) = F(p,h)$ . Since  $\mathcal{T}_F$  is a php-tree, it follows that v is not a root and so cannot be labeled by n.

Thus, **D** may pick any possible  $h \in H_n$  as an answer to n and apply CMT<sub>h</sub>. Since she is guaranteed not to encounter question n anymore, we can construct a winning anti-strategy for **D** by modifying one for n-1. As we have already seen, this allows **D** to win against any  $s \ge s_0 + 1$ .

Hence, we assume there is a loose pair (p, h) for  $\mathcal{T}_F$ . From this point onward, the proof proceeds in a way similar to the one of 5.11, where we use derivations to construct covers-by-two. The difference is that at this point we have some additional structural information which we can leverage.

We leave these derivations to Appendix 7.3.

We finish this section by stating another open question.

**Definition 5.19.** Let  $n \geq 3$  and  $P_n = \{0, ..., n\}$ ,  $H_n = \{0, ..., n-1\}$ . Let (s, F) be **P**'s strategy and  $\alpha$  be a partial one-to-one mapping from  $P_n$  to  $H_n$ .

We say that **D**'s anti-strategy against (s, F) is **compatible with**  $\alpha$  iff for any question p and any answer h given according to **D**'s anti-strategy the set  $\alpha \cup \{(p, h)\}$  is a partial one-to-one mapping.

Corollary 5.20. Let  $n \geq 3$  and (s, F) be **P**'s strategy for n. Then, there is a partial one-to-one mapping  $\alpha$  from  $P_n$  to  $H_n$  of size n-3 for which there is a winning-antistrategy for **D** against (s, F) which is compatible with  $\alpha$ . Moreover,  $\alpha$  does not depend on s.

Question 5.21. Let  $n \geq 3$  and (s, F) as above. Let  $\alpha$  be an arbitrary one-to-one mapping from  $P_n$  to  $H_n$  of size n-3. Is there a winning anti-strategy for **D** against (s, F) which is compatible with  $\alpha$ ?

# 6 Acknowledgement

The authors deeply thank Jan Krajíček for his suggestions, which greatly improved the presentation of this article. We also would like to thank Ondřej Ježil for his comments on  $\S 5$ , which sharpened our understandings of our toycase. We are also grateful for Neil Thapen for his comments on  $\mathcal{G}_2$  and suggestions on further readings.

The first author (Eitetsu Ken) was supported by JSPS KAKENHI Grant Number 22KJ1121, Grant-in-Aid for JSPS Fellows, and FoPM program at the University of Tokyo.

The second author (Mykyta Narusevych) was supported by the Charles University project PRIMUS/21/SCI/014, Charles University Research Centre program No. UNCE/24/SCI/022 and GA UK project No. 246223

# 7 Appendix

# 7.1 Our formulation of $T_2^2(R)$ -proofs and its propositional translation

Let  $\widetilde{\mathcal{L}}_{BA}$  be the language of bounded arithmetics adopted in [9] to which we add:

- the unary function symbol len(s),
- the binary function symbols  $(s)_i$  and s \* x.

Their interpretations on the standard model is defined as follows.

We encode a finite sequence  $\sigma = (a_1, \ldots, a_l)$  of numbers as follows; Let  $i \in [l]$ . Consider the binary representation  $b_1^{(i)} \cdots b_{k_i}^{(i)}$  of  $a_i$ . Let  $\tilde{a}_i$  be the number having the following binary representation;

$$\widetilde{a}_i = 1b_1^{(i)} \cdots 1b_{k_i}^{(i)}.$$

We encode  $\sigma$  by the number  $\widetilde{\sigma}$ , which is determined by the following binary representation;

$$\widetilde{\sigma} = \widetilde{a}_1 00 \widetilde{a}_2 \cdots 00 \widetilde{a}_l.$$

It is clear that  $\sigma \mapsto \widetilde{\sigma}$  is injective. We adopt the convention that numbers out of the range of  $\widetilde{\cdot}$  code the empty sequence.

Based on this, we define:

- If  $s = \widetilde{\sigma}$  for some  $\sigma = (a_1, \dots, a_l)$ , then len(s) := l,  $(s)_i := a_i$  for  $i \in [len(s)]$  and  $(s)_i := 0$  for  $i \notin [len(s)]$ .  $s * x := (a_1, \dots, a_l, x)$ .
- Otherwise, len(s) = 0,  $(s)_i := 0$ , s \* x := (x).

We consider the theory  $\widetilde{T_2^2}(R)$  of the language  $\widetilde{\mathcal{L}}_{BA} \cup \{R\}$ , where R is a fresh binary predicate symbol; whose axioms are listed below:

- 1. BASIC given in [9].
- 2. len(0) = 0.
- 3. len(s \* x) = len(s) + 1.

- 4.  $len(s) \leq |s|$ .
- 5.  $(s)_i < s$ .
- 6.  $(s*x)_{len(s)+1} = x$ .
- 7.  $1 \le i \land i \le len(s) \to (s * x)_i = (s)_i$ .
- 8.  $s \le s * x$ .
- 9. x < s \* x.
- 10.  $|s * x| \le |s| + 2(|x| + 1) + 2$ .
- 11.  $\widetilde{\Sigma}_{2}^{b}(R)$ -Induction, where  $\widetilde{\Sigma}_{i}^{b}(R)$  and  $\widetilde{\Pi}_{i}^{b}(R)$   $(i \in \mathbb{N})$  are classes of bounded formulae defined as follows:
  - $\widetilde{\Sigma_0^b}(R)$  and  $\widetilde{\Pi_0^b}(R)$  are the class of all sharply bounded formulae (of the extended language  $\widetilde{\mathcal{L}}_{BA} \cup \{R\}$ ).
  - $\widetilde{\Sigma_{i+1}^b}(R)$  is the class collecting bounded formulae of the following form;

$$\exists x \le t. \varphi(x),$$

where  $\varphi \in \widetilde{\Pi_i^b}(R)$ . Similarly for  $\widetilde{\Pi_{i+1}^b}(R)$ .

It is routine to verify the following:

#### **Proposition 7.1.** The following hold:

1. For a formula  $\varphi(x,y)$ ,

$$\widetilde{T_2^2}(R) \vdash (\forall x \le t. \forall y \le u. \ \varphi(x,y)) \leftrightarrow \forall p \le (0*t)*u. \ \varphi((p)_1,(p)_2).$$

2. For any formula of the form  $\forall x \leq |t| . \exists y \leq u. \varphi(x, y)$ , where  $\varphi(x, y) \in \bigcup_{i=0}^{1} \widetilde{\Pi_{i}^{b}}(R)$ , the following holds;

$$\widetilde{T_2^2}(R) \vdash \forall x \leq |t| . \exists y \leq u. \varphi(x,y) \leftrightarrow \exists s \leq t \# 16u^2. \ \varphi(x,(s)_x).$$

3. For any  $\Sigma_2^b(R)$ -formula  $\varphi$  in the sense of [9], there exists  $\widetilde{\Sigma_2^b}(R)$ -formula  $\widetilde{\varphi}$  such that

$$\widetilde{T_2^2}(R) \vdash \varphi \leftrightarrow \widetilde{\varphi}.$$

Hence,  $\widetilde{T_2^2}(R)$  is an extension of  $T_2^2(R)$  of usual formalization. Furthermore, it is also straightforward to see that it is actually a conservative extension of  $T_2^2(R)$ . (The all function symbols we added are representing polynomial time functions, and their basic properties such as the additional axioms above can be proven in  $S_2^1$  under appropriate formalizations of them. See [9].)

Now, one-sided sequent-calculus formulation of  $\widetilde{T_2^2}(R)$  can be considered. We adopt the conventions and formalization given in [4], that is, we only treat formulae of negation normal form and adopt the following derivation rules:

• Initial Sequent;

$$\frac{}{\Gamma, L, \overline{L}}$$
 (where L is a literal)

• V-Rule;

$$\frac{\Gamma, \varphi_{i_0}}{\Gamma}$$
 (where  $\varphi_1 \vee \varphi_2 \in \Gamma$ ,  $i_0 = 1, 2$ )

• ∃-Rule:

$$\frac{\Gamma, \varphi(u)}{\Gamma}$$
 (where  $\exists x. \varphi(x) \in \Gamma$ )

• ∃≤-Rule;

$$\frac{\Gamma, \varphi(u) \qquad \Gamma, u \leq t}{\Gamma} \quad \text{(where } \exists x \leq t. \varphi(x) \in \Gamma, \text{ and } u \text{ is a term)}$$

•  $\land$ -Rule;

$$\frac{\Gamma, \varphi_1 \qquad \Gamma, \varphi_2}{\Gamma} \quad \text{(where } \varphi_1 \land \varphi_2 \in \Gamma\text{)}$$

•  $\forall$ -Rule:

$$\frac{\Gamma, \varphi(a)}{\Gamma} \quad \text{(where } \forall x. \varphi(x) \in \Gamma, \text{ and } a \text{ is an eigenvariable)}$$

•  $\forall \leq$ -Rule;

$$\frac{\Gamma, \overline{a \leq t}, \varphi(a)}{\Gamma} \quad \text{(where } \forall x \leq t. \varphi(x) \in \Gamma, \text{ and } a \text{ is an eigenvariable)}$$

• Axiom of  $\widetilde{T_2^2}(R)$ ;

$$\frac{\Gamma, \overline{\varphi}}{\Gamma}$$
 (where  $\varphi$  is a substitution instance of one of open axioms of  $\widetilde{T_2^2}(R)$ )

•  $\widetilde{\Sigma_2^b}(R)$ -Indunction;

$$\frac{\Gamma, \varphi(0) \qquad \Gamma, \overline{\varphi(a)}, \varphi(a+1) \qquad \Gamma, \overline{\varphi(t)}}{\Gamma} \qquad \text{(where } \varphi \in \widetilde{\Sigma_2^b}(R), \ a \text{ is an eigenvariable)}$$

Furthermore, we consider the following first-order version of  $ontoPHP_n^{n+1}$ :

**Definition 7.2.** Let  $ontoPHP_x^{x+1}(R)$  be the following sequent of  $\widetilde{\mathcal{L}}_{BA}$ -formulae:

$$\{\exists i \leq x + 1. \forall j \leq x. \ \neg R(i, j), \\ \exists i \leq x + 1. \exists i' \leq x + 1. \exists j \leq x. (R(i, j) \land R(i', j) \land i \neq i')), \\ \exists j \leq x. \forall i \leq x + 1. \ \neg R(i, j), \\ \exists j \leq x. \exists j' \leq x. \exists i \leq x + 1. (R(i, j) \land R(i', j) \land j \neq j')) \}.$$

Now, Proposition 2.8 can be proved by a straightforward application of the Paris-Wilkie translation argument. (See [11] for a presentation under a similar convention to ours. For the theoretical background and context, see section 8.2 of [17].)

## 7.2 An equivalent formulation of $G_2$

In this section, we see that the auxiliary information  $\vec{A}$  in the formulation of  $\mathcal{G}_2$  is not essential.

**Definition 7.3.** Let  $n, C \in \mathbb{N}$  and T be an (n, C)-tree.  $\mathcal{G}'_2(n, C, T)$  is the following game: the same as  $\mathcal{G}_2(n, C, T)$  except:

- A possible position is a map L such that:
  - 1.  $dom(L) \subseteq T$ .
  - 2. For  $v \in \text{dom}(L)$ ,  $L(v) \in (\mathcal{M}_n)_{<|n|^C \times height(v)}$ .
  - 3. dom(L) is closed downwards under  $\subseteq$ .
  - 4.  $v \subseteq w \in \text{dom}(L)$ , if M = L(v) and M' = L(w), then they satisfy  $M \subseteq M'$ .

- $L_0 := \emptyset$ .
- When **Prover** plays  $\langle o, x, B \rangle$ , he plays a pair  $\langle o, x \rangle$  instead. The transitions are made in the same way as  $\mathcal{G}_2$ , just disregarding the auxiliary information  $\vec{A}$ ,  $\vec{\alpha}$ , and B.

As  $\mathcal{G}_2$ , the game  $\mathcal{G}_2'$  is also determined. The proof is in exactly the same lines as Corollary 4.7 and we omit it. The notions of **Prover**'s oblivious strategies, **Delayer**'s strategies, and which beats which are defined analogously with Definitions 4.10-4.12, disregarding the auxiliary information  $\vec{A}$ . Then  $\mathcal{G}_2$  and  $\mathcal{G}_2'$  are equivalent in the following sense:

#### **Proposition 7.4.** The following are equivalent:

- 1. There exists C > 0 such that, for sufficiently large n, there exists an (n, C)-tree T such that **Prover** has an oblivious winning strategy for  $\mathcal{G}_2(n, C, T)$ .
- 2. The same condition for  $\mathcal{G}'_2(n, C, T)$ .

*Proof.* (2)  $\Leftarrow$  (1) is easier. Let C and n be as claimed in (2). Then there exists T such that **Prover** has an oblivious winning strategy  $(f'_1, f'_2)$  for  $\mathcal{G}'_2(n, C, T)$ . We observe that it induces **Prover**'s oblivious winning strategy for  $\mathcal{G}_2(n, C, T)$ . Indeed, set:

$$f_1(v, M, \vec{A}) := f'_1(v, M),$$
  
 $f_2(v, M, \vec{A}, M') := \langle o, x, 1 \rangle, \text{ where } \langle o, x \rangle := f'_2(v, M, M').$ 

Then  $(f_1, f_2)$  beats arbitrary **Delayer**'s strategy g for  $\mathcal{G}_2(n, C, T)$  since the play depends only on  $g \upharpoonright_D$ , where

$$D := \{(L, Q) \mid \text{For each } v \in \text{dom}(L), L(v) \text{ is of the form } \langle M, 1, \dots, 1 \rangle \},\$$

and  $(f'_1, f'_2)$  beats **Delayer**'s strategy g' for  $\mathcal{G}'_2$  induced by  $g \upharpoonright_D$ :

$$g'(L,Q) := g(\tilde{L},Q),$$

where  $dom(\tilde{L}) = dom(L)$  and  $\tilde{L}(v) := \langle M, \vec{1} \rangle$  with  $|\vec{1}| = height(v)$ .

Now, we consider the converse. Let C be as claimed in (1). There exists N such that, for all  $n \geq N$ , there exists an (n, C)-tree T such that **Prover** has an oblivious winning strategy for  $\mathcal{G}_2(n, C, T)$ . Let  $C' \geq C$  be a constant and  $N' \geq N$  be large enough such that

$$\forall n \ge N'. \ 2^{|n|^C} (2^{|n|^C} + 1) + 2^{|n|^C} \le 2^{|n|^{C'}}.$$

We show that, for all  $n \geq N'$ , there exists an (n, C')-tree T' such that **Prover** has an oblivious winning strategy  $(f'_1, f'_2)$  for  $\mathcal{G}'_2(n, C', T')$ . Take T for n assured by (1). **Prover** has an oblivious winning strategy  $(f_1, f_2)$  for it. To define T', let

$$[\![k,A]\!] := A(2^{|n|^C} + 1) + k \in [2^{|n|^{C'}}]$$

for  $k, A \in [2^{|n|^C}]$ . Note that  $[\cdot, \cdot]$  is injective on  $[2^{|n|^C}] \times [2^{|n|^C}]$ . Now, define T' as follows:

$$T' := \left\{ (\llbracket v_1, A_1 \rrbracket, \dots, \llbracket v_h, A_h \rrbracket) \in [2^{|n|^{C'}}]^{\leq C} \mid (v_1, \dots, v_h) \in T, \& \forall j \in [h]. \ A_j \in [2^{|n|^C}] \right\}.$$

For each  $v' = ([v_1, A_1], \dots, [v_h, A_h]) \in T'$ , let

$$\vec{A}(v') := (A_1, \dots, A_h).$$

Then the following  $(f'_1, f'_2)$  is the desired oblivious winning strategy of **Prover** for  $\mathcal{G}'_2(n, C', T')$ .

$$f'_{1}(v', M) := f_{1}(v, M, \vec{A}(v'))$$

$$f'_{2}(v', M, M') := \begin{cases} \langle o, [\![x, B]\!] \rangle & \text{(if } o = 1) \\ \langle o, ([\![x_{1}, A_{1}]\!], \dots, [\![x_{h}, A_{h}]\!]) \rangle & \text{(if } o \text{ is } 2 \text{ or } 3, \ x = (x_{i})_{i=1}^{h} \in T, \text{ and } \vec{A}(v') = (A_{i})_{i}.) \end{cases}$$

where  $\langle o, x, B \rangle := f_2(v, M, \vec{A}(v'), M').$ 

#### 7.3 Derivation of labelings

We are given **P**'s strategy (s, F) for  $\mathcal{G}_2(n)$  with  $s \ge \max\{s_0 + 1, 2(n-1) + 2\}$  and we know that there are winning anti-strategies for **D** against any (s', F') for  $\mathcal{G}_2(n-1)$  and (s'', F'') for  $\mathcal{G}_2(n-2)$  with  $s', s'' \ge s_0$ .

We have also argued that the associated php-tree  $\mathcal{T}_F$  is complete, F has no loops and there is a loose pair (p, h) for  $\mathcal{T}_F$ .

Finally, the root of  $\mathcal{T}_F$  is  $n \in P_n$ .

Without loss of generality let (p,h) be  $(0,0) \in P_n \times H_n$ . Let **D** use FH<sub>0</sub>. If  $0 \in P_n$  is guaranteed not to appear as **P**'s question, we construct an anti-strategy for **D** against any  $s \geq s_0$ . So we can assume there is  $p' \in P_n$  and  $h' \in H_n$  with  $p' \neq 0 \in P_n$  and  $h' \neq 0 \in H_n$  such that F(p',h') = 0.

Without loss of generality let (p', h') = (1, 1).

Let **D** use  $FH_{0,1}$ . Similarly as above, we may assume there are  $p'' \neq 0, 1$  and  $h'' \neq 0, 1$  such that either F(p'', h'') = 0 or F(p'', h'') = 1. Here we use the fact that we have winning anti-strategies for **D** for  $\mathcal{G}_2(n-2)$ . Without loss of generality let (p'', h'') = (2, 2).

We start by considering a subtree  $\mathcal{T}'$  of  $\mathcal{T}_F$  formed as follows: start at the root  $n \in P_n$  and proceed without using edges labeled as  $0, 1, 2 \in H_n$ . Since  $\mathcal{T}_F$  is complete, eventually an (n-3)-long branch of  $\mathcal{T}_F$  is built which can also be viewed as a globally consistent path of  $\mathcal{G}_F$  of length n-3. Denote this branch as P. Denote P' and P'' branches of length n-2 with P as their initial segment and last edges labeled by 1 or  $2 \in H_n$ .  $\mathcal{T}'$ 




Figure 27 - representation of  $\mathcal{T}'$ 

is then defined as a collection of all the maximal branches of  $\mathcal{T}_F$  which have P' or P'' as their initial segments.  $\mathcal{T}'$  has a natural tree structure which is illustrated in Figure 27.

Blank circles represent nodes for which we have not yet fixed any particular labeling. The proof proceeds by subsequently deriving more and more information in the form of labeling of nodes. Derivations proceed similarly as in Theorem 5.11.

Without loss of generality, we assume the top-most blank node as above is labeled as  $3 \in P_n$  (notice that such a node cannot be labeled as  $0, 1, 2 \in P_n$  as this would contradict the fact that (0,0) is loose for  $\mathcal{T}_F$ ).

Using the fact that (0,0) is loose we also derive that nodes labeled by  $0 \in P_n$  must appear on a branch of  $\mathcal{T}'$  somewhere below an edge labeled as  $0 \in H_n$ .

We can also derive F(3,2) = 2, otherwise F(3,2) equals 1 and since F(1,1) = 0 we get that (0,0) cannot be loose for  $\mathcal{T}_F$ .

The resulting configuration is shown in Figure 28.

We need to consider two different cases.

1. F(2,0) = 1. We can then label the rest of the blank nodes of  $\mathcal{T}'$  using the fact that  $\mathcal{T}_F$  is a complete symmetric php-tree.

All the derivations and their order are represented by Roman numerals in Figure 29.

Let  $F(0,1) = a \in P_n$ . Based upon the value of a we get the following covers-by-two.




Figure 28




Figure 29 - sequence of derivations of labeling of  $\mathcal{T}'$ 

- $a \ge 3$  is shown in Figure 30.
- a = 2 is shown in Figure 31.
- a = 1 is shown in Figure 32.




Figure 30 - cover-by-two for case  $a \ge 3$ 




Figure 31 - cover-by-two for case a=2




Figure 32 - cover-by-two for case a=1

2. F(2,0) = 0. The resulting labeling of  $\mathcal{T}'$  is illustrated in Figure 33, where we additionally derive that F(0,1) = 1:




Figure 33

At this point, the only possible values for  $a \in P_n$  so that F(3,1) = a are 1 or 2.

(a) a=1. We arrive at Figure 34, where additional labelings and the order of their derivations are represented by Roman numerals as before.





Figure 34







Figure 35 - cover-by-two derived from F(3,1) = 1

This leads to a cover-by-two as is shown in Figure 35.

(b) a = 2. Additional derivations and their order are illustrated in Figure 36.



This leads to a cover-by-two as is shown in Figure 37.

This finishes the case analysis. Notice that all the winning anti-strategies produced by covers-by-two as above allow  $\mathbf{D}$  to win against any possible s.




Figure 37 - cover-by-two for case a=2

### References

- [1] Ajtai, M. (1988). The complexity of the pigeonhole principle, in: *Proc. IEEE 29th Annual Symp. on Foundation of Computer Science*, 346-355.
- [2] Ajtai, M. (1994). The complexity of the Pigeonhole Principle. Combinatorica, 14, 417-433. doi:10.1007/BF01302964
- [3] Ajtai, M. (1994). The independence of the modulo p counting principles, in: *Proc. of the 26th Annual ACM Symposium on Theory of Computing, ACM Press. 2*, 402-411.
- [4] Arai, T. (2020). Ordinal Analysis with an Introduction to Proof Theory. Springer Singapore, Logic in Asia: Studia Logica Library.
- [5] Atserias, A., & Dalmau, V. (2008). A combinatorial characterization of resolution width, *Journal of Computer and System Sciences*, 74(3), 323-334. https://doi.org/10.1016/j.jcss.2007.06.025.
- [6] Atserias, A., & Müller, M. (2015). Partially definable forcing and bounded arithmetic. *Arch. Math. Logic*, 54, no. 1-2, 1-33.
- [7] Beame, P., Impagliazzo, R., Krajíček, J., Pitassi, T. & Pudlak, P. (1994). Lower bounds on Hilbert's Nullstellensatz and propositional proofs, *Proceedings 35th Annual Symposium on Foundations of Computer Science*, 794-806. doi: 10.1109/SFCS.1994.365714.
- [8] Beame, P., & Riis, S. (1998). More on the relative strength of counting principles, in Proof Complexity and Feasible Arithmetics, P. Beame, & S. Buss (Eds.), DIMACS Series in Discrete Mathematics and Theoretical Computer Science, vol.39, American Mathematical Society, Providence, RI, 13-35.
- [9] Buss, S. Bounded arithmetic. (1986). Naples: Bibliopolis, Studies in Proof Theory, Lecture Notes 3.

- [10] Buss, S., Impagliazzo, R., Krajíček, J., Pudlák, P., Razborov, A. A., & Sgall, J. (1996). Proof complexity in algebraic systems and bounded depth Frege systems with modular counting, *Computational Complexity*, 6(3), 256-298.
- [11] Buss, S. Bounded Arithmetic and Constant Depth Frege Proofs, in *Complexity of Computations and Proofs*, J. Krajicek (Ed.), Quaderni di matematica, vol 13, Dipartimento de Matematica della Seconda Universita di Naoli (2004), 153-174.
- [12] Buss, S., Kołodziejczyk, L., & Thapen, N. (2014). Fragments of approximate counting, *The Journal of Symbolic Logic*, 79(2), 496-525.
- [13] Chiari, M., & Krajíček, J. (1998). Witnessing functions in bounded arithmetic and search problems. *Journal of Symbolic Logic*, 63(3), 1095-1115.
- [14] Ken, E. (2022). On some  $\Sigma_0^B$ -generalizations of the pigeonhole and the modular counting principles over  $V^0$ , arXiv:2203.10237v6 (preprint).
- [15] Krajíček, J. (1995). Bounded arithmetic, propositional logic, and complexity theory. New York, NY: Cambridge University Press, Encyclopedia of Mathematics and Its Applications 60.
- [16] Krajíček, J. (2006). Forcing with Random Variables and Proof Complexity. In Logical Approaches to Computational Barriers, Beckmann, A., Berger, U., Löwe, B., Tucker, J.V. (eds), CiE 2006. Lecture Notes in Computer Science, vol 3988. Springer, Berlin, Heidelberg.
- [17] Krajíček, J. (2019). *Proof complexity*. Cambridge, UK: Cambridge University Press, Encyclopedia of Mathematics and Its Applications 170.
- [18] Krajíček, J., Skelley, A., & Thapen, N. (2007). NP search problems in low fragments of bounded arithmetic, J. Symbolic Logic, 72(2), 649-672.
- [19] Krajíček, J., Pudlák, P., & Woods, A. (1995). An exponential lower bound to the size of bounded depth Frege proofs of the pigeonhole principle, *Random Structures and Algorithms*, 7(1), 15-39. doi:10.1002/rsa.3240070103
- [20] Narusevych, M. (2022). Models of Bounded Arithmetic, ECCC.
- [21] Paris, J., & Wilkie, A. (1985). Counting problems in bounded arithmetic. In: Di Prisco, C.A. (eds), *Methods in Mathematical Logic*. Lecture Notes in Mathematics, vol 1130. Springer, Berlin, Heidelberg.
- [22] Pitassi, T., Beame, P., & Impagliazzo, R. (1993). Exponential lower bounds for the pigeonhole principle, *Computational Complexity*, 3(2), 97-140. doi:10.1145/129712.129733

- [23] Pudlák, P. Lengths of proofs, in *Handbook of Proof Theory*, S. Buss (Ed.), Studies in Logic and the Foundations of Mathematics 137, Elsevier, Amsterdam (1998), 548-637.
- [24] Pudlák, P. (2013). Logical foundations of mathematics and computational complexity: a gentle introduction, Cham: Springer, Springer Monographs in Mathematics.
- [25] Riis, S. (2001). A complexity gap for tree resolution. Comput. complex, 10, 179-209.