

The algebraic and geometric classification of compatible Pre-Lie algebras^{*†}

Hani Abdelwahab[‡], Ivan Kaygorodov[§] & Abdenacer Makhlouf[¶]

Abstract: *In this paper, we develop a method to obtain the algebraic classification of compatible pre-Lie algebras from the classification of pre-Lie algebras of the same dimension. We use this method to obtain the algebraic classification of complex 2-dimensional compatible pre-Lie algebras. As a byproduct, we obtain the classification of complex 2-dimensional compatible commutative associative, compatible associative and compatible Novikov algebras. In addition, we consider the geometric classification of varieties of cited algebras, that is the description of its irreducible components.*

Keywords: *compatible algebra, compatible associative algebra, compatible pre-Lie algebra, algebraic classification, geometric classification.*

MSC2020: 17A30 (primary); 17D25, 14L30 (secondary).

CONTENTS

Introduction	2
1. The algebraic classification of compatible algebras	3
1.1. The algebraic classification of algebras	3
1.2. 2-dimensional pre-Lie algebras	5
1.3. The algebraic classification of compatible pre-Lie algebras	5
1.4. The algebraic classification of compatible commutative associative algebras	15
1.5. The algebraic classification of compatible associative algebras	16
1.6. The algebraic classification of compatible Novikov algebras	16
2. The geometric classification of compatible algebras	17
2.1. Degenerations and the geometric classification of algebras	17
2.2. The geometric classification of compatible commutative associative algebras	19
2.3. The geometric classification of compatible associative algebras	20
2.4. The geometric classification of compatible Novikov algebras	20
2.5. The geometric classification of compatible pre-Lie algebras	21
References	22

^{*}The authors thank Amir Fernández Ouaridi for sharing some useful software programs.

[†]The work is supported by FCT UIDB/00212/2020, UIDP/00212/2020 and 2022.14950.CBM (Programa PESSOA).

[‡]Department of Mathematics, Faculty of Science, Mansoura University, Mansoura, Egypt; hania-mar1985@gmail.com

[§]CMA-UBI, Universidade da Beira Interior, Covilhã, Portugal; kaygorodov.ivan@gmail.com

[¶]IRIMAS - Département de Mathématiques, University of Haute Alsace, Mulhouse, France; abdenacer.makhlouf@uha.fr

INTRODUCTION

The algebraic classification (up to isomorphism) of algebras of small dimensions from a certain variety defined by a family of polynomial identities is a classic problem in the theory of non-associative algebras. Another interesting approach to studying algebras of a fixed dimension is to study them from a geometric point of view (that is, to study the degenerations and deformations of these algebras). The results in which the complete information about degenerations of a certain variety is obtained are generally referred to as the geometric classification of the algebras of these varieties. There are many results related to the algebraic and geometric classification of Jordan, Lie, Leibniz, Zinbiel, and other algebras (see, [1–7, 16–18, 23, 25, 27, 28] and references in [24, 26, 32]).

Pre-Lie algebras, also known as right symmetric algebras, appeared in some papers by Gerstenhaber, Koszul, and Vinberg in the 1960s. It is a generalization of associative algebras and the most popular non-associative subvariety of Lie-admissible algebras. They have various applications in geometry and physics [8]; recently some connections between them and trees, braces and F -manifold algebras were established in [9, 30, 37, 38]. Novikov algebras are pre-Lie algebras with an additional identity. They were introduced in papers by Gel'fand and Dorfman, Balinskii and Novikov (about Novikov algebras see [15] and references therein). Let Ω be a variety of algebras. We say that an algebra $(\mathbf{A}, \cdot, *)$ is a compatible Ω -algebra, if and only if (\mathbf{A}, \cdot) , $(\mathbf{A}, *)$ and $(\mathbf{A}, \cdot + *)$ are Ω -algebras. Compatible Lie algebras are considered in the study of the classical Yang-Baxter equation [21], integrable equations of the principal chiral model type [20], elliptic theta functions [33], and other areas. The study of non-Lie compatible algebras is also very popular. So, cohomology and deformations were studied for compatible Lie algebras [31], compatible associative algebras [10], compatible Hom-Lie algebras [12], compatible L_∞ -algebras [11], compatible 3-Lie algebras [19], compatible dendriform algebras [13], and so on. Free compatible algebras were studied in the associative algebra case in [14] and in the Lie algebra case in [22, 29]. Compatible associative structures on matrix algebras were studied in [34–36]. General compatible structures have been studied from an operadic point of view in [39, 40]. At this moment there is only one paper about the classification of small dimensional compatible algebras. Namely, all 4-dimensional nilpotent compatible Lie algebras were classified in [28].

The main goal of the present paper is to obtain the algebraic and geometric description of the variety of complex 2-dimensional compatible pre-Lie algebras. To do so, we first determine all such 2-dimensional algebra structures, up to isomorphism (what we call the algebraic classification), and then proceed to determine the geometric properties of the corresponding variety, namely its dimension and description of the irreducible components (the geometric classification). As some corollaries, we have the algebraic and geometric classification of complex 2-dimensional compatible commutative associative, compatible associative and compatible Novikov algebras.

Our main results regarding the algebraic classification are summarized below.

Theorem A1. *There are infinitely many isomorphism classes of complex 2-dimensional compatible pre-Lie algebras, described explicitly in Theorem 7 in terms of 6 three-parameter families, 14 two-parameter families, 13 one-parameter families, and 8 additional isomorphism classes.*

Theorem A2. *There are infinitely many isomorphism classes of complex 2-dimensional compatible commutative associative algebras, described explicitly in Theorem 8 in terms of 1 three-parameter family, 3 two-parameter families, 8 one-parameter families, and 6 additional isomorphism classes.*

Theorem A3. *There are infinitely many isomorphism classes of complex 2-dimensional compatible associative algebras, described explicitly in Theorem 10 in terms of 1 three-parameter family, 3 two-parameter families, 10 one-parameter families, and 10 additional isomorphism classes.*

Theorem A4. *There are infinitely many isomorphism classes of complex 2-dimensional compatible Novikov algebras, described explicitly in Theorem 12 in terms of 3 three-parameter family, 7 two-parameter families, 11 one-parameter families, and 7 additional isomorphism classes.*

The geometric part of our work aims to generalize previously obtained results about the geometric classification of 2-dimensional Novikov [7] and pre-Lie [6] algebras. Our main results regarding the geometric classification are summarized below.

Theorem G1. *The variety of complex 2-dimensional compatible pre-Lie algebras has dimension 7. It is defined by 2 rigid algebras, 1 one-parametric family of algebras, 7 two-parametric family of algebras, and 4 three-parametric families of algebras and can be described as the closure of the union of $\mathrm{GL}_2(\mathbb{C})$ -orbits of the algebras given in Theorem 22.*

Theorem G2. *The variety of complex 2-dimensional compatible commutative associative algebras has dimension 7. It is defined by 1 two-parametric family of algebras and 1 three-parametric families of algebras and can be described as the closure of the union of $\mathrm{GL}_2(\mathbb{C})$ -orbits of the algebras given in Theorem 19.*

Theorem G3. *The variety of complex 2-dimensional compatible associative algebras has dimension 7. It is defined by 2 rigid algebras, 2 two-parametric family of algebras, and 1 three-parametric families of algebras and can be described as the closure of the union of $\mathrm{GL}_2(\mathbb{C})$ -orbits of the algebras given in Theorem 20.*

Theorem G4. *The variety of complex 2-dimensional compatible Novikov algebras has dimension 7. It is defined by 1 rigid algebra, 3 two-parametric family of algebras, and 1 three-parametric families of algebras and can be described as the closure of the union of $\mathrm{GL}_2(\mathbb{C})$ -orbits of the algebras given in Theorem 21.*

1. THE ALGEBRAIC CLASSIFICATION OF COMPATIBLE ALGEBRAS

All the algebras below will be over \mathbb{C} and all the linear maps will be \mathbb{C} -linear. For simplicity, every time we write the multiplication table of an algebra the products of basic elements whose values are zero are omitted.

1.1. The algebraic classification of algebras. In this paper, we work with compatible pre-Lie algebras with two multiplications. Let us review the method we will use to obtain the algebraic classification for the variety of compatible pre-Lie algebras (the present method, in the case of Poisson algebras, is given with more details in [2, 3]).

Definition 1. *An algebra is called a pre-Lie algebra if it satisfies the identity*

$$(x \cdot y) \cdot z - x \cdot (y \cdot z) = (y \cdot x) \cdot z - y \cdot (x \cdot z).$$

Definition 2. A compatible pre-Lie algebra is a vector space \mathbf{A} equipped with two multiplication \cdot and another multiplication $*$, such that, (\mathbf{A}, \cdot) , $(\mathbf{A}, *)$ and $(\mathbf{A}, \cdot + *)$ are pre-Lie algebras. These two operations are required to satisfy the following identities.

$$\begin{aligned} (x \cdot y) \cdot z - x \cdot (y \cdot z) &= (y \cdot x) \cdot z - y \cdot (x \cdot z), \\ (x * y) * z - x * (y * z) &= (y * x) * z - y * (x * z), \\ (x * y) \cdot z - x * (y \cdot z) + (x \cdot y) * z - x \cdot (y * z) &= (y * x) \cdot z - y * (x \cdot z) + (y \cdot x) * z - y \cdot (x * z). \end{aligned}$$

The main examples of compatible pre-Lie algebras are the following: compatible commutative associative, compatible associative and compatible Novikov algebras.

Definition 3. Let (\mathbf{A}, \cdot) be a pre-Lie algebra. Define $Z^2(\mathbf{A}, \mathbf{A})$ to be the set of all bilinear maps $\theta : \mathbf{A} \times \mathbf{A} \rightarrow \mathbf{A}$ such that:

$$\begin{aligned} \theta(\theta(x, y), z) - \theta(x, \theta(y, z)) &= \theta(\theta(y, x), z) - \theta(y, \theta(x, z)), \\ \theta(x, y) \cdot z - \theta(x, y \cdot z) + \theta(x \cdot y, z) - x \cdot \theta(y, z) &= \theta(y, x) \cdot z - \theta(y, x \cdot z) + \theta(y \cdot x, z) - y \cdot \theta(x, z). \end{aligned}$$

Then $Z^2(\mathbf{A}, \mathbf{A}) \neq \emptyset$ since $\theta = 0 \in Z^2(\mathbf{A}, \mathbf{A})$.

Now, for $\theta \in Z^2(\mathbf{A}, \mathbf{A})$, let us define a multiplication \bullet_θ on \mathbf{A} by $x \bullet_\theta y = \theta(x, y)$ for all $x, y \in \mathbf{A}$. Then $(\mathbf{A}, \cdot, \bullet_\theta)$ is a compatible pre-Lie algebra. Conversely, if $(\mathbf{A}, \cdot, \bullet)$ is a compatible pre-Lie algebra, then there exists $\theta \in Z^2(\mathbf{A}, \mathbf{A})$ such that $(\mathbf{A}, \cdot, \bullet_\theta) \cong (\mathbf{A}, \cdot, \bullet)$. To see this, consider the bilinear map $\theta : \mathbf{A} \times \mathbf{A} \rightarrow \mathbf{A}$ defined by $\theta(x, y) = x \bullet y$ for all x, y in \mathbf{A} . Then $\theta \in Z^2(\mathbf{A}, \mathbf{A})$ and $(\mathbf{A}, \cdot, \bullet_\theta) = (\mathbf{A}, \cdot, \bullet)$.

Let (\mathbf{A}, \cdot) be a pre-Lie algebra. The automorphism group, $\text{Aut}(\mathbf{A})$, of \mathbf{A} acts on $Z^2(\mathbf{A}, \mathbf{A})$ by

$$(\theta * \phi)(x, y) = \phi^{-1}(\theta(\phi(x), \phi(y))),$$

for $\phi \in \text{Aut}(\mathbf{A})$, and $\theta \in Z^2(\mathbf{A}, \mathbf{A})$.

Lemma 4. Let (\mathbf{A}, \cdot) be a pre-Lie algebra and $\theta, \vartheta \in Z^2(\mathbf{A}, \mathbf{A})$. Then $(\mathbf{A}, \cdot, \bullet_\theta)$ and $(\mathbf{A}, \cdot, \bullet_\vartheta)$ are isomorphic if and only if there is a $\phi \in \text{Aut}(\mathbf{A})$ with $\theta * \phi = \vartheta$.

Proof. If $\theta * \phi = \vartheta$, then $\phi : (\mathbf{A}, \cdot, \bullet_\vartheta) \rightarrow (\mathbf{A}, \cdot, \bullet_\theta)$ is an isomorphism since $\phi(\vartheta(x, y)) = \theta(\phi(x), \phi(y))$. On the other hand, if $\phi : (\mathbf{A}, \cdot, \bullet_\vartheta) \rightarrow (\mathbf{A}, \cdot, \bullet_\theta)$ is an isomorphism of compatible pre-Lie algebras, then $\phi \in \text{Aut}(\mathbf{A})$ and $\phi(x \bullet_\vartheta y) = \phi(x) \bullet_\theta \phi(y)$. Hence

$$\vartheta(x, y) = \phi^{-1}(\theta(\phi(x), \phi(y))) = (\theta * \phi)(x, y)$$

and therefore $\theta * \phi = \vartheta$. □

Consequently, we have a procedure to classify the compatible pre-Lie algebras with given associated pre-Lie algebra (\mathbf{A}, \cdot) . It consists of three steps:

- (1) Compute $Z^2(\mathbf{A}, \mathbf{A})$.
- (2) Find the orbits of $\text{Aut}(\mathbf{A})$ on $Z^2(\mathbf{A}, \mathbf{A})$.
- (3) Choose a representative θ from each orbit and then construct the compatible pre-Lie algebra $(\mathbf{A}, \cdot, \bullet_\theta)$.

1.2. 2-dimensional pre-Lie algebras.

Lemma 5. *Let \mathcal{C} be a nonzero 2-dimensional pre-Lie algebra. Then \mathcal{C} is isomorphic to one and only one of the following algebras:*

$$\begin{aligned}
\mathcal{C}_{01} & : e_1 \cdot e_1 = e_1 + e_2, \quad e_2 \cdot e_1 = e_2. \\
\mathcal{C}_{02} & : e_1 \cdot e_1 = e_1 + e_2, \quad e_1 \cdot e_2 = e_2. \\
\mathcal{C}_{03} & : e_1 \cdot e_1 = e_2. \\
\mathcal{C}_{04} & : e_2 \cdot e_1 = e_1. \\
\mathcal{C}_{05}^\alpha & : e_1 \cdot e_1 = e_1, \quad e_1 \cdot e_2 = \alpha e_2. \\
\mathcal{C}_{06}^\alpha & : e_1 \cdot e_1 = e_1, \quad e_1 \cdot e_2 = \alpha e_2, \quad e_2 \cdot e_1 = e_2. \\
\mathcal{C}_{07} & : e_1 \cdot e_1 = e_1, \quad e_2 \cdot e_2 = e_2. \\
\mathcal{C}_{08} & : e_1 \cdot e_1 = e_1, \quad e_1 \cdot e_2 = 2e_2, \quad e_2 \cdot e_1 = \frac{1}{2}e_1 + e_2, \quad e_2 \cdot e_2 = e_2.
\end{aligned}$$

Lemma 6. *The description of the group of automorphisms of every 2-dimensional pre-Lie algebra is given below.*

- (1) *If $\phi \in \text{Aut}(\mathcal{C}_{01})$, then $\phi(e_1) = e_1 + ve_2$ and $\phi(e_2) = e_2$, for $v \in \mathbb{C}$.*
- (2) *If $\phi \in \text{Aut}(\mathcal{C}_{02})$, then $\phi(e_1) = e_1 + ve_2$ and $\phi(e_2) = e_2$, for $v \in \mathbb{C}$.*
- (3) *If $\phi \in \text{Aut}(\mathcal{C}_{03})$, then $\phi(e_1) = \xi e_1 + ve_2$ and $\phi(e_2) = \xi^2 e_2$, for $\xi \in \mathbb{C}^*$ and $v \in \mathbb{C}$.*
- (4) *If $\phi \in \text{Aut}(\mathcal{C}_{04})$, then $\phi(e_1) = \xi e_1$ and $\phi(e_2) = e_2$, for $\xi \in \mathbb{C}^*$.*
- (5) *If $\phi \in \text{Aut}(\mathcal{C}_{05}^{\alpha \neq 1})$, then $\phi(e_1) = e_1$ and $\phi(e_2) = \xi e_2$, for $\xi \in \mathbb{C}^*$.*
- (6) *If $\phi \in \text{Aut}(\mathcal{C}_{05}^1)$, then $\phi(e_1) = e_1 + ve_2$ and $\phi(e_2) = \xi e_2$, for $\xi \in \mathbb{C}^*$ and $v \in \mathbb{C}$.*
- (7) *If $\phi \in \text{Aut}(\mathcal{C}_{06}^{\alpha \neq 0})$, then $\phi(e_1) = e_1$ and $\phi(e_2) = \xi e_2$, for $\xi \in \mathbb{C}^*$.*
- (8) *If $\phi \in \text{Aut}(\mathcal{C}_{06}^0)$, then $\phi(e_1) = e_1 + ve_2$ and $\phi(e_2) = \xi e_2$, for $\xi \in \mathbb{C}^*$ and $v \in \mathbb{C}$.*
- (9) *If $\phi \in \text{Aut}(\mathcal{C}_{07})$, then $\phi \in \mathbb{S}_2$, i.e. $\phi(e_1) = e_1, \phi(e_2) = e_2$ or $\phi(e_1) = e_2, \phi(e_2) = e_1$.*
- (10) *If $\phi \in \text{Aut}(\mathcal{C}_{08})$, then $\phi(e_1) = e_1, \phi(e_2) = e_2$ or $\phi(e_1) = -e_1 + 4e_2, \phi(e_2) = e_2$.*

1.3. The algebraic classification of compatible pre-Lie algebras. The main aim of the present section is to prove the following results.

Theorem 7. *Let \mathcal{C} be a nonzero 2-dimensional compatible pre-Lie algebra. Then \mathcal{C} is isomorphic to one and only one of the following algebras:*

$$\begin{aligned}
\mathcal{C}_{01} & : e_1 * e_1 = e_1 + e_2, \quad e_2 * e_1 = e_2. \\
\mathcal{C}_{02} & : e_1 * e_1 = e_1 + e_2, \quad e_1 * e_2 = e_2. \\
\mathcal{C}_{03} & : e_1 * e_1 = e_2. \\
\mathcal{C}_{04} & : e_2 * e_1 = e_1. \\
\mathcal{C}_{05}^\alpha & : e_1 * e_1 = e_1, \quad e_1 * e_2 = \alpha e_2. \\
\mathcal{C}_{06}^\alpha & : e_1 * e_1 = e_1, \quad e_1 * e_2 = \alpha e_2, \quad e_2 * e_1 = e_2. \\
\mathcal{C}_{07} & : e_1 * e_1 = e_1, \quad e_2 * e_2 = e_2. \\
\mathcal{C}_{08} & : e_1 * e_1 = e_1, \quad e_1 * e_2 = 2e_2, \quad e_2 * e_1 = \frac{1}{2}e_1 + e_2, \quad e_2 * e_2 = e_2. \\
\mathcal{C}_{09}^{\alpha, \beta \neq 0} & : \begin{cases} e_1 \cdot e_1 = e_1 + e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = \beta e_2, \quad e_2 * e_1 = \alpha e_2. \end{cases} \\
\mathcal{C}_{10}^{\alpha, \beta} & : \begin{cases} e_1 \cdot e_1 = e_1 + e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1 + \beta e_2, & e_2 * e_1 = \alpha e_2. \end{cases} \\
\mathcal{C}_{11}^{\alpha, \beta \neq \alpha} & : \begin{cases} e_1 \cdot e_1 = e_1 + e_2, & e_1 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = \beta e_2. \end{cases}
\end{aligned}$$

$$\begin{aligned}
\mathcal{C}_{12}^{\alpha,\beta} &: \begin{cases} e_1 \cdot e_1 = e_1 + e_2, & e_1 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha e_1 + \beta e_2, & e_1 * e_2 = \alpha e_2. \end{cases} \\
\mathcal{C}_{13}^{\alpha} &: \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = e_2, & e_2 * e_1 = \alpha e_2. \end{cases} \\
\mathcal{C}_{14}^{\alpha} &: \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = e_1 + \alpha e_2, & e_2 * e_1 = e_2. \end{cases} \\
\mathcal{C}_{15}^{\alpha} &: \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_2. \end{cases} \\
\mathcal{C}_{16}^{\alpha} &: \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = e_1, & e_1 * e_2 = \alpha e_2. \end{cases} \\
\mathcal{C}_{17}^{\alpha} &: \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = e_1 + \alpha e_2, & e_1 * e_2 = e_2. \end{cases} \\
\mathcal{C}_{18}^{\alpha} &: \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_2, & e_1 * e_2 = e_1, & e_2 * e_1 = e_1, & e_2 * e_2 = e_2. \end{cases} \\
\mathcal{C}_{19}^{\alpha} &: \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_2, & e_2 * e_1 = e_1, & e_2 * e_2 = 2e_2. \end{cases} \\
\mathcal{C}_{20}^{\alpha,\beta} &: \begin{cases} e_2 \cdot e_1 = e_1, \\ e_2 * e_1 = \alpha e_1, & e_2 * e_2 = e_1 + \beta e_2. \end{cases} \\
\mathcal{C}_{21}^{\alpha,\beta} &: \begin{cases} e_2 \cdot e_1 = e_1, \\ e_2 * e_1 = \alpha e_1, & e_2 * e_2 = \beta e_2. \end{cases} \\
\mathcal{C}_{22}^{\alpha \neq 0, \beta} &: \begin{cases} e_2 \cdot e_1 = e_1, \\ e_1 * e_2 = \alpha e_1, & e_2 * e_1 = \beta e_1, & e_2 * e_2 = e_1 + \alpha e_2. \end{cases} \\
\mathcal{C}_{23}^{\alpha \neq 0, \beta} &: \begin{cases} e_2 \cdot e_1 = e_1, \\ e_1 * e_2 = \alpha e_1, & e_2 * e_1 = \beta e_1, & e_2 * e_2 = \alpha e_2. \end{cases} \\
\mathcal{C}_{24}^{\alpha,\beta,\gamma} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \alpha e_2, \\ e_1 * e_1 = \beta e_1 + e_2, & e_1 * e_2 = \gamma e_2. \end{cases} \\
\mathcal{C}_{25}^{\alpha,\beta,\gamma} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \alpha e_2, \\ e_1 * e_1 = \beta e_1, & e_1 * e_2 = \gamma e_2. \end{cases} \\
\mathcal{C}_{26}^{\alpha} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \frac{1}{2}e_2, \\ e_1 * e_1 = 2\alpha e_1, & e_1 * e_2 = \alpha e_2, & e_2 * e_2 = e_1. \end{cases} \\
\mathcal{C}_{27}^{\alpha,\beta} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \frac{1}{2}e_2, \\ e_1 * e_1 = 2\alpha e_1, & e_1 * e_2 = e_1 + \alpha e_2, & e_2 * e_1 = 2e_1 & e_2 * e_2 = \beta e_1 + e_2. \end{cases} \\
\mathcal{C}_{28} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = e_2, \\ e_2 * e_1 = e_1, & e_2 * e_2 = e_2. \end{cases} \\
\mathcal{C}_{29}^{\alpha} &: \begin{cases} e_1 \cdot e_1 = e_1, \\ e_1 * e_1 = \alpha e_1, & e_2 * e_2 = e_2. \end{cases} \\
\mathcal{C}_{30}^{\alpha,\beta} &: \begin{cases} e_1 \cdot e_1 = e_1, \\ e_1 * e_1 = \alpha e_1 + \beta e_2, & e_1 * e_2 = e_1, & e_2 * e_1 = e_1, & e_2 * e_2 = e_2. \end{cases} \\
\mathcal{C}_{31}^{\alpha,\beta,\gamma} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \alpha e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \beta e_1 + e_2, & e_1 * e_2 = \gamma e_2, & e_2 * e_1 = \beta e_2. \end{cases} \\
\mathcal{C}_{32}^{\alpha,\beta,\gamma} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \alpha e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \beta e_1, & e_1 * e_2 = \gamma e_2, & e_2 * e_1 = \beta e_2. \end{cases} \\
\mathcal{C}_{33} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_2 \cdot e_1 = e_2, \\ e_1 * e_2 = e_1, & e_2 * e_2 = e_2. \end{cases} \\
\mathcal{C}_{34}^{\alpha,\beta} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = \alpha e_2, & e_2 * e_1 = \alpha e_2, & e_2 * e_2 = e_1 + \beta e_2. \end{cases}
\end{aligned}$$

$$\begin{aligned}
\mathcal{C}_{35}^{\alpha} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = \alpha e_2, & e_2 * e_1 = \alpha e_2, & e_2 * e_2 = e_2. \end{cases} \\
\mathcal{C}_{36}^{\alpha, \beta} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = 2e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1 + \beta e_2, & e_1 * e_2 = 2\alpha e_2, & e_2 * e_1 = e_1 + \alpha e_2, & e_2 * e_2 = 2e_2. \end{cases} \\
\mathcal{C}_{37}^{\alpha} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = 2e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = e_1 + 2\alpha e_2, & e_2 * e_1 = 2e_1 + \alpha e_2, & e_2 * e_2 = e_2. \end{cases} \\
\mathcal{C}_{38}^{\alpha, \beta, \gamma} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = (\gamma + \beta - \alpha)e_1 - \beta e_2, & e_1 * e_2 = \alpha e_1 + \beta e_2, \\ e_2 * e_1 = \alpha e_1 + \beta e_2, & e_2 * e_2 = -\alpha e_1 + \gamma e_2. \end{cases} \\
\mathcal{C}_{39}^{\alpha, \beta \neq \alpha} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_2 * e_2 = \beta e_2. \end{cases} \\
\mathcal{C}_{40}^{\alpha, \beta, \gamma} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = 2e_2, & e_2 \cdot e_1 = \frac{1}{2}e_1 + e_2, & e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha e_1 + \beta e_2, & e_1 * e_2 = 2\alpha e_2, & e_2 * e_1 = \gamma e_1 + \alpha e_2, & e_2 * e_2 = 2\gamma e_2. \end{cases} \\
\mathcal{C}_{41}^{\alpha, \beta \neq 0} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = 2e_2, \\ e_2 \cdot e_1 = \frac{1}{2}e_1 + e_2, & e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = \beta e_1 + 2\alpha e_2, \\ e_2 * e_1 = (2\beta + \frac{1}{2}\alpha)e_1 + \alpha e_2, & e_2 * e_2 = \frac{1}{2}\beta e_1 + (\alpha + \beta)e_2. \end{cases}
\end{aligned}$$

All algebras are non-isomorphic, except

$$\begin{aligned}
\mathcal{C}_{24}^{1, \beta, \gamma} &\cong \mathcal{C}_{25}^{1, \beta, \gamma}, & \mathcal{C}_{31}^{0, \beta, \gamma} &\cong \mathcal{C}_{32}^{0, \beta, \gamma}, & \mathcal{C}_{38}^{\alpha, \beta, \gamma} &\cong \mathcal{C}_{38}^{\beta, \alpha, -\alpha + \beta + \gamma}, \\
\mathcal{C}_{39}^{\alpha, \beta} &\cong \mathcal{C}_{39}^{\beta, \alpha}, & \mathcal{C}_{40}^{\alpha, \beta, \gamma} &\cong \mathcal{C}_{40}^{4\gamma - \alpha, \beta - 8\alpha + 16\gamma, \gamma}, & \mathcal{C}_{41}^{\alpha, \beta} &\cong \mathcal{C}_{41}^{\alpha + 4\beta, -\beta}.
\end{aligned}$$

1.3.1. *Compatible pre-Lie algebras defined on \mathcal{C}_{01} .* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{01}, \mathcal{C}_{01})$, the compatible pre-Lie algebra structures defined on \mathcal{C}_{01} are of the form:

$$\begin{cases} e_1 \cdot e_1 = e_1 + e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha_1 e_1 + \alpha_2 e_2, & e_1 * e_2 = \alpha_3 e_2, & e_2 * e_1 = \alpha_1 e_2. \end{cases}$$

Then we have the following cases:

- If $\alpha_3 \neq 0$, then choose $\nu = -\frac{\alpha_2}{\alpha_3}$ and obtain the parametric family:

$$\mathcal{C}_{09}^{\alpha, \beta \neq 0} := \begin{cases} e_1 \cdot e_1 = e_1 + e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = \beta e_2, & e_2 * e_1 = \alpha e_2. \end{cases}$$

The algebras $\mathcal{C}_{09}^{\alpha, \beta}$ and $\mathcal{C}_{09}^{\alpha', \beta'}$ are isomorphic if and only if $(\alpha, \beta) = (\alpha', \beta')$.

- If $\alpha_3 = 0$, we obtain the parametric family:

$$\mathcal{C}_{10}^{\alpha, \beta} := \begin{cases} e_1 \cdot e_1 = e_1 + e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1 + \beta e_2, & e_2 * e_1 = \alpha e_2. \end{cases}$$

The algebras $\mathcal{C}_{10}^{\alpha, \beta}$ and $\mathcal{C}_{10}^{\alpha', \beta'}$ are isomorphic if and only if $(\alpha, \beta) = (\alpha', \beta')$.

1.3.2. *Compatible pre-Lie algebras defined on \mathcal{C}_{02} .* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{02}, \mathcal{C}_{02})$, the compatible pre-Lie algebra structures defined on \mathcal{C}_{02} are of the form:

$$\begin{cases} e_1 \cdot e_1 = e_1 + e_2, & e_1 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha_1 e_1 + \alpha_2 e_2, & e_1 * e_2 = \alpha_3 e_2. \end{cases}$$

Then we have the following cases:

- If $\alpha_1 \neq \alpha_3$, then choose $\nu = \frac{\alpha_2}{\alpha_1 - \alpha_3}$ and obtain the parametric family:

$$\mathcal{C}_{11}^{\alpha, \beta \neq \alpha} := \begin{cases} e_1 \cdot e_1 = e_1 + e_2, & e_1 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = \beta e_2. \end{cases}$$

The algebras $\mathcal{C}_{11}^{\alpha, \beta}$ and $\mathcal{C}_{11}^{\alpha', \beta'}$ are isomorphic if and only if $(\alpha, \beta) = (\alpha', \beta')$.

- If $\alpha_1 = \alpha_3$, we obtain the parametric family:

$$\mathcal{C}_{12}^{\alpha, \beta} := \begin{cases} e_1 \cdot e_1 = e_1 + e_2, & e_1 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha e_1 + \beta e_2, & e_1 * e_2 = \alpha e_2. \end{cases}$$

The algebras $\mathcal{C}_{12}^{\alpha, \beta}$ and $\mathcal{C}_{12}^{\alpha', \beta'}$ are isomorphic if and only if $(\alpha, \beta) = (\alpha', \beta')$.

1.3.3. *Compatible pre-Lie algebras defined on \mathcal{C}_{03} .* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{03}, \mathcal{C}_{03})$, the compatible pre-Lie algebra structures defined on \mathcal{C}_{03} are of the forms:

- (1) $\begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha_1 e_1 + \alpha_2 e_2, & e_1 * e_2 = \alpha_3 e_2, & e_2 * e_1 = \alpha_1 e_2. \end{cases}$
- (2) $\begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha_4 e_1 + \alpha_5 e_2, & e_1 * e_2 = \alpha_6 e_2. \end{cases}$
- (3) $\begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha_7 e_1 + \alpha_8 e_2, & e_1 * e_2 = \alpha_9 e_1, & e_2 * e_1 = \alpha_9 e_1, & e_2 * e_2 = \alpha_9 e_2. \end{cases}$
- (4) $\begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha_{10} e_1 + \alpha_{11} e_2, & e_1 * e_2 = 2\alpha_{10} e_2, & e_2 * e_1 = \alpha_{12} e_1 + \alpha_{10} e_2, & e_2 * e_2 = 2\alpha_{12} e_2. \end{cases}$

We may assume $\alpha_4 \alpha_9 \alpha_{12} \neq 0$.

First, we consider the first form. Then we have the following cases:

- If $\alpha_3 \neq 0$, then choose $\xi = \alpha_3^{-1}, \nu = -\alpha_2 \alpha_3^{-2}$ and obtain the parametric family:

$$\mathcal{C}_{13}^{\alpha} := \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = e_2, & e_2 * e_1 = \alpha e_2. \end{cases}$$

The algebras $\mathcal{C}_{13}^{\alpha}$ and $\mathcal{C}_{13}^{\alpha'}$ are isomorphic if and only if $\alpha = \alpha'$.

- If $\alpha_3 = 0, \alpha_1 \neq 0$, then choose $\xi = \alpha_1^{-1}$ and obtain the parametric family:

$$\mathcal{C}_{14}^{\alpha} := \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = e_1 + \alpha e_2, & e_2 * e_1 = e_2. \end{cases}$$

The algebras $\mathcal{C}_{14}^{\alpha}$ and $\mathcal{C}_{14}^{\alpha'}$ are isomorphic if and only if $\alpha = \alpha'$.

- If $\alpha_1 = \alpha_3 = 0$, then we obtain the parametric family:

$$\mathcal{C}_{15}^{\alpha} := \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_2. \end{cases}$$

The algebras $\mathcal{C}_{15}^{\alpha}$ and $\mathcal{C}_{15}^{\alpha'}$ are isomorphic if and only if $\alpha = \alpha'$.

Second, we consider the second form. Then we have the following cases:

- If $\alpha_4 \neq \alpha_6$, then choose $\xi = \alpha_4^{-1}, \nu = \frac{\alpha_5}{\alpha_4 - \alpha_6}$ and obtain the parametric family:

$$\mathcal{C}_{16}^{\alpha} := \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = e_1, & e_1 * e_2 = \alpha e_2. \end{cases}$$

The algebras $\mathcal{C}_{16}^{\alpha}$ and $\mathcal{C}_{16}^{\alpha'}$ are isomorphic if and only if $\alpha = \alpha'$.

- If $\alpha_4 = \alpha_6$, then choose $\xi = \alpha_4^{-1}$ and obtain the parametric family:

$$\mathcal{C}_{17}^\alpha := \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = e_1 + \alpha e_2, \quad e_1 * e_2 = e_2. \end{cases}$$

The algebras \mathcal{C}_{17}^α and $\mathcal{C}_{17}^{\alpha'}$ are isomorphic if and only if $\alpha = \alpha'$.

Third, we consider the third form. Then, we choose $\xi = \alpha_9^{-\frac{1}{2}}, \nu = -\frac{\alpha_7 \alpha_9^{-\frac{3}{2}}}{2}$ and obtain the parametric family:

$$\mathcal{C}_{18}^\alpha := \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_2, \quad e_1 * e_2 = e_1, \quad e_2 * e_1 = e_1, \quad e_2 * e_2 = e_2. \end{cases}$$

The algebras \mathcal{C}_{18}^α and $\mathcal{C}_{18}^{\alpha'}$ are isomorphic if and only if $\alpha = \alpha'$.

Finally, we consider the fourth form. Then, we choose $\xi = \alpha_{12}^{-\frac{1}{2}}, \nu = -\alpha_{10} \alpha_{12}^{-\frac{3}{2}}$ and obtain the parametric family:

$$\mathcal{C}_{19}^\alpha := \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_2, \quad e_2 * e_1 = e_1, \quad e_2 * e_2 = 2e_2. \end{cases}$$

The algebras \mathcal{C}_{19}^α and $\mathcal{C}_{19}^{\alpha'}$ are isomorphic if and only if $\alpha = \alpha'$.

1.3.4. *Compatible pre-Lie algebras defined on \mathcal{C}_{04} .* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{04}, \mathcal{C}_{04})$, the compatible pre-Lie algebra structures defined on \mathcal{C}_{04} are of the form:

$$(1) \begin{cases} e_2 \cdot e_1 = e_1, \\ e_2 * e_1 = \alpha_1 e_1, \quad e_2 * e_2 = \alpha_2 e_1 + \alpha_3 e_2. \end{cases}$$

$$(2) \begin{cases} e_2 \cdot e_1 = e_1, \\ e_1 * e_2 = \alpha_4 e_1, \quad e_2 * e_1 = \alpha_5 e_1, \quad e_2 * e_2 = \alpha_6 e_1 + \alpha_4 e_2. \end{cases}$$

We may assume $\alpha_4 \neq 0$.

First, we study the first form. Then we have the following cases:

- If $\alpha_2 \neq 0$, then choose $\xi = \alpha_2$ and obtain the parametric family:

$$\mathcal{C}_{20}^{\alpha, \beta} := \begin{cases} e_2 \cdot e_1 = e_1, \\ e_2 * e_1 = \alpha e_1, \quad e_2 * e_2 = e_1 + \beta e_2. \end{cases}$$

The algebras $\mathcal{C}_{20}^{\alpha, \beta}$ and $\mathcal{C}_{20}^{\alpha', \beta'}$ are isomorphic if and only if $(\alpha, \beta) = (\alpha', \beta')$.

- If $\alpha_2 = 0$, we obtain the parametric family:

$$\mathcal{C}_{21}^{\alpha, \beta} := \begin{cases} e_2 \cdot e_1 = e_1, \\ e_2 * e_1 = \alpha e_1, \quad e_2 * e_2 = \beta e_2. \end{cases}$$

The algebras $\mathcal{C}_{21}^{\alpha, \beta}$ and $\mathcal{C}_{21}^{\alpha', \beta'}$ are isomorphic if and only if $(\alpha, \beta) = (\alpha', \beta')$.

Second, we consider the second form. Then we have the following cases:

- If $\alpha_6 \neq 0$, then choose $\xi = \alpha_6$ and obtain the parametric family:

$$\mathcal{C}_{22}^{\alpha \neq 0, \beta} := \begin{cases} e_2 \cdot e_1 = e_1, \\ e_1 * e_2 = \alpha e_1, \quad e_2 * e_1 = \beta e_1, \quad e_2 * e_2 = e_1 + \alpha e_2. \end{cases}$$

The algebras $\mathcal{C}_{22}^{\alpha, \beta}$ and $\mathcal{C}_{22}^{\alpha', \beta'}$ are isomorphic if and only if $(\alpha, \beta) = (\alpha', \beta')$.

- If $\alpha_6 = 0$, we obtain the parametric family:

$$\mathcal{C}_{23}^{\alpha \neq 0, \beta} := \begin{cases} e_2 \cdot e_1 = e_1, \\ e_1 * e_2 = \alpha e_1, \quad e_2 * e_1 = \beta e_1, \quad e_2 * e_2 = \alpha e_2. \end{cases}$$

The algebras $\mathcal{C}_{23}^{\alpha, \beta}$ and $\mathcal{C}_{23}^{\alpha', \beta'}$ are isomorphic if and only if $(\alpha, \beta) = (\alpha', \beta')$.

1.3.5. *Compatible pre-Lie algebras defined on $\mathcal{C}_{05}^{\alpha \neq 0, \frac{1}{2}, 1}$.* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{05}^{\alpha}, \mathcal{C}_{05}^{\alpha})$, the compatible pre-Lie algebra structures defined on $\mathcal{C}_{05}^{\alpha}$ are of the form:

$$\begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \alpha e_2, \\ e_1 * e_1 = \alpha_1 e_1 + \alpha_2 e_2, & e_1 * e_2 = \alpha_3 e_2. \end{cases}$$

Then we have the following cases:

- If $\alpha_2 \neq 0$, then choose $\xi = \alpha_2$ and obtain the parametric family:

$$\mathcal{C}_{24}^{\alpha, \beta, \gamma} := \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \alpha e_2, \\ e_1 * e_1 = \beta e_1 + e_2, & e_1 * e_2 = \gamma e_2. \end{cases}$$

The algebras $\mathcal{C}_{24}^{\alpha, \beta, \gamma}$ and $\mathcal{C}_{24}^{\alpha', \beta', \gamma'}$ are isomorphic if and only if $(\alpha, \beta, \gamma) = (\alpha', \beta', \gamma')$.

- If $\alpha_2 = 0$, we obtain the parametric family:

$$\mathcal{C}_{25}^{\alpha, \beta, \gamma} := \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \alpha e_2, \\ e_1 * e_1 = \beta e_1, & e_1 * e_2 = \gamma e_2. \end{cases}$$

The algebras $\mathcal{C}_{25}^{\alpha, \beta, \gamma}$ and $\mathcal{C}_{25}^{\alpha', \beta', \gamma'}$ are isomorphic if and only if $(\alpha, \beta, \gamma) = (\alpha', \beta', \gamma')$.

1.3.6. *Compatible pre-Lie algebras defined on $\mathcal{C}_{05}^{\frac{1}{2}}$.* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{05}^{\frac{1}{2}}, \mathcal{C}_{05}^{\frac{1}{2}})$, the compatible pre-Lie algebra structures defined on $\mathcal{C}_{05}^{\frac{1}{2}}$ are of the form:

$$\begin{aligned} (1) & \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \frac{1}{2} e_2, \\ e_1 * e_1 = \alpha_1 e_1 + \alpha_2 e_2, & e_1 * e_2 = \alpha_3 e_2. \end{cases} \\ (2) & \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \frac{1}{2} e_2, \\ e_1 * e_1 = 2\alpha_4 e_1, & e_1 * e_2 = \alpha_4 e_2, \quad e_2 * e_2 = \alpha_5 e_1. \end{cases} \\ (3) & \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \frac{1}{2} e_2, \\ e_1 * e_1 = 2\alpha_6 e_1, & e_1 * e_2 = \alpha_7 e_1 + \alpha_6 e_2, \quad e_2 * e_1 = 2\alpha_7 e_1 \quad e_2 * e_2 = \alpha_8 e_1 + \alpha_7 e_2. \end{cases} \end{aligned}$$

We may assume $\alpha_5 \alpha_7 \neq 0$.

If the compatible pre-Lie algebra structures defined on $\mathcal{C}_{05}^{\frac{1}{2}}$ is of the first form, then we obtain the algebras $\mathcal{C}_{24}^{\frac{1}{2}, \beta, \gamma}$ and $\mathcal{C}_{25}^{\frac{1}{2}, \beta, \gamma}$.

Assume now that the compatible pre-Lie algebra structures defined on $\mathcal{C}_{05}^{\frac{1}{2}}$ is of the second form. Then, choose $\xi = \alpha_5^{-\frac{1}{2}}$ and obtain the parametric family:

$$\mathcal{C}_{26}^{\alpha} := \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \frac{1}{2} e_2, \\ e_1 * e_1 = 2\alpha e_1, & e_1 * e_2 = \alpha e_2, \quad e_2 * e_2 = e_1. \end{cases}$$

The algebras $\mathcal{C}_{26}^{\alpha}$ and $\mathcal{C}_{26}^{\alpha'}$ are isomorphic if and only if $\alpha = \alpha'$.

Finally, assume that the compatible pre-Lie algebra structures defined on $\mathcal{C}_{05}^{\frac{1}{2}}$ is of the third form. Then, choose $\xi = \alpha_7^{-1}$ and obtain the parametric family:

$$\mathcal{C}_{27}^{\alpha,\beta} := \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \frac{1}{2}e_2, \\ e_1 * e_1 = 2\alpha e_1, & e_1 * e_2 = e_1 + \alpha e_2, \quad e_2 * e_1 = 2e_1 \quad e_2 * e_2 = \beta e_1 + e_2. \end{cases}$$

The algebras $\mathcal{C}_{27}^{\alpha,\beta}$ and $\mathcal{C}_{27}^{\alpha',\beta'}$ are isomorphic if and only if $(\alpha, \beta) = (\alpha', \beta')$.

1.3.7. *Compatible pre-Lie algebras defined on \mathcal{C}_{05}^1 .* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{05}^1, \mathcal{C}_{05}^1)$, the compatible pre-Lie algebra structures defined on \mathcal{C}_{05}^1 are of the form:

$$(1) \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha_1 e_1 + \alpha_2 e_2, & e_1 * e_2 = \alpha_3 e_2. \end{cases}$$

$$(2) \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha_4 e_1, & e_1 * e_2 = \alpha_4 e_2, \quad e_2 * e_1 = \alpha_5 e_1, \quad e_2 * e_2 = \alpha_5 e_2. \end{cases}$$

We may assume $\alpha_5 \neq 0$.

First, suppose that the compatible pre-Lie algebra structures defined on \mathcal{C}_{05}^1 is of the first form. Then we have the following cases:

- If $\alpha_1 \neq \alpha_3$, then choose $\nu = \frac{\alpha_2}{\alpha_1 - \alpha_3}$ and obtain the parametric family $\mathcal{C}_{25}^{1,\beta,\gamma \neq \beta}$.
- If $\alpha_1 = \alpha_3$, then we will consider the following two cases.
 - If $\alpha_2 \neq 0$, then choose $\xi = \alpha_2$ and obtain the parametric family $\mathcal{C}_{24}^{1,\beta,\beta}$.
 - If $\alpha_2 = 0$, then we obtain the parametric family $\mathcal{C}_{25}^{1,\beta,\beta}$.

Assume now that the compatible pre-Lie algebra structures defined on \mathcal{C}_{05}^1 is of the second form. Then, choose $\xi = \alpha_5^{-1}$, $\nu = -\alpha_4 \alpha_5^{-1}$ and obtain the algebra:

$$\mathcal{C}_{28} := \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = e_2, \\ e_2 * e_1 = e_1, & e_2 * e_2 = e_2. \end{cases}$$

1.3.8. *Compatible pre-Lie algebras defined on \mathcal{C}_{05}^0 .* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{05}^0, \mathcal{C}_{05}^0)$, the compatible pre-Lie algebra structures defined on \mathcal{C}_{05}^0 are of the form:

$$(1) \begin{cases} e_1 \cdot e_1 = e_1, \\ e_1 * e_1 = \alpha_1 e_1 + \alpha_2 e_2, \quad e_1 * e_2 = \alpha_3 e_2. \end{cases}$$

$$(2) \begin{cases} e_1 \cdot e_1 = e_1, \\ e_1 * e_1 = \alpha_4 e_1, \quad e_2 * e_2 = \alpha_5 e_2. \end{cases}$$

$$(3) \begin{cases} e_1 \cdot e_1 = e_1, \\ e_1 * e_1 = \alpha_6 e_1 + \alpha_7 e_2, \quad e_1 * e_2 = \alpha_8 e_1, \quad e_2 * e_1 = \alpha_8 e_1, \quad e_2 * e_2 = \alpha_8 e_2. \end{cases}$$

We may assume $\alpha_5 \alpha_8 \neq 0$.

If the compatible pre-Lie algebra structures defined on \mathcal{C}_{05}^0 is of the first form, then we obtain the algebras $\mathcal{C}_{24}^{0,\beta,\gamma}$ and $\mathcal{C}_{25}^{0,\beta,\gamma}$.

Assume now that the compatible pre-Lie algebra structures defined on \mathcal{C}_{05}^0 are of the second form. Then, choose $\xi = \alpha_5^{-1}$ and obtain the parametric family:

$$\mathcal{C}_{29}^{\alpha} := \begin{cases} e_1 \cdot e_1 = e_1, \\ e_1 * e_1 = \alpha e_1, \quad e_2 * e_2 = e_2. \end{cases}$$

The algebras $\mathcal{C}_{29}^{\alpha}$ are $\mathcal{C}_{29}^{\alpha'}$ are isomorphic if and only if $\alpha = \alpha'$.

Finally, assume that the compatible pre-Lie algebra structures defined on \mathcal{C}_{05}^0 is of the third form. Then, choose $\xi = \alpha_8^{-1}$ and obtain the parametric family:

$$\mathcal{C}_{30}^{\alpha,\beta} := \begin{cases} e_1 \cdot e_1 = e_1, \\ e_1 * e_1 = \alpha e_1 + \beta e_2, \quad e_1 * e_2 = e_1, \quad e_2 * e_1 = e_1, \quad e_2 * e_2 = e_2. \end{cases}$$

The algebras $\mathcal{C}_{30}^{\alpha,\beta}$ and $\mathcal{C}_{30}^{\alpha',\beta'}$ are isomorphic if and only if $(\alpha, \beta) = (\alpha', \beta')$.

1.3.9. *Compatible pre-Lie algebras defined on $\mathcal{C}_{06}^{\alpha \neq 0, 1, 2}$.* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{06}^\alpha, \mathcal{C}_{06}^\alpha)$, the compatible pre-Lie algebra structures defined on \mathcal{C}_{06}^α are of the form:

$$\begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \alpha e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha_1 e_1 + \alpha_2 e_2, & e_1 * e_2 = \alpha_3 e_2, & e_2 * e_1 = \alpha_1 e_2. \end{cases}$$

Then we have the following cases:

- If $\alpha_2 \neq 0$, then choose $\xi = \alpha_2$ and obtain the parametric family:

$$\mathcal{C}_{31}^{\alpha,\beta,\gamma} := \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \alpha e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \beta e_1 + e_2, & e_1 * e_2 = \gamma e_2, & e_2 * e_1 = \beta e_2. \end{cases}$$

The algebras $\mathcal{C}_{31}^{\alpha,\beta,\gamma}$ and $\mathcal{C}_{31}^{\alpha',\beta',\gamma'}$ are isomorphic if and only if $(\alpha, \beta, \gamma) = (\alpha', \beta', \gamma')$.

- If $\alpha_2 = 0$, we obtain the parametric family:

$$\mathcal{C}_{32}^{\alpha,\beta,\gamma} := \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = \alpha e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \beta e_1, & e_1 * e_2 = \gamma e_2, & e_2 * e_1 = \beta e_2. \end{cases}$$

The algebras $\mathcal{C}_{32}^{\alpha,\beta,\gamma}$, $\mathcal{C}_{32}^{\alpha',\beta',\gamma'}$ are isomorphic if and only if $(\alpha, \beta, \gamma) = (\alpha', \beta', \gamma')$.

1.3.10. *Compatible pre-Lie algebras defined on \mathcal{C}_{06}^0 .* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{06}^0, \mathcal{C}_{06}^0)$, the compatible pre-Lie algebra structures defined on \mathcal{C}_{06}^0 are of the forms:

$$(1) \begin{cases} e_1 \cdot e_1 = e_1, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha_1 e_1 + \alpha_2 e_2, & e_1 * e_2 = \alpha_3 e_2, & e_2 * e_1 = \alpha_1 e_2. \end{cases}$$

$$(2) \begin{cases} e_1 \cdot e_1 = e_1, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha_4 e_1, & e_1 * e_2 = \alpha_5 e_1, & e_2 * e_1 = \alpha_4 e_2, & e_2 * e_2 = \alpha_5 e_2. \end{cases}$$

We may assume $\alpha_5 \neq 0$.

Let us first consider the first form. Then we have the following cases:

- If $\alpha_3 \neq 0$, then $\nu = -\frac{\alpha_2}{\alpha_3}$ choose and obtain the parametric family $\mathcal{C}_{32}^{0,\beta,\gamma \neq 0}$.
- If $\alpha_3 = 0$, then we consider the following two cases.
 - If $\alpha_2 \neq 0$, then choose $\xi = \alpha_2$ and obtain the parametric family $\mathcal{C}_{31}^{0,\beta,0}$.
 - If $\alpha_2 = 0$, then we obtain the parametric family $\mathcal{C}_{32}^{0,\beta,0}$.

Now, we consider the second form. Then choose $\xi = \alpha_5^{-1}$, $\nu = -\alpha_4 \alpha_5^{-1}$ and obtain the algebra:

$$\mathcal{C}_{33} := \begin{cases} e_1 \cdot e_1 = e_1, & e_2 \cdot e_1 = e_2, \\ e_1 * e_2 = e_1, & e_2 * e_2 = e_2. \end{cases}$$

1.3.11. *Compatible pre-Lie algebras defined on \mathcal{C}_{06}^1 .* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{06}^1, \mathcal{C}_{06}^1)$, the compatible pre-Lie algebra structures defined on \mathcal{C}_{06}^1 are of the forms:

$$(1) \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha_1 e_1 + \alpha_2 e_2, & e_1 * e_2 = \alpha_3 e_2, & e_2 * e_1 = \alpha_1 e_2. \end{cases}$$

$$(2) \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha_4 e_1, & e_1 * e_2 = \alpha_4 e_2, & e_2 * e_1 = \alpha_4 e_2, & e_2 * e_2 = \alpha_5 e_1 + \alpha_6 e_2. \end{cases}$$

We may assume $(\alpha_5, \alpha_6) \neq (0, 0)$.

If the compatible pre-Lie algebra structures defined on \mathcal{C}_{06}^1 is of the first form, then we obtain the algebras $\mathcal{C}_{31}^{1,\beta,\gamma}$ and $\mathcal{C}_{32}^{1,\beta,\gamma}$.

Assume now that the compatible pre-Lie algebra structures defined on \mathcal{C}_{06}^1 is of the second form. Then we have the following cases:

- If $\alpha_5 \neq 0$, then choose $\xi = \alpha_5^{-\frac{1}{2}}$ and obtain the parametric family:

$$\mathcal{C}_{34}^{\alpha,\beta} := \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = \alpha e_2, & e_2 * e_1 = \alpha e_2, & e_2 * e_2 = e_1 + \beta e_2. \end{cases}$$

The algebras $\mathcal{C}_{34}^{\alpha,\beta}$ and $\mathcal{C}_{34}^{\alpha',\beta'}$ are isomorphic if and only if $(\alpha, \beta) = (\alpha', \beta')$.

- If $\alpha_5 = 0$, then choose $\xi = \alpha_6^{-1}$ and obtain the parametric family:

$$\mathcal{C}_{35}^{\alpha} := \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = \alpha e_2, & e_2 * e_1 = \alpha e_2, & e_2 * e_2 = e_2. \end{cases}$$

The algebras $\mathcal{C}_{35}^{\alpha}$ and $\mathcal{C}_{35}^{\alpha'}$ are isomorphic if and only if $\alpha = \alpha'$.

1.3.12. *Compatible pre-Lie algebras defined on \mathcal{C}_{06}^2 .* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{06}^2, \mathcal{C}_{06}^2)$, the compatible pre-Lie algebra structures defined on \mathcal{C}_{06}^2 are of the forms:

$$(1) \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = 2e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha_1 e_1 + \alpha_2 e_2, & e_1 * e_2 = \alpha_3 e_2, & e_2 * e_1 = \alpha_1 e_2. \end{cases}$$

$$(2) \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = 2e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha_4 e_1 + \alpha_5 e_2, & e_1 * e_2 = 2\alpha_4 e_2, & e_2 * e_1 = \alpha_6 e_1 + \alpha_4 e_2, & e_2 * e_2 = 2\alpha_6 e_2. \end{cases}$$

$$(3) \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = 2e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha_7 e_1, & e_1 * e_2 = \alpha_8 e_1 + 2\alpha_7 e_2, & e_2 * e_1 = 2\alpha_8 e_1 + \alpha_7 e_2, & e_2 * e_2 = \alpha_8 e_2. \end{cases}$$

We may assume $\alpha_6 \alpha_8 \neq 0$.

If the compatible pre-Lie algebra structures defined on \mathcal{C}_{06}^2 is of the first form, then we obtain the algebras $\mathcal{C}_{31}^{2,\beta,\gamma}$ and $\mathcal{C}_{32}^{2,\beta,\gamma}$.

Assume now that the compatible pre-Lie algebra structures defined on \mathcal{C}_{06}^2 are of the second form. Then, choose $\xi = \alpha_6^{-1}$ and obtain the parametric family:

$$\mathcal{C}_{36}^{\alpha,\beta} := \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = 2e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1 + \beta e_2, & e_1 * e_2 = 2\alpha e_2, & e_2 * e_1 = e_1 + \alpha e_2, & e_2 * e_2 = 2e_2. \end{cases}$$

The algebras $\mathcal{C}_{36}^{\alpha,\beta}$ and $\mathcal{C}_{36}^{\alpha',\beta'}$ are isomorphic if and only if $(\alpha, \beta) = (\alpha', \beta')$.

Finally, assume that the compatible pre-Lie algebra structures defined on \mathcal{C}_{06}^2 are of the third form. Then, choose $\xi = \alpha_8^{-1}$ and obtain the parametric family:

$$\mathcal{C}_{37}^\alpha := \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = 2e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = e_1 + 2\alpha e_2, & e_2 * e_1 = 2e_1 + \alpha e_2, & e_2 * e_2 = e_2. \end{cases}$$

The algebras \mathcal{C}_{37}^α and $\mathcal{C}_{37}^{\alpha'}$ are isomorphic if and only if $\alpha = \alpha'$.

1.3.13. *Compatible pre-Lie algebras defined on \mathcal{C}_{07} .* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{07}, \mathcal{C}_{07})$, the compatible pre-Lie algebra structures defined on \mathcal{C}_{07} are of the forms:

$$(1) \begin{cases} e_1 \cdot e_1 = e_1, & e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = (\alpha_3 + \alpha_2 - \alpha_1)e_1 - \alpha_2 e_2, & e_1 * e_2 = \alpha_1 e_1 + \alpha_2 e_2, \\ e_2 * e_1 = \alpha_1 e_1 + \alpha_2 e_2, & e_2 * e_2 = -\alpha_1 e_1 + \alpha_3 e_2. \end{cases}$$

$$(2) \begin{cases} e_1 \cdot e_1 = e_1, & e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha_4 e_1, & e_2 * e_2 = \alpha_5 e_2. \end{cases}$$

We may assume that $\alpha_4 \neq \alpha_5$.

Suppose first that the compatible pre-Lie algebra structures defined on \mathcal{C}_{07} has the first form. Then we obtain the algebras:

$$\mathcal{C}_{38}^{\alpha, \beta, \gamma} := \begin{cases} e_1 \cdot e_1 = e_1, & e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = (\gamma + \beta - \alpha)e_1 - \beta e_2, & e_1 * e_2 = \alpha e_1 + \beta e_2, \\ e_2 * e_1 = \alpha e_1 + \beta e_2, & e_2 * e_2 = -\alpha e_1 + \gamma e_2. \end{cases}$$

The algebras $\mathcal{C}_{38}^{\alpha, \beta, \gamma}$ and $\mathcal{C}_{38}^{\alpha', \beta', \gamma'}$ are isomorphic if and only if

$$(\alpha, \beta, \gamma) = (\alpha', \beta', \gamma') \text{ or } (\alpha, \beta, \gamma) = (\beta', \alpha', -\alpha' + \beta' + \gamma').$$

Now, assume that the compatible pre-Lie algebra structures defined on \mathcal{C}_{07} has the second form. Then we obtain the algebras:

$$\mathcal{C}_{39}^{\alpha, \beta \neq \alpha} := \begin{cases} e_1 \cdot e_1 = e_1, & e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_2 * e_2 = \beta e_2. \end{cases}$$

The algebras $\mathcal{C}_{39}^{\alpha, \beta}$ and $\mathcal{C}_{39}^{\alpha', \beta'}$ are isomorphic if and only if $(\alpha, \beta) = (\alpha', \beta')$ or $(\alpha, \beta) = (\beta', \alpha')$.

1.3.14. *Compatible pre-Lie algebras defined on \mathcal{C}_{08} .* From the computation of $Z_{\text{CPL}}^2(\mathcal{C}_{08}, \mathcal{C}_{08})$, the compatible pre-Lie algebra structures defined on \mathcal{C}_{08} are of the forms:

$$(1) \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = 2e_2, & e_2 \cdot e_1 = \frac{1}{2}e_1 + e_2, & e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha_1 e_1 + \alpha_2 e_2, & e_1 * e_2 = 2\alpha_1 e_2, & e_2 * e_1 = \alpha_3 e_1 + \alpha_1 e_2, & e_2 * e_2 = 2\alpha_3 e_2. \end{cases}$$

$$(2) \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = 2e_2, \\ e_2 \cdot e_1 = \frac{1}{2}e_1 + e_2, & e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha_4 e_1, & e_1 * e_2 = \alpha_5 e_1 + 2\alpha_4 e_2, \\ e_2 * e_1 = (2\alpha_5 + \frac{1}{2}\alpha_4)e_1 + \alpha_4 e_2, & e_2 * e_2 = \frac{1}{2}\alpha_5 e_1 + (\alpha_4 + \alpha_5)e_2. \end{cases}$$

We may assume $\alpha_5 \neq 0$. Then we obtain the following compatible pre-Lie algebras:

$$\mathcal{C}_{40}^{\alpha, \beta, \gamma} := \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = 2e_2, & e_2 \cdot e_1 = \frac{1}{2}e_1 + e_2, & e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha e_1 + \beta e_2, & e_1 * e_2 = 2\alpha e_2, & e_2 * e_1 = \gamma e_1 + \alpha e_2, & e_2 * e_2 = 2\gamma e_2. \end{cases}$$

$$\mathcal{C}_{41}^{\alpha, \beta \neq 0} := \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = 2e_2, & e_2 \cdot e_1 = \frac{1}{2}e_1 + e_2, & e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = \beta e_1 + 2\alpha e_2, \\ e_2 * e_1 = (2\beta + \frac{1}{2}\alpha)e_1 + \alpha e_2, & e_2 * e_2 = \frac{1}{2}\beta e_1 + (\alpha + \beta)e_2. \end{cases}$$

Moreover, the algebras $\mathcal{C}_{40}^{\alpha,\beta,\gamma}$ and $\mathcal{C}_{40}^{\alpha',\beta',\gamma'}$ are isomorphic if and only if

$$(\alpha', \beta', \gamma') = (\alpha, \beta, \gamma) \text{ or } (\alpha', \beta', \gamma') = (4\gamma - \alpha, \beta - 8\alpha + 16\gamma, \gamma).$$

Also, the algebras $\mathcal{C}_{41}^{\alpha,\beta}$ and $\mathcal{C}_{41}^{\alpha',\beta'}$ are isomorphic if and only if

$$(\alpha', \beta') = (\alpha, \beta) \text{ or } (\alpha', \beta') = (\alpha + 4\beta, -\beta).$$

1.4. The algebraic classification of compatible commutative associative algebras.

It is easy to see that each commutative pre-Lie algebra is associative. Hence, we can choose only commutative compatible pre-Lie algebras from the Theorem 7.

Theorem 8. *Let \mathcal{C} be a nonzero 2-dimensional compatible commutative associative algebra. Then \mathcal{C} is isomorphic to one and only one and only one of the following algebras:*

$$\begin{aligned} \mathcal{C}_{03} & : e_1 * e_1 = e_2. \\ \mathcal{C}_{05}^0 & : e_1 * e_1 = e_1. \\ \mathcal{C}_{06}^1 & : e_1 * e_1 = e_1, e_1 * e_2 = e_2, e_2 * e_1 = e_2. \\ \mathcal{C}_{07} & : e_1 * e_1 = e_1, e_2 * e_2 = e_2. \\ \mathcal{C}_{13}^1 & : \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = e_1, e_1 * e_2 = e_2, e_2 * e_1 = e_2. \end{cases} \\ \mathcal{C}_{15}^\alpha & : \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_2. \end{cases} \\ \mathcal{C}_{16}^0 & : \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = e_1. \end{cases} \\ \mathcal{C}_{18}^\alpha & : \begin{cases} e_1 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_2, e_1 * e_2 = e_1, e_2 * e_1 = e_1, e_2 * e_2 = e_2. \end{cases} \\ \mathcal{C}_{24}^{0,\beta,0} & : \begin{cases} e_1 \cdot e_1 = e_1, \\ e_1 * e_1 = \beta e_1 + e_2. \end{cases} \\ \mathcal{C}_{25}^{0,\beta,0} & : \begin{cases} e_1 \cdot e_1 = e_1, \\ e_1 * e_1 = \beta e_1. \end{cases} \\ \mathcal{C}_{29}^\alpha & : \begin{cases} e_1 \cdot e_1 = e_1, \\ e_1 * e_1 = \alpha e_1, e_2 * e_2 = e_2. \end{cases} \\ \mathcal{C}_{30}^{\alpha,\beta} & : \begin{cases} e_1 \cdot e_1 = e_1, \\ e_1 * e_1 = \alpha e_1 + \beta e_2, e_1 * e_2 = e_1, e_2 * e_1 = e_1, e_2 * e_2 = e_2. \end{cases} \\ \mathcal{C}_{31}^{1,\beta,\beta} & : \begin{cases} e_1 \cdot e_1 = e_1, e_1 \cdot e_2 = e_2, e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \beta e_1 + e_2, e_1 * e_2 = \beta e_2, e_2 * e_1 = \beta e_2. \end{cases} \\ \mathcal{C}_{32}^{1,\beta,\beta} & : \begin{cases} e_1 \cdot e_1 = e_1, e_1 \cdot e_2 = e_2, e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \beta e_1, e_1 * e_2 = \beta e_2, e_2 * e_1 = \beta e_2. \end{cases} \\ \mathcal{C}_{34}^{\alpha,\beta} & : \begin{cases} e_1 \cdot e_1 = e_1, e_1 \cdot e_2 = e_2, e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1, e_1 * e_2 = \alpha e_2, e_2 * e_1 = \alpha e_2, e_2 * e_2 = e_1 + \beta e_2. \end{cases} \\ \mathcal{C}_{35}^\alpha & : \begin{cases} e_1 \cdot e_1 = e_1, e_1 \cdot e_2 = e_2, e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1, e_1 * e_2 = \alpha e_2, e_2 * e_1 = \alpha e_2, e_2 * e_2 = e_2. \end{cases} \\ \mathcal{C}_{38}^{\alpha,\beta,\gamma} & : \begin{cases} e_1 \cdot e_1 = e_1, e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = (\gamma + \beta - \alpha) e_1 - \beta e_2, e_1 * e_2 = \alpha e_1 + \beta e_2, \\ e_2 * e_1 = \alpha e_1 + \beta e_2, e_2 * e_2 = -\alpha e_1 + \gamma e_2. \end{cases} \\ \mathcal{C}_{39}^{\alpha,\beta \neq \alpha} & : \begin{cases} e_1 \cdot e_1 = e_1, e_2 \cdot e_2 = e_2, \\ e_1 * e_1 = \alpha e_1, e_2 * e_2 = \beta e_2. \end{cases} \end{aligned}$$

All algebras are non-isomorphic, except $\mathcal{C}_{38}^{\alpha,\beta,\gamma} \cong \mathcal{C}_{38}^{\beta,\alpha,-\alpha+\beta+\gamma}$, $\mathcal{C}_{39}^{\alpha,\beta} \cong \mathcal{C}_{39}^{\beta,\alpha}$.

1.5. The algebraic classification of compatible associative algebras.

Definition 9. A compatible associative algebra is a vector space \mathbf{A} equipped with two multiplication \cdot and another multiplication $*$, such that, (\mathbf{A}, \cdot) , $(\mathbf{A}, *)$ and $(\mathbf{A}, \cdot + *)$ are associative algebras. These two operations are required to satisfy the following identities.

$$\begin{aligned} (x \cdot y) \cdot z &= x \cdot (y \cdot z), \\ (x * y) * z &= x * (y * z), \\ (x * y) \cdot z + (x \cdot y) * z &= x * (y \cdot z) + x \cdot (y * z). \end{aligned}$$

Theorem 10. Let \mathcal{C} be a nonzero 2-dimensional compatible associative algebra. Then \mathcal{C} is isomorphic to one and only one compatible commutative associative algebra listed in Theorem 8 or one of the following algebras:

$$\begin{aligned} \mathcal{C}_{05}^1 &: e_1 * e_1 = e_1, e_1 * e_2 = e_2. \\ \mathcal{C}_{06}^0 &: e_1 * e_1 = e_1, e_2 * e_1 = e_2. \\ \mathcal{C}_{25}^{1,\beta,\beta} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = e_2, \\ e_1 * e_1 = \beta e_1, & e_1 * e_2 = \beta e_2. \end{cases} \\ \mathcal{C}_{28} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_1 \cdot e_2 = e_2, \\ e_2 * e_1 = e_1, & e_2 * e_2 = e_2. \end{cases} \\ \mathcal{C}_{32}^{0,\beta,0} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \beta e_1, & e_2 * e_1 = \beta e_2. \end{cases} \\ \mathcal{C}_{33} &: \begin{cases} e_1 \cdot e_1 = e_1, & e_2 \cdot e_1 = e_2, \\ e_1 * e_2 = e_1, & e_2 * e_2 = e_2. \end{cases} \end{aligned}$$

1.6. The algebraic classification of compatible Novikov algebras.

Definition 11. A compatible Novikov algebra is a vector space \mathbf{A} equipped with two multiplication \cdot and another multiplication $*$, such that, (\mathbf{A}, \cdot) , $(\mathbf{A}, *)$ and $(\mathbf{A}, \cdot + *)$ are Novikov algebras. These two operations are required to satisfy the following identities.

$$\begin{aligned} (x \cdot y) \cdot z - x \cdot (y \cdot z) &= (y \cdot x) \cdot z - y \cdot (x \cdot z), \\ (x \cdot y) \cdot z &= (x \cdot z) \cdot y, \\ (x * y) * z - x * (y * z) &= (y * x) * z - y * (x * z), \\ (x * y) * z &= (x * z) * y, \\ (x * y) \cdot z - x * (y \cdot z) + (x \cdot y) * z - x \cdot (y * z) &= (y * x) \cdot z - y * (x \cdot z) + (y \cdot x) * z - y \cdot (x * z), \\ (x * y) \cdot z + (x \cdot y) * z &= (x * z) \cdot y + (x \cdot z) * y. \end{aligned}$$

Theorem 12. Let \mathcal{C} be a nonzero 2-dimensional compatible Novikov algebra. Then \mathcal{C} is isomorphic to one and only one compatible commutative associative algebra listed in Theorem 8 or one of the following algebras:

$$\begin{aligned} \mathcal{C}_{01} &: e_1 * e_1 = e_1 + e_2, e_2 * e_1 = e_2. \\ \mathcal{C}_{04} &: e_2 * e_1 = e_1. \\ \mathcal{C}_{06}^{\alpha \neq 1} &: e_1 * e_1 = e_1, e_1 * e_2 = \alpha e_2, e_2 * e_1 = e_2. \\ \mathcal{C}_{09}^{\alpha,\beta \neq 0} &: \begin{cases} e_1 \cdot e_1 = e_1 + e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1, & e_1 * e_2 = \beta e_2, e_2 * e_1 = \alpha e_2. \end{cases} \\ \mathcal{C}_{10}^{\alpha,\beta} &: \begin{cases} e_1 \cdot e_1 = e_1 + e_2, & e_2 \cdot e_1 = e_2, \\ e_1 * e_1 = \alpha e_1 + \beta e_2, & e_2 * e_1 = \alpha e_2. \end{cases} \end{aligned}$$

$$\begin{array}{l}
\mathcal{C}_{13}^{\alpha \neq 1} \\
\mathcal{C}_{14}^{\alpha} \\
\mathcal{C}_{20}^{\alpha, 0} \\
\mathcal{C}_{21}^{\alpha, 0} \\
\mathcal{C}_{22}^{\alpha \neq 0, \beta} \\
\mathcal{C}_{23}^{\alpha \neq 0, \beta} \\
\mathcal{C}_{31}^{(\alpha, \beta, \gamma) \neq (1, \beta, \beta)} \\
\mathcal{C}_{32}^{(\alpha, \beta, \gamma) \neq (1, \beta, \beta)} \\
\mathcal{C}_{33}
\end{array}
: \left\{ \begin{array}{l}
e_1 \cdot e_1 = e_2, \\
e_1 * e_1 = \alpha e_1, \quad e_1 * e_2 = e_2, \quad e_2 * e_1 = \alpha e_2. \\
e_1 \cdot e_1 = e_2, \\
e_1 * e_1 = e_1 + \alpha e_2, \quad e_2 * e_1 = e_2. \\
e_2 \cdot e_1 = e_1, \\
e_2 * e_1 = \alpha e_1, \quad e_2 * e_2 = e_1. \\
e_2 \cdot e_1 = e_1, \\
e_2 * e_1 = \alpha e_1. \\
e_2 \cdot e_1 = e_1, \\
e_1 * e_2 = \alpha e_1, \quad e_2 * e_1 = \beta e_1, \quad e_2 * e_2 = e_1 + \alpha e_2. \\
e_2 \cdot e_1 = e_1, \\
e_1 * e_2 = \alpha e_1, \quad e_2 * e_1 = \beta e_1, \quad e_2 * e_2 = \alpha e_2. \\
e_1 \cdot e_1 = e_1, \quad e_1 \cdot e_2 = \alpha e_2, \quad e_2 \cdot e_1 = e_2, \\
e_1 * e_1 = \beta e_1 + e_2, \quad e_1 * e_2 = \gamma e_2, \quad e_2 * e_1 = \beta e_2. \\
e_1 \cdot e_1 = e_1, \quad e_1 \cdot e_2 = \alpha e_2, \quad e_2 \cdot e_1 = e_2, \\
e_1 * e_1 = \beta e_1, \quad e_1 * e_2 = \gamma e_2, \quad e_2 * e_1 = \beta e_2. \\
e_1 \cdot e_1 = e_1, \quad e_2 \cdot e_1 = e_2, \\
e_1 * e_2 = e_1, \quad e_2 * e_2 = e_2.
\end{array} \right.$$

All algebras are non-isomorphic, except $\mathcal{C}_{31}^{0, \beta, \gamma} \cong \mathcal{C}_{32}^{0, \beta, \gamma}$.

2. THE GEOMETRIC CLASSIFICATION OF COMPATIBLE ALGEBRAS

2.1. Degenerations and the geometric classification of algebras. Let us introduce the techniques used to obtain the geometric classification for an arbitrary variety of compatible Ω -algebras. Given a complex vector space \mathbb{V} of dimension n , the set of bilinear maps

$$\text{Bil}(\mathbb{V} \times \mathbb{V}, \mathbb{V}) \cong \text{Hom}(\mathbb{V}^{\otimes 2}, \mathbb{V}) \cong (\mathbb{V}^*)^{\otimes 2} \otimes \mathbb{V}$$

is a vector space of dimension n^3 . The set of pairs of bilinear maps

$$\text{Bil}(\mathbb{V} \times \mathbb{V}, \mathbb{V}) \oplus \text{Bil}(\mathbb{V} \times \mathbb{V}, \mathbb{V}) \cong (\mathbb{V}^*)^{\otimes 2} \otimes \mathbb{V} \oplus (\mathbb{V}^*)^{\otimes 2} \otimes \mathbb{V}$$

which is a vector space of dimension $2n^3$. This vector space has the structure of the affine space \mathbb{C}^{2n^3} in the following sense: fixed a basis e_1, \dots, e_n of \mathbb{V} , then any pair with multiplication (μ, μ') , is determined by some parameters $c_{ij}^k, c'_{ij}{}^k \in \mathbb{C}$, called structural constants, such that

$$\mu(e_i, e_j) = \sum_{p=1}^n c_{ij}^k e_k \quad \text{and} \quad \mu'(e_i, e_j) = \sum_{p=1}^n c'_{ij}{}^k e_k,$$

which corresponds to a point in the affine space \mathbb{C}^{2n^3} . Then a set of bilinear pairs \mathcal{S} corresponds to an algebraic variety, i.e., a Zariski closed set, if there are some polynomial equations in variables $c_{ij}^k, c'_{ij}{}^k$ with zero locus equal to the set of structural constants of the bilinear pairs in \mathcal{S} . Given the identities defining a particular class of compatible Ω -algebras, we can obtain a set of polynomial equations in variables $c_{ij}^k, c'_{ij}{}^k$. This class of n -dimensional compatible Ω -algebras is a variety. Denote it by \mathcal{T}_n . Now, consider the following action of $\text{GL}(\mathbb{V})$ on \mathcal{T}_n :

$$(g * (\mu, \mu'))(x, y) := (g\mu(g^{-1}x, g^{-1}y), g\mu'(g^{-1}x, g^{-1}y))$$

for $g \in \text{GL}(\mathbb{V})$, $(\mu, \mu') \in \mathcal{T}_n$ and for any $x, y \in \mathbb{V}$. Observe that the $\text{GL}(\mathbb{V})$ -orbit of (μ, μ') , denoted $\mathcal{O}((\mu, \mu'))$, contains all the structural constants of the bilinear pairs isomorphic to the compatible Ω -algebras with structural constants (μ, μ') .

A geometric classification of a variety of algebras consists on describing the irreducible components of the variety. Recall that any affine variety can be represented as a finite union of its irreducible components uniquely. Note that describing the irreducible components of \mathcal{T}_n gives us the rigid algebras of the variety, which are those bilinear pairs with an open $\text{GL}(\mathbb{V})$ -orbit. This is due to the fact that a bilinear pair is rigid in a variety if and only if the closure of its orbit is an irreducible component of the variety. For this reason, the following notion is convenient. Denote by $\overline{\mathcal{O}((\mu, \mu'))}$ the closure of the orbit of $(\mu, \mu') \in \mathcal{T}_n$.

Definition 13. Let T and T' be two n -dimensional compatible Ω -algebras of a fixed class corresponding to the variety \mathcal{T}_n and $(\mu, \mu'), (\lambda, \lambda') \in \mathcal{T}_n$ be their representatives in the affine space, respectively. The algebra T is said to degenerate to T' , and we write $T \rightarrow T'$, if $(\lambda, \lambda') \in \overline{\mathcal{O}((\mu, \mu'))}$. If $T \not\cong T'$, then we call it a proper degeneration. Conversely, if $(\lambda, \lambda') \notin \overline{\mathcal{O}((\mu, \mu'))}$ then we say that T does not degenerate to T' and we write $T \not\rightarrow T'$.

Furthermore, for a parametric family of algebras we have the following notion.

Definition 14. Let $T(*) = \{T(\alpha) : \alpha \in I\}$ be a family of n -dimensional compatible Ω -algebras of a fixed class corresponding to \mathcal{T}_n and let T' be another algebra. Suppose that $T(\alpha)$ is represented by the structure $(\mu(\alpha), \mu'(\alpha)) \in \mathcal{T}_n$ for $\alpha \in I$ and T' is represented by the structure $(\lambda, \lambda') \in \mathcal{T}_n$. We say that the family $T(*)$ degenerates to T' , and write $T(*) \rightarrow T'$, if $(\lambda, \lambda') \in \overline{\{\mathcal{O}((\mu(\alpha), \mu'(\alpha)))\}_{\alpha \in I}}$. Conversely, if $(\lambda, \lambda') \notin \overline{\{\mathcal{O}((\mu(\alpha), \mu'(\alpha)))\}_{\alpha \in I}}$ then we call it a non-degeneration, and we write $T(*) \not\rightarrow T'$.

Observe that T' corresponds to an irreducible component of \mathcal{T}_n (more precisely, $\overline{T'}$ is an irreducible component) if and only if $T \not\rightarrow T'$ for any n -dimensional compatible Ω -algebra T and $T(*) \not\rightarrow T'$ for any parametric family of n -dimensional compatible Ω -algebras $T(*)$. To prove a particular algebra corresponds to an irreducible component, we will use the next ideas. Firstly, since $\dim \mathcal{O}((\mu, \mu')) = n^2 - \dim \mathcal{D}\text{er}(T)$, then if $T \rightarrow T'$ and $T \not\cong T'$, we have that $\dim \mathcal{D}\text{er}(T) < \dim \mathcal{D}\text{er}(T')$, where $\mathcal{D}\text{er}(T)$ denotes the Lie algebra of derivations of T . Secondly, to prove degenerations, let T and T' be two compatible Ω -algebras represented by the structures (μ, μ') and (λ, λ') from \mathcal{T}_n , respectively. Let $c_{ij}^k, c_{ij}'^k$ be the structure constants of (λ, λ') in a basis e_1, \dots, e_n of \mathbb{V} . If there exist n^2 maps $\alpha_i^j(t) : \mathbb{C}^* \rightarrow \mathbb{C}$ such that $E_i(t) = \sum_{j=1}^n \alpha_i^j(t) e_j$ ($1 \leq i \leq n$) form a basis of \mathbb{V} for any $t \in \mathbb{C}^*$ and the structure constants $c_{ij}^k(t), c_{ij}'^k(t)$ of (μ, μ') in the basis $E_1(t), \dots, E_n(t)$ satisfy $\lim_{t \rightarrow 0} c_{ij}^k(t) = c_{ij}^k$ and $\lim_{t \rightarrow 0} c_{ij}'^k(t) = c_{ij}'^k$, then $T \rightarrow T'$. In this case, $E_1(t), \dots, E_n(t)$ is called a parametrized basis for $T \rightarrow T'$. Thirdly, to prove non-degenerations we may use a remark that follows from this lemma, see [3].

Lemma 15. Consider two compatible Ω -algebras T and T' . Suppose $T \rightarrow T'$. Let C be a Zariski closed in \mathcal{T}_n that is stable by the action of the invertible upper (lower) triangular matrices. Then if there is a representation (μ, μ') of T in C , then there is a representation (λ, λ') of T' in C .

In order to apply this lemma, we will give the explicit definition of the appropriate stable Zariski closed C in terms of the variables $c_{ij}^k, c_{ij}'^k$ in each case.

Remark 16. *Moreover, let T and T' be two compatible Ω -algebras represented by the structures (μ, μ') and (λ, λ') from \mathcal{T}_n . Suppose $T \rightarrow T'$. Then if $\mu, \mu', \lambda, \lambda'$ represents algebras T_0, T_1, T'_0, T'_1 in the affine space \mathbb{C}^{n^3} of algebras with a single multiplication, respectively, we have $T_0 \rightarrow T'_0$ and $T_1 \rightarrow T'_1$. So for example, $(0, \mu)$ can not degenerate in $(\lambda, 0)$ unless $\lambda = 0$.*

Fourthly, to prove $T(*) \rightarrow T'$, suppose that $T(\alpha)$ is represented by the structure $(\mu(\alpha), \mu'(\alpha)) \in \mathcal{T}_n$ for $\alpha \in I$ and T' is represented by the structure $(\lambda, \lambda') \in \mathcal{T}_n$. Let c_{ij}^k, c_{ij}^{lk} be the structure constants of (λ, λ') in a basis e_1, \dots, e_n of \mathbb{V} . If there is a pair of maps $(f, (a_i^j))$, where $f: \mathbb{C}^* \rightarrow I$ and $a_i^j: \mathbb{C}^* \rightarrow \mathbb{C}$ are such that $E_i(t) = \sum_{j=1}^n a_i^j(t) e_j$ ($1 \leq i \leq n$) form a basis of \mathbb{V} for any $t \in \mathbb{C}^*$ and the structure constants $c_{ij}^k(t), c_{ij}^{lk}(t)$ of $(\mu(f(t)), \mu'(f(t)))$ in the basis $E_1(t), \dots, E_n(t)$ satisfy $\lim_{t \rightarrow 0} c_{ij}^k(t) = c_{ij}^k$ and $\lim_{t \rightarrow 0} c_{ij}^{lk}(t) = c_{ij}^{lk}$, then $T(*) \rightarrow T'$. In this case $E_1(t), \dots, E_n(t)$ and $f(t)$ are called a parametrized basis and a parametrized index for $T(*) \rightarrow T'$, respectively. Fifthly, to prove $T(*) \not\rightarrow T'$, we may use an analogous of Remark 16 for parametric families that follows from Lemma 17.

Lemma 17. *Consider the family of compatible Ω -algebras $T(*)$ and the compatible Ω -algebra T' . Suppose $T(*) \rightarrow T'$. Let C be a Zariski closed in \mathcal{T}_n that is stable by the action of the invertible upper (lower) triangular matrices. Then if there is a representation $(\mu(\alpha), \mu'(\alpha))$ of $T(\alpha)$ in C for every $\alpha \in I$, then there is a representation (λ, λ') of T' in C .*

Finally, the following remark simplifies the geometric problem.

Remark 18. *Let (μ, μ') and (λ, λ') represent two compatible Ω -algebras. Suppose $(\lambda, 0) \notin \mathcal{O}((\mu, 0))$, (resp. $(0, \lambda') \notin \mathcal{O}((0, \mu'))$), then $(\lambda, \lambda') \notin \mathcal{O}((\mu, \mu'))$. As we construct the classification of a given class of compatible Ω -algebras from a certain class of algebras with a single multiplication that remains unchanged, this remark becomes very useful.*

2.2. The geometric classification of compatible commutative associative algebras.

The main result of the present section is the following theorem.

Theorem 19. *The variety of complex 2-dimensional compatible commutative associative algebras has dimension 7 and it has 2 irreducible components defined by $\mathcal{O}(\mathcal{C}_{39}^{\alpha, \beta})$ and $\mathcal{O}(\mathcal{C}_{38}^{\alpha, \beta, \gamma})$. In particular, there are no rigid algebras in this variety.*

Proof. Thanks to Theorem 8, we have the algebraic classification of 2-dimensional compatible commutative associative algebras. After carefully checking the dimensions of orbit closures of the more important for us algebras, we have

$$\dim \mathcal{O}(\mathcal{C}_{38}^{\alpha, \beta, \gamma}) = 7 \quad \text{and} \quad \dim \mathcal{O}(\mathcal{C}_{39}^{\alpha, \beta}) = 6.$$

$\mathcal{C}_{38}^* \not\rightarrow \mathcal{C}_{39}^{\alpha, \beta}$ due to the following relation

$$\mathcal{R} = \{c_{22}^1 = c_{12}^1 = 0, c_{22}^2 c_{12}^{\prime 2} + c_{12}^2 c_{21}^{\prime 1} + c_{11}^1 c_{22}^{\prime 2} = c_{11}^{\prime 1} c_{22}^2 + c_{11}^1 c_{21}^{\prime 1} + c_{12}^2 c_{22}^{\prime 2}\}.$$

Thanks to [27], we have $\mathcal{C}_{07} \rightarrow \{\mathcal{C}_{03}, \mathcal{C}_{05}^0, \mathcal{C}_{06}^1\}$. All necessary degenerations are given below

$$\begin{array}{c} \mathcal{C}_{38}^{0,0,t^{-1}} \xrightarrow{(te_1, te_2)} \mathcal{C}_{07}, \quad \mathcal{C}_{39}^{-it^{-1}, it^{-1}} \xrightarrow{(ite_1 - ite_2, -t^2 e_1 - t^2 e_2)} \mathcal{C}_{13}^1, \quad \mathcal{C}_{38}^{0,0,\alpha} \xrightarrow{(ite_1 - ite_2, -t^2 e_1 - t^2 e_2)} \mathcal{C}_{15}^\alpha, \\ \mathcal{C}_{38}^{\frac{i}{2t}, 0, \frac{i}{t}} \xrightarrow{(ite_1 - ite_2, -t^2 e_1 - t^2 e_2)} \mathcal{C}_{16}^0, \quad \mathcal{C}_{38} \xrightarrow{\left(-\frac{1+at^2}{4t^2}, -\frac{1+at^2}{4t^2}, \frac{at^2-3}{4t^2}\right)} \xrightarrow{(ite_1 - ite_2, -t^2 e_1 - t^2 e_2)} \mathcal{C}_{18}^\alpha, \end{array}$$

$$\begin{aligned}
\mathcal{C}_{38}^{0,-t,\beta+t} &\xrightarrow{(e_1, te_2)} \mathcal{C}_{24}^{0,\beta,0}, \quad \mathcal{C}_{38}^{0,0,\beta} \xrightarrow{(e_1, te_2)} \mathcal{C}_{25}^{0,\beta,0}, \quad \mathcal{C}_{39}^{\alpha,t^{-1}} \xrightarrow{(e_1, te_2)} \mathcal{C}_{29}^{\alpha}, \quad \mathcal{C}_{39}^{-t+\beta,\beta} \xrightarrow{(e_1+e_2, te_2)} \mathcal{C}_{31}^{1,\beta,\beta}, \\
&\quad \mathcal{C}_{38}^{-\alpha-\beta t+t^{-1},-\beta t,t^{-1}} \xrightarrow{(e_1+e_2, te_2)} \mathcal{C}_{30}^{\alpha,\beta}, \quad \mathcal{C}_{38}^{0,\beta,0} \xrightarrow{(e_1+e_2, te_2)} \mathcal{C}_{32}^{1,\beta,\beta}, \\
&\quad \mathcal{C}_{38}^{-t^{-2},\alpha-(1+\beta t)t^{-2},(1+\beta t)t^{-2}} \xrightarrow{(e_1+e_2, te_2)} \mathcal{C}_{34}^{\alpha,\beta}, \quad \mathcal{C}_{38}^{\frac{1+2\alpha t}{2t},\alpha,0} \xrightarrow{(e_1+e_2, -te_1+te_2)} \mathcal{C}_{35}^{\alpha}.
\end{aligned}$$

□

2.3. The geometric classification of compatible associative algebras. The main result of the present section is the following theorem.

Theorem 20. *The variety of complex 2-dimensional compatible associative algebras has dimension 7 and it has 4 irreducible components defined by $\overline{\mathcal{O}(\mathcal{C}_{28})}$, $\overline{\mathcal{O}(\mathcal{C}_{33})}$, $\overline{\mathcal{O}(\mathcal{C}_{39}^{\alpha,\beta})}$ and $\overline{\mathcal{O}(\mathcal{C}_{38}^{\alpha,\beta,\gamma})}$. In particular, there are two rigid algebras in this variety.*

Proof. Thanks to Theorem 10, we have the algebraic classification of 2-dimensional compatible associative algebras. After carefully checking the dimensions of orbit closures of the more important for us algebras, we have

$$\dim \mathcal{O}(\mathcal{C}_{38}^{\alpha,\beta,\gamma}) = 7, \quad \dim \mathcal{O}(\mathcal{C}_{39}^{\alpha,\beta}) = 6, \quad \dim \mathcal{O}(\mathcal{C}_{28}) = \dim \mathcal{O}(\mathcal{C}_{34}) = 4.$$

Thanks to Theorem 20, we have that $\mathcal{C}_{38}^{\alpha,\beta,\gamma}$ and $\mathcal{C}_{39}^{\alpha,\beta}$ give irreducible components in the variety of 2-dimensional compatible commutative associative algebras. The rest of the necessary degenerations are given below.

$$\mathcal{C}_{28} \xrightarrow{(e_2, te_1)} \mathcal{C}_{05}^1, \quad \mathcal{C}_{33} \xrightarrow{(e_2, te_1)} \mathcal{C}_{06}^0, \quad \mathcal{C}_{28} \xrightarrow{(e_1+\beta e_2, te_2)} \mathcal{C}_{25}^{1,\beta,\beta}, \quad \mathcal{C}_{33} \xrightarrow{(e_1+\beta e_2, te_2)} \mathcal{C}_{32}^{0,\beta,0}.$$

□

2.4. The geometric classification of compatible Novikov algebras. The main result of the present section is the following theorem.

Theorem 21. *The variety of complex 2-dimensional compatible Novikov algebras has dimension 7 and it has 6 irreducible components defined by $\overline{\mathcal{O}(\mathcal{C}_{33})}$, $\overline{\mathcal{O}(\mathcal{C}_{09}^{\alpha,\beta})}$, $\overline{\mathcal{O}(\mathcal{C}_{22}^{\alpha,\beta})}$, $\overline{\mathcal{O}(\mathcal{C}_{39}^{\alpha,\beta})}$, $\overline{\mathcal{O}(\mathcal{C}_{31}^{\alpha,\beta,\gamma})}$ and $\overline{\mathcal{O}(\mathcal{C}_{38}^{\alpha,\beta,\gamma})}$. In particular, there is one rigid algebra in this variety.*

Proof. Thanks to Theorem 12, we have the algebraic classification of 2-dimensional compatible Novikov algebras. After carefully checking the dimensions of orbit closures of the more important for us algebras, we have

$$\begin{aligned}
\dim \mathcal{O}(\mathcal{C}_{38}^{\alpha,\beta,\gamma}) &= 7, \\
\dim \mathcal{O}(\mathcal{C}_{39}^{\alpha,\beta}) = \dim \mathcal{O}(\mathcal{C}_{09}^{\alpha,\beta}) = \dim \mathcal{O}(\mathcal{C}_{22}^{\alpha,\beta}) = \dim \mathcal{O}(\mathcal{C}_{31}^{\alpha,\beta,\gamma}) &= 6, \\
\dim \mathcal{O}(\mathcal{C}_{33}) &= 4.
\end{aligned}$$

Thanks to Theorem 21, we have that $\mathcal{C}_{38}^{\alpha,\beta,\gamma}$ and $\mathcal{C}_{39}^{\alpha,\beta}$ give irreducible components in the variety of 2-dimensional compatible commutative associative algebras. Algebras $\mathcal{C}_{09}^{\alpha,\beta}$, $\mathcal{C}_{22}^{\alpha,\beta}$, and $\mathcal{C}_{31}^{\alpha,\beta,\gamma}$ have a one-dimensional subalgebra with zero multiplication (concerning both multiplications), but \mathcal{C}_{33} has not. Hence, \mathcal{C}_{33} is not in the orbit closure of $\mathcal{C}_{09}^{\alpha,\beta}$, $\mathcal{C}_{22}^{\alpha,\beta}$, and \mathcal{C}_{31} . The rest of the necessary degenerations are given below.

$$\begin{aligned}
& \mathcal{C}_{31}^{t, \frac{t+2}{2t}, -\frac{1}{2}} \xrightarrow{(te_1 + \frac{2t}{3}e_2, \frac{t^2}{2}e_1 + t^2e_2)} \mathcal{C}_{01}, \quad \mathcal{C}_{31}^{0,0,t^{-1}} \xrightarrow{(t^2e_1 + e_2, te_1 - t^2e_2)} \mathcal{C}_{04}, \\
& \mathcal{C}_{31}^{0,t^{-1},at^{-1}} \xrightarrow{(te_1 - \alpha^{-1}t^2e_2, te_2)} \mathcal{C}_{06}^{\alpha \neq 0}, \quad \mathcal{C}_{09}^{\frac{(2\alpha+t)\beta}{2\beta+t}, -\frac{\beta t}{2\beta+t}} \xrightarrow{(\frac{2\beta+t}{2\beta}e_1 - \frac{(2\beta+t)^2}{2\beta t}e_2, \frac{(2\beta+t)t}{4\beta^2}te_1)} \mathcal{C}_{10}^{\alpha, \beta \neq 0}, \\
& \mathcal{C}_{31}^{1,at^{-1},t^{-1}} \xrightarrow{(te_1 - t^2e_2, -t^3e_2)} \mathcal{C}_{13}^{\alpha}, \quad \mathcal{C}_{31}^{1,t^{-1},0} \xrightarrow{(te_1 + \alpha^{-1}te_2, \alpha^{-1}t^2e_2)} \mathcal{C}_{14}^{\alpha \neq 0}, \\
& \mathcal{C}_{31}^{t^{-1}-t,0,-1+\alpha t^{-1}} \xrightarrow{(t^2e_1 + \frac{t^2}{1-\alpha t}e_2, te_1 + \frac{t^3}{1-\alpha t}e_2)} \mathcal{C}_{20}^{\alpha,0}, \quad \mathcal{C}_{31}^{t^{-1}-t,0,-\alpha t^{-1}} \xrightarrow{(t^2e_1 - \alpha^{-1}te_2, te_1 - \alpha^{-1}te_2)} \mathcal{C}_{21}^{\alpha,0}, \\
& \mathcal{C}_{31}^{t^{-1}-t,at^{-1},\beta t^{-1}} \xrightarrow{(t^2e_1 - \beta^{-1}te_2, te_1 - \beta^{-1}t^2e_2)} \mathcal{C}_{23}^{\alpha,\beta}, \quad \mathcal{C}_{31}^{\alpha,\beta,\gamma} \xrightarrow{(e_1, t^{-1}e_2)} \mathcal{C}_{32}^{\alpha,\beta,\gamma}.
\end{aligned}$$

□

2.5. The geometric classification of compatible pre-Lie algebras. The main result of the present section is the following theorem.

Theorem 22. *The variety of complex 2-dimensional compatible pre-Lie algebras has dimension 7 and it has 14 irreducible components defined by*

$$\begin{aligned}
& \overline{\mathcal{O}(\mathcal{C}_{28})}, \overline{\mathcal{O}(\mathcal{C}_{33})}, \overline{\mathcal{O}(\mathcal{C}_{37}^{\alpha})}, \overline{\mathcal{O}(\mathcal{C}_{09}^{\alpha,\beta})}, \overline{\mathcal{O}(\mathcal{C}_{11}^{\alpha,\beta})}, \overline{\mathcal{O}(\mathcal{C}_{22}^{\alpha,\beta})}, \overline{\mathcal{O}(\mathcal{C}_{27}^{\alpha,\beta})}, \overline{\mathcal{O}(\mathcal{C}_{36}^{\alpha,\beta})}, \overline{\mathcal{O}(\mathcal{C}_{39}^{\alpha,\beta})}, \\
& \overline{\mathcal{O}(\mathcal{C}_{41}^{\alpha,\beta})}, \overline{\mathcal{O}(\mathcal{C}_{24}^{\alpha,\beta,\gamma})}, \overline{\mathcal{O}(\mathcal{C}_{31}^{\alpha,\beta,\gamma})}, \overline{\mathcal{O}(\mathcal{C}_{38}^{\alpha,\beta,\gamma})} \text{ and } \overline{\mathcal{O}(\mathcal{C}_{40}^{\alpha,\beta,\gamma})}.
\end{aligned}$$

In particular, there are two rigid algebras in this variety.

Proof. Thanks to Theorem 7, we have the algebraic classification of 2-dimensional compatible pre-Lie algebras. After carefully checking the dimensions of orbit closures of the more important for us algebras, we have

$$\begin{aligned}
\dim \mathcal{O}(\mathcal{C}_{24}^{\alpha,\beta,\gamma}) &= \dim \mathcal{O}(\mathcal{C}_{31}^{\alpha,\beta,\gamma}) = \dim \mathcal{O}(\mathcal{C}_{38}^{\alpha,\beta,\gamma}) = \dim \mathcal{O}(\mathcal{C}_{40}^{\alpha,\beta,\gamma}) = 7, \\
\dim \mathcal{O}(\mathcal{C}_{09}^{\alpha,\beta}) &= \dim \mathcal{O}(\mathcal{C}_{11}^{\alpha,\beta}) = \dim \mathcal{O}(\mathcal{C}_{22}^{\alpha,\beta}) = \dim \mathcal{O}(\mathcal{C}_{27}^{\alpha,\beta}) = \\
& \dim \mathcal{O}(\mathcal{C}_{36}^{\alpha,\beta}) = \dim \mathcal{O}(\mathcal{C}_{39}^{\alpha,\beta}) = \dim \mathcal{O}(\mathcal{C}_{41}^{\alpha,\beta}) = 6, \\
& \dim \mathcal{O}(\mathcal{C}_{37}^{\alpha}) = 5. \\
\dim \mathcal{O}(\mathcal{C}_{28}) &= \dim \mathcal{O}(\mathcal{C}_{33}) = 4.
\end{aligned}$$

All necessary reasons for non-degenerations are listed below.

Non-degenerations reasons			
\mathcal{C}_{24}^*	$\not\rightarrow$	$\mathcal{C}_{22}^{\alpha,\beta}, \mathcal{C}_{26}^{\alpha}, \mathcal{C}_{27}^{\alpha,\beta}, \mathcal{C}_{28}$	$\mathcal{R} = \{ c_{21}^1 = c_{21}^2 = c_{22}^1 = c_{22}^2 = c_{22}^1 = c_{22}^2 = 0 \}$
\mathcal{C}_{31}^*	$\not\rightarrow$	$\mathcal{C}_{33}, \mathcal{C}_{36}^{\alpha,\beta}, \mathcal{C}_{37}^{\alpha}$	$\mathcal{R} = \{ c_{22}^1 = c_{23}^2 = c_{22}^1 = c_{22}^2 = 0 \}$
\mathcal{C}_{36}^*	$\not\rightarrow$	$\mathcal{C}_{37}^{\alpha}$	$\mathcal{R} = \{ c_{22}^1 = c_{12}^1 = c_{12}^1 = 0 \}$
\mathcal{C}_{40}^*	$\not\rightarrow$	$\mathcal{C}_{41}^{\alpha,\beta}$	$\mathcal{R} = \{ c_{22}^1 = c_{12}^1 = c_{12}^1 = 0, 2c_{11}^1 = c_{12}^1 \}$

The rest of the non-degenerations between indicated algebras follows from Remark 18.

Thanks to Theorem 21, we have that $\mathcal{C}_{33}, \mathcal{C}_{09}^{\alpha,\beta}, \mathcal{C}_{22}^{\alpha,\beta}, \mathcal{C}_{39}^{\alpha,\beta}, \mathcal{C}_{31}^{\alpha,\beta,\gamma}$, and $\mathcal{C}_{38}^{\alpha,\beta,\gamma}$ give irreducible components in the variety of 2-dimensional compatible Novikov algebras. Hence, each 2-dimensional compatible Novikov algebras is on the orbit closure of these algebras. The rest of the necessary degenerations are given below.

$$\begin{aligned}
& \mathcal{C}_{24}^{1,1+t^{-1},-1+t^{-1}} \xrightarrow{(te_1+(-1+t)e_2, t^2e_1+te_2)} \mathcal{C}_{02}, \mathcal{C}_{24}^{1,t^{-1},at^{-1}} \xrightarrow{(te_1+\frac{t^2}{1-a}e_2, t^2e_1+e_2)} \mathcal{C}_{05}^{\alpha \neq 1}, \\
\mathcal{C}_{40}^{t^{-1},0,\frac{1}{2t}} \xrightarrow{(te_1, te_2)} \mathcal{C}_{08}, \mathcal{C}_{11}^{\alpha,\alpha+t} \xrightarrow{(e_1+\beta t^{-1}e_2, e_2)} \mathcal{C}_{12}^{\alpha,\beta}, \mathcal{C}_{24}^{-1+\alpha,t^{-1},at^{-1}} \xrightarrow{(te_1-\frac{t^2}{\alpha-1}e_2, t^2e_1-t^3e_2)} \mathcal{C}_{16}^{\alpha \neq 1}, \\
& \mathcal{C}_{24}^{-2,\alpha+t^{-1},-\alpha+t^{-1}} \xrightarrow{(te_1-\alpha^{-1}te_2, t^2e_1+2\alpha^{-1}t^2e_2)} \mathcal{C}_{17}^{\alpha \neq 0}, \\
& \mathcal{C}_{40}^{-\alpha-2t^{-2},-6\alpha-6t^{-2},-\frac{4+\alpha t^2}{6t^2}} \xrightarrow{(te_1-3te_2, \frac{t^2}{2}e_1-3t^2e_2)} \mathcal{C}_{19}^{\alpha}, \\
& \mathcal{C}_{24}^{-1-t^{-2},-(\beta+t)t^{-2},(t-\alpha)t^{-2}} \xrightarrow{(-t^3e_1+\frac{t^4+t^6}{1+\alpha t-\beta t}e_2, -t^2e_1+\frac{t^5}{\beta t-\alpha t-1}e_2)} \mathcal{C}_{20}^{\alpha,\beta}, \\
& \mathcal{C}_{24}^{-1-t^{-2},-\beta t^{-2},-\alpha t^{-2}} \xrightarrow{(-t^3e_1+\frac{t^3+t^5}{\alpha-\beta}e_2, -t^2e_1-\frac{t^4}{\alpha-\beta}e_2)} \mathcal{C}_{21}^{\alpha,\beta}, \mathcal{C}_{27}^{\alpha-5t^2,t^{-2}} \xrightarrow{(e_1+2t^2e_2, te_2)} \mathcal{C}_{26}^{\alpha}, \\
& \mathcal{C}_{24}^{\alpha,\beta,\gamma} \xrightarrow{(e_1, t^{-1}e_2)} \mathcal{C}_{25}^{\alpha,\beta,\gamma}.
\end{aligned}$$

□

REFERENCES

- [1] Abdelwahab H., Barreiro E., Calderón A., Fernández Ouaridi A., The algebraic classification and degenerations of nilpotent Poisson algebras, *Journal of Algebra*, 615 (2023), 243–277.
- [2] Abdelwahab H., Fernández Ouaridi A., Kaygorodov I., Degenerations of Poisson-type algebras, arXiv:2403.17193.
- [3] Abdelwahab H., Fernández Ouaridi A., Martín González C., Degenerations of Poisson algebras, *Journal of Algebra and Its Applications*, 2023, DOI: 10.1142/S0219498825500872.
- [4] Alvarez M., Kaygorodov I., The algebraic and geometric classification of nilpotent weakly associative and symmetric Leibniz algebras, *Journal of Algebra*, 588 (2021), 278–314.
- [5] Alvarez M., Fehlberg Júnior R., Kaygorodov I., The algebraic and geometric classification of Zinbiel algebras, *Journal of Pure and Applied Algebra*, 226 (2022), 11, 107106.
- [6] Beneš T., Burde D., Degenerations of pre-Lie algebras, *Journal of Mathematical Physics*, 50 (2009), 11, 112102.
- [7] Beneš T., Burde D., Classification of orbit closures in the variety of three-dimensional Novikov algebras, *Journal of Algebra and Its Applications*, 13 (2014), 2, 1350081.
- [8] Burde D., Left-symmetric algebras, or pre-Lie algebras in geometry and physics, *Central European Journal of Mathematics*, 4 (2006), 3, 323–357.
- [9] Chapoton F., Livernet M., Pre-Lie algebras and the rooted trees operad, *International Mathematics Research Notices*, (2001), 8, 395–408.
- [10] Chtioui T., Das A., Mabrouk S., (Co)homology of compatible associative algebras, *Communications in Algebra*, 52 (2024), 2, 582–603.
- [11] Das A., Compatible L_∞ -algebras, *Journal of Algebra*, 610 (2022), 241–269.
- [12] Das A., Cohomology and deformations of compatible Hom-Lie algebras, *Journal of Geometry and Physics*, 192 (2023), 104951.
- [13] Das A., Guo S., Qin Y., L_∞ -structures and cohomology theory of compatible \mathcal{O} -operators and compatible dendriform algebras, *Journal of Mathematical Physics*, 65 (2024), 3, 031701.
- [14] Dotsenko V., Compatible associative products and trees, *Algebra & Number Theory*, 3 (2009), 5, 567–586.
- [15] Dotsenko V., Ismailov N., Umirbaev U., Polynomial identities in Novikov algebras, *Mathematische Zeitschrift*, 303 (2023), 3, 60.
- [16] Ehret Q., Makhlof A., On classification and deformations of Lie-Rinehart superalgebras, *Communications in Mathematics*, 30 (2022), 2, 67–100.
- [17] Eick B., Moede T., Computing subalgebras and \mathbb{Z}_2 -gradings of simple Lie algebras over finite fields, *Communications in Mathematics*, 30 (2022), 2, 37–50.
- [18] Fernández Ouaridi A., Kaygorodov I., Khrypchenko M., Volkov Yu., Degenerations of nilpotent algebras, *Journal of Pure and Applied Algebra*, 226 (2022), 3, 106850.

- [19] Hou S., Sheng Y., Zhou Y., Deformations, cohomologies and abelian extensions of compatible 3-Lie algebras, *Journal of Geometry and Physics*, 202 (2024), 105218.
- [20] Golubchik I., Sokolov V., Compatible Lie brackets and integrable equations of the principal chiral field model type, *Functional Analysis and Its Applications*, 36 (2002), 3, 172–181
- [21] Golubchik I., Sokolov V., Compatible Lie brackets and the Yang-Baxter equation, *Theoretical and Mathematical Physics*, 146 (2006), 2, 159–169
- [22] González D'León R., On the free Lie algebra with multiple brackets, *Advances in Applied Mathematics*, 79 (2016), 37–97.
- [23] Grunewald F., O'Halloran J., Varieties of nilpotent Lie algebras of dimension less than six, *Journal of Algebra*, 112 (1988), 315–325.
- [24] Kaygorodov I., Non-associative algebraic structures: classification and structure, *Communications in Mathematics*, 32 (2024), 3, 1–62.
- [25] Kaygorodov I., Khrypchenko M., Lopes S., The geometric classification of nilpotent algebras, *Journal of Algebra*, 633 (2023), 857–886.
- [26] Kaygorodov I., Khrypchenko M., Páez-Guillán P., The geometric classification of non-associative algebras: a survey, preprint
- [27] Kaygorodov I., Volkov Y., The variety of two-dimensional algebras over an algebraically closed field, *Canadian Journal of Mathematics*, 71 (2019), 4, 819–842.
- [28] Ladra M., Leite da Cunha B., Lopes S., A classification of nilpotent compatible Lie algebras, [arXiv:2406.04036](https://arxiv.org/abs/2406.04036).
- [29] Liu F., Combinatorial bases for multilinear parts of free algebras with two compatible brackets, *Journal of Algebra*, 323 (2010), 1, 132–166.
- [30] Liu J., Sheng Y., Bai C., F -manifold algebras and deformation quantization via pre-Lie algebras, *Journal of Algebra*, 559 (2020), 467–495.
- [31] Liu J., Sheng Y., Bai C., Maurer-Cartan characterizations and cohomologies of compatible Lie algebras, *Science China Mathematics*, 66 (2023), 6, 1177–1198.
- [32] Lopes S., Noncommutative algebra and representation theory: symmetry, structure & invariants, *Communications in Mathematics*, 32 (2024), 2, 63–117.
- [33] Odesskii A., Sokolov V., Compatible Lie brackets related to elliptic curve, *Journal of Mathematical Physics*, 47 (2006), 1, 013506.
- [34] Odesskii A., Sokolov V., Algebraic structures connected with pairs of compatible associative algebras, *International Mathematics Research Notices*, (2006), 43734.
- [35] Odesskii A., Sokolov V., Integrable matrix equations related to pairs of compatible associative algebras, *Journal of Physics A*, 39 (2006), 40, 12447–12456.
- [36] Odesskii A., Sokolov V., Pairs of compatible associative algebras, classical Yang-Baxter equation and quiver representations, *Communications in Mathematical Physics*, 278 (2008), 1, 83–99.
- [37] Shalev A., Smoktunowicz A., From braces to pre-Lie rings, *Proceedings of the American Mathematical Society*, 152 (2024), 4, 1545–1559.
- [38] Smoktunowicz A., On the passage from finite braces to pre-Lie rings, *Advances in Mathematics*, 409 (2022), part B, 108683.
- [39] Strohmayr H., Operads of compatible structures and weighted partitions, *Journal of Pure and Applied Algebra*, 212 (2008), 11, 2522–2534.
- [40] Zhang H., Gao X., Guo L., Compatible structures of nonsymmetric operads, Manin products and Koszul duality, *Applied Categorical Structures*, 32 (2024), 1, 2.