

SMALL BALL PROBABILITIES FOR THE PASSAGE TIME IN PLANAR FIRST-PASSAGE PERCOLATION

DOR ELBOIM

ABSTRACT. We study planar first-passage percolation with independent weights whose common distribution is supported on $(0, \infty)$ and is absolutely continuous with respect to Lebesgue measure. We prove that the passage time from x to y denoted by $T(x, y)$ satisfies

$$\max_{a \geq 0} \mathbb{P}(T(x, y) \in [a, a + 1]) \leq \frac{C}{\sqrt{\log \|x - y\|}},$$

answering a question posed by of Ahlberg and de la Riva [1]. This estimate recovers earlier results on the fluctuations of the passage time by Newman and Piza [11], Pemantle and Peres [12] and Chatterjee [5].

1. INTRODUCTION

First-passage percolation is a model for a random metric space, formed by a random perturbation of an underlying base space. Since its introduction by Hammersley–Welsh in 1965 [9], it has been studied extensively in the probability and statistical physics literature. We refer to [10] for general background and to [3] for more recent results.

We study first-passage percolation on the square lattice $(\mathbb{Z}^2, E(\mathbb{Z}^2))$, in an independent and identically distributed (IID) random environment. The model is specified by a *weight distribution* G , which is a probability measure on the non-negative reals. It is defined by assigning each edge $e \in E(\mathbb{Z}^2)$ a random passage time t_e with distribution G , independently between edges. Then, each finite path p in \mathbb{Z}^2 is assigned the random passage time

$$T(p) := \sum_{e \in p} t_e,$$

yielding a random metric T on \mathbb{Z}^2 by setting the *passage time between* $u, v \in \mathbb{Z}^2$ to

$$T(u, v) := \inf_p T(p), \tag{1.1}$$

where the infimum ranges over all finite paths connecting u and v . Any path achieving the infimum is termed a *geodesic* between u and v . A unique geodesic exists when G is atomless and will be denoted $\gamma(u, v)$. The focus of first-passage percolation is the study of the large-scale properties of the random metric T and its geodesics.

Ahlberg and de la Riva [1, Equation (11) and Section 7] raised the problem of proving that the passage time between far away points is unlikely to be in any given interval of constant length. More precisely, they conjectured that for all $C > 0$ one has

$$\max_{a > 0} \mathbb{P}(T(0, (n, 0)) \in [a, a + C]) \rightarrow 0, \quad n \rightarrow \infty.$$

In our main result below we establish a quantitative version of this conjecture for the class of absolutely continuous weight distributions.

Theorem 1.1. *Suppose that the weight distribution G is absolutely continuous. Then, there exists a constant $C > 0$ depending on G such that for any $x \in \mathbb{Z}^2$ with $\|x\| \geq 2$*

$$\max_{a \geq 0} \mathbb{P}(T(0, x) \in [a, a + 1]) \leq \frac{C}{\sqrt{\log \|x\|}}. \quad (1.2)$$

1.1. previous works. Pemantle and Peres [12] proved an estimate similar to (1.2) when the weight distribution G is exponential. The proof relies on the memoryless property of the exponential distribution. We are not aware of a similar result for any other weight distribution.

The estimate in (1.2) is closely related to anti-concentration bounds for the passage time. The first result of this kind is by Newman and Piza [11] who proved that

$$\text{Var}(T(0, x)) \geq c \log \|x\|. \quad (1.3)$$

This was later extended to a wider class of weight distributions by Auffinger and Damron [2]. Chatterjee [5, Theorem 2.6] proved the stronger result that the fluctuations of the passage time are at least of order $\sqrt{\log \|x\|}$ by showing that there exists $c > 0$ such that for all $x \in \mathbb{Z}^2$ and all $a < b$ with $b - a \leq c\sqrt{\log \|x\|}$,

$$\mathbb{P}(T(0, x) \notin [a, b]) \geq c. \quad (1.4)$$

This was generalized to a wider class of weight distributions by Damron, Hanson, Houdré and Xu [6] and independently by Bates and Chatterjee [4]. Observe that both (1.3) and (1.4) follow from Theorem 1.1 for the class of absolutely continuous weight distributions.

Finally, let us remark that the perturbation argument used in this paper is similar to that of Chatterjee [5] with one notable difference. In [5] the author considers a single perturbation of the weights while in here we are continuously perturbing the weights. We then show that this continuous perturbation quickly pushes the passage time out of any given interval.

1.2. acknowledgments. We thank Barbara Dembin, Ron Peled, Daniel Ahlberg and Daniel de la Riva for fruitful discussions about this problem.

2. A MERMIN–WAGNER TYPE ESTIMATE

The following lemma is the main technical tool required for the proof of the main theorem. The lemma is a Mermin–Wagner type estimate and is taken from [7, Lemma 2.12]. A similar lemma was used also in [8].

Lemma 2.1. *Suppose that G is absolutely continuous distribution on \mathbb{R} . Then, there exist*

- *A Borel set $S_G \subseteq \mathbb{R}$ with $G(S_G) = 1$.*
- *Borel subsets $(B_\delta)_{\delta > 0}$ of S_G with $\lim_{\delta \downarrow 0} G(B_\delta) = 1$,*
- *For each $\tau \in \mathbb{R}$, a bijection $g_\tau : S_G \rightarrow S_G$.*

such that the following holds:

- (1) *The function $g_\tau(s)$ is increasing both in s and in τ .*
- (2) *g_0 is the identity function on S_G and for any $\tau_1, \tau_2 \in \mathbb{R}$ we have $g_{\tau_1} \circ g_{\tau_2} = g_{\tau_1 + \tau_2}$.*
- (3) *For any $\tau \in [0, 1]$ and $s \in B_\delta$ we have $g_\tau(s) \geq s + \delta\tau$.*
- (4) *For any integer $n \geq 1$, a vector $\tau \in \mathbb{R}^n$ and a Borel set $A \subset \mathbb{R}^n$ we have*

$$\mathbb{P}(X \in A) \leq e^{\|\tau\|_2^2/2} \sqrt{\mathbb{P}((g_{\tau_i}(X_i))_{i=1}^n \in A) \mathbb{P}((g_{-\tau_i}(X_i))_{i=1}^n \in A)} \quad (2.1)$$

where $X = (X_1, \dots, X_n)$ is a vector of i.i.d. random variables with distribution G .

Proof sketch. Let us briefly explain the main ideas in the proof of the lemma while the full argument is given in [7]. First, we consider the case when $G = N(0, 1)$ is the standard normal distribution. In this case the functions g_τ are simply taken to be $g_\tau(s) := s + \tau$. With this choice the inequality (2.1) translates to

$$\mathbb{P}(X \in A) \leq e^{\|\tau\|_2^2/2} \sqrt{\mathbb{P}(X + \tau \in A)\mathbb{P}(X - \tau \in A)} \quad (2.2)$$

which follows from a simple application of Cauchy-Schwarz inequality (see [7, Claim 2.13]).

For a general continuous distribution G , we consider the (unique) increasing function $h : \mathbb{R} \rightarrow S_G$ with $h(N) \sim G$ where N is a standard normal variable. Then, we define the function g_τ by $g_\tau(s) := h(h^{-1}(s) + \tau)$ and prove that this function satisfies the requirements (the inequality (2.1) follows immediately as it is the push forward of (2.2) by h).

Finally, let us note that the statement of [7, Lemma 2.12] does not include parts (1) and (2) of Lemma 2.1. However, these parts follow immediately from our definition of the perturbation function $g_\tau(s) = h(h^{-1}(s) + \tau)$. \square

3. PROOF OF THEOREM 1.1

Let $x \in \mathbb{Z}^2$ with $\|x\|$ is sufficiently large and let $n := \|x\|$. Let Λ_k be the set of edges with both endpoints in the annulus $[-2^{k+1}, 2^{k+1}]^2 \setminus [-2^k, 2^k]^2$ or one endpoint in the annulus and the other in $[-2^k, 2^k]$. Let $k_0 := \lfloor \log_2 \sqrt{n} \rfloor$ and $k_1 := \lfloor \log_2 n \rfloor - 1$.

For $r \in [-2, 2]$ we define the perturbation function $\tau_r : E(\mathbb{Z}^d) \rightarrow [0, \infty)$ by

$$\tau_r(e) := \frac{r}{2^k \sqrt{\log n}},$$

for all $e \in \Lambda_k$ with $k_0 \leq k \leq k_1$. We let $\tau_r(e) = 0$ for edges outside of $\bigcup_{k=k_0}^{k_1} \Lambda_k$.

We define the modified environment by $t_{e,r} := g_{\tau_r(e)}(t_e)$. We denote by $T_r(p)$ the weight a path p in the modified environment and by $T_r(x, y)$ the passage between x and y in the modified environment. In our proof we continuously vary r and analyse the effect it has on the passage time $T(0, x)$.

It is important that for all $r \in [-1, 1]$ we have

$$\|\tau_r\|_2^2 \leq \sum_{k=k_0}^{k_1} \frac{|\Lambda_k|}{4^k \log n} \leq \frac{C(k_1 - k_0)}{\log n} \leq C$$

so that in Lemma 2.1 the factor $e^{\|\tau\|_2^2/2}$ remains constant.

We begin by showing that the weight of paths in scale k is increased by this perturbation. To this end let \mathcal{P}_k be the set of paths of length 2^k whose edges are in Λ_k and let δ_0 such that $G(B_{\delta_0}) \geq 0.999$.

Lemma 3.1. *For any $k_0 \leq k \leq k_1$ and $r \in [0, 1]$ we have that*

$$\mathbb{P}\left(\exists p \in \mathcal{P}_k, T_r(p) - T(p) \leq \frac{\delta_0 r}{2\sqrt{\log n}}\right) \leq e^{-2^k}.$$

Proof. Let $k_0 \leq k \leq k_1$ and $r \in [0, 1]$. Recall that if $t_e \in B_{\delta_0}$ then

$$t_{e,r} = g_{\tau_r(e)}(t_e) \geq t_e + \delta_0 \tau_r(e) = t_e + \frac{\delta_0 r}{2^k \sqrt{\log n}}.$$

Thus, for a fixed path $p \in \mathcal{P}_k$ we have

$$\begin{aligned} \mathbb{P}\left(T_r(p) - T(p) \leq \frac{\delta_0 r}{2\sqrt{\log n}}\right) &\leq \mathbb{P}(|\{e \in p : t_e \in B_{\delta_0}\}| \leq 2^k/2) \leq \\ &\mathbb{P}(\text{Bin}(2^k, 0.999) \leq 2^k/2) \leq 8^{-2^k}. \end{aligned}$$

Moreover, we have that $|\mathcal{P}_k| \leq C4^k 3^{2^k}$ as we can first choose the starting point of $p \in \mathcal{P}_k$ and then choose each step of the path. The lemma follows from a union bound over $p \in \mathcal{P}_k$ as long as n is sufficiently large. \square

Corollary 3.2. *For any $k_0 \leq k \leq k_1$, $s \in [-1, 1]$ and $r \in [0, 1]$ we have that*

$$\mathbb{P}\left(\exists p \in \mathcal{P}_k, T_{s+r}(p) - T_s(p) \leq \frac{\delta_0 r}{2\sqrt{\log n}}\right) \leq C e^{-2^{k-1}}.$$

Proof. The corollary follows from Lemma 3.1 using Lemma 2.1. Let us note that this will not be the main use of Lemma 2.1 and that this part can be done using different arguments.

The set $A \subseteq \mathbb{R}^{\Lambda_k}$ from Lemma 2.1 will be

$$A := \left\{ (z_e)_{e \in \Lambda_k} : \exists p \in \mathcal{P}_k, \sum_{e \in p} g_{\tau_{s+r}(e)}(z_e) - g_{\tau_s(e)}(z_e) \leq \frac{\delta_0 r}{2\sqrt{\log n}} \right\}.$$

By Lemma 2.1 and the fact that $\|\tau_s\|_2 \leq C$ we have that

$$\mathbb{P}((t_e)_{e \in \Lambda_k} \in A) \leq C \sqrt{\mathbb{P}((g_{-\tau_s(e)}(t_e))_{e \in \Lambda_k} \in A)}.$$

It is easy to check (using part (2) of Lemma 2.1) that the event on the right hand side of the last inequality is precisely the event of Lemma 3.1 while the event on the left hand side is the event of the corollary. \square

Lemma 3.3. *There exists $C > 0$ depending only on G such that for any $a \geq 0$ we have*

$$\int_{-1}^1 \mathbb{P}(T_r(0, x) \in [a, a+1]) dr \leq \frac{C}{\sqrt{\log n}}.$$

Proof. Let $r_0 := 8/(\delta_0 \sqrt{\log n})$ and let $R := r_0 \mathbb{Z} \cap [-1, 1]$. Define the event

$$\Omega := \bigcap_{k=k_0}^{k_1} \bigcap_{r \in R} \left\{ \forall p \in \mathcal{P}_k, T_{r+r_0}(p) - T_r(p) > \frac{4}{\log n} \right\}.$$

Using Corollary 3.2 and a union bound we obtain

$$\mathbb{P}(\Omega^c) \leq C \sum_{k=k_0}^{k_1} \sum_{r \in R} e^{-2^{k-1}} \leq e^{-c\sqrt{n}}. \quad (3.1)$$

Next, we write

$$\int_{-1}^1 \mathbb{P}(T_r(0, x) \in [a, a+1]) dr \leq \mathbb{P}(\Omega^c) + \int_{-1}^1 \mathbb{P}(\Omega, T_r(0, x) \in [a, a+1]) dr. \quad (3.2)$$

By (3.1), it suffices to bound the second term on the right hand side of (3.2). To this end, note that any path p connecting 0 to x will have a subpath in \mathcal{P}_k for any $k_0 \leq k \leq k_1$. It follows that on Ω for any path p connecting 0 and x and for all $r \in R$ we have $T_{r+r_0}(p) - T_r(p) \geq 4(k_1 - k_0)/\log n \geq 2$. Hence, on Ω we have $T_{r+r_0}(0, x) - T_r(0, x) \geq 2$ for all $r \in R$. Thus, using also that $T_r(0, x)$ is increasing in r we obtain that on Ω there is at most one element $r \in R$

for which $T^r(0, x) \in [a, a + 1]$ and that $|\{r \in [-1, 1] : T^r(0, x) \in [a, a + 1]\}| \leq C/\sqrt{\log n}$, where in here $|\cdot|$ denotes the Lebesgue measure of the set. Hence, using Fubini's theorem

$$\int_0^1 \mathbb{P}(\Omega, T_r(0, x) \in [a, a + 1]) dr = \mathbb{E} \left[\mathbb{1}_\Omega |\{r \in [0, 1] : T_r(0, x) \in [a, a + 1]\}| \right] \leq \frac{C}{\sqrt{\log n}}.$$

Substituting this estimate into (3.2) finishes the proof of the lemma. \square

We can now prove Theorem 1.1.

Proof of Theorem 1.1. We claim that for any $r \in [0, 1]$ and $a > 0$ we have

$$\mathbb{P}(T(0, x) \in [a, a + 1]) \leq C \sqrt{\mathbb{P}(T_r(0, x) \in [a, a + 1]) \mathbb{P}(T_{-r}(0, x) \in [a, a + 1])}. \quad (3.3)$$

Unfortunately, this is not a direct use of Lemma 2.1 since $T(0, x)$ depends on infinitely many random variables while in Lemma 2.1 there are finitely many. Let us briefly explain how to overcome this issue using an approximation argument. For $R > n$ define the restricted passage time $T^R(0, x)$ exactly as in (1.1) but when the infimum is over paths restricted to stay in the box $[-R, R]^2$. Note that the restricted passage time depends only on the variables in the box $[-R, R]^2$. Similarly, we define the restricted passage times $T_r^R(0, x)$ and $T_{-r}^R(0, x)$ in the modified environments. Using Lemma 2.1 we obtain for all $R > n$

$$\mathbb{P}(T^R(0, x) \in [a, a + 1]) \leq e^{\|\tau_r\|^2/2} \sqrt{\mathbb{P}(T_r^R(0, x) \in [a, a + 1]) \mathbb{P}(T_{-r}^R(0, x) \in [a, a + 1])}. \quad (3.4)$$

Moreover, it is easy to check that for all $\epsilon > 0$ there exists R such that

$$\mathbb{P}(T^R(0, x) \neq T(0, x)) \leq \mathbb{P}(\gamma(0, x) \not\subseteq [-R, R]^2) \leq \epsilon.$$

Similarly, for R sufficiently large

$$\mathbb{P}(T_r^R(0, x) \neq T_r(0, x)) \leq \epsilon \quad \text{and} \quad \mathbb{P}(T_{-r}^R(0, x) \neq T_{-r}(0, x)) \leq \epsilon.$$

We can therefore replace the restricted passage times in (3.4) with the non restricted passage times and this will change the probabilities by at most ϵ . Taking ϵ to zero finishes the proof of (3.3).

Integrating (3.3) over $r \in [0, 1]$ and using Cauchy-Schwartz inequality we obtain

$$\mathbb{P}(T(0, x) \in [a, a + 1]) \leq C \sqrt{\int_0^1 \mathbb{P}(T_r(0, x) \in [a, a + 1]) dr \int_0^1 \mathbb{P}(T_{-r}(0, x) \in [a, a + 1]) dr}.$$

By Lemma 3.3 the right hand side of the last equation is bounded by $C/\sqrt{\log n}$ which finishes the proof of the theorem. \square

REFERENCES

- [1] Daniel Ahlberg and Daniel de la Riva. Is' being above the median'a noise sensitive property? *arXiv preprint arXiv:2308.16388*, 2023.
- [2] Antonio Auffinger and Michael Damron. Differentiability at the edge of the percolation cone and related results in first-passage percolation. *Probability Theory and Related Fields*, 156(1):193–227, 2013.
- [3] Antonio Auffinger, Michael Damron, and Jack Hanson. *50 years of first-passage percolation*, volume 68 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2017.
- [4] Erik Bates and Sourav Chatterjee. Fluctuation lower bounds in planar random growth models. 2020.
- [5] Sourav Chatterjee. A general method for lower bounds on fluctuations of random variables. *The Annals of Probability*, 47(4):2140–2171, 2019.
- [6] Michael Damron, Jack Hanson, Christian Houdré, and Chen Xu. Lower bounds for fluctuations in first-passage percolation for general distributions. 2020.

- [7] Barbara Dembin, Dor Elboim, and Ron Peled. Coalescence of geodesics and the BKS midpoint problem in planar first-passage percolation. *arXiv preprint arXiv:2204.02332*, 2022.
- [8] Barbara Dembin, Dor Elboim, and Ron Peled. On the influence of edges in first-passage percolation on \mathbb{Z}^d . *arXiv preprint arXiv:2307.01162*, 2023.
- [9] J. M. Hammersley and D. J. A. Welsh. First-passage percolation, subadditive processes, stochastic networks, and generalized renewal theory. In *Proc. Internat. Res. Semin., Statist. Lab., Univ. California, Berkeley, Calif*, pages 61–110. Springer-Verlag, New York, 1965.
- [10] Harry Kesten. Aspects of first passage percolation. In *École d'été de probabilités de Saint-Flour, XIV—1984*, volume 1180 of *Lecture Notes in Math.*, pages 125–264. Springer, Berlin, 1986.
- [11] Charles M Newman and Marcelo ST Piza. Divergence of shape fluctuations in two dimensions. *The Annals of Probability*, pages 977–1005, 1995.
- [12] Robin Pemantle and Yuval Peres. Planar first-passage percolation times are not tight. *Probability and phase transition*, pages 261–264, 1994.

DOR ELBOIM

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, NEW JERSEY, UNITED STATES.

Email address: delboim@ias.edu