Local wealth condensation for yard-sale models with wealth-dependent biases

Christoph Börgers¹ and Claude Greengard²

¹ Department of Mathematics, Tufts University, Medford, MA

² Two Sigma Investments, LP, New York, NY, and

Courant Institute of Mathematical Sciences, New York University, New York, NY

Abstract. In Chakraborti's *yard-sale model* of an economy [6], identical agents engage in pairwise trades, resulting in wealth exchanges that conserve each agent's expected wealth. Doob's martingale convergence theorem immediately implies almost sure wealth condensation, *i.e.*, convergence to a state in which a single agent owns the entire economy. If some pairs of agents are not allowed to trade with each other, the martingale convergence theorem still implies *local* wealth condensation, *i.e.*, convergence to a state in which some agents are wealthy, while all their trading partners are impoverished. In this note, we propose a new, more elementary proof of this result. Unlike the proof based on the martingale convergence theorem, our argument applies to models with a wealth-acquired advantage, and even to certain models with a *poverty*-acquired advantage.

1 Introduction

In Chakraborti's *yard-sale model* [6] of the economy, *N* identical agents engage in pairwise trades. After ℓ trades, the *i*-th agent owns the fraction $X_{\ell}^{i} \in (0, 1)$ of the total economy. We write

$$X_{\ell} = [X_{\ell}^{i}]_{1 \le i \le N}, \quad \ell = 0, 1, 2, \dots$$

and note that

$$\sum_{i=1}^{N} X_{\ell}^{i} = 1 \quad \text{for all } \ell.$$

We think of X_0 as deterministic and given. The X_{ℓ} with $\ell \ge 1$ are random. In the original version of the model proposed by Chakraborti, they are defined inductively as follows. Choose a number $b \in (0,1)$. Given $X_{\ell-1}$, choose a uniformly distributed random pair of integers

$$(i_{\ell}, j_{\ell}) \in \{(i, j) : 1 \le i, j \le N, i \ne j\},\$$

and define $(\mu_{\ell}, \nu_{\ell}) = (i_{\ell}, j_{\ell})$ if $X_{\ell-1}^{i_{\ell}} \leq X_{\ell-1}^{j_{\ell}}$, and $(\mu_{\ell}, \nu_{\ell}) = (j_{\ell}, i_{\ell})$ otherwise, so that in any case $X_{\ell-1}^{\mu_{\ell}} \leq X_{\ell-1}^{\nu_{\ell}}$. Choose a random $s_{\ell} \in \{-1, 1\}$ with $P(s_{\ell} = 1) = P(s_{\ell} = -1) = \frac{1}{2}$. Assume that (i_{ℓ}, j_{ℓ}) and s_{ℓ} are independent of each other, and independent of the X_k , (i_k, j_k) , and s_k with $k < \ell$. Set

$$X_{\ell}^{\mu_{\ell}} = X_{\ell-1}^{\mu_{\ell}} - s_{\ell} b X_{\ell-1}^{\mu_{\ell}}, \quad X_{\ell}^{\nu_{\ell}} = X_{\ell-1}^{\nu_{\ell}} + s_{\ell} b X_{\ell-1}^{\mu_{\ell}},$$
(1)
and $X_{\ell}^{i} = X_{\ell-1}^{i}$ for $i \notin \{\mu_{\ell}, \nu_{\ell}\}.$

Thus a fraction b of the poorer agent's wealth is transferred from one agent to the other, with the direction of transfer determined by a fair coin flip. This model is known to have the following striking property.

Theorem 1 (Yard-Sale Convergence Theorem). *The vectors* X_{ℓ} *converge to a canonical basis vector of* \mathbb{R}^N *almost surely.*

In the limit, one agent comes to own everything. This sort of maximal inequality is called *wealth condensation*. In the yard-sale model, wealth condensation is the inescapable result of random, statistically unbiased interactions. Chakraborti first observed this fact numerically [6]. For a version of the model in which there is a continuum of agents, rather than a finite number, an analogous result was presented by Boghosian *et al* [2].

Chorro [7] pointed out that Theorem 1 is an immediate consequence of Doob's martingale convergence theorem: For a fixed *i*, the sequence $\{X_{\ell}^i\}_{\ell=0,1,2,...}$ is a bounded martingale, and therefore must converge almost surely. It is clear from the definition of the model that almost sure convergence of the sequence $\{X_{\ell}\}_{\ell=0,1,2,...}$ implies almost sure convergence to a canonical basis vector. Since $E(X_{\ell}^i) = E(X_0^i)$ for all ℓ and *i*, we also conclude

$$P\left(\lim_{\ell \to \infty} X_{\ell}^{i} = 1\right) = X_{0}^{i}.$$
(2)

Many variations can be analyzed similarly. For instance, the distribution of the pairs (i_{ℓ}, j_{ℓ}) need not be uniform. Different agents can be assumed to have different degrees of risk tolerance [5], and their risk tolerance may even be history-dependent. Thus the number b in (1) might be replaced by random numbers $B_{\ell} \in [\delta, 1)$, where $\delta \in (0, 1)$ is a fixed given number. The B_{ℓ} may depend on $X_0, \ldots, X_{\ell-1}$ and $(\mu_1, \nu_1), \ldots, (\mu_{\ell}, \nu_{\ell})$. If B_{ℓ} is smaller, agent μ_{ℓ} is more risk-averse during the ℓ -th trade. We do need the assumption $B_{\ell} \ge \delta$, so that risk-taking never wholly disappears. Obviously, but remarkably, eq. (2) still holds for the modified model; risk aversion does not affect an agent's likelihood of ending up owning the entire economy.

An especially interesting variation is obtained when trades are allowed only among some, but not all, pairs of agents [4]. One can interpret this as an implementation of the model on an undirected graph in which the vertices are agents, and there is an edge between vertices *i* and *j* if and only if agents *i* and *j* are allowed to trade. In Chakrobarti's original model, the graph is *complete*, *i.e.*, any two vertices are connected by an edge. When the graph is *incomplete*, there is *local* wealth condensation [4, 11]; *i.e.*, if $a_{ij} \neq 0$, then $\min\{X_{\ell}^i, X_{\ell}^j\} \rightarrow 0$ as $\ell \rightarrow \infty$. See Fig. 1 for illustration. This variation, too, can be analyzed easily using the martingale convergence theorem.

Cardoso *et al.* [5] proposed an altogether different approach to proving the yard-sale convergence theorem, based on the Gini index. Their analysis relies on what they call the *fair rule hypothesis* [5, Equation (8)]. For wealth-conserving models, it is the martingale property. Most, though not all, examples in [5] are wealth-conserving. The main result in [5] is that the Gini index is monotonically increasing, and stationary if and only if it assumes its maximal value.

Arguments based on the martingale property break down when the coin flips governing who benefits from a trade are taken to be biased. Boghosian *et al.* [1] allowed a wealth-acquired advantage: $s_{\ell} = 1$ with some probability $p \in [\frac{1}{2}, 1)$. One should certainly expect




Figure 1: Outcomes of three simulations of a yard-sale process on the same graph over many trades. The sizes of the gray dots indicate the eventual wealth of the agents. There are two wealthy agents in the left panel, three in the middle panel, and four in the right panel. The vertices representing wealthy agents are not directly connected by edges, so wealthy agents never trade with each other.

wealth condensation for $p > \frac{1}{2}$, considering that it occurs even for $p = \frac{1}{2}$. More surprisingly, Moukarzel *et al.* [12] gave a heuristic argument showing that even with a very slight *poverty*-acquired advantage, there can be wealth condensation. In these models, $\{X_{\ell}^i\}_{\ell \ge 0}$ is no longer a martingale, nor a sub- or super-martingale.

In this paper, which generalizes our preprint [3], we give a new probabilistic analysis of the results sketched above, not relying on the martingale property. We use $||X_{\ell}||^2$ as a measure of concentration, and analyze its time evolution. For any non-zero vector $X \in \mathbb{R}^N$, the ratio $||X||_2/||X||_1$ is ≤ 1 , and can be viewed as a measure of concentration. In fact, this ratio is equal to 1 if and only if X is "maximally concentrated," namely, a non-zero multiple of a canonical basis vector. This observation has found many uses, for instance in quantum physics [9], political science [10], ecology [13], and antitrust regulation [15]. It also underlies the idea of adding an L^1 penalty term in regression, the "lasso" method [14], to obtain sparse (namely, concentrated) solutions.

2 Mathematical statement of our model

We will state our precise mathematical assumptions without repeating the economic motivation already given in the introduction. We assume that we are given

- an integer $N \ge 2$,
- a vector $X_0 \in (0, 1)^N$ with $\sum_{i=1}^N X_0^i = 1$,
- an adjacency matrix $A = [a_{ij}]_{1 \le i,j \le N}$ with $a_{ii} = 0$ for all $i, a_{ij} = a_{ji} \in \{0,1\}$ for $i \ne j$, and $a_{ij} = a_{ji} = 1$ for least one pair (i, j),
- a probability distribution π on the set

$$\mathcal{E} = \{(i, j) : 1 \le i, j \le N, a_{ij} = 1\}$$

assigning to each element of \mathcal{E} a positive probability,

- a sequence of pairs $(i_{\ell}, j_{\ell}), \ell \ge 1$, independent of each other with probability distribution π ,

- sequences of random numbers $\{U_\ell\}_{\ell\geq 1}$, $\{V_\ell\}_{\ell\geq 1}$, and $\{W_\ell\}_{\ell\geq 1}$, uniformly distributed in (0,1) and independent of each other and of the pairs (i_ℓ, j_ℓ) ,
- Borel measurable functions

$$f_{\ell}: \left(\mathbb{R}^{N}\right)^{\ell} \times \mathcal{E}^{\ell} \times (0,1) \to [\delta,1), \quad \ell \geq 1,$$

where $\delta \in (0,1)$ is a fixed given number, independent of ℓ ,

- Borel measurable functions

$$g_{\ell}: \left(\mathbb{R}^{N}\right)^{\ell} \times \mathcal{E}^{\ell} \times (0,1) \to (0,1), \quad \ell \geq 1.$$

Given these data, we define random vectors $X_{\ell} \in (0,1)^N$ with $\sum_{i=1}^N X_{\ell}^i = 1$, $\ell = 1, 2, ...$, inductively as follows. Given $X_{\ell-1}$, set $(\mu_{\ell}, \nu_{\ell}) = (i_{\ell}, j_{\ell})$ if $X_{\ell-1}^{i_{\ell}} \leq X_{\ell-1}^{j_{\ell}}$, and $(\mu_{\ell}, \nu_{\ell}) = (j_{\ell}, i_{\ell})$ otherwise. In order to allow history dependence of B_{ℓ} and p_{ℓ} , yet leave room for randomness, we define

$$B_{\ell} = f_{\ell}(X_0, X_1, \dots, X_{\ell-1}, i_1, j_1, i_2, j_2, \dots, i_{\ell}, j_{\ell}, U_{\ell}) \in [\delta, 1)$$
(3)

and

$$p_{\ell} = g_{\ell}(X_0, X_1, \dots, X_{\ell-1}, i_1, j_1, i_2, j_2, \dots, i_{\ell}, j_{\ell}, V_{\ell}) \in (0, 1).$$
(4)

The assumption that U_{ℓ} and V_{ℓ} in eqs. (3) and (4) are uniformly distributed is not restrictive, since for any distribution Φ on \mathbb{R} , there exists a Borel-measurable function φ so that $\varphi(Z)$ has distribution Φ if $Z \in (0, 1)$ is uniformly distributed [8, Theorem 1.2.2]. Let further

$$s_{\ell} = \begin{cases} 1 & \text{if } W_{\ell} \le p_{\ell}, \\ -1 & \text{otherwise.} \end{cases}$$

Then define

$$X_{\ell}^{\mu_{\ell}} = X_{\ell-1}^{\mu_{\ell}} - s_{\ell} B_{\ell} X_{\ell-1}^{\mu_{\ell}}, \quad X_{\ell}^{\nu_{\ell}} = X_{\ell-1}^{\nu_{\ell}} + s_{\ell} B_{\ell} X_{\ell-1}^{\mu_{\ell}},$$
and $X_{\ell}^{i} = X_{\ell-1}^{i}$ for $i \notin \{\mu_{\ell}, \nu_{\ell}\}.$
(5)

3 Conditional expected change in the Euclidean norm of the wealth vector in a single trade

The key step in our analysis is to calculate the conditional expected change in the Euclidean norm of the vector X_{ℓ} in one trade:

$$E\left(\|X_{\ell}\|^{2} - \|X_{\ell-1}\|^{2} | X_{0}, \dots, X_{\ell-1}, \mu_{\ell}, \nu_{\ell}, B_{\ell}, p_{\ell}\right)$$

$$= p_{\ell}\left(\left(X_{\ell-1}^{\mu_{\ell}} - B_{\ell}X_{\ell-1}^{\mu_{\ell}}\right)^{2} + \left(X_{\ell-1}^{\nu_{\ell}} + B_{\ell}X_{\ell-1}^{\mu_{\ell}}\right)^{2}\right) + \left(1 - p_{\ell}\right)\left(\left(X_{\ell-1}^{\mu_{\ell}} + B_{\ell}X_{\ell-1}^{\mu_{\ell}}\right)^{2} + \left(X_{\ell-1}^{\nu_{\ell}} - B_{\ell}X_{\ell-1}^{\mu_{\ell}}\right)^{2}\right) - \left(\left(X_{\ell-1}^{\mu_{\ell}}\right)^{2} + \left(X_{\ell-1}^{\nu_{\ell}}\right)^{2}\right)$$

$$= (4p_{\ell} - 2)B_{\ell}X_{\ell-1}^{\mu_{\ell}}\left(X_{\ell-1}^{\nu_{\ell}} - X_{\ell-1}^{\mu_{\ell}}\right) + 2\left(B_{\ell}X_{\ell-1}^{\mu_{\ell}}\right)^{2}.$$
(6)

For fair coin tosses $(p_{\ell} = \frac{1}{2} \text{ for all } \ell)$, the left summand in (6) is zero, and therefore (6) is positive. For coin tosses with a wealth-acquired advantage $(p_{\ell} > \frac{1}{2})$, (6) is of course positive as well. In order to demonstrate condensation even in some cases with a poverty-acquired advantage $(p_{\ell} < \frac{1}{2})$, we assume

$$(4p_{\ell}-2)\left(X_{\ell-1}^{\mathbf{v}_{\ell}}-X_{\ell-1}^{\mu_{\ell}}\right) \ge -\delta X_{\ell-1}^{\mu_{\ell}} \quad \text{almost surely.}$$

$$\tag{7}$$

Note that this is an assumption on the functions g_{ℓ} . Since $-\delta X_{\ell-1}^{\mu_{\ell}} \ge -B_{\ell} X_{\ell-1}^{\mu_{\ell}}$, it then follows that (6) is $\ge (B_{\ell} X_{\ell-1}^{\mu_{\ell}})^2$. We summarize our result in the following lemma.

Lemma 1. Inequality (7) implies

$$E\left(\|X_{\ell}\|^{2} - \|X_{\ell-1}\|^{2} | X_{0}, \dots, X_{\ell-1}, \mu_{\ell}, \nu_{\ell}, B_{\ell}, p_{\ell}\right) \ge \left(B_{\ell} X_{\ell-1}^{\mu_{\ell}}\right)^{2}.$$
(8)

4 Convergence of the amount of wealth at stake in a transaction

Lemma 2. Inequality (7) implies $\lim_{\ell \to \infty} X_{\ell-1}^{\mu_{\ell}} = 0$ almost surely.

Proof. Let $\varepsilon > 0$. We will show that almost surely, $(B_{\ell}X_{\ell-1}^{\mu_{\ell}})^2 \ge \varepsilon$ for at most finitely many *n*. By the Borel-Cantelli lemma, it is sufficient to show that

$$\sum_{\ell=1}^{\infty} P\left(\left(B_{\ell} X_{\ell-1}^{\mu_{\ell}} \right)^2 \ge \varepsilon \right) < \infty.$$

Since

$$E\left(\left(B_{\ell}X_{\ell-1}^{\mu_{\ell}}\right)^{2}\right) \geq \varepsilon^{2}P\left(\left(B_{\ell}X_{\ell-1}^{\mu_{\ell}}\right)^{2} \geq \varepsilon\right),$$

it suffices to prove

$$\sum_{\ell=1}^{\infty} E\left((B_{\ell} X_{\ell-1}^{\mu_{\ell}})^2 \right) < \infty.$$

Taking (unconditional) expectations in (8), we find

$$E((B_{\ell}X_{\ell-1}^{\mu_{\ell}})^2) \leq E(\|X_{\ell}\|^2 - \|X_{\ell-1}\|^2).$$

and therefore, for all k,

$$\sum_{\ell=1}^{k} E\left((B_{\ell} X_{\ell-1}^{\mu_{\ell}})^2 \right) \le E\left(\|X_k\|^2 \right) - E\left(\|X_0\|^2 \right) \le E\left(\sum_{i=1}^{N} \left(X_k^i \right)^2 \right) \le E\left(\sum_{i=1}^{N} X_k^i \right) = 1.$$

We conclude that $(B_{\ell}X_{\ell-1}^{\mu_{\ell}})^2 \to 0$ almost surely, and since $B_{\ell} \ge \delta > 0$, this implies $X_{\ell-1}^{\mu_{\ell}} \to 0$ almost surely.

5 Local wealth condensation

We write

$$\mathcal{C} = \left\{ X \in [0,1]^N : \sum_{i=1}^N X^i = 1, \ X^i X^j a_{ij} = 0 \text{ for all } (i,j) \text{ with } 1 \le i, j \le N \right\}.$$

A wealth distribution in C has the property that any agent with a positive amount of wealth is connected only to agents with zero wealth. Local wealth condensation means convergence of the X_{ℓ} to C.

Theorem 2 (Generalized Yard-Sale Convergence Theorem). Inequality (7) implies

$$\lim_{\ell \to \infty} \operatorname{dist}(X_{\ell}, \mathcal{C}) = 0 \quad almost \ surely.$$
(9)

Here dist *denotes the distance in any norm on* \mathbb{R}^N *.*

Proof. Equation (9) means

$$\lim_{\ell \to \infty} \max\left\{\min\left\{X_{\ell-1}^i, X_{\ell-1}^j\right\} : 1 \le i, j \le N, a_{ij} = 1\right\} = 0 \text{ almost surely}$$

If this were not the case, then with positive probability, there would be an $\varepsilon > 0$ so that

$$\max\left\{\min\left\{X_{\ell-1}^i, X_{\ell-1}^j\right\} : 1 \le i, j \le N, a_{ij} = 1\right\} \ge \varepsilon.$$

infinitely often. This would imply that with positive probability, $X_{\ell-1}^{\mu_{\ell}} \ge \varepsilon$ infinitely often, in contradiction with Lemma 2.

6 Discussion

The yard-sale convergence theorem shows that (extreme) inequality can be the result of pure chance, and does not require agents of varying ability or industriousness. This is why it is interesting.

In this paper, we have tried to strike a balance between clarity and generality. We could, for instance, have allowed the choice of (i_{ℓ}, j_{ℓ}) to be history-dependent, as long as for every $(i, j) \in \mathcal{E}$, the probability $P((i_{\ell}, j_{\ell}) = (i, j))$ is uniformly bounded away from zero, so that no pair of agents drops out altogether in the limit as $\ell \to \infty$. Further, instead of $B_{\ell} \ge \delta$, we could have assumed that $P(B_{\ell} \ge \delta)$ is uniformly bounded away from zero. These generalizations didn't seem to add insight justifying the expense of more opaque notation.

Numerical experiments and heuristic arguments [12] suggest that wealth condensation even occurs for a fixed $p_{\ell} = p$ very slightly smaller than $\frac{1}{2}$. However, (7) does not hold then, so our theorem does not extend to this case.

It is plausible and supported by numerical simulations that eq. (9) can be replaced by the slightly stronger statement

$$\exists X \in \mathcal{C} \quad \lim_{\ell \to \infty} X_{\ell} = X.$$
⁽¹⁰⁾

For a complete graph, C consists of isolated points only, namely, the canonical basis vectors in \mathbb{R}^N . Conditions (9) and (10) are equivalent in that case. For an incomplete graph, Calways contains non-isolated points, and the equivalence of (9) and (10) is therefore no longer obvious. However, for p = 1/2, the argument based on the martingale convergence theorem does prove (10) even for an incomplete graph.

These limitations notwithstanding, the argument given in this paper appears to be both the simplest and the most general rigorous mathematical proof of the yard-sale convergence theorem for a finite number of agents.

References

- B. BOGHOSIAN, A. DEVITT-LEE, M. JOHNSON, J. MARCQ, AND H. WANG, Oligarchy as a phase transition: The effect of wealth-attained advantage in a Fokker-Planck description of asset exchange, Physica A, 476 (2017), pp. 15–37.
- [2] B. BOGHOSIAN, M. JOHNSON, AND J. MARCQ, An H theorem for Boltzmann's equation for the yard-sale model of asset exchange, J. Stat. Phys., 161 (2015), pp. 1339–1350.
- [3] C. BÖRGERS AND C. GREENGARD, A new probabilistic analysis of the yard-sale model, ArXiv e-print 2308.01485v1, 2023.
- [4] R. BUSTOS-GUAJARDO AND C. F. MOUKARZEL, Yard-sale exchange on networks: wealth sharing and wealth appropriation, J. Stat. Mech.: Theory Exp., P12009, (2012).
- [5] B.-H. F. CARDOSO, S. GONÇALVES, AND J. R. IGLESIAS, Why equal opportunities lead to maximum inequality? The wealth condensation paradox generally solved, Chaos Solit. Fractals, 168, 113181 (2023).
- [6] A. CHAKRABORTI, Distributions of money in model markets of economy, Int. J. Mod. Phs. C, 13 (2002), pp. 1315–1321.
- [7] C. CHORRO, A simple probabilistic approach of the yard-sale model, Stat. Probab. Lett., 112 (2016), pp. 35–40.
- [8] R. DURRETT, Probability: Theory and Examples, Cambridge University Press, 2019.
- [9] B. KRAMER AND A. MACKINNON, *Localization: theory and experiment*, Rep. Prog. Phys., 56 (1993), pp. 1469–1564.
- [10] M. LAAKSO AND R. TAAGEPERA, "Effective" number of parties: a measure with applications to West Europe, Comp. Polit. Stud., 12 (1979), pp. 3–27.
- [11] H. G. LEE AND D.-S. LEE, Scaling in local to global condensation of wealth on sparse networks, Phys. Rev. E, 108, doi: 10.1103/PhysRevE.108.064303 (2023).
- [12] C. F. MOUKARZEL, S. CONÇALVES, J. R. IGLESIAS, M. RODRÍGUEZ-ACHACH, AND R. HUERTA-QUINTANILLA, Wealth condensation in a multiplicative random asset exchange model, Eur. Phys. J. Special Topics, 143 (2007), pp. 75–79.
- [13] E. H. SIMPSON, *Measurement of diversity*, Nature, 163 (1949).

- [14] R. TIBSHIRANI, *Regression shrinkage and selection via the lasso*, J. Royal Stat. Soc. B, 58 (1996), pp. 267–288.
- [15] U.S. DEPARTMENT OF JUSTICE, *Herfindahl-Hirschmann Index*, https://www.justice.gov/atr/herfindahl-hirschman-index.