

INEQUALITIES FOR k -REGULAR PARTITIONS

BERNHARD HEIM AND MARKUS NEUHAUSER

ABSTRACT. We build upon the work by Bessenrodt and Ono, as well as Beckwith and Bessenrodt concerning the combined additive and multiplicative behavior of the k -regular partition functions $p_k(n)$. Our focus is on addressing the solutions of the Bessenrodt–Ono inequality

$$p_k(a)p_k(b) > p_k(a+b).$$

We determine the sets E_k and F_k consisting of all pairs (a, b) , where we have equality or the opposite inequality. Bessenrodt and Ono previously determined the exception sets E_∞ and F_∞ for the partition function $p(n)$. We prove by induction that $E_k = E_\infty$ and $F_k = F_\infty$ if and only if $k \geq 10$. Beckwith and Bessenrodt used analytic methods to consider $2 \leq k \leq 6$, while Alanazi, Gagola, and Munagi studied the case $k = 2$ using combinatorial methods. Finally, we present a precise and comprehensive conjecture on the log-concavity of the k -regular partition function extending previous speculations by Craig and Pun. The case $k = 2$ was recently proven by Dong and Ji.

1. INTRODUCTION

In this paper we investigate partition functions [An98, On04] and provide a complete solution for the Bessenrodt–Ono inequality [BO16] for k -regular partition functions $p_k(n)$ [Ha71]. We extend the previous results for $2 \leq k \leq 6$ by Beckwith and Bessenrodt [BB16] to $k \geq 2$, proving that for $k \geq 10$, the solutions coincide with the solutions of the partition function $p(n)$.

Let $\{\alpha(n)\}_n$ denote a sequence of positive integers. The sequence satisfies the Bessenrodt–Ono inequality for a pair (a, b) of positive integers if

$$(1.1) \quad \Delta(a, b) := \alpha(a)\alpha(b) - \alpha(a+b) > 0.$$

For partitions $\alpha(0) = \alpha(1) = 1$ and due to symmetry, we only consider pairs with $1 < a \leq b$. Let E denote the set of pairs where $\Delta(a, b) = 0$ and F the set, where $\Delta(a, b) < 0$. For k -regular partition functions, we use the symbols $\Delta_k(a, b)$, E_k and F_k , where $k \in \{2, 3, 4, \dots\} \cup \{\infty\}$.

2020 *Mathematics Subject Classification*. Primary 05A17, 11P82; Secondary 05A20.

Key words and phrases. Bessenrodt–Ono inequality, Generating functions, k -regular partitions.

Let us recall the definition of the k -regular partition function $p_k(n)$. A partition of a positive integer n is a finite nonincreasing sequence of positive integers that sum to n . These integers are called the parts of the partition. The partition function $p(n)$ is the number of partitions of n ([An98], Section 1.1). Let $k \geq 2$. The k -regular partition function $p_k(n)$ is the number of partitions of n in which none of the parts is a multiple of k . Note, that $p_k(n)$ is also equal to the number of partitions with each part appearing at most $k - 1$ times. Therefore, the number of partitions of n with odd parts is equal to the number of partitions with different parts. Based on the results of this paper and the fact that $p_k(n) = p(n)$ for $k > n$, we define $p_\infty(n) := p(n)$.

In 2016, Bessenrodt and Ono [BO16] discovered the following property of the partition function $p_\infty(n)$.

Theorem 1.1 (Bessenrodt, Ono 2016). *Let $1 < a \leq b$. Then the Bessenrodt–Ono inequality (1.1) is satisfied for the partition function $p_\infty(n)$, except for the pairs*

$$\begin{aligned} E_\infty &= \{(2, 6), (2, 7), (3, 4)\}, \\ F_\infty &= \{(2, 2), (2, 3), (2, 4), (2, 5), (3, 3), (3, 5)\}. \end{aligned}$$

Bessenrodt and Ono provided an analytic proof ([BO16], Section 2) utilizing deep results by Lehmer [Le39] and Rademacher [Ra37] for the partition function, built on the circle method by Hardy and Ramanujan. A second combinatorial proof was found by Alanazi, Gagola, and Munagi [AGM17]. Their sophisticated proof involved an injective map and an induction. A third algebraic proof was provided in [HN22] using induction, and was also generalized to plane partitions [HNT23].

Beckwith and Bessenrodt [BB16] demonstrated that k -regular partition functions for $2 \leq k \leq 6$ also fulfill the Bessenrodt–Ono inequality (see Section 2 for their result).

Moreover, many more sequences turned out to satisfy (1.1). Chern, Fu, and Tang [CFT18] generalized and proved the Bessenrodt–Ono inequality (BO) for k -colored partitions. Hou and Jagadeesan [HJ18] extended on the numbers of partitions with ranks in a given residue class modulo 3 and Males [Ma20] did so for general modulus. A modified BO was presented in [HN19]. Dawsey and Masri [DM19] proved BO for the Andrews spt -function, and Gajdzica did so for A -partitions. In [HNT20], the polynomization of the k -colored partitions [CFT18] was initiated, leading to a continuous version of the Chern, Fu, Tang result [CFT18] for k -regular partitions. The proof is utilizing an Euler-type recursion formula for the involved polynomials and their derivatives. This polynomization approach and the underlying proof

strategy were also successfully applied by Li [Li23] for the overpartition function and by Gajdzica et al. ([GHN24], Theorem 1.3) for A -partition functions.

Recently, several interesting and important papers have explored the asymptotic behavior leading to log-concavity properties and the Bessenrodt–Ono inequality, as well as their relations. We refer to Bringmann, Franke, and Heim ([BFH24], Theorem 1.3 and Theorem 1.5), Benfield and Roy ([BR24], Theorem 1.1) and Gajdzica, Miska, and Ulas ([GMU24], Theorem 2.1 and Theorem 3.1).

2. RESULTS ON k -REGULAR PARTITIONS

Let $k \geq 2$. The generating function for the number of k -regular partitions is given by

$$\sum_{n=0}^{\infty} p_k(n) q^n = \prod_{n=1}^{\infty} \frac{(1 - q^{kn})}{(1 - q^n)}.$$

Note that

$$\prod_{n=1}^{\infty} \frac{(1 - q^{2n})}{(1 - q^n)} = \prod_{n=1}^{\infty} (1 - q^{2n-1})^{-1},$$

which implies that the number of 2-regular partitions of n is equal to the number of partitions of n with odd parts. We also refer to Glaisher’s generalization for arbitrary k ([An98], Corollary 1.3).

Let $g_k(n) := \sigma(n) - k\sigma(n/k)$, where $\sigma(n) := \sum_{d|n} d$ and $\sigma(x) = 0$ if $x \notin \mathbb{N}$. Then

$$\sum_{n=0}^{\infty} p_k(n) q^n = \exp\left(\sum_{n=1}^{\infty} g_k(n) \frac{q^n}{n}\right).$$

This follows from the identity $\prod_{n=1}^{\infty} (1 - q^n) = \exp(-\sum_{n=1}^{\infty} \sigma(n) q^n/n)$ and implies

$$n p_k(n) = \sum_{\ell=1}^n g_k(\ell) p_k(n - \ell),$$

with initial value $p_k(0) = 1$. Beckwith and Bessenrodt proved, in the same spirit as [BO16], that the k -regular partition function for $2 \leq k \leq 6$ satisfies (1.1). They utilized a Rademacher-type formula for $p_k(n)$ given by Hagsis [Ha71] and determined for each k the exception sets E_k and F_k . We recall the following: Let a, b, k be positive integers with $1 < a \leq b$ and let $k \geq 2$. Then we denote by E_k the set of all pairs (a, b) with equality in (1.1) and F_k the set of all pairs (a, b) with

$$p_k(a) p_k(b) < p_k(a + b).$$

k	Elements of E_k	Elements of F_k
2	(3, 3), (3, 5), (3, 6), (3, 7), (3, 8), (4, 15), (4, 16), (4, 17), (5, 6), (5, 7), (5, 8)	(2, b), $b \geq 2$, (3, 4), (4, 4), (4, 5), (4, 6), (4, 7), (4, 8), (4, 9), (4, 10), (4, 11), (4, 12), (4, 13), (4, 14), (5, 5)
3	(2, 2), (3, 10)	(2, 3), (3, 3), (3, 4), (3, 5), (3, 6), (3, 7), (3, 8), (3, 9), (3, 11), (3, 13)
4	(2, 2), (2, 3), (2, 5), (3, 3), (3, 4), (4, 4)	(2, 4)
5	(2, 3), (2, 4)	(2, 2), (2, 5), (3, 3), (3, 5)
6	(2, 4), (2, 5), (2, 6)	(2, 2), (2, 3), (3, 3)

TABLE 1. Exceptions to the Bessenrodt–Ono inequality for $2 \leq k \leq 6$

Theorem 2.1 (Beckwith, Bessenroth 2016). *Let a, b, k be integers with $1 < a \leq b$ and $2 \leq k \leq 6$. Then the Bessenrodt–Ono inequality*

$$(2.1) \quad p_k(a) p_k(b) > p_k(a + b)$$

is satisfied for all pairs, except for the pairs presented in Table 1 for E_k and F_k . Therefore, let n_k and m_k be given by the following table:

k	2	3	4	5	6
n_k	3	2	2	2	2
m_k	22	17	9	9	9

Then for any $a, b \geq n_k$ and $a + b \geq m_k$, the Bessenrodt–Ono inequality (2.1) is valid.

Further, Alanzi, Gagola, and Munagi [AGM17] demonstrated that their combinatorial proof of the Bessenrodt–Ono inequality can, in principle, be adapted to the k -regular case. They provided a proof for $k = 2$ and indicated that the number of cases increase rapidly for larger k .

We extend the table of Beckwith and Bessenroth for E_k and F_k to $7 \leq k \leq 10$. This leads to Table 2.

k	Elements of E_k	Elements of F_k
7	(2, 5), (2, 7), (3, 5)	(2, 2), (2, 3), (2, 4), (3, 3)
8	(3, 4), (3, 5)	(2, 2), (2, 3), (2, 4), (2, 5), (3, 3)
9	(2, 6), (3, 4)	(2, 2), (2, 3), (2, 4), (2, 5), (3, 3), (3, 5)
10	(2, 6), (2, 7), (3, 4)	(2, 2), (2, 3), (2, 4), (2, 5), (3, 3), (3, 5)

TABLE 2. Exceptions to the Bessenrodt–Ono inequality for $7 \leq k \leq 10$

We observe that $E_k = E_\infty$ and $F_k = F_\infty$ for $k = 10$ and prove that this even holds true for all $k \geq 10$.

Theorem 2.2. *Let $a, b, k \geq 2$ be integers, and $p_k(n)$ be the k -regular partition function. Then the Bessenrodt–Ono inequality*

$$p_k(a)p_k(b) > p_k(a+b)$$

holds true for all pairs (a, b) except for pairs in E_k , where equality is in place and F_k , where $p_k(a)p_k(b) < p_k(a+b)$. For $2 \leq k \leq 10$, the sets E_k and F_k are given by Tables 1 and 2. For all $k \geq 10$ we have

$$E_k = E_\infty \text{ and } F_k = F_\infty.$$

3. PROOF OF THEOREM 2.2

We closely follow the proof scheme ([HN22], Section 2) proposed to prove Bessenrodt–Ono inequalities.

First, we estimate the arithmetic function $g_k(n)$ and the k -regular partition function $p_k(n)$ with the following upper and lower bounds, independent of k .

Lemma. *Let $k \geq 2$. Then*

$$(3.1) \quad \begin{aligned} g_k(n) &\leq n(1 + \ln(n)), \\ p_k(n) &> 2\sqrt{2n/3+1/4-3/2}. \end{aligned}$$

Proof. The upper bound for $g_k(n) \leq \sigma(n)$ follows easily by integral comparison with the bound for $n > 1$. The lower bound of $p_k(n)$ cannot be directly related to $p(n)$ as given in ([HN22], equation (2.2)), and requires some additional investigation.

The lower bound of $p_k(n)$ follows from

$$p_k(n) \geq 4 \lfloor \sqrt{2n/3+1/2-1/2} \rfloor - \lfloor \lfloor \sqrt{2n/3+1/4-1/2} \rfloor / k \rfloor > 4^{(k-1)} (\sqrt{2n/3+1/4-3/2}) / k.$$

Here $\lfloor x \rfloor$ denotes the usual floor symbol. Consider partitions of n using only parts from $\{1, 2, \dots, \lfloor \sqrt{n/3} \rfloor\} \setminus k\mathbb{N}$ with multiplicity at most 3. To any admissible submultiset, we can assign a partition. The largest value we obtain is

$$\begin{aligned} 3 \sum_{m=1}^{\lfloor \sqrt{2n/3+1/4}-1/2 \rfloor} m &\leq \frac{3}{2} \left(\sqrt{2n/3+1/4} - 1/2 + 1 \right) \left(\sqrt{2n/3+1/4} - 1/2 \right) \\ &= 3 \frac{2n/3 + 1/4 - 1/4}{2} = n. \end{aligned}$$

Also, we have removed $\lfloor \frac{1}{k} \lfloor \sqrt{\frac{2n}{3} + \frac{1}{4}} - \frac{3}{2} \rfloor \rfloor$ elements from the set. This demonstrates the claim. \square

Remark. Note that the proof shows that we could even choose as a lower bound

$$p_k(n) \geq 2^{\lfloor \sqrt{2n/3+1/4}-1/2 \rfloor}.$$

3.1. Proof of Theorem 2.2 for $k > 3$. Let $A := 2$ and $B := 10$. Let $n \geq B$. We say the statement $S(n)$ holds true if for all partitions $n = a + b$ with $a, b \geq A$:

$$\Delta_k(a, b) := p_k(a)p_k(b) - p_k(a+b) > 0.$$

We assume that $n > N_0 > 1$ and $S(m)$ hold true for all $B \leq m \leq n-1$. For $B \leq m \leq N_0$, we demonstrate $S(m)$ through a direct computer calculation with PARI/GP. It will turn out that we can choose $N_0 = 2938$.

We obtain the following expressions for $\Delta_k(a, b) = L + R$:

$$\begin{aligned} L &:= - \sum_{\ell=1}^b \frac{g_k(\ell+a)}{a+b} p_k(b-\ell), \\ R &:= \sum_{\ell=1}^a \left(\frac{g_k(\ell)}{a} p_k(a-\ell) p_k(b) - \frac{g_k(\ell)}{a+b} p_k(a+b-\ell) \right). \end{aligned}$$

Further, we will refine the right sum R into

$$R = R_1 + R_2 + R_3.$$

3.1.1. *Left sum L .* We have $L > -b p_k(b) (1 + \ln(a+b))$.

3.1.2. *Right sum R .* Let

$$\ell_0 := a - \max\{B-b, A\} + 1.$$

Therefore, $\ell_0 = a - \max\{9 - b, 1\}$. Let

$$R_1 := \sum_{\ell=1}^1 f_\ell(a, b), \quad R_2 := \sum_{\ell=2}^{\ell_0-1} f_\ell(a, b), \quad R_3 := \sum_{\ell=\ell_0}^a f_\ell(a, b),$$

$$f_\ell(a, b) := \frac{g_k(\ell)}{a} p_k(a - \ell) p_k(b) - \frac{g_k(\ell)}{a + b} p_k(a + b - \ell).$$

The sum R_1 . We obtain the lower bound:

$$R_1 > \frac{b}{2a^2} p_k(a - 1) p_k(b).$$

The sum R_2 . The second sum: $R_2 > 0$.

The sum R_3 .

We split the third sum again into three parts: $R_3 = R_{31} + R_{32} + R_{33}$, where

$$(3.2) \quad R_{31} := \sum_{\ell=\ell_0}^{a-A} f_\ell(a, b) > 5(4 - p_\infty(8))(1 + \ln(a)),$$

$$R_{32} := \sum_{\ell=a-A+1}^{a-1} f_\ell(a, b) > -p_k(b)(1 + \ln(a)),$$

$$R_{33} := \sum_{\ell=a}^a f_\ell(a, b) \geq 0.$$

We first use that $p_k(9 - b)p_k(b) \geq p_k(9)$, which leads to at most 5 summands. The estimation in (3.2) follows from $p_k(a - \ell) \geq p_k(2)$. For the estimation, we use that

$$p_k(b) \geq p_k(2) = 2, \quad \frac{1}{a + b} < \frac{1}{a}, \quad \text{and } p_k(a + b - \ell) \leq p_k(8) \leq p_\infty(8).$$

We obtain

$$R_3 > -23b p_k(b)(1 + \ln(a + b)).$$

3.1.3. *Final step.* Putting everything together leads to:

$$(3.3) \quad \Delta_k(a, b) > \frac{b p_k(b)}{2a^2} \left(-48a^2(1 + \ln(2a)) + 2\sqrt{2^{(a-1)/3+1/4-3/2}} \right).$$

As the final step, we used the property (3.1). For $a \rightarrow \infty$ we can immediately observe that this is positive since the second term grows in a faster than $a \mapsto a^2(1 + \ln(2a))$. In fact, the expression (3.3) is positive for all $a \geq 1470$. Note that if $a < 1470$, then $a + b \leq 2a \leq N_0$.

Note that for $k > N_0$, we have $p_k(n) = p_\infty(n)$ for all $n \leq N_0$. Therefore, we only have to check the inequality for $k \leq N_0$ and $n \leq N_0$. Hence, $\Delta_k(a, b) > 0$, which proves the Theorem.

Remark. Note that we can also prove the cases $k \in \{2, 3\}$ directly with our method from [HN22], but then more adjustments are necessary as we cannot directly use ([HN22], (2.8)) anymore. For $k = 2$, we can choose $A = 3$ and $B = 22$ with $N_0 = 3662$, and for $k = 3$, we can choose $A = 2$ and $B = 17$ with $N_0 = 3776$. The crucial step is to adjust the estimate of R_{31} in (3.2) for these cases.

4. LOG-CONCAVITY

In this final section, we provide further evidence, that the k -regular partition function and the partition function $p_\infty(n)$ share similar properties up to small ks . We indicate that, as in the case of the Bessenrodt–Ono inequality, this may also apply to the log-concave property. Nicolas [Ni78] and DeSalvo and Pak [DP15] proved that the partition function $p_\infty(n)$ is log-concave at n :

$$p(n)_\infty^2 \geq p_\infty(n-1)p_\infty(n+1)$$

if and only if $n \geq 25$ or n is even. For k -regular partitions, building on Hagis' Rademacher-type formula and a new method developed by Griffin, Ono, Rolén, and Zagier [GORZ19], Craig and Pun [CP20] proved that the k -regular partition functions $p_k(n)$ satisfy higher-order Turán inequalities for sufficiently large n dependent on k . Let N_k be the smallest number, such that $\{p_k(n)\}_n$ is log-concave at n

$$p_k(n)^2 \geq p_k(n-1)p_k(n+1)$$

for all $n \geq N_k$. We calculated all exceptions for $2 \leq k \leq 100$ and $1 \leq n \leq 1000$. For $2 \leq k \leq 20$, we have provided them in Table 3, where for $k = 3$, the exceptions are all odd $1 \leq n \leq 57$. This is compatible with the numbers conjectured by Craig and Pun ([CP21], Remark 1.3) for $2 \leq k \leq 4$. Recently, Dong and Ji [DJ24] proved the conjecture for $k = 2$.

Similar to the Bessenrodt–Ono inequality, we believe that the set of exceptions stabilizes for large k and is the same as the one of $p_\infty(n)$. This set is given by all odd numbers $1 \leq n \leq 25$. We put $N_\infty := 26$.

Conjecture. *Let $k \geq 2$. Let N_k be the smallest integer, such that $\{p_k(n)\}_n$ is log-concave at every $n \geq N_k$. Then $N_k = N_\infty$ if $k \geq 30$.*

This is in accordance with Remark 1.3 given by Craig and Pun [CP21]. Since $p_k(n) = p_\infty(n)$ for $k > n$ we also have:

Corollary. *Let the conjecture hold true. Let $k \geq 30$. Then the exceptions n are the same as for $\{p_\infty(n)\}_n$.*

$n \setminus k$	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
1		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
2	•																		
3		•		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
4	•		•																
5		•		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
6																			
7		•		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
8	•		•																
9		•		•	•	•		•	•	•	•	•	•	•	•	•	•	•	•
10																			
11	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
12																			
13	•	•	•	•		•	•	•	•	•	•	•	•	•	•	•	•	•	•
14	•																		
15		•		•		•		•	•	•	•	•	•	•	•	•	•	•	•
16	•		•																
17	•	•		•		•	•	•	•	•	•	•	•	•	•	•	•	•	•
18																			
19	•	•		•		•	•	•	•	•	•	•	•	•	•	•	•	•	•
20	•																		
21		•		•		•		•		•	•	•		•	•	•	•	•	•
22																			
23	•	•		•		•		•		•		•	•	•	•	•	•	•	•
24																			
25		•		•		•		•		•		•		•	•	•	•	•	•
26	•																		
27		•		•		•		•		•		•		•		•		•	
28																			
29	•	•		•		•		•		•		•		•		•		•	
30																			
31		•		•		•		•		•		•							
32	•																		
33		•		•		•		•											
34																			
35		•		•		•													
36																			
37		•		•															
38																			
39		•		•															
40																			
41		•		•															
42																			
43		•																	
44																			
45		•																	

TABLE 3. Exceptions of log-concavity at n for $2 \leq k \leq 20$ and $1 \leq n \leq 45$

REFERENCES

- [AGM17] A. A. Alanazi, S. M. Gagola III, A. O. Munagi: *Combinatorial proof of a partition inequality of Bessenrodt–Ono*. *Ann. Comb.* **21** (2017), 331–337.
- [An98] G. E. Andrews: *The Theory of Partitions*. Cambridge Univ. Press, Cambridge (1998).
- [BB16] O. Beckwith, C. Bessenrodt: *Multiplicative properties of the number of k -regular partitions*. *Ann. Comb.* **20** No. 2 (2016), 231–250.
- [BR24] B. Benfield, A. Roy: *Log-concavity and the multiplicative properties of restricted partition functions*. arXiv:2404.03153 (2024).
- [BO16] C. Bessenrodt, K. Ono: *Maximal multiplicative properties of partitions*. *Ann. Comb.* **20** No. 1 (2016), 59–64.
- [BFH24] K. Bringmann, J. Franke, B. Heim: *Asymptotics of commuting ℓ -tuples in symmetric groups and log-concavity*. arXiv:2401.05874v1 (2024).
- [CFT18] S. Chern, S. Fu, D. Tang: *Some inequalities for k -colored partition functions*. *Ramanujan J.* **46** (2018), 713–725.
- [CP20] W. Craig, A. Pun: *Higher order Turán inequalities for k -regular partitions*. *Séminaire Loth. de Comb.* **84B** Article #23 (2020).
- [CP21] W. Craig, A. Pun: *A note on the higher order Turán inequalities for k -regular partitions*. *Research in Number Theory* **7** Article #5 (2021).
- [DM19] M. L. Dawsey, R. Masri: *Effective bounds for the Andrews spt -function*. *Forum Math.* **31** No. 3 (2019), 743–767.
- [DP15] S. DeSalvo, I. Pak: *Log-concavity of the partition function*. *Ramanujan J.* **38** (2015), 61–73.
- [DJ24] J. Dong, K. Ji: *Higher order Turán inequalities for the distinct partition function*. *Journal of Number Theory* **260** (2024), 71–102.
- [GHN24] K. Gajdzica, B. Heim, M. Neuhauser: *Polynomization of the Bessenrodt–Ono type inequalities for A -partition functions*. *Ann. Comb.* (2024). <https://doi.org/10.1007/s00026-024-00692-4>
- [GMU24] K. Gajdzica, P. Miska, M. Ulas: *On a general approach to Bessenrodt–Ono type inequalities and log-concavity properties*. *Ann. Comb.* (2024). <https://doi.org/10.1007/s00026-024-00700-7>
- [GORZ19] M. Griffin, K. Ono, L. Rolin, D. Zagier: *Jensen polynomials for the Riemann zeta function and other sequences*. *Proc. Natl. Acad. Sci., USA* **116** No. 23 (2019), 11103–11110.
- [Ha71] P. Hagsis: *Partitions with a restriction on the multiplicity of the summands*. *Trans. Amer. Math. Soc.* **155** (1971), 375–384.
- [HN19] B. Heim, M. Neuhauser: *Variants of a partition inequality of Bessenrodt–Ono*. *Research in Number Theory* **5** Article #32 (2019).
- [HN22] B. Heim, M. Neuhauser: *Proof of the Bessenrodt–Ono inequality by induction*. *Research in Number Theory* **8** Article #3 (2022).
- [HNT20] B. Heim, M. Neuhauser, R. Tröger: *Polynomization of the Bessenrodt–Ono inequality*. *Ann. Comb.* **24** (2020), 697–709.
- [HNT23] B. Heim, M. Neuhauser, R. Tröger: *Inequalities for plane partitions*. *Ann. Comb.* **27** (2023), 87–108.

- [HJ18] E. Hou, M. Jagadeesan: *Dyson's partition ranks and their multiplicative extension*. Ramanujan J. **45** No. 3 (2018), 817–839.
- [Le39] D. H. Lehmer: *On the remainders and convergence of the series for the partition functions*. Trans. Amer. Math. Soc. **46** (1939), 362–373.
- [Li23] X. Li: *Polynomization of the Liu–Zhang inequality for the overpartition function*. Ramanujan J. **62** (2023), 797–817.
- [Ma20] J. Males: *Asymptotic equidistribution and convexity for partition ranks*. Ramanujan J. **54** No. 2 (2021), 397–413.
- [Ni78] J.-L. Nicolas: *Sur les entiers N pour lesquels il y a beaucoup des groupes abéliens d'ordre N* . Ann. Inst. Fourier **28** No. 4 (1978), 1–16.
- [On04] K. Ono: *The Web of Modularity: Arithmetic of the Coefficients of Modular Forms and q -series*. CBMS Regional Conference Series in Mathematics **102**, American Mathematical Society, Providence, RI (2004).
- [Ra37] H. Rademacher: *A convergent series for the partition function $p(n)$* . Proc. Natl. Acad. Sci. USA **23** (1937), 78–84.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, DIVISION OF MATHEMATICS,
UNIVERSITY OF COLOGNE, WEYERTAL 86–90, 50931 COLOGNE, GERMANY

LEHRSTUHL A FÜR MATHEMATIK, RWTH AACHEN UNIVERSITY, 52056 AACHEN, GERMANY
Email address: bheim@uni-koeln.de

KUTAISI INTERNATIONAL UNIVERSITY, 5/7, YOUTH AVENUE, KUTAISI, 4600 GEORGIA

LEHRSTUHL A FÜR MATHEMATIK, RWTH AACHEN UNIVERSITY, 52056 AACHEN, GERMANY
Email address: markus.neuhauser@kiu.edu.ge