

2d QCD and Integrability

Part II: Generalized QCD

Federico Ambrosino¹ and Shota Komatsu²

¹Deutsches Elektronen-Synchrotron DESY, Notkestr. 85, 22607 Hamburg, Germany

²Department of Theoretical Physics, CERN, 1211 Meyrin, Switzerland

Abstract

We extend our study of integrable structures in large N_c QCD₂ to a broad class of theories called the generalized QCD, which are given by the Lagrangian $\mathcal{L} \propto \text{tr} B \wedge F - \text{tr} V(B)$ coupled to quarks in the fundamental representation. We recast the Bethe-Salpeter equation for the meson spectrum into a TQ-Baxter equation and determine a transfer matrix in a closed form for any given polynomial $V(B)$. Using this reformulation, we derive the asymptotic expansion of energy levels and wavefunctions. Examining the analytic structure of the spectrum as a function of the coefficients of $V(B)$, we determine the region of couplings where the theory admits a positive and discrete spectrum of mesons. Furthermore, we uncover a multi-sheeted structure with infinitely many multi-critical points, where several mesons become simultaneously massless. Lastly, we illustrate that this structure persists in the large-representation limit of the generalized QCD with the SU(2) gauge group.

June 18, 2024

Contents

1	Introduction	2
2	Generalized QCD in the Large N limit	4
2.1	Bound state equation	6
3	Generalized 't Hooft equation as a TQ-system	9
3.1	ν -space representation	9
3.2	Discreteness of the spectrum and instabilities	10
3.3	TQ-equation	12
3.4	Inhomogenous extension	13
3.5	Solving the TQ-equation: general strategy	16
3.5.1	Weak-coupling expansion of the spectral sum.	16
3.5.2	Asymptotic expansion of eigenvalues	17
3.5.3	Asymptotic expansion of wavefunctions	19
3.6	Solving the TQ-equation: cubic coupling	20
3.6.1	Constructing asymptotic solutions	20
3.6.2	Determining the eigenvalues	22
4	Analytical structure of the spectrum and critical points	24
4.1	Multicritical points from collision-of-roots	25
4.2	Critical points beyond the first sheet	29
4.3	A case study: cubic potential	29
4.3.1	Real couplings	30
4.3.2	Analytical continuation and tricritical point	32
4.3.3	Singularities beyond the cuts	36
5	Mesons spectrum in the large representation limit	40
6	Conclusion	47
A	Recurrence relation for integrals	48
B	Extracting the spectral determinants for cubic potential	49
C	Derivation of the bound state equation at large representation	51

1 Introduction

Understanding strong-coupling dynamics of gauge theory is one of the most important challenges in modern physics with diverse theoretical and phenomenological implications. In particular, essential mechanisms of how the Yang-Mills theory exhibits color confinement through the formation of chromoelectric fluxtubes and how the dynamics of the fluxtubes controls physical observables, such as the meson and glueball spectrum, still remain elusive.

To make progress on this front, in this paper we continue our exploration of toy models that are amenable to analytical methods but exhibit qualitative features similar to more realistic, yet complicated examples of four-dimensional gauge theories. Specifically, we study a class of theories in $1 + 1$ dimensions called *generalized QCD* in the large N_c limit. These are generalizations of the two-dimensional QCD (also known as the 't Hooft model) and are described by the *generalized Yang-Mills*, with Lagrangian given by $\mathcal{L} \propto \text{tr} B \wedge F - \text{tr} V(B)$, coupled to quarks in the fundamental representation of the gauge group.

In our previous paper [1], building on the results of [2], we demonstrated that the integral equation that determines the spectrum of mesons in the 't Hooft model can be reformulated into a finite difference equation that takes the form of the TQ-Baxter equations — the equations that determine the spectrum of integrable systems. Based on this reformulation, we obtained systematic asymptotic expansions of the mass spectrum and the wavefunctions.

As with the 't Hooft model, for a class of potentials $V(B)$ the generalized QCD at large N_c has stable mesonic bound states with a discrete spectrum. Thanks to the simplification that occurs at large N_c , the mass spectrum can be computed explicitly by solving integral equations that generalize the one for the 't Hooft model. In this paper, we show that these generalized equations can also be recast into the TQ-Baxter equation, which turns out to be remarkably simple and universal:

$$Q(\nu + 2i) + Q(\nu - 2i) - 2Q(\nu) = T(\nu)Q(\nu), \quad (1.1)$$

where the dependence on the potential $V(B)$ is contained entirely in the transfer matrix $T(\nu)$, which can be computed in a closed form for any given potential. See (3.17) for details. As in the 't Hooft model, this reformulation enables systematic asymptotic expansions of the mass spectrum and the wavefunctions.

Furthermore, the reformulation in terms of the TQ-Baxter equation and an associated

inhomogeneous Fredholm equation allow us to explore detailed analytic structures of the meson spectrum as a function of parameters of the theory. A similar study for the 't Hooft model revealed an intricate structure of the spectrum in the complex quark-mass plane [1,2]: it has a multi-sheeted structure with infinitely many branch points, each of which signals a tachyonic instability of one of the mesons. The first of such branch points corresponds to a massless quark point where the chiral symmetry becomes exact. The IR phase of that point is described by a CFT with the $U(1)_N$ current algebra [3,4]. This suggests the correspondence between the branch points in the complex mass plane and actual *critical points* of the theory. In this paper, we extend these results to the generalized QCD_2 . A novelty in the present case is that we can also study the dependence on the coefficients of the potential $V(B)$. We found that the theory develops instability for a certain range of parameters and the onset of the instability is signaled by “critical points” at which one of the mesons becomes massless (and beyond which it becomes tachyonic). The connection to the inhomogeneous Fredholm equation provides a simple and systematic criterion for the stability, bypassing the case-by-case numerical analysis performed in [5]. In addition, we found that, when the coefficients of the potential are appropriately tuned, several mesons can become simultaneously massless, realizing “multi-critical points”. As in the 't Hooft model, we conjecture that these “critical points” correspond to actual critical points of the theory. Unlike the standard QCD_2 , the IR phases of the generalized QCD_2 have not been sufficiently explored in the literature and our results strongly motivate such analysis.

These findings extend and strengthen a surprising connection between QCD_2 and integrable systems, initially found in [2] and further explored in our previous paper [1]. For theories with fundamental quarks studied in this paper, one might argue that this is simply a technical improvement since it is basically a rewriting of the already-known integral equation (although there are numerous advantages in doing so as we emphasized above). However such a rewriting could be truly beneficial for theories with quarks in the adjoint representation of the gauge group (often referred to as *adjoint QCD*). There have been renewed interest in these theories since they could serve as toy models of confining fluxtubes with nontrivial dynamics [6–20]. Furthermore, there is the possibility that the confining fluxtube becomes integrable at a certain point in the parameter space [21–25]. However, unlike theories with fundamental quarks, there is no simple integral equation that determines the spectrum of these theories. One possible strategy to overcome this difficulty may be to *assume* integrability and try to write down a consistent TQ-Baxter equation directly, based on basic properties of the theory such as the symmetry and also on educated guesses. Such a strategy

worked quite well for $\mathcal{N} = 4$ supersymmetric Yang-Mills theory in four dimensions [26] and it may not be so far-fetched to hope for the same for the adjoint QCD₂. To achieve this, it would be important to understand better the connection between TQ-equations and physics of QCD-like theories in two dimensions. This is one of the main reasons for studying generalized QCD₂ in this paper. Also, the deformation of the Yang-Mills Lagrangian by a potential $V(B)$ may be important for finding an aforementioned integrable point of the adjoint QCD₂.

We also perform a similar analysis for the *large representation limit* of generalized Yang-Mills with the SU(2) gauge group coupled to quarks in the spin J representation of SU(2). This is the limit introduced recently for the 't Hooft model in [27], in which J is sent to infinity keeping $\lambda_J \equiv g_{\text{YM}}^2 J$ fixed. A similar double-scaling limit was studied extensively in the context of large charge limits of conformal field theory (see e.g [28–34] and references therein for the large charge double scaling limits and [35–41] for the large representation limits of line defects) since it simplifies the dynamics of interacting conformal field theories and makes it accessible through analytic methods while still maintaining important qualitative features of the original theory. As in those examples, it was found in [27] that the dynamics of the 't Hooft model simplifies in the large representation limit and the meson spectrum can be determined by *exactly the same* integral equation as in the large N_c limit. We extend this result to the generalized QCD₂. Our result also provides evidence that the structures found in the large N_c limit (the connection to integrability and the analytic structure in the complex quark-mass plane) may persist even at finite N_c .

The outline of the paper is the following: In **Section 2**, we review the generalized QCD₂ and the derivation of the integral equation determining the mass spectrum of the mesons. In **Section 3**, we rewrite the integral equation into the TQ-Baxter equation. In **Section 4** we analyze the analytical structure of the spectrum of mesons as a function for generic $V(B)$ and complex quark masses, and we study in detail the example of a cubic potential $V(B)$ that demonstrates all the relevant features of the generic case. In **Section 5**, we generalize the analysis to the large representation limit of the generalized QCD₂ with gauge group $SU(2)$. Finally we discuss future directions in **Section 6**.

2 Generalized QCD in the Large N limit

Generalized QCD₂ is described by the Lagrangian [5]:

$$\mathcal{L} = \frac{N}{8\pi} \text{tr} B \wedge F - \text{tr} \frac{N}{4\pi} g^2 V(B) + \text{tr} \bar{\Psi} (i\gamma^\mu D_\mu - m) \Psi, \quad (2.1)$$

where both the field strength $F = dA$ and the B -field are in the adjoint representation of $SU(N)$, for simplicity we consider a single flavor of quarks Ψ in the fundamental representation¹. Classical power counting implies that *any* potential for the scalar field B is allowed in two dimensions. In this work, we take the potential for the B -field to be given by the formal power series²:

$$V(B) = \sum_{n=2}^{\infty} v_n \text{tr} B^n. \quad (2.2)$$

Integrating out the field B gives: $dA = 2g^2 V'(B)$, therefore, by fixing $V(B) = \frac{1}{8\pi} B^2$, the usual Yang-Mills Lagrangian is retrieved. While one could expect that the theory could admit a discrete and positive spectrum of mesons when the potential $V(B)$ is bounded from below, one of the results of this paper is that this is not a sufficient condition. This is due due to instabilities produced by the odd powers in (2.2). Instead, the more conservative choice $v_{2n+1} = 0, v_{2n} > 0$, guarantees positivity and discreteness of the spectrum. At the same time, one of the surprising facts we illustrate in this paper is that also for unbounded-from below potentials, there exist regions of the parameter space where the theory admits a positive spectrum in the large N limit.

Lightcone gauge. As for the standard 't Hooft model [42], upon going to lightcone coordinates³ $x_{\pm} = \frac{1}{\sqrt{2}}(x_0 \pm x_1)$, and fixing lightcone gauge $A_- = 0, A_+ = A$, the lagrangian presentation (2.1) simplifies and reduces to:

$$\mathcal{L} = \frac{N}{8\pi} \text{tr} B \partial_- A - \text{tr} \frac{N}{4\pi} g^2 \sum_{n=2}^{\infty} v_n \text{tr} B^n + \text{tr} \bar{\Psi}_a (i\gamma_{\mu} \partial^{\mu} - m_a - \gamma_- A) \Psi_a. \quad (2.3)$$

Here, we can regard all the $v_n \text{tr} B^n$ terms as interaction vertices, so the only Feynman rules are:

$$\begin{aligned} \longrightarrow &= \frac{p_-}{2p_+ p_- - m^2 + i\epsilon}, & \begin{array}{c} \nearrow \\ \searrow \end{array} \text{---} &= \frac{-8\pi}{k_-}, & \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} &= \frac{inv_n}{4\pi}. \end{aligned} \quad (2.4)$$

Henceforth in an arbitrary Feynman diagram, any propagator from a B^n vertex must be connected with a quark line. In the large N limit, the only Feynman diagrams that survive are the planar ones; an example of a planar contribution to the quark self energy is reported

¹The generalization to multi-flavors is analogous to the one for the 't Hooft model [42].

²This is not the most generic form of a self-interaction term: also power of traces would be allowed.

³The conventions on lightcone are the same of §2.1 of [1].




Figure 2: Diagrammatic representation of Bethe-Salpeter equation (2.9).

schematically illustrated in Figure 2, that in this model takes the form:

$$\left[2p_+ - m^2 \left(\frac{1}{p_-} + \frac{1}{p_- - q_-} \right) \right] \phi(p, p-q) = (\Gamma(q) + \Gamma(p-q)) \phi(q) + \int_0^{q_-} dk_- K(k, p; q, q) \phi(k) \quad (2.9)$$

In this model, (2.9) can be expressed in the following contour-integral form [5]:

$$\begin{aligned} & \left[2\pi^2 \lambda - (\alpha + 1) \left(\frac{1}{x} + \frac{1}{1-x} \right) \right] \phi(x) = \\ & = -2\pi \oint_{\gamma_\infty} \frac{dz}{2\pi i} V'(z) \left[iz \frac{z^2 + \pi^2}{(z^2 - \pi^2)^2} \frac{2}{\pi} \arctan\left(\frac{\pi}{iz}\right) \left(\frac{1}{x} + \frac{1}{1-x} \right) \phi(x) + \right. \\ & \quad \left. + 2 \frac{z^2}{(z^2 - \pi^2)^2} \int_0^1 dy \frac{1}{(x-y)^2} \exp\left(\frac{2}{\pi} \arctan\left(\frac{\pi}{iz}\right) \log\left(\frac{x(1-y)}{y(1-x)}\right)\right) \phi(y) \right], \end{aligned} \quad (2.10)$$

where: $x = p_-/q_-$ is the momentum fraction, $\alpha = \pi m^2/g^2 - 1$, and $\phi(x)$ is the reduced meson wavefunction: $\phi(x) = \int dp_+ \Phi(p, r)$ ($\Phi(p, r) = \langle \Omega | \bar{\psi} \psi | \Phi \rangle$). The contour integral is along a curve γ_∞ enclosing all the singularities but the one at $z = \infty$. This, as 't Hooft equation, is an eigenproblem in $x \in [0, 1]$ for the mesons wavefunctions $\phi(x)$ and masses μ^2 .

Explicit form. We found closed-form expressions for the result of the contour integrations. The first term in the square bracket can be readily evaluated:

$$\begin{aligned} \alpha_n &= 2\pi \oint \frac{dz}{2\pi i} in z^n \frac{z^2 + \pi^2}{(z^2 - \pi^2)^2} \frac{2}{\pi} \arctan\left(\frac{\pi}{iz}\right) = \\ &= \begin{cases} 0 & n \text{ odd} \\ 2\pi^{n-1} n^2 (-1 + H(\frac{n-1}{2}) + \log(4)) & n \text{ even} \end{cases}, \end{aligned} \quad (2.11)$$

where $H(n)$ are the Harmonic numbers⁴.

⁴The combination $(H(\frac{n-1}{2}) + \log(4))$ is a rational number for $n \in 2\mathbb{Z}$.

Denoting $m = n - 2$ and employing the variable:

$$\Lambda := \log\left(\frac{x(1-y)}{y(1-x)}\right), \quad (2.12)$$

the second term is expressed as⁵:

$$\begin{aligned} p_n(\Lambda) &= 2\pi \oint \frac{dz}{2\pi i} \frac{n z^{n+1}}{(z^2 - \pi^2)^2} \exp\left(\Lambda \frac{2}{\pi} \arctan\left(\frac{\pi}{iz}\right)\right) = \\ &= 4\pi n \sum_{q=0}^{\lfloor \frac{m}{2} \rfloor} \sum_{p=0}^{\lfloor \frac{m}{2} \rfloor - 1 - q} \frac{(-2i)^{m-2p-2q} (q+1) \pi^{2q+2p}}{(m-2q-2p)!} \Lambda^{m-2q-2p} \kappa(p), \\ \kappa(p) &= \sum_{k_0, \dots, k_p=0}^{2p} \binom{2p}{k_0, \dots, k_p} \prod_{t=2}^p \left(\frac{1}{2t+1}\right)^{k_t} \delta\left(\sum_{l=0}^p k_l - 2p\right) \delta\left(\sum_{l=1}^p l k_l - p\right). \end{aligned} \quad (2.13)$$

From this analytical expression⁶, it is evident that for any n even (odd) $p_n(\Lambda)$ is a polynomial with real (imaginary) coefficients of degree $n - 2$ in Λ containing only even (odd) powers thereof. The imaginary coefficients of the odd-terms guarantee simultaneously the hermiticity of the equation and the following parity of the eigenfunctions $\phi_n(x) = (-1)^n \phi_n(1-x)$.

Using the formulae above, the generalized 't Hooft equation is:

$$\mu^2 \phi(x) = \left(\alpha + 1 - \sum_{n=1}^{\infty} v_{2n} \alpha_{2n}\right) \left(\frac{1}{x} + \frac{1}{1-x}\right) \phi(x) - \int_0^1 dy \frac{\sum_{n=2}^{\infty} v_n p_n(\Lambda)}{(x-y)^2} \phi(y). \quad (2.14)$$

We report here the first few terms as computed from the expressions above:

$$\begin{aligned} \sum_{n=1}^{\infty} v_{2n} \alpha_{2n} &= 8\pi v_2 + \frac{160}{3} \pi^3 v_4 + \frac{744}{5} \pi^5 v_6 + \frac{31616}{105} \pi^7 v_8 + \frac{32440}{63} \pi^9 v_{10} + \dots \\ \sum_{n=2}^{\infty} v_n p_n(\Lambda) &= 8\pi v_2 - 24i\pi v_3 \Lambda + 32\pi v_4 (-\Lambda^2 + \pi^2) - \frac{40i\pi}{3} v_5 (-2\Lambda^3 + 7\pi^2 \Lambda) + \dots \end{aligned} \quad (2.15)$$

As expected, by setting $v_n = \frac{1}{8\pi} \delta_{2,n}$ we recover the familiar 't Hooft equation of QCD₂:

$$\mu^2 \phi(x) = \alpha \left(\frac{1}{x} + \frac{1}{1-x}\right) \phi(x) - \int_0^1 dy \frac{1}{(x-y)^2} \phi(y). \quad (2.16)$$

⁵In practice, rather than using the explicit expression we exhibit in the text, it is more convenient to directly evaluate the integral below by taking the residue at infinity for any specific n .

⁶This was also observed empirically in [5].

A direct comparison between (2.14) and (2.16) illustrates that the self-interaction potential for the B -field at the level of the mesons effectively *renormalizes* the mass of the quarks α , to:

$$\alpha \rightarrow \tilde{\alpha} = \alpha + 1 - \sum v_{2n} \alpha_{2n}, \quad (2.17)$$

and modifies the kernel (that is an effective inter-quark potential) by adding logarithmic corrections.

3 Generalized 't Hooft equation as a TQ-system

3.1 ν -space representation

Following the approach initiated for the 't Hooft model [1,2], we perform a Fourier transform with respect to the rapidity variables $\theta = \frac{1}{2} \log \frac{x}{1-x}$:

$$\phi(y) = \int_{-\infty}^{+\infty} \frac{d\nu}{2\pi} \left(\frac{x}{1-x} \right)^{-i\nu/2} \Psi(\nu), \quad \Psi(\nu) = \int_0^1 \frac{dx}{2x(1-x)} \left(\frac{x}{1-x} \right)^{i\nu/2} \phi(x). \quad (3.1)$$

In ν -space, the generalized 't Hooft equation (2.14) takes the form ($2\pi^2\lambda = \mu^2$):

$$\lambda \int_{-\infty}^{+\infty} d\nu' \frac{\pi(\nu - \nu') \Psi(\nu')}{2 \sinh\left(\frac{\pi}{2}(\nu - \nu')\right)} = \underbrace{\left[\frac{2\tilde{\alpha}}{\pi} + \sum_{n=2}^{\infty} v_n p_n(\mathcal{I}) \right]}_{=:f(\nu)} \Psi(\nu), \quad (3.2)$$

where the polynomial $p_n(\mathcal{I})$ is obtained by replacing in $p_n(\Lambda)$ of (2.13) powers of Λ^n with \mathcal{I}_{n+2} :

$$\mathcal{I}_{n+2} = \left(\frac{\left(\frac{x}{1-x}\right)^{\frac{i\nu}{2}}}{\frac{2}{\pi}(x-1)x} \right)^{-1} \int_0^1 dy \frac{\left(\frac{y}{1-y}\right)^{\frac{i\nu}{2}} \left(\log\left(\frac{y}{1-y}\right) - \log\left(\frac{x}{1-x}\right) \right)^n}{(x-y)^2}. \quad (3.3)$$

E.g. the first few terms are (cf. (2.15)):

$$\sum_{n=2}^{\infty} v_n p_n(\mathcal{I}) = 8\pi v_2 \mathcal{I}_2 - 24\pi i v_3 \mathcal{I}_3 + 32\pi v_4 (-\mathcal{I}_4 + \pi^2 \mathcal{I}_2) - \frac{40i\pi}{3} v_5 (-2\mathcal{I}_5 + 7\pi^2 \mathcal{I}_3) + \dots \quad (3.4)$$

Computing the principal parts integrals in (3.3) is made straightforward by the following observation⁷:

$$\mathcal{I}_{n+2} = \left(\frac{2}{i}\right)^n \frac{d^n}{d\nu^n} \mathcal{I}_2, \quad \mathcal{I}_2 = \nu \coth\left(\frac{\pi\nu}{2}\right), \quad (3.6)$$

as \mathcal{I}_2 coincides with the integral computed in eq. (3.3) of [1] for the standard QCD₂. We list a few of them to give a flavor of the resulting expressions:

$$\mathcal{I}_3 = \frac{i(\pi\nu - \sinh(\pi\nu))}{\sinh^2\left(\frac{\pi\nu}{2}\right)}, \mathcal{I}_4 = 2\pi \frac{2 - \pi\nu \coth\left(\frac{\pi\nu}{2}\right)}{\sinh^2\left(\frac{\pi\nu}{2}\right)}, \mathcal{I}_5 = 2i\pi^2 \frac{3\sinh(\pi\nu) - \pi\nu(\cosh(\pi\nu) + 2)}{\sinh^4\left(\frac{\pi\nu}{2}\right)} \quad (3.7)$$

Henceforth, we can straightforwardly compute the function $f(\nu)$ given any potential $V(B)$ expressed via a polynomial form⁸:

$$f(\nu) = \alpha + 1 + \sum_{n=1}^{\infty} v_{2n} \alpha_{2n} + \sum_{n=2}^{\infty} v_n p_n(\mathcal{I}). \quad (3.8)$$

As an example, taken $V(B)$ to be a quartic polynomial, the resulting expression for $f(\nu)$ is:

$$\begin{aligned} f(\nu) = \frac{2}{\pi} \left(\alpha + 1 - 8\pi v_2 - \frac{160\pi^3}{3} v_4 \right) + \frac{8\pi}{\tanh\left(\frac{\pi\nu}{2}\right)} (v_2\nu + 4\pi^2 v_4\nu - 6v_3) \\ + \frac{8\pi^2}{\sinh^2\left(\frac{\pi\nu}{2}\right)} \left(3v_3\nu + 8\pi v_4\nu \coth\left(\frac{\pi\nu}{2}\right) - 16v_4 \right) \end{aligned} \quad (3.9)$$

3.2 Discreteness of the spectrum and instabilities

Although for any potential there is an associated bound state equation (3.2), arbitrary choices of $V(B)$ will not always lead to positive mass spectrum for the mesons⁹. This was firstly empirically observed for the case of a cubic potential in [5], where, upon numerical direct solution of (2.14), it was found that for high values of v_3 the model exhibits tachyonic bound states. Below we use the reformulation of the eigenproblem (2.14) in ν -space in order to

⁷If one neglects the principal part integration, this follows by iteration from the definition of \mathcal{I}_n :

$$\frac{d}{d\nu} \mathcal{I}_{n+2} = \frac{i}{2} \left(\frac{\left(\frac{x}{1-x}\right)^{\frac{i\nu}{2}}}{\frac{2}{\pi}(x-1)x} \right)^{-1} \int_0^1 dy \left(\log\left(\frac{y}{(1-y)}\right) - \log\left(\frac{x}{(1-x)}\right) \right) \frac{\left(\frac{y}{1-y}\right)^{\frac{i\nu}{2}} \log^n\left(\frac{y(1-x)}{x(1-y)}\right)}{(x-y)^2} = \frac{i}{2} \mathcal{I}_{n+3}. \quad (3.5)$$

A more thorough proof is provided in Appendix A where we compute them as Fourier transform of distributions.

⁸In ν -space the hermiticity of (3.2) for $v_i \in \mathbb{R}$ is evident from the fact that in the products $v_{2n+1} \mathcal{I}_{2m+1}$ are always real.

⁹This is a generalization of the condition $\alpha > -1$ in 't Hooft model that ensures a stable positive spectrum of mesons. In QCD₂ this condition corresponds to imposing reality of the quark masses: $\pi \frac{m^2}{g^2} - 1 = \alpha$.

systematically analyze the analytical structures and instabilities of the generalized 't Hooft model and illustrates a rich and interesting analytical structure.

Exactly as discussed for the 't Hooft model [1], the generalized 't Hooft equation in ν -space takes the form of a homogeneous Fredholm equation of the second kind:

$$\phi(\nu) = \lambda \int_{-\infty}^{+\infty} d\nu' \kappa(\nu, \nu') \phi(\nu'), \quad \kappa(\nu, \nu') := \frac{\pi(\nu - \nu')}{2 \sinh\left(\frac{\pi}{2}(\nu - \nu')\right)} \frac{1}{\sqrt{f(\nu) f(\nu')}}. \quad (3.10)$$

where $f(\nu)$ is defined in (3.2). The finiteness of the $L^2(\mathbb{R}^2)$ of $\kappa(\nu, \nu')$, guaranteed by $f(\nu \in \mathbb{R}) \neq 0$, ensures that this self-adjoint kernel defines a Hilbert-Schmidt integral operator with discrete and non-degenerate spectrum. Since from (3.2) follows that $\Psi(\nu)$ has poles at the location of zeros of $f(\nu)$, when the latter has zeros on the real axis, the eigenfunctions are not normalizable and do not correspond to physical bound states.

Henceforth, upon studying the zeros of $f(\nu)$ in the complex plane of the couplings, one determines the regions where the generalized QCD produces a physical confining theory. We defer to section 4 a detailed discussion of the rich analytical structure of generalized QCD as function of the (complexified) couplings.

We conclude this subsection by remarking that the region of coupling space where the meson spectrum is positive and real does not correspond to imposing that $V(B)$ is bounded from below as one could expect. Even potentials that are unbounded from below, such as polynomials with odd leading powers, surprisingly admit physical regions in the coupling space where the spectrum is real and discrete. A prototypical example of this feature, is the one of a cubic $V(B)$, analyzed thoroughly in section 4.3. Moreover, even for potentials that are bounded from below, for large values of any of the odd couplings v_{2n+1} , the spectrum develops instabilities, and becomes tachyonic at large values thereof. Indeed, already at the classical level, a potential with odd couplings does not have their minimum at $B \equiv 0$ and tachyonic modes are ultimately just a manifestation of the instability of the vacuum at the origin. Naturally, this classical analysis receives corrections from quantum effects: from the discussion in section 4, one can verify that even for a potential of the form $V(B) = v_2 B^2 + v_3 B^3 + v_4 B^4$, even though $B = 0$ is never the true minimum for $v_3 < 0$, there is a critical value \bar{v}_3 (depending on v_2 and v_4) such that for any $|v_3| < \bar{v}_3$ the model has a real and discrete spectrum of mesons.

Normalization and discrete symmetries. Having identified where the model admits a discrete meson spectrum, in this paragraph we restrict to such coupling range. The normal-

ization of the eigenfunction in ν -space is fixed by (cf. eq. 3.11 of [1]):

$$\lambda_n \delta_{nm} = \lambda_n \int_0^1 dx \phi_n(x) \phi_m^*(x) = \int_{-\infty}^{+\infty} \frac{d\mu}{2\pi} \frac{f(\mu)}{2\pi} \Psi_n(\mu) \Psi_m^*(\mu). \quad (3.11)$$

Convergence of this norm imposes that eigenfunctions decay at least polynomially fast on the real axis. Furthermore, since the function $f(\nu)$ enjoys the discrete symmetry:

$$f(\nu) = \check{f}(-\nu) \equiv f(-\nu) \Big|_{v_{2n+1} \rightarrow -v_{2n+1}}, \quad (3.12)$$

the solutions Ψ_n will be also eigenfunction thereof:

$$\Psi(\nu) = (-1)^n \check{\Psi}(-\nu), \quad \check{\Psi}(\nu) \equiv \Psi(\nu) \Big|_{v_{2n+1} \rightarrow -v_{2n+1}}. \quad (3.13)$$

This discrete symmetry is a direct generalization of the parity symmetry $f(\nu) \rightarrow f(-\nu)$ of 't Hooft model, and we consequently still refer to that as *parity*.

3.3 TQ-equation

In the following, we choose the generalized YM potential $V(B)$ such that $f(\nu \in \mathbb{R}) \neq 0$. According to the discussion of the previous subsection 3.2, this ensures that the model admits a physical discrete spectrum of mesons with real masses and normalizable eigenfunction. In particular, the right hand side $f(\nu)\Psi(\nu)$ of (3.2) does not have any singularities in the strip¹⁰ $|\text{Im } \nu| < 2$ where we can analytically continue both sides of (3.2). Then, following [1, 2], we introduce the *Q-functions*:

$$Q(\nu) = \sinh\left(\frac{\pi\nu}{2}\right) \left[\frac{2\tilde{\alpha}}{\pi} + \sum_{n=2}^{\infty} v_n p_n(\mathcal{I}) \right] \Psi(\nu) = \sinh\left(\frac{\pi\nu}{2}\right) f(\nu) \Psi(\nu), \quad (3.14)$$

that are defined to be regular in the strip, to decay less-than exponentially at $|\nu| \rightarrow \infty$. Moreover, these *Q-functions* satisfy the related integral equation:

$$Q(\nu) = \lambda \sinh\left(\frac{\pi}{2}\nu\right) \int_{-\infty}^{+\infty} d\nu' \frac{\pi(\nu - \nu')}{2 \sinh\left(\frac{\pi}{2}(\nu - \nu')\right)} \frac{Q(\nu')}{\sinh\left(\frac{\pi}{2}\nu'\right) f(\nu')}, \quad (3.15)$$

¹⁰The discussion of this subsection follows closely the approach of section 3.2 of [1], and we refer the reader to that paper for details concerning all the derivations.

together with the *quantization* conditions

$$Q(0) = Q(\pm 2i) = 0. \quad (3.16)$$

This is completely analogous to the case of 't Hooft model; in particular, a computation identical to the one presented in section 3.2 of [1] shows that $\Psi(\nu)$ is a solution of (3.2) decaying polynomially at infinity if and only if the associated $Q(\nu)$ (through (3.14)), analytical in $|\text{Im } \nu| < 2$, solves the following TQ-Baxter equation:

$$\boxed{Q(\nu + 2i) + Q(\nu - 2i) - 2Q(\nu) = \frac{-4\pi\lambda}{f(\nu)} Q(\nu) = \frac{-4\pi\lambda Q(\nu)}{\frac{2\tilde{\alpha}}{\pi} + \sum_{n=2}^{\infty} v_n p_n(\mathcal{I})}}, \quad (3.17)$$

subject to the quantization conditions (3.16), and decaying less-than-exponentially at infinity in the whole strip.

Thus, for any given generalized Yang-Mills potential, the only modification to the TQ-Baxter equation is contained inside the single function $f(\nu)$, computed in closed form (through equations (3.8), (2.13),(2.11)) for given any polynomial $V(B)$.

Asymptotic decay. The asymptotic decay of the eigenfunctions can be further refined upon studying the large- ν limit of (3.17). Assuming the asymptotic behavior $Q(\nu) \asymp e^{\pi k \nu} q(\nu)$, with $q(\nu)$ periodic by $2i$, and bounded by any exponential, (3.17) is:

$$0 = Q(\nu + 2i) + Q(\nu - 2i) - 2Q(\nu) + \frac{4\pi\lambda}{8\pi\nu \sum_{n=1}^{\infty} n^2 v_{2n}} Q(\nu) \asymp -4e^{\pi k \nu} \sin^2(\pi k) q(\nu), \quad (3.18)$$

from which one deduces the stronger conditions of exponential decay of $\Psi(\nu)$ at $|\text{Re } \nu| \rightarrow \infty$ in the strip $(-2i, 2i)$, that refines the polynomial decay guaranteed by square integrability¹¹.

3.4 Inhomogenous extension

The TQ-system (3.17) admits solutions with the given decay at infinity and quantization only at the discrete spectrum $\lambda = \lambda_n$ of the integral operator (3.10). The TQ-Baxter equation is a simultaneous problem for Q and λ and, as such, it can be regarded as hard to solve analytically as the problem with started with. However, in [1, 2] a notable conceptual and

¹¹This result could have been imported directly from [1] given that in eq. (3.32) the r.h.s. does not depend on λ .

practical simplification specific to this problem was obtained upon considering an auxiliary *inhomogeneous* problem, obtained by modifying (3.2) and (3.16), so that (3.17) admits solutions for *any* $\lambda \in \mathbb{R}$. This turned out to be extremely useful as it allowed one to solve the TQ-equation in a two-step procedure: we first solve the simpler inhomogeneous problem, and then use this solutions to obtaine the discrete spectrum of the homogeneous problem. This ultimately leads to a fastly-convergent asymptotic analytic expansion for the spectral data in a systematic way. We refer the readers to section 3.2, 3.3 of [1] and [2] for a more thorough discussion.

Remarkably, the same discussion and technique also apply to the generalized QCDs. We extend (3.10) to an inhomogeneous problem by:

$$\frac{F(\nu)}{\sqrt{f(\nu)}} = \phi(\nu|\lambda) - \lambda \int_{-\infty}^{\infty} d\nu' \kappa(\nu, \nu') \phi(\nu'|\lambda), \quad (3.19)$$

$$F_+(\nu) = \frac{\nu}{\sinh(\frac{\pi}{2}\nu)}, \quad F_-(\nu) = \frac{1}{\sinh(\frac{\pi}{2}\nu)}. \quad (3.20)$$

The inhomogeneous terms (3.20) are the same as in [1, 2] and have definite parity under the discrete symmetry (3.13). Each of those drives a unique solution of the inhomogeneous integral equation depending on the continuous parameter λ :

$$F_{\pm}(\nu) = f(\nu) \Psi_{\pm}(\nu|\lambda) - \lambda \int_{-\infty}^{+\infty} d\nu' \frac{\pi(\nu - \nu')}{2 \sinh(\frac{\pi}{2}(\nu - \nu'))} \Psi_{\pm}(\nu'|\lambda), \quad (3.21)$$

whose solutions inherits definite parity:

$$\Psi_{\pm}(\pm\nu) = \pm \check{\Psi}_{\pm}(\nu). \quad (3.22)$$

The *inhomogeneous* Q-function defined by (3.14):

$$Q_{\pm}(\nu|\lambda) = \sinh\left(\frac{\pi\nu}{2}\right) f(\nu) \Psi_{\pm}(\nu|\lambda) = \mp Q_{\pm}(-\nu|\lambda) \quad (3.23)$$

are also shown to satisfy the same TQ-equation of the homogeneous problem (3.17):

$$Q_{\pm}(\nu + 2i|\lambda) + Q_{\pm}(\nu - 2i|\lambda) - 2Q_{\pm}(\nu|\lambda) = \frac{-4\pi\lambda}{f(\nu)} Q_{\pm}(\nu|\lambda), \quad (3.24)$$

but at any values of $\lambda \in \mathbb{R}$, upon assuming the same asymptotic, but subject to modified

quantization conditions¹²:

$$Q_+(\pm 2i) = \pm 2i, \quad Q_+(0) = 0, \quad Q_-(\pm 2i) = Q_-(0) = 1. \quad (3.25)$$

The Wronskian constructed with the two solutions $Q_\pm(\nu|\lambda)$ results to be a constant:

$$W(\nu|\lambda) = Q_+(\nu + i|\lambda)Q_-(\nu - i|\lambda) - Q_+(\nu - i|\lambda)Q_-(\nu + i|\lambda) = W(i|\lambda) = 2i. \quad (3.26)$$

Importantly, as for the 't Hooft model, the *complete integrability* of the kernel of the integral equation (3.2), entails that the the spectral determinants

$$D_+(\lambda) = \prod_{n=0}^{\infty} \left(1 - \frac{\lambda}{\lambda_{2n}}\right) \exp\left(\frac{\lambda}{\lambda_{2n}}\right), \quad D_-(\lambda) = e^{2\lambda} \prod_{n=0}^{\infty} \left(1 - \frac{\lambda}{\lambda_{2n+1}}\right) \exp\left(\frac{\lambda}{\lambda_{2n+1}}\right), \quad (3.27)$$

can be expressed in terms of the solutions of the inhomogeneous problem via the following TBA-like integral relations (cf. section 3.3 of [1] and Appendix B of [2]):

$$\begin{aligned} \partial_\lambda \log(D_+(\lambda)D_-(\lambda)) &= C_0 + \int_{-\infty}^{+\infty} d\nu \frac{\pi}{2} \left[\frac{Q_+(\nu|\lambda) \partial_\nu Q_-(\nu|\lambda) - Q_-(\nu|\lambda) \partial_\nu Q_+(\nu|\lambda)}{f(\nu)} + R^{(0)}(\nu) \right], \\ \partial_\lambda \log\left(\frac{D_+(\lambda)}{D_-(\lambda)}\right) &= - \int_{-\infty}^{+\infty} d\nu \frac{\pi^2}{2} \frac{Q_+(\nu|\lambda) Q_-(\nu|\lambda)}{\sinh(\pi\nu) f(\nu)}, \end{aligned} \quad (3.28)$$

These integrals are convergent (up to fixing a function $R^{(0)}(\nu)$ that is potential-dependent) as a consequence of the less-than-exponential asymptotic growth of the $Q_\pm(\nu)$.

The spectral determinants (3.27) encode in their series expansion in λ the spectral sums:

$$\log D_\pm(\lambda) = (1 \mp 1)\lambda - \sum_{s=2}^{+\infty} s^{-1} G_\pm^{(s)} \lambda^s, \quad G_+^{(s)} = \sum_{n=0}^{\infty} \frac{1}{\lambda_{2n}^s}, \quad G_-^{(s)} = \sum_{n=0}^{\infty} \frac{1}{\lambda_{2n+1}^s}, \quad (3.29)$$

and have their only zeros in correspondence with the eigenvalues of the homogeneous problem. The constant C_0 and the function $R^{(0)}(\nu)$ in (3.28) are fixed case-by-case to ensure convergence of the integral together with the correct normalization imposed by the lowest order in λ in (3.29), $(1 \mp 1)\lambda$, that is universal.

¹²Note that this changes the analytical properties of the solutions $\Psi(\nu|\lambda)$ compared to $\Psi_n(\nu)$, allowing for poles at the boundary of the strip.

3.5 Solving the TQ-equation: general strategy

We now generalize the approaches in [1, 2], and derive systematic weak-coupling and asymptotic expansions.

3.5.1 Weak-coupling expansion of the spectral sum.

Let us first discuss the weak-coupling expansion of the spectral sums (3.29).

The function $f(\nu)$ contains linear powers of ν , and trigonometric functions of ν invariant under $\nu \rightarrow \nu \pm 2i$. By collecting the trigonometric parts multiplying the linear term in ν , the transfer matrix can always be cast in the form:

$$Q(\nu + 2i|\lambda) + Q(\nu - 2i|\lambda) - 2Q(\nu|\lambda) = \frac{-2\lambda T_N(\nu)}{\nu + T_D(\nu)} Q(\nu|\lambda) \quad (3.30)$$

with:

$$T_D(\nu \pm 2i) = T_D(\nu), \quad T_N(\nu \pm 2i) = T_N(\nu). \quad (3.31)$$

In the case of 't Hooft model, this corresponds to $v_n = \delta_{2n} \frac{1}{8\pi}$, where $T_N(\nu) = 2\pi \tanh\left(\frac{\pi\nu}{2}\right)$, $T_D = \frac{2\alpha}{\pi} \tanh\left(\frac{\pi\nu}{2}\right)$, for generic potentials, the expressions become soon quite lengthy. For instance already with a cubic coupling one finds

$$T_N(\nu) = \frac{2}{8v_2 \coth\left(\frac{\pi\nu}{2}\right) + 24\pi v_3 \operatorname{csch}^2\left(\frac{\pi\nu}{2}\right)}, \quad T_D(\nu) = \frac{\frac{2}{\pi}(1 + \alpha + 8\pi v_2) - 48\pi v_3 \coth\left(\frac{\pi\nu}{2}\right)}{8\pi v_2 \coth\left(\frac{\pi\nu}{2}\right) + 24\pi^2 v_3 \operatorname{csch}^2\left(\frac{\pi\nu}{2}\right)} \quad (3.32)$$

One can show¹³ that two solutions of the TQ-system (3.17) are given by:

$$Q_1(\nu|\lambda) = \frac{i}{2} (\nu + T_D(\nu)) \operatorname{M} \left(1 + \frac{i}{2} (\nu + T_D(\nu)), 2, -i\lambda T_N(\nu) \right), \quad (3.34)$$

$$Q_2(\nu|\lambda) = \Gamma \left(1 + \frac{i}{2} (\nu + T_D(\nu)) \right) \operatorname{U} \left(1 + \frac{i}{2} (\nu + T_D(\nu)), 2, -i\lambda T_N(\nu) \right). \quad (3.35)$$

Due the appearance of poles inside in $\nu + T_D(\nu)$ and $T_N(\nu)$, and of the logarithmic branch

¹³This follows from the identities in formulas 13.4.1 and 13.4.15 of [43] satisfied by the confluent hypergeometric functions that we report here specialized to our case for convenience:

$$\begin{aligned} \frac{\Gamma(a)}{a-1} \left[\operatorname{U}(a-1, 2, z) + a(a-1)\operatorname{U}(a+1, 2, z) - 2(a-1)\operatorname{U}(a, 2, z) \right] &= \frac{z}{a-1} \Gamma(a) \operatorname{U}(a, 2, z) \\ (a-2)\operatorname{M}(a-1, 2, z) + a\operatorname{M}(a+1, 2, z) - 2(a-1)\operatorname{M}(a, 2, z) &= \frac{z}{a-1} (a-1)\operatorname{M}(a, 2, z) \end{aligned} \quad (3.33)$$

cut, those two functions do not correspond directly to solutions of our problem, as they do not have the correct analytical structure in the strip $|\text{Im } \nu| < 2$. The latter of those problems, is solved analogously to [1, 2] by modifying (3.34) to:

$$\begin{aligned}
M_+(\nu|\lambda) &= (\nu + T_D(\nu)) e^{\frac{i}{2}\lambda T_N(\nu)} M\left(1 + \frac{i}{2}(\nu + T_D(\nu)), 2, -i\lambda T_N(\nu)\right) \\
&= (\nu + T_D(\nu)) e^{\frac{i}{2}\lambda T_N(\nu)} \sum_{n=0}^{\infty} \frac{\left(1 + \frac{i}{2}(\nu + T_D(\nu))\right)_n}{n!(n+1)!} (-i\lambda T_N(\nu))^n \\
M_-(\nu|\lambda) &= \frac{1}{2} \left(e^{\frac{i}{2}\lambda T_N(\nu)} \Sigma(\nu|\lambda) + e^{\frac{i}{2}\lambda \tilde{T}_N(-\nu)} \tilde{\Sigma}(-\nu|\lambda) \right), \\
\Sigma(\nu|\lambda) &= 1 + \sum_{n=1}^{\infty} \frac{\left(\frac{i}{2}(\nu + T_D(\nu))\right)_n}{n!(n-1)!} (-i\lambda T_N(\nu))^n \left[\psi\left(\frac{1}{2}\right) - \psi(n) - \psi(n+1) \right. \\
&\quad \left. + \psi\left(n + \frac{i}{2}(\nu + T_D(\nu))\right) \right],
\end{aligned} \tag{3.36}$$

that are convergent series in powers of λ , solving (3.24) without branch cuts, and with definite parity

$$M_{\pm}(\nu) = \mp \check{M}_{\pm}(-\nu) = \mp M_{\pm}(-\nu) \Big|_{v_{2n+1} \rightarrow -v_{2n+1}}. \tag{3.37}$$

Then, the two combinations

$$Q_{\pm}(\nu|\lambda) = A_{\pm}(\lambda, \nu) M_{\pm}(\nu|\lambda) + i\lambda T_D(\nu) B_{\pm}(\lambda, \nu) M_{\mp}(\nu|\lambda) \tag{3.38}$$

are two independent solutions of the inhomogenous TQ-system (3.24), provided that we can fix the two functions A_{\pm} and B_{\pm} that must be invariant under $\nu \rightarrow \nu \pm 2i$ such that at each order in an expansion in series of λ , they cancel the poles of M_{\pm} . For any given potential and for any given order in λ , this is a well defined problem we can uniquely solve. Henceforth, this expansion produces at each order in λ an exact solution of the inhomogenous TQ system (3.24), that we can use to extract the spectral sums (3.29) via:

$$\begin{aligned}
G_{\pm}^{(s)} + G_{\pm}^{(s)} &= \frac{1}{s-1} \left[\frac{\partial^{s-1}}{\partial \lambda^{s-1}} \int_{-\infty}^{+\infty} d\nu \frac{\pi}{2} \frac{Q_+(\nu|\lambda) \partial_{\nu} Q_-(\nu|\lambda) - Q_-(\nu|\lambda) \partial_{\nu} Q_+(\nu|\lambda)}{f(\nu)} \right]_{\lambda=0} \\
G_{\pm}^{(s)} - G_{\pm}^{(s)} &= \frac{-1}{s-1} \left[\frac{\partial^{s-1}}{\partial \lambda^{s-1}} \int_{-\infty}^{+\infty} d\nu \frac{\pi^2}{2} \frac{Q_+(\nu|\lambda) Q_-(\nu|\lambda)}{\sinh(\pi\nu) f(\nu)} \right]_{\lambda=0}
\end{aligned} \tag{3.39}$$

3.5.2 Asymptotic expansion of eigenvalues

We next derive an asymptotic expansion for the spectrum by employing the same strategy used in §5.3 of [1] (see also [2]). The leading order behavior in $-\lambda \rightarrow \infty$ of (3.36) is (see

formulas 13.5.1, 13.5.2 of [43]) $M_{\pm}(\nu|\lambda) \sim (-\lambda)^{-\frac{i}{2}(\nu+T_D(\nu))} + \mathcal{O}(\lambda^{-1})$. Then, for at asymptotic values of λ , we can look for solutions of (3.24) with the following structure:

$$S(\nu|\lambda) = (-\lambda)^{-\frac{i\nu}{2} - \frac{i}{2}T_D(\nu)} \sum_{n=0}^{\infty} S_n(\nu) \lambda^{-n}, \quad (3.40)$$

where a factor $(-\lambda)^{-\frac{i}{2}T_D(\nu)}$ is periodic in $\nu \rightarrow \nu \pm 2i$. By plugging the ansatz (3.40) into the TQ equation, one obtains, order by order in λ , an equation for $S_n(\nu)$ that admits solutions in the form $S_n(\nu) = S_0(\nu)\sigma_n(\nu)$. As a result, one obtains

$$S(\nu|\lambda) = (-\lambda)^{-\frac{i\nu}{2}} S_0(\nu) \sum_{n=0}^{\infty} \frac{(1 + \frac{i}{2}\nu + \frac{i}{2}T_D(\nu))_n (\frac{i}{2}\nu + \frac{i}{2}T_D(\nu))_n}{n! (iT_N(\nu))^n}, \quad (3.41)$$

where $(a)_n$ is the Pochhammer symbol $(a)_k = \Gamma(a+k)/\Gamma(a)$. Henceforth, to completely determine (3.40), it is enough to solve the equation

$$S_0(\nu + 2i) = \frac{2T_N(\nu)}{\nu + T_D(\nu)} S_0(\nu), \quad (3.42)$$

whose solutions are determined on a case-by-case basis, depending on the analytical structure of the transfer matrix, see for instance the discussion in [1]. A solution to (3.42) can be easily determined, if one relaxes the analyticity requirements:

$$\tilde{S}_0(\nu) = 2^{\frac{i\nu}{2}} (iT_N(\nu))^{-\frac{1}{2}(i\nu)} \Gamma\left(\frac{iT_N(\nu)}{2} + \frac{i\nu}{2} + 1\right). \quad (3.43)$$

Even though this fails to be analytical in the strip¹⁴, it provides an indirect proof for (3.40): substituting this $\tilde{S}_0(\nu)$ in (3.40) one sees explicitly that the series in (3.41) coincides with the asymptotic expansion at $\lambda \rightarrow -\infty$ of (3.35) [43], which we know to satisfy the inhomogeneous TQ system.

Knowing $S_0(\nu)$ allows us construct the asymptotic solutions of (3.24) as:

$$Q_{\pm}(\nu|\lambda) = P_{\pm}(\nu|\lambda)S(\nu|\lambda) \mp \check{P}_{\pm}(\nu|\lambda)\check{S}(-\nu|\lambda), \quad (3.44)$$

where $P_{\pm}(\nu|\lambda)$, that must be invariant under $\nu \rightarrow \nu \pm 2i$, can be fixed by requiring the resulting $Q_{\pm}(\nu|\lambda)$ are analytical in the strip $|\text{Im } \nu| \leq 2$. In order to determine the eigenvalues λ_n

¹⁴Depending on the potential, slight modification of this solution can be more analytically appealing. For instance, in the case of non-vanishing v_2 , it is more convenient to construct product expansions akin to the one in footnote 27 of [1].




Figure 3: Confront between the asymptotic expression for the wavefunction and the numerical solutions for $\alpha = 0.5$, $v_3 = 0.0026$.

of the homogeneous system (3.17), it is enough to find the zeros of the spectral determinants $D_{\pm}(\lambda)$ through the integral formulae (3.28).

This procedure is readily applied to arbitrary choices of the potential $V(B)$. In section 3.6 we illustrate it thoroughly for the specific case of a cubic potential.

3.5.3 Asymptotic expansion of wavefunctions

The solutions (3.36) provide also an asymptotic expansion for the wavefunctions valid at $\nu \gg 1$. This is retrieved by evaluating (3.23) at the spectral values $\lambda = \lambda_n$:

$$\Psi_n(\nu) \sim \frac{\mathcal{Q}(\nu, \lambda_n)}{\sinh \frac{\pi\nu}{2} f(\nu)} = \frac{\nu + t_D(\nu)}{\sinh \frac{\pi\nu}{2} f(\nu)} e^{\frac{i}{2} \lambda_n t_N(\nu)} \text{M} \left(1 + \frac{i}{2} (\nu + t_D(\nu)), 2, -i \lambda_n t_N(\nu) \right), \quad (3.45)$$

where $t_{N/D}(\nu) = T_{D/N}(\nu \gg 1)$. For potentials containing only even powers, the solutions at negative values of ν can be simply retrieved by parity. Instead for potentials containing also odd powers, it is convenient to consider symmetrization/antisymmetrization of the numerical solutions under $\nu \rightarrow -\nu$ and $v_{2n+1} \rightarrow -v_{2n+1}$.

Specializing the formula above to the cubic potential case, and setting $v_2 = \frac{1}{8\pi}$, we find:

$$\begin{aligned} \Psi_n(\nu) &= \frac{2 \sinh \left(\frac{\pi\nu}{2} \right)}{48\pi^2 v_3 + \sinh(\pi\nu)} \exp \left(\frac{i\pi\lambda(\cosh(\pi\nu) - 1)}{48\pi^2 v_3 + \sinh(\pi\nu)} \right) \\ &\times \text{M} \left(1 + i \frac{2(\cosh(\pi\nu) - 1)\alpha + 48v_3\pi^3\nu + \pi(\nu - 48v_3\pi) \sinh(\pi\nu)}{2\pi(48\pi^2 v_3 + \sinh(\pi\nu))}, 2, -\frac{2i\pi\lambda(\cosh(\pi\nu) - 1)}{48\pi^2 v_3 + \sinh(\pi\nu)} \right) \end{aligned} \quad (3.46)$$

We tested this analytical expression against the numerical results in Figure 3.

3.6 Solving the TQ-equation: cubic coupling

We now describe the details of the general procedure to derive the asymptotic expansion, outlined above, by taking a cubic potential as an illustrative example:

$$V(B) = \frac{1}{8\pi}B^2 + v_3B^3, \quad f(\nu, \alpha, v_3) = \frac{2\alpha}{\pi} + \nu \coth\left(\frac{\pi\nu}{2}\right) + 24\pi v_3 \frac{\pi\nu - \sinh(\pi\nu)}{\sinh^2\left(\frac{\pi\nu}{2}\right)}. \quad (3.47)$$

3.6.1 Constructing asymptotic solutions

Following the discussion in section 3.5.2, two asymptotic solutions at $\lambda \rightarrow -\infty$ that solve the inhomogeneous TQ-equation:

$$Q_{\pm}(\nu + 2i) + Q_{\pm}(\nu - 2i) - 2Q(\nu) = \frac{-4\pi\lambda}{f(\nu, \alpha, v_3)} \quad (3.48)$$

and that are eigenstate of *parity*, can be expressed as:

$$\begin{aligned} Q_{\pm}(\nu|\lambda) &= P_{\pm}(z, y|\lambda)S(\nu|\lambda) \mp \check{P}_{\pm}(-z, -y|\lambda)\check{S}(-\nu|\lambda), \\ S(\nu|\lambda) &= (-\lambda)^{-\frac{i\nu}{2}} S_0(\nu) \sum_{n=0}^{\infty} \frac{\left(1 + \frac{i}{2}\nu + \frac{i}{2}T_D(z, y)\right)_n \left(\frac{i}{2}\nu + \frac{i}{2}T_D(z, y)\right)_n}{n! (2\pi i \lambda T_N(z, y))^n}, \end{aligned} \quad (3.49)$$

where we found convenient to use the variables

$$z = \tanh\left(\frac{\pi\nu}{2}\right), \quad y = \operatorname{csch}(\pi\nu), \quad (3.50)$$

that are invariant under $\nu \rightarrow \nu \pm 2i$, to express (3.32):

$$T_N(z, y) = \frac{z}{1 + 48\pi^2 v_3 y}, \quad T_D(z, y) = \frac{\frac{2\alpha z}{\pi} - 48\pi v_3}{1 + 48\pi^2 v_3 y}. \quad (3.51)$$

The function $S_0(\nu)$ satisfies the recursion relation:

$$S_0(\nu + 2i) = \frac{2T_N(z, y)}{\nu + T_D(z, y)} S_0(\nu). \quad (3.52)$$

We express a solution of (3.52) as:

$$S_0(\nu) = (2\pi)^{-\frac{1}{2} - \frac{i\nu}{2}} \frac{G\left(2 + \frac{i\nu}{2}\right) G\left(\frac{1}{2} - \frac{i\nu}{2}\right)}{G\left(1 - \frac{i\nu}{2}\right) G\left(\frac{3}{2} + \frac{i\nu}{2}\right)} \sigma(\nu) \quad (3.53)$$

where $\sigma(\nu)$ solves¹⁵:

$$\sigma(\nu + 2i) = \left(\frac{-48\pi v_3 + 2\alpha z/\pi}{\nu} + 1 + 48\pi^2 v_3 y \right)^{-1} \sigma(\nu). \quad (3.54)$$

For what follows, an explicit form¹⁶ of the solution to this equation is not needed. We will only employ the defining recursion (3.54), together with the fact that the resulting $S(\nu)$ is analytical in $\text{Im } \nu \in [-2i, 2i]$.

Removing the singularities. While the functions $Q_{\pm}(\nu|\lambda)$ in (3.53) solve the TQ system, they are not yet analytical in the strip. There are two sources of singularities: poles of increasing order at $z = 0$, where $T_N(z, y)$ vanishes, and singularities due to poles of $T_D(z, y)$, that are located at: $1 + 48\pi^2 v_3 y = 0$. In order to remove those unwanted singularities, at each order in λ , we fix the function $P(z, y|\lambda)$ such that the singularities cancel in the product. This can be done uniquely at each order by taking:

$$\begin{aligned} P_{\pm}(z, y|\lambda) &= C_{\pm}(\lambda) \left[\sum_k R_{\pm}^{(k)}(z^{-1}, y, \log(-\lambda)) \lambda^{-k} \right] \left[\sum_k T^{(k)}(y) \lambda^{-k} \right] \\ &:= C_{\pm}(\lambda) R_{\pm}(z^{-1}|\lambda) T(z, y|\lambda), \end{aligned} \quad (3.56)$$

where R_{\pm} and T remove respectively the singularities of the first and second kind; the function $C_{\pm}(\lambda)$ is fixed by the normalization of the Q functions. The procedure is analogous to what done for the 't Hooft model in section 5.3.1 of [1] (see also [2] for the case $\alpha = 0$), to which we refer the reader for more details. Using the combinations:

$$L = \log(-2\pi\lambda) + \gamma_E, \quad \beta = \alpha + 1, \quad (3.57)$$

¹⁵All the normalizations has been fixed so that at $v_3 = 0$, we recover the same function of the 't Hooft model (eq. (5.26) of [1]).

¹⁶A solution for this equation that is regular on the real axis has a representation in terms of Gamma functions:

$$\tilde{\sigma}(\nu) = \frac{(1 + 48\pi^2 v_3 y)^{-1 - \frac{i\nu}{2}} \Gamma\left(1 + \frac{i\nu}{2} + \frac{i}{2} T_D(z, y)\right)}{\Gamma\left(1 + \frac{i\nu}{2}\right)}, \quad (3.55)$$

that yet fails to be analytical in the entire strip $[-2i, 2i]$. A full solution can be constructed by employing the Weierstrass representation for the $\Gamma(x)$ as discussed in footnote 27 of [1].

one obtains that function R_{\pm} is a polynomial in the variable $c = i\pi z^{-1}$, whose first non-trivial orders are:

$$\begin{aligned} \frac{(-\lambda)^{\frac{1\pm 1}{2}} R_{\pm}(c, L)}{\sqrt{i} C_{\pm}(\lambda)} &= 1 \mp \frac{1}{\lambda^2} \left[\pm \frac{6iv_3\beta c^2}{\pi^3} + \frac{\beta - 384\pi^4 v_3^2 c}{4\pi^4} \right] + \mathcal{O}(\lambda^{-3}) \\ C_{\pm}(\lambda) &= \exp \left[\frac{\beta}{2\pi^2 \lambda} - \frac{(\beta - 6(8\pi^2 v_3)^2)L - \beta^2}{2\pi^4 \lambda^2} + \mathcal{O}(\lambda^{-3}) \right]. \end{aligned} \quad (3.58)$$

Computing either terms is straightforward; we omit them here as they get soon rather lengthy.

Analogously, we proceed order by order in removing the singularities in the strip $[-2i, 2i]$ due to zeros of the equation $1 + 48\pi^2 v_3 y = 0$. that has three solutions in this strip at: $\nu_0, \pm i \mp \nu_0$ with $\nu_0 = -\frac{1}{\pi} \operatorname{arcsinh}(48\pi^2 v_3)$. It turns out that all this poles are removed by $T(z, y|\lambda)$ being a polynomial in the variables \mathbf{t} and \mathbf{a} :

$$\mathbf{t} = i\pi z \frac{\mathbf{a}}{1 + 48\pi^2 v_3 y}, \quad \mathbf{a} = \alpha - 24\pi^2 v_3 / z \quad (3.59)$$

$$\begin{aligned} T(\mathbf{t}; \mathbf{a}, \lambda) &= 1 - \frac{\mathbf{a}\mathbf{t}}{2\pi^4 \lambda} + \frac{\mathbf{a}^2 \mathbf{t}}{8\pi^8 \lambda^2} (\mathbf{t} - 2\pi^2) - \frac{\mathbf{a}^3 \mathbf{t}}{48\pi^{12} \lambda^3} (\mathbf{t}^2 - 6\pi^2 \mathbf{t} + 10\pi^4) + \\ &\quad + \frac{\mathbf{a}^4 \mathbf{t}}{384\pi^{16} \lambda^4} (\mathbf{t}^3 - 12\pi^2 \mathbf{t}^2 + 52\pi^4 \mathbf{t} - 84\pi^6) + \mathcal{O}(\lambda^{-5}). \end{aligned} \quad (3.60)$$

This is the same function $T(\chi; \alpha, \lambda)$ in eq. (5.37) of [1] that we determined for the 't Hooft model¹⁷, up to the replacements $\chi \rightarrow \mathbf{t}, \alpha \rightarrow \mathbf{a}$.

3.6.2 Determining the eigenvalues

Having constructed the inhomogeneous solutions $Q_{\pm}(\nu|\lambda)$ via the formula (3.49), we can extract from them the spectral determinants $D_{\pm}(\lambda)$ applying the integral formulas (3.28). This is done following the same strategy thoroughly explained in Appendix B of [1], and we defer more details of hwo this is applied to the case in hand to Appendix B. There we determine explicitly all the coefficients $d_n^{(\pm)}$ in the asymptotic expansion at $\lambda \rightarrow -\infty$ of the spectral determinants:

$$D_{\pm}(\lambda < 0) = \sum_{n=0}^{\infty} \frac{d_n^{(\pm)}}{\lambda^n} \quad (3.61)$$

up to arbitrary order in λ .

This expression has still to be analytically continued to $\lambda > 0$. This is done by averaging

¹⁷In particular all the considerations made there on its closed form expression also hold here.

among the analytical continuation in the upper- and lower- half plane $D_{\pm}^{(\pm)}(\lambda) = D_{\pm}(-e^{\pm i\pi}\lambda)$ via the *continuation formula* [1, 2]:

$$D_{\pm}(\lambda > 0) \asymp \frac{1}{2}(D_{\pm}^{(-)}(\lambda) + D_{\pm}^{(+)}(\lambda)) = \delta_{\pm} e^{\Phi_{\pm}(\lambda)} \cos\left(\frac{\pi}{2} \Xi_{\pm}(\lambda)\right),$$

$$\Phi_{\pm}(\lambda) = \frac{1}{2}(\mathcal{D}_{\pm}(l + i\pi) + \mathcal{D}_{\pm}(l - i\pi)), \quad \Xi_{\pm}(\lambda) = \frac{1}{i\pi}(\mathcal{D}_{\pm}(l + i\pi) - \mathcal{D}_{\pm}(l - i\pi)),$$
(3.62)

where $\mathcal{D}_{\pm}(\lambda) = \log(D_{\pm}(\lambda))$, $l = \log(\lambda)$ and δ_{\pm} is an influential integration constant. Thus, the eigenvalues are determined by the quantization condition:

$$\Xi_{+}(\lambda_{2n}) = \Xi_{-}(\lambda_{2n+1}) = 2n + 1 \quad n \in \mathbb{N}. \quad (3.63)$$

Upon keeping terms up to order $\mathcal{O}(\lambda^{-4})$ in (3.60) (3.58), in appendix B we illustrate how




Figure 4: Comparison between the numerical eigenvalues and the analytical expression (3.64), at $\alpha = -0.5$. For this value of the mass, the expected critical point is $v_3^{(\text{II})} \simeq -0.0058$.

to derive the following result

$$n = 2\lambda_n - \frac{2\alpha \log(2\lambda_n)}{\pi^2} + \frac{\alpha^2}{\pi^4 \lambda_n} - \frac{3}{4} - \frac{2\alpha \log(4\pi e^{\gamma_E})}{\pi^2} + \frac{1}{\pi^3} \mathcal{I}(\alpha, v_3)$$

$$+ \frac{1}{2\pi^6 \lambda_n^2} \left(\alpha^3 + (-1)^n \pi^2 (\alpha + 1) + (-1)^n v_3^2 \left(C_1 + \frac{288 \log^2(\lambda_n)}{\pi^2} + C_2 \log(\lambda_n) \right) \right) + \mathcal{O}(\lambda_n^{-3}),$$
(3.64)

where the constants C_1 and C_2 are

$$\begin{aligned}
C_1 &= -288(2\alpha + 3) + \frac{(\alpha + 1)}{\pi^2} \left[144(2\gamma_E^2 + 1 + 2\log^2(2) + \log(4) + \right. \\
&\quad \left. + \gamma_E(2 + 4\log(2\pi)) + 2\log(\pi)(1 + \log(4\pi)) \right], \\
C_2 &= \frac{288(2\gamma_E + 1 + \log(4) + 2\log(\pi))}{\pi^2}.
\end{aligned} \tag{3.65}$$

and $\mathcal{I}(\alpha, v_3)$ is a complicated integral reported in (B.4). Upon setting $v_3 = 0$ we recover¹⁸ the same result of [1], and in particular this matches the known WKB expansion of 't Hooft model.

The asymptotic expansion (3.64) provides a new (exact) WKB-like expansion for generalized QCD, which, to our knowledge, was never discussed in the literature before. One can then solve the (truncated) equation (3.64) for any given values of n . In Figure 4 we report the first energy level so obtained and we compare them with the numerical eigenvalues obtained by direct solution of the ν -space 't Hooft equation by discretization.

Both results illustrate, as expected a positive discrete spectrum of mesons in the region predicted by the discussion in section 3.2, with a massless mode emerging at its boundary in correspondance with some particular value of the couplings. The emergence this massless pion away from the chiral limit, is not an accident and could be predicted analytically, and this is precisely the subject of the next section 4.

4 Analytical structure of the spectrum and critical points

The simple form of 't Hooft equation in ν -variables (3.2) allows for a straightforward qualitative analysis of the main features of the model in the complexified space of couplings. In particular, the analytical structure of the function $f(\nu)$ completely determines the analytical structure of the eigenvalues $\lambda_n(\alpha, \mathbf{v})$ as functions of the complexified couplings. In what follows, in order to emphasize the dependence on the complexified couplings, we denote $f(\nu)$ by $f(\nu, \alpha, \mathbf{v})$.

As already observed in section 3.2, the zeros of $f(\nu, \alpha, \mathbf{v})$ coincide with the poles of the eigenfunction $\Psi_n(\nu)$, so that whenever they do not lie on the real line, the model will admit a discrete spectrum of mesons with normalizable eigenfunctions. Among the zeros of $f(\nu, \alpha, \mathbf{v})$ the higher order zeros, where also the derivatives thereof vanish are special

¹⁸In particular note that the integral in the appendix reduces to the integral expression of the 't Hooft model, by setting $v_3 = 0$: $\mathcal{I}(\alpha, 0) = \int_{-\infty}^{+\infty} d\nu \frac{(\sinh(\pi\nu) - \pi\nu) \sinh(\frac{\pi\nu}{2})}{\nu \cosh^2 \frac{\pi\nu}{2} (\alpha \sinh(\frac{\pi\nu}{2}) + \frac{\pi\nu}{2} \cosh(\frac{\pi\nu}{2}))}$.

and distinct points. Already in the case of the 't Hooft model [1], which can be recovered by setting all the v_k to zero, we found that the location of double zeros of $f(\nu, \alpha)$ in the complex plane on ν determines infinitely many square-root branch points at some specific values $\alpha_k \in \mathbb{C}$ where one of the mesons becomes massless. At finite N_c these critical points are likely to become non-unitary interacting CFTs. This interesting structure extends and enriches greatly in the generalized Yang-Mills theory with polynomial potentials. In this section we describe the general features of the analytical structure, and of the type of criticality that we encounter in this very broad class of models by varying the polynomial potential $V(B)$. The reader can refer to the concrete example of a cubic self-interaction that we discuss thoroughly in section 4.3, that illustrates all the main features of the generic case.

\mathcal{PT} -symmetry and reality of the spectrum. It is important to note that physical choices of $V(B)$ shall not be limited to real values for the coefficients. Since the B -field is odd under \mathcal{PT} -symmetry (or equivalently, charge-conjugation), potentials with imaginary odd couplings, $v_{2n+1} \in i\mathbb{R}$, provide examples of \mathcal{PT} -symmetric quantum field theories. It is by now well-established that non-hermitian field theories enjoying unbroken \mathcal{PT} -symmetry, can have real spectra; we refer the reader to the classical papers [44–46] (and references therein) for a thorough discussion on the topic. However, \mathcal{PT} -symmetry can also be spontaneously broken, in which case the spectrum contains complex conjugate pairs, and in generalized QCD both cases happen (recently also observed in the context of non-unitary minimal models in [47–49]), see for instance section 4.3.2.

4.1 Multicritical points from collision-of-roots

By tuning the parameters of the potential $V(B)$, one can make $f(\nu, \alpha, \mathbf{v})$ to have zeros of higher order¹⁹. As we explain below, a n -th order zero corresponds to a n -th root-branch point of the eigenvalues, if it is not possible to deform the contour of integration to avoid the pinching singularity. At each of these n -th branch-points the eigenvalues behave as $\lambda \sim (\alpha - \alpha^{(n)})^{1/n}$, (and similarly as functions of all the other couplings), and correspond to multicritical points where exactly $n - 1$ mesons become simultaneously massless.

Collision-of-roots point and branch points. In general, roots of $f(\nu, \alpha, \mathbf{v})$, do not correspond to branch-point singularities of the spectrum, but only to poles of $\Psi(\nu)$. This is due to the fact that if the colliding roots are away from the integration contour, the integral

¹⁹For the physical potential, the order of the zeros is bounded from above by $\deg V(B)$.




Figure 5: Collision-of-roots points may or may not lead to singularities.

eigenproblem (3.2) is still well-defined. On the other hand, if two roots collide in such a way that the contour of integration is trapped in between, the spectrum will develop a branch cut²⁰. Only the collision-of-roots of type Figure 5a would lead to a singular point of λ , while the one of Figure 5b is not a singularity on this sheet of the complex plane since the contour is not trapped by the collision. However it will be a singularity on another sheet of the complex plane as we will shortly discuss.

Type of singular points. The precise type of singularity associated to this critical point is inferred by studying the monodromy of the zeros of the function f as we meromorphically continue the parameter α along a closed curve around any of the values of the critical values $\alpha^{(n)}$. If $\nu^{(n)}$ is the position of the n -th order zeros at $\alpha = \alpha^{(n)}$, as we move in the parameter space to $\alpha \neq \alpha^{(n)}$, the degeneracy of the roots will be generically split and we will be left with exactly n non degenerate roots $\nu_1 \cdots \nu_n$. Then it is easy to see that upon the analytical continuation of α along a closed curve winding around $\alpha^{(n)}$ a single time, $f(\nu, \alpha, \mathbf{v})$ encompasses a monodromy that permutes the roots by: $\nu_i \rightarrow \nu_{i+1}$. Because of this non-trivial monodromy, each root of these root may or may not cross the integration contour, if this happens, then the integral eigenproblem will differ by the respective residue terms.

The simplest case is the one of a square-root cut, as in Figure 5: in this case after a 2π analytical continuation $\alpha \rightarrow \tilde{\alpha}$, the eigenproblem will also undergo a non-trivial monodromy

²⁰This is very well known in the context of the S-matrix bootstrap where many singularities of these kind arise, cfr. e.g. [50].




Figure 6: Complex plane before and after the a 2π analytical continuation with non-trivial monodromy.

transformation, that act by adding two further residue terms:

$$\begin{aligned}
 f(\nu)\Psi(\nu) &= \tilde{\lambda}(\alpha) \int_{-\infty}^{+\infty} d\nu' \frac{\pi(\nu - \nu')}{2 \sinh \frac{\pi}{2}(\nu - \nu')} \Psi(\nu') \\
 &+ \tilde{\lambda}(\tilde{\alpha}) \text{Res} \left(\frac{\pi(\nu - \nu')}{2 \sinh \frac{\pi}{2}(\nu - \nu')} \Psi(\nu'; \alpha_0, \mathbf{v}_0), \nu' = \nu_1 \right) \\
 &- \tilde{\lambda}(\tilde{\alpha}) \text{Res} \left(\frac{\pi(\nu - \nu')}{2 \sinh \frac{\pi}{2}(\nu - \nu')} \Psi(\nu'; \alpha_0, \mathbf{v}_0), \nu' = \nu_2 \right).
 \end{aligned} \tag{4.1}$$

This indicates $\alpha^{(\text{II})}$ is a square-root branch point of the eigenvalues. This point coincides with the boundary of the allowed region for having a positive spectrum of mesons, and indeed we find a corresponding critical point where one of the meson becomes massless upon direct solution (analytical or numerical) of the spectral problem. The critical point that we found by solving asymptotically the spectrum in Figure 4, is exactly of this nature. To illustrate a more complicated case, let us examine the case of a zero of order 3. As we thoroughly explain in section 4.3 the typical situation is the one of figure Figure 7a, where the contour cannot be continuously deformed to avoid a pinching-of-roots singularity at some special values of the couplings $\alpha^{(\text{III})}, v_3^{(\text{III})}$. Then, it is easy to see that the monodromy of α under a 6π monodromy along a curve encircling the collision-of-root point $\alpha^{(\text{III})}$ is the one described in Figure 7b, Figure 7c and Figure 7d, i.e. a third order zero leads to a cubic-root branch cut of the spectrum and a tricritical point where two of the meson masses turn simultaneously to zero (cfr. Figure 17). It is straightforward from the graphical representation in Figure 7 to write down, on each of the three sheets represented there, the corresponding bound state




Figure 7: Monodromy of the roots around a 3rd order zero in the complex plane.

equation as we have done in (4.1). The generalization to arbitrary order of root is completely straightforward.

Geometry of the phase space. For any given polynomial $V(B)$ of degree n , the singularities of type p -th root will form a codimension $n - p$ locus in the phase space of the theory given that they generically solve a system of p equations. In particular, n -th-root branch points form a set of infinitely many isolated points in the space of couplings, $(n - 1)$ -th-root branch points form set of lines that may intersect at n -th root branch points, and so on. The typical structure of the first sheet of the phase space is the one in Figure 15. Each of the several connected components of these loci, lives in a unique sheet of the complex plane. On a given sheet, only the collision-of-roots which pinches the contour leads to singularities, while the others are singular points on other sheets, as we discuss more later. These critical loci correspond to the boundary of regions where the spectrum of mesons is positive and discrete. At the special points in the space of couplings, several mesons become massless (as in Figure 10 and Figure 12).




Figure 8: Collision-of-roots away from the contour are singularities in higher-sheet of the complex ν' -plane

4.2 Critical points beyond the first sheet

As anticipated already in the previous subsection, the singularities on the first sheet are only one of the many connected components forming the singular loci of the parameter space. Indeed, while collision-of-roots happening away from the integration contour are not singularity on the first sheet, they are singularities on other sheets of the complex plane: As we analytically continue beyond the cuts on the first sheet, the contour of integration for the integral equation picks up additional contributions from poles that crossed the contour (see (4.1) and Figure 7). After this happens, if these poles collide with other poles on the complex plane, they lead to singularities on the corresponding sheet. See e.g. Figure 8 (and the corresponding situation on the first sheet Figure 5b).

By comparing them with the analytical prediction or with the numerics, one can check that those singularities correspond to critical points where one of the higher mesons (instead of the lowest one) becomes tachyonic and triggers a critical phase transition.

4.3 A case study: cubic potential

In this section, we analyze the spectrum and phases of generalized QCD with a cubic potential: $V(B) = v_2 B^2 + v_3 B^3$ that is never bounded from below except for $v_3 = 0$. The latter is a purely quadratic potential and is the simplest example of a gYM potential $V(B) = v_2 B^2$. In this case we have

$$f(\nu, \alpha, v_2) = \frac{2(\alpha + 1 - 8\pi v_2)}{\pi} + 8\pi v_2 \nu \coth\left(\frac{\pi\nu}{2}\right) = 8\pi v_2 \left(\bar{\alpha} + \nu \coth\left(\frac{\pi\nu}{2}\right)\right). \quad (4.2)$$

Clearly, the problem depends only on the single variable $\bar{\alpha} = \frac{\alpha - 8\pi v_2 + 1}{8\pi v_2}$ and the entire structure is equivalent to the usual QCD₂ analyzed in [1, 2], upon the replacement $\bar{\alpha} \rightarrow \alpha$: the spectrum

has a square root branch cut in $\bar{\alpha} = -1$ for any values of v_2 , and the energy levels are real and positive for $v_2 > 0 (< 0)$ for $\bar{\alpha} > -1 (< -1)$.

For $v_3 \neq 0$, only two of the three couplings (α, v_2, v_3) are physically relevant and we can always canonically normalize $v_2 = \frac{1}{8\pi}$. From (3.8), we obtain

$$f(\nu, \alpha, v_3) = \frac{2\alpha}{\pi} + \nu \coth\left(\frac{\pi\nu}{2}\right) + 24\pi v_3 \frac{\pi\nu - \sinh(\pi\nu)}{\sinh^2\left(\frac{\pi\nu}{2}\right)}. \quad (4.3)$$

To identify regions in the parameter space where this model exhibits a real and positive spectrum of mesons, below we systematically study zeros of $f(\nu)$ in the complex space of the couplings (α, v_3) .

4.3.1 Real couplings

Let us start by considering real values of the quark masses corresponding to $\alpha \geq -1$. For arbitrary values of α and $v_3 \in \mathbb{R}$, zeros of $f(\nu)$ are all simple. Given any fixed value of α , there are two critical values $\pm v_3^{(\text{II})}(\alpha)$, where the two zeros closest to the real axis degenerate to a double zero at $\nu = \pm \nu^{(\text{II})}(\alpha) \in \mathbb{R}$:

$$f\left(\pm \nu^{(\text{II})}(\alpha), \alpha, \pm v_3^{(\text{II})}(\alpha)\right) = \partial_\nu f\left(\pm \nu^{(\text{II})}(\alpha), \alpha, \pm v_3^{(\text{II})}(\alpha)\right) = 0. \quad (4.4)$$

Solutions to this equation give a curve in Figure 10 in the plane (α, v_3) . As shown in Figure 9, while for $|v_3| < v_3^{(\text{II})}(\alpha)$ there are no real zeros of $f(\nu, \alpha, v_3)$, when $|v_3| > v_3^{(\text{II})}(\alpha)$ exactly two of them lie on the real axis. Therefore, the analytical structure of the bound state equation (3.2) implies that the spectrum of mesons is positive in a shaded region of Figure 10 bounded by the curve ²¹ (4.4). For any $v_3 \neq 0$, $f(\nu, -1, v_3 \neq 0)$ has two real zeros, (one of which is always at $\nu = 0$), and therefore the eigenfunction are not-normalizable. So, for $m^2 = 0$ the massless meson becomes unstable at $v_3 \neq 0$. So that, for points laying outside of the region bounded by the curve, the eigenfunction are not normalizable and the spectrum becomes tachyonic.

It is worth stressing that, although the cubic potential is not bounded from below, in the large N_c approximation, the mesons can have a well-defined positive spectrum.

²¹Analytic solutions for the coupled equations (4.4) are not available. However near $\alpha = -1$, one can study the solution $v_3^* = v_3^*(\alpha)$ analytically by linearizing (4.4) in ν since we know that for small $\alpha \rightarrow -1$, both $v_3^{(\text{II})}, \nu^{(\text{II})} \rightarrow 0$. The result of the linear approximation, $v_3^*(\alpha) \simeq \pm \frac{\sqrt{\alpha+1}}{8\sqrt{6}\pi^2}, \nu^*(\alpha) \simeq \pm \frac{\sqrt{6}\sqrt{\alpha+1}}{\pi}$, exhibit many of the features we demonstrate in the text.




Figure 9: Positions of the zeros (black) and poles (white) of $f(\nu, \alpha = 0.5, v_3)$ in the ν -complex plane for different values of v_3 . For $|v_3| < v_3^{(\text{II})}(0.5) \simeq 0.01176$ (Figure 9a, 9f) there are two distinct real zeros that degenerate for $v_3 = v_3^{(\text{II})}(0.5)$ (Figure 9b, 9e) at $\nu^{(\text{II})}(0.5) \simeq 0.974336$. In all the physical region $|v_3| < v_3^{(\text{II})}(0.5)$, there are no real zeros of $f(\nu, 0.5, v_3)$.




Figure 10: The shaded region correspond to the values of the couplings where the model has a positive spectrum of mesons, while the curve identifies the critical points where the theory has a massless mode.

Massless meson. The critical points $\left(v_3^{(\text{II})}(\bar{\alpha}), \bar{\alpha}\right)$ solving (4.4) are square-root branch points of the eigenvalues $\lambda(\alpha, v_3)$, at which the lowest meson becomes massless. One can infer the branch-cut structure by studying the monodromy of the roots under analytical continuation of the quark mass α along a curve that encircles $\bar{\alpha}$ an odd number of times. As illustrated in Figure 11, it exchanges the two simple zeros $\bar{\nu}_{\pm}$ of $f(\nu, \alpha, v_3)$, which eventually degenerate into $\nu^{(\text{II})}(\bar{\alpha})$.




Figure 11: We fix $\bar{\alpha} \simeq 2.12$, its corresponding positive critical point is $v_3^{(\text{II})}(\bar{\alpha}) \simeq 0.2$. The figure illustrates the positions of the zeros (black) of $f(\nu, \alpha(\theta), v_3^{(\text{II})}(\bar{\alpha}))$, for values of $\alpha(\theta) = 2.4 - 0.4e^{i2\pi\theta}$ that encircle the point $\bar{\alpha}$. Along this analytical continuation in the complex plane the two zeros are exchanged.


This shows that, by tuning the strength of the potential v_3 , for any real and positive value of the quark masses α , one can always produce a massless meson²². If we set $\alpha = -1$, where the quark are massless, the curve (4.4) degenerates into a point $v_3^{(\text{II})} = 0$.

We tested these results against the numerical solutions of the ν -space eigenproblem (3.2), as explained in Appendix D. The numerical solutions in Figure 12 illustrate a typical confining spectrum of mesons with Regge-like trajectories. The ground state becomes massless in correspondance with the critical point identified by the collision of zero of $f(\nu, \alpha, v_3)$.

4.3.2 Analytical continuation and tricritical point

In the previous section, we analyzed the spectrum of mesons for real values of the quark masses ($\alpha \geq -1$) and cubic couplings. We showed that at any fixed $v_3 > 0$, there is a square root branch-cut in the α plane at $\alpha = \alpha^{(\text{II})}(v_3)$, corresponding to a critical point in the first sheet, where the first meson becomes massless. In this section we are interested in the analytic continuation to negative values of α . Because of the branch cuts in the $\alpha, v_3 \in \mathbb{R}$ (see

²²This agrees with some preliminary exploration of [5] for some specific values of the coupling. Note also that this meson is not associated with chiral symmetry as in [1].



(a) First five eigenfunctions with cubic potential for $v_3 = 0.0026$.

(b) First 4 energy levels for $v_3 \in [0, f_3^*]$ spanning the whole physical spectrum.

Figure 12: Energy levels and eigenfunctions for $\alpha = 0.5$, $v_3^{(\text{II})}(0.5) = f_3^* \simeq 0.01176$.

Figure 10), this requires an extra care. Below, we first analytically continue v_3 to imaginary values keeping $\alpha > 1$, and we study the region $\alpha < -1$ afterwards.

Analytical continuation to imaginary couplings. At complex values of $v_3 \in \mathbb{C}$, the Hamiltonian is no longer hermitian, and the spectrum will be generically complex. Yet, if restrict to purely imaginary values of v_3 , i.e. we take the self-interaction of the B -field to be:

$$V(B) = \frac{1}{8\pi} B^2 + i|v_3| B^3, \quad (4.5)$$

the corresponding bound-state equation (3.2) enjoys \mathcal{PT} -symmetry²³, since the B field is odd under \mathcal{PT} symmetry as discussed in general in section 4. At fixed $\alpha > -1$, for $v_3 \in i\mathbb{R}$, there are no collision-of-root singularities in the first sheet, and no zeros of $f(\nu, \alpha, v_3)$ on the real line. So we expect for this region of the parameter to have a positive spectrum of mesons. Indeed, by solving numerically the eigensystem at imaginary values of v_3 in a neighborhood of the real axis, we find that the spectrum remains real and positive as shown in Figure 13. This means that \mathcal{PT} -symmetry is not spontaneously broken for $\alpha > -1$ at least in the neighborhood of the real axis of v_3 .

Critical points at $\alpha < -1$. The analytical continuation to imaginary values of v_3 allow us to also continue the spectrum to $\alpha < -1$ without crossing any of the branch cuts. More concretely, we first continue to²⁴ $v_3 = i\mathbb{R}^+$ at $\alpha > -1$, and only then we continue from

²³The structure of the potential here considered can be regarded as a field theory analogous of the quantum mechanical system $\mathcal{H} = p^2 + m^2 x^2 + ix^3$ considered e.g. in [44].

²⁴ $|v_3|$ has to be large enough to guarantee any pinching of root singularity on the contour, as we will discuss below.




Figure 13: Numerical solutions for the first 5 levels.




Figure 14: First two roots of $f(\nu', \alpha, v_3)$ upon analytical continuation. In (a) and (b), at fixed $\alpha > -1$, we analytically continue in v_3 to $v_3 \in i\mathbb{R}^+$. The red arrows illustrate the shift of the first two roots under this continuation. In (c), we then plot the result of the analytical continuation from $\alpha > -1$ to $\alpha < -1$ at fixed value $v_3 \in i\mathbb{R}$. The root crosses the real axis exactly at $\alpha = -1$. When the root crosses the real axis, the integration contour is deformed to avoid the singularity. For $iv_3 > 0$ it is analogous but with the root in the upper half plane moving in the lower half plane.

$\alpha > -1$ to $\alpha < -1$. In doing that, exactly one of the roots crosses the real axis, (as $f(\nu, -1, v_3)$ has always a root at $\nu = 0$), yet, as long as one makes sure that there are no collision of roots on the contour, one can deform the ν' integration contour in (3.2), to avoid the singular point. This is illustrated in Figure 14.

Collision-of-roots singularities happens at $\alpha < -1$, whenever, upon varying v_3 the first two roots in Figure 14 pinch the contour. This happens, for any $\alpha < -1$, at two isolated points on the imaginary axis $\pm v_3^{(\text{II})}(\alpha)$, determined by the solutions of the equation $f(\nu, \alpha, v_3) = 0$ and $\partial_\nu f(\nu, \alpha, v_3) = 0$ at fixed $\alpha < -1$. Upon decreasing α , one finds solutions $v_3^{(\text{II})}(\alpha) \in i\mathbb{R}$ only for $\alpha > \alpha^{(\text{III})} \simeq -1.27649$, while for $\alpha > \alpha^{(\text{III})}$ the corresponding solutions $v_3^{(\text{II})}(\alpha)$ split in the complex plane (red curves in Figure 15). The collision-of-roots singularity




Figure 15: Critical points for $v_3 \in \mathbb{R}^+$ or $v_3 \in i\mathbb{R}^+$ and $\alpha \in \mathbb{R}$. For $\alpha > -1$, the blue curves are loci of critical points lying at $\text{Im}(v_3) = 0$ and are exactly the one studied in Section 4.3.1 (cfr. Figure 10). The blue curve at $\alpha < -1$, is located at $\text{Re}(v_3) = 0$ while the red ones correspond to critical points for the complex spectrum. The black dot corresponds to the tricritical point.

ties are critical points in the spectrum where the first meson becomes massless. In Figure 15, we report the loci of critical points at $\alpha \in \mathbb{R}$ and v_3 in the upper half plane (the lower half plane of v_3 is symmetrical). The red curves correspond to the solutions for $\alpha > \alpha^{(\text{III})}$ and are indeed located in a region of the parameter space where the 't Hooft equation is neither hermitian nor \mathcal{PT} -symmetric, and the masses of mesons are $\lambda_n \sim M_n^2$ are complex. The critical points in this region correspond to points where one of the complex eigenvalues turns to zero. Any analytical continuation performed along a path on the first sheet of v_3 and α that does not intersect any of the blue or red curves and does not ends in the region shaded in red, leads to a discrete and positive spectrum of mesons for $\alpha > \alpha^{(\text{III})}$. Correspondingly, the \mathcal{PT} symmetry is not spontaneously broken on the first sheet²⁵.

²⁵Upon going around the cubic-root branch cut at $\alpha = \alpha^{(\text{III})}$ (i.e. in the region shaded with red in Figure 15) \mathcal{PT} -symmetry is broken spontaneously *non-critically* by the eigenfunctions. The same phenomenon was also recently illustrated for the non-unitary version of tricritical Ising model (the minimal model $M(2,7)$) in [48, 49]. Non-critical \mathcal{PT} -breaking means that in this region of parameter space, two of the meson masses degenerate and become complex conjugate, but with none of them turning to zero. Hence, in this case the transition to a spontaneously broken phase of \mathcal{PT} -symmetry phase is non-critical as there are no associated massless mode in the spectrum.




Figure 16: Integration contour in the ν' complex plane for $\alpha < \alpha^{(\text{III})}$ at fixed $v_3 = v_3^{(\text{III})}$. As α approaches the tricritical point, the contour is trapped between three singularities.

Tricritical point. As illustrated in the previous subsection, the line of critical points in the first sheet of the (α, v_3) plane for $\alpha < -1$, terminates at the special point:

$$\alpha^{(\text{III})} \simeq -1.27649, \quad v_3^{(\text{III})} \simeq \pm 0.00344328 i, \quad (4.6)$$

where $f(\nu)$ has a triple zero. Here the contour has a singularity due to the collision of three roots simultaneously as shown in Figure 16. The monodromy of the roots under analytical continuation in α , is exactly the one illustrated in Figure 7. Henceforth, according to the general discussion of section 4, one expects that the theory admits a description in terms of a tricritical CFT where two of the energy levels turn to zero simultaneously as $\lambda \sim \sqrt[3]{\alpha - \alpha^{(\text{III})}}$. In Figure 17 the result of the numerical spectrum after analytical continuation; we refer to Appendix D for a detailed explanation of the method we employed to analytically continue to $\alpha < -1$.

4.3.3 Singularities beyond the cuts

Each of the critical point we discussed is associated to a branch cut in the complex plane of the coupling. As a generalization to what happens in the 't Hooft model [1], as we analytically continue the model to complex values of α and v_3 , we probe a very rich and complicated multi-sheeted structure.

Critical points in the second sheet. For fixed $v_3 \in \mathbb{R}$, by analytically continuing in α through the square root cut, the two roots inevitably cross the integration contour, and the eigenproblem differs by a further contribution due to the contour integration around those singularities, as in Figure 6. In the second sheet of these cuts, we find infinitely many other




Figure 17: First 4 meson masses as a function of α at fixed $v_3 = v_3^{(\text{III})}$. The first two levels turn to zero at $\alpha = \alpha^{(\text{III})}$.


critical points where, at complex values of α , one of the roots that took a monodromy after analytical continuation collides with one of the higher singularities (e.g. Figure 18b). On each of the corresponding sheets, the k -th meson becomes massless, rather than the first energy level λ_0 . This is a generalization of what already discussed for the 't Hooft model [1], but at generic values of $v_3 \in \mathbb{R}$.

Tricritical points in higher sheets. A similar argument shows the existence of infinitely many tricritical points after the analytic continuation in α through the cubic root branch cut at (4.6). On the second sheet relative to this cut, one finds two infinite towers of cubic branch points, each associated with distinct values of v_3 and located respectively in the second and third sheet of the α -plane. We report them in Figure 19, as determined by the points corresponding to triple zeros of $f(\nu, \alpha, v_3)$ in the complex plane.


The two infinite towers are distinguished by the sign of the imaginary part of the corresponding $v_3^{(\text{III})}$ ²⁶. Each of those tricritical points is associated to a point where two of the higher meson masses, rather than the first and second, turn to zero simultaneously.

Second sheet critical points at imaginary v_3 . As one considers the region $v_3 \in i\mathbb{R}$, a new type of critical points in the second sheet arise. Following the same strategy of the previous subsection, it is not difficult to show that for any value of $\alpha \in \mathbb{R}$, there are infinitely

²⁶This can also be understood from the square root branch cut in the v_3 - α plane, present for any value of α , in correspondence of which the value of v_3 changes sign.



(a) Singularity in the second sheet of the α plane after analytical continuation through the cut along the real axis, for various values of v_3 .



(b) Collision of the first singularity, after analytical continuation with one of the higher.

Figure 18: Singularities in the second sheet of the α plane continued through the cut at fixed $v_3 \in \mathbb{R}$.





Figure 19: Position of the tricritical points in the complex α -plane. Each of them is associated with a distinct value of $v_3^{(\text{III})}$.



(a) Phase space of the theory for $\alpha > -1$ in the second sheet of the α plane, and $v_3 \in i\mathbb{R}$. In the shaded region, the model present a physical and confining spectrum of mesons.

(b) First collision of roots of $f(\nu, \alpha, v_3)$ at $v_3^{(\text{II},0)}$.

Figure 20: Collision of roots for imaginary v_3 and corresponding phase-space.

many double zeros $\pm v_3^{(\text{II},k)}(\alpha) \in i\mathbb{R}$, $k \in \mathbb{N}$, strictly ordered as:

$$\left| v_3^{(\text{II},0)}(\alpha) \right| \leq \left| v_3^{(\text{II},1)}(\alpha) \right| \leq \left| v_3^{(\text{II},2)}(\alpha) \right| \leq \dots, \quad (4.7)$$

where $f(\nu, \alpha, v_3)$ has a double zero at $\nu^{(\text{II},k)}(\alpha) \in i\mathbb{R}$:

$$f\left(\pm \nu^{(\text{II},k)}(\alpha), \alpha, \pm v_3^{(\text{II},k)}(\alpha)\right) = \partial_\nu f\left(\pm \nu^{(\text{II},k)}(\alpha), \alpha, \pm v_3^{(\text{II},k)}(\alpha)\right) = 0, \quad (4.8)$$

At the first of those critical points $\pm \nu_3^{(\text{II},0)}$, the two closest roots of $f(\nu)$ in the upper-half plane (that are always in the strip $\nu \in [0, 2i]$) collide. At fixed α , none of these collision-of-root point is a singularity in the first sheet. But, if one first analytically continue in α along a closed curve enclosing a the square root branch point, then $v_3^{(\text{II},k)}(\alpha)$ becomes a singular point where the first eigenvalues closes the gap and breaks spontaneously \mathcal{PT} -symmetry. This is exactly identical to what discussed in Section 6.3.1 of [1] (see Figure 16 there).

Note that the line of critical points at $\alpha < -1$ remains singular also in this second sheet, the resulting phase-space of the theory is illustrated in Figure 20a, where is interesting to note that these new locus of critical point intersects the one discussed in the previous subsection exactly at the tricritical point.

This can be generalized to any of the $\nu^{(\text{II},k)}$. It turns out that, at any fixed α , $v_3^{(\text{II},0)}(\alpha)$ is also a branch point of infinite degree in the v_3 plane. Indeed upon analytically continue in

v_3 around any closed curve encircling a single time $v_3^{(\Pi,k)}(\alpha)$, one observes a monodromy of the roots of the form $\nu_i \rightarrow \nu_{i\pm 1}$ where ν_i is the position of the roots of $f(\nu, \alpha, v_3)$ (with the exception of the one corresponding to $-v_3^{(\Pi,0)}(\alpha)$) and \pm depends on the sign of $\text{Re}(\nu_i)$. This is illustrated in Figure 21. Then, it is easy to see that under this analytical continuation, one can always find a sheet where $\nu^{(\Pi,k)}$ becomes a critical point and the corresponding k -th meson closes the gap and breaks spontaneously \mathcal{PT} -symmetry.

5 Mesons spectrum in the large representation limit

In this section we consider a theory with a finite number of colors N , but of fermions in a very large representation of the gauge group. This limit was also considered in [27] for the 't Hooft model. Here we illustrate that analogous results can be derived also for the generalized QCD case. Specifically, we consider an $SU(2)$ gauge theory coupled to a single flavor of quarks in the isospin J representation, and consider a scaling limit where $J \rightarrow \infty$.

Effective Hamiltonian for fermions. For our scopes, it is best to work with an Hamiltonian formulation of the theory. The starting point is the generalized YM Lagrangian in lightcone gauge, coupled to quarks in a generic representation of the gauge group. Not specifying yet $N = 2$ the Hamiltonian is:

$$\mathcal{L} = \frac{N}{4\pi} \text{tr} B \partial_- A_+ - \frac{Ng^2}{4\pi} \sum_{n=2}^{\infty} \text{tr} v_n B^n + \bar{\psi}_a (i\partial - m_a) \psi_a - \text{tr} A_+^A (\bar{\psi}_a \gamma_- T^A \psi_a), \quad (5.1)$$

and we take the generators T^A , $A = 1, \dots, N^2 - 1$ in a generic large dimensional representation of $SU(N)$. Integrating out the A_+ field in the path integral leads to the insertion of a delta function distribution setting

$$\frac{N}{4\pi} \partial_- B^A - \bar{\psi}_a \gamma_- T^A \psi_a = 0, \quad (5.2)$$

and leads to

$$\mathcal{L} = -\frac{Ng^2}{4\pi} \sum_{n=2}^{\infty} \text{tr}_{\text{adj}} \left(\frac{4\pi}{N} \right)^n v_n (\partial_-^{-1} \bar{\psi}_a \gamma_- T \psi_a)^n + \text{tr} \bar{\psi}_a (i\partial - m_a) \psi_a, \quad (5.3)$$




Figure 21: Monodromy of the roots under analytical continuation in v_3 around $v_3^{(\text{II},0)}$.

where with tr_{adj} we mean that the trace has to be taken in the adjoint representation:

$$\text{tr}_{\text{adj}} \left(\partial_-^{-1} \bar{\psi}_a \gamma_- T \psi_a \right)^n = \text{tr} [F^{A_1} \dots F^{A_n}] \prod_{i=1}^n \left(\partial_-^{-1} \bar{\psi}_a \gamma_- T^{A_i} \psi_a \right), \quad A_i = +, -, 3, \quad (5.4)$$

where F^A are the generators in the adjoint representation of $SU(N)$.

The component ψ_+ is non-dynamical because the field has no time derivatives in the lightcone gauge Lagrangian; it is fixed by the equations of motion to:

$$\psi_+ = \frac{im}{2} \partial_-^{-1} \psi_-. \quad (5.5)$$

Henceforth, one reduces the Lagrangian (2.1) to the following effective action for the only dynamical quark component ψ_- :

$$\mathcal{L}_{\text{eff}} = -\frac{Ng^2}{4\pi} \sum_{n=2}^{\infty} \text{tr}_{\text{adj}} \left(\frac{4\pi}{N} \right)^n v_n \left(\partial_-^{-1} \bar{\psi}_a \gamma_- T \psi_a \right)^n + \text{tr} \bar{\psi}_a (i\cancel{\partial} - m_a) \psi_a, \quad (5.6)$$

The standard quantization:

$$[\pi_-(x), \psi_-(y)] = \delta(x_- - y_-), \quad \pi(x) = \frac{\partial \mathcal{L}_{\text{eff}}}{\partial \psi_-(x)} = i\sqrt{2} \psi_+^\dagger(x), \quad (5.7)$$

leads to the effective Hamiltonian density:

$$\mathcal{H}_{\text{eff}} = -\frac{Ng^2}{4\pi} \sum_{n=2}^{\infty} \left(\frac{4\pi}{N} \right)^n v_n \text{tr}_{\text{adj}} \left(\partial_-^{-1} \bar{\psi}_a \gamma_- T \psi_a \right)^n - i\frac{m^2}{4} \text{tr} \psi_-^\dagger \partial_-^{-1} \psi_-, \quad (5.8)$$

The quarks depend on the two lightcone coordinates $\psi(y) = \psi(y_-, y_+)$, while in the potential only derivatives with respect to the y_- coordinate appear. To simplify the notation, we will momentarily drop the explicit “-” label of the coordinates, with the understanding that inside the argument of the fermions $\psi(y) \equiv \psi(y_-, y_+)$, and everywhere else $y \equiv y_-$. Using the identities in Appendix A of [27], each of the quark bilinears that appears in (5.8) can readily expressed as²⁷:

$$\partial_x^{-1} \psi_-^\dagger(x) \psi_-(x) = \int dy \partial_y^{-1} \delta(x-y) \psi_-^\dagger(y) T^a \psi_-(y) = \frac{1}{2} \int dy \text{sgn}(x-y) \psi_-^\dagger(x) T^a \psi_-(y). \quad (5.9)$$

²⁷A similar rewriting was also employed in this context in [51].

Then, the effective Hamiltonian is ($y_1 := x, y_0 = y_n$):

$$\begin{aligned}
H_{\text{eff}} = & -\frac{g^2}{4\pi} \sum_{n=2}^{\infty} \text{tr} (2\pi)^n v_n \left(\prod_{k=1}^n \int dy_k \text{sgn}(y_k - y_{k-1}) \psi_{-}^{\dagger}(y_k) T^{A_k} \psi_{-}(y_k) \right) \text{Tr} [F^{A_1} \dots F^{A_n}] \\
& - \text{tr} i \frac{m_a^2}{8} \int dx dy \psi_{-}^{\dagger}(x) \text{sgn}(x - y) \psi_{-}(y)
\end{aligned} \tag{5.10}$$

Large representation limit. Let us specialize to the group $SU(2)$. In the spin J representation, we have

$$\begin{aligned}
\psi_{-}^{\dagger} T^{+} \psi_{-}(x) &= \sum_{m=-J}^J \sqrt{(J-m)(J+m+1)} \psi_{m+1,-}^{*}(x) \psi_{-}^m(x), \\
\psi_{-}^{\dagger} T^{-} \psi_{-}(x) &= \sum_{m=-J}^J \sqrt{(J+m)(J-m+1)} \psi_{m,-}^{*}(x) \psi_{-}^{m+1}(x), \\
\psi_{-}^{\dagger} T^3 \psi_{-}(x) &= \sum_{m=-J}^J m \psi_{m,-}^{*}(x) \psi_{-}^m(x).
\end{aligned} \tag{5.11}$$

Following the approach of [27], in the large J limit $\frac{m}{J}$ is approximated by a continuous variable $\frac{m}{J} = \cos \theta$. Rather than carrying a representation index m , in this limit, the quarks depend explicitly on the continuous variable θ :

$$\psi_{-}(\theta, x) := \sqrt{J} \psi_{-}^m(x), \tag{5.12}$$

where, the additional factor \sqrt{J} guarantees the correct scaling of the canonical quantization in the limit $J \rightarrow \infty$:

$$\{\psi_{-}^{\dagger}(\theta, x), \psi_{-}(\theta', y)\} = J \left(\frac{1}{J \sin \theta} \delta(\theta - \theta') \right) \delta(x - y) \tag{5.13}$$

Accordingly, we must replace the trace over the representation by its continuous limit:

$$\sum_{m=-J}^J \rightarrow J \int_0^{\pi} d\theta \sin \theta, \tag{5.14}$$

Using the continuous fields $\psi_-(\theta, x)$, one derives [27] the following large J scaling of (5.11):

$$\begin{aligned}\psi_-^\dagger(x) T^\pm \psi_-(x) &\rightarrow J \int_0^\pi d\theta \sin^2 \theta \psi_-^*(\theta, x) \psi_-(\theta, x), \\ \psi_-^\dagger(x) T^3 \psi_-(x) &\rightarrow J \int_0^\pi d\theta \sin \theta \cos \theta \psi_-^*(\theta, x) \psi_-(\theta, x).\end{aligned}\tag{5.15}$$

Now, let us assume that the potential truncates at a certain order $\bar{n} \in 2\mathbb{N}$ (the reason for fixing \bar{n} even will be clear in a moment), and that the none of v_n scales with J . Then, because of the overall trace, the Hamiltonian is order J , and admits a double-scaling limit at $J \rightarrow \infty$ where the combination

$$4(2\pi)^{\bar{n}-1} g^2 J^{\bar{n}} = \lambda\tag{5.16}$$

is taken fixed. Taking $v_n = \mathcal{O}(J^0)$ suppresses any term $V(B)$ with $n < \bar{n}$ ²⁸. Under this conditions, in the large spin limit, (5.10) is:

$$\begin{aligned}H_{\text{eff}} &= -\frac{v_{\bar{n}}\lambda}{8} \text{tr} \left(\prod_{k=1}^{\bar{n}} \int dy_k \text{sgn}(y_k - y_{k-1}) \psi_{a,-}^\dagger(y_k) T^{A_k} \psi_{a,-}(y_k) \right) \text{tr} [F^{A_1} \dots F^{A_{\bar{n}}}] \\ &\quad - \text{tr} i \frac{m_a^2}{4} \int dx dy \psi_{a,-}^\dagger(x) \text{sgn}(x - y) \psi_{a,-}(y),\end{aligned}\tag{5.17}$$

where we intend that all the representation indexes are contracted through integrals over the respective θ variables and that we take the trace thereof ($\theta_1 = \theta_n$) to ensure gauge invariance. For the special case of $SU(2)$, only traces of an even number of adjoint generators are non-vanishing²⁹. Henceforth, odd powers in the potential of $V(B)$ do not contribute (this is the reason for setting $\bar{n} \in 2\mathbb{N}$) to in an $SU(2)$ theory.

As in [27], we introduce the bi-local field:

$$\langle y, \phi | \mathbf{M} | x, \theta \rangle = M(x, y; \theta, \phi) = \psi_-(\theta, x) \psi_-^*(\phi, y).\tag{5.18}$$

We will also use the notation $\mathbf{M}(x, y)$ whenever we want to specify the positions while keeping the short hand matrix notation. This notation is convenient for expressing contraction of bilinear fields in the continuum limit $\int d \cos \theta A(x, y, \theta, \phi) B(z, t, \theta, \lambda)$ in the short-hand matrix multiplication notation $\mathbf{A}(x, y) \mathbf{B}(z, t)$. After we have substituted the adjoint repre-

²⁸ Any of them can be retained by imposing a scaling limit on the $v_n = \mathcal{O}(J^{\bar{n}-n})$ at large J . Since the generalization to this case is straightforward, here we consider the simplest case where only the highest power contributes at large spin.

²⁹ We normalize the generators in the adjoint representation such that the killing form is $\kappa^{ab} = \begin{pmatrix} 0 & 1/2 & 0 \\ 1/2 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$.

sentation traces, the interaction term of \mathcal{H}_{eff} , keeping only the symmetric terms in θ , and indicating the integration measure as $d\mathbf{y} d\boldsymbol{\theta} = \frac{1}{\bar{n}!} \prod_{i=1}^{\bar{n}} d\theta_i \sin \theta_i dy_i$, takes the form :

$$\int d\mathbf{y} d\boldsymbol{\theta} \prod_{i=1}^{\bar{n}} \text{sgn}(y_k - y_{k-1}) \left[M(y_1, y_2; \theta_1, \theta_2) \widehat{M}(y_2, y_3, \theta_2, \theta_3) M(y_3, y_4, \theta_3, \theta_4) \widehat{M}(y_4, y_5; \theta_4, \theta_5) \cdots \right. \\ \left. + \text{perm.} \right], \quad (5.19)$$

where $\widehat{M}(y_i, y_j, \theta_i, \theta_j) := M(y_i, y_j, \theta_i, \theta_j) \cos(\theta_i - \theta_j)$. These quantities are conveniently expressed through the matrix notation introduced in (5.18):

$$\text{tr} \int \prod_{i=1}^{\bar{n}} (dy_i \text{sgn}(y_k - y_{k-1})) \left(\mathbf{M}(y_1, y_2) \widehat{\mathbf{M}}(y_2, y_3) \mathbf{M}(y_3, y_4) \cdots \right). \quad (5.20)$$

Then, for an $SU(2)$ theory, at large spin, the Hamiltonian (5.8) is:

$$H_{\text{eff}} = \frac{im^2}{8} \int dx dy \text{sgn}(x - y) M(x, y; \theta, \theta) + \\ + \frac{\lambda v_{\bar{n}}}{8} \int \prod_{i=1}^{\bar{n}} (dy_i \text{sgn}(y_k - y_{k-1})) \text{tr} \left(\mathbf{M}(y_1, y_2) \widehat{\mathbf{M}}(y_2, y_3) \mathbf{M}(y_3, y_4) \cdots + \text{perm.} \right), \quad (5.21)$$

Bound state equation. The fermions satisfy the same anti-commutation relation of [27], and therefore their same considerations apply here³⁰: in particular the bilinear quantum operators \mathbf{M} satisfy a W_{∞}^{2J+1} algebra (cf. §3.2, Appendix C,D of [27] and references therein). It has been shown in [27, 52, 53] that, in the case of 't Hooft model at large N or large J , the fluctuation along the co-adjoint orbit around a classical value of \mathbf{M} can be identified with the meson wavefunction. The equation of motion for those fluctuations give the t' Hooft equation determining the spectrum of the mesons. At large spin, this is a very efficient way to obtain the bound state equation, as this method does not rely on planar perturbation theory. We illustrate this in Appendix C where we compute the equation of motions for the

³⁰Adding a potential for the gauge part of the action, does not modify the quantization of the fermions.

fluctuation \mathbf{W} around the classical solution:

$$\begin{aligned}
& -i\partial_+ W(p, q; \theta, \phi) - \frac{m^2}{4} \left(\frac{1}{p} + \frac{1}{q} \right) W(p, q; \theta, \phi) \\
& - \frac{\lambda(-i)^n n v_n}{4\pi} \int_0^{p-q} dk I_n(k, q) (W(k, q-k, \theta, \phi) \cos(\theta - \phi) - W(p, q; \theta, \phi)) = 0.
\end{aligned} \tag{5.22}$$

In order to get an equation for a gauge-invariant quantity, we take the trace over the representation indexes:

$$\phi(x) \equiv \phi(x, 1-x) := \int d \cos \theta W(x, 1-x; \theta, \theta), \quad x := \frac{p}{r}, \quad q := p-r. \tag{5.23}$$

Recall the the functions $\phi(x)$ retain a residual dependence on the x_+ component. Going into its Fourier space $\phi(x; p_+) \equiv \phi(x)$, and restoring the $-$ label on the momenta, (C.11) takes the form:

$$4r_+ r_- \phi(x) = m^2 \left(\frac{1}{x} + \frac{1}{1-x} \right) \phi(x) - \frac{\lambda}{\pi} \int_0^1 dy K(y, x; 1-x, 1-x) (\phi(y) - \phi(x)) \tag{5.24}$$

that, is exactly equivalent to the generalized Bethe-Salpeter equation (2.9) [5], with K and Γ computed using the potential $V(B) = B^{\bar{n}}$.

We can also obtain arbitrary potentials in the large representation limit (see also footnote 28), if we scale the coefficients v_n in the potential with appropriate powers of J . Specifically, once a double scaling limit of the YM coupling g^2 and J is fixed as in (5.16): $4(2\pi)^{\bar{n}-1} g^2 J^{\bar{n}} = \lambda$, imposing that of each of the other coefficients scales as $v_m = \mathcal{O}(J^{\bar{n}-m})$ at large J guarantees that all the combinations

$$4(2\pi)^{m-1} g^2 v_m J^m = \lambda_m \tag{5.25}$$

are finite and fixed in the limit $J \rightarrow \infty$. Henceforth, at large spin J , an arbitrary potential (that now depends also explicitly on J), $V(B, J)$ leads to a mass spectrum μ^2 of mesons determined by the same generalized 't Hooft equation (2.14):

$$2\pi\mu^2\phi(x) = \left(m^2 - \sum_{n=1}^{\infty} \lambda_{2n} \alpha_{2n} \right) \left(\frac{1}{x} + \frac{1}{1-x} \right) \phi(x) - \int_0^1 dy \frac{\sum_{n=1}^{\infty} \lambda_{2n} p_{2n}(\Lambda)}{(x-y)^2} \phi(y), \tag{5.26}$$

where the summation is only over the even coefficients as a consequence of fixing the specific case of the $SU(2)$ algebra.

This implies that all the results we have derived for the meson spectra at large N are equally valid in this large representation limit. This generalizes the result of [27], obtained for the specific case of standard QCD_2 , corresponding to setting $v_n = \frac{1}{8\pi}\delta_{n,2}$. In particular, also in this limit, the theory exhibits the integrable structure.

6 Conclusion

In this paper, we studied the meson spectrum of the generalized QCD_2 at large N_c . We demonstrated that the integral equation determining the meson spectrum can be recast into the TQ-Baxter equation. The end result (3.17) turned out to be remarkably simple and universal; the dependence on the potential of the generalized Yang-Mills $V(B)$ is contained entirely in the form of the transfer matrix $T(\nu)$ that can be computed explicitly for any given potential. Based on the reformulation, we analyzed the spectrum as a function of the coefficients of $V(B)$ and uncovered a multi-sheeted structure with infinitely many branch points, each of which signals the emergence of a massless meson. These results extend previous findings for the 't Hooft model [1, 2] to much a broader class of theories. There are numerous future directions worth exploring, many of which have already been mentioned in our previous paper [1]. Below we list a few more which were not mentioned there:

- We showed that the spectrum of the generalized QCD_2 contains infinitely many branch points in the complex plane of quark masses. As is the case with the 't Hooft model, it is likely that these branch points correspond to physical critical points of the theory at which the infrared phase is described by a conformal field theory. It would be interesting to see if these branch points persist at finite N_c and, if so, to understand what their precise CFT description is by extending the analysis of [3, 4].
- As already mentioned multiple times, it would be interesting to study theories with quarks in the adjoint representation of the gauge group and analyze the spectrum of glueballs and confining fluxtubes. Developing analytic approaches to these theories is a challenging yet important direction. Another direction worth pursuing is to perform the numerical analysis for the generalized Yang-Mills coupled to adjoint quarks, extending the existing results in the literature for the adjoint QCD_2 [9, 10, 13, 14, 17, 19, 20].
- It would be interesting to find examples of four-dimensional gauge theories for which the meson spectrum can be computed exactly. One promising target is $\mathcal{N} = 4$ supersymmetric Yang-Mills (SYM) theory on the Coulomb branch. Although $\mathcal{N} = 4$ SYM

is conformal, going to the Coulomb branch allows us introduce massive excitations i.e. W-bosons. When the symmetry breaking pattern is $U(N) \rightarrow U(N-1) \times U(1)$, these W-bosons transform under the fundamental representation of the residual $U(N-1)$ gauge group and form stable mesonic bound states in the large N limit. As shown in [54] based on a judicious use of the Regge theory, a part of the spectrum of these bound states can be determined from the cusp anomalous dimension, which in turn can be computed exactly using the planar integrability of $\mathcal{N} = 4$ SYM [26]. More recently, direct evidence for the integrability on the Coulomb branch was obtained in [55]. It would be interesting to explore this direction further and determine the full spectrum of these mesonic bound states more directly from integrability. The first step may be to derive the TQ-Baxter equation governing the spectrum of these bound states at weak 't Hooft coupling. Also interesting would be to find a similar setup in the fishnet limit of $\mathcal{N} = 4$ SYM [56–60].

- There is another potential point of contact between the analysis of this paper and four-dimensional gauge theories. As discussed in Section 2.2 of [61], the dynamics in the large spin sector of four-dimensional Yang-Mills theory with (massless) dynamical quarks can be described by an effective two-dimensional QCD with higher derivative corrections, when the gauge coupling is sufficiently small (but finite). Thus, it is interesting to extend our analysis and make predictions on the large-spin dynamics of four-dimensional gauge theories.

Acknowledgement

We thank Bruno Balthazar, Aleksey Cherman, Gabriel Cuomo, Diego Delmastro, Sergei Dubovsky, John Donahue, Matijn François, Jaume Gomis, Zohar Komargodski, Sergei Lukyanov, Marcos Mariño, Giuseppe Mussardo, Alessio Miscioscia, Alba Grassi, Yifan Wang, Alexander Zamolodchikov for useful discussions.

A Recurrence relation for integrals

In this Appendix we provide a more in-depth proof for the formula (3.6). The starting point is the observation that the principal part integrals:

$$J_{n+2} = \int_0^1 \frac{\left(\frac{y}{1-y}\right)^{\frac{i\nu}{2}} \left(\log\left(\frac{1-y}{y}\right)\right)^n}{(x-y)^2}, \quad (\text{A.1})$$

can be computed as Fourier transforms of distributions, once represented through rapidity variables $\theta = \frac{1}{2} \log\left(\frac{y}{1-y}\right)$:

$$\begin{aligned}
J_{n+2} &= \int_{-\infty}^{+\infty} d\theta e^{i\theta\nu} \left(-2 \frac{\cosh^2(\theta)}{\sinh^2(\theta - \varphi)} \right) (2\theta)^n = \left(\frac{2}{i}\right)^n \frac{d^n}{d\nu^n} J_2 \\
&= \left(\frac{2}{i}\right)^n \frac{d^n}{d\nu^n} \left[\left(\frac{\left(\frac{x}{1-x}\right)^{\frac{i\nu}{2}}}{\frac{2}{\pi}x(1-x)} \right) \mathcal{I}_2 \right] \\
&= \left(\frac{2}{i}\right)^n \left(\frac{\left(\frac{x}{1-x}\right)^{\frac{i\nu}{2}}}{\frac{2}{\pi}x(1-x)} \right) \frac{d^n}{d\nu^n} \mathcal{I}_2 + \sum_{p=0}^{n-1} (-1)^{n-p+1} \binom{n}{p} \log^{n-p} \left(\frac{x}{1-x} \right) J_p
\end{aligned} \tag{A.2}$$

where \mathcal{I}_2 is the one defined in (3.3). This can be equivalently expressed as:

$$\left(\frac{2}{i}\right)^n \frac{d^n}{d\nu^n} \mathcal{I}_2 = \left(\frac{\left(\frac{x}{1-x}\right)^{\frac{i\nu}{2}}}{\frac{2}{\pi}x(1-x)} \right)^{-1} \sum_{p=0}^n (-1)^{n-p} \binom{n}{p} \log^{n-p} \left(\frac{x}{1-x} \right) J_{p+2}, \tag{A.3}$$

but, using that:

$$\int_0^1 \frac{\left(\frac{1-y}{y}\right)^{\frac{i\nu}{2}} \left(\log\left(\frac{1-y}{y}\right) - \log\left(\frac{1-x}{x}\right) \right)^n}{(x-y)^2} = \sum_{p=0}^n (-1)^{n-p} \binom{n}{p} \log^{n-p} \left(\frac{1-x}{x} \right) J_{p+2}, \tag{A.4}$$

from (A.3) we obtain (3.6):

$$\left(\frac{2}{i}\right)^n \frac{d^n}{d\nu^n} \mathcal{I}_2 = \mathcal{I}_{n+2}. \tag{A.5}$$

B Extracting the spectral determinants for cubic potential

Having constructed the inhomogeneous solutions $Q_{\pm}(\nu|\lambda)$ via the formula (3.49), we can extract from them the spectral determinants $D_{\pm}(\lambda)$ applying the integral formulas (3.28). This is done following the same strategy thoroughly explained in Appendix B of [1], and we will therefore not repeat all the details here. The crucial point is using the analogous of eq. (B.1) of [1]:

$$\frac{1}{f(\nu)} = \frac{2 \tanh\left(\frac{\pi\nu}{2}\right) (\alpha(-\cosh(\pi\nu)) + \alpha - 24\pi^3\nu v_3 + 24\pi^2 v_3 \sinh(\pi\nu))}{\underbrace{\nu(2\alpha \cosh(\pi\nu) - 2\alpha + 48\pi^3\nu v_3 + \pi \sinh(\pi\nu)(\nu - 48\pi v_3))}_{:=1/g(\nu)}} + \frac{1}{\nu \coth\left(\frac{\pi\nu}{2}\right)} \tag{B.1}$$

to write:

$$\begin{aligned}
\partial_\lambda \log(D_+(\lambda)D_-(\lambda)) &= 2 + \underbrace{\int_{-\infty}^{+\infty} d\nu \left[\frac{Q_+(\nu|\lambda) \partial_\nu Q_-(\nu|\lambda) - Q_-(\nu|\lambda) \partial_\nu Q_+(\nu|\lambda)}{\frac{\pi\nu}{2} \coth\left(\frac{\pi\nu}{2}\right)} + \frac{\tanh\left(\frac{\pi\nu}{2}\right)}{\nu} \right]}_{:=\mathcal{I}_\times^{(1)}} + \\
&\quad + \underbrace{\int_{-\infty}^{+\infty} d\nu \frac{(Q_+(\nu|\lambda) \partial_\nu Q_-(\nu|\lambda) - Q_-(\nu|\lambda) \partial_\nu Q_+(\nu|\lambda))}{g(\nu)}}_{:=\mathcal{I}_\times^{(2)}}, \\
\partial_\lambda \log\left(\frac{D_+(\lambda)}{D_-(\lambda)}\right) &= \underbrace{\int_{-\infty}^{+\infty} d\nu \frac{Q_+(\nu|\lambda)Q_-(\nu|\lambda)}{\sinh(\pi\nu)\frac{\pi\nu}{2} \coth\left(\frac{\pi\nu}{2}\right)}}_{:=\mathcal{I}_\prime^{(1)}} + \underbrace{\int_{-\infty}^{+\infty} d\nu \frac{Q_+(\nu|\lambda)Q_-(\nu|\lambda)}{\sinh(\pi\nu)g(\nu)}}_{:=\mathcal{I}_\prime^{(2)}}.
\end{aligned} \tag{B.2}$$

Now, following the discussion in [1], we reduce the evaluation of $\mathcal{I}_\times^{(1)}, \mathcal{I}_\prime^{(1)}$ to its residue at $\nu = i$ making use of the recursion relation:

$$\sigma(\nu \sim -i) = \left(\frac{-48\pi v_3 + 2\alpha z/\pi}{\nu} + 1 + 48\pi^2 v_3 y \right) \sigma(\nu \sim i) \tag{B.3}$$

. Moreover this simplification hold for the the entire term $\partial_\lambda \log\left(\frac{D_+(\lambda)}{D_-(\lambda)}\right)$, if one is only interested ultimately in deriving asymptotic expansions for the energy levels λ_n at lowest order in λ . Instead, as in [1], the term $\mathcal{I}_\times^{(2)}$ requires more care (the same applies for $\mathcal{I}_\prime^{(1)}$ at next to leading order in λ of the λ_n). This is conveniently split into two distinct contributions:

$$\begin{aligned}
\mathcal{I}_\times^{(2)} &= \frac{-i}{\pi\lambda} \mathcal{I}(\alpha, v_3) + \int_{-\infty}^{+\infty} d\nu \tilde{\mathcal{I}}(\nu) \\
\mathcal{I}(\alpha, v_3) &= \int_{-\infty}^{+\infty} d\nu \left[\frac{\alpha^2 (\sinh(\pi\nu) - \pi\nu)}{\frac{\pi\nu}{2} \cosh^2\left(\frac{\pi\nu}{2}\right) f(\nu)} + \frac{24\pi^2 \alpha v_3 (\pi^2 \nu^2 - 2 \cosh(\pi\nu) + 2\pi\nu \coth\left(\frac{\pi\nu}{2}\right) - 2)}{\frac{\pi\nu}{2} \cosh^2\left(\frac{\pi\nu}{2}\right) f(\nu)} + \right. \\
&\quad \left. + \frac{72\pi^4 v_3^2 (-2\pi^2 \nu^2 \cosh(\pi\nu) + \cosh(2\pi\nu) - 1) \operatorname{csch}^4\left(\frac{\pi\nu}{2}\right)}{\frac{\pi\nu}{2} \cosh^2\left(\frac{\pi\nu}{2}\right) f(\nu)} \right].
\end{aligned} \tag{B.4}$$

The term $\int_{-\infty}^{+\infty} d\nu \tilde{\mathcal{I}}(\nu)$ can be again reduced to its residue at $\nu = i$. The integral expression $\mathcal{I}(\alpha, v_3)$ is the analogous in this problem of the integral in equation 5.41 of [1]. In particular, the integral develops a singularity, due to collision of roots in the denominator, exactly at the values of (α, v_3) where the theory has a critical point.

C Derivation of the bound state equation at large representation

In this appendix we follow closely [27, 52, 53]; we do not repeat many of the technical points that are carefully explained in Appendix D of [27] to which we refer the reader for more details.

The starting point is the action for a field for \mathbf{M} , that can be borrowed from the standard theory of quantization of the coadjoint orbits by adding a *kinetic term* $S_{\text{kin}}(\mathbf{M})$ ³¹ to the Hamiltonian \mathcal{H}_{eff} :

$$S[\mathbf{M}] = S_{\text{kin}}(\mathbf{M}) - \frac{im^2}{8} \int dx^+ dy_1^- \text{tr} \mathbf{S} \mathbf{M} + \frac{\lambda v_n}{8} \int dx^+ \int \prod_{i=1}^{\bar{n}} (dy_i \text{sgn}(y_k - y_{k-1})) \text{tr} \left(\mathbf{M}(y_1, y_2) \widehat{\mathbf{M}}(y_2, y_3) \mathbf{M}(y_3, y_4) \cdots + \text{perm.} \right), \quad (\text{C.1})$$

where $S(x, y, \theta, \phi) = \text{sgn}(x_- - y_-) \frac{1}{\sin(\theta)} \delta(\theta - \phi)$. This generalizes the action used in [27] only in the last term in (C.1).

We chose a coadjoint orbit with vanishing $U(1)$ charge, of whose a representative is $M_0(p, q, \theta, \phi) = 2\pi\Theta(p_-)\delta(p_- - q_-)\frac{1}{\sin\theta}\delta(\theta - \phi)$, where

$$M(p, q, \theta, \phi) = \int_{-\infty}^{+\infty} dx_- dy_- e^{ipx_-} e^{iqy_-} M(x, y, \theta, \phi). \quad (\text{C.2})$$

Fluctuation \mathbf{W} around the coadjoint orbit are be parametrized as: (eq. 3.26 of [27]):

$$\mathbf{M} = \mathbf{M}_0 + \frac{1}{\sqrt{J}}[\mathbf{W}, \mathbf{M}_0] - \frac{1}{2J}[\mathbf{W}, [\mathbf{W}, \mathbf{M}_0]] + \mathcal{O}(J^{-3/2}). \quad (\text{C.3})$$

The action (C.1) is order J , and therefore we must take the $\mathcal{O}(1/J)$ term under the variation (C.3) to get the finite leading order action for the fluctuations. The first two terms of (C.1), are directly borrowed the result from Appendix D of [27]:

$$\delta S_{\text{kin}} = i \frac{1}{J} \int dx^+ \int_{p, q > 0} dp dq \text{tr} \mathbf{W}^{+-}(p, q) \partial_+ \mathbf{W}^{-+}(q, p), \quad (\text{C.4})$$

$$\delta S_{\text{mass}} = -\frac{m^2}{4J} \int_{k, q > 0} dp dq \left(\frac{1}{k} + \frac{1}{p} \right) \text{tr} \mathbf{W}^{+-}(k, p) \mathbf{W}^{-+}(p, k),$$

³¹The details of this are not important for our discussion, we refer the interested reader to [27].

where, for clarity of comparison, we use the notation³² : $W^{\pm\pm}(p, q) = W(\pm p, \pm q)$. The interaction term of H_{eff} receives three contributions (up to permutations of θ_i s) at order $\mathcal{O}(1/J)$. Schematically:

$$\text{I} : \mathbf{M}_0 \frac{1}{2J} \widehat{\mathbf{W}} \widehat{\mathbf{W}} \left(\underbrace{\mathbf{M}_0 \widehat{\mathbf{M}}_0 \cdots}_{(\bar{n}-2) \text{ terms}} \right), \quad \text{II} : \widehat{\mathbf{M}}_0 \frac{1}{2J} \mathbf{W} \mathbf{W} \left(\underbrace{\mathbf{M}_0 \widehat{\mathbf{M}}_0 \cdots}_{(\bar{n}-2) \text{ terms}} \right) \simeq \text{I}, \quad (\text{C.5})$$

the II and the I are equal contributions up to permutations of the θ_i s; a third, inequivalent contribution is:

$$\text{III} : \frac{1}{\sqrt{J}} \mathbf{W} \frac{1}{\sqrt{J}} \widehat{\mathbf{W}} \left(\underbrace{\mathbf{M}_0 \widehat{\mathbf{M}}_0 \cdots}_{(\bar{n}-2) \text{ terms}} \right) \quad (\text{C.6})$$

All this contribution to the variation of (C.1) can be computed as a generalization of Appendix D of [27]. The contributions I, II are evaluated to³³:

$$-i \int_{p, q > 0} dp dq \int_0^{p-q} dk \int d^{\bar{n}-1} \mathbf{k} \frac{-i \operatorname{sgn}(k_1)}{2\pi(k-k_1)} \frac{-i \operatorname{sgn}(k_2)}{2\pi(k_1-k_2)} \cdots \frac{-i \operatorname{sgn}(k_{\bar{n}-1})}{2\pi(k_{\bar{n}-1}-k_1)} \operatorname{tr} \mathbf{W}^{-+}(p, q) \mathbf{W}^{+-}(q, p) \quad (\text{C.7})$$

all the $\bar{n} + 1$ integrations over $dp dq$ but the two remaining above over $dp dq$ are removed by the $\bar{n} - 3 \delta(\theta_i - \theta_{i+1})$ inside each of the \mathbf{M}_0 , this also cancel all the $\cos(\theta_i - \theta_{i+1})$ trigonometric factors. the integral over the k_i is exactly the *master integral* (2.7) introduced in section 2:

$$\int_{p, q > 0} dp dq \int_0^{p-q} dk I_n(k, q) \operatorname{tr} \mathbf{W}^{-+}(p, p) \mathbf{W}^{+-}(q, p). \quad (\text{C.8})$$

Analogously, the third contribution is evaluated as:

$$\int_{p, q > 0} dp dq \int_0^{p-q} dk I_n(k, q) \operatorname{tr} \left[\mathbf{W}^{+-}(q-k, q) \widehat{\mathbf{W}}^{-+}(p, q) + \mathbf{W}^{-+}(k, q-k) \widehat{\mathbf{W}}^{+-}(q, p) \right]. \quad (\text{C.9})$$

Once we sum over all the permutations of I, II and III, taking into account the combinatorial

³²The same of [27] up to $+(-) \leftrightarrow p(n)$.

³³The form of the integral over the k_i can be readily obtained (for instance) from a direct generalizing eq. D.17 of [27]: in each of the two integral the sign can be recognized to be $\operatorname{sgn}(k)$, and $1/k^2$ (Fourier transform of $|x|$) is replaced for each term of our problem with $\frac{i}{(k_{i-1}-k_i)} \frac{i}{(k_i-k_{i+1})}$ (Fourier transform of $\operatorname{sgn}(x)$), with $k_0 = k, k_{\bar{n}} = q$

factor $\frac{1}{n!}$, we obtain the following expression for the interaction term of \mathcal{H}_{eff} at order $\mathcal{O}(1/J)$:

$$\begin{aligned} \frac{-\lambda}{4J} v_n (-i)^{\bar{n}} \bar{n} \int_{q,p>0} dp dq \int_0^{p-q} dk I_n(k, q) \text{tr} \left[\left(\widehat{\mathbf{W}}^{+-}(q-k, k) - \mathbf{W}^{+-}(q, p) \right) \mathbf{W}^{-+}(p, q) + \right. \\ \left. \left(\widehat{\mathbf{W}}^{-+}(k, q-k) - \mathbf{W}^{-+}(p, q) \right) \mathbf{W}^{+-}(q, p) \right] \end{aligned} \quad (\text{C.10})$$

Varying the action w.r.t. $W^{+-}(p, q; \theta, \phi)$ we obtain the following equation of motions:

$$\begin{aligned} -i\partial_+ W^{-+}(p, q; \theta, \phi) - \frac{m^2}{4} \left(\frac{1}{p} + \frac{1}{q} \right) W^{-+}(p, q; \theta, \phi) \\ - \frac{\lambda(-i)^n n v_n}{4\pi} \int_0^{p-q} dk I_n(k, q) \left(W^{-+}(k, q-k, \theta, \phi) \cos(\theta - \phi) - W^{-+}(p, q; \theta, \phi) \right) = 0. \end{aligned} \quad (\text{C.11})$$

In the main text we report (C.11) by replacing $W^{-+} \mapsto W$ in (5.22) to simplify the notation.

D Numerical methods for analytical continuation

In this appendix we illustrate the numerical method we employ to solve numerically the 't Hooft equation in ν -space (3.2) in the complex space of the couplings. As discussed in the main text (cfr. Section 4), the integral eigenvalue problem exhibits a multi-sheeted structure in the complex plane, arising from collision-of-roots singularities inside the kernel of (3.2) due to the poles of $\Psi(\nu)$. Compared to the same integral equation in x -space, the problem in Fourier space, is particularly convenient to solve numerically even in the complex space of the couplings. To solve this eigenproblem numerically, we employ an orthonormal basis of Hermite polynomials:

$$\langle n|x \rangle = H_n(x) = \frac{(-1)^n e^{\frac{x^2}{2}}}{\sqrt{2^n \pi^{1/2} n!}} \frac{d^n}{dx^n} e^{-x^2}, \quad \langle m|n \rangle = \delta_{nm}, \quad (\text{D.1})$$

that share many properties with the eingefunction $\Psi_n(x)$ having exactly n zeros and definite parity in $x \in (-\infty, +\infty)$.

In this basis the generalized 't Hooft equation (3.2) reduces to the eingeproblem for the following ‘Hamiltonian’:

$$\langle m|\mathcal{H}|n \rangle = \mathcal{H}_{mn} = \int_{-\infty}^{+\infty} dx \int_{-\infty}^{+\infty} dy \frac{H_m(x) H_n(y)}{f(x)} \frac{\pi(x-y)}{2 \sinh\left(\frac{\pi}{2}(x-y)\right)}, \quad (\text{D.2})$$




Figure 22: The integration contour in the x -plane of (D.2) is deformed under analytical continuation from $\alpha > -1$ to $\alpha < -1$ at fixed $-iv_3 > 0$.

that can be straightforwardly solved upon truncation of the basis. The eigenvalues e_k of \mathcal{H}_{nm} are related to the masses of the mesons λ_k by inversion: $\lambda_k = e_k^{-1}$.

This numerical method is particularly akin to analytical continuation. Indeed, whenever upon analytical continuation of the couplings α, \mathbf{v} , one (or more) of the zeros of $f(\nu)$ crosses the integration contour, one can just modify the Hamiltonian \mathcal{H}_{mn} simply by including the corresponding residue terms.

To exemplify, consider the case of $V(B) = \frac{1}{8\pi}B^2 + i|v_3|B^3$. Fixed any $v_3 > 0$, for $\alpha > -1$ no zeros of $f(\nu)$ lie on the integration contour, while upon analytically continue in α to values $\alpha < -1$, exactly one of the zero crosses the real axis, and therefore the contour has to be deformed to avoid this singular point as in Figure 22. Then, the dx integral in (D.2) will pick a further contribution due a residue term³⁴:

$$\mathcal{H}_{mn} \rightarrow \mathcal{H}_{mn} - 2\pi i H_m(x_0) \text{Res} [f^{-1}(x, \alpha, v_3), x = x_0] \int_{-\infty}^{+\infty} dy H_n(y) \frac{\pi(x_0 - y)}{2 \sinh\left(\frac{\pi}{2}(x_0 - y)\right)}, \quad (\text{D.3})$$

where $x_0 = x_0(\alpha, v_3)$ is the position of the zero of $f(x, \alpha, v_3)$ that has crossed the integration contour.

Analogous manipulations make straightforward the numerical analysis of (3.2) among the various sheets in the couplings-space. In practice, already truncating the basis at $n_{max} \sim 30$, reproduces the first 4 eigenvalues of [2] up to an accuracy of 10^{-5} , that used a basis of 400 elements to diagonalize the x -space for the simplest case of $\mathbf{v} = \mathbf{0}$ and $\alpha = 0$. This is yet another illustration of the efficiency of solving the ν -space problem, rather than the original one in x -space, also at the level of numerical methods.

³⁴Note that the sign of the residue comes from the contour orientation in Figure 22

References

- [1] F. Ambrosino and S. Komatsu, *2d QCD and Integrability, Part I: 't Hooft model*, [arXiv:2312.15598](#).
- [2] V. A. Fateev, S. L. Lukyanov, and A. B. Zamolodchikov, *On mass spectrum in 't Hooft's 2D model of mesons*, *J. Phys. A* **42** (2009) 304012, [[arXiv:0905.2280](#)].
- [3] D. Delmastro, J. Gomis, and M. Yu, *Infrared phases of 2d QCD*, *JHEP* **02** (2023) 157, [[arXiv:2108.02202](#)].
- [4] D. Delmastro and J. Gomis, *RG flows in 2d QCD*, *JHEP* **09** (2023) 158, [[arXiv:2211.09036](#)].
- [5] M. R. Douglas, K. Li, and M. Staudacher, *Generalized two-dimensional QCD*, *Nucl. Phys. B* **420** (1994) 118–140, [[hep-th/9401062](#)].
- [6] S. Dalley and I. R. Klebanov, *String spectrum of (1+1)-dimensional large N QCD with adjoint matter*, *Phys. Rev. D* **47** (1993) 2517–2527, [[hep-th/9209049](#)].
- [7] D. Kutasov, *Two-dimensional QCD coupled to adjoint matter and string theory*, *Nucl. Phys. B* **414** (1994) 33–52, [[hep-th/9306013](#)].
- [8] J. Boorstein and D. Kutasov, *Symmetries and mass splittings in QCD in two-dimensions coupled to adjoint fermions*, *Nucl. Phys. B* **421** (1994) 263–277, [[hep-th/9401044](#)].
- [9] G. Bhanot, K. Demeterfi, and I. R. Klebanov, *(1+1)-dimensional large N QCD coupled to adjoint fermions*, *Phys. Rev. D* **48** (1993) 4980–4990, [[hep-th/9307111](#)].
- [10] K. Demeterfi, I. R. Klebanov, and G. Bhanot, *Glueball spectrum in a (1+1)-dimensional model for QCD*, *Nucl. Phys. B* **418** (1994) 15–29, [[hep-th/9311015](#)].
- [11] A. V. Smilga, *Instantons and fermion condensate in adjoint QCD in two-dimensions*, *Phys. Rev. D* **49** (1994) 6836–6848, [[hep-th/9402066](#)].
- [12] F. Lenz, M. A. Shifman, and M. Thies, *Quantum mechanics of the vacuum state in two-dimensional QCD with adjoint fermions*, *Phys. Rev. D* **51** (1995) 7060–7082, [[hep-th/9412113](#)].

- [13] E. Katz, G. Marques Tavares, and Y. Xu, *Solving 2D QCD with an adjoint fermion analytically*, *JHEP* **05** (2014) 143, [[arXiv:1308.4980](#)].
- [14] E. Katz, G. Marques Tavares, and Y. Xu, *A solution of 2D QCD at Finite N using a conformal basis*, [arXiv:1405.6727](#).
- [15] A. Cherman, T. Jacobson, Y. Tanizaki, and M. Ünsal, *Anomalies, a mod 2 index, and dynamics of 2d adjoint QCD*, *SciPost Phys.* **8** (2020), no. 5 072, [[arXiv:1908.09858](#)].
- [16] Z. Komargodski, K. Ohmori, K. Roumpedakis, and S. Seifnashri, *Symmetries and strings of adjoint QCD₂*, *JHEP* **03** (2021) 103, [[arXiv:2008.07567](#)].
- [17] R. Dempsey, I. R. Klebanov, and S. S. Pufu, *Exact symmetries and threshold states in two-dimensional models for QCD*, *JHEP* **10** (2021) 096, [[arXiv:2101.05432](#)].
- [18] F. K. Popov, *Supersymmetry in QCD₂ coupled to fermions*, *Phys. Rev. D* **105** (2022), no. 7 074005, [[arXiv:2202.04017](#)].
- [19] R. Dempsey, I. R. Klebanov, L. L. Lin, and S. S. Pufu, *Adjoint Majorana QCD₂ at finite N*, *JHEP* **04** (2023) 107, [[arXiv:2210.10895](#)].
- [20] R. Dempsey, I. R. Klebanov, S. S. Pufu, and B. T. Søgaaard, *Lattice Hamiltonian for Adjoint QCD₂*, [arXiv:2311.09334](#).
- [21] S. Dubovsky, *A Simple Worldsheet Black Hole*, *JHEP* **07** (2018) 011, [[arXiv:1803.00577](#)].
- [22] J. C. Donahue and S. Dubovsky, *Confining Strings, Infinite Statistics and Integrability*, *Phys. Rev. D* **101** (2020), no. 8 081901, [[arXiv:1907.07799](#)].
- [23] J. C. Donahue and S. Dubovsky, *Classical Integrability of the Zigzag Model*, *Phys. Rev. D* **102** (2020), no. 2 026005, [[arXiv:1912.08885](#)].
- [24] J. C. Donahue and S. Dubovsky, *Quantization of the zigzag model*, *JHEP* **08** (2022) 047, [[arXiv:2202.11746](#)].
- [25] M. Asrat, *(1+1)D QCD with heavy adjoint quarks*, *Phys. Rev. D* **107** (2023), no. 10 106022, [[arXiv:2212.02162](#)].
- [26] N. Beisert et al., *Review of AdS/CFT Integrability: An Overview*, *Lett. Math. Phys.* **99** (2012) 3–32, [[arXiv:1012.3982](#)].

- [27] A. Kaushal, N. S. Prabhakar, and S. R. Wadia, *Meson spectrum of $SU(2)$ QCD_{1+1} with Quarks in Large Representations*, [arXiv:2307.15015](#).
- [28] A. Bourget, D. Rodriguez-Gomez, and J. G. Russo, *A limit for large R -charge correlators in $\mathcal{N} = 2$ theories*, *JHEP* **05** (2018) 074, [[arXiv:1803.00580](#)].
- [29] G. Arias-Tamargo, D. Rodriguez-Gomez, and J. G. Russo, *The large charge limit of scalar field theories and the Wilson-Fisher fixed point at $\epsilon = 0$* , *JHEP* **10** (2019) 201, [[arXiv:1908.11347](#)].
- [30] G. Arias-Tamargo, D. Rodriguez-Gomez, and J. G. Russo, *Correlation functions in scalar field theory at large charge*, *JHEP* **01** (2020) 171, [[arXiv:1912.01623](#)].
- [31] G. Badel, G. Cuomo, A. Monin, and R. Rattazzi, *The Epsilon Expansion Meets Semiclassics*, *JHEP* **11** (2019) 110, [[arXiv:1909.01269](#)].
- [32] M. Watanabe, *Accessing large global charge via the ϵ -expansion*, *JHEP* **04** (2021) 264, [[arXiv:1909.01337](#)].
- [33] S. Giombi and J. Hyman, *On the large charge sector in the critical $O(N)$ model at large N* , *JHEP* **09** (2021) 184, [[arXiv:2011.11622](#)].
- [34] J. a. Caetano, S. Komatsu, and Y. Wang, *Large Charge 't Hooft Limit of $\mathcal{N} = 4$ Super-Yang-Mills*, [arXiv:2306.00929](#).
- [35] G. Cuomo, Z. Komargodski, M. Mezei, and A. Raviv-Moshe, *Spin impurities, Wilson lines and semiclassics*, *JHEP* **06** (2022) 112, [[arXiv:2202.00040](#)].
- [36] O. Aharony, G. Cuomo, Z. Komargodski, M. Mezei, and A. Raviv-Moshe, *Phases of Wilson Lines in Conformal Field Theories*, *Phys. Rev. Lett.* **130** (2023), no. 15 151601, [[arXiv:2211.11775](#)].
- [37] O. Aharony, G. Cuomo, Z. Komargodski, M. Mezei, and A. Raviv-Moshe, *Phases of Wilson Lines: Conformality and Screening*, [arXiv:2310.00045](#).
- [38] D. Rodriguez-Gomez and J. G. Russo, *Wilson loops in large symmetric representations through a double-scaling limit*, *JHEP* **08** (2022) 253, [[arXiv:2206.09935](#)].
- [39] M. Beccaria, S. Giombi, and A. A. Tseytlin, *Wilson loop in general representation and RG flow in 1D defect QFT*, *J. Phys. A* **55** (2022), no. 25 255401, [[arXiv:2202.00028](#)].

- [40] D. Rodriguez-Gomez and J. G. Russo, *Defects in scalar field theories, RG flows and dimensional disentangling*, *JHEP* **11** (2022) 167, [[arXiv:2209.00663](#)].
- [41] I. Carreño Bolla, D. Rodriguez-Gomez, and J. G. Russo, *RG flows and stability in defect field theories*, *JHEP* **05** (2023) 105, [[arXiv:2303.01935](#)].
- [42] G. 't Hooft, *A two-dimensional model for mesons*, *Nuclear Physics B* **75** (1974), no. 3 461–470.
- [43] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. Dover, New York, ninth dover printing, tenth gpo printing ed., 1964.
- [44] C. M. Bender and S. Boettcher, *Real spectra in non-hermitian hamiltonians having pt symmetry*, *Physical Review Letters* **80** (jun, 1998) 5243–5246.
- [45] C. M. Bender, *Introduction to pt -symmetric quantum theory*, *Contemporary Physics* **46** (jul, 2005) 277–292.
- [46] C. M. Bender, D. C. Brody, and H. F. Jones, *Extension of PT symmetric quantum mechanics to quantum field theory with cubic interaction*, *Phys. Rev. D* **70** (2004) 025001, [[hep-th/0402183](#)]. [Erratum: *Phys.Rev.D* 71, 049901 (2005)].
- [47] M. Lencsés, A. Miscioscia, G. Mussardo, and G. Takács, *Multicriticality in Yang-Lee edge singularity*, *JHEP* **02** (2023) 046, [[arXiv:2211.01123](#)].
- [48] M. Lencsés, A. Miscioscia, G. Mussardo, and G. Takács, *\mathcal{PT} breaking and RG flows between multicritical Yang-Lee fixed points*, *JHEP* **09** (2023) 052, [[arXiv:2304.08522](#)].
- [49] M. Lencsés, A. Miscioscia, G. Mussardo, and G. Takács, *Ginzburg-Landau description for multicritical Yang-Lee models*, [arXiv:2404.06100](#).
- [50] R. J. Eden, P. V. Landshoff, D. I. Olive, and J. C. Polkinghorne, *The analytic S -matrix*. Cambridge Univ. Press, Cambridge, 1966.
- [51] I. Pesando, *Generalized QCD in two-dimensions via the bilocal method*, *Mod. Phys. Lett. A* **9** (1994) 2927–2936, [[hep-th/9408018](#)].
- [52] A. DHAR, G. MANDAL, and S. R. WADIA, *W_∞ coherent states and path-integral derivation of bosonization of non-relativistic fermions in one dimension*, *Modern Physics Letters A* **08** (Dec., 1993) 3557–3568.

- [53] A. Dhar, G. Mandal, and S. R. Wadia, *String field theory of two dimensional qcd: a realization of w_∞ algebra*, *Physics Letters B* **329** (June, 1994) 15–26.
- [54] S. Caron-Huot and J. M. Henn, *Solvable Relativistic Hydrogenlike System in Supersymmetric Yang-Mills Theory*, *Phys. Rev. Lett.* **113** (2014), no. 16 161601, [[arXiv:1408.0296](#)].
- [55] V. Ivanovskiy, S. Komatsu, V. Mishnyakov, N. Terziev, N. Zagraev, and K. Zarembo, *Vacuum Condensates on the Coulomb Branch*, [arXiv:2405.19043](#).
- [56] A. B. Zamolodchikov, *'FISHNET' DIAGRAMS AS A COMPLETELY INTEGRABLE SYSTEM*, *Phys. Lett. B* **97** (1980) 63–66.
- [57] O. Gürdoğan and V. Kazakov, *New Integrable 4D Quantum Field Theories from Strongly Deformed Planar $\mathcal{N} = 4$ Supersymmetric Yang-Mills Theory*, *Phys. Rev. Lett.* **117** (2016), no. 20 201602, [[arXiv:1512.06704](#)]. [Addendum: *Phys.Rev.Lett.* 117, 259903 (2016)].
- [58] J. a. Caetano, O. Gürdoğan, and V. Kazakov, *Chiral limit of $\mathcal{N} = 4$ SYM and ABJM and integrable Feynman graphs*, *JHEP* **03** (2018) 077, [[arXiv:1612.05895](#)].
- [59] F. Loebbert, J. Miczajka, D. Müller, and H. Münkler, *Massive Conformal Symmetry and Integrability for Feynman Integrals*, *Phys. Rev. Lett.* **125** (2020), no. 9 091602, [[arXiv:2005.01735](#)].
- [60] F. Loebbert and J. Miczajka, *Massive Fishnets*, *JHEP* **12** (2020) 197, [[arXiv:2008.11739](#)].
- [61] L. F. Alday and J. M. Maldacena, *Comments on operators with large spin*, *JHEP* **11** (2007) 019, [[arXiv:0708.0672](#)].