

K -STABILITY OF \mathcal{Z} -STABLE C^* -ALGEBRAS

SHANSHAN HUA

ABSTRACT. We provide a shorter new proof of the fact that \mathcal{Z} -stable C^* -algebras are K_1 -surjective using the Rørdam-Winter picture of the Jiang-Su algebra \mathcal{Z} . Consequently, we recapture the K -stability of \mathcal{Z} -stable C^* -algebras.

1. INTRODUCTION

Operator algebraic K -theory is defined using equivalence classes of unitaries and projections in matrix amplifications over a C^* -algebra. By Bott periodicity, the K -theory of a unital C^* -algebra A is given by the higher homotopy groups of $\mathcal{U}_\infty(A)$, where $\pi_{2n}(\mathcal{U}_\infty(A)) \cong K_1(A)$ and $\pi_{2n+1}(\mathcal{U}_\infty(A)) \cong K_0(A)$ for $n \in \mathbb{N}$, see [12, Proposition 11.4.1] for instance. On the other hand, non-stable K -theory studies the homology theory given by the higher homotopy groups of the unitary group $\mathcal{U}(A)$ based at 1_A . The K -theory of stable C^* -algebras can be computed without taking matrix amplifications and thus coincides with their non-stable K -theory. However, K -theory and non-stable K -theory are different in general, even in the case of commutative C^* -algebras. Take $C(S^3)$ as an example, where we have $K_1(C(S^3)) = \mathbb{Z}$, but $\pi_0(\mathcal{U}(C(S^3))) = 0$.

In many important cases, even without stability, the K -theory and non-stable K -theory agree, see [9] for noncommutative irrational tori and [21] for simple C^* -algebras with real rank zero and stable rank one. In general, exact characterisations of when stable and non-stable K -theory coincide are not known, even at the level of π_0 . Blackadar raised a question in [1] of when a C^* -algebra is K_1 -injective or K_1 -surjective, namely when the canonical homomorphism from $\pi_0(\mathcal{U}(A))$ to $K_1(A)$ is injective or surjective. A stronger notion called K -stability is defined by Thomsen in [16], which means that for any $n \in \mathbb{N}$, the canonical inclusion $\mathcal{U}(M_n \otimes A) \rightarrow \mathcal{U}_\infty(A)$ induces isomorphisms between higher homotopy groups of $\mathcal{U}(M_n \otimes A)$ and the K -theory of A . C^* -algebras are K -stable under certain structural conditions, such as having stable rank one in [9] or being simple and purely infinite in [21]. However, it remains open whether simple C^* -algebras are automatically K -stable or K_1 -isomorphic.

Recently, non-stable K -theory has shown its importance in the new abstract approach of classification for C^* -algebras initiated in [14]. In the history of the Elliott classification program, a central object is the Jiang-Su algebra \mathcal{Z} , which was originally constructed in [8] as the inductive limit of dimension drop algebras, with explicitly defined connecting maps. Tensorial absorption of \mathcal{Z} excludes the exotic C^* -algebras constructed in [19], [10], [17] and is needed to obtain a satisfying classification theorem for simple separable nuclear C^* -algebras by K -theory and traces. Though this is a different tensorial absorption property from stability, it does imply similar nice K -theoretical properties. For simple \mathcal{Z} -stable exact C^* -algebras, a dichotomy in [4] shows that they are either purely infinite or stably finite.

This work forms part of the author's DPhil thesis at the University of Oxford.

Rørdam proved in 2004 in [11] that simple stably finite \mathcal{Z} -stable C^* -algebras necessarily have stable rank one, by extensively studying structural properties of \mathcal{Z} -stable C^* -algebras. Then combining results in [21] and [9], one can conclude that \mathcal{Z} -stability implies K -stability for simple C^* -algebras.

The result that \mathcal{Z} -stable C^* -algebras are K -stable in the general setting is actually proven earlier by Jiang in his unpublished paper [7], which in particular implies K_1 -injectivity and K_1 -surjectivity for \mathcal{Z} -stable C^* -algebras. In the new abstract approach to the classification theorem in [2], K_1 -injectivity for certain non-simple \mathcal{Z} -stable C^* -algebras plays an important role in [2, Section 5.4] to obtain the uniqueness theorem for maps. Since the result in [7] remains unpublished, a short and self-contained new proof for K_1 -injectivity of \mathcal{Z} -stable C^* -algebras was provided in [2, Section 4.2]. The proof takes advantage of a modern description of \mathcal{Z} , given by Rørdam and Winter in [13], around ten years after its introduction. They realize \mathcal{Z} as a stationary inductive limit of the so-called generalized dimension drop algebras, whose fibres are UHF-algebras instead of finite matrix algebras. This should be seen as an alternative definition for \mathcal{Z} , since many fundamental properties of \mathcal{Z} can be readily obtained from this viewpoint, for instance in reproving strong self-absorption of \mathcal{Z} in [15] and Winter's localization of the Elliott conjecture in [20], which is vital in the Gong-Lin-Niu theorem [5].

In this paper, we will recapture the rest of Jiang's K -stability results. Jiang's proof relies on explicit constructions of homomorphisms to produce inverse maps between higher homotopy groups when proving K -stability, which then implies K_1 -injectivity and K_1 -surjectivity of \mathcal{Z} -stable C^* -algebras. In comparison, we first provide a short new proof for K_1 -surjectivity of \mathcal{Z} -stable C^* -algebras in Section 3, using the Rørdam-Winter picture in the spirit of the K_1 -injectivity proof of [2]. The extra flexibility is given by UHF-fibers of generalized dimension drop algebras, since M_{n^∞} -stable C^* -algebras are shown to have nice K -theoretical properties in Lemma 3.1 and [2, Lemma 4.9], which is not true in general when tensoring by finite matrix algebras. With K_1 -injectivity and K_1 -surjectivity of \mathcal{Z} -stable C^* -algebras, we are able to recapture K -stability in Theorem 4.2, without needing to construct explicit maps.

The result in the paper is becoming increasingly important. For instance, in Toms' recent work [18] on higher homotopy groups of Cuntz classes for simple approximately divisible C^* -algebras of real rank zero, K_1 -surjectivity plays an important role, and it is likely that this will continue to be the case in further work in this direction.

2. PRELIMINARIES

For a unital C^* -algebra A , we will denote the set of unitaries in A by $\mathcal{U}(A)$. For $n \geq 1$, we will use the notation 1_n for the unit of $M_n \otimes A$ and $\mathcal{U}_n(A)$ for the set of unitaries in $M_n \otimes A$. The set of unitaries in $\mathcal{U}_n(A)$ that are homotopic to 1_n is denoted by $\mathcal{U}_n^0(A)$. If A is a non-unital C^* -algebra, we denote by \tilde{A} the unitization of A , with canonically associated maps $\iota : A \rightarrow \tilde{A}$ and $q : \tilde{A} \rightarrow \mathbb{C}$.

Definition 2.1. A unital C^* -algebra A is called *K_1 -surjective* (*K_1 -injective*) if the group homomorphism $\mathcal{U}(A)/\mathcal{U}^0(A) \rightarrow K_1(A)$, $[u] \mapsto [u]_1$, is surjective (injective).

In the non-unital case, a C^* -algebra A is called *K_1 -surjective* (*K_1 -injective*) if \tilde{A} is K_1 -surjective (K_1 -injective).

We will also describe briefly the different descriptions of \mathcal{Z} . For coprime integers p, q , the *prime dimension drop algebra* is defined to be

$$(2.1) \quad \mathcal{Z}_{p,q} = \{f \in C([0, 1], M_p \otimes M_q) : f(0) \in M_p \otimes 1_q, f(1) \in 1_p \otimes M_q\}.$$

The Jiang-Su algebra \mathcal{Z} was originally constructed in [8] as an inductive limit of such $\mathcal{Z}_{p,q}$, together with delicately defined connecting maps. In fact, it is the unique inductive limit of prime dimension drop algebras that is unital, simple, monotracial and $K_*(A) \cong K_*(\mathbb{C})$ by [8, Theorem 6.2].

The Rørdam-Winter picture of \mathcal{Z} uses the *generalized prime dimension drop algebra* corresponding to infinite coprime supernatural numbers \mathfrak{p} and \mathfrak{q} , which is defined by

$$(2.2) \quad \mathcal{Z}_{\mathfrak{p},\mathfrak{q}} = \{f \in C([0, 1], M_{\mathfrak{p}} \otimes M_{\mathfrak{q}}) : f(0) \in M_{\mathfrak{p}} \otimes 1_{\mathfrak{q}}, f(1) \in 1_{\mathfrak{p}} \otimes M_{\mathfrak{q}}\}.$$

A unital endomorphism φ on a unital C*-algebra A is said to be *trace-collapsing* if $\tau \circ \varphi = \tau' \circ \varphi$ for any $\tau, \tau' \in T(A)$. Then the following theorem is proven.

Theorem 2.2. ([13, Theorem 3.4]) *Let $\mathfrak{p}, \mathfrak{q}$ be infinite coprime supernatural numbers. Then there exists a trace-collapsing unital endomorphism φ on $\mathcal{Z}_{\mathfrak{p},\mathfrak{q}}$. For any such φ , the Jiang-Su algebra \mathcal{Z} is isomorphic to the stationary inductive limit of the sequence*

$$\mathcal{Z}_{\mathfrak{p},\mathfrak{q}} \xrightarrow{\varphi} \mathcal{Z}_{\mathfrak{p},\mathfrak{q}} \xrightarrow{\varphi} \mathcal{Z}_{\mathfrak{p},\mathfrak{q}} \xrightarrow{\varphi} \cdots.$$

As noted in the introduction, this viewpoint for \mathcal{Z} is used to reprove K_1 -injectivity of unital \mathcal{Z} -stable C*-algebras in [2, Theorem 4.8]. This quickly extends to the non-unital case, which recaptures Jiang's K_1 -injectivity result in [7, Theorem 2].

Theorem 2.3. *If A is a C*-algebra, then $A \otimes \mathcal{Z}$ is K_1 -injective.*

Proof. The unital case is proven in [2, Theorem 4.8]. When A is non-unital, take $u \in \mathcal{U}((A \otimes \mathcal{Z})^\sim)$ with $[u]_1 = 0 \in K_1(A \otimes \mathcal{Z})$. Then $[u]_1 = 0 \in K_1(\tilde{A} \otimes \mathcal{Z})$ and $(q \otimes \text{id}_{\mathcal{Z}}(u)) = \lambda 1_{\mathcal{Z}}$ for some $\lambda \in \mathbb{T}$. Since $\tilde{A} \otimes \mathcal{Z}$ is K_1 -injective, there is a path of unitaries $(u_t)_{t \in [0,1]} \subseteq \mathcal{U}(\tilde{A} \otimes \mathcal{Z})$ such that $u_0 = 1_{\tilde{A} \otimes \mathcal{Z}}$ and $u_1 = u$. For $t \in [0,1]$, take $v_t = (1_{\tilde{A}} \otimes (q \otimes \text{id}_{\mathcal{Z}})(u_t))^* u_t$, then $(v_t)_{t \in [0,1]}$ is a path of unitaries in $\mathcal{U}(\tilde{A} \otimes \mathcal{Z})$ with $v_0 = 1_{\tilde{A} \otimes \mathcal{Z}}$ and $v_1 = \bar{\lambda}u$. Since $(q \otimes \text{id}_{\mathcal{Z}})(v_t) = 1_{\mathcal{Z}}$ for $t \in [0,1]$, then $(v_t)_{t \in [0,1]} \subseteq \mathcal{U}((A \otimes \mathcal{Z})^\sim)$, which implies $1_{(A \otimes \mathcal{Z})^\sim} \sim_h \bar{\lambda}u \sim_h u$ in $(A \otimes \mathcal{Z})^\sim$. \square

Since the approach in [2, Theorem 4.8] inspires the results of this paper, we will use Theorem 2.3 freely in later sections.

3. K_1 -SURJECTIVITY OF \mathcal{Z} -STABLE C*-ALGEBRAS

We start by proving K_1 -surjectivity of UHF-absorbing C*-algebras.

Lemma 3.1. *If A is a unital C*-algebra and $n \geq 2$, then $A \otimes M_{n^\infty}$ is K_1 -surjective.*

Proof. Take $B = A \otimes M_{n^\infty}$ and $[u]_1 \in K_1(B)$, where $u \in \mathcal{U}_m(B)$ for some $m \in \mathbb{N}$. By enlarging m if necessary, we may assume that $m = n^k$ for some $k \in \mathbb{N}$. Since there exists an isomorphism between M_{n^∞} and $M_{n^\infty} \otimes M_{n^k}$ that is approximately unitarily equivalent to the first-factor embedding, there exists a *-isomorphism $\theta : B \rightarrow B \otimes M_{n^k}$ such that θ is approximately unitarily equivalent to $\text{id}_B \otimes 1_{n^k}$.

Take $v = \theta^{-1}(u) \in \mathcal{U}(B)$, then there exists $w \in \mathcal{U}_{n^k}(B)$ such that

$$(3.1) \quad \|u - w(v \otimes 1^{n^k})w^*\| = \|\theta(v) - w(v \otimes 1^{n^k})w^*\| < 1.$$

Then u is homotopic to $w(v \otimes 1^{n^k})w^*$ in $\mathcal{U}_{n^k}(B)$ and thus $[u]_1 = [v^{n^k}]_1 \in K_1(B)$. \square

Lemma 3.1 extends to non-unital UHF-absorbing C*-algebras by a direct application of the following lemma, which will also be applied to non-unital \mathcal{Z} -stable C*-algebras later. The proof is standard and we include it for completeness.

Lemma 3.2. *Let A, B be C^* -algebras with A non-unital and B unital. If $\tilde{A} \otimes B$ is K_1 -surjective and B is K_1 -injective, then $A \otimes B$ is K_1 -surjective, where \otimes is the minimal tensor product.*

Proof. By [3, Lemma 1] for example, we have the following split short exact sequence,

$$(3.2) \quad 0 \longrightarrow A \otimes B \xrightarrow{\iota \otimes \text{id}_B} \tilde{A} \otimes B \xrightarrow{q \otimes \text{id}_B} B \longrightarrow 0,$$

which induces a split short exact sequence for K_1 -groups by split-exactness of K_1 , see [12, Proposition 8.2.5]. Then $(\iota \otimes \text{id}_B)^\sim$ is injective and for any $u \in \mathcal{U}(\tilde{A} \otimes B)$, it is in $\mathcal{U}((A \otimes B)^\sim)$ if and only if $(q \otimes \text{id}_B)(u) \in \mathbb{C}1_B$.

Take any $x \in K_1(A \otimes B) \subseteq K_1(\tilde{A} \otimes B)$, then there exists $v_0 \in \mathcal{U}(\tilde{A} \otimes B)$ such that $x = [v_0]_1$ since $\tilde{A} \otimes B$ is assumed to be K_1 -surjective. Then

$$(3.3) \quad [(q \otimes \text{id}_B)(v_0)]_1 = K_1(q \otimes \text{id}_B)(x) = 0 \in K_1(B),$$

which implies $(q \otimes \text{id}_B)(v_0) \in \mathcal{U}^0(B)$ by K_1 -injectivity of B . Therefore there exists $v_1 \in \mathcal{U}^0(\tilde{A} \otimes B)$ with $(q \otimes \text{id}_B)(v_0) = (q \otimes \text{id}_B)(v_1)$. We take $u = v_1^* v_0 \in \mathcal{U}(\tilde{A} \otimes B)$ so that $[u]_1 = x$ and $(q \otimes \text{id}_B)(u) = 1_B$, which implies that $u \in \mathcal{U}((A \otimes B)^\sim)$. \square

Corollary 3.3. *If A is a C^* -algebra and $n \geq 2$, then $A \otimes M_{n\infty}$ is K_1 -surjective.*

Proof. In the unital case, the result follows from Lemma 3.1. If A is non-unital, since $\tilde{A} \otimes M_{n\infty}$ is K_1 -surjective and $M_{n\infty}$ is unital and K_1 -injective, by Lemma 3.2, we get K_1 -surjectivity of $A \otimes M_{n\infty}$. \square

Now we prove the following main result of the section. We will denote the *suspension* of a C^* -algebra A by $SA = C_0((0, 1)) \otimes A$.

Lemma 3.4. *If A is a unital C^* -algebra, then $A \otimes \mathcal{Z}_{2\infty, 3\infty}$ is K_1 -surjective.*

Proof. We view $A \otimes \mathcal{Z}_{2\infty, 3\infty}$ as a C^* -subalgebra of $C([0, 1], A \otimes M_{6\infty})$, where

$$(3.4) \quad A \otimes \mathcal{Z}_{2\infty, 3\infty} = \{f \in C([0, 1], A \otimes M_{6\infty}) : f(0) \in A \otimes M_{2\infty}, f(1) \in A \otimes M_{3\infty}\}.$$

Then we have the following short exact sequence,

$$(3.5) \quad 0 \rightarrow A \otimes SM_{6\infty} \xrightarrow{\iota} A \otimes \mathcal{Z}_{2\infty, 3\infty} \xrightarrow{q} (A \otimes M_{2\infty}) \oplus (A \otimes M_{3\infty}) \rightarrow 0,$$

where $q(f) = (f(0), f(1))$ for any $f \in A \otimes \mathcal{Z}_{2\infty, 3\infty}$.

Take $u \in \mathcal{U}_m(A \otimes \mathcal{Z}_{2\infty, 3\infty})$ for some $m \in \mathbb{N}$ so that $u(0) \in \mathcal{U}_m(A \otimes M_{2\infty})$ and $u(1) \in \mathcal{U}_m(A \otimes M_{3\infty})$. By Theorem 3.1, both $A \otimes M_{2\infty}$ and $A \otimes M_{3\infty}$ are K_1 -surjective. Then there exists $w_0 \in \mathcal{U}(A \otimes M_{2\infty})$ and $w_1 \in \mathcal{U}(A \otimes M_{3\infty})$ such that $[u(0)]_1 = [w_0]_1 \in K_1(A \otimes M_{2\infty})$ and $[u(1)]_1 = [w_1]_1 \in K_1(A \otimes M_{3\infty})$.

Since u is a continuous path in $\mathcal{U}_m(A \otimes M_{6\infty})$ connecting $u(0)$ and $u(1)$, then

$$(3.6) \quad [w_0]_1 = [u(0)]_1 = [u(1)]_1 = [w_1]_1 \in K_1(A \otimes M_{6\infty}),$$

where we regard $A \otimes M_{2\infty}$ and $A \otimes M_{3\infty}$ as C^* -subalgebras of $A \otimes M_{6\infty}$. Since $A \otimes M_{6\infty}$ is unital and UHF-absorbing, then it is K_1 -injective by [2, Lemma 4.9]. Thus there exists a unitary path $w \in C([0, 1]) \otimes A \otimes M_{6\infty}$ with $w(0) = w_0 \in A \otimes M_{2\infty}$ and $w(1) = w_1 \in A \otimes M_{3\infty}$. Thus $w \in \mathcal{U}(A \otimes \mathcal{Z}_{2\infty, 3\infty})$ with $q(w) = (w_0, w_1)$ and we have

$$(3.7) \quad [q(u)]_1 = [q(w)]_1 \in K_1(A \otimes M_{2\infty}) \oplus K_1(A \otimes M_{3\infty}).$$

Then $[u(w \oplus 1_{m-1})^*]_1 \in \ker(K_1(q)) = \text{Im}(K_1(\iota))$ by half-exactness of K_1 , see [12, Proposition 8.2.4]. Since $A \otimes SM_{6\infty}$ is K_1 -surjective by Corollary 3.3, there exists $z \in \mathcal{U}((A \otimes SM_{6\infty})^\sim) \subseteq$

$A \otimes \mathcal{Z}_{2^\infty, 3^\infty}$ such that $[u(w \oplus 1_{m-1})^*]_1 = K_1(\iota)([z]_1)$. Thus $[u]_1 = [zw]_1 \in K_1(A \otimes \mathcal{Z}_{2^\infty, 3^\infty})$. \square

For infinite coprime supernatural numbers $\mathfrak{p}, \mathfrak{q}$, then $\mathcal{Z}_{\mathfrak{p}, \mathfrak{q}}$ is \mathcal{Z} -stable by [6, Theorem 4.6], since fibers of $\mathcal{Z}_{\mathfrak{p}, \mathfrak{q}}$ are \mathcal{Z} -stable. Then a direct application of Jiang's result shows that $A \otimes \mathcal{Z}_{2^\infty, 3^\infty}$ is K_1 -surjective. Whereas in our approach, Lemma 3.4 is the last ingredient needed to reprove K_1 -surjectivity of \mathcal{Z} -stable C*-algebras.

Theorem 3.5. ([7, Theorem 2]) *If A is a C*-algebra, then $A \otimes \mathcal{Z}$ is K_1 -surjective.*

Proof. In the unital case, this follows from Theorem 2.2, Theorem 3.4 and the fact that K_1 -surjectivity is preserved by inductive limits. For non-unital A , since \mathcal{Z} is K_1 -injective by Theorem 2.3, then $A \otimes \mathcal{Z}$ is K_1 -surjective by Lemma 3.2. \square

In fact, all we need in the proof is that \mathcal{Z} is an inductive limit of generalized dimension drop algebras, regardless of the specific connecting maps.

4. HOMOTOPY GROUPS FOR \mathcal{Z} -STABLE C*-ALGEBRAS

In this section, we prove K -stability for both unital and non-unital \mathcal{Z} -stable C*-algebras. In order to cover the non-unital case, we need to extend the definition for unitary groups. For any C*-algebra A , we denote the minimal proper unitization of A by \tilde{A} , where $\tilde{A} = A \oplus \mathbb{C}$ if A is unital. The canonical maps associated to \tilde{A} are $\iota : A \rightarrow \tilde{A}$ and $q : \tilde{A} \rightarrow \mathbb{C}$. Then the generalized unitary group $\mathcal{V}(A) = \{u \in \mathcal{U}(\tilde{A}) : q(u) = 1\}$ is a topological group and $\mathcal{V}(A) \cong \mathcal{U}(A)$ as topological groups when A is unital. We use the notation $\mathcal{V}_m(A)$ for $\mathcal{V}(M_m \otimes A)$, where $m \geq 1$.

In the unital case, we define $\mathcal{U}_\infty(A)$ to be the inductive limit of groups $\mathcal{U}_m(A)$ with metric preserving connecting maps. Thus $\mathcal{U}_\infty(A)$ is equipped with a well-defined metric from finite stages. For a general C*-algebra, take $\mathcal{V}_\infty(A) = \{u \in \mathcal{U}_\infty(\tilde{A}) : q(u) = 1\}$, where q is the matrix amplification of $q : \tilde{A} \rightarrow \mathbb{C}$. Then there are embeddings $i_{m, m'} : \mathcal{V}_m(A) \rightarrow \mathcal{V}_{m'}(A)$ for $1 \leq m \leq m' \leq \infty$ and it is standard to check that maps $\pi_n(\mathcal{V}_m(A)) \rightarrow \pi_n(\mathcal{U}_m(\tilde{A}))$ induced by inclusions $\iota_m : \mathcal{V}_m(A) \rightarrow \mathcal{U}_m(\tilde{A})$ are isomorphisms for $n \geq 1$ and $m \in \mathbb{N} \cup \{\infty\}$.

By Bott periodicity of K -theory (see [12, Proposition 11.4.1] for example), for any $n \geq 0$, the canonical maps induce the following isomorphisms,

$$(4.1) \quad \pi_n(\mathcal{V}_\infty(A)) \cong \begin{cases} K_0(A) & \text{if } n \text{ is odd,} \\ K_1(A) & \text{if } n \text{ is even.} \end{cases}$$

The definition for K -stability is given by Thomsen using quasi-unitaries in [16]. Since $\mathcal{V}(A)$ is homeomorphic to the topological group of quasi-unitaries in A defined in [16], the following definition for K -stability indeed coincides with Thomsen's.

Definition 4.1. ([16, Definition 3.1]) Let A be a C*-algebra, then it is called K -stable if for any $m \geq 1$ and $n \geq 0$, the induced maps $\pi_n(i_{m, m+1})$ are isomorphisms.

Now we prove the main theorem of the section.

Theorem 4.2. ([7, Theorem 2.8 and Theorem 3]) *Let A be a \mathcal{Z} -stable C*-algebra, then A is K -stable. Moreover, for any integer $n \geq 0$, the following isomorphisms are induced by the canonical embedding $i_{1, \infty} : \mathcal{V}(A) \rightarrow \mathcal{V}_\infty(A)$,*

$$(4.2) \quad \pi_n(\mathcal{V}(A)) \cong \begin{cases} K_0(A) & \text{if } n \text{ is odd,} \\ K_1(A) & \text{if } n \text{ is even.} \end{cases}$$

Proof. When A is \mathcal{Z} -stable, then A is K_1 -injective and K_1 -surjective by Theorem 2.3 and Theorem 3.5, which implies $\pi_0(\mathcal{U}(\tilde{A})) \cong K_1(\tilde{A}) \cong \pi_0(\mathcal{U}_\infty(\tilde{A}))$ by (4.1). Since $\pi_0(\iota_0)$ and $\pi_0(\iota_\infty)$ are isomorphisms, then $i_{1,\infty}$ induces an isomorphism through

$$(4.3) \quad \pi_0(\mathcal{V}(A)) \cong \pi_0(\mathcal{U}(\tilde{A})) \cong \pi_0(\mathcal{U}_\infty(\tilde{A})) \cong \pi_0(\mathcal{V}_\infty(A)) = K_1(A).$$

Since \mathcal{Z} -stability of A implies that of $M_m \otimes A$, then the maps $i_{m,\infty}$ induce $\pi_0(\mathcal{V}_m(A)) \cong K_1(A)$ for $m \geq 1$ by the argument above and thus all the maps $\pi_0(i_{m,m+1})$ are isomorphisms.

For $n \geq 1$, there is a canonical identification between $\pi_n(\mathcal{V}(A))$ and $\pi_0(\mathcal{V}(S^n A))$ by [16, Lemma 2.3], where $S^n A = C_0((0, 1)^n) \otimes A$ is the n^{th} suspension of A . When A is \mathcal{Z} -stable, then $S^n(M_m(A))$ is \mathcal{Z} -stable for any $m, n \geq 1$. Thus by (4.3), the maps $i_{m,\infty}$ induce the following isomorphisms for $m \geq 1$,

$$(4.4) \quad \pi_n(\mathcal{V}_m(A)) = \pi_0(\mathcal{V}(S^n(M_m(A)))) \cong K_1((S^n(M_m(A)))) \cong K_{n+1}(A).$$

Thus $\pi_n(i_{m,m+1})$ are isomorphisms for $m \geq 1$ and $n \geq 0$, which implies K -stability of A . Further by Bott periodicity of K -theory, for $m \geq 1$ and $n \geq 0$,

$$(4.5) \quad \pi_n(\mathcal{V}_m(A)) \cong \begin{cases} K_0(A) & \text{if } n \text{ is odd,} \\ K_1(A) & \text{if } n \text{ is even.} \end{cases}$$

In particular, we get (4.2). □

REFERENCES

- [1] B. Blackadar. Rational C^* -algebras and nonstable K -theory. In *Proceedings of the Seventh Great Plains Operator Theory Seminar (Lawrence, KS, 1987)*, volume 20, pages 285–316, 1990.
- [2] J. R. Carrión, J. Gabe, C. Schafhauser, A. Tikuisis, and S. White. Classifying $*$ -homomorphisms I: Unital simple nuclear C^* -algebras. *arXiv:2307.06480 [math.OA]*, 2023.
- [3] P. de la Harpe, A. G. Robertson, and A. Valette. On exactness of group C^* -algebras. *Quart. J. Math. Oxford Ser. (2)*, 45(180):499–513, 1994.
- [4] G. Gong, X. Jiang, and H. Su. Obstructions to \mathcal{Z} -stability for unital simple C^* -algebras. *Canad. Math. Bull.*, 43(4):418–426, 2000.
- [5] G. Gong, H. Lin, and Z. Niu. A classification of finite simple amenable \mathcal{Z} -stable C^* -algebras, I: C^* -algebras with generalized tracial rank one. *C. R. Math. Acad. Sci. Soc. R. Can.*, 42(3):63–450, 2020.
- [6] I. Hirshberg, M. Rørdam, and W. Winter. $C_0(X)$ -algebras, stability and strongly self-absorbing C^* -algebras. *Math. Ann.*, 339(3):695–732, 2007.
- [7] X. Jiang. Nonstable K -theory for \mathcal{Z} -stable C^* -algebras. *arXiv:math/9707228 [math.OA]*, 1997.
- [8] X. Jiang and H. Su. On a simple unital projectionless C^* -algebra. *Amer. J. Math.*, 121(2):359–413, 1999.
- [9] M. A. Rieffel. The homotopy groups of the unitary groups of noncommutative tori. *J. Operator Theory*, 17(2):237–254, 1987.
- [10] M. Rørdam. A simple C^* -algebra with a finite and an infinite projection. *Acta Math.*, 191(1):109–142, 2003.
- [11] M. Rørdam. The stable and the real rank of \mathcal{Z} -absorbing C^* -algebras. *Internat. J. Math.*, 15(10):1065–1084, 2004.
- [12] M. Rørdam, F. Larsen, and N. Laustsen. *An introduction to K -theory for C^* -algebras*, volume 49 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2000.
- [13] M. Rørdam and W. Winter. The Jiang-Su algebra revisited. *J. Reine Angew. Math.*, 642:129–155, 2010.
- [14] C. Schafhauser. A new proof of the Tikuisis-White-Winter theorem. *J. Reine Angew. Math.*, 759:291–304, 2020.
- [15] A. Schemaitat. The Jiang-Su algebra is strongly self-absorbing revisited. *J. Funct. Anal.*, 282(6):Paper No. 109347, 39, 2022.
- [16] K. Thomsen. Nonstable K -theory for operator algebras. *K-Theory*, 4(3):245–267, 1991.
- [17] A. Toms. On the independence of K -theory and stable rank for simple C^* -algebras. *J. Reine Angew. Math.*, 578:185–199, 2005.
- [18] A. S. Toms. Higher homotopy groups of Cuntz classes. *Adv. Math.*, 443, 2024.

- [19] J. Villadsen. Simple C^* -algebras with perforation. *J. Funct. Anal.*, 154(1):110–116, 1998.
- [20] W. Winter. Localizing the Elliott conjecture at strongly self-absorbing C^* -algebras. *J. Reine Angew. Math.*, 692:193–231, 2014.
- [21] S. Zhang. Matricial structure and homotopy type of simple C^* -algebras with real rank zero. *J. Operator Theory*, 26(2):283–312, 1991.

Shanshan Hua, Mathematical Institute, University of Oxford, Oxford, OX2 6GG, UK.
Email address: `shanshan.hua@maths.ox.ac.uk`