

New characterization of weighted inequalities involving superposition of Hardy integral operators

Amiran Gogatishvili¹ and Tuğçe Ünver^{2*}

¹ Institute of Mathematics of the Czech Academy of Sciences, Žitná 25, 115 67 Praha 1, Czech Republic

²Department of Mathematics, Kirikkale University, 71450 Yahsihan, Kirikkale, Turkey

*Corresponding author. E-mail: tugceunver@kku.edu.tr, ORCID:0000-0003-0414-8400

Contributing author: gogatish@math.cas.cz, ORCID:0000-0003-3459-0355

ABSTRACT. Let $1 \leq p < \infty$ and $0 < q, r < \infty$. We characterize the validity of the inequality for the composition of the Hardy operator,

$$\left(\int_a^b \left(\int_a^x \left(\int_a^t f(s) ds \right)^q u(t) dt \right)^{\frac{x}{q}} w(x) dx \right)^{\frac{1}{r}} \leq C \left(\int_a^b f(x)^p v(x) dx \right)^{\frac{1}{p}}$$

for all non-negative measurable functions f on (a, b) , $-\infty \leq a < b \leq \infty$. We construct a more straightforward discretization method than those previously presented in the literature, and we provide some new scales of weight characterizations of this inequality in both discrete and continuous forms and we obtain previous characterizations as the special case of the parameter.

Keywords: weighted Hardy inequality, iterated operators, Copson operator, Hardy operator, inequalities for monotone functions

2010 Mathematics Subject Classification: 26D10, 26D15

1 Introduction and the main results

Let $-\infty \leq a < b \leq \infty$. Denote by $\mathfrak{M}^+(a, b)$ the set of all non-negative measurable functions on (a, b) and $\mathfrak{M}^\uparrow(a, b)$ is the class of non-decreasing elements of $\mathfrak{M}^+(a, b)$.

In operator theory, weighted inequalities involving operator composition may be found in various topics. Let $0 < q, r < \infty$ and $1 \leq p < \infty$. The validity of inequalities

$$\left(\int_0^\infty \left(\int_0^x \left(\int_t^\infty h(s) ds \right)^q u(t) dt \right)^{\frac{x}{q}} w(x) dx \right)^{\frac{1}{r}} \leq C \left(\int_0^\infty h(x)^p v(x) dx \right)^{\frac{1}{p}}, \quad (1.1)$$

and

$$\left(\int_0^\infty \left(\int_0^x \left(\int_0^t h(s) ds \right)^q u(t) dt \right)^{\frac{x}{q}} w(x) dx \right)^{\frac{1}{r}} \leq C \left(\int_0^\infty h(x)^p v(x) dx \right)^{\frac{1}{p}}, \quad (1.2)$$

for all $h \in \mathfrak{M}^+(0, \infty)$ are crucial because many classical inequalities can be reduced to them. For example, duality techniques reduce the embeddings between Lorentz-type spaces, Morrey-type spaces, and Cesàro-type spaces to the weighted iterated inequalities (see, e.g., [4, 5, 7, 9,

34]). On the other hand, characterizations of weighted bilinear Hardy and Copson inequalities reduce to the characterizations of iterated Hardy inequalities (see, e.g., [1, 23, 33]).

Various approaches have been used to handle inequalities (1.1) and (1.2) resulting in conditions of a different nature. Inequality (1.1) is investigated thoroughly. The most recent results are presented in the paper [20]. Alternative characterizations of inequalities (1.1) and (1.2) are found in [19]. Detailed information on the development and history of this inequality may be found in the recent paper [6].

Note that, when $q = 1$ using Fubini's Theorem, inequality (1.2) reduces to the weighted Hardy-type inequality involving kernel, that is,

$$\left(\int_0^\infty \left(\int_0^x \left(\int_s^x u(t) dt \right) h(s) ds \right)^r w(x) dx \right)^{\frac{1}{r}} \leq C \left(\int_0^\infty h(x)^p v(x) dx \right)^{\frac{1}{p}}, \quad h \in \mathfrak{M}^+(0, \infty). \quad (1.3)$$

Inequality (1.3) was completely characterized in [26, 3, 28, 32] when $1 \leq r, p < \infty$. However, for a long period, there was no adequate characterization in the case when $0 < r < 1 \leq p < \infty$. Several attempts have been made to tackle this case (see, e.g., [32, 25, 29, 13]); in some works, necessary and sufficient conditions did not match, while in others, characterization had a discrete form or involved auxiliary functions. Hence, it was not easily verifiable. Finally, in [22], the missing integral conditions were provided.

In the general cases, (1.2) is characterized in [8], but the conditions are in a non-standard form. It was also considered in [30], but the conditions are not applicable because they involve auxiliary functions. Recently, in [24], a more complicated discretization method is used to establish a characterization of the same inequality that involves iteration of the Copson operators and is restricted to non-degenerate weights, and the case $p = 1$ is presented without a proof. In our approach, the case $p > 1$ is not separated from $p = 1$.

Recently, in [6], with a new and simpler discretization technique that requires neither parameter restrictions nor non-degeneracy conditions, characterization of (1.1) is given. We adapt this approach to the specific demands of the inequality considered in this paper. Our technique allows us to obtain a scale of characterization which was not possible before; moreover, we obtain the previous characterizations of inequality (1.2) as a special case (more, precisely for $\beta = 0$, see Theorem 1.1).

We would like to point out that the characterization of Hardy inequality involving non-decreasing functions, that is,

$$\left(\int_0^\infty \left(\int_0^x f(s)u(s) ds \right)^q w(x) dx \right)^{\frac{1}{q}} \leq C \left(\int_0^\infty f(x)^p v(x) dx \right)^{\frac{1}{p}}, \quad f \in \mathfrak{M}^\uparrow(0, \infty), \quad (1.4)$$

is obtained directly from inequality (1.2) without any further work (see the proof of Theorem 1.2) as a direct outcome of our main theorem (see Theorem 1.1).

We should also mention that in [12, 13], using reduction techniques, (1.4) is reduced to inequality (1.3). However, as we have already mentioned, at that point of time the characterizations of the reduced inequalities were not known. Corollary 3.2 from [13] provides a characterization of (1.4) but the result is non-standard. The earlier works on inequality (1.4) can be found in [17, 27, 14, 15].

We should note that Theorem 1.1 allowed us to give a scale of characterizations of (1.4), and we would like to provide the characterization here to integrate all relevant parameter choices into a single theorem for the reader's convenience (see, Theorem 1.2).

As one can see in Section 3, the discretization method transforms the inequality at hand equivalently to discrete inequalities that involve local characterizations of inequalities having low-order iterations. For this very reason, our aim in this paper is to revisit inequality (1.2) on (a, b) where $-\infty \leq a < b \leq \infty$.

Let $-\infty \leq a < b \leq \infty$ and a weight be a positive measurable function on (a, b) . The principal goal of this study is to determine the scale of necessary and sufficient conditions on weights u, v, w on (a, b) for which

$$\left(\int_a^b \left(\int_a^x \left(\int_a^t f(s) ds \right)^q u(t) dt \right)^{\frac{r}{q}} w(x) dx \right)^{\frac{1}{r}} \leq C \left(\int_a^b f(x)^p v(x) dx \right)^{\frac{1}{p}} \quad (1.5)$$

holds for $f \in \mathfrak{M}^+(a, b)$, with exponents $1 \leq p < \infty$ and $0 < q, r < \infty$. It is worth noting that if $p < 1$, inequality (1.5) only holds for trivial functions.

Let us first review the essential notations and conventions before presenting our main results. The left and right sides of the inequality numbered by $(*)$ are denoted by $\text{LHS}(*)$ and $\text{RHS}(*)$, respectively. We put $0 \cdot \infty = \infty / \infty = 0 / 0 = 0$. The symbol $A \lesssim B$ means a constant $c > 0$ exists such that $A \leq cB$ where c depends only on the parameters p, q, r . If both $A \lesssim B$ and $B \lesssim A$, then we write $A \approx B$.

For $1 \leq p < \infty$, non-negative measurable functions v and w on (a, b) , and $x, y \in [a, b]$, denote by

$$V_p(x, y) := \begin{cases} \left(\int_x^y v^{-\frac{1}{p-1}} \right)^{\frac{p-1}{p}}, & 1 < p < \infty, \\ \text{ess sup}_{s \in (x, y)} v(s)^{-1}, & p = 1, \end{cases} \quad (1.6)$$

and

$$\mathcal{W}(x) = \int_x^b w(s) ds.$$

Now, we are ready to formulate our main results.

Theorem 1.1. *Let $1 \leq p < \infty$, $0 < q, r < \infty$, $-\infty < \beta < 1$ and let u, v, w be weights on (a, b) such that $0 < \mathcal{W}(t) < \infty$ for all $t \in (a, b)$. Then inequality (1.5) holds for all $f \in \mathfrak{M}^+(a, b)$ and only if*

(i) $p \leq r$, $p \leq q$ and

$$C_1 := \text{ess sup}_{x \in (a, b)} \left(\int_x^b \mathcal{W}(t)^{-\beta} w(t) \left(\int_x^t \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} dt \right)^{\frac{1}{r}} V_p(a, x) < \infty. \quad (1.7)$$

Moreover, the best constant C in inequality (1.5) satisfies $C \approx C_1$.

(ii) $r < p \leq q$,

$$C_2 := \left(\int_a^b \mathcal{W}(x)^{\frac{(1-\beta)p}{p-r}-1} w(x) \text{ess sup}_{t \in (a, x)} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{rp}{q(p-r)}} V_p(a, t)^{\frac{pr}{p-r}} dx \right)^{\frac{p-r}{pr}} < \infty,$$

and

$$C_3 := \left(\int_a^b \left(\int_x^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_x^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \mathcal{W}(x)^{-\beta} w(x)$$

$$\times \operatorname{ess\,sup}_{t \in (a,x)} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a,t)^{\frac{pr}{p-r}} dx \Big)^{\frac{p-r}{pr}} < \infty. \quad (1.8)$$

Moreover, the best constant C in inequality (1.5) satisfies $C \approx C_2 + C_3$.

(iii) $q < p \leq r$, $C_1 < \infty$ and

$$C_4 := \sup_{x \in (a,b)} \mathcal{W}(x)^{\frac{1-\beta}{r}} \left(\int_a^x \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(a,t)^{\frac{pq}{p-q}} dt \right)^{\frac{p-q}{pq}} < \infty,$$

where C_1 is defined in (1.7). Moreover, the best constant C in inequality (1.5) satisfies $C \approx C_1 + C_4$.

(iv) $r < p$, $q < p$, $C_3 < \infty$ and

$$C_5 := \left(\int_a^b \mathcal{W}(x)^{\frac{(1-\beta)p}{p-r}-1} w(x) \right. \\ \left. \times \left(\int_a^x \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(a,t)^{\frac{pq}{p-q}} dt \right)^{\frac{r(p-q)}{q(p-r)}} dx \right)^{\frac{p-r}{pq}} < \infty,$$

where C_3 is defined in (1.8). Moreover, the best constant C in inequality (1.5) satisfies $C \approx C_3 + C_5$.

Theorem 1.2. *Let $0 < p, q < \infty$, $-\infty < \beta < 1$ and u, v, w be weights on (a, b) such that $0 < \mathcal{W}(t) < \infty$ for all $t \in (a, b)$. Then inequality*

$$\left(\int_a^b \left(\int_a^x f(s)u(s)ds \right)^q w(x)dx \right)^{\frac{1}{q}} \leq C \left(\int_a^b f(x)^p v(x)dx \right)^{\frac{1}{p}} \quad (1.9)$$

holds for all $f \in \mathfrak{M}^\uparrow(a, b)$ if and only if

(i) $p \leq q$, $p \leq 1$ and

$$\mathcal{C}_1 := \operatorname{ess\,sup}_{x \in (a,b)} \left(\int_x^b \mathcal{W}(t)^{-\beta} w(t) \left(\int_x^t \mathcal{W}^{\frac{\beta}{q}} u \right)^q dt \right)^{\frac{1}{q}} \left(\int_x^b v \right)^{-\frac{1}{p}} < \infty. \quad (1.10)$$

Moreover, the best constant C in inequality (1.9) satisfies $C \approx \mathcal{C}_1$.

(ii) $q < p \leq 1$,

$$\mathcal{C}_2 := \left(\int_a^b \mathcal{W}(x)^{\frac{(1-\beta)p}{p-q}-1} w(x) \operatorname{ess\,sup}_{t \in (a,x)} \left(\int_t^x \mathcal{W}^{\frac{\beta}{q}} u \right)^{\frac{pq}{p-q}} \left(\int_t^b v \right)^{-\frac{q}{p-q}} dx \right)^{\frac{p-q}{pq}} < \infty,$$

and

$$\mathcal{C}_3 := \left(\int_a^b \left(\int_x^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_x^s \mathcal{W}^{\frac{\beta}{q}} u \right)^q ds \right)^{\frac{q}{p-q}} \mathcal{W}(x)^{-\beta} w(x) \right. \\ \left. \times \operatorname{ess\,sup}_{t \in (a,x)} \left(\int_t^x \mathcal{W}^{\frac{\beta}{q}} u \right)^q \left(\int_t^b v \right)^{-\frac{q}{p-q}} dx \right)^{\frac{p-q}{pq}} < \infty. \quad (1.11)$$

Moreover, the best constant C in inequality (1.9) satisfies $C \approx \mathcal{C}_2 + \mathcal{C}_3$.

(iii) $1 < p \leq q$, $C_1 < \infty$ and

$$\mathcal{C}_4 := \sup_{x \in (a,b)} \mathcal{W}(x)^{\frac{1-\beta}{q}} \left(\int_a^x \left(\int_t^x \mathcal{W}^{\frac{\beta}{q}} u \right)^{\frac{1}{p-1}} \mathcal{W}(t)^{\frac{\beta}{q}} u(t) \left(\int_t^b v \right)^{-\frac{1}{p-1}} dt \right)^{\frac{p-1}{p}} < \infty,$$

where \mathcal{C}_1 is defined in (1.10). Moreover, the best constant C in inequality (1.9) satisfies $C \approx \mathcal{C}_1 + \mathcal{C}_4$.

(iv) $q < p$, $1 < p$, $\mathcal{C}_3 < \infty$ and

$$\begin{aligned} \mathcal{C}_5 := & \left(\int_a^b \mathcal{W}^{\frac{(1-\beta)p}{p-q}-1} w(x) \right. \\ & \left. \times \left(\int_a^x \left(\int_t^x \mathcal{W}^{\frac{\beta}{q}} u \right)^{\frac{1}{p-1}} \mathcal{W}(t)^{\frac{\beta}{q}} u(t) \left(\int_t^b v \right)^{-\frac{1}{p-1}} dt \right)^{\frac{q(p-1)}{p-q}} dx \right)^{\frac{p-q}{pq}} < \infty, \end{aligned}$$

where \mathcal{C}_3 is defined in (1.11). Moreover, the best constant C in inequality (1.9) satisfies $C \approx \mathcal{C}_3 + \mathcal{C}_5$.

Proofs of Theorem 1.1 and Theorem 1.2 will be given in Section 4.

2 Preliminary Results

In this section, we cover the foundations of discretization and several new results that will be employed frequently throughout the proof of the main theorem.

Definition 2.1. Let $N \in \mathbb{Z} \cup \{-\infty\}$, $M \in \mathbb{Z} \cup \{+\infty\}$, $N < M$, and $\{a_k\}_{k=N}^M$ be a sequence of positive numbers. We say that $\{a_k\}_{k=N}^M$ is *geometrically decreasing* if

$$\sup \left\{ \frac{a_{k+1}}{a_k}, \quad N \leq k \leq M \right\} < 1.$$

Lemma 2.2. [10] Let $\alpha > 0$ and $n \in \mathbb{Z} \cup \{-\infty\}$. If $\{\tau_k\}_{k=n}^{\infty}$ is a geometrically decreasing sequence, then,

$$\sup_{n \leq k < \infty} \tau_k \sum_{i=n}^k a_i \approx \sup_{n \leq k < \infty} \tau_k a_k, \quad (2.1)$$

$$\sum_{k=n}^{\infty} \tau_k \left(\sum_{i=n}^k a_i \right)^{\alpha} \approx \sum_{k=n}^{\infty} \tau_k a_k^{\alpha}, \quad (2.2)$$

and

$$\sum_{k=n}^{\infty} \tau_k \sup_{n \leq i \leq k} a_i \approx \sum_{k=n}^{\infty} \tau_k a_k, \quad (2.3)$$

for all non-negative sequences $\{a_k\}_{k=n}^{\infty}$.

Lemma 2.3. Let $\alpha > 0$ and $n \in \mathbb{Z} \cup \{-\infty\}$. Assume that $\{x_k\}_{k=n}^{\infty} \subset (a, b)$ is a strictly increasing sequence. If $\{\tau_k\}_{k=n+1}^{\infty}$ is a geometrically decreasing sequence, then,

$$\sup_{n+1 \leq k < \infty} \tau_k \left(\int_{x_n}^{x_k} g \right) \approx \sup_{n+1 \leq k < \infty} \tau_k \left(\int_{x_{k-1}}^{x_k} g \right), \quad (2.4)$$

$$\sum_{k=n+1}^{\infty} \tau_k \left(\int_{x_n}^{x_k} g \right)^\alpha \approx \sum_{k=n+1}^{\infty} \tau_k \left(\int_{x_{k-1}}^{x_k} g \right)^\alpha, \quad (2.5)$$

and

$$\sum_{k=n+1}^{\infty} \tau_k \operatorname{ess\,sup}_{s \in (x_n, x_k)} g(s) \approx \sum_{k=n+1}^{\infty} \tau_k \operatorname{ess\,sup}_{s \in (x_{k-1}, x_k)} g(s), \quad (2.6)$$

for all non-negative measurable g on (a, b) .

Proof. For each $n \in \mathbb{Z} \cup \{-\infty\}$, we can write

$$\int_{x_n}^{x_k} g = \sum_{i=n+1}^k \int_{x_{i-1}}^{x_i} g.$$

Then (2.4) and (2.5) are direct consequences of (2.1) and (2.2), respectively.

Similarly, for each $n \in \mathbb{Z} \cup \{-\infty\}$, we have

$$\operatorname{ess\,sup}_{s \in (x_n, x_k)} g(s) = \sup_{n+1 \leq i \leq k} \operatorname{ess\,sup}_{s \in (x_{i-1}, x_i)} g(s),$$

so that applying (2.3), we obtain (2.6). \square

Lemma 2.4. *Let $\alpha > 0$ and $n \in \mathbb{Z} \cup \{-\infty\}$. Assume that $\{x_k\}_{k=n}^{\infty} \subset (a, b)$ is a strictly increasing sequence, $\{\tau_k\}_{k=n+1}^{\infty}$ is a geometrically decreasing sequence, and $\{\sigma_k\}_{k=n}^{\infty}$ is a positive non-decreasing sequence. Then*

$$\sup_{n+1 \leq k < \infty} \tau_k \sup_{n \leq i < k} \left(\int_{x_i}^{x_k} g \right)^\alpha \sigma_i \approx \sup_{n+1 \leq k < \infty} \tau_k \left(\int_{x_{k-1}}^{x_k} g \right)^\alpha \sigma_{k-1} \quad (2.7)$$

and

$$\sum_{k=n+1}^{\infty} \tau_k \sup_{n \leq i < k} \left(\int_{x_i}^{x_k} g \right)^\alpha \sigma_i \approx \sum_{k=n+1}^{\infty} \tau_k \left(\int_{x_{k-1}}^{x_k} g \right)^\alpha \sigma_{k-1} \quad (2.8)$$

hold for all non-negative measurable g on (a, b) .

Proof. Let us start with the equivalency (2.7). Since $\{\tau_k\}_{k=n+1}^{\infty}$ is a geometrically decreasing sequence, interchanging supremum and (2.4) give

$$LHS(2.7) = \sup_{n+1 \leq i < \infty} \sigma_i \sup_{i+1 \leq k < \infty} \tau_k \left(\int_{x_i}^{x_k} g \right)^\alpha \approx \sup_{n+1 \leq i < \infty} \sigma_i \sup_{i+1 \leq k < \infty} \tau_k \left(\int_{x_{k-1}}^{x_k} g \right)^\alpha.$$

Interchanging supremum once again and monotonicity of $\{\sigma_k\}_{k=n}^{\infty}$ results in

$$LHS(2.7) \approx \sup_{n+1 \leq k < \infty} \tau_k \left(\int_{x_{k-1}}^{x_k} g \right)^\alpha \sup_{n \leq i \leq k-1} \sigma_i = RHS(2.7).$$

Let us now tackle (2.8). Monotonicity of $\{\sigma_k\}_{k=n}^{\infty}$ gives that

$$LHS(2.8) \leq \sum_{k=n+1}^{\infty} \tau_k \sup_{n \leq i < k} \left(\sum_{j=i}^{k-1} \sigma_j^\alpha \int_{x_j}^{x_{j+1}} g \right)^\alpha = \sum_{k=n+1}^{\infty} \tau_k \left(\sum_{j=n}^{k-1} \sigma_j^\alpha \int_{x_j}^{x_{j+1}} g \right)^\alpha.$$

Then, using (2.2), we have the following upper estimate

$$LHS(2.8) \leq \sum_{k=n+1}^{\infty} \tau_k \left(\sum_{j=n+1}^k \sigma_{j-1}^{\frac{1}{\alpha}} \int_{x_{j-1}}^{x_j} g \right)^{\alpha} \approx RHS(2.8).$$

On the other hand, the reverse estimate is clear, and the proof is complete. \square

Definition 2.5. Let w be a non-negative measurable function on (a, b) such that $0 < \mathcal{W}(t) < \infty$ for all $t \in (a, b)$. A strictly increasing sequence $\{x_k\}_{k=N}^{\infty} \subset [a, b]$ is said to be a discretizing sequence of the function \mathcal{W} , if it satisfies $\mathcal{W}(x_k) \approx 2^{-k}$, $N \leq k < \infty$. If $N > -\infty$ then $x_N := a$, otherwise $x_{-\infty} := \lim_{k \rightarrow -\infty} x_k = a$.

Remark 2.6. Let w be a non-negative measurable function on (a, b) such that $0 < \mathcal{W}(t) < \infty$ for all $t \in (a, b)$. The Darboux property of continuous functions implies that the discretizing sequence exists for \mathcal{W} . We can define the discretizing sequence in the following way:

$$N = \sup \left\{ k \in \mathbb{Z} : 2^{-k} \geq \mathcal{W}(t) \text{ for every } t \in (a, b) \right\}.$$

If the set is empty then $N = -\infty$. If $\lim_{t \rightarrow a+} \mathcal{W}(t) < \infty$, then $N > -\infty$ and $x_N = a$ and for $k > N$

$$x_k = \sup \left\{ t \in (a, b) : 2^{-k} \leq \mathcal{W}(t) \right\}.$$

It is worth noting that if $N = -\infty$, then $N + 1$ is also $-\infty$.

Lemma 2.7. Let $\alpha > 0$ and $N \in \mathbb{Z} \cup \{-\infty\}$. Assume that w is a weight on (a, b) such that $0 < \mathcal{W}(t) < \infty$ for all $t \in (a, b)$, and $\{x_k\}_{k=N}^{\infty}$ is a discretizing sequence of the function \mathcal{W} . Then for any $n : N \leq n$,

$$\int_{x_n}^b \mathcal{W}(x)^{\alpha-1} w(x) h(x) dx \approx \sum_{k=n+1}^{\infty} 2^{-k\alpha} h(x_k) \quad (2.9)$$

and

$$\operatorname{ess\,sup}_{x \in (x_n, b)} \mathcal{W}(x)^{\alpha} h(x) \approx \sup_{n+1 \leq k} 2^{-k\alpha} h(x_k) \quad (2.10)$$

hold for all non-negative and non-decreasing h on (a, b) .

Proof. Monotonicity of h and properties of the discretizing sequence $\{x_k\}_{k=N}^{\infty}$ yield

$$\begin{aligned} LHS(2.9) &= \sum_{k=n+1}^{\infty} \int_{x_{k-1}}^{x_k} h(x) \mathcal{W}(x)^{\alpha-1} w(x) dx \lesssim \sum_{k=n+1}^{\infty} h(x_k) \int_{x_{k-1}}^{x_k} d[-\mathcal{W}(x)^{\alpha}] \\ &\approx \sum_{k=n+1}^{\infty} 2^{-k\alpha} h(x_k) = RHS(2.9), \end{aligned}$$

and, conversely

$$LHS(2.9) \geq \sum_{k=n+1}^{\infty} \int_{x_k}^{x_{k+1}} h(x) \mathcal{W}(x)^{\alpha-1} w(x) dx \gtrsim \sum_{k=n+1}^{\infty} h(x_k) \int_{x_k}^{x_{k+1}} d[-\mathcal{W}(x)^{\alpha}]$$

$$\approx \sum_{k=n+1}^{\infty} 2^{-k\alpha} h(x_k) = RHS(2.9).$$

Thus, (2.9) holds.

On the other hand, similarly,

$$LHS(2.10) = \sup_{n+1 \leq k < \infty} \operatorname{ess\,sup}_{x \in (x_{k-1}, x_k)} \mathcal{W}(x)^\alpha h(x) \approx \sup_{n+1 \leq k < \infty} 2^{-k\alpha} \operatorname{ess\,sup}_{x \in (x_{k-1}, x_k)} h(x) = RHS(2.10)$$

holds. \square

Theorem 2.8. [11, Theorem 4.5] Let $p, q \in (0, \infty)$, $N, M \in \mathbb{Z} \cup \{-\infty, \infty\}$, $N < M$. Assume that $\{w_k\}_{k=N}^M$ and $\{v_k\}_{k=N}^M$ be two sequences of positive real numbers. Then inequality

$$\left(\sum_{k=N}^M a_k^q v_k^q \right)^{\frac{1}{q}} \leq C \left(\sum_{k=N}^M a_k^p w_k^p \right)^{\frac{1}{p}} \quad (2.11)$$

holds for every sequence $\{a_k\}_{k=N}^M$ of non-negative real numbers if and only if either

(i) $p \leq q$ and

$$L_1 = \sup_{N \leq k \leq M} \{v_k w_k^{-1}\} < \infty,$$

or

(ii) $p > q$ and

$$L_2 = \left(\sum_{k=N}^M v_k^{\frac{pq}{p-q}} w_k^{-\frac{pq}{p-q}} \right)^{\frac{p-q}{pq}} < \infty.$$

Moreover, if we denote by C the optimal embedding constant in (2.11), then

$$C = \begin{cases} L_1 & \text{in the case (i),} \\ L_2 & \text{in the case (ii).} \end{cases}$$

Theorem 2.9. [11, Theorem 4.6] Assume that $0 < p, q < \infty$, $N, M \in \mathbb{Z} \cup \{-\infty, \infty\}$, $N < M$, $\{a_k\}_{k=N}^M$ and $\{b_k\}_{k=N}^M$ are sequences of non-negative real numbers. Then the inequality

$$\left(\sum_{k=N}^M \left(\sum_{i=N}^k x_i b_i \right)^q a_k \right)^{\frac{1}{q}} \leq C \left(\sum_{k=N}^M x_k^p \right)^{\frac{1}{p}} \quad (2.12)$$

holds for every sequence $\{x_k\}_{k=N}^M$ of non-negative real numbers if and only if

(i) either $p \leq 1$, $p \leq q$ and

$$H_1 = \sup_{N \leq k \leq M} \left\{ \left(\sum_{i=k}^M a_i \right)^{\frac{1}{q}} b_k \right\} < \infty,$$

(ii) or $q < p \leq 1$ and

$$H_2 = \left(\sum_{k=N}^M a_k \left(\sum_{i=k}^M a_i \right)^{\frac{q}{p-q}} \sup_{N \leq i \leq k} b_i^{\frac{qp}{p-q}} \right)^{\frac{p-q}{pq}} < \infty,$$

(iii) or $1 < p, q < p$ and

$$H_3 = \left(\sum_{k=N}^M a_k \left(\sum_{i=k}^M a_i \right)^{\frac{q}{p-q}} \left(\sum_{i=N}^k b_i^{\frac{p}{p-1}} \right)^{\frac{q(p-1)}{p-q}} \right)^{\frac{p-q}{pq}} < \infty,$$

(iv) or $1 < p \leq q$ and

$$H_4 = \sup_{N \leq k \leq M} \left\{ \left(\sum_{i=k}^M a_i \right)^{\frac{1}{q}} \left(\sum_{i=N}^k b_i^{\frac{p}{p-1}} \right)^{\frac{p-1}{p}} \right\} < \infty.$$

Moreover, if we denote by C the best constant in (2.12), then

$$C \approx \begin{cases} H_1 & \text{in the case (i),} \\ H_2 & \text{in the case (ii),} \\ H_3 & \text{in the case (iii),} \\ H_4 & \text{in the case (iv).} \end{cases}$$

Moreover, the multiplicative constants in all the equivalences above depend only on p, q .

3 Discrete Characterization

We begin this section by observing that inequality (1.5) is equivalent to two other discrete inequalities, and we present the characterization in discrete form, which is noteworthy on its own.

Let us start with the discretization of inequality (1.5). To this end we need the following notation, denote by $M(x_{k-1}, x_k)$ the best constant of weighted Hardy inequality, that is,

$$B(x_{k-1}, x_k) := \sup_{h \in \mathfrak{M}^+(x_{k-1}, x_k)} \frac{\left(\int_{x_{k-1}}^{x_k} \left(\int_{x_{k-1}}^t h(s) ds \right)^q u(t) dt \right)^{\frac{1}{q}}}{\left(\int_{x_{k-1}}^{x_k} h(t)^p v(t) dt \right)^{\frac{1}{p}}}$$

and using the classical characterizations of weighted Hardy inequalities (see, [21, 31]), we have for $1 \leq p < \infty, 0 < q < \infty$

$$B(x_{k-1}, x_k) \approx \begin{cases} \operatorname{ess\,sup}_{t \in (x_{k-1}, x_k)} \left(\int_t^{x_k} u \right)^{\frac{1}{q}} V_p(x_{k-1}, t) & \text{if } p \leq q, \\ \left(\int_{x_{k-1}}^{x_k} \left(\int_t^{x_k} u \right)^{\frac{q}{p-q}} u(t) V_p(x_{k-1}, t)^{\frac{pq}{p-q}} dt \right)^{\frac{p-q}{pq}} & \text{if } q < p. \end{cases} \quad (3.1)$$

Furthermore, observe that $V_p(x_{k-1}, x_k)$, $N+1 \leq k$ defined in (1.6) can be expressed as

$$\sup_{g \in \mathfrak{M}^+(x_{k-1}, x_k)} \frac{\int_{x_{k-1}}^{x_k} g(t) dt}{\left(\int_{x_{k-1}}^{x_k} g(t)^p v(t) dt \right)^{\frac{1}{p}}} = V_p(x_{k-1}, x_k).$$

Lemma 3.1. *Let $1 \leq p < \infty$, $0 < q, r < \infty$ and let u, v, w be weights on (a, b) such that $0 < \mathcal{W}(t) < \infty$ for all $t \in (a, b)$, and $\{x_k\}_{k=N}^{\infty} \subset [a, b]$ is a discretizing sequence of the function \mathcal{W} . Then, there exists a positive constant C such that inequality (1.5) holds for all $f \in \mathfrak{M}^+(a, b)$ if and only if there exist positive constants \mathcal{C}' and \mathcal{C}'' such that*

$$\left(\sum_{k=N+1}^{\infty} 2^{-k} \left(\int_{x_{k-1}}^{x_k} \left(\int_{x_{k-1}}^t f \right)^q u(t) dt \right)^{\frac{r}{q}} \right)^{\frac{1}{r}} \leq \mathcal{C}' \left(\sum_{k=N+1}^{\infty} \int_{x_{k-1}}^{x_k} f^p v \right)^{\frac{1}{p}} \quad (3.2)$$

and

$$\left(\sum_{k=N+1}^{\infty} 2^{-k} \left(\int_a^{x_k} f \right)^r \left(\int_{x_k}^{x_{k+1}} u \right)^{\frac{r}{q}} \right)^{\frac{1}{r}} \leq \mathcal{C}'' \left(\sum_{k=N+1}^{\infty} \int_{x_{k-1}}^{x_k} f^p v \right)^{\frac{1}{p}} \quad (3.3)$$

hold for all $f \in \mathfrak{M}^+(a, b)$. Moreover, $C \approx \mathcal{C}' + \mathcal{C}''$.

Proof. Applying (2.9) with $\alpha = 1$ and $h(x) = \left(\int_a^x \left(\int_a^t f \right)^q u(t) dt \right)^{\frac{r}{q}}$, we have that

$$LHS(1.5) \approx \left(\sum_{k=N+1}^{\infty} 2^{-k} \left(\int_a^{x_k} \left(\int_a^t f \right)^q u(t) dt \right)^{\frac{r}{q}} \right)^{\frac{1}{r}}.$$

Since, $\{2^{-k}\}$ is geometrically decreasing, using (2.5), we obtain that

$$\begin{aligned} LHS(1.5) &\approx \left(\sum_{k=N+1}^{\infty} 2^{-k} \left(\int_{x_{k-1}}^{x_k} \left(\int_a^t f \right)^q u(t) dt \right)^{\frac{r}{q}} \right)^{\frac{1}{r}} \\ &\approx \left(\sum_{k=N+1}^{\infty} 2^{-k} \left(\int_{x_{k-1}}^{x_k} \left(\int_{x_{k-1}}^t f \right)^q u(t) dt \right)^{\frac{r}{q}} \right)^{\frac{1}{r}} \\ &\quad + \left(\sum_{k=N+2}^{\infty} 2^{-k} \left(\int_a^{x_{k-1}} f \right)^r \left(\int_{x_{k-1}}^{x_k} u \right)^{\frac{r}{q}} \right)^{\frac{1}{r}}. \end{aligned}$$

Then, it is clear that there exists a positive constant C such that inequality (1.5) holds for all $f \in \mathfrak{M}^+(a, b)$ if and only if there exist positive constants \mathcal{C}' and \mathcal{C}'' such that (3.2) and (3.3) hold for all $f \in \mathfrak{M}^+(a, b)$. Moreover, $C \approx \mathcal{C}' + \mathcal{C}''$. \square

Lemma 3.2. *Let $1 \leq p < \infty$, $0 < q, r < \infty$ and let u, v be weights on (a, b) . Assume that $N \in \mathbb{Z} \cup \{-\infty\}$ and $\{x_k\}_{k=N}^{\infty}$ is a strictly increasing sequence in $[a, b]$. Then there exists a positive constant \mathcal{C}' such that inequality (3.2) holds for all $f \in \mathfrak{M}^+(a, b)$ if and only if there exists a positive constant C' such that*

$$\left(\sum_{k=N+1}^{\infty} 2^{-k} a_k^r B(x_{k-1}, x_k)^r \right)^{\frac{1}{r}} \leq C' \left(\sum_{k=N+1}^{\infty} a_k^p \right)^{\frac{1}{p}} \quad (3.4)$$

holds for every sequence of non-negative numbers $\{a_k\}_{k=N+1}^{\infty}$. Moreover the best constants \mathcal{C}' and C' , respectively, in (3.2) and (3.4) satisfy $\mathcal{C}' \approx C'$.

Proof. Assume that (3.2) holds. By the definition of $B(x_{k-1}, x_k)$, there exist non-negative measurable functions h_k , $N+1 \leq k$ on (a, b) such that

$$\text{supp } h_k \subset [x_{k-1}, x_k], \quad \int_{x_{k-1}}^{x_k} h_k^p v = 1, \quad \left(\int_{x_{k-1}}^{x_k} \left(\int_{x_{k-1}}^t h_k \right)^q u(t) dt \right)^{\frac{1}{q}} \gtrsim B(x_{k-1}, x_k).$$

Thus, inserting $f = \sum_{m=N+1}^{\infty} a_m h_m$, where $\{a_m\}_{m=N+1}^{\infty}$ is any sequence of non-negative numbers, into (3.2), (3.4) follows. Moreover, $C' \lesssim \mathcal{C}'$.

Conversely, observe first that for each $h \in \mathfrak{M}^+(x_{k-1}, x_k)$,

$$B(x_{k-1}, x_k) \geq \left(\int_{x_{k-1}}^{x_k} \left(\int_{x_{k-1}}^t h(s) ds \right)^q u(t) dt \right)^{\frac{1}{q}} \left(\int_{x_{k-1}}^{x_k} h(t)^p v(t) dt \right)^{-\frac{1}{p}}.$$

Then (3.2) follows by, inserting $a_k = \left(\int_{x_{k-1}}^{x_k} h^p v \right)^{\frac{1}{p}}$ in (3.4) and $\mathcal{C}' \leq C'$. Further, we have $\mathcal{C}' \approx C'$. \square

Lemma 3.3. *Let $1 \leq p < \infty$, $0 < q, r < \infty$ and let u, v be weights on (a, b) . Assume that $N \in \mathbb{Z} \cup \{-\infty\}$ and $\{x_k\}_{k=N}^{\infty}$ is a strictly increasing sequence in $[a, b]$. Then there exists a positive constant \mathcal{C}'' such that inequality (3.3) holds for all $f \in \mathfrak{M}^+(a, b)$ if and only if there exists a positive constant C'' such that*

$$\left(\sum_{k=N+1}^{\infty} 2^{-k} \left(\int_{x_k}^{x_{k+1}} u \right)^{\frac{r}{q}} \left(\sum_{j=N+1}^k a_j V_p(x_{j-1}, x_j) \right)^r \right)^{\frac{1}{r}} \leq C'' \left(\sum_{k=N+1}^{\infty} a_k^p \right)^{\frac{1}{p}} \quad (3.5)$$

holds for every sequence of non-negative numbers $\{a_k\}_{k=N+1}^{\infty}$. Moreover the best constants \mathcal{C}'' and C'' , respectively, in (3.3) and (3.5) satisfy $\mathcal{C}'' \approx C''$.

Proof. Suppose that (3.3) holds for all $f \in \mathfrak{M}^+(a, b)$. Using (3.5), there exist non-negative measurable functions g_k , $N+1 \leq k$ on (x_{k-1}, x_k) such that

$$\text{supp } g_k \subset [x_{k-1}, x_k], \quad \int_{x_{k-1}}^{x_k} g_k^p v = 1, \quad \int_{x_{k-1}}^{x_k} g_k \gtrsim V_p(x_{k-1}, x_k).$$

Thus, inserting $f = \sum_{m=N+1}^{\infty} a_m g_m$, where $\{a_m\}_{m=N+1}^{\infty}$ is any sequence of non-negative numbers, into (3.3), (3.5) follows. Moreover, $C'' \lesssim \mathcal{C}''$ holds.

Conversely, taking $a_k = \left(\int_{x_{k-1}}^{x_k} f^p v \right)^{\frac{1}{p}}$ in (3.5) and using the estimate

$$V_p(x_{k-1}, x_k) \geq \int_{x_{k-1}}^{x_k} f(t) dt \left(\int_{x_{k-1}}^{x_k} f(t)^p v(t) dt \right)^{-\frac{1}{p}}$$

gives (3.3). Additionally, $\mathcal{C}'' \leq C''$ holds. Consequently $\mathcal{C}'' \approx C''$ follows. \square

Theorem 3.4. *Let $1 \leq p < \infty$, $0 < q, r < \infty$, $-\infty < \beta < 1$ and let u, v, w be weights on (a, b) such that $0 < \mathcal{W}(t) < \infty$ for all $t \in (a, b)$, and $\{x_k\}_{k=N+1}^{\infty}$ be the discretizing sequence of \mathcal{W} . Then inequality (3.4) holds for every sequence of non-negative numbers $\{a_k\}_{k=N+1}^{\infty}$ if and only if*

(i) $p \leq r$, $p \leq q$ and

$$\mathcal{A}_1 := \sup_{N+1 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \operatorname{ess\,sup}_{t \in (x_{k-1}, x_k)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(x_{k-1}, t) < \infty,$$

(ii) $r < p \leq q$ and

$$\mathcal{A}_2 := \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)\frac{p}{p-r}} \operatorname{ess\,sup}_{t \in (x_{k-1}, x_k)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(x_{k-1}, t)^{\frac{p-r}{p-r}} \right)^{\frac{p-r}{pr}} < \infty,$$

(iii) $q < p \leq r$ and

$$\mathcal{A}_3 := \sup_{N+1 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \left(\int_{x_{k-1}}^{x_k} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(x_{k-1}, t)^{\frac{pq}{p-q}} dt \right)^{\frac{p-q}{pq}} < \infty,$$

(iv) $r < p$, $q < p$ and

$$\mathcal{A}_4 := \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)\frac{p}{p-r}} \left(\int_{x_{k-1}}^{x_k} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(x_{k-1}, t)^{\frac{pq}{p-q}} dt \right)^{\frac{r(p-q)}{q(p-r)}} \right)^{\frac{p-r}{pr}} < \infty.$$

Moreover, the best constant C' in inequality (3.4) satisfies

$$C' = \begin{cases} \mathcal{A}_1 & \text{in the case (i),} \\ \mathcal{A}_2 & \text{in the case (ii),} \\ \mathcal{A}_3 & \text{in the case (iii),} \\ \mathcal{A}_4 & \text{in the case (iv).} \end{cases}$$

Proof. The result follows easily by combining Theorem 2.8 for suitable parameters and weights with (3.1) and the fact that $\mathcal{W}(t) \approx 2^{-k}$, for every $t \in [x_{k-1}, x_k]$. \square

Theorem 3.5. *Let $1 \leq p < \infty$, $0 < q, r < \infty$, $-\infty < \beta < 1$ and let u, v, w be weights on (a, b) such that $0 < \mathcal{W}(t) < \infty$ for all $t \in (a, b)$, and $\{x_k\}_{k=N+1}^{\infty}$ be the discretizing sequence of \mathcal{W} . Then inequality (3.5) holds for every sequence of non-negative numbers $\{a_k\}_{k=N+1}^{\infty}$ if and only if*

(i) $p \leq r$ and

$$\mathcal{B}_1 := \sup_{N+1 \leq k < \infty} \left(\sum_{i=k}^{\infty} 2^{-i(1-\beta)} \left(\int_{x_i}^{x_{i+1}} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \right)^{\frac{1}{r}} V_p(a, x_k) < \infty,$$

(ii) $r < p$ and

$$\begin{aligned} \mathcal{B}_2 := & \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)} \left(\int_{x_k}^{x_{k+1}} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \right. \\ & \left. \times \left(\sum_{i=k}^{\infty} 2^{-i(1-\beta)} \left(\int_{x_i}^{x_{i+1}} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \right)^{\frac{r}{p-r}} V_p(a, x_k)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} < \infty. \end{aligned}$$

Moreover, the best constant C''' in inequality (3.5) satisfies

$$C''' = \begin{cases} \mathcal{B}_1 & \text{in the case (i),} \\ \mathcal{B}_2 & \text{in the case (ii).} \end{cases}$$

Proof. We will apply Theorem 2.9 with suitable parameters and weights $a_k = 2^{-k} \left(\int_{x_k}^{x_{k+1}} u \right)^{\frac{r}{q}}$ and $b_k = V_p(x_{k-1}, x_k)$. We need to treat the cases when $p > 1$ and $p = 1$, separately. Note that when $p > 1$, since $\{x_k\}_{k=N+1}^{\infty}$ is the discretizing sequence of \mathcal{W} , we have for each $k : N+1 \leq k$ that

$$\sum_{i=N+1}^{\infty} V_p(x_{i-1}, x_i)^{\frac{p}{p-1}} = \sum_{i=N+1}^{\infty} \int_{x_{i-1}}^{x_i} v(s)^{-\frac{1}{p-1}} ds = V_p(a, x_k)^{\frac{p-1}{p}}.$$

Thus, applying [Theorem 2.9, (iv)] when $1 < p \leq r$ and [Theorem 2.9, (iii)] when $1 < p, r < p$, the result follows.

On the other hand, if $p = 1$, for each $k : N+1 \leq k$,

$$\sup_{N+1 \leq i \leq k} V_1(x_{i-1}, x_i) = \sup_{N+1 \leq i \leq k} \operatorname{ess\,sup}_{s \in (x_{i-1}, x_i)} v(s)^{-1} = V_1(a, x_k).$$

Therefore, applying [Theorem 2.9, (ii)] when $r < p = 1$, we have $C'' \approx \mathcal{B}_2$.

Lastly, if $p = 1 \leq r$, then applying [Theorem 2.9, (i)], and the fact that $\mathcal{W}(t) \approx 2^{-i}$, for every $t \in [x_i, x_{i+1}]$ we have

$$\begin{aligned} C'' &\approx \sup_{N+1 \leq k < \infty} \left(\sum_{i=k}^{\infty} 2^{-i} \left(\int_{x_i}^{x_{i+1}} u \right)^{\frac{r}{q}} \right)^{\frac{1}{r}} V_1(x_{k-1}, x_k) \\ &\approx \sup_{N+1 \leq k < \infty} \left(\sum_{i=k}^{\infty} 2^{-i(1-\beta)} \left(\int_{x_i}^{x_{i+1}} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \right)^{\frac{1}{r}} V_1(x_{k-1}, x_k). \end{aligned}$$

Finally, interchanging supremum yields that

$$\begin{aligned} C'' &\approx \sup_{N+1 \leq k < \infty} V_1(x_{k-1}, x_k) \sup_{k \leq m < \infty} \left(\sum_{i=m}^{\infty} 2^{-i(1-\beta)} \left(\int_{x_i}^{x_{i+1}} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \right)^{\frac{1}{r}} \\ &= \sup_{N+1 \leq m < \infty} \left(\sum_{i=m}^{\infty} 2^{-i(1-\beta)} \left(\int_{x_i}^{x_{i+1}} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \right)^{\frac{1}{r}} \sup_{N+1 \leq k \leq m} V_1(x_{k-1}, x_k) = \mathcal{B}_1. \end{aligned}$$

□

We can formulate the discrete characterization of inequality (1.5).

Theorem 3.6. *Let $1 \leq p < \infty$, $0 < q, r < \infty$ and let u, v, w be weights on (a, b) such that $0 < \mathcal{W}(t) < \infty$ for all $t \in (a, b)$, and $\{x_k\}_{k=N+1}^{\infty}$ be the discretizing sequence of \mathcal{W} . Then inequality (1.5) holds for all $f \in \mathfrak{M}^+(a, b)$ if and only if*

- (i) $p \leq r$, $p \leq q$ and $\mathcal{A}_1 + \mathcal{B}_1 < \infty$,
- (ii) $r < p \leq q$ and $\mathcal{A}_2 + \mathcal{B}_2 < \infty$,
- (iii) $q < p \leq r$ and $\mathcal{A}_3 + \mathcal{B}_1 < \infty$,
- (iv) $r < p$, $q < p$ and $\mathcal{A}_4 + \mathcal{B}_2 < \infty$.

Moreover the best constant C in (1.5) satisfies

$$C = \begin{cases} \mathcal{A}_1 + \mathcal{B}_1 & \text{in the case (i),} \\ \mathcal{A}_2 + \mathcal{B}_2 & \text{in the case (ii),} \\ \mathcal{A}_3 + \mathcal{B}_1 & \text{in the case (iii),} \\ \mathcal{A}_4 + \mathcal{B}_2 & \text{in the case (iv).} \end{cases}$$

Proof. According to Lemmas [3.1-3.3], inequality (1.5) holds if and only if inequalities (3.4) and (3.5) hold. Moreover, the best constant C in (1.5) satisfies $C \approx C' + C''$, where C' and C'' are the best constants in inequalities (3.4) and (3.5), respectively. The result follows from the combination of Theorem 3.4 and Theorem 3.5. \square

4 Proofs

Proof of Theorem 1.1

(i) Let $1 \leq p \leq \min\{r, q\}$. We have from [Theorem 3.6, (i)] that $C \approx \mathcal{A}_1 + \mathcal{B}_1$. We will prove that $C_1 \approx \mathcal{A}_1 + \mathcal{B}_1$. First, we will show that $\mathcal{A}_1 + \mathcal{B}_1 \approx A_1 + \mathcal{B}_1$, where

$$A_1 := \sup_{N+1 \leq k} 2^{-\frac{k(1-\beta)}{r}} \operatorname{ess\,sup}_{t \in (a, x_k)} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t).$$

It is clear that $\mathcal{A}_1 \leq A_1$. On the other hand, observe that

$$\begin{aligned} A_1 &= \sup_{N+1 \leq k} 2^{-\frac{k(1-\beta)}{r}} \sup_{N+1 \leq i \leq k} \operatorname{ess\,sup}_{t \in (x_{i-1}, x_i)} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t) \\ &\approx \sup_{N+1 \leq k} 2^{-\frac{k(1-\beta)}{r}} \sup_{N+1 \leq i \leq k} \operatorname{ess\,sup}_{t \in (x_{i-1}, x_i)} \left(\int_t^{x_i} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t) \\ &\quad + \sup_{N+2 \leq k} 2^{-\frac{k(1-\beta)}{r}} \sup_{N+1 \leq i < k} \left(\int_{x_i}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, x_i). \end{aligned}$$

Then, interchanging the supremum in the first term and applying (2.7) with $n = N + 2$, for the second term, we have that

$$\begin{aligned} A_1 &\approx \sup_{N+1 \leq k} 2^{-\frac{k(1-\beta)}{r}} \operatorname{ess\,sup}_{t \in (x_{k-1}, x_k)} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t) \\ &\quad + \sup_{N+2 \leq k} 2^{-\frac{k(1-\beta)}{r}} \left(\int_{x_{k-1}}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, x_{k-1}). \end{aligned}$$

Note that, for any $k \geq N + 2$, we have

$$V_p(a, t) \approx V_p(a, x_{k-1}) + V_p(x_{k-1}, t), \quad \text{for every } t \in (x_{k-1}, x_k). \quad (4.1)$$

Then, in view of (4.1),

$$\begin{aligned} A_1 &\approx \sup_{N+1 \leq k} 2^{-\frac{k(1-\beta)}{r}} \operatorname{ess\,sup}_{t \in (x_{k-1}, x_k)} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(x_{k-1}, t) \\ &\quad + \sup_{N+2 \leq k} 2^{-\frac{k(1-\beta)}{r}} \left(\int_{x_{k-1}}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, x_{k-1}) \\ &\lesssim \mathcal{A}_1 + \mathcal{B}_1. \end{aligned}$$

Then we have that $\mathcal{A}_1 + \mathcal{B}_1 \leq A_1 + \mathcal{B}_1 \lesssim \mathcal{A}_1 + \mathcal{B}_1$.

It remains to show that $A_1 + \mathcal{B}_1 \approx C_1$. Applying (2.10) with $\alpha = \frac{1-\beta}{r}$,

$$A_1 \approx \operatorname{ess\,sup}_{x \in (a,b)} \left(\int_x^b w \right)^{\frac{1-\beta}{r}} \operatorname{ess\,sup}_{t \in (a,x)} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t)$$

holds, and interchanging supremum gives that

$$\begin{aligned} A_1 &\approx \operatorname{ess\,sup}_{t \in (a,b)} V_p(a, t) \operatorname{ess\,sup}_{x \in (t,b)} \left(\int_x^b w \right)^{\frac{1-\beta}{r}} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} \\ &\leq \operatorname{ess\,sup}_{t \in (a,b)} V_p(a, t) \left(\int_t^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_t^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{1}{r}} = C_1. \end{aligned} \quad (4.2)$$

On the other hand, applying (2.9) with $\alpha = 1 - \beta$, then using (2.5) with $n = k$, we obtain for any $k \geq N$ that

$$\begin{aligned} \int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds &\approx \sum_{i=k+1}^{\infty} 2^{-i(1-\beta)} \left(\int_{x_k}^{x_i} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \\ &\approx \sum_{i=k}^{\infty} 2^{-i(1-\beta)} \left(\int_{x_i}^{x_{i+1}} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}}. \end{aligned} \quad (4.3)$$

Therefore, in view of (4.3),

$$\begin{aligned} \mathcal{B}_1 &\approx \sup_{N+1 \leq k} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{1}{r}} V_p(a, x_k) \\ &\leq \sup_{N+1 \leq k} \operatorname{ess\,sup}_{t \in (x_{k-1}, x_k)} \left(\int_t^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_t^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{1}{r}} V_p(a, t) = C_1. \end{aligned} \quad (4.4)$$

Thus, combining (4.2) with (4.4), we have that $A_1 + \mathcal{B}_1 \lesssim C_1$.

Conversely, using (4.1), we have

$$\begin{aligned} C_1 &\approx \sup_{N+1 \leq k} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{1}{r}} V_p(x_{k-1}, x_k) \\ &\quad + \sup_{N+1 \leq k} 2^{-\frac{k(1-\beta)}{r}} \operatorname{ess\,sup}_{t \in (x_{k-1}, x_k)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(x_{k-1}, t) \\ &\quad + \sup_{N+1 \leq k} \operatorname{ess\,sup}_{t \in (x_{k-1}, x_k)} \left(\int_t^{x_k} \mathcal{W}(s)^{-\beta} w(s) \left(\int_t^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{1}{r}} V_p(x_{k-1}, t) \\ &\quad + \sup_{N+2 \leq k} \left(\int_{x_{k-1}}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_{k-1}}^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{1}{r}} V_p(a, x_{k-1}). \end{aligned}$$

Then, using (4.3), we arrive at

$$C_1 \lesssim \sup_{N+1 \leq k} 2^{-\frac{k(1-\beta)}{r}} \operatorname{ess\,sup}_{t \in (x_{k-1}, x_k)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t)$$

$$\begin{aligned}
& + \sup_{N+1 \leq k} \left(\sum_{i=k}^{\infty} 2^{-i(1-\beta)} \left(\int_{x_i}^{x_{i+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \right)^{\frac{1}{r}} V_p(a, x_k) \\
& \leq A_1 + \mathcal{B}_1.
\end{aligned} \tag{4.5}$$

As a result, we arrive at the conclusion that the best constant C in (1.5) satisfies $C \approx C_1$.

(ii) Let $r < p \leq q$. Then, we have from [Theorem 3.6, (ii)] that the best constant in (1.5) satisfies $C \approx \mathcal{A}_2 + \mathcal{B}_2$. We will start by showing that $\mathcal{A}_2 + \mathcal{B}_2 \approx A_2 + B_2$, where

$$A_2 := \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta) \frac{p}{p-r}} \operatorname{ess\,sup}_{t \in (a, x_k)} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, t)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}}$$

and

$$\begin{aligned}
B_2 & := \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)} \left(\int_{x_k}^{x_{k+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \right. \\
& \quad \times \left. \left(\sum_{i=k+2}^{\infty} 2^{-i(1-\beta)} \left(\int_{x_i}^{x_{i+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \right)^{\frac{r-r}{p-r}} V_p(a, x_k)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}}.
\end{aligned} \tag{4.6}$$

We have $\mathcal{A}_2 \leq A_2$ by the definitions of \mathcal{A}_2 and A_2 . On the other hand,

$$\begin{aligned}
\mathcal{B}_2 & \approx B_2 + \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta) \frac{p}{p-r}} \left(\int_{x_k}^{x_{k+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, x_k)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\
& \quad + \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta) \frac{p}{p-r}} \left(\int_{x_k}^{x_{k+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \left(\int_{x_{k+1}}^{x_{k+2}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r^2}{q(p-r)}} V_p(a, x_k)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\
& \lesssim B_2 + \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta) \frac{p}{p-r}} \left(\int_{x_k}^{x_{k+2}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, x_k)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\
& \lesssim B_2 + \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta) \frac{p}{p-r}} \operatorname{ess\,sup}_{t \in (a, x_k)} \left(\int_t^{x_{k+2}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, t)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\
& \lesssim B_2 + A_2
\end{aligned} \tag{4.7}$$

holds, hence $\mathcal{A}_2 + \mathcal{B}_2 \lesssim A_2 + B_2$.

Next, we will show that $A_2 \lesssim \mathcal{A}_2 + \mathcal{B}_2$. Observe that,

$$\begin{aligned}
A_2 & = \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta) \frac{p}{p-r}} \sup_{N+1 \leq i \leq k} \operatorname{ess\,sup}_{t \in (x_{i-1}, x_i)} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, t)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\
& \approx \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta) \frac{p}{p-r}} \sup_{N+1 \leq i \leq k} \operatorname{ess\,sup}_{t \in (x_{i-1}, x_i)} \left(\int_t^{x_i} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, t)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\
& \quad + \left(\sum_{k=N+2}^{\infty} 2^{-k(1-\beta) \frac{p}{p-r}} \sup_{N+1 \leq i < k} \left(\int_{x_i}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, x_i)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}}.
\end{aligned}$$

Applying (2.3) for the first term and (2.8) for the second term, we obtain that

$$\begin{aligned} A_2 &\approx \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)\frac{p}{p-r}} \operatorname{ess\,sup}_{t \in (x_{k-1}, x_k)} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, t)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\ &\quad + \left(\sum_{k=N+2}^{\infty} 2^{-k(1-\beta)\frac{p}{p-r}} \left(\int_{x_{k-1}}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, x_{k-1})^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}}. \end{aligned}$$

Using (4.1) we arrive at

$$A_2 \lesssim \mathcal{A}_2 + \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)\frac{p}{p-r}} \left(\int_{x_k}^{x_{k+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, x_k)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \lesssim \mathcal{A}_2 + \mathcal{B}_2.$$

Furthermore, it is clear from the definitions of \mathcal{B}_2 and B_2 that $B_2 \leq \mathcal{B}_2$. Then, we have $A_2 + B_2 \lesssim \mathcal{A}_2 + \mathcal{B}_2$, as well. Consequently, $C \approx A_2 + B_2$ holds.

Next, we will prove that $A_2 + B_2 \approx C_2 + C_3$. First of all, applying (2.9) with $\alpha = \frac{p(1-\beta)}{p-r}$ and

$$h(x) = \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^x \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, t)^{\frac{pr}{p-r}}, \quad x \in (a, b),$$

it is clear that

$$A_2 \approx \left(\int_a^b \mathcal{W}(x)^{\frac{p(1-\beta)}{p-r}-1} w(x) \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^x \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, t)^{\frac{pr}{p-r}} dx \right)^{\frac{p-r}{pr}} = C_2. \quad (4.8)$$

On the other hand, using (4.3),

$$\begin{aligned} B_2 &\approx \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)} \left(\int_{x_{k+2}}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_{k+2}}^s \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \right. \\ &\quad \times \left. \left(\int_{x_k}^{x_{k+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, x_k)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\ &\leq \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)} \left(\int_{x_{k+2}}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_{k+2}}^s \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \right. \\ &\quad \times \left. \operatorname{ess\,sup}_{t \in (a, x_{k+1})} \left(\int_t^{x_{k+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\ &\approx \left(\sum_{k=N+1}^{\infty} \int_{x_{k+1}}^{x_{k+2}} \mathcal{W}(x)^{-\beta} w(x) dx \left(\int_{x_{k+2}}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_{k+2}}^s \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \right. \\ &\quad \times \left. \operatorname{ess\,sup}_{t \in (a, x_{k+1})} \left(\int_t^{x_{k+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\ &\leq \left(\sum_{k=N+1}^{\infty} \int_{x_{k+1}}^{x_{k+2}} \mathcal{W}(x)^{-\beta} w(x) \left(\int_x^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_x^s \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \right. \\ &\quad \times \left. \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^x \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} dx \right)^{\frac{p-r}{pr}} \end{aligned}$$

$$\leq C_3. \tag{4.9}$$

Combination of (4.8) and (4.9) yield that $A_2 + B_2 \lesssim C_2 + C_3$.

Conversely,

$$\begin{aligned} C_3 &= \left(\sum_{k=N+1}^{\infty} \int_{x_{k-1}}^{x_k} \left(\int_x^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_x^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \mathcal{W}(x)^{-\beta} w(x) \right. \\ &\quad \times \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} dx \Big)^{\frac{p-r}{pr}} \\ &\approx \left(\sum_{k=N+1}^{\infty} \int_{x_{k-1}}^{x_k} \left(\int_x^{x_k} \mathcal{W}(s)^{-\beta} w(s) \left(\int_x^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \mathcal{W}(x)^{-\beta} w(x) \right. \\ &\quad \times \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} dx \Big)^{\frac{p-r}{pr}} \\ &+ \left(\sum_{k=N+1}^{\infty} \int_{x_{k-1}}^{x_k} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_x^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \mathcal{W}(x)^{-\beta} w(x) \right. \\ &\quad \times \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} dx \Big)^{\frac{p-r}{pr}} \\ &\approx \left(\sum_{k=N+1}^{\infty} \int_{x_{k-1}}^{x_k} \left(\int_x^{x_k} \mathcal{W}(s)^{-\beta} w(s) \left(\int_x^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \mathcal{W}(x)^{-\beta} w(x) \right. \\ &\quad \times \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} dx \Big)^{\frac{p-r}{pr}} \\ &+ \left(\sum_{k=N+1}^{\infty} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) ds \right)^{\frac{r}{p-r}} \int_{x_{k-1}}^{x_k} \left(\int_x^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r^2}{q(p-r)}} \mathcal{W}(x)^{-\beta} w(x) \right. \\ &\quad \times \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} dx \Big)^{\frac{p-r}{pr}} \\ &+ \left(\sum_{k=N+1}^{\infty} \int_{x_{k-1}}^{x_k} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \mathcal{W}(x)^{-\beta} w(x) \right. \\ &\quad \times \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} dx \Big)^{\frac{p-r}{pr}}. \end{aligned}$$

Since

$$\left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) ds \right)^{\frac{r}{p-r}} \approx 2^{-k(1-\beta)\frac{r}{p-r}},$$

$$\begin{aligned} C_3 &\approx \left(\sum_{k=N+1}^{\infty} \int_{x_{k-1}}^{x_k} \left(\int_x^{x_k} \mathcal{W}(s)^{-\beta} w(s) \left(\int_x^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \mathcal{W}(x)^{-\beta} w(x) \right. \\ &\quad \times \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} dx \Big)^{\frac{p-r}{pr}} \end{aligned}$$

$$\begin{aligned}
& + \left(\sum_{k=N+1}^{\infty} 2^{-k \frac{(1-\beta)r}{p-r}} \int_{x_{k-1}}^{x_k} \left(\int_x^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r^2}{q(p-r)}} \mathcal{W}(x)^{-\beta} w(x) \right. \\
& \quad \times \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} dx \Big)^{\frac{p-r}{pr}} \\
& + \left(\sum_{k=N+1}^{\infty} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \right. \\
& \quad \times \int_{x_{k-1}}^{x_k} \mathcal{W}(x)^{-\beta} w(x) \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} dx \Big)^{\frac{p-r}{pr}} \\
& =: C_{3,1} + C_{3,2} + C_{3,3}.
\end{aligned}$$

Note that

$$\int_{x_{k-1}}^{x_k} \left(\int_x^{x_k} \mathcal{W}^{-\beta} w \right)^{\frac{r}{p-r}} \mathcal{W}(x)^{-\beta} w(x) dx \approx \left(\int_{x_{k-1}}^{x_k} \mathcal{W}^{-\beta} w \right)^{\frac{p}{p-r}} \approx 2^{-k(1-\beta) \frac{p}{p-r}}. \quad (4.10)$$

In view of (4.10), we get

$$\begin{aligned}
C_{3,1} & \leq \left(\sum_{k=N+1}^{\infty} \int_{x_{k-1}}^{x_k} \left(\int_x^{x_k} \mathcal{W}^{-\beta} w \right)^{\frac{r}{p-r}} \mathcal{W}(x)^{-\beta} w(x) \right. \\
& \quad \times \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, t)^{\frac{pr}{p-r}} dx \Big)^{\frac{p-r}{pr}} \\
& \leq \left(\sum_{k=N+1}^{\infty} \int_{x_{k-1}}^{x_k} \left(\int_x^{x_k} \mathcal{W}^{-\beta} w \right)^{\frac{r}{p-r}} \mathcal{W}(x)^{-\beta} w(x) dx \right. \\
& \quad \times \operatorname{ess\,sup}_{t \in (a, x_k)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, t)^{\frac{pr}{p-r}} \Big)^{\frac{p-r}{pr}} \\
& \approx A_2
\end{aligned}$$

and

$$\begin{aligned}
C_{3,2} & \leq \left(\sum_{k=N+1}^{\infty} 2^{-k \frac{(1-\beta)r}{p-r}} \int_{x_{k-1}}^{x_k} \mathcal{W}(x)^{-\beta} w(x) \right. \\
& \quad \times \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r^2}{q(p-r)}} \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} dx \Big)^{\frac{p-r}{pr}} \\
& \leq \left(\sum_{k=N+1}^{\infty} 2^{-k \frac{(1-\beta)r}{p-r}} \int_{x_{k-1}}^{x_k} \mathcal{W}(x)^{-\beta} w(x) \operatorname{ess\,sup}_{t \in (a, x)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, t)^{\frac{pr}{p-r}} dx \right)^{\frac{p-r}{pr}} \\
& \leq \left(\sum_{k=N+1}^{\infty} 2^{-k \frac{(1-\beta)r}{p-r}} \int_{x_{k-1}}^{x_k} \mathcal{W}(x)^{-\beta} w(x) dx \operatorname{ess\,sup}_{t \in (a, x_k)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, t)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\
& \approx \left(\sum_{k=N+1}^{\infty} 2^{-k \frac{(1-\beta)p}{p-r}} \operatorname{ess\,sup}_{t \in (a, x_k)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, t)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}}
\end{aligned}$$

$$= A_2.$$

Furthermore, using (4.10) once more,

$$\begin{aligned}
C_{3,3} &\lesssim \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \right. \\
&\quad \times \operatorname{ess\,sup}_{t \in (a, x_k)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} \Big)^{\frac{p-r}{pr}} \\
&= \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \right. \\
&\quad \times \sup_{N+1 \leq i \leq k} \operatorname{ess\,sup}_{t \in (x_{i-1}, x_i)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} \Big)^{\frac{p-r}{pr}} \\
&\approx \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \right. \\
&\quad \times \sup_{N+1 \leq i \leq k} \operatorname{ess\,sup}_{t \in (x_{i-1}, x_i)} \left(\int_t^{x_i} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} \Big)^{\frac{p-r}{pr}} \\
&+ \left(\sum_{k=N+2}^{\infty} 2^{-k(1-\beta)} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \right. \\
&\quad \times \sup_{N+1 \leq i < k} \left(\int_{x_i}^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, x_i)^{\frac{pr}{p-r}} \Big)^{\frac{p-r}{pr}} \\
&=: I + II.
\end{aligned}$$

Since, the sequence $\{a_k\}_{k=N+1}^{\infty}$, with

$$a_k =: 2^{-k(1-\beta)} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}}$$

is geometrically decreasing, (2.3) yields that

$$\begin{aligned}
I &\approx \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \right. \\
&\quad \times \operatorname{ess\,sup}_{t \in (x_{k-1}, x_k)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} \Big)^{\frac{p-r}{pr}}.
\end{aligned}$$

Let $y_k \in [x_{k-1}, x_k]$, $N+1 \leq k$ be such that

$$\operatorname{ess\,sup}_{t \in (x_{k-1}, x_k)} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, t)^{\frac{pr}{p-r}} \lesssim \left(\int_{y_k}^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, y_k)^{\frac{pr}{p-r}}.$$

Then, we have

$$I \lesssim \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \right)^{\frac{p-r}{pr}}$$

$$\times \left(\int_{y_k}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, y_k)^{\frac{pr}{p-r}} \frac{p-r}{pr}.$$

Thus, (4.3) ensures that

$$\begin{aligned} I &\lesssim \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)} \left(\sum_{i=k}^{\infty} 2^{-i(1-\beta)} \left(\int_{x_i}^{x_{i+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \right)^{\frac{r}{p-r}} \right. \\ &\quad \times \left. \left(\int_{y_k}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, y_k)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\ &\leq \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)} \left(\sum_{i=k}^{\infty} 2^{-i(1-\beta)} \left(\int_{y_i}^{y_{i+2}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \right)^{\frac{r}{p-r}} \right. \\ &\quad \times \left. \left(\int_{y_k}^{y_{k+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, y_k)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}}. \end{aligned}$$

Moreover,

$$2^{-k} \approx \int_{x_k}^b w \leq \int_{y_k}^b w \leq \int_{x_{k-1}}^b w \approx 2^{-(k-1)}, \quad N+1 \leq k.$$

As a result, $\{y_k\}_{k=N+1}^{\infty}$ is a discretizing sequence of \mathcal{W} , as well. This fact together with (4.7) yield $I \lesssim \mathcal{B}_2 \lesssim A_2 + B_2$.

On the other hand, applying (2.8) with

$$\tau_k = 2^{-k(1-\beta)} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}}, \quad \sigma_k = V_p(a, x_k)^{\frac{pr}{p-r}}, \quad \alpha = \frac{r}{q},$$

gives

$$\begin{aligned} II &\approx \left(\sum_{k=N+2}^{\infty} 2^{-k(1-\beta)} \left(\int_{x_k}^b \mathcal{W}(s)^{-\beta} w(s) \left(\int_{x_k}^s \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} ds \right)^{\frac{r}{p-r}} \right. \\ &\quad \times \left. \left(\int_{x_{k-1}}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, x_{k-1})^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}}. \end{aligned}$$

Lastly, using (4.3) and (4.7), we obtain that

$$\begin{aligned} II &\approx \left(\sum_{k=N+2}^{\infty} 2^{-k(1-\beta)} \left(\sum_{i=k}^{\infty} 2^{-i(1-\beta)} \left(\int_{x_i}^{x_{i+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} \right)^{\frac{r}{1-r}} \left(\int_{x_{k-1}}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} V_p(a, x_{k-1})^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\ &\leq A_2 + B_2. \end{aligned}$$

Therefore,

$$C_3 \lesssim A_2 + B_2. \quad (4.11)$$

Finally, combination of (4.8) and (4.11) yield, $C_2 + C_3 \approx A_2 + B_2$. Accordingly, the best constant C in (1.5) satisfies $C \approx C_2 + C_3$.

(iii) Let $q < p \leq r$. According to [Theorem 3.6, (iii)], the best constant in (1.5) satisfies $C \approx \mathcal{A}_3 + \mathcal{B}_1$. We will begin our proof by showing that $\mathcal{A}_3 + \mathcal{B}_1 \approx \mathcal{A}_3 + \mathcal{B}_1$, where

$$\mathcal{A}_3 := \sup_{N+1 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \left(\int_a^{x_k} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(a, t)^{\frac{pq}{p-q}} dt \right)^{\frac{p-q}{pq}}.$$

It is clear that $\mathcal{A}_3 \leq A_3$, the proof of this part is complete if we show that $A_3 \lesssim \mathcal{A}_3 + \mathcal{B}_1$. Assume that $\max\{\mathcal{A}_3, \mathcal{B}_1\} < \infty$. Then,

$$\left(\int_{x_{k-1}}^{x_k} \left(\int_t^{x_k} \mathcal{W} \frac{\beta q}{r} u \right)^{\frac{q}{p-q}} \mathcal{W}(t) \frac{\beta q}{r} u(t) V_p(a, t)^{\frac{pq}{p-q}} dt \right)^{\frac{p-q}{pq}} < \infty, \quad k \geq N+1,$$

holds. Also, for each $t \in [x_{k-1}, x_k]$, $k \geq N+1$, we have

$$\left(\int_t^{x_k} \mathcal{W} \frac{\beta q}{r} u \right)^{\frac{1}{q}} V_p(a, t) \lesssim \left(\int_t^{x_k} \left(\int_s^{x_k} \mathcal{W} \frac{\beta q}{r} u \right)^{\frac{q}{p-q}} \mathcal{W}(s) \frac{\beta q}{r} u(s) V_p(a, s)^{\frac{pq}{p-q}} ds \right)^{\frac{p-q}{pq}}.$$

Therefore, we have

$$\lim_{t \rightarrow x_k^-} \left(\int_t^{x_k} \mathcal{W} \frac{\beta q}{r} u \right)^{\frac{1}{q}} V_p(a, t) = 0. \quad (4.12)$$

In that case, integration by parts gives

$$\begin{aligned} & \left(\int_a^{x_k} \left(\int_t^{x_k} \mathcal{W} \frac{\beta q}{r} u \right)^{\frac{q}{p-q}} \mathcal{W}(t) \frac{\beta q}{r} u(t) V_p(a, t)^{\frac{pq}{p-q}} dt \right)^{\frac{p-q}{pq}} \\ & \approx \left(\int_a^{x_k} \left(\int_t^{x_k} \mathcal{W} \frac{\beta q}{r} u \right)^{\frac{p}{p-q}} d[V_p(a, t)^{\frac{pq}{p-q}}] \right)^{\frac{p-q}{pq}} + \lim_{t \rightarrow a^+} \left(\int_t^{x_k} \mathcal{W} \frac{\beta q}{r} u \right)^{\frac{1}{q}} V_p(a, t) \\ & \approx \left(\sum_{i=N+1}^k \int_{x_{i-1}}^{x_i} \left(\int_t^{x_i} \mathcal{W} \frac{\beta q}{r} u \right)^{\frac{p}{p-q}} d[V_p(a, t)^{\frac{pq}{p-q}}] \right)^{\frac{p-q}{pq}} \\ & \quad + \left(\sum_{i=N+1}^{k-1} \left(\int_{x_i}^{x_k} \mathcal{W} \frac{\beta q}{r} u \right)^{\frac{p}{p-q}} \int_{x_{i-1}}^{x_i} d[V_p(a, t)^{\frac{pq}{p-q}}] \right)^{\frac{p-q}{pq}} \\ & \quad + \lim_{t \rightarrow a^+} \left(\int_t^{x_k} \mathcal{W} \frac{\beta q}{r} u \right)^{\frac{1}{q}} V_p(a, t). \end{aligned}$$

Moreover, Minkowski's inequality with $\frac{p}{p-q} > 1$ yields that

$$\begin{aligned} & \left(\sum_{i=N+1}^{k-1} \left(\int_{x_i}^{x_k} \mathcal{W} \frac{\beta q}{r} u \right)^{\frac{p}{p-q}} \int_{x_{i-1}}^{x_i} d[V_p(a, t)^{\frac{pq}{p-q}}] \right)^{\frac{p-q}{pq}} \\ & = \left(\sum_{i=N+1}^{k-1} \left(\sum_{j=i}^{k-1} \int_{x_j}^{x_{j+1}} \mathcal{W} \frac{\beta q}{r} u \right)^{\frac{p}{p-q}} \int_{x_{i-1}}^{x_i} d[V_p(a, t)^{\frac{pq}{p-q}}] \right)^{\frac{p-q}{pq}} \\ & \leq \left(\sum_{j=N+1}^{k-1} \left(\int_{x_j}^{x_{j+1}} \mathcal{W} \frac{\beta q}{r} u \right) \left(\sum_{i=N+1}^j \int_{x_{i-1}}^{x_i} d[V_p(a, t)^{\frac{pq}{p-q}}] \right)^{\frac{p-q}{p}} \right)^{\frac{1}{q}} \\ & = \left(\sum_{j=N+1}^{k-1} \left(\int_{x_j}^{x_{j+1}} \mathcal{W} \frac{\beta q}{r} u \right) \left(\int_a^{x_j} d[V_p(a, t)^{\frac{pq}{p-q}}] \right)^{\frac{p-q}{p}} \right)^{\frac{1}{q}}. \end{aligned}$$

Then, we arrive at

$$\left(\int_a^{x_k} \left(\int_t^{x_k} \mathcal{W} \frac{\beta q}{r} u \right)^{\frac{q}{p-q}} \mathcal{W}(t) \frac{\beta q}{r} u(t) V_p(a, t)^{\frac{pq}{p-q}} dt \right)^{\frac{p-q}{pq}}$$

$$\begin{aligned}
&\lesssim \left(\sum_{i=N+1}^k \int_{x_{i-1}}^{x_i} \left(\int_t^{x_i} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{p}{p-q}} d[V_p(a, t)^{\frac{pq}{p-q}}] \right)^{\frac{p-q}{pq}} \\
&\quad + \left(\sum_{j=N+1}^{k-1} \left(\int_{x_j}^{x_{j+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right) \left(\int_a^{x_j} d[V_p(a, t)^{\frac{pq}{p-q}}] \right)^{\frac{p-q}{p}} \right)^{\frac{1}{q}} \\
&\quad + \lim_{t \rightarrow a^+} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t). \tag{4.13}
\end{aligned}$$

Now, we are in a position to find the upper estimate for A_3 . Using (4.13), we have that

$$\begin{aligned}
A_3 &\lesssim \sup_{N+1 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \left(\sum_{i=N+1}^k \int_{x_{i-1}}^{x_i} \left(\int_t^{x_i} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{p}{p-q}} d[V_p(a, t)^{\frac{pq}{p-q}}] \right)^{\frac{p-q}{pq}} \\
&\quad + \sup_{N+2 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \left(\sum_{j=N+1}^{k-1} \left(\int_{x_j}^{x_{j+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right) \left(\int_a^{x_j} d[V_p(a, t)^{\frac{pq}{p-q}}] \right)^{\frac{p-q}{p}} \right)^{\frac{1}{q}} \\
&\quad + \sup_{N+1 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \lim_{t \rightarrow a^+} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t).
\end{aligned}$$

Further, (2.1) yields,

$$\begin{aligned}
A_3 &\lesssim \sup_{N+1 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \left(\int_{x_{k-1}}^{x_k} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{p}{p-q}} d[V_p(a, t)^{\frac{pq}{p-q}}] \right)^{\frac{p-q}{pq}} \\
&\quad + \sup_{N+1 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \left(\int_{x_k}^{x_{k+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} \left(\int_a^{x_k} d[V_p(a, t)^{\frac{pq}{p-q}}] \right)^{\frac{p-q}{pq}} \\
&\quad + \sup_{N+1 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \lim_{t \rightarrow a^+} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t) \\
&=: A_{3,1} + A_{3,2} + A_{3,3}.
\end{aligned}$$

Integrating by parts again, we have that

$$A_{3,1} \lesssim \sup_{N+1 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \left(\int_{x_{k-1}}^{x_k} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(a, t)^{\frac{pq}{p-q}} dt \right)^{\frac{p-q}{pq}}.$$

Thus, (4.1) gives,

$$\begin{aligned}
A_{3,1} &\lesssim \sup_{N+1 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \left(\int_{x_{k-1}}^{x_k} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(x_{k-1}, t)^{\frac{pq}{p-q}} dt \right)^{\frac{p-q}{pq}} \\
&\quad + \sup_{N+2 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \left(\int_{x_{k-1}}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, x_{k-1}) \\
&\lesssim \mathcal{A}_3 + \mathcal{B}_1. \tag{4.14}
\end{aligned}$$

Additionally,

$$A_{3,2} \lesssim \sup_{N+1 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \left(\int_{x_k}^{x_{k+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, x_k) \leq \mathcal{B}_1. \tag{4.15}$$

Lastly, we will find a suitable upper estimate for $A_{3,3}$. To this end, we will treat the cases $N = -\infty$ and $N > -\infty$, separately. Observe that, if $N = -\infty$, since $x_i \rightarrow a$ if $i \rightarrow -\infty$, we have for any k ,

$$\begin{aligned} \lim_{t \rightarrow a+} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t) &= \lim_{i \rightarrow -\infty} \left(\int_{x_i}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, x_i) \\ &\leq \sup_{i < k} \left(\int_{x_i}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, x_i). \end{aligned} \quad (4.16)$$

Then, then using (4.16) together with (2.7), we get

$$\begin{aligned} A_{3,3} &\lesssim \sup_{k \in \mathbb{Z}} 2^{-\frac{k(1-\beta)}{r}} \sup_{i < k} \left(\int_{x_i}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, x_i) \\ &\approx \sup_{k \in \mathbb{Z}} 2^{-\frac{k(1-\beta)}{r}} \left(\int_{x_{k-1}}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, x_{k-1}) \lesssim \mathcal{B}_1. \end{aligned}$$

On the other hand, if $N > -\infty$

$$\begin{aligned} \lim_{t \rightarrow a+} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t) &\leq \operatorname{ess\,sup}_{t \in (a, x_{N+1})} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t) \\ &\approx \operatorname{ess\,sup}_{t \in (a, x_{N+1})} \left(\int_t^{x_{N+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t) \\ &\quad + \left(\int_{x_{N+1}}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, x_{N+1}). \end{aligned} \quad (4.17)$$

Additionally, it is easy to see that

$$\operatorname{ess\,sup}_{\tau \in (x, y)} \left(\int_{\tau}^y \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(x, \tau) \leq \left(\int_x^y \left(\int_t^y \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(x, t)^{\frac{pq}{p-q}} dt \right)^{\frac{p-q}{pq}}. \quad (4.18)$$

First, using (4.17), we get

$$\begin{aligned} A_{3,3} &\lesssim \sup_{N+1 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \operatorname{ess\,sup}_{t \in (a, x_{N+1})} \left(\int_t^{x_{N+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t) \\ &\quad + \sup_{N+2 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \sup_{N+1 \leq i < k} \left(\int_{x_i}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, x_i). \end{aligned}$$

Then, using (4.18) for the first term and applying (2.7) for the second term, we get

$$\begin{aligned} A_{3,3} &\lesssim \sup_{N+1 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \left(\int_{x_{k-1}}^{x_k} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(a, t)^{\frac{pq}{p-q}} dt \right)^{\frac{p-q}{pq}} \\ &\quad + \sup_{N+2 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \left(\int_{x_{k-1}}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, x_{k-1}). \end{aligned}$$

Finally, using (4.1), we arrive at

$$A_{3,3} \lesssim \mathcal{A}_3 + \sup_{N+2 \leq k < \infty} 2^{-\frac{k(1-\beta)}{r}} \left(\int_{x_{k-1}}^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, x_{k-1}) \lesssim \mathcal{A}_3 + \mathcal{B}_1.$$

Consequently, we have for any $N \in \mathbb{Z} \cup \{-\infty\}$, $A_{3,3} \lesssim \mathcal{A}_3 + \mathcal{B}_1$. Combining the last estimate with (4.14) and (4.15), we arrive at $A_3 \lesssim \mathcal{A}_3 + \mathcal{B}_1$, hence, $C \approx \mathcal{A}_3 + \mathcal{B}_1$.

Let us now continue the proof by showing $A_3 + \mathcal{B}_1 \approx C_1 + C_4$.

To this end, taking $\alpha = \frac{1-\beta}{r}$ and

$$h(x) = \left(\int_a^x \left(\int_t^x \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(a, t)^{\frac{pq}{p-q}} dt \right)^{\frac{p-q}{pq}}, \quad x \in (a, b)$$

in (2.10), we have that $A_3 \approx C_4$. Moreover, we have already shown in (4.4) and (4.5) that $\mathcal{B}_1 \lesssim C_1 \lesssim A_1 + \mathcal{B}_1$. Furthermore, (4.18) yields that $A_1 \lesssim A_3$. Consequently, we have $A_3 + \mathcal{B}_1 \lesssim C_4 + C_1 \lesssim A_3 + \mathcal{B}_1$, which is the desired estimate.

(iv) Let $\max\{r, q\} < p$. Then, using [Theorem 3.6, (iv)], we have that $C \approx \mathcal{B}_2 + \mathcal{A}_4$. First of all, we will show that $\mathcal{B}_2 + \mathcal{A}_4 \approx B_2 + A_4$, where

$$A_4 := \left(\sum_{k=N+1}^{\infty} 2^{-k \frac{p(1-\beta)}{p-r}} \left(\int_a^{x_k} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(a, t)^{\frac{pq}{p-q}} dt \right)^{\frac{r(p-q)}{q(p-r)}} \right)^{\frac{p-r}{pr}},$$

and B_2 is defined in (4.6).

It is clear that $\mathcal{A}_4 \leq A_4$. We have already shown in (4.7) that $\mathcal{B}_2 \lesssim A_2 + B_2$. Moreover, analogously as in the previous proof, using (4.18), one can easily see that $A_2 \lesssim A_4$. Thus, $\mathcal{B}_2 + \mathcal{A}_4 \lesssim B_2 + A_4$ follows.

It remains to prove that $B_2 + A_4 \lesssim \mathcal{B}_2 + \mathcal{A}_4$. Assume that $\max\{A_4, \mathcal{B}_2\} < \infty$. Then, using the same steps as in the previous case, we can see that (4.12) holds; therefore, (4.13) is true in this case, as well.

Applying (4.13) combined with (2.5), we obtain that

$$\begin{aligned} A_4 &\lesssim \left(\sum_{k=N+1}^{\infty} 2^{-k \frac{p(1-\beta)}{p-r}} \left(\int_{x_{k-1}}^{x_k} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{p}{p-q}} d \left[V_p(a, t)^{\frac{pq}{p-q}} \right] \right)^{\frac{r(p-q)}{q(p-r)}} \right)^{\frac{p-r}{pr}} \\ &\quad + \left(\sum_{k=N+2}^{\infty} 2^{-k \frac{p(1-\beta)}{p-r}} \left(\int_{x_{k-1}}^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} \left(\int_a^{x_{k-1}} d \left[V_p(a, t)^{\frac{pq}{p-q}} \right] \right)^{\frac{r(p-q)}{q(p-r)}} \right)^{\frac{p-r}{pr}} \\ &\quad + \left(\sum_{k=N+1}^{\infty} 2^{-k \frac{p(1-\beta)}{p-r}} \left[\lim_{t \rightarrow a^+} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{1}{q}} V_p(a, t) \right]^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\ &=: A_{4,1} + A_{4,2} + A_{4,3} \end{aligned}$$

holds.

As in the proof of the previous case, using integration by parts in combination with (4.1), we have that

$$A_{4,1} \lesssim \left(\sum_{k=N+1}^{\infty} 2^{-k \frac{p(1-\beta)}{p-r}} \left(\int_{x_{k-1}}^{x_k} \left(\int_t^{x_k} \mathcal{W}^{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(a, t)^{\frac{pq}{p-q}} dt \right)^{\frac{r(p-q)}{q(p-r)}} \right)^{\frac{p-r}{pr}}$$

$$\begin{aligned}
&\approx \mathcal{A}_4 + \left(\sum_{k=N+2}^{\infty} 2^{-k \frac{p(1-\beta)}{p-r}} \left(\int_{x_{k-1}}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, x_{k-1})^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\
&\approx \mathcal{A}_4 + \left(\sum_{k=N+1}^{\infty} 2^{-k(1-\beta)} \left(\int_{x_k}^{x_{k+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r}{q}} 2^{-k \frac{r(1-\beta)}{p-r}} \left(\int_{x_k}^{x_{k+1}} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{r^2}{q(p-r)}} V_p(a, x_k)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\
&\lesssim \mathcal{A}_4 + \mathcal{B}_2.
\end{aligned} \tag{4.19}$$

On the other hand, it is clear that

$$A_{4,2} \lesssim \left(\sum_{k=N+2}^{\infty} 2^{-k \frac{p(1-\beta)}{p-r}} \left(\int_{x_{k-1}}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, x_{k-1})^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \lesssim \mathcal{B}_2. \tag{4.20}$$

Furthermore, if $N = -\infty$, using (4.16), and then applying (2.8), we get

$$\begin{aligned}
A_{4,3} &\lesssim \left(\sum_{k=-\infty}^{\infty} 2^{-k \frac{p(1-\beta)}{p-r}} \sup_{i < k} \left(\int_{x_i}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, x_i)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \\
&\approx \left(\sum_{k=-\infty}^{\infty} 2^{-k \frac{p(1-\beta)}{p-r}} \left(\int_{x_{k-1}}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, x_{k-1})^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \lesssim \mathcal{B}_2.
\end{aligned}$$

If $N > -\infty$, (4.17) together with (4.18) yields,

$$\begin{aligned}
A_{4,3} &\lesssim \left(\sum_{k=N+1}^{\infty} 2^{-k \frac{p(1-\beta)}{p-r}} \left(\int_{x_{k-1}}^{x_k} \left(\int_t^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(a, t)^{\frac{pq}{p-q}} dt \right)^{\frac{r(p-q)}{q(p-r)}} \right)^{\frac{p-r}{pr}} \\
&+ \left(\sum_{k=N+2}^{\infty} 2^{-k \frac{p(1-\beta)}{p-r}} \sup_{N+1 \leq i < k} \left(\int_{x_i}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, x_i)^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}}.
\end{aligned}$$

Applying (4.1) to the first term and (2.8) to the second term, we have

$$A_{4,3} \lesssim \mathcal{A}_4 + \left(\sum_{k=N+2}^{\infty} 2^{-k \frac{p(1-\beta)}{p-r}} \left(\int_{x_{k-1}}^{x_k} \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{pr}{q(p-r)}} V_p(a, x_{k-1})^{\frac{pr}{p-r}} \right)^{\frac{p-r}{pr}} \lesssim \mathcal{A}_4 + \mathcal{B}_2.$$

Thus, for any $N \in \mathbb{Z} \cup \{-\infty\}$, we arrive at $A_{4,3} \lesssim \mathcal{A}_4 + \mathcal{B}_2$. This fact, combined with (4.19) and (4.20) yields $A_4 \lesssim \mathcal{A}_4 + \mathcal{B}_2$. Since $B_2 \leq \mathcal{B}_2$, we have $A_4 + B_2 \lesssim \mathcal{A}_4 + \mathcal{B}_2$ and consequently, $C \approx B_2 + A_4$.

Now, we will show that $B_2 + A_4 \approx C_3 + C_5$. Applying (2.9) with $\alpha = \frac{p(1-\beta)}{p-r}$ and

$$h(x) = \left(\int_a^x \left(\int_t^x \mathcal{W}_{\frac{\beta q}{r}} u \right)^{\frac{q}{p-q}} \mathcal{W}(t)^{\frac{\beta q}{r}} u(t) V_p(a, t)^{\frac{pq}{p-q}} dt \right)^{\frac{r(p-q)}{q(p-r)}}, \quad x \in (a, b),$$

it is clear that $A_4 \approx C_5$. We have also shown in (4.9) that $B_2 \lesssim C_3$. Hence, it remains to show that $C_3 \lesssim A_4 + B_2$. To this end, we can use (4.18), and obtain $A_2 \lesssim A_4$. Moreover, we know from (4.11) that $C_3 \lesssim A_2 + B_2$. Consequently, $C_3 \lesssim A_4 + B_2$ holds and the proof is complete. \square

Proof of Theorem 1.2 We will prove that inequality (1.9) holds for all $f \in \mathfrak{M}^\uparrow(a, b)$ if and only if inequality

$$\left(\int_a^b \left(\int_a^x \left(\int_a^t h(\tau) d\tau \right)^{\frac{1}{p}} u(t) dt \right)^q w(x) dx \right)^{\frac{p}{q}} \leq C^p \int_a^b h(x) \left(\int_x^b v \right) dx \quad (4.21)$$

holds for all $h \in \mathfrak{M}^+(a, b)$.

Assume that (1.9) holds for all $f \in \mathfrak{M}^\uparrow(a, b)$. Substituting $f(x) = \left(\int_a^x h \right)^{\frac{1}{p}}$, $x \in (a, b)$ for $h \in \mathfrak{M}^+(a, b)$ in (1.9) and applying Fubini's theorem on the right-hand side, (4.21) follows.

Conversely, assume that (4.21) holds for all $h \in \mathfrak{M}^+(a, b)$. Since any $f \in \mathfrak{M}^\uparrow(a, b)$, even if $f(0) > 0$, can be approximated pointwise from below by a function of the form $f(x)^p = \int_a^x h$, $x \in (a, b)$, then the validity of (4.21) yields (1.9). Therefore, the result follows from Theorem 1.1. □

Acknowledgments

T. Ünver would like to express her gratitude to the Institute of Mathematics of the Czech Academy of Sciences for hosting her, providing her with a perfect working atmosphere, and supplying technical support.

We would like to thank the anonymous referees for their thorough and critical reading of the paper and their numerous useful comments and suggestions, which helped improve it substantially.

Declarations

- Funding:

The research of A. Gogatishvili was partially supported by the grant project 23-04720S of the Czech Science Foundation (GAČR), The Institute of Mathematics, CAS is supported by RVO:67985840, by Shota Rustaveli National Science Foundation (SRNSF), grant no: FR21-12353, and by the grant Ministry of Education and Science of the Republic of Kazakhstan (project no. AP14869887). The research of T. Ünver was supported by the grant of The Scientific and Technological Research Council of Turkey (TUBITAK), Grant No: 1059B192000075.

- Conflict of interest:

The authors have no competing interests to declare relevant to this article's content.

References

- [1] M. I. Aguilar Cañestro, P. Ortega Salvador and C. Ramírez Torreblanca, Weighted bilinear Hardy inequalities. *J. Math. Anal. Appl.* 387 (2012), no. 1, 320–334. <https://doi.org/10.1016/j.jmaa.2011.08.078>
- [2] G. Bennett, Some elementary inequalities. III, *Quart. J. Math. Oxford Ser. (2)*, 42 (1991) 149–174. <https://doi.org/10.1093/qmath/42.1.149>
- [3] S. Bloom and R. Kerman, Weighted norm inequalities for operators of Hardy type, *Proc. Amer. Math. Soc.*, 113 (1991), 135–141. <https://doi.org/10.2307/2048449>

- [4] V. I. Burenkov and R. Oinarov, Necessary and sufficient conditions for boundedness of the Hardy-type operator from a weighted Lebesgue space to a Morrey-type space, *Math. Inequal. Appl.* 16 (2013), 1–19. <https://doi.org/10.7153/mia-16-01>
- [5] A. Gogatishvili, M. Křepela, L. Pick, and F. Soudský, Embeddings of Lorentz-type spaces involving weighted integral means, *J. Funct. Anal.*, 273 (2017), 2939–2980. <https://doi.org/10.1016/j.jfa.2017.06.008>
- [6] A. Gogatishvili, Z. Mihula, L. Pick, H. Turčinová, and T. Ünver, Weighted inequalities for a superposition of the Copson operator and the Hardy operator, *J. Fourier Anal. Appl.*, 28 (2022), Paper No. 24. 24 pp. <https://doi.org/10.1007/s00041-022-09918-6>
- [7] A. Gogatishvili, R. Ch. Mustafayev, and T. Ünver, Embeddings between weighted Copson and Cesàro function spaces, *Czechoslovak Math. J.*, 67(142) (2017), 1105–1132. <https://doi.org/10.21136/CMJ.2017.0424-16>
- [8] A. Gogatishvili and R. Ch. Mustafayev, Weighted iterated Hardy-type inequalities, *Math. Inequal. Appl.*, 20 (2017), 683–728. <https://doi.org/10.7153/mia-20-45>
- [9] A. Gogatishvili, R. Ch. Mustafayev, and T. Ünver, Embedding relations between weighted complementary local Morrey-type spaces and weighted local Morrey-type spaces, *Eurasian Math. J.*, 8 (2017), 34–49.
- [10] A. Gogatishvili and L. Pick, Discretization and anti-discretization of rearrangement-invariant norms, *Publ. Mat.*, 47 (2003), 311–358. https://doi.org/10.5565/PUBLMAT_47203_02
- [11] A. Gogatishvili, L. Pick, and T. Ünver, Weighted inequalities involving Hardy and Copson operators, *J. Funct. Anal.*, 283 (12) (2022), Paper No. 109719. 50 pp. <https://doi.org/10.1016/j.jfa.2022.109719>
- [12] A. Gogatishvili and V. D. Stepanov, Reduction theorems for operators on the cones of monotone functions, *J. Math. Anal. Appl.*, 405 (2013), 156–172. <https://doi.org/10.1016/j.jmaa.2013.03.046>
- [13] A. Gogatishvili and V. D. Stepanov, Reduction theorems for weighted integral inequalities on the cone of monotone functions, *Uspekhi Mat. Nauk*, 68 (2013), 3–68. <https://doi.org/10.1070/rm2013v068n04abeh004849>
- [14] M. L. Goldman, Order-sharp estimates for Hardy-type operators on the cones of functions with properties of monotonicity, *Eurasian Math. J.*, 3 (2012), 53–84.
- [15] M. L. Goldman, Some equivalent criteria for the boundedness of Hardy-type operators on the cone of quasimonotone functions, *Eurasian Math. J.*, 4 (2013), 43–63.
- [16] K.-G. Grosse-Erdmann, The blocking technique, weighted mean operators and Hardy’s inequality, *Lecture Notes in Mathematics*, Volume 1679 of Springer-Verlag, Berlin, 1998. <https://doi.org/10.1007/BFb0093486>
- [17] H. P. Heinig and V. D. Stepanov, Weighted Hardy inequalities for increasing functions, *Canad. J. Math.*, 45 (1993), 104–116. <https://doi.org/10.4153/CJM-1993-006-3>

- [18] V. Kabaila, On imbedding the space $L_p(\mu)$ into $L_r(\nu)$, *Litovsk. Mat. Sb.*, 21 (1981), 143–148.
- [19] A. Kalybay, Weighted estimates for a class of quasilinear integral operators. *Sib. Math. J.* 60(2) (2019), 291–303. <https://doi.org/10.33048/smzh.2019.60.209>
- [20] A. Kalybay and R. Oinarov, On weighted inequalities for a class of quasilinear integral operators, *Banach J. Math. Anal.* 17 (2023), no. 1, Paper No. 3, 18 pp. <https://doi.org/10.1007/s43037-022-00226-1>
- [21] A. Kufner, L.-E. Persson, and N. Samko, *Weighted inequalities of Hardy type*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, second ed., 2017.
- [22] M. Křepela, Boundedness of Hardy-type operators with a kernel: integral weighted conditions for the case $0 < q < 1 \leq p < \infty$, *Rev. Mat. Complut.*, 30 (2017), 547–587. <https://doi.org/10.1007/s13163-017-0230-9>
- [23] M. Křepela, Iterating bilinear Hardy inequalities. *Proc. Edinb. Math. Soc. (2)* 60 (2017), no. 4, 955–971. <https://doi.org/10.1017/S0013091516000602>
- [24] M. Křepela and L. Pick, Weighted inequalities for iterated Copson integral operators, *Studia Math.*, 253 (2020), 163–197. <https://doi.org/10.4064/sm181016-5-5>
- [25] Q. Lai, Weighted modular inequalities for Hardy type operators, *Proc. London Math. Soc. (3)*, 79 (1999), 649–672. <https://doi.org/10.1112/S0024611599012010>
- [26] F. J. Martín-Reyes and E. Sawyer, Weighted inequalities for Riemann-Liouville fractional integrals of order one and greater, *Proc. Amer. Math. Soc.*, 106 (1989), 727–733. <https://doi.org/10.2307/2047428>
- [27] E. A. Myasnikov, L.-E. Persson, and V. D. Stepanov, On the best constants in certain integral inequalities for monotone functions, *Acta Sci. Math. (Szeged)*, 59 (1994), 613–624.
- [28] R. Oinarov, Two-sided estimates for the norm of some classes of integral operators, *Proc. Steklov Inst. Math.*, 204 (1994), 205–214.
- [29] D. V. Prokhorov, On a weighted inequality of Hardy type, *Dokl. Akad. Nauk*, 453 (2013), 265–267. <https://doi.org/10.1134/s1064562413060161>
- [30] D. V. Prokhorov and V. D. Stepanov, On weighted Hardy inequalities in mixed norms, *Proc. Steklov Inst. Math.*, 283 (2013), 149–164. Translation of *Tr. Mat. Inst. Steklova* 283 (2013), 155–170. <https://doi.org/10.1134/S0081543813080117>
- [31] G. Sinnamon and V. D. Stepanov, The weighted Hardy inequality: new proofs and the case $p = 1$, *J. London Math. Soc. (2)*, 54 (1996), 89–101. <https://doi.org/10.1112/jlms/54.1.89>
- [32] V. D. Stepanov, Weighted norm inequalities of Hardy type for a class of integral operators, *J. London Math. Soc. (2)*, 50 (1994), 105–120. <https://doi.org/10.1112/jlms/50.1.105>

- [33] V. D. Stepanov and G. E. Shambilova, On iterated and bilinear integral Hardy-type operators. *Math. Inequal. Appl.* 22 (2019), 1505–1533. <https://doi.org/10.7153/mia-2019-22-105>
- [34] T. Ünver, Embeddings between weighted Cesàro function spaces, *Math. Inequal. Appl.*, 23 (2020), 925–942. <https://doi.org/10.7153/mia-2020-23-72>