ON EXISTENCE OF SADOVSKII VORTEX PATCH: A TOUCHING PAIR OF SYMMETRIC COUNTER-ROTATING UNIFORM VORTEX

KYUDONG CHOI, IN-JEE JEONG, AND YOUNG-JIN SIM

ABSTRACT. The Sadovskii vortex patch is a traveling wave for the two-dimensional incompressible Euler equations consisting of an odd symmetric pair of vortex patches touching the symmetry axis. Its existence was first suggested by numerical computations of Sadovskii in [J. Appl. Math. Mech., 1971], and has gained significant interest due to its relevance in inviscid limit of planar flows via Prandtl–Batchelor theory and as the asymptotic state for vortex ring dynamics. In this work, we prove the existence of a Sadovskii vortex patch, by solving the energy maximization problem under the exact impulse condition and an upper bound on the circulation.

Contents

1. Introduction	2
1.1. Significance of Sadovskii vortex patch	3
1.2. Technical statements	4
1.3. Key ideas	6
1.4. Further technical discussions	10
1.5. Open problems	12
1.6. Outline of the paper	13
2. Preliminaries	13
2.1. Notations	13
2.2. Some technical estimates	13
3. Existence of maximizers of patch-type	15
3.1. Variational problem on exact impulse and mass bound	15
3.2. Existence of maximizers	16
3.3. Estimate on maximal energy	26
4. Maximizers under small impulse with unit mass bound	27
4.1. Vanishing of flux constant	28
4.2. Touching of pair of uniform vortices	29
4.3. Estimates when flux constant vanishes due to small impulse	34
5. Variational problem on exact impulse without mass bound	37
5.1. Variational problem without mass bound	37
5.2. Scaling property	38
5.3. Proof of Theorem 5.1	39
Acknowledgments	40
References	40

Date: July 2, 2024.

1

1. Introduction

We consider the Cauchy problem for the 2D incompressible Euler equations in vorticity form:

(1.1)
$$\begin{cases} \partial_t \theta + \mathbf{u} \cdot \nabla \theta = 0, \\ \mathbf{u} = k * \theta, \\ \theta|_{t=0} = \theta_0, \end{cases}$$

where $\theta: [0, \infty) \times \mathbb{R}^2 \to \mathbb{R}$. Here the kernel k is given as

$$k(\mathbf{x}) = (2\pi)^{-1}(-x_2, x_1)/|\mathbf{x}|^2$$

implying that $\mathbf{u} = (u^1, u^2) = \nabla^{\perp} \psi$ where $\nabla^{\perp} = (\partial_{x_2}, -\partial_{x_1})$ with the stream function $\psi = \psi[\theta]$ defined by

$$\psi := (-\Delta)_{\mathbb{R}^2}^{-1} \theta = -\left(\frac{1}{2\pi} \log |\cdot|\right) * \theta.$$

In this paper, we are concerned with traveling waves of (1.1) with the following properties: The function $\theta:[0,\infty)\times$ $\mathbb{R}^2 \to \mathbb{R}$ given by

$$\theta(t, \mathbf{x}) = (\mathbf{1}_A - \mathbf{1}_{A_-})(\mathbf{x} - (W, 0)t)$$

solves (1.1) for some constant W > 0 and for a bounded open set $A \subset \{(x_1, x_2) \in \mathbb{R}^2 : x_2 > 0\}$ satisfying that, for some R > 0, we have

$$\{\mathbf{x} \in \mathbb{R}^2 : |\mathbf{x}| < R\} \subset \overline{A \cup A_-}$$

where A_- is defined as $A_- := \{(x_1, x_2) \in \mathbb{R}^2 : (x_1, -x_2) \in A\}$. In other words, we deal with vortex patch solutions whose support in the upper half plane touches the horizontal axes and travels with the constant velocity (W, 0) without change in its profile. Hereafter, we will refer to such a solution as a Sadovskii vortex patch in honor of Sadovskii's contribution [78] in 1971. While there has been extensive research on such touching vortex patches (including [21, 71, 74, 80, 81, 102, 76, 89]), most of it has focused on their properties rather than proving their existence. We shall defer the detailed historical discussion to Subsection 1.1 below.

Our main result in this paper gives the existence of a Sadovskii vortex patch as a bounded open set in the upper half plane $\mathbb{R}^2_+ := \{(x_1, x_2) \in \mathbb{R}^2 : x_2 > 0\}.$

Theorem 1.1. There exists a bounded open set $\Omega \subset \mathbb{R}^2_+$ satisfying the following properties:

(i) (Touching) There exists a constant R > 0 such that

$$\{ \mathbf{x} \in \mathbb{R}^2 : |\mathbf{x}| < R \} \subset \overline{\Omega \cup \Omega}$$

- (ii) (Impulse normalization) We have $\int_{\Omega} x_2 d\mathbf{x} = 1$. (iii) (Traveling wave) The $(L^1 \cap L^{\infty})(\mathbb{R}^2)$ function $\theta_0 : \mathbb{R}^2 \to \mathbb{R}$ defined by

$$\theta_0 := \mathbf{1}_{\rm O} - \mathbf{1}_{\rm O}$$

generates a traveling wave in the sense that, for the constant W > 0 given by

$$W:=\frac{1}{3}\int_{\mathbb{R}^2}\left|\mathbf{u}_0\right|^2d\mathbf{x}\quad where\ \mathbf{u}_0:=k*\theta_0,$$

the function $\theta:[0,\infty)\times\mathbb{R}^2\to\mathbb{R}$ defined by

$$\theta(t, \mathbf{x}) := \theta_0(\mathbf{x} - (W, 0)t)$$

solves the 2D incompressible Euler equations (1.1).

(iv) (Steiner symmetric with continuous boundary) There exists a continuous function $l:[0,\infty)\to[0,\infty)$ such that

$$\{(x_1, x_2) \in \overline{\Omega} : x_2 = s\} = \{(x_1, s) \in \overline{\mathbb{R}^2_+} : |x_1| \le l(s)\} \text{ for each } s \ge 0.$$

The rest of the introduction is organized as follows. We begin with describing the relevance of Sadovskii vortex patch in fluid dynamics in Subsection 1.1. Then, in Subsection 1.2, we divide the statement of Theorem 1.1 into three theorems, which together gives the main result. Subsection 1.3 provides key ideas that goes into the proofs. Important technical discussions and open problems are given in Subsections 1.4 and 1.5, respectively.




FIGURE 1. Setup of Gol'dshtik (1962), Childress (1966), and Sadovskii (1971). The steady vorticity is 0 and 1 in $\mathbb{R}^2_+ \setminus \Omega$ and Ω , respectively, and the problem is to find the dashed curve.

1.1. Significance of Sadovskii vortex patch.

Counter-rotating vortex pairs are one of the simplest flow patterns and easily observed in various situations, most notably in the wake of any flying objects. There have been numerous studies of counter-rotating vortex pairs from experimental, computational, and theoretical point of view. An excellent review of the literature is given in Leweke–Dizès–Williams [65] (see also Saffman [79]). While Sadovskii vortex can be thought of just an example of such vortex pairs, it has several distinctive characters as we shall explain below.

History of Sadovskii vortex. The Prandtl-Batchelor theorem states that, if the vanishing viscosity limit of a twodimensional flow has closed streamlines, then the limiting vorticity is piecewise constant in each region separated by the streamlines. Therefore, there has been significant interest in understanding potential shape of limiting closed streamlines in various flow geometries. The case of a streamline intersecting the fluid boundary is of particular interest. This was precisely the motivation of Sadovskii (1971) [78]: in the upper half-plane \mathbb{R}^2_+ , he considered the problem of determining a region Ω of uniform vorticity corresponding to a steady flow which is irrotational in $\mathbb{R}^2_+ \backslash \Omega$. He assumed that the steady velocity is asymptotic to $-We_1$ at infinity, which is precisely (1.2). The key assumption on Ω was $\Omega \cap \mathbb{R}^2_+ = [-1, 1]$, and the problem was to determine the streamline connecting (-1, 0) with (1, 0) in the interior of \mathbb{R}^2_+ (see Figure 1). Sadovskii was able to numerically compute the family of such streamlines, parameterized by the strength of the vortex sheet (size of the velocity jump) across this interior streamline. Sadovskii's original computation indicated that the case when the vortex sheet is absent is distinguished by the fact that the corresponding streamline is orthogonal to the domain boundary; see [78, Figure 3]. After this work, the term Sadovskii vortex was widely used to denote equilibrium inviscid flows involving uniform vorticity regions, potentially with vortex sheets, in various geometries ([20, 22, 41, 71, 3, 95]). Therefore, to avoid confusion, in the current work we reserve the phrase Sadovskii vortex patch to denote a uniformly translating vortex patch, touching the boundary of the half-plane, without an accompanying vortex sheet. Sadovskii himself wrote [78, p. 734] that "An analytical proof of the existence and uniqueness (or otherwise) of the solution of this system does not seem possible." While many authors have studied the properties of Sadovskii vortex patch since then ([100, 21, 20, 22, 41, 51, 62, 71, 74, 80, 81, 76, 89, 102]), the rigorous proof of its existence seems to be still open. In particular, it is not clear whether the boundary of the vortex region defines a smooth curve up to $\partial \mathbb{R}^2_+$, the property assumed by Sadovskii and most of the subsequent authors. The paper [100, Table I] includes numerical computations for a Sadovskii vortex patch, which shows that when the maximum height from the axis is normalized to be 1, the length touching the x_1 -axis is approximately 3.398, the area (or mass) is about 2.47, and the vertical center of mass is around 0.415. Lastly, we point out that essentially the same problem of Sadovskii was considered earlier by Gol'dshtik (1962, [47]) and Childress (1966, [23]), although these authors were using certain approximations of the Euler equations to compute the solutions.

Sadovskii vortex as the endpoint of the bifurcation diagram. There exist a large set of counter-rotating vortex pairs, even if we restrict them to the class of symmetric and uniformly-translating ones. In what follows, we shall further assume that the uniformly translating vortex pairs have uniform normalized vorticity (i.e. vortex patch). We further assume for simplicity that the vortex is odd symmetric with respect to the x_1 -axis and non-negative on the upper half-plane \mathbb{R}^2_+ . The important physical parameters are the circulation Γ (=mass), traveling velocity W, and the impulse μ . (Note that all of them are positive under our setting.) Using scale invariance of the incompressible Euler equations, one may normalize one of them: our choice is to normalize the impulse to be 1, as seen in the statement of Theorem 1.1. In the upper half plane, a point vortex (vorticity given as the Dirac delta measure at a point) is formally a traveling wave solution of the Euler equations, and one may de-singularize it to obtain uniformly traveling vortex patches with $\|\omega\|_{L^\infty} = O(\epsilon^{-1})$ and Γ , W, $\mu = O(1)$ for sufficiently small $\epsilon > 0$ ([10, 15, 98]). Normalizing such a vortex

to be of unit vorticity can give a uniformly traveling vortex patch with $\Gamma = \mu = 1$ and $W = O(\epsilon)$. As $\epsilon \to 0^+$, the patches are asymptotically circular and $W \to 0^+$. Then, one may find a branch of uniformly traveling patch solutions by increasing W and keeping $\mu = 1$. Numerical studies show that there is a critical threshold at which the patch becomes a Sadovskii vortex and bifurcation cannot be continued further ([74, 67, 65]). This phenomenon seems to show relevance of Sadovskii vortex patch in dynamics of solutions; we expand this point further in Subsections 1.3 and 1.4 below.

Sadovskii vortex in three-dimensional flows. In three-dimensional incompressible inviscid flows, a counter-rotating pair of vortex rings can approach each other, increasing their distance from the symmetry axis and at the same time giving rise to various instabilities. This "head-on collision" scenario of vortex rings was the subject of intense numerical and computational studies ([60, 61, 82, 34, 84, 66, 90, 75, 73, 19, 49, 86, 84, 83, 17]). Employing the axisymmetric Euler equations and imposing exact odd symmetry (this is equivalent with posing the Euler equations on the upper half-space \mathbb{R}^3_+), one can suppress various instabilities and ask the asymptotic behavior of the axial cross section of the rings for large times.

Based on arguments based on variational principles, Childress [25, 26] argued that the growth of the vorticity maximum $\|\omega(t,\cdot)\|_{L^\infty}$ as well as the distance of the vortex from the symmetry axis can grow at most by $t^{4/3}$ as $t\to\infty$. (The exponent 4/3 can be explained by dimensional arguments based on kinetic energy conservation.) Later, Childress and his collaborators ([28, 27]) performed numerical computations and provided theoretical and computational evidence that the optimal $t^{4/3}$ growth is indeed achieved by vortex rings whose axial cross sections are asymptotic to the Sadovskii vortex after a time-dependent rescaling. As long as the cross section of the vortex ring is sufficiently localized, this is reasonable since far away from the symmetry axis, the axisymmetric Biot–Savart kernel converges to that for the two-dimensional Euler equations ([24]). Furthermore, note the numerical computations done by Carley [17]. Here, it seems that the appearance of this limiting asymptotic profile is not depending sensitively on the shape of the initial vortex ring, and in particular, it is not needed that the initial ring is "attached" to the boundary of the upper half-space. Indeed, this dynamical appearance of the Sadovskii vortex patch was observed earlier and explicitly by Shariff–Leonard–Ferziger in 2008 [83] for the head-on collision of two Hill's spherical vortices, see [83, Figures 8, 9]. So far, the best known lower bound in time on the vorticity growth is slower than linear-in-time ([30, 50]) and it seems that understanding stability of the Sadovskii vortex in the two-dimensional case is the natural first step towards establishing the sharp growth $t^{4/3}$.

1.2. **Technical statements.** In this subsection, in order to more easily convey the ideas needed for the proof of our main result (Theorem 1.1), we divide Theorem 1.1 into the following three theorems: Variational method (Theorem 1.2), Touching property (Theorem 1.4), and Boundary behavior (Theorem 1.6).

To find a traveling wave solution, for some constant W > 0, we are looking for a steady solution (\mathbf{U}, ω) that solves

(1.2)
$$\begin{cases} \mathbf{U} \cdot \nabla \omega = 0 & \text{in } \mathbb{R}^2, \\ \mathbf{U} = k * \omega - (W, 0), \end{cases}$$

so that the solution (\mathbf{u}, θ) in the form of

(1.3)
$$\mathbf{u}(t, \mathbf{x}) = \mathbf{U}(\mathbf{x} - (W, 0)t) + (W, 0),$$

(1.4)
$$\theta(t, \mathbf{x}) = \omega(\mathbf{x} - (W, 0)t),$$

solves (1.1). The equations (1.2) admits a vortex patch solution of a form

$$\omega = \mathbf{1}_{\{\Psi>0\}} = \begin{cases} 1 & \text{for } \Psi > 0, \\ 0 & \text{for } \Psi \le 0 \end{cases} \quad \text{in } \mathbb{R}^2_+,$$

with the odd symmetry $\omega(x_1, -x_2) = -\omega(x_1, x_2)$ for each $(x_1, x_2) \in \mathbb{R}^2_+$, and $\mathbf{U} = \nabla^{\perp} \Psi$ where

$$\Psi(\mathbf{x}) := \psi[\omega](\mathbf{x}) - Wx_2 - \gamma$$

for some constant $\gamma \ge 0$ which is referred to as the flux constant in the literature. One may easily observe that touching of a vortex patch implies $\gamma = 0$. We will prove that such a solution with $\gamma = 0$ can be found, and it is a Sadovskii vortex patch.

¹See dynamical appearance of Sadovskii-like vortex in https://people.bath.ac.uk/ensmjc/Research/Vortex/headon.html.

We will obtain Theorem 1.1 by the following steps. In Theorem 1.2, we will find a solution ω of (1.2) satisfying that, with the constants W > 0 and $\gamma = 0$, we have

$$\omega = \mathbf{1}_{\{\mathbf{x} \in \mathbb{R}^2_+ : \psi[\omega](\mathbf{x}) - Wx_2 > 0\}}$$
 a.e. in \mathbb{R}^2_+ .

In Theorem 1.4, for the function ω we have found above, we will show that there exists R > 0 such that we have

[Touching]
$$\{ \mathbf{x} \in \mathbb{R}^2_+ : |\mathbf{x}| < R \} \subset \{ \mathbf{x} \in \mathbb{R}^2_+ : \psi[\omega](\mathbf{x}) - Wx_2 > 0 \}.$$

In Theorem 1.6 we will say that the boundary of the set $\{\psi[\omega] - Wx_2 > 0\}$ is continuous.

In Theorem 1.2 below, we use a variational method to prove that there exists a patch-type solution of (1.2) which consequently generates a traveling wave solution of (1.1) with the traveling speed W > 0 and the constant $\gamma = 0$. In specific, we will solve the variational problem of finding a maximizer ω of the kinetic energy with a fixed constraint on the impulse $\int_{\mathbb{R}^2} x_2 \omega \, d\mathbf{x}$. The proof of Theorem 1.2 can be found in the proof of Theorem 5.1.

Theorem 1.2. Define K by

$$K:=\left\{\omega\in L^1(\mathbb{R}^2)\ :\ \omega=\mathbf{1}_A-\mathbf{1}_{A_-}\ \ a.e.\ in\ \mathbb{R}^2\quad for\ some\ open\ bounded\ set\ A\subset\mathbb{R}^2_+\ \ satisfying\ \int_Ax_2\ d\mathbf{x}=1\right\}$$

and define $S \subset K$ by the set of maximizers of the kinetic energy

$$\int_{\mathbb{R}^2} |\mathbf{u}|^2 d\mathbf{x}, \quad \text{where } \mathbf{u} := k * \omega \quad \text{for each } \omega \in K.$$

Then the maximal kinetic energy in the class K is finite, and the set S is nonempty. Moreover, there exist absolute constants $C_0, C_1, C_2 \in (0, \infty)$ satisfying $C_1 < C_2$ such that, for each $\omega \in S$, there exists a constant $\tau \in \mathbb{R}$ and a set $\Omega \subset \mathbb{R}^2_+$ which is open bounded, satisfying the following properties:

(i) By denoting ω_{τ} as the translation of ω of amount $(\tau,0)$ by $\omega_{\tau}(x_1,x_2) := \omega(x_1-\tau,x_2)$, we have

$$\omega_{\tau} = \mathbf{1}_{\Omega} - \mathbf{1}_{\Omega_{-}}$$
 a.e. in \mathbb{R}^2 .

Moreover, we get

$$\Omega = \left\{ \mathbf{x} \in \mathbb{R}_+^2 : \psi[\omega_\tau](\mathbf{x}) - W x_2 > 0 \right\}$$

with the constant W > 0 given by

$$W := \frac{1}{3} \int_{\mathbb{R}^2} |\mathbf{u}|^2 d\mathbf{x} \quad where \, \mathbf{u} := k * \omega.$$

(ii) There exists a function $l:(0,\infty)\to [0,\infty)$ such that we have

$$\left\{ \, (x_1,x_2) \in \Omega \, : \, x_2 = s \, \right\} \, = \, \left\{ \, (x_1,s) \in \mathbb{R}_+^2 \, : \, |x_1| < l(s) \, \right\} \quad for \, each \, s > 0.$$

Moreover, we have

$$\Omega \subset \{ \mathbf{x} \in \mathbb{R}^2 : |\mathbf{x}| < C_0 \} \quad and \quad C_1 < |\Omega| < C_2$$

where $|\cdot|$ denotes the (Lebesgue) measure in \mathbb{R}^2 .

Remark 1.3. In Theorem 1.2, the traveling speed is proportional to the maximal kinetic energy. Moreover, observe that the condition $\int_A x_2 d\mathbf{x} = 1$ in the definition of K can be replaced by $\int_A x_2 d\mathbf{x} = \mu$ for any $\mu \in (0, \infty)$. Indeed, the scaling

$$\omega'(\mathbf{x}) := \omega(\mu^{-1/3}\mathbf{x}) \quad \text{for } \omega \in S$$

gives a one-to-one correspondence between the sets of maximizers. Then we get the same results with adjusted constants W', C'_0 , C'_1 , $C'_2 > 0$ under the relations

$$W' = \frac{1}{3\mu} \int_{\mathbb{R}^2} |\mathbf{u}'|^2 d\mathbf{x} = W\mu^{1/3}, \quad C_0' = C_0\mu^{1/3}, \quad C_1' = C_1\mu^{2/3}, \quad C_2' = C_2\mu^{2/3}.$$

Theorem 1.4 below says that, for each traveling vortex patch that is symmetrically concentrated to a vertical line with odd symmetry with respect to the horizontal line and travels with a constant velocity (W, 0), the fluid velocity at the center of the dipolar vortex has the direction (1, 0), with the speed greater than 2W. Its proof will be deduced from the proof of Lemma 4.5.

Theorem 1.4. Let W > 0 be any constant. Let $\Omega \subset \mathbb{R}^2_+$ be an open bounded set satisfying that $\int_{\Omega} x_2 d\mathbf{x} > 0$, and for some function $l:(0,\infty) \to [0,\infty)$ assume that

$$\{(x_1, x_2) \in \Omega : x_2 = s\} = \{(x_1, s) \in \mathbb{R}^2_+ : |x_1| < l(s)\} \text{ for each } s > 0,$$

and that the function $\theta:[0,\infty)\times\mathbb{R}^2\to\mathbb{R}$ given by

$$\theta(t, \mathbf{x}) := (\mathbf{1}_{\Omega} - \mathbf{1}_{\Omega_{-}})(\mathbf{x} - (W, 0)t)$$

solves (1.1). Then for the velocity function $\mathbf{u}:[0,\infty)\times\mathbb{R}^2\to\mathbb{R}^2$ defined by $\mathbf{u}=k*\theta$, we have

$$u^{1}(t,(Wt,0)) = u^{1}(0,(0,0)) > 2W$$
 and $u^{2}(t,(Wt,0)) = u^{2}(0,(0,0)) = 0$ for each $t > 0$.

Remark 1.5. [Proof of Theorem 1.1 (i)-(iii) assuming Theorem 1.2 and Theorem 1.4] By Theorem 1.2, there exists a function ω having the odd symmetry with respect to x_1 -axis such that we have $\int_{\mathbb{R}^2} x_2 \omega \, d\mathbf{x} = 1$ and the relation

$$\omega = \mathbf{1}_{\{\psi[\omega] - Wx_2 > 0\}}$$
 a.e. in \mathbb{R}^2_+

where the set $\{\psi[\omega] - Wx_2 > 0\} \cap \mathbb{R}^2_+$ is symmetrically concentrated along x_2 -axis. Note that ω solves (1.2) and generates a traveling wave solution of (1.1). Then by Theorem 1.4, we have

$$\left(\frac{\partial}{\partial x_2}\psi[\omega]\right)(0,0) = u^1(0,(0,0)) > 2W.$$

Due to the continuity of $\nabla \psi[\omega]$ in \mathbb{R}^2 , there exists R > 0 such that

$$\left(\frac{\partial}{\partial x_2}\psi[\omega]\right) > 2W \quad \text{in} \quad B_R(0,0) := \{ \mathbf{x} \in \mathbb{R}^2 : |\mathbf{x}| < R \}.$$

This implies that we obtain

$$B_R(0,0) \cap \mathbb{R}^2_+ \subset \{ \psi[\omega] - Wx_2 > 0 \} \cap \mathbb{R}^2_+,$$

since the following inequality holds for each $\mathbf{x} \in B_R(0,0) \cap \mathbb{R}^2_+$:

$$\psi[\omega](\mathbf{x}) = \int_0^{x_2} \left(\frac{\partial}{\partial x_2} \psi[\omega]\right) (x_1, s) \, ds > 2W x_2 > W x_2.$$

Here we used the fact that $\psi[\omega](x_1,0) = 0$ for any $x_1 \in \mathbb{R}$, which is a consequence of the odd-symmetry of ω .

Lastly, Theorem 1.6 below implies Theorem 1.1 (iv). For its proof, see the proof of Theorem 4.2.

Theorem 1.6. Let S be the set of functions in Theorem 1.2. Choose any $\omega \in S$ with the corresponding function $l:(0,\infty) \to [0,\infty)$. Then l is continuous in $(0,\infty)$ and the limit

$$\lim_{s \searrow 0} l(s) \in (0, \infty) \quad exists.$$

Moreover, there exists $L \in (0, \infty)$ such that, for each $\epsilon \in (0, L)$, we have $l \in C^{1,r}([\epsilon, L])$ for each $r \in (0, 1)$.

1.3. Key ideas.

Let us describe the key ideas that are involved in the proof of the main theorem.

Steady Euler equations and vorticity function. We recall the steady Euler equation (1.2) in \mathbb{R}^2_+ , for an arbitrary traveling speed W > 0. This equation allows a solution of the form

$$\omega = f(\Psi)$$
 in \mathbb{R}^2

for a function f called a *vorticity function* where the adjusted stream function Ψ is given by

$$\Psi(\mathbf{x}) := \psi[\omega](\mathbf{x}) - Wx_2 - \gamma$$
 for some constant $\gamma \ge 0$

with the relation $U = \nabla^{\perp}\Psi$. The constant γ is referred to as a *flux constant*. We extend the odd-symmetric solution ω of (1.2) as

$$\omega(x_1, -x_2) = -\omega(x_1, x_2)$$
 for each $(x_1, x_2) \in \mathbb{R}^2_+$.

In this paper, we put the Heaviside step function

$$(1.5) f(\cdot) = \mathbf{1}_{\{\cdot > 0\}},$$

(which is the limit of $f(t) = t_+^{1/(p-1)}$ as p goes to infinity²) as the vorticity function to obtain an odd-symmetric vortex patch solution given by

(1.6)
$$\omega = f(\Psi) = \mathbf{1}_{\{\psi[\omega] - W, x_2 - \gamma > 0\}} \quad \text{in } \mathbb{R}^2_+.$$

In Theorem 3.3, we prove that such a solution exists as a maximizer of the kinetic energy among functions under some constraints on their impulse and mass. Each maximizer is symmetrically concentrated about x_2 -axis in the upper half plane \mathbb{R}^2_+ (so-called *Steiner symmetric about* x_2 -axis; see Definition 3.2 for details).

Touching implies zero flux constant ($\gamma = 0$). We here discuss by which conditions a solution (1.6) would be a Sadovskii vortex patch. We observe that any solution with positive flux constant $\gamma > 0$ cannot touch x_1 -axis. Indeed, if $\gamma > 0$, then for all sufficiently small $x_2 > 0$ we have

$$\mathbf{x} = (x_1, x_2) \notin \{ \mathbf{x} \in \mathbb{R}^2_+ : \psi[\omega](\mathbf{x}) - Wx_2 - \gamma > 0 \},$$

due to $\psi(x_1, x_2)\Big|_{x_2=0} \equiv 0$. In other words, for ω to generate a Sadovskii vortex patch, we should have $\gamma = 0$. However, it is not clear that $\gamma = 0$ is a sufficient condition for *touching* of the dipole, as the distribution of the stream function $\psi - Wx_2$ is unknown.

Zero flux constant ($\gamma = 0$) **implies touching when symmetrically concentrated about** x_2 -axis. In this paper, we show that any vortex patch solution, which has zero flux constant and is symmetrically concentrated about x_2 -axis, always touches x_1 -axis. Unlike [8](also see [2, Section 6]) which used the moving plane method [45] to produce a Lamb-Chaplygin dipole, we directly compare the distribution of ψ and Wx_2 . More specifically, we want to compare the traveling speed W > 0 and the horizontal velocity $u^1 = \psi_{x_2}[\omega]$ to guarantee that

$$\psi_{x_2} - W > 0$$
 near the origin,

so that we get

$$B \cap \mathbb{R}^2_+ \subset \{ \psi - Wx_2 > 0 \}$$

for some ball $B \subset \mathbb{R}^2$ centered on x_1 -axis. For the comparison, we first split the dipole as

$$\omega = \mathbf{1}_A - \mathbf{1}_{A_-} = \omega_+ + \omega_-, \quad \mathbf{u} = \nabla^{\perp} \psi[\omega_+] + \nabla^{\perp} \psi[\omega_-]$$

where ω_+ is the nonnegative (upper) part of ω whose support is in \mathbb{R}^2_+ , and obtain the following relation that, on x_1 -axis,

$$u^1 = 2\psi_{x_2}[\omega_-]$$

thanks to the symmetry between A and A_- . We may observe that the eddy flow generated by the lower part ω_- makes the upper part ω_+ propagate in the direction (W,0), and vice versa. That is, the traveling speed W is the average value of the nonlocal velocity $\psi_{x_2}[\omega_-]$ in the set supp $(\omega_+) \subset \mathbb{R}^2_+$, i.e. we may compare the traveling speed given by

$$W = \int_{\text{supp}(\omega_{+})} \psi_{x_{2}}[\omega_{-}] d\mathbf{x}$$

and the horizontal velocity on x_1 -axis, given by $u^1(x_1,0) = 2\psi_{x_2}[\omega_-](x_1,0)$, if we can estimate the distribution of the nonlocal velocity $\psi_{x_2}[\omega_-]$ in the domain $\overline{\mathbb{R}^2_+}$. In Lemma 4.5, we will prove that the nonlocal velocity $\psi_{x_2}[\omega_-]$ has certain monotonicity and attains its unique maximum at the origin among all points in $\overline{\mathbb{R}^2_+}$ if the set supp (ω) is symmetrically concentrated about x_2 -axis. Therefore we obtain

$$u^1(0,0) > 2W$$
.

which means that the speed at the center of the dipole is more than twice as fast as the traveling speed of the entire dipole. Therefore, $\gamma = 0$ is an equivalent condition for *touching*, when the dipole vortex is symmetrically concentrated about x_2 -axis. From now on, we introduce the variational method to obtain a solution with $\gamma = 0$.

²cf. Lamb–Chaplygin dipole for p = 2 (e.g. see [2]).

Variational problem on exact impulse and mass upper bound. It is classical to use a variational method when seeking a steady solution to the Euler equations. In this paper, in some admissible set, we shall find a maximizer $\omega \in L^1(\mathbb{R}^2)$ of the kinetic energy

$$E(\omega) = \frac{1}{4} \int_{\mathbb{R}^2} |\mathbf{u}|^2 d\mathbf{x}, \quad \mathbf{u} = k * \omega.$$

Given a triple of parameters

$$(\mu, \nu, \lambda)$$
 where $\mu, \nu, \lambda > 0$,

the admissible set $K_{\mu,\nu,\lambda}$ will be taken as the set of functions having the simple forms $\omega = \lambda \cdot (\mathbf{1}_A - \mathbf{1}_{A_-})$ for some $A \subset \mathbb{R}^2_+$, with the prescribed impulse condition

$$\int_{\mathbb{R}^2} x_2 \omega \ d\mathbf{x} = \mu$$

and the upper bound of mass

$$\|\omega\|_{L^1(\mathbb{R}^2)} \leq \nu$$
.

Here, λ is the strength of the vortex patch. We denote the set of maximizers as $S_{\mu,\nu,\lambda}$ and the maximal energy as $I_{\mu,\nu,\lambda}$ (see Remark 3.1 for detailed discussions). The simple scaling argument of Remark 3.1-(v) says that we can kill two parameters of the triple (μ, ν, λ) . For instance, the variational problem of parameters (μ, ν, λ) is equivalent to the problem of (M, 1, 1) where $M = \lambda^{\frac{1}{2}} \nu^{-\frac{3}{2}} \mu$. Indeed, the sets $S_{\mu,\nu,\lambda}$ and $S_{M,1,1}$ are bijective under some scaling map. Therefore we may assume $\nu = \lambda = 1$ and regard the impulse μ as the only parameter of our variational problem.

Existence of maximizer. Theorem 3.3 says that, in the variational problem of the parameter $(\mu, 1, 1)$, there exists a maximizer, and any maximizer $\omega \in S_{\mu,1,1}$ has the form (1.6):

$$\omega = \mathbf{1}_{\{\psi[\omega] - Wx_2 - \gamma > 0\}}$$
 a.e. in \mathbb{R}^2_+ for some constants $W > 0, \ \gamma \ge 0$

with the odd-symmetry in \mathbb{R}^2 . The proof consists of several steps:

(i) In Lemma 3.6, we prove the existence of maximizers having the form of a simple function $\mathbf{1}_A - \mathbf{1}_{A_-}$. We first find a weak subsequential limit $\omega' \in L^2(\mathbb{R}^2)$ of an energy-maximizing sequence ω_n in the larger admissible set: the set of odd-symmetric bounded functions ω satisfying that $0 \le \omega \le 1$ in \mathbb{R}^2_+ with the upper bounds of impulse and mass

$$\int_{\mathbb{R}^2} x_2 \omega \ d\mathbf{x} \le \mu, \quad \|\omega\|_{L^1(\mathbb{R}^2)} \le 1.$$

We may assume that ω' and ω_n satisfy the Steiner symmetrization condition, since we have

$$E(\omega^*) \ge E(\omega)$$

if ω^* is the Steiner symmetrization of ω about x_2 -axis. We will see that the sequence of kinetic energy $E(\omega_n)$ converges to $E(\omega')$, implying that ω' is a maximizer of kinetic energy in the larger admissible set. Then we prove that ω' has the full impulse $\int_{\mathbb{R}^2} x_2 \omega' d\mathbf{x} = \mu$ and has the form $\omega' = \mathbf{1}_A$ in \mathbb{R}^2_+ .

(ii) Proposition 3.11 says that each maximizer ω solves (1.2) in the sense that, for some constant $W, \gamma \geq 0$, the relation (1.6) holds. Knowing that ω is a simple function $\mathbf{1}_A - \mathbf{1}_{A_-}$ for some measurable set $A \subset \mathbb{R}^2_+$, we first prove that there exist some constant $W, \gamma \geq 0$ such that

$$A = \{ \mathbf{x} \in \mathbb{R}^2 : \psi[\omega](\mathbf{x}) - Wx_2 - \gamma > 0 \}.$$

For this, we generate a perturbation of $\omega = \mathbf{1}_A - \mathbf{1}_{A_-}$ in the larger admissible set. We shall construct a function $\eta \in L_c^{\infty}(\mathbb{R}^2_+)$ satisfying that

$$\eta \le 0 \text{ in } A, \quad \eta \ge 0 \text{ in } A^c, \quad \text{and } \int_{\mathbb{R}^2} \eta = \int_{\mathbb{R}^2} x_2 \eta = 0.$$

Then for small $\epsilon > 0$, we have $0 \le \mathbf{1}_A + \epsilon \cdot \eta \le 1$, and the function $\mathbf{1}_A + \epsilon \cdot \eta$ has the same mass and impulse with $\mathbf{1}_A$. The perturbed kinetic energy E_{ϵ} of the odd-extension of $(\mathbf{1}_A + \epsilon \eta)$ is maximized at $\epsilon = 0$ so as to

$$\frac{d}{d\epsilon}E_{\epsilon}\Big|_{\epsilon=0} = \int_{\mathbb{R}^2_+} \psi[\omega] \cdot \eta \ d\mathbf{x} \le 0.$$

We will use this inequality to prove that $A = \{\psi[\omega] - Wx_2 - \gamma > 0\}$ for some constants W, γ , and the construction of η is the key part of the proof. We will define η so that the above inequality gives

$$\int_{\mathbb{R}^2} (\psi[\omega] - Wx_2 - \gamma) h \, d\mathbf{x} \le 0$$

for an arbitrary function h satisfying that $h \le 0$ in A and $h \ge 0$ in A^c , which would eventually gives us the result. Meanwhile, the emergence of the constants W, γ from the formula of η are necessary. However, we observe that W, γ are determined by the boundary points of the set $\{\psi[\omega] - Wx_2 - \gamma > 0\}$ (see (3.2)), while we yet only know that A is a measurable set. It means that the classical definition of (topological) boundary points is not applicable, and we shall define boundary-like points of general measurable sets which are called *the exceptional points* (see Definition 3.12). This approach was developed in [29].

Small impulse implies $\gamma = 0$. For the variational problem with parameters $(\mu, 1, 1)$, Lemma 4.4 and the estimate (4.3) in Lemma 4.6 guarantee that, if μ is small enough, then we can get $\gamma = 0$, i.e. *touching*, and prove that maximizers cannot achieve the full mass. It confirms our heuristic intuition for the case of $\lambda = 1$: if the impulse imposed on the maximizers is small enough compared to the mass, each maximizer would rather be concentrated in a bounded region close to the axis than be stretched widely and thin, allowing for mass loss. On the other hand, one may infer that if the impulse dominates over the mass sufficiently, the maximizers would move away from the axis, that is, $\gamma > 0$ will be large enough.

Estimates using small impulse: admissible set without mass bound. Another result of dealing with small impulse, in addition to *touching*, is that we can find maximizers of the variational problems where the mass bound is omitted. In the below, we explain the reason why obtaining the following estimate:

$$\sup_{\omega \in S_{\mu,1,1}} \|\omega\|_{L^1} \lesssim \mu^{2/3} \quad \text{ for small } \mu > 0$$

is sufficient to guarantee the existence of the maximizer of the variational problem without mass bound. The above estimate follows from (4.5) in Lemma 4.6, and we will shortly give the key idea of the estimate.

(i) Variational problem without mass bound and prior estimate: We first consider the variational problem of maximizing the kinetic energy in an admissible set given as the set of L^1 simple functions $\omega = \mathbf{1}_A - \mathbf{1}_{A_-}$ having the constraint only on the impulse

$$\int_{\mathbb{R}^2} x_2 \omega \, d\mathbf{x} = \mu \quad \text{ for } \mu > 0,$$

where the set of its maximizer functions is denoted by $S_{\mu,\infty}$, and the maximal energy is denoted by $I_{\mu,\infty}$. Assuming that the set $S_{1,\infty}$ is nonempty, we can obtain $\omega \in S_{\mu,\infty}$ from each $\omega' \in S_{1,\infty}$ by the scaling $\omega(\cdot) = \omega'(\mu^{-1/3}(\cdot))$, and get the priori estimate

(ii) Difficulty: It is nontrivial that the set $S_{1,\infty}$ is nonempty because it is not clear that

$$I_{\mu,\infty} < \infty$$
.

Indeed, the kinetic energy is bounded using $\|\omega\|_{L^1}$ from the inequality (2.5):

$$E(\omega) \lesssim \|\omega\|_{L^1} \|\omega\|_{L^2}^{1/2} \Big(\int_{\mathbb{R}^2} x_2 \omega \, d\mathbf{x} \Big)^{1/2},$$

and it is not clear that

$$\sup_{\omega \in S_{1,\infty}} ||\omega||_{L^1} < \infty.$$

Although the kinetic energy of each function in the admissible set is finite, the maximal energy may be infinite, and there might not be any maximizer in the admissible set.

(iii) Small impulse estimate guarantees existence of maximizers: In this paper, in Lemma 4.6, we obtain the estimation : for small $\mu > 0$,

(1.8)
$$\sup_{\omega \in S_{\mu,1,1}} \|\omega\|_{L^1} \lesssim \mu^{2/3}.$$

One may notice that this estimate coincides well with the priori estimate. From now on, we explain how the inequality above would imply that the set $S_{1,\infty}$ is nonempty (For the formal proof, see Section 5). In the meantime, we will observe that the power 2/3 is the critical number to solve the variational problem. First, to prove $S_{1,\infty} \neq \emptyset$, it suffices to show that there exists $v_1 > 0$ such that

(1.9)
$$S_{1,\nu,1} = S_{1,\nu_1,1}$$
 for any $\nu > \nu_1$.

It means that, for fixed impulse, any energy maximizer does not attain arbitrarily large mass, and that the maximal energy $I_{1,\nu,1}$ stops increasing for large $\nu > 0$. It would imply that $I_{1,\infty} = I_{1,\nu_1,1} < \infty$ and therefore $S_{1,\nu_1,1} \subset S_{1,\infty} \neq \emptyset$. To show (1.9), we need to show that, for some $\nu_1 > 0$ and each $\nu > \nu_1$, it holds that

$$\sup_{\omega \in S_{1,\nu,1}} ||\omega||_{L^1} \leq \nu_1,$$

or equivalently

$$\sup_{\omega \in S_{\nu^{-3/2},1,1}} \|\omega\|_{L^1} \ \le \ \frac{\nu_1}{\nu}.$$

If we put $\mu := v^{-3/2}$, we obtain the inequality

$$\sup_{\omega \in S_{\mu,1,1}} ||\omega||_{L^1} \le v_1 \cdot \mu^{2/3}$$

which must holds for small $\mu > 0$. Therefore, we note that the estimate (1.8) is strong enough to prove that the set $S_{1,\infty}$ is nonempty.

(iv) To obtain (1.8) for small impulse: In Lemma 4.6, we obtain (1.8) by several steps. We sketch the strategy of the proof here: for each $\omega \in S_{\mu,1,1}$ with the traveling speed W > 0 and the flux constant $\gamma \ge 0$, we have

$$\|\omega\|_{L^1} \le \left| \left\{ \mathbf{x} \in \mathbb{R}_+^2 : \psi[\omega](\mathbf{x}) > W x_2 \right\} \right|$$

where $|\cdot|$ denotes the (Lebesgue) measure in \mathbb{R}^2 . One may find it inevitable to find the lower bound of W in terms of small $\mu > 0$ for estimating the measure of the set { $\psi[\omega] > Wx_2$ }, knowing that function $\psi[\omega]$ can be controlled by impulse μ (see Lemma 2.1 and Lemma 2.3). We can show that

$$u^{1/3} \le W$$

which follows by the Pohožaev identity (3.5) in Lemma 3.21 (see [7]):

$$I_{\mu,1,1} = \frac{3}{4}W\mu + \frac{1}{2}\gamma||\omega||_{L^1} \quad \text{(where we know } \gamma = 0 \ \text{ for small } \mu \text{)}$$

and by the lower bound estimate of maximal energy

$$I_{\mu,1,1} \gtrsim \mu^{4/3}$$
 (thanks to a scaling)

given in Lemma 4.3.

1.4. Further technical discussions.

Variational methods for vortex dipoles. A vortex dipole is a symmetric pair of vortices with opposite signs that move along the symmetry axis. This type of dipole is often referred to as a *counter-rotating vortex pair*. It is a model of coherent vortex structures in large-scale geophysical flows (for experimental works, see [54], [39] and references therein). The existence via variational method goes back to Kelvin in 1880 [93], Arnold [4] in 1966, and Benjamin in 1976 [5]. In particular, we consider the *vorticity method*, a variational principle based on the *vorticity*. This method for vortex pairs was further developed by Turkington [94] and Burton [7]. We refer to the work of Norbury [72] and Yang [101] for the stream function method.

Our variational setting is based on the vorticity method of Friedman–Turkington [42] in 1981 which was developed for vortex rings. Following [42], we consider the class of vorticity

$$\int_{\mathbb{R}^2_+} \omega \quad \text{is smaller than or equal to given } v > 0.$$

After fixing ν (e.g. $\nu=1$), as in [42, 2, 29], the maximizers for small impulse $\mu=\int_{\mathbb{R}^2_+} x_2\omega > 0$ strictly lose part of its mass (i.e. $\int_{\mathbb{R}^2_+} \omega < 1$). Then we prove that the parameter ν on mass becomes unnecessary. In other words, the maximizers are characterized by a single constraint on impulse (when small). Our Sadovskii vortex patch is obtained as such a maximizer in $K_{\mu,1,1}$ (i.e. $\omega \in S_{\mu,1,1}$ for $\mu \ll 1$). Thus the Sadovskii patch can be considered as the limiting case $\mu \to 0$ of the one parameter family $\mu > 0$.

Approximating the point vortex dipole. The opposite limit $(\mu \to \infty \text{ in } S_{\mu,1,1})$ can be understood in the following sense. By scaling, we observe

$$\omega \in K_{\mu,1,1} \iff \hat{\omega} \in K_{1,1,(1/\varepsilon^2)}$$

via the transform

$$\hat{\omega}(\mathbf{x}) = \frac{1}{\varepsilon^2} \omega(\frac{\mathbf{x}}{\varepsilon})$$
 with $\mu = 1/\varepsilon$.

The class $K_{1,1,(1/\varepsilon^2)}$ consists of patches of strength $1/\varepsilon^2$ with the unit vertical center of mass $(1 = \int_{\mathbb{R}^2_+} x_2 \omega / \int_{\mathbb{R}^2_+} \omega)$. As ε goes to 0, any sequence $\{\omega_{\varepsilon}\}$ from the class $K_{1,1,(1/\varepsilon^2)}$ converges to the Dirac point mass: in other words,

(1.10)
$$\omega_{\varepsilon} \longrightarrow \delta_{(0,1)} - \delta_{(0,-1)} \quad \text{in} \quad \mathbb{R}^2.$$

The study of approximations to the point dipole, known as the *desingularization problem*, has been a subject of interest and research for a long time (e.g. see the classical book [64] of Lamb). We refer to [94, 102, 88, 58, 13] and references therein. In general, for small ε , the boundary is smooth and closed to a circle. For an analogue in vortex ring, see the Helmholtz's rings of small cross-section [55]. One may wonder interaction between vortex dipoles/rings. For instance, we refer to [85, 36, 52] and references therein for leapfrogging phenomena.

Lamb–Chaplygin dipole and Hill's spherical vortex. While we do not know the exact shape of dipoles including our Sadovskii vortex patch, there are other interesting examples which can be explicitly written. One is the Lamb–Chaplygin dipole [64] while the other is the Hill's spherical vortex [57]. We also refer to [56], [70], [1] and references therein. The *Lamb–Chaplygin dipole* is an exact solution to (1.2) with zero flux constant $\gamma = 0$ and the vorticity function given by the Lipschitz function

$$(1.11) f(\cdot) = \max\{\cdot, 0\}$$

(see [64]). Its support is simply a closed ball centered at the origin, and the dipole is symmetrically placed with opposite signs with respect to x_1 -axis. Thanks to the structure of the vorticity function (1.11), the moving plane method [45] can be applied to characterize the dipole. A variational characterization of the dipole can be found in [8, 9] (also see [2] for stability). For Hill's spherical vortex, we refer to [29] and references therein.

Stability questions. The variational method produces an extremizer of a conserved quantity in an appropriate admissible class. Since Arnold's work [4], it is quite classical to deduce (Lyapunov) stability for steady solutions characterized by critical points of the energy (also see the explanation in [43]). The simplest example would be the circular patch (Rankine vortex), whose stability was rigorously proved by [97]. (Strictly speaking, patch solutions are not regular enough for Arnold's argument to be directly applicable, cf. [87, 33].) For vortex pairs, orbital stability was first obtained in [11] for a certain solutions by the vorticity method. Stability of Lamb–Chaplygin dipoles was proved in [2] (also see [98, 31]). The recent paper of Wang [98], which is a refinement of Burton [10], proved stability an approximation for point vortex (1.10). We also refer to [16, 14] and references therein. All the stability results are restricted on *symmetric* perturbations. In general, non-symmetric disturbances may increase the corresponding energy as explained in [80, p2342]. Lastly, uniformly rotating patch solutions are commonly referred to as "V-states". For existence, rigidity and stability, we refer to [6, 96, 92, 59, 18, 48, 32] and references therein.

Vortex atmosphere is touching. In the steady Euler equations (1.2), we consider a solution ω satisfying the Steiner symmetry condition and having a form $\mathbf{1}_A - \mathbf{1}_{A_-}$ where

$$A = \{ \psi[\omega] - Wx_2 - \gamma > 0 \}$$

for some constant $W>0, \ \gamma\geq 0$. Then each streamline $\{\mathbf{x}\in\overline{\mathbb{R}^2_+}: \Psi(\mathbf{x})=C\}$, generated by the adjusted stream function

$$\Psi := \psi[\omega] - Wx_2 - \gamma$$

and characterized by $C \in \mathbb{R}$, is a integral curve of the flow velocity. We note that if a single streamline forms a closed curve and surrounds any bounded subset of $\overline{\mathbb{R}^2_+}$, the set travels along with the traveling vortex patch without change in its profile. By the *vortex atmosphere* we mean the union of all such subsets of \mathbb{R}^2_+ . One may observe that the set $\{\mathbf{x} \in \mathbb{R}^2_+ : \psi(\mathbf{x}) - Wx_2 > 0\}$ is the vortex atmosphere from the observations that the set $\{\mathbf{x} \in \mathbb{R}^2_+ : \psi(\mathbf{x}) - Wx_2 - C > 0\}$ is not bounded for each C < 0, and is bounded for $C \ge 0$ (see the proof of Lemma 3.18). As we can prove that the set $\{\psi - Wx_2 > 0\}$ touches the axis of symmetry (regardless of whether $\gamma > 0$ or $\gamma = 0$), we may conclude that any vortex atmosphere touches the axis of symmetry even when the flux constant γ of the solution is positive. In other words, any traveling dipole vortex patch which is symmetrically concentrated about x_2 -axis carries its certain neighborhood fluid including a part of the axis of symmetry.

1.5. Open problems.

We clarify several open problems regarding Sadovskii vortex.

Detailed shape. Due to limitations of the vortex method, unfortunately we are unable to determine many conjectured properties on the shape of the Sadovskii vortex patch.

Conjecture 1. The bounded open set Ω defining a Sadovskii vortex patch (as described in Theorem 1.1) is connected, convex, and makes the right angle with the horizontal axis.

It seems that in general, one needs certain regularity of the corresponding vorticity function in order to prove connectedness of maximizers from variational method (e.g. see [40, 12]). Our discontinuous vorticity function f in (1.5) is not regular enough to deduce Ω consists of a single connected component.

Moreover, while several references claim to prove that the touching angle of Ω with the horizontal angle should be 90 degrees using formal expansions of the velocity on the boundary ([100, 101, 81]), it is not even clear that the boundary function l(s) given in Theorem 1.1 defines an angle; it might keep oscillating as $s \to 0^+$, although the size of the oscillations should go to zero from continuity of l up to the boundary. However, under the assumptions that the angle is well-defined and l is *smooth up to the boundary* (not merely continuous), it can be made rigorous that the angle must be either 0, 90, or 180 degrees (cf. [37]). We also remark that proving the existence of V-states (uniformly rotating vortex patches) having 90 degree corners seems to remain open ([53, 44]).

Furthermore, the existence of Sadovskii vortex accompanying vortex sheets on its boundary seems to be a very difficult open problem.

Variational properties. There is a possibility that uniqueness still holds for small impulse $\mu > 0$ which would give stability for symmetric perturbations as in the case of Lamb–Chaplygin dipole [2] and Hill's vortex ring [29]. Formally, we have the following conjecture.

Conjecture 2. There exists a universal constant $\mu^* > 0$ such that, for all $0 < \mu \le \mu^*$, the set of maximizers $S_{\mu,1,1}$ coincides with x_1 -translation and rescaling of a single patch, which then can be referred to as the Sadovskii vortex patch. Furthermore, for all $\mu > \mu^*$, $S_{\mu,1,1}$ only contains patches which are separated from the horizontal axis.

This conjecture is supported by the fact that so far no one was able to numerically find two different shapes of Sadovskii vortex patch, and if true, it would have significant consequences for the initial value problem of the 2D Euler equations. Furthermore, this would also explain why one could not continue, even at the numerical level, the bifurcation curve consisting of vortex dipoles beyond Sadovskii vortex patch. We remark that the analogous conjecture for the vortex rings was established recently in [29].

Related to this, Pierrehumbert [76] and Saffman-Szeto [80, p2342] discussed the following more general problem.

Question 1. Is there a traveling wave $\omega = \mathbf{1}_{\Omega} - \mathbf{1}_{\Omega_{-}}$ such that Ω is a bounded open set of \mathbb{R}^{2}_{+} satisfying

$$|\Omega| = 1$$
 while $\mu := \int_{\Omega} x_2 d\mathbf{x} < \mu^*$?

It is possible that there are no such traveling wave patches, even if we do not require them to be maximizers of the energy in a suitable class. We note that Pierrehumbert calculated up to $\mu = 0.29$ and anticipated that $\mu^* \simeq 0.26$ is the possible limit. It almost matches to the limiting number in [100, Table I], after scaling.

1.6. **Outline of the paper.** The rest of the paper is organized as follows. In Section 2, we fix notations and collect several technical estimates. Existence of maximizers of patch-type for the variational problem is established in Section 3. Then in Section 4, we show that when the impulse is sufficiently small, the corresponding flux constant is zero and therefore the maximizing patch touches the axis. Lastly, in Section 5, we consider the variational problem without a mass bound on the admissible class, and show that the maximizer coincides with that of the variational problem with a mass bound.

2. Preliminaries

2.1. Notations.

• We use the following notations for the norms in \mathbb{R}^2_+ ,

$$||f||_1 := ||f||_{L^1(\mathbb{R}^2_+)}$$
 (Mass), $||f||_2 := ||f||_{L^2(\mathbb{R}^2_+)}$, $||x_2f||_1 := \int_{\mathbb{R}^2} x_2 |f(\mathbf{x})| d\mathbf{x}$ (Impulse).

• For each R > 0 and $\mathbf{x} \in \mathbb{R}^2$, we define the open ball

$$B_R(\mathbf{x}) := \{ \mathbf{y} \in \mathbb{R}^2 : |\mathbf{y} - \mathbf{x}| < R \}.$$

• For each measurable set $A \in \mathbb{R}^2$, we define the characteristic function $\mathbf{1}_A : \mathbb{R}^2 \to \mathbb{R}$ as

$$\mathbf{1}_{A}(\mathbf{x}) := \begin{cases} 1 & \text{if } \mathbf{x} \in A, \\ 0 & \text{if } \mathbf{x} \in A^{c} \end{cases}$$

• For each function $f: \mathbb{R}^2_+ \to \mathbb{R}$, we can define its odd extension to \mathbb{R}^2 , denoted by $\widetilde{f}: \mathbb{R}^2 \to \mathbb{R}$ and defined as

$$\widetilde{f}(x_1, x_2) := \begin{cases} f(x_1, x_2) & \text{if } x_2 > 0, \\ -f(x_1, -x_2) & \text{if } x_2 < 0, \\ 0 & \text{if } x_2 = 0. \end{cases}$$

2.2. Some technical estimates.

We will collect some elementary lemmas which will be used in the sequel. In \mathbb{R}^2_+ , we denote the Green's function $G: \mathbb{R}^2_+ \times \mathbb{R}^2_+ \to \mathbb{R}$ defined as

(2.1)
$$G(\mathbf{x}, \mathbf{y}) := \frac{1}{4\pi} \log \left(1 + \frac{4x_2 y_2}{|\mathbf{x} - \mathbf{y}|^2} \right) > 0, \quad \text{for } \mathbf{x} \neq \mathbf{y}.$$

The following lemma implies that the Green function is integrable for each $x \in \mathbb{R}^2_+$, in the domain $\{y \in \mathbb{R}^2_+ : y_2 < \alpha\}$ for any $\alpha > 0$. This lemma will be used later in the proof of Lemma 4.4 and Lemma 4.6.

Lemma 2.1. There exists a constant C > 0 such that, for each $\alpha > 0$ and $\mathbf{x} = (x_1, x_2) \in \mathbb{R}^2_+$, we have

(2.2)
$$\int_{0 < y_2 \le \alpha} G(\mathbf{x}, \mathbf{y}) \, d\mathbf{y} \le C \, x_2^{1/2} \alpha^{3/2}.$$

Proof. For each $\alpha > 0$ and $\mathbf{x} = (x_1, x_2) \in \mathbb{R}^2_+$, we have

$$\int_{0 < y_2 \le \alpha} G(\mathbf{x}, \mathbf{y}) \, d\mathbf{y} \, \leq \, \frac{1}{4\pi} \int_{0 < y_2 \le \alpha} \log \left(1 + \frac{4\alpha x_2}{y_1^2} \right) d\mathbf{y} \, \leq \, \frac{\alpha (4\alpha x_2)^{\frac{1}{2}}}{4\pi} \int_{\mathbb{R}} \log \left(1 + \frac{1}{t^2} \right) dt \, \lesssim \, x_2^{1/2} \alpha^{3/2},$$

using the fact that

$$\int_0^\infty \log\left(1 + \frac{1}{x^2}\right) d\mathbf{x} < \infty$$

which is the consequence of integration by parts.

Definition 2.2. For each $\omega \in (L^1 \cap L^2)(\mathbb{R}^2_+)$, we can define the function $\mathcal{G}[\omega] : \mathbb{R}^2_+ \to \mathbb{R}$ as

$$\mathcal{G}[\omega](\mathbf{x}) := \int_{\mathbb{R}^2_+} G(\mathbf{x}, \mathbf{y}) \omega(\mathbf{y}) \ d\mathbf{y}.$$

The function $\mathcal{G}[\omega]$ is well defined by the following lemma, which is a slight variation of [2, Proposition 2.1]. We will prove that, for each $\omega \in (L^1 \cap L^2)(\mathbb{R}^2_+)$ with $||x_2\omega||_1 < \infty$, we get $\mathcal{G}[\omega] \in L^{\infty}(\mathbb{R}^2_+)$.

Lemma 2.3. There exists a constant $C_1 \in (0, \infty)$ such that, for each $\omega \in (L^1 \cap L^2)(\mathbb{R}^2_+)$, we have

(2.3)
$$\left| \int_{\mathbb{R}^2} G(\mathbf{x}, \mathbf{y}) \omega(\mathbf{y}) \, d\mathbf{y} \right| \leq C_1 \, x_2^{1/2} ||\omega||_1^{1/2} ||\omega||_2^{1/2} \quad \text{for each } \mathbf{x} = (x_1, x_2) \in \mathbb{R}^2_+.$$

If we additionally assume that $||x_2\omega||_1 < \infty$, then there exists an absolute constant $C_2 \in (0, \infty)$ such that

$$\left| \int_{\mathbb{R}^{2}_{+}} G(\mathbf{x}, \mathbf{y}) \omega(\mathbf{y}) \, d\mathbf{y} \right| \leq C_{2} \left(\|\omega\|_{1}^{1/3} \|\omega\|_{2}^{1/3} \|x_{2}\omega\|_{1}^{1/3} + \|\omega\|_{2}^{1/2} \|x_{2}\omega\|_{1}^{1/2} \right) \quad \text{for each } \mathbf{x} = (x_{1}, x_{2}) \in \mathbb{R}^{2}_{+},$$

and that

$$\left| \int_{\mathbb{R}^{2}_{+}} \int_{\mathbb{R}^{2}_{-}} G(\mathbf{x}, \mathbf{y}) \omega(\mathbf{x}) \omega(\mathbf{y}) \, d\mathbf{x} d\mathbf{y} \right| \leq C_{2} \, \|\omega\|_{1} \|\omega\|_{2}^{1/2} \|x_{2}\omega\|_{1}^{1/2}.$$

Proof. The estimates (2.3) and (2.5) were borrowed from [2, Proposition 2.1]. To show (2.4), assume that $||x_2\omega||_1 < \infty$ and split the integral as

$$\left| \int_{\mathbb{R}^2_+} G(\mathbf{x}, \mathbf{y}) \omega(\mathbf{y}) \, d\mathbf{y} \right| \leq \left| \int_{|x-y| \geq x_7/2} \left| + \left| \int_{|x-y| < x_7/2} \right| \right| = (\mathrm{I}) + (\mathrm{II}).$$

For (I), using the inequality,

$$\log(1+t) \le t$$
 for each $t > 0$,

we have

(I)
$$\leq \int_{|\mathbf{x}-\mathbf{y}| > |\mathbf{x}_2|/2} \frac{x_2 y_2}{\pi |\mathbf{x} - \mathbf{y}|^2} |\omega(\mathbf{y})| d\mathbf{y} \leq \frac{4}{\pi x_2} ||x_2 \omega||_1 =: A.$$

On the other hand, by (2.3), we obtain

(I)
$$\lesssim x_2^{1/2} ||\omega||_1^{1/2} ||\omega||_2^{1/2} =: B.$$

In sum, we have

(I)
$$\lesssim A^{\frac{1}{3}}B^{\frac{2}{3}} \lesssim ||\omega||_1^{1/3}||\omega||_2^{1/3}||x_2\omega||_1^{1/3}.$$

For (II), we put $\mathcal{B} := B_{x_2/2}(\mathbf{x})$ and apply (2.3) to have

$$(II) = \left| \int_{\mathbb{R}^2} G(\mathbf{x}, \mathbf{y}) \Big(\omega \cdot \mathbf{1}_{\mathcal{B}} \Big) (\mathbf{y}) \, d\mathbf{y} \right| \le C x_2^{1/2} \|\omega\|_{L^1(\mathcal{B})}^{1/2} \|\omega\|_{L^2(\mathcal{B})}^{1/2}$$

Here observe that

$$\|\omega\|_{L^{1}(\mathcal{B})} \leq \frac{2}{x_{2}} \int_{\mathcal{B}} |y_{2}\omega(\mathbf{y})| d\mathbf{y} \leq \frac{2}{x_{2}} \|x_{2}\omega\|_{1}$$

due to the definition of \mathcal{B} . Therefore we have

(II)
$$\lesssim \|\omega\|_2^{1/2} \|x_2\omega\|_1^{1/2}$$

which completes the proof of (2.4).

The following lemma gives the relations between $\mathcal{G}[\omega]$ and ω . For the regularity argument of $\mathcal{G}[\omega]$, we refer to [68, Lemma 8.1], [69, Appendix 2.3] (see also [46], [38]).

Lemma 2.4. For each $\omega \in (L^1 \cap L^\infty)(\mathbb{R}^2_+)$, the function $\widetilde{\mathcal{G}}[\omega]$, which is the odd extension of $\mathcal{G}[\omega]$ in Definition 2.2, satisfies $\widetilde{\mathcal{G}}[\omega] \in C^{1,r}(\mathbb{R}^2)$ for each $r \in (0,1)$ and

$$\widetilde{\mathcal{G}[\omega]} \in H^2_{loc}(\mathbb{R}^2) \text{ with } -\Delta \, \widetilde{\mathcal{G}[\omega]} = \widetilde{\omega} \text{ a.e. in } \mathbb{R}^2$$

where $\widetilde{\omega}$ is the odd extension of ω to \mathbb{R}^2 .

Proof. It is due to the relation

$$\widetilde{\mathcal{G}[\omega]} = \psi[\widetilde{\omega}] \quad \text{in } \mathbb{R}^2$$

and potential estimates of ψ .

We also obtain the asymptotic behavior of $\mathcal{G}[\omega]$ and a different representation of the kinetic energy which were borrowed from [2, Proposition 2.2]).

Lemma 2.5. For each $\omega \in (L^1 \cap L^2)(\mathbb{R}^2_+)$ with $||x_2\omega||_1 < \infty$, we have

$$\lim_{|\mathbf{x}|\to\infty} \mathcal{G}[\omega](\mathbf{x}) = 0.$$

Moreover, we have

$$\frac{1}{2} \int_{\mathbb{R}^2} \mathcal{G}[\omega] \omega \, d\mathbf{x} = \frac{1}{4} \int_{\mathbb{R}^2} \left| \mathbf{u} \right|^2 \, d\mathbf{x}$$

where $\mathbf{u} = k * \widetilde{\omega}$. Here $\widetilde{\omega}$ is the odd extension of ω to \mathbb{R}^2 , and the kernel k is given in the equation 1.1.

Proof. By (2.4) in Lemma 2.3, there exists a constant C > 0 such that, for any $\omega \in (L^1 \cap L^2)(\mathbb{R}^2_+)$ with $||x_2\omega||_1 < \infty$ and for the sequence $\{\omega_n\} = \{\omega \cdot \mathbf{1}_{B_n(0)}\}$, we have

$$\left|\mathcal{G}[\omega](\mathbf{x})\right| \leq \left|\mathcal{G}[\omega - \omega_n](\mathbf{x})\right| + \left|\mathcal{G}[\omega_n](\mathbf{x})\right| \leq C\left(C_n + \frac{|\mathbf{x}|}{(|\mathbf{x}| - n)^2}||x_2\omega||_1\right) \quad \text{for each } |\mathbf{x}| > n,$$

where

$$C_n := \|\omega - \omega_n\|_1^{1/3} \|\omega - \omega_n\|_2^{1/3} \|x_2(\omega - \omega_n)\|_1^{1/3} + \|\omega - \omega_n\|_2^{1/2} \|x_2(\omega - \omega_n)\|_1^{1/2}.$$

Therefore we have

$$\limsup_{|\mathbf{x}|\to\infty} |\mathcal{G}[\omega](\mathbf{x})| \leq C \cdot C_n \quad \text{ for each } n \geq 1.$$

Observe that $C_n \to 0$ as $n \to \infty$. As we take the limit $n \to \infty$, we obtain

$$\lim_{|\mathbf{x}|\to\infty} \mathcal{G}[\omega](\mathbf{x}) = 0.$$

This finishes the proof.

3. Existence of maximizers of patch-type

In this section, we will use the variational method to prove that there exists a patch-type maximizer of the kinetic energy in the admissible class whose setting is from [42] (see also [2]). Each maximizer will generate a traveling wave solution of (1.1). The main result of this section is Theorem 3.3.

3.1. Variational problem on exact impulse and mass bound.

Given $\mu, \nu, \lambda > 0$, we define an admissible set

$$K_{\mu,\gamma,\lambda} := \left\{ \omega \in L^1(\mathbb{R}^2_+) : \omega = \lambda \cdot \mathbf{1}_A \text{ a.e. where } A \subset \mathbb{R}^2_+ \text{ is (Lebesgue) measurable,} \right\}$$

$$||x_2\omega||_1 = \mu, \quad ||\omega||_1 \le \nu$$

on which its energy functional $E(\cdot): K_{\mu,\nu,\lambda} \to \mathbb{R}$ is defined as

$$E(\omega) := \frac{1}{2} \int_{\mathbb{R}^2_+} \mathcal{G}[\omega](\mathbf{x}) \omega(\mathbf{x}) d\mathbf{x} = \frac{1}{2} \int_{\mathbb{R}^2_+} \int_{\mathbb{R}^2_+} G(\mathbf{x}, \mathbf{y}) \omega(\mathbf{y}) \omega(\mathbf{x}) d\mathbf{y} d\mathbf{x}.$$

Note that, if ω has a bounded support, then Lemma 2.5 implies that

$$E(\omega) = \frac{1}{4} \int_{\mathbb{R}^2} |\mathbf{u}|^2 d\mathbf{x}$$

where $\mathbf{u} = k * \widetilde{\omega}$ with k from (1.1) and $\widetilde{\omega}$ is the odd extension of ω . We formulate the maximization problem by

$$I_{\mu,\nu,\lambda} := \sup_{\omega \in K_{\mu,\nu,\lambda}} E(\omega)$$

and define the set of maximizers as $S_{\mu,\nu,\lambda} := \{ \omega \in K_{\mu,\nu,\lambda} : E(\omega) = I_{\mu,\nu,\lambda} \}$.

Remark 3.1.

(i) Recall that, for each $(\mathbf{x}, \mathbf{y}) \in \mathbb{R}^2_+ \times \mathbb{R}^2_+$ with $\mathbf{x} \neq \mathbf{y}$, we have

$$G(\mathbf{x}, \mathbf{y}) = G(\mathbf{y}, \mathbf{x}) > 0.$$

It follows that, for each $\omega \in K_{\mu,\nu,\lambda}$, we have $E(\omega) \ge 0$ and $I_{\mu,\nu,\lambda} > 0$. Moreover, we get $I_{\mu,\nu,\lambda} < \infty$ by (2.5) in Lemma 2.3.

(ii) If $f_{\tau}: \mathbb{R}^2 \to \mathbb{R}^2$ denotes the translation of some fixed amount $\tau \in \mathbb{R}$ along the x_1 -axis, i.e.

$$f_{\tau}(x_1, x_2) = (x_1 + \tau, x_2),$$

then we have

$$G(\mathbf{x}, \mathbf{y}) = G(f_{\tau}(\mathbf{x}), f_{\tau}(\mathbf{y})).$$

Therefore, if $\omega \in K_{\mu,\nu,\lambda}$, then any translation $\omega_{\tau} := \omega \circ f_{\tau}$ satisfies

$$\omega_{\tau} \in K_{\mu,\nu,\lambda}$$
 and $E(\omega) = E(\omega_{\tau})$.

(iii) If $g_h : \mathbb{R}^2 \to \mathbb{R}^2$ denotes the lift by some fixed amount h > 0 along the x_2 -axis, i.e.

$$g_h(x_1, x_2) = (x_1, x_2 + h),$$

then for each $\omega \in K_{\mu,\nu,\lambda}$, we have

$$E(\omega) < E(\omega \circ g_{-h}).$$

(iv) For each $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2_+$ with $\mathbf{x} \neq \mathbf{y}$, we have

$$G(\mathbf{x}, \mathbf{y}) = G(r\mathbf{x}, r\mathbf{y})$$

for any r > 0. By this property, for each $\omega \in (L^1 \cap L^\infty)(\mathbb{R}^2_+)$ with $||x_2\omega||_1 < \infty$, the scaling map given by $\hat{\omega}(\mathbf{x}) := \omega(r\mathbf{x})$ gives the relation

$$E(\hat{\omega}) = r^{-4}E(\omega)$$
 for each $r > 0$.

(v) As in [42] (and also in [2]), we can reduce the maximization problem to the case containing a single parameter. Indeed, for each $\omega \in K_{\mu,\nu,\lambda}$, consider the scaling

$$\hat{\omega}(\mathbf{x}) = \lambda^{-1} \cdot \omega(\lambda^{-1/2} v^{1/2} \mathbf{x}).$$

Observe that $\hat{\omega} \in K_{M,1,1}$ where $M = \lambda^{1/2} v^{-3/2} \mu > 0$ with the energy $E(\hat{\omega}) = v^{-2} E(\omega)$. Here, given $\mu, \nu, \lambda > 0$, the map $\omega \mapsto \hat{\omega}$ is a bijective map from $K_{\mu,\nu,\lambda}$ onto $K_{M,1,1}$, and from $K_{\mu,\nu,\lambda}$ onto $K_{M,1,1}$. Therefore, the two maximization problems below are congruent with each other,

$$I_{\mu,\nu,\lambda} = \sup_{\omega \in K_{\mu,\nu,\lambda}} E(\omega)$$
 and $I_{M,1,1} = \sup_{\omega \in K_{M,1,1}} E(\omega)$,

with the relation $I_{M,1,1} = v^{-2}I_{\mu,\nu,\lambda}$. We hereafter abbreviate the notations as

$$K_M := K_{M,1,1}, \quad I_M := I_{M,1,1}, \quad S_M := S_{M,1,1}.$$

3.2. Existence of maximizers.

For convenience, we start with a definition that will be frequently used in a sequel.

Definition 3.2. We say that a nonnegative measurable function $f: \mathbb{R}^2_+ \to \mathbb{R}$ satisfies the Steiner symmetry condition if

$$f(x_1, x_2) = f(-x_1, x_2)$$
 for each $\mathbf{x} \in \mathbb{R}^2_+$,

and for each fixed $x_2 > 0$, the function $f(x_1, x_2)$ is non-increasing in $x_1 \ge 0$. We also say that a measurable set $A \subset \mathbb{R}^2_+$ satisfies the Steiner symmetry condition if the function $g: \mathbb{R}^2_+ \to \mathbb{R}$ given by $g:= \mathbf{1}_A$ satisfies the Steiner symmetry condition.

The main theorem of Subsection 3.2 is Theorem 3.3 below. We will prove Theorem 3.3 at the end of Subsection 3.2 after the proof of several key lemmas. Each lemma is quite classical and stems from [29, 2].

Theorem 3.3. For any M > 0, we have the following properties.

- (i) The set S_M , which is the set of maximizers in K_M , is nonempty.
- (ii) For each $\omega \in S_M$, there exist some constants W > 0, $\gamma \ge 0$ that are uniquely determined by ω satisfying

(3.1)
$$\omega = \mathbf{1}_{\left\{\mathbf{x} \in \mathbb{R}^2_+ : \mathcal{G}[\omega](\mathbf{x}) - Wx_2 - \gamma > 0\right\}} \quad a.e. \quad in \quad \mathbb{R}^2_+.$$

(iii) Each $\omega \in S_M$ has a bounded support and satisfies the Steiner symmetry condition in Definition 3.2 up to translation. More precisely, for each $\omega \in S_M$, there exist $\tau \in \mathbb{R}$ and an open bounded set $A \subset \mathbb{R}^2_+$ satisfying the Steiner symmetry condition such that, for the translation ω_{τ} given by $\omega_{\tau}(\cdot) := \omega(\cdot + (\tau, 0))$, we have

$$\omega_{\tau} = \mathbf{1}_{A}$$
 a.e. in \mathbb{R}^{2}_{+} .

(iv) For each $\omega \in S_M$, if we have $||\omega||_1 < 1$, then we get $\gamma = 0$, where γ is given in (ii).

Note that we have the following corollary by the scaling in Remark 3.5.

Corollary 3.4. For any μ , ν , $\lambda > 0$, the set $S_{\mu,\nu,\lambda}$ is nonempty. For each maximizer $\omega \in S_{\mu,\nu,\lambda}$, there exist some constants W > 0, $\gamma \geq 0$ that are uniquely determined by ω satisfying

$$\omega = \lambda \cdot \mathbf{1}_{\Omega}$$
 a.e. in \mathbb{R}^2_+ where $\Omega = \{ \mathbf{x} \in \mathbb{R}^2_+ : \mathcal{G}[\omega](\mathbf{x}) - Wx_2 - \gamma > 0 \}.$

Moreover, each maximizer satisfies the Steiner symmetry condition in Definition 3.2 up to translation. In addition, if we have $\|\omega\|_1 < \nu$, then we get $\gamma = 0$.

Remark 3.5. Let $\mu, \nu, \lambda > 0$. Recall that the scaling map $\omega \mapsto \hat{\omega}$ given by

$$\hat{\omega}(\mathbf{x}) = \lambda^{-1} \cdot \omega(\lambda^{-1/2} \nu^{1/2} \mathbf{x})$$

is a bijective map from $S_{\mu,\nu,\lambda}$ onto S_M where $M = \lambda^{1/2} \nu^{-3/2} \mu > 0$. Moreover, under the scaling map $\omega \mapsto \hat{\omega}$, for the constants $W, \hat{W} > 0$ and $\gamma, \hat{\gamma} \ge 0$ satisfying

$$\omega = \lambda \cdot \mathbf{1}_{\left\{\mathcal{G}[\omega] - Wx_2 - \gamma > 0\right\}} \quad \text{and} \quad \hat{\omega} = \mathbf{1}_{\left\{\mathcal{G}[\hat{\omega}] - \hat{W}x_2 - \hat{\gamma} > 0\right\}} \quad \text{a.e. in } \mathbb{R}^2_+,$$

one can easily check the relations

$$\hat{W} = W \cdot \lambda^{-1/2} \cdot \nu^{-1/2}$$
 and $\hat{\gamma} = \gamma \cdot \nu^{-1}$.

Existence of maximizers in a larger set.

For the proof of Theorem 3.3, for each M > 0 we define a larger admissible set $\widetilde{K}_M \supset K_M$ as

$$\widetilde{K}_M := \left\{ \omega \in L^1(\mathbb{R}^2_+) : 0 \le \omega \le 1 \text{ a.e. in } \mathbb{R}^2_+, \|x_2\omega\|_1 \le M, \|\omega\|_1 \le 1 \right\},$$

and

$$\widetilde{I}_M := \sup_{\omega \in \widetilde{K}_M} E(\omega) \geq I_M, \qquad \widetilde{S}_M := \left\{ \omega \in \widetilde{K}_M : E(\omega) = \widetilde{I}_M \right\}.$$

Note that K_M is a set of characteristic functions having full impulse M, and \widetilde{K}_M is a set of nonnegative, bounded functions for which the bound of impulse is M. We will prove the existence of a maximizer in \widetilde{K}_M and show that the maximizer lies actually on K_M . This would imply that $S_M = \widetilde{S}_M$, and therefore S_M is nonempty.

Lemma 3.6. For each M > 0, the set \widetilde{S}_M is nonempty. In other words, there exists $\omega \in \widetilde{K}_M$ such that

$$E(\omega) = \widetilde{I}_M.$$

Proof. Consider a maximizing sequence $\{\omega_n\} \subset \widetilde{K}_M$, i.e.

$$E(\omega_n) \to \widetilde{I}_M$$
 as $n \to \infty$.

Since $\|\omega_n\|_{L^2(\mathbb{R}^2_+)} \leq \|\omega_n\|_1^{1/2} \leq 1$ for all $n \geq 1$, there exists a subsequence $\{\omega_{n_k}\} \subset \{\omega_n\}$ and $\omega \in L^2(\mathbb{R}^2_+)$ such that ω_{n_k} converges weakly to ω in $L^2(\mathbb{R}^2_+)$. For convenience, we assume $\omega_n \to \omega$ in $L^2(\mathbb{R}^2_+)$ as $n \to \infty$. It implies that $\omega \in \widetilde{K}_M$. In other words, we have

$$0 \le \omega \le 1$$
 a.e. in \mathbb{R}^2_+ , $||x_2\omega||_1 \le M$, and $||\omega||_1 \le 1$.

It remains to prove that $|E(\omega_n) - E(\omega)| \to 0$, which would imply that $E(\omega) = \widetilde{I}_M$ and $\omega \in \widetilde{S}_M$. The convergence $E(\omega_n) \to E(\omega)$ is obtained from the following lemmas, which are borrowed from [2]. Lemma 3.7 is a standard argument for the Steiner symmetrization which is well-known from previous works; see, e.g., [40, Appendix I], [94, p.1053].

Lemma 3.7. [2, Proposition 3.1] For $\omega \ge 0$ satisfying $\omega \in L^1 \cap L^2(\mathbb{R}^2_+)$ and $x_2\omega \in L^1(\mathbb{R}^2_+)$, there exists $\omega^* \ge 0$ such that $\omega^*(x_1, x_2) = \omega^*(-x_1, x_2)$, and for each x > 0, the function $\omega^*(x_1, x_2)$ is non-increasing in $x_1 > 0$. Moreover,

$$\|\omega^*\|_q = \|\omega\|_q \quad 1 \le q \le 2,$$

 $\|x_2\omega^*\|_1 = \|x_2\omega\|_1,$
 $E(\omega^*) \ge E(\omega).$

Remark 3.8. ω^* in Lemma 3.7 is defined to satisfy $|\{\omega > \alpha\}| = |\{\omega^* > \alpha\}|$ for all $\alpha \ge 0$. It implies that $0 \le \omega^* \le 1$. In particular, ω^* can be chosen to satisfy the Steiner symmetry condition in Definition 3.2.

By Lemma 3.7, we may assume that the maximizing sequence ω_n and the weak limit ω satisfy the Steiner symmetry condition in Definition 3.2. Then the convergence $E(\omega_n) \to E(\omega)$ is obtained by the Lemma 3.9 below.

Lemma 3.9. [2, Lemma 3.5] Let $\{\omega_n\}$ be a sequence such that

$$\sup_{n\geq 1} \left\{ \|\omega_n\|_{L^1 \cap L^2} + \|x_2 \omega_n\|_{L^1} \right\} < \infty,$$

$$\omega_n \rightharpoonup \omega \quad in L^2(\mathbb{R}^2_+) \quad as \ n \to \infty.$$

Assume that each ω_n satisfies the Steiner symmetry condition. Then, $E(\omega_n) \to E(\omega)$ as $n \to \infty$.

As we obtained that $E(\omega) = \widetilde{I}_M$, we complete the proof of Lemma 3.6.

Maximizers in smaller set.

Lemma 3.10. For each M > 0, we have

$$\widetilde{S}_M = S_M$$
.

Proof. Let M > 0. It suffices to show that $\widetilde{S}_M \subset K_M$. As the set \widetilde{S}_M is nonempty by Lemma 3.6, we choose any

We will first prove that $||x_2\omega||_1 = M$. Suppose that we have $||x_2\omega||_1 < M$. Consider the translation of ω along x_2 -axis of amount $\tau > 0$, say ω_{τ} . Then for small $\tau > 0$, we still have $\|x_2\omega_{\tau}\|_1 < M$, and by Remark3.1-(iii) we have

$$E(\omega_{\tau}) > E(\omega)$$
,

which contradicts our assumption that $\omega \in \widetilde{S}_M$. Therefore we have $||x_2\omega||_1 = M$.

It remains to show that ω is a characteristic function. In other words, we will show that

$$\left| \{ 0 < \omega < 1 \} \right| = 0,$$

which would imply that, for some measurable set $\Omega \subset \mathbb{R}^2_+$, we have $\omega = \mathbf{1}_{\Omega}$ a.e. in \mathbb{R}^2_+ . We suppose that

$$|\{0 < \omega < 1\}| > 0.$$

Then there exists a small constant $\delta_0 > 0$ such that

$$|\{\delta_0 < \omega < 1 - \delta_0\}| > 0.$$

Observe that $|\{\delta_0 < \omega < 1 - \delta_0\}| < \infty$ due to $||\omega||_1 \le 1$. We can choose two functions $h_1, h_2 \in (L^1 \cap L^\infty)(\mathbb{R}^2_+)$ such that

$$supp(h_i) \subset \{\delta_0 < \omega < 1 - \delta_0\}$$

for each $i \in \{1, 2\}$ satisfying

$$\int_{\mathbb{R}^{2}_{+}} h_{1} d\mathbf{x} = 1, \quad \int_{\mathbb{R}^{2}_{+}} x_{2} h_{1} d\mathbf{x} = 0,$$

$$\int_{\mathbb{R}^2_+} h_2 \ d\mathbf{x} = 0, \quad \int_{\mathbb{R}^2_+} x_2 h_2 \ d\mathbf{x} = 1.$$

So far, note that $\delta_0 > 0$ and $h_1, h_2 \in L^1 \cap L^\infty$ are fixed. For an arbitrary $\delta \in (0, \delta_0)$, we consider an arbitrary function $h \in (L^1 \cap L^\infty)(\mathbb{R}^2_+)$ such that $||x_2h||_1 < \infty$ and

$$h \ge 0$$
 on $\{0 \le \omega \le \delta\}$,
 $h \le 0$ on $\{1 - \delta \le \omega \le 1\}$.

We denote

$$\eta := h - \left(\int_{\mathbb{R}^2_+} h \, d\mathbf{x} \right) h_1 - \left(\int_{\mathbb{R}^2_+} x_2 h \, d\mathbf{x} \right) h_2.$$

Then we get $\eta \in (L^1 \cap L^{\infty})(\mathbb{R}^2_+)$ and

$$\int_{\mathbb{R}^2_+} \eta \ d\mathbf{x} = \int_{\mathbb{R}^2_+} x_2 \eta \ d\mathbf{x} = 0.$$

(Claim 1) There exists a small constant $\epsilon_0 > 0$ such that, for each $\epsilon \in (0, \epsilon_0)$, we have $\omega + \epsilon \eta \in \widetilde{K}_M$.

First, we will show that there exists a small constant $\epsilon_0 > 0$ such that, for each $\epsilon \in (0, \epsilon_0)$, we have $0 \le \omega + \epsilon \eta \le 1$. Observe that, on $\{0 \le \omega \le \delta_0\}$, we have $h_1 = h_2 = 0$ and thus $\eta = h \ge 0$ due to our assumption on h. Therefore $\omega + \epsilon \eta \ge 0$ on $\{0 \le \omega \le \delta_0\}$. And by choosing small $\epsilon_0 > 0$, we have

$$0 \le \omega + \epsilon \eta \le \delta_0 + \epsilon_0 \|\eta\|_{L^{\infty}} \le 1$$
 for each $\epsilon \in (0, \epsilon_0)$,

on the set $\{0 \le \omega \le \delta_0\}$. Similarly, for each $\epsilon \in (0, \epsilon_0)$, we have $0 \le \omega + \epsilon \eta \le 1$ on the set $\{1 - \delta_0 \le \omega \le 1\}$. Lastly, by choosing $\epsilon_0 < \delta_0 / \|\eta\|_{L^{\infty}}$, we obtain

$$0 \le \omega + \epsilon \eta \le 1$$
 for each $\epsilon \in (0, \epsilon_0)$

on the set $\{\delta_0 < \omega < 1 - \delta_0\}$. As we have $0 \le \omega + \epsilon \eta \le 1$ for each $\epsilon < \epsilon_0$, we have

$$\|\omega + \epsilon \eta\|_1 = \int (\omega + \epsilon \eta) d\mathbf{x} = \int \omega d\mathbf{x} \le 1 \quad \text{and} \quad \|x_2(\omega + \epsilon \eta)\|_1 = \int x_2(\omega + \epsilon \eta) d\mathbf{x} = \int x_2 \omega d\mathbf{x} \le M,$$

and therefore $\omega + \epsilon \eta \in \widetilde{K}_M$. This completes the proof of (*Claim 1*).

Using the assumption that $\omega \in \widetilde{S}_M$, the following function $F:[0,\epsilon_0)\to \mathbb{R}$ attains its maximum at $\epsilon=0$,

$$F(\epsilon) := E(\omega + \epsilon \eta) = \frac{1}{2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} G(\mathbf{x}, \mathbf{y}) \Big(\omega(\mathbf{y}) + \epsilon \eta(\mathbf{y}) \Big) \Big(\omega(\mathbf{x}) + \epsilon \eta(\mathbf{x}) \Big) d\mathbf{y} d\mathbf{x}.$$

Observe that F is a polynomial of degree 2. Clearly, $F'(0) \le 0$. By defining the constants

$$W := \int_{\mathbb{R}^2_+} \mathcal{G}[\omega] h_2 d\mathbf{x} \quad \text{and} \quad \gamma := \int_{\mathbb{R}^2_+} \mathcal{G}[\omega] h_1 d\mathbf{x},$$

we have

$$0 \geq F'(0) = \int_{\mathbb{R}^{2}_{+}} \int_{\mathbb{R}^{2}_{+}} G(\mathbf{x}, \mathbf{y}) \omega(\mathbf{x}) \eta(\mathbf{y}) \, d\mathbf{y} d\mathbf{x}$$

$$= \int_{\mathbb{R}^{2}_{+}} \mathcal{G}[\omega](\mathbf{y}) \, \eta(\mathbf{y}) \, d\mathbf{y}$$

$$= \int_{\mathbb{R}^{2}_{+}} \mathcal{G}[\omega](\mathbf{y}) \left[h(\mathbf{y}) - \left(\int_{\mathbb{R}^{2}_{+}} h(\mathbf{x}) \, d\mathbf{x} \right) h_{1}(\mathbf{y}) - \left(\int_{\mathbb{R}^{2}_{+}} x_{2} h(\mathbf{x}) \, d\mathbf{x} \right) h_{2}(\mathbf{y}) \right] d\mathbf{y}$$

$$= \int_{\mathbb{R}^{2}_{+}} \mathcal{G}[\omega](\mathbf{x}) \, h(\mathbf{x}) \, d\mathbf{x} - \left(\int_{\mathbb{R}^{2}_{+}} \mathcal{G}[\omega](\mathbf{y}) \, h_{2}(\mathbf{y}) \, d\mathbf{y} \right) \cdot \int_{\mathbb{R}^{2}_{+}} x_{2} h(\mathbf{x}) \, d\mathbf{x} - \left(\int_{\mathbb{R}^{2}_{+}} \mathcal{G}[\omega](\mathbf{y}) \, h_{1}(\mathbf{y}) \, d\mathbf{y} \right) \cdot \int_{\mathbb{R}^{2}_{+}} h(\mathbf{x}) \, d\mathbf{x}$$

$$= \int_{\mathbb{R}^{2}_{+}} \left(\mathcal{G}[\omega](\mathbf{x}) - W x_{2} - \gamma \right) h(\mathbf{x}) \, d\mathbf{x}$$

Then we split the integral into three terms:

$$0 \geq \int_{\mathbb{R}^2_+} \left(\mathcal{G}[\omega] - W x_2 - \gamma \right) \cdot h \; d\mathbf{x} = \int_{0 \leq \omega \leq \delta} + \int_{\delta < \omega < 1 - \delta} + \int_{1 - \delta \leq \omega \leq 1}.$$

Recall that, for each $\delta \in (0, \delta_0)$, the function $h \in (L^1 \cap L^\infty)(\mathbb{R}^2_+)$ is arbitrarily chosen just to satisfy

$$h \ge 0$$
 on $\{0 \le \omega \le \delta\}$ and $h \le 0$ on $\{1 - \delta \le \omega \le 1\}$.

Therefore, for any $\delta \in (0, \delta_0)$, we obtain that

$$\begin{cases} \mathcal{G}[\omega](x) - Wx_2 - \gamma \le 0 & \text{a.e.} & \text{on } \{0 \le \omega \le \delta\}, \\ \mathcal{G}[\omega](x) - Wx_2 - \gamma = 0 & \text{a.e.} & \text{on } \{\delta < \omega < 1 - \delta\}, \\ \mathcal{G}[\omega](x) - Wx_2 - \gamma \ge 0 & \text{a.e.} & \text{on } \{1 - \delta \le \omega \le 1\}. \end{cases}$$

Due to $\{0 < \omega < 1\} = \bigcup_{\delta \in (0,\delta_0)} \{\delta < \omega < 1 - \delta\}, \text{ we get }$

$$\begin{cases} \mathcal{G}[\omega](x) - Wx_2 - \gamma \le 0 & \text{a.e.} & \text{on } \{ \omega = 0 \}, \\ \mathcal{G}[\omega](x) - Wx_2 - \gamma = 0 & \text{a.e.} & \text{on } \{ 0 < \omega < 1 \}, \\ \mathcal{G}[\omega](x) - Wx_2 - \gamma \ge 0 & \text{a.e.} & \text{on } \{ \omega = 1 \}. \end{cases}$$

So we obtain $\{0 < \omega < 1\} \subset \{\mathcal{G}[\omega] - Wx_2 - \gamma = 0\}$. Define $\Psi(\mathbf{x}) := \mathcal{G}[\omega](\mathbf{x}) - Wx_2 - \gamma$. It remains to prove that

$$\{\Psi = 0\} \subset \{\omega = 0\}$$
 a.e. in \mathbb{R}^2_+ ,

which would imply that

$$\{0 < \omega < 1\} \subset \{\omega = 0\}$$
 a.e. in \mathbb{R}^2_+ ,

which contradicts to our assumption that $|\{0 < \omega < 1\}| > 0$.

(Claim 2) For each $\omega \in \widetilde{S}_M$, we have $\{ \Psi = 0 \} \subset \{ \omega = 0 \}$ a.e. in \mathbb{R}^2_+ .

By Lemma 2.4, we have $\Psi \in H^2_{loc}(\mathbb{R}^2_+)$ and $-\Delta \Psi = \omega$ in \mathbb{R}^2_+ . Therefore we have

$$\{ \Psi = 0 \} \subset \{ \nabla \Psi = 0 \} \subset \{ \Delta \Psi = 0 \} = \{ \omega = 0 \}.$$

This completes the proof of (Claim 2) and we finish the proof of Lemma 3.10.

Every maximizer generates a traveling wave solution.

Note that Lemma 3.6 and Lemma 3.10 imply that the set S_M is nonempty. Proposition 3.11 says that each maximizer in S_M generates a traveling wave solution to (1.1).

Proposition 3.11. For any M>0 and for each $\omega\in S_M$, there exist some constants $W,\gamma\geq 0$ that are uniquely determined by ω satisfying that

$$\omega = \mathbf{1}_{\left\{\mathbf{x} \in \mathbb{R}^2_+ : \mathcal{G}[\omega](\mathbf{x}) - Wx_2 - \gamma > 0\right\}} \quad a.e. \quad in \quad \mathbb{R}^2_+.$$

Proof. Let M > 0. We choose any $\omega = \mathbf{1}_{\Omega} \in S_M$ for some measurable set $\Omega \subset \mathbb{R}^2_+$. For the proof, we first assume the existence of the constants $W, \gamma \in \mathbb{R}$ satisfying that

$$\Omega \ = \ \big\{ \mathcal{G}[\omega] - W x_2 - \gamma > 0 \big\}.$$

In (Claim 1), we will prove that W, γ are unique. The existence and non-negativity of such constants will be proved later in (Claim 2) and (Claim 3) each.

(Claim 1) If there exist some constants $W, \gamma \in \mathbb{R}$ satisfying that $\Omega = \{ \mathcal{G}[\omega] - Wx_2 - \gamma > 0 \}$, then they are uniquely determined by ω .

Assume that we have $\Omega = \{ \mathcal{G}[\omega] - Wx_2 - \gamma > 0 \}$ for some constants $W, \gamma \in \mathbb{R}$. We choose any two points $\mathbf{y}, \mathbf{z} \in \partial \Omega \cap \mathbb{R}^2_+$ such that $y_2 \neq z_2$. Observe that

$$\partial\Omega\cap\mathbb{R}^2_+\subset\left\{\mathcal{G}[\omega]-Wx_2-\gamma=0\right\}$$

by the continuity of $G[\omega]$. Therefore

$$\mathcal{G}[\omega](\mathbf{y}) - Wy_2 - \gamma = \mathcal{G}[\omega](\mathbf{z}) - Wz_2 - \gamma = 0$$

and thus

(3.2)
$$W = \frac{\mathcal{G}[\omega](\mathbf{y}) - \mathcal{G}[\omega](\mathbf{z})}{y_2 - z_2}, \quad \gamma = -\left(\frac{z_2}{y_2 - z_2}\right) \mathcal{G}[\omega](\mathbf{y}) + \left(\frac{y_2}{y_2 - z_2}\right) \mathcal{G}[\omega](\mathbf{z}).$$

It implies that W, γ are explicitly expressed by ω , so they are unique. This completes the proof of (Claim 1).

(*Claim 2*) There exist some constants $W, \gamma \in \mathbb{R}$ such that we have $\Omega = \{ \mathcal{G}[\omega] - Wx_2 - \gamma > 0 \}$.

The main part of the proof is to generate perturbation of $\omega = \mathbf{1}_{\Omega}$ in the set \widetilde{K}_M to discover the specific form of Ω . As we can see in the proof of the uniqueness of $W, \gamma \geq 0$, these constants depend on the boundary points of Ω . However, we yet only know that Ω is a measurable set, which means that we cannot use the definition of boundary points as usual. We naturally want to define boundary-like points for each measurable set which are called *exceptional points*. This notion appeared in several works including [35, 91, 63, 29].

Definition 3.12. [29, Definition 5.4] Let $U \subset \mathbb{R}^N$ be open. For any (Lebesgue) measurable set $E \subset U$, the density $D_e(E)$ of E is the collection of $\mathbf{x} \in U$ such that

$$\liminf_{r \searrow 0} \frac{|B_r(\mathbf{x}) \cap E|}{|B_r(\mathbf{x})|} = 1.$$

Similarly, define the dispersion $D_i(E)$ of E by the collection of $\mathbf{x} \in U$ such that

$$\limsup_{r\searrow 0}\frac{|B_r(\mathbf{x})\cap E|}{|B_r(\mathbf{x})|}=0.$$

Set $\mathcal{E}(E) := U \setminus (D_e(E) \cup D_i(E))$, the set of exceptional points.

The following two propositions say that the set of exceptional points has zero measure, but in most cases, it is a nonempty set. The proof of Proposition 3.13 can be found in [99, Theorem 7.13], and the proof of Proposition 3.14 is given in [29, Appendix B].

Proposition 3.13. [29, Lemma 5.5] $E = D_e(E)$ and $U \setminus E = D_i(E)$ a.e.

Proposition 3.14. [29, Lemma 5.6] Let $U \subset \mathbb{R}^N$ be a non-empty connected open set. Then for any measurable set $E \subset U$ with $|E|, |U \setminus E| \in (0, \infty]$, we have $\mathcal{E}(E) \neq \emptyset$.

By Proposition 3.14, we can obtain exceptional points $\mathbf{y}, \mathbf{z} \in \mathbb{R}^2_+$ s.t. $y_2 > z_2$, and $\{r_n\} \setminus 0$ such that

$$\begin{aligned} & \left| B_{r_n}(\mathbf{y}) \cap \Omega \right|, \ \left| B_{r_n}(\mathbf{y}) \cap \Omega^c \right| & \in (0, \infty), \\ & \left| B_{r_n}(\mathbf{z}) \cap \Omega \right|, \ \left| B_{r_n}(\mathbf{z}) \cap \Omega^c \right| & \in (0, \infty), \quad \text{for each } n \ge 1. \end{aligned}$$

To show this, we apply Proposition 3.14. We first choose l > 0 such that both $\Omega \cap \{x_2 > l\}$ and $\Omega \cap \{x_2 < l\}$ have finite and positive measures. In order to obtain such $\mathbf{y} \in \mathbb{R}^2_+$, put

$$U:=\{x_2>l\},\quad E:=\Omega\cap\{x_2>l\}$$

and choose $\mathbf{y} \in \mathcal{E}(E)$ so that $y_2 > l$. Then for any decreasing sequence of positive numbers $\{r_n\} \setminus 0$, we can simply check that

$$|B_{r_n}(\mathbf{y}) \cap \Omega|, |B_{r_n}(\mathbf{y}) \cap \Omega^c| \in (0, \infty), \text{ for each } n \ge 1.$$

Similarly, we can choose $\mathbf{z} \in \mathbb{R}^2_+$ such that $0 < z_2 < l$ and do the same process.

Define sequences of sets of positive measures near $\mathbf{y}, \mathbf{z} \in \mathbb{R}^2$ as

$$Y_n^+ := B_{r_n}(\mathbf{y}) \cap \Omega, \qquad Y_n^- := B_{r_n}(\mathbf{y}) \cap \Omega^c,$$

 $Z_n^+ := B_{r_n}(\mathbf{z}) \cap \Omega, \qquad Z_n^- := B_{r_n}(\mathbf{z}) \cap \Omega^c.$

and sequences of compactly supported and bounded functions as

$$f_n^{\pm} := \frac{1}{|Y_n^{\pm}|} \mathbf{1}_{Y_n^{\pm}}, \quad g_n^{\pm} := \frac{1}{|Z_n^{\pm}|} \mathbf{1}_{Z_n^{\pm}}.$$

These functions will play as approximations of Dirac delta at \mathbf{y} and \mathbf{z} each, as $n \to \infty$. Recalling the proof of (*Claim 1*) (see the identity (3.2)), define the constants

$$W := \left(\frac{1}{y_2 - z_2}\right) \mathcal{G}[\omega](\mathbf{y}) - \left(\frac{1}{y_2 - z_2}\right) \mathcal{G}[\omega](\mathbf{z}) \in \mathbb{R}, \qquad \gamma := -\left(\frac{z_2}{y_2 - z_2}\right) \mathcal{G}[\omega](\mathbf{y}) + \left(\frac{y_2}{y_2 - z_2}\right) \mathcal{G}[\omega](\mathbf{z}) \in \mathbb{R}.$$

As $n \to \infty$, we have

$$\int x_2 f_n^{\pm}(\mathbf{x}) d\mathbf{x} \to y_2 \quad \text{and} \quad \int x_2 g_n^{\pm}(\mathbf{x}) d\mathbf{x} \to z_2.$$

We denote $y_n^{\pm} := \int x_2 f_n^{\pm}(\mathbf{x}) d\mathbf{x} \in \mathbb{R}$ and $z_n^{\pm} := \int x_2 g_n^{\pm}(\mathbf{x}) d\mathbf{x} \in \mathbb{R}$. Moreover, by the continuity of $\mathcal{G}[\omega]$,

$$\int \mathcal{G}[\omega] f_n^{\pm} d\mathbf{x} \to \mathcal{G}[\omega](\mathbf{y}) \quad \text{and} \quad \int \mathcal{G}[\omega] g_n^{\pm} d\mathbf{x} \to \mathcal{G}[\omega](\mathbf{z})$$

as $n \to \infty$. In order to construct sequences converging to W and γ , define the sequences of coefficients as

$$a_n^{\pm} := \frac{y_n^{\mp}}{y_n^{\mp} - z_n^{\pm}}, \quad b_n^{\pm} := \frac{z_n^{\pm}}{y_n^{\mp} - z_n^{\pm}}, \quad c_n^{\pm} := \frac{1}{y_n^{\pm} - z_n^{\mp}}.$$

Note that they are all positive for large $n \ge 1$. Assume that $a_n^{\pm}, b_n^{\pm}, c_n^{\pm} > 0$, for all $n \ge 1$. Using these sequences, we construct functions as

$$h_{n,1}^{\pm} := a_n^{\pm} g_n^{\pm} - b_n^{\pm} f_n^{\mp}, \quad h_{n,2}^{\pm} := c_n^{\pm} (f_n^{\pm} - g_n^{\mp}).$$

These functions will be used to construct the perturbation for $\omega = \mathbf{1}_{\Omega}$ in the set \widetilde{K}_{M} . Observe that

$$\int h_{n,1}^{\pm} d\mathbf{x} = 1, \quad \int x_2 h_{n,1}^{\pm} d\mathbf{x} = 0,$$
$$\int h_{n,2}^{\pm} d\mathbf{x} = 0, \quad \int x_2 h_{n,2}^{\pm} d\mathbf{x} = 1.$$

Observe that the signs + and - in each coefficient and function are chosen to satisfy

$$h_{n,1}^+, h_{n,2}^+ \ge 0$$
 on Ω ,
 $h_{n,1}^+, h_{n,2}^+ \le 0$ on Ω^c ,

and

$$h_{n,1}^-, h_{n,2}^- \le 0$$
 on Ω ,
 $h_{n,1}^-, h_{n,2}^- \ge 0$ on Ω^c .

Now we are ready to prove that $\Omega = \{ \mathcal{G}[\omega] - Wx_2 - \gamma > 0 \}$ by using perturbation of the function $\omega = \mathbf{1}_{\Omega}$ in the set \widetilde{K}_M . Consider any function $h \in (L^1 \cap L^{\infty})(\mathbb{R}^2_+)$ such that $||x_2h||_1 < \infty$ and

$$h \ge 0$$
 on Ω^c , $h \le 0$ on Ω .

For any fixed function h, we choose the sign superscripts for $h_{n,1}^{\pm}$ and $h_{n,2}^{\pm}$ as follows. We define $h_{n,1}$ and $h_{n,2}$ to satisfy

$$h_{n,1} := \begin{cases} h_{n,1}^+ & \text{if} & \int h \ge 0, \\ h_{n,1}^- & \text{if} & \int h < 0, \end{cases}$$

and

$$h_{n,2} := \begin{cases} h_{n,2}^+ & \text{if} & \int x_2 h \ge 0, \\ h_{n,2}^- & \text{if} & \int x_2 h < 0. \end{cases}$$

Moreover we define

$$W_n := \int \mathcal{G}[\omega] h_{n,2} d\mathbf{x}, \quad \gamma_n := \int \mathcal{G}[\omega] h_{n,1} d\mathbf{x}.$$

Observe that $W_n \to W$ and $\gamma_n \to \gamma$ as $n \to \infty$. We set

$$\eta_n := h - \left(\int h \, d\mathbf{x}\right) h_{n,1} - \left(\int x_2 h \, d\mathbf{x}\right) h_{n,2}.$$

Naturally $\eta_n \le 0$ on Ω and $\eta_n \ge 0$ on Ω^c . Moreover, observe that $\int \eta_n = \int x_2 \eta_n = 0$. As $\eta_n \in L^{\infty}$, by the similar arguments in the proof of (*Claim 1*) of 3.10, we may show that there exists a small $\epsilon_0 > 0$ such that

$$\omega + \epsilon \eta_n \subset \widetilde{K}_M$$
 for each $\epsilon \in (0, \epsilon_0)$,

and using the fact that $E(\cdot)$ is maximized by ω in the admissible set \widetilde{K}_M , we obtain

$$0 \ge \frac{d}{d\epsilon} E(\omega + \epsilon \eta_n) \Big|_{\epsilon=0} = \int \mathcal{G}[\omega] \, \eta_n \, d\mathbf{x} = \int \left(\mathcal{G}[\omega] - W_n x_2 - \gamma_n \right) h \, d\mathbf{x},$$

for each $n \ge 1$. We can take the limit $n \to \infty$ to obtain

$$0 \geq \int (\mathcal{G}[\omega] - Wx_2 - \gamma) h \, d\mathbf{x}.$$

Since h is arbitrary, we have

$$G[\omega] - Wx_2 - \gamma \ge 0$$
 a.e. on Ω ,
 $G[\omega] - Wx_2 - \gamma \le 0$ a.e. on Ω^c .

Put $\Psi := \mathcal{G}[\omega] - Wx_2 - \gamma$ and observe

$$\{ \Psi < 0 \} \subset \Omega^c \subset \{ \Psi \le 0 \}.$$

Since $-\Delta \Psi = \omega$ a.e. in \mathbb{R}^2_+ , we have

$$\{\Psi = 0\} \subset \{\omega = 0\} = \Omega^c \text{ a.e.}$$

and thus $\{ \Psi \leq 0 \} = \Omega^c$. In conclusion, we have

$$\Omega = \{ \Psi > 0 \} = \{ \mathcal{G}[\omega] - Wx_2 - \gamma > 0 \}$$
 a.e.

This completes the proof of (*Claim 2*).

(*Claim 3*) The constants $W, \gamma \in \mathbb{R}$ are non-negative.

If $\gamma < 0$, observe that

$$\left\{\mathbf{x} \in \mathbb{R}^2_+ : |W|x_2 < -\gamma\right\} \subset \left\{\mathbf{x} \in \mathbb{R}^2_+ : Wx_2 < -\gamma\right\} = \left\{\mathcal{G}[\omega] - Wx_2 - \gamma > 0\right\} = \Omega.$$

It implies that $|\Omega| = \infty$, which contradicts to $|\Omega| \le 1$. Therefore we have $\gamma \ge 0$. If W < 0, we have

$$\left\{ \, \mathbf{x} \in \mathbb{R}_+^2 \, : \, -Wx_2 > \gamma \, \right\} \, \subset \, \left\{ \, \mathcal{G}[\omega] - Wx_2 - \gamma > 0 \, \right\} \, = \, \Omega,$$

which also contradicts to $|\Omega| \le 1$. It follows that $W \ge 0$. This completes the proof of (*Claim 3*) and we finish the proof of Proposition 3.11.

Lemma 3.15 below says that, if any maximizer in K_M does not attain the full mass, then the corresponding flux constant γ vanishes.

Lemma 3.15. For any M > 0 and for each $\omega \in S_M$, if we have $\|\omega\|_1 < 1$, then we get

$$\gamma = 0$$

where $\gamma = \gamma(\omega) \ge 0$ is given in Proposition 3.11.

Proof. Let M > 0. We choose any $\omega = \mathbf{1}_{\Omega} \in S_M$ where we have

$$\Omega = \left\{ \mathcal{G}[\omega] - Wx_2 - \gamma > 0 \right\}$$

where the constants $W, \gamma \ge 0$ are given in Proposition 3.11. We assume that $\|\omega\|_1 < 1$. We now follow the same process in the proof of (*Claim 2*) in Proposition 3.11 in a way that, instead of using η_n , we use the following function

$$\hat{\eta}_n := h - \Big(\int\limits_{23} x_2 h \ d\mathbf{x}\Big) h_{n,2}$$

with the same definitions for $h, h_{n,2}$ in the proof of (Claim 2) in Proposition 3.11. As $\|\omega\|_1 < 1$, we have

$$\|\omega + \epsilon \hat{\eta}_n\|_1 \le \|\omega\|_1 + \epsilon \|\hat{\eta}_n\|_1 \le 1$$

for small $\epsilon > 0$. Moreover, as $\hat{\eta}_n \in L^{\infty}$, for small $\epsilon > 0$ we have $0 \le \omega + \epsilon \hat{\eta}_n \le 1$, and due to $\int x_2 \hat{\eta}_n d\mathbf{x} = 0$, we get

$$||x_2(\omega + \epsilon \hat{\eta}_n)||_1 = \int_{\mathbb{R}^2_+} x_2 \omega \, d\mathbf{x} + \epsilon \int_{\mathbb{R}^2_+} x_2 \hat{\eta}_n \, d\mathbf{x} = M.$$

This implies that there exists a small $\epsilon_0 > 0$ such that

$$\omega + \epsilon \hat{\eta}_n \in \widetilde{K}_M$$

for each $\epsilon \in (0, \epsilon_0)$. Then we have

$$0 \ge \frac{d}{d\epsilon} E(\omega + \epsilon \hat{\eta}_n) = \int (\mathcal{G}[\omega] - W_n x_2) h \, d\mathbf{x}$$

with the same definition for W_n with the proof of (Claim 2) in Proposition 3.11. It follows that

$$\omega = \mathbf{1}_{\{\mathcal{G}[\omega]-Wx_2>0\}}$$
 a.e. in \mathbb{R}^2_+ .

By the uniqueness of the constant $\gamma \ge 0$ given in Proposition 3.11, we complete the proof of Lemma 3.15.

Positivity of traveling speed.

Remark 3.16. For any $\omega \in (L^1 \cap L^\infty)(\mathbb{R}^2_+)$, whose odd-extension $\widetilde{\omega}$ is a traveling wave solution of the equation (1.1), in the form

$$\omega(\mathbf{x}) = \omega_0(\mathbf{x} - (W, 0)t)$$

for some $\omega_0 \in (L^1 \cap L^\infty)(\mathbb{R}^2_+)$ having bounded support with some constant $W \in \mathbb{R}$, its traveling speed W is determined by ω_0 by the identity in [94, p.1062]:

$$(3.3) W = \|\omega_0\|_1^{-1} \cdot \int_{\mathbb{R}^2} \mathcal{G}[\omega_0]_{x_2} \cdot \omega_0 \, d\mathbf{x} = \|\omega_0\|_1^{-1} \cdot \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{1}{2\pi} \frac{x_2 + y_2}{|\mathbf{x} - \mathbf{y}^*|^2} \, \omega_0(\mathbf{y}) \, \omega_0(\mathbf{x}) \, d\mathbf{y} d\mathbf{x}$$

where $\mathbf{y}^* = (y_1, -y_2)$. Indeed, for the first equality, observe that

$$W\int_{\mathbb{R}^2_+}\omega_0\,d\mathbf{x}=\frac{d}{dt}\int_{\mathbb{R}^2_+}x_1\omega\,d\mathbf{x}=\int_{\mathbb{R}^2_+}x_1\omega_t\,d\mathbf{x}=-\int_{\mathbb{R}^2_+}x_1(\mathbf{u}\cdot\nabla\omega)\,d\mathbf{x}=\int_{\mathbb{R}^2_+}u_1\cdot\omega\,d\mathbf{x}$$

where we used integration by parts. The second equality is obtained by considering obvious cancellation.

Lemma 3.17. For any M > 0 and for each $\omega \in S_M$, we have

where $W = W(\omega)$ is given in Proposition 3.11.

Proof. Let M > 0 and choose any $\omega \in S_M$. We suppose W = 0. Then $\gamma > 0$ due to the assumption $\|\omega\|_1 \le 1$. By Lemma 2.5, it is clear that the set $\{\mathcal{G}[\omega] - \gamma > 0\}$ is bounded. Then we can derive W > 0 by the identity (3.3) in Remark 3.16, which contradicts our assumption that W = 0.

Bounded support of maximizers.

In the following lemma, we show that each maximizer ω has a bounded support.

Lemma 3.18. For any M > 0 and for each ω in S_M , the set $\{G[\omega] - Wx_2 - \gamma > 0\}$ is bounded, for the corresponding constant W, γ given in Proposition 3.11. Moreover, there exists a constant C > 0 such that, for any M > 0 and for each $\omega \in S_M$, we have

$$\left|\frac{\mathcal{G}[\omega](\mathbf{x})}{x_2}\right| \leq C\left(\epsilon^{-1}\|\omega\|_1 + \epsilon^{1/3}\|\omega\|_{L^1\left([x_1 - \epsilon, x_1 + \epsilon] \times (0, \infty)\right)}^{1/3}\right),$$

for any $\epsilon > 0$ and $\mathbf{x} = (x_1, x_2) \in \mathbb{R}^2_+$.

Remark 3.19. We can deduce from the estimate (3.4) in Lemma 3.18 below that $|\mathcal{G}[\omega](\mathbf{x})/x_2| \to 0$ as $|x_1| \to \infty$ where the convergence is uniform in x_2 . This estimate will be used later in the proofs of Theorem 4.2 and Lemma 4.7.

Proof of Lemma 3.18. Let M > 0. As S_M is nonempty, we choose any $\omega \in S_M$. By Proposition 3.11, we have

$$\omega = \mathbf{1}_{\{\mathcal{G}[\omega] - Wx_2 - \gamma > 0\}} \quad \text{in } \mathbb{R}^2_+$$

for some unique constants $W = W(\omega) > 0$ and $\gamma = \gamma(\omega) \ge 0$. By (2.4) in Lemma 2.3, we have $\mathcal{G}[\omega] \in L^{\infty}(\mathbb{R}^2_+)$. Therefore we obtain

$$\left\{ \mathcal{G}[\omega] - Wx_2 - \gamma > 0 \right\} \subset \left\{ \mathcal{G}[\omega] - Wx_2 > 0 \right\} \subset \left\{ x_2 < \frac{\|\mathcal{G}[\omega]\|_{L^{\infty}}}{W} \right\}.$$

It remains to prove that there exists a constant L > 0 such that

$$\left\{ \mathcal{G}[\omega] - Wx_2 - \gamma > 0 \right\} \subset \left\{ |x_1| \le L \right\}.$$

Observe that

$$\{\mathcal{G}[\omega] - Wx_2 - \gamma > 0\} \subset \{\mathcal{G}[\omega]/x_2 > W\}.$$

By the mean value theorem, for each $\mathbf{x} \in \mathbb{R}^2_+$, there exists a constant $c = c(\mathbf{x}) \in (0, x_2)$ such that

$$\frac{\mathcal{G}[\omega](\mathbf{x})}{x_2} = \frac{\mathcal{G}[\omega](\mathbf{x}) - \mathcal{G}[\omega](x_1, 0)}{x_2} = \mathcal{G}[\omega]_{x_2}(x_1, c).$$

Here we have

$$\mathcal{G}[\omega]_{x_2}(x_1,c) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \frac{(y_2-c)}{(y_1-x_1)^2 + (y_2-c)^2} \, \widetilde{\omega}(\mathbf{y}) \, d\mathbf{y}$$

where $\widetilde{\omega}$ is the odd-extension of ω to \mathbb{R}^2 . Note that, for each $\epsilon > 0$, we have

$$\left| \mathcal{G}[\omega]_{x_2}(x_1,c) \right| \lesssim \left| \int_{B^c} + \int_{B} \frac{1}{|(y_1,y_2) - (x_1,c)|} |\widetilde{\omega}(\mathbf{y})| \, d\mathbf{y} \right|$$

where $B := B_{\epsilon}(x_1, c)$. The first integral satisfies

$$\int_{B^c} \frac{|\widetilde{\omega}(\mathbf{y})|}{|(y_1, y_2) - (x_1, c)|} d\mathbf{y} \le \frac{2}{\epsilon} ||\omega||_1.$$

For the second integral, by Hölder's inequality, we have

$$\int_{B} \frac{|\widetilde{\omega}(\mathbf{y})|}{|(y_{1}, y_{2}) - (x_{1}, c)|} d\mathbf{y} \lesssim \left(\int_{0}^{\epsilon} r^{-1/2} dr \right)^{2/3} ||\omega||_{L^{3}(B)} = \left(\int_{0}^{\epsilon} r^{-1/2} dr \right)^{2/3} ||\omega||_{L^{1}(B)}^{1/3} \lesssim \epsilon^{1/3} ||\omega||_{L^{1}([x_{1} - \epsilon, x_{1} + \epsilon] \times (0, \infty))}^{1/3}.$$

In sum, we obtain (3.4) as we have

$$\left|\frac{\mathcal{G}[\omega](\mathbf{x})}{x_2}\right| \leq C_0 \left(\epsilon^{-1} \|\omega\|_1 + \epsilon^{1/3} \|\omega\|_{L^1\left([x_1 - \epsilon, x_1 + \epsilon] \times (0, \infty)\right)}^{1/3}\right)$$

for some absolute constant $C_0 > 0$. Moreover, note that the right-hand side does not depend on x_2 . Here we put

$$\epsilon = \frac{3C_0||\omega||_1}{W}$$

to have

$$\left|\frac{\mathcal{G}[\omega](\mathbf{x})}{x_2}\right| \leq \frac{W}{3} + C_0 \epsilon^{1/3} \|\omega\|_{L^1([x_1 - \epsilon, x_1 + \epsilon] \times (0, \infty))}^{1/3}.$$

Due to $\omega \in L^1(\mathbb{R}^2_+)$, the last term tends to 0 as $|x_1| \to \infty$, uniformly in x_2 . So there exists a constant $L = L(\omega) > 0$ such that, for any $\mathbf{x} \in \mathbb{R}^2_+$,

$$|x_1| > L \implies \left| \frac{\mathcal{G}[\omega](\mathbf{x})}{x_2} \right| \le \frac{W}{3} + \frac{W}{3} < W,$$

and thus

$$\left\{ \mathcal{G}[\omega] - Wx_2 - \gamma > 0 \right\} \subset \left\{ \mathcal{G}[\omega]/x_2 > W \right\} \subset \left\{ |x_1| \le L \right\}.$$

This completes the proof.

Remark 3.20. In the proof above, we essentially showed that, for any $0 \le f \in (L^1 \cap L^\infty)(\mathbb{R}^2_+)$ with $||x_2 f||_1 < \infty$ and for each W > 0, the set $\{ \mathbf{x} \in \mathbb{R}^2_+ : \mathcal{G}[f](\mathbf{x}) - Wx_2 > 0 \}$ is bounded in \mathbb{R}^2_+ .

Proof of Theorem 3.3. Let M > 0. By Lemma 3.6 and Lemma 3.10, the set S_M is nonempty. Then for any $\omega \in S_M$, by Proposition 3.11, we have

$$\omega = 1_{\{\mathcal{G}[\omega] - Wx_2 - \gamma > 0\}}$$
 a.e. in \mathbb{R}^2_+

for some constants W, $\gamma \ge 0$ which are uniquely determined by ω , and by Lemma 3.17, we have W > 0. Since $E(\omega)$ is the absolute maximum of $E(\cdot)$, we may assume that ω satisfies the Steiner symmetry condition in Definition 3.2 up to translation. Lemma 3.18 implies that ω has a bounded support, up to measure zero set. Lastly, by Lemma 3.15, if we have $\|\omega\|_1 < 1$, then we get $\gamma = 0$.

3.3. Estimate on maximal energy.

Using the specific form of the maximizers given in Theorem 3.3, we obtain the following crucial identity concerning the maximal energy, traveling speed, and impulse.

Lemma 3.21. For any M > 0, we have

(3.5)
$$I_{M} = \frac{3}{4}WM + \frac{1}{2}\gamma ||\omega||_{1},$$

for each $\omega \in S_M$ with the constants $W = W(\omega) > 0$, $\gamma = \gamma(\omega) \ge 0$ in Theorem 3.3.

Remark 3.22. The identity in the above lemma can be essentially found in Lemma 9 of [7]. It is a so-called Pohožaev-type identity (see [77]). We present the proof below for completeness. For vortex rings, we refer to Lemma 3.2 of [42] (also see Proposition 5.13 of [29]).

Proof of Lemma 3.21. Let M > 0. As S_M is nonempty, we take any $\omega \in S_M$. Put $\omega = \mathbf{1}_{\{\mathcal{G}[\omega] - Wx_2 - \gamma > 0\}}$ for the constants $W = W(\omega) > 0$ and $\gamma = \gamma(\omega) \ge 0$ in Theorem 3.3. Let $\widetilde{\omega}$ be the odd extension of ω to \mathbb{R}^2 and denote $\psi := \widetilde{\mathcal{G}[\omega]}$, the odd extension of $\mathcal{G}[\omega]$. We first make the following claim.

(*Claim*) We have $\int_{\mathbb{R}^2} \widetilde{\omega} (\mathbf{x} \cdot \nabla \psi) d\mathbf{x} = 0$.

We first observe that, for any scalar function $f \in C^2(\mathbb{R}^2)$, we have

$$\Delta f \cdot (\mathbf{x} \cdot \nabla f) = \nabla \cdot \left(\frac{1}{2}x_1(f_{x_1}^2 - f_{x_2}^2) + x_2 f_{x_1} f_{x_2}, \frac{1}{2}x_2(-f_{x_1}^2 + f_{x_2}^2) + x_1 f_{x_1} f_{x_2}\right).$$

Knowing that $\widetilde{\omega}$ is compactly supported by Lemma 3.18 and $\psi \in H^2_{loc}(\mathbb{R}^2)$ with the relation $-\Delta \psi = \widetilde{\omega}$ by Lemma 2.4, the approximation to smooth function in the bounded domain in \mathbb{R}^2 gives

$$\int_{\mathbb{R}^2} \widetilde{\omega} \left(\mathbf{x} \cdot \nabla \psi \right) d\mathbf{x} = 0.$$

This completes the proof of (Claim).

Since the function $\widetilde{\omega}$ ($\mathbf{x} \cdot \nabla \psi$) is an even function, we get

$$\int_{\mathbb{R}^2_+} \omega \left(\mathbf{x} \cdot \nabla \mathcal{G}[\omega] \right) d\mathbf{x} = 0.$$

Here, by the relation $\omega = \mathbf{1}_{\{\mathcal{G}[\omega] - Wx_2 - \gamma > 0\}}$, observe that

$$\omega \cdot (\mathcal{G}[\omega] - Wx_2 - \gamma) = (\mathcal{G}[\omega] - Wx_2 - \gamma)_+$$

where $s_+ := \max\{s, 0\}$, $s \in \mathbb{R}$. We define $K := (\mathcal{G}[\omega] - Wx_2 - \gamma)_+$, and using the chain rule (see [46, Theorem 7.8]), we observe that $K \in H^1_{loc}(\mathbb{R}^2_+)$ with the relation

$$\nabla K = \omega \cdot (\nabla \mathcal{G}[\omega] - (0, W)).$$

Note that we have

$$0 = \int_{\mathbb{R}^2_+} \omega \left(\mathbf{x} \cdot \nabla \mathcal{G}[\omega] \right) d\mathbf{x} = \int_{\mathbb{R}^2_+} \mathbf{x} \cdot \nabla K \, d\mathbf{x} + \int_{\mathbb{R}^2_+} W x_2 \omega \, d\mathbf{x}$$

where

$$\int_{\mathbb{R}^2} \mathbf{x} \cdot \nabla K \, d\mathbf{x} = -2 \int_{\mathbb{R}^2} K \, d\mathbf{x}$$

using approximation to smooth function and integration by parts. As we know that

$$\int_{\mathbb{R}^2_+} K \, d\mathbf{x} = \int_{\mathbb{R}^2_+} \omega \cdot \left(\mathcal{G}[\omega] - W x_2 - \gamma \right) d\mathbf{x} = 2I_M - W M - \gamma ||\omega||_1 \quad \text{and} \quad \int_{\mathbb{R}^2_+} W x_2 \omega \, d\mathbf{x} = W M,$$

we get (3.5).

By the scaling given in Remark 3.1-(v) and Remark 3.5, we have the following corollary.

Corollary 3.23. *For any* μ , ν , $\lambda > 0$, *we have*

(3.6)
$$I_{\mu,\nu,\lambda} = \frac{3}{4} W \mu + \frac{1}{2} \gamma ||\omega||_1,$$

for each $\omega \in S_{\mu,\nu,\lambda}$ with the constants $W = W(\omega) > 0$, $\gamma = \gamma(\omega) \ge 0$ in Corollary 3.4.

Remark 3.24. By Theorem 3.3, for any M > 0, the set of maximizers S_M is nonempty, and that each $\omega \in S_M$ satisfies

$$\omega = \mathbf{1}_{\{g[\omega]-Wx_2-\gamma>0\}}$$
 a.e. in \mathbb{R}^2_+

for some constants $W = W(\omega) > 0$, $\gamma = \gamma(\omega) \ge 0$ that are uniquely determined by ω . Hereafter, if there is no confusion, we will simply assume that

$$\omega = \mathbf{1}_{\{\mathcal{G}[\omega]-Wx_2-\gamma>0\}} \quad \text{in } \mathbb{R}^2_+.$$

That is, we dismiss measure zero sets and assume that each maximizer ω is a characteristic function on the open bounded set $\{G[\omega] - Wx_2 - \gamma > 0\}$.

4. Maximizers under small impulse with unit mass bound

In this section, we will prove that, if the constraint on the impulse of the variational problem is sufficiently small, then the flux constant γ of each maximizer vanishes. Accordingly, each traveling vortex patch, generated from any maximizer, touches the horizontal line from which a continuous boundary streamline emerges. The main results of this section are given in Theorem 4.1 and Theorem 4.2 below. In addition, as we obtain $\gamma = 0$ due to a small impulse M, we have the relation

$$I_M = \frac{3}{4}WM$$

which is deduced from the identity (3.5) in Lemma 3.21, and the relation says that $W = W(\omega)$ is completely determined by the choice of M. It leads to various estimates for quantities concluding mass and traveling speed, using the M only. These results are given in Lemma 4.6 and Lemma 4.7, which will be used in Section 5.

Theorem 4.1. There exists an absolute constant $M_1 > 0$ such that, for any $M \in (0, M_1)$ and for each maximizer $\omega = \mathbf{1}_{\Omega} \in S_M$ satisfying the Steiner symmetry condition in Definition 3.2, there exists $\epsilon > 0$ such that

$$B_{\epsilon}(0,0) \subset \overline{\Omega \cup \Omega_{-}}$$

Theorem 4.2 below says that the boundary streamline of the touching vortex patch continuously runs to its end-point on x_1 -axis.

Theorem 4.2. Let $M_1 > 0$ be the constant in Theorem 4.1. For any $M \in (0, M_1)$ and for each $\omega \in S_M$ satisfying the Steiner symmetry condition in Definition 3.2, with the constant $W = W(\omega) > 0$ in Theorem 3.3, the following holds:

(i) Define $A := \{ y_2 > 0 : \omega(0, y_2) = 1 \}$. Then the set A is a nonempty, bounded, and open subset of $(0, \infty)$, and there exists $\epsilon > 0$ such that $(0, \epsilon) \subset A$. In addition, we can define a function $l : (0, \infty) \to [0, \infty)$ such that

$$\{y_1 \in \mathbb{R} : \omega(y_1, y_2) = 1\} = (-l(y_2), l(y_2)) \text{ for } y_2 \in A, \text{ and } l(y_2) = 0 \text{ for } y_2 \in (0, \infty) \setminus A.$$

(ii) The function l is continuous in $(0, \infty)$ and lies in $C^{1,r}$ locally in A, for each $r \in (0, 1)$. Furthermore, for each $y_2 \in A$, we have

$$\mathcal{G}[\omega](l(y_2), y_2) = Wy_2.$$

(iii) There exists a unique constant $a \in (0, \infty)$ such that

$$\widetilde{\mathcal{G}[\omega]}_{x_2}(a,0) = W.$$

Moreover the limit $\lim_{y_2\to 0^+} l(y_2)$ exists and is equal to the number $a\in (0,\infty)$.

4.1. Vanishing of flux constant.

In Lemma 4.4 we will prove $\gamma = 0$ given that the impulse of the maximizer is small enough. Lemma 4.3 below will be used in the proof of Lemma 4.4.

Lemma 4.3. For any $M \in (0, 1)$, we have

$$I_M \geq M^{4/3} \cdot I_1$$
.

Proof. By Theorem 3.3, the set S_1 of maximizers is nonempty. Choose any $f \in S_1$ so that

$$||f||_1 \le 1$$
, $||x_2f||_1 = 1$.

After the scaling $g(\mathbf{x}) = f(M^{-1/3}\mathbf{x})$, we have

$$||g||_1 = M^{2/3} ||f||_1 \le M^{2/3} < 1$$
, $||x_2g||_1 = M||x_2f||_1 = M$, and $E(g) = M^{4/3}E(f) = M^{4/3}I_1$,

so we have $g \in K_M$ and

$$I_M \ge E(g) = M^{4/3} I_1.$$

Lemma 4.4. There exists a constant $M_1 > 0$ such that, for any $M \in (0, M_1)$ and for each $\omega \in S_M$, we have

$$\gamma = 0$$
,

where $\gamma = \gamma(\omega) \ge 0$ is given in Theorem 3.3.

Proof. Let M > 0. As S_M is a nonempty set, we take any $\omega \in S_M$ with the constants $W = W(\omega) > 0$, $\gamma = \gamma(\omega) \ge 0$ given in Theorem 3.3. Then for any L > 0, we split the integral as

$$\|\omega\|_1 = \int_{0 < x_2 \le L} \omega(\mathbf{x}) \, d\mathbf{x} + \int_{x_2 > L} \omega(\mathbf{x}) \, d\mathbf{x}$$

and observe that

$$M = \|x_2\omega\|_1 \ge \int_{x_2>L} x_2\omega(\mathbf{x}) \, d\mathbf{x} \ge L \int_{x_2>L} \omega(\mathbf{x}) \, d\mathbf{x}$$

and so

$$\|\omega\|_1 \leq \int_{0 < x_2 \leq L} \omega(\mathbf{x}) \ d\mathbf{x} + ML^{-1}.$$

On the other hand, for any $\alpha > 0$, we have

$$(4.1) \qquad \int_{\alpha < x_2 \le 2\alpha} \omega \, d\mathbf{x} \le \int_{\alpha < x_2 \le 2\alpha} \omega \cdot \frac{\mathcal{G}[\omega]}{W x_2} \, d\mathbf{x} \le \frac{1}{W\alpha} \int_{\alpha < x_2 \le 2\alpha} \mathcal{G}[\omega] \omega \, d\mathbf{x}.$$

Using the symmetry of G and Lemma 2.1, we obtain

$$\int_{\alpha < x_2 \le 2\alpha} \mathcal{G}[\omega] \omega \, d\mathbf{x} = \int_{\mathbb{R}^2_+} \left[\int_{\alpha < x_2 \le 2\alpha} G(\mathbf{y}, \mathbf{x}) \omega(\mathbf{x}) \, d\mathbf{x} \right] \omega(\mathbf{y}) \, d\mathbf{y} \, \lesssim \, \alpha^{3/2} \int_{\mathbb{R}^2_+} y_2^{1/2} \omega(\mathbf{y}) \, d\mathbf{y} \, \leq \, \alpha^{3/2} \|x_2 \omega\|_1^{1/2} \|\omega\|_1^{1/2} \le \alpha^{3/2} M^{1/2},$$

and so we get

$$\int_{\alpha < x_2 \le 2\alpha} \omega \, d\mathbf{x} \, \lesssim \, \frac{M^{1/2}}{W} \alpha^{1/2}.$$

Therefore we have

$$(4.2) ||\omega||_1 \leq \int_{0 < x_2 \leq L} \omega(\mathbf{x}) d\mathbf{x} + ML^{-1} = \sum_{n \geq 0} \int_{L/2^{n+1} < x_2 \leq L/2^n} \omega(\mathbf{x}) d\mathbf{x} + ML^{-1} \lesssim \frac{M^{1/2}}{W} L^{1/2} + ML^{-1}.$$

As the choice of L > 0 is arbitrary, we put $L = W^{2/3}M^{1/3}$ to obtain

$$\|\omega\|_1 \lesssim \left(\frac{M}{W}\right)^{2/3}.$$

Knowing that $\|\omega\|_1 \lesssim (M/W)^{2/3}$, we will show that the following never holds: there exists a sequence of impulse $\{M_n\} \searrow 0$, and a sequence $\omega_n \subset S_{M_n}$, together with a sequence of positive flux constant $\gamma_n > 0$.

To prove it, we assume that such sequences $\{M_n\}$, $\{\omega_n\}$ and $\{\gamma_n\}$ exist. We may assume that $M_n < 1$ for each $n \ge 1$, and we fix any $n \ge 1$. By the assumption $\gamma_n > 0$, we have $\|\omega_n\|_1 = 1$ by Lemma 3.15. Therefore, by the relation $\|\omega\|_1 \le (M/W)^{2/3}$ and Lemma 3.21, we get

$$W_n \lesssim M_n$$
, and $I_{M_n} \lesssim M_n^2 + \gamma_n$.

Moreover, as $M_n \in (0, 1)$, by Lemma 4.3 we have the relation

$$M_n^{4/3} \lesssim M_n^2 + \gamma_n$$
.

Take N>1 large enough so that we have $M_n^{4/3} \lesssim \gamma_n$ for each $n \geq N$. Recall that we used the relation $\omega \leq \omega \cdot \mathcal{G}[\omega]/Wx_2$ in the inequality (4.1). As $M_n^{4/3} \lesssim \gamma_n$, we can instead use the relation $\omega_n \leq \omega_n \cdot \mathcal{G}[\omega_n]/\gamma_n$ to obtain that

$$\int_{\alpha < x_2 \le 2\alpha} \omega_n \ d\mathbf{x} \ \lesssim \ M_n^{-5/6} \alpha^{3/2}, \quad \text{ for any } \ \alpha > 0.$$

Similarly with (4.2), we then obtain the following relation:

$$1 = \|\omega_n\|_1 \le \sum_{k \ge 0} \int_{L/2^{k+1} < x_2 \le L/2^k} \omega_n(\mathbf{x}) \, d\mathbf{x} + M_n L^{-1} \lesssim M_n^{-5/6} L^{3/2} + M_n L^{-1}, \quad \text{for any } L > 0.$$

Then by choosing $L = M_n^{11/15}$, we have

$$1 \lesssim M_n^{4/15}$$

which contradicts to our assumption that $\{M_n\} \setminus 0$. This completes the proof.

4.2. Touching of pair of uniform vortices.

By Lemma 4.4, for small M > 0 and for each maximizer $\omega \in S_M$, the corresponding flux constant γ vanishes such that we have

$$\omega = \mathbf{1}_{\{G[\omega] - Wx_2 > 0\}}$$

for the traveling speed W > 0. We now assume that the set $\Omega := \{ \mathcal{G}[\omega] - Wx_2 > 0 \} \subset \mathbb{R}^2_+$ satisfies the Steiner symmetry condition, i.e. Ω is concentrated along x_2 -axis. In order to prove that the set Ω touches x_1 -axis at the origin, we have to prove

$$\frac{\partial}{\partial x_2} (\mathcal{G}[\omega](x_1, x_2) - Wx_2) > 0$$
 for any small $|\mathbf{x}| > 0$.

In other words, we have to estimate the horizontal velocity u^1 of the fluid near the center of the dipolar vortex. Lemma 4.5 below gives us a strong estimate, which would guarantee that vanishing of γ is an equivalent condition to touching of a vortex dipole patch.

Central speed estimation.

Lemma 4.5. For some constants W > 0 and $\gamma \ge 0$, if a function ω is given by the relation $\omega = \mathbf{1}_{\Omega}$ for some open bounded set $\Omega \subset \mathbb{R}^2_+$ satisfying the Steiner symmetry condition in Definition 3.2 and the identity

$$\Omega = \{ \mathbf{x} \in \mathbb{R}^2_+ : \mathcal{G}[\omega](\mathbf{x}) - Wx_2 - \gamma > 0 \},$$

we have

$$\widetilde{\mathcal{G}[\omega]}_{x_2}(0,0) > 2W$$

where $\widetilde{\mathcal{G}[\omega]}$ is the odd extention of $\mathcal{G}[\omega]$ to \mathbb{R}^2 .

Proof. Let $\omega = \mathbf{1}_{\{\mathbf{x} \in \mathbb{R}^2_+ : G[\omega](\mathbf{x}) - W, x_2 - \gamma > 0\}}$ with some constants W > 0 and $\gamma \ge 0$. For each $\mathbf{x} \in \mathbb{R}^2$, we put

$$\widetilde{\mathcal{G}}[\omega]_{x_2}(\mathbf{x}) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \frac{-x_2 + y_2}{|\mathbf{x} - \mathbf{y}|^2} \widetilde{\omega}(\mathbf{y}) d\mathbf{y} = \frac{1}{2\pi} \int_{\mathbb{R}^2_+} \frac{-x_2 + y_2}{|\mathbf{x} - \mathbf{y}|^2} \omega(\mathbf{y}) d\mathbf{y} + \frac{1}{2\pi} \int_{\mathbb{R}^2_+} \frac{x_2 + y_2}{|\mathbf{x} - \mathbf{y}^*|^2} \omega(\mathbf{y}) d\mathbf{y} := u_l^1(\mathbf{x}) + u_{nl}^1(\mathbf{x})$$

where $\mathbf{y}^* = (y_1, -y_2)$. By the relation $\widetilde{\mathcal{G}[\omega]}_{x_2}(0,0) = 2u_{nl}^1(0,0)$, we need to show that

$$u_{nl}^1(0,0) > W.$$

Note that the identity (3.3) in Remark 3.16 implies that

$$W = \|\omega\|_1^{-1} \cdot \int_{\mathbb{R}^2} u_{nl}^1 \cdot \omega \, d\mathbf{x},$$

which means that W is the average of u_{nl}^1 in the set supp (ω) . We will claim that, assuming the Steiner symmetry condition of ω , the function u_{nl}^1 attains its unique maximum at $\mathbf{x} = (0,0)$ among all points in \mathbb{R}^2_+ , which would imply that

$$u_{nl}^{1}(0,0) > u_{nl}^{1}(\mathbf{x})$$
 for all $\mathbf{x} \in \text{supp}(\omega) \setminus \{(0,0)\}$

and therefore $u_{nl}^1(0,0) > W$.

(*Claim 1*) For any $(x_1, x_2) \in \overline{\mathbb{R}^2_+}$, we have $u_{nl}^1(x_1, x_2) = u_{nl}^1(-x_1, x_2)$. Moreover, for fixed $x_2 \ge 0$, the function $u_{nl}^1(x_1, x_2)$ decreases in $x_1 \ge 0$.

Let $(x_1, x_2) \in \overline{\mathbb{R}^2_+}$ and assume that $x_1 \neq 0$. We have

$$u_{nl}^{1}(x_1, x_2) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \frac{y_2 + x_2}{y_1^2 + (y_2 + x_2)^2} \omega(y_1 + x_1, y_2) d\mathbf{y}.$$

For any fixed $y_2 > 0$ satisfying that $\omega(0, y_2) = 1$, observe that there exists a constant $l = l(y_2) > 0$ such that

$$\{y_1 \in \mathbb{R} : \omega(y_1, y_2) = 1\} = (-l(y_2), l(y_2))$$

Put $A := \{ y_2 > 0 : \omega(0, y_2) = 1 \}$, which is a nonempty open set, and we get

$$u_{nl}^{1}(x_{1}, x_{2}) = \frac{1}{2\pi} \int_{A} \int_{-l(y_{2})-x_{1}}^{l(y_{2})-x_{1}} \frac{y_{2} + x_{2}}{y_{1}^{2} + (y_{2} + x_{2})^{2}} dy_{1} dy_{2}.$$

Observe that, for fixed $y_2 \in A$, the following function

$$f(x_1) := \int_{-l(y_2)-x_1}^{l(y_2)-x_1} \frac{y_2 + x_2}{y_1^2 + (y_2 + x_2)^2} dy_1, \quad x_1 \in \mathbb{R}$$

satisfies $f(x_1) = f(-x_1)$ and decreases in $x_1 \ge 0$. This completes the proof of (Claim 1).

(*Claim 2*) The function $g(x_2) := u_{nl}^1(0, x_2)$ decreases in $x_2 \ge 0$.

For each $x_2 \ge 0$,

$$g(x_2) = \frac{1}{2\pi} \int_{\mathbb{R}^2_+} \frac{y_2 + x_2}{y_1^2 + (y_2 + x_2)^2} \omega(\mathbf{y}) \, d\mathbf{y}.$$

As in the proof of (Claim 1), we have

$$g(x_2) = \frac{1}{2\pi} \int_A \int_{-l(y_2)}^{l(y_2)} \frac{y_2 + x_2}{y_1^2 + (y_2 + x_2)^2} dy_1 dy_2 = \frac{1}{\pi} \int_A \arctan\left(\frac{l(y_2)}{y_2 + x_2}\right) dy_2.$$

It implies that $g(x_2)$ decreases strictly in $x_2 \ge 0$. This completes the proof of (Claim 2).

By (Claim 1) and (Claim 2), we have

$$u_{nl}^{1}(0,0) > \|\omega\|_{1}^{-1} \cdot \int_{\mathbb{R}^{2}_{+}} u_{nl}^{1} \cdot \omega \, d\mathbf{x} = W,$$

and so we have $\widetilde{\mathcal{G}[\omega]}_{x_2}(0,0) > 2W$. This completes the proof.

Proof of Theorem 4.1.

Lemma 4.5 implies that each maximizer $\omega \in S_M$ touches x_1 -axis if and only if we get $\gamma = 0$. Recall Lemma 4.4 which says that a sufficient condition for γ to vanish is the smallness condition of the impulse M > 0. We will prove Theorem 4.1 here.

Proof of Theorem 4.1. Let $M_1 > 0$ be the constant in Lemma 4.4. For any $M \in (0, M_1)$, as the set S_M is nonempty, we choose any $\omega \in S_M$. By Lemma 4.4, we have $\gamma = 0$ and thus

$$\omega = \mathbf{1}_{\{G[\omega] - Wx_2 > 0\}},$$

where W, γ are the constants in Theorem 3.3. It suffices to show that there exists $\epsilon > 0$ such that

$$B_{\epsilon}(0,0) \cap \mathbb{R}^2_+ \subset \left\{ \mathcal{G}[\omega] - Wx_2 > 0 \right\}.$$

By Lemma 4.5, we have

$$\widetilde{\mathcal{G}[\omega]}_{x_2}(0,0) > 2W.$$

By $\widetilde{\mathcal{G}[\omega]} \in C^1(\mathbb{R}^2)$, there exists a small $\epsilon > 0$ such that

$$\widetilde{\mathcal{G}[\omega]}_{x_2} > 2W$$
 in $B_{\epsilon}(0,0)$.

Then for each $\mathbf{x} = (x_1, x_2) \in B_{\epsilon}(0, 0) \cap \mathbb{R}^2_+$, we have

$$\mathcal{G}[\omega](\mathbf{x}) = \int_0^{x_2} \mathcal{G}[\omega]_{x_2}(x_1, s) \, ds > 2Wx_2 > Wx_2,$$

and therefore $\mathbf{x} \in \{ \mathcal{G}[\omega] - Wx_2 > 0 \}$. This completes the proof.

Proof of Theorem 4.2.

In the proof below, we obtain the continuity of the boundary of each vortex patch by the continuity of the stream function and the implicit function theorem for which we prove the monotonicities of the stream function in $x_1 > 0$ and the horizontal speed u^1 on the axis of symmetry $\{x_2 = 0\}$.

Proof of Theorem 4.2. Let $M_1 > 0$ be the constant in Lemma 4.4. For any $M \in (0, M_1)$, as the set S_M is nonempty, we choose any $\omega = \mathbf{1}_{\{\mathcal{G}[\omega] - Wx_2 > 0\}} \in S_M$ where $W = W(\omega) > 0$ is the constant in Theorem 3.3. Note that $\gamma = \gamma(\omega) = 0$ by Lemma 4.4. We assume that ω satisfies the Steiner symmetry condition. To prove (i), we define a set

$$A := \{ y_2 > 0 : \omega(0, y_2) = 1 \} = \{ y_2 > 0 : \mathcal{G}[\omega](0, y_2) - Wy_2 > 0 \}$$

which is clearly a nonempty set. We observe that, by Theorem 4.1, there exists $\epsilon > 0$ satisfying $(0, \epsilon) \subset A$. Moreover, the set $\{\mathcal{G}[\omega] - Wx_2 > 0\}$ is open bounded set in \mathbb{R}^2_+ by Theorem 3.3, the set A is open bounded subset of $(0, \infty)$. By the Steiner symmetry condition of ω , we can define a function $l:(0,\infty) \to [0,\infty)$ satisfying that, for each $y_2 \in A$,

$$l(y_2) > 0$$
 and $\{y_1 \in \mathbb{R} : \omega(y_1, y_2) = 1\} = (-l(y_2), l(y_2)),$

and $l \equiv 0$ in $(0, \infty)\backslash A$. This proves (i). For the proof of (ii), we first make the following claims.

(Claim 1) For each $y_2 \in \overline{A}$, we have $\mathcal{G}[\omega](l(y_2), y_2) = Wy_2$.

Let $y_2 \in A$. For any $y_1 \in (0, l(y_2))$, we have $\omega(y_1, y_2) = 1$, and so $\mathcal{G}[\omega](y_1, y_2) > Wy_2$. If $y_1 \in (l(y_2), \infty)$, then $\omega(y_1, y_2) = 0$, and so we have $\mathcal{G}[\omega](y_1, y_2) \leq Wy_2$. By the continuity of $\mathcal{G}[\omega]$, we have

$$\mathcal{G}[\omega](l(y_2), y_2) = Wy_2.$$

For each $y'_2 \in \partial A$, we have $l(y'_2) = 0$ by definition of l, and a similar argument can be done to obtain that

$$\mathcal{G}[\omega](0,y_2')-Wy_2'=0.$$

This proves (Claim 1).

(Claim 2) For each $\mathbf{x} = (x_1, x_2) \in \mathbb{R}^2_+$ satisfying $x_1 > 0$, we have $\mathcal{G}[\omega]_{x_1}(\mathbf{x}) < 0$.

For each $\mathbf{x} = (x_1, x_2) \in \mathbb{R}^2_+$ satisfying $x_1 > 0$, observe that

$$2\pi \cdot \mathcal{G}[\omega]_{x_{1}}(\mathbf{x}) = \int_{\mathbb{R}^{2}_{+}} \omega(\mathbf{y})(y_{1} - x_{1}) \left[\frac{1}{|\mathbf{y} - \mathbf{x}|^{2}} - \frac{1}{|\mathbf{y}^{*} - \mathbf{x}|^{2}} \right] d\mathbf{y}$$

$$= \int_{A} \int_{-l(y_{2})}^{l(y_{2})} \frac{4x_{2}y_{2} (y_{1} - x_{1})}{|\mathbf{y} - \mathbf{x}|^{2} |\mathbf{y}^{*} - \mathbf{x}|^{2}} dy_{1} dy_{2}$$

$$= \int_{A} \int_{-l(y_{2}) - x_{1}}^{l(y_{2}) - x_{1}} \frac{y_{1} \cdot (4x_{2}y_{2})}{(y_{1}^{2} + (y_{2} - x_{2})^{2}) \cdot (y_{1}^{2} + (y_{2} + x_{2})^{2})} dy_{1} dy_{2}$$

$$= \int_{A} \left[F(l(y_{2}) - x_{1}) - F(-l(y_{2}) - x_{1}) \right] dy_{2}$$

where, for each fixed $y_2 \in A \setminus \{x_2\}$, the function $F : \mathbb{R} \to \mathbb{R}$ is defined as

$$F(s) := \int_0^s \frac{y_1 \cdot (4x_2y_2)}{(y_1^2 + (y_2 - x_2)^2) \cdot (y_1^2 + (y_2 + x_2)^2)} \, dy_1.$$

Note that F is an even function in $s \in \mathbb{R}$ by the odd symmetry of the integrand in $y_1 \in \mathbb{R}$. Moreover, we observe that F(s) increases strictly in $|s| \ge 0$, since the integrand is positive in $y_1 > 0$. For each $y_2 \in A \setminus \{x_2\}$, we have $I(y_2) > 0$ and so we get

$$|l(y_2) - x_1| < |-l(y_2) - x_1|$$
, implying that $F(l(y_2) - x_1) < F(-l(y_2) - x_1)$.

This gives $\mathcal{G}[\omega]_{x_1}(\mathbf{x}) < 0$ and completes the proof of (*Claim 2*).

We will now prove (ii). First we want to prove that $l \in C^{1,r}_{loc}(A)$ for each $r \in (0,1)$. We define a C^1 function $\Psi: \mathbb{R}^2_{\perp} \to \mathbb{R}$ as

$$\Psi(\mathbf{x}) := \mathcal{G}[\omega](\mathbf{x}) - Wx_2.$$

For each $y_2 \in A$, by (Claim 1) and (Claim 2), we have $\Psi(l(y_2), y_2) = 0$ and $\Psi_{x_1}(l(y_2), y_2) < 0$. Then the implicit function theorem can be applied to Ψ at each point $(l(y_2), y_2)$. That is, for each $y_2 \in A$, there exists a small neighborhood $U \subset A$ containing y_2 such that we have

$$\Psi(f(s), s) = 0, \quad s \in U$$

for some function $f:U\to\mathbb{R}$ which is continuously differentiable in U and satisfies $f(y_2)=l(y_2)$. In (Claim 2), we proved that $\Psi(x_1, x_2)$ is strictly decreasing in $x_1 > 0$, and it follows that f(s) = l(s) for each $s \in U$. In conclusion, l is continuously differentiable in A. To show that $l' \in C^r_{loc}(A)$ for each $r \in (0, 1)$, we fix any $r \in (0, 1)$. It suffices to show that, for any $s \in A$ and $\epsilon > 0$ satisfying that $[s - \epsilon, s + \epsilon] \subset A$, we have

$$\sup_{0<|s-t|\leq\epsilon}\frac{|l'(s)-l'(t)|}{|s-t|^r}<\infty.$$

By the formula $\Psi(l(s), s) = 0$, we have

$$l'(s) = -\frac{\Psi_{x_2}(l(s), s)}{\Psi_{x_1}(l(s), s)}.$$

Fix $s \in A$ and $\epsilon > 0$ to satisfy $[s - \epsilon, s + \epsilon] \subset A$. Then for any $t \in [s - \epsilon, s + \epsilon]$, we have

$$|l'(s) - l'(t)| \lesssim \frac{\left| (\nabla \Psi)(l(t), t) \right|}{\left| \Psi_{x_1}(l(s), s) \cdot \Psi_{x_1}(l(t), t) \right|} \cdot \left| (\nabla \Psi)(l(s), s) - (\nabla \Psi)(l(t), t) \right|.$$

We deal with the denominator first. As the continuous function $g(y) := \Psi_{x_1}(l(y), y)$ satisfies g < 0 in the domain A, there exists a constant $C_0 > 0$ such that $g < -C_0$ in $[s - \epsilon, s + \epsilon]$. Therefore we have

$$\left| \ l'(s) - l'(t) \ \right| \ \lesssim \left| (\nabla \Psi)(l(t), t) \ \right| \cdot \left| \ (\nabla \Psi)(l(s), s) - (\nabla \Psi)(l(t), t) \ \right|$$

By Lemma 2.4, the relation $\nabla \Psi = \nabla \mathcal{G}[\omega] - (0, W)$ leads to $\|\nabla \Psi\|_{L^{\infty}} \lesssim \|\nabla \mathcal{G}[\omega]\|_{L^{\infty}} + W < \infty$, so we finally get

$$|l'(s) - l'(t)| \le |(\nabla \Psi)(l(s), s) - (\nabla \Psi)(l(t), t)|.$$

As $\nabla \Psi \in C^r(\mathbb{R}^2_+)$ by Lemma 2.4, we have

$$|l'(s) - l'(t)| \lesssim_r |(l(s) - l(t))^2 + (s - t)^2|^{r/2} \le |s - t|^r \cdot ||l'||_{L^{\infty}([s - \epsilon, s + \epsilon])}^2 + 1|^{r/2}$$

where $||l'||_{L^{\infty}([s-\epsilon,s+\epsilon])} < \infty$ due to $l' \in C(A)$. Therefore we have $l' \in C^r_{loc}(A)$.

It remains to show that $l \in C((0, \infty))$. Note that, at each point in A and $(0, \infty) \setminus \overline{A}$, the continuity of l is trivially obtained using the fact that A is an open set. For each $y_2 \in (0, \infty) \cap \partial A$, we observe that

$$G[\omega](0, y_2) = Wy_2$$
 and $l(y_2) = 0$

by (Claim 1). We need to show that $\lim_{s\to y_2} l(s) = 0$. As ω is boundedly supported in \mathbb{R}^2_+ , we have

$$\limsup_{s\to v_2}l(s)<\infty.$$

If l(s) does not converges to 0 as $s \to y_2$, there exists a sequence $\{s_n\} \to y_2$ and a number $L \in (0, \infty)$ such that $\lim_{n \to \infty} l(s_n) = L$. By the continuity of $\mathcal{G}[\omega]$, we have

$$\mathcal{G}[\omega](L, y_2) - Wy_2 = 0,$$

which means that $\mathcal{G}[\omega](0, y_2) = \mathcal{G}[\omega](L, y_2)$. It contradicts to (*Claim 2*). In sum, we have $l \in C((0, \infty))$.

Finally, we now prove (iii), starting from the following claim.

(*Claim 3*) $\widetilde{\mathcal{G}}[\omega]_{x_2}(s,0)$ strictly decreases in $s \ge 0$.

For $s \ge 0$, we have

$$\widetilde{\mathcal{G}[\omega]}_{x_2}(s,0) = \frac{1}{\pi} \int_{\mathbb{R}^2_+} \frac{y_2}{(y_1 - s)^2 + y_2^2} \omega(\mathbf{y}) \, d\mathbf{y} = \frac{1}{\pi} \int_A \int_{-l(y_2) - s}^{l(y_2) - s} \frac{y_2}{y_1^2 + y_2^2} dy_1 dy_2.$$

For fixed $y_2 \in A$, observe that the integrand

$$\int_{-l(y_2)-s}^{l(y_2)-s} \frac{y_2}{y_1^2 + y_2^2} dy_1$$

strictly decreases in $s \ge 0$. This completes the proof of (*Claim 3*).

Note that we have

$$\widetilde{\mathcal{G}[\omega]}_{x_2}(0,0) > 2W$$

by Lemma 4.5. By (3.4) in Lemma 3.18, there exists $K \in (0, \infty)$ such that we have

$$x_1 > K \Rightarrow \widetilde{\mathcal{G}[\omega]}_{x_2}(x_1, 0) = \lim_{x_2 \searrow 0} \frac{\mathcal{G}[\omega](x_1, x_2)}{x_2} \leq W.$$

As $\widetilde{\mathcal{G}[\omega]}_{x_2}(s,0)$ decreases strictly in $s \ge 0$, there exists a unique constant $a \in (0,\infty)$ satisfying

$$\widetilde{\mathcal{G}[\omega]}_{x_2}(a,0) = W.$$

It remains to show that $\lim_{s\to 0^+} l(s) = a$. As ω is boundedly supported, we have

$$\limsup_{s \searrow 0} l(s) < \infty.$$

Suppose that l(s) does not converge to a as $s \setminus 0$. Then there exists a sequence $s'_n \setminus 0$ and a number $a' \in (0, \infty)$ satisfying that

$$\lim_{n\to\infty}l(s_n') = a' \neq a.$$

In the relation

$$W = \frac{\mathcal{G}[\omega](\ l(s_n'),\ s_n')}{s_n'},$$

due to $\mathcal{G}[\omega](l(s'_n), 0) = 0$, we can use the mean value theorem for each $n \ge 1$ to find a sequence $0 < s''_n < s'_n$ such that

$$W = \widetilde{\mathcal{G}[\omega]}_{x_2}(l(s'_n), s''_n).$$

Letting $n \to \infty$ gives $W = \mathcal{G}[\omega]_{x_2}(a', 0)$, which contradicts to $a' \neq a$. This completes the proof of (iii).

4.3. Estimates when flux constant vanishes due to small impulse.

Under the small impulse condition, knowing that $\gamma = 0$ by Lemma 4.4, we can obtain several estimates depending only on M.

Lemma 4.6. There exists an absolute constant $C \ge 1$ such that the following hold:

Let $M_1 > 0$ be the constant in Lemma 4.4. For any $M \in (0, M_1)$ and for each $\omega \in S_M$, we have

$$\frac{1}{C} \cdot M^{2/3} \le ||\omega||_1 \le C \cdot M^{2/3},$$

$$\frac{1}{C} \cdot M^{4/3} \leq I_M \leq C \cdot M^{4/3},$$

$$\frac{1}{C} \cdot M^{1/3} \leq W \leq C \cdot M^{1/3},$$

where $W = W(\omega) > 0$ is given in Theorem 3.3.

Proof. Let $M_1 > 0$ be the constant in Lemma 4.4 and choose $M \in (0, M_1)$. As S_M is nonmepty, we choose any $\omega \in S_M$ with the constants $W = W(\omega) > 0$ and $\gamma = \gamma(\omega) \ge 0$ given in Theorem 3.3. Note that $\gamma = 0$ by Lemma 4.4. We will prove each inequality separately.

(Claim 1)
$$M^{4/3} \lesssim I_M$$
 and $M^{1/3} \lesssim W$.

Lemma 4.3 says that $M^{4/3} \lesssim I_M$ given that M < 1. Using the identity $I_M = (3/4)WM$ in Lemma 3.21, we obtain $M^{1/3} \lesssim W$. This completes the proof of (*Claim 1*).

(Claim 2)
$$\|\omega\|_1 \lesssim M^{2/3}$$
.

Before we estimate $\|\omega\|_1$, recall that $\mathcal{G}[\omega] \in L^{\infty}(\mathbb{R}^2)$ by (2.4) in Lemma 2.3 and that we have

$$(4.7) \qquad \left\{ \mathcal{G}[\omega] - Wx_2 > 0 \right\} \subset \left\{ x_2 < \|\mathcal{G}[\omega]\|_{L^{\infty}(\mathbb{R}^2_+)} / W \right\} \quad \text{which implies} \quad \sup_{\mathbf{x} \in \text{supp}(\omega)} x_2 \le \frac{\|\mathcal{G}[\omega]\|_{L^{\infty}(\mathbb{R}^2_+)}}{W}.$$

With the fact that $\|\omega\|_2 = \|\omega\|_1^{1/2}$, the estimate (2.4) in Lemma 2.3 gives

$$\|\mathcal{G}[\omega]\|_{L^{\infty}(\mathbb{R}^{2}_{+})} \lesssim \|\omega\|_{1}^{1/2}M^{1/3} + \|\omega\|_{1}^{1/4}M^{1/2}.$$

Using the relation $M^{1/3} \lesssim W$ in (Claim 1), we have

$$\sup_{\mathbf{x} \in \text{supp}(\omega)} x_2 \lesssim \|\omega\|_1^{1/2} + \|\omega\|_1^{1/4} M^{1/6}.$$

Put $K := \sup_{\mathbf{x} \in \text{supp}(\omega)} x_2$ and observe that

$$\|\omega\|_1 = \int_{Y_2 \le K} \omega \, d\mathbf{x}$$
 where $K \le \|\omega\|_1^{1/2} + \|\omega\|_1^{1/4} M^{1/6}$.

We will now use Lemma 2.1 to estimate $\|\omega\|_1$ in terms of K. Observe that

$$\|\omega\|_1 = \sum_{n>0} \int_{K/2^{n+1} < x_2 \le K/2^n} \omega(\mathbf{x}) d\mathbf{x}.$$

For any $\alpha > 0$, we have

$$\int_{\alpha < x_2 \le 2\alpha} \omega \, d\mathbf{x} \, \le \, \int_{\alpha < x_2 \le 2\alpha} \omega \cdot \frac{\mathcal{G}[\omega]}{W x_2} \, d\mathbf{x} \, \le \, \frac{1}{W\alpha} \int_{\alpha < x_2 \le 2\alpha} \mathcal{G}[\omega] \omega \, d\mathbf{x} \, \lesssim \, \frac{1}{M^{1/3}\alpha} \int_{\alpha < x_2 \le 2\alpha} \mathcal{G}[\omega] \omega \, d\mathbf{x}.$$

Using the symmetry of G and Lemma 2.1, we obtain

$$\int_{\alpha < x_2 \le 2\alpha} \mathcal{G}[\omega] \omega \ d\mathbf{x} = \int_{\mathbb{R}^2_+} \left[\int_{\alpha < x_2 \le 2\alpha} G(\mathbf{y}, \mathbf{x}) \omega(\mathbf{x}) \ d\mathbf{x} \right] \omega(\mathbf{y}) \ d\mathbf{y} \ \lesssim \ \alpha^{3/2} \int_{\mathbb{R}^2_+} y_2^{1/2} \omega(\mathbf{y}) \ d\mathbf{y} \ \leq \ \alpha^{3/2} M^{1/2} \|\omega\|_1^{1/2}.$$

Therefore we have

$$\int_{\alpha < x_2 < 2\alpha} \omega \, d\mathbf{x} \, \lesssim \, \alpha^{1/2} M^{1/6} \|\omega\|_1^{1/2}.$$

Then

$$||\omega||_1 = \sum_{n \geq 0} \int_{K/2^{n+1} < x_2 \leq K/2^n} \omega(\mathbf{x}) \, d\mathbf{x} \lesssim M^{1/6} ||\omega||_1^{1/2} \sum_{n \geq 0} \left(\frac{K}{2^{n+1}}\right)^{1/2} \lesssim K^{1/2} M^{1/6} ||\omega||_1^{1/2}.$$

In sum, we obtain

$$\|\omega\|_1 \ \lesssim \ M^{1/6} \|\omega\|_1^{1/2} \big(\|\omega\|_1^{1/2} + \|\omega\|_1^{1/4} M^{1/6} \big)^{1/2}$$

and therefore

$$\|\omega\|_1^{3/4} \lesssim M^{1/3} \|\omega\|_1^{1/4} + M^{1/2}.$$

Under the substitution $t := M^{-1/6} ||\omega||_1^{1/4}$, we obtain

$$t^3 \lesssim t + 1$$

and thus $t \lesssim 1$. In sum, we get

$$M^{-1/6} \|\omega\|_1^{1/4} \lesssim 1$$
 and so $\|\omega\|_1 \lesssim M^{2/3}$.

(Claim 3) $I_M \lesssim M^{4/3}$, $\|\mathcal{G}[\omega]\|_{L^{\infty}(\mathbb{R}^2_+)} \lesssim M^{2/3}$, and $W \lesssim M^{1/3}$.

Using the relation $\|\omega\|_1 \lesssim M^{2/3}$ and the fact that $\|\omega\|_2 = \|\omega\|_1^{1/2}$, the estimate (2.5) in Lemma 2.3 gives

$$I_M \lesssim M^{1/2} ||\omega||_1^{5/4} \lesssim M^{4/3},$$

and the estimate (2.4) in Lemma 2.3 gives

$$\|\mathcal{G}[\omega]\|_{L^{\infty}(\mathbb{R}^{2}_{+})} \lesssim \|\omega\|_{1}^{1/2}M^{1/3} + \|\omega\|_{1}^{1/4}M^{1/2} \lesssim M^{2/3}.$$

And lastly, using Lemma 3.21, we obtain

$$W = \frac{4I_M}{3M} \lesssim M^{1/3}.$$

It completes the proof of (4.4), (4.5), and (4.6).

(Claim 4) $M^{2/3} \lesssim ||\omega||_1$.

Using the relation $\|\mathcal{G}[\omega]\|_{L^{\infty}(\mathbb{R}^{2}_{+})} \lesssim M^{2/3}$ and $M^{1/3} \lesssim W$, the inequality (4.7) implies that

(4.8)
$$\sup_{\mathbf{x} \in \text{supp}(\omega)} x_2 \leq \frac{\|\mathcal{G}[\omega]\|_{L^{\infty}(\mathbb{R}^2_+)}}{W} \lesssim M^{1/3}.$$

Therefore, we can observe that

$$M = \int_{x_2 \leq M^{1/3}} x_2 \omega(\mathbf{x}) d\mathbf{x} \leq M^{1/3} ||\omega||_1,$$

so we get $M^{2/3} \lesssim ||\omega||_1$.

Combining some estimates in Lemma 4.6, we can roughly estimate the size of the compact support of maximizers in S_M .

Lemma 4.7. There exists a constant C > 0 satisfying the following:

Let $M_1 > 0$ be the constant in Lemma 4.4. For any $M \in (0, M_1)$ and for each $\omega \in S_M$ satisfying the Steiner symmetry condition in Definition 3.2, we have

$$\sup_{\mathbf{x} \in \text{supp}(\omega)} |\mathbf{x}| \leq C \cdot M^{1/3},$$

Proof. Let $M \in (0, M_1)$. As S_M is nonmepty, we choose any $\omega = \mathbf{1}_{\{\mathcal{G}[\omega] - Wx_2 > 0\}} \in S_M$ with the constant $W = W(\omega) > 0$ in Theorem 3.3. We assume that ω satisfies the Steiner symmetry condition. First, we recall the relation

$$\sup_{\mathbf{x} \in \operatorname{supp}(\omega)} x_2 \leq \frac{\|\mathcal{G}[\omega]\|_{L^{\infty}}}{W} \lesssim M^{1/3}$$

which was obtained by the inequality (4.8) in the proof of Lemma 4.6. It remains to show that

$$\sup_{\mathbf{x}\in\operatorname{supp}(\omega)}|x_1|\lesssim M^{1/3}.$$

We first observe that

$$\operatorname{supp}(\omega) \subset \left\{ \frac{\mathcal{G}[\omega](\mathbf{x})}{x_2} > W \right\}$$

and recall the Lemma 3.18 to have

$$\left|\frac{\mathcal{G}[\omega](\mathbf{x})}{x_2}\right| \leq C_0 \left(\epsilon^{-1} \|\omega\|_1 + \epsilon^{1/3} \|\omega\|_{L^1\left([x_1 - \epsilon, x_1 + \epsilon] \times (0, \infty)\right)}^{1/3}\right)$$

which holds for any $\mathbf{x} \in \mathbb{R}^2_+$ and $\epsilon > 0$. Here $C_0 > 0$ is an absolute constant. We fix $\epsilon = 3C_0 ||\omega||_1 / W$ to obtain

$$\left|\frac{\mathcal{G}[\omega](\mathbf{x})}{x_2}\right| \leq \frac{W}{3} + C_0 \epsilon^{1/3} \|\omega\|_{L^1\left([x_1 - \epsilon, x_1 + \epsilon] \times (0, \infty)\right)}^{1/3}.$$

Suppose $x_1 > \epsilon$ and observe that

$$\|\omega\|_{L^{1}\left([x_{1}-\epsilon,x_{1}+\epsilon]\times(0,\infty)\right)} = \int_{0}^{\infty} \left[\int_{x_{1}-\epsilon}^{x_{1}+\epsilon} \omega(t,x_{2}) dt\right] dx_{2} \le \frac{1}{2} \|\omega\|_{1} \cdot \int_{x_{1}-\epsilon}^{x_{1}+\epsilon} \frac{1}{t} dt = \frac{1}{2} \|\omega\|_{1} \ln\left[\frac{x_{1}+\epsilon}{x_{1}-\epsilon}\right] dt$$

where we have used the relation

$$t \cdot \omega(t, x_2) \le \int_0^\infty \omega(s, x_2) ds$$
, for any $t, x_2 > 0$

which follows from the Steiner symmetry condition on ω . If we additionally assume $x_1 > 2\epsilon$, we have

$$\|\omega\|_{L^1\left([x_1-\epsilon,x_1+\epsilon]\times(0,\infty)\right)} \le 2\epsilon x_1^{-1}\|\omega\|_1$$

and therefore

$$\left| \frac{\mathcal{G}[\omega](\mathbf{x})}{x_2} \right| \leq \frac{W}{3} + C_1 \epsilon^{2/3} ||\omega||_1^{1/3} x_1^{-1/3}$$

for the constant $C_1 := 2^{1/3}C_0$. If we suppose that

$$x_1 > C_2 \frac{\epsilon^2}{W^3} ||\omega||_1$$
 where $C_2 := (3C_1)^3$,

we get

$$\left|\frac{\mathcal{G}[\omega](\mathbf{x})}{x_2}\right| \leq \frac{2}{3}W,$$

and therefore $\mathbf{x} \in (\text{supp}(\omega))^c$. In sum, we have the following.

$$\operatorname{supp}(\omega) \subset \left\{ \mathbf{x} \in \mathbb{R}_+^2 : |x_1| > \max \left\{ 2\epsilon, C_2 \frac{\epsilon^2}{W^3} ||\omega||_1 \right\} \right\}^c, \quad \text{where } \epsilon = 3C_0 ||\omega||_1 / W.$$

Using the estimates (4.3) and (4.5) in Lemma 4.6, we get

$$\epsilon \lesssim M^{1/3}$$
 and so $\frac{\epsilon^2}{W^3} ||\omega||_1 \lesssim M^{1/3}$.

This implies that

$$\sup_{\mathbf{x}\in\operatorname{supp}(\omega)}|x_1| \lesssim M^{1/3}.$$

5. VARIATIONAL PROBLEM ON EXACT IMPULSE WITHOUT MASS BOUND

5.1. Variational problem without mass bound.

For M > 0, we then consider the following admissible set without mass bound,

$$K_{M,\infty} := \left\{ \omega \in L^1(\mathbb{R}^2_+) : \omega = \mathbf{1}_A \text{ a.e. where } A \subset \mathbb{R}^2_+ \text{ is (Lebesgue) measurable, } \|x_2 \omega\|_1 = M \right\},$$

with the maximal energy

$$I_{M,\infty} := \sup_{\omega \in K_{M,\infty}} E(\omega)$$

and the set of maximizers

$$S_{M,\infty} := \{ \omega \in K_{M,\infty} : E(\omega) = I_{M,\infty} \}.$$

Observe that this new admissible set $K_{M,\infty}$ does not contain any upper bound condition for mass. We note that the elementary estimate (2.5) in Lemma 2.3:

$$\left| \int_{\mathbb{R}^2_+} \int_{\mathbb{R}^2_+} G(\mathbf{x}, \mathbf{y}) \omega(\mathbf{x}) \omega(\mathbf{y}) \, d\mathbf{x} d\mathbf{y} \right| \leq \|\omega\|_1 \|\omega\|_2^{1/2} \|x_2 \omega\|_1^{1/2}$$

does not guarantee that $I_{M,\infty} < \infty$. In this section, we will prove that $I_{M,\infty} < \infty$ and $S_{M,\infty} \neq \emptyset$, which is quite surprising. Moreover, this new problem inherits all results from the previous variational problems. The main result of this section is given in Theorem 5.1 below, and this section is dedicated to its proof.

Theorem 5.1. We have the following properties:

- (i) $I_{1,\infty} < \infty$.
- (ii) $S_{1,\infty}$ is nonempty.
- (iii) Each $\omega \in S_{1,\infty}$ satisfies the relation

$$\omega = \mathbf{1}_{\{\mathcal{G}[\omega]-Wx_2>0\}}$$
 a.e. in \mathbb{R}^2_+ , for the traveling speed $W := \frac{4}{3} I_{1,\infty}$.

(iv) Each $\omega \in S_{1,\infty}$ has a bounded support and satisfies the Steiner symmetry condition in Definition 3.2 up to translation. More precisely, for each $\omega \in S_{1,\infty}$, there exist $\tau \in \mathbb{R}$ and an open bounded set $A \subset \mathbb{R}^2_+$ satisfying the Steiner symmetry condition such that, for the translation ω_{τ} given by $\omega_{\tau}(\cdot) := \omega(\cdot + (\tau, 0))$, we have

$$\omega_{\tau} = \mathbf{1}_{A}$$
 a.e. in \mathbb{R}^{2}_{+} .

(v) There exists an absolute constant $C \ge 1$ such that, for each $\omega \in S_{1,\infty}$, we have

$$\sup_{\mathbf{x} \in \text{supp}(\omega)} |\mathbf{x}| \leq C \quad and \quad \frac{1}{C} \leq ||\omega||_1 \leq C.$$

Remark 5.2. The key idea of the proof of Theorem 5.1 is to show that

$$S_{1,\infty} = S_{1,\nu,1}$$
 for sufficiently large $\nu > 0$.

Remark 5.3. In the statements (i)-(iv) of Theorem 5.1, we can replace $I_{1,\infty}$, $S_{1,\infty}$ with $I_{M,\infty}$, $S_{M,\infty}$ for each M > 0 by the scaling: The scaling map $\phi : \omega \mapsto \hat{\omega}$ given by

$$\hat{\omega}(\mathbf{x}) = \omega(M^{-1/3}\mathbf{x})$$

maps the set $S_{1,\infty}$ to the set $S_{M,\infty}$ bijectively, with the scaled traveling speed and the maximal energy given by

$$W_M := W \cdot M^{1/3}$$
 and $I_{M,\infty} = I_{1,\infty} \cdot M^{4/3}$.

5.2. Scaling property.

Recall the estimate (4.3) in Lemma 4.6, which gives

$$||\omega||_1 \leq M^{2/3}$$
.

As the impulse M is small enough, each maximizer cannot achieve the full mass (which is 1) to attain the maximal energy. Using this property, we can show the following two lemmas.

Lemma 5.4. There exists a constant $M_2 > 0$ satisfying the following:

For each $M \in (0, M_2)$ and any $\alpha > 1$, we have

$$S_M = S_{M,\alpha,1}$$
.

Proof. Denote the constant C > 1 in Lemma 4.6 as $C_0 > 1$. Choose any small constant $M_2 \in (0, 1)$ satisfying that

$$M_2 \le M_1$$
 and $C_0 \cdot M_2^{2/3} \le 1$.

Let $M \in (0, M_2)$ and $\alpha > 1$. As $S_{M,\alpha,1}$ is nonempty, choose any $\omega \in S_{M,\alpha,1}$ and consider the scaling map

$$\hat{\omega}(\mathbf{x}) = \omega(\alpha^{1/2}\mathbf{x})$$

to attain

$$\|\hat{\omega}\|_1 = \alpha^{-1} \|\omega\|_1 \le 1, \quad \|x_2\hat{\omega}\|_1 = \alpha^{-3/2} \|x_2\omega\|_1 = \alpha^{-3/2} M$$

and so $\hat{\omega} \in K_{\alpha^{-3/2}M}$. By the bijectivity of the scaling map from $S_{M,\alpha,1}$ onto $S_{\alpha^{-3/2}M}$, we get $\hat{\omega} \in S_{\alpha^{-3/2}M}$. Here, as $\alpha^{-3/2}M < M_1$, we can use the estimate (4.3) in Lemma 4.6 to have

$$\|\hat{\omega}\|_1 \leq C_0 \cdot (\alpha^{-3/2} M)^{2/3},$$

and therefore

$$\|\omega\|_1 \leq C_0 \cdot M^{2/3} < 1.$$

This implies that $\omega \in K_M$. Using the assumption that $\omega \in S_{M,\alpha,1}$ and the fact that $K_M \subset K_{M,\alpha,1}$, we obtain $\omega \in S_M$. In sum, we have $S_{M,\alpha,1} \subset S_M$, which leads to $S_{M,\alpha,1} = S_M$.

Corollary 5.5. Let $M_2 > 0$ be the constant in Lemma 5.4. Then for any M, v > 0 satisfying $Mv^{-3/2} < M_2$, we have

$$S_{M,v,1} = S_{M,\alpha,1}, \quad for \ all \ \alpha > v.$$

Proof. Let M, v, $\alpha > 0$ and assume that $Mv^{-3/2} < M_2$ and $\alpha > v$. Note that the scaling map $\phi : \omega \mapsto \hat{\omega}$ given by

$$\hat{\omega}(\mathbf{x}) = \omega(\nu^{1/2}\mathbf{x})$$

maps the set $S_{M,\nu,1}$ onto the set $S_{\nu^{-3/2}M}$ bijectively and the set $S_{M,\alpha,1}$ onto the set $S_{\nu^{-3/2}M,\alpha/\nu,1}$ bijectively. As $M\nu^{-3/2} < M_2$ and $\alpha/\nu > 1$, by Lemma 5.4 we have

$$S_{v^{-3/2}M} = S_{v^{-3/2}M, \alpha/\nu, 1},$$

and therefore

$$S_{M,\nu,1} \ = \ S_{M,\alpha,1}.$$

5.3. Proof of Theorem 5.1.

Proof of Theorem 5.1. We first prove (i),(ii) by the following claim.

(Claim) There exists a constant $\nu \in (0, \infty)$ such that we have

$$I_{1,\infty} = I_{1,\nu,1} = I_{1,\alpha,1}$$
 and $S_{1,\infty} = S_{1,\nu,1} = S_{1,\alpha,1}$ for each $\alpha > \nu$.

First, observe that

$$K_{1,\infty} = \bigcup_{\tau>0} K_{1,\tau,1}.$$

Then we have

$$I_{1,\infty} = \sup \left\{ E(\omega) : \omega \in \bigcup_{\tau > 0} K_{1,\tau,1} \right\} = \sup \left[\bigcup_{\tau > 0} \left\{ E(\omega) : \omega \in K_{1,\tau,1} \right\} \right] \leq \limsup_{\tau \to \infty} I_{1,\tau,1}.$$

For the constant $M_2 > 0$ in Lemma 5.4, we fix any v > 0 to satisfy $v^{-3/2} < M_2$. Then by Corollary 5.5, we get

$$I_{1,\nu,1} = I_{1,\alpha,1}$$
 and $S_{1,\nu,1} = S_{1,\alpha,1}$

for any $\alpha > \nu$. So we obtain

$$I_{1,\infty} \leq I_{1,\nu,1} < \infty$$
.

Moreover, observe that $I_{1,\nu,1} \leq I_{1,\infty}$ by the relation $K_{1,\nu,1} \subset K_{1,\infty}$. So we get

$$I_{1,\infty} = I_{1,\nu,1} = I_{1,\alpha,1}$$
 for each $\alpha > \nu$.

Using this, we observe that $S_{1,\nu,1} \subset S_{1,\infty}$, with relation $S_{1,\nu,1} = S_{1,\alpha,1}$ for each $\alpha > \nu$. In addition, we can show $S_{1,\nu,1} \supset S_{1,\infty}$ in the following way. We choose any $\omega \in S_{1,\infty}$ and put $\alpha' := \max\{\nu, \|\omega\|_1\}$ to get

$$\omega \in K_{1,\alpha',1}$$
 and $E(\omega) = I_{1,\infty} = I_{1,\nu,1} = I_{1,\alpha',1}$,

so we obtain $\omega \in S_{1,\alpha',1}$. By the relation $S_{1,\nu,1} = S_{1,\alpha',1}$ due to $\alpha' > \nu$, we have $\omega \in S_{1,\nu,1}$. Therefore we obtain

$$S_{1,\infty} = S_{1,\nu,1} = S_{1,\alpha,1}$$
 for each $\alpha > \nu$.

It completes the proof of the claim.

Note that the above claim implies that $I_{1,\infty} < \infty$ and the set $S_{1,\infty}$ is nonempty. By Corollary 3.4, for each $\omega \in S_{1,\infty}$ there exists $W = W(\omega) > 0$ and $\gamma = \gamma(\omega) \ge 0$ satisfying that

$$\omega = \mathbf{1}_{\{\mathcal{G}[\omega] - Wx_2 - \gamma > 0\}},$$

where we have $\gamma = 0$ due to the relations $\omega \in S_{1,\infty} = S_{1,\nu,1} = S_{1,2\nu,1}$ and $\|\omega\|_1 \le \nu < 2\nu$. Moreover, Corollary 3.4 guarantees that ω may be assumed to satisfy the Steiner symmetry condition up to translation. By Corollary 3.23, we obtain the relation

$$I_{1,\infty}=\frac{3}{4}W,$$

which implies that the traveling speed is unique in the class $S_{1,\infty}$. Moreover, for the scaled maximizer $\hat{\omega} \in S_{\nu^{-3/2}}$, we can apply Lemma 4.6 and Lemma 4.7 to obtain that

$$\frac{1}{C_0} \left(v^{-3/2} \right)^{2/3} \leq \|\hat{\omega}\|_1 \leq C_0 \left(v^{-3/2} \right)^{2/3} \quad \text{ and } \sup_{\mathbf{x} \in \text{supp} (\hat{\omega})} |\mathbf{x}| \leq C_1 \cdot \left(v^{-3/2} \right)^{1/3}$$

for the absolute constants $C_0 > 1$, $C_1 > 0$ in Lemma 4.6 and Lemma 4.7 each. Therefore we have

$$\frac{1}{C_0} \leq \|\omega\|_1 \leq C_0 \quad \text{and} \quad \sup_{\mathbf{x} \in \text{supp}(\omega)} |\mathbf{x}| \leq C_1.$$

Note that both inequalities hold for each $\omega \in S_{1,\infty}$. It completes the proof of Theorem 5.1.

ACKNOWLEDGMENTS

KC has been supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT)(grant No. 2022R1A4A1032094, RS-2023-00274499). IJ has supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT) (No. 2022R1C1C1011051).

When we were finalizing the manuscript, we noted the preprint arXiv:2406.09849 by Huang–Tong which obtained the existence of a Sadovskii vortex patch using a fixed-point approach. It is not clear to us whether their vortex patch coincides with the one in Theorem 1.1. They have a better description of the patch boundary while our variational construction of the Sadovskii vortex as the kinetic maximizer is natural in view of its observed dynamical stability.

REFERENCES

- [1] Ken Abe. Existence of vortex rings in Beltrami flows. Comm. Math. Phys., 391(2):873-899, 2022.
- [2] Ken Abe and Kyudong Choi. Stability of Lamb dipoles. Arch. Ration. Mech. Anal., 244(3):877-917, 2022.
- [3] Y.A. Antipov and A.Y. Zemlyanova. Sadovskii vortex in a wedge and the associated riemann-hilbert problem on a torus. arXiv:2010.08118.
- [4] V. I. Arnold. Sur la géométrie différentielle des groupes de Lie de dimension infinie et ses applications à l'hydrodynamique des fluides parfaits. Ann. Inst. Fourier (Grenoble), 16:319–361, 1966.
- [5] T. B. Benjamin. The alliance of practical and analytical insights into the nonlinear problems of fluid mechanics. pages 8–29. Lecture Notes in Math., 503, 1976.
- [6] Jacob Burbea. Motions of vortex patches. Lett. Math. Phys., 6(1):1–16, 1982.
- [7] G. R. Burton. Steady symmetric vortex pairs and rearrangements. Proc. Roy. Soc. Edinburgh Sect. A, 108(3-4):269–290, 1988.
- [8] G. R. Burton. Uniqueness for the circular vortex-pair in a uniform flow. Proc. Roy. Soc. London Ser. A, 452:2343-2350, 1996.
- [9] G. R. Burton. Isoperimetric properties of Lamb's circular vortex-pair. J. Math. Fluid Mech., 7:S68–S80, 2005.
- [10] G. R. Burton. Compactness and stability for planar vortex-pairs with prescribed impulse. J. Differential Equations, 270:547–572, 2021.
- [11] G. R. Burton, H. J. Nussenzveig Lopes, and M. C. Lopes Filho. Nonlinear stability for steady vortex pairs. <u>Comm. Math. Phys.</u>, 324:445–463, 2013.
- [12] Luis A. Caffarelli and Avner Friedman. The shape of axisymmetric rotating fluid. J. Functional Analysis, 35(1):109-142, 1980.
- [13] Daomin Cao, Shanfa Lai, and Weicheng Zhan. Traveling vortex pairs for 2D incompressible Euler equations. <u>Calc. Var. Partial Differential</u> Equations, 60(5):Paper No. 190, 16, 2021.
- [14] Daomin Cao, Guolin Qin, Weicheng Zhan, and Changjun Zou. Existence and stability of smooth traveling circular pairs for the generalized surface quasi-geostrophic equation. Int. Math. Res. Not. IMRN, (6):4761–4804, 2023.
- [15] Daomin Cao, Guolin Qin, Weicheng Zhan, and Changjun Zou. Global solutions for the generalized SQG equation and rearrangements. <u>Trans. Amer. Math. Soc.</u>, 376(3):2181–2211, 2023.
- [16] Daomin Cao, Jie Wan, and Guodong Wang. Nonlinear orbital stability for planar vortex patches. <u>Proc. Amer. Math. Soc.</u>, 147(2):775–784, 2019.
- [17] Michael Carley. A triangulated vortex method for the axisymmetric Euler equations. J. Comput. Phys., 180(2):616–641, 2002.
- [18] Angel Castro, Diego Córdoba, and Javier Gómez-Serrano. Uniformly rotating analytic global patch solutions for active scalars. <u>Ann. PDE</u>, 2(1):Art. 1, 34, 2016.
- [19] M. Cheng, J. Lou, and T. T. Lim. Numerical simulation of head-on collision of two coaxial vortex rings. <u>Fluid Dyn. Res.</u>, 50(6):065513, 24, 2018
- [20] S. Chernyshenko. Stratified Sadovskii flow in a channel. J. Fluid Mech., 250:423-431, 1993.
- [21] S. I. Chernyshenko. Asymptotics of a steady separated flow around a body with large Reynolds numbers. Prikl. Mat. Mekh., 52(6):958–966,
- [22] S. I. Chernyshenko. Cdensity-stratified sadovskii flow in a channel. Fluid Dyn, 28:524–528, 1993.
- [23] Stephen Childress. Solutions of Euler's equations containing finite eddies. Phys. Fluids, 9:860-872, 1966.
- [24] Stephen Childress. Nearly two-dimensional solutions of Euler's equations. The Physics of Fluids, 30(4):944–953, 04 1987.
- [25] Stephen Childress. Models of vorticity growth in Euler flows I, Axisymmetric flow without swirl and II, Almost 2-D dynamics. <u>AML reports</u> 05-07 and 06-07, courant institute of mathematical sciences, 2007.
- [26] Stephen Childress. Growth of anti-parallel vorticity in Euler flows. Phys. D, 237(14-17):1921–1925, 2008.
- [27] Stephen Childress and Andrew D. Gilbert. Eroding dipoles and vorticity growth for Euler flows in \mathbb{R}^3 : the hairpin geometry as a model for finite-time blowup. Fluid Dyn. Res., 50(1):011418, 40, 2018.
- [28] Stephen Childress, Andrew D. Gilbert, and Paul Valiant. Eroding dipoles and vorticity growth for Euler flows in \mathbb{R}^3 : axisymmetric flow without swirl. J. Fluid Mech., 805:1–30, 2016.
- [29] Kyudong Choi. Stability of Hill's spherical vortex. Comm. Pure Appl. Math., 77(1):52–138, 2024.
- [30] Kyudong Choi and In-Jee Jeong. On vortex stretching for anti-parallel axisymmetric flows. Amer. J. Math., to appear.
- [31] Kyudong Choi and In-Jee Jeong. Infinite growth in vorticity gradient of compactly supported planar vorticity near Lamb dipole. Nonlinear Anal. Real World Appl., 65:Paper No. 103470, 20, 2022.
- [32] Kyudong Choi and In-Jee Jeong. Stability and instability of Kelvin waves. <u>Calc. Var. Partial Differential Equations</u>, 61(6):Paper No. 221, 38, 2022
- [33] Kyudong Choi and Deokwoo Lim. Stability of radially symmetric, monotone vorticities of 2D Euler equations. Calc. Var. Partial Differential Equations, 61(4):Paper No. 120, 27, 2022.
- [34] C. Chu, C. Wang, C. Chang, R. Chang, and W. Chang. Head-on collision of two coaxial vortex rings: Experiment and computation. <u>Journal</u> of Fluid Mechanics, 296:39–71, 1995.

- [35] H. T. Croft. Three lattice-point problems of Steinhaus, Quart. J. Math. Oxford Ser. (2), 33(129):71–83, 1982.
- [36] Juan Dávila, Manuel del Pino, Monica Musso, and Juncheng Wei. Leapfrogging vortex rings for the three-dimensional incompressible euler equations. Communications on Pure and Applied Mathematics, to appear.
- [37] Tarek M. Elgindi and In-Jee Jeong. On singular vortex patches, I: Well-posedness issues. Mem. Amer. Math. Soc., 283(1400):v+89, 2023.
- [38] Lawrence C. Evans. Partial differential equations, volume 19 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, second edition, 2010.
- [39] J.B. Flor and G. J. F. Van Heijst. An experimental study of dipolar vortex structures in a stratified fluid. J. Fluid Mech., 279:101–133, 1994.
- [40] L. E. Fraenkel and M. S. Berger. A global theory of steady vortex rings in an ideal fluid. Acta Math., 132:13-51, 1974.
- [41] Daniel V. Freilich and Stefan G. Llewellyn Smith. The Sadovskii vortex in strain. J. Fluid Mech., 825:479–501, 2017.
- [42] A. Friedman and B. Turkington. Vortex rings: existence and asymptotic estimates. Trans. Amer. Math. Soc., 268:1–37, 1981.
- [43] Thierry Gallay and Vladimír Šverák. Arnold's variational principle and its application to the stability of planar vortices. <u>Anal. PDE</u>, 17(2):681–722, 2024.
- [44] Claudia García and Susanna V. Haziot. Global bifurcation for corotating and counter-rotating vortex pairs. <u>Comm. Math. Phys.</u>, 402(2):1167–1204, 2023.
- [45] B. Gidas, W. M. Ni, and L. Nirenberg. Symmetry and related properties via the maximum principle. Comm. Math. Phys., 68:209–243, 1979.
- [46] David Gilbarg and Neil S. Trudinger. <u>Elliptic partial differential equations of second order</u>. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [47] M. A. Goldshtik. A mathematical model of discontinuous incompressible fluid flows. Dokl. Akad. Nauk SSSR, 147(6):1310–1313, 1962.
- [48] Javier Gómez-Serrano, Jaemin Park, Jia Shi, and Yao Yao. Symmetry in stationary and uniformly rotating solutions of active scalar equations. Duke Math. J., 170(13):2957–3038, 2021.
- [49] Hui Guan, Zhi-Jun Wei, Elizabeth Rumenova Rasolkova, and Chui-Jie Wu. Numerical simulations of two coaxial vortex rings head-on collision. Adv. Appl. Math. Mech., 8(4):616–647, 2016.
- [50] Stephen Gustafson, Evan Miller, and Tai-Peng Tsai. Growth rates for anti-parallel vortex tube Euler flows in three and higher dimensions. arXiv:2303.12043.
- [51] Ummu Habibah and Yasuhide Fukumoto. A counter-rotating vortex pair in inviscid fluid. <u>AIP Conference Proceedings</u>, 1913(1):020022, 12 2017.
- [52] Zineb Hassainia, Taoufik Hmidi, and Nader Masmoudi. Rigorous derivation of the leapfrogging motion for planar euler equations. arXiv:2311.15765.
- [53] Zineb Hassainia, Nader Masmoudi, and Miles H. Wheeler. Global bifurcation of rotating vortex patches. <u>Comm. Pure Appl. Math.</u>, 73(9):1933–1980, 2020.
- [54] G. J. F. Van Heijst and J. B. Flor. Dipole formation and collisions in a stratified fluid. Nature, 340:212-215, 1989.
- [55] H. Helmholtz. Über Integrale der hydrodynamischen Gleichungen, welche den Wirbelbewegungen entsprechen. <u>J. Reine Angew. Math.</u>, 55:25–55, 1858.
- [56] W. M. Hicks. Researches in vortex motion. part iii: On spiral or gyrostatic vortex aggregates. Phil. Trans. R. Soc. A., 192:33–99, 1899.
- [57] M. J. M. Hill. On a spherical vortex. Philos. Trans. Roy. Soc. London Ser. A, 185:213-245, 1894.
- [58] Taoufik Hmidi and Joan Mateu. Existence of corotating and counter-rotating vortex pairs for active scalar equations. <u>Comm. Math. Phys.</u>, 350(2):699–747, 2017.
- [59] Taoufik Hmidi, Joan Mateu, and Joan Verdera. Boundary regularity of rotating vortex patches. <u>Arch. Ration. Mech. Anal.</u>, 209(1):171–208, 2013
- [60] T. Kambe and T. Minota. Acoustic wave radiated by head-on collision of two vortex rings. Proc. R. Soc. Lond. A, 386:277-308, 1983.
- [61] T. Kambe and U. Mya Oo. An axisymmetric viscous vortex motion and its acoustic emission. J. Phys. Soc. Japan, 53:2263-2273, 1984.
- [62] Sun-Chul Kim. Asymptotic study of Navier-Stokes flows. <u>Trends in Mathematics, information center for Mathematical sciences</u>, 6:29–33, 2003.
- [63] V. I. Kolyada. On the metric Darboux property. Anal. Math., 9(4):291–312, 1983.
- [64] H. Lamb. Hydrodynamics. Cambridge Univ. Press., 3rd ed. edition, 1906.
- [65] Thomas Leweke, Stéphane Le Dizès, and Charles H. K. Williamson. Dynamics and instabilities of vortex pairs. In Annual review of fluid mechanics. Vol. 48, volume 48 of Annu. Rev. Fluid Mech., pages 507–541. Annual Reviews, Palo Alto, CA, 2016.
- [66] T. T. Lim and T. B. Nickels. Instability and reconnection in the head-on collision of two vortex rings. Nature, 357:225-227, 1992.
- [67] P. Luzzatto-Fegiz and C. H. K. Williamson. Determining the stability of steady two-dimensional flows through imperfect velocity-impulse diagrams. J. Fluid Mech., 706:323–350, 2012.
- [68] Andrew J. Majda and Andrea L. Bertozzi. <u>Vorticity and incompressible flow</u>, volume 27 of <u>Cambridge Texts in Applied Mathematics</u>. Cambridge University Press, Cambridge, 2002.
- [69] Carlo Marchioro and Mario Pulvirenti. Mathematical theory of incompressible nonviscous fluids, volume 96 of Applied Mathematical Sciences. Springer-Verlag, New York, 1994.
- [70] H. K. Moffatt. Degree of knottedness of tangled vortex lines. <u>J. Fluid Mech.</u>, 35:117–129, 1969.
- [71] D. W. Moore, P. G. Saffman, and S. Tanveer. The calculation of some Batchelor flows: The Sadovskii vortex and rotational corner flow. <u>The Physics of Fluids</u>, 31(5):978–990, 05 1988.
- [72] J. Norbury. Steady planar vortex pairs in an ideal fluid. Comm. Pure Appl. Math., 28:679–700, 1975.
- [73] Y. Oshima. Head-on collision of two vortex rings. J. Phys. Soc. Japan, 44:328–331, 1978.
- [74] Edward A. Overman, II. Steady-state solutions of the Euler equations in two dimensions. II. Local analysis of limiting *V*-states. <u>SIAM J. Appl. Math.</u>, 46(5):765–800, 1986.
- [75] A. J. Peace and N. Riley. A viscous vortex pair in ground effect. J. Fluid Mech., 129:409-426, 1983.
- [76] R. T. Pierrehumbert. A family of steady, translating vortex pairs with distributed vorticity. Journal of Fluid Mechanics, 99(1):129–144, 1980.

- [77] S. I. Pohožaev. On the eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$. Dokl. Akad. Nauk SSSR, 165:36–39, 1965.
- [78] V.S. Sadovskii. Vortex regions in a potential stream with a jump of bernoulli's constant at the boundary: Pmm vol. 35, no. 5, 1971, pp. 773–779. Journal of Applied Mathematics and Mechanics, 35(5):729–735, 1971.
- [79] P. G. Saffman. Vortex dynamics. Cambridge Monographs on Mechanics and Applied Mathematics. Cambridge University Press, New York, 1992.
- [80] P. G. Saffman and R. Szeto. Equilibrium shapes of a pair of equal uniform vortices. Phys. Fluids, 23(12):2339–2342, 1980.
- [81] P. G. Saffman and S. Tanveer. The touching pair of equal and opposite uniform vortices. Phys. Fluids, 25(11):1929–1930, 1982.
- [82] K. Shariff and A. Leonard. Vortex Rings. Ann. Rev. Fluid Mech., 24:235–279, 1992.
- [83] Karim Shariff, Anthony Leonard, and Joel H. Ferziger. A contour dynamics algorithm for axisymmetric flow. <u>J. Comput. Phys.</u>, 227(21):9044–9062, 2008.
- [84] Karim Shariff, Anthony Leonard, Norman J. Zabusky, and Joel H. Ferziger. Acoustics and dynamics of coaxial interacting vortex rings. <u>Fluid</u> Dynamics Research, 3(1):337–343, 1988.
- [85] Banavara N. Shashikanth and Jerrold E. Marsden. Leapfrogging vortex rings: Hamiltonian structure, geometric phases and discrete reduction. Fluid Dynam. Res., 33(4):333–356, 2003.
- [86] M. J. Shelley, D. I. Meiron, and S. A. Orszag. Dynamical aspects of vortex reconnection of perturbed anti-parallel vortex tubes. <u>J. Fluid Mech.</u>, 246:613–652, 1993.
- [87] $\overline{\text{T. C.}}$ Sideris and L. Vega. Stability in L^1 of circular vortex patches. Proc. Amer. Math. Soc., 137:4199–4202, 2009.
- [88] Didier Smets and Jean Van Schaftingen. Desingularization of vortices for the Euler equation. Arch. Ration. Mech. Anal., 198(3):869–925, 2010
- [89] F.T. Smith. Concerning inviscid solutions for large-scale separated flows. J. Engrg. Math., 20:271-292, 1986.
- [90] S. Stanaway, K. Shariff, and F. Hussain. Head-on collision of viscous vortex rings. <u>Proceedings of the Summer Program, Center for Turbulence Research</u>, 1988.
- [91] András Szenes. Exceptional points for Lebesgue's density theorem on the real line. Adv. Math., 226(1):764–778, 2011.
- [92] Yun Tang. Nonlinear stability of vortex patches. Trans. Amer. Math. Soc., 304(2):617-638, 1987.
- [93] W. Thomson (Lord Kelvin). Maximum and minimum energy in vortex motion, Nature 22, no. 574, 618–620 (1880). In <u>Mathematical and</u> Physical Papers 4, pages 172–183. Cambridge: Cambridge University Press, 1910.
- [94] B. Turkington. On steady vortex flow in two dimensions. I, II. Comm. Partial Differential Equations, 8:999–1030, 1031–1071, 1983.
- [95] Christopher TW and Llewellyn Smith SG. Hollow vortex in a corner. Journal of Fluid Mechanics., 908, 2021.
- [96] Y. H. Wan. The stability of rotating vortex patches. Comm. Math. Phys., 107:1-20, 1986.
- [97] Y. H. Wan and M. Pulvirenti. Nonlinear stability of circular vortex patches. Comm. Math. Phys., 99:435-450, 1985.
- [98] Guodong Wang. On concentrated traveling vortex pairs with prescribed impulse. Trans. Amer. Math. Soc., 377(4):2635–2661, 2024.
- [99] Richard L. Wheeden and Antoni Zygmund. Measure and integral. Pure and Applied Mathematics (Boca Raton). CRC Press, Boca Raton, FL, second edition, 2015. An introduction to real analysis.
- [100] H. M. Wu, E. A. Overman, II, and N. J. Zabusky. Steady-state solutions of the Euler equations in two dimensions: rotating and translating V-states with limiting cases. I. Numerical algorithms and results. J. Comput. Phys., 53(1):42–71, 1984.
- [101] J. Yang. Existence and asymptotic behavior in planar vortex theory. Math. Models Methods Appl. Sci., 1:461–475, 1991.
- [102] Joseph Yang and Toshi Kubota. The steady motion of a symmetric, finite core size, counterrotating vortex pair. <u>SIAM J. Appl. Math.</u>, 54(1):14–25, 1994.

Department of Mathematical Sciences, Ulsan National Institute of Science and Technology, 50 UNIST-gil, Eonyang-eup, Ulju-gun, Ulsan 44919, Republic of Korea.

Email address: kchoi@unist.ac.kr

DEPARTMENT OF MATHEMATICAL SCIENCES AND RIM, SEOUL NATIONAL UNIVERSITY, 1 GWANAK-RO, GWANAK-GU, SEOUL 08826, AND SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY, REPUBLIC OF KOREA.

Email address: injee_j@snu.ac.kr

DEPARTMENT OF MATHEMATICAL SCIENCES, ULSAN NATIONAL INSTITUTE OF SCIENCE AND TECHNOLOGY, 50 UNIST-GIL, EONYANG-EUP, ULJU-GUN, ULSAN 44919, REPUBLIC OF KOREA.

Email address: yj.sim@unist.ac.kr