

Note on a vector-host epidemic model with spatial structure¹

Mingxin Wang²

School of Mathematics and Information Science, Henan Polytechnic University, Jiaozuo 454003,
China

Abstract. Magal, Webb and Wu [Nonlinearity 31, 5589-5614 (2018)] studied the outbreak of Zika in Rio De Janerio model, and provided complete dynamic properties. In this note we will use a very simple approach to provide a complete proof.

Keywords: Epidemic models; Equilibrium solutions; Global stabilities.

AMS Subject Classification (2020): 35K57; 92D30

1 Introduction

The Zika virus has caused significant public health impacts, including congenital microcephaly and neurodevelopmental disorders after pregnancy infection. Suppose that individuals are living in a bounded smooth domain $\Omega \subset \mathbb{R}^n$. Let H_i, V_u and V_i be the densities of infected hosts, uninfected vectors, and infected vectors, respectively. Then the model proposed in [1] to study the outbreak of Zika in Rio De Janerio is the following reaction-diffusion system

$$\begin{cases} \partial_t H_i - \nabla \cdot d_1(x) \nabla H_i = -\rho(x) H_i + \sigma_1(x) H_u(x) V_i, & x \in \Omega, t > 0, \\ \partial_t V_u - \nabla \cdot d_2(x) \nabla V_u = -\sigma_2(x) V_u H_i + \beta(x) (V_u + V_i) - \mu(x) (V_u + V_i) V_u, & x \in \Omega, t > 0, \\ \partial_t V_i - \nabla \cdot d_2(x) \nabla V_i = \sigma_2(x) V_u H_i - \mu(x) (V_u + V_i) V_i, & x \in \Omega, t > 0, \\ \frac{\partial H_i}{\partial n} = \frac{\partial V_u}{\partial n} = \frac{\partial V_i}{\partial n} = 0, & x \in \partial\Omega, t > 0, \\ (H_i(\cdot, 0), V_u(\cdot, 0), V_i(\cdot, 0)) = (H_{i0}, V_{u0}, V_{i0}) \in C(\bar{\Omega}; \mathbb{R}_+^3), \end{cases} \quad (1)$$

where $d_1, d_2 \in C^{1+\alpha}(\bar{\Omega})$ and $\rho, \beta, \sigma_1, \sigma_2, \mu \in C^\alpha(\bar{\Omega})$ are strictly positive, and $H_u \in C^\alpha(\bar{\Omega})$ is nonnegative and non-trivial. The flux of new infected humans is given by $\sigma_1(x) H_u(x) V_i(x, t)$ in which $H_u(x)$ is the density of susceptible population depending on the spatial location x . The initial functions satisfy $\frac{\partial H_{i0}}{\partial n} = \frac{\partial V_{u0}}{\partial n} = \frac{\partial V_{i0}}{\partial n} = 0$ on $\partial\Omega$.

The model (1) has been thoroughly studied by Magal, Webb and Wu [2, 3]. In [2], the authors found the basic reproduction number R_0 and show that R_0 is a threshold parameter: if $R_0 \leq 1$ the disease free equilibrium is globally stable; if $R_0 > 1$ the model has a unique globally stable positive equilibrium. In [3], it was shown that the basic reproduction number R_0 can be defined as the spectral radius of a product of a local basic reproduction number R and strongly positive compact linear operators with spectral radii one.

In this note we will use a very simple approach to prove the main results of [2].

¹This work was supported by NSFC Grant 12171120.

²Corresponding author. *E-mail:* mxwang@hpu.edu.cn

2 Existence of positive equilibrium solutions

The equilibrium problem of (1) is

$$\begin{cases} -\nabla \cdot d_1(x) \nabla H_i^* = -\rho(x) H_i^* + \sigma_1(x) H_u(x) V_i^*, & x \in \Omega, \\ -\nabla \cdot d_2(x) \nabla V_u^* = -\sigma_2(x) V_u^* H_i^* + \beta(x) (V_u^* + V_i^*) - \mu(x) (V_u^* + V_i^*) V_u^*, & x \in \Omega, \\ -\nabla \cdot d_2(x) \nabla V_i^* = \sigma_2(x) V_u H_i - \mu(x) (V_u^* + V_i^*) V_i^*, & x \in \Omega, \\ \frac{\partial H_i^*}{\partial n} = \frac{\partial V_u^*}{\partial n} = \frac{\partial V_i^*}{\partial n} = 0, & x \in \partial\Omega. \end{cases} \quad (2)$$

Let $\lambda(\beta)$ be the principal eigenvalue of

$$\begin{cases} -\nabla \cdot d_2(x) \nabla \phi - \beta(x) \phi = \lambda \phi, & x \in \Omega, \\ \frac{\partial \phi}{\partial n} = 0, & x \in \partial\Omega. \end{cases}$$

If $\lambda(\beta) \geq 0$, then $(0, 0, 0)$ is the only nonnegative solution of (2). In fact, let (H_i^*, V_u^*, V_i^*) be a nonnegative solution of (2) and set $V_* = V_u^* + V_i^*$. Then

$$\begin{cases} -\nabla \cdot d_2(x) \nabla V_* = \beta(x) V_* - \mu(x) V_*^2, & x \in \Omega, \\ \frac{\partial V_*}{\partial n} = 0, & x \in \partial\Omega. \end{cases} \quad (3)$$

It is well known that (3) has no positive solution when $\lambda(\beta) \geq 0$. Therefore, $V_* = 0$, i.e., $V_u^* = V_i^* = 0$, and then $H_i^* = 0$ by the first equation of (2).

If $\lambda(\beta) < 0$, then the problem (3) has a unique positive solution $V_*(x)$ and $V_*(x)$ is globally asymptotically stable. Thus, $(0, V_*(x), 0)$ is a nontrivial and nonnegative solution of (2). It is easy to see that $(0, V_*(x), 0)$ is the only possible semi-trivial nonnegative solution of (2).

Assume $\lambda(\beta) < 0$ and let $V_*(x)$ be the unique positive solution of (3). To investigate positive solutions of (2) is equivalent to study positive solutions (H_i^*, V_i^*) with $V_i^* \leq V_*$ of

$$\begin{cases} -\nabla \cdot d_1(x) \nabla H_i^* = -\rho(x) H_i^* + \sigma_1(x) H_u(x) V_i^*, & x \in \Omega, \\ -\nabla \cdot d_2(x) \nabla V_i^* = \sigma_2(x) (V_*(x) - V_i^*) H_i^* - \mu(x) V_*(x) V_i^*, & x \in \Omega, \\ \frac{\partial H_i^*}{\partial n} = \frac{\partial V_i^*}{\partial n} = 0, & x \in \partial\Omega \end{cases} \quad (4)$$

with $V_i^* \leq V_*$. Clearly, $(0, 0)$ is the unique trivial nonnegative solution of (4). The linearized eigenvalue problem of (4) at $(0, 0)$ is

$$\begin{cases} -\nabla \cdot d_1(x) \nabla \phi + \rho(x) \phi - \sigma_1(x) H_u(x) \varphi = \lambda \phi, & x \in \Omega, \\ -\nabla \cdot d_2(x) \nabla \varphi - \sigma_2(x) V_*(x) \phi + \mu(x) V_*(x) \varphi = \lambda \varphi, & x \in \Omega, \\ \frac{\partial \phi}{\partial n} = \frac{\partial \varphi}{\partial n} = 0, & x \in \partial\Omega \end{cases} \quad (5)$$

By the results of [4], (5) has a unique principle eigenvalue $\lambda(V_*)$ with eigenfunction $(\phi, \varphi) : \phi, \varphi > 0$. Moreover, $\lambda(V_*)$ has the same sign with $R_0 - 1$, where R_0 is given in [2].

Theorem 2.1. *The problem (4) has a positive solution (H_i^*, V_i^*) if and only if $\lambda(V_*) < 0$. Moreover, the positive solution (H_i^*, V_i^*) of (4) is unique and satisfies $V_i^* < V_*$ when it exists. Therefore, (2) has a positive solution (H_i^*, V_i^*, V_u) if and only if $\lambda(V_*) < 0$, and (H_i^*, V_i^*, V_u) is unique and takes the form $(H_i^*, V_i^*, V_* - V_i^*)$ when it exists.*

Proof. Step 1: We prove that if (4) has a positive solution (H_i^*, V_i^*) , then $\lambda(V_*) < 0$. Define

$$\mathcal{L}\psi = \begin{pmatrix} -\nabla \cdot d_1(x)\nabla\psi_1 + \rho(x)\psi_1, \\ -\nabla \cdot d_2(x)\nabla\psi_2 + \mu(x)V_*(x)\psi_2 \end{pmatrix}, \quad A(x) = \begin{pmatrix} \lambda(V_*) & \sigma_1(x)H_u(x) \\ \sigma_2(x)V_*(x) & \lambda(V_*) \end{pmatrix}.$$

Then \mathcal{L} is reversible and \mathcal{L}^{-1} is a strongly positive compact operator. By (5), $(\phi, \varphi) = (\mathcal{L}^{-1}A(x))(\phi, \varphi)$. If $\lambda(V_*) \geq 0$, then the operator $T = \mathcal{L}^{-1}A(x)$ is also a strongly positive compact operator, and $r(T) = 1$ as $(\phi, \varphi) = (\mathcal{L}^{-1}A(x))(\phi, \varphi)$.

As (H_i^*, V_i^*) is a positive solution of (4), then we have

$$\begin{pmatrix} H_i^* \\ V_i^* \end{pmatrix} = T \begin{pmatrix} H_i^* \\ V_i^* \end{pmatrix} - \mathcal{L}^{-1} \begin{pmatrix} \lambda(V_*) & 0 \\ \sigma_2(x)V_i^* & \lambda(V_*) \end{pmatrix} \begin{pmatrix} H_i^* \\ V_i^* \end{pmatrix},$$

and

$$\mathcal{L}^{-1} \begin{pmatrix} \lambda(V_*) & 0 \\ \sigma_2(x)V_i H_i & \lambda(V_*) \end{pmatrix} \begin{pmatrix} H_i^* \\ V_i^* \end{pmatrix} \geq \mathcal{L}^{-1} \begin{pmatrix} 0 \\ \sigma_2(x)V_i H_i \end{pmatrix} =: \begin{pmatrix} 0 \\ \chi(x) \end{pmatrix},$$

where $\chi(x) > 0$. This is impossible by the conclusion [5, Theorem 3.2 (iv)].

Step 2: We prove that if $\lambda(V_*) < 0$, then (4) has at least one positive solution (H_i^*, V_i^*) . Let \bar{H}_i^* be the unique positive solution of the linear problem

$$\begin{cases} -\nabla \cdot d_1(x)\nabla\bar{H}_i^* + \rho(x)\bar{H}_i^* = \sigma_1(x)H_u(x)V_*(x), & x \in \Omega, \\ \frac{\partial\bar{H}_i^*}{\partial n} = 0, & x \in \partial\Omega. \end{cases}$$

Then (\bar{H}_i^*, V_*) is an upper solution of (4). Let (ϕ, φ) be the positive eigenfunction corresponding to $\lambda(V_*)$ and $\varepsilon > 0$ be a constant. It is easy to verify that $\varepsilon(\phi, \varphi)$ is a lower solution of (4) and $\varepsilon(\phi, \varphi) \leq (\bar{H}_i^*, V_*)$ provided $\varepsilon > 0$ is suitable small. By the upper and lower solutions method, (4) has at least one positive solution.

Step 3: We shall prove the uniqueness of positive solution of (4). Let (H_i^*, V_i^*) be a positive solution of (4). Then $\lambda(V_*) < 0$ by Step 1. Consider the linear problem

$$\begin{cases} -\nabla \cdot d_2(x)\nabla U = \sigma_2(x)(V_*(x) - U)H_i^* - \mu(x)V_*(x)U, & x \in \Omega, \\ \frac{\partial U}{\partial n} = 0, & x \in \partial\Omega. \end{cases} \quad (6)$$

Obviously, V_* is a strict upper solution of (6), and there exists $0 < \varepsilon \ll 1$ such that $\varepsilon\varphi \leq V_*$ and $\varepsilon(\sigma_2 V_* \phi + \lambda(V_*)\varphi) \leq \sigma_2(V_* - \varepsilon\phi)H_i^*$ in Ω . Then we can verify that $\varepsilon\varphi$ is a lower solution of (6). Thus (6) has a solution $U : \varepsilon\varphi \leq U < V_*$. Since V_i^* solves (6), we see $V_i^* = U < V_*$ by the uniqueness. It then follows from the first equation of (4) that $H_i^* < \bar{H}_i^*$.

Noticing that (\bar{H}_i^*, V_*) is an upper solution of (4). By the monotone iterative method, (4) has a positive solution $(\hat{H}_i^*, \hat{V}_i^*)$ and $(H_i^*, V_i^*) \leq (\hat{H}_i^*, \hat{V}_i^*) \leq (\bar{H}_i^*, V_*)$. We shall prove $(H_i^*, V_i^*) = (\hat{H}_i^*, \hat{V}_i^*)$. Clearly, we can find $\zeta > 1$ such that $\zeta(H_i^*, V_i^*) \geq (\hat{H}_i^*, \hat{V}_i^*)$ in Ω . Set

$$\underline{\zeta} = \inf\{\zeta \geq 1 : \zeta(H_i^*, V_i^*) \geq (\hat{H}_i^*, \hat{V}_i^*) \text{ in } \Omega\}.$$

Then $\underline{\zeta}$ is well defined, $\underline{\zeta} \geq 1$ and $\underline{\zeta}(H_i^*, V_i^*) \geq (\hat{H}_i^*, \hat{V}_i^*)$ in Ω . We shall prove $\underline{\zeta} = 1$. Assume $\underline{\zeta} > 1$. Let $u = \underline{\zeta}H_i^* - \hat{H}_i^*$ and $v = \underline{\zeta}V_i^* - \hat{V}_i^*$. Then $u, v \geq 0$, and by the carefully calculations,

$$\begin{cases} -\nabla \cdot d_1(x)\nabla u + \rho(x)u = \sigma_1(x)H_u(x)v, & x \in \Omega, \\ -\nabla \cdot d_2(x)\nabla v + (\sigma_2H_i^* + \mu(x)V_*)v > \sigma_2(V_* - \hat{V}_i^*)u, & x \in \Omega, \\ \frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = 0, & x \in \partial\Omega \end{cases} \quad (7)$$

It follows that $u, v > 0$ in $\bar{\Omega}$ by the maximum principle. We can find $\varepsilon > 0$ such that $(u, v) \geq \varepsilon(H_i^*, V_i^*)$, i.e., $(\underline{\zeta} - \varepsilon)(H_i^*, V_i^*) \geq (\hat{H}_i^*, \hat{V}_i^*)$ in Ω . This is contradicts to the definition of $\underline{\zeta}$. Then $\underline{\zeta} = 1$, i.e., $(H_i^*, V_i^*) = (\hat{H}_i^*, \hat{V}_i^*)$. The proof is complete. \square

3 Stabilities of nonnegative solutions of (2)

Theorem 3.1. *Let (H_i, V_u, V_i) be the unique positive solution of (1). Then the following holds.*

(i) *If $\lambda(\beta) < 0$ and $\lambda(V_*) < 0$, then*

$$\lim_{t \rightarrow \infty} (H_i, V_u, V_i) = (H_i^*, V_* - V_i^*, V_i^*) = (H_i^*, V_u^*, V_i^*) \text{ in } [C^2(\bar{\Omega})]^3. \quad (8)$$

(ii) *If $\lambda(\beta) < 0$ and $\lambda(V_*) \geq 0$, then*

$$\lim_{t \rightarrow \infty} (H_i, V_u, V_i) = (0, V_*, 0) \text{ in } [C^2(\bar{\Omega})]^3. \quad (9)$$

(iii) *If $\lambda(\beta) \geq 0$, then*

$$\lim_{t \rightarrow \infty} (H_i, V_u, V_i) = (0, 0, 0) \text{ in } [C^2(\bar{\Omega})]^3. \quad (10)$$

Proof. (i) Let $\varepsilon > 0$ and $\lambda(V_* \pm \varepsilon)$ be the principal eigenvalue of

$$\begin{cases} -\nabla \cdot d_1(x)\nabla \phi + \rho(x)\phi - \sigma_1(x)H_u(x)\varphi = \lambda\phi, & x \in \Omega, \\ -\nabla \cdot d_2(x)\nabla \varphi - \sigma_2(x)(V_*(x) \pm \varepsilon)\phi + \mu(x)(V_*(x) \mp \varepsilon)\varphi = \lambda\varphi, & x \in \Omega, \\ \frac{\partial \phi}{\partial n} = \frac{\partial \varphi}{\partial n} = 0, & x \in \partial\Omega. \end{cases} \quad (11_{\pm\varepsilon})$$

As $\lambda(V_*) < 0$, there is $0 < \varepsilon_0 \ll 1$ such that, when $0 < \varepsilon \leq \varepsilon_0$, $\lambda(V_* \pm \varepsilon) < 0$, and the problems

$$\begin{cases} -\nabla \cdot d_1(x)\nabla H_i = -\rho(x)H_i + \sigma_1(x)H_u(x)V_i, & x \in \Omega, \\ -\nabla \cdot d_2(x)\nabla V_i = \sigma_2(x)(V_*(x) \pm \varepsilon - V_i)^+ H_i - \mu(x)(V_*(x) \mp \varepsilon)V_i, & x \in \Omega, \\ \frac{\partial H_i}{\partial n} = \frac{\partial V_i}{\partial n} = 0, & x \in \partial\Omega \end{cases} \quad (12_{\pm\varepsilon})$$

has a unique positive solution $(H_{i,\pm\varepsilon}^*, V_{i,\pm\varepsilon}^*)$, and $V_{i,\pm\varepsilon}^* < V_* \pm \varepsilon$ and

$$\lim_{\varepsilon \rightarrow 0} (H_{i,\pm\varepsilon}^*, V_{i,\pm\varepsilon}^*) = (H_i^*, V_i^*) \quad \text{in } [C^2(\bar{\Omega})]^2. \quad (13)$$

Let (H_i, V_u, V_i) be the unique solution of (1), and set $V = V_u + V_i$. Then V satisfies

$$\begin{cases} V_t - \nabla \cdot d_2(x) \nabla V = \beta(x)V - \mu(x)V^2, & x \in \Omega, t > 0, \\ \frac{\partial V}{\partial n} = 0, & x \in \partial\Omega, t > 0, \\ V(x, 0) = V_u(x, 0) + V_i(x, 0) > 0, & x \in \bar{\Omega}, \end{cases}$$

and $\lim_{t \rightarrow \infty} V = V_*$ in $C^2(\bar{\Omega})$. For any given $0 < \varepsilon \leq \varepsilon_0$, there exists $T_\varepsilon \gg 1$ such that

$$0 < V_*(x) - \varepsilon \leq V(x, t) \leq V_*(x) + \varepsilon, \quad \forall t \geq T_\varepsilon, x \in \bar{\Omega}. \quad (14)$$

Let (\bar{H}_i, \bar{V}_i) and $(\underline{H}_i, \underline{V}_i)$ be the unique positive solutions of

$$\begin{cases} \partial_t \bar{H}_i - \nabla \cdot d_1(x) \nabla \bar{H}_i = -\rho(x)\bar{H}_i + \sigma_1(x)H_u(x)\bar{V}_i, & x \in \Omega, t > T_\varepsilon, \\ \partial_t \bar{V}_i - \nabla \cdot d_2(x) \nabla \bar{V}_i = \sigma_2(x)(V_*(x) + \varepsilon - \bar{V}_i)^+ \bar{H}_i - \mu(x)(V_*(x) - \varepsilon)\bar{V}_i, & x \in \Omega, t > T_\varepsilon, \\ \frac{\partial \bar{H}_i}{\partial n} = \frac{\partial \bar{V}_i}{\partial n} = 0, & x \in \partial\Omega, t > T_\varepsilon, \\ \bar{H}_i(x, T_\varepsilon) = kH_{i,+ \varepsilon}^*, \quad \bar{V}_i(x, T_\varepsilon) = kV_{i,+ \varepsilon}^*, & x \in \bar{\Omega}, \end{cases} \quad (15)$$

and

$$\begin{cases} \partial_t \underline{H}_i - \nabla \cdot d_1(x) \nabla \underline{H}_i = -\rho(x)\underline{H}_i + \sigma_1(x)H_u(x)\underline{V}_i, & x \in \Omega, t > T_\varepsilon, \\ \partial_t \underline{V}_i - \nabla \cdot d_2(x) \nabla \underline{V}_i = \sigma_2(x)(V_*(x) - \varepsilon - \underline{V}_i)\underline{H}_i - \mu(x)(V_*(x) + \varepsilon)\underline{V}_i, & x \in \Omega, t > T_\varepsilon, \\ \frac{\partial \underline{H}_i}{\partial n} = \frac{\partial \underline{V}_i}{\partial n} = 0, & x \in \partial\Omega, t > T_\varepsilon, \\ \underline{H}_i(x, T_\varepsilon) = \delta\phi_{-\varepsilon}, \quad \underline{V}_i(x, T_\varepsilon) = \delta\varphi_{-\varepsilon}, & x \in \bar{\Omega}, \end{cases}$$

respectively, where $(\phi_{-\varepsilon}, \varphi_{-\varepsilon})$ is the positive eigenfunction corresponding to $\lambda(V_*; -\varepsilon)$, $k \gg 1$ and $0 < \delta \ll 1$ are constants such that

$$(kH_{i,+ \varepsilon}^*, kV_{i,+ \varepsilon}^*) \geq (H_i(x, T_\varepsilon), V_i(x, T_\varepsilon)), \quad (\delta\phi_{-\varepsilon}, \delta\varphi_{-\varepsilon}) \leq (H_i(x, T_\varepsilon), V_i(x, T_\varepsilon)).$$

It is clear that $(kH_{i,+ \varepsilon}^*, kV_{i,+ \varepsilon}^*)$ is an upper solution of $(12_{+\varepsilon})$ and $(\delta\phi_{-\varepsilon}, \delta\varphi_{-\varepsilon})$ is a lower solution of $(12_{-\varepsilon})$. Hence, $(\bar{H}_i, \bar{V}_i) \geq (\underline{H}_i, \underline{V}_i)$, and (\bar{H}_i, \bar{V}_i) and $(\underline{H}_i, \underline{V}_i)$ are decreasing and increasing in t , respectively. Then

$$\lim_{t \rightarrow \infty} (\bar{H}_i, \bar{V}_i) = (H_{i,+ \varepsilon}^*, V_{i,+ \varepsilon}^*), \quad \lim_{t \rightarrow \infty} (\underline{H}_i, \underline{V}_i) = (H_{i,- \varepsilon}^*, V_{i,- \varepsilon}^*) \quad \text{in } [C^2(\bar{\Omega})]^2 \quad (16)$$

by the uniform estimate and compact arguments (cf. [6, Theorems 2.11, 3.14]).

On the other hand, using (14) and the comparison principle we deduce that (H_i, V_i) satisfies $(\bar{H}_i, \bar{V}_i) \geq (H_i, V_i) \geq (\underline{H}_i, \underline{V}_i)$ for $t \geq T_\varepsilon$ and $x \in \bar{\Omega}$. Hence, by (16),

$$(H_{i,- \varepsilon}^*, V_{i,- \varepsilon}^*) \leq \liminf_{t \rightarrow \infty} (H_i, V_i) \leq \limsup_{t \rightarrow \infty} (H_i, V_i) \leq (H_{i,+ \varepsilon}^*, V_{i,+ \varepsilon}^*) \quad \text{in } [C^2(\bar{\Omega})]^2.$$

This combines with (13) gives $\lim_{t \rightarrow \infty} (H_i, V_i) = (H_i^*, V_i^*)$ in $[C^2(\bar{\Omega})]^2$. As we have shown that $\lim_{t \rightarrow \infty} (V_u + V_i) = V_*$, it is to see that (8) holds.

(ii) Let $\lambda(V_* + \varepsilon)$ be the principal eigenvalue of $(11_{+\varepsilon})$. If $\lambda(V_*) > 0$, then $\lambda(V_* + \varepsilon) > 0$ when $0 < \varepsilon \ll 1$. So $(12_{+\varepsilon})$ has no positive solution. Let (\bar{H}_i, \bar{V}_i) be the unique positive solution of (15) with initial data $\bar{H}_i(x, T_\varepsilon) = C_1, \bar{V}_i(x, T_\varepsilon) = C_2$, where C_1 and C_2 are large positive constants, for example,

$$C_1 \geq \max_{\bar{\Omega}} \max\{V_*(x) + \varepsilon, H_i(x, T_\varepsilon)\}, \quad C_2 \geq \max_{\bar{\Omega}} \max\{C_1 \sigma_1(x) H_u(x) / \rho(x), V_i(x, T_\varepsilon)\}.$$

Then (C_1, C_2) is an upper solution of $(12_{+\varepsilon})$, and (\bar{H}_i, \bar{V}_i) is decreasing in t . Similar to the above, limit $\lim_{t \rightarrow \infty} (\bar{H}_i, \bar{V}_i)$ exists and is a nonnegative solution of $(12_{+\varepsilon})$. Certainly, $\lim_{t \rightarrow \infty} (\bar{H}_i, \bar{V}_i) = (0, 0)$ since $(12_{+\varepsilon})$ has no positive solution. Then, by the comparison principle, $\lim_{t \rightarrow \infty} (H_i, V_i) = (0, 0)$. This together with $\lim_{t \rightarrow \infty} (V_u + V_i) = V_*$ gives (9).

If $\lambda(V_*) = 0$, then $\lambda(V_* + \varepsilon) < 0$ and $(12_{+\varepsilon})$ has a unique positive solution $(H_{i,+\varepsilon}^*, V_{i,+\varepsilon}^*)$ and $\lim_{\varepsilon \rightarrow 0} (H_{i,+\varepsilon}^*, V_{i,+\varepsilon}^*) = (0, 0)$ in $[C^2(\bar{\Omega})]^2$ since (4) has no positive solution in the present case. Similar to the above, $\lim_{t \rightarrow \infty} (H_i, V_i) = (0, 0)$ and (9) holds.

(iii) In such case, (3) has no positive solution and $\lim_{t \rightarrow \infty} (V_u + V_i) = 0$. Therefore, $\lim_{t \rightarrow \infty} H_i = 0$ by the first equation of (1). The proof is complete. \square

References

- [1] W.E. Fitzgibbon, J.J. Morgan and G.F. Webb, An outbreak vector-host epidemic model with spatial structure: the 2015-2016 Zika outbreak in Rio De Janeiro. *Theor. Biol. Med. Modell.* 14, 2-17 (2017)
- [2] P. Magal, G. Webb and Y. Wu, On a vector-host epidemic model with spatial structure. *Nonlinearity* 31, 5589-5614 (2018)
- [3] P. Magal, G. Webb and Y. Wu, On the basic reproduction number of reaction-diffusion epidemic models. *SIAM J. Appl. Math.* 79, 284-304 (2019)
- [4] G. Sweers, Strong positivity in $C(\bar{\Omega})$ for elliptic systems. *Math. Z.* 209, 251-271 (1992)
- [5] H. Amann, Fixed point equations and nonlinear eigenvalue problems in ordered Banach spaces. *SIAM Rev.* 18, 620-709 (1976)
- [6] M.X. Wang, *Nonlinear Second Order Parabolic Equations*. Boca Raton: CRC Press, 2021.