

Note on a vector-host epidemic model with spatial structure¹

Mingxin Wang²

School of Mathematics and Information Science, Henan Polytechnic University, Jiaozuo 454003, China

Abstract. Magal, Webb and Wu [Nonlinearity 31, 5589-5614 (2018)] studied the model describing outbreak of Zika in Rio De Janerio, and provided a complete analysis of dynamical properties for the solutions. In this note we first use a very simple approach to prove their results, and then investigate the modified version of the model concerned in their paper, with Neumann boundary condition replaced by Dirichlet boundary condition.

Keywords: Epidemic models; Equilibrium solutions; Global stabilities.

AMS Subject Classification (2020): 35B40, 35K57, 35Q92

1 Introduction

Vector-borne diseases such as chikungunya, dengue, malaria, West Nile virus, yellow fever and Zika virus pose a major threat to global public health [1, 2, 3, 4]. In recent years, many authors have proposed reaction-diffusion models to study the transmission of diseases in spatial settings. Among them, Fitzgibbon et al. [5] proposed and studied a spatially correlated bounded vector-host epidemic model that qualitatively described the 2015-2016 Zika epidemic in Rio de Janeiro. The global dynamics of the spatial model were further established based on the basic reproduction number. Suppose that individuals are living in a bounded smooth domain $\Omega \subset \mathbb{R}^n$, and ν is the outward normal vector of $\partial\Omega$. Let $H_i(x, t)$, $V_u(x, t)$ and $V_i(x, t)$ be the densities of infected hosts, uninfected vectors, and infected vectors at position x and time t , respectively. Then the model proposed in [5] to study the outbreak of Zika in Rio De Janerio is the following reaction-diffusion system

$$\begin{cases} \partial_t H_i - \nabla \cdot d_1(x) \nabla H_i = -\rho(x) H_i + \sigma_1(x) H_u(x) V_i, & x \in \Omega, t > 0, \\ \partial_t V_u - \nabla \cdot d_2(x) \nabla V_u = -\sigma_2(x) V_u H_i + \beta(x) (V_u + V_i) - \mu(x) (V_u + V_i) V_u, & x \in \Omega, t > 0, \\ \partial_t V_i - \nabla \cdot d_2(x) \nabla V_i = \sigma_2(x) V_u H_i - \mu(x) (V_u + V_i) V_i, & x \in \Omega, t > 0, \\ \partial_\nu H_i = \partial_\nu V_u = \partial_\nu V_i = 0, & x \in \partial\Omega, t > 0, \\ (H_i(\cdot, 0), V_u(\cdot, 0), V_i(\cdot, 0)) = (H_{i0}, V_{u0}, V_{i0}) \in C(\bar{\Omega}; \mathbb{R}_+^3), & \end{cases} \quad (1.1)$$

where $d_1, d_2 \in C^{1+\alpha}(\bar{\Omega})$ and $\rho, \beta, \sigma_1, \sigma_2, \mu \in C^\alpha(\bar{\Omega})$ are strictly positive, and $H_u \in C^\alpha(\bar{\Omega})$ is nonnegative and non-trivial. The flux of new infected humans is given by $\sigma_1(x) H_u(x) V_i(x, t)$ in which $H_u(x)$ is the density of susceptible population depending on the spatial location x . The initial functions satisfy $\partial_\nu H_{i0} = \partial_\nu V_{u0} = \partial_\nu V_{i0} = 0$ on $\partial\Omega$.

¹This work was supported by NSFC Grant 12171120.

²E-mail: mxwang@hpu.edu.cn

When establishing the model (1.1), the main assumption is that the susceptible human population is (almost) not affected by the epidemic during a relatively short period of time and therefore the flux of new infected is (almost) constant. Such a functional response mainly permits to take care of realistic density of population distributed in space.

The model (1.1) has been thoroughly studied by Magal, Webb and Wu [6, 7]. In [6], they found the basic reproduction number \mathcal{R}_0 and show that \mathcal{R}_0 is a threshold parameter: if $\mathcal{R}_0 \leq 1$ the disease free equilibrium is globally stable; if $\mathcal{R}_0 > 1$ the model has a unique globally stable positive equilibrium. In [7], it was shown that the basic reproduction number \mathcal{R}_0 can be defined as the spectral radius of a product of a local basic reproduction number \mathcal{R} and strongly positive compact linear operators with spectral radii one.

In this note we first use a very simple approach to prove the main results of [6], and then study the other version of (1.1) coupled with homogeneous Dirichlet boundary condition

$$\begin{cases} \partial_t H_i - \nabla \cdot d_1(x) \nabla H_i = -\rho(x)H_i + \sigma_1(x)H_u(x)V_i, & x \in \Omega, t > 0, \\ \partial_t V_u - \nabla \cdot d_2(x) \nabla V_u = -\sigma_2(x)V_u H_i + \beta(x)(V_u + V_i) - \mu(x)(V_u + V_i)V_u, & x \in \Omega, t > 0, \\ \partial_t V_i - \nabla \cdot d_2(x) \nabla V_i = \sigma_2(x)V_u H_i - \mu(x)(V_u + V_i)V_i, & x \in \Omega, t > 0, \\ H_i = V_u = V_i = 0, & x \in \partial\Omega, t > 0, \\ H_i(x, 0) = H_{i0}(x), V_u(x, 0) = V_{u0}(x), V_i(x, 0) = V_{i0}(x), \end{cases} \quad (1.2)$$

where the initial data $H_{i0}, V_{u0}, V_{i0} \in C(\bar{\Omega})$, and are positive in Ω and satisfy $H_{i0} = V_{u0} = V_{i0} = 0$ on $\partial\Omega$, all parameters are same as in (1.1). It is easy to see that the solutions (H_i, V_u, V_i) of (1.1) and (1.2) are unique, positive and bounded.

For the sake of convenience, we denote $\nabla \cdot d_1(x) \nabla = \mathcal{L}_1$ and $\nabla \cdot d_2(x) \nabla = \mathcal{L}_2$.

2 Preliminaries on the equilibrium solutions of (1.1) and (1.2)

Equilibrium problems of (1.1) and (1.2) are the following boundary value problems

$$\begin{cases} -\mathcal{L}_1 H_i^* = -\rho(x)H_i^* + \sigma_1(x)H_u(x)V_i^*, & x \in \Omega, \\ -\mathcal{L}_2 V_u^* = -\sigma_2(x)V_u^* H_i^* + \beta(x)(V_u^* + V_i^*) - \mu(x)(V_u^* + V_i^*)V_u^*, & x \in \Omega, \\ -\mathcal{L}_2 V_i^* = \sigma_2(x)V_u^* H_i^* - \mu(x)(V_u^* + V_i^*)V_i^*, & x \in \Omega, \\ \mathcal{B}[H_i^*] = \mathcal{B}[V_u^*] = \mathcal{B}[V_i^*] = 0, & x \in \partial\Omega \end{cases} \quad (2.1\mathcal{B})$$

with $\mathcal{B}[W] = \partial_\nu W$ and $\mathcal{B}[W] = W$, respectively. Let $\lambda^{\mathcal{B}}(\beta)$ be the principal eigenvalue of

$$\begin{cases} -\mathcal{L}_2 \varphi - \beta(x)\varphi = \lambda\varphi, & x \in \Omega, \\ \mathcal{B}[\varphi] = 0, & x \in \partial\Omega. \end{cases}$$

Assume that (H_i^*, V_u^*, V_i^*) is a nonnegative solution of (2.1 \mathcal{B}) and set $V_{\mathcal{B}} = V_u^* + V_i^*$. Then $V_{\mathcal{B}}$ satisfies

$$\begin{cases} -\mathcal{L}_2 V_{\mathcal{B}} = \beta(x)V_{\mathcal{B}} - \mu(x)V_{\mathcal{B}}^2, & x \in \Omega, \\ \mathcal{B}[V_{\mathcal{B}}] = 0, & x \in \partial\Omega. \end{cases} \quad (2.2_{\mathcal{B}})$$

If $\lambda^{\mathcal{B}}(\beta) \geq 0$, then (2.2 $_{\mathcal{B}}$) has no positive solution. Therefore, $V_{\mathcal{B}} = 0$, i.e., $V_u^* = V_i^* = 0$, and then $H_i^* = 0$ by the first equation of (2.1 $_{\mathcal{B}}$). So $(0, 0, 0)$ is the only nonnegative solution of (2.1 $_{\mathcal{B}}$). If $\lambda^{\mathcal{B}}(\beta) < 0$, then (2.2 $_{\mathcal{B}}$) has a unique positive solution $V_{\mathcal{B}}$ and $V_{\mathcal{B}}$ is globally asymptotically stable.

Assume $\lambda^{\mathcal{B}}(\beta) < 0$ and let $V_{\mathcal{B}}$ be the unique positive solution of (2.2 $_{\mathcal{B}}$). To investigate positive solutions of (2.1 $_{\mathcal{B}}$) is equivalent to study positive solutions (H_i^*, V_i^*) of

$$\begin{cases} -\mathcal{L}_1 H_i^* = -\rho(x)H_i^* + \sigma_1(x)H_u(x)V_i^*, & x \in \Omega, \\ -\mathcal{L}_2 V_i^* = \sigma_2(x)(V_{\mathcal{B}}(x) - V_i^*)H_i^* - \mu(x)V_{\mathcal{B}}(x)V_i^*, & x \in \Omega, \\ \mathcal{B}[H_i^*] = \mathcal{B}[V_i^*] = 0, & x \in \partial\Omega \end{cases} \quad (2.3_{\mathcal{B}})$$

satisfying $V_i^* < V_{\mathcal{B}}$. Clearly, $(0, 0)$ is the unique trivial nonnegative solution of (2.3 $_{\mathcal{B}}$). The linearized eigenvalue problem of (2.3 $_{\mathcal{B}}$) at $(0, 0)$ is

$$\begin{cases} -\mathcal{L}_1 \phi_1 + \rho(x)\phi_1 - \sigma_1(x)H_u(x)\phi_2 = \lambda\phi_1, & x \in \Omega, \\ -\mathcal{L}_2 \phi_2 - \sigma_2(x)V_{\mathcal{B}}(x)\phi_1 + \mu(x)V_{\mathcal{B}}(x)\phi_2 = \lambda\phi_2, & x \in \Omega, \\ \mathcal{B}[\phi_1] = \mathcal{B}[\phi_2] = 0, & x \in \partial\Omega. \end{cases} \quad (2.4_{\mathcal{B}})$$

By the results of [8], the problem (2.4 $_{\mathcal{B}}$) has a unique principal eigenvalue $\lambda^{\mathcal{B}}(V_{\mathcal{B}})$ with positive eigenfunction $\phi = (\phi_1, \phi_2)^T$. Moreover, when $\mathcal{B}[W] = \partial_{\nu}W$ (the Neumann boundary condition), $\lambda^{\mathcal{B}}(V_{\mathcal{B}})$ has the same sign with $\mathcal{R}_0 - 1$, where \mathcal{R}_0 is given in [6].

Theorem 2.1. *Assume $\lambda^{\mathcal{B}}(\beta) < 0$. If (2.3 $_{\mathcal{B}}$) has a positive solution (H_i^*, V_i^*) , then $\lambda^{\mathcal{B}}(V_{\mathcal{B}}) < 0$.*

Proof. Define an operator

$$\mathcal{L}_{\mathcal{B}}\psi = \begin{pmatrix} -\mathcal{L}_1\psi_1 + \rho(x)\psi_1, \\ -\mathcal{L}_2\psi_2 + \mu(x)V_{\mathcal{B}}(x)\psi_2 \end{pmatrix}, \quad \psi_i \in C^2(\bar{\Omega}), \mathcal{B}[\psi_i]_{\partial\Omega} = 0, i = 1, 2,$$

and a matrix

$$A_{\mathcal{B}}(x) = \begin{pmatrix} \lambda^{\mathcal{B}}(V_{\mathcal{B}}) & \sigma_1(x)H_u(x) \\ \sigma_2(x)V_{\mathcal{B}}(x) & \lambda^{\mathcal{B}}(V_{\mathcal{B}}) \end{pmatrix}.$$

Then $\mathcal{L}_{\mathcal{B}}$ is reversible and $\mathcal{L}_{\mathcal{B}}^{-1}$ is strongly positive and compact. On the contrary we assume $\lambda^{\mathcal{B}}(V_{\mathcal{B}}) \geq 0$. Then the operator $T_{\mathcal{B}} := \mathcal{L}_{\mathcal{B}}^{-1}A_{\mathcal{B}}(x)$ is also strongly positive and compact, and $r(T_{\mathcal{B}}) = 1$ as $\phi_1, \phi_2 > 0$ and $\phi = (\phi_1, \phi_2)^T$ satisfies $\phi = \mathcal{L}_{\mathcal{B}}^{-1}A_{\mathcal{B}}(x)[\phi]$ by (2.4 $_{\mathcal{B}}$).

On the other hand, since (H_i^*, V_i^*) is a positive solution of (2.3 $_{\mathcal{B}}$), we have

$$r(T_{\mathcal{B}}) \begin{pmatrix} H_i^* \\ V_i^* \end{pmatrix} = \begin{pmatrix} H_i^* \\ V_i^* \end{pmatrix} = T_{\mathcal{B}} \begin{pmatrix} H_i^* \\ V_i^* \end{pmatrix} - \mathcal{L}_{\mathcal{B}}^{-1} \begin{pmatrix} \lambda^{\mathcal{B}}(V_{\mathcal{B}}) & 0 \\ \sigma_2(x)V_i^* & \lambda^{\mathcal{B}}(V_{\mathcal{B}}) \end{pmatrix} \begin{pmatrix} H_i^* \\ V_i^* \end{pmatrix},$$

and

$$\mathcal{L}_{\mathcal{B}}^{-1} \begin{pmatrix} \lambda^{\mathcal{B}}(V_{\mathcal{B}}) & 0 \\ \sigma_2(x)V_i^* & \lambda^{\mathcal{B}}(V_{\mathcal{B}}) \end{pmatrix} \begin{pmatrix} H_i^* \\ V_i^* \end{pmatrix} \geq \mathcal{L}_{\mathcal{B}}^{-1} \begin{pmatrix} 0 \\ \sigma_2(x)V_i^*H_i^* \end{pmatrix} =: \begin{pmatrix} 0 \\ \chi(x) \end{pmatrix},$$

where $\chi(x) > 0$. This is impossible by the conclusion [9, Theorem 3.2 (iv)]. \square

3 Dynamical properties of (1.1)—main results in [6] and proofs

In this section we focus on the model (1.1) and write the boundary operator \mathcal{B} as \mathcal{N} to represent the Neumann boundary condition and $V_{\mathcal{B}} = V_{\mathcal{N}}$. Clearly, $\lambda^{\mathcal{N}}(\beta) < 0$ since $\beta(x) > 0$.

Theorem 3.1. *Then the problem (2.3 $_{\mathcal{N}}$) has a positive solution (H_i^*, V_i^*) if and only if $\lambda^{\mathcal{N}}(V_{\mathcal{N}}) < 0$. Moreover, (H_i^*, V_i^*) is unique and satisfies $V_i^* < V_{\mathcal{N}}$ when it exists. Therefore, (2.1 $_{\mathcal{N}}$) has a positive solution (H_i^*, V_u^*, V_i^*) if and only if $\lambda^{\mathcal{N}}(V_{\mathcal{N}}) < 0$, and (H_i^*, V_u^*, V_i^*) is unique and takes the form $(H_i^*, V_{\mathcal{N}} - V_i^*, V_i^*)$ when it exists.*

Proof. By Theorem 2.1, it suffices to prove that if $\lambda^{\mathcal{N}}(V_{\mathcal{N}}) < 0$, then (2.3 $_{\mathcal{N}}$) has a unique positive solution (H_i^*, V_i^*) and $V_i^* < V_{\mathcal{N}}$. To this aim, we prove the following general conclusion.

- There is $0 < \varepsilon_0 \ll 1$ such that, when $|\varepsilon| \leq \varepsilon_0$, the problem

$$\begin{cases} -\mathcal{L}_1 H_i = -\rho(x)H_i + \sigma_1(x)H_u(x)V_i, & x \in \Omega, \\ -\mathcal{L}_2 V_i = \sigma_2(x)(V_{\mathcal{N}}(x) + \varepsilon - V_i)^+ H_i - \mu(x)(V_{\mathcal{N}}(x) - \varepsilon)V_i, & x \in \Omega, \\ \partial_\nu H_i = \partial_\nu V_i = 0, & x \in \partial\Omega \end{cases} \quad (3.1_\varepsilon)$$

has a unique positive solution $(H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$, and $V_{i,\varepsilon}^* < V_{\mathcal{N}} + \varepsilon$.

Existence of positive solution of (3.1 $_\varepsilon$). Let $\lambda^{\mathcal{N}}(V_{\mathcal{N}}; \varepsilon)$ be the principal eigenvalue of

$$\begin{cases} -\mathcal{L}_1 \phi_1 + \rho(x)\phi_1 - \sigma_1(x)H_u(x)\phi_2 = \lambda\phi_1, & x \in \Omega, \\ -\mathcal{L}_2 \phi_2 - \sigma_2(x)(V_{\mathcal{N}}(x) + \varepsilon)\phi_1 + \mu(x)(V_{\mathcal{N}}(x) - \varepsilon)\phi_2 = \lambda\phi_2, & x \in \Omega, \\ \partial_\nu \phi_1 = \partial_\nu \phi_2 = 0, & x \in \partial\Omega, \end{cases} \quad (3.2_\varepsilon)$$

(the existence and uniqueness of $\lambda^{\mathcal{N}}(V_{\mathcal{N}}; \varepsilon)$ is given in [8]). As $\lambda^{\mathcal{N}}(V_{\mathcal{N}}) < 0$, there is $0 < \varepsilon_0 \ll 1$ such that, when $|\varepsilon| \leq \varepsilon_0$, we have $\lambda^{\mathcal{N}}(V_{\mathcal{N}}; \varepsilon) < 0$, and $V_{\mathcal{N}}(x) \pm \varepsilon > 0$, $\varepsilon^2 \mu(x) < \beta(x)V_{\mathcal{N}}(x)$ in $\bar{\Omega}$. Let \bar{H}_i^* be the unique positive solution of the linear problem

$$\begin{cases} -\mathcal{L}_1 \bar{H}_i^* + \rho(x)\bar{H}_i^* = \sigma_1(x)H_u(x)(V_{\mathcal{N}}(x) + \varepsilon), & x \in \Omega, \\ \partial_\nu \bar{H}_i^* = 0, & x \in \partial\Omega. \end{cases}$$

Then $(\bar{H}_i^*, V_{\mathcal{N}} + \varepsilon)$ is a strict upper solution of (3.1 $_\varepsilon$). Let (ϕ_1, ϕ_2) be the positive eigenfunction corresponding to $\lambda^{\mathcal{N}}(V_{\mathcal{N}}; \varepsilon)$. It is easy to verify that $\delta(\phi_1, \phi_2)$ is a lower solution of (3.1 $_\varepsilon$) and $\delta(\phi_1, \phi_2) \leq (\bar{H}_i^*, V_{\mathcal{N}} + \varepsilon)$ provided $\delta > 0$ is suitably small. By the upper and lower solutions method, (3.1 $_\varepsilon$) has at least one positive solution $(H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$, and $V_{i,\varepsilon}^* < V_{\mathcal{N}} + \varepsilon$, $H_{i,\varepsilon}^* < \bar{H}_i^*$.

Uniqueness of positive solutions of (3.1 $_\varepsilon$). Let $(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*)$ be another positive solution of (3.1 $_\varepsilon$). We can find $0 < s < 1$ such that $s(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \leq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ in $\bar{\Omega}$. Set

$$\bar{s} = \sup\{0 < s \leq 1 : s(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \leq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*) \text{ in } \bar{\Omega}\}.$$

Then $0 < \bar{s} \leq 1$ and $\bar{s}(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \leq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ in $\bar{\Omega}$. We shall prove $\bar{s} = 1$. If $\bar{s} < 1$, then $W := H_{i,\varepsilon}^* - \bar{s}\hat{H}_{i,\varepsilon}^* \geq 0$, $Z := V_{i,\varepsilon}^* - \bar{s}\hat{V}_{i,\varepsilon}^* \geq 0$. Since $\bar{s}\hat{V}_{i,\varepsilon}^* \leq V_{i,\varepsilon}^* < V_{\mathcal{N}} + \varepsilon$ and $\hat{V}_{i,\varepsilon}^* > \bar{s}\hat{V}_{i,\varepsilon}^*$, we have

$$(V_{\mathcal{N}} + \varepsilon - V_{i,\varepsilon}^*)^+ = V_{\mathcal{N}} + \varepsilon - V_{i,\varepsilon}^*, \quad (V_{\mathcal{N}} + \varepsilon - \hat{V}_{i,\varepsilon}^*)^+ < V_{\mathcal{N}} + \varepsilon - \bar{s}\hat{V}_{i,\varepsilon}^*,$$

$$V_{\mathcal{N}}(x) + \varepsilon - V_{i,\varepsilon}^* - (V_{\mathcal{N}}(x) + \varepsilon - \hat{V}_{i,\varepsilon}^*)^+ > \bar{s}\hat{V}_{i,\varepsilon}^* - V_{i,\varepsilon}^* = -Z.$$

After careful calculation, it derives that

$$\begin{cases} -\mathcal{L}_1 W = -\rho(x)W + \sigma_1(x)H_u(x)Z, \\ -\mathcal{L}_2 Z > -[\mu(x)(V_{\mathcal{N}}(x) - \varepsilon) + \sigma_2(x)\hat{H}_{i,\varepsilon}^*]Z + \sigma_2(x)(V_{\mathcal{N}}(x) + \varepsilon - V_{i,\varepsilon}^*)W. \end{cases}$$

Notice $V_{\mathcal{N}}(x) + \varepsilon - V_{i,\varepsilon}^* > 0$ and $W, Z \geq 0$. It follows that $W, Z > 0$ in $\bar{\Omega}$ by the maximum principle.

Then there exists $0 < \tau < 1 - \bar{s}$ such that $(W, Z) \geq \tau(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*)$, i.e., $(\bar{s} + \tau)(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \leq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ in $\bar{\Omega}$. This contradicts the definition of \bar{s} . Hence $\bar{s} = 1$, i.e., $(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \leq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ in $\bar{\Omega}$. Certainly, $\hat{V}_{i,\varepsilon}^* < V_{\mathcal{N}} + \varepsilon$, and $(V_{\mathcal{N}}(x) + \varepsilon - \hat{V}_{i,\varepsilon}^*)^+ = V_{\mathcal{N}}(x) + \varepsilon - \hat{V}_{i,\varepsilon}^*$.

On the other hand, we can find $\zeta > 1$ such that $\zeta(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \geq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ in Ω . Set

$$\underline{\zeta} = \inf\{\zeta \geq 1 : \zeta(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \geq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*) \text{ in } \bar{\Omega}\}.$$

Then $\underline{\zeta}$ is well defined, $\underline{\zeta} \geq 1$ and $\underline{\zeta}(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \geq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ in Ω . If $\underline{\zeta} > 1$, then $P := \underline{\zeta}\hat{H}_{i,\varepsilon}^* - H_{i,\varepsilon}^* \geq 0$ and $Q := \underline{\zeta}\hat{V}_{i,\varepsilon}^* - V_{i,\varepsilon}^* \geq 0$, and (P, Q) satisfies

$$\begin{cases} -\mathcal{L}_1 P = -\rho(x)P + \sigma_1(x)H_u(x)Q, \\ -\mathcal{L}_2 Q > -[\mu(x)(V_{\mathcal{N}}(x) - \varepsilon) + \sigma_2(x)H_{i,\varepsilon}^*]Q + \sigma_2(x)(V_{\mathcal{N}}(x) + \varepsilon - \hat{V}_{i,\varepsilon}^*)P. \end{cases}$$

As $V_{\mathcal{N}}(x) + \varepsilon - \hat{V}_{i,\varepsilon}^* > 0$ and $P, Q \geq 0$. Then $P, Q > 0$ in $\bar{\Omega}$, and there exists $0 < r < \underline{\zeta} - 1$ such that $(P, Q) \geq r(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*)$, i.e., $(\underline{\zeta} - r)(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \geq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ in $\bar{\Omega}$. This contradicts the definition of $\underline{\zeta}$. Hence $\underline{\zeta} = 1$ and $(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \geq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$. The uniqueness is proved.

Taking $\varepsilon = 0$ in the above, we obtain the desired conclusion. \square

Theorem 3.2. *Let (H_i, V_u, V_i) be the unique positive solution of (1.1). Then the following hold.*

(i) *If $\lambda^{\mathcal{N}}(V_{\mathcal{N}}) < 0$, then*

$$\lim_{t \rightarrow \infty} (H_i, V_u, V_i) = (H_i^*, V_{\mathcal{N}} - V_i^*, V_i^*) = (H_i^*, V_u^*, V_i^*) \text{ in } [C^2(\bar{\Omega})]^3. \quad (3.3)$$

(ii) *If $\lambda^{\mathcal{N}}(V_{\mathcal{N}}) \geq 0$, then*

$$\lim_{t \rightarrow \infty} (H_i, V_u, V_i) = (0, V_{\mathcal{N}}, 0) \text{ in } [C^2(\bar{\Omega})]^3. \quad (3.4)$$

Proof. (i) Let (H_i, V_u, V_i) be the unique solution of (1.1), and set $V = V_u + V_i$. Then we have

$$\begin{cases} \partial_t V - \mathcal{L}_2 V = \beta(x)V - \mu(x)V^2, & x \in \Omega, t > 0, \\ \partial_\nu V = 0, & x \in \partial\Omega, t > 0, \\ V(x, 0) = V_u(x, 0) + V_i(x, 0) > 0, & x \in \bar{\Omega}, \end{cases}$$

and $\lim_{t \rightarrow \infty} V(x, t) = V_{\mathcal{N}}(x)$ in $C^2(\bar{\Omega})$. For any given $0 < \varepsilon \leq \varepsilon_0$, there exists $T_\varepsilon \gg 1$ such that

$$0 < V_{\mathcal{N}}(x) - \varepsilon \leq V(x, t) \leq V_{\mathcal{N}}(x) + \varepsilon, \quad \forall x \in \bar{\Omega}, t \geq T_\varepsilon.$$

Hence, (H_i, V_i) satisfies

$$\begin{cases} \partial_t H_i - \mathcal{L}_1 H_i = -\rho(x)H_i + \sigma_1(x)H_u(x)V_i, & x \in \Omega, t > T_\varepsilon, \\ \partial_t V_i - \mathcal{L}_2 V_i \leq \sigma_2(x)(V_{\mathcal{N}}(x) + \varepsilon - V_i)^+ H_i - \mu(x)(V_{\mathcal{N}}(x) - \varepsilon)V_i, & x \in \Omega, t > T_\varepsilon, \\ \partial_\nu H_i = \partial_\nu V_i = 0, & x \in \partial\Omega, t > T_\varepsilon. \end{cases}$$

Let $(H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ be the unique positive solution of (3.1 $_\varepsilon$), and (ϕ_1, ϕ_2) be the positive eigenfunction corresponding to $\lambda^{\mathcal{N}}(V_{\mathcal{N}}; \varepsilon)$. We can take constants $k \gg 1$ and $0 < \delta \ll 1$ such that

$$k(H_{i,\varepsilon}^*(x), V_{i,\varepsilon}^*(x)) \geq (H_i(x, T_\varepsilon), V_i(x, T_\varepsilon)), \quad \delta(\phi_1(x), \phi_2(x)) \leq (H_i(x, T_\varepsilon), V_i(x, T_\varepsilon)), \quad x \in \bar{\Omega}.$$

Then $k(H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ and $\delta(\phi_1, \phi_2)$ are the upper and lower solutions of (3.1 $_\varepsilon$), respectively. Let $(H_{i,\varepsilon}, V_{i,\varepsilon})$ be the unique positive solution of

$$\begin{cases} \partial_t H_i - \mathcal{L}_1 H_i = -\rho(x)H_i + \sigma_1(x)H_u(x)V_i, & x \in \Omega, t > 0, \\ \partial_t V_i - \mathcal{L}_2 V_i = \sigma_2(x)(V_{\mathcal{N}}(x) + \varepsilon - V_i)^+ H_i - \mu(x)(V_{\mathcal{N}}(x) - \varepsilon)V_i, & x \in \Omega, t > 0, \\ \partial_\nu H_i = \partial_\nu V_i = 0, & x \in \partial\Omega, t > 0, \\ (H_i(x, 0), V_i(x, 0)) = k(H_{i,\varepsilon}^*, V_{i,\varepsilon}^*), & x \in \bar{\Omega}. \end{cases} \quad (3.5)$$

Then $(H_{i,\varepsilon}(x, t), V_{i,\varepsilon}(x, t))$ is decreasing in t , and

$$\delta(\phi_1(x), \phi_2(x)) \leq (H_{i,\varepsilon}(x, t), V_{i,\varepsilon}(x, t)), \quad (H_i(x, t + T_\varepsilon), V_i(x, t + T_\varepsilon)) \leq (H_{i,\varepsilon}(x, t), V_{i,\varepsilon}(x, t))$$

by the comparison principle. Moreover, $\lim_{t \rightarrow \infty} (H_{i,\varepsilon}(x, t), V_{i,\varepsilon}(x, t)) = (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ in $[C^2(\bar{\Omega})]^2$ by the uniform estimate and compact arguments (cf. [10, Theorems 2.11, 3.14]) and the uniqueness of positive solutions of (3.1 $_\varepsilon$). So $\limsup_{t \rightarrow \infty} (H_i(x, t), V_i(x, t)) \leq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ and, by letting $\varepsilon \rightarrow 0$,

$$\limsup_{t \rightarrow \infty} (H_i(x, t), V_i(x, t)) \leq (H_i^*, V_i^*) \quad \text{uniformly in } \bar{\Omega}. \quad (3.6)$$

On the other hand, as $V_i^*(x) < V_{\mathcal{N}}(x)$ in $\bar{\Omega}$, there exists $0 < r_0 \leq \varepsilon_0$ such that $V_i^*(x) < V_{\mathcal{N}}(x) - 2r$ in $\bar{\Omega}$ for all $0 < r < r_0$. For such a r , using (3.6) and $\lim_{t \rightarrow \infty} V(x, t) = V_{\mathcal{N}}(x)$ in $C^2(\bar{\Omega})$, we can find $T_r \gg 1$ such that

$$V_i(x, t) \leq V_i^*(x) + r < V_{\mathcal{N}}(x) - r, \quad 0 < V_{\mathcal{N}}(x) - r \leq V(x, t) \leq V_{\mathcal{N}}(x) + r, \quad \forall x \in \bar{\Omega}, t \geq T_r.$$

Consequently, (H_i, V_i) satisfies

$$\begin{cases} \partial_t H_i - \mathcal{L}_1 H_i = -\rho(x)H_i + \sigma_1(x)H_u(x)V_i, & x \in \Omega, t > T_r, \\ \partial_t V_i - \mathcal{L}_2 V_i \geq \sigma_2(x)(V_{\mathcal{N}}(x) - r - V_i)H_i - \mu(x)(V_{\mathcal{N}}(x) + r)V_i \\ \quad = \sigma_2(x)(V_{\mathcal{N}}(x) - r - V_i)^+ H_i - \mu(x)(V_{\mathcal{N}}(x) + r)V_i, & x \in \Omega, t > T_r, \\ \partial_\nu H_i = \partial_\nu V_i = 0, & x \in \partial\Omega, t > T_r, \\ H_i(x, T_r) > 0, \quad 0 < V_i(x, T_r) < V_{\mathcal{N}}(x) - r, & x \in \bar{\Omega}. \end{cases}$$

Let (ϕ_1, ϕ_2) be the positive eigenfunction corresponding to $\lambda^{\mathcal{N}}(V_{\mathcal{N}}; -r)$. Similar to the above, $\delta(\phi_1, \phi_2)$ is a lower solution of (3.1 $_{-r}$) and $\delta(\phi_1(x), \phi_2(x)) \leq (H_i(x, T_r), V_i(x, T_r))$ in $\bar{\Omega}$ provided $0 < \delta \ll 1$. Let $(H_{i,-r}, V_{i,-r})$ be the unique positive solution of (3.5) with $\varepsilon = -r$ and $(H_i(x, 0), V_i(x, 0)) = \delta(\phi_1(x), \phi_2(x))$. Then

$$(H_i(x, t + T_r), V_i(x, t + T_r)) \geq (H_{i,-r}(x, t), V_{i,-r}(x, t)), \quad x \in \Omega, \quad t > 0,$$

and $(H_{i,-r}, V_{i,-r})$ is increasing in t by the comparison principle. Similar to the above,

$$\lim_{t \rightarrow \infty} (H_{i,-r}(x, t), V_{i,-r}(x, t)) = (H_{i,-r}^*, V_{i,-r}^*) \quad \text{in } [C^2(\bar{\Omega})]^2,$$

where $(H_{i,-r}^*, V_{i,-r}^*)$ is the unique positive solution of (3.1 $_{-r}$). Thus, $\liminf_{t \rightarrow \infty} (H_i, V_i) \geq (H_{i,-r}^*, V_{i,-r}^*)$, and then $\liminf_{t \rightarrow \infty} (H_i, V_i) \geq (H_i^*, V_i^*)$ by letting $r \rightarrow 0$. This combined with (3.6) yields $\lim_{t \rightarrow \infty} (H_i, V_i) = (H_i^*, V_i^*)$ uniformly in $\bar{\Omega}$. Using the fact that $\lim_{t \rightarrow \infty} (V_u + V_i) = V_{\mathcal{N}}$ in $C^2(\bar{\Omega})$ and the uniform estimate (cf. [10, Theorems 2.11, 3.14]), we see that (3.3) holds.

(ii) Let $\lambda^{\mathcal{N}}(V_{\mathcal{N}}; \varepsilon)$ be the principal eigenvalue of (3.2 $_{\varepsilon}$). If $\lambda^{\mathcal{N}}(V_{\mathcal{N}}) > 0$, then $\lambda^{\mathcal{N}}(V_{\mathcal{N}}; \varepsilon) > 0$ when $0 < \varepsilon \ll 1$. So (3.1 $_{\varepsilon}$) has no positive solution. Let $(H_{i,\varepsilon}, V_{i,\varepsilon})$ be the unique positive solution of (3.5) with initial data (C, K) , where C and K are suitably large positive constants, for example,

$$K > \max_{\Omega} \max\{V_{\mathcal{N}}(x) + \varepsilon, V_i(x, T_{\varepsilon})\}, \quad C > \max_{\Omega} \max\{K\sigma_1(x)H_u(x)/\rho(x), H_i(x, T_{\varepsilon})\}.$$

Then (C, K) is an upper solution of (3.1 $_{\varepsilon}$), and $(H_{i,\varepsilon}, V_{i,\varepsilon})$ is decreasing in t . Similar to the above, $\lim_{t \rightarrow \infty} (H_{i,\varepsilon}, V_{i,\varepsilon}) = (0, 0)$ since (3.1 $_{\varepsilon}$) has no positive solution. Then, by the comparison principle, $\lim_{t \rightarrow \infty} (H_i, V_i) = (0, 0)$. This together with $\lim_{t \rightarrow \infty} (V_u + V_i) = V_{\mathcal{N}}$ gives the limit (3.4).

If $\lambda^{\mathcal{N}}(V_{\mathcal{N}}) = 0$, then $\lambda^{\mathcal{N}}(V_{\mathcal{N}}; \varepsilon) < 0$ and (3.1 $_{\varepsilon}$) has a unique positive solution $(H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ for any $\varepsilon > 0$. Obviously, $\lim_{\varepsilon \rightarrow 0} (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*) = (0, 0)$ in $[C^2(\bar{\Omega})]^2$ since (3.1 $_0$), i.e., (2.3 $_{\mathcal{N}}$) has no positive solution in the present case. Similar to the above, $\lim_{t \rightarrow \infty} (H_i, V_i) = (0, 0)$ and (3.4) holds. The proof is complete. \square

4 Dynamical properties of (1.2)

In the present situation we write the boundary operator \mathcal{B} as \mathcal{D} to represent the Dirichlet boundary condition and $V_{\mathcal{B}} = V_{\mathcal{D}}$. The results and arguments are similar to that in Section 3.

Theorem 4.1. *Assume $\lambda^{\mathcal{D}}(\beta) < 0$. Then the problem (2.3 $_{\mathcal{D}}$) has a positive solution (H_i^*, V_i^*) if and only if $\lambda^{\mathcal{D}}(V_{\mathcal{D}}) < 0$. Moreover, (H_i^*, V_i^*) is unique and satisfies $V_i^* < V_{\mathcal{D}}$ when it exists. Therefore, (2.1 $_{\mathcal{D}}$) has a positive solution (H_i^*, V_u^*, V_i^*) if and only if $\lambda^{\mathcal{D}}(V_{\mathcal{D}}) < 0$, and (H_i^*, V_u^*, V_i^*) is unique and takes the form $(H_i^*, V_{\mathcal{D}} - V_i^*, V_i^*)$ when it exists.*

Proof. By Theorem 2.1, it suffices to prove that if $\lambda^{\mathcal{D}}(V_{\mathcal{D}}) < 0$, then (2.3 $_{\mathcal{D}}$) has a unique positive solution (H_i^*, V_i^*) and $V_i^* < V_{\mathcal{D}}$. To this aim, we prove the following general conclusion.

- Let $\varphi(x)$, with $\max_{\Omega} \|\varphi\| = 1$, be the positive eigenfunction corresponding to $\lambda^{\mathcal{D}}(\beta)$. Then there is $0 < \varepsilon_0 \ll 1$ such that, when $|\varepsilon| \leq \varepsilon_0$, the problem

$$\begin{cases} -\mathcal{L}_1 H_i^* = -\rho(x)H_i^* + \sigma_1(x)H_u(x)V_i^*, & x \in \Omega, \\ -\mathcal{L}_2 V_i^* = \sigma_2(x)(V_{\mathcal{D}}(x) + \varepsilon\varphi(x) - V_i^*)^+ H_i^* - \mu(x)(V_{\mathcal{D}}(x) - \varepsilon\varphi(x))V_i^*, & x \in \Omega, \\ H_i^* = V_i^* = 0, & x \in \partial\Omega \end{cases} \quad (4.1_{\varepsilon})$$

has a unique positive solution $(H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$, and $V_{i,\varepsilon}^*(x) < V_{\mathcal{D}}(x) + \varepsilon\varphi(x)$.

Existence of positive solution of (4.1 $_{\varepsilon}$). Let $\lambda^{\mathcal{D}}(V_{\mathcal{D}}; \varepsilon)$ be the principal eigenvalue of

$$\begin{cases} -\mathcal{L}_1 \phi_1 + \rho(x)\phi_1 - \sigma_1(x)H_u(x)\phi_2 = \lambda\phi_1, & x \in \Omega, \\ -\mathcal{L}_2 \phi_2 - \sigma_2(x)(V_{\mathcal{D}}(x) + \varepsilon\varphi(x))\phi_1 + \mu(x)(V_{\mathcal{D}}(x) - \varepsilon\varphi(x))\phi_2 = \lambda\phi_2, & x \in \Omega, \\ \phi_1 = \phi_2 = 0, & x \in \partial\Omega. \end{cases}$$

As $\lambda^{\mathcal{D}}(V_{\mathcal{D}}) < 0$, there is $0 < \varepsilon_0 \ll 1$ such that, when $|\varepsilon| \leq \varepsilon_0$, we have $\lambda^{\mathcal{D}}(V_{\mathcal{D}}; \varepsilon) < 0$, and (cf. [11, Lemma 2.1])

$$V_{\mathcal{D}}(x) \pm \varepsilon\varphi(x) > 0, \quad (\varepsilon\varphi(x))^2 \mu(x) < \beta(x)V_{\mathcal{D}}(x) + \varepsilon\varphi(x)(\lambda^{\mathcal{D}}(V_{\mathcal{D}}) + \beta(x)) \quad \text{in } \Omega.$$

Let \bar{H}_i^* be the unique positive solution of the linear problem

$$\begin{cases} -\mathcal{L}_1 \bar{H}_i^* + \rho(x)\bar{H}_i^* = \sigma_1(x)H_u(x)(V_{\mathcal{D}}(x) + \varepsilon\varphi(x)), & x \in \Omega, \\ \bar{H}_i^* = 0, & x \in \partial\Omega. \end{cases}$$

Then $(\bar{H}_i^*, V_{\mathcal{D}} + \varepsilon\varphi(x))$ is a strict upper solution of (4.1 $_{\varepsilon}$). Let (ϕ_1, ϕ_2) be the positive eigenfunction corresponding to $\lambda^{\mathcal{D}}(V_{\mathcal{D}}; \varepsilon)$. It is easy to verify that $\delta(\phi_1, \phi_2)$ is a lower solution of (4.1 $_{\varepsilon}$) and $\delta(\phi_1, \phi_2) \leq (\bar{H}_i^*, V_{\mathcal{D}} + \varepsilon\varphi(x))$ provided $\delta > 0$ is suitably small. By the upper and lower solutions method, (4.1 $_{\varepsilon}$) has at least one positive solution $(H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$, and $V_{i,\varepsilon}^* < V_{\mathcal{D}} + \varepsilon\varphi(x)$, $H_{i,\varepsilon}^* < \bar{H}_i^*$.

Uniqueness of positive solutions of (4.1 $_{\varepsilon}$). Let $(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*)$ be another positive solution of (4.1 $_{\varepsilon}$). We can find $0 < s < 1$ such that $s(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \leq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ in Ω . Set

$$\bar{s} = \sup\{0 < s \leq 1 : s(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \leq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*) \text{ in } \Omega\}.$$

Then \bar{s} is well defined, $0 < \bar{s} \leq 1$ and $\bar{s}(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \leq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ in Ω . We shall prove $\bar{s} = 1$. If $\bar{s} < 1$, then $W := H_{i,\varepsilon}^* - \bar{s}\hat{H}_{i,\varepsilon}^* \geq 0$, $Z := V_{i,\varepsilon}^* - \bar{s}\hat{V}_{i,\varepsilon}^* \geq 0$. Since $\bar{s}\hat{V}_{i,\varepsilon}^* \leq V_{i,\varepsilon}^* < V_{\mathcal{D}} + \varepsilon\varphi(x)$ and $\bar{s} < 1$, we have

$$\begin{aligned} (V_{\mathcal{D}} + \varepsilon\varphi(x) - V_{i,\varepsilon}^*)^+ &= V_{\mathcal{D}} + \varepsilon\varphi(x) - V_{i,\varepsilon}^*, \quad (V_{\mathcal{D}} + \varepsilon\varphi(x) - \hat{V}_{i,\varepsilon}^*)^+ < V_{\mathcal{D}} + \varepsilon\varphi(x) - \bar{s}\hat{V}_{i,\varepsilon}^*, \\ V_{\mathcal{D}}(x) + \varepsilon\varphi(x) - V_{i,\varepsilon}^* - (V_{\mathcal{D}}(x) + \varepsilon\varphi(x) - \hat{V}_{i,\varepsilon}^*)^+ &> \bar{s}\hat{V}_{i,\varepsilon}^* - V_{i,\varepsilon}^* = -Z. \end{aligned}$$

By the careful calculations,

$$\begin{cases} -\mathcal{L}_1 W = -\rho(x)W + \sigma_1(x)H_u(x)Z, \\ -\mathcal{L}_2 Z > -[\mu(x)(V_{\mathcal{D}}(x) - \varepsilon\varphi(x)) + \sigma_2(x)\hat{H}_{i,\varepsilon}^*]Z + \sigma_2(x)(V_{\mathcal{D}}(x) + \varepsilon\varphi(x) - V_{i,\varepsilon}^*)W. \end{cases}$$

Notice $V_{\mathcal{D}}(x) + \varepsilon\varphi(x) - V_{i,\varepsilon}^* > 0$ and $W, Z \geq 0$. It follows that $W, Z > 0$ in Ω by the maximum principle. Then there exists $0 < r < 1 - \bar{s}$ such that $(W, Z) \geq r(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*)$, i.e., $(\bar{s} + r)(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \leq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ in Ω . This contradicts the definition of \bar{s} . Hence $\bar{s} = 1$, i.e., $(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \leq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ in Ω . Certainly, $\hat{V}_{i,\varepsilon}^* < V_{\mathcal{D}} + \varepsilon\varphi(x)$, and $(V_{\mathcal{D}}(x) + \varepsilon\varphi(x) - \hat{V}_{i,\varepsilon}^*)^+ = V_{\mathcal{D}}(x) + \varepsilon\varphi(x) - \hat{V}_{i,\varepsilon}^*$.

Same as the proof of Theorem 3.1 we can prove $(\hat{H}_{i,\varepsilon}^*, \hat{V}_{i,\varepsilon}^*) \geq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$.

Take $\varepsilon = 0$ in the above, we obtain the desired conclusion. \square

Theorem 4.2. *Let (H_i, V_u, V_i) be the unique positive solution of (1.2). Then the following holds.*

(i) *If $\lambda^{\mathcal{D}}(\beta) < 0$ and $\lambda^{\mathcal{D}}(V_{\mathcal{D}}) < 0$, then $\lim_{t \rightarrow \infty} (H_i, V_u, V_i) = (H_i^*, V_{\mathcal{D}} - V_i^*, V_i^*) = (H_i^*, V_u^*, V_i^*)$ in $[C^2(\bar{\Omega})]^3$.*

(ii) *If $\lambda^{\mathcal{D}}(\beta) < 0$ and $\lambda^{\mathcal{D}}(V_{\mathcal{D}}) \geq 0$, then $\lim_{t \rightarrow \infty} (H_i, V_u, V_i) = (0, V_{\mathcal{D}}, 0)$ in $[C^2(\bar{\Omega})]^3$.*

(iii) *If $\lambda^{\mathcal{D}}(\beta) \geq 0$, then $\lim_{t \rightarrow \infty} (H_i, V_u, V_i) = (0, 0, 0)$ in $[C^2(\bar{\Omega})]^3$.*

Proof. (i) Let (H_i, V_u, V_i) be the unique solution of (1.2), and set $V = V_u + V_i$. Then we have

$$\begin{cases} \partial_t V - \mathcal{L}_2 V = \beta(x)V - \mu(x)V^2, & x \in \Omega, t > 0, \\ V = 0, & x \in \partial\Omega, t > 0, \\ V(x, 0) = V_u(x, 0) + V_i(x, 0) > 0, & x \in \bar{\Omega}, \end{cases}$$

and $\lim_{t \rightarrow \infty} V(x, t) = V_{\mathcal{D}}(x)$ in $C^2(\bar{\Omega})$. Let $\varphi(x)$, with $\max_{\bar{\Omega}} \|\varphi\| = 1$, be the positive eigenfunction corresponding to $\lambda^{\mathcal{D}}(\beta)$. We first prove that for any given $0 < \varepsilon \leq \varepsilon_0$, there exists $T_\varepsilon \gg 1$ such that

$$0 < V_{\mathcal{D}}(x) - \varepsilon\varphi(x) < V(x, t) < V_{\mathcal{D}}(x) + \varepsilon\varphi(x), \quad \forall x \in \Omega, t \geq T_\varepsilon. \quad (4.2)$$

The idea of the proof of (4.2) comes from that of [11, Lemma 2.1]. In fact, set $W(x, t) = V(x, t) - V_{\mathcal{D}}(x)$ and $Z(x) = \varepsilon\varphi(x)$. Then $W(x, t) \rightarrow 0$ in $C^1(\bar{\Omega})$ as $t \rightarrow \infty$. Thanks to $Z \in C^1(\bar{\Omega})$ and $\partial_\nu Z|_{\partial\Omega} < 0$, there exist $\tau > 0$ and $T_1 \gg 1$ such that $\partial_\nu(W(x, t) - Z(x))|_{\partial\Omega} \geq \tau$ for all $t \geq T_1$. This together with $(W(x, t) - Z(x))|_{\partial\Omega} = 0$ asserts that there exists $\Omega_0 \Subset \Omega$ such that $W(x, t) - Z(x) < 0$ in $\Omega \setminus \Omega_0$. As $Z > 0$ in $\bar{\Omega}_0$ and $W(x, t) \rightarrow 0$ in $C(\bar{\Omega}_0)$ as $t \rightarrow \infty$, we can find $T_2 \gg 1$ such that $W(x, t) - Z(x) < 0$ in $\bar{\Omega}_0$. Thus, $W(x, t) - Z(x) < 0$, i.e., $V(x, t) < V_{\mathcal{D}}(x) + \varepsilon\varphi(x)$ in Ω for $t \geq \max\{T_1, T_2\}$. Similarly, we can prove $V_{\mathcal{D}}(x) - \varepsilon\varphi(x) < V(x, t)$ in Ω when t is large.

Making use of (4.2), we see that (H_i, V_i) satisfies

$$\begin{cases} \partial_t H_i - \mathcal{L}_1 H_i = -\rho(x)H_i + \sigma_1(x)H_u(x)V_i, & x \in \Omega, t > T_\varepsilon, \\ \partial_t V_i - \mathcal{L}_2 V_i \leq \sigma_2(x)(V_{\mathcal{D}}(x) + \varepsilon\varphi(x) - V_i)^+ H_i - \mu(x)(V_{\mathcal{D}}(x) - \varepsilon\varphi(x))V_i, & x \in \Omega, t > T_\varepsilon, \\ H_i = V_i = 0, & x \in \partial\Omega, t > T_\varepsilon. \end{cases}$$

Let $(H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ be the unique positive solution of (4.1 $_\varepsilon$), and (ϕ_1, ϕ_2) be the positive eigenfunction corresponding to $\lambda^{\mathcal{D}}(V_{\mathcal{D}}; \varepsilon)$. According to [11, Lemma 2.1], we can take constants $k \gg 1$ and $0 < \delta \ll 1$ such that

$$k(H_{i,\varepsilon}^*(x), V_{i,\varepsilon}^*(x)) \geq (H_i(x, T_\varepsilon), V_i(x, T_\varepsilon)), \quad \delta(\phi_1(x), \phi_2(x)) \leq (H_i(x, T_\varepsilon), V_i(x, T_\varepsilon)).$$

Then $k(H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ and $\delta(\phi_1, \phi_2)$ are the upper and lower solutions of (4.1 $_\varepsilon$), respectively. Let $(H_{i,\varepsilon}, V_{i,\varepsilon})$ be the unique positive solution of

$$\begin{cases} \partial_t H_i - \mathcal{L}_1 H_i = -\rho(x)H_i + \sigma_1(x)H_u(x)V_i, & x \in \Omega, t > 0, \\ \partial_t V_i - \mathcal{L}_2 V_i = \sigma_2(x)(V_D(x) + \varepsilon - V_i)^+ H_i - \mu(x)(V_D(x) - \varepsilon)V_i, & x \in \Omega, t > 0, \\ H_i = V_i = 0, & x \in \partial\Omega, t > 0, \\ (H_i(x, 0), V_i(x, 0)) = k(H_{i,\varepsilon}^*, V_{i,\varepsilon}^*), & x \in \bar{\Omega}. \end{cases}$$

Then $(H_{i,\varepsilon}(x, t), V_{i,\varepsilon}(x, t))$ is decreasing in t , and

$$\delta(\phi_1(x), \phi_2(x)) \leq (H_{i,\varepsilon}(x, t), V_{i,\varepsilon}(x, t)), \quad (H_i(x, t + T_\varepsilon), V_i(x, t + T_\varepsilon)) \leq (H_{i,\varepsilon}(x, t), V_{i,\varepsilon}(x, t))$$

by the comparison principle. Therefore, $\lim_{t \rightarrow \infty} (H_{i,\varepsilon}(x, t), V_{i,\varepsilon}(x, t)) = (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ in $[C^2(\bar{\Omega})]^2$ by the uniform estimate and compact arguments (cf. [10, Theorems 2.11, 3.14]) and the uniqueness of positive solutions of (4.1 $_\varepsilon$). So $\limsup_{t \rightarrow \infty} (H_i(x, t), V_i(x, t)) \leq (H_{i,\varepsilon}^*, V_{i,\varepsilon}^*)$ and, by letting $\varepsilon \rightarrow 0$,

$$\limsup_{t \rightarrow \infty} (H_i(x, t), V_i(x, t)) \leq (H_i^*, V_i^*) \quad \text{uniformly in } \bar{\Omega}.$$

On the other hand, let $W(x) = V_D(x) - V_i^*(x)$. Then $W(x) > 0$ as $V_i^*(x) < V_D(x)$ in Ω . Through the direct calculations we have

$$-\mathcal{L}_2 W + [\sigma_2(x)H_i^* + \mu(x)V_D(x)]W = \beta(x)V_D > 0.$$

By the Hopf boundary lemma, $\partial_\nu W|_{\partial\Omega} < 0$. Making use of [11, Lemma 2.1], we can find a $0 < r_0 \leq \varepsilon_0$ such that $W > 2r\varphi$, i.e.,

$$V_i^*(x) < V_D(x) - 2r\varphi(x) \quad \text{in } \Omega \quad \forall 0 < r < r_0.$$

For such r , as $\lim_{\varepsilon \rightarrow 0} V_{i,\varepsilon}^* = V_i^*$ in $C^2(\bar{\Omega})$ (the uniform estimate and compact arguments), similar to the proof of (4.2), we can find $0 < \bar{\varepsilon} < \varepsilon_0$ such that

$$V_{i,\bar{\varepsilon}}^*(x) < V_i^*(x) + \frac{r}{2}\varphi(x) \quad \text{in } \Omega.$$

Owing to $\lim_{t \rightarrow \infty} V_{i,\bar{\varepsilon}}(x, t) = V_{i,\bar{\varepsilon}}^*$ in $C^2(\bar{\Omega})$, by (4.2), there exists $\bar{T}_{\bar{\varepsilon}} \gg 1$ such that

$$V_{i,\bar{\varepsilon}}(x, t) < V_{i,\bar{\varepsilon}}^*(x) + \frac{r}{2}\varphi(x) < V_i^*(x) + r\varphi(x) < V_D(x) - r\varphi(x) \quad \text{in } \Omega, \quad \forall t \geq \bar{T}_{\bar{\varepsilon}}.$$

Therefore, when $t \geq T_{\bar{\varepsilon}} + \bar{T}_{\bar{\varepsilon}}$, we have $V_i(x, t) \leq V_{i,\bar{\varepsilon}}(x, t - T_{\bar{\varepsilon}}) < V_D(x) - r\varphi(x)$ in Ω . This combined with (4.2) indicates that there exists $T_r \gg 1$ such that

$$V_i(x, t) < V_D(x) - r\varphi(x), \quad 0 < V_D(x) - r\varphi(x) \leq V(x, t) \leq V_D(x) + r\varphi(x)$$

for all $t \geq T_r$ and $x \in \Omega$. Consequently, (H_i, V_i) satisfies

$$\begin{cases} \partial_t H_i - \mathcal{L}_1 H_i = -\rho(x)H_i + \sigma_1(x)H_u(x)V_i, & x \in \Omega, t > T_r, \\ \partial_t V_i - \mathcal{L}_2 V_i \geq \sigma_2(x)(V_D(x) - r\varphi - V_i)H_i - \mu(x)(V_D(x) + r\varphi)V_i \\ \quad = \sigma_2(x)(V_D(x) - r\varphi - V_i)^+ H_i - \mu(x)(V_D(x) + r\varphi)V_i, & x \in \Omega, t > T_r, \\ H_i = V_i = 0, & x \in \partial\Omega, t > T_r, \\ H_i(x, T_r) > 0, \quad 0 < V_i(x, T_r) < V_D(x) - r\varphi, & x \in \bar{\Omega}. \end{cases}$$

The remaining proof is the same as that of Theorem 3.2(i), and the details are omitted.

(ii) The proof is the same as that of Theorem 3.2(ii), and the details are also omitted.

(iii) In such case, (2.3 \mathcal{D}) has no positive solution and $\lim_{t \rightarrow \infty} (V_u + V_i) = 0$. Therefore, $\lim_{t \rightarrow \infty} H_i = 0$ by the first equation of (1.2). The proof is complete. \square

Define boundary operators $\mathcal{B}_j[W] = a_j(x)\partial_\nu W + b_j(x)W$, where either $a_j(x) \equiv 0, b_j(x) \equiv 1$ or $a_j(x) \equiv 1, b_j(x) \geq 0$ with $b_j \in C^{1+\alpha}(\partial\Omega)$, $j = 1, 2$. It should be mentioned that when the boundary conditions in (1.1) and (1.2) are replaced by $\mathcal{B}_1[H_i] = 0, \mathcal{B}_2[V_u] = \mathcal{B}_2[V_i] = 0$, then the conclusions of this paper are still true and the proofs are similar.

Acknowledgment: The author greatly appreciates the suggestions provided by Dr. Lei Li and Dr. Wenjie Ni in the proof of (4.2).

References

- [1] PAHO, Chikungunya, <http://www.paho.org/hq/index.php?option=comtopics&view=article&id=343&Itemid=40931&lang=en>.
- [2] PAHO, Zika virus infection, <http://www.paho.org/hq/index.php?option=comcontent&view=article&id=11585&Itemid=41688&lang=en>.
- [3] WHO, Dengue and severe dengue, <https://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue>.
- [4] WHO, EPI-WIN digest 7-Zika virus disease, <https://www.who.int/publications/m/item/epi-win-digest-7-zika-virus-disease>.
- [5] W.E. Fitzgibbon, J.J. Morgan and G.F. Webb, An outbreak vector-host epidemic model with spatial structure: the 2015-2016 Zika outbreak in Rio De Janeiro. *Theor. Biol. Med. Modell.* 14, 2-17 (2017).
- [6] P. Magal, G.F. Webb and Y.X. Wu, On a vector-host epidemic model with spatial structure. *Nonlinearity* 31, 5589-5614 (2018).
- [7] P. Magal, G.F. Webb and Y.X. Wu, On the basic reproduction number of reaction-diffusion epidemic models. *SIAM J. Appl. Math.* 79, 284-304 (2019).
- [8] G. Sweers, Strong positivity in $C(\overline{\Omega})$ for elliptic systems. *Math. Z.* 209, 251-271 (1992).
- [9] H. Amann, Fixed point equations and nonlinear eigenvalue problems in ordered Banach spaces. *SIAM Rev.* 18, 620-709 (1976).
- [10] M. X. Wang, *Nonlinear Second Order Parabolic Equations*. Boca Raton: CRC Press, 2021.
- [11] M.X. Wang and P.Y.H. Pang, *Nonlinear Second Order Elliptic Equations*. Heidelberg: Springer, 2024.