

# A proof of Sugawara's conjecture on Hasse-Weber ray class invariants

Patrick Morton

June 11, 2024

## Abstract

In this paper a proof is given of Sugawara's conjecture from 1936, that the ray class field of conductor  $\mathfrak{f}$  over an imaginary quadratic field  $K$  is generated over  $K$  by a single primitive  $\mathfrak{f}$ -division value of the  $\tau$ -function, first defined by Weber and then modified by Hasse in his 1927 paper giving a new foundation of complex multiplication.

## 1 Introduction.

Hasse's well-known paper [9] contains the first complete proof of Kronecker's *Jugendtraum* which only uses modular functions of level one and the Weierstrass  $\wp$ -function. An earlier, different proof was given by Takagi and Fueter [7, 8, 24], using modular functions of level four which were also defined in terms of the  $\wp$ -function. As part of his proof, Hasse shows that the ray class field  $K_{\mathfrak{m}}$  over an imaginary quadratic field  $K$  of a given conductor  $\mathfrak{m} \neq 1$  is generated over  $K$  by  $j(\mathfrak{k})$  and  $\tau(\mathfrak{k}^*)$ :

$$K_{\mathfrak{m}} = K(j(\mathfrak{k}), \tau(\mathfrak{k}^*));$$

here  $\mathfrak{k}^*$  is a ray class in the ray class group of  $K$  modulo  $\mathfrak{m}$ ,  $\mathfrak{m}\mathfrak{k}^{-1}$  is the ordinary ideal class containing  $\mathfrak{k}^*$ ,  $j(w)$  is Klein's  $j$ -function, and  $\tau(\mathfrak{k}^*) = \tau(\rho, \mathfrak{a})$  is Weber's  $\tau$ -function, for an element  $\rho$  and ideals  $\mathfrak{r}, \mathfrak{a}$  of  $K$  satisfying

$$\rho \cong \frac{\mathfrak{r}\mathfrak{a}}{\mathfrak{m}}, \text{ integral } \mathfrak{r} \in \mathfrak{k}^*, \mathfrak{a} \in \mathfrak{k},$$

and  $(\mathfrak{r}, \mathfrak{m}) = 1$ . (See below and [9, p. 138].) The value  $\tau(\rho, \mathfrak{a})$  is independent of the ideals  $\mathfrak{a} \in \mathfrak{k}$  and  $\mathfrak{r} \in \mathfrak{k}^*$  and the choice of the generator  $\rho$ , and therefore depends only on the ray class  $\mathfrak{k}^*$ .

In [26, p. 572], Weber defines the  $\tau$ -function as

$$\tau(u, \mathfrak{a}) = \begin{cases} \frac{g_2 g_3}{G} \wp(u), & g_2 g_3 \neq 0, \\ \frac{\wp(u)^2}{g_2}, & g_3 = 0, \\ \frac{\wp(u)^3}{g_3}, & g_2 = 0; \end{cases}$$

where  $\wp(u) = \wp(u, \mathbf{a})$ ,  $\wp'(u)^2 = 4\wp(u)^3 - g_2\wp(u) - g_3$ , and

$$16G = g_2^3 - 27g_3^2 = \Delta.$$

On the other hand, Hasse, in [9, p. 127], sets

$$\tau(u, \mathbf{a}) = \begin{cases} -2^7 3^5 \frac{g_2 g_3}{\Delta} \cdot \wp(u), & g_2 g_3 \neq 0, \\ 2^8 3^4 \frac{g_2^2}{\Delta} \cdot \wp(u)^2, & g_3 = 0, \\ -2^9 3^6 \frac{g_3}{\Delta} \cdot \wp(u)^3, & g_2 = 0. \end{cases}$$

See also [4, p. 34]. Except for the constant factors in front, this is the same normalization that is given in [21, p.135]. I use Hasse's normalization in this paper, though I am always assuming  $g_2 g_3 \neq 0$ , i.e., the corresponding quadratic field is not  $\mathbb{Q}(\omega)$  ( $\omega = (-1 + \sqrt{-3})/2$ ) or  $\mathbb{Q}(i)$ . (We can also eliminate quadratic fields  $K$  with class number 1, since the conjecture to be proved is obviously true for them.) In this case, note that the factor  $\lambda$  multiplying  $\wp(u)$  satisfies

$$\lambda^6 = \frac{12^6 j(\mathbf{a})^2 (j(\mathbf{a}) - 1728)^3}{\Delta}, \quad \lambda = -2^7 3^5 \frac{g_2 g_3}{\Delta}.$$

As has been pointed out in [11, 12], Hasse asked Hecke whether the ray class field  $K_{\mathfrak{f}}$  of conductor  $\mathfrak{f} \neq 1$  could be generated over  $K$  by  $\tau(\mathfrak{f}^*)$  alone, and mentioned this question also in [10, p. 85]. See the discussion in [6, pp. 88-91], especially on p. 91. This was conjectured by Sugawara to be the case in his papers [22, 23], which give a partial answer to this question. This was also essentially conjectured by Hasse in [10, p. 85]. Hasse's question and Sugawara's papers are referred to in [5, p. 60]. Also see [20, p. 132].

Sugawara showed that the answer to Hasse's question is *yes*, if one of three conditions holds for the modulus  $\mathfrak{m}$ :

$$4 \mid \mathfrak{m}, \quad \mathfrak{m} \neq 4. \tag{1.1}$$

$$\text{There is a prime divisor } \mathfrak{p} \text{ of } 2 \text{ in } K \text{ with } \mathfrak{p}^2 \mid \mathfrak{m} \text{ and } \varphi(\mathfrak{m}) \geq 6. \tag{1.2}$$

$$\Psi(\mathfrak{m}) = N(\mathfrak{m}) \prod_{\mathfrak{p} \mid \mathfrak{m}} \left(1 - \frac{2}{N(\mathfrak{p})}\right) \geq 5. \tag{1.3}$$

In this paper I will complete the proof of Sugawara's conjecture, starting with his conditions (1.1)-(1.3). Progress on this conjecture has also been made by Jung, Koo, Shin and Yoon, who prove in several papers [11, 12] that Sugawara's conjecture is true for ideals  $\mathfrak{m} = (n)$ , where  $n$  is a natural number. In [11, 12] the methods are analytic, whereas the arguments I give here are algebraic and arithmetic, though they do rely on some results for modular functions in several previous papers. Also see [13], where the coordinates of torsion points on specially constructed elliptic curves are shown to generate ray class fields.

The proof I give here proceeds by cases. Sugawara's conditions show that his conjecture is true for all but finitely many ideals  $\mathfrak{m}$ , for a fixed imaginary

quadratic number field  $K$ . On the other hand, this leaves open the question for several infinite families of pairs  $(K, \mathfrak{m})$ . We note the following concerning the possible remaining pairs.

Condition (1.3) shows that the conjecture is true for any prime ideal  $\mathfrak{m} = \mathfrak{p}$  whose norm is at least 7, and also for any  $\mathfrak{m}$  divisible by such a prime ideal, if  $\mathfrak{m}$  is not divisible by a first degree prime divisor of 2. The same condition (with the same restriction on prime divisors of 2) applies if  $\mathfrak{m}$  is divisible by any inert prime other than 2 in  $K/\mathbb{Q}$ . If a prime ideal  $\mathfrak{p}$  of degree 1 does divide  $(2, \mathfrak{m})$ , then  $\mathfrak{p}^2 \mid \mathfrak{m}$  in order for  $\mathfrak{m}$  to be a conductor. In this case, if  $\mathfrak{m} = \mathfrak{p}^2 \mathfrak{a}$ , where  $(\mathfrak{a}, 2) = 1$  and  $\varphi(\mathfrak{a}) \geq 3$ , then by (2), Sugawara's conjecture is true. The same holds if  $\mathfrak{m} = \mathfrak{p}^e \mathfrak{a}$  with  $e = 3$  and  $\mathfrak{a} \neq 1$  or  $e \geq 4$ . Thus, we restrict ourselves to prime divisors of 2, 3 and 5 in building the ideal  $\mathfrak{m}$ , and we have the following possibilities, where  $\wp_l$  denotes a prime divisor of degree 1 of  $l$  and (2) is inert:

$$\begin{aligned} \mathfrak{m} &= (2), \wp_2^2 \cong 2, \wp_2^3, \wp_2^2 \wp_2'^2 \cong 4; \\ \mathfrak{m} &= \wp_3 \wp_3', \wp_3^2, (2) \wp_3 \wp_3', \wp_2^2 \wp_3; \\ \mathfrak{m} &= \wp_5, \wp_3 \wp_5, \wp_3 \wp_3' \wp_5. \end{aligned} \tag{1.4}$$

In the cases with an unramified first degree prime ideal, the discriminant  $d_K$  of  $K$  satisfies  $d_K \equiv 1 \pmod{8}$  (for  $\wp_2$ );  $d_K \equiv 1 \pmod{3}$  (for  $\wp_3$ ); and  $d_K \equiv 1, 4 \pmod{5}$  (for  $\wp_5$ ). Five of the above cases are covered by the results of [11, 12] and six are not; namely, the cases

$$\mathfrak{m} = \wp_2^3, \wp_3^2, \wp_2^2 \wp_3, \wp_5, \wp_3 \wp_5, \wp_3 \wp_3' \wp_5$$

require a new argument. In particular, the fifth and sixth cases in this list are by far the most difficult. (Note that the ideal  $\mathfrak{m} = (2) \wp_3$  is not a possibility because it is not a conductor, since  $\mathbb{K}_{\mathfrak{m}} \supseteq \mathbb{K}_{(2)}$  and  $\varphi(\mathfrak{m})/2 = \varphi((2)) = [\mathbb{K}_{(2)} : \Sigma]$ .)

Converting to algebraic notation, the  $\tau$ -invariants for  $\mathfrak{m}$  and  $K \neq \mathbb{Q}(\sqrt{-3})$  or  $\mathbb{Q}(\sqrt{-4})$  can be written in the form

$$\tau(\mathfrak{k}^*) = -2^7 3^5 \frac{g_2 g_3}{\Delta} X(P) = h(P),$$

where  $P$  is a primitive  $\mathfrak{m}$ -division point with  $X$ -coordinate  $X(P)$  on an elliptic curve  $E$  in Weierstrass normal form with complex multiplication by  $R_K$  and invariants  $g_2, g_3, \Delta$ . It is clear that the expressions  $h(P)$  are independent of the particular model of the elliptic curve  $E$  which is used to compute the torsion points  $P$ . (See [21, p. 135].)

The general method presented here is to take an elliptic  $E$  of a special form which is determined in each case by the ideal  $\mathfrak{m}$ , and give explicit formulas for enough points in  $E[\mathfrak{m}]$  to be able to determine the  $\tau$ -invariants for  $\mathfrak{m}$ . We take  $E$  to be either: the Legendre normal form  $E_2$ , the Deuring normal form  $E_3$ , or the Tate normal form  $E_n$  for a point of order  $n$ , where  $n \in \{4, 5, 9, 12\}$ . In each case, the curve  $E$  is defined by certain parameters lying either in the Hilbert class field of  $K$  or in an abelian extension of small conductor over  $K$ .

In many of the cases, the prime ideal factorization of one of these parameters plays a decisive role in our arguments. Most of these factorizations have been determined in previous papers [14, 17, 18, 19, 1]. In the last two cases in (1.4) the ray class invariants for  $\mathfrak{m}$  are computed by finding the points of order 3 on  $E_5$ , which leads to some interesting arithmetic relationships.

In each case we must show that the  $\tau$ -invariants  $\tau(\mathfrak{k}^*)$  for ray classes  $\mathfrak{k}^*$  corresponding to the ideal class  $\mathfrak{k}$  and  $j$ -invariant  $j(\mathfrak{k})$  are distinct, as  $\mathfrak{k}$  runs over all ideal classes in the ring of integers  $R_K$  of  $K$ . Hasse shows in [10, pp. 83-85] that the ray class polynomial

$$T_{\mathfrak{m}}(t, j(\mathfrak{k})) = \prod_{\mathfrak{k}^* \text{ for } \mathfrak{k}} (t - \tau(\mathfrak{k}^*)) \in K[t, j(\mathfrak{k})],$$

whose roots are the  $\tau$ -invariants corresponding to a given ideal class, has coefficients in the Hilbert class field  $\Sigma = K_1$  and is irreducible over this field. Thus, the set of  $\tau$ -invariants for the ideal class  $\mathfrak{k}$  is a complete set of conjugates over  $\Sigma$ , and different ray class polynomials are conjugate by automorphisms of  $\Sigma/K$ . We will show that the invariants  $\tau(\mathfrak{k}^*)$  for a given class  $\mathfrak{k}$  determine  $j(\mathfrak{k})$ . To do this we will often assume that the automorphism  $\psi \in \text{Gal}(K_{\mathfrak{m}}/K)$  takes  $j(\mathfrak{k})$  to  $j(\mathfrak{k})^{\psi} = j(\mathfrak{k}')$  and leaves invariant the set  $\{\tau(\mathfrak{k}^*) \mid \mathfrak{k}^* \text{ for } \mathfrak{k}\}$  taken as a whole. Then our task is to show that  $\psi$  restricted to the Hilbert class field  $\Sigma$  is 1, implying that  $j(\mathfrak{k}) = j(\mathfrak{k}')$  and  $\mathfrak{k} = \mathfrak{k}'$ . Since the  $\tau$  invariants corresponding to a given class  $\mathfrak{k}$  are distinct [9, Satz 20], this proves the assertion. In Hasse's papers, he takes the ideal class corresponding to  $\mathfrak{k}^*$  to be the class containing  $\frac{\mathfrak{m}}{\mathfrak{k}^*}$ , i.e., the ideal class containing the ideals  $\frac{\mathfrak{m}}{\mathfrak{r}}, \mathfrak{r} \in \mathfrak{k}^*$ . It is clear, however, that we can take any correspondence between ray classes and ideal classes, as long as all ray classes *belonging* to a given ideal class *correspond* to a single class  $\mathfrak{k}$  and this correspondence is preserved by automorphisms over  $K$ .

Combining the results proved here with Suagawara's results [22, 23] gives the following.

**Main Theorem.** *If  $K_{\mathfrak{f}}$  is the ray class field with conductor  $\mathfrak{f} \neq 1$  over the imaginary quadratic field  $K$ , then  $K_{\mathfrak{f}} = K(\tau(\mathfrak{k}^*))$  is generated over  $K$  by a single  $\tau$ -invariant for the ideal  $\mathfrak{f}$ .*

In the cases when  $d_K \equiv 1$  modulo 8 or 3, I also show that the  $\tau$ -invariant for any of the respective ideals  $\wp_2, \wp_2\wp_2', \wp_3$  generates the Hilbert class field  $\Sigma = K_1$  of  $K$ . Thus, for these two families of quadratic fields, the  $j$ -invariant is not needed to generate the abelian extensions of  $K$ . See Theorems 2 and 7 in Sections 2 and 5.

## 2 The case $\mathfrak{m} = (2)$ .

We begin with the easiest case (from a computational point of view). We consider the Legendre normal form

$$E_2 : Y^2 = X(X - 1)(X - a),$$

which we assume to have complex multiplication by the quadratic field  $K$ , whose discriminant  $d_K$  satisfies  $d_K \equiv 5 \pmod{8}$ . Its associated Weierstrass normal form is

$$E' : Y_1^2 = 4X_1^3 - g_2X_1 - g_3,$$

where  $X_1 = X - \frac{a+1}{3}$ ,

$$g_2 = \frac{4}{3}(a^2 - a + 1), \quad g_3 = \frac{4}{27}(a+1)(a-2)(2a-1),$$

and

$$\Delta = 16a^2(a-1)^2.$$

Thus

$$j(E_2) = j(a) = \frac{2^8(a^2 - a + 1)^3}{a^2(a-1)^2},$$

and

$$j(a) - 1728 = \frac{64(a+1)^2(a-2)^2(2a-1)^2}{a^2(a-1)^2}.$$

We compute the ray class invariants for  $\mathfrak{m} = (2)$  and a given ideal class  $\mathfrak{k}$  for which  $j(a) = j(\mathfrak{k})$  to be

$$\begin{aligned} \tau_0 &= \frac{-2^7 3^5 g_2 g_3}{\Delta} \left(0 - \frac{a+1}{3}\right) = \frac{128(a^2 - a + 1)(a+1)^2(a-2)(2a-1)}{a^2(a-1)^2}; \\ \tau_1 &= \frac{-2^7 3^5 g_2 g_3}{\Delta} \left(1 - \frac{a+1}{3}\right) = \frac{128(a^2 - a + 1)(a+1)(a-2)^2(2a-1)}{a^2(a-1)^2}; \\ \tau_2 &= \frac{-2^7 3^5 g_2 g_3}{\Delta} \left(a - \frac{a+1}{3}\right) = \frac{-128(a^2 - a + 1)(a+1)(a-2)(2a-1)^2}{a^2(a-1)^2}. \end{aligned}$$

We note that

$$\tau_0 + \tau_1 + \tau_2 = 0 \tag{2.1}$$

and the  $\tau_i$  are roots of the cubic polynomial

$$F(X, j(\mathfrak{k})) = X^3 - 3j(\mathfrak{k})(j(\mathfrak{k}) - 1728)X + 2j(\mathfrak{k})(j(\mathfrak{k}) - 1728)^2.$$

Hasse proved in [10] that this polynomial is irreducible over the Hilbert class field  $\mathbb{K}_1$  of  $K$ . Furthermore, the formulas for the coefficients imply that

$$\frac{1}{\tau_0} + \frac{1}{\tau_1} + \frac{1}{\tau_2} = \frac{3}{2(j(\mathfrak{k}) - 1728)}. \tag{2.2}$$

This immediately implies that if  $\mathfrak{k}, \mathfrak{k}'$  are two ideal classes in  $K$  for which the sets  $\{\tau_0, \tau_1, \tau_2\}$  and  $\{\tau'_0, \tau'_1, \tau'_2\}$  coincide, then  $j(\mathfrak{k}) = j(\mathfrak{k}')$  and therefore  $\mathfrak{k} = \mathfrak{k}'$ . This implies, by Hasse's argument [10, p. 85], that  $K(\tau_i) = \mathbb{K}_{\mathfrak{m}}$ , so that Sugawara's conjecture is true for  $\mathfrak{m} = (2)$  and  $d_K \equiv 5 \pmod{8}$ .

We also note the relation

$$\frac{\tau_2 - \tau_0}{\tau_1 - \tau_0} = a.$$

It follows that  $a \in \mathbb{K}_{(2)}$  and the above formulas show that  $K(a) = \mathbb{K}_{(2)}$ .

**Theorem 1.** *If the Legendre normal form  $E_2$  has complex multiplication by the ring of integers in the quadratic field  $K = \mathbb{Q}(\sqrt{d_K})$ , where  $d_K \equiv 5 \pmod{8}$ , then the parameter  $a$  generates the ray class field  $K_{(2)}$  of conductor  $\mathfrak{m} = (2)$  over  $K$ . Also,  $K_{(2)}$  is generated over  $K$  by a single  $\tau$ -invariant for the ideal  $(2)$ .*

Now the same formulas hold if  $4 \mid d_K$  and  $d_K < -4$ , in which case  $(2) = \mathfrak{p}^2$  and  $\varphi(\mathfrak{p}^2) = 2$ . In this case there are two ray class invariants  $\tau(\mathfrak{k}^*)$  for each ideal class  $\mathfrak{k}$ . This implies that the polynomial  $F(x, j(\mathfrak{k}))$  is reducible over  $K(j(\mathfrak{k}))$  and factors into a linear times an irreducible quadratic. If the two ray class invariants  $(\text{mod } (2))$  for two ideal classes agree, i.e.  $\{\tau_i, \tau_k\} = \{\tau'_i, \tau'_k\}$ , corresponding to  $\mathfrak{k}$  and  $\mathfrak{k}'$ , then the third ray class invariants (corresponding to  $\mathfrak{m} = \mathfrak{p}$ ) must also agree, by (2.1), and then (2.2) implies that  $j(\mathfrak{k}) = j(\mathfrak{k}')$ . Thus, Sugawara's conjecture also holds when  $\mathfrak{m} = \mathfrak{p}^2 = (2)$ . This establishes the conjecture for the first two possibilities in the first line of Section 1, (1.4).

Lastly, suppose that  $(2) = \mathfrak{p}_1\mathfrak{p}_2$  in  $K$ , so that  $d_K \equiv 1 \pmod{8}$ . Since  $\varphi(\mathfrak{p}_1) = \varphi(\mathfrak{p}_2) = \varphi(\mathfrak{p}_1)\varphi(\mathfrak{p}_2) = 1$ , there is one ray class invariant each for  $\mathfrak{m} = \mathfrak{p}_1, \mathfrak{p}_2$ , and  $\mathfrak{p}_1\mathfrak{p}_2 = (2)$ . Each of these invariants lies in  $K_1$ . Together, they generate  $K_1$ , by (2.2).

I conjecture the following.

**Conjecture 1.** *If  $d_K \equiv 1 \pmod{8}$  and  $(2) = \mathfrak{p}_1\mathfrak{p}_2$ , the invariants for  $\mathfrak{m} = \mathfrak{p}_1, \mathfrak{p}_2$  have degree  $2h(d_K)$  over  $\mathbb{Q}$ , so that each generates  $K_1$  over  $\mathbb{Q}$ ; while the invariant for  $\mathfrak{p}_1\mathfrak{p}_2$  has degree  $h(d_K)$  over  $\mathbb{Q}$  and generates  $K_1$  over  $K$ .*

We can approach this conjecture by showing that the discriminant of  $F(X, j(\mathfrak{k}))$  is negative, when  $\mathfrak{k} = \mathfrak{o}$  is the principal class. This discriminant is

$$\text{disc}(F(X, j(\mathfrak{k}))) = 2^8 3^6 j^2 (j - 1728)^3, \quad j = j(\mathfrak{k}).$$

If this discriminant is negative, then two of the invariants  $\tau_i(\mathfrak{k})$  are complex and one is real. The complex invariants would have to be the invariants for  $\mathfrak{m} = \mathfrak{p}_1, \mathfrak{p}_2$ , which are interchanged by complex conjugation. Hence, the real invariant is the invariant for  $\mathfrak{m} = \mathfrak{p}_1\mathfrak{p}_2$ .

**Lemma 1.** *If  $d_K \equiv 1 \pmod{8}$ , then the  $j$ -invariant  $j(\mathfrak{o})$  of the principal class  $\mathfrak{o}$  in  $R_K$  satisfies  $j(\mathfrak{o}) < 1728$ .*

*Proof.* This will follow from the formula

$$j(\mathfrak{o}) - 1728 = \frac{64(a+1)^2(a-2)^2(2a-1)^2}{a^2(a-1)^2} = R(a)^2,$$

where

$$R(a) = \frac{8(a+1)(a-2)(2a-1)}{a(a-1)}.$$

From [14, Eq. (7.1), p. 1979, (i)], we have

$$j(\mathfrak{o}) - 1728 = j(\mathfrak{p}_1)^\tau - 1728 = R(a)^{2\tau} = R\left(-\frac{\xi^4}{\pi^4}\right)^{2\tau},$$

where  $\tau = \left(\frac{\Sigma/K}{\mathfrak{p}_1}\right)$ ,  $a = -\frac{\xi^4}{\pi^4}$ , and  $\pi^4 + \xi^4 = 1$ ,  $\pi \cong \mathfrak{p}_1$ ,  $\xi \cong \mathfrak{p}_2$ . For this value of  $a$ , we have

$$\begin{aligned} R\left(-\frac{\xi^4}{\pi^4}\right) &= \frac{8(\pi^4 - \xi^4)(\pi^4 + 2\xi^4)(2\pi^4 + \xi^4)}{\pi^4\xi^4(\pi^4 + \xi^4)} \\ &= \frac{8(\pi^4 - \xi^4)(1 + \xi^4)(1 + \pi^4)}{\pi^4\xi^4}. \end{aligned}$$

Also from [14], the automorphism  $\sigma : \xi \rightarrow \frac{\xi+1}{\xi-1} = \pi^{\tau^2}$  is complex conjugation, for which  $\pi^\sigma = \xi^{\sigma\tau^{-2}\sigma} = \xi^{\tau^2}$ . Hence

$$\begin{aligned} R\left(-\frac{\xi^4}{\pi^4}\right)^{\tau\sigma} &= R\left(-\frac{\xi^4}{\pi^4}\right)^{\sigma\tau^{-1}} \\ &= -\left(\frac{8(\pi^4 - \xi^4)(1 + \xi^4)(1 + \pi^4)}{\pi^4\xi^4}\right)^{\tau^2\tau^{-1}} \\ &= -R\left(-\frac{\xi^4}{\pi^4}\right)^\tau. \end{aligned}$$

This implies that  $R(a)^\tau$  is pure imaginary, hence  $j(\mathfrak{o}) - 1728 < 0$ , proving the lemma.  $\square$

**Theorem 2.** *If  $d_K \equiv 1 \pmod{8}$  and  $(2) = \mathfrak{p}_1\mathfrak{p}_2$  in  $K = \mathbb{Q}(\sqrt{-d})$ , the  $\tau$ -invariant  $\tau(\mathfrak{k}^*)$  for any of the ideals  $\mathfrak{p}_1$ ,  $\mathfrak{p}_2$  or  $\mathfrak{p}_1\mathfrak{p}_2$  and any ideal class  $\mathfrak{k}$  generates the Hilbert class field  $K_1 = \Sigma$  over  $K$ .*

*Proof.* We have to show that each of the invariants  $\tau_i$  generates  $\Sigma$  over  $K$ . By [14, p. 1979, (ii)], we can take  $a = \pi^4$ , where  $(\pi) = \mathfrak{p}_1$ , as in the proof of Lemma 1.

Assume that  $\tau_0 = \tau'_0$  for two ideal classes  $\mathfrak{k}$  and  $\mathfrak{k}'$ . Let  $\alpha \in \text{Gal}(\Sigma/K)$  be such that  $j(\mathfrak{k}') = j(\mathfrak{k})^\alpha$ . Suppose that  $\tau_0 = \tau'_0$ . Setting  $x = a = \pi^4$  and  $y = a' = \pi'^4 \cong \pi^4$  yields that

$$\begin{aligned} \tau_0 - \tau'_0 &= \frac{(-y+x)(xy-1)}{x^2(x-1)^2y^2(y-1)^2} [2x^4y^3 + 2x^3y^4 - 4x^4y^2 - 7x^3y^3 - 4x^2y^4 \\ &\quad + 2x^4y + 10x^3y^2 + 10x^2y^3 + 2xy^4 - 7x^3y - 8x^2y^2 - 7xy^3 \\ &\quad + 2x^3 + 10x^2y + 10xy^2 + 2y^3 - 4x^2 - 7xy - 4y^2 + 2x + 2y]. \end{aligned}$$

The terms in square brackets whose total degrees in  $x$  and  $y$  are at least 3 are divisible by  $\pi^{12}$ . If the expression in square brackets is zero, then

$$-4x^2 - 7xy - 4y^2 + 2x + 2y \equiv 0 \pmod{\pi^{12}}.$$

Now, the powers of  $\pi$  dividing the terms in this expression are, respectively, 10, 8, 10, 5, 5. It follows that  $2x + 2y \equiv 0 \pmod{\pi^8}$  and therefore

$$a + a' = \pi^4 + \pi'^4 = \pi^4 + \pi^{4\alpha} \equiv 0 \pmod{\mathfrak{p}_1^7}.$$

Hence,  $a - a' = a + a' - 2a' \equiv 0 \pmod{\pi^5}$ . But then

$$\begin{aligned} j(\mathfrak{k}) - j(\mathfrak{k}') &= \frac{2^8(a^2 - a + 1)^3}{a^2(a-1)^2} - \frac{2^8(a'^2 - a' + 1)^3}{a'^2(a'-1)^2} \\ &= \frac{2^8(aa' - a + 1)(aa' - a' + 1)(a + a' - 1)(aa' - 1)(a - a')(aa' - a - a')}{a^2(a-1)^2 a'^2(a'-1)^2} \\ &= \frac{2^8}{a^2(a-1)^2} \frac{(a - a')(aa' - a - a')}{a'^2(a'-1)^2} \mathbf{A}, \end{aligned}$$

where

$$\mathbf{A} = [(aa' - a + 1)(aa' - a' + 1)(a + a' - 1)(aa' - 1)].$$

Now  $\frac{256}{a^2(a-1)^2} = \frac{256}{\pi^8 \xi^8}$  is a unit. Also,  $\frac{(a-a')(aa'-a-a')}{a'^2(a'-1)^2}$  is divisible by  $\pi^5 \pi^7 / \pi^8 = \pi^4$ , so

$$j(\mathfrak{k}) \equiv j(\mathfrak{k}') = j(\mathfrak{k})^\alpha \pmod{\mathfrak{p}_1^4}.$$

But the discriminant of the class equation  $H_{d_K}(X)$  is not divisible by 2, since  $d_K \equiv 1 \pmod{8}$ , which implies that  $\alpha = 1$  and  $j(\mathfrak{k}') = j(\mathfrak{k})^\alpha = j(\mathfrak{k})$ .

This argument applies under the assumption that the expression in square brackets is 0. Otherwise,  $y = x$  or  $y = 1/x$ , which implies  $a = a'$  or  $a = 1/a'$ , and we conclude again that  $j(\mathfrak{k}') = j(\mathfrak{k})$ .

Next suppose that  $\tau_0 = \tau'_1$ . Setting  $x = a = \pi^4$  and  $y = a' = \pi'^4 \cong \pi^4$  yields that

$$\begin{aligned} \tau_0 - \tau'_1 &= \frac{1}{x^2(x-1)^2 y^2(y-1)^2} (y-1+x)(xy-x+1)[2x^4 y^3 - 2x^3 y^4 - 2x^4 y^2 \\ &\quad + x^3 y^3 + 4x^2 y^4 - x^3 y^2 - 6x^2 y^3 - 2xy^4 + 2x^2 y^2 + xy^3 + 8x^2 y + 2y^3 \\ &\quad - 4x^2 - 2y^2 - xy^2]. \end{aligned}$$

The terms in square brackets whose total degrees in  $x$  and  $y$  are at least 4 are divisible by  $\pi^{16}$ . If the expression in square brackets is zero, then

$$8x^2 y + 2y^3 - 4x^2 - 2y^2 - xy^2 \equiv 0 \pmod{\pi^{16}}.$$

But the powers of  $\pi$  dividing the terms in this expression are, respectively, 15, 13, 10, 9 and 12. Thus the left side is exactly divisible by  $\pi^9$  and this congruence is impossible. Hence we must have  $(y-1+x)(xy-x+1) = 0$ , so that  $y = 1-x$  or  $y = \frac{x-1}{x}$ . Both expressions are anharmonic transformations in  $a$  which fix the  $j$ -invariant. Hence  $j(\mathfrak{k}') = j(\mathfrak{k})$ .

A similar analysis applies if  $\tau_1 = \tau'_1$  or  $\tau_1 = \tau'_2$ . Combining this with the previous case, which also applies to  $\tau_1 = \tau'_0$ , shows that the conjugates of  $\tau_1$  over  $K$  are distinct, hence  $K(\tau_1) = \Sigma$ .



If we apply similar reasoning as in the previous case to the relations  $\tau_0 = \tau'_2$  or  $\tau_2 = \tau'_2$  we find that there are two terms in  $x$  and  $y$  (in square brackets) with minimum valuation; and these terms are  $2x - 2y$  for the first case and  $2x + 2y$  for the second, both with valuation 5. In both cases the next smallest valuation is 8, which occurs for the terms  $-xy$  and  $9xy$ . We conclude that

$$2a \pm 2a' \equiv 0 \pmod{\pi^8},$$

giving that

$$a \pm a' \equiv 0 \pmod{\pi^7}.$$

Now the same argument as in the case  $\tau_0 = \tau'_0$ , applied to the difference  $j(\mathfrak{k}) - j(\mathfrak{k}')$ , shows that

$$j(\mathfrak{k}) \equiv j(\mathfrak{k}') \pmod{\mathfrak{p}_1^4}$$

and hence  $j(\mathfrak{k}) = j(\mathfrak{k}')$  and  $\mathfrak{k} = \mathfrak{k}'$ .

It follows that the sets  $S(\mathfrak{k}) = \{\tau_0, \tau_1, \tau_2\}$  are disjoint for different ideal classes  $\mathfrak{k}$ . Since the conjugates over  $K$  of a  $\tau$ -invariant for any of the ideals  $\mathfrak{p}_1, \mathfrak{p}_2$ , or  $\mathfrak{p}_1\mathfrak{p}_2$  are also  $\tau$ -invariants for the same ideals, we conclude that the conjugates of any  $\tau(\mathfrak{k}^*)$  for one of these ideals are distinct. This proves the theorem.  $\square$

### 3 The case $\mathfrak{m} = (4)$ .

In this section we assume  $\mathfrak{m} = (4) = \mathfrak{p}_1^2\mathfrak{p}_2^2$  and the discriminant  $d_K \equiv 1 \pmod{8}$ . We compute the ray class invariants for divisors of  $\mathfrak{m} = (4)$  using the Tate normal form  $E_4$ , on which  $(0, 0)$  has order 4:

$$E_4 : Y^2 + XY + bY = X^3 + bX^2, \quad b = \frac{1}{\alpha^4} = \frac{\beta^4 - 16}{16\beta^4},$$

where  $16\alpha^4 + 16\beta^4 = \alpha^4\beta^4$  and  $\mathbb{Q}(\alpha) = \mathbb{Q}(\beta) = \mathbb{Q}(\beta^4) = \Sigma = \mathbb{K}_1$ , as in [lm]. The arguments in [14] yield specific values of  $\alpha$  and  $\beta$  for which  $E_4$  has complex multiplication by  $R_K$  in this case. The coefficients of the Weierstrass normal form of  $E_4$  are

$$g_2 = \frac{(\beta^4 + 4\beta^3 + 8\beta^2 - 16\beta + 16)(\beta^4 - 4\beta^3 + 8\beta^2 + 16\beta + 16)}{192\beta^8},$$

$$g_3 = -\frac{(\beta^4 + 16)(\beta^2 + 4\beta - 4)(\beta^2 - 4\beta - 4)(\beta^4 + 24\beta^2 + 16)}{13824\beta^{12}};$$

with

$$\Delta = \frac{(\beta - 2)^4(\beta + 2)^4(\beta^2 + 4)^4}{4096\beta^{20}}.$$

Thus,

$$j(E_4) = j(\beta) = \frac{(\beta^4 + 4\beta^3 + 8\beta^2 - 16\beta + 16)^3(\beta^4 - 4\beta^3 + 8\beta^2 + 16\beta + 16)^3}{(\beta - 2)^4(\beta + 2)^4(\beta^2 + 4)^4\beta^4},$$

$$j(\beta) - 1728 = \frac{(\beta^4 + 16)^2(\beta^2 + 4\beta - 4)^2(\beta^2 - 4\beta - 4)^2(\beta^4 + 24\beta^2 + 16)^2}{(\beta - 2)^4(\beta + 2)^4(\beta^2 + 4)^4\beta^4}.$$

We will see that here, the (primitive and nonprimitive) ray class invariants for divisors of  $\mathfrak{m}$  which are not divisors of (2) are roots of the polynomial

$$F(x, j(\mathfrak{k})) = X^6 - 15j(j - 1728)X^4 - 40j(j - 1728)^2X^3 - 45j^2(j - 1728)^2X^2 - 24j^2(j - 1728)^3X - j^2(j - 1728)^3(5j - 55296), \quad j = j(\mathfrak{k}).$$

This implies that the ray class invariants  $\tau_i = \tau(\mathfrak{k}_i^*)$  for  $\mathfrak{m} = \mathfrak{p}_1^2, \mathfrak{p}_2^2, \mathfrak{p}_1\mathfrak{p}_2, \mathfrak{p}_1\mathfrak{p}_2^2$  and  $\mathfrak{m} = (4)$  satisfy

$$\sum_{i=0}^5 \frac{1}{\tau_i} = \frac{-24}{5j(\mathfrak{k}) - 55296}. \quad (3.1)$$

We label the  $X$ -coordinates of points of order 4 on  $E_4$  in order as

$$X_0 = 0, \quad X_1 = -\frac{(\beta - 2)(\beta^2 + 4)}{8\beta^3}, \quad X_2 = -2b = -\frac{(\beta^4 - 16)}{8\beta^4},$$

$$X_3 = X_1(i\beta) = \frac{-(\beta + 2i)(\beta^2 - 4)}{8\beta^3}, \quad X_4 = X_1(-\beta) = -\frac{(\beta + 2)(\beta^2 + 4)}{8\beta^3},$$

$$X_5 = X_1(-i\beta) = \frac{-(\beta - 2i)(\beta^2 - 4)}{8\beta^3}.$$

See [14, pp. 1971-1974]. The corresponding ray class invariants are given by

$$\tau_i = -2^7 3^5 \frac{g_2 g_3}{\Delta} \left( X_i + \frac{4b + 1}{12} \right), \quad b = \frac{\beta^4 - 16}{16\beta^4}.$$

A calculation shows that these six quantities are the roots of the polynomial  $F(X, j(\mathfrak{k}))$  given above. In particular,

$$\tau_0 = -2^7 3^5 \frac{g_2 g_3}{\Delta} \frac{4b + 1}{12} = -2^7 3^5 \frac{g_2 g_3}{\Delta} \frac{5\beta^4 - 16}{48\beta^4}.$$

This gives

$$\frac{\tau_3 - \tau_0}{\tau_0} = -6 \frac{\beta(\beta - 2)(\beta + 2)(\beta + 2i)}{5\beta^4 - 16}, \quad (3.2)$$

$$\frac{\tau_5 - \tau_0}{\tau_0} = -6 \frac{\beta(\beta - 2)(\beta + 2)(\beta - 2i)}{5\beta^4 - 16}. \quad (3.3)$$

We also note that the map  $\sigma : \beta \rightarrow 2\frac{\beta+2}{\beta-2}$  induces the permutation

$$\tau_0^\sigma = \tau_4, \quad \tau_1^\sigma = \tau_2, \quad \tau_3^\sigma = \tau_5.$$

By the results of [14, Prop. 8.2],  $\sigma$  is an automorphism of  $K_1 = \mathbb{Q}(\beta)$ . It is also clear that  $\tau_0, \tau_1, \tau_2, \tau_4 \in K_1$ , while  $\tau_3, \tau_5 \in K_1(i) = K_m$ , the ray class field of conductor  $\mathfrak{m} = (4)$ , in the case that  $2 = \mathfrak{p}_1\mathfrak{p}_2$ .

Since

$$\frac{\tau_1 - \tau_0}{\tau_2 - \tau_0} = \frac{\beta}{\beta + 2},$$

we see that  $K(\tau_0, \tau_1, \tau_2) = K_1 = \Sigma$ .

**Theorem 3.** *The invariants  $\tau_0, \tau_1, \tau_2, \tau_4$  are the roots of the polynomial*

$$\begin{aligned} G(X, \tau_3, \tau_5) = & 4X^4 + 4(\tau_3 + \tau_5)X^3 - 6(\tau_3^2 + 6\tau_3\tau_5 + \tau_5^2)X^2 \\ & + 4(\tau_3 + \tau_5)(\tau_3^2 + 5\tau_3\tau_5 + \tau_5^2)X - \tau_3^4 - 6\tau_3^3\tau_5 - 6\tau_3^2\tau_5^2 - 6\tau_3\tau_5^3 - \tau_5^4. \end{aligned}$$

*Proof.* Taking the resultant

$$\begin{aligned} \text{Res}_\beta((5\beta^4 - 16)(\tau_3 - X) + 6\beta(\beta - 2)(\beta + 2)(\beta + 2i)X, \\ (5\beta^4 - 16)(\tau_5 - X) + 6\beta(\beta - 2)(\beta + 2)(\beta - 2i)X) \end{aligned}$$

yields

$$2^{26}3^4X^4G(X, \tau_3, \tau_5).$$

By the above relations (3.2) and (3.3) we see that  $G(\tau_0, \tau_3, \tau_5) = 0$ . Applying the map  $\sigma$  gives that  $G(\tau_4, \tau_5, \tau_3) = G(\tau_4, \tau_3, \tau_5) = 0$ , since the coefficients of  $G$  are symmetric in  $\tau_3$  and  $\tau_5$ . Similarly, the relations

$$\begin{aligned} \frac{\tau_3 - \tau_1}{\tau_1} &= \frac{12(1+i)\beta(\beta-2)(\beta+2i)}{\beta^4 - 12\beta^3 + 24\beta^2 - 48\beta + 16}, \\ \frac{\tau_5 - \tau_1}{\tau_1} &= \frac{12(1-i)\beta(\beta-2)(\beta-2i)}{\beta^4 - 12\beta^3 + 24\beta^2 - 48\beta + 16}, \end{aligned}$$

and the resultant

$$\begin{aligned} \text{Res}_\beta((\beta^4 - 12\beta^3 + 24\beta^2 - 48\beta + 16)(\tau_3 - X) - 12(1+i)\beta(\beta-2)(\beta+2i)X, \\ (\beta^4 - 12\beta^3 + 24\beta^2 - 48\beta + 16)(\tau_5 - X) - 12(1-i)\beta(\beta-2)(\beta-2i)X) \\ = 2^{26}3^4X^4G(X, \tau_3, \tau_5) \end{aligned}$$

yield that  $\tau_1$  is a root of  $G(X, \tau_3, \tau_5)$ . Applying  $\sigma$  yields that  $\tau_2$  is also a root. This proves the theorem.  $\square$

Note that

$$\text{disc}(G(X, \tau_3, \tau_5)) = -2^8(\tau_3 + 5\tau_5)^3(5\tau_3 + \tau_5)^3(\tau_3 - \tau_5)^6.$$

Furthermore,

$$(\tau_3 + 5\tau_5)(5\tau_3 + \tau_5) = (j(\beta) - 1728)R(\beta),$$

where

$$R(\beta) = \frac{36(\beta^4 + 4\beta^3 + 8\beta^2 - 16\beta + 16)^2(\beta^4 - 4\beta^3 + 8\beta^2 + 16\beta + 16)^2}{\beta^4(\beta + 2)^4(\beta - 2)^4}$$

is zero exactly when  $j(\beta) = 0$ , which is excluded since we are assuming  $d_K \neq -3, -4$ . Thus, in all cases we are considering,  $G(X, \tau_3, \tau_5)$  has distinct roots.

This theorem implies Sugawara's conjecture for  $\mathfrak{m} = \mathfrak{p}_1^2 \mathfrak{p}_2^2$ . Namely, the invariants  $\tau_3, \tau_5$  are the ray class invariants corresponding to a given ideal class  $\mathfrak{k}$ , since these are the only invariants for which  $K(j(\mathfrak{k}), \tau(\mathfrak{k}^*)) = K_{\mathfrak{m}} = K(j(\mathfrak{k}), i)$ . The other four invariants lie in  $K_1 = K(\beta)$ . For this note that  $\frac{\varphi(\mathfrak{m})}{2} = 2$ . Now, if  $\mathfrak{k}, \mathfrak{k}'$  are two ideal classes for which  $\{\tau_3, \tau_5\} = \{\tau'_3, \tau'_5\}$ , then the theorem implies that  $S = \{\tau_i : i = 0, 1, 2, 4\}$  coincides with  $S' = \{\tau'_i : i = 0, 1, 2, 4\}$ . Hence, (3.1) implies that  $j(\mathfrak{k}) = j(\mathfrak{k}')$  and  $\mathfrak{k} = \mathfrak{k}'$ .

If  $(2) = \mathfrak{p}$  and  $\mathfrak{m} = \mathfrak{p}^2$ , then  $\varphi(\mathfrak{m}) = 12$ , in which case the conjecture follows immediately from (1.2) in Section 1.

Finally, if  $(2) = \mathfrak{p}^2$ , then  $\mathfrak{m} = \mathfrak{p}^4$  and  $\varphi(\mathfrak{m}) = 8 > 6$ , so this case also follows from (1.2) in Section 1. See [23].

So far we have the following:

**Theorem 4.** *For all divisors  $\mathfrak{m}$  of the ideals (2) or (4), for which  $K_{\mathfrak{m}} \neq K_1$ , Sugawara's conjecture holds, namely,  $K_{\mathfrak{m}} = K(\tau(\mathfrak{k}^*))$  is generated over the quadratic field  $K$  by a single ray class invariant for the modulus  $\mathfrak{m}$ .*

## 4 The case $\mathfrak{m} = \wp_2^3$ .

In this section we consider the case  $d_K \equiv 1 \pmod{8}$  and  $\mathfrak{m} = \wp_2^3$ , to align with the notation in [1].

On the same curve we appealed to in Section 3,

$$E_4(b) : Y^2 + XY + bY = X^3 + bX^2, \quad b = \frac{1}{16} - \frac{1}{\beta^4},$$

the point  $Q = (-2b, 2b\beta_1\beta_3)$  has order 4, where

$$\beta_1\beta_3 = \frac{\beta + 2i}{2\beta} \frac{\beta - 2i}{2\beta} = \frac{\beta^2 + 4}{4\beta^2}.$$

Note that  $\xi = \beta/2$  and  $\pi = \beta/(\zeta_8^j \alpha)$  (for some odd integer  $j$ ; see [14, pp. 1978, 1984]) satisfy  $\pi^4 + \xi^4 = 1$  and are the same quantities that we encountered in the proof of Lemma 1 in Section 2.

If  $P = (x, y)$  is a point on  $E_4(b)$  for which  $2P = Q$ , then  $P$  has order 8. Hence, if  $x = X(P)$ , we have  $X(2P) = -2b$  or

$$0 = \frac{(x^4 - bx^2 - 2b^2x - b^3)}{(x+b)(4x^2 + x + b)} + 2b = \frac{x^4 + 8bx^3 + b(8b+1)x^2 + 2b^2x + b^3}{(x+b)(4x^2 + x + b)}.$$

We now solve

$$f_8(x) = x^4 + 8bx^3 + b(8b+1)x^2 + 2b^2x + b^3 = 0,$$

with  $b = \frac{1}{16} - \frac{1}{\beta^4} = \frac{1}{16} - \frac{1}{16\xi^4}$ . We have

$$\begin{aligned} f_8(x-2b) &= x^4 + (-16b^2 + b)x^2 + (32b^3 - 2b^2)x - 16b^4 + b^3 \\ &= x^4 + px^2 + qx + r, \end{aligned}$$

with

$$p = \frac{(\xi^4 - 1)}{16\xi^8}, \quad q = -\frac{(\xi^4 - 1)^2}{128\xi^{12}}, \quad r = \frac{(\xi^4 - 1)^3}{4096\xi^{16}}.$$

Then the cubic resolvent of  $f_8(x)$  factors completely:

$$x^3 - 2px^2 + (p^2 - 4r)x + q^2 = (x - \theta_1)(x - \theta_2)(x - \theta_3),$$

with roots

$$\begin{aligned} \theta_1 &= \frac{(\xi^4 - 1)}{16\xi^8} = -\frac{\pi^4}{16\xi^8}, \\ \theta_2 &= -\frac{(\xi^2 + 1)(\xi^2 - 1)^2}{32\xi^8} = -\frac{\pi^4(1 - \xi^2)}{32\xi^8}, \\ \theta_3 &= \frac{(\xi^2 - 1)(\xi^2 + 1)^2}{32\xi^8} = -\frac{\pi^4(\xi^2 + 1)}{32\xi^8}. \end{aligned}$$

Thus, the roots of  $x^4 + px^3 + qx^2 + r = 0$  are

$$\begin{aligned} x_1 &= \frac{1}{2} \left( \sqrt{-\theta_1} + \sqrt{-\theta_2} + \sqrt{-\theta_3} \right) \\ &= \frac{\pi^2}{8\xi^4} + \frac{\pi^2\sqrt{2}\sqrt{1-\xi^2}}{16\xi^4} + \frac{\pi^2\sqrt{2}\sqrt{\xi^2+1}}{16\xi^4}; \\ x_2 &= \frac{1}{2} \left( \sqrt{-\theta_1} - \sqrt{-\theta_2} - \sqrt{-\theta_3} \right) \\ &= \frac{\pi^2}{8\xi^4} - \frac{\pi^2\sqrt{2}\sqrt{1-\xi^2}}{16\xi^4} - \frac{\pi^2\sqrt{2}\sqrt{\xi^2+1}}{16\xi^4}; \\ x_3 &= \frac{1}{2} \left( -\sqrt{-\theta_1} + \sqrt{-\theta_2} - \sqrt{-\theta_3} \right) \\ &= -\frac{\pi^2}{8\xi^4} + \frac{\pi^2\sqrt{2}\sqrt{1-\xi^2}}{16\xi^4} - \frac{\pi^2\sqrt{2}\sqrt{\xi^2+1}}{16\xi^4}; \\ x_4 &= \frac{1}{2} \left( -\sqrt{-\theta_1} - \sqrt{-\theta_2} + \sqrt{-\theta_3} \right) \\ &= -\frac{\pi^2}{8\xi^4} - \frac{\pi^2\sqrt{2}\sqrt{1-\xi^2}}{16\xi^4} + \frac{\pi^2\sqrt{2}\sqrt{\xi^2+1}}{16\xi^4}. \end{aligned}$$

These roots correspond to points  $P_i$  for which  $X(P_i) = x_i - 2b$  are the roots of  $f_8(x) = 0$ . Since

$$x_1 + x_2 = \frac{\pi^2}{4\xi^4}, \quad x_1x_2 = \frac{\pi^6}{64\xi^8},$$

$x_1$  and  $x_2$  are roots of

$$m(X) = X^2 - \frac{\pi^2}{4\xi^4}X + \frac{\pi^6}{64\xi^8} \in \Sigma[X].$$

Similarly,  $x_3$  and  $x_4$  are roots of

$$\tilde{m}(x) = X^2 + \frac{\pi^2}{4\xi^4}X - \frac{\pi^6}{64\xi^8} = X^2 + (x_1 + x_2)X - x_1x_2.$$

The discriminant of  $m(X)$  is

$$\text{disc}(m(X)) = \frac{\pi^4}{16\xi^8} - 4\frac{\pi^6}{64\xi^8} = \frac{\pi^4(1 - \pi^2)}{16\xi^8},$$

so the roots of  $m(X)$  lie in  $\Sigma(\sqrt{1 - \pi^2})$ . Furthermore,  $\sqrt{1 - \pi^2}\sqrt{1 + \pi^2} = \sqrt{1 - \pi^4} = \xi^2$ , so  $\Sigma(\sqrt{1 - \pi^2}) = \Sigma(\sqrt{1 + \pi^2}) = \mathbb{K}_{\wp^2} = F$ , by [1, Thm. 1]. Similarly,

$$\text{disc}(\tilde{m}(X)) = \frac{\pi^4}{16\xi^8} + 4\frac{\pi^6}{64\xi^8} = \frac{\pi^4(1 + \pi^2)}{16\xi^8},$$

and a root of  $\tilde{m}(x)$  also generates  $F/\Sigma$ .

Note the relations

$$\begin{aligned} X(P_1) + X(P_2) &= x_1 + x_2 - 4b = \frac{\pi^2}{4\xi^4} - \frac{4\pi^4}{16(\pi^4 - 1)} \\ &= \frac{-\pi^2 - \pi^4}{4(\pi^4 - 1)} \\ &= -\frac{\pi^2}{4(\pi^2 - 1)}; \\ X(P_3) + X(P_4) &= x_3 + x_4 - 2b = -\frac{\pi^2}{4\xi^4} - \frac{4\pi^4}{16(\pi^4 - 1)} \\ &= \frac{\pi^2 - \pi^4}{4(\pi^4 - 1)} \\ &= -\frac{\pi^2}{4(\pi^2 + 1)}. \end{aligned}$$

Setting  $\lambda = -2^7 3^5 \frac{g_2 g_3}{\Delta}$ , it follows that

$$\begin{aligned}\tau(\mathfrak{k}_1^*) + \tau(\mathfrak{k}_2^*) &= \lambda \left( X(P_1) + X(P_2) + 2 \frac{4b+1}{12} \right) \\ &= \lambda \left( \frac{-\pi^2 - \pi^4}{4(\pi^4 - 1)} + \frac{1}{6} \left( \frac{\pi^4}{4(\pi^4 - 1)} + 1 \right) \right) \\ &= -\lambda \frac{\pi^4 + 6\pi^2 + 4}{24(\pi^4 - 1)},\end{aligned}$$

and

$$\begin{aligned}\tau(\mathfrak{k}_3^*) + \tau(\mathfrak{k}_4^*) &= \lambda \left( X(P_3) + X(P_4) + 2 \frac{4b+1}{12} \right) \\ &= \lambda \left( \frac{\pi^2 - \pi^4}{4(\pi^4 - 1)} + \frac{1}{6} \left( \frac{\pi^4}{4(\pi^4 - 1)} + 1 \right) \right) \\ &= -\lambda \frac{\pi^4 - 6\pi^2 + 4}{24(\pi^4 - 1)}.\end{aligned}$$

This gives that

$$\frac{\tau(\mathfrak{k}_3^*) + \tau(\mathfrak{k}_4^*)}{\tau(\mathfrak{k}_1^*) + \tau(\mathfrak{k}_2^*)} = \frac{\pi^4 - 6\pi^2 + 4}{\pi^4 + 6\pi^2 + 4}. \quad (4.1)$$

Now I claim that the pairs  $\{\tau(\mathfrak{k}_1^*), \tau(\mathfrak{k}_2^*)\}$  and  $\{\tau(\mathfrak{k}_3^*), \tau(\mathfrak{k}_4^*)\}$  are the  $\tau$ -invariants for the ideals  $\wp_2'^3$  and  $\wp_2'^3 \wp_2$ . They cannot be the invariants for proper divisors of these ideals, because each of these pairs generates  $K_{\wp_2'^3}$  over  $\Sigma$ , and the invariants for  $\wp_2'^2$  and  $\wp_2'^2 \wp_2$  lie in  $\Sigma$ , since their  $\varphi$ -values divided by 2 are both 1. They cannot be the invariants for larger ideals dividing (8), such as  $\mathfrak{m}' = \wp_2'^3 \wp_2^2$ , because this ideal satisfies Sugawara's condition (1.2) and  $\varphi(\wp_2'^3 \wp_2^2) = 8 > 6$ , so that the  $\tau$ -invariant for  $\mathfrak{m}'$  generates  $K_{\mathfrak{m}'}$ , which has degree 4 over  $\Sigma$ . Furthermore, the field  $K_{\wp_2'^3}$  is disjoint from  $K_{\wp_2'^3}$  over  $\Sigma$ . Finally, these are the only two ideals dividing (8) whose corresponding conductors are  $\wp_2'^3$ , and each of these ideals has two invariants. Thus, since  $m(x)$  and  $\tilde{m}(x)$  are quadratic and irreducible over  $\Sigma$ , by the above calculations, each of the aforementioned pairs is a pair of conjugates over  $\Sigma$  and corresponds to one of the ideals  $\mathfrak{m}_1 = \wp_2'^3$  and  $\mathfrak{m}_2 = \wp_2'^3 \wp_2$ .

These considerations allow use to prove Sugawara's conjecture for the ideal  $\mathfrak{m}_1$ . Suppose the polynomial  $T_{\mathfrak{m}_1}(X, \mathfrak{k})$  equals  $T_{\mathfrak{m}_1}(X, \bar{\mathfrak{k}})$  for two ideal classes  $\mathfrak{k}, \bar{\mathfrak{k}}$ . Then there is an automorphism  $\sigma$  of  $\Sigma/K$  for which  $j(\mathfrak{k})^\sigma = j(\bar{\mathfrak{k}})$ , but  $T_{\mathfrak{m}_1}(X, \mathfrak{k})^\sigma = T_{\mathfrak{m}_1}(X, \bar{\mathfrak{k}})$ . Since  $\sigma$  fixes  $K$ , it also fixes the ideals  $\mathfrak{m}_1$  and  $\mathfrak{m}_2$ . Let  $\psi$  be an extension of  $\sigma$  to the field  $\Sigma_{\wp_2'^3}$ . Then we have either that

$$\begin{aligned}\{\tau(\mathfrak{k}_1^*), \tau(\mathfrak{k}_2^*)\}^\psi &= \{\tau(\bar{\mathfrak{k}}_1^*), \tau(\bar{\mathfrak{k}}_2^*)\}, \\ \{\tau(\mathfrak{k}_3^*), \tau(\mathfrak{k}_4^*)\}^\psi &= \{\tau(\bar{\mathfrak{k}}_3^*), \tau(\bar{\mathfrak{k}}_4^*)\}\end{aligned}$$

or

$$\begin{aligned}\{\tau(\mathfrak{k}_1^*), \tau(\mathfrak{k}_2^*)\}^\psi &= \{\tau(\bar{\mathfrak{k}}_3^*), \tau(\bar{\mathfrak{k}}_4^*)\}, \\ \{\tau(\mathfrak{k}_3^*), \tau(\mathfrak{k}_4^*)\}^\psi &= \{\tau(\bar{\mathfrak{k}}_1^*), \tau(\bar{\mathfrak{k}}_2^*)\}.\end{aligned}$$

Hence, denoting the conjugate of  $\pi$  corresponding to  $j(\bar{\mathfrak{k}})$  by  $\bar{\pi} = \pi^\sigma$ , (8) yields two possibilities:

$$\frac{\pi^4 - 6\pi^2 + 4}{\pi^4 + 6\pi^2 + 4} = \frac{\bar{\pi}^4 - 6\bar{\pi}^2 + 4}{\bar{\pi}^4 + 6\bar{\pi}^2 + 4} \quad (4.2)$$

or

$$\frac{\pi^4 - 6\pi^2 + 4}{\pi^4 + 6\pi^2 + 4} = \frac{\bar{\pi}^4 + 6\bar{\pi}^2 + 4}{\bar{\pi}^4 - 6\bar{\pi}^2 + 4}. \quad (4.3)$$

The possibility (4.2) implies that

$$\frac{\pi^2 + \frac{4}{\pi^2} - 6}{\pi^2 + \frac{4}{\pi^2} + 6} = \frac{\bar{\pi}^2 + \frac{4}{\bar{\pi}^2} - 6}{\bar{\pi}^2 + \frac{4}{\bar{\pi}^2} + 6}.$$

Since both sides of this equation are linear fractional, this gives that

$$\pi^2 + \frac{4}{\pi^2} = \bar{\pi}^2 + \frac{4}{\bar{\pi}^2}. \quad (4.4)$$

However,

$$x^2 + \frac{4}{x^2} - y^2 - \frac{4}{y^2} = \frac{(xy - 2)(xy + 2)(-y + x)(x + y)}{x^2y^2},$$

and (4.4) implies that  $\bar{\pi}$  equals one of  $\pi, -\pi, 2/\pi$  or  $-2/\pi$ . The last two are impossible, since  $\bar{\pi} \cong \wp_2$  and  $2/\pi \cong \wp'_2$ . if  $\bar{\pi} = -\pi$ , then  $\pi$  and  $-\pi$  would be conjugates and  $b_d(x)$ , the minimal polynomial of  $\pi$ , would satisfy  $b_d(-x) = b_d(x)$  and be a polynomial in  $x^2$ . But then  $\pi^2$  would have degree less than the degree of  $\pi$  over  $\mathbb{Q}$ , contradicting the fact that  $\mathbb{Q}(\pi^2) = \mathbb{Q}(\pi^4) = \Sigma$ . (See [14, Thm. 8.1].) Hence, (4.2) implies that  $\bar{\pi} = \pi$ .

To show (4.3) cannot happen, we note the identity

$$\frac{x^2 - 6x + 4}{x^2 + 6x + 4} - \frac{y^2 + 6y + 4}{y^2 - 6y + 4} = -12 \frac{(x + y)(xy + 4)}{(x^2 + 6x + 4)(y^2 - 6y + 4)}.$$

Thus, (4.3) would imply that  $\bar{\pi}^2 = -\pi^2$  or  $-4/\pi^2$ , and both are impossible because  $i = \sqrt{-1} \notin \Sigma$ .

Now,  $\bar{\pi} = \pi$  implies that  $\bar{\beta} = 2\bar{\xi} = \pm 2\xi = \pm\beta$ ; and the fact that the  $j$ -invariant  $j(E_4)$  is a rational function in  $\beta^2$  (see Section 3) shows that  $j(\bar{\mathfrak{k}}) = j(\mathfrak{k})$ . This proves that  $T(X, \mathfrak{k}) = T(X, \bar{\mathfrak{k}})$  can only happen if  $j(\mathfrak{k}) = j(\bar{\mathfrak{k}})$  and therefore  $\mathfrak{k} = \bar{\mathfrak{k}}$ . Therefore, Sugawara's conjecture holds for the ideals  $\mathfrak{m} = \wp_2^3$  and (by complex conjugation)  $\wp_2^3$ .

With this, we have proved Sugawara's conjecture for all four possibilities in the first line of (1.4). Note that if  $(2) = \wp_2^2$ , then  $\varphi(\wp_2^2) = 2 = \varphi(\wp_2^3)/2$ , so



$\mathfrak{m} = \wp_2^3$  is not the conductor of  $K_{\mathfrak{m}} = K_{\wp_2^2}$ . Hence, we do not have to consider this case when 2 is ramified.

**Remark.** We note the cross-ratio

$$\begin{aligned} \frac{(x_1 - x_3)(x_2 - x_4)}{(x_1 - x_4)(x_2 - x_3)} &= \frac{\xi^6 + 2\pi^4 + \xi^4 - \xi^2 - 1}{-\xi^6 + 2\pi^4 + \xi^4 + \xi^2 - 1} \\ &= \frac{\xi^6 + \pi^4 - \xi^2}{-\xi^6 + \pi^4 + \xi^2} \\ &= \frac{\xi^6 + 1 - \xi^4 - \xi^2}{-\xi^6 + 1 - \xi^4 + \xi^2} \\ &= \frac{(\xi^2 + 1)(\xi - 1)^2(\xi + 1)^2}{-(\xi - 1)(\xi + 1)(\xi^2 + 1)^2} = \frac{1 - \xi^2}{1 + \xi^2}. \end{aligned}$$

Since the  $\tau$ -invariants corresponding to the points  $P_i$  are

$$\begin{aligned} \tau(\mathfrak{k}_i^*) &= -2^7 3^5 \frac{g_2 g_3}{\Delta} \left( X(P_i) + \frac{4b+1}{12} \right) \\ &= -2^7 3^5 \frac{g_2 g_3}{\Delta} \left( x_i - 2b + \frac{4b+1}{12} \right), \end{aligned}$$

the cross-ratio of the  $x_i$  equals the cross-ratio of the invariants  $\tau(\mathfrak{k}_i^*)$ .

## 5 The case $\mathfrak{m} = (3)$ .

For the next three sections we work on the Deuring normal form of an elliptic curve:

$$E_3 : Y^2 + \alpha XY + Y = X^3.$$

The points  $P_1 = (0, 0)$  and

$$P_2 = \left( \frac{-3\beta}{\alpha(\beta-3)}, \frac{\beta-3\omega^i}{\beta-3} \right), \quad (5.1)$$

where  $27\alpha^3 + 27\beta^3 = \alpha^3\beta^3$  and  $\omega = \frac{-1+\sqrt{-3}}{2}$ , are points of order 3 on  $E_3$ . For any discriminant  $d_K \equiv 1 \pmod{3}$ , there are  $\alpha, \beta$  which generate (separately) the Hilbert class field  $\Sigma$  over  $\mathbb{Q}$ , and for which  $E_3$  has complex multiplication by  $R_K$ . See [17]. Let  $(3) = \wp_3 \wp_3' = \mathfrak{p}_1 \mathfrak{p}_2$  in  $R_K$ . In addition  $(\alpha, 3) = \wp_3'$  and  $(\beta, 3) = \wp_3$ .

The Weierstrass normal form of the curve  $E_3$  is

$$E' : Y^2 = 4X^3 - g_2 X - g_3,$$

where

$$g_2 = \frac{1}{12}(\alpha^4 - 24\alpha), \quad g_3 = \frac{-1}{216}(\alpha^6 - 36\alpha^3 + 216);$$

and  $\Delta = \alpha^3 - 27$ . Thus,

$$j(E_3) = \frac{\alpha^3(\alpha^3 - 24)^3}{\alpha^3 - 27} = \frac{\beta^3(\beta^3 + 216)}{(\beta^3 - 27)^3} = j(\mathfrak{k}),$$

for some ideal class  $\mathfrak{k}$ , where the expression in  $\beta$  is obtained by using  $\alpha^3 = \frac{27\beta^3}{\beta^3 - 27}$ . The ray class invariant

$$\tau(\mathfrak{k}^*) = -2^7 3^5 \frac{g_2 g_3}{\Delta} \left( X(P) + \frac{\alpha^2}{12} \right)$$

is the invariant for a suitable ray class  $\mathfrak{k}^*$  for the modulus  $\mathfrak{m} = \mathfrak{p}_1, \mathfrak{p}_2$ , or  $(3) = \mathfrak{p}_1 \mathfrak{p}_2$  in  $K$ . Let

$$\begin{aligned} \tau_1 &= -2^7 3^5 \frac{g_2 g_3}{\Delta} \left( 0 + \frac{\alpha^2}{12} \right) \\ &= \frac{\alpha^3(\alpha^3 - 24)(\alpha^6 - 36\alpha^3 + 216)}{\alpha^3 - 27}, \\ \tau_2 &= \tau(\mathfrak{k}^*) = -2^7 3^5 \frac{g_2 g_3}{\Delta} \left( \frac{-3\beta}{\alpha(\beta - 3)} + \frac{\alpha^2}{12} \right) \\ &= \frac{(\alpha^3 - 24)(\alpha^6 - 36\alpha^3 + 216)}{\alpha^3 - 27} \frac{\alpha^3(\beta - 3) - 36\beta}{(\beta - 3)} \end{aligned}$$

be the  $\tau$ -invariants for the points  $P_1$  and  $P_2$  in (5.1). These two invariants clearly lie in  $\Sigma$ , so they are the invariants for the ideals  $\mathfrak{p}_1, \mathfrak{p}_2$ . Replacing  $\beta$  by  $\omega\beta$ , respectively  $\omega^2\beta$  results in replacing the point  $P$  by the points

$$\begin{aligned} P_3 &= \left( \frac{-3\omega\beta}{\alpha(\omega\beta - 3)}, \frac{\omega\beta - 3\omega^i}{\omega\beta - 3} \right), \\ P_4 &= \left( \frac{-3\omega^2\beta}{\alpha(\omega^2\beta - 3)}, \frac{\omega^2\beta - 3\omega^i}{\omega^2\beta - 3} \right); \end{aligned}$$

which also lie in  $E_3[3]$ . The corresponding  $\tau$ -invariants are

$$\begin{aligned} \tau_3 &= \frac{(\alpha^3 - 24)(\alpha^6 - 36\alpha^3 + 216)}{\alpha^3 - 27} \frac{\alpha^3(\omega\beta - 3) - 36\omega\beta}{(\omega\beta - 3)}, \\ \tau_4 &= \frac{(\alpha^3 - 24)(\alpha^6 - 36\alpha^3 + 216)}{\alpha^3 - 27} \frac{\alpha^3(\omega^2\beta - 3) - 36\omega^2\beta}{(\omega^2\beta - 3)}, \end{aligned}$$

which lie in  $\mathbf{K}_3 = \Sigma(\omega)$  and are conjugate over  $\Sigma$ . These are the invariants for  $\mathfrak{m} = (3)$ .

A computation shows that

$$\frac{\tau_2 - \tau_1}{\tau_1} = \frac{12}{\alpha^2} X(P_2) = \frac{-36\beta}{\alpha^3(\beta - 3)}.$$

Replacing  $\beta$  by  $\omega\beta$  and  $\omega^2\beta$  yields

$$\begin{aligned}\frac{\tau_3 - \tau_1}{\tau_1} &= \frac{12}{\alpha^2} X(P_3) = \frac{-36\beta}{\alpha^3(\beta - 3\omega^2)} \\ \frac{\tau_4 - \tau_1}{\tau_1} &= \frac{12}{\alpha^2} X(P_4) = \frac{-36\beta}{\alpha^3(\beta - 3\omega)}.\end{aligned}$$

Taking quotients yields that

$$\frac{\tau_2 - \tau_1}{\tau_3 - \tau_1} = \frac{\beta - 3\omega^2}{\beta - 3}, \quad \frac{\tau_2 - \tau_1}{\tau_4 - \tau_1} = \frac{\beta - 3\omega}{\beta - 3}.$$

These are the  $Y$ -coordinates of the points  $P_2, -P_2$ . Replacing  $\beta$  again by  $\omega^i\beta$  for  $i = 1, 2$  shows that the  $Y$ -coordinates of all points in  $E_3[3]$  are contained in  $\mathbb{K} = \mathbb{Q}(\tau_1, \tau_2, \tau_3, \tau_4)$ .

Multiplying the two  $Y$  coordinates above shows that

$$\frac{\beta - 3\omega^2}{\beta - 3} \frac{\beta - 3\omega}{\beta - 3} = \frac{\beta^2 + 3\beta + 9}{(\beta - 3)^2} \in \mathbb{K}.$$

Now, by the equation for  $E_3$ , we have for  $P_2 = (x, y)$  that  $y^2 + y = x^3 - \alpha xy$ . However, using  $\alpha^3 = \frac{27\beta^3}{\beta^3 - 27}$  yields by (5.1) that

$$x^3 = \frac{-27\beta^3}{\alpha^3(\beta - 3)^3} = -\frac{\beta^2 + 3\beta + 9}{(\beta - 3)^2} \in \mathbb{K}.$$

This implies by the equation for  $E_3$  that  $\alpha x = \frac{-3\beta}{\beta - 3} \in \mathbb{K}$ . This gives, finally, that  $\beta \in \mathbb{K}$ , and the above formulas imply that  $j(\mathfrak{f}), \omega \in \mathbb{K}$ , as well. Hence  $\mathbb{K} = \mathbb{K}_3$ .

**Theorem 5.** *The ray class field  $\mathbb{K}_3$  over  $K = \mathbb{Q}(\sqrt{d_K})$ , with  $d_K \equiv 1 \pmod{3}$ , is generated over  $\mathbb{Q}$  by the ray class invariants  $\tau(\mathfrak{f}^*)$  for the divisors  $\mathfrak{m} \neq 1$  of (3) corresponding to any single absolute ideal class  $\mathfrak{f}$ .*

We also have

$$\frac{\tau_2}{\tau_1} + \frac{\tau_3}{\tau_1} + \frac{\tau_4}{\tau_1} = \frac{-1}{3\beta^2} ((\beta + 6)^2 + (\beta + 6\omega^2)^2 + (\beta + 6\omega)^2) = -1;$$

hence,

$$\tau_2 + \tau_3 + \tau_4 = -\tau_1. \tag{5.2}$$

This shows that  $\mathbb{K}_3 = K(\tau_2, \tau_3, \tau_4)$ . It remains to show that  $\tau_2 \in K(\tau_3, \tau_4)$  and that  $\Sigma = K(\tau_1) = K(\tau_2)$ .

Computing the other elementary functions of the  $\tau_i$  on Maple yields:

$$\begin{aligned}\sum_{m \neq n} \tau_m \tau_n &= -6j(\mathfrak{f})(j(\mathfrak{f}) - 1728), \\ \sum_{m \neq n \neq l} \tau_m \tau_n \tau_l &= -8j(\mathfrak{f})(j(\mathfrak{f}) - 1728)^2, \\ \prod_{i=1}^4 \tau_i &= -3j(\mathfrak{f})^2(j(\mathfrak{f}) - 1728)^2.\end{aligned}$$

Hence the polynomial satisfied by the  $\tau_i$  is

$$F(X, j(\mathfrak{k})) = X^4 - 6j(\mathfrak{k})(j(\mathfrak{k}) - 1728)X^2 + 8j(\mathfrak{k})(j(\mathfrak{k}) - 1728)^2X - 3j(\mathfrak{k})^2(j(\mathfrak{k}) - 1728)^2,$$

which gives the relation

$$\frac{3}{8} \sum_{i=1}^4 \frac{1}{\tau_i} = \frac{1}{j(\mathfrak{k})}. \quad (5.3)$$

Using this relation we see that if the  $\tau_i$  ( $1 \leq i \leq 4$ ) are the same for two different ideal classes  $\mathfrak{k}, \mathfrak{k}'$ , then  $j(\mathfrak{k}) = j(\mathfrak{k}')$ , so that  $\mathfrak{k} = \mathfrak{k}'$ .

Note the cross ratio

$$(\tau_4, \tau_3; \tau_2, \tau_1) = \frac{(\tau_4 - \tau_2)(\tau_3 - \tau_1)}{(\tau_3 - \tau_2)(\tau_4 - \tau_1)} = -\omega. \quad (5.4)$$

From above we have

$$\frac{\tau_4 - \tau_1}{\tau_3 - \tau_1} = \frac{\beta - 3\omega^2}{\beta - 3\omega}.$$

Using the fact that  $\sigma_1 : \beta \rightarrow \frac{3(\beta+6)}{\beta-3}$  is an automorphism of  $\Sigma/\mathbb{Q}$  fixing  $\omega$  and interchanging the pairs  $(\tau_1, \tau_2), (\tau_3, \tau_4)$ , we obtain

$$\frac{\tau_3 - \tau_2}{\tau_4 - \tau_2} = \frac{\sigma_1(\beta) - 3\omega^2}{\sigma_1(\beta) - 3\omega} = -\omega^2 \frac{\beta - 3\omega}{\beta - 3\omega^2};$$

from which the above cross-ratio follows. The cross-ratio and (5.2) then yield the following equation satisfied by  $x = \tau_2$ :

$$\begin{aligned} 0 &= (2\tau_3 + x + \tau_4)(\tau_4 - x) + \omega(2\tau_4 + x + \tau_3)(\tau_3 - x) \\ &= \omega^2 x^2 - 2(\tau_3 + \omega\tau_4)x + \omega(\tau_3 - \omega\tau_4)^2. \end{aligned}$$

Multiplying by  $\omega$  gives the equation

$$x^2 - 2(\omega\tau_3 + \omega^2\tau_4)x + (\omega\tau_3 - \omega^2\tau_4)^2 = 0, \quad (5.5)$$

whose discriminant is  $16\tau_3\tau_4$ . Multiplying this equation by the equation obtained by replacing  $\omega$  by  $\omega^2$  gives the following quartic equation:

$$\begin{aligned} G(x; \tau_3, \tau_4) &= x^4 + 2(\tau_3 + \tau_4)x^3 + (3\tau_3^2 - 8\tau_3\tau_4 + 3\tau_4^2)x^2 \\ &\quad + 2(\tau_3 + \tau_4)(\tau_3^2 - 5\tau_3\tau_4 + \tau_4^2)x + (\tau_3^2 + \tau_3\tau_4 + \tau_4^2)^2. \end{aligned}$$

The second root of the quadratic equation (16) is  $\tilde{\tau} = \left(\frac{\beta-6}{\beta}\right)^2 \tau_1$ , since

$$\tau_2 + \tilde{\tau} = 2(\omega\tau_3 + \omega^2\tau_4). \quad (5.6)$$

*Proof of Sugawara's conjecture for  $\mathfrak{m} = (3) = \mathfrak{p}_1\mathfrak{p}_2$ .*

Now suppose  $\mathfrak{k}$  and  $\mathfrak{k}'$  are two ideal classes for which  $\{\tau_3, \tau_4\} = \{\tau'_3, \tau'_4\}$ . Then the automorphism  $\psi : j(\mathfrak{k}) \rightarrow j(\mathfrak{k}')$  over  $K$  satisfies  $\psi(\beta) = \beta'$ , for some

root  $\beta'$  of  $p_{d_K}(x)$ , the minimal polynomial of  $\beta$  over  $\mathbb{Q}$ . (See [17].) We may extend this automorphism to  $\mathbb{K}_3 = \Sigma(\omega)$  by fixing  $\omega$ . Then  $\tau'_3 = \tau_3^\psi, \tau'_4 = \tau_4^\psi$ .

Assume first that  $\tau'_3 = \tau_3$  and  $\tau'_4 = \tau_4$ . Equation (5.5) shows that the roots  $\tau'_2, \left(\frac{\beta'-6}{\beta'}\right)^2 \tau'_1$  have to coincide with the roots  $\tau_2, \left(\frac{\beta-6}{\beta}\right)^2 \tau_1$ . If  $\tau'_2 = \tau_2$ , then  $\tau'_1 = \tau_1$  by the relation  $\tau_2 + \tau_1 = -(\tau_3 + \tau_4) = \tau'_2 + \tau'_1$  (or using the cross-ratio). In that case  $j(\mathfrak{k}) = j(\mathfrak{k}')$  by (5.3), so  $\mathfrak{k} = \mathfrak{k}'$ . Assume instead that

$$\begin{aligned}\tau'_2 &= \left(\frac{\beta-6}{\beta}\right)^2 \tau_1, \\ \tau_2 &= \left(\frac{\beta'-6}{\beta'}\right)^2 \tau'_1, \text{ or} \\ \tau'_1 &= \left(\frac{\beta'}{\beta'-6}\right)^2 \tau_2.\end{aligned}$$

Since  $\psi$  is an automorphism, the cross-ratio (5.4) implies that

$$(\tau_4, \tau_3; \tau'_2, \tau'_1) = \frac{(\tau_4 - \tau'_2)(\tau_3 - \tau'_1)}{(\tau_3 - \tau'_2)(\tau_4 - \tau'_1)} = -\omega.$$

However, replacing  $\tau'_2$  and  $\tau'_1$  by the above expressions in terms of  $\tau_1$  and  $\tau_2$  gives

$$(\tau_4, \tau_3; \tau'_2, \tau'_1) + \omega = \frac{-\omega^2(\beta\beta' - 3\beta - 3\beta' + 36)(3\beta' + \beta - 12)(\beta - 3)}{(\beta\beta' - 3\omega^2\beta' - 3\beta - 18\omega)(\omega\beta' - \beta - \beta' - 6\omega)(3\omega - \beta + 9)}.$$

Setting the numerator in this expression equal to zero and solving for  $\beta'$  yields that

$$\beta' = \frac{-\beta}{3} + 4, \text{ or } \beta' = \frac{3(\beta - 12)}{\beta - 3}.$$

The first relation is impossible, since  $\beta' = \beta^\psi$  is a conjugate of  $\beta$ , but  $\frac{-1}{3}\beta$  is not an algebraic integer, using that  $(\beta, 3) = \wp_3 = \mathfrak{p}_1$  from [17, Lemma 2.3, Prop. 3.2]. Thus we must have

$$\beta^\psi = \psi(\beta) = \frac{3(\beta - 12)}{\beta - 3}.$$

But we know that  $\beta^{\sigma_1} = \frac{3(\beta+6)}{\beta-3}$ , and thus

$$\beta^{\sigma_1\psi} = \psi \circ \sigma_1(\beta) = 6 - \beta.$$

Now  $\psi \in \text{Gal}(\mathbb{K}_3/K)$ , by assumption, and  $\sigma_1 \notin \text{Gal}(\mathbb{K}_3/K)$ , since  $\sigma_1$  switches the ideals  $\wp_3 = \mathfrak{p}_1$  and  $\wp'_3 = \mathfrak{p}_2$ . For this, note that  $\sigma_1(\beta) = 3 + \frac{27}{\beta-3}$  and  $\beta = 3 + \gamma^3$ , where  $(\gamma) = \wp_3$  in  $\Sigma$ . (Use [17, Thm 3.4(i), p. 868] and the automorphism  $\phi = \sigma_1 \circ (\Sigma/K, \wp_2)$  which switches  $\alpha$  and  $\beta$  from [17, Prop. 3.2, p.865].) It follows that  $\psi\sigma_1 = \sigma_1\psi \notin \text{Gal}(\mathbb{K}_3/K)$ , either. On the other hand

$$\beta^{\sigma_1\psi} - 3 = 3 - \beta \cong \wp_3^3 \cong \beta - 3.$$

This would show that  $\psi\sigma_1$  does *not* switch  $\wp_3$  and  $\wp'_3$ , giving a contradiction. Hence, this case does not occur and we have the desired conclusion  $\mathfrak{k} = \mathfrak{k}'$ .

Now assume that  $\tau'_3 = \tau_3^\psi = \tau_4$  and  $\tau'_4 = \tau_3$ . Then we have

$$(\tau_3, \tau_4; \tau'_2, \tau'_1) = \frac{(\tau_3 - \tau'_2)(\tau_4 - \tau'_1)}{(\tau_4 - \tau'_2)(\tau_3 - \tau'_1)} = -\omega,$$

which implies that

$$(\tau_4, \tau_3; \tau'_2, \tau'_1) = -\omega^2. \quad (5.7)$$

This leads to the conjugate equation of (5.5), which is

$$x^2 - 2(\omega^2\tau_3 + \omega\tau_4)x + (\omega^2\tau_3 - \omega\tau_4)^2 = 0.$$

This equation also arises from (5.5) by applying the automorphism  $\sigma_1$ , which switches  $\tau_3$  and  $\tau_4$ . Therefore, its roots are  $\tau_2^{\sigma_1} = \tau_1$  and

$$\left(\frac{\beta-6}{\beta}\right)^{2\sigma_1} \tau_1^{\sigma_1} = \left(\frac{\beta-12}{\beta+6}\right)^2 \tau_2.$$

Once again we have two cases, according as  $\tau'_1 = \tau_1$  or  $\tau'_1 = \left(\frac{\beta-12}{\beta+6}\right)^2 \tau_2$ . The first case implies as before that  $\tau'_2 = \tau_2$ , so that  $j(\mathfrak{k}) = j(\mathfrak{k}')$  from (5.3). Otherwise we have

$$\tau'_1 = \left(\frac{\beta-12}{\beta+6}\right)^2 \tau_2, \quad \tau'_2 = \left(\frac{\beta'+6}{\beta'-12}\right)^2 \tau_1.$$

From (5.7) we find that the numerator of  $(\tau_4, \tau_3; \tau'_2, \tau'_1) + \omega^2 = 0$  is

$$\nu = (2\omega+1)[(\beta'^2+3\beta'+63)\beta^3-9(\beta'^2-6\beta'+90)\beta^2+27(\beta'-12)^2\beta-27(\beta'-12)^2].$$

It follows from  $\nu = 0$  and (9) =  $\wp_3^2\wp_3'^2$  that  $\wp_3'^2 \mid (\beta'^2+3\beta'+63)\beta^3$ . However,  $\wp_3' \nmid \beta$ ; and

$$\beta'^2+3\beta'+63 = (\beta'-3)^2+9(\beta'-3)+81$$

is also not divisible by  $\wp_3'$ , since  $\beta'-3 = \gamma'^3 \cong \wp_3^3$ . This contradiction shows that this case cannot occur.

This proves the following.

**Theorem 6.** *For the field  $K = \mathbb{Q}(\sqrt{d_K})$ , with  $d_K \equiv 1 \pmod{3}$ , and  $\mathfrak{m} = (3) = \wp_3\wp_3'$ , the ray class invariants corresponding to different ideal classes  $\mathfrak{k}$  are distinct. Hence, the ray class field  $K_{\mathfrak{m}} = K(\tau(\mathfrak{k}^*))$  is generated over  $K$  by a single ray class invariant  $\tau(\mathfrak{k}^*)$  ( $= \tau_3$  or  $\tau_4$ ) for the conductor  $\mathfrak{m}$ .*

If  $\mathfrak{m} = (3) = \mathfrak{p}^2$ , then there are  $\frac{\varphi(\mathfrak{m})}{2} = 3$  ray class invariants for each ideal class, so the argument is the same as in the case  $\mathfrak{m} = \mathfrak{q}^2 = (2)$  in Section 2. If the invariants for  $\mathfrak{k}, \mathfrak{k}'$  are the same, then by (5.2), the single ray class invariants for these ideal classes and  $\mathfrak{m} = \mathfrak{p}$  are also equal, and then (5.3) shows that  $j(\mathfrak{k}) = j(\mathfrak{k}')$ . Thus, Sugawara's conjecture holds for  $\mathfrak{m} = (3) = \mathfrak{p}^2$ . In this case,

note that the conditions  $\alpha, \beta \in \Sigma$  no longer obviously apply, but the algebraic formulas, including (5.2), remain valid.

Note that if  $3 \cong \mathfrak{p}$ , then  $N(\mathfrak{p}) = 9$  implies that Sugawara's second argument (1.3) applies. This also follows immediately from (5.3), since the 4 ray class invariants are conjugate over  $K_1$  in this case.

This raises the question whether  $\tau_1$  and  $\tau_2$  individually generate  $K_1$  over  $K$ , when  $\mathfrak{m} = \mathfrak{p}_1$  or  $\mathfrak{p}_2$  and  $(3) = \mathfrak{p}_1\mathfrak{p}_2$ . In the case of  $\tau_1$ , we have

$$\tau_1 = \frac{\alpha^3(\alpha^3 - 24)(\alpha^6 - 36\alpha^3 + 216)}{\alpha^3 - 27} = g(\alpha^3);$$

$$g(x) = \frac{x(x - 24)(x^2 - 36x + 216)}{x - 27}.$$

Now factor the difference  $g(x) - g(y)$  for  $x = \alpha^3, y = \alpha'^3$ :

$$(x-27)(y-27)(g(x)-g(y)) = (x-y)(x^3y+x^2y^2+xy^3+3h(x,y)), \quad h(x,y) \in \mathbb{Z}[x,y].$$

If  $x \neq y$ , then the cofactor in this equation is 0, which implies

$$xy(x^2 + xy + y^2) \equiv 0 \pmod{\mathfrak{p}_1}.$$

Since  $\mathfrak{p}_1 = \wp_3 \nmid xy$ , this gives that

$$x^3 - y^3 = (x - y)(x^2 + xy + y^2) \equiv 0 \pmod{\mathfrak{p}_1}.$$

Hence  $\alpha^9 \equiv \alpha'^9 \pmod{\mathfrak{p}_1}$ . We know that

$$j(\mathfrak{k}^*) = \frac{\alpha^3(\alpha^3 - 24)^3}{\alpha^3 - 27} \equiv \alpha^9 \pmod{\mathfrak{p}_1},$$

which implies

$$j(\mathfrak{k}^*) \equiv \alpha^9 \equiv \alpha'^9 \equiv j(\mathfrak{k}'^*) \pmod{\mathfrak{p}_1}.$$

But  $j(\mathfrak{k}^*)$  and  $j(\mathfrak{k}'^*)$  are roots of the class equation  $H_{-d}(X)$ , whose discriminant is not divisible by 3. It follows from the last congruence that  $j(\mathfrak{k}^*) = j(\mathfrak{k}'^*)$  and  $\mathfrak{k}^* = \mathfrak{k}'^*$ . If  $x = y$ , then  $\alpha^3 = \alpha'^3$  immediately gives  $j(\mathfrak{k}^*) = j(\mathfrak{k}'^*)$  and the same conclusion.

Now the automorphism  $\sigma_1$  of  $K_1/\mathbb{Q}$  satisfies  $\tau_1^{\sigma_1} = \tau_2$ , so  $K_1 = K(\tau_1) = K(\tau_2)$ .

**Theorem 7.** *If  $d_K \equiv 1 \pmod{3}$  and  $(3) = \mathfrak{p}_1\mathfrak{p}_2$ , then the  $\tau$ -invariants for both  $\mathfrak{p}_1$  and  $\mathfrak{p}_2$  are distinct, so that the Hilbert class field  $K_1$  of  $K$  is generated by a single  $\tau$ -invariant for either  $\mathfrak{p}_1$  or  $\mathfrak{p}_2$ .*

## 6 The case $m \mid 9$ .

The Tate normal form for a point of order 9 is

$$E_9(t) : Y^2 + (1+t^2-t^3)XY + (1-t)(1-t+t^2)t^2Y = X^3 + (1-t)(1-t+t^2)t^2X^2. \quad (6.1)$$

Its  $j$ -invariant is

$$j(E_9) = \frac{(t^3 - 3t^2 + 1)^3(t^9 - 9t^8 + 27t^7 - 48t^6 + 54t^5 - 45t^4 + 27t^3 - 9t^2 + 1)^3}{t^9(t-1)^9(t^2-t+1)^3(t^3-6t^2+3t+1)}. \quad (6.2)$$

This can be verified using the polynomial

$$f_9(a, b) = a^5 - 6a^4 + a^3b + 15a^3 - 6a^2b + 3ab^2 - b^3 - 19a^2 + 9ab - 3b^2 + 12a - 4b - 3,$$

since  $f_9(a, b) = 0$  is the condition that  $P = (0, 0)$  represents a point of order 9 on the curve

$$Y^2 + aXY + bY = X^3 + bX^2.$$

(See [15, pp. 248-250]. Note that the formula for  $j(E_n)$  should have a minus sign.) The curve  $f_9(a, b) = 0$  has genus 0 and is parametrized by

$$a = 1 + t^2 - t^3, \quad b = (1-t)(1-t+t^2)t^2.$$

Now let

$$g(t) = \frac{t^3 - 3t^2 + 1}{t(t-1)}.$$

Noting that

$$t^9 - 9t^8 + 27t^7 - 48t^6 + 54t^5 - 45t^4 + 27t^3 - 9t^2 + 1 = (t^3 - 3t^2 + 1)^3 - 24t^3(t-1)^3$$

and

$$\begin{aligned} (t^2 - t + 1)^3 &= (t^3 - 3t^2 + 1)^2 + 3(t^3 - 3t^2 + 1)t(t-1) + 9t^2(t-1)^2, \\ t^3 - 6t^2 + 3t + 1 &= t^3 - 3t^2 + 1 - 3(t^2 - t), \end{aligned}$$

it follows that

$$j(E_9(t)) = \frac{g(t)^3(g(t)^3 - 24)^3}{g(t)^3 - 27}. \quad (6.3)$$

This implies that  $E_9$  is isomorphic to the Deuring normal form

$$E_3(\alpha) : Y^2 + \alpha XY + Y = X^3, \quad \alpha = g(t),$$

whose  $j$ -invariant is

$$j(E_3) = \frac{\alpha^3(\alpha^3 - 24)^3}{\alpha^3 - 27}.$$



Now by Proposition 3.6(ii) of [16] and the remark thereafter, a point  $P = (\xi, \eta)$  on  $E_3(\alpha)$  satisfies  $3P = \pm(0, 0)$  whenever its  $X$ -coordinate satisfies  $x^3 - (3 + \alpha)x^2 + \alpha x + 1 = 0$ . But this equation implies the relation

$$\alpha = \frac{\xi^3 - 3\xi^2 + 1}{\xi(\xi - 1)} = g(\xi).$$

Hence, the point  $P = (\xi, \eta)$  is a point of order 9 on  $E_3(\alpha)$ .

We have the discriminant formula

$$\text{disc}(x^3 - (3 + \alpha)x^2 + \alpha x + 1) = (\alpha^2 + 3\alpha + 9)^2.$$

In order to decide whether Sugawara's conjecture is true in the case  $(3) = \wp_3 \wp_3'$  and  $\mathfrak{m} = \wp_3'^2$ , we let  $\alpha \in \Sigma$  be as in Section 5. Then  $\alpha - 3 \cong \wp_3'^3$  and  $(\alpha) = \wp_3' \mathfrak{a}$ , with  $(\mathfrak{a}, 3) = 1$ , by [17]. It follows that the above discriminant is relatively prime to  $\wp_3$ . Furthermore,

$$(\alpha^2 + 3\alpha + 9)(\alpha - 3) = \alpha^3 - 27 = \frac{27\alpha^3}{\beta^3} \cong \wp_3'^6 \mathfrak{c},$$

for some integral ideal  $\mathfrak{c}$  prime to  $(3)$ . In fact  $\mathfrak{c} = (1)$ , since

$$(\alpha^3 - 27)(\beta^3 - 27) = \alpha^3 \beta^3 - 27\alpha^3 - 27\beta^3 + 27^2 = 3^6.$$

Thus, it is clear that  $(\alpha^2 + 3\alpha + 9) \cong \wp_3'^3$ .

Assuming  $k(x) = x^3 - (3 + \alpha)x^2 + \alpha x + 1$  is irreducible over  $\Sigma$ , its root  $\xi$  generates a cyclic cubic extension of  $\Sigma$ ; its conjugates over  $\Sigma$  are  $\frac{1}{1-\xi}$  and  $\frac{\xi-1}{\xi}$ . Also, since  $g_2, g_3$  and  $\Delta$  for the curve  $E_3(\alpha)$  lie in  $\Sigma$ , the fact that the invariant

$$\begin{aligned} \tau(\mathfrak{k}^*) &= -2^7 3^5 \frac{g_2 g_3}{\Delta} \left( \xi + \frac{\alpha^2}{12} \right) \\ &= \frac{\alpha(\alpha^3 - 24)(\alpha^6 - 36\alpha^3 + 216)}{\alpha^3 - 27} (\alpha^2 + 12\xi) \end{aligned}$$

lies in  $\mathbb{K}_{(9)}$  implies that  $L = \Sigma(\xi) \subset \mathbb{K}_{(9)}$ . By the previous paragraph, the field  $L$  has conductor  $\mathfrak{f} = \wp_3'^2$  over  $K$ , and  $\wp_3$  is unramified in  $L/K$ . This shows that  $L = \mathbb{K}_{\wp_3'^2}$ , since  $\frac{\varphi_{\mathbb{K}}(\wp_3'^2)}{2} = 3$ . This also shows that  $L$  is the inertia field for the prime  $\wp_3$  in  $\mathbb{K}_{(9)}/K$ , since any subfield of  $\mathbb{K}_{(9)}/\Sigma$  not contained in  $L$  must have a conductor which is divisible by  $\wp_3$ . It remains to show that  $k(x)$  is irreducible over  $\Sigma$ .

This may be shown using the Newton polygon for the shifted polynomial

$$k\left(x + \frac{\alpha}{3} + 1\right) = x^3 - \frac{\alpha^2 + 3\alpha + 9}{3}x - \frac{(2\alpha + 3)(\alpha^2 + 3\alpha + 9)}{27}, \quad (6.4)$$

for a prime divisor  $\mathfrak{p}$  of  $\wp_3'$  in  $\Sigma$ . I claim that the additive valuation  $w_{\mathfrak{p}}$  of the last two coefficients is 2. For the coefficient of  $x$ , this follows from the above remarks, since

$$w_{\mathfrak{p}}\left(\frac{\alpha^2 + 3\alpha + 9}{3}\right) = 3 - 1 = 2.$$

For the constant term,

$$w_p \left( \frac{(2\alpha + 3)(\alpha^2 + 3\alpha + 9)}{27} \right) = w_p(2\alpha + 3) = w_p(9 + 2\gamma^3) = 2,$$

where  $\alpha = 3 + \gamma^3$ , with  $\gamma \cong \wp'_3$ , by results of [17]. It follows that the Newton polygon for the polynomial in (6.4) is the line segment joining the points  $(0, 2)$  and  $(3, 0)$ , since  $(1, 2)$  and  $(2, \infty)$  lie above this line segment. The slope of this segment is  $-2/3$ , which implies the irreducibility of  $k(x)$  over the completion  $\Sigma_p$ . (See [25, pp. 76-77].)

Now let

$$\begin{aligned} \tau_1 = \tau(\mathfrak{k}_1^*) &= -2^7 3^5 \frac{g_2 g_3}{\Delta} \left( \xi + \frac{\alpha^2}{12} \right) \\ &= \frac{\alpha(\alpha^3 - 24)(\alpha^6 - 36\alpha^3 + 216)(\alpha^2 + 12\xi)}{\alpha^3 - 27}, \\ \tau_2 = \tau(\mathfrak{k}_2^*) &= -2^7 3^5 \frac{g_2 g_3}{\Delta} \left( \frac{1}{1 - \xi} + \frac{\alpha^2}{12} \right) \\ &= \frac{\alpha(\alpha^3 - 24)(\alpha^6 - 36\alpha^3 + 216)(\alpha^2 \xi - \alpha^2 - 12)}{(\xi - 1)(\alpha^3 - 27)}, \\ \tau_3 = \tau(\mathfrak{k}_3^*) &= -2^7 3^5 \frac{g_2 g_3}{\Delta} \left( \frac{\xi - 1}{\xi} + \frac{\alpha^2}{12} \right) \\ &= \frac{\alpha(\alpha^3 - 24)(\alpha^6 - 36\alpha^3 + 216)(\alpha^2 \xi + 12\xi - 12)}{\xi(\alpha^3 - 27)}, \end{aligned}$$

be the three  $\tau$ -invariants corresponding to a fixed  $j$ -invariant  $j(\mathfrak{k})$  (and ideal class  $\mathfrak{k}$ ). These invariants are roots of the polynomial

$$T_m(X, \mathfrak{k}) = X^3 - c_1 X^2 + c_2 X - c_3 \in \Sigma[X],$$

where

$$\begin{aligned} c_1 &= \frac{3\alpha(\alpha^3 - 24)(\alpha^6 - 36\alpha^3 + 216)(\alpha^2 + 4\alpha + 12)}{\alpha^3 - 27}, \\ c_2 &= \frac{3\alpha^3(\alpha^3 - 24)^2(\alpha^6 - 36\alpha^3 + 216)^2(\alpha^3 + 8\alpha^2 + 24\alpha + 48)}{(\alpha^3 - 27)^2}, \\ c_3 &= \frac{\alpha^3(\alpha^3 - 24)^3(\alpha^6 - 36\alpha^3 + 216)^3(\alpha^6 + 12\alpha^5 + 36\alpha^4 + 144\alpha^3 - 1728)}{(\alpha^3 - 27)^3}. \end{aligned}$$

The  $c_i$  are the sums of the products of the  $\tau_j$  taken  $i$  at a time. The discriminant of  $T_m(X, \mathfrak{k})$  is

$$\text{disc}(T_m(X, \mathfrak{k})) = 12^6 \frac{\alpha^6(\alpha^3 - 24)^6(\alpha^6 - 36\alpha^3 + 216)^6}{(\alpha - 3)^6(\alpha^2 + 3\alpha + 9)^4}.$$

$T_{\mathfrak{m}}(X, \mathfrak{k})$  is irreducible over  $\Sigma$  by Hasse's results [9]. Since its discriminant is a square, any of its roots generates a cyclic cubic extension, and since the  $\tau_i \in K_{\wp_3'}$ , we clearly have  $\Sigma(\tau_i) = K_{\wp_3'}$ . Furthermore,

$$\frac{\tau_1 - \tau_2}{\tau_2 - \tau_3} = -\xi, \quad \frac{\tau_2 - \tau_3}{\tau_3 - \tau_1} = \frac{1}{\xi - 1}, \quad \frac{\tau_3 - \tau_1}{\tau_1 - \tau_2} = \frac{1 - \xi}{\xi}; \quad (6.5)$$

$$\frac{\tau_1 - \tau_3}{\tau_3 - \tau_2} = \xi - 1, \quad \frac{\tau_3 - \tau_2}{\tau_2 - \tau_1} = \frac{-1}{\xi}, \quad \frac{\tau_2 - \tau_1}{\tau_1 - \tau_3} = \frac{\xi}{1 - \xi}. \quad (6.6)$$

The values of these ratios are the negatives and the negative reciprocals of the roots of  $k(x)$ . Suppose that  $T_1(X) = T(X, \mathfrak{k}_1) = T(X, \mathfrak{k}_2) = T_2(X)$  for two different ideal classes  $\mathfrak{k}_1, \mathfrak{k}_2$ , corresponding to different conjugates  $\alpha_1, \alpha_2$  of  $\alpha$  over  $K$  and corresponding roots  $\xi_1, \xi_2$ . Then the negatives and negative reciprocals of the roots of  $k_1(x) = x^3 - (3 + \alpha_1)x^2 + \alpha_1x + 1$  must coincide with the corresponding expressions in the roots of  $k_2(x) = x^3 - (3 + \alpha_2)x^2 + \alpha_2x + 1$ . If, for example,  $-\xi_1 = \frac{1}{\xi_2 - 1}$ , then  $\xi_1 = \frac{1}{1 - \xi_2}$  is a conjugate of  $\xi_2$  over  $\Sigma$ , whence it follows that  $\alpha_1 = \alpha_2$ . The same holds if the ratios in (6.5) for  $T_1(X)$  coincide with a permutation of the same ratios for  $T_2(X)$ . The ratios in (6.6) are the reciprocals of the ratios in (6.5), so that a similar statement holds if the ratios in (6.6) for the polynomials  $T_i(X)$  are permutations of each other. Now suppose that  $-\xi_1 = \frac{\xi_2}{1 - \xi_2}$ . Then  $\frac{1}{\xi_1} = \frac{\xi_2 - 1}{\xi_2}$  is a conjugate of  $\xi_2$ , from which it would follow that  $x^3 k_1(1/x) = x^3 + \alpha_1 x^2 - (3 + \alpha_1)x + 1$  coincides with  $k_2(x) = x^3 - (3 + \alpha_2)x^2 + \alpha_2x + 1$ ; hence,  $\alpha_1 + \alpha_2 = -3$ . But this is impossible, since this would imply

$$0 = \alpha_1 + \alpha_2 + 3 = (3 + \gamma_1^3) + (3 + \gamma_2^3) + 3 = 9 + \gamma_1^3 + \gamma_2^3,$$

where  $\gamma_1, \gamma_2 \in \Sigma$  and  $(\gamma_1) = (\gamma_2) = \wp_3'$ , implying that  $\wp_3'^3 \mid 9$ .

This shows that the ratios in (6.5) for  $T_1(X)$  cannot coincide with the ratios in (6.6) for  $T_2(X)$ . Hence, we must have  $\alpha_1 = \alpha_2$  and therefore  $j(\mathfrak{k}_1) = j(\mathfrak{k}_2)$ . This proves that the polynomials  $T_{\mathfrak{m}}(X, \mathfrak{k})$  are distinct for different ideal classes.

**Theorem 8.** *If  $\mathfrak{m} = \mathfrak{p}_1^2$ , where  $(3) = \mathfrak{p}_1 \mathfrak{p}_2$ , then the ray class field  $K_{\mathfrak{m}} = K(\tau_i)$  is generated over  $K$  by a single  $\tau$ -invariant for the conductor  $\mathfrak{m}$ .*

Note that the case  $\mathfrak{m} = \mathfrak{p}^2 = (3)$  has been handled in Section 5, and the case of any higher power of a first degree prime divisor  $\mathfrak{p}$  of 3 is taken care of by Sugawara's condition (1.3).

## 7 The case $\mathfrak{m} = (2)\wp_3\wp_3'$ .

On the curve  $E_3(\alpha)$ , the doubling formula is

$$X(2P) = \frac{x(x^3 - \alpha x - 2)}{4x^3 + (\alpha x + 1)^2}.$$

Then  $P = (x, y)$  on  $E_3$  satisfies  $2P = \pm(0, 0)$  and  $P \neq \pm(0, 0)$  if and only if  $x^3 - \alpha x - 2 = 0$ . If this condition holds, then  $P$  has order 6 on  $E_3$ . We have

$$\alpha = \frac{x^3 - 2}{x} = \frac{(-x)^3 + 2}{-x},$$

and it follows from the result of [2, Prop. 13] that

$$x = -2c(w/3), \frac{1}{c(w/6)} \text{ or } \frac{1}{c_1(w/6)}, \quad (7.1)$$

where  $c(\tau)$  is Ramanujan's cubic continued fraction and  $c_1(\tau) = c(\tau + \frac{3}{2})$ ; and  $w/3$  is the basis quotient of a suitable integral ideal. Each of these values lies in the ring class field  $\Omega_2 = \mathbf{K}_2$ , by [2, pp. 20, 27].

Given the factorization

$$Y^2 + \alpha x Y + Y - x^3 = Y^2 + (x^3 - 1)Y - x^3 = (Y - 1)(Y + x^3) = 0$$

we set  $P = (x, 1)$  and  $-P = (x, -x^3)$ . Assuming  $2P = (0, 0)$ , this yields that

$$3P = 2P + P = (0, 0) + (x, 1) = \left( \left( \frac{1}{x} \right)^2 + \alpha \left( \frac{1}{x} \right) - x, y_1 \right) = \left( \frac{-1}{x^2}, y_1 \right).$$

The alternative would be that  $2P = (0, -1)$ , in which case

$$3P = 2P + P = (0, -1) + (x, 1) = \left( \left( \frac{2}{x} \right)^2 + \alpha \left( \frac{2}{x} \right) - x, y_2 \right) = (x, y_2);$$

and this would imply that  $3P = -P$ , which is false. Hence  $3P = \left( \frac{-1}{x^2}, \frac{-1}{x^3} \right)$  has order 2, so that

$$\bar{\tau} = -2^7 3^5 \frac{g_2 g_3}{\Delta} \left( \frac{-1}{x^2} + \frac{\alpha^2}{12} \right)$$

is the  $\tau$ -invariant corresponding to  $3P$ . By Theorem 1 and the fact that the  $\tau$ -invariants are independent of the choice of the elliptic curve, we know that  $\bar{\tau}$  generates  $\mathbf{K}_2/\Sigma$  and therefore so does the root  $x$ , since  $\alpha, g_2, g_3, \Delta$  lie in  $\Sigma$  and  $x \in \mathbf{K}_2$ . Hence,  $t(X) = X^3 - \alpha X - 2$  is irreducible over  $\Sigma$ . This also follows from [2, Thm. 1].

Now assume  $\mathfrak{m} = (6) = (2)\wp_3\wp_3'$ , where 2 is inert in  $K$ . Consider the point of order 6:

$$\begin{aligned} Q &= (x, 1) + \left( \frac{3\beta}{\alpha(\beta - 3)}, \frac{\beta - 3\omega}{\beta - 3} \right) \\ &= (X(Q), Y(Q)), \end{aligned}$$

where

$$X(Q) = \frac{3\beta[-\alpha^2(\beta^3 - 27)x^2 - 3\alpha\beta(2\beta + 3\omega^2)(\beta - 3\omega)x + 9\beta^2(-\beta + 9\omega + 3)]}{\alpha(\beta^2 + 3\beta + 9)(\alpha\beta x - 3\alpha x + 3\beta)^2}.$$

Using that the reciprocal of  $\alpha\beta x - 3\alpha x + 3\beta$  is

$$\rho = \frac{(\beta^2 + 3\beta + 9)(\beta - 3)^2\alpha^2}{729\beta^3}x^2 - \frac{(\beta^3 - 27)\alpha}{243\beta^2}x - \frac{(2\beta^2 - 12\beta - 9)}{81\beta},$$

this yields that

$$\begin{aligned} X(Q) &= \rho^2 \frac{3\beta[-\alpha^2(\beta^3 - 27)x^2 - 3\alpha\beta(2\beta + 3\omega^2)(\beta - 3\omega)x + 9\beta^2(-\beta + 9\omega + 3)]}{\alpha(\beta^2 + 3\beta + 9)} \\ &= -\frac{(\beta^2\omega + \beta^2 - 3\beta - 9\omega)\alpha}{9\beta}x^2 + \frac{(\beta\omega + \beta - 3)}{3}x \\ &\quad + \frac{\beta(2\beta^2\omega + 2\beta^2 + 6\beta\omega - 3\beta - 9\omega - 9)}{\alpha(\beta^2 + 3\beta + 9)} \\ &= \left( -\frac{(\beta^2 - 9)\alpha}{9\beta}x^2 + \frac{\beta}{3}x + \frac{\beta(2\beta^2 + 6\beta - 9)}{\alpha(\beta^2 + 3\beta + 9)} \right)\omega \\ &\quad - \frac{(\beta^2 - 3\beta)\alpha}{9\beta}x^2 + \frac{(\beta - 3)}{3}x + \frac{\beta(2\beta^2 - 3\beta - 9)}{\alpha(\beta^2 + 3\beta + 9)}. \end{aligned}$$

Since  $\{1, x, x^2\}$  is a basis of  $\mathbf{K}_2/\Sigma$  and  $\{1, \omega\}$  is a basis of  $\mathbf{K}_3/\Sigma$ , it is clear that  $\{1, x, x^2, \omega, \omega x, \omega x^2\}$  is a basis of  $\mathbf{K}_6/\Sigma$  ( $\mathbf{K}_2\mathbf{K}_3 = \mathbf{K}_6$  because the degrees match). Hence, the above representation shows that  $X(Q)$  does not lie in any of the subfields  $\mathbf{K}_2, \mathbf{K}_3$  or  $\Sigma$ . The trace of  $X(Q)$  to  $\mathbf{K}_2$  is

$$\begin{aligned} \text{Tr}_{\mathbf{K}_6/\mathbf{K}_2}(X(Q)) &= -\left( -\frac{(\beta^2 - 9)\alpha}{9\beta}x^2 + \frac{\beta}{3}x + \frac{\beta(2\beta^2 + 6\beta - 9)}{\alpha(\beta^2 + 3\beta + 9)} \right) \\ &\quad + 2\left( -\frac{(\beta^2 - 3\beta)\alpha}{9\beta}x^2 + \frac{(\beta - 3)}{3}x + \frac{\beta(2\beta^2 - 3\beta - 9)}{\alpha(\beta^2 + 3\beta + 9)} \right) \\ &= -\frac{(\beta - 3)^2\alpha}{9\beta}x^2 + \frac{(\beta - 6)}{3}x + \frac{\beta(2\beta^2 - 12\beta - 9)}{(\beta^2 + 3\beta + 9)\alpha}. \end{aligned}$$

From this expression we compute the trace of  $X(Q)$  to  $\Sigma$  to be

$$\begin{aligned} \text{Tr}_{\mathbf{K}_6/\Sigma}(X(Q)) &= -\frac{(\beta - 3)^2\alpha}{9\beta}\text{Tr}_{\mathbf{K}_2/\Sigma}(x^2) + \frac{(\beta - 6)}{3}\text{Tr}_{\mathbf{K}_2/\Sigma}(x) \\ &\quad + 3\frac{\beta(2\beta^2 - 12\beta - 9)}{(\beta^2 + 3\beta + 9)\alpha} \\ &= -\frac{(\beta - 3)^2\alpha}{9\beta}(2\alpha) + 0 + 3\frac{\beta(2\beta^2 - 12\beta - 9)}{(\beta^2 + 3\beta + 9)\alpha} \\ &= \frac{-9\beta(2\beta + 3)}{\alpha(\beta^2 + 3\beta + 9)}. \end{aligned}$$

Now we have

$$\begin{aligned}\mathrm{Tr}_{K_6/\Sigma}\left(X(Q) + \frac{\alpha^2}{12}\right) &= \frac{-9\beta(2\beta+3)}{\alpha(\beta^2+3\beta+9)} + \frac{\alpha^2}{2} \\ &= \frac{(\alpha^3\beta^2 + 3\alpha^3\beta + 9\alpha^3 - 36\beta^2 - 54\beta)}{2\alpha(\beta^2+3\beta+9)} \\ &= -\frac{9\beta(\beta^2-6\beta-18)}{2\alpha(\beta^2+3\beta+9)(\beta-3)}.\end{aligned}$$

It follows from this that

$$\begin{aligned}\mathrm{Tr}_{K_6/\Sigma}(\tau(\mathfrak{k}^*)) &= -2^7 3^5 \frac{g_2 g_3}{\Delta} \mathrm{Tr}_{K_6/\Sigma}\left(X(Q) + \frac{\alpha^2}{12}\right) \\ &= -54 \frac{(\alpha^3-24)(\alpha^6-36\alpha^3+216)\beta(\beta^2-6\beta-18)}{(\beta^3-27)(\alpha^3-27)} \\ &= \frac{6(\beta^2-6\beta-18)^2\beta(\beta^4+6\beta^3+54\beta^2-108\beta+324)(\beta^3+216)}{(\beta^3-27)^3}.\end{aligned}$$

Now assume that the ray class invariants for two ray classes  $\mathfrak{k}_1^*, \mathfrak{k}_2^*$  are the same. Since the six invariants for  $\mathfrak{m}$  are conjugate over  $\Sigma$ ,  $\tau(\mathfrak{k}_1^*)$  and  $\tau(\mathfrak{k}_2^*)$  are conjugate and their traces to  $\Sigma$  must be equal. Set

$$f(x) = \frac{(x^2-6x-18)^2x(x^4+6x^3+54x^2-108x+324)(x^3+216)}{(x^3-27)^3}.$$

Thus,  $f(x) = f(y)$ , for  $x = \beta_1, y = \beta_2$ , implies that

$$\begin{aligned}0 &= (x^3-27)^3(y^3-27)^3(f(x)-f(y)) \\ &= (-y+x)(xy-3x-3y-18)(x^{10}y^8+x^9y^9+x^8y^{10}+3q(x,y)),\end{aligned}\quad (7.2)$$

where  $q(x,y) \in \mathbb{Z}[x,y]$  and the third factor is irreducible over  $\mathbb{Z}$ . If the third factor is 0, we reduce it modulo  $\wp'_3$  and find that

$$x^{10}y^8+x^9y^9+x^8y^{10}+3q(x,y) \equiv x^8y^8(x^2+xy+y^2) \equiv x^8y^8(x-y)^2 \pmod{\wp'_3}.$$

Since the left side of this congruence is 0, we have  $\wp'_3 \mid (x-y)$  in  $\Sigma$ , since  $(xy, \wp'_3) = 1$  and  $\wp'_3$  is unramified in  $\Sigma/K$ . But this means that  $\wp'_3$  divides  $\beta_1 - \beta_2$ , which is impossible, since the discriminant of the minimal polynomial of  $\beta$  over  $K$  is not divisible by  $\wp'_3$ , by [17, p.880, Eq. (4.27)]. (Apply the automorphism switching  $\alpha$  and  $\beta$  and  $\wp_3$  and  $\wp'_3$  in that equation.) Hence, the third factor in (7.2) is not 0. The second factor cannot be 0, either, since this would imply that  $\beta_2 = \frac{3\beta_1+18}{\beta_1-3} = \alpha_1^\tau$ , by [17, Thm. 3.2, p. 865]. Hence, the first factor must be 0, which gives that  $\beta_1 = \beta_2$  and  $j(\mathfrak{k}_1) = j(\mathfrak{k}_2)$ . (See the formula for  $j(E_3)$  in Section 5.) This proves the result we want.

**Theorem 9.** *If  $\mathfrak{m} = (6) = (2)\wp_3\wp'_3$ , where 2 is inert in  $K$ , then the ray class field  $K_{\mathfrak{m}}$  is generated by a single  $\tau$ -invariant for the ideal  $\mathfrak{m}$ .*

Note that the  $\tau$ -invariant for the point  $Q$  is certainly an invariant for the ideal  $\mathfrak{m}$ , since the quantity  $X(Q)$  generates  $K_{\mathfrak{m}}$ . This proves Sugawara's conjecture for the first three cases in line 2 of (1.4). We will postpone the discussion of the fourth case to a later section. We have also now established the main results of [11] using our methods. Next we turn to ideals divisible by some  $\wp_5$ .

## 8 The case $\mathfrak{m} = \wp_5$ .

Let

$$E_5(b) : Y^2 + (1+b)XY + bY = X^3 + bX^2$$

be the Tate normal form for a point of order 5. We use the computations and results of [18]. If  $\mathfrak{m} = \wp_5'$ , where  $(5) = \wp_5 \wp_5'$  in  $K = \mathbb{Q}(\sqrt{-d})$ , then  $d \equiv \pm 1 \pmod{5}$ . If  $b = r(w/5)^5$ , where  $w$  is a suitable integer in  $R_K$  and  $r(\tau)$  is the Rogers-Ramanujan continued fraction, then  $E_5(b)$  has complex multiplication by the ring of integers  $R_K$  of  $K$ . See [18, Thm. 2.1]. By choosing  $w$  appropriately, we can arrange for  $\wp_5'$  to be either of the prime ideals dividing  $(5)$ .

The Weierstrass normal form of  $E_5(b)$  is

$$Y_1^2 = 4X_1^3 - g_2X_1 - g_3,$$

where  $X_1 = X + \frac{1}{12}(b^2 + 6b + 1)$ ,

$$\begin{aligned} g_2 &= g_2(\mathfrak{f}) = \frac{1}{12}(b^4 + 12b^3 + 14b^2 - 12b + 1), \\ g_3 &= \frac{-1}{216}(b^2 + 1)(b^4 + 18b^3 + 74b^2 - 18b + 1); \\ \Delta &= -b^5(b^2 + 11b - 1). \end{aligned}$$

Thus,

$$j(E_5) = j(\mathfrak{f}) = -\frac{(b^4 + 12b^3 + 14b^2 - 12b + 1)^3}{b^5(b^2 + 11b - 1)}.$$

Two of the  $X$ -coordinates of points in  $E_5[5]$  are  $X = 0, -b$ . Hence, we let

$$\begin{aligned} \tau_0 &= -2^7 3^5 \frac{g_2 g_3}{\Delta} \left(0 + \frac{1}{12}(b^2 + 6b + 1)\right) \\ &= -\frac{(b^4 + 12b^3 + 14b^2 - 12b + 1)(b^2 + 1)(b^4 + 18b^3 + 74b^2 - 18b + 1)(b^2 + 6b + 1)}{b^5(b^2 + 11b - 1)}; \end{aligned}$$

and

$$\begin{aligned} \tau_1 &= -2^7 3^5 \frac{g_2 g_3}{\Delta} \left(-b + \frac{1}{12}(b^2 + 6b + 1)\right) \\ &= -\frac{(b^4 + 12b^3 + 14b^2 - 12b + 1)(b^2 + 1)(b^4 + 18b^3 + 74b^2 - 18b + 1)(b^2 - 6b + 1)}{b^5(b^2 + 11b - 1)}. \end{aligned}$$

Note that the only difference between  $\tau_0$  and  $\tau_1$  is in the final factor of the numerator, so that

$$\frac{\tau_0}{\tau_1} = \frac{b^2 + 6b + 1}{b^2 - 6b + 1}.$$

From [18, Thm. 4.6] we know that the ray class field  $K_{\mathfrak{f}} = F_1$  of conductor  $\mathfrak{f} = \wp_5'$  over  $K$  is generated over the Hilbert class field  $\Sigma$  (and even over  $\mathbb{Q}$ ) by the quantity  $b = r(w/5)^5$ . Also,  $\sigma : b \rightarrow -1/b$  is the nontrivial automorphism of  $F_1/\Sigma$ . The expressions for the  $\tau_i$  imply easily that  $\tau_0^\sigma = \tau_1$ , so that  $\tau_0$  and  $\tau_1$  are quadratic over  $\Sigma$  and lie in  $F_1$ . Furthermore,  $\tau_0 \neq \tau_1$ , since

$$\frac{\tau_0}{\tau_1} - 1 = \frac{12b}{b^2 - 6b + 1}$$

and  $b \neq 0$ .

We want to show that  $K(\tau_0) = K(\tau_1) = F_1$ . We compute that

$$(\tau_0 - \tau_1)^2 = 12b^2 \frac{j(\mathfrak{k})(j(\mathfrak{k}) - 1728)}{g_2(\mathfrak{k})}. \quad (8.1)$$

and

$$\frac{\tau_0 - \tau_1}{\tau_0 + \tau_1} = \frac{6b}{b^2 + 1}. \quad (8.2)$$

It follows that if  $\{\tau_0, \tau_1\} = \{\tau'_0, \tau'_1\}$  for two different ideal classes  $\mathfrak{k}, \mathfrak{k}'$ , corresponding to the values  $b, b'$ , then

$$\frac{6b}{b^2 + 1} = \pm \frac{6b'}{(b')^2 + 1}.$$

This equation easily implies that  $b'$  is given by one of the possibilities  $b' = b, -1/b, -b, 1/b$ . If  $b' = b, -1/b$  it follows that  $j(\mathfrak{k}') = j(\mathfrak{k})$  and  $\mathfrak{k} = \mathfrak{k}'$ . It remains to eliminate the cases  $b' = -b, 1/b$ , which are equivalent, since  $j(\mathfrak{k}) = j(b)$  is invariant under  $\sigma$ .

If, without loss of generality,  $b' = -b$ , then (8.1) implies that

$$\frac{j(\mathfrak{k}')(j(\mathfrak{k}') - 1728)}{g_2(\mathfrak{k}')} = \frac{j(\mathfrak{k})(j(\mathfrak{k}) - 1728)}{g_2(\mathfrak{k})}$$

or, with easily understood notation,

$$\begin{aligned} 0 &= g_2(b)j(-b)(j(-b) - 1728) - g_2(-b)j(b)(j(b) - 1728) \\ &= -\frac{P(b)}{3b^9(b^2 + 11b - 1)^2(b^2 - 11b - 1)^2}, \end{aligned}$$

where

$$\begin{aligned} P(b) &= (b^2 + 1)^2(b^4 + 12b^3 + 14b^2 - 12b + 1)(b^4 - 12b^3 + 14b^2 + 12b + 1) \\ &\quad \times (b - 1)(b + 1)(19b^8 - 2264b^6 - 8886b^4 - 2264b^2 + 19) \\ &\quad \times (b^8 - 26b^6 - 11934b^4 - 26b^2 + 1). \end{aligned}$$



Now, the roots of the first two factors yield  $j(b) = 1728, 0$ , which are excluded, since they imply  $d_K = -4, -3$ . The roots of the remaining quartic, as well as  $b = \pm 1$ , yield values of  $j$  which are not algebraic integers, as can easily be checked. We just have to eliminate the last two factors as possibilities. Let

$$\begin{aligned} f_1(x) &= 19b^8 - 2264b^6 - 8886b^4 - 2264b^2 + 19, \\ f_2(x) &= b^8 - 26b^6 - 11934b^4 - 26b^2 + 1 = b^4 m\left(b - \frac{1}{b}\right), \\ m(x) &= x^4 - 22x^2 - 11984. \end{aligned}$$

The roots of  $f_1(x)$  are not algebraic integers, but  $b$  is a unit, so this polynomial cannot have  $b$  as a root. Furthermore, if  $f_2(x)$  occurred as the minimal polynomial of  $b$ ,  $m(x)$  would be the minimal polynomial of  $z = b - 1/b \in \Sigma$ . However, the discriminant of  $m(x)$  is divisible by  $5^2$ , and it is easily checked that  $5^2 \mid d_L$ , where  $L$  is a root field of  $m(x)$ . But this is impossible, since 5 does not ramify in  $K$  or in  $\Sigma/K$ . Thus, the roots of  $f_2(x)$  also cannot occur in the present situation.

This shows that  $b'$  cannot be  $-b$  or  $1/b$ , and therefore  $b' = b, -1/b$  and  $\mathfrak{k} = \mathfrak{k}'$ . This gives the following.

**Theorem 10.** *For  $\mathfrak{m} = \wp_5$  or  $\wp_5'$ , where  $(5) = \wp_5 \wp_5'$  in  $K$ , Sugawara's conjecture holds, namely,  $K_{\mathfrak{m}} = K(\tau(\mathfrak{k}^*))$  is generated over the quadratic field  $K$  by a single ray class invariant for the modulus  $\mathfrak{m}$ .*

If  $\mathfrak{m} = \mathfrak{p}$ , where  $\mathfrak{p}^2 = (5)$ , then the above computations and arguments all hold, except for the argument which eliminated  $f_2(x)$ . In this case the ray class field  $\Sigma_{\mathfrak{p}} = \mathbb{Q}(b)$  is normal over  $\mathbb{Q}$ . (See the discussion in [19, Section 5, pp. 123-129] for this case.) However, the polynomial  $f_2$  splits modulo 41 into distinct linears and irreducible quadratics. Since a normal polynomial splits into irreducible factors of the same degree mod  $p$ , for all primes not dividing the discriminant, this fact shows that  $f_2(x)$  is not normal and hence can be eliminated as a possibility. Thus, Sugawara's conjecture also holds in this case.

## 9 The case $\mathfrak{m} = \wp_3 \wp_5'$ .

For this case we might think of using the Tate normal form for a point of order 15. However, the coefficients of the defining equation for  $E_{15}$  are unwieldy, so it turns out to be more convenient to again make use of the curve

$$E_5(b) : Y^2 + (1+b)XY + bY = X^3 + bX^2, \quad (9.1)$$

where, as in [18],  $b \in K_{\wp_5'}$ , and  $E_5(b)$  has the Weierstrass normal form

$$E' : Y_1^2 = 4X_1^3 - g_2X_1 - g_3,$$

with  $X_1 = X + \frac{1}{12}(b^2 + 6b + 1)$  and

$$\begin{aligned} g_2 &= \frac{1}{12}(b^4 + 12b^3 + 14b^2 - 12b + 1), \\ g_3 &= \frac{-1}{216}(b^2 + 1)(b^4 + 18b^3 + 74b^2 - 18b + 1); \\ \Delta &= -b^5(b^2 + 11b - 1). \end{aligned}$$

The doubling formula on  $E_5(b)$  is

$$X(2P) = F(X) = \frac{X^4 - (b^2 + b)X^2 - 2b^2X - b^3}{4X^3 + (b^2 + 6b + 1)X^2 + 2b(b + 1)X + b^2}, \quad X = X(P);$$

so setting  $F(X) = X, X = X(P)$ , yields the polynomial whose roots are the  $X$ -coordinates of points  $P$  of order 3 on  $E_5(b)$ :

$$g(X) = 3X^4 + (b^2 + 6b + 1)X^3 + (3b^2 + 3b)X^2 + 3b^2X + b^3. \quad (9.2)$$

Assuming that  $-d \equiv 1, 4 \pmod{15}$ , the ideals  $\wp_3, \wp'_3$ , and  $\wp_3\wp'_3 = (3)$  are associated to 1, 1, and 2  $\tau$ -invariants, respectively. There are four roots of  $g(X)$ , so these correspond to these three ideals in some permutation.

Now we solve  $g(X) = 0$ . First, we shift to eliminate the  $X^2$  term:

$$\frac{1}{3}g\left(X - \frac{b^2 + 6b + 1}{12}\right) = X^4 + pX^3 + qX + r, \quad (9.3)$$

where

$$\begin{aligned} p &= -\frac{1}{24}(b^4 + 12b^3 + 14b^2 - 12b + 1) = -\frac{g_2}{2}, \\ q &= \frac{(b^2 + 1)(b^4 + 18b^3 + 74b^2 - 18b + 1)}{216} = -g_3, \\ r &= -\frac{(b^4 + 12b^3 + 14b^2 - 12b + 1)^2}{6912} = \frac{-g_2^2}{48}. \end{aligned}$$

The cubic resolvent of (31) is

$$k(y) = y^3 - 2py^2 + (p^2 - 4r)y + q^2 = y^3 + g_2y^2 + \frac{g_2^2}{3}y + g_3^2,$$

for which we have

$$k\left(y - \frac{g_2}{3}\right) = y^3 - \frac{g_2^3 - 27g_3^2}{27} = y^3 - \frac{\Delta}{27}.$$

Hence, one root of  $k(y)$  is

$$\begin{aligned}
\theta_1 &= -\frac{g_2}{3} + \frac{\Delta^{1/3}}{3} \\
&= \frac{-1}{36}(b^4 + 12b^3 + 14b^2 - 12b + 1) - \frac{1}{3}b^{5/3}(b^2 + 11b - 1)^{1/3} \\
&= \frac{-1}{36}(b^4 + 12b^3 + 14b^2 - 12b + 1) - \frac{1}{3}b^2 \left(b - \frac{1}{b} + 11\right)^{1/3} \\
&= \frac{-b^2}{36} \left(b^2 + 12b + 14 - \frac{12}{b} + \frac{1}{b^2}\right) - \frac{1}{3}b^2(z + 11)^{1/3}.
\end{aligned} \tag{9.4}$$

Now put  $z + 11 = \rho^3$ . Note that  $z + 11 \cong \wp_5^{\prime 3}$  and  $\rho \cong \wp_5^{\prime}$ . This gives that

$$\begin{aligned}
-\theta_1 &= \frac{b^2}{36}(z^2 + 12z + 16) + \frac{1}{3}b^2\rho = \frac{b^2}{36}(z^2 + 12z + 16 + 12\rho) \\
&= \frac{b^2}{36}(\rho^6 - 10\rho^3 + 12\rho + 5) \\
&= \frac{b^2}{36}(\rho^2 + 2\rho + 5)(\rho^2 - \rho - 1)^2,
\end{aligned} \tag{9.5}$$

and therefore

$$\sqrt{-\theta_1} = \frac{b}{6}(\rho^2 - \rho - 1)\sqrt{\rho^2 + 2\rho + 5}. \tag{9.6}$$

The other two roots of  $k(y)$  are obtained by replacing  $\Delta^{1/3}$  in (9.4) by  $\Delta^{1/3}\omega^i$ , or  $\rho$  by  $\omega^i\rho$  in (9.6), for  $i = 1, 2$ , giving

$$\begin{aligned}
\sqrt{-\theta_2} &= \frac{b}{6}(\omega^2\rho^2 - \omega\rho - 1)\sqrt{\omega^2\rho^2 + 2\omega\rho + 5} \\
\sqrt{-\theta_3} &= \frac{b}{6}(\omega\rho^2 - \omega^2\rho - 1)\sqrt{\omega\rho^2 + 2\omega^2\rho + 5}.
\end{aligned}$$

The roots of  $g(X)$  are

$$\begin{aligned}
X(P_1) = x_1 &= -\frac{b^2 + 6b + 1}{12} + \frac{1}{2}(\sqrt{-\theta_1} + \sqrt{-\theta_2} + \sqrt{-\theta_3}), \\
X(P_2) = x_2 &= -\frac{b^2 + 6b + 1}{12} + \frac{1}{2}(\sqrt{-\theta_1} - \sqrt{-\theta_2} - \sqrt{-\theta_3}), \\
X(P_3) = x_3 &= -\frac{b^2 + 6b + 1}{12} + \frac{1}{2}(-\sqrt{-\theta_1} + \sqrt{-\theta_2} - \sqrt{-\theta_3}), \\
X(P_4) = x_4 &= -\frac{b^2 + 6b + 1}{12} + \frac{1}{2}(-\sqrt{-\theta_1} - \sqrt{-\theta_2} + \sqrt{-\theta_3}).
\end{aligned}$$

Therefore, the corresponding  $\tau$ -invariants are:

$$\begin{aligned}\tau(\mathfrak{k}_1^*) &= \lambda \left( X(P_1) + \frac{b^2 + 6b + 1}{12} \right) = \frac{\lambda}{2} (\sqrt{-\theta_1} + \sqrt{-\theta_2} + \sqrt{-\theta_3}), \\ \tau(\mathfrak{k}_2^*) &= \lambda \left( X(P_2) + \frac{b^2 + 6b + 1}{12} \right) = \frac{\lambda}{2} (\sqrt{-\theta_1} - \sqrt{-\theta_2} - \sqrt{-\theta_3}), \\ \tau(\mathfrak{k}_3^*) &= \lambda \left( X(P_3) + \frac{b^2 + 6b + 1}{12} \right) = \frac{\lambda}{2} (-\sqrt{-\theta_1} + \sqrt{-\theta_2} - \sqrt{-\theta_3}), \\ \tau(\mathfrak{k}_4^*) &= \lambda \left( X(P_4) + \frac{b^2 + 6b + 1}{12} \right) = \frac{\lambda}{2} (-\sqrt{-\theta_1} - \sqrt{-\theta_2} + \sqrt{-\theta_3}),\end{aligned}$$

where

$$\begin{aligned}\lambda &= -2^7 3^5 \frac{g_2 g_3}{\Delta} \\ &= -12 \frac{(b^4 + 12b^3 + 14b^2 - 12b + 1)(b^2 + 1)(b^4 + 18b^3 + 74b^2 - 18b + 1)}{b^5(b^2 + 11b - 1)}.\end{aligned}$$

We know that these  $\tau$ -invariants lie in  $K_{(3)} = \Sigma(\omega)$ . By our earlier arguments for  $\mathfrak{m} = (3)$  we also know two of them are conjugate and generate  $\Sigma(\omega)$  over  $K$ . The other two must therefore lie in  $K_{\wp_3} = K_{\wp_3'} = \Sigma$ . Consider the sum

$$\tau(\mathfrak{k}_1^*) + \tau(\mathfrak{k}_2^*) = \lambda \sqrt{-\theta_1} = \frac{\lambda b}{6} (\rho^2 - \rho - 1) \sqrt{\rho^2 + 2\rho + 5}. \quad (9.7)$$

The quantity  $\lambda b$  lies in  $K_{\wp_5'}$  and  $z \in \Sigma$ , so that (9.5) and (9.7) imply that  $\rho \in K_{\wp_5'} K_{(3)} = K_{\wp_5'}(\omega)$ . But this extension has degree 4 over  $\Sigma$ , so the cubic  $Y^3 - (z + 11)$  must have a root in  $\Sigma$ . Hence, we may assume  $\rho \in \Sigma$ . With this assumption I claim that  $\sqrt{\rho^2 + 2\rho + 5} \in K_{\wp_5'}$ . Using  $z = b - 1/b$  we have

$$\begin{aligned}\lambda b &= -12 \frac{b^6(z^2 + 12z + 16)(b + 1/b)(z^2 + 18z + 76)}{b^6(z + 11)} \\ &= -12 \left( b + \frac{1}{b} \right) \frac{(z^2 + 12z + 16)(z^2 + 18z + 76)}{z + 11};\end{aligned}$$

therefore,

$$(\lambda b)^2 = 144 \left( b + \frac{1}{b} \right)^2 \left( \frac{(z^2 + 12z + 16)(z^2 + 18z + 76)}{z + 11} \right)^2.$$

But  $(b + 1/b)^2 = z^2 + 4$ , so that  $(\lambda b)^2 \in \Sigma$  and  $\lambda b$  is a Kummer element for  $K_{\wp_5'}/\Sigma$ . It follows from (9.5) that

$$\begin{aligned}(\tau(\mathfrak{k}_1^*) + \tau(\mathfrak{k}_3^*))^2 &= (\tau(\mathfrak{k}_2^*) + \tau(\mathfrak{k}_4^*))^2 \\ &= (\lambda \sqrt{-\theta_2})^2 = \frac{(\lambda b)^2}{36} (z^2 + 12z + 16 + 12\rho\omega); \\ (\tau(\mathfrak{k}_1^*) + \tau(\mathfrak{k}_4^*))^2 &= (\tau(\mathfrak{k}_2^*) + \tau(\mathfrak{k}_3^*))^2 \\ &= (\lambda \sqrt{-\theta_3})^2 = \frac{(\lambda b)^2}{36} (z^2 + 12z + 16 + 12\rho\omega^2).\end{aligned}$$

Now  $\rho\omega$  and  $\rho\omega^2$  are conjugates over  $\Sigma$ , whence it follows that  $\sqrt{-\theta_2}$  and  $\pm\sqrt{-\theta_3}$  are conjugate over  $K_{\wp'_5} = \Sigma(b)$ . Choosing signs so that  $\sqrt{-\theta_2}$  and  $\sqrt{-\theta_3}$  are conjugate, it follows from the relation

$$\sqrt{-\theta_2}\sqrt{-\theta_3} = \frac{g_3}{\sqrt{-\theta_1}}$$

(with the sign of  $\sqrt{-\theta_1}$  chosen correctly) that  $\tau(\mathfrak{t}_1^*)$  and  $\tau(\mathfrak{t}_2^*)$  are fixed by the automorphism  $\alpha = (b, \omega) \rightarrow (b, \omega^2)$  of  $K_{\wp'_5}K_{(3)}$  over  $\Sigma$ , while  $\tau(\mathfrak{t}_3^*)$  and  $\tau(\mathfrak{t}_4^*)$  are interchanged. Since the squares above lie in  $\Sigma(\omega) \setminus \Sigma$ , it follows that  $\{\tau(\mathfrak{t}_1^*), \tau(\mathfrak{t}_2^*)\}$  must be the pair of invariants which lies in  $\Sigma$ . It follows that either

$$\wp_3 P_1 = O \text{ or } \wp'_3 P_1 = O \text{ on } E_5(b).$$

Also,  $(\lambda b)\sqrt{\rho^2 + 2\rho + 5} \in \Sigma$ , so  $\sqrt{\rho^2 + 2\rho + 5}$  must also be a Kummer element for  $K_{\wp'_5}/\Sigma$ .

As a corollary of the discussion so far, we have the following fact, which follows from  $\rho^3 = z + 11 = -(\eta(w/5)/\eta(w))^6$  (see [18]).

**Theorem 11.** *If  $-d \equiv 1, 4 \pmod{15}$ , then for some  $i \in \{0, 1, 2\}$  we have that*

$$\rho = -\omega^i \left( \frac{\eta(w/5)}{\eta(w)} \right)^2 \in \Sigma = K_1,$$

where  $w = \frac{v + \sqrt{d_K}}{2} \in R_K$  satisfies  $5^2 \mid N(w)$ , as in [18, Thm. 1.1]; and  $\Sigma(\sqrt{\rho^2 + 2\rho + 5}) = K_{\wp'_5}$ .

Our next task is to find primitive  $\mathfrak{m}$ -division points on the curve  $E_5(b)$ , where  $\mathfrak{m} = \wp_3 \wp'_5$ .

The kernel of multiplication by  $\wp'_5$  on  $E_5(b)$  is

$$\ker(\wp'_5) = \{(0, 0), (0, -b), (-b, 0), (-b, b^2)\} = \{Q_1, Q_2, Q_3, Q_4\},$$

since the associated  $\tau$ -invariants generate  $K_{\wp'_5}$ . Hence the points  $Q_i + P_1$  satisfy  $\wp_3 \wp'_5(Q_i + P_1) = O$  or  $\wp'_3 \wp'_5(Q_i + P_1) = O$ , depending on whether  $\wp_3(P_1) = O$  or  $\wp'_3(P_1) = O$ . Assume the former. Setting  $y_1 = Y(P_1)$ , we have that

$$Q_1 + P_1 = (0, 0) + (X(P_1), Y(P_1)) = (0, 0) + (x_1, y_1);$$

so that

$$\tilde{x}_1 = X(Q_1 + P_1) = \left( \frac{y_1}{x_1} \right)^2 + (1+b) \frac{y_1}{x_1} - b - 0 - x_1 = \frac{-by_1}{x_1^2}.$$

Hence, we have that

$$\begin{aligned} \frac{x_1}{b^2} \tilde{x}_1^2 - \frac{b+1}{b} \tilde{x}_1 &= \frac{y_1^2 + (1+b)x_1 y_1}{x_1^3} \\ &= \frac{-by_1 + x_1^3 + bx_1^2}{x_1^3} = \frac{1}{x_1} \tilde{x}_1 + 1 + \frac{b}{x_1}; \end{aligned}$$

and  $\tilde{x}_1 = X(Q_1 + P_1)$  satisfies the equation

$$\frac{x_1}{b^2}X^2 - \left(\frac{b+1}{b} + \frac{1}{x_1}\right)X - 1 - \frac{b}{x_1} = 0. \quad (9.8)$$

Similarly, we find

$$\tilde{x}_2 = X(Q_2 + P_1) = \frac{by_1 + b^2}{x_1^2} + \frac{b + b^2}{x_1},$$

so that

$$\tilde{x}_1 + \tilde{x}_2 = \frac{b^2}{x_1^2} + \frac{b + b^2}{x_1}$$

is the coefficient of  $X$  in (9.8) after dividing through by  $x_1/b^2$ ; and

$$\tilde{x}_1\tilde{x}_2 = -\frac{b^2}{x_1} - \frac{b^3}{x_1^2},$$

which is the constant term in (9.8) after dividing by  $x_1/b^2$ . Now the coefficients of (9.8) involve the quantities  $b$  and

$$x_1 = -\frac{b^2 + 6b + 1}{12} + \frac{\tau(\mathfrak{k}_1^*)}{\lambda},$$

and therefore lie in  $\Sigma(b) = \mathbb{K}_{\varphi'_5}$  (recall that  $\tau(\mathfrak{k}_1^*) \in \Sigma$ ). Thus,  $\tilde{x}_1, \tilde{x}_2$  are roots of the equation

$$h_1(X) = X^2 - \left(\frac{b^2}{x_1^2} + \frac{b + b^2}{x_1}\right)X - \frac{b^2}{x_1} - \frac{b^3}{x_1^2} \in \mathbb{K}_{\varphi'_5}[X],$$

which I claim is irreducible over  $\mathbb{K}_{\varphi'_5}$ . Its roots satisfy

$$\begin{aligned} \lambda \left( \tilde{x}_1 + \frac{b^2 + 6b + 1}{12} \right) &= \tau(\bar{\mathfrak{k}}_1^*), \\ \lambda \left( \tilde{x}_2 + \frac{b^2 + 6b + 1}{12} \right) &= \tau(\bar{\mathfrak{k}}_2^*), \end{aligned}$$

which are the  $\tau$ -invariants for  $\mathfrak{m} = \varphi_3\varphi'_5$  corresponding to the points  $Q_1 + P_1, Q_2 + P_1$ . Since these invariants satisfy a quadratic equation over  $\mathbb{K}_{\varphi'_5}$  and must generate  $\mathbb{K}_{\varphi'_5\varphi_3}$  over  $\Sigma$ , they are conjugate over  $\mathbb{K}_{\varphi'_5}$ . It follows that  $h_1(X)$  is irreducible over  $\mathbb{K}_{\varphi'_5}$ . Furthermore, since the roots

$$\begin{aligned} \tilde{x}_3 &= X(Q_3 + P_1) = \frac{b^2y_1 - bx_1^2 - b^2x_1}{(x_1 + b)^2}, \\ \tilde{x}_4 &= X(Q_4 + P_1) = -\frac{b^2y_1 + bx_1^2 + (b^3 + 2b^2)x_1 + b^3}{(x_1 + b)^2}, \end{aligned}$$

are related by the same linear transformation to the invariants  $\tau(\bar{\mathfrak{k}}_i^*)$ , ( $i = 3, 4$ ), as are  $\tilde{x}_1, \tilde{x}_2$  to their invariants, the roots  $\tilde{x}_3, \tilde{x}_4$  must satisfy the equation

$$h_2(X) = X^2 + \left( \frac{b(2x_1^2 + (b^2 + 3b)x_1 + b^2)}{(x_1 + b)^2} \right) X + \frac{b^2 x_1}{x_1 + b},$$

where

$$\lambda^{-4} T_{\mathfrak{m}} \left( \lambda X + \lambda \frac{b^2 + 6b + 1}{12}, \mathfrak{k} \right) = h_1(X) h_2(X), \quad \mathfrak{m} = \wp_5' \wp_3. \quad (9.9)$$

This implies that  $h_2(X)$  is also irreducible over  $\mathbb{K}_{\wp_5'}$ . As a corollary of this discussion we note:

**Theorem 12.** *If the torsion point  $P_1$  on  $E_5(b)$  satisfies  $\wp_3(P_1) = O$ , its coordinates  $P_1 = (x_1, y_1)$  generate  $\mathbb{K}_{\wp_5' \wp_3}$  over  $\mathbb{Q}$ .*

*Proof.* We have that  $\mathbb{K}_{\wp_5'}(y_1) = \mathbb{K}_{\wp_5' \wp_3}$ , since  $\tilde{x}_1 = -by_1/x_1^2$  generates this field over  $\mathbb{K}_{\wp_5'}$  (recalling that  $\tau(\bar{\mathfrak{k}}_1^*) \in \Sigma \Rightarrow x_1 \in \mathbb{K}_{\wp_5'}$ ). Moreover, if  $\mathbb{P}_{\mathfrak{m}}$  denotes  $(R_K/\mathfrak{m})^\times$ , then

$$\mathbb{P}_{\wp_5' \wp_3} / \langle -1 \rangle \cong (\mathbb{P}_{\wp_5'} \times \mathbb{P}_{\wp_3}) / \langle (-1, -1) \rangle$$

is cyclic of order 4, generated by  $(2, -1)$ , from which it follows that  $\mathbb{K}_{\wp_5' \wp_3} / \Sigma$  is a cyclic quartic extension. Since  $y_1 \in \mathbb{K}_{\wp_5' \wp_3} \setminus \mathbb{K}_{\wp_5'}$ , this gives that  $\mathbb{K}_{\wp_5' \wp_3} = \Sigma(y_1)$ . Furthermore, by the defining equation for  $E_5(b)$ , we have

$$y_1^2 + x_1 y_1 - x_1^3 = b(-x_1 y_1 - y_1 + x_1^2).$$

The right side is clearly nonzero, so that  $b \in \mathbb{Q}(x_1, y_1)$ . Now the fact that  $\mathbb{Q}(b) = \mathbb{K}_{\wp_5'}$  yields the assertion of the theorem. (See [18, Thm. 4.6, p. 1196].)  $\square$

Now assume that

$$T_{\mathfrak{m}}(X, \mathfrak{k}) = T_{\mathfrak{m}}(X, \mathfrak{k}') = T_{\mathfrak{m}}(X, \mathfrak{k})^\psi$$

for two ideal classes  $\mathfrak{k}, \mathfrak{k}'$ , where  $j(\mathfrak{k}') = j(\mathfrak{k})^\psi$  and  $\psi$  is an automorphism of  $\Sigma/K$ . Assume  $\psi$  has been extended to an automorphism of  $\mathbb{K}_{\wp_5' \wp_3}/K$ , and denote the images  $\alpha^\psi = \alpha'$  under  $\psi$  by primes. It is clear that  $(0, 0) \in E_5(b)$  maps to  $(0, 0) \in E_5(b')$  and  $P_1 = (x_1, y_1)$  maps to  $P_1' = (x_1', y_1')$ . The field  $\mathbb{K}_{\wp_5'}$  is normal over  $K$ , so the polynomials  $h_1(X), h_2(X)$  above are mapped to the pair of irreducible polynomials  $h_1^\psi(X), h_2^\psi(X)$  over  $\mathbb{K}_{\wp_5'}$ .

Assume first that  $\{\tau(\bar{\mathfrak{k}}_1'^*), \tau(\bar{\mathfrak{k}}_2'^*)\} = \{\tau(\bar{\mathfrak{k}}_1^*), \tau(\bar{\mathfrak{k}}_2^*)\}$ . Then the differences

$$\begin{aligned} \tau(\bar{\mathfrak{k}}_1'^*) - \tau(\bar{\mathfrak{k}}_2'^*) &= \lambda(\tilde{x}_1 - \tilde{x}_2) \\ \tau(\bar{\mathfrak{k}}_1'^*) - \tau(\bar{\mathfrak{k}}_2'^*) &= \lambda'(\tilde{x}_1' - \tilde{x}_2') \end{aligned}$$

imply that

$$\lambda(\tilde{x}_1 - \tilde{x}_2) = \pm \lambda'(\tilde{x}_1' - \tilde{x}_2'). \quad (9.10)$$

There is a similar relation for  $\tilde{x}_3, \tilde{x}_4$  and their images under  $\psi$ . Now

$$\frac{\tau(\tilde{\mathfrak{k}}_1^*) - \tau(\tilde{\mathfrak{k}}_2^*)}{\tau(\tilde{\mathfrak{k}}_3^*) - \tau(\tilde{\mathfrak{k}}_4^*)} = \frac{\tilde{x}_1 - \tilde{x}_2}{\tilde{x}_3 - \tilde{x}_4} = -\frac{(x_1 + b)^2}{bx_1^2}.$$

Hence, (9.10) gives that

$$\frac{(x_1 + b)^2}{bx_1^2} = \pm \frac{(x'_1 + b')^2}{b'x_1'^2}. \quad (9.11)$$

We now recall that  $z + 11 \cong \wp_5^3$  (see [18, p. 1193]). Let  $\mathfrak{q}$  be the product of prime ideals dividing  $\wp_5'$  in  $K_{\wp_5'}$ , so that  $\mathfrak{q}^2 = \wp_5'$ . Since  $z = b - 1/b$  and  $b$  is a unit, this gives that  $b^2 + 11b - 1 \equiv (b - 57)^2 \equiv 0 \pmod{\mathfrak{q}^6}$ , so  $b \equiv 2 \pmod{\mathfrak{q}}$ . Furthermore, from (9.2) the congruence

$$g(X) \equiv 3X^4 + 2X^3 + 3X^2 + 2X + 3 \equiv 3(X + 1)^4 \pmod{\mathfrak{q}}$$

implies that  $x_1 \equiv -1 \pmod{\mathfrak{q}}$ . This gives that

$$\frac{(x_1 + b)^2}{bx_1^2} \equiv \frac{(-1 + 2)^2}{2(-1)^2} \equiv 3 \pmod{\mathfrak{q}}. \quad (9.12)$$

Since this congruence also holds for  $\frac{(x'_1 + b')^2}{b'x_1'^2}$  (and  $\mathfrak{q}$  is invariant under  $\psi$ ), this shows that only the plus sign in (9.11) can hold. It follows from

$$\frac{(x_1 + b)^2}{bx_1^2} = \frac{(x'_1 + b')^2}{b'x_1'^2} \quad (9.13)$$

that  $b/b' = b/b^\psi = \mathbf{B}^2$ ,  $\mathbf{B} \in K_{\wp_5'}$ .

This suggests the following conjecture. For its statement recall that a quadratic discriminant can be written as  $d_K = \prod_{p|d_K} p^*$ , where  $p^* = (-1)^{(p-1)/2}p$ , if  $p$  is odd, and  $2^*$  is one of the possibilities  $2^* = -4, 8, -8$ .

**Conjecture 2.** *Assume that  $-d \equiv 1, 4 \pmod{15}$  and the 2-factor of  $d_K = \prod_{p|d} p^*$  is not  $2^* = -4$ . If  $b/b^\psi \in (K_{\wp_5'}^\times)^2$  for some  $\psi \in \text{Gal}(K_{\wp_5'}/K)$ , then  $\psi = 1$ . Moreover, there is a unique  $\psi \in \text{Gal}(K_{\wp_5'}/K)$  for which  $b/b^\psi = -\mathbf{B}^2$ , with  $\mathbf{B} \in K_{\wp_5'}$ . Namely,  $\psi : b \rightarrow -1/b$  is the unique automorphism in  $\text{Gal}(K_{\wp_5'}/K)$  with this property.*

The assumption on  $2^*$  is equivalent to the assertion that  $\sqrt{-1}$  does not lie in the genus field of  $K$ .

If (9.13) holds, then

$$\frac{(x_1 + b)^2}{bx_1^2} = \frac{(x'_1 + b')^2}{b'x_1'^2} = \left( \frac{(x_1 + b)^2}{bx_1^2} \right)^\psi$$



implies that  $\frac{(x_1+b)^2}{bx_1^2} = A$  lies in the fixed field  $L$  of  $\langle \psi \rangle$  inside  $K_{\phi'_5}$ . This is the case if

$$\{\tau(\bar{\mathfrak{k}}_1^*), \tau(\bar{\mathfrak{k}}_2^*)\} = \{\tau(\bar{\mathfrak{k}}_1^*), \tau(\bar{\mathfrak{k}}_2^*)\}.$$

If, on the other hand,

$$\{\tau(\bar{\mathfrak{k}}_1^*), \tau(\bar{\mathfrak{k}}_2^*)\} = \{\tau(\bar{\mathfrak{k}}_3^*), \tau(\bar{\mathfrak{k}}_4^*)\},$$

then we have, by the congruence condition (9.12), the equation

$$\frac{(x_1+b)^2}{bx_1^2} = -\frac{b'x_1'^2}{(x'_1+b')^2}. \quad (9.14)$$

Let  $\bar{\psi}$  denote the automorphism  $\bar{\psi} : b \rightarrow -1/b$  in  $\text{Gal}(K_{\phi'_5}/\Sigma)$ . Setting  $\tau = \tau_1(\bar{\mathfrak{k}}_1^*) \in \Sigma$ , we have that  $(b\lambda)^{\bar{\psi}} = -b\lambda$  and

$$\begin{aligned} x_1^{\bar{\psi}} &= \left( -\frac{b^2+6b+1}{12} + \frac{\tau}{\lambda} \right)^{\bar{\psi}} \\ &= -\frac{b^2-6b+1}{12b^2} + \frac{\tau}{b^2\lambda} \\ &= \frac{1}{b^2}(x_1+b). \end{aligned}$$

Hence,

$$(x_1+b)^{\bar{\psi}} = \frac{1}{b^2}(x_1+b) - \frac{1}{b} = \frac{x_1}{b^2}$$

and

$$\left( \frac{(x_1+b)^2}{bx_1^2} \right)^{\bar{\psi}} = -\frac{bx_1^2}{(x_1+b)^2}.$$

Using the fact that  $\bar{\psi}$  commutes with  $\psi$ , (9.14) implies that

$$\left( \frac{(x_1+b)^2}{bx_1^2} \right)^{\psi} = \left( \frac{(x_1+b)^2}{bx_1^2} \right)^{\bar{\psi}}. \quad (9.15)$$

In either case, (9.13) and (9.15) show that

$$\begin{aligned} \xi &= \frac{(x_1+b)^2}{bx_1^2} + \left( \frac{(x_1+b)^2}{bx_1^2} \right)^{\bar{\psi}} \\ &= \frac{(x_1+b)^2}{bx_1^2} - \frac{bx_1^2}{(x_1+b)^2} \\ &= \frac{(bx_1^2+b^2+2bx_1+x_1^2)(-bx_1^2+b^2+2bx_1+x_1^2)}{bx_1^2(x_1+b)^2} \end{aligned}$$

is fixed by  $\psi$  and therefore lies in  $\Sigma \cap L$ . We compute the following polynomial satisfied by  $\xi$ . First, note that  $\frac{(x_1+b)^2}{bx_1^2}$  is a root of the resultant

$$\begin{aligned} \text{Res}_{x_1}(g(x_1), bx_1^2X - (x_1+b)^2) &= b^9(bX^4 + (6b-1)X^3 + 9bX^2 - (b^2+6b)X + b) \\ &= b^9g_1(X, b). \end{aligned}$$

Since  $b$  is a unit, it is clear that  $\frac{(x_1+b)^2}{bx_1^2}$  is also a unit and  $\xi$  is an algebraic integer. Now let

$$\begin{aligned} F(X) &= -g_1(X, b)g_1\left(X, \frac{-1}{b}\right) \\ &= X^8 + (12+z)X^7 + (53+6z)X^6 + (96+8z)X^5 + (-z^2-12z+9)X^4 \\ &\quad - (96+8z)X^3 + (53+6z)X^2 - (12+z)X + 1, \end{aligned}$$

where  $z = b - \frac{1}{b} \in \Sigma$ . This polynomial also has  $\frac{(x_1+b)^2}{bx_1^2}$  as a root and has coefficients in  $\Sigma$ . Since this polynomial is reverse reciprocal in  $X$ , we can write it as a polynomial in  $X - \frac{1}{X}$ :

$$\begin{aligned} F(X) &= G\left(X - \frac{1}{X}, z\right), \\ G(X, z) &= X^4 + (12+z)X^3 + (57+6z)X^2 + (132+11z)X - z^2 + 117. \end{aligned}$$

Now the polynomial  $G(X, z)$  has  $X = \xi$  as a root. We write  $G(X, z)$  as a polynomial in  $z_1 = z + 11$ :

$$G(X, z) = -z_1^2 + (X^3 + 6X^2 + 11X + 22)z_1 + X^4 + X^3 - 9X^2 + 11X - 4,$$

so that  $z_1 = z + 11$  is a root of

$$G_1(\xi, Z) = -Z^2 + (\xi^3 + 6\xi^2 + 11\xi + 22)Z + \xi^4 + \xi^3 - 9\xi^2 + 11\xi - 4 = 0.$$

Since  $\xi$  lies in  $\Sigma \cap L$ ,  $z_1$  is at most quadratic over the latter field, i.e.,  $[\Sigma : \Sigma \cap L] \leq 2$ . On the other hand,  $z_1^\psi$  must also be a root of this polynomial. Suppose that  $z_1^\psi \neq z_1$ . Then

$$z_1 + z_1^\psi = \xi^3 + 6\xi^2 + 11\xi + 22. \quad (9.16)$$

But  $z_1 = z + 11 \cong \wp_5^3$  implies, since  $\psi$  fixes  $K$  and therefore  $\wp_5'$ , that  $z_1^\psi \cong \wp_5^3$ , as well. Hence, we have that

$$\xi^3 + 6\xi^2 + 11\xi + 22 \equiv 0 \pmod{\wp_5^3};$$

and  $G_1(\xi, z_1) = 0$  implies that its constant term satisfies

$$\xi^4 + \xi^3 - 9\xi^2 + 11\xi - 4 \equiv 0 \pmod{\wp_5^3}.$$

Now we argue 5-adically. The unique root of  $X^3 + 6X^2 + 11X + 22 = 0$  in  $\mathbb{Q}_5$  is

$$\alpha = 1 + 2 \cdot 5 + 2 \cdot 5^2 + 4 \cdot 5^6 + 2 \cdot 5^7 + 2 \cdot 5^8 + \dots,$$

since

$$X^3 + 6X^2 + 11X + 22 \equiv (X^2 + 2X + 3)(X + 4) \pmod{5}.$$

On the other hand,

$$\xi^4 + \xi^3 - 9\xi^2 + 11\xi - 4 = (\xi + 4)(\xi - 1)^3 \equiv (\xi + 4)^4 \pmod{\wp'_5},$$

and we conclude that  $\xi \equiv 1$  modulo each prime divisor of  $\wp'_5$  in  $\Sigma$  and therefore  $\xi \equiv 1 \pmod{\wp'_5}$ . Now the expansion of  $\alpha$  shows that the unique root of  $f(X) = X^3 + 6X^2 + 11X + 22$  modulo  $\wp_5^3$  which is congruent to 1 mod  $\wp'_5$  is  $X \equiv 61$ . For example, we have

$$f(x) - f(y) = (x - y)(x^2 + xy + y^2 + 6x + 6y + 11),$$

so since  $\xi \equiv 1 \pmod{\wp'_5}$ , we have

$$0 \equiv f(\xi) - f(61) = (\xi - 61)(\xi^2 + 67\xi + 4098) \pmod{\wp_5^3},$$

where  $\xi^2 + 67\xi + 4098 \equiv 1 \pmod{\wp'_5}$  is relatively prime to  $\wp'_5$ . Thus we must have

$$\xi \equiv 61 \pmod{\wp_5^3}.$$

But now consider the equation

$$0 = G_1(\xi, z_1) = -z_1^2 + (\xi^3 + 6\xi^2 + 11\xi + 22)z_1 + \xi^4 + \xi^3 - 9\xi^2 + 11\xi - 4.$$

The quadratic and linear terms in  $z_1$  on the right are each divisible by  $\wp_5^6$ . This gives that

$$\xi^4 + \xi^3 - 9\xi^2 + 11\xi - 4 \equiv 0 \pmod{\wp_5^6}. \quad (9.17)$$

However,  $\xi \equiv 61 + h \pmod{\wp_5^6}$ , where  $\wp_5^3 \mid h$ . Setting

$$k(x) = x^4 + x^3 - 9x^2 + 11x - 4 = (x + 4)(x - 1)^3,$$

we have

$$k(x + h) = h^4 + (4x + 1)h^3 + (6x^2 + 3x - 9)h^2 + (4x^3 + 3x^2 - 18x + 11)h + k(x),$$

where the coefficient of  $h$  is

$$a_3 = 4x^3 + 3x^2 - 18x + 11 \equiv (4x + 1)(x + 4)^2 \pmod{5}.$$

Thus,

$$\begin{aligned} k(\xi) &\equiv k(61 + h) \equiv (4 \cdot 61^3 + 3 \cdot 61^2 - 18 \cdot 61 + 11)h + k(61) \\ &\equiv (2^4 \cdot 3^3 \cdot 5^3 \cdot 17)h + k(61) \\ &\equiv k(61) = 2^6 \cdot 3^3 \cdot 5^4 \cdot 13 \pmod{\wp_5^6}, \end{aligned}$$

which is not zero modulo  $\wp_5^6$ , contradicting (9.17)!

This contradiction shows that  $z^\psi = z$  and therefore  $j(\mathfrak{f}') = j(\mathfrak{f})^\psi = j(\mathfrak{f})$ . This completes the proof of Sugawara's conjecture in this case. It is immaterial whether we take the ideal  $\wp_3$  or  $\wp'_3$  in this argument, since we have only argued 5-adically. Thus, we have proved

**Theorem 13.** *If  $\mathfrak{m} = \wp_3\wp_5'$  or  $\wp_3'\wp_5'$ , where  $d_K \equiv 1, 4 \pmod{15}$ , then the ray class field  $K_{\mathfrak{m}}$  is generated by a single  $\tau$ -invariant for the ideal  $\mathfrak{m}$ .*

It remains to consider the cases when  $(3) = \wp_3^2$  or  $(5) = \wp_5^2$  in  $K$ . In case  $(5) = \wp_5^2$ , we appeal to the result of [19, Prop. 5.1, pp. 124-125] and its proof, which applies to any fundamental discriminant  $d_K$  divisible by 5, and which shows that  $b = r^5(w_i/5)$  (denoted  $\rho_i, i = 1, 2$  in [19]) has the same properties with respect to the curve  $E_5(b)$  as in the cases discussed above. Namely:  $E_5(b)$  has complex multiplication by  $R_K$ , the ring of integers in  $K = \mathbb{Q}(\sqrt{-d})$ ;  $\mathbb{K}_{\wp_5} = K(b)$  and  $\psi : b \rightarrow -1/b$  is the non-trivial automorphism of the quadratic extension  $\mathbb{K}_{\wp_5}/\Sigma$ ; and  $b - 1/b + 11 \cong \wp_5^3$  in  $\Sigma$ . The same arguments above apply, up to and including the first sentence after equation (9.16) (with  $\wp_5' = \wp_5$ ). In this case the final argument does not lead to a contradiction, since  $5^4 \cong \wp_5^8$ . However, the constant term of  $G_1(X, z)$  gives that

$$-k(\xi) = -(\xi + 4)(\xi - 1)^3 = z_1 z_1^\psi \cong \wp_5^6.$$

Let  $\bar{\mathfrak{p}} \mid \wp_5$  be a prime divisor of  $\wp_5$  in  $\Sigma$ . The ideal  $\bar{\mathfrak{p}}^6$  exactly divides  $\wp_5^6$ , since  $\wp_5$  is unramified in  $\Sigma/K$ . Now  $\bar{\mathfrak{p}}$  divides at least one of  $\xi + 4$  and  $\xi - 1$  and therefore both, since  $(\xi + 4) - (\xi - 1) = 5$ . Furthermore,  $\bar{\mathfrak{p}}^2$  divides at least one of these factors – otherwise, the left side would only be divisible by  $\bar{\mathfrak{p}}^4$ . But  $\bar{\mathfrak{p}}^2 \mid 5$ , so  $\bar{\mathfrak{p}}^2$  must divide both factors, which leads to a contradiction, since then the left side would be divisible by  $\bar{\mathfrak{p}}^8$ . Hence,  $z_1^\psi = z_1$  and the same conclusion follows.

Finally, suppose that  $(3) = \wp_3^2$  in  $K$ . Here,

$$[\mathbb{K}_{(3)} : \Sigma] = \frac{\varphi(\wp_3^2)}{2} = 3,$$

so that  $\mathbb{K}_{(3)}$  is a cubic extension of  $\Sigma$ . In this case, there are three  $\tau$ -invariants corresponding to  $\mathfrak{m} = \wp_3^2$  and one invariant corresponding to  $\mathfrak{m} = \wp_3$ . Thus, one of the four  $\tau$ -invariants lies in  $\Sigma$  and the other three generate  $\mathbb{K}_{(3)}$  over  $\Sigma$ . I claim that  $\rho^3 = z + 11$ , where  $\rho$  generates  $\mathbb{K}_{(3)}$  over  $\Sigma$ . If  $\rho$  were an element of  $\Sigma$ , then since  $\omega \in \Sigma$ , the square-roots  $\sqrt{-\theta_i}$  would be quadratic over  $\mathbb{K}_{\wp_5'}$  (or  $\mathbb{K}_{\wp_5}$ , if  $5 \mid d_K$ ), and their sums could not generate the cubic extension  $\mathbb{K}_{\wp_5'}\mathbb{K}_{(3)}$  of  $\mathbb{K}_{\wp_5'}$ .

Let  $\sigma$  be the generator of  $\text{Gal}(\mathbb{K}_{(3)}/\Sigma)$ , for which  $\rho^\sigma = \omega\rho$ , and define

$$\sqrt{-\theta_1}^\sigma = \sqrt{-\theta_2}, \quad \sqrt{-\theta_2}^\sigma = \sqrt{-\theta_3}.$$

Then  $\sqrt{-\theta_3}^\sigma = \sqrt{-\theta_1}^{\sigma^3} = \sqrt{-\theta_1}$ . Extending  $\sigma$  to  $\mathbb{K}_{\wp_5'}\mathbb{K}_{(3)}$  so that it fixes  $\mathbb{K}_{\wp_5'}$ , it follows that

$$\tau(\mathfrak{k}_1^*) = \frac{\lambda}{2} \text{Tr}_{\mathbb{K}_{\wp_5'}}(\sqrt{-\theta_1}) \in \mathbb{K}_{\wp_5'},$$

which implies that  $\tau(\mathfrak{k}_1^*)$  cannot generate  $\mathbb{K}_{(3)}$  and must therefore lie in  $\Sigma$ . Now the rest of the calculations are the same. Hence, we have proved the following.

**Theorem 14.** *If  $\mathfrak{m} = \wp_3\wp_5$ , where  $\wp_3, \wp_5$  are first degree prime divisors of 3, 5 in  $K$ , then the ray class field  $K_{\mathfrak{m}}$  is generated by a single  $\tau$ -invariant for the ideal  $\mathfrak{m}$ .*

Before proceeding to the next case we consider the following example.

**Example.** Let  $K = \mathbb{Q}(\sqrt{-11})$ . Even though this field has class number 1, it is interesting to use the arithmetic of this section to compute the various  $\tau$ -invariants. Using  $j(\mathfrak{o}) = -32^3$ , we find that  $b$  is a root of

$$B(x) = x^4 + 4x^3 + 46x^2 - 4x + 1,$$

and we take

$$b = -1 + \sqrt{-11} + \frac{3 - \sqrt{-11}}{4} \sqrt{-2\sqrt{-11} + 6}.$$

With this value of  $b$  we compute the roots of  $g(X) = 0$  in  $K_{\wp_5'} = K(b) = K(\sqrt{-2\sqrt{-11} + 6})$  to be

$$\begin{aligned} x_1 &= -\frac{2}{3} + \frac{\sqrt{-11}}{6} + \frac{7 - \sqrt{-11}}{24} \sqrt{-2\sqrt{-11} + 6}, \\ x_2 &= -\frac{5}{6} - \frac{\sqrt{-11}}{3} + \frac{1 + \sqrt{-11}}{8} \sqrt{-2\sqrt{-11} + 6}. \end{aligned}$$

Now  $\tilde{x}_1, \tilde{x}_2$  are roots of the polynomial

$$\begin{aligned} h_1(X) &= X^2 + \left( -\frac{31}{4} - \frac{7\sqrt{-11}}{4} + \frac{13 + 7\sqrt{-11}}{8} \sqrt{-2\sqrt{-11} + 6} \right) X \\ &\quad - \frac{63}{2} - 15\sqrt{-11} + \frac{23 + 47\sqrt{-11}}{8} \sqrt{-2\sqrt{-11} + 6}; \end{aligned}$$

while  $\tilde{x}_3, \tilde{x}_4$  are roots of

$$\begin{aligned} h_2(X) &= X^2 + \left( \frac{-15 + \sqrt{-11}}{4} + \frac{11 + \sqrt{-11}}{8} \sqrt{-2\sqrt{-11} + 6} \right) X \\ &\quad - \frac{11 + 5\sqrt{-11}}{4} + \frac{1 + 2\sqrt{-11}}{4} \sqrt{-2\sqrt{-11} + 6}. \end{aligned}$$

Note that the polynomials  $h_1, h_2$  are not conjugate over  $K$ , but the polynomials

$$\begin{aligned} \tilde{h}_1(X) &= \lambda^2 h_1 \left( \frac{X}{\lambda} - \frac{b^2 + 6b + 1}{12} \right) \\ &= X^2 + \left( -19712 - 16128\sqrt{-11} + 5376\sqrt{-11} \sqrt{-2\sqrt{-11} + 6} \right) X \\ &\quad - 2684616704 - 1271660544\sqrt{-11} + 423886848\sqrt{-11} \sqrt{-2\sqrt{-11} + 6} \end{aligned}$$

and

$$\begin{aligned}\tilde{h}_2(X) &= \lambda^2 h_2 \left( \frac{X}{\lambda} - \frac{b^2 + 6b + 1}{12} \right) \\ &= X^2 + \left( -19712 - 16128\sqrt{-11} - 5376\sqrt{-11}\sqrt{-2\sqrt{-11} + 6} \right) X \\ &\quad - 2684616704 - 1271660544\sqrt{-11} - 423886848\sqrt{-11}\sqrt{-2\sqrt{-11} + 6}\end{aligned}$$

are conjugate over  $K$ , and their product is the polynomial

$$\begin{aligned}T_{\mathfrak{m}}(X, \mathfrak{k}) &= X^4 + (-39424 - 32256\sqrt{-11})X^3 \\ &\quad + (-5934415872 - 2543321088\sqrt{-11})X^2 \\ &\quad + (-44563506921472 + 36461051117568\sqrt{-11})X \\ &\quad + 1277724870452445184 + 2874880958518001664\sqrt{-11}.\end{aligned}$$

This question of conjugacy is the source of the difficulty in proving Sugawara's conjecture. The roots of  $T_{\mathfrak{m}}(X, \mathfrak{k})$  are the  $\tau$ -invariants  $\tau(\mathfrak{k}_i^*)$  for the ideal  $\mathfrak{m} = \wp_3\wp_5'$  and the four ray classes mod  $\mathfrak{m}$ . Note in this case that

$$z = -2 + 2\sqrt{-11}, \quad \rho = \frac{-3 + \sqrt{-11}}{2}, \quad \rho^2 + 2\rho + 5 = \frac{3 - \sqrt{-11}}{2},$$

where  $(\rho) = \wp_5'$  in  $K$ . (In Theorem 11, the value  $i = 0$ , so  $\rho = -\eta(w/5)^2/\eta(w)^2$ , where  $w = (33 + \sqrt{-11})/2$ .) Taking the product of  $T_{\mathfrak{m}}(X, \mathfrak{k})$  with its image under complex conjugation yields the interesting polynomial

$$\begin{aligned}T(X) &= T_{\wp_3\wp_5'}(X, \mathfrak{k})T_{\wp_3\wp_5}(X, \mathfrak{k}) \\ &= X^8 - (2^{10} \cdot 7 \cdot 11)X^7 + (2^{21} \cdot 7^2 \cdot 11)X^6 + (2^{30} \cdot 7^5 \cdot 11^2)X^5 \\ &\quad + (2^{40} \cdot 7^4 \cdot 11^2 \cdot 271)X^4 - (2^{50} \cdot 5 \cdot 7^5 \cdot 11^3 \cdot 29)X^3 \\ &\quad - (2^{60} \cdot 7^6 \cdot 11^3 \cdot 883)X^2 + (2^{71} \cdot 7^8 \cdot 11^5)X \\ &\quad + 2^{80} \cdot 7^8 \cdot 11^4 \cdot 907.\end{aligned}$$

If we were to replace  $x_1$  with the root  $x_2$  of  $g(X)$  we would obtain the polynomial  $\tilde{T}(X) = T_{\wp_3\wp_5'}(X, \mathfrak{k})T_{\wp_3\wp_5}(X, \mathfrak{k})$ .

## 10 The case $\mathfrak{m} = \wp_3\wp_3'\wp_5$ .

We can use the same points  $P_i, Q_j$  and the calculations from Section 9 to deal with the case  $\mathfrak{m} = \wp_3\wp_3'\wp_5$ .

By the results of Section 5 and the arguments preceding Theorem 11, the invariants  $\tau(\mathfrak{k}_3^*)$  and  $\tau(\mathfrak{k}_4^*)$  generate  $\mathbf{K}_{(3)} = \Sigma(\omega)$  and

$$\wp_3\wp_3'P_3 = \wp_3\wp_3'P_4 = O \quad \text{on } E_5(b).$$

The points  $P_3, P_4$  are primitive (3)-division points; otherwise the associated  $\tau$ -invariants would lie in  $\Sigma$ , which we showed in Section 9 is not the case. Hence, the points  $P_i + Q_j$ , for  $i = 3, 4$  and  $1 \leq j \leq 4$ , are primitive  $\mathfrak{m}$ -division points. We also know that  $\varphi(\mathfrak{m}) = 16$ , so that

$$[\mathbf{K}_{\mathfrak{m}} : \Sigma] = 8;$$

$\mathbf{K}_{\mathfrak{m}}$  is quadratic over  $\mathbf{K}_{\wp_3 \wp'_3}$  and quartic over  $\mathbf{K}_{(3)} = \Sigma(\omega)$ . The  $X$ -coordinates of the points  $P_3 + Q_j$  are obtained by replacing  $x_1$  by  $x_3$  in the formulas for  $\tilde{x}_j$  in Section 9, where  $P_3 = (x_3, y_3)$ :

$$\begin{aligned}\tilde{x}_5 &= X(P_3 + Q_1) = \frac{-by_3}{x_3^2}, \\ \tilde{x}_6 &= X(P_3 + Q_2) = \frac{by_3 + b^2}{x_3^2} + \frac{b + b^2}{x_3}, \\ \tilde{x}_7 &= X(P_3 + Q_3) = \frac{b^2y_3 - bx_3^2 - b^2x_3}{(x_3 + b)^2}, \\ \tilde{x}_8 &= X(P_3 + Q_4) = -\frac{b^2y_3 + bx_3^2 + (b^3 + 2b^2)x_3 + b^3}{(x_3 + b)^2}.\end{aligned}$$

The coordinates  $X(P_4 + Q_j)$  are formed by replaicng  $x_3$  by  $x_4$  in these formulas. See the discussion just before Theorem 11. Now the quantity

$$\tau(\bar{\mathfrak{F}}_5^*) = \lambda \left( \tilde{x}_5 + \frac{b^2 + 6b + 1}{12} \right) = \lambda \left( \frac{-by_3}{x_3^2} + \frac{b^2 + 6b + 1}{12} \right)$$

must generate  $\mathbf{K}_{\mathfrak{m}}$  over  $\Sigma$ , while the quantity we considered in Section 9,

$$\tau(\mathfrak{F}_3^*) = \lambda \left( x_3 + \frac{b^2 + 6b + 1}{12} \right) = \frac{\lambda}{2} \left( -\sqrt{-\theta_1} - \sqrt{-\theta_2} + \sqrt{-\theta_3} \right),$$

generates  $\mathbf{K}_{(3)} = \Sigma(\omega)$  over  $\Sigma$ . It follows that  $x_3 \in \mathbf{K}_{\wp'_3}(\omega)$  and  $\mathbf{K}_{\mathfrak{m}} = \Sigma(b, x_3, y_3)$ .

Thus, the trace of  $\tilde{x}_5$  to  $F(\omega) = \mathbf{K}_{\wp'_3}(\omega)$  is

$$\mathrm{Tr}_{\mathbf{K}_{\mathfrak{m}}/F(\omega)}(\tilde{x}_5) = \tilde{x}_5 + \tilde{x}_6 = \frac{(b^2 + b)x_3 + b^2}{x_3^2},$$

since

$$\tilde{x}_5 \tilde{x}_6 = -\frac{b^2(x_3 + b)}{x_3^2},$$

as in Section 9, so that  $\tilde{x}_5, \tilde{x}_6$  satisfy a quadratic polynomial with coefficients in  $F(\omega)$ :

$$h_3(X) = X^2 - \frac{(b^2 + b)x_3 + b^2}{x_3^2}X - \frac{b^2(x_3 + b)}{x_3^2}.$$

Hence,

$$\mathrm{Tr}_{\mathbf{K}_{\mathfrak{m}}/F}(\tilde{x}_5) = \frac{(b^2 + b)x_3 + b^2}{x_3^2} + \frac{(b^2 + b)x_4 + b^2}{x_4^2},$$

since  $\tau(\mathfrak{k}_3^*)$  and  $\tau(\mathfrak{k}_4^*)$  are conjugates over  $F = \mathbb{K}_{\varphi'_5}$ . Similarly,  $\tilde{x}_7, \tilde{x}_8$  are roots of the polynomial

$$h_4(X) = X^2 + \frac{b(2x_3^2 + (b^2 + 3b)x_3 + b^2)}{(x_3 + b)^2}X + \frac{b^2x_3}{x_3 + b},$$

which is irreducible over  $F(\omega)$ . Hence, setting

$$\begin{aligned} s_1 &= \tilde{x}_5 + \tilde{x}_6 = \frac{b^2}{x_3^2} + \frac{b + b^2}{x_3}, \\ s_2 &= \tilde{x}_7 + \tilde{x}_8 = -\frac{b(2x_3^2 + (b^2 + 3b)x_3 + b^2)}{(x_3 + b)^2}, \end{aligned}$$

we have

$$s_1 - s_2 = \tilde{x}_5 + \tilde{x}_6 - \tilde{x}_7 - \tilde{x}_8 \quad (10.1)$$

$$= b \frac{(b^2 + 1)x_3^3 + (3b^2 + 3b)x_3^2 + (3b^3 + 3b^2)x_3 + b^3}{3x_3^2(x_3 + b)^2} = u(x_3, b), \quad (10.2)$$

on reducing the numerator modulo  $g(x_3)$ . With the automorphism  $\bar{\psi} : b \rightarrow -1/b$  from Section 9, extended to  $\mathbb{K}_{\varphi'_5}(\omega)$  to fix  $\omega$ , a similar calculation to the displayed lines following (9.14), using that  $\tau = \tau(\mathfrak{k}_3^*) \in \mathbb{K}_{(3)} = \Sigma(\omega)$ , shows that

$$x_3^{\bar{\psi}} = \frac{1}{b^2}(x_3 + b).$$

Furthermore, with

$$\tau(\bar{\mathfrak{k}}_5^*) + \tau(\bar{\mathfrak{k}}_6^*) = \lambda \left( s_1 + \frac{b^2 + 6b + 1}{6} \right),$$

we have

$$\begin{aligned} (\tau(\bar{\mathfrak{k}}_5^*) + \tau(\bar{\mathfrak{k}}_6^*))^{\bar{\psi}} &= \lambda^{\bar{\psi}} \left( s_1 + \frac{b^2 + 6b + 1}{6} \right)^{\bar{\psi}} \\ &= \lambda b^2 \left( s_1^{\bar{\psi}} + \frac{b^2 - 6b + 1}{6b^2} \right) \\ &= \lambda \left( b^2 \frac{(1-b)x_3 + b}{(x_3 + b)^2} + \frac{b^2 - 6b + 1}{6} \right) \\ &= \lambda \left( \frac{(b^2 - b^3)x_3 + b^3 - 2b(x_3 + b)^2}{(x_3 + b)^2} + \frac{b^2 + 6b + 1}{6} \right) \\ &= \lambda \left( -\frac{b(2x_3^2 + (b^2 + 3b)x_3 + b^2)}{(x_3 + b)^2} + \frac{b^2 + 6b + 1}{6} \right) \\ &= \lambda \left( s_2 + \frac{b^2 + 6b + 1}{6} \right) \\ &= \tau(\bar{\mathfrak{k}}_7^*) + \tau(\bar{\mathfrak{k}}_8^*). \end{aligned}$$



Therefore, the traces to  $\mathbb{K}_{\wp'_5}(\omega)$  of the  $\tau$ -invariants for  $\mathbf{m}$  are

$$S = \{\lambda(s_1 + \beta), \lambda(s_2 + \beta), \lambda(s_3 + \beta), \lambda(s_4 + \beta)\},$$

where

$$s_3 = s_1^\alpha, s_4 = s_2^\alpha, \alpha : (b, \omega) \rightarrow (b, \omega^2); \beta = \frac{b^2 + 6b + 1}{6}.$$

The traces in  $S$  are the respective images of  $\lambda(s_1 + \beta)$  by the automorphisms in

$$\text{Gal}(\mathbb{K}_{\wp'_5}(\omega)/\Sigma) = \{1, \bar{\psi}, \alpha, \bar{\psi}\alpha\}.$$

**Lemma 2.** *The conjugates over  $\Sigma$  of the trace  $\tau(\bar{\mathfrak{k}}_5) + \tau(\bar{\mathfrak{k}}_6) = \lambda(s_1 + \beta)$  are distinct.*

*Proof.* If  $\lambda(s_1 + \beta) = \lambda(s_3 + \beta)$ , then  $s_1 = s_3$ , or

$$\begin{aligned} 0 = s_1 - s_3 &= \frac{(b^2 + b)x_3 + b^2}{x_3^2} - \frac{(b^2 + b)x_4 + b^2}{x_4^2} \\ &= \frac{-b(x_3 - x_4)[(b + 1)x_3x_4 + b(x_3 + x_4)]}{x_3^2x_4^2}. \end{aligned}$$

Now the roots of  $g(X)$  are distinct, since

$$\text{disc}(g(X)) = -27b^{10}(b^2 + 11b - 1)^2 = -27\Delta^2.$$

Hence,  $x_3 \neq x_4$ . It follows that  $(b + 1)x_3x_4 + b(x_3 + x_4) = 0$ . But

$$\begin{aligned} &\text{Res}_{x_4}(\text{Res}_{x_3}((b + 1)x_3x_4 + b(x_3 + x_4), g(x_3)), g(x_4)) \\ &= b^{28}(b^4 + 2b^3 - 32b^2 + 14b - 1)(b^4 + 3b^3 + 2b^2 - 10b + 1)^2. \end{aligned}$$

However, the discriminants of these quartics are  $2^{12} \cdot 19 \cdot 103$  and  $-7^3 \cdot 701$ , neither of which is divisible by 5. Therefore, their product cannot be 0 and  $s_1 \neq s_3$ . From  $(s_2 - s_4)^{\bar{\psi}} = \frac{1}{b^2}(s_1 - s_3)$  we also know  $s_2 \neq s_4$ . (See the proof of Lemma 3 below.)

Now suppose that  $s_1 = s_2$ . Then

$$\begin{aligned} 0 = s_1 - s_2 &= \frac{(b^2 + b)x_3 + b^2}{x_3^2} + \frac{b(2x_3^2 + (b^2 + 3b)x_3 + b^2)}{(x_3 + b)^2} \\ &= \frac{b}{x_3^2(x_3 + b)^2} \\ &\quad \times (2x_3^4 + (b^2 + 4b + 1)x_3^3 + (3b^2 + 3b)x_3^2 + (b^3 + 3b^2)x_3 + b^3). \end{aligned}$$

In this case we have

$$\begin{aligned} &\text{Res}_{x_3}(2x_3^4 + (b^2 + 4b + 1)x_3^3 + (3b^2 + 3b)x_3^2 + (b^3 + 3b^2)x_3 + b^3, g(x_3)) \\ &= b^{14}(b^2 + b - 1)(b^2 + 11b - 1). \end{aligned}$$

Since the roots of  $x^2 + x - 1$  are real and  $K \subset \mathbb{Q}(b)$ , this polynomial cannot have  $b$  as a root. Hence  $s_1 \neq s_2$ . Applying  $\alpha$  gives that  $s_3 \neq s_4$ .

Finally, suppose  $s_1 = s_4$ . In this case

$$\begin{aligned} 0 = s_1 - s_4 &= \frac{(b^2 + b)x_3 + b^2}{x_3^2} + \frac{b(2x_4^2 + (b^2 + 3b)x_4 + b^2)}{(x_4 + b)^2} \\ &= \frac{b}{x_3^2(x_4 + b)^2} \\ &\quad \times ((2x_4^2 + (b^2 + 3b)x_4 + b^2)x_3^2 + ((b + 1)x_4^2 + (2b^2 + 2b)x_4 + b^3 + b^2)x_3 \\ &\quad + b^3 + 2b^2x_4 + bx_4^2) \\ &= \frac{b}{x_3^2(x_4 + b)^2} k(x_3, x_4). \end{aligned}$$

We compute that

$$\begin{aligned} &\text{Res}_{x_4}(\text{Res}_{x_3}(k(x_3, x_4), g(x_3)), g(x_4)) \\ &= b^{55}(b^2 + b - 1)(b^2 + 11b - 1)^3 \\ &\quad \times (b^{10} + 15b^9 + 47b^8 + 44b^7 + 1014b^6 + 58b^5 - 1014b^4 + 44b^3 - 47b^2 + 15b - 1). \end{aligned}$$

The tenth degree factor  $L(b)$  has a nonsolvable Galois group, since  $L(b) = b^5 G(b - \frac{1}{b})$ , where

$$G(x) = x^5 + 15x^4 + 52x^3 + 104x^2 + 1160x + 176$$

has Galois group  $S_5$ . It follows that  $s_1 \neq s_4$  and  $s_2 \neq s_3$ . This proves the lemma.  $\square$

**Lemma 3.** *The cross-ratio  $\kappa = (s_1, s_2; s_3, s_4) = \frac{(s_1 - s_3)(s_2 - s_4)}{(s_1 - s_4)(s_2 - s_3)}$  lies in the Hilbert class field  $\Sigma$ . We also have*

$$s_1 - s_2 + s_3 - s_4 = \eta b, \quad \eta \in \Sigma.$$

*Proof.* We compute that

$$\kappa^\alpha = (s_3, s_4; s_1, s_2) = \frac{(s_3 - s_1)(s_4 - s_2)}{(s_3 - s_2)(s_4 - s_1)} = \kappa.$$

Using  $\lambda^{\bar{\psi}} = b^2\lambda$ , we find that

$$\begin{aligned} (s_1 - s_3)^{\bar{\psi}} &= \left( \frac{\lambda s_1 - \lambda s_3}{\lambda} \right)^{\bar{\psi}} \\ &= \frac{\lambda s_2 - \lambda s_4}{b^2 \lambda} = \frac{1}{b^2} (s_2 - s_4). \end{aligned}$$

The same idea yields that

$$(s_2 - s_4)^{\bar{\psi}} = \frac{1}{b^2}(s_1 - s_3), \quad (s_1 - s_4)^{\bar{\psi}} = \frac{1}{b^2}(s_2 - s_3), \quad (s_2 - s_3)^{\bar{\psi}} = \frac{1}{b^2}(s_1 - s_4).$$

It follows that

$$\kappa^{\bar{\psi}} = \frac{b^{-4}(s_2 - s_4)(s_1 - s_3)}{b^{-4}(s_2 - s_3)(s_1 - s_4)} = \kappa.$$

Thus,  $\kappa$  is fixed by  $\text{Gal}(\mathbf{K}_{\rho_5'}(\omega)/\Sigma)$ . This proves the first assertion. The second follows from the calculation

$$\begin{aligned} \eta^{\bar{\psi}} &= \left( \frac{s_1 - s_2 + s_3 - s_4}{b} \right)^{\bar{\psi}} = \frac{\frac{-1}{b^2}(s_1 - s_2) + \frac{-1}{b^2}(s_3 - s_4)}{\frac{-1}{b}} \\ &= \frac{s_1 - s_2 + s_3 - s_4}{b} = \eta, \end{aligned}$$

together with the fact that the quantity  $\eta$  is obviously fixed by the automorphism  $\alpha$ . Hence,  $\eta \in \Sigma$ .  $\square$

**Lemma 4.** *We have that  $\rho^4 - 2\rho^3 - \rho^2 - 10\rho + 25 = \nu^2\rho^2$ , where  $\nu \in \Sigma$  is an algebraic integer.*

*Proof.* On one hand, we have

$$\begin{aligned} \sqrt{-\theta_2}\sqrt{-\theta_3} &= \frac{b^2}{36}(\omega^2\rho^2 - \omega\rho - 1)(\omega\rho^2 - \omega^2\rho - 1) \\ &\quad \times \sqrt{(\omega^2\rho^2 + 2\omega\rho + 5)(\omega\rho^2 + 2\omega^2\rho + 5)} \\ &= \frac{b^2}{36}(\rho^4 + \rho^3 + 2\rho^2 - \rho + 1)\sqrt{\rho^4 - 2\rho^3 - \rho^2 - 10\rho + 25}. \quad (10.3) \end{aligned}$$

We also have

$$\begin{aligned} \sqrt{-\theta_2}\sqrt{-\theta_3} &= \frac{g_3}{\sqrt{-\theta_1}} \\ &= \frac{-b^2}{36} \left( b + \frac{1}{b} \right) \frac{z^2 + 18z + 76}{(\rho^2 - \rho - 1)\sqrt{\rho^2 + 2\rho + 5}} \\ &= \frac{-b^2}{36} \frac{z^2 + 18z + 76}{\rho^2 - \rho - 1} \frac{\sqrt{z^2 + 4}}{\sqrt{\rho^2 + 2\rho + 5}}. \end{aligned}$$

The last equation and (10.3) give that

$$\begin{aligned} (\rho^4 + \rho^3 + 2\rho^2 - \rho + 1)\sqrt{\rho^4 - 2\rho^3 - \rho^2 - 10\rho + 25} &= -\frac{z^2 + 18z + 76}{\rho^2 - \rho - 1} \frac{\sqrt{z^2 + 4}}{\sqrt{\rho^2 + 2\rho + 5}} \\ &= -\frac{(\rho^2 - \rho - 1)(\rho^4 + \rho^3 + 2\rho^2 - \rho + 1)}{\rho^2 - \rho - 1} \frac{\sqrt{z^2 + 4}}{\sqrt{\rho^2 + 2\rho + 5}} \\ &= -(\rho^4 + \rho^3 + 2\rho^2 - \rho + 1) \frac{\sqrt{z^2 + 4}}{\sqrt{\rho^2 + 2\rho + 5}}. \end{aligned}$$

Hence,

$$\sqrt{\rho^4 - 2\rho^3 - \rho^2 - 10\rho + 25} = -\frac{\sqrt{z^2 + 4}}{\sqrt{\rho^2 + 2\rho + 5}}.$$

Now we know that both square roots on the right side of this equation are Kummer elements for  $K_{\rho'_5}/\Sigma$ , so the quotient lies in  $\Sigma$ . Hence,

$$\rho^4 - 2\rho^3 - \rho^2 - 10\rho + 25 = \gamma^2, \quad \gamma \in \Sigma.$$

But  $\rho \cong \rho'_5$  also lies in  $\Sigma$  and  $\rho^2$  divides the left side of this equation. It follows that  $\gamma = \nu\rho$ , with  $\nu \in R_\Sigma$ . This proves the lemma.  $\square$

**Corollary 1.** *For the quantity  $\nu$  in Lemma 3 we have*

$$\nu^2 + 12 = \left(\rho - 1 + \frac{5}{\rho}\right)^2.$$

*Proof.* This follows directly from

$$\nu^2 = \frac{\rho^4 - 2\rho^3 - \rho^2 - 10\rho + 25}{\rho^2} = \frac{(\rho^2 - \rho + 5)^2 - 12\rho^2}{\rho^2}.$$

$\square$

**Lemma 5.** *If  $5 \nmid d_K$ , then  $\mathbb{Q}(\rho) = \Sigma$ ; while if  $5 \mid d_K$ , then  $K(\rho) = \Sigma$ .*

*Proof.* This is clear from  $z = \rho^3 - 11$  and  $j = -\frac{(z^2 + 12z + 16)^3}{z + 11}$ , using that  $\mathbb{Q}(z) = \Sigma$  from [18, Prop. 3.2] when  $5 \nmid d_K$ ; and  $K(z) = \Sigma$  from [19, Prop. 5.1] when  $5 \mid d_K$ .  $\square$

**Lemma 6.** *We have:*

(a) *For any root  $x$  of  $g(X) = 0$ ,*

$$\frac{1}{x} = \frac{-1}{b^3}(3x^3 + (b^2 + 6b + 1)x^2 + (3b^2 + 3b)x + 3b^2)$$

and

$$\frac{1}{x + b} = \frac{1}{b^5}(3x^3 + (b^2 + 3b + 1)x^2 + (-b^3 + 2b)x + b^4 + b^2).$$

(b)

$$\begin{aligned} s_1 = f_1(x_3) &= \frac{(b^2 + b)x_3 + b^2}{x_3^2} = \frac{-3(b-2)}{b^2}x_3^3 - \frac{(b^3 + 4b^2 - 8b - 2)}{b^2}x_3^2 \\ &\quad - \frac{(4b^2 + 3b - 5)}{b}x_3 - 6b + 3. \end{aligned}$$

(c)

$$\begin{aligned}
s_2 = f_2(x_3) &= \frac{-b(2x_3^2 + (b^2 + 3b)x_3 + b^2)}{(x_3 + b)^2} \\
&= \frac{3(2b + 1)}{b^3}x_3^3 + \frac{(2b^3 + 10b^2 + 5b + 1)}{b^3}x_3^2 - \frac{(b^3 + b^2 - 5b - 2)}{b^2}x_3 \\
&\quad - \frac{(b^2 - 3b - 1)}{b}.
\end{aligned}$$

*Proof.* This follows by straightforward calculation, using the formula in (a) for  $x = x_3$ .  $\square$

Note that the formulas for  $s_3 = f_1(x_4)$  and  $s_4 = f_2(x_4)$  are the same as the formulas in (b) and (c), with  $x_3$  replaced by  $x_4$ .

**Proposition 1.**

$$s_1 - s_2 - s_3 + s_4 = (x_3 - x_4) \left( \frac{-(\rho + 1)\sqrt{z^2 + 4}}{2} + \frac{(\rho^3 - 2\rho - 7)\sqrt{\rho^2 + 2\rho + 5}}{2} \right) \tag{10.4}$$

$$= (x_3 - x_4) \left( \frac{(\rho^3 + \nu\rho^2 + (\nu - 2)\rho - 7)\sqrt{\rho^2 + 2\rho + 5}}{2} \right). \tag{10.5}$$

*Proof.* Write  $s_1 = f_1(x_3)$ ,  $s_3 = f_1(x_4)$ ,  $s_2 = f_2(x_3)$ ,  $s_4 = f_2(x_4)$ . We start by noting the polynomial identity

$$\begin{aligned}
s_1 - s_2 - s_3 + s_4 &= f_1(x_3) - f_2(x_3) - f_1(x_4) + f_2(x_4) \\
&= \frac{(x_3 - x_4)}{b^3} [(x_3 + x_4 + 3)b^4 + (6x_3 + 6x_4 + 2)b^3 \\
&\quad + (3x_3^2 + 3x_4^2 + 2x_3 + 2x_4 + 3x_3x_4)b^2 \\
&\quad + (3x_3 + 3x_4 + 2)b + 3x_3^2 + 3x_4^2 + x_3 + x_4 + 3x_3x_4]
\end{aligned}$$

Now substitute

$$x_3 + x_4 = -\frac{b^2 + 6b + 1}{6} - \sqrt{-\theta_1}, \quad x_3^2 + x_4^2 = (x_3 + x_4)^2 - 2x_3x_4$$

into the last expression. This yields

$$\begin{aligned}
s_1 - s_2 - s_3 + s_4 &= \frac{-(x_3 - x_4)}{b^3} 3b\sqrt{-\theta_1} \\
&\quad + \frac{(x_3 - x_4)}{b^3} \frac{(b^2 + 1)(b^4 + 12b^3 + 2b^2 - 18b + 1 + 36x_3x_4 + 36\theta_1)}{12}.
\end{aligned} \tag{10.6}$$

Putting  $36\theta_1 = -b^2(z^2 + 12z + 16 + 12\rho)$  from (33) with  $z = b - \frac{1}{b}$  in (51) gives

$$s_1 - s_2 - s_3 + s_4 = \frac{-(x_3 - x_4)}{b^3} \left( 3b\sqrt{-\theta_1} + (b^2 + 1)((\rho + 1)b^2 + \frac{b}{2} - 3x_3x_4) \right). \quad (10.7)$$

Now use the result of the calculation

$$\begin{aligned} x_3x_4 &= \left( -\frac{b^2 + 6b + 1}{12} - \frac{1}{2}\sqrt{-\theta_1} + \frac{(\sqrt{-\theta_2} - \sqrt{-\theta_3})}{2} \right) \\ &\quad \times \left( -\frac{b^2 + 6b + 1}{12} - \frac{1}{2}\sqrt{-\theta_1} - \frac{(\sqrt{-\theta_2} - \sqrt{-\theta_3})}{2} \right) \\ &= \left( \frac{b^2 + 6b + 1}{12} \right)^2 + \frac{b^2 + 6b + 1}{12}\sqrt{-\theta_1} - \frac{\theta_1}{4} - \frac{(\sqrt{-\theta_2} - \sqrt{-\theta_3})^2}{4} \\ &= \left( \frac{b^2 + 6b + 1}{12} \right)^2 + \frac{b^2 + 6b + 1}{12}\sqrt{-\theta_1} - \frac{\theta_1}{4} + \frac{\theta_2 + \theta_3 + 2\sqrt{-\theta_2}\sqrt{-\theta_3}}{4}, \end{aligned}$$

with

$$\theta_2 + \theta_3 = \frac{-b^2}{36}(2z^2 + 24z + 32 + 12\omega\rho + 12\omega^2\rho) = \frac{-b^2}{36}(2z^2 + 24z + 32 - 12\rho)$$

in (10.7), giving

$$\begin{aligned} s_1 - s_2 - s_3 + s_4 &= \frac{-(x_3 - x_4)}{b^3} \\ &\quad \times \left( \frac{(-b^4 - 6b^3 - 2b^2 + 6b - 1)\sqrt{-\theta_1}}{4} + \frac{(b^2 + 1)(-3\sqrt{-\theta_2}\sqrt{-\theta_3} + (\rho + 1)b^2)}{2} \right) \\ &= (x_3 - x_4) \left( \frac{(b^4 + 6b^3 + 2b^2 - 6b + 1)\sqrt{-\theta_1}}{4b^3} - \frac{(b^2 + 1)(-3\frac{\sqrt{-\theta_2}\sqrt{-\theta_3}}{b^2} + (\rho + 1))}{2b} \right). \end{aligned}$$

Now we use

$$\begin{aligned} \frac{(b^4 + 6b^3 + 2b^2 - 6b + 1)}{b^2} &= z^2 + 6z + 4, \quad b + \frac{1}{b} = \sqrt{z^2 + 4}, \\ \sqrt{-\theta_1} &= \frac{b}{6}(\rho^2 - \rho - 1)\sqrt{\rho^2 + 2\rho + 5}, \\ \frac{-3\sqrt{-\theta_2}\sqrt{-\theta_3}}{b^2} &= \frac{(\rho^4 + \rho^3 + 2\rho^2 - \rho + 1)\sqrt{z^2 + 4}}{12\sqrt{\rho^2 + 2\rho + 5}}, \end{aligned}$$

(the last from the proof of Lemma 4) to write the factor in the large parenthesis as

$$\mathbf{A} = -\frac{2(\rho + 1)\sqrt{z^2 + 4}\sqrt{\rho^2 + 2\rho + 5} - z(\rho^4 + \rho^3 + 2\rho^2) + (z^2 + 7z + 4)\rho + z^2 + 5z + 4}{4\sqrt{\rho^2 + 2\rho + 5}}.$$

The final terms in the numerator of  $\mathbf{A}$  factor on setting  $z = \rho^3 - 11$ :

$$-z(\rho^4 + \rho^3 + 2\rho^2) + (z^2 + 7z + 4)\rho + z^2 + 5z + 4 = -2(\rho^3 - 2\rho - 7)(\rho^2 + 2\rho + 5),$$

giving

$$\begin{aligned} A &= -\frac{2(\rho+1)\sqrt{z^2+4}\sqrt{\rho^2+2\rho+5}-2(\rho^3-2\rho-7)(\rho^2+2\rho+5)}{4\sqrt{\rho^2+2\rho+5}} \\ &= \frac{-(\rho+1)\sqrt{z^2+4}+(\rho^3-2\rho-7)\sqrt{\rho^2+2\rho+5}}{2}. \end{aligned}$$

This proves (10.4). Equation (10.5) follows from (10.4) on using

$$\sqrt{z^2+4} = -\sqrt{\rho^2+2\rho+5}\sqrt{\rho^4-2\rho^3-\rho^2-10\rho+25} = -\nu\rho\sqrt{\rho^2+2\rho+5}$$

by Lemma 4 and its proof.  $\square$

**Corollary 2.** *The quantity  $s_1 - s_2 - s_3 + s_4 \neq 0$ .*

*Proof.* This follows from (10.4), since

$$\begin{aligned} (\rho+1)^2((\rho^3-11)^2+4) - (\rho^3-2\rho-7)^2(\rho^2+2\rho+5) &= 12(\rho-2)(\rho^2+2\rho+5) \\ \text{and } \rho = -1 \pm 2i \text{ gives } z = \pm 2i \text{ and } j = 1728, \text{ which is excluded.} &\quad \square \end{aligned}$$

A similar calculation leads to the following result.

**Proposition 2.**

$$\begin{aligned} s_1 + s_2 - s_3 - s_4 &= (x_3 - x_4) \\ &\times \left( \frac{-(\rho^4 + 3\rho^3 - 15\rho - 23)\sqrt{\rho^2 + 2\rho + 5} + (\rho^2 + 4\rho + 5)\sqrt{z^2 + 4}}{2\sqrt{\rho^2 + 2\rho + 5}} \right) \end{aligned} \quad (10.8)$$

$$= -(x_3 - x_4) \frac{(\rho^4 + 3\rho^3 - 15\rho - 23) + (\rho^2 + 4\rho + 5)\nu\rho}{2}. \quad (10.9)$$

*Proof.* This follows from

$$\begin{aligned} s_1 + s_2 - s_3 - s_4 &= f_1(x_3) + f_2(x_3) - f_1(x_4) - f_2(x_4) \\ &= \frac{-(x_3 - x_4)}{b^3} B_1, \\ B_1 &= (x_3 + x_4 + 5)b^4 + (2x_3 + 2x_4 + 4)b^3 \\ &\quad + (3x_3^2 + 3x_4^2 - 18x_3 - 18x_4 + 3x_3x_4 - 10)b^2 \\ &\quad + (-12x_3^2 - 12x_4^2 - 7x_3 - 7x_4 - 12x_3x_4 - 2)b \\ &\quad - 3x_3^2 - 3x_4^2 - x_3 - x_4 - 3x_3x_4, \end{aligned}$$

using the same substitutions as in the proof of Proposition 1. This leads to

$$B = \frac{-1}{b^3} B_1 = \frac{-(\rho^4 + 3\rho^3 - 15\rho - 23)\sqrt{\rho^2 + 2\rho + 5} + (\rho^2 + 4\rho + 5)\sqrt{z^2 + 4}}{2\sqrt{\rho^2 + 2\rho + 5}}.$$

$\square$

Now we find a formula for the quotient of the sums in Propositions 1 and 2. A straightforward calculation establishes the following result.

**Theorem 15.** *If  $A$  and  $B$  are the respective cofactors of  $(x_3 - x_4)$  in (10.4) and (10.8), we have*

$$\frac{s_1 + s_2 - s_3 - s_4}{s_1 - s_2 - s_3 + s_4} = \frac{B}{A} = \frac{(-2\rho + 3)\sqrt{\rho^2 + 2\rho + 5} + \sqrt{z^2 + 4}}{(\rho - 2)(\rho^2 + 2\rho + 5)} \quad (10.10)$$

$$= \frac{(3 - (\nu + 2)\rho)\sqrt{\rho^2 + 2\rho + 5}}{(\rho - 2)(\rho^2 + 2\rho + 5)}. \quad (10.11)$$

**Remark.** It is clear that the quotient in (10.10) is the quotient of sums and differences of the traces of the eight  $\tau$ -invariants for  $\wp_3\wp_3'\wp_5'$  in  $\mathbb{K}_{(3)\wp_5'}/\mathbb{K}_{\wp_5'}(\omega)$ .

Now assume that  $\psi \in \text{Gal}(\mathbb{K}_{(3)\wp_5'}/K)$  is an automorphism fixing the set of  $\tau$ -invariants for  $\wp_3\wp_3'\wp_5'$ . Then the set of traces of these invariants to  $\mathbb{K}_{\wp_5'}(\omega)$  is also fixed, and

$$\{\lambda(s_1 + \beta), \lambda(s_2 + \beta), \lambda(s_3 + \beta), \lambda(s_4 + \beta)\}^\psi$$

is a permutation of

$$S = \{\lambda(s_1 + \beta), \lambda(s_2 + \beta), \lambda(s_3 + \beta), \lambda(s_4 + \beta)\}.$$

This permutation must respect the orbits of  $S$  under the automorphisms  $\alpha, \bar{\psi}, \alpha\bar{\psi}$ , since they commute with  $\psi$ . The numerator on the left side of (10.10) is the difference of the orbit sums of  $S$  under  $\bar{\psi}$  divided by  $\lambda$ , while the denominator is the difference of orbit sums under  $\alpha\bar{\psi}$  divided by  $\lambda$ . Hence, the quotient is preserved up to sign, and we have the relation

$$\left( \frac{(3 - (\nu + 2)\rho)\sqrt{\rho^2 + 2\rho + 5}}{(\rho - 2)(\rho^2 + 2\rho + 5)} \right)^\psi = \pm \frac{(3 - (\nu + 2)\rho)\sqrt{\rho^2 + 2\rho + 5}}{(\rho - 2)(\rho^2 + 2\rho + 5)}.$$

Now  $(\rho) = \wp_5'$  (see the line just before (9.5)) implies that  $\rho^\psi = a\rho$ , for some unit  $a \in \Sigma$ . Furthermore,  $\sqrt{\rho^2 + 2\rho + 5}^\psi = \varepsilon\sqrt{\rho^2 + 2\rho + 5}$ , with  $\varepsilon \in \Sigma$ , since both square-roots are Kummer elements for  $\mathbb{K}_{\wp_5'}/\Sigma$ . This implies the relation

$$\frac{(3 - (\nu^\psi + 2)a\rho)}{(a\rho - 2)} \frac{1}{\varepsilon} = \pm \frac{(3 - (\nu + 2)\rho)}{(\rho - 2)}.$$

This implies that  $\varepsilon \equiv \pm 1 \pmod{(\rho)}$ , i.e.,  $\pmod{\wp_5'}$ . On the other hand, we also have

$$(\rho^2 + 2\rho + 5)^\psi = a^2\rho^2 + 2a\rho + 5 = \varepsilon^2(\rho^2 + 2\rho + 5),$$

or

$$(a^2 - \varepsilon^2)\rho^2 + 2(a - \varepsilon^2)\rho + 5 - 5\varepsilon^2 = 0.$$



Dividing this equation by  $\rho$  gives

$$(a^2 - \varepsilon^2)\rho + 2(a - \varepsilon^2) + 5\frac{1 - \varepsilon^2}{\rho} = 0,$$

hence

$$a \equiv \varepsilon^2 \equiv 1 \pmod{(\rho)}$$

and

$$\rho^\psi = a\rho \equiv \rho \pmod{(\rho)^2}. \quad (10.12)$$

Now we appeal to the formula

$$j(\mathfrak{k}) = j = -\frac{(z^2 + 12z + 16)^3}{z + 11}.$$

Putting  $z = \rho^3 - 11$  implies that

$$j(\mathfrak{k}) = j = -\frac{(\rho^6 - 10\rho^3 + 5)^3}{\rho^3} = -\left(\rho^5 - 10\rho^2 + \frac{5}{\rho}\right)^3.$$

Since  $5/\rho$  is an algebraic integer, this gives further that

$$j^\psi - j = -\left(\rho^{5\psi} - 10\rho^{2\psi} + \frac{5}{\rho^\psi} - \left(\rho^5 - 10\rho^2 + \frac{5}{\rho}\right)\right)\Xi, \quad (10.13)$$

for some algebraic integer  $\Xi$ . Now by (10.12),

$$\rho^{5\psi} \equiv \rho^5, \quad \rho^{2\psi} \equiv \rho^2 \pmod{(\rho)^2},$$

and

$$\frac{5}{\rho^\psi} - \frac{5}{\rho} = 5\frac{\rho - \rho^\psi}{\rho\rho^\psi} \equiv 0 \pmod{(\rho)},$$

since  $\rho\rho^\psi \cong \rho^2$ . Thus, (10.13) yields that

$$j(\mathfrak{k})^\psi \equiv j(\mathfrak{k}) \pmod{\wp'_5};$$

which implies  $\psi|_\Sigma = 1$  when  $5 \nmid d_K$ , since the discriminant of  $H_{d_K}(X)$  is not divisible by 5 in this case. This proves that  $j(\mathfrak{k})^\psi = j(\mathfrak{k})$  and the polynomials  $T_{\mathfrak{m}}(X, \mathfrak{k})$  are distinct for different ideal classes  $\mathfrak{k}$ .

This proves:

**Theorem 16.** *If  $K = \mathbb{Q}(\sqrt{d})$ , where  $d = d_k \equiv 1, 4 \pmod{15}$ , then Sugawara's conjecture holds for  $K$  and the ideal  $\mathfrak{m} = \wp_3\wp'_3\wp'_5$ , i.e.,  $K_{\mathfrak{m}}$  is generated over  $K$  by a single  $\tau$ -invariant for the ideal  $\mathfrak{m}$ .*

Note that the ideal  $\mathfrak{m} = \wp_3^2\wp_5$  satisfies Sugawara's condition (1.3), so there is no need to consider fields in which 3 ramifies.

## 10.1 Proof of the conjecture when $(5) = \wp_5^2$ .

All the above arguments apply in the case that  $(5) = \wp_5^2$ , except for the final conclusion. In this case, namely, when  $5 \mid d_K$ , the conjugates of  $j(\mathfrak{k})$  can be congruent to each other modulo prime divisors of  $\wp_5$ . To handle this case we set

$$R(\rho, \nu) = \frac{(3 - (\nu + 2)\rho)^2}{(\rho - 2)^2(\rho^2 + 2\rho + 5)}.$$

This is the square of the expression in (10.11). Then we must show:

$$\psi \in \text{Gal}(\mathbf{K}_m/K) \wedge R(\rho, \nu)^\psi = R(\rho, \nu) \Rightarrow \psi|_\Sigma = 1. \quad (10.14)$$

As before, we know  $\rho^\psi = a\rho$  for some unit  $a$  and  $(\rho^2 + 2\rho + 5)^\psi = \varepsilon^2(\rho^2 + 2\rho + 5)$ , where  $\varepsilon \in \Sigma$  and

$$a \equiv 1, \quad \varepsilon \equiv \pm 1 \pmod{(\rho)}.$$

In this case, dividing the equation

$$a^2\rho^2 + 2a\rho + 5 = \varepsilon^2(\rho^2 + 2\rho + 5)$$

by  $\rho$  yields that

$$(a^2 - \varepsilon^2)\rho + 2(a - \varepsilon^2) + 5\frac{(1 - \varepsilon^2)}{\rho} = 0,$$

and implies the congruence

$$a \equiv \varepsilon^2 \pmod{\rho^2}.$$

**Proposition 3.** *There are at most 2 automorphisms  $\psi \in \text{Gal}(\Sigma/K)$  for which  $R(\rho, \nu)^\psi = R(\rho, \nu)$ .*

*Proof.* Expanding  $R(\rho, \nu)$  and using Corollary 1 to Lemma 4 gives that

$$\begin{aligned} r = R(\rho, \nu) &= \frac{(3 - (\nu + 2)\rho)^2}{(\rho - 2)^2(\rho^2 + 2\rho + 5)} \\ &= \frac{2(2\rho - 3)\rho\nu}{(\rho - 2)^2(\rho^2 + 2\rho + 5)} + \frac{\rho^4 - 2\rho^3 + 3\rho^2 - 22\rho + 34}{(\rho - 2)^2(\rho^2 + 2\rho + 5)}. \end{aligned}$$

Squaring and using Corollary 1 again shows that

$$\begin{aligned} r^2 = R(\rho, \nu)^2 &= \frac{4(\rho^4 - 2\rho^3 + 3\rho^2 - 22\rho + 34)\rho(2\rho - 3)\nu}{(\rho - 2)^4(\rho^2 + 2\rho + 5)^2} \\ &\quad + \frac{\rho^8 - 4\rho^7 + 26\rho^6 - 136\rho^5 + 281\rho^4 - 452\rho^3 + 1532\rho^2 - 3056\rho + 2056}{(\rho - 2)^4(\rho^2 + 2\rho + 5)^2}. \end{aligned}$$

Solving for  $\nu$  in terms of  $r$  in the first equation yields that

$$\nu = \frac{\rho^4 - 2\rho^3 + \rho^2 - 12\rho + 20}{2(2\rho - 3)\rho}r + \frac{-\rho^4 + 2\rho^3 - 3\rho^2 + 22\rho - 34}{2(2\rho - 3)\rho}.$$

Now substitute for  $\nu$  in the expression for  $r^2$ :

$$\begin{aligned} r^2 &= \frac{(2\rho^6 + 8\rho^4 - 52\rho^3 + 10\rho^2 - 84\rho + 340)r}{(\rho^2 + 2\rho + 5)^2(\rho - 2)^2} \\ &\quad + \frac{-\rho^6 + 10\rho^4 + 16\rho^3 - 25\rho^2 - 80\rho - 64}{(\rho^2 + 2\rho + 5)^2(\rho - 2)^2} \\ &= \frac{2(\rho^2 + 2\rho + 5)(\rho^4 - 2\rho^3 + 3\rho^2 - 22\rho + 34)r}{(\rho^2 + 2\rho + 5)^2(\rho - 2)^2} - \frac{(\rho^3 - 5\rho - 8)^2}{(\rho^2 + 2\rho + 5)^2(\rho - 2)^2}. \end{aligned}$$

Rearranging gives the polynomial identity

$$\begin{aligned} 0 = F(r, \rho) &= (\rho^2 + 2\rho + 5)^2(\rho - 2)^2 r^2 \\ &\quad - 2(\rho^2 + 2\rho + 5)(\rho^4 - 2\rho^3 + 3\rho^2 - 22\rho + 34)r \\ &\quad + (\rho^3 - 5\rho - 8)^2. \end{aligned} \tag{10.15}$$

If  $r = 1$ , then a computation using the above formula for  $\nu$  above gives

$$\nu = \frac{-\rho^2 + 5\rho - 7}{(2\rho - 3)\rho},$$

and substituting yields

$$0 = R\left(\rho, -\frac{\rho^2 - 5\rho + 7}{\rho(2\rho - 3)}\right) - 1 = -4\frac{\rho^4 - \rho^3 - \rho^2 - 9\rho + 11}{(\rho^2 + 2\rho + 5)(2\rho - 3)^2}.$$

Thus,  $\rho$  would satisfy the irreducible equation

$$\rho^4 - \rho^3 - \rho^2 - 9\rho + 11 = 0.$$

But this is impossible, since  $(\rho) = \wp_5 \nmid 11$ .

Rewriting equation (10.15) as a polynomial in  $\rho$  gives us an equation satisfied by  $\rho$  over the field  $K(r)$ , where  $r = R(\rho, \nu)$ :

$$\begin{aligned} 0 = F(r, \rho) &= (r^2 - 2r + 1)\rho^6 + (2r^2 - 8r - 10)\rho^4 + (-20r^2 + 52r - 16)\rho^3 \\ &\quad + (r^2 - 10r + 25)\rho^2 + (-20r^2 + 84r + 80)\rho + 100r^2 - 340r + 64. \end{aligned} \tag{10.16}$$

Dividing through in (10.16) by the leading coefficient gives an equation with constant term

$$c = \frac{100r^2 - 340r + 64}{(r - 1)^2} = \frac{4(5r - 1)(5r - 16)}{(r - 1)^2}.$$

Since

$$r = \frac{(3 - (\nu + 2)\rho)^2}{(\rho - 2)^2(\rho^2 + 2\rho + 5)} \cong \frac{\mathfrak{a}}{\mathfrak{b}\wp_5},$$

where  $\mathfrak{a}, \mathfrak{b}$  are integral ideals which are relatively prime to  $\wp_5$ , it is clear that  $c = \frac{4(5r-1)(5r-16)}{(r-1)^2}$  is exactly divisible by  $\wp_5^2 \cong \rho^2$ . Furthermore, the coefficients

of the scaled equation are integral for  $\wp_5$ , and its roots are therefore integral for  $\wp_5$ . Also, since the conjugates of  $\rho$  are  $a_\psi\rho \cong \rho$ , there can be at most two such conjugates which are roots of (10.16). Hence, there can be at most two automorphisms  $1, \psi$ , for which  $\rho^\psi$  is a root of (10.16). Therefore, either  $\psi = 1$  is the only such automorphism; or  $\psi^2 = 1$  and  $\Sigma$  is quadratic over  $K(r)$ . This proves the proposition.  $\square$

Assume that  $R(\rho, \nu)^\psi = R(\rho, \nu)$ , where  $\psi$  has order 2. Then  $\rho^\psi = a\rho$  implies that  $\rho^{\psi^2} = a^\psi a\rho = \rho$ , giving that  $a^\psi = \frac{1}{a}$ .

From the proofs of Proposition 5.1 and Theorem 5.3 in [19, pp. 124-129] (which are also valid for fundamental discriminants  $d_K = -5d$  in place of the discriminants  $-5l$  considered in that paper), we know that, for some  $w \in R_K$

$$\lambda = \left( \frac{\eta(w/5)}{\eta(w)} \right)^6 = -\rho^3$$

is conjugate to the value

$$\lambda^{\sigma_{\wp_5}} = \frac{5^3}{\lambda}$$

over  $K$ , where  $\sigma_{\wp_5} = \left( \frac{\Sigma/K}{\wp_5} \right)$ . (See Theorem 11 in Section 9.) Since the cube roots of unity are not in  $\Sigma$ , this implies that

$$\rho^{\sigma_{\wp_5}} = \frac{5}{\rho}.$$

Now the automorphism  $\sigma = \sigma_{\wp_5}$  acts on  $a = \frac{\rho^\psi}{\rho}$  by

$$a^\sigma = \frac{(5/\rho)^\psi}{5/\rho} = \frac{\rho}{\rho^\psi} = \frac{1}{a}. \quad (10.17)$$

But we also have  $a^\psi = \frac{1}{a}$  from above. Now there are two cases.

*Case 1:*  $K(a) = \Sigma$ . In this case  $a^\psi = a^\sigma$  implies that  $\psi = \sigma = \sigma_{\wp_5}$ . This implies further, using the notation of (10.16), that

$$0 = F(r, \rho)^\psi = F(r, \rho^\sigma) = F\left(r, \frac{5}{\rho}\right).$$

Thus,  $\rho$  must be a root of the resultant

$$\begin{aligned} \text{Res}_r(F(r, \rho), \rho^6 F\left(r, \frac{5}{\rho}\right)) \\ &= 256(\rho^2 - 5)^2(\rho^2 + 2\rho + 5)^4(11\rho^4 - 14\rho^3 - 86\rho^2 - 70\rho + 275) \\ &\times (11\rho^8 - 164\rho^7 + 829\rho^6 - 1980\rho^5 + 3685\rho^4 - 9900\rho^3 + 20725\rho^2 \\ &\quad - 20500\rho + 6875). \end{aligned}$$

$\rho$  cannot be a root of the irreducible quartic or octic factor, since  $\rho$  is an algebraic integer. Further,  $\rho^2 + 2\rho + 5 \neq 0$ , since the discriminant of this quadratic is  $-16$  and not divisible by 5. Thus,  $\rho^2 - 5 = 0$ . A simple computation shows this is only possible for the discriminant  $d_K = -20$ : in this case

$$\begin{aligned} \text{Res}_\rho(\rho^2 - 5, z + 11 - \rho^3) &= z^2 + 22z - 4, \\ \text{Res}_z(z^2 + 22z - 4, (z + 11)j + (z^2 + 12z + 16)^3) \\ &= -5^3(j^2 - 1264000j - 681472000) = -5^3 H_{-20}(j). \end{aligned}$$

Hence,  $K = \mathbb{Q}(\sqrt{-5})$  and  $\Sigma = K(\sqrt{5}) = K(\sqrt{-1})$ ; since  $\wp_5$  splits in  $\Sigma$ , we have  $\sigma_{\wp_5} = \left(\frac{\Sigma/K}{\wp_5}\right) = 1$ , so  $\psi \neq \sigma_{\wp_5}$ . Notice that for  $\rho = \sqrt{5}$  the nontrivial automorphism  $\psi$  of  $\Sigma/K$  satisfies  $a = \frac{\rho^\psi}{\rho} = -1$ , which is certainly not congruent to 1 (mod  $\rho$ ). Thus,  $\psi$  does not fix  $r = R(\rho, \nu)$ , proving (10.14). This is an instance of Case 2, which we consider next.

*Case 2:*  $K(a) \neq \Sigma$ . Assume that  $\psi \neq \sigma_{\wp_5}$  fixes  $r = R(\rho, \nu)$ , so that  $F(r, \rho) = 0$ , as in Case 1. Then  $\psi$  also fixes

$$s = r^\sigma = R(\rho^\sigma, \nu^\sigma), \quad \sigma = \sigma_{\wp_5}.$$

By Lemma 4,

$$\begin{aligned} (\nu^2 \rho^2)^\sigma &= (\rho^4 - 2\rho^3 - \rho^2 - 10\rho + 25)^\sigma \\ &= \left(\frac{5}{\rho}\right)^4 - 2\left(\frac{5}{\rho}\right)^3 - \left(\frac{5}{\rho}\right)^2 - 10\left(\frac{5}{\rho}\right) + 25 \\ &= \frac{5^2}{\rho^4}(5^2 - 10\rho - \rho^2 - 2\rho^3 + \rho^4) \\ &= \frac{5^2}{\rho^4} \times \nu^2 \rho^2 = \frac{5^2}{\rho^2} \nu^2. \end{aligned}$$

This yields that  $\nu^\sigma = \pm \nu$ .

Assuming first that  $\nu^\sigma = \nu$ , we obtain that

$$s = R(\rho^\sigma, \nu) = R\left(\frac{5}{\rho}, \nu\right) = \frac{(3\rho - 5\nu - 10)^2 \rho^2}{(5(-5 + 2\rho)^2(\rho^2 + 2\rho + 5))}.$$

Proceeding as we did in Proposition 3 with  $R(\rho, \nu)$ , we compute that

$$\begin{aligned} 0 = F_2(s, \rho) &= (100s^2 - 340s + 64)\rho^6 + (-100s^2 + 420s + 400)\rho^5 \\ &\quad + (25s^2 - 250s + 625)\rho^4 + (-2500s^2 + 6500s - 2000)\rho^3 \\ &\quad + (1250s^2 - 5000s - 6250)\rho^2 + 15625s^2 - 31250s + 15625. \end{aligned}$$

Next, we compute the resultant

$$\text{Res}_r(F(r, \rho), F(r, a\rho)) = 256\rho^2(a - 1)^2 A,$$

where

$$A = (a^{10} + 2a^9 + a^8)\rho^{18} + \dots + 540000$$

is a polynomial in  $a$  and  $\rho$ . Similarly, we compute

$$\text{Res}_s(F_2(s, \rho), F_2(s, a\rho)) = 160000\rho^4(a-1)^2B,$$

where

$$B = 21600a^{10}\rho^{18} + \dots + 152587890625$$

is also a polynomial in  $a$  and  $\rho$ . Now we compute

$$\begin{aligned} \text{Res}_a(A, B) &= \rho^{80}(\rho^2 + 2\rho + 5)^{20}(11\rho^4 - 14\rho^3 - 86\rho^2 - 70\rho + 275) \\ &\quad \times (11\rho^8 - 164\rho^7 + 829\rho^6 - 1980\rho^5 + 3685\rho^4 - 9900\rho^3 + 20725\rho^2 \\ &\quad - 20500\rho + 6875) \\ &\quad \times (2488869\rho^8 - 9142380\rho^7 - 12557677\rho^6 + 46638270\rho^5 - 3856021\rho^4 \\ &\quad + 233191350\rho^3 - 313941925\rho^2 - 1142797500\rho + 1555543125) \\ &\quad \times P, \end{aligned}$$

where  $P$  is a product of 7 primitive, irreducible polynomials of degrees 12, 20, 24:

$$\begin{aligned} P &= (6543999\rho^{12} - 23871960\rho^{11} - 93034852\rho^{10} + 328442160\rho^9 + 296712789\rho^8 \\ &\quad + 1001170920\rho^7 - 9112922960\rho^6 + 5005854600\rho^5 + 7417819725\rho^4 + 41055270000\rho^3 \\ &\quad - 58146782500\rho^2 - 74599875000\rho + 102249984375) \\ &\quad \times (47989326\rho^{20} - 687672216\rho^{19} + \dots + 468645761718750) \\ &\quad \times (18251706\rho^{20} - 636597576\rho^{19} + \dots + 178239316406250) \\ &\quad \times (195570225\rho^{20} - 944115600\rho^{19} + \dots + 1909865478515625) \\ &\quad \times (1363919525\rho^{20} - 15974832650\rho^{19} + \dots + 13319526611328125) \\ &\quad \times (125164944\rho^{24} - 1106740704\rho^{23} + \dots + 30557847656250000) \\ &\quad \times (872908496\rho^{24} - 5290763616\rho^{23} + \dots + 213112425781250000). \end{aligned}$$

For each factor of this resultant (with degree at least 4), the leading coefficient does not divide the next coefficient, from which we conclude that  $\rho$  cannot be a root of any of these factors. It follows that  $\text{Res}_a(A, B) \neq 0$ , and therefore at least one of  $A$  or  $B$  is nonzero, giving that  $a = 1$ . Hence,  $\psi = 1$  in  $\text{Gal}(\Sigma/K)$ .

A similar calculation works if  $\nu^\sigma = -\nu$ . In this case, we have that

$$t = R(\rho^\sigma, -\nu) = R\left(\frac{5}{\rho}, -\nu\right) = \frac{(3\rho + 5\nu - 10)^2\rho^2}{(5(-5 + 2\rho))^2(\rho^2 + 2\rho + 5)}.$$

This leads as in Proposition 3 to the equation

$$\begin{aligned} 0 &= F_3(t, \rho) = (100t^2 - 340t + 64)\rho^6 + (-100t^2 + 420t + 400)\rho^5 \\ &\quad + (25t^2 - 250t + 625)\rho^4 + (-2500t^2 + 6500t - 2000)\rho^3 \\ &\quad + (1250t^2 - 5000t - 6250)\rho^2 + 15625t^2 - 31250t + 15625, \end{aligned}$$

which is the same as the equation  $0 = F_2(s, \rho)$ . This case also leads to  $a = 1$  and  $\psi = 1$ .

The only other alternative is that  $\psi = \sigma_{\wp_5}$ , and in this case the same calculation as in Case 1 shows that  $\psi = 1$ . This completes the proof of (10.14) and shows that Theorem 16 also holds in the ramified case, when  $(5) = \wp_5^2$  in  $K$ .

In connection with the result of this section, I put forward the following conjecture.

**Conjecture 3.** *Assume  $5 \mid d_K$ ,  $(5) = \wp_5^2$  and  $1 \neq \psi \in \text{Gal}(\Sigma/K)$ . If  $\rho^\psi = a\rho$ , then  $a \not\equiv 1 \pmod{\wp_5}$  in  $\Sigma = K_1$ .*

It can be shown in the situation of this conjecture that  $a = \varepsilon^2$  is a square in  $\Sigma$ . The congruence just after equation (10.14) shows that Sugawara's conjecture for this case would also follow immediately from Conjecture 3.

## 11 $E_{12}$ and the case $\mathfrak{m} = \wp_2^2 \wp_3$ .

The Tate normal form for a point of order 12 can be given in the form

$$\begin{aligned} E_{12}(t) : Y^2 + \frac{3t^4 + 4t^3 - 2t^2 + 4t - 1}{(t-1)^3}XY - \frac{t(t+1)(t^2+1)(3t^2+1)}{(t-1)^4}Y \\ = X^3 - \frac{t(t+1)(t^2+1)(3t^2+1)}{(t-1)^4}X^2. \end{aligned}$$

Denote the nontrivial coefficients in this equation by

$$a_1 = \frac{3t^4 + 4t^3 - 2t^2 + 4t - 1}{(t-1)^3}, \quad a_3 = a_2 = -\frac{t(t+1)(t^2+1)(3t^2+1)}{(t-1)^4}.$$

For the curve  $E_{12}(t)$  we have

$$\begin{aligned} g_2 &= \frac{(3t^4 + 6t^2 - 1)(3t^{12} + 234t^{10} + 249t^8 + 60t^6 - 27t^4 - 6t^2 - 1)}{12(t-1)^{12}}, \\ g_3 &= \frac{(3t^8 + 24t^6 + 6t^4 - 1)}{216(t-1)^{18}} \\ &\quad \times (9t^{16} - 1584t^{14} - 3996t^{12} - 3168t^{10} + 30t^8 + 528t^6 - 12t^4 + 1), \\ \Delta &= \frac{(t+1)^{12}t^6(t^2+1)^3(3t^2+1)^4(3t^2-1)}{(t-1)^{24}}. \end{aligned}$$

Its  $j$ -invariant is

$$j(E_{12}(t)) = \frac{(3t^4 + 6t^2 - 1)^3(3t^{12} + 234t^{10} + 249t^8 + 60t^6 - 27t^4 - 6t^2 - 1)^3}{t^6(t-1)^{12}(t^2+1)^3(3t^2+1)^4(3t^2-1)}.$$

Note that  $j_{12}(t) = j(E_{12}(t))$  is invariant under the map  $t^2 \rightarrow A(-t^2)$ , where

$$A(x) = \frac{x-1}{3x+1}.$$

If  $E_{12}(t)$  has complex multiplication by  $R_K$ , then  $j_{12}(t)$  is an algebraic integer, and putting  $t = 1/t'$  shows that  $t' = 1/t$  is an algebraic integer.

Using the standard formulas, the point  $P = (0, 0)$  satisfies

$$X(2P) = \frac{-b_8}{b_6} = -\frac{a_2 a_3^2}{a_3^2} = \frac{t(t+1)(t^2+1)(3t^2+1)}{(t-1)^4},$$

with corresponding  $Y$ -coordinate

$$Y(2P) = -\frac{t^2(t^2+1)(t+1)^2(3t^2+1)^2}{(t-1)^7}.$$

The point

$$2P = \left( \frac{t(t+1)(t^2+1)(3t^2+1)}{(t-1)^4}, -\frac{t^2(t^2+1)(t+1)^2(3t^2+1)^2}{(t-1)^7} \right)$$

has order 6. Furthermore, the point

$$4P = \left( \frac{(t^2+1)t(t+1)}{(t-1)^2}, -\frac{2t^2(t+1)^2(t^2+1)^2}{(t-1)^5} \right)$$

has order 3. Hence

$$6P = 2P + 4P = \left( \frac{(t+1)(3t^2+1)}{4(t-1)}, -\frac{(t+1)^2(3t^2+1)^2}{8(t-1)^3} \right)$$

has order 2. We also have the points

$$3P = P + 2P = \left( -\frac{t(t+1)(3t^2+1)}{(t-1)^3}, \frac{t^2(t+1)^2(3t^2+1)}{(t-1)^4} \right),$$

$$5P = P + 4P = \left( -\frac{2t(t+1)(t^2+1)(3t^2+1)}{(t-1)^5}, \frac{t(t+1)^2(t^2+1)(3t^2+1)^2}{(t-1)^7} \right),$$

which have order 4 and 12, respectively.

The  $\tau$ -invariant for  $\mathfrak{m} = \wp_2^2 \wp_3$  and a given point  $Q$  is given by

$$\tau(\mathfrak{k}^*) = -2^7 3^5 \frac{g_2 g_3}{\Delta} \left( X(Q) + \frac{a_1^2 + 4a_2}{12} \right).$$

Let  $\tau_k$  denote the  $\tau$ -invariant for the point  $kP$ .

**Lemma 7.** *Assume that  $E_{12}(t)$  has complex multiplication by  $R_K$ , where  $\wp_2, \wp_3$  are first degree prime divisors of 2 and 3 in  $R_K$ , and  $P = (0, 0)$  is a primitive  $\mathfrak{a}$ -division point on  $E_{12}(t)$  for the ideal  $\mathfrak{a} = \wp_2^2 \wp_3$ . The invariants  $\tau_2, \tau_3, \tau_4, \tau_6$  lie in  $\Sigma$ , and  $t$  generates  $K_{\wp_2^2 \wp_3}$  over  $\Sigma$ . Moreover, the invariants  $\tau_i$ , for  $1 \leq i \leq 6$  are all algebraic integers.*



*Proof.* For the ideal  $\mathfrak{a} = \wp_2^2 \wp_3$ , there are always  $\mathfrak{a}$ -division points on an elliptic curve  $E$  with  $R_K$  as its multiplier ring. This is because, by [4, p. 42], the number of primitive (“echten”)  $\mathfrak{a}$ -division points is

$$\begin{aligned} & \sum_{\mathfrak{d}|\mathfrak{a}} \mu(\mathfrak{a}/\mathfrak{d})N(\mathfrak{d}) \\ &= \mu(1)N(\wp_2^2 \wp_3) + \mu(\wp_2)N(\wp_2 \wp_3) + \mu(\wp_3)N(\wp_2^2) + \mu(\wp_2 \wp_3)N(\wp_2) \\ &= 12 - 6 - 4 + 2 = 4. \end{aligned}$$

If a point  $\tilde{P}$  is one such division point on  $E$ , then  $\tilde{P}$  has order 12 in  $E[12]$ , so there must be a value of  $t$  and an isomorphism  $E \rightarrow E_{12}(t)$  for which  $P \rightarrow P = (0, 0)$ , and  $P$  is a primitive  $\mathfrak{a}$ -division point on  $E_{12}(t)$ . (See [15, p. 250], where the assignment  $y \rightarrow u^3 y + vx + d$  in the proof of Lemma 4 should be  $y \rightarrow u^3 y + vu^2 x + d$ .) Now let  $\mathcal{K}$  be the elliptic function field defined by  $E_{12}(t)$ . Since  $N(\mathfrak{a}) = 12 = \text{ord}(P)$  and  $\mathfrak{a}P = O$ , the field  $\mathcal{K}^{\mathfrak{a}}$  is the fixed field of the translation group

$$\text{Gal}(\mathcal{K}/\mathcal{K}^{\mathfrak{a}}) = \langle \sigma_{\mathfrak{o}, \mathfrak{p}} \rangle$$

generated by the translation  $\sigma_{\mathfrak{o}, \mathfrak{p}}$  taking  $\mathfrak{o}$  to the prime divisor  $\mathfrak{p}$  of  $\mathcal{K}$  corresponding to  $P = (0, 0)$ . This gives an isogeny  $\phi : E_{12} \rightarrow E_{12}^{\mathfrak{a}}$  for which  $\ker(\phi) = \{Q \in E_{12} : \mathfrak{a}Q = O\} = \langle P \rangle$  is generated by  $P$ . (In Deuring’s terminology [3],  $\mathcal{K}^{\mathfrak{a}}$  has the same multiplier-ring  $R_K$  as  $\mathcal{K}$ , so that  $\phi : E_{12}(t) \rightarrow E_{12}^{\mathfrak{a}}$  is a Heegner point on  $X_0(12)$ .)

The following containments follow from these considerations:

$$2P \in \ker(\wp_2 \wp_3), \quad 3P \in \ker(\wp_2^2), \quad 4P \in \ker(\wp_3), \quad 6P \in \ker(\wp_2).$$

Since the conductors of each of these ideals is  $\mathfrak{f} = 1$ , it follows that  $\tau_2, \tau_3, \tau_4, \tau_6$  all lie in  $\Sigma$ . On the other hand,  $\tau_1$  and  $\tau_5$  do not lie in  $\Sigma$ , since the  $\tau$ -invariants for  $\mathfrak{m} = \wp_2^2 \wp_3$  must generate the quadratic extension  $\mathbb{K}_{\mathfrak{m}}/\Sigma$ .

Note that

$$\begin{aligned} \frac{\tau_1 - \tau_6}{\tau_4 - \tau_6} &= -\frac{(3t^2 + 1)(t - 1)}{(1 + t)^3}, \\ \frac{\tau_2 - \tau_3}{\tau_6 - \tau_3} &= \frac{4t^2}{t^2 - 1}, \\ \frac{\tau_3 - \tau_6}{\tau_3 - \tau_4} &= \frac{3t^2 + 1}{4t^2}. \end{aligned}$$

It follows from the above arguments that  $t^2 \in \Sigma$  but  $t \notin \Sigma$ , so that  $t$  is a Kummer element for  $\mathbb{K}_{\wp_2^2 \wp_3}$  over  $\Sigma$ . Hence, the nontrivial automorphism of  $\mathbb{K}_{\wp_2^2 \wp_3}/\Sigma$  is  $\rho : t \rightarrow -t$ .

Finally,  $\tau_3, \tau_4$ , and  $\tau_6$  are algebraic integers by the formulas for  $F(X, \mathfrak{k})$  in Sections 2, 3 and 5, since they correspond to points of orders 4, 3 and 2, respectively. Also,  $\tau_1, \tau_2$  and  $\tau_5$  are algebraic integers by Hasse’s results [9, eq. (35), p. 134], since they are invariants for the ideals  $\wp_2^2 \wp_3, \wp_2 \wp_3$  and  $\wp_2^2 \wp_3$ , respectively.  $\square$

**Corollary 3.** For the value of  $t$  in this Lemma,  $[K(t) : K] = [K_{\wp_2^2 \wp_3} : K] = 2h(d_K)$ .

*Proof.* This statement holds because  $j(E_{12}(t)) \in K(t)$  and  $K(j(E_{12}(t)), t) = \Sigma(t) = K_{\wp_2^2 \wp_3}$ .  $\square$

Now assume that  $\psi$  is an automorphism of  $K_{\wp_2^2 \wp_3}/K$  for which  $j(\mathfrak{t}') = j(\mathfrak{t})^\psi$  and  $\{\tau_1, \tau_5\}^\psi = \{\tau_1, \tau_5\}$ . If the fixed field of  $\langle \psi \rangle$  is  $L$ , and  $L \subseteq \Sigma$ , with  $\psi \neq 1$ , then the nontrivial automorphism  $\rho : t \rightarrow -t$  of  $K_{\wp_2^2 \wp_3}/\Sigma$  lies in  $\langle \psi \rangle$ . In that case, for some  $i$ ,  $\tau_1^{\psi^i} = \tau_1^\rho = \tau_5$ .

**Lemma 8.** In  $K_{\wp_2^2 \wp_3}$  we have  $t \cong \mathfrak{q}_3^{-1}$ , where  $\mathfrak{q}_3^2 = \wp_3$ , i.e.,  $\mathfrak{q}_3$  is the product of the prime divisors of  $K_{\wp_2^2 \wp_3}$  lying over  $\wp_3$ . The same holds if  $(3) = \wp_3^2$ .

*Proof.* First assume that  $(3) = \wp_3 \wp_3'$  in  $R_K$ . Put  $t = u/\sqrt{3}$  in  $j_{12}(t) = j(E_{12}(t))$ . This gives that

$$j_{12}(t) = \frac{(u^4 + 6u^2 - 3)^3(u^{12} + 234u^{10} + 747u^8 + 540u^6 - 729u^4 - 486u^2 - 243)^3}{(u^2 - 3)^{12}u^6(u^2 + 3)^3(u^2 + 1)^4(u^2 - 1)}.$$

Hence,  $u$  is an algebraic integer, so that  $u^2 = 3t^2$  is an algebraic integer. This shows that the denominator of  $t$  is divisible at most by prime divisors of  $(3) = \mathfrak{q}_3^2 \wp_3'$ . It follows that no prime divisor of  $\wp_3'$  can divide this denominator, and furthermore, at most the first power of any prime divisor of  $\mathfrak{q}_3$  can divide the denominator. If  $\sigma \in \text{Gal}(K_{\wp_2^2 \wp_3}/K)$ , then  $t^\sigma = t\alpha$ , for some  $\alpha \in \Sigma$ , since  $t^\sigma$  is also a Kummer element for  $K_{\wp_2^2 \wp_3}/\Sigma$ . But since the reciprocals of  $t$  and  $t^\sigma$  are algebraic integers, the numerators in their ideal factorizations are both 1, so that the prime divisors of  $t^\sigma/t = \alpha$  occur in the numerator and denominator at most to the first power. I claim that at least one prime ideal divisor of  $\mathfrak{q}_3$  divides the denominator of  $t$ . If not,  $t^2 \in \Sigma$  would be a unit and the prime divisors of  $\wp_3$  would not be ramified in  $\Sigma(t)/\Sigma$ , which is false. But since  $\alpha \in \Sigma$  is only divisible by (ramified) prime divisors of  $\wp_3$ , the ideal  $(\alpha)$  must be a square in the group of ideals in  $K_{\wp_2^2 \wp_3}$ . This can only be the case if  $(t^\sigma) = (t)$ . Since the prime divisors of  $\mathfrak{q}_3$  are conjugate to each other over  $K$ , this shows that all of them must divide the denominator of  $t$ . If  $(3) = \wp_3^2$ , then the denominator of  $t^2$  is divisible by at most the square of a prime divisor of  $\wp_3$  in  $\Sigma$ . But if the square of  $\mathfrak{q}$  divides the denominator, then  $\mathfrak{q}$  would not be ramified in  $K_{\wp_2^2 \wp_3}/\Sigma$ , since it divides an odd prime. Hence, exactly the first power of  $\mathfrak{q}$  divides the denominator. The rest of the argument is the same. This proves the lemma.  $\square$

**Corollary 4.** If  $\sigma \in \text{Gal}(K_{\wp_2^2 \wp_3}/K)$ , then  $t^\sigma = \alpha t$ , where  $\alpha$  is a unit in  $\Sigma$ .

Now we use the fact that both

$$\begin{aligned}\tau_1 + \tau_5 &= \frac{2(3t^4 + 6t^2 - 1)(3t^{12} + 234t^{10} + 249t^8 + 60t^6 - 27t^4 - 6t^2 - 1)}{(t^2 - 1)^{12}t^6(t^2 + 1)^3(3t^2 + 1)^4(3t^2 - 1)} \\ &\quad \times (3t^8 + 24t^6 + 6t^4 - 1)(3t^8 - 42t^4 - 24t^2 - 1) \\ &\quad \times (9t^{16} - 1584t^{14} - 3996t^{12} - 3168t^{10} + 30t^8 + 528t^6 - 12t^4 + 1) \\ &= \frac{N_1(t)}{D_1(t)}\end{aligned}$$

and

$$\begin{aligned}\frac{\tau_1 + \tau_5}{\tau_1 - \tau_5} &= -\frac{3t^8 - 42t^4 - 24t^2 - 1}{12t(t^2 - 1)(3t^2 + 1)(t^2 + 1)} \\ &= \frac{N_2(t)}{D_2(t)}\end{aligned}$$

are invariant (up to sign) under  $\psi : t \rightarrow \alpha t$ . Assuming first that  $\left(\frac{N_2(t)}{D_2(t)}\right)^\psi = \frac{N_2(t)}{D_2(t)}$ , we compute the resultant

$$\begin{aligned}R_t(a) &= \text{Res}_t \left( N_1(\alpha t)D_1(t) - N_1(t)D_1(\alpha t), \frac{1}{t}(N_2(\alpha t)D_2(t) - N_2(t)D_2(\alpha t)) \right) \\ &= 2^{572}3^{170}(a-1)^{108}(a+1)^{50}a^{362}(a^8 - 68a^6 - 570a^4 - 68a^2 + 1)^2 \\ &\quad \times (a^8 - 24a^7 + 28a^6 - 168a^5 + 582a^4 - 168a^3 + 28a^2 - 24a + 1)^2 \\ &\quad \times (a^{16} + 72a^{14} - 1412a^{12} + 3960a^{10} + 11142a^8 + 3960a^6 - 1412a^4 + 72a^2 + 1)^2 \\ &\quad \times (3a^2 - 1)^6(a^2 - 3)^6(a^2 + 3)^8(3a^2 + 1)^8(a^2 + 1)^{12} \times R_{192}(a)^2,\end{aligned}$$

where  $R_{192}(a) \in \mathbb{Z}[a]$  is an irreducible polynomial of degree 192 with leading coefficient 11664 =  $2^4 \cdot 3^6$  and coefficient of  $a$  equal to  $-2970432 = -2^6 \cdot 3^5 \cdot 191$ . It follows that the unit  $\alpha$  is a root of  $R_t(a)$ , but cannot be a root of  $R_{192}(a)$  or of any of the factors in the last line of the resultant formula, other than  $a^2 + 1$ . The three nontrivial factors (of degrees 8, 8 and 16) are symmetric in  $a$ , and in these cases  $x = \alpha + 1/\alpha$  satisfies the polynomial, respectively, given by

$$\begin{aligned}A_1(x) &= x^4 - 72x^2 - 432, \\ A_2(x) &= x^4 - 24x^3 + 24x^2 - 96x + 528, \\ A_3(x) &= x^8 + 64x^6 - 1824x^4 + 10240x^2 + 256.\end{aligned}$$

The discriminants of  $A_1(x)$  and  $A_2(x)$  are

$$\text{disc}(A_1(x)) = -2^4 \cdot 3^9, \quad \text{disc}(A_2(x)) = -2^{36} \cdot 3^3.$$

Since  $\alpha + \alpha^{-1}$  lies in  $\Sigma$ , which is normal over  $\mathbb{Q}$ , it would follow that  $\sqrt{-3} \in \Sigma$  if  $\alpha + \alpha^{-1}$  is a root of  $A_1(x)$  or  $A_2(x)$ . But this is impossible in the case that

(3) =  $\wp_3\wp_3'$  is unramified in  $K$ . The same argument holds for  $(\alpha + \alpha^{-1})^2$  and  $A_3(x) = f_3(x^2)$ , where  $\text{disc}(f_3(x)) = 2^{40} \cdot 3^7 \cdot 11^2$ .

If (3) =  $\wp_3^2$ , then the Galois groups of  $A_1(x)$  and  $A_2(x)$  are both  $D_4$ , and

$$\text{Gal}(A_3(x)/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^3 \rtimes D_4$$

is a semi-direct product of order 64. The last is impossible, since the normal closure of  $\mathbb{Q}(\alpha + \alpha^{-1})/\mathbb{Q}$  is abelian over the quadratic field  $K$ , and is thus at most quadratic over the root field  $\mathbb{Q}(\alpha + \alpha^{-1})$ . Hence,  $\alpha + \alpha^{-1}$  cannot be a root of  $A_3(x)$ . If it were a root of  $A_1(x)$  or  $A_2(x)$ , then since the normal closure  $L$  of  $\mathbb{Q}(\alpha + \alpha^{-1}) \subseteq \Sigma$  has Galois group  $D_4$ , it would have to contain the field  $K$ ; otherwise  $L$  and  $K$  would be linearly disjoint and  $\text{Gal}(LK/K) \cong \text{Gal}(L/\mathbb{Q})$  would not be abelian. In that case only 2 and 3 can divide the discriminant of  $K$ . The only possibility is then  $K = \mathbb{Q}(\sqrt{-6})$ , with discriminant  $d_K = -24$ . (This because  $\mathbb{Q}(\sqrt{-3})$  is excluded and  $\mathbb{Q}(\sqrt{-2})$  has class number 1.) In this case we have

$$H_{-24}(X) = X^2 - 4834944X + 14670139392$$

and from the formula for  $j(E_{12}(t))$  and  $[\mathbb{K}_{\wp_2^2\wp_3} : \mathbb{Q}] = 8$  we find that the minimal polynomial of  $t$  can only be

$$f(t) = 9t^8 - 36t^6 - 18t^4 + 12t^2 + 1.$$

But then

$$\begin{aligned} \text{Res}_y(f(xy), f(y)) &= 3^{16}(x^8 + 60x^6 + 134x^4 + 60x^2 + 1)^2 \\ &\quad \times (x^4 - 10x^2 + 1)^4(x^4 + 6x^2 + 1)^4(x-1)^8(x+1)^8. \end{aligned}$$

This shows that  $\alpha$  cannot be a root of either of the 8-th degree polynomials dividing  $R_t(a)$ .

Now suppose that  $\alpha$  is a root of  $a^2 + 1 = 0$ . Then  $\alpha = \pm i$  and  $\psi(t) = \alpha t$  implies that  $\psi(t^2) = -t^2$ . Now computing  $\frac{N_2(\alpha t)}{D_2(\alpha t)}$  and setting it equal to  $\pm \frac{N_2(t)}{D_2(t)}$ , we find that  $t$  is a root of the 20-th degree polynomial

$$\begin{aligned} P_1(t) &= (-9t^{10} + 3it^8 + 126t^6 + 30it^4 - 21t^2 - i) \\ &\quad \times (9t^{10} + 3it^8 - 126t^6 + 30it^4 + 21t^2 - i) \\ &= -(81t^{20} - 2259t^{16} + 16434t^{12} - 4398t^8 + 381t^4 + 1) = -P_2(t^4). \end{aligned}$$

But  $\text{Gal}(P_2(t)/\mathbb{Q}) \cong S_5$  is not solvable, showing that this is impossible. Hence  $\alpha$  cannot be a root of  $a^2 + 1$ .

On the other hand, if  $\left(\frac{N_2(t)}{D_2(t)}\right)^\psi = -\frac{N_2(t)}{D_2(t)}$ , then

$$\text{Res}_t \left( N_1(at)D_1(t) - N_1(t)D_1(at), \frac{1}{t}(N_2(at)D_2(t) + N_2(t)D_2(at)) \right) = R_t(-a)$$

is obtained by replacing  $a$  by  $-a$  in the above resultant, and the argument is the same.

Hence, whether 3 is ramified in  $K$  or not, the only possibilities are  $\alpha = \pm 1$ , in which case  $\psi : t \rightarrow \pm t$  fixes  $L = \Sigma$ . Hence, we find that  $j(\mathfrak{t}') = j(\mathfrak{t})^\psi = j(\mathfrak{t})$ . Thus, Sugawara's conjecture holds for these fields and  $\mathfrak{m} = \wp_2^2 \wp_3$ .

**Theorem 17.** *If  $\mathfrak{m} = \wp_2^2 \wp_3$ , where  $\wp_2, \wp_3$  are first degree prime ideals in  $K$ , then  $K(\tau_1) = \Sigma_{\wp_2^2 \wp_3}$  and Sugawara's conjecture holds for  $K$  and  $\mathfrak{m}$ .*

This completes the proof of Sugawara's Conjecture for all ideals  $\mathfrak{m}$  of  $K$  listed in (1.4). All other ideals satisfy one of Sugawara's conditions (1.1)-(1.3), and for these Sugawara's arguments in [22, 23] apply. This completes the proof of the Main Theorem.

## References

- [1] S.J. Akkarapakam, P. Morton, Periodic points of algebraic functions related to a continued fraction of Ramanujan, arXiv:2210.00659v2, New York J. Math. **30** (2024), 783-827.
- [2] S.J. Akkarapakam, P. Morton, On Ramanujan's cubic continued fraction, arXiv:2311.06591v2, submitted.
- [3] M. Deuring, Die Typen der Multiplikatorenringe elliptischer Funktionenkörper, Abh. Math. Sem. Hamb. **14** (1941), 197-272.
- [4] M. Deuring, Algebraische Begründung der komplexen Multiplikation, Abh. Math. Sem. Hamb. **16** (1947), 32-47.
- [5] M. Deuring, *Die Klassenkörper der komplexen Multiplikation*, Enzyklopädie der Mathematischen Wissenschaften I2, Vol. 23 (B. G. Teubner, 1958).
- [6] G. Frei, F. Lemmermeyer, P. J. Roquette, eds., *Emil Artin and Helmut Hasse: The Correspondence 1923-1958*, Springer, 2014.
- [7] R. Fueter, Abelsche Gleichungen in quadratisch-imaginären Zahlkörpern, Math. Annalen **75** (1914).
- [8] R. Fueter, *Vorlesungen über die singulären Moduln und die komplexe Multiplikation der elliptischen Funktionen, erster und zweiter Teil*, B. G. Teubner, Leipzig, 1924 and 1927.
- [9] Hasse, H. Neue Begründung der komplexen Multiplikation. I. Einordnung in die allgemeine Klassenkörpertheorie, J. reine angew. Math. **157** (1927), 115-139; paper 33 in Mathematische Abhandlungen, Bd. 2, Walter de Gruyter, Berlin, 1975, pp. 3-27.

- [10] Hasse, H. Neue Begründung der komplexen Multiplikation. II. Aufbau ohne Benützung der allgemeinen Klassenkörpertheorie, J. reine angew. Math. **165** (1931), 64-88; paper 34 in *Mathematische Abhandlungen*, Bd. 2, Walter de Gruyter, Berlin, 1975, pp. 28-52.
- [11] J. K. Koo, D. H. Shin, D. S. Yoon, Generation of ray class fields modulo 2, 3, 4 or 6 by using the Weber function, J. Korean Math. Soc. **55** (2018), 343-372.
- [12] H. Y. Jung, J. K. Koo, D. H. Shin, On a problem of Hasse and Ramachandra, *Open Mathematics* **17** (2019), 131-140.
- [13] H. Y. Jung, J. K. Koo, D. H. Shin, Class fields generated by coordinates of elliptic curves, *Open Mathematics* **20** (2022), 1145-1158.
- [14] R. Lynch, P. Morton, The quartic Fermat equation in Hilbert class fields of imaginary quadratic fields, *Int. J. of Number Theory* **11** (2015), 1961-2017.
- [15] P. Morton, Explicit identities for invariants of elliptic curves, *J. Number Theory* **120** (2006), 234-271.
- [16] P. Morton, The cubic Fermat equation and complex multiplication on the Deuring normal form, *Ramanujan J.* **25** (2011), 247-275.
- [17] P. Morton, Solutions of the cubic Fermat equation in ring class fields of imaginary quadratic fields (as periodic points of a 3-adic algebraic function), *Int. J. Number Theory* **12** (2016), 853-902.
- [18] P. Morton, Solutions of diophantine equations as periodic points of  $p$ -adic algebraic functions, II: the Rogers-Ramanujan continued fraction, *New York J. Math.* **25** (2019), 1178-1213.
- [19] P. Morton, The Hasse invariant of the Tate normal form  $E_5$  and the class number of  $\mathbb{Q}(\sqrt{-5l})$ , *J. Number Theory* **227** (2021), 94-143.
- [20] K. Ramachandra, Some applications of Kronecker's limit formulas, *Annals of Math.* **80** (1964), 104-148.
- [21] J. H. Silverman, *Advanced Topics in the Arithmetic of Elliptic Curves*, Graduate Texts in Mathematics, Vol. 151, Springer, 1994.
- [22] M. Sugawara, Zur komplexen Multiplikation, I, J. reine angew. Math. **174** (1936), 189-191.
- [23] M. Sugawara, Zur komplexen Multiplikation, II, J. reine angew. Math. **175** (1936), 65-68.
- [24] T. Takagi, Über eine Theorie des relativ Abel'schen Zahlkörpers, *Journal of the College of Science, Imperial Univ. of Tokyo* **41** (1920), 1-133. In *Collected Papers*, 2nd edition, Springer, 1990, 73-166.

- [25] B. L. van der Waerden, *Modern Algebra*, vol. 1, 2nd edition, Frederick Ungar Publishing Co., New York.
- [26] H. Weber, *Lehrbuch der Algebra, III: Elliptische Funktionen und algebraische Zahlen*, 2nd edition, Friedrich Vieweg und Sohn, Braunschweig, 1908.

Department of Mathematical Sciences, Indiana University – Purdue University  
at Indianapolis (Soon to be renamed Indiana University at Indianapolis)

402 N. Blackford St., Indianapolis, Indiana, 46202

*e-mail:* pmorton@iu.edu