

# Precompactness in matrix weighted Bourgain-Morrey spaces

Tengfei Bai and Jingshi Xu

*Hainan Normal University and Guilin University of Electronic Technology*

**Abstract.** In this paper, we introduce matrix weighted Bourgain-Morrey spaces and obtain two sufficient conditions for precompact sets in matrix weighted Bourgain-Morrey spaces. We prove that the dyadic average operator is bounded on some matrix weighted Bourgain-Morrey spaces. With this result, we obtain the necessity for precompact sets in some matrix weighted Bourgain-Morrey spaces.

## 1. Introduction

In 1931, Kolmogorov [20] first discovered the characterization of precompact sets in  $L^p([0, 1])$  for  $p \in (1, \infty)$ . After that, there are many criteria for compactness of sets in Lebesgue spaces, which are called the Kolmogorov-Riesz compactness theorems on the Lebesgue spaces. More details and the history, we refer the reader to [14].

Inspired by [14], Clop-Cruz [7] obtained a compactness criterion in scalar weighted Lebesgue spaces  $L^p(\omega)$  for  $1 < p < \infty$  with a scalar weight  $\omega$  in Muckenhoupt class  $A_p$ . In [13], Guo and Zhao improved the result in [7] and obtained a compactness criterion in  $L^p(\omega)$  for  $p \in (0, \infty)$  with locally integrable weight  $\omega$ . In [21], Liu, Yang and Zhuo proved the Kolmogorov-Riesz compactness theorem in matrix weighted Lebesgue spaces  $L^p(W)$  with  $1 < p < \infty$ .

The theory of matrix weighted function spaces goes back to [32]. Indeed, in 1958, Wiener and Masani [32, Section 4] studied the matrix weighted  $L^2(W)$  for the prediction theory for multivariate stochastic processes. In [29], Treil and Volberg introduced matrix class  $\mathcal{A}_2$ . Nazarov and Treil [23] and Volberg [30] extended  $\mathcal{A}_2$  to  $\mathcal{A}_p$  with  $p \in (1, \infty)$ . In [11], Goldberg showed that the matrix  $\mathcal{A}_p$  condition leads to  $L^p$  boundedness of a Hardy-Littlewood maximal function and obtained the boundedness of matrix weighted singular integral operators in Lebesgue spaces  $L^p$ ,  $1 < p < \infty$ . In [24, 25, 26], Roudenko introduced the matrix-weighted homogeneous Besov spaces  $\dot{B}_p^{s,q}(W)$  and matrix-weighted sequence Besov spaces  $\dot{b}_p^{s,q}(W)$  and showed their equivalence via  $\varphi$ -transform and wavelets. In [9], Frazier and Roudenko introduced the

---

*Mathematics Subject Classification:* 46B50, 46E40, 42B35, 46E30.

*Key words and phrases:* Kolmogorov-Riesz theorem, matrix weight, precompactness, Bourgain-Morrey space.

matrix class  $\mathcal{A}_p$ , ( $0 < p \leq 1$ ) and studied the continuous and discrete matrix-weighted Besov spaces  $\dot{B}_p^{s,q}(W)$  and  $\dot{b}_p^{s,q}(W)$  with  $0 < p \leq 1$ . In [10], Frazier and Roudenko introduced the homogeneous matrix-weighted Triebel-Lizorkin spaces  $\dot{F}_p^{s,q}(W)$  for  $s \in \mathbb{R}$ ,  $0 < p < \infty$ ,  $0 < q \leq \infty$  and obtained the Littlewood-Paley characterizations of matrix-weighted Lebesgue spaces  $L^p(W)$  and matrix-weighted Sobolev space  $L_k^p(W)$  for  $k \in \mathbb{N}$ ,  $1 < p < \infty$ . In [30], Volberg also introduced an analogue condition for the matrix class  $\mathcal{A}_\infty$ . In [3, 4, 5], Bu, Hytönen, Yang, Yuan studied the matrix weighted Besov-type and Triebel-Lizorkin-type spaces. Specifically, they introduced a new concept of the  $\mathcal{A}_p$ -dimension  $\tilde{d}$ , which is useful to the proof of main results of this paper. In [2], Bu, Hytönen, Yang, Yuan obtained several new characterizations of  $\mathcal{A}_{p,\infty}$ -matrix weights. For many other results on the matrix class  $\mathcal{A}_p$  and matrix weighted function spaces, we refer the reader to [6, 8, 9, 10, 11, 24, 25, 31].

Bourgain [1] introduced a special case of Bourgain-Morrey spaces to study the Stein-Tomas (Strichartz) estimate. In [16], Hatano, Nogayama, Sawano, and Hakim researched the Bourgain-Morrey spaces from the viewpoints of harmonic analysis and functional analysis. In [19], Hu, Li and Yang introduced the Triebel-Lizorkin-Bourgain-Morrey spaces which connect Bourgain-Morrey spaces and global Morrey spaces.

Motivated by above literature, we will introduce matrix weighted Bourgain-Morrey spaces and research precompact sets in these spaces. The paper is organized as follows. In Section 2, dyadic cubes, the matrix class  $\mathcal{A}_p$ ,  $\mathcal{A}_p$ -dimension  $\tilde{d}$ , matrix weighted Bourgain-Morrey spaces are given. The first result lies in Section 3. Specifically, a sufficient condition for totally bounded set in matrix weighted spaces  $M_p^{t,r}(W)$  with  $1 \leq p < r < \infty$  or  $1 \leq p \leq t < r = \infty$  is obtained in Theorem 3.2. As a application, we get a criterion for totally bounded set in degenerate Bourgain-Morrey spaces with matrix weight. The second result (Theorem 4.1) is replacing the translation operator by the average operator. Note that the translation operator is not bounded on  $L^p(W)$  and  $M_p^{t,r}(W)$  in general. We prove that dyadic average operator is bounded on matrix weighted Bourgain-Morrey spaces with some conditions in Theorem 4.5. Using this result, we obtain the Kolmogorov-Riesz compactness theorem in matrix weighted Bourgain-Morrey spaces.

Throughout this paper, we let  $c, C$  denote constants that are independent of the main parameters involved but whose value may differ from line to line. Let  $\mathbb{N} = \{1, 2, 3, \dots\}$  and  $\mathbb{N}_0 = \{0, 1, 2, 3, \dots\}$ . Let  $\mathbb{Z}$  be the set of integers. Let  $\chi_E$  be the characteristic function of the set  $E \subset \mathbb{R}^n$ . By  $A \lesssim B$  we mean that  $A \leq CB$  with some positive constant  $C$  independent of appropriate quantities. By  $A \approx B$ , we mean that  $A \lesssim B$  and  $B \lesssim A$ .

## 2. Preliminaries

For  $j \in \mathbb{Z}$ ,  $m \in \mathbb{Z}^n$ , let  $Q_{j,m} := \prod_{i=1}^n [2^{-j}m_i, 2^{-j}(m_i + 1))$ . For a cube  $Q$ ,  $\ell(Q)$  stands for the length of cube  $Q$ . We denote by  $\mathcal{D}$  the family of all dyadic cubes in  $\mathbb{R}^n$ , while  $\mathcal{D}_j$  is the set of all dyadic cubes with  $\ell(Q) = 2^{-j}$ ,  $j \in \mathbb{Z}$ . Let  $x_Q$  be the left corner of  $Q \in \mathcal{D}$ . For  $\lambda > 0$ , let  $\lambda Q$  be the cube with the same center of  $Q$  and the edge length  $\lambda \ell(Q)$ . For  $k \in \mathbb{N}$ , let  $k_{\text{pa}}Q$  be the  $k$ -th dyadic parent of  $Q$ , which is the dyadic cube in  $\mathcal{D}$  satisfying  $Q \subset k_{\text{pa}}Q$  and  $\ell(k_{\text{pa}}Q) = 2^k \ell(Q)$ .

**2.1. Matrix weights** First, we recall some basic concepts and results from the theory of matrix weights.

For any  $d, n \in \mathbb{N}$ , denote by  $M_{d,n}(\mathbb{C})$  the set of all  $d \times n$  complex-valued matrices.  $M_{d,d}(\mathbb{C})$  is simply denoted by  $M_d(\mathbb{C})$ . The zero matrix in  $M_{d,n}(\mathbb{C})$  (or  $M_d(\mathbb{C})$ ) is denoted by  $O_{d,n}$  (or  $O_d$ ). Denote by  $A^*$  the conjugate transpose of  $A \in M_{d,n}(\mathbb{C})$ . A matrix  $A \in M_d(\mathbb{C})$  is called a *Hermitian matrix* if  $A = A^*$  and is called a *unitary matrix* if  $A^*A = I_d$  where  $I_d$  is the identity matrix. We denote a diagonal matrix by  $\text{diag}(\lambda_1, \dots, \lambda_d) = \text{diag}(\lambda_i)$ .

For a vector  $x \in \mathbb{C}^d$ , let  $|x| = (\sum_{i=1}^d |x_i|^2)^{1/2}$ . For  $1 \leq p < \infty$ , let  $|x|_p = (\sum_{i=1}^d |x_i|^p)^{1/p}$ . For  $p = \infty$ , let  $|x|_\infty = \max(x_1, \dots, x_d)$ . In the finite dimension space, the norms are equivalent. That is, for  $1 \leq p < q \leq \infty$ ,

$$|x|_q \leq |x|_p \leq d^{1/p} |x|_\infty \leq d^{1/p} |x|_q. \quad (2.1)$$

For  $A \in M_d(\mathbb{C})$ , let

$$\|A\| := \sup_{\vec{z} \in \mathbb{C}^d, |\vec{z}|=1} |A\vec{z}|.$$

We say that a matrix  $A \in M_d(\mathbb{C})$  is *positive definite* if, for any  $\vec{x} \in \mathbb{C}^d \setminus \{\vec{0}\}$ ,  $\vec{z}^* A \vec{z} > 0$ . And a matrix  $A \in M_d(\mathbb{C})$  is called *nonnegative definite* if, for any  $\vec{x} \in \mathbb{C}^d$ ,  $\vec{z}^* A \vec{z} \geq 0$ .

From [18, Theorem 4.1.4], any nonnegative definite matrix is always Hermitian. Hence any nonnegative definite matrix is self-adjoint.

Let  $A \in M_d(\mathbb{C})$  be a positive definite matrix and have eigenvalues  $\{\lambda_i\}_{i=1}^d$ . From [18, Theorem 2.5.6(c)], there exists a unitary matrix  $U \in M_d(\mathbb{C})$  such that

$$A = U \text{diag}(\lambda_1, \dots, \lambda_d) U^*. \quad (2.2)$$

Moreover, by [18, Theorem 4.1.8], we find  $\{\lambda_i\}_{i=1}^d \subset (0, \infty)^d$ . The following definition is based on these conclusions.

**DEFINITION 2.1.** Let  $A \in M_d(\mathbb{C})$  be a positive definite matrix with positive eigenvalues  $\{\lambda_i\}_{i=1}^d$ . For any  $\alpha \in \mathbb{R}$ , define

$$A^\alpha := U \text{diag}(\lambda_1^\alpha, \dots, \lambda_m^\alpha) U^*, \quad (2.3)$$

where  $U$  is the same as in (2.2).

REMARK 2.2. From [17, p. 408], we obtain that  $A^\alpha$  is independent of the choices of the order of  $\{\lambda_i\}_{i=1}^m$  and  $U$ , and hence  $A^\alpha$  is well defined.

Now, we recall some concepts of matrix weights.

DEFINITION 2.3. A matrix-valued function  $W : \mathbb{R}^n \rightarrow M_d(\mathbb{C})$  is called a *matrix weight* if  $W$  satisfies that

- (i) for any  $x \in \mathbb{R}^n$ ,  $W(x)$  is nonnegative definite;
- (ii) for almost every  $x \in \mathbb{R}^n$ ,  $W(x)$  is invertible;
- (iii) the entries of  $W$  is locally integrable;

DEFINITION 2.4. Let  $p \in (0, \infty)$ ,  $W$  be a matrix weight. Suppose that  $E \subset \mathbb{R}^n$  is a bounded measurable set satisfying  $0 < |E| < \infty$ . Then the matrix  $A_E \in M_d(\mathbb{C})$  is called a reducing operator of order  $p$  for  $W$  if  $A_E$  is positive definite and, for any  $\vec{z} \in \mathbb{C}^d$ ,

$$|A_E \vec{z}| \approx \left( \frac{1}{|E|} \int_E |W^{1/p}(x) \vec{z}|^p dx \right)^{1/p}, \quad (2.4)$$

where the positive equivalence constants depend only on  $d$  and  $p$ .

Next we recall the concepts of scalar weight class  $A_p$  (see [12, Definitions 7.1.1, 7.1.3]) and matrix weight class  $\mathcal{A}_p$  (see [26] for  $1 < p < \infty$ , [9] for  $0 < p \leq 1$ ).

DEFINITION 2.5. A weight  $\omega$  is a nonnegative locally integrable function on  $\mathbb{R}^n$  such that  $0 < \omega(x) < \infty$  for almost all  $x \in \mathbb{R}^n$ .

A weight  $\omega$  is called an  $A_1$  weight if  $\mathcal{M}(\omega)(x) \leq c\omega(x)$  for almost all  $x \in \mathbb{R}^n$  where  $\mathcal{M}$  is the Hardy-Littlewood maximal operator.

For  $1 < p < \infty$ , a weight  $\omega$  is said to be of class  $A_p$  if

$$\sup_{Q \text{ cubes in } \mathbb{R}^n} \left( \frac{1}{|Q|} \int_Q \omega(x) dx \right) \left( \frac{1}{|Q|} \int_Q \omega(x)^{-1/(p-1)} dx \right)^{p-1} < \infty.$$

For  $1 < p < \infty$ , a matrix weight  $W \in \mathcal{A}_p(\mathbb{R}^n)$  if and only if

$$\sup_Q \frac{1}{|Q|} \int_Q \left( \frac{1}{|Q|} \int_Q \|W^{1/p}(x)W^{-1/p}(y)\|^{p'} dy \right)^{p/p'} dx < \infty,$$

where  $p' = p/(p-1)$  is the conjugate index of  $p$ , and the supremum is taken over all cubes  $Q \subset \mathbb{R}^n$ .

For  $0 < p \leq 1$ , a matrix weight  $W \in \mathcal{A}_p(\mathbb{R}^n)$  if and only if

$$\sup_Q \operatorname{ess\,sup}_{y \in Q} \frac{1}{|Q|} \int_Q \|W^{1/p}(x)W^{-1/p}(y)\|^p dx < \infty.$$

We write  $\mathcal{A}_p := \mathcal{A}_p(\mathbb{R}^n)$  for brevity.

Given any matrix weight  $W$  and  $0 < p < \infty$ , there exists (see e.g., [11, Proposition 1.2]) for  $p > 1$  and [9, p.1237] for  $0 < p \leq 1$ ) a sequence  $\{A_Q\}_{Q \in \mathcal{D}}$  of positive definite  $d \times d$  matrices such that

$$c_1 |A_Q \vec{y}| \leq \left( \frac{1}{|Q|} \int_Q |W^{1/p}(x) \vec{y}|^p dx \right)^{1/p} \leq c_2 |A_Q \vec{y}|,$$

with positive constants  $c_1, c_2$  independent of  $\vec{y} \in \mathbb{C}^d$  and  $Q \in \mathcal{D}$ . In this case, we call  $\{A_Q\}_{Q \in \mathcal{D}}$  a sequence of reducing operators of order  $p$  for  $W$ .

**DEFINITION 2.6.** A matrix weight  $W$  is called a doubling matrix weight of order  $p > 0$  if the scalar measures  $w_{\vec{y}}(x) = |W^{1/p}(x) \vec{y}|^p$ , for  $\vec{y} \in \mathbb{C}^d$ , are uniformly doubling: there exists  $c > 0$  such that for all cubes  $Q \subset \mathbb{R}^n$  and all  $\vec{y} \in \mathbb{C}^d$ ,

$$\int_{2Q} w_{\vec{y}}(x) dx \leq c \int_Q w_{\vec{y}}(x) dx.$$

If  $c = 2^\beta$  is the smallest constant for which this inequality holds, we say that  $\beta$  is the doubling exponent of  $W$ .

From [15, Proposition 2.10], we know that  $\beta$  is always not less than  $n$ .

In [3], Bu, Hytönen, Yang and Yuan introduced the  $\mathcal{A}_p$ -dimension of matrix weights, which will be used in Theorems 4.5 and 4.6.

**DEFINITION 2.7.** Let  $0 < p < \infty$ ,  $\tilde{d} \in \mathbb{R}$ . A matrix weight  $W$  has the  $\mathcal{A}_p$ -dimension  $\tilde{d}$ , denoted by  $W \in \mathbb{D}_{p, \tilde{d}}(\mathbb{R}^n, \mathbb{C}^n)$ , if there exists a positive constant  $C$  such that for any cube  $Q \subset \mathbb{R}^n$  and any  $i \in \mathbb{N}_0$ ,

$$\operatorname{ess\,sup}_{y \in 2^i Q} \frac{1}{|Q|} \int_Q \left\| W^{1/p}(x) W^{-1/p}(y) \right\|^p dx \leq C 2^{i \tilde{d}}, \quad \text{for } 0 < p \leq 1, \quad (2.5)$$

or,

$$\frac{1}{|Q|} \int_Q \left( \frac{1}{|2^i Q|} \int_{2^i Q} \left\| W^{1/p}(x) W^{-1/p}(y) \right\|^{p'} dy \right)^{p/p'} dx \leq C 2^{i \tilde{d}}, \quad \text{for } 1 < p < \infty,$$

where  $1/p + 1/p' = 1$ .

We denote  $\mathbb{D}_{p, \tilde{d}}(\mathbb{R}^n, \mathbb{C}^n)$  simply by  $\mathbb{D}_{p, \tilde{d}}$ .

**LEMMA 2.8** (Corollary 2.32, [3]). *Let  $0 < p < \infty$ , let  $W \in \mathcal{A}_p$  have the  $\mathcal{A}_p$ -dimension  $\tilde{d} \in [0, n)$ , and let  $\{A_Q\}_{\text{cube } Q}$  be the reducing operator of order  $p$  for  $W$ . (i) If  $1 < p < \infty$ , let  $\tilde{W} := W^{-1/(p-1)}$  (which belongs to  $\mathcal{A}_{p'}$ ) have the  $\mathcal{A}_{p'}$ -dimension*

$\tilde{d}$ . Then there exists a positive constant  $C$  such that, for any cubes  $Q$  and  $R$  of  $\mathbb{R}^n$ ,

$$\|A_Q A_R^{-1}\| \leq C \max \left( \left[ \frac{\ell(R)}{\ell(Q)} \right]^{\tilde{d}/p}, \left[ \frac{\ell(Q)}{\ell(R)} \right]^{\tilde{d}/p'} \right) \left[ 1 + \frac{|x_Q - x_R|}{\max(\ell(Q), \ell(R))} \right]^{\tilde{d}/p + \tilde{d}/p'}.$$

(ii) If  $0 < p \leq 1$ , then there exists a positive constant  $C$  such that, for any cubes  $Q$  and  $R$  of  $\mathbb{R}^n$ ,

$$\|A_Q A_R^{-1}\| \leq C \max \left( \left[ \frac{\ell(R)}{\ell(Q)} \right]^{\tilde{d}/p}, 1 \right) \left[ 1 + \frac{|x_Q - x_R|}{\max(\ell(Q), \ell(R))} \right]^{\tilde{d}/p}.$$

## 2.2. Matrix weighted Bourgain-Morrey spaces

DEFINITION 2.9. Let  $0 < p < \infty$ ,  $d \in \mathbb{N}$  and  $W : \mathbb{R}^n \rightarrow M_d(\mathbb{C})$  be a matrix weight. The space  $L_{\text{loc}}^p(W)$  collects all measurable functions  $\vec{f} : \mathbb{R}^n \rightarrow \mathbb{C}^d$  such that for each compact set  $K$ ,

$$\|\vec{f}\chi_K\|_{L^p(W)} = \left( \int_K |W^{1/p}(x)\vec{f}(x)|^p dx \right)^{1/p} < \infty.$$

Let  $\Omega \subset \mathbb{R}^n$  be an open set. The space  $L^p(W, \Omega)$  collects all measurable functions  $\vec{f} : \Omega \rightarrow \mathbb{C}^d$  such that

$$\|\vec{f}\|_{L^p(W, \Omega)} = \left( \int_{\Omega} |W^{1/p}(x)\vec{f}(x)|^p dx \right)^{1/p} < \infty.$$

DEFINITION 2.10. Let  $\mathcal{D} = \{Q_{j,k}\}_{j \in \mathbb{Z}, k \in \mathbb{Z}^n}$  be the standard dyadic system. Let  $0 < p < t < r < \infty$  or  $0 < p \leq t < r = \infty$ . Let  $W : \mathbb{R}^n \rightarrow M_d(\mathbb{C})$  be a matrix weight. Define  $M_p^{t,r}(W)$  as the set of all  $\vec{f} \in L_{\text{loc}}^p(W)$  such that

$$\|\vec{f}\|_{M_p^{t,r}(W)} := \left\| \left\{ W(Q_{j,k})^{1/t-1/p} \left( \int_{Q_{j,k}} |W^{1/p}(y)\vec{f}(y)|^p dy \right)^{1/p} \right\}_{j \in \mathbb{Z}, k \in \mathbb{Z}^n} \right\|_{\ell^r} < \infty,$$

where  $W(Q_{j,k}) = \int_{Q_{j,k}} \|W(y)\| dy$ .

Let  $\{A_Q\}_{Q \in \mathcal{D}}$  be the reducing operator of order  $p$  for  $W$ . Define  $M_p^{t,r}(\{A_Q\})$  as the set of all  $\vec{f} \in L_{\text{loc}}^p(\{W\})$  such that

$$\begin{aligned} & \|\vec{f}\|_{M_p^{t,r}(\{A_Q\})} \\ & := \left\| \left\{ (\|A_{Q_{j,k}}\|^p |Q_{j,k}|)^{1/t-1/p} \left( \int_{Q_{j,k}} |W^{1/p}(x)\vec{f}(y)|^p dy \right)^{1/p} \right\}_{j \in \mathbb{Z}, k \in \mathbb{Z}^n} \right\|_{\ell^r} < \infty. \end{aligned}$$

REMARK 2.11. Let  $\{\vec{e}_i\}_{i=1}^d$  be the standard unit basis for  $\mathbb{C}^d$ . Since  $W$  is a matrix weight, then for each  $x \in \mathbb{R}^n$ ,  $W(x)$  is a nonnegative matrix and self-adjoint. Hence we have

$$\begin{aligned} \|W(x)\| &= \|W^{2/p}(x)\|^{p/2} \approx [\text{tr}(W^{2/p}(x))]^{p/2} \\ &= \left( \sum_{i=1}^d |W^{1/p}(x)\vec{e}_i|^2 \right)^{p/2} \approx \sum_{i=1}^d |W^{1/p}(x)\vec{e}_i|^p. \end{aligned}$$

From this and (2.4), we obtain

$$\begin{aligned} W(Q_{j,k}) &= \int_{Q_{j,k}} \|W(y)\| dy \approx \int_{Q_{j,k}} \sum_{i=1}^d |W^{1/p}(x)\vec{e}_i|^p dy \\ &\approx \sum_{i=1}^d |A_{Q_{j,k}}\vec{e}_i|^p |Q_{j,k}| \approx \|A_{Q_{j,k}}\|^p |Q_{j,k}|. \end{aligned}$$

Thus  $M_p^{t,r}(W)$  is same with  $M_p^{t,r}(\{A_Q\})$  in meaning of the equivalent quasi-norms.

If  $d = 1$ ,  $W \equiv 1$ ,  $M_p^{t,r}(W)$  is the classical Bourgain-Morrey space  $M_p^{t,r}$  in [16]. We define the scalar weighted Bourgain-Morrey space  $M_p^{t,r}(\omega)$  by

$$\|f\|_{M_p^{t,r}(\omega)} := \left\| \left\{ (\omega(Q_{j,k}))^{1/t-1/p} \left( \int_{Q_{j,k}} |f(y)|^p \omega(y) dy \right)^{1/p} \right\}_{j \in \mathbb{Z}, k \in \mathbb{Z}^n} \right\|_{\ell^r} < \infty. \quad (2.6)$$

That is  $M_p^{t,r}(\omega)$  is the case  $d = 1$  of matrix weighted Bourgain-Morrey space  $M_p^{t,r}(W)$ .

The following Lemma is proved in [22, 28].

LEMMA 2.12. *Let  $1 \leq p < \infty$  and  $W : \mathbb{R}^n \rightarrow M_d(\mathbb{C})$  be a matrix weight. Let  $\Omega \subset \mathbb{R}^n$  be an open set. Then matrix weighted Lebesgue space  $L^p(W, \Omega)$  is a Banach space.*

THEOREM 2.13. *Let  $1 \leq p < \infty$  and  $W : \mathbb{R}^n \rightarrow M_d(\mathbb{C})$  be a matrix weight. Then the space  $L_{\text{loc}}^p(W)$  is complete.*

PROOF. Let  $\{\vec{f}_k\}_{k=1}^\infty$  be a Cauchy sequence in  $L_{\text{loc}}^p(W)$ . That is, for any compact set  $K$ , any  $\varepsilon > 0$ , there exists  $N > 0$  such that if  $j, k > N$ ,  $\|(\vec{f}_j - \vec{f}_k)\chi_K\|_{L^p(W)} < \varepsilon$ . Then for any compact set  $K$ ,  $\{\vec{f}_k\chi_K\}_{k=1}^\infty$  is a Cauchy sequence in  $L^p(W)$ . Since  $L^p(W)$  is complete (Lemma 2.12), there exists  $\vec{f}\chi_K$  in  $L^p(W)$  such that  $\vec{f}_k\chi_K \rightarrow \vec{f}\chi_K$ . By the Fatou lemma, we have

$$\|\vec{f}\chi_K\|_{L^p(W)} \leq \liminf_{k \rightarrow \infty} \|\vec{f}_k\chi_K\|_{L^p(W)} < \infty.$$

By the dominated convergence theorem, for  $j, k > N$ , we obtain

$$\|(\vec{f} - \vec{f}_k)\chi_K\|_{L^p(W)} = \lim_{j \rightarrow \infty} \|(\vec{f}_j - \vec{f}_k)\chi_K\|_{L^p(W)} < \varepsilon.$$

Hence  $\vec{f}_k \rightarrow \vec{f}$  in  $L^p_{\text{loc}}(W)$  and the space  $L^p_{\text{loc}}(W)$  is complete.  $\square$

In what follows, the symbol  $\hookrightarrow$  always stands for continuous embedding.

**PROPOSITION 1.** *Let  $W : \mathbb{R}^n \rightarrow M_d(\mathbb{C})$  be a matrix weight.*

(i) *If  $0 < p < t < r_1 < r_2 \leq \infty$ , then*

$$M_p^{t,r_1}(W) \hookrightarrow M_p^{t,r_2}(W).$$

(ii) *If  $0 < p_1 < p_2 < t < r < \infty$  or  $0 < p_1 < p_2 \leq t < r = \infty$ , then*

$$M_{p_2}^{t,r}(W) \hookrightarrow M_{p_1}^{t,r}(W).$$

(iii) *If  $0 < p \leq t < r = \infty$ , then*

$$L^t(W) \hookrightarrow M_p^{t,r}(W) \hookrightarrow L^p_{\text{loc}}(W).$$

**PROOF.** (i) It comes from  $\ell^{r_1} \hookrightarrow \ell^{r_2}$  since  $0 < r_1 < r_2 \leq \infty$ .

(ii) It comes from the Hölder inequality. Indeed, for each  $Q \in \mathcal{D}$ ,

$$\begin{aligned} & \left( \frac{1}{\overline{W}(Q)} \int_Q |W^{1/p_1}(y)\vec{f}(y)|^{p_1} dy \right)^{1/p_1} \\ &= \left( \frac{1}{\overline{W}(Q)} \int_Q |W^{1/p_1-1/p_2}(y)W^{1/p_2}(y)\vec{f}(y)|^{p_1} dy \right)^{1/p_1} \\ &\leq \left( \frac{1}{\overline{W}(Q)} \int_Q \|W^{1/p_1-1/p_2}(y)\|^{p_1} |W^{1/p_2}(y)\vec{f}(y)|^{p_1} dy \right)^{1/p_1} \\ &\leq \left( \frac{1}{\overline{W}(Q)} \int_Q \|W^{1/p_1-1/p_2}(y)\|^{p_1 p_2 / (p_2 - p_1)} dy \right)^{1/p_1 - 1/p_2} \\ &\quad \times \left( \frac{1}{\overline{W}(Q)} \int_Q |W^{1/p_2}(y)\vec{f}(y)|^{p_2} dy \right)^{1/p_2} \\ &= \left( \frac{1}{\overline{W}(Q)} \int_Q \|W(y)\| dy \right)^{1/p_1 - 1/p_2} \left( \frac{1}{\overline{W}(Q)} \int_Q |W^{1/p_2}(y)\vec{f}(y)|^{p_2} dy \right)^{1/p_2} \\ &= \left( \frac{1}{\overline{W}(Q)} \int_Q |W^{1/p_2}(y)\vec{f}(y)|^{p_2} dy \right)^{1/p_2}. \end{aligned}$$

Hence,

$$\|\vec{f}\|_{M_{p_1}^{t,r}(W)} \leq \|\vec{f}\|_{M_{p_2}^{t,r}(W)}.$$



Thus we prove (ii).

(iii) The first embedding comes from the fact that

$$L^t(W) = M_t^{t,\infty}(W) \hookrightarrow M_p^{t,\infty}(W).$$

For any compact set  $K \subset \mathbb{R}^n$ , there exist at most  $2^n$  dyadic cubes  $Q_j$  such that  $K \subset \cup_{j=1}^{2^n} Q_j$ . Hence

$$\begin{aligned} \|\vec{f}\chi_K\|_{L^p(W)} &\leq \sum_{j=1}^{2^n} \frac{|Q_j|^{1/t-1/p}}{|Q_j|^{1/t-1/p}} \|\vec{f}\chi_{Q_j}\|_{L^p(W)} \\ &\leq \|\vec{f}\|_{M_p^{t,\infty}(W)} \sum_{j=1}^{2^n} \frac{1}{|Q_j|^{1/t-1/p}} < \infty. \end{aligned}$$

This shows that  $M_p^{t,r}(W) \hookrightarrow L_{\text{loc}}^p(W)$ . □

**PROPOSITION 2.** *The matrix weighted Bourgain-Morrey space  $M_p^{t,r}(W)$  is a Banach space when  $1 \leq p < t < r < \infty$  or  $1 \leq p \leq t < r = \infty$ .*

**PROOF.** The argument are standard, see, for example, [27, Theorem 2.4]. We only show that  $M_p^{t,r}(W)$  is complete since others are simple.

Let  $\{\vec{f}_j\}_{j=1}^\infty$  be a Cauchy sequence in  $M_p^{t,r}(W)$ . By Proposition 1, we have that  $\{\vec{f}_j\}_{j=1}^\infty$  is also a Cauchy sequence in  $L_{\text{loc}}^p(W)$ . Since  $L_{\text{loc}}^p(W)$  is complete, there exists a vector function  $\vec{f}$  in  $L_{\text{loc}}^p(W)$  such that  $\vec{f}_j \rightarrow \vec{f}$ . By the Fatou lemma,

$$\|\vec{f}\|_{M_p^{t,r}(W)} = \|\liminf_{j \rightarrow \infty} \vec{f}_j\|_{M_p^{t,r}(W)} \leq \liminf_{j \rightarrow \infty} \|\vec{f}_j\|_{M_p^{t,r}(W)} < \infty.$$

Thus  $\vec{f} \in M_p^{t,r}(W)$ . Therefore,

$$\limsup_{j \rightarrow \infty} \|\vec{f} - \vec{f}_j\|_{M_p^{t,r}(W)} \leq \limsup_{j \rightarrow \infty} \left( \liminf_{k \rightarrow \infty} \|\vec{f}_k - \vec{f}_j\|_{M_p^{t,r}(W)} \right) = 0.$$

Thus, the sequence  $\{\vec{f}_j\}_{j=1}^\infty$  is convergent to  $\vec{f}$  in  $M_p^{t,r}(W)$ . □

Recall that a matrix weight  $W$  is almost everywhere invertible. Hereafter, we define  $w(x) := \|W^{-1}(x)\|^{-1}$  when a matrix weight  $W$  is invertible at  $x \in \mathbb{R}^n$ .

**LEMMA 2.14** (Proposition 3.2,[8]). *For a matrix weight  $W : \mathbb{R}^n \rightarrow M_d(\mathbb{C})$ , we have  $0 < w(x) \leq \|W(x)\| < \infty$  for a.e.  $x \in \mathbb{R}^n$ . Furthermore,  $W$  satisfies a two weight, degenerate ellipticity condition: for  $\xi \in \mathbb{R}^d$ ,*

$$w(x)|\xi|^p \leq |W^{1/p}(x)\xi|^p \leq \|W(x)\||\xi|^p. \quad (2.7)$$

**REMARK 2.15.** In [8, Proposition 3.2], it is assumed that  $\xi \in \mathbb{R}^d$  but it also works for  $\xi \in \mathbb{C}^d$ .

PROPOSITION 3. *Let  $1 \leq p < t < r < \infty$  or  $1 \leq p \leq t < r = \infty$ . Let  $W$  be a matrix weight and  $\vec{f}, \vec{g} \in M_p^{t,r}(W)$ . Then  $\|\vec{f} - \vec{g}\|_{M_p^{t,r}(W)} = 0$  if and only if  $\vec{f}(x) = \vec{g}(x)$  a.e..*

PROOF. Clearly, if  $\vec{f}(x) = \vec{g}(x)$  a.e., then  $\|\vec{f} - \vec{g}\|_{M_p^{t,r}(W)} = 0$ . Then we apply Lemma 2.14 to prove the converse. By the degenerate ellipticity condition (2.7), we have

$$\begin{aligned} 0 &= \|\vec{f} - \vec{g}\|_{M_p^{t,r}(W)} \\ &\geq \left\| \left\{ W(Q_{j,k})^{1/t-1/p} \left( \int_{Q_{j,k}} |\vec{f}(y) - \vec{g}(y)|^p w(y) dy \right)^{1/p} \right\}_{j \in \mathbb{Z}, k \in \mathbb{Z}^n} \right\|_{\ell^r}. \end{aligned}$$

Since  $w(y) > 0$  a.e., it follows that  $\vec{f}(y) - \vec{g}(y) = 0$  a.e.  $\square$

### 3. Sufficient conditions for precompact sets

DEFINITION 3.1. (i) Suppose that  $(X, \rho)$  is a metric space. Let  $A$  be a set of  $X$  and let  $\varepsilon > 0$ . A set  $E \subset X$  is called an  $\varepsilon$ -net for  $A$  if every point  $a \in A$ , there exists a point  $e \in E$  such that  $\rho(a, e) < \varepsilon$ .

(ii) A set  $A$  is called totally bounded if, for each  $\varepsilon > 0$ , it possesses a finite  $\varepsilon$ -net.

(iii) A subset in a topological space is precompact if its closure is compact.

It is well known that in a complete metric space, a set is precompact if and only if it is totally bounded.

THEOREM 3.2. *Let  $1 \leq p < t < r < \infty$  or  $1 \leq p \leq t < r = \infty$ . Let  $W : \mathbb{R}^n \rightarrow M_d(\mathbb{C})$  be a matrix weight. A subset  $\mathcal{F} \subset M_p^{t,r}(W)$  is totally bounded if the following conditions are valid:*

(i)  $\mathcal{F}$  uniformly vanishes at infinity, that is,

$$\lim_{R \rightarrow \infty} \sup_{\vec{f} \in \mathcal{F}} \|\vec{f} \chi_{B^c(0,R)}\|_{M_p^{t,r}(W)} = 0;$$

(ii)  $\mathcal{F}$  is equicontinuous, that is,

$$\lim_{b \rightarrow 0} \sup_{\vec{f} \in \mathcal{F}} \sup_{y \in B(0,b)} \|\vec{f} - \tau_y \vec{f}\|_{M_p^{t,r}(W)} = 0. \quad (3.1)$$

Here and what follows,  $\tau_y$  denotes the translation operator:  $\tau_y \vec{f}(x) := \vec{f}(x - y)$ .

REMARK 3.3. In this Remark, we will prove that the conditions (i) and (ii) in Theorem 3.2 together imply that the set  $\mathcal{F}$  is bounded. Indeed, choose  $b > 0$  such

that for all  $h \in \mathbb{R}^n$ ,  $|h| \leq b$ , all  $\vec{f} \in \mathcal{F}$ ,

$$\|\vec{f} - \tau_y \vec{f}\|_{M_p^{t,r}(W)} \leq 1.$$

Choose  $R > 0$  such that for all  $\vec{f} \in \mathcal{F}$ ,

$$\|\vec{f}\chi_{B^c(0,R)}\|_{M_p^{t,r}(W)} \leq 1.$$

Fix  $h$  with  $|h| = b$ . Then for all  $\vec{f} \in \mathcal{F}$ ,  $x \in \mathbb{R}^n$ , we have

$$\begin{aligned} \|\vec{f}\chi_{B(x,R)}\|_{M_p^{t,r}(W)} &\leq \|(\vec{f} - \tau_h \vec{f})\chi_{B(x,R)}\|_{M_p^{t,r}(W)} + \|\tau_h \vec{f}\chi_{B(x,R)}\|_{M_p^{t,r}(W)} \\ &= \|(\vec{f} - \tau_h \vec{f})\chi_{B(x,R)}\|_{M_p^{t,r}(W)} + \|\vec{f}\chi_{B(x+h,R)}\|_{M_p^{t,r}(W)} \\ &\leq 1 + \|\vec{f}\chi_{B(x+h,R)}\|_{M_p^{t,r}(W)}. \end{aligned}$$

Hence, by induction,

$$\|\vec{f}\chi_{B(0,R)}\|_{M_p^{t,r}(W)} \leq N + \|\vec{f}\chi_{B(Nh,R)}\|_{M_p^{t,r}(W)}.$$

Now choose  $N \geq 1$  such that  $Nb = N|h| > 2R$ . Then  $B(Nh, R) \subset B^c(0, R)$ . Hence

$$\begin{aligned} \|\vec{f}\|_{M_p^{t,r}(W)} &\leq \|\vec{f}\chi_{B(0,R)}\|_{M_p^{t,r}(W)} + \|\vec{f}\chi_{B^c(0,R)}\|_{M_p^{t,r}(W)} \\ &\leq N + \|\vec{f}\chi_{B(Nh,R)}\|_{M_p^{t,r}(W)} + \|\vec{f}\chi_{B^c(0,R)}\|_{M_p^{t,r}(W)} \leq N + 2. \end{aligned}$$

This proves that  $\mathcal{F}$  is bounded.

PROOF OF THEOREM 3.2. Assume that  $\mathcal{F} \subset M_p^{t,r}(W)$  satisfies (i) and (ii). Given  $\varepsilon > 0$  small enough, to prove the total boundedness of  $\mathcal{F}$ , it suffices to find a finite  $\varepsilon$ -net of  $\mathcal{F}$ . Denote by  $R_i := [-2^i, 2^i]^n$  for  $i \in \mathbb{Z}$ . Then from condition (i), there exist a positive integer  $m$  large enough such that

$$\sup_{\vec{f} \in \mathcal{F}} \|\vec{f} - \vec{f}\chi_{R_m}\|_{M_p^{t,r}(W)} < \varepsilon. \quad (3.2)$$

Moreover, by condition (ii), there exists a negative integer  $a$  such that

$$\sup_{\vec{f} \in \mathcal{F}} \sup_{y \in R_a} \|\vec{f} - \tau_y \vec{f}\|_{M_p^{t,r}(W)} < \varepsilon. \quad (3.3)$$

There exists a sequence  $\{Q_j\}_{j=1}^N$  of disjoint cubes in  $\mathcal{D}_{-a}$  such that  $R_m = \cup_{j=1}^N Q_j$ , where  $N = 2^{(m+1-a)n}$ .

For any  $\vec{f} \in \mathcal{F}$  and  $x \in \mathbb{R}^n$ , let

$$\Phi(\vec{f})(x) := \begin{cases} \vec{f}_{Q_j} := \frac{1}{|Q_j|} \int_{Q_j} \vec{f}(y) dy, & x \in Q_j, j = 1, 2, \dots, N, \\ \vec{0}, & \text{otherwise.} \end{cases}$$

Then for each fixed  $x \in \mathbb{R}^n$ , we have

$$\begin{aligned} \left| W(x) \left( \vec{f}(x) - \vec{f}_{Q_j} \right) \right| \chi_{Q_j}(x) &= \left| \frac{1}{|Q_j|} \int_{Q_j} W(x) \left( \vec{f}(x) - \vec{f}(y) \right) dy \right| \chi_{Q_j}(x) \\ &\leq \frac{1}{|Q_j|} \int_{Q_j} \left| W(x) \left( \vec{f}(x) - \vec{f}(y) \right) \right| dy \chi_{Q_j}(x). \end{aligned} \quad (3.4)$$

We split  $\|(\vec{f} - \Phi(\vec{f}))\chi_{R_m}\|_{M_p^{t,r}(W)}$  into three parts:

$$\begin{aligned} &\left\{ \sum_{k \in \mathbb{Z}} \sum_{Q \in \mathcal{D}_k} W(Q)^{r/t-r/p} \left( \int_Q \left| W(x) \left( \vec{f}(x)\chi_{R_m} - \Phi(\vec{f})(x) \right) \right|^p dx \right)^{r/p} \right\}^{1/r} \\ &\leq \left\{ \sum_{k > -a} \sum_{Q \in \mathcal{D}_k} W(Q)^{r/t-r/p} \left( \int_Q \left| W(x) \left( \vec{f}(x)\chi_{R_m} - \vec{f}_{Q_j} \right) \right|^p dx \right)^{r/p} \right\}^{1/r} \\ &\quad + \left\{ \sum_{k=-m}^{-a} \sum_{Q \in \mathcal{D}_k} W(Q)^{r/t-r/p} \left( \int_Q \left| W(x) \left( \vec{f}(x)\chi_{R_m} - \Phi(\vec{f})(x) \right) \right|^p dx \right)^{r/p} \right\}^{1/r} \\ &\quad + \left\{ \sum_{k < -m} \sum_{Q \in \mathcal{D}_k} W(Q)^{r/t-r/p} \left( \int_Q \left| W(x) \left( \vec{f}(x)\chi_{R_m} - \Phi(\vec{f})(x) \right) \right|^p dx \right)^{r/p} \right\}^{1/r} \\ &=: S_1 + S_2 + S_3. \end{aligned}$$

We first estimate  $S_1$ . By (3.4), Jensen's inequality ( $p \geq 1$ ), the Fubini theorem, we have

$$\begin{aligned} S_1 &\leq \left\{ \sum_{k > -a} \sum_{Q \in \mathcal{D}_k, Q \subset R_m} W(Q)^{r/t-r/p} \right. \\ &\quad \times \left. \left( \int_Q \left| \frac{1}{|Q_j|} \int_{Q_j} \left| W(x) \left( \vec{f}(x) - \vec{f}(y) \right) \right| dy \chi_{Q_j}(x) \right|^p dx \right)^{r/p} \right\}^{1/r} \\ &\leq \left\{ \sum_{k > -a} \sum_{Q \in \mathcal{D}_k, Q \subset R_m} W(Q)^{r/t-r/p} \right. \\ &\quad \times \left. \left( \frac{1}{|Q_j|} \int_{Q_j} \int_Q \left| W(x) \left( \vec{f}(x) - \vec{f}(y) \right) \right|^p dx dy \right)^{r/p} \right\}^{1/r} \end{aligned}$$

$$\begin{aligned}
&= \left\{ \sum_{k > -a} \sum_{Q \in \mathcal{D}_k, Q \subset R_m} W(Q)^{r/t-r/p} \right. \\
&\quad \times \left. \left( \frac{1}{|Q_j|} \int_Q \int_{Q_j} |W(x) (\vec{f}(x) - \vec{f}(y))|^p dy dx \right)^{r/p} \right\}^{1/r} \\
&= 2^{-an/p} \left\{ \sum_{k > -a} \sum_{Q \in \mathcal{D}_k, Q \subset R_m} W(Q)^{r/t-r/p} \right. \\
&\quad \times \left. \left( \int_Q \int_{x-Q_j} |W(x) (\vec{f}(x) - \vec{f}(x-y))|^p dy dx \right)^{r/p} \right\}^{1/r} \tag{3.5}
\end{aligned}$$

where  $x - Q_j := \{x - y : y \in Q_j\}$ . Since  $x \in Q \subset Q_j$  for some  $j \in \{1, 2, \dots, N\}$ , we have  $x - Q_j \subset R_a$ . Hence by (3.3), we obtain

$$\begin{aligned}
(3.5) &\leq 2^{-an/p} |Q_j|^{1/p} \sup_{y \in R_a} \left\{ \sum_{k > -a} \sum_{Q \in \mathcal{D}_k, Q \subset R_m} W(Q)^{r/t-r/p} \right. \\
&\quad \times \left. \left( \int_Q |W(x) (\vec{f}(x) - \vec{f}(x-y))|^p dx \right)^{r/p} \right\}^{1/r} \\
&\leq \varepsilon.
\end{aligned}$$

As for  $S_2$ , for each  $Q \in \mathcal{D}_k$  where  $k = -m, -m+1, \dots, -a$ , there are  $2^{(-k-a)n}$  cubes  $Q_j \in \mathcal{D}_{-a}$  such that  $\cup Q_j = Q$ . Denote by  $Q_{j_\ell}$ ,  $\ell = 1, 2, \dots, 2^{(-k-a)n}$  these cubes. Then by (3.4), Jensen's inequality ( $p \geq 1$ ), the Fubini theorem, we obtain

$$\begin{aligned}
S_2 &= \left\{ \sum_{k=-m}^{-a} \sum_{Q \in \mathcal{D}_k, Q \subset R_m} W(Q)^{r/t-r/p} \right. \\
&\quad \times \left. \left( \sum_{\ell=1}^{2^{(-k-a)n}} \int_{Q_{j_\ell}} |W(x) (\vec{f}(x)\chi_{R_m} - \vec{f}_{Q_{j_\ell}})|^p dx \right)^{r/p} \right\}^{1/r} \\
&\leq \left\{ \sum_{k=-m}^{-a} \sum_{Q \in \mathcal{D}_k, Q \subset R_m} W(Q)^{r/t-r/p} \right. \\
&\quad \times \left. \left( \sum_{\ell=1}^{2^{(-k-a)n}} \int_{Q_{j_\ell}} \left| \frac{1}{|Q_{j_\ell}|} \int_{Q_{j_\ell}} |W(x) (\vec{f}(x) - \vec{f}(y))|^p dy \right| dx \right)^{r/p} \right\}^{1/r}
\end{aligned}$$

$$\begin{aligned}
&\leq \left\{ \sum_{k=-m}^{-a} \sum_{Q \in \mathcal{D}_k, Q \subset R_m} W(Q)^{r/t-r/p} \right. \\
&\quad \times \left. \left( \sum_{\ell=1}^{2^{(-k-a)n}} \frac{1}{|Q_{j_\ell}|} \int_{Q_{j_\ell}} \int_{Q_{j_\ell}} |W(x) (\vec{f}(x) - \vec{f}(y))|^p dx dy \right)^{r/p} \right\}^{1/r} \\
&= 2^{-an/p} \left\{ \sum_{k=-m}^{-a} \sum_{Q \in \mathcal{D}_k, Q \subset R_m} W(Q)^{r/t-r/p} \right. \\
&\quad \times \left. \left( \sum_{\ell=1}^{2^{(-k-a)n}} \int_{Q_{j_\ell}} \int_{Q_{j_\ell}} |W(x) (\vec{f}(x) - \vec{f}(y))|^p dy dx \right)^{r/p} \right\}^{1/r} \\
&= 2^{-an/p} \left\{ \sum_{k=-m}^{-a} \sum_{Q \in \mathcal{D}_k, Q \subset R_m} W(Q)^{r/t-r/p} \right. \\
&\quad \times \left. \left( \sum_{\ell=1}^{2^{(-k-a)n}} \int_{Q_{j_\ell}} \int_{x-Q_{j_\ell}} |W(x) (\vec{f}(x) - \vec{f}(x-y))|^p dy dx \right)^{r/p} \right\}^{1/r} \tag{3.6}
\end{aligned}$$

where  $x - Q_{j_\ell} := \{x - y : y \in Q_{j_\ell}\}$ . Note that  $x - Q_{j_\ell} \subset R_a$  when  $x \in Q_{j_\ell}$ . By (3.3), we have

$$\begin{aligned}
(3.6) &\leq 2^{-an/p} 2^{an/p} \sup_{y \in R_a} \left\{ \sum_{k=-m}^{-a} \sum_{Q \in \mathcal{D}_k, Q \subset R_m} W(Q)^{r/t-r/p} \right. \\
&\quad \times \left. \left( \sum_{\ell=1}^{2^{(-k-a)n}} \int_{Q_{j_\ell}} |W(x) (\vec{f}(x) - \vec{f}(x-y))|^p dx \right)^{r/p} \right\}^{1/r} \\
&\leq \varepsilon.
\end{aligned}$$

As for  $S_3$ , for each set  $\mathcal{D}_k$ , there are only  $2^n$  cubes  $Q \in \mathcal{D}_k$  such that  $Q \cap R_m \neq \emptyset$ . We denote by  $Q_\ell$ ,  $\ell = 1, 2, \dots, 2^n$  these cubes. And for each cube  $Q_\ell$ , there are  $2^{(m-a)n}$  cubes  $Q_j \in \mathcal{D}_{-a}$  such that  $\cup_j Q_j = Q_\ell$ . Then

$$S_3 = \left\{ \sum_{k < -m} \sum_{\ell=1}^{2^n} W(Q_\ell)^{r/t-r/p} \left( \int_{Q_\ell} |W(x) (\vec{f}(x)\chi_{R_m} - \vec{\Phi}(\vec{f})(x))|^p dx \right)^{r/p} \right\}^{1/r}$$

$$= \left\{ \sum_{k < -m} \sum_{\ell=1}^{2^n} W(Q_\ell)^{r/t-r/p} \left( \sum_{j=1}^{2^{(m-a)n}} \int_{Q_j} |W(x) (\vec{f}(x)\chi_{R_m} - \vec{f}_{Q_j})|^p dx \right)^{r/p} \right\}^{1/r}.$$

Similarly as  $S_2$ , from (3.4), Jensen's inequality ( $p \geq 1$ ), the Fubini theorem, and (3.3), we get that  $S_3 \leq \varepsilon$ . Together with the estimates of  $S_1, S_2, S_3$ , we obtain

$$\left\{ \sum_{k \in \mathbb{Z}} \sum_{Q \in \mathcal{D}_k} W(Q)^{r/t-r/p} \left( \int_Q |W(x)(\vec{f}(x)\chi_{R_m} - \Phi(\vec{f})(x))|^p dx \right)^{r/p} \right\}^{1/r} \leq 3\varepsilon. \quad (3.7)$$

Note that

$$\begin{aligned} \|\vec{f} - \Phi(\vec{f})\|_{M_p^{t,r}(W)} &\leq \|(\vec{f} - \Phi(\vec{f}))\chi_{R_m}\|_{M_p^{t,r}(W)} + \|(\vec{f} - \Phi(\vec{f}))\chi_{R_m^c}\|_{M_p^{t,r}(W)} \\ &= \|(\vec{f} - \Phi(\vec{f}))\chi_{R_m}\|_{M_p^{t,r}(W)} + \|\vec{f}\chi_{R_m^c}\|_{M_p^{t,r}(W)}. \end{aligned}$$

Thus via (3.2) and (3.7), we have

$$\sup_{\vec{f} \in \mathcal{F}} \|\vec{f} - \Phi(\vec{f})\|_{M_p^{t,r}(W)} \leq 4\varepsilon. \quad (3.8)$$

From (3.8), it suffices to show that  $\Phi(\mathcal{F})$  is totally bounded in  $M_p^{t,r}(W)$ .

From Remark 3.3, we have

$$\sup_{\vec{f} \in \mathcal{F}} \|\Phi(\vec{f})\|_{M_p^{t,r}(W)} \leq \sup_{\vec{f} \in \mathcal{F}} \|\Phi(\vec{f}) - \vec{f}\|_{M_p^{t,r}(W)} + \sup_{\vec{f} \in \mathcal{F}} \|\vec{f}\|_{M_p^{t,r}(W)} < \infty.$$

Thus, for any  $\vec{f} \in \mathcal{F}$ ,

$$|W^{1/p}(x)\Phi(\vec{f})(x)| < \infty \quad \text{a.e. } x \in \mathbb{R}^n. \quad (3.9)$$

Since  $W$  is a matrix weight, by (i), (ii) of Definition 2.3, we have for almost everywhere  $x \in \mathbb{R}^n$ ,  $W(x)$  is a positive definite matrix. Therefore, we obtain

$$|\Phi(\vec{f})(x)| < \infty \quad \text{a.e. } x \in \mathbb{R}^n. \quad (3.10)$$

which implies  $|\vec{f}_{Q_j}| < \infty$ ,  $j = 1, 2, \dots, N$ . From this and the entries of  $W$  is locally integrable, we see that  $\Phi$  is a map from  $\mathcal{F}$  to  $\mathcal{B}$ , a finite dimensional Banach subspace of  $M_p^{t,r}(W)$ . Note that  $\Phi(\mathcal{F}) \subset \mathcal{B}$  is bounded, and hence is totally bounded. The proof of Theorem 3.2 is complete.  $\square$

Then we give an application in degenerate Bourgain-Morrey-Sobolev spaces with matrix weights.

**DEFINITION 3.4.** Let  $1 \leq p < t < r < \infty$  or  $1 \leq p \leq t < r = \infty$ . Let  $W : \mathbb{R}^n \rightarrow M_n(\mathbb{C})$  be a matrix weight and set the scalar weight  $\omega := \|W\|$ . We

define the degenerate Bourgain-Morrey-Sobolev spaces  $\mathcal{W}^{1,p,t,r}(W)$  by the set of all Lebesgue measurable functions on  $\mathbb{R}^n$  such that

$$\|f\|_{\mathcal{W}^{1,p,t,r}(W)} := \|f\|_{M_p^{t,r}(\omega)} + \|\nabla f\|_{M_p^{t,r}(W)} < \infty$$

where  $\nabla f = (\partial_{x_1} f, \dots, \partial_{x_n} f)^T$  is the gradient of  $f$  and

$$\|f\|_{M_p^{t,r}(\omega)} := \left\| \left\{ \omega(Q_{j,k})^{1/t-1/p} \left( \int_{Q_{j,k}} |f(y)|^p \omega(y) dy \right)^{1/p} \right\}_{j \in \mathbb{Z}, k \in \mathbb{Z}^n} \right\|_{\ell^r},$$

is the norm of the scalar weighted Bourgain-Morrey space in (2.6).

LEMMA 3.5 (p. 262, [33]). *A set  $\mathcal{F}$  in metric space  $X$  is totally bounded if and only if it is Cauchy-precompact, that is, every sequence has a Cauchy subsequence.*

COROLLARY 3.6. *Let  $1 \leq p < t < r < \infty$  or  $1 \leq p \leq t < r = \infty$ . Let  $W : \mathbb{R}^n \rightarrow M_n(\mathbb{C})$  be a matrix weight. A subset  $\mathcal{F} \subset \mathcal{W}^{1,p,t,r}(W)$  is totally bounded if the following conditions are valid:*

(i)  $\mathcal{F}$  uniformly vanishes at infinity, that is,

$$\lim_{R \rightarrow \infty} \sup_{\vec{f} \in \mathcal{F}} \|\vec{f} \chi_{B^c(0,R)}\|_{\mathcal{W}^{1,p,t,r}(W)} = 0;$$

(ii)  $\mathcal{F}$  is equicontinuous, that is,

$$\lim_{a \rightarrow 0} \sup_{\vec{f} \in \mathcal{F}} \sup_{y \in B(0,a)} \|\vec{f} - \tau_y \vec{f}\|_{\mathcal{W}^{1,p,t,r}(W)} = 0.$$

PROOF. Note that  $\mathcal{F} \subset \mathcal{W}^{1,p,t,r}(W)$  satisfies conditions (i)-(ii) if and only if  $\mathcal{F} \subset M_p^{t,r}(\omega)$  satisfies conditions (i)-(ii) of Theorem 3.2 ( $d = 1$ ) and  $\nabla \mathcal{F} := \{\nabla f : f \in \mathcal{F}\} \subset M_p^{t,r}(W)$  satisfies (i)-(ii) of Theorem 3.2 ( $d = n$ ). Hence by Theorem 3.2, we obtain that both  $\mathcal{F} \subset M_p^{t,r}(\omega)$  and  $\nabla \mathcal{F} \subset M_p^{t,r}(W)$  are totally bounded. Then Corollary 3.6 follows from Lemma 3.5.  $\square$

#### 4. Dyadic average operator conditions for precompact sets

In this section, we give a criterion for precompactness by dyadic average operator. Let  $\mathcal{F}$  be a subset of  $M_p^{t,r}(W)$ .  $\mathcal{F}$  is equicontinuous by means of the dyadic average operator if

$$\lim_{a \rightarrow -\infty, a \in \mathbb{Z}} \sup_{\vec{f} \in \mathcal{F}} \|\vec{f} - E_{d,a} \vec{f}\|_{M_p^{t,r}(W)} = 0, \quad (4.1)$$

where

$$E_{d,a} \vec{f}(x) := \sum_{Q \in \mathcal{D}_{-a}} \frac{\chi_Q(x)}{|Q|} \int_Q \vec{f}(y) dy. \quad (4.2)$$



Recall that  $x_Q$  is the left corner of  $Q \in \mathcal{D}$ . Then for  $x \in Q \in \mathcal{D}_{-a}$

$$E_{d,a}\vec{f}(x) = \frac{\chi_{Q_{-a,0}}(x - x_Q)}{|Q_{-a,0}|} \int_{Q_{-a,0}} \vec{f}(y - x_Q) dy.$$

Then we will prove that (3.1) is stronger than (4.1). Indeed, suppose that condition (3.1) holds. For any  $\varepsilon > 0$ , there exists a cube  $R$  with center 0 and side length  $2^{a+1} > 0$  such that

$$\sup_{\vec{f} \in \mathcal{F}} \sup_{y \in R} \|\vec{f} - \tau_y \vec{f}\|_{M_p^{t,r}(W)} < \varepsilon. \quad (4.3)$$

If  $x, y \in Q \in \mathcal{D}_{-a}$ , then  $x - y \in R$ . By Jensen's inequality ( $p \geq 1$ ), Hölder's inequality ( $r/p \geq 1$ ), (4.3), we obtain

$$\begin{aligned} \|\vec{f} - E_{d,a}\vec{f}\|_{M_p^{t,r}(W)} &= \left( \sum_{Q \in \mathcal{D}} W(Q)^{r/t-r/p} \left( \int_Q \left| \frac{1}{|Q_{-a,0}|} \right. \right. \right. \\ &\quad \times \left. \left. \int_{Q_{-a,0}} W^{1/p}(x) \left( \vec{f}(x) - \chi_{Q_{-a,0}}(x - x_Q) \vec{f}(y - x_Q) \right) dy \right|^p dx \right)^{r/p} \right)^{1/r} \\ &\leq \left( \sum_{Q \in \mathcal{D}} W(Q)^{r/t-r/p} \left( \frac{1}{|Q_{-a,0}|} \int_{Q_{-a,0}} \right. \right. \\ &\quad \times \left. \left. \int_Q \left| W^{1/p}(x) \left( \vec{f}(x) - \chi_{Q_{-a,0}}(x - x_Q) \vec{f}(y - x_Q) \right) \right|^p dx dy \right)^{r/p} \right)^{1/r} \\ &\leq \left( \sum_{Q \in \mathcal{D}} W(Q)^{r/t-r/p} \frac{1}{|Q_{-a,0}|} \int_{Q_{-a,0}} \right. \\ &\quad \times \left. \left( \int_Q \left| W^{1/p}(x) \left( \vec{f}(x) - \chi_{Q_{-a,0}}(x - x_Q) \vec{f}(y - x_Q) \right) \right|^p dx \right)^{r/p} dy \right)^{1/r} \\ &\leq \frac{1}{|Q_{-a,0}|^{1/r}} |Q_{-a,0}|^{1/r} \sup_{y \in R} \left( \sum_{Q \in \mathcal{D}} W(Q)^{r/t-r/p} \right. \\ &\quad \times \left. \left( \int_Q \left| W^{1/p}(x) \left( \vec{f}(x) - \vec{f}(x - y) \right) \right|^p dx \right)^{r/p} \right)^{1/r} \\ &\leq \varepsilon, \end{aligned}$$

modified when  $r = \infty$ . Hence we prove that (3.1) is stronger than (4.1).

Replacing the translation operator by the average operator, we have the following result.

**THEOREM 4.1.** *Let  $1 \leq p < t < r < \infty$  or  $1 \leq p \leq t < r = \infty$ . Let  $W : \mathbb{R}^n \rightarrow M_d(\mathbb{C})$  be a matrix weight. A subset  $\mathcal{F} \subset M_p^{t,r}(W)$  is totally bounded if the following conditions are valid:*

(i)  $\mathcal{F}$  is bounded, that is,

$$\sup_{\vec{f} \in \mathcal{F}} \|\vec{f}\|_{M_p^{t,r}(W)} < \infty;$$

(ii)  $\mathcal{F}$  uniformly vanishes at infinity, that is,

$$\lim_{R \rightarrow \infty} \sup_{\vec{f} \in \mathcal{F}} \|\vec{f} \chi_{B^c(0,R)}\|_{M_p^{t,r}(W)} = 0;$$

(iii)  $\mathcal{F}$  is equicontinuous by means of the dyadic average operator, that is, for any

$$\lim_{a \rightarrow -\infty, a \in \mathbb{Z}} \sup_{\vec{f} \in \mathcal{F}} \|\vec{f} - E_{d,a} \vec{f}\|_{M_p^{t,r}(W)} = 0.$$

where  $E_{d,a}$  is same as (4.2).

**PROOF.** Assume that  $\mathcal{F} \subset M_p^{t,r}(W)$  satisfies (i)-(iii). Given  $\varepsilon > 0$  small enough, to prove the total boundedness of  $\mathcal{F}$ , it suffices to find a finite  $\varepsilon$ -net of  $\mathcal{F}$ . Denote by  $R_i := [-2^i, 2^i]^n$  for  $i \in \mathbb{Z}$ . Then from condition (ii), there exists a positive integer  $m$  large enough such that

$$\sup_{\vec{f} \in \mathcal{F}} \|\vec{f} - \vec{f} \chi_{R_m}\|_{M_p^{t,r}(W)} < \varepsilon. \quad (4.4)$$

Moreover, by condition (iii), there exists an integer  $a < 0$  such that

$$\sup_{\vec{f} \in \mathcal{F}} \|\vec{f} - E_{d,a} \vec{f}\|_{M_p^{t,r}(W)} < \varepsilon. \quad (4.5)$$

There exists a sequence  $\{Q_j\}_{j=1}^N$  of disjoint cubes in  $\mathcal{D}_{-a}$  such that  $R_m = \cup_{i=1}^N Q_j$ , where  $N = 2^{(m+1-a)n}$ .

For any  $\vec{f} \in \mathcal{F}$  and  $x \in \mathbb{R}^n$ , let

$$\Phi(\vec{f})(x) := \begin{cases} \vec{f}_{Q_j} := \frac{1}{|Q_j|} \int_{Q_j} \vec{f}(y) dy, & x \in Q_j, j = 1, 2, \dots, N, \\ \vec{0}, & \text{otherwise.} \end{cases}$$

Note that for  $x \in R_m$ ,

$$\Phi(\vec{f})(x) = E_{d,a} \vec{f}(x).$$

Hence via (4.4) and (4.5), we have

$$\begin{aligned}
\|\vec{f} - \Phi(\vec{f})\|_{M_p^{t,r}(W)} &\leq \|(\vec{f} - \Phi(\vec{f}))\chi_{R_m}\|_{M_p^{t,r}(W)} + \|(\vec{f} - \Phi(\vec{f}))\chi_{R_m^c}\|_{M_p^{t,r}(W)} \\
&= \|(\vec{f} - E_{d,a}\vec{f})\chi_{R_m}\|_{M_p^{t,r}(W)} + \|\vec{f}\chi_{R_m^c}\|_{M_p^{t,r}(W)} \\
&\leq \|(\vec{f} - E_{d,a}\vec{f})\|_{M_p^{t,r}(W)} + \varepsilon \\
&\leq 2\varepsilon.
\end{aligned} \tag{4.6}$$

From (4.6), it suffices to show that  $\Phi(\mathcal{F})$  is totally bounded in  $M_p^{t,r}(W)$ . And we have proved that  $\Phi(\mathcal{F})$  is totally bounded in  $M_p^{t,r}(W)$  in Theorem 3.2. Thus the proof of Theorem 4.1 is complete.  $\square$

LEMMA 4.2 (Lemma 4.5, [8]). *Let  $1 \leq p < \infty$  and  $W \in \mathcal{A}_p$ . Then  $\|W\|$  and  $\|W^{-1}\|^{-1}$  are scalar  $A_p$  weights.*

The following is a vector-valued extension of the Lebesgue differentiation theorem on matrix weighted spaces  $M_p^{t,r}(W)$ . For any cube  $Q$ ,  $\vec{f} \in L_{\text{loc}}^1(\mathbb{R}^n)$ , define

$$E_Q(\vec{f})(x) = \frac{1}{|Q|} \int_Q \vec{f}(y) dy.$$

THEOREM 4.3. *Let  $1 \leq p < t < r < \infty$  or  $1 \leq p \leq t < r = \infty$ . If  $W \in \mathcal{A}_p$ , then for any  $\vec{f} \in M_p^{t,r}(W)$ ,*

$$\lim_{\ell(Q) \rightarrow 0} |E_Q \vec{f}(x) - \vec{f}(x)| = 0 \quad \text{a.e. } x \in \mathbb{R}^n.$$

PROOF. First, for any  $\vec{f} = (f_1, f_2, \dots, f_d)^T \in M_p^{t,r}(W)$ , it suffices to show that  $f_i \in L_{\text{loc}}^1(\mathbb{R}^n)$  for each  $i = 1, 2, \dots, d$ . Since  $W \in \mathcal{A}_p$ , then by (2.3)

$$|\vec{f}|^p = |W^{-1/p} W^{1/p} \vec{f}|^p \leq \|W^{-1/p}\|^p |W^{1/p} \vec{f}|^p = \|W^{-1}\| |W^{1/p} \vec{f}|^p.$$

It follows that

$$|\vec{f}|^p \|W^{-1}\|^{-1} \leq |W^{1/p} \vec{f}|^p.$$

From Lemma 4.2, we conclude that  $\|W^{-1}\|^{-1}$  is a scalar  $A_p$  weight, hence  $|\vec{f}| \in L_{\text{loc}}^1(\mathbb{R}^n)$ . Indeed, since any compact set  $K \subset \mathbb{R}^n$  can be contained in a finite set of dyadic cubes, it suffices to show that for any  $Q \in \mathcal{D}$ ,  $|\vec{f}| \in L^1(Q)$ . By Hölder's inequality, we have

$$\begin{aligned}
\int_Q |\vec{f}| dx &= \int_Q |\vec{f}| \|W^{-1}\|^{-1/p} \|W^{-1}\|^{1/p} dx \\
&\leq \left( \int_Q |\vec{f}|^p \|W^{-1}\|^{-1} dx \right)^{1/p} \left( \int_Q \|W^{-1}\|^{p'/p} dx \right)^{1/p'}
\end{aligned}$$

$$\begin{aligned} &\leq C_{Q,W,p} \left( \int_Q |W^{1/p} \vec{f}|^p dx \right)^{1/p} \\ &\lesssim \|\vec{f}\|_{M_p^{t,r}(W)} < \infty. \end{aligned}$$

Hence,  $f_i \in L_{\text{loc}}^1(\mathbb{R}^n)$  for each  $i = 1, 2, \dots, d$ . By the classical Lebesgue differentiation theorem, for each  $1 \leq i \leq d$ , we have

$$\lim_{\ell(Q) \rightarrow 0} |E_Q f_i(x) - f_i(x)| = 0 \quad \text{a.e. } x \in \mathbb{R}^n.$$

Theorem 4.3 comes from the fact that for any  $x \in \mathbb{R}^n$ ,

$$\left| E_Q \vec{f}(x) - \vec{f}(x) \right| \leq d^{1/2} \max_i |E_Q f_i(x) - f_i(x)|. \quad \square$$

Next we need the boundedness of dyadic average operator. Given any collection  $\mathcal{Q}$  of pairwise disjoint cubes  $Q \subset \mathbb{R}^n$ , define the averaging operator  $A_{\mathcal{Q}}$  by

$$A_{\mathcal{Q}} \vec{f}(x) = \sum_{Q \in \mathcal{Q}} \frac{\chi_Q(x)}{|Q|} \int_Q \vec{f}(y) dy.$$

LEMMA 4.4 (Proposition 4.7,[8]). *Let  $1 \leq p < \infty$ . Let  $W : \mathbb{R}^n \rightarrow M_d(\mathbb{C})$  be a matrix weight. Then  $W \in \mathcal{A}_p$  if and only if it satisfies*

$$\|A_{\mathcal{Q}} \vec{f}\|_{L^p(W)} \leq C_{n,d,p,W} \|\vec{f}\|_{L^p(W)}.$$

THEOREM 4.5. *Let  $W \in \mathcal{A}_p$  has the  $\mathcal{A}_p$ -dimension  $\tilde{d} \in [0, n)$  with the doubling exponent  $\beta$ .*

(i) *Let  $1 \leq p < t < r < \infty$ . If  $(\tilde{d} - n + \beta)r/p + n - \beta r/t < 0$ , then for each  $a \in \mathbb{Z}$ ,  $E_{d,a}$  is a bounded operator on matrix weighted Bourgain-Morrey spaces  $M_p^{t,r}(W)$ .*

(ii) *Let  $1 \leq p \leq t < r = \infty$ . If  $\tilde{d}/p - n/p - \beta(1/t - 1/p) \leq 0$ , then for each  $a \in \mathbb{Z}$ ,  $E_{d,a}$  is a bounded operator on matrix weighted Bourgain-Morrey spaces  $M_p^{t,r}(W)$ .*

PROOF. (i) For a fixed  $a \in \mathbb{Z}$ , we have

$$\begin{aligned} \|E_{d,a} \vec{f}\|_{M_p^{t,r}(W)} &\leq \left( \sum_{j \leq -a} \sum_{Q \in \mathcal{D}_j} W(Q)^{r/t - r/p} \left( \int_Q |W^{1/p}(x) E_{d,a} \vec{f}(x)|^p dx \right)^{r/p} \right)^{1/r} \\ &\quad + \left( \sum_{j > -a} \sum_{Q \in \mathcal{D}_j} W(Q)^{r/t - r/p} \left( \int_Q |W^{1/p}(x) E_{d,a} \vec{f}(x)|^p dx \right)^{r/p} \right)^{1/r} \\ &=: S_1 + S_2. \end{aligned}$$

By Lemma 4.4, we obtain

$$\begin{aligned} S_1 &\leq C_{n,d,p,W} \left( \sum_{j \leq -a} \sum_{Q \in \mathcal{D}_j} W(Q)^{r/t-r/p} \left( \int_Q |W^{1/p}(x) \vec{f}(x)|^p dx \right)^{r/p} \right)^{1/r} \\ &\lesssim \|\vec{f}\|_{M_p^{t,r}(W)}. \end{aligned}$$

When  $j > -a$ , we denote by  $(j+a)_{\text{pa}}Q \in \mathcal{D}_{-a}$  the  $(j+a)$ -th dyadic parent of  $Q \in \mathcal{D}_j$ . Let  $\{A_Q\}_{Q \in \mathcal{D}}$  be the reducing operator of order  $p$  for  $W$ . By Lemmas 2.8, we have

$$\|A_Q A_{(j+a)_{\text{pa}}Q}^{-1}\| \leq c 2^{(j+a)\tilde{d}/p}. \quad (4.7)$$

In  $S_2$ , using (2.5) and (4.7), we have

$$\begin{aligned} S_2^r &= \sum_{j > -a} \sum_{Q \in \mathcal{D}_j} W(Q)^{r/t-r/p} \left( \int_Q |W^{1/p}(x) E_{d,a} \vec{f}(x)|^p dx \right)^{r/p} \\ &= \sum_{j > -a} \sum_{Q \in \mathcal{D}_j} W(Q)^{r/t-r/p} \\ &\quad \times \left( \frac{|Q|}{|Q|} \int_Q \left| W^{1/p}(x) \frac{1}{|(j+a)_{\text{pa}}Q|} \int_{(j+a)_{\text{pa}}Q} \vec{f}(y) dy \right|^p dx \right)^{r/p} \\ &\approx \sum_{j > -a} \sum_{Q \in \mathcal{D}_j} W(Q)^{r/t-r/p} |Q|^{r/p} \left( \left| A_Q \frac{1}{|(j+a)_{\text{pa}}Q|} \int_{(j+a)_{\text{pa}}Q} \vec{f}(y) dy \right| \right)^r \\ &\lesssim \sum_{j > -a} \sum_{Q \in \mathcal{D}_j} W(Q)^{r/t-r/p} |Q|^{r/p} \\ &\quad \times \left( 2^{(j+a)\tilde{d}/p} \left| A_{(j+a)_{\text{pa}}Q} \frac{1}{|(j+a)_{\text{pa}}Q|} \int_{(j+a)_{\text{pa}}Q} \vec{f}(y) dy \right| \right)^r \\ &\approx \sum_{j > -a} \sum_{Q \in \mathcal{D}_j} W(Q)^{r/t-r/p} |Q|^{r/p} 2^{(j+a)\tilde{d}r/p} \\ &\quad \times \left( \frac{1}{|(j+a)_{\text{pa}}Q|} \int_{(j+a)_{\text{pa}}Q} |W^{1/p}(x) \frac{1}{|(j+a)_{\text{pa}}Q|} \int_{(j+a)_{\text{pa}}Q} \vec{f}(y) dy|^p dx \right)^{r/p} \\ &= \sum_{j > -a} \sum_{Q \in \mathcal{D}_j} W(Q)^{r/t-r/p} 2^{-jn r/p} 2^{(j+a)\tilde{d}r/p} \\ &\quad \times \left( \frac{1}{|(j+a)_{\text{pa}}Q|} \int_{(j+a)_{\text{pa}}Q} |W^{1/p}(x) E_{d,a} \vec{f}(x)|^p dx \right)^{r/p} \end{aligned}$$

$$\begin{aligned}
&= \sum_{j > -a} \sum_{Q \in \mathcal{D}_j} W(Q)^{r/t-r/p} 2^{-jnr/p} 2^{(j+a)\tilde{d}r/p} 2^{-anr/p} \\
&\quad \times \left( \int_{(j+a)_{pa}Q} |W^{1/p}(x) E_{d,a} \vec{f}(x)|^p dx \right)^{r/p}.
\end{aligned}$$

Remark that given  $S \in \mathcal{D}_{-a}$ , there are  $2^{(j+a)n}$  cubes  $R$  such that  $(j+a)R = S$ . Hence, by Lemma 4.4 and  $(\tilde{d} - n + \beta)r/p + n - \beta r/t < 0$ , we have

$$\begin{aligned}
&\sum_{j > -a} \sum_{Q \in \mathcal{D}_j} W(Q)^{r/t-r/p} 2^{-jnr/p} 2^{(j+a)\tilde{d}r/p} 2^{-anr/p} \\
&\quad \times \left( \int_{(j+a)_{pa}Q} |W^{1/p}(x) E_{d,a} \vec{f}(x)|^p dx \right)^{r/p} \\
&= \sum_{j > -a} 2^{-jnr/p} 2^{(j+a)\tilde{d}r/p} 2^{-anr/p} \\
&\quad \times \sum_{Q \in \mathcal{D}_j} W(Q)^{r/t-r/p} \left( \int_{(j+a)_{pa}Q} |W^{1/p}(x) E_{d,a} \vec{f}(x)|^p dx \right)^{r/p} \\
&\leq \sum_{j > -a} 2^{-jnr/p} 2^{(j+a)\tilde{d}r/p} 2^{-anr/p} 2^{(j+a)n} 2^{-(j+a)\beta(r/t-r/p)} \\
&\quad \times \sum_{S \in \mathcal{D}_{-a}} W(S)^{r/t-r/p} \left( \int_S |W^{1/p}(x) E_{d,a} \vec{f}(x)|^p dx \right)^{r/p} \\
&\lesssim \|\vec{f}\|_{M_p^{t,r}(W)}^r,
\end{aligned}$$

where in the second step we use the following fact. Since  $W((j+a)_{pa}Q) \leq 2^{(j+a)\beta}W(Q)$  and  $1/t - 1/p < 0$ , we have

$$W(Q)^{1/t-1/p} < 2^{-(j+a)\beta(1/t-1/p)} W((j+a)_{pa}Q)^{1/t-1/p}.$$

(ii) Fix  $Q_{j,k} \in \mathcal{D}$ . If  $j \leq -a$ , then by Lemma 4.4, we have

$$\begin{aligned}
&W(Q_{j,k})^{1/t-1/p} \left( \int_{Q_{j,k}} |W^{1/p}(x) E_{d,a} \vec{f}(x)|^p dx \right)^{1/p} \\
&\lesssim W(Q_{j,k})^{1/t-1/p} \left( \int_{Q_{j,k}} |W^{1/p}(x) \vec{f}(x)|^p dx \right)^{1/p}.
\end{aligned}$$

If  $j > -a$ , we denote by  $(j+a)_{\text{pa}}Q_{j,k} \in \mathcal{D}_{-a}$  the  $(j+a)$ -th dyadic parent of  $Q_{j,k} \in \mathcal{D}_j$ .

$$\begin{aligned}
 & W(Q_{j,k})^{1/t-1/p} \left( \int_{Q_{j,k}} |W^{1/p}(x)E_{d,a}\vec{f}(x)|^p dx \right)^{1/p} \\
 &= W(Q_{j,k})^{1/t-1/p} \left( \frac{|Q_{j,k}|}{|Q_{j,k}|} \int_{Q_{j,k}} \left| W^{1/p}(x) \frac{1}{|(j+a)_{\text{pa}}Q_{j,k}|} \int_{(j+a)_{\text{pa}}Q_{j,k}} \vec{f}(y) dy \right|^p dx \right)^{1/p} \\
 &\approx W(Q_{j,k})^{1/t-1/p} |Q_{j,k}|^{1/p} \left| A_{Q_{j,k}} \frac{1}{|(j+a)_{\text{pa}}Q_{j,k}|} \int_{(j+a)_{\text{pa}}Q_{j,k}} \vec{f}(y) dy \right| \\
 &\lesssim W(Q_{j,k})^{1/t-1/p} |Q_{j,k}|^{1/p} 2^{(j+a)\bar{d}/p} \left| A_{(j+a)_{\text{pa}}Q_{j,k}} \frac{1}{|(j+a)_{\text{pa}}Q_{j,k}|} \int_{(j+a)_{\text{pa}}Q_{j,k}} \vec{f}(y) dy \right| \\
 &\approx W(Q_{j,k})^{1/t-1/p} 2^{-jn/p} 2^{(j+a)\bar{d}/p} \left( \frac{1}{|(j+a)_{\text{pa}}Q_{j,k}|} \right. \\
 &\quad \left. \times \int_{(j+a)_{\text{pa}}Q_{j,k}} \left| W^{1/p}(x) \frac{1}{|(j+a)_{\text{pa}}Q_{j,k}|} \int_{(j+a)_{\text{pa}}Q_{j,k}} \vec{f}(y) dy \right|^p dx \right)^{1/p}.
 \end{aligned}$$

Then by Lemma 4.4 and  $\bar{d}/p - n/p - \beta(1/t - 1/p) \leq 0$ , we obtain

$$\begin{aligned}
 & W(Q_{j,k})^{1/t-1/p} \left( \int_{Q_{j,k}} |W^{1/p}(x)E_{d,a}\vec{f}(x)|^p dx \right)^{1/p} \\
 &\lesssim W(Q_{j,k})^{1/t-1/p} 2^{-jn/p} 2^{(j+a)\bar{d}/p} \\
 &\quad \times \left( \frac{1}{|(j+a)_{\text{pa}}Q_{j,k}|} \int_{(j+a)_{\text{pa}}Q_{j,k}} |W^{1/p}(x)\vec{f}(x)|^p dx \right)^{1/p} \\
 &= W(Q_{j,k})^{1/t-1/p} 2^{-jn/p} 2^{(j+a)\bar{d}/p} 2^{-an/p} \left( \int_{(j+a)_{\text{pa}}Q_{j,k}} |W^{1/p}(x)\vec{f}(x)|^p dx \right)^{1/p} \\
 &\lesssim 2^{(j+a)(\bar{d}/p-n/p)} 2^{-(j+a)\beta(1/t-1/p)} W((j+a)_{\text{pa}}Q_{j,k})^{1/t-1/p} \\
 &\quad \times \left( \int_{(j+a)_{\text{pa}}Q_{j,k}} |W^{1/p}(x)\vec{f}(x)|^p dx \right)^{1/p} \\
 &\lesssim W((j+a)_{\text{pa}}Q_{j,k})^{1/t-1/p} \left( \int_{(j+a)_{\text{pa}}Q_{j,k}} |W^{1/p}(x)\vec{f}(x)|^p dx \right)^{1/p}
 \end{aligned}$$

where in the third step we use the following fact. Since  $W((j+a)_{\text{pa}}Q) \leq 2^{(j+a)\beta} W(Q)$

and  $1/t - 1/p \leq 0$ , we have

$$W(Q)^{1/t-1/p} \leq 2^{-(j+a)\beta(1/t-1/p)} W((j+a)_{\text{pa}}Q)^{1/t-1/p}.$$

Finally, take the supremum over the cubes  $Q_{j,k} \in D$ , we obtain

$$\|E_{d,a}\vec{f}\|_{M_p^{t,\infty}(W)} \lesssim \|\vec{f}\|_{M_p^{t,\infty}(W)}. \quad \square$$

Finally, we have the following Kolmogorov-Riesz compactness theorem for matrix weighted Bourgain-Morrey spaces.

**THEOREM 4.6.** *Let  $1 \leq p < t < r < \infty$ . Let  $W \in \mathcal{A}_p$  has the  $\mathcal{A}_p$ -dimension  $\tilde{d} \in [0, n)$  with the doubling exponent  $\beta$ . Let  $(\tilde{d} - n + \beta)r/p + n - \beta r/t < 0$ . A subset  $\mathcal{F}$  of  $M_p^{t,r}(W)$  is totally bounded if and only if the following conditions hold:*

(i)  $\mathcal{F}$  is bounded, that is,

$$\sup_{\vec{f} \in \mathcal{F}} \|\vec{f}\|_{M_p^{t,r}(W)} < \infty;$$

(ii)  $\mathcal{F}$  uniformly vanishes at infinity, that is,

$$\lim_{R \rightarrow \infty} \sup_{\vec{f} \in \mathcal{F}} \|\vec{f}\chi_{B^c(0,R)}\|_{M_p^{t,r}(W)} = 0;$$

(iii)  $\mathcal{F}$  is equicontinuous by means of the dyadic average operator, that is, for any

$$\lim_{a \rightarrow -\infty, a \in \mathbb{Z}} \sup_{\vec{f} \in \mathcal{F}} \|\vec{f} - E_{d,a}\vec{f}\|_{M_p^{t,r}(W)} = 0.$$

where  $E_{d,a}$  is same as (4.2).

**PROOF.** The sufficiency is due to Theorem 4.1. Now we prove the necessity. Assume that  $\mathcal{F}$  of  $M_p^{t,r}(W)$  is totally bounded. For any given  $\varepsilon > 0$ , there exists  $\{\vec{f}_k\}_{k=1}^{N_0} \subset \mathcal{F}$  such that  $\{\vec{f}_k\}_{k=1}^{N_0}$  is an  $\varepsilon$ -net of  $\mathcal{F}$ , that is, for any  $\vec{f} \in \mathcal{F}$ , there exists  $\vec{f}_k, k \in \{1, 2, \dots, N_0\}$  such that  $\|\vec{f} - \vec{f}_k\|_{M_p^{t,r}(W)} < \varepsilon$ .

Clearly, (i) is true.

For any  $\varepsilon > 0$  and  $k \in \{1, 2, \dots, N_0\}$ , since  $r < \infty$ , there exists  $R_k > 0$  such that

$$\|\vec{f}_k\chi_{B^c(0,R_k)}\|_{M_p^{t,r}(W)} < \varepsilon.$$

Taking  $R = \max_{k \in \{1, \dots, N_0\}} \{R_k\}$ , we have  $\|\vec{f}_k\chi_{B^c(0,R)}\|_{M_p^{t,r}(W)} < \varepsilon$ . Then for any  $\vec{f} \in \mathcal{F}$ ,

$$\|\vec{f}\chi_{B^c(0,R)}\|_{M_p^{t,r}(W)} \leq \|(\vec{f} - \vec{f}_k)\chi_{B^c(0,R)}\|_{M_p^{t,r}(W)} + \|\vec{f}_k\chi_{B^c(0,R)}\|_{M_p^{t,r}(W)} < 2\varepsilon.$$

Thus we prove that

$$\lim_{R \rightarrow \infty} \sup_{\vec{f} \in \mathcal{F}} \|\vec{f}\chi_{B^c(0,R)}\|_{M_p^{t,r}(W)} = 0.$$



Therefore, we prove that  $\mathcal{F}$  satisfies condition (ii).

As for (iii), for each  $1 \leq k \leq N_0$ , by Theorem 4.5, we have for each  $a \in \mathbb{Z}$ ,

$$\|\vec{f}_k - E_{d,a}\vec{f}_k\|_{M_p^{t,r}(W)} \leq (1+c)\|\vec{f}_k\|_{M_p^{t,r}(W)}.$$

Thus using the dominated convergence theorem to obtain that there exists  $a_0 \in \mathbb{Z}$  such that for any  $a \leq a_0$ ,

$$\max_{1 \leq k \leq N_0} \|\vec{f}_k - E_{d,a}\vec{f}_k\|_{M_p^{t,r}(W)} < \varepsilon.$$

Now if  $a \leq a_0$ , then

$$\begin{aligned} & \|\vec{f} - E_{d,a}\vec{f}\|_{M_p^{t,r}(W)} \\ & \leq \|E_{d,a}\vec{f} - E_{d,a}\vec{f}_k\|_{M_p^{t,r}(W)} + \|E_{d,a}\vec{f}_k - \vec{f}_k\|_{M_p^{t,r}(W)} + \|\vec{f}_k - \vec{f}\|_{M_p^{t,r}(W)} \\ & \leq c\|\vec{f} - \vec{f}_k\|_{M_p^{t,r}(W)} + \varepsilon + \varepsilon \\ & \lesssim \varepsilon. \end{aligned} \quad \square$$

REMARK 4.7. Theorem 4.6 is not true for  $1 < p < t < r = \infty$ . Indeed, let  $d = 1, W \equiv 1, 0 < p < t < r = \infty$ . Let  $f(x) = |x|^{-n/t}$ . Then

$$\|f\|_{M_p^{t,\infty}} = \sup_{Q \in \mathcal{D}} |Q|^{1/t-1/p} \left( \int_Q |y|^{-np/t} dy \right)^{1/p} \approx 1.$$

Let  $f_j = \chi_{B(0,2^j)} f$  for  $j \in \mathbb{N}$ . Note that  $f$  and  $f_j$  are radial and symmetric functions. Let  $x = (x_1, 0, \dots, 0)$  where  $x_1 = 2^j + 2 \times 2^j$ . Let  $s = x_1/10$ . Then  $B(x, s) \subset B(0, 2^j)^c$ . Then

$$\begin{aligned} & \|f - f_j\|_{M_p^{t,\infty}} \\ & \geq cs^{-n(1/t-1/p)} \left( \int_{B(x,s)} |y|^{-np/t} dy \right)^{1/p} \\ & \geq cs^{-n(1/t-1/p)} \left( \int_{|x|-|s|}^{|x|+|s|} \eta^{-np/t} \eta^{n-1} d\eta \right)^{1/p} \\ & = c \left( \frac{|x|}{10} \right)^{-n(1/t-1/p)} \left( \left( 1 + \frac{1}{10} \right)^{n-np/t} - \left( 1 - \frac{1}{10} \right)^{n-np/t} \right)^{1/p} |x|^{n(1/t-1/p)} \\ & = c > 0. \end{aligned}$$

This shows that the totally bounded set  $\mathcal{F} = \{f\}$  of  $M_p^{t,\infty}$  is not uniformly vanishes at infinity.

ACKNOWLEDGEMENT(S). The corresponding author Jingshi Xu is supported by the National Natural Science Foundation of China (Grant No. 12161022) and the Science and Technology Project of Guangxi (Guike AD23023002).

### References

- [1] J. BOURGAIN, On the restriction and multiplier problems in  $R^3$ , In *Geometric aspects of functional analysis*, Proceedings of the Israel seminar (GAFA), Springer-Verlag, 1991, 179–191.
- [2] F. BU, T. HYTÖNEN, D. YANG, AND W. YUAN, New characterizations and properties of matrix  $a_\infty$  weights, arXiv:2311.05974 (2023).
- [3] F. BU, T. HYTÖNEN, D. YANG, AND W. YUAN, Matrix-Weighted Besov-Type and Triebel–Lizorkin-Type Spaces I:  $A_p$ -Dimensions of Matrix Weights and  $\varphi$ -Transform Characterizations, arXiv:2304.00292 (2023).
- [4] F. BU, T. HYTÖNEN, D. YANG, AND W. YUAN, Matrix-Weighted Besov-Type and Triebel–Lizorkin-Type Spaces II: Sharp Boundedness of Almost Diagonal Operators, arXiv:2312.13548 (2023).
- [5] F. BU, T. HYTÖNEN, D. YANG, AND W. YUAN, Matrix-Weighted Besov-Type and Triebel–Lizorkin-Type Spaces III: Characterizations of Molecules and Wavelets, Trace Theorems, and Boundedness of Pseudo-Differential Operators and Calderón–Zygmund Operators, arXiv:2312.13549 (2023).
- [6] F. BU, D. YANG, AND W. YUAN, Real-variable characterizations and their applications of matrix-weighted Besov spaces on spaces of homogeneous type, *Math. Z.* **305** (2023), 81.
- [7] A. CLOP, V. CRUZ, Weighted estimates for Beltrami equations, *Ann. Acad. Sci. Fenn., Math.* **38** (2013), 91–113.
- [8] D. CRUZ-URIBE, K. MOEN, AND S. RODNEY, Matrix  $A_p$  weights, degenerate Sobolev spaces, and mappings of finite distortion, *J. Geom. Anal.* **26** (2016), 2797–2830.
- [9] M. FRAZIER, S. ROUDENKO, Matrix-weighted Besov spaces and conditions of  $A_p$  type for  $0 < p \leq 1$ , *Indiana Univ. Math. J.* **53** (2004), 1225–1254.
- [10] M. FRAZIER, S. ROUDENKO, Littlewood–Paley theory for matrix-weighted function spaces, *Math. Ann.* **380** (2021), 487–537.
- [11] M. GOLDBERG, Matrix  $A_p$  weights via maximal functions, *Pac. J. Math.* **211** (2003), 201–220.
- [12] L. GRAFAKOS, *Classical Fourier analysis*, Grad. Texts Math. 249, Springer, 2014.
- [13] W. GUO, G. ZHAO, On relatively compact sets in quasi-Banach function spaces, *Proc. Am. Math. Soc.* **148** (2020), 3359–3373.
- [14] H. HANCHE-OLSEN, H. HOLDEN, The Kolmogorov–Riesz compactness theorem, *Expo. Math.* **28** (2010), 385–394.
- [15] D. D. HAROSKE, P. SKANDERA, Embeddings of doubling weighted Besov spaces, In *Function spaces X, Proceedings of the 10th international conference*, Warszawa: Polish Academy of Sciences, Institute of Mathematics, 2014, 105–119.
- [16] N. HATANO, T. NOGAYAMA, Y. SAWANO, AND D. I. HAKIM, Bourgain–Morrey spaces and their applications to boundedness of operators, *J. Funct. Anal.* **284** (2023), 52.
- [17] R. A. HORN, C. R. JOHNSON, *Topics in matrix analysis*, Cambridge University Press, 1994.
- [18] R. A. HORN, C. R. JOHNSON, *Matrix analysis*, Cambridge University Press, 2013.
- [19] P. HU, Y. LI, AND D. YANG, Bourgain–Morrey spaces meet structure of Triebel–Lizorkin spaces, *Math. Z.* **304** (2023), 49.
- [20] A. KOLMOGOROFF, Über Kompaktheit der Funktionenmengen bei der Konvergenz im Mittel, *Nachr. Ges. Wiss. Göttingen, Math.-Phys. Kl.* **1931** (1931), 60–63.

- [21] S. LIU, D. YANG, AND C. ZHUO, Matrix weighted Kolmogorov-Riesz's compactness theorem, *Front. Math. (Beijing)* **18** (2023), 1167–1189.
- [22] D. D. MONTICELLI, S. RODNEY, AND R. L. WHEEDEN, Boundedness of weak solutions of degenerate quasilinear equations with rough coefficients, *Differ. Integral Equ.* **25** (2012), 143–200.
- [23] F. L. NAZAROV, S. R. TREIL', The hunt for a Bellman function: Applications to estimates of singular integral operators and to other classical problems in harmonic analysis, *St. Petersburg. Math. J.* **8** (1996), 32–162.
- [24] S. ROUDENKO, Matrix-weighted Besov spaces, *Trans. Am. Math. Soc.* **355** (2003), 273–314.
- [25] S. ROUDENKO, Duality of matrix-weighted Besov spaces, *Stud. Math.* **160** (2004), 129–156.
- [26] S. ROUDENKO, *The theory of function spaces with matrix weights*, ProQuest LLC, Ann Arbor, MI. Thesis (Ph.D.) Michigan State University, 2002.
- [27] Y. SAWANO, *Theory of Besov spaces*, Dev. Math. 56, Singapore: Springer, 2018.
- [28] E. T. SAWYER, R. L. WHEEDEN, Degenerate Sobolev spaces and regularity of subelliptic equations, *Trans. Am. Math. Soc.* **362** (2010), 1869–1906.
- [29] S. TREIL, A. VOLBERG, Wavelets and the angle between past and future, *J. Funct. Anal.* **143** (1997), 269–308.
- [30] A. VOLBERG, Matrix  $A_p$  weights via  $S$ -functions, *J. Am. Math. Soc.* **10** (1997), 445–466.
- [31] Q. WANG, D. YANG, AND Y. ZHANG, Real-variable characterizations and their applications of matrix-weighted Triebel-Lizorkin spaces, *J. Math. Anal. Appl.* **529** (2024), 37.
- [32] N. WIENER, P. MASANI, The prediction theory of multivariate stochastic processes. I. The regularity condition. - II. The linear predictor, *Acta Math.* **98** (1957), 111–150,
- [33] S. WILLARD, *General topology*, Mineola, NY: Dover Publications, reprint of the 1970 original edition, 2004.

*Present Address:*

TENGFEE BAI  
 COLLEGE OF MATHEMATICS AND STATISTICS  
 HAINAN NORMAL UNIVERSITY  
 HAIKOU, HAINAN 571158, CHINA  
*e-mail*: 202311070100007@hainnu.edu.cn

JINGSHI XU  
 SCHOOL OF MATHEMATICS AND COMPUTING SCIENCE  
 GUILIN UNIVERSITY OF ELECTRONIC TECHNOLOGY  
 GUILIN 541004, CHINA  
 CENTER FOR APPLIED MATHEMATICS OF GUANGXI (GUET)  
 GUILIN 541004, CHINA  
 GUANGXI COLLEGES AND UNIVERSITIES KEY LABORATORY OF DATA ANALYSIS AND COMPUTATION  
 GUILIN 541004, CHINA  
*e-mail*: jingshixu@126.com