Global well-posedness, scattering and blow-up for the energy-critical, Schrödinger equation with general nonlinearity in the radial case*

Jun Wang^a, Zhaoyang Yin a,b†

^aDepartment of Mathematics, Sun Yat-sen University, Guangzhou, 510275, China
 ^bSchool of Science, Shenzhen Campus of Sun Yat-sen University, Shenzhen, 518107, China

Abstract

In this paper, we study the well-posedness theory and the scattering asymptotics for the energy-critical, Schrödinger equation with general nonlinearity

$$\begin{cases} i\partial_t u + \Delta u + f(u) = 0, \ (x, t) \in \mathbb{R}^N \times \mathbb{R}, \\ u|_{t=0} = u_0 \in H^1(\mathbb{R}^N), \end{cases}$$

where $f: \mathbb{C} \to \mathbb{C}$ satisfies Sobolev critical growth condition. Using contraction mapping method and concentration compactness argument, we obtain the well-posedness theory in proper function spaces and scattering asymptotics. This paper generalizes the conclusions in [17](Invent. Math).

Keywords: Schrödinger equation, General nonlinearity, Well-posedness, Scattering

2010 MSC: 35Q55, 35R11, 37K05, 37L50

1 Introduction and main results

This paper studies the well-posedness theory and the scattering asymptotics for the energy-critical, focusing, Schrödinger equation with general nonlinearity

$$\begin{cases}
i\partial_t u + \Delta u + f(u) = 0, & (x, t) \in \mathbb{R}^N \times \mathbb{R}, \\
u|_{t=0} = u_0 \in H^1(\mathbb{R}^N),
\end{cases}$$
(1.1)

where $f: \mathbb{C} \to \mathbb{C}$ satisfies Sobolev critical growth condition.

In [17], Kenig and Merle studied the \dot{H}^1 critical non-linear Schrödinger equation

$$\begin{cases} i\partial_t u + \Delta u \pm |u|^{\frac{4}{N-2}} u = 0, & (x,t) \in \mathbb{R}^N \times \mathbb{R}, \\ u|_{t=0} = u_0 \in \dot{H}^1(\mathbb{R}^N). \end{cases}$$

^{*}This work was partially supported by NNSFC (No. 12171493).

[†]Corresponding author. wangj937@mail2.sysu.edu.cn (J. Wang), mcsyzy@mail.sysu.edu.cn (Z. Yin)

Here the - sign corresponds to the defocusing problem, while the + sign corresponds to the focusing problem. They obtained the global well-posedness, scattering and blow-up results in the radial case and $3 \le N \le 5$. Recently, Oh and Wang in [21] considered global well-posedness for one-dimensional cubic nonlinear Schrödinger equation by introducing a new function space. For N=2, [6] established global well-posedness results in the defocusing case, posed on the two-dimensional unit ball. For high dimensions case, Tao et.al in [22] established global well-posedness and scattering for solutions to the defocusing mass-critical nonlinear Schrödinger equation. For more well-posedness results, please refer to [1, 7–9, 11, 15, 24].

Unlike other articles on well-posedness and scattering, this paper presents for the first time results on general critical nonlinear terms. This idea mainly comes from the variational method. In recent decades, many researchers have been interested in semi-linear Schrödinger problems with general nonlinear terms. From a mathematical perspective, it is very meaningful. In order to obtain the solution of the equation, it is necessary to impose appropriate conditions on the nonlinear term. The most classic condition is the Ambrosetti-Rabinowitz(AR) condition:

There exists
$$\mu > 2$$
 such that $G(t) \ge \mu g(t) > 0$ for all $t > 0$,

where $G(t) = \int_0^t g(s)ds$. As is well known, (AR) condition plays a crucial role in getting a bounded (PS)-sequence, so we can use the Nehari manifold method to obtain the solution of the equation. As we can see, the (AR) condition ensures that nonlinearity has good properties. However, we use another weaker Berestycki-Lions(BL) condition than (AR) in this paper, namely

There exists
$$\xi > 0$$
 such that $G(\xi) > 0$.

As mentioned in [2], under the (BL) condition, we can use Pohozaev manifold to obtain the existence of standing wave solutions (even the existence of infinite solutions), which is crucial for obtaining variational estimates in Section 6.

Now, we give the most general assumption of growth, i.e., assume that $f: \mathbb{C} \to \mathbb{C}$ is C^1 and satisfies:

$$(F_1)$$
 $\lim_{|z| \to 0^+} \frac{|f(z)\overline{z}|}{|z|^2} = 0.$

$$(F_2)$$
 $\limsup_{|z|\to\infty} \frac{|f(z)\overline{z}|}{|z|^{2^*}} < \infty$, where $2^* = \frac{2N}{N-2}$.

$$(F_3) \quad \lim_{|\nabla z| \to 0^+} \frac{|\nabla (f(z))|}{|\nabla z|} = 0.$$

$$(F_4)$$
 $\limsup_{|\nabla z| \to \infty} \frac{|\nabla (f(z))|}{|z|^{2^*-2}||\nabla z|} < \infty.$

Remark 1.1. By (F_1) , (F_2) , (F_3) , (F_4) , there exist ε , C > 0 such that

$$\frac{|f(z)\overline{z}|}{|z|^2} < \varepsilon, \ \frac{|\nabla(f(z))|}{|\nabla z|} < \varepsilon$$

and

$$\frac{|f(z)\overline{z}|}{|z|^{2^*}} < C, \ \frac{|\nabla(f(z))|}{|z|^{2^*-2}||\nabla z|} < C.$$

Hence, it follows from $f \in C^1$ that there exist C_1 , C_2 , C_3 , $C_4 > 0$ such that

$$|f(z)| \le C_1|z| + C_2|z|^{2^*-1}, |\nabla(f(z))| \le C_3|\nabla z| + C_4|z|^{2^*-2}|\nabla z|.$$

Remark 1.2. Let $f(u) = |u|^{\frac{4}{N-2}}u + |u|^{\frac{2}{N-2}}u$, then clearly

$$|f(u)| \le |u|^{\frac{N+2}{N-2}} + |u|^{\frac{N}{N-2}}, |\partial_z f(u)| \le C|u|^{\frac{4}{N-2}} + |u|^{\frac{2}{N-2}}, |\partial_{\bar{z}} f(u)| \le C|u|^{\frac{4}{N-2}} + |u|^{\frac{2}{N-2}}.$$

Moreover, for $3 \le N \le 4$,

$$\frac{|\partial_z f(u) - \partial_z f(v)|}{|\partial_{\bar{z}} f(u) - \partial_{\bar{z}} f(v)|} \le C|u - v| \cdot \left\{ |u|^{\frac{6-N}{N-2}} + |v|^{\frac{6-N}{N-2}} + |u|^{\frac{4-N}{N-2}} + |v|^{\frac{4-N}{N-2}} \right\}.$$

Also,

$$|f(u) - f(v)| \le |u - v| \left\{ |u|^{\frac{4}{N-2}} + |v|^{\frac{4}{N-2}} + |u|^{\frac{2}{N-2}} + |v|^{\frac{2}{N-2}} \right\}.$$

In addition,

$$\nabla_{x}(f(u(x))) - \nabla_{x}(f(v(x)))$$

$$= (\nabla f)(u(x))\nabla u - (\nabla f)(v(x))\nabla v$$

$$= (\nabla f)(u(x))\nabla u - (\nabla f)(u(x))\nabla v + {\nabla f(u(x))} - \nabla f(v(x))}\nabla v,$$

so

$$|\nabla_x f(u(x)) - \nabla_x f(v(x))| \leq C|u|^{\frac{4}{N-2}} |\nabla u - \nabla v| + C|\nabla v| \left\{ |u|^{\frac{6-N}{N-2}} + |v|^{\frac{6-N}{N-2}} \right\} |u - v| + C|u|^{\frac{2}{N-2}} |\nabla u - \nabla v| + C|\nabla v| \left\{ |u|^{\frac{6-N}{N-2}} + |v|^{\frac{4-N}{N-2}} \right\} |u - v|.$$

In the subsequent proof, energy conservation is necessary. If the nonlinear term is of power type, there is obviously energy conservation. This paper considers more general nonlinear terms. Inspired by [13], we need to make the following new assumption:

$$(F_5)$$
 For all $z \in \mathbb{C}$, $f(\bar{z}) = \overline{f(z)}$. For all $z \in \mathbb{C}$ and all $\omega \in \mathbb{C}$ with $|\omega| = 1$, $f(\omega z) = \omega f(z)$.

It follows from (F_5) that f(z) is of the form $f(z) = z\tilde{f}(|z|)$ where \tilde{f} is a real function defined on $(0, \infty)$. Moreover, note that f is continuous, one can define a function F from \mathbb{C} to \mathbb{R} by

$$F(\rho) := 2 \int_0^{\rho} \sigma \tilde{f}(\sigma) d\sigma \qquad \text{for all } \rho \ge 0,$$
$$F(z) := F(|z|) \qquad \text{for all } z \in \mathbb{C}.$$

The function f can be recovered from F by the relation

$$f(z) = \frac{\partial F(z)}{\partial \bar{z}}.$$

Hence, by using Propositions 3.3 and 3.4 in [13], we can obtain the conservation of the L^2 -norm and of the energy in differential form for the regularized equation. That is, if the solution u of (1.1) has sufficient decay at infinity and smoothness, it satisfies the conservation of mass

$$M(u(t)) := ||u(t)||_{L^2} = ||u_0||_{L^2}$$
(1.2)

and the conservation of energy

$$E(u(t)) = E(u_0), \tag{1.3}$$

where E(u(t)) is defined by

$$E(u(t)) := \frac{1}{2} \int_{\mathbb{R}^N} |\nabla u(t, x)|^2 dx - \int_{\mathbb{R}^N} F(u(t)) dx, \ F(u) = F(|u|) = 2 \int_0^{|u|} s\tilde{f}(s) ds$$

and the energy space is H^1 .

The main results of this paper are as follows.

Theorem 1.1. Assume $u_0 \in H^1_{rad}(\mathbb{R}^N)$, $N \geq 3$ and $(F_1) - (F_5)$ hold, then there exists a unique solution u to (1.1) satisfying

$$u \in C([0,T], H^1_{rad}(\mathbb{R}^N)) \cap L^{\frac{2(N+2)}{N-2}}_{loc}([0,T], W^{1,\frac{2N(N+2)}{N^2+4}}).$$

Moreover, there exists the globally solution u to (1.1) if $||u_0||_{L^2}$ is small enough.

Remark 1.3. The conclusion of Theorem 1.1 also holds for defocusing cases, please refer to the proof of Lemma 3.2 for details.

Theorem 1.2. Assume that $N \geq 3$ and $u_0 \in H^1_{rad}$ and $(F_1) - (F_5)$ hold. Let u be such that the corresponding solution to (1.1) exists on the maximal time T^* . If $E(u_0) < 0$, then one of the following statements holds true:

- (1) u(t) blows up in finite time in the sense that $T^* < +\infty$ must hold.
- (2) u(t) blows up infinite time such that

$$\sup_{t \ge 0} \|\nabla u(\cdot, t)\|_{L^2} = \infty. \tag{1.4}$$

Remark 1.4. Under the assumption of Theorem 1.1, we can define a maximal interval $I(u_0) = (t_0 - T_-(u_0), t_0 + T_+(u_0))$, with $T_\pm(u_0) > 0$, where the solution is defined. If $T_+(u_0) < \infty$, then by standard finite blow-up criterion, we know that

$$||u||_{L^{\frac{2(N+2)}{N-2}}_{[t_0,t_0+T_+(u_0)]}W^{1,\frac{2N(N+2)}{N^2+4}}} = +\infty,$$

the corresponding result holds for $T_{-}(u_0)$.

Theorem 1.3. Let $(F_1) - (F_6)$ hold, N = 3, 4. Assume that

$$E(u_0) < E(W), \int_{\mathbb{R}^N} |\nabla u_0|^2 dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx$$

and u_0 is radial. Then there exist $u_{0,+}, u_{0,-}$ in H^1 such that

$$\lim_{t \to +\infty} \|u(t) - e^{it\Delta} u_{0,+}\|_{H^1} = 0, \ \lim_{t \to -\infty} \|u(t) - e^{it\Delta} u_{0,-}\|_{H^1} = 0.$$

Remark 1.5. Theorem 1.3 is a part of Corollary 9.1. In fact, according to the analysis in Remark 5.1, we only need to prove that the conclusion holds when t tends towards positive infinity, and the other half of the conclusion can be obtained similarly.

In section 2, we provide some notations and some important lemma in the proof of main theorems. In the next section, we aim to prove theorem 1.1, that is well-posedness. In section 4, we get the blow up solutions in infinite time. Finally, we will obtain the scattering asymptotics. To achieve this goal, we need some new variational estimates and compactness results, that is, sections 6, 7 and 8.

2 Preliminary

In this section, we provide some notations and some important lemma in the proof of main theorems.

Lemma 2.1. (Strichartz estimate [9, 16]). We say that a pair of exponents (q, r) is admissible if $\frac{2}{q} + \frac{N}{r} = \frac{N}{2}$ and $2 \le q, r \le \infty$. Then, if $2 \le r \le \frac{2N}{N-2}$ $(N \ge 3)(\text{or } 2 \le r < \infty, N = 2)$ and $2 \le r \le \infty, N = 1$) we have

$$||e^{it\Delta}h||_{L^qL^r_-} \le C||h||_{L^2},$$

ii)
$$\left\| \int_{-\infty}^{+\infty} e^{i(t-\tau)\Delta} g(-,\tau) d\tau \right\|_{L_t^q L_x^r} + \left\| \int_0^t e^{i(t-\tau)\Delta} g(-,\tau) d\tau \right\|_{L_t^q L_x^r} \le C \|g\|_{L_t^{q'} L_x^{r'}},$$

iii)
$$\left\| \int_{-\infty}^{+\infty} e^{it\Delta} g(-,\tau) d\tau \right\|_{L^{2}_{x}} \le C \|g\|_{L^{q'}_{t}L''_{x}}.$$

Remark 2.1. In the estimate ii) in Lemma 2.1, one can actually show: ([16]) ii')

$$\left\| \int_{-\infty}^{+\infty} e^{i(t-\tau)\Delta} g(-,\tau) d\tau \right\|_{L^q_t L^r_x} \le C \|g\|_{L^{m'}_t L^{n'}_x},$$

where (q, r), (m, n) are any pair of admissible indices as in i) of Lemma 2.1.

Lemma 2.2. (Sobolev embedding). For $v \in C_0^{\infty}(\mathbb{R}^{N+1})$, we have

$$||v||_{L_t^{\frac{2(N+2)}{N-2}}L_x^{\frac{2(N+2)}{N-2}}} \le C||\nabla_x v||_{L_t^{\frac{2(N+2)}{N-2}}L_x^{\frac{2N(N+2)}{N^2+4}}} (N \ge 3).$$

(Note that $\frac{2(N+2)}{N-2} = q$, $\frac{2N(N+2)}{N^2+4} = r$ is admissible.)

Lemma 2.3. ([12]) Let $F(z) = |z|^k z$ with $k > 0, s \ge 0$ and $1 < p, p_1 < \infty, 1 < q_1 \le \infty$ satisfying $\frac{1}{p} = \frac{1}{p_1} + \frac{k}{q_1}$. If k is an even integer or k is not an even integer with $[s] \le k$, then there exists C > 0 such that for all $u \in \mathscr{S}$,

$$||F(u)||_{\dot{W}^{s,p}} \le C||u||_{L^{q_1}}^k ||u||_{\dot{W}^{s,p_1}}.$$

A similar estimate holds with $\dot{W}^{s,p}$, \dot{W}^{s,p_1} -norms replaced by $W^{s,p}$, W^{s,p_1} norms.

Lemma 2.4. ([5]) Let $N \geq 1$ and $f : \mathbb{R}^N \to \mathbb{R}$ satisfy $\nabla f \in W^{1,\infty}(\mathbb{R}^N)$. Then, for all $u \in H^{\frac{1}{2}}(\mathbb{R}^N)$, it holds that

$$\left| \int_{\mathbb{R}^N} \bar{u}(x) \nabla f(x) \cdot \nabla u(x) dx \right| \leqslant C \left(\left\| |\nabla|^{\frac{1}{2}} u \right\|_{L^2}^2 + \|u\|_{L^2} \left\| |\nabla|^{\frac{1}{2}} u \right\|_{L^2} \right),$$

with some constant C > 0 depending only on $\|\nabla f\|_{W^{1,\infty}}$ and N.

Definition 2.1. Let $v_0 \in H^1$, $v(t) = e^{it\Delta}v_0$ and let $\{t_n\}$ be a sequence, with $\lim_{n \to \infty} t_n = \bar{t} \in [-\infty, +\infty]$. We say that u(x,t) is a non-linear profile associated with $(v_0, \{t_n\})$ if there exists an interval I, with $\bar{t} \in I$ (if $\bar{t} = \pm \infty$, $I = [a, +\infty)$ or $(-\infty, a]$) such that u is a solution of (1.1) in I and

$$\lim_{n \to \infty} \|u(-, t_n) - v(-, t_n)\|_{H^1} = 0.$$

Next, we give some definitions that play a crucial role in the proof of concentration compactness in section 6.

Definition 2.2. (i) We call scale, every sequence $\mathbf{h} = (h_n)_{n \geqslant 0}$ of positive numbers and core, every sequence $\mathbf{z} = (z_n)_{n \geqslant 0} = (t_n, x_n)_{n \geqslant 0} \subset \mathbb{R} \times \mathbb{R}^N$.

(ii) We say that two pairs (\mathbf{h}, \mathbf{z}) and $(\mathbf{h}', \mathbf{z}')$ are orthogonal if

$$\frac{h_n}{h'_n} + \frac{h'_n}{h_n} + \left| \frac{t_n - t'_n}{(h_n)^2} \right| + \left| \frac{x_n - x'_n}{h_n} \right| \to +\infty, \ n \to \infty.$$

Definition 2.3. (i) A pair (q,r) is L^2 -admissible, if $r \in [2, \frac{2N}{N-2})$ and q satisfies

$$\frac{2}{q} + \frac{N}{r} = \frac{N}{2}.$$

(ii) A pair (q,r) is H^1 -admissible, if $r \in [\frac{2N}{N-2}, +\infty)$ and q satisfies

$$\frac{2}{q} + \frac{N}{r} = \frac{N-2}{2}.$$

The main result proved in [14, 23] is the following:

Proposition 2.1. Let (q,r) be an L^2 -admissible pair. There exists C=C(r), such that

$$\left\| e^{it\Delta} h \right\|_{L_t^q L_x^r} \le C \|\varphi\|_{L^2(\mathbb{R}^N)} \tag{2.1}$$

for every $\varphi \in L^2(\mathbb{R}^N)$.

A direct consequence of Proposition 2.1, via Sobolev's inequality, is the following.

Proposition 2.2. Let (q,r) be an H^1 -admissible pair. There exists C=C(r), such that

$$\left\| e^{it\Delta} h \right\|_{L^q_t L^r_x} \le C \|\nabla \varphi\|_{L^2(\mathbb{R}^N)} \tag{2.2}$$

for every $\varphi \in \dot{H}^1(\mathbb{R}^N)$.

Definition 2.4. Let (\mathbf{h}, \mathbf{z}) be a pair of scales-cores, such that the quantity $\frac{t_n}{h_n^2}$ has a limit in $[-\infty, +\infty]$, when n goes to the infinity. Let V be a solution of linear Schrödinger equation (7.5). We say that U is the nonlinear profile associated to $\mathscr{W} = (V, \mathbf{h}, \mathbf{z})$ if U is the solution of the nonlinear Schrödinger equation (1.1) satisfying

$$\left\| (U - V) \left(-\frac{t_n}{h_n^2} \right) \right\|_{H^1(\mathbb{R}^N)} \to 0, \ n \to \infty.$$

Notations:

- ullet Throughout this paper, we use C to denote the universal constant and C may change line by line.
 - We also use notation C(B) (or C_B) to denote a constant depends on B.
- We use usual L^p spaces and Sobolev spaces H^1 . p' for the dual index of $p \in (1, +\infty)$ in the sense that $\frac{1}{p'} + \frac{1}{p} = 1$.
 - We use notation $A \subseteq B$ to denote A is a open subset of B.

3 Well-posedness

First, we get the existence and uniqueness of solution to (1.1) by using contraction mapping method.

Lemma 3.1. Assume $u_0 \in H^1_{rad}(\mathbb{R}^N)$, $N \geq 3$, then there exists a unique solution u to (1.1) satisfying

$$u \in C([0,T], H^1_{rad}(\mathbb{R}^N)) \cap L^{\frac{2(N+2)}{N-2}}_{loc}([0,T], W^{1,\frac{2N(N+2)}{N^2+4}}).$$

Proof. (1.1) is equivalent to the integral equation

$$u(t) = e^{it\Delta}u_0 + i \int_0^t e^{i(t-t')\Delta} f(u)dt'.$$

Now, we consider the complete metric space

$$B(T,\rho) = \left\{ C([0,T], H^1(\mathbb{R}^N)) \cap L_T^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}} : \|u\|_{L_T^{\infty}H^1} + \|u\|_{L_T^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}} \le \rho \right\}$$

equipped with the distance

$$d_B(u,v) := \|u - v\|_{L_T^{\infty}L^2} + \|u - v\|_{L_T^{\frac{2(N+2)}{N-2}}L^{\frac{2N(N+2)}{N^2+4}}}$$

and

$$\Phi_{u_0}(v) = e^{it\Delta}u_0 + i \int_0^t e^{i(t-t')\Delta} f(v)dt'.$$

Next, we prove that $\Phi_{u_0}(v): B(T,\rho) \to B(T,\rho)$ and is a contraction. In fact, by (F_1) and (F_2) , we obtain

$$|f(v)| \le C_1|v| + C_2|v|^{2^*-2}|v|, |\nabla(f(z))| \le C_1|\nabla z| + C_2|z|^{2^*-2}|\nabla z|.$$

Hence, it follows from i) in lemma 2.1, Remark 2.1, lemma 2.2 and Hölder inequality that

$$\begin{split} & \| \Phi_{u_0}(v) \|_{L_T^\infty H^1 \cap L_T^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}} \\ &= \left\| e^{it\Delta} u_0 + i \int_0^t e^{i(t-t')\Delta} f(v) dt' \right\|_{L_T^\infty H^1 \cap L_T^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}} \\ &\leq C \| u_0 \|_{H^1} + \left\| \int_0^t e^{i(t-t')\Delta} f(v) dt' \right\|_{L_T^\infty H^1 \cap L_T^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}} \\ &\leq C \| u_0 \|_{H^1} + \left\| \int_0^t e^{i(t-t')\Delta} v dt' \right\|_{L_T^\infty H^1} + \left\| \int_0^t e^{i(t-t')\Delta} v \cdot |v|^{\frac{4}{N-2}} dt' \right\|_{L_T^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}} \\ &\leq C \| u_0 \|_{H^1} + \| v \|_{L_T^1 H^1} + C \left\| v \cdot |v|^{\frac{4}{N-2}} \right\|_{L_T^2 W^{1,\frac{2N}{N+2}}} \\ &\leq C \| u_0 \|_{H^1} + T \| v \|_{L_T^\infty H^1} + C \left\| v \|_{W^{1,\frac{2N(N+2)}{N^2+4}} \cdot \| v \|_{W^{\frac{N-2}{N-2}} U^{\frac{N-2}{N-2}}} \right\|_{L_T^2} \\ &\leq C \| u_0 \|_{H^1} + T \| v \|_{L_T^\infty H^1} + C \| v \|_{L_T^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}} \cdot \| v \|_{L_T^{\frac{N-2}{N-2}} L^{\frac{2(N+2)}{N-2}} L^{\frac{2(N+2)}{N-2}}} \\ &\leq C \| u_0 \|_{H^1} + T \| v \|_{L_T^\infty H^1} + C \| v \|_{L_T^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}} \cdot \| v \|_{L_T^{\frac{N-2}{N-2}} L^{\frac{2(N+2)}{N-2}} L^{\frac{2(N+2)}{N-2}}} \\ &\leq C \| u_0 \|_{H^1} + T \| v \|_{L_T^\infty H^1} + C \| v \|_{L_T^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}}} \cdot \| v \|_{L_T^{\frac{N-2}{N-2}} L^{\frac{2(N+2)}{N-2}} L^{\frac{2(N+2)}{N-2}}} \right\|_{L_T^{\frac{N+2}{N-2}} L^{\frac{N+2}{N-2}}} \\ &\leq C \| u_0 \|_{H^1} + T \| v \|_{L_T^\infty H^1} + C \| v \|_{L_T^{\frac{N+2}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}} \cdot \| v \|_{L_T^{\frac{N+2}{N-2}} L^{\frac{2(N+2)}{N-2}} L^{\frac{2(N+2)}{N-2}}} \right\|_{L_T^{\frac{N+2}{N-2}} L^{\frac{N+2}{N-2}}} \\ &\leq C \| u_0 \|_{H^1} + T \| v \|_{L_T^\infty H^1} + C \| v \|_{L_T^{\frac{N+2}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}} \cdot \| v \|_{L_T^{\frac{N+2}{N-2}} L^{\frac{N+2}{N-2}}} \right\|_{L_T^{\frac{N+2}{N-2}} L^{\frac{N+2}{N-2}}}$$

Choosing T and ρ such that $C(\|u_0\|_{H^1} + T\rho + \rho^{\frac{N+2}{N-2}}) \leq \rho$, then $\Phi_{u_0}(v)$ maps $B(T,\rho)$ to $B(T,\rho)$.

Now we prove that $\Phi_{u_0}(v)$ is a contraction map for sufficiently small T. Let $u, v \in B(T, \rho)$, by Remark 2.1 and lemma 2.3, then we have

$$\begin{split} &d_{B}(\Phi_{u_{0}}(u),\Phi_{u_{0}}(v))\\ &= \|\Phi_{u_{0}}(u)-\Phi_{u_{0}}(v)\|_{L^{\infty}_{T}L^{2}} + \|\Phi_{u_{0}}(u)-\Phi_{u_{0}}(v)\|_{L^{2(N+2)}_{T}L^{\frac{2N(N+2)}{N^{2}+4}}}\\ &\leq \left\|\int_{0}^{t}e^{i(t-t')\Delta}|u-v|dt'\right\|_{L^{\infty}_{T}L^{2}} + \left\|\int_{0}^{t}e^{i(t-t')\Delta}|u\cdot|u|^{\frac{4}{N-2}}-v\cdot|v|^{\frac{4}{N-2}}|dt'\|_{L^{\infty}_{T}L^{2}}\\ &+ \left\|\int_{0}^{t}e^{i(t-t')\Delta}(|u-v|+|u\cdot|u|^{\frac{4}{N-2}}-v\cdot|v|^{\frac{4}{N-2}}|)dt'\right\|_{L^{\frac{2(N+2)}{N-2}}_{T}L^{\frac{2N(N+2)}{N^{2}+4}}}\\ &\leq C\|u-v\|_{L^{1}_{T}L^{2}} + \|u\cdot|u|^{\frac{4}{N-2}}-v\cdot|v|^{\frac{4}{N-2}}\|_{L^{2}_{T}L^{\frac{2N}{N+2}}} + \|u\cdot|u|^{\frac{4}{N-2}}-v\cdot|v|^{\frac{4}{N-2}}\|_{L^{2}_{T}L^{\frac{2N}{N+2}}}\\ &+ \|u-v\|_{L^{\infty}_{T}L^{\frac{2N}{N+2}}} \\ &\leq CT\|u-v\|_{L^{\infty}_{T}L^{2}} + C\|u\cdot|u|^{\frac{4}{N-2}}-v\cdot|v|^{\frac{4}{N-2}}\|_{L^{2}_{T}L^{\frac{2N}{N+2}}} + CT\|u-v\|_{L^{\infty}_{T}H^{1}}\\ &\leq CT\|u-v\|_{L^{\infty}_{T}H^{1}} + C\|u-v\|_{L^{\frac{2(N+2)}{N-2}}} \frac{2N(N+2)}{L^{\frac{2N}{N-2}}} \left(\|u\|_{L^{\frac{4-2}{N-2}}}^{\frac{4-2}{N-2}} \frac{2N(N+2)}{L^{\frac{2N}{N-2}}} \frac{2N(N+2)}{L^{\frac{2N}{N-2}}} \frac{2N(N+2)}{L^{\frac{2N}{N-2}}} \right)\\ &\leq C\|u-v\|_{L^{\frac{2(N+2)}{N-2}}} \frac{2N(N+2)}{L^{\frac{2N}{N-2}}} \left(\|u\|_{L^{\frac{4-2}{N-2}}}^{\frac{4-2}{N-2}} \frac{2N(N+2)}{L^{\frac{2N}{N-2}}} \frac{1}{L^{\frac{2N(N+2)}{N-2}}} \frac{2N(N+2)}{L^{\frac{2N}{N-2}}} \frac{1}{L^{\frac{2N(N+2)}{N-2}}} \frac{1}{L^$$

$$CT \|u - v\|_{L_T^{\infty} H^1}$$

$$\leq C(T + \rho^{\frac{4}{N-2}}) d_B(u, v).$$

Choosing T and ρ such that $C(T + \rho^{\frac{4}{N-2}}) \leq \frac{1}{2}$, the above estimate implies that $\Phi_{u_0}(v)$ is a contraction. Therefore, $\Phi_{u_0}(v)$ has a fixed point in $B(T, \rho)$.

The above lemma implies local well-posedness for (1.1). Next, we will show the global well-posedness.

Lemma 3.2. Let $N \geq 3$. Then for any $u_0 \in H^1_{rad}$, there exists the globally solution u to (1.1) if $||u_0||_{L^2}$ is small enough.

Proof. Assume that T^* be the maximal existence time. We will show that $T^* = \infty$ by contradiction. Let $T^* < \infty$, then according to the local wellposedness, it holds

$$||u||_{L_{T*}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}} = \infty.$$
(3.2)

Case 1. Consider defocusing case. Using the conservation laws (1.2) and (1.3), for any $t < T^*$, the solution u satisfies

$$\frac{1}{2}||u(t)||_{H^1}^2 \le \frac{1}{2}||u(t)||_{L^2}^2 + E(u) = \frac{1}{2}||u_0||_{L^2}^2 + E(u_0).$$

From the estimate (3.1), we have

$$||u||_{L_{T^*}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}}^{\frac{2N(N+2)}{N^2+4}} \le C||u_0||_{H^1} + T||v||_{L_T^{\infty}H^1} + C||u||_{L_T^{\frac{N+2}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}}^{\frac{2(N+2)}{N-2}} \le C\left((||u_0||_{L^2}^2 + E(u_0))^{\frac{1}{2}} + T\left(||u_0||_{L^2}^2 + E(u_0)\right)^{\frac{1}{2}} + ||u||_{L_T^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}}^{\frac{2(N+2)}{N^2+4}}\right).$$

Note that $||u||_{L_T^{\frac{N+2}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}}^{\frac{N+2}{N-2}} \leq \rho$, so for sufficiently small T depending only on $||u_0||_{L^2}^2 + E(u_0)$, we have

$$||u||_{L^{\frac{2(N+2)}{N-2}}_{(T_{j-1},T_j)}W^{1,\frac{2N(N+2)}{N^2+4}}} \le C \left(||u_0||_{L^2}^2 + E(u_0)\right)^{\frac{1}{2}},$$

where $T_j - T_{j-1} = T$. If $T^* < \infty$, then there exists $M \in \mathbb{N}$ such that $(M-1)T < T^* \leq MT$. Let $T_j = (j-1)T, j = 1, \dots, M-1$ and $T_M = T^*$. Then we have

$$||u||_{L_{(0,T^*)}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}}^{\frac{2(N+2)}{N-2}} \leq \sum_{1\leq j\leq M} ||u||_{L_{(T_{j-1},T_{j})}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}}^{\frac{2(N+2)}{N-2}}$$
$$\leq \left(MC\left(||u_{0}||_{L^{2}}^{2} + E(u_{0})\right)^{\frac{1}{2}}\right)^{p} < \infty.$$

This contradicts to (3.2).

Case 2. Consider focusing case. By $(F_1) - (F_5)$,

$$E(u_0) = E(u(t)) = \frac{1}{2} \int_{\mathbb{R}^N} |\nabla u(t, x)|^2 dx - \int_{\mathbb{R}^N} F(u(t)) dx$$

$$\geq \frac{1}{2} ||\nabla u||_{L^2}^2 - \varepsilon ||\nabla u||_{L^2}^2 - C_{\varepsilon} ||\nabla u||_{L^2}^{2^*},$$

which implies $||u||_{H^1} \leq CE(u_0)$, so we get

$$||u(t)||_{H^1}^2 \le ||u(t)||_{L^2}^2 + CE(u) = \frac{1}{2}||u_0||_{L^2}^2 + CE(u_0).$$

Similarly to the proof of the Case 1, which completes the proof of the lemma. \Box

Proof of Theorem 1.1. Combining Lemmas 3.1 and 3.2, Theorem 1.1 holds.

4 Blow up solutions

In this section, we will investigate the blow up solutions of (1.1). Let $\psi \in C_0^{\infty}(\mathbb{R}^N)$ be radial and satisfy

$$\psi(r) = \begin{cases} \frac{1}{2}r^2, & \text{for } r \leqslant 1\\ 0, & \text{for } r \geqslant 2 \end{cases} \text{ and } \psi''(r) \leqslant 1 \text{ for } r = |x| \geqslant 0.$$

For a fixed R > 0, we define the rescaled function $\psi_R : \mathbb{R}^N \to \mathbb{R}$ by setting

$$\psi_R(r) := R^2 \psi\left(\frac{r}{R}\right). \tag{4.1}$$

Next we will show that

$$1 - \psi_R''(r) \geqslant 0, \quad 1 - \frac{\psi_R'(r)}{r} \geqslant 0, \quad N - \Delta \psi_R(r) \ge 0 \quad \text{ for all } r \geqslant 0.$$
 (4.2)

Indeed, this first inequality follows from $\psi_R''(r) = \psi''(\frac{r}{R}) \le 1$. We obtain the second inequality by integrating the first inequality on [0, r] and using that $\psi_R'(0) = 0$. Finally, we see that last inequality follows from

$$N - \Delta \psi_R(r) = 1 - \psi_R''(r) + (N - 1) \left(1 - \frac{1}{r} \psi_R'(r) \right) \ge 0.$$

Besides (4.2), ψ_R admits the following properties, which can be easily checked. We define

$$\mathcal{M}_{\psi_R}[u(t)] := 2\mathrm{Im} \int_{\mathbb{R}^N} \overline{u}(t) \nabla \psi_R \cdot \nabla u(t) dx = 2\mathrm{Im} \int_{\mathbb{R}^N} \overline{u}(t) \partial_j \psi_R \partial_j u(t) dx.$$

Define the self-adjoint differential operator

$$\Gamma_{\psi_R} := i \left(\nabla \cdot \nabla \psi_R + \nabla \psi_R \cdot \nabla \right),$$

which acts on functions according to

$$\Gamma_{\psi_R} f = i \left(\nabla \cdot \left(\left(\nabla \psi_R \right) f \right) + \left(\nabla \psi_R \right) \cdot \left(\nabla f \right) \right).$$

It's easy to check that

$$\mathcal{M}_{\psi_R}[u(t)] = \langle u(t), \Gamma_{\psi_R} u(t) \rangle$$
.

Next, we show the following useful lemma.

Lemma 4.1. Let $N \ge 3$, and $u \in H^1_{rad}$ is a solution of (1.1). Let ψ_R be as in (4.1), T^* be the maximal existence time of solution u(t) in H^1_{rad} . Then for sufficiently large R, it holds

$$\frac{d}{dt}\mathcal{M}_{\psi_R}[u(t)] \le 8E(u(t)), \quad t \in [0, T^*).$$

Proof. By taking the derivative of $\mathcal{M}_{\psi_R}[u(t)]$ with respect to time t and using the equation of u(t), for any $t \in [0, T)$, it follows that

$$\frac{d}{dt}\mathcal{M}_{\psi_R}[u(t)] = \langle u(t), [-\Delta, i\Gamma_{\psi_R}] u(t) \rangle + \langle -f(u), i\Gamma_{\psi_R} u(t) \rangle + \langle u(t), i\Gamma_{\psi_R} f(u) \rangle
= I_1 + I_2 + I_3,$$

where $[X, Y] \equiv XY - YX$ denotes the commutator of operators X and Y. According to the localized radial virial estimate in [5], we obtain

$$\begin{split} I_1 &= \langle u(t), [-\Delta, i\Gamma_{\psi_R}] \, u(t) \rangle \leq 8 \, \|\nabla u\|_{L^2}^2 + CR^{-2}, \\ I_2 &= \langle -f(u), i\Gamma_{\psi_R} u(t) \rangle \\ &= -\langle f(u), (\nabla \cdot ((\nabla \psi_R) \, u) + (\nabla \psi_R) \cdot (\nabla u)) \rangle \\ &\leq 2 \int_{\mathbb{R}^N} \nabla \psi_R \cdot \nabla \left(F(u) \right) dx \\ &= -2 \int_{\mathbb{R}^N} (\Delta \psi_R) \, F(u) dx \\ &= -2(N+1) \int_{\mathbb{R}^N} F(u) dx - 2 \int_{|x| \geq R} (\Delta \psi_R - N) F(u) dx \\ &\leq -2(N+1) \int_{\mathbb{R}^N} F(u) dx + 2 \int_{|x| \geq R} (N - \Delta \psi_R) (\varepsilon |u|^2 + C_\varepsilon |u|^{2^*}) dx \\ &\leq -2(N+1) \int_{\mathbb{R}^N} F(u) dx + 2CR^{-2} \|N - \Delta \psi_R\|_{L^\infty} \|\nabla u\|_{L^2(|x| \geq R)}^2, \\ I_3 &= \langle u(t), i\Gamma_{\psi_R} f(u) \rangle \\ &= -\langle u(t), (\nabla \cdot ((\nabla \psi_R) \, f(u)) + (\nabla \psi_R) \cdot (\nabla f(u))) \rangle \\ &\leq 2 \int_{\mathbb{R}^N} \nabla \psi_R \cdot \nabla \left(F(u) \right) dx \\ &= -2 \int_{\mathbb{R}^N} (\Delta \psi_R) \, F(u) dx \end{split}$$

$$= -2(N+1) \int_{\mathbb{R}^N} F(u) dx - 2 \int_{|x| \ge R} (\Delta \psi_R - N) F(u) dx$$

$$\leq -2(N+1) \int_{\mathbb{R}^N} F(u) dx + 2C \int_{|x| \ge R} (N - \Delta \psi_R) (|u|^2 + |u|^{2^*}) dx$$

$$\leq -2(N+1) \int_{\mathbb{R}^N} F(u) dx + 2CR^{-2} ||N - \Delta \psi_R||_{L^{\infty}} ||\nabla u||_{L^2(|x| \ge R)}^2,$$

Therefore,

$$\frac{d}{dt}\mathcal{M}_{\psi_R}[u(t)] \leq 8 \|\nabla u\|_{L^2}^2 + CR^{-2} - 4(N+1) \int_{\mathbb{R}^N} F(u) dx + 4CR^{-2} \|N - \Delta \psi_R\|_{L^\infty} \|\nabla u\|_{L^2(|x| \geq R)}^2,$$

where the constant C > 0 is independent of R. When R > 1 is sufficiently large, then

$$\frac{d}{dt}\mathcal{M}_{\psi_R}[u(t)] \le 8E(u(t)) = 8E(u_0).$$

Proof of Theorem 1.2. By lemma 3.1, for a radial initial function $\varphi \in H^1_{rad}$, (1.1) admits a local solution $u \in H^1_{rad}$. If $T^* < \infty$, then we are done. If $T^* = \infty$, we show (1.4). We suppose that u(t) exists for all times $t \geq 0$, i.e. $T^* = \infty$. It follows from Lemma 4.1 and conservation of mass, for $R \gg 1$ large enough,

$$\frac{d}{dt}\mathcal{M}_{\psi_R}[u(t)] \le 8E(u_0) := -A^* < 0, \quad t \ge 0.$$

From this, we infer that

$$\mathcal{M}_{\psi_R}[u(t)] \le -A^*t + \mathcal{M}_{\psi_R}[u_0], \quad t \ge 0.$$

On the one hand, let $T_0 = \frac{2|\mathcal{M}_{\psi_R}[u_0]|}{A^*} > 0$, then for any $t \geq T_0$, we have

$$\mathcal{M}_{\psi_R}[u(t)] \le -\frac{1}{2}A^*t < 0.$$
 (4.3)

On the other hand, by Lemma 2.4 and the conservation of mass, we see that for any $t \in [0, +\infty)$,

$$|\mathcal{M}_{\psi_{R}}[u(t)]| \leq C(\psi_{R}) \left(\left\| |\nabla|^{\frac{1}{2}} u(t) \right\|_{L^{2}}^{2} + \|u(t)\|_{L^{2}} \left\| |\nabla|^{\frac{1}{2}} u(t) \right\|_{L^{2}} \right)$$

$$\leq C(\psi_{R}) \left(\|\nabla u\|_{L^{2}} \|u\|_{L^{2}} + \|u(t)\|_{L^{2}}^{\frac{3}{2}} \|\nabla u\|_{L^{2}}^{\frac{1}{2}} \right)$$

$$\leq C(\psi_{R}) \|\nabla u\|_{L^{2}}^{2},$$

where we have used the interpolation estimate

$$\left\| |\nabla|^{\frac{1}{2}} u \right\|_{L^{2}}^{2} \le C \left\| \nabla u \right\|_{L^{2}} \left\| u \right\|_{L^{2}}.$$

This combined with (4.3) yields that for any $t \geq T_0$,

$$A^*t \le -2\mathcal{M}_{\psi_R}[u(t)] \le C \left\| \nabla u \right\|_{L^2}^2.$$

This shows that

$$\|\nabla u(t)\|_{L^2}^2 \ge Ct, \quad t \ge T_0.$$

It means that

$$\sup_{t>0} \|\nabla u(\cdot,t)\|_{L^2} = \infty.$$

5 Perturbation theory

In this section, we will study the long-time perturbation theory.

Proposition 5.1. (Perturbation theory) Let $\tilde{u}: I \times \mathbb{R}^N \to \mathbb{C}$ be a solution to the perturbed Schrödinger equation with general nonlinearity

$$i\partial_t \tilde{u} + \Delta \tilde{u} + f(\tilde{u}) = e$$

for some function e. Suppose that

$$\|\tilde{u}\|_{L^{\infty}H^{1}(I\times\mathbb{R}^{N})} \le E,\tag{5.1}$$

$$\|\tilde{u}\|_{L^{\frac{2(N+2)}{N-2}}L^{\frac{2(N+2)}{N-2}}(I\times\mathbb{R}^N)} \le L \tag{5.2}$$

for some E, L > 0. Let $u_0 \in H^1(\mathbb{R}^N)$ with $||u_0||_{L^2(\mathbb{R}^N)} \leq M$ for some M > 0 and let $t_0 \in I$. There exists $\varepsilon_0 = \varepsilon_0(E, L, M) > 0$ such that if

$$||u_0 - \tilde{u}(t_0)||_{H^1} \le \varepsilon, \tag{5.3}$$

$$||e||_{L_t^\infty H^1 \cap L_t^2 L^{\frac{2N}{N+2}}} \le \varepsilon \tag{5.4}$$

for $0 < \varepsilon < \varepsilon_0$, then the unique global solution u to (1.1) with $u(t_0) = u_0$ satisfies

$$||u - \tilde{u}||_{L_I^{\infty} H^1 \cap L_I^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}} \le C(E, L, M)\varepsilon, \tag{5.5}$$

where C(E, L, M) is a non-decreasing function of E, L, and M.

Proof. Without loss of generality, we may assume $t_0 = 0 \in I$. From Theorem 1.1 we know that u exists globally and

$$||u||_{L_I^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}} \le \rho,$$

so we need to get (5.5).

Let

$$w := u - \tilde{u} \text{ and } A(t) := \|w\|_{L^{\infty}_{(I \cap [-T,T])}H^1 \cap L^{\frac{2(N+2)}{N-2}}_{(I \cap [-T,T])}W^{1,\frac{2N(N+2)}{N^2+4}}}.$$

Note that

$$i\partial_t \tilde{w} + \Delta \tilde{w} - f(\tilde{u}) + f(\tilde{u} + w) = -e,$$

which is equivalent to the integral equation

$$w(t) = e^{it\Delta}w_0 + i \int_0^t e^{i(t-t')\Delta} [-f(\tilde{u}) + f(\tilde{u} + w)]dt' + i \int_0^t e^{i(t-t')\Delta}edt'.$$

Then by Strichartz, Hölder, (5.1), (5.2), (5.3), (5.4), we obtain

$$\begin{split} & \left\|w(t)\right\|_{L^{\infty}_{(I\cap[-T,T])}H^{1}\cap L^{\frac{2(N+2)}{N-2}}_{(L(\Gamma)-T,T])}W^{1,\frac{2N(N+2)}{N^{2}+4}}} \\ & = \left\|e^{it\Delta}w_{0} + i\int_{0}^{t}e^{i(t-t')\Delta}[-f(\tilde{u}) + f(\tilde{u}+w)]dt' + i\int_{0}^{t}e^{i(t-t')\Delta}edt'\right\|_{L^{\infty}_{T}H^{1}\cap L^{\frac{2(N+2)}{N^{2}+4}}} \\ & \leq C\|w_{0}\|_{H^{1}} + \left\|\int_{0}^{t}e^{i(t-t')\Delta}edt'\right\|_{L^{\infty}_{T}H^{1}} + \left\|\int_{0}^{t}e^{i(t-t')\Delta}wdt'\right\|_{L^{\infty}_{T}H^{1}} \\ & + \left\|\int_{0}^{t}e^{i(t-t')\Delta}[-\tilde{u}\cdot|\tilde{u}|^{\frac{4}{N-2}} + (\tilde{u}+w)\cdot|\tilde{u}+w|^{\frac{4}{N-2}}]dt'\right\|_{L^{\frac{2(N+2)}{N^{2}+4}}} \\ & \leq C\|w_{0}\|_{H^{1}} + \|w\|_{L^{1}_{T}H^{1}} + \|e\|_{L^{1}_{T}H^{1}} + C\left\||w|\cdot(|\tilde{u}+w|^{\frac{4}{N-2}} + |\tilde{u}|^{\frac{4}{N-2}})\right\|_{L^{2}_{T}W^{1,\frac{2N}{N+2}}} \\ & \leq C\|w_{0}\|_{H^{1}} + T\|w\|_{L^{\infty}_{T}H^{1}} + T\|e\|_{L^{\infty}_{T}H^{1}} + C\left\||w|\cdot|\tilde{u}+w|^{\frac{4}{N-2}}\right\|_{L^{2}_{T}W^{1,\frac{2N}{N+2}}} \\ & \leq C\|w_{0}\|_{H^{1}} + T\|w\|_{L^{\infty}_{T}H^{1}} + T\|e\|_{L^{\infty}_{T}H^{1}} + C\|w\|_{L^{\frac{2(N+2)}{N^{2}}}_{T^{\frac{N-2}{N^{2}}}}W^{1,\frac{2N(N+2)}{N^{2}+4}}\cdot\|\tilde{u}\|^{\frac{4-2}{N^{2}}}_{L^{\frac{2(N+2)}{N^{2}}}_{T^{\frac{N-2}{N^{2}}}} \\ & \leq C\|w_{0}\|_{H^{1}} + T\|w\|_{L^{\infty}_{T}H^{1}} + T\|e\|_{L^{\infty}_{T}H^{1}} + C\|w\|_{L^{\frac{2(N+2)}{N^{2}}}_{T^{\frac{N-2}{N^{2}}}}W^{1,\frac{2N(N+2)}{N^{2}+4}}\cdot\|\tilde{u}\|^{\frac{4-2}{N^{2}}}_{L^{\frac{2(N+2)}{N^{2}}}_{T^{\frac{N-2}{N^{2}}}} \\ & \leq C\|w_{0}\|_{H^{1}} + T\|w\|_{L^{\infty}_{T}H^{1}} + T\|e\|_{L^{\infty}_{T}H^{1}} + C\|w\|_{L^{\frac{2(N+2)}{N^{2}}}_{T^{\frac{N-2}{N^{2}}}}W^{1,\frac{2N(N+2)}{N^{2}+4}}\cdot\|\tilde{u}\|^{\frac{4-2}{N^{2}}}_{L^{\frac{2(N+2)}{N^{2}}}_{T^{\frac{N-2}{N^{2}}}} \\ & \leq C\|w_{0}\|_{H^{1}} + T\|w\|_{L^{\infty}_{T}H^{1}} + T\|e\|_{L^{\infty}_{T}H^{1}} + C\|w\|_{L^{\frac{2(N+2)}{N^{2}}}_{T^{\frac{N-2}{N^{2}}}}W^{1,\frac{2N(N+2)}{N^{2}+4}}\cdot\|\tilde{u}\|^{\frac{4-2}{N^{2}}}_{L^{\frac{N-2}{N^{2}}}} \\ & \leq C\|w_{0}\|_{H^{1}} + T\|w\|_{L^{\infty}_{T}H^{1}} + T\|e\|_{L^{\infty}_{T}H^{1}} + C\|w\|_{L^{\frac{2(N+2)}{N^{2}}}W^{1,\frac{2N(N+2)}{N^{2}+4}}}\cdot\|\tilde{u}\|^{\frac{4-2}{N^{2}}}_{L^{\frac{N-2}{N^{2}}} \\ & \leq C\|w_{0}\|_{H^{1}} + T\|w\|_{L^{\infty}_{T}H^{1}} + T\|e\|_{L^{\infty}_{T}H^{1}} + C\|w\|_{L^{\infty}_{T}H^{1}} + C\|w\|_{$$

where all space time norms are over $(I \cap [-T, T]) \times \mathbb{R}^N$. Using the standard continuity argument to remove the restriction to [-T, T], we derive (5.5).

Remark 5.1. Note that $f(u) \in L_T^{\infty} H^1 \cap L_T^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}$ and hence

$$\left\| \int_{t}^{\infty} e^{i(t-t')\Delta} f(u) dt' \right\|_{H^{1}} \to 0, \ t \to +\infty.$$

Then, $u(t) = e^{i(t-a)\Delta}u_0 + \int_a^t e^{i(t-t')\Delta}f(u)dt'$ and hence $u^+ = e^{-ia\Delta}u_0 + \int_a^\infty e^{-it'\Delta}f(u)dt'$ has the desired property. In fact note that the argument used at the beginning of the proof of Proposition 5.1 shows that it suffices to assume u to be a solution of (1.1) in $I' \times \mathbb{R}^N$, $I' \in I$, such that $\|u\|_{L_r^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}} < \infty$.

6 Some variational estimates

In this section, let us first recall some variational results. If we look for standing wave solutions (form $\Phi(t,x) = e^{-imt}u(x)$) about the following equation

$$i\partial_t \Phi + \Delta \Phi + f(\Phi) = 0, \ (x, t) \in \mathbb{R}^N \times \mathbb{R},$$
 (6.1)

then (6.1) becomes non-linear elliptic equation

$$\Delta u + f(u) = 0, \ (x, t) \in \mathbb{R}^N \times \mathbb{R}. \tag{6.2}$$

As described in [2], if we add condition (F_6) under the assumptions of $(F_1) - (F_5)$, that is,

 (F_6) There exists $\xi > 0$ such that $F(\xi) = \int_0^{\xi} f(s)ds > 0$.

then we obtain the existence of a solution of (6.2), i.e., (6.2) possesses a solution u such that

- i) u > 0 on \mathbb{R}^N .
- ii) u is spherically symmetric: u(x) = u(r), where r = |x|, and u decreases with respect to r.
 - iii) $u \in C^2(\mathbb{R}^N)$.
 - iv) u together with its derivatives up to order 2 have exponential decay at infinity:

$$|D^{\alpha}u(x)| \le Ce^{-\delta|x|}, \quad x \in \mathbb{R}^N,$$

for some $C, \delta > 0$ and for $|\alpha| \leq 2$.

Now, we assume that the solution of (6.2) is W(x). The equation (6.2) gives

$$\int_{\mathbb{R}^N} |\nabla W|^2 dx = \int_{\mathbb{R}^N} F(u) dx, \ F(u) = \int_0^u f(s) ds.$$

Moreover, note that W(x) is a solution of (6.2), so we have the following Pohozaev identity

$$\frac{N-2}{2} \int_{\mathbb{R}^N} |\nabla W|^2 dx = N \int_{\mathbb{R}^N} F(W) dx.$$

Hence,

$$E(W) = \frac{1}{2} \int_{\mathbb{R}^N} |\nabla W|^2 dx - \int_{\mathbb{R}^N} F(W) dx = \frac{1}{N} \int_{\mathbb{R}^N} |\nabla W|^2 dx.$$

Lemma 6.1. Assume u satisfies

$$\|\nabla u\|_{L^2}^2 < \|\nabla W\|_{L^2}^2.$$

Moreover, let $E(u) \leq (1 - \delta_0)E(W)$, where $\delta_0 > 0$. Then, there exists $\delta_1 > 0$, $\bar{\delta} > 0$ such that

$$\int_{\mathbb{R}^N} |\nabla u|^2 dx \le (1 - \delta_1) \int_{\mathbb{R}^N} |\nabla W|^2 dx, \tag{6.3}$$

$$\int_{\mathbb{R}^N} (|\nabla u|^2 - F(u)) dx \ge \bar{\delta} \int_{\mathbb{R}^N} |\nabla u|^2 dx, \tag{6.4}$$

$$\int_{\mathbb{R}^N} (|\nabla u|^2 - f(u)\overline{u}) dx \ge \bar{\delta} \int_{\mathbb{R}^N} |\nabla u|^2 dx, \tag{6.5}$$

$$E(u) \ge 0. \tag{6.6}$$

Proof. In order to get (6.4), by $(F_1) - (F_2)$, we have, for any $\varepsilon > 0$,

$$F(u) \le \varepsilon |u|^2 + C_{\varepsilon} |u|^{2^*}.$$

Hence, it holds

$$\int_{\mathbb{R}^{N}} (|\nabla u|^{2} - F(u)) dx \geq \int_{\mathbb{R}^{N}} |\nabla u|^{2} dx - \varepsilon ||u||_{L^{2}}^{2} - C_{\varepsilon} ||u||_{L^{2}}^{2^{*}}$$

$$= ||\nabla u||_{L^{2}}^{2} - \varepsilon ||\nabla u||_{L^{2}}^{2} - C_{\varepsilon} ||\nabla u||_{L^{2}}^{2^{*}}$$

$$:= \bar{\delta} \int_{\mathbb{R}^{N}} |\nabla u|^{2} dx,$$

which implies (6.4) holds. Similarly, it can be proven that (6.7) is established. To prove (6.3), by $E(u) \leq (1 - \delta_0)E(W)$ and $\|\nabla u\|_{L^2}^2 < \|\nabla W\|_{L^2}^2$, we have

$$\frac{1}{2} \int_{\mathbb{R}^N} |\nabla u|^2 dx - \int_{\mathbb{R}^N} F(u) dx \le (1 - \delta_0) E(W) = \frac{1 - \delta_0}{2N} \int_{\mathbb{R}^N} |\nabla W|^2 dx,$$

SO

$$\begin{split} \int_{\mathbb{R}^{N}} |\nabla u|^{2} dx & \leq & \frac{1}{2} \int_{\mathbb{R}^{N}} |\nabla u|^{2} dx + \frac{1 - \delta_{0}}{2N} \int_{\mathbb{R}^{N}} |\nabla W|^{2} dx + \int_{\mathbb{R}^{N}} F(u) dx \\ & \leq & \frac{1}{2} \int_{\mathbb{R}^{N}} |\nabla W|^{2} dx + \frac{1 - \delta_{0}}{2N} \int_{\mathbb{R}^{N}} |\nabla W|^{2} dx + \varepsilon \|\nabla u\|_{L^{2}}^{2} + C_{\varepsilon} \|\nabla u\|_{L^{2}}^{2^{*}} \\ & \leq & \frac{1}{2} \int_{\mathbb{R}^{N}} |\nabla W|^{2} dx + \frac{1 - \delta_{0}}{2N} \int_{\mathbb{R}^{N}} |\nabla W|^{2} dx + \varepsilon \|\nabla W\|_{L^{2}}^{2} + C_{\varepsilon} \|\nabla W\|_{L^{2}}^{\frac{2^{*}}{2}} \\ & := & (1 - \delta_{1}) \int_{\mathbb{R}^{N}} |\nabla W|^{2} dx. \end{split}$$

According to the proof of (6.4), (6.6) obviously holds, this completes the proof.

Remark 6.1. From Lemma 6.1, we know that the selection of W is not arbitrary. In fact, we need to choose W such that $\int_{\mathbb{R}^N} |\nabla W|^2 dx$ is small. Moreover, according to the conditions of the nonlinear term and f is odd, there are actually infinite standing wave solutions for equation (1.1), see [3]. In this case, we only need to take the solution that minimizes $\int_{\mathbb{R}^N} |\nabla W|^2 dx$.

Corollary 6.1. Assume that $u \in H^1$ and $\int_{\mathbb{R}^N} |\nabla u|^2 dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx$, then $E(u) \geq 0$.

Proof. If $E(u) \geq E(W) = \frac{1}{N} \int_{\mathbb{R}^N} |\nabla W|^2 dx$, this is obvious. If E(u) < E(W), the claim follows from (6.6).

Theorem 6.1. (Energy trapping). Let u be a solution of the (1.1), with $t_0 = 0$, $u|_{t=0} = u_0$ such that for $\delta_0 > 0$,

$$\int_{\mathbb{R}^N} |\nabla u_0|^2 \, dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx, \ E(u_0) < (1 - \delta_0) E(W).$$

Let $I(0 \in I)$ be the maximal interval of existence. Let δ_0 , $\bar{\delta}$ be as in Lemma 6.1. Then, for each $t \in I$, we have

$$\int_{\mathbb{R}^N} |\nabla u(t)|^2 dx \le (1 - \delta_1) \int_{\mathbb{R}^N} |\nabla W|^2 dx,\tag{6.7}$$

$$\int_{\mathbb{R}^N} (|\nabla u(t)|^2 - F(u(t))) dx \ge \bar{\delta} \int_{\mathbb{R}^N} |\nabla u(t)|^2 dx, \tag{6.8}$$

$$\int_{\mathbb{R}^N} (|\nabla u(t)|^2 - f(u(t))\overline{u}(t))dx \ge \bar{\delta} \int_{\mathbb{R}^N} |\nabla u(t)|^2 dx, \tag{6.9}$$

$$E(u(t)) \ge 0. \tag{6.10}$$

Proof. By energy conservation, $E(u(t)) = E(u_0)$, $t \in I$ and the theorem follows directly from Lemma 6.1 and a continuity argument.

Corollary 6.2. Let u, u_0 be as in Theorem 6.1. Then for all $t \in I$ we have $E(u(t)) \simeq \int_{\mathbb{R}^N} |\nabla u(t)|^2 dx \simeq \int_{\mathbb{R}^N} |\nabla u_0|^2 dx$, with comparability constants which depend only on N.

Proof. $E(u(t)) \leq \int_{\mathbb{R}^N} |\nabla u(t)|^2 dx$, but by the proof of (6.4) there exists $\widetilde{\delta}$ such that

$$E(u(t)) = \frac{1}{2} \int_{\mathbb{R}^N} |\nabla u(t, x)|^2 dx - \int_{\mathbb{R}^N} F(u(t, x)) dx$$
$$\geq \widetilde{\delta} \int_{\mathbb{R}^N} |\nabla u(t)|^2 dx,$$

so the first equivalence follows. For the second one note that

$$E(u(t)) = E(u_0) \simeq \int_{\mathbb{R}^N} |\nabla u_0|^2 dx,$$

by the first equivalence when t=0.

7 Concentration compactness

The main purpose of this section is to prove the following concentration compactness lemma, which plays a crucial role in the proof of the theorem in the next section.

Lemma 7.1. (Concentration compactness) Let $\{v_{0,n}\} \in H^1, \|v_{0,n}\|_{H^1} \leq \rho$. Assume that $\|e^{it\Delta}v_{0,n}\|_{L^{\frac{2(N+2)}{N-2}}_{t,x}} \geq \rho > 0$, where ρ is as in Lemma 3.1. Then there exists a sequence $\{V_{0,j}\}_{j=1}^{\infty}$ in H^1 , a subsequence of $\{v_{0,n}\}$ (which we still call $\{v_{0,n}\}$) and a triple $(\lambda_{j,n}; x_{j,n}; t_{j,n}) \in \mathbb{R}^+ \times \mathbb{R}^N \times \mathbb{R}$, with

$$\frac{\lambda_{j,n}}{\lambda_{j',n}} + \frac{\lambda_{j',n}}{\lambda_{j,n}} + \frac{|t_{j,n} - t_{j',n}|}{\lambda_{j,n}^2} + \frac{|x_{j,n} - x_{j',n}|}{\lambda_{j,n}} \to \infty$$

as $n \to \infty$ for $j \neq j'$ (we say that $(\lambda_{j,n}; x_{j,n}; t_{j,n})$ is orthogonal if this property is verified) such that

$$||V_{0,1}||_{H^1} \ge \alpha_0(\rho) > 0. \tag{7.1}$$

If $V_{i}^{l}(x,t)=e^{it\Delta}V_{0,j}$, then, given $\varepsilon_{0}>0$, there exists $J=J\left(\varepsilon_{0}\right)$ and

$$\{w_n\}_{n=1}^{\infty} \in H^1 \text{ so that } v_{0,n} = \sum_{j=1}^{J} \frac{1}{\lambda_{j,n}^{\frac{N-2}{2}}} V_j^l \left(\frac{x - x_{j,n}}{\lambda_{j,n}}, \frac{-t_{j,n}}{\lambda_{j,n}^2}\right) + w_n$$
 (7.2)

with $\|e^{it\Delta}w_n\|_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,+\infty)}W^{1,\frac{2N(N+2)}{N^2+4}}} \le \varepsilon_0$, for n large

$$\int_{\mathbb{R}^N} |\nabla v_{0,n}|^2 dx = \sum_{j=1}^J \int_{\mathbb{R}^N} |\nabla V_{0,j}|^2 dx + \int_{\mathbb{R}^N} |\nabla w_n|^2 dx + o(1), \ n \to \infty,$$
 (7.3)

$$E(v_{0,n}) = \sum_{j=1}^{J} E(V_j^l(\frac{-t_{j,n}}{\lambda_{j,n}^2})) + E(w_n) + o(1), \ n \to \infty.$$
 (7.4)

The proof of this lemma originates from Keraani [18], but we need to modify the proof since this paper considers general nonlinear terms. Firstly, we consider linear equation

$$\begin{cases}
i\partial_t u + \Delta u = 0, & (x,t) \in \mathbb{R}^N \times \mathbb{R}, \\
u(0,x)|_{t=0} = u_0(x) \in H^1(\mathbb{R}^N).
\end{cases}$$
(7.5)

Lemma 7.2. Let $(\varphi_n)_{n\geq 0}$ be a bounded sequence in $H^1(\mathbb{R}^N)$. Let $(v_n)_{n\geqslant 0}$ be the sequence of solutions to (7.5) with initial data $v_n(0,x)=\varphi_n(x)$. Then there exist a subsequence (v'_n) of (v_n) , a sequence $(\mathbf{h}^j)_{j\geqslant 1}$ of scales, a sequence $(\mathbf{z}^j)_{j\geqslant 1}$ of cores and a sequence $(V^j)_{j\geqslant 1}$ of solutions to (7.5), such that

- (i) the pairs $(\mathbf{h}^j, \mathbf{z}^j)$ are pairwise orthogonal;
- (ii) for every $l \geq 1$,

$$v_n'(x,t) = \sum_{j=1}^l \frac{1}{(h_n^j)^{\frac{N-2}{2}}} V^j \left(\frac{x - x_n^j}{h_n^j}, \frac{t - t_n^j}{(h_n^j)^2} \right) + w_n^l(x,t), \tag{7.6}$$

with

$$\limsup_{n \to \infty} \|w_n^l\|_{L^q(\mathbb{R}, L^r(\mathbb{R}^N))} \to 0, \ l \to \infty$$
 (7.7)

for every H^1 -admissible pair (q,r) (defined in Definition 2.3), and, for every $l \ge 1$,

$$\int_{\mathbb{R}^{N}} |\nabla v_{n}'|^{2} dx = \sum_{j=1}^{l} \int_{\mathbb{R}^{N}} |\nabla V^{j}|^{2} dx + \int_{\mathbb{R}^{N}} |\nabla w_{n}^{l}|^{2} dx + o(1), \quad n \to \infty.$$
 (7.8)

Proof. As stated by [17]. It is based on the "refined Sobolev inequality" (N=3)

$$||h||_{L^6(\mathbb{R}^3)} \le C ||\nabla h||_{L^2(\mathbb{R}^3)}^{1/3} ||\nabla h||_{\dot{B}_{2,\infty}^0}^{2/3},$$

where $\dot{B}_{2,\infty}^0$ is the standard Besov space [4]. The proof of this lemma is completely similar to [17, 18], so we omit it here.

Lemma 7.3. Let (v'_n) be the subsequence of (v_n) given by Lemma 7.2. Let (u'_n) be the sequence of solutions to (1.1), with the same Cauchy data at t = 0 as v'_n . Then

$$u'_n(x,t) = \sum_{j=1}^l \frac{1}{(h_n^j)^{\frac{N-2}{2}}} U^j \left(\frac{x - x_n^j}{h_n^j}, \frac{t - t_n^j}{\left(h_n^j\right)^2} \right) + w_n^l(x,t) + r_n^l(x,t), \tag{7.9}$$

with

$$\limsup_{n\to\infty} \left(\sup_{t\in\mathbb{R}} E\left(r_n^l(t,\cdot)\right) + \left\|r_n^l\right\|_{L_{t,x}^{\frac{2(N+2)}{N-2}}(\mathbb{R}\times\mathbb{R}^N)} + \left\|\nabla r_n^l\right\|_{L_{t,x}^{\frac{2(2+N)}{N}}(\mathbb{R}\times\mathbb{R}^N)} \right) \to 0, \ l\to\infty,$$

where V^j , \mathbf{h}^j , \mathbf{z}^j , w_n^l are as in (7.6) and U^j is the nonlinear profile associated to $(V^j, \mathbf{h}^j, \mathbf{z}^j)$ (defined in Definition 2.4).

Proof. Let us first restate the problem. Let $(\varphi_n)_{n\geqslant 0}$ be a bounded sequence in $H^1(\mathbb{R}^N)$, such that $\limsup_{n\to\infty} \|u_n(0,.)\|_{H^1(\mathbb{R}^N)} < \rho$ (where ρ is given by Lemma 3.2). Let $(v_n)_{n\geqslant 0}$ (resp. $(u_n)_{n\geqslant 0}$) the sequence of solutions to (7.5) (resp. to (1.1)) with initial data $(\varphi_n)_{n\geqslant 0}$. Lemma 7.2 provides a decomposition of $(v_n)_{n\geqslant 0}$ for a subsequence (v'_n) of (v_n) in the form

$$v'_n(x,t) = \sum_{j=1}^l \frac{1}{(h_n^j)^{\frac{N-2}{2}}} V^j \left(\frac{x - x_n^j}{h_n^j}, \frac{t - t_n^j}{\left(h_n^j\right)^2} \right) + w_n^l(x,t),$$

where (V^j) is a family of solutions to (7.5), $(\mathbf{h}^j, \mathbf{z}^j)_{j \geq 1}$ is a pairwise orthogonal family of scales-cores, and the remainder w_n^l satisfies

$$\limsup_{n\to\infty} \|w_n^l\|_{L^{\frac{2(N+2)}{N-2}}_{t,x}(\mathbb{R}\times\mathbb{R}^N)}\to 0,\ l\to\infty.$$

Also, the following almost orthogonality identity holds

$$\int_{\mathbb{R}^N} |\nabla v_n'|^2 dx = \sum_{j=1}^l \int_{\mathbb{R}^N} |\nabla V^j|^2 dx + \int_{\mathbb{R}^N} |\nabla w_n^l|^2 dx + o(1), \quad n \to \infty.$$

Let (u'_n) be the sequence of solutions of (7.5), with the same Cauchy data at t=0 as v'_n and U^j the nonlinear profile associated to $(V^j, \mathbf{h}^j, \mathbf{z}^j)$ for every $j \geq 1$. Observe that, in view of (7.8), $\int_{\mathbb{R}^N} (V^j) dx < \rho^2$ and then the nonlinear profile U^j is globally well defined. We set

$$r_n^l(x,t) = u_n'(x,t) - \sum_{j=1}^l \frac{1}{(h_n^j)^{\frac{N-2}{2}}} U^{(j)} \left(\frac{x - x_n^j}{h_n^j}, \frac{t - t_n^j}{\left(h_n^j\right)^2} \right) - w_n^l(x,t).$$

Our purpose is to prove that

$$\limsup_{n \to \infty} \left(\sup_{t \in \mathbb{R}} E\left(r_n^l(t,.)\right) + \left\| r_n^l \right\|_{L_{t,x}^{\frac{2(N+2)}{N-2}}(\mathbb{R} \times \mathbb{R}^N)} + \left\| \nabla r_n^l \right\|_{L_{t,x}^{\frac{2(2+N)}{N}}(\mathbb{R} \times \mathbb{R}^N)} \right) \to 0, \ l \to \infty.$$

The following notations will be used

$$\begin{split} &p(z) = f(z), \\ &V_n^j(x,t) = \frac{1}{(h_n^j)^{\frac{N-2}{2}}} V^j \left(\frac{x - x_n^j}{h_n^j}, \frac{t - t_n^j}{\left(h_n^j\right)^2} \right), \\ &U_n^j(x,t) = \frac{1}{(h_n^j)^{\frac{N-2}{2}}} U^j \left(\frac{x - x_n^j}{h_n^j}, \frac{t - t_n^j}{\left(h_n^j\right)^2} \right), \\ &W_n^l = \sum_{j=1}^l U_n^j, \\ &f_n^l = \sum_{j=1}^l p\left(U_n^j \right) - p\left(\sum_{j=1}^l U_n^j + w_n^l + r_n^l \right). \end{split}$$

The function r_n^l provided by (7.9) satisfies the equation

$$\begin{cases} i\partial_t r_n^l + \Delta r_n^l = f_n^l \\ r_n^l(x,0) = \sum_{j=1}^l (V_n^j - U_n^j)(x,0). \end{cases}$$

We also introduce the norm

$$|||g|||_I = ||\nabla g||_{L_{t,x}^{\frac{2(N+2)}{N}}(I \times \mathbb{R}^N)} + ||g||_{L_{t,x}^{\frac{2(N+2)}{N-2}}(I \times \mathbb{R}^N)}.$$

Note that, by Strichartz estimates (2.1) and (2.2), we get

$$|||e^{it\Delta}\varphi|||_{\mathbb{R}} \le C||\varphi||_{H^1(\mathbb{R}^N)}$$

for every $\varphi \in H^1(\mathbb{R}^N)$. In the rest of the paper we note, for every $a \in \mathbb{R}$,

$$\gamma_n^l(a) = \left\| \nabla r_n^l(\cdot, a) \right\|_{L^2(\mathbb{R}^N)}. \tag{7.10}$$

Recall that our purpose is to prove that

$$\limsup_{n \to \infty} \left(\sup_{t \in \mathbb{R}} E\left(r_n^l(\cdot, t)\right) + |||r_n^l|||_{\mathbb{R}} \right) \to 0, \ l \to \infty.$$

The following lemma is a combination of Strichartz estimates and energy inequalities, see [10, 18, 19].

Lemma 7.4. Let $v \in C([a,b], \dot{H}^1(\mathbb{R}^N))$ be a solution on I = [a,b] of the nohomogeneous Schrödinger equation

$$i\partial_t v + \Delta v = f$$

with $\nabla f \in L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^N)$. Then we have

$$|||v|||_I + \sup_{t \in I} ||\nabla v(t)||_{L^2(\mathbb{R}^N)} \le C \left(||\nabla v(a)||_{L^2(\mathbb{R}^N)} + ||\nabla f||_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^N)} \right).$$

Applying Lemma 7.4 to r_n^l on I = [a, b], it holds

$$|||r_n^l|||_I + \sup_{t \in I} ||\nabla r_n^l(t)||_{L^2(\mathbb{R}^N)} \le C \left(\gamma_n^l(a) + ||\nabla f_n^l||_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^N)} \right).$$
(7.11)

Next, we estimate

$$\|\nabla f_{n}^{l}\|_{L^{\frac{2(N+2)}{N+4}}(I\times\mathbb{R}^{N})}$$

$$= \|\nabla \left[\sum_{j=1}^{l} p\left(U_{n}^{j}\right) - p\left(\sum_{j=1}^{l} U_{n}^{j} + w_{n}^{l} + r_{n}^{l}\right)\right]\|_{L^{\frac{2(N+2)}{N+4}}(I\times\mathbb{R}^{N})}$$

$$\leq \|\nabla \left[\sum_{j=1}^{l} p\left(U_{n}^{j}\right) - p\left(W_{n}^{l}\right)\right]\|_{L^{\frac{2(N+2)}{N+4}}(I\times\mathbb{R}^{N})} + \|\nabla \left(p\left(W_{n}^{l} + w_{n}^{l}\right) - p\left(W_{n}^{l}\right)\right)\|_{L^{\frac{2(N+2)}{N+4}}(I\times\mathbb{R}^{N})}$$

$$+ \|\nabla \left(p\left(W_{n}^{l} + w_{n}^{l} + r_{n}^{l}\right) - p\left(W_{n}^{l} + w_{n}^{l}\right)\right)\|_{L^{\frac{2(N+2)}{N+4}}(I\times\mathbb{R}^{N})}.$$

$$(7.12)$$

Furthermore, a combination of Leibnitz formula, Hölder inequality and $(F_1) - (F_4)$ gives

$$\begin{split} & \left\| \nabla \left(p \left(W_n^l + w_n^l + r_n^l \right) - p \left(W_n^l + w_n^l \right) \right) \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^N)} \\ & \leq & C \left\| |\nabla r_n^l| \left(\left(W_n^l + w_n^l + r_n^l \right)^{\frac{4}{N-2}} + \left(W_n^l + w_n^l \right)^{\frac{4}{N-2}} \right) \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^N)} + \varepsilon \| \nabla r_n^l \|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^N)} \\ & \leq & C \left\| |\nabla r_n^l| \left(W_n^l + w_n^l + r_n^l \right)^{\frac{4}{N-2}} \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^N)} + \left\| |\nabla r_n^l| \left(W_n^l + w_n^l \right)^{\frac{4}{N-2}} \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^N)} \\ & + \varepsilon \| \nabla r_n^l \|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^N)} \cdot \left\| \left(W_n^l + w_n^l + r_n^l \right)^{\frac{4}{N-2}} \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^N)} + \varepsilon \| \nabla r_n^l \|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^N)} \\ & + \left\| |\nabla r_n^l| \left(W_n^l + w_n^l \right)^{\frac{4}{N-2}} \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^N)} \end{split}$$

$$\leq C \left\| \nabla r_{n}^{l} \right\|_{L^{\frac{2(N+2)}{N}}(I \times \mathbb{R}^{N})} \cdot \left\| W_{n}^{l} + w_{n}^{l} + r_{n}^{l} \right\|_{L^{\frac{2(N+2)}{N-2}}(I \times \mathbb{R}^{N})}^{\frac{4}{N-2}} + \varepsilon \left\| \nabla r_{n}^{l} \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^{N})}^{\frac{4}{N-2}} \\ + \left\| \left| \nabla r_{n}^{l} \right| \left(W_{n}^{l} + w_{n}^{l} \right)^{\frac{4}{N-2}} \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^{N})}^{\frac{4}{N-2} - \alpha} \\ \leq C \left\| \nabla r_{n}^{l} \right\|_{L^{\frac{2(N+2)}{N}}(I \times \mathbb{R}^{N})} \cdot \sum_{\alpha=1}^{\frac{4}{N-2}} \left\| W_{n}^{l} + w_{n}^{l} \right\|_{L^{\frac{2(N+2)}{N-2}}(I \times \mathbb{R}^{N})}^{\frac{4}{N-2} - \alpha} \cdot \left\| (r_{n}^{l}) \right\|_{L^{\frac{2(N+2)}{N-2}}(I \times \mathbb{R}^{N})}^{\frac{2(N+2)}{N-2}(I \times \mathbb{R}^{N})} \\ + \left\| \left| \nabla r_{n}^{l} \right| \left(W_{n}^{l} + w_{n}^{l} \right)^{\frac{4}{N-2}} \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^{N})}^{\frac{4}{N-2}} + \varepsilon \left\| \nabla r_{n}^{l} \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^{N})}^{\frac{6-N}{N-2}} \left(W_{n}^{l} + w_{n}^{l} \right) \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^{N})}^{\frac{2(N+2)}{N+4}} \\ + \varepsilon \left\| \nabla r_{n}^{l} \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^{N})}^{\frac{4}{N-2} - \alpha} \cdot \left\| r_{n}^{l} \right\|_{I}^{\frac{\alpha+1}{N-2} + C} \left\| r_{n}^{l} \right\|_{I} \left\| W_{n}^{l} + w_{n}^{l} \right\|_{I}^{\frac{6-N}{N-2}}^{\frac{6-N}{N-2}} \left\| W_{n}^{l} + w_{n}^{l} \right\|_{L^{\frac{2(N+2)}{N-2}}(I \times \mathbb{R}^{N})}^{\frac{2(N+2)}{N-2}} \\ + \varepsilon \left\| \nabla r_{n}^{l} \right\|_{I}^{\frac{2(N+2)}{N-2} - \alpha} \left\| r_{n}^{l} \right\|_{I}^{\frac{\alpha+1}{N-2} - \alpha}^{\frac{1}{N-2}} \left\| r_{n}^{l} \right\|_{I}^{\frac{1}{N-2}}^{\frac{1}{N-2}} \left\| W_{n}^{l} + w_{n}^{l} \right\|_{I}^{\frac{2(N+2)}{N-2}}^{\frac{1}{N-2}} \left\| W_{n}^{l} + w_{n}^{l} \right\|_{L^{\frac{2(N+2)}{N-2}}(I \times \mathbb{R}^{N})}^{\frac{1}{N-2}} \\ + \varepsilon \left\| \nabla r_{n}^{l} \right\|_{I}^{\frac{2(N+2)}{N-2}}^{\frac{1}{N-2}} \left\| r_{n}^{l} \right\|_{I}^{\frac{\alpha+1}{N-2} - \alpha}^{\frac{1}{N-2}} \left\| r_{n}^{l} \right\|_{I}^{\frac{\alpha+1}{N-2}}^{\frac{1}{N-2}} \left\| r_{n}^{l} \right\|_{I}^{\frac{2(N+2)}{N-2}}^{\frac{1}{N-2}} \left$$

We set

$$\delta_{n}^{l} = \left\| \nabla \left[\sum_{j=1}^{l} p\left(U_{n}^{j}\right) - p\left(W_{n}^{l}\right) \right] \right\|_{L^{\frac{2(N+2)}{N+4}}(\mathbb{R} \times \mathbb{R}^{N})} + \left\| \nabla \left(p\left(W_{n}^{l} + w_{n}^{l}\right) - p\left(W_{n}^{l}\right) \right) \right\|_{L^{\frac{2(N+2)}{N+4}}(\mathbb{R} \times \mathbb{R}^{N})}.$$

$$(7.14)$$

Combining (7.11), (7.12), (7.13) and (7.14), we have

We prepare several propositions. The first one gives a uniform bound on $\|W_n^l + w_n^l\|_{\mathbb{R}}$.

Proposition 7.1. There exists C > 0, such that, for every $l \ge 1$,

$$\lim \sup_{n \to \infty} |||W_n^l + w_n^l||_{\mathbb{R}} \le C. \tag{7.16}$$

In view of (7.15) and Proposition 7.1, we get for every $l \ge 1$ and $n \ge N(l)$,

$$|||r_n^l|||_I + \sup_{t \in I} ||\nabla r_n^l(t)||_{L^2(\mathbb{R}^N)}$$

$$\leq C \left(\sum_{\alpha=1}^{\frac{4}{N-2}} \| |r_n^l| \|_I^{\alpha+1} + C \| |r_n^l| \|_I \| W_n^l + w_n^l) \|_{L^{\frac{2(N+2)}{N-2}}(I \times \mathbb{R}^N)} + C \gamma_n^l(a) + C \delta_n^l \right) . (7.17)$$

The second proposition shows that under a suitable finite partition of \mathbb{R} , one can absorb the linear term in $||r_n^l||_I$ in the right-hand side of (7.17).

Proposition 7.2. For every $\varepsilon > 0$, there exists an n-dependent finite partition of \mathbb{R}_+

$$\mathbb{R}_{+} = \underbrace{\left[0, a_{n}^{1}\right]}_{I_{n}^{1}} \cup \underbrace{\left[a_{n}^{1}, a_{n}^{2}\right]}_{I_{n}^{2}} \cup \cdots \cup \underbrace{\left[a_{n}^{p-1}, +\infty\right]}_{I_{n}^{p}}, \tag{7.18}$$

such that

$$\limsup_{n \to \infty} \left\| W_n^l + w_n^l \right\|_{L^{\frac{2(N+2)}{N-2}}(I_n^i \times \mathbb{R}^N)} \leqslant \varepsilon \tag{7.19}$$

for every $1 \le i \le p$ and every $l \ge 1$.

The third proposition proves the smallness of δ_n^l .

Proposition 7.3.

$$\lim_{n \to \infty} \sup \delta_n^l \to 0, \ l \to \infty. \tag{7.20}$$

Let us first assume the validity of Propositions 7.1, 7.2, 7.3, and use them to achieve the proof of Lemma 7.3. Applying (7.17) on an interval I_n^i , provided by Proposition 7.2, it follows that

$$|||r_n^l|||_{I_n^i} + \sup_{t \in I_n^i} ||\nabla r_n^l(t)||_{L^2(\mathbb{R}^N)} \le C \left(\gamma_n^l \left(a_n^i \right) + \delta_n^l + 2\varepsilon ||r_n^l||_{I_n^i} + \sum_{\alpha=2}^{\frac{N+2}{N-2}} |||r_n^l||_{I_n^i}^{\alpha} \right)$$
(7.21)

for every $l \geq 1$ and $n \geq N(l)$. If we choose ε so that $C\varepsilon < \frac{1}{2}$, we obtain

$$|||r_n^l|||_{I_n^i} + \sup_{t \in I_n^i} ||\nabla r_n^l(t)||_{L^2(\mathbb{R}^N)} \le C \left(\gamma_n^l \left(a_n^i \right) + \delta_n^l + \sum_{\alpha=2}^{\frac{N+2}{N-2}} |||r_n^l|||_{I_n^i}^{\alpha} \right). \tag{7.22}$$

For i = 1, (7.22) reads

$$|||r_n^l|||_{I_n^1} + \sup_{t \in I_n^1} ||\nabla r_n^l(t)||_{L^2(\mathbb{R}^N)} \le C \left(\gamma_n^l(0) + \delta_n^l + \sum_{\alpha=2}^{\frac{N+2}{N-2}} |||r_n^l|||_{I_n^1}^{\alpha} \right).$$
 (7.23)

Recall that, in view of definition of nonlinear profiles and γ_n^l , we have

$$\gamma_n^l(0) \leqslant \sum_{j=1}^l \left\| (U - V) \left(-\frac{t_n}{h_n^2} \right) \right\|_{H^1(\mathbb{R}^N)} \to 0, \ n \to \infty$$
 (7.24)

for every $l \geq 1$. Now we need the following classical lemma, see [18].

Lemma 7.5. Let M = M(t) be a positive continuous function on [0, T], such that M(0) = 0 and, for every $t \in [0, T]$, we have

$$M(t) \leqslant C \left(a + \sum_{\alpha=2}^{\frac{N+2}{N-2}} M^{\alpha}(t) \right),$$

with $0 < a < a_0 = a_0(C)$. Then we have, for every $t \in [0, T]$,

$$M(t) \leq 2Ca$$
.

From (7.20) and (7.24), it follows that, for every l large enough, there exists N(l), such that

$$\gamma_n^l(0) + \delta_n^l \le a_0(C)$$

for every $n \geq N(l)$. We denote by M_n^l the function defined on $I_n^1 = [0, a_n^1]$ by

$$M_n^l(s) = \||r_n^l||_{[0,s]} + \frac{s}{a_n^1} \sup_{t \in I_n^1} \|\nabla r_n^l(s)\|_{L^2(\mathbb{R}^N)}.$$

It is clear that (7.22) still holds if we replace $I_n^1 = [0, a_n^1]$ by [0, s] for every $s \in I_n^1$. Thus,

$$M_n^l(s) \le C \left(\gamma_n^l(0) + \delta_n^l + \sum_{\alpha=2}^{\frac{N+2}{N-2}} \left(M_n^l \right)^{\alpha} (s) \right).$$

Hence, M_n^l satisfies the conditions of Lemma 7.5 for l large and $n \ge N(l)$, and so

$$M_n^l\left(a_n^1\right) = \||r_n^l\||_{I_n^1} + \sup_{t \in I_n^1} \|\nabla r_n^l(t)\|_{L^2(\mathbb{R}^N)} \le 2C\left(\gamma_n^l(0) + \delta_n^l\right)$$
(7.25)

for l large and $n \geq N(l)$. Summing (7.20), (7.24), and (7.25), it follows that

$$\limsup_{n \to \infty} \left(\||r_n^l||_{I_n^1} + \sup_{t \in I_n^1} \|\nabla r_n^l(t)\|_{L^2(\mathbb{R}^N)} \right) \to 0, \ l \to \infty.$$

On the other hand, in view of notation (7.10), we have

$$\gamma_n^l\left(a_n^1\right) \le \sup_{t \in I_n^1} \|\nabla r_n^l(t)\|_{L^2(\mathbb{R}^N)},$$

which implies

$$\limsup_{n \to \infty} \gamma_n^l \left(a_n^1 \right) \to 0, \ l \to \infty.$$

This fact allows us to repeat the same argument on $I_n^2 = [a_n^1, a_n^2]$. We get, similarly,

$$|||r_n^l||_{I_n^2} + \sup_{t \in I_n^2} ||\nabla r_n^l(t)||_{L^2(\mathbb{R}^N)} \le C \left(\gamma_n^l(a_n^1) + \delta_n^l\right),$$

SO

$$\limsup_{n \to \infty} \left(\||r_n^l||_{I_n^2} + \sup_{t \in I_n^2} ||\nabla r_n^l(t)||_{L^2(\mathbb{R}^N)} \right) \to 0, \ l \to \infty.$$

By iterating this process, we obtain that, for every $1 \le i \le p$,

$$\limsup_{n\to\infty} \left(\||r_n^l||_{I_n^i} + \sup_{t\in I_n^i} \|\nabla r_n^l(t)\|_{L^2(\mathbb{R}^N)} \right) \to 0, \ l\to\infty.$$

Since p does not depend on n and l, we get

$$\limsup_{n\to\infty} \left(\||r_n^l||_{\mathbb{R}_+} + \sup_{t\in\mathbb{R}_+} \|\nabla r_n^l(t)\|_{L^2(\mathbb{R}^N)} \right) \to 0, \ l\to\infty.$$

A similar decomposition to (7.18) for \mathbb{R}_{-} can be provided. Arguing as above we prove

$$\limsup_{n \to \infty} \left(\||r_n^l||_{\mathbb{R}_-} + \sup_{t \in \mathbb{R}_-} \|\nabla r_n^l(t)\|_{L^2(\mathbb{R}^N)} \right) \to 0, \ l \to \infty.$$

Hence, we get

$$\limsup_{n\to\infty} \left(\||r_n^l||_{\mathbb{R}} + \sup_{t\in\mathbb{R}} \|\nabla r_n^l(t)\|_{L^2(\mathbb{R}^N)} \right) \to 0, \ l\to\infty.$$

Since

$$E\left(r_n^l(t)\right) \le \|\nabla r_n^l(t)\|_{L^2(\mathbb{R}^N)}^2 \left(1 + \|\nabla r_n^l(t)\|_{L^2(\mathbb{R}^N)}^{\frac{4}{N-2}}\right) \to 0, \ l \to \infty,$$

it follows that

$$\limsup_{n \to \infty} \left(\||r_n^l||_{\mathbb{R}} + \sup_{t \in \mathbb{R}} E\left(r_n^l(t)\right) \right) \to 0, \ l \to \infty.$$

To complete the proof of the lemma, we need to prove Propositions 7.1, 7.2, and 7.3. The proof of Propositions 7.1 and 7.2 can be found in [18]. Since this paper considers general nonlinear terms, we need to modify the proof of the Proposition 7.3. The proof is divided into two parts. In the first one we prove that, for every $l \ge 1$,

$$\left\| \nabla \left(\sum_{j=1}^{l} p\left(U_n^j \right) - p\left(\sum_{j=1}^{l} U_n^j \right) \right) \right\|_{L^{\frac{2(N+2)}{N+4}}(\mathbb{R} \times \mathbb{R}^N)} \to 0, \ n \to \infty.$$
 (7.26)

In the second and main one, we prove

$$\limsup_{n \to \infty} \left\| \nabla \left(p \left(W_n^l + w_n^l \right) - p \left(W_n^l \right) \right) \right\|_{L^{\frac{2(N+2)}{N+4}}(\mathbb{R} \times \mathbb{R}^N)} \to 0, \ l \to \infty. \tag{7.27}$$

Part 1. Arguing in the same way as in the proof of Lemma 2.7 and Proposition 3.4 in [18], we concludes the proof of (7.26).

Part 2. By Leibnitz formula and Hölder inequality we get

$$\left\| \nabla \left(p \left(W_n^l + w_n^l \right) - p \left(W_n^l \right) \right) \right\|_{L^{\frac{2(N+2)}{N+4}}(\mathbb{R} \times \mathbb{R}^N)}$$

$$\leq C \left\| |\nabla W_{n}^{l}| \left(W_{n}^{l} + w_{n}^{l} \right)^{\frac{6-N}{N-2}} |w_{n}^{l}| \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^{N})} + C \left\| |\nabla w_{n}^{l}| \left(W_{n}^{l} \right)^{\frac{4}{N-2}} \right\|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^{N})} \\
+ \varepsilon \| \nabla w_{n}^{l} \|_{L^{\frac{2(N+2)}{N-4}}(I \times \mathbb{R}^{N})} \\
\leq C \| \nabla W_{n}^{l} \|_{L^{\frac{2(N+2)}{N}}(I \times \mathbb{R}^{N})} \| W_{n}^{l} + w_{n}^{l} \|_{L^{\frac{6-N}{N-2}}(I \times \mathbb{R}^{N})}^{\frac{6-N}{N-2}} \| w_{n}^{l} \|_{L^{\frac{2(N+2)}{N-2}}(I \times \mathbb{R}^{N})} \\
+ C \| |W_{n}^{l} \nabla w_{n}^{l}| \left(W_{n}^{l} \right)^{\frac{6-N}{N-2}} \|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^{N})} + \varepsilon \| \nabla w_{n}^{l} \|_{L^{\frac{2(N+2)}{N+4}}(I \times \mathbb{R}^{N})} \\
\leq C \left(\| w_{n}^{l} \|_{L^{\frac{2(N+2)}{N-2}}(\mathbb{R} \times \mathbb{R}^{N})} \| W_{n}^{l} + w_{n}^{l} \|_{\mathbb{R}}^{\frac{4}{N-2}} + \| W_{n}^{l} \|_{\mathbb{R}}^{\frac{6-N}{N-2}} \| W_{n}^{l} \nabla \left(w_{n}^{l} \right) \|_{L^{\frac{N+2}{N-1}}(\mathbb{R} \times \mathbb{R}^{N})} \\
+ o(1). \tag{7.28}$$

Putting together (7.7), (7.16) and (7.28), it follows that if we prove

$$\limsup_{n \to \infty} \left\| W_n^l \nabla \left(w_n^l \right) \right\|_{L^{\frac{N+2}{N-1}}(\mathbb{R} \times \mathbb{R}^N)} \to 0, \ l \to \infty,$$

then the proof of (7.27) is done. On the other hand, the convergence of the the series $\sum_{j\geq 1}\|U^j\|_{L^{\frac{2(N+2)}{N-2}}(\mathbb{R}\times\mathbb{R}^N)}^{\frac{2(N+2)}{N-2}} \text{ implies that, for every } \varepsilon>0, \text{ there exists } l(\varepsilon), \text{ such that}$

$$\sum_{j>l(\varepsilon)} \left\| U^j \right\|_{L^{\frac{2(N+2)}{N-2}}(\mathbb{R}\times\mathbb{R}^N)}^{\frac{2(N+2)}{N-2}} \leq \varepsilon^{\frac{2(N+2)}{N-2}}.$$

In particular,

$$\lim_{n \to \infty} \left\| \left(\sum_{j=l(\varepsilon)}^{l} U_n^j \right) \nabla \left(w_n^l \right) \right\|_{L^{\frac{N+2}{N-1}}(\mathbb{R} \times \mathbb{R}^N)}^{\frac{2(N+2)}{N-2}} \\
\leq \left(\sum_{j \ge l(\varepsilon)} \left\| U^j \right\|_{L^{\frac{2(N+2)}{N-2}}(\mathbb{R} \times \mathbb{R}^N)}^{\frac{2(N+2)}{N-1}} \right) \lim_{n \to \infty} \sup_{n \to \infty} \left\| \nabla w_n^l \right\|_{L^{\frac{2(N+2)}{N-2}}(\mathbb{R} \times \mathbb{R}^N)}^{\frac{2(N+2)}{N-2}} \\
\leq C\varepsilon^{\frac{2(N+2)}{N-2}}. \tag{7.29}$$

The last inequality follows from the fact that $\|\nabla w_n^l\|_{L^{\frac{2(N+2)}{N}}(\mathbb{R}\times\mathbb{R}^N)}$ is uniformly bounded. Thereby,

$$\limsup_{n \to \infty} \left\| W_n^l \nabla \left(w_n^l \right) \right\|_{L^{\frac{N+2}{N-1}}(\mathbb{R} \times \mathbb{R}^N)} \le \limsup_{n \to \infty} \left\| W_n^{l(\varepsilon)} \nabla \left(w_n^l \right) \right\|_{L^{\frac{N+2}{N-1}}(\mathbb{R} \times \mathbb{R}^N)} + C\varepsilon,$$

for $l \ge l(\varepsilon)$. Hence, our problem is reduced to prove that

$$\limsup_{n \to \infty} \left\| W_n^{l_0} \nabla \left(w_n^l \right) \right\|_{L^{\frac{N+2}{N-1}}(\mathbb{R} \times \mathbb{R}^N)} \to 0, \ l \to \infty$$

for every fixed $l_0 \geq 1$. Since $W_n^{l_0} = \sum_{i=1}^{l_0} U_n^i$, we have to show that

$$\limsup_{n \to \infty} \left\| U_n^j \nabla \left(w_n^l \right) \right\|_{L^{\frac{N+2}{N-1}}(\mathbb{R} \times \mathbb{R}^N)} \to 0, \ l \to \infty$$

for every $l_0 \ge j \ge 1$. A change of variables $x = h_n^j y + x_n, t = (h_n^j)^2 s + t_n$ gives

$$\left\| U_n^j \nabla \left(w_n^l \right) \right\|_{L^{\frac{N+2}{N-1}}(\mathbb{R} \times \mathbb{R}^N)} = \left\| U^j \nabla \left(\widetilde{w}_n^l \right) \right\|_{L^{\frac{N+2}{N-1}}(\mathbb{R} \times \mathbb{R}^N)},$$

where

$$\widetilde{w}_l^n(y,s) = (h_n^j)^{\frac{N-2}{2}} w_n^l \left(h_n^j y + x_n, \left(h_n^j \right)^2 s + t_n \right).$$

Observe that

$$\left\|w_n^l\right\|_{L^{\frac{2(N+2)}{N-2}}(\mathbb{R}\times\mathbb{R}^N)} = \left\|\widetilde{w}_n^l\right\|_{L^{\frac{2(N+2)}{N-2}}(\mathbb{R}\times\mathbb{R}^N)} \quad \text{and} \quad \left\|\nabla w_n^l\right\|_{L^{\frac{2(N+2)}{N}}(\mathbb{R}\times\mathbb{R}^N)} = \left\|\nabla\widetilde{w}_n^l\right\|_{L^{\frac{2(N+2)}{N}}(\mathbb{R}\times\mathbb{R}^N)}.$$

By density, we can take $U^j \in C_0^{\infty}$. Using Hölder inequality, it is enough to prove

$$\limsup_{n \to \infty} \left\| \nabla \widetilde{w}_n^l \right\|_{L^2(\mathscr{B})} \to 0, \ l \to \infty, \tag{7.30}$$

where \mathscr{B} is a fixed compact of $\mathbb{R}_t \times \mathbb{R}_x^N$. This fact will follow from the following Lemma which can be founded in [18].

Lemma 7.6. Let \mathscr{B} be a compact set of $\mathbb{R}_t \times \mathbb{R}_x^N$. Then, for every $\varepsilon > 0$, there exists a constant C_{ε} such that

$$\|\nabla v\|_{L^2(\mathscr{B})} \le C_{\varepsilon} \|v\|_{L^{\frac{2(N+2)}{N-2}}(\mathbb{R}\times\mathbb{R}^N)} + \varepsilon \|\nabla v(0,\cdot)\|_{L^2(\mathbb{R}^N)}$$

for every v(t,x) solution to linear Schrödinger equation (7.5).

Applying Lemma 7.6 and (7.7), we know that (7.30) holds. This concludes the proof of Lemma 7.3.

Proof of Lemma 7.1. (7.1) is a consequence of the proof of Corollary 1.9 in [18], here, we use the hypothesis $\|e^{it\Delta}v_{0,n}\|_{L^{\frac{2(N+2)}{N-2}}_{t,x}} \ge \rho > 0$. (7.4) follows from the orthogonality of $(\lambda_{j,n}; x_{j,n}; t_{j,n})$ as in the proof of (7.3). The rest of the lemma is contained in the proof of Lemmas 7.2 and 7.3.

8 Compactness of critical element

Let us consider the statement:

(SC) For all $u_0 \in H^1(\mathbb{R}^N)$, with

$$\int_{\mathbb{D}^N} |\nabla u_0|^2 dx < \int_{\mathbb{D}^N} |\nabla W|^2 dx \text{ and } E(u_0) < E(W),$$

if u is the corresponding solution to the (1.1), with maximal interval of existence I, then $I=(-\infty,+\infty)$ and $\|u\|_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,+\infty)}W^{1,\frac{2N(N+2)}{N^2+4}}}<\infty$.

We say that $(SC)(u_0)$ holds if for this particular u_0 (such as, take $u|_{t=0} = u_0$), with

$$\int_{\mathbb{R}^N} |\nabla u_0|^2 dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx \text{ and } E(u_0) < E(W)$$

and u the corresponding solution to the (CP), with maximal interval of existence I, we have $I=(-\infty,+\infty)$ and $\|u\|_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,+\infty)}W^{1,\frac{2N(N+2)}{N^2+4}}}<\infty$.

Note that, because of Lemma 3.2, if $\|\nabla u_0\|_{L^2} \leq \bar{\delta}$, (SC) (u_0) holds. Thus, in light of Corollary 6.2, there exists $\eta_0 > 0$ such that, if u_0 is as in (SC) and $E(u_0) < \eta_0$, then (SC) (u_0) holds. Moreover, for any u_0 as in (SC), $E(u_0) \geq 0$, in light of Theorem 6.1. Thus, there exists a number E_C , with $\eta_0 \leq E_C \leq E(W)$, such that, if u_0 is as in (SC) and $E(u_0) < E_C$, (SC) (u_0) holds and E_C is optimal with this property. For the rest of this section we will assume that $E_C < E(W)$. We now prove that there exits a critical element $u_{0,C}$ at the critical level of energy E_C so that (SC) $(u_{0,C})$ does not hold and from the minimality, this element has a compactness property up to the symmetry of this equation. This is in fact a general principle which follows from the concentration compactness ideas. More precisely,

Lemma 8.1. There exists $u_{0,C}$ in H^1 , with

$$E(u_{0,C}) = E_C < E(W), \quad \int_{\mathbb{R}^N} |\nabla u_{0,C}|^2 dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx$$

such that, if u_C is the solution of (1.1) with data $u_{0,C}$, and maximal interval of existence $I, 0 \in \stackrel{\circ}{I}$, then $||u_C||_{L_{I}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}} = +\infty$.

Lemma 8.2. Assume u_C is as in Lemmas 8.1 and $\|u_C\|_{L_{I_+}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}} = +\infty$, where $I_+ = (0, +\infty) \cap I$. Then there exists $x(t) \in \mathbb{R}^N$ and $\lambda(t) \in \mathbb{R}^+$, for $t \in I_+$, such that

$$K = \left\{ v(x,t) : v(x,t) = \frac{1}{\lambda(t)^{\frac{N-2}{2}}} u_C \left(\frac{x - x(t)}{\lambda(t)}, t \right) \right\}$$

has the property that \overline{K} is compact in H^1 . A corresponding conclusion is reached if

$$||u_C||_{L_{I_{-}}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}} = +\infty,$$

where $I_{-}=(-\infty,0)\cap I$.

Lemma 8.3. Let $\{z_{0,n}\} \in H^1$, with

$$\int_{\mathbb{R}^N} |\nabla z_{0,n}|^2 dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx \text{ and } E(z_{0,n}) \to E_C$$

and with $\|e^{it\Delta}z_{0,n}\|_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,+\infty)}W^{1,\frac{2N(N+2)}{N^2+4}}} \ge \rho$, where ρ as in Lemma 3.1. Let $\{V_{0,j}\}$ be as in

Lemma 7.1. Assume that one of the two hypothesis

$$\underline{\lim}_{n \to \infty} E(V_1^l(-\frac{t_{1,n}}{\lambda_{1,n}^2})) < E_C$$
(8.1)

or after passing to a subsequence, we have that, with $s_n = -\frac{t_{1,n}}{\lambda_{1,n}^2}$, $E(V_1^l(s_n)) \to E_C$, and $s_n \to s_* \in [-\infty, +\infty]$, and if U_1 is the non-linear profile (see Definition 2.1 and 2.4) associated to $(V_{0,1}, \{s_n\})$ we have that the maximal interval of existence of U_1 is $I = (-\infty, +\infty)$ and $\|U_1\|_{L^{2(N+2)}_{(-\infty, +\infty)}W^{1,\frac{2N(N+2)}{N^2+4}}} < \infty$ and

$$\underline{\lim}_{n \to \infty} E(V_1^l(-\frac{t_{1,n}}{\lambda_{1,n}^2})) = E_C. \tag{8.2}$$

Then (after passing to a subsequence), for n large, if z_n is the solution of (1.1) with data at t = 0 equal to $z_{0,n}$, then $(SC)(z_{0,n})$ holds.

Let us first assume the validity of Lemma 8.3 and use it (together with Lemma 7.1) to establish Lemmas 8.1 and 8.2.

Proof of Lemma 8.1. According to the definition of E_C , and the assumption that $E_C < E(W)$, we can find $u_{0,n} \in H^1$, with

$$\int_{\mathbb{R}^N} |\nabla u_{0,n}|^2 \, dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx, \ E(u_{0,n}) \to E_C$$

and such that if u_n is the solution of (1.1) with data at t=0, $u_{0,n}$ and maximal interval of existence $I_n=(-T_-\left(u_{0,n}\right),T_+\left(u_{0,n}\right))$, then $\left\|e^{it\Delta}u_{0,n}\right\|_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,+\infty)}W^{1,\frac{2N(N+2)}{N^2+4}}}\geq \rho>0$, where ρ is as in Lemma 3.1 and $\left\|u_n\right\|_{L^{\frac{2(N+2)}{N-2}}_{I_n}W^{1,\frac{2N(N+2)}{N^2+4}}}=+\infty$ (Here we are also using Lemma 2.1 and Lemma 3.1). Note that, since $E_C< E(W)$, there exists $\delta_0>0$ such that

$$E(u_{0,n}) \le (1 - \delta_0) E(W), \ \forall n.$$

Because of Theorem 6.1, we can find $\bar{\delta}$ so that

$$\int_{\mathbb{R}^N} |\nabla u_n(t)|^2 dx \le (1 - \bar{\delta}) \int_{\mathbb{R}^N} |\nabla W|^2 dx \text{ for all } t \in I_n, \ \forall n.$$

Apply now Lemma 7.1 for $\varepsilon_0 > 0$ and Lemma 8.3. We then have, for $J = J(\varepsilon_0)$, that

$$u_{0,n} = \sum_{j=1}^{J} \frac{1}{\lambda_{j,n}^{\frac{N-2}{2}}} V_j^l \left(\frac{x - x_{j,n}}{\lambda_{j,n}}, \frac{-t_{j,n}}{\lambda_{j,n}^2} \right) + w_n, \tag{8.3}$$

$$\int_{\mathbb{R}^N} |\nabla u_{0,n}|^2 dx = \sum_{i=1}^J \int_{\mathbb{R}^N} |\nabla V_{0,i}|^2 dx + \int_{\mathbb{R}^N} |\nabla w_n|^2 dx + o(1), \tag{8.4}$$

$$E(u_{0,n}) = \sum_{j=1}^{J} E\left(V_j^l \left(\frac{-t_{j,n}}{\lambda_{j,n}^2}\right)\right) + E(w_n) + o(1).$$
(8.5)

Note that because of (8.4) we have, for all n large, that

$$\int_{\mathbb{R}^N} |\nabla w_n|^2 dx \le (1 - \frac{\bar{\delta}}{2}) \int_{\mathbb{R}^N} |\nabla W|^2 dx \text{ and } \int_{\mathbb{R}^N} |\nabla V_{0,j}|^2 dx \le (1 - \frac{\bar{\delta}}{2}) \int_{\mathbb{R}^N} |\nabla W|^2 dx.$$

From Corollary 6.1 it now follows that $E(V_j^l(-\frac{t_{j,n}}{\lambda_{jn}^2})) \geq 0$ and $E(w_n) \geq 0$. From this and (8.5) it follows that

$$E(V_1^l(-\frac{t_{1,n}}{\lambda_{1,n}^2})) \le E(u_{0,n}) + o(1)$$

and hence $\lim_{n\to\infty} E(V_1^l(-\frac{t_{1,n}}{\lambda_{1,n}^2})) \leq E_C$. If the left-hand side is strictly less than E_C , Lemma 8.3 gives us a contradiction with the choice of $u_{0,n}$, for n large (after passing to a subsequence). Hence, the left-hand side must equal E_C .

Let then U_1 be the non-linear profile associated to $(V_1^l, \{s_n\})$, with $s_n = -\frac{t_{1,n}}{\lambda_{1,n}^2}$ (after passing to a subsequence). We first note that we must have J = 1. This is because (8.5) and $E(u_{0,n}) \to E_C$, $E(V_1^l(-s_n)) \to E_C$ now imply that

$$E(w_n) \to 0 \text{ and } E(V_j^l(-\frac{t_{j,n}}{\lambda_{j,n}^2})) \to 0, j = 2, \dots, J.$$

Using (6.4) and the argument in the proof of Corollary 6.2, we have

$$\sum_{j=2}^{J} \int_{\mathbb{R}^{N}} |\nabla V_{j}^{l}(-\frac{t_{j,n}}{\lambda_{j,n}^{2}})|^{2} dx + \int_{\mathbb{R}^{N}} |\nabla w_{n}|^{2} dx \to 0.$$

We then have, since $\int_{\mathbb{R}^N} |\nabla V_j^l(-\frac{t_{j,n}}{\lambda_{j,n}^2})|^2 dx = \int_{\mathbb{R}^N} |\nabla V_{0,j}|^2 dx$ that $V_{0,j} = 0, j = 2, \ldots, J$ and $\int_{\mathbb{R}^N} |\nabla w_n|^2 dx \to 0$. Hence (8.3) becomes $u_{0,n} = \frac{1}{\lambda_{1,n}^{\frac{N-2}{2}}} V_1^l \left(\frac{x-x_{1,n}}{\lambda_{1,n}}, s_n\right) + w_n$. Let $v_{0,n} = \lambda_{1,n}^{\frac{N-2}{2}} u_{0,n} \left(\lambda_{1,n} \left(x+x_{1,n}\right)\right)$ and note that scaling gives us that $v_{0,n}$ verifies the same hypothesis as $u_{0,n}$. Moreover, $\widetilde{w}_n = \lambda_{1,n}^{\frac{N-2}{2}} w_n \left(\lambda_{1,n} (x+x_{1,n})\right)$ still verifies $\int_{\mathbb{R}^N} |\nabla \widetilde{w}_n|^2 dx \to 0$. Thus

$$v_{0,n} = V_1^l(s_n) + \widetilde{w}_n, \ \int_{\mathbb{R}^N} |\nabla \widetilde{w}_n|^2 dx \to 0.$$

Let us return to U_1 , the non-linear profile associated to $(V_{0,1},\{s_n\})$ and let

$$I_1 = (T_-(U_1), T_+(U_1))$$

be its maximal interval of existence. Note that, by definition of non-linear profile, we have

$$\int_{\mathbb{R}^{N}} |\nabla U_{1}(s_{n})|^{2} dx = \int_{\mathbb{R}^{N}} |\nabla V_{1}^{l}(s_{n})|^{2} dx + o(1) \text{ and } E(U_{1}(s_{n})) = E(V_{1}^{l}(s_{n})) + o(1)$$

Note that in this case $E(V_1^l(s_n)) = E_C + o(1)$ and

$$\int_{\mathbb{R}^{N}} \left| \nabla V_{1}^{l} \left(s_{n} \right) \right|^{2} dx = \int_{\mathbb{R}^{N}} \left| \nabla V_{0,1} \right|^{2} dx = \int_{\mathbb{R}^{N}} \left| \nabla u_{0,n} \right|^{2} dx + o(1) < \int_{\mathbb{R}^{N}} \left| \nabla W \right|^{2} dx$$

for n large by Theorem 6.1. Let's fix $\bar{s} \in I_1$. Then $E(U_1(s_n)) = E(U_1(\bar{s}))$, so that

$$E\left(U_1(\bar{s})\right) = E_C.$$

Moreover, $\int_{\mathbb{R}^N} |\nabla U_1(s_n)|^2 dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx$ for n large and hence by (6.7)

$$\int_{\mathbb{R}^N} |\nabla U_1(\bar{s})|^2 dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx.$$

If $||U_1||_{L_{I_1}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}} < +\infty$, Remark 1.4 gives us that $I_1 = (-\infty, +\infty)$ and we then obtain a contradiction from Lemma 8.3. Thus,

$$||U_1||_{L_{I_1}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}} = +\infty$$

and we then set $u_C = U_1$ (after a translation in time to make $\bar{s} = 0$).

Proof of Lemma 8.2. We argue by contradiction. For brevity of notation, let us set $u(x,t) = u_C(x,t)$. If not, there exists $\eta_0 > 0$ and a sequence $\{t_n\}_{n=1}^{\infty}$, $t_n \geq 0$ such that, for all $\lambda_0 \in \mathbb{R}^+$, $x_0 \in \mathbb{R}^N$, we have

$$\left\| \frac{1}{\lambda_0^{\frac{N-2}{2}}} u\left(\frac{x - x_0}{\lambda_0}, t_n\right) - u\left(x, t_{n'}\right) \right\|_{H^1} \ge \eta_0, \quad \text{for } n \ne n'.$$
 (8.6)

Note that (after passing to a subsequence, so that $t_n \to \bar{t} \in [0, T_+(u_0)]$), we must have $\bar{t} = T_+(u_0)$, in view of the continuity of the flow in H^1 , as guaranteed by Lemma 3.1. Note that, in view of Lemma 3.1 we must also have $\|e^{it\Delta}u(t_n)\|_{L^{2(N+2)}_{(n+\infty)}W^{1,\frac{2N(N+2)}{N^2+4}}} \ge \rho$.

Step 1. Let us apply Lemma 7.1 to $v_{0,n} = u(t_n)$ with $\varepsilon_0 > 0$. We will show that J = 1. Indeed, if $\lim_{n \to \infty} E(V_1^l(-\frac{t_{1,n}}{\lambda_{1,n}^2})) < E_C$, then by Theorem 6.1, we have

$$\int_{\mathbb{R}^N} |\nabla u(t)|^2 dx \le (1 - \delta_1) \int_{\mathbb{R}^N} |\nabla W|^2 dx \text{ for all } t \in I_+$$

and $E(u(t)) = E(u_0) = E_C < E(W)$, by Lemma 8.3 we obtain that (SC)(u) holds. So $||u||_{L_{I_+}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}} < +\infty$, which contradicts the hypothesis. Therefore, it follows that

 $\underline{\lim}_{n\to\infty} E(V_1^l(-\frac{t_{1,n}}{\lambda_{1,n}^2})) = E_C$. Similar to the proof of Lemma 8.1, we get

$$J = 1, \ \int_{\mathbb{R}^N} |\nabla w_n|^2 \, dx \to 0.$$

Thus, we have

$$u(t_n) = \frac{1}{\lambda_{1,n}^{\frac{N-2}{2}}} V_1^l \left(\frac{x - x_{1,n}}{\lambda_{1,n}}, \frac{-t_{1,n}}{\lambda_{1,n}^2} \right) + w_n, \quad \int_{\mathbb{R}^N} |\nabla w_n|^2 dx \to 0.$$
 (8.7)

Step 2. we prove that $s_n = \frac{-t_{1,n}}{\lambda_{1,n}^2}$ must be bounded. In fact, note that

$$e^{it\Delta}u(t_n) = \lambda_{1,n}^{-\frac{N-2}{2}}V_1^l\left(\frac{x - x_{1,n}}{\lambda_{1,n}}, \frac{t - t_{1,n}}{(\lambda_{1,n})^2}\right) + e^{it\Delta}w_n.$$

On the other hand, assume $\frac{t_{1,n}}{\lambda_{1,n}^2} \leq -C_0$, where C_0 is a large positive constant. Then, since

$$\|e^{it\Delta}w_n\|_{L^{\frac{2(N+2)}{N-2}}_{(0,+\infty)}W^{1,\frac{2N(N+2)}{N^2+4}}} < \frac{\rho}{2} \text{ for } n \text{ large enough}$$

and

$$\left\| \lambda_{1,n}^{-\frac{N-2}{2}} V_1^l \left(\frac{x - x_{1,n}}{\lambda_{1,n}}, \frac{t - t_{1,n}}{\left(\lambda_{1,n}\right)^2} \right) \right\|_{L_{\frac{2(N+2)}{N-2}}} \leq \left\| V_1^l(y,s) \right\|_{L_{\frac{2(N+2)}{N-2}}} \leq \frac{\rho}{2}$$

for C_0 large, which contradicts $\left\|e^{it\Delta}u\left(t_n\right)\right\|_{L_{(0,+\infty)}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}} \geq \rho$.

On the other hand, assume that $\frac{t_{1,n}}{\lambda_{1,n}^2} \geq C_0$, for a large positive constant C_0 , n large, we have

$$\left\| \lambda_{1,n}^{-\frac{N-2}{2}} V_1^l \left(\frac{x - x_{1,n}}{\lambda_{1,n}}, \frac{t - t_{1,n}}{(\lambda_{1,n})^2} \right) \right\|_{L_{(-\infty,0)}^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}} \le \left\| V_1^l(y,s) \right\|_{L_{(-\infty,-C_0)}^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}} \le \frac{\rho}{2}$$

for C_0 large. Hence, $\|e^{it\Delta}u(t_n)\|_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,0)}W^{1,\frac{2N(N+2)}{N^2+4}}} \leq \rho$, for n large. By Lemma 3.1, we know that $\|u\|_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,t_n)}W^{1,\frac{2N(N+2)}{N^2+4}}} \leq \rho$, which gives us a contradiction because of $t_n \to T_+(u_0)$. Thus $\left|\frac{t_{1,n}}{\lambda_{1,n}^2}\right| \leq C_0$ and after passing to a subsequence,

$$\frac{t_{1,n}}{\lambda_{1,n}^2} \to t_0 \in (-\infty, +\infty).$$

Step 3. By (8.6) and (8.7), for $n \neq n'$ large (independently of λ_0, x_0), it holds

$$\left\| \frac{1}{\lambda_0^{\frac{N-2}{2}}} \frac{1}{\lambda_{1,n}^{\frac{N-2}{2}}} V_1^l \left(\frac{\frac{x-x_0}{\lambda_0} - x_{1,n}}{\lambda_{1,n}}, \frac{-t_{1,n}}{(\lambda_{1,n})^2} \right) - \frac{1}{(\lambda_{1,n'})^{\frac{N-2}{2}}} V_1^l \left(\frac{x-x_{1,n'}}{\lambda_{1,n'}}, \frac{-t_{1,n'}}{(\lambda_{1,n'})^2} \right) \right\|_{H^1} \ge \eta_0/2$$

or

$$\left\| \left(\frac{\lambda_{1,n'}}{\lambda_{1,n}\lambda_0} \right)^{\frac{N-2}{2}} V_1^l \left(\frac{y\lambda_{1,n'}}{\lambda_0\lambda_{1,n}} + \widetilde{x}_{n,n'} - \widetilde{x}_0, -\frac{t_{1,n}}{\left(\lambda_{1,n}\right)^2} \right) - V_1^l \left(y, -\frac{t_{1,n'}}{\lambda_{1,n'}^2} \right) \right\|_{H^1} \geq \frac{\eta_0}{2},$$

where $\widetilde{x}_{n,n'}$ is a suitable point in \mathbb{R}^N and $\lambda_0, \widetilde{x}_0$ are arbitrary. But if we choose $\lambda_0 = \frac{\lambda_{1,n'}}{\lambda_{1,n}}, \widetilde{x}_0 = x_{n,n'}$, then $-\frac{t_{1,n}}{(\lambda_{1,n})^2} \to -t_0$ and $\frac{-t_{1,n'}}{(\lambda_{1,n'})^2} \to -t_0$. So $\|0\|_{H^1} \ge \frac{\eta_0}{2}$, which reaches a contradiction.

Thus, to complete the proofs of Lemmas 8.1 and 8.2 we only need to provide the proof of Lemma 8.3.

Proof of Lemma 8.3. Let us assume first that (8.2) holds and set

$$A = \int_{\mathbb{R}^N} |\nabla W|^2 dx, A' = \int_{\mathbb{R}^N} |\nabla W|^2 dx, M = ||U_1||_{L_{(-\infty, +\infty)}^{\frac{2(N+2)}{N-2}}, W^{1, \frac{2N(N+2)}{N^2+4}}}.$$

Arguing (for some $\varepsilon_0 > 0$ in Lemma 7.1) as in the proof of Lemmas 8.1, we see that

$$\underline{\lim_{n \to \infty}} E(V_1^l(-\frac{t_{1,n}}{\lambda_{1,n}^2})) = E_C \text{ and } E_C < E(W),$$

which imply that J=1, $\int_{\mathbb{R}^N} |\nabla w_n|^2 dx \to 0$. Moreover, if

$$v_{0,n} = \lambda_{1,n}^{\frac{N-2}{2}} z_{0,n}(\lambda_{1,n}(x+x_{1,n})), \ \widetilde{w}_n = \lambda_{1,n}^{\frac{N-2}{2}} w_n(\lambda_{1,n}(x+x_{1,n})), \ s_n = -\frac{t_{1,n}}{\lambda_{1,n}^2},$$

we have $\int_{\mathbb{R}^N} |\nabla \widetilde{w}_n|^2 dx \to 0$ and $v_{0,n} = V_1^l(s_n) + \widetilde{w}_n$, while

$$\left\| e^{it\Delta} v_{0,n} \right\|_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,+\infty)} W^{1,\frac{2N(N+2)}{N^2+4}}} \ge \delta, \ \int_{\mathbb{R}^N} \left| \nabla v_{0,n} \right|^2 dx < \int_{\mathbb{R}^N} \left| \nabla W \right|^2 dx, \ E\left(v_{0,n}\right) \to E_C.$$

By definition of non-linear profile, we know that

$$\int_{\mathbb{R}^N} |\nabla V_1^l(s_n) - \nabla U_1(s_n)|^2 dx = o(1).$$

We then have

$$v_{0,n} = U_1(s_n) + \widetilde{\widetilde{w}}_n, \int_{\mathbb{R}^N} |\nabla \widetilde{\widetilde{w}}_n|^2 dx \to 0.$$

Moreover, as in the proof of Lemma 8.1, $E(U_1(0)) = E_C$ and $\int_{\mathbb{R}^N} |\nabla U_1(t)|^2 dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx$ for all t. We now apply Proposition 5.1, with $\varepsilon_0 < \varepsilon_0(M, A, A', N)$ and n large, with $\tilde{u} = U_1, e \equiv 0, t_0 = 0, u_0 = v_{0,n}$. This case now follows.

Next, assume that (8.1) holds, the proof is divided into five steps.

Step 1 We prove that for $j \geq 2$, we also have $\underline{\lim}_{n \to \infty} E(V_j^l(-\frac{t_{j,n}}{\lambda_{j,n}^2})) < E_C$. In fact, up to a subsequence, assume $\lim_{n \to \infty} E(V_1^l(-\frac{t_{1,n}}{\lambda_{1,n}})) < E_C$. Due to (7.3), it holds

$$\int_{\mathbb{R}^N} |\nabla z_{0,n}|^2 dx \ge \sum_{j=1}^J \int_{\mathbb{R}^N} |\nabla V_{0,j}|^2 dx + o(1)$$

and since $E_C < E(W)$, for n large we have $E(z_{0,n}) \le (1 - \delta_0)E(W)$, by Lemma 6.1,

$$\int_{\mathbb{R}^N} |\nabla z_{0,n}|^2 dx \le (1 - \delta_1) \int_{\mathbb{R}^N} |\nabla W|^2 dx \text{ and } \int_{\mathbb{R}^N} |\nabla V_{0,j}|^2 dx \le (1 - \delta_1) \int_{\mathbb{R}^N} |\nabla W|^2 dx.$$

Similarly, $\int_{\mathbb{R}^N} |\nabla w_n|^2 dx \leq (1 - \delta_1) \int_{\mathbb{R}^N} |\nabla W|^2 dx$. By Corollary 6.1, we have

$$E(V_j^l(-\frac{t_{j,n}}{\lambda_{j,n}^2})) \ge 0, \ E(w_n) \ge 0.$$

Moreover, using (7.1) and the proof of Corollary 6.2, we have

$$E(V_1^l(-\frac{t_{1,n}}{\lambda_{1,n}^2})) \ge C \int_{\mathbb{R}^N} |\nabla V_{0,1}|^2 dx \ge c\alpha_0 = \overline{\alpha_0} > 0 \text{ for } n \text{ large.}$$

By (7.4), it holds

$$E(z_{0,n}) \ge \overline{\alpha_0} + \sum_{j=2}^{J} E(V_j^l(-\frac{t_{j,n}}{\lambda_{j,n}^2})) + o(1)$$
 for n large,

so the claim follows from $E\left(z_{0,n}\right) \to E_{C}.$

Step 2 We show that (after passing to a subsequence so that, for each j, $\lim_{n} E(V_j^l(-\frac{t_{j,n}}{\lambda_{j,n}^2}))$ exists and $\lim_{n}(-\frac{t_{j,n}}{\lambda_{j,n}^2}) = \overline{s_j} \in [-\infty, +\infty]$ exists) if U_j is the non-linear profile associated to $(V_j^l, \{-\frac{t_{j,n}}{\lambda_{j,n}^2}\})$, then U_j satisfies (SC). Indeed, according to the definition of non-linear profile and Step 1, it follows that $E(U_j) < E_C$ because of $\lim_{n \to \infty} E(V_j^l(-\frac{t_{j,n}}{\lambda_{j,n}^2})) < E_C$. Moreover, since

$$\int_{\mathbb{R}^N} |\nabla V_j^l(-\frac{t_{j,n}}{\lambda_{j,n}^2})|^2 dx \le (1-\delta_1) \int_{\mathbb{R}^N} |\nabla W|^2 dx,$$

the definition of non-linear profile and Theorem 6.1, if $\bar{t} \in I_j$ (the maximal interval for U_j), we have $\int_{\mathbb{R}^N} |\nabla U_j(\bar{t})|^2 dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx$. By the definition of E_C , our claim follows. Note that the argument in the proof of Proposition 5.1 also gives that $||U_j||_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,+\infty)}W^{1,\frac{2N(N+2)}{N^2+4}}} < +\infty$.

Step 3 We claim that there exists j_0 so that, for $j \geq j_0$ we have

$$||U_j||_{L_{(-\infty,+\infty)}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}}^{\frac{2(N+2)}{N-2}} \le C||V_{0,j}||_{H^1}^{\frac{2(N+2)}{N-2}}.$$
(8.8)

In fact, from (7.3), for fixed J we see that (choosing n large)

$$\sum_{j=1}^{J} \int_{\mathbb{R}^{N}} |\nabla V_{0,j}|^{2} dx \le \int_{\mathbb{R}^{N}} |\nabla z_{0,n}|^{2} dx + o(1) \le 2 \int_{\mathbb{R}^{N}} |\nabla W|^{2} dx.$$

Thus, for $j \geq j_0$, we have

$$\int_{\mathbb{D}^N} |\nabla V_{0,j}|^2 \, dx \le \widetilde{\delta},$$

where $\tilde{\delta}$ is so small that $\|e^{it\Delta}V_{0,j}\|_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,+\infty)}W^{1,\frac{2N(N+2)}{N^2+4}}} \leq \rho$, with ρ as in Lemma 3.1. From the definition of non-linear profile, it then follows that $\|U_j\|_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,+\infty)}W^{1,\frac{2N(N+2)}{N^2+4}}} \leq 2\rho$, and using the integral equation

$$u(t) = e^{it\Delta}u_0 + i \int_0^t e^{i(t-t')\Delta} f(u)dt'.$$

So $||U_j(0)||_{H^1} \le C ||V_{0,j}||_{H^1}$ and $||U_j||_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,+\infty)}W^{1,\frac{2N(N+2)}{N^2+4}}} \le C ||V_{0,j}||_{H^1}$, which implies (8.8).

Step 4 For $\varepsilon_0 > 0$, to be chosen, define now

$$H_{n,\varepsilon_0} = \sum_{j=1}^{J(\varepsilon_0)} \frac{1}{\lambda_{j,n}^{\frac{N-2}{2}}} U_j\left(\frac{x - x_{j,n}}{\lambda_{j,n}}, \frac{t - t_{j,n}}{\lambda_{j,n}^2}\right),$$

then it follows that

$$||H_{n,\varepsilon_0}||_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,+\infty)}L^{\frac{2N(N+2)}{N-2}}} \le C_0,$$
(8.9)

uniformly in ε_0 , for $n \geq n(\varepsilon_0)$. In fact,

$$\begin{split} & \|H_{n,\varepsilon_0}\|_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,+\infty)}L^{\frac{2(N+2)}{N-2}}} \\ & = \iint \left[\sum_{j=1}^{J(\varepsilon_0)} \frac{1}{\lambda_{j,n}^{\frac{N-2}{2}}} U_j \left(\frac{x-x_{j,n}}{\lambda_{j,n}}, \frac{t-t_{j,n}}{\lambda_{j,n}^2} \right) \right]^{\frac{2(N+2)}{N-2}} \\ & \leq C_{J(\varepsilon_0)} \sum_{j'\neq j} \iint \left| \frac{1}{\lambda_{j,n}^{\frac{N-2}{2}}} U_j \left(\frac{x-x_{j,n}}{\lambda_{j,n}}, \frac{t-t_{j,n}}{\lambda_{j,n}^2} \right) \right| \cdot \left| \frac{1}{\lambda_{j',n}^{\frac{N-2}{2}}} U_{j'} \left(\frac{x-x_{j',n}}{\lambda_{j',n}}, \frac{t-t_{j',n}}{\lambda_{j',n}^2} \right) \right|^{\frac{N+6}{N-2}} \\ & + \sum_{j=1}^{J(\varepsilon_0)} \iint \left| \frac{1}{\lambda_{j,n}^{\frac{N-2}{2}}} U_j \left(\frac{x-x_{j,n}}{\lambda_{j,n}}, \frac{t-t_{j,n}}{\lambda_{j,n}^2} \right) \right|^{\frac{2(N+2)}{N-2}} \\ & = I + II. \end{split}$$

By the orthogonality of $(\lambda_{j,n}; x_{j,n}; t_{j,n})$, we know that II $\to 0$ for n large(see Keraani [18]). Hence, for n large we have II \leq I. Since (7.3), it follows that

$$I \leq \sum_{j=1}^{j_0} \|U_j\|_{L_{(-\infty,+\infty)}}^{\frac{2(N+2)}{N-2}} L_{\frac{2N(N+2)}{N-2}}^{\frac{2(N+2)}{N-2}} + \sum_{j=j_0}^{J(\varepsilon_0)} \|U_j\|_{L_{(-\infty,+\infty)}}^{\frac{2(N+2)}{N-2}} L_{\frac{2N(N+2)}{N-2}}^{\frac{2(N+2)}{N-2}} L_{\frac{N-2}{N-2}}^{\frac{2(N+2)}{N-2}} \\ \leq \sum_{j=1}^{j_0} \|U_j\|_{L_{(-\infty,+\infty)}}^{\frac{2(N+2)}{N-2}} L_{\frac{2N(N+2)}{N-2}}^{\frac{2N(N+2)}{N-2}} + C \sum_{j=j_0}^{J(\varepsilon_0)} \|V_{0,j}\|_{H^1}^{\frac{2(N+2)}{N-2}} \\ \leq \frac{C_0}{2},$$

where j_0 is defined as in (8.8). For $\varepsilon_0 > 0$, to be chosen, define

$$R_{n,\varepsilon_0} = |H_{n,\varepsilon_0}|^{\frac{4}{N-2}} H_{n,\varepsilon_0} - \sum_{j=1}^{J(\varepsilon_0)} \left| \frac{1}{\lambda_{j,n}^{\frac{N-2}{2}}} U_j \left(\frac{x - x_{j,n}}{\lambda_{j,n}}, \frac{t - t_{j,n}}{\lambda_{j,n}^2} \right) \right|^{\frac{4}{N-2}} \frac{1}{\lambda_{j,n}^{\frac{N-2}{2}}} U_j \left(\frac{x - x_{j,n}}{\lambda_{j,n}}, \frac{t - t_{j,n}}{\lambda_{j,n}^2} \right).$$

using the arguments of Keraani [18], we get

For
$$n = n(\varepsilon_0)$$
 large, $\|\nabla R_{n,\varepsilon_0}\|_{L^2_t L^{\frac{2N}{N+2}}_x} \to 0$ as $n \to \infty$.

Step 5 Finally, we apply Proposition 5.1 to obtain our purpose. Let

$$\widetilde{u} = H_{n,\varepsilon_0}, \ e = R_{n,\varepsilon_0},$$

where ε_0 is still to be determined. Recall that

$$z_{0,n} = \sum_{j=1}^{J(\varepsilon_0)} \frac{1}{\lambda_{i,n}^{\frac{N-2}{2}}} V_j^l \left(\frac{x - x_{j,n}}{\lambda_{j,n}}, \frac{-t_{j,n}}{\lambda_{j,n}^2} \right) + w_n,$$

where $\|e^{it\Delta}w_n\|_{L^{\frac{2(N+2)}{N-2}}_{L(-\infty,+\infty)}L^{\frac{2N(N+2)}{N-2}}} \le \varepsilon_0$. By the definition of non-linear profile, we now have

$$z_{0,n}(x) = H_{n,\varepsilon_0}(x,0) + \widetilde{w}_n(x),$$

where, for n large $\|e^{it\Delta}\widetilde{w}_n\|_{L^{\frac{2(N+2)}{N-2}}_{(-\infty,+\infty)}L^{\frac{2N(N+2)}{N-2}}} \leq 2\varepsilon_0$. Moreover, according to the orthogonality of $(\lambda_{j,n}; x_{j,n}; t_{j,n})$ and Corollary 6.2, for $n = n(\varepsilon_0)$ large, it holds

$$\int_{\mathbb{R}^{N}} |\nabla H_{n,\varepsilon_{0}}(t)|^{2} dx \leq 2 \sum_{j=1}^{J(\varepsilon_{0})} \int_{\mathbb{R}^{N}} |\nabla U_{j}(\frac{t-t_{j,n}}{\lambda_{j,n}^{2}})|^{2} dx \leq 4C \sum_{j=1}^{J(\varepsilon_{0})} \int_{\mathbb{R}^{N}} |\nabla V_{0,j}|^{2} dx$$

and

$$\sum_{j=1}^{J(\varepsilon_0)} \int_{\mathbb{R}^N} |\nabla V_{0,j}|^2 dx \le \int_{\mathbb{R}^N} |\nabla z_{0,n}|^2 dx + \int_{\mathbb{R}^N} |\nabla z_{0,n}|^2 dx + o(1) \le 2 \int_{\mathbb{R}^N} |\nabla W|^2 dx.$$

Let $M = C_0$ with C_0 as in 8.8,

$$A = \widetilde{C} \int_{\mathbb{R}^N} |\nabla W|^2 dx, A' = A + \int_{\mathbb{R}^N} |\nabla W|^2 dx, \varepsilon_0 < \frac{\varepsilon_0(M, A, A', N)}{2},$$

where $\varepsilon_0(M, A, A', N)$ is defined as in Proposition 5.1. Fix ε_0 and choose n so large that $\|\nabla R_{n,\varepsilon_0}\|_{L^\infty_T H^1 \cap L^2_T L^{\frac{2N}{N+2}}} < \varepsilon_0$ and so that all the above properties hold. Then Proposition 5.1 indicates that the conclusion is valid in the case when (8.1) holds.

Remark 8.1. Assume that $\{z_{0,n}\}$ in Lemma 7.1 are all radial. Then $V_{0,j}$, w_n can be chosen to be radial and we can choose $x_{j,n} \equiv 0$. This follows directly from Keraani's proof [18]. If we then define (SC) and E_C by restricting only to radial functions, we obtain a u_C as in Lemma 8.1 which is radial, and we can establish Lemma 8.2 with $x(t) \equiv 0$.

9 Rigidity theorem

In this section we will prove the following:

Theorem 9.1. Assume that $u_0 \in H^1$ is such that

$$E(u_0) < E(W), \int_{\mathbb{R}^N} |\nabla u_0|^2 dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx.$$

Assume that u be the solution of (1.1) and $u|_{t=0} = u_0$ with maximal interval of existence $(-T_-(u_0), T_+(u_0))$. If there exists $\lambda(t) > 0$, for $t \in [0, T_+(u_0))$, with the property that

$$K = \left\{ v(x,t) = \frac{1}{\lambda(t)^{\frac{N-2}{2}}} u\left(\frac{x}{\lambda(t)}, t\right) : t \in [0, T_{+}(u_{0})) \right\}$$

is such that \overline{K} is compact in H^1 . Then $T_+(u_0) = +\infty$, $u_0 \equiv 0$.

Remark 9.1. We conjecture that Theorem 9.1 remains true if $v(x,t) = \frac{1}{\lambda(t)^{\frac{N-2}{2}}} u\left(\frac{x-x(t)}{\lambda(t)},t\right)$, with $x(t) \in \mathbb{R}^N$, $t \in [0, T_+(u_0))$. In other words, for "energy subcritical" initial data, compactness up to the invariances of the equation, for solutions, is only true for $u \equiv 0$.

We start out with a special case of the strengthened form of Theorem 9.1, i.e.,

Lemma 9.1. Assume that $u, v, \lambda(t), x(t)$ are as in Remark 9.1, that $|x(t)| \leq C_0$ and that $\lambda(t) \geq A_0 > 0$. Then the conclusion of Theorem 9.1 holds. Moreover, if $T_+(u_0) < +\infty$, the hypothesis $|x(t)| \leq C_0$ is not needed.

In the next lemma we will collect some useful facts:

Lemma 9.2. Let u, v be as in Lemma 9.1.

i) Let $\delta_0 > 0$ be such that $E(u_0) \leq (1 - \delta_0)E(W)$. Then for all $t \in [0, T_+(u_0))$, we have

$$\int_{\mathbb{R}^{N}} |\nabla u(t)|^{2} dx \leq (1 - \delta_{1}) \int_{\mathbb{R}^{N}} |\nabla W|^{2} dx,$$

$$\int_{\mathbb{R}^{N}} (|\nabla u|^{2} - F(u)) dx \geq \bar{\delta} \int_{\mathbb{R}^{N}} |\nabla u|^{2} dx,$$

$$\int_{\mathbb{R}^{N}} (|\nabla u|^{2} - f(u)\overline{u}) dx \geq \bar{\delta} \int_{\mathbb{R}^{N}} |\nabla u|^{2} dx,$$

$$C_{1,\delta_{0}} \int_{\mathbb{R}^{N}} |\nabla u_{0}|^{2} dx \leq E(u_{0}) \leq C_{2} \int_{\mathbb{R}^{N}} |\nabla u_{0}|^{2} dx,$$

$$E(u(t)) = E(u_{0}),$$

$$C_{1,\delta_{0}} \int_{\mathbb{R}^{N}} |\nabla u_{0}|^{2} dx \leq \int_{\mathbb{R}^{N}} |\nabla u(t)|^{2} dx \leq C_{2} \int_{\mathbb{R}^{N}} |\nabla u_{0}|^{2} dx.$$

ii)
$$\int_{\mathbb{R}^N} |\nabla v(t)|^2 dx \le C_2 \int_{\mathbb{R}^N} |\nabla W|^2 dx,$$

$$||v(t)||_{L_x^{2^*}}^2 \le C_3 \int_{\mathbb{R}^N} |\nabla W|^2 dx.$$

iii) For all $x_0 \in \mathbb{R}^N$

$$\int_{\mathbb{R}^N} \frac{|v(x,t)|^2}{|x-x_0|^2} dx \le C_4 \int_{\mathbb{R}^N} |\nabla W|^2 dx.$$

iv) For each $\varepsilon_0 > 0$, there exists $R(\varepsilon_0) > 0$, such that, for $0 \le t < T_+(u_0)$, we have

$$\int_{|x|>R(\varepsilon_0)} \left(|\nabla v|^2 dx + F(v) + \frac{|v|^2}{|x|^2} \right) dx \le \varepsilon_0.$$

Proof. Using Theorem 6.1, Corollary 6.2 and Sobolev embedding, it is easy to see that i) and ii) hold. iii) follows from Hardy's inequality. using Sobolev embedding and the Hardy inequality, follows from the compactness of \overline{K} .

The next lemma is a localized virial identity about general nonlinear equation. The proof idea comes from Merle [20].

Lemma 9.3. Let $\psi \in C_0^{\infty}(\mathbb{R}^N)$, $t \in [0, T_+(u_0))$. Then:

i)
$$\frac{d}{dt} \int_{\mathbb{R}^N} |u|^2 \psi dx = 2 \operatorname{Im} \int_{\mathbb{R}^N} \bar{u} \nabla u \nabla \psi dx$$
 ii)

$$\frac{d^2}{dt^2} \int_{\mathbb{R}^N} |u|^2 \psi dx = 4 \int_{\mathbb{R}^N} \Delta \psi |\nabla u|^2 dx + \frac{1}{2} \int_{\mathbb{R}^N} \nabla |u|^2 \nabla (\Delta \psi) dx + \int_{\mathbb{R}^N} \Delta \psi [F(u) - f(u)\bar{u}] dx.$$

Proof. By (1.1) and direct calculation, we get

$$\frac{d}{dt} \int_{\mathbb{R}^N} \psi(x) |u(t,x)|^2 dx = 2 \int_{\mathbb{R}^N} \operatorname{Re} \psi \frac{\partial u}{\partial t} \bar{u} dx$$

$$= 2 \int_{\mathbb{R}^N} \operatorname{Re} \left[i \Delta u + i f(u) \right] \bar{u} \psi dx$$

$$= -2 \operatorname{Im} \int_{\mathbb{R}^N} \Delta u \bar{u} \psi dx$$

$$= 2 \operatorname{Im} \int_{\mathbb{R}^N} \nabla u \bar{u} \nabla \psi dx$$

because of $\operatorname{Im} \int_{\mathbb{R}^N} \nabla u \overline{\nabla u} \psi dx = 0$ and part (i) follows.

Using (i), we know that

$$\begin{split} \frac{d^2}{dt^2} \int_{\mathbb{R}^N} \psi(x) |u(t,x)|^2 dx &= 2 \left[\operatorname{Im} \int_{\mathbb{R}^N} \nabla \psi \bar{u} \nabla \frac{\partial u}{\partial t} dx + \operatorname{Im} \int_{\mathbb{R}^N} \nabla \psi \frac{\overline{\partial u}}{\partial t} \nabla u dx \right] \\ &= 2 \left[2 \operatorname{Im} \int_{\mathbb{R}^N} \nabla \psi \frac{\overline{\partial u}}{\partial t} \nabla u dx - \operatorname{Im} \int_{\mathbb{R}^N} \Delta \psi \bar{u} \frac{\partial u}{\partial t} dx \right]. \end{split}$$

On the one hand,

$$-\operatorname{Im} \int_{\mathbb{R}^{N}} \Delta \psi \bar{u} \frac{\partial u}{\partial t} dx = -\operatorname{Re} \int_{\mathbb{R}^{N}} \Delta \psi \bar{u} \left(\Delta u + f(u)\right) dx$$

$$= -\int_{\mathbb{R}^N} \Delta \psi f(u) \bar{u} dx + \int_{\mathbb{R}^N} \Delta \psi |\nabla u|^2 dx + \frac{1}{2} \int_{\mathbb{R}^N} \nabla |u|^2 \nabla (\Delta \psi) dx.$$

On the other hand, note that

$$-2\operatorname{Re}\int_{\mathbb{R}^{N}}\nabla\psi\Delta u\overline{\nabla u}dx = 2\operatorname{Re}\int_{\mathbb{R}^{N}}\Delta\psi\nabla u\overline{\nabla u}dx + 2\operatorname{Re}\int_{\mathbb{R}^{N}}\nabla\psi\overline{\Delta u}\nabla udx$$
$$= 2\operatorname{Re}\int_{\mathbb{R}^{N}}\Delta\psi\nabla u\overline{\nabla u}dx + 2\operatorname{Re}\int_{\mathbb{R}^{N}}\nabla\psi\overline{\Delta u}\overline{\nabla u}dx$$
$$= 2\operatorname{Re}\int_{\mathbb{R}^{N}}\Delta\psi\nabla u\overline{\nabla u}dx + 2\operatorname{Re}\int_{\mathbb{R}^{N}}\nabla\psi\Delta u\overline{\nabla u}dx,$$

SO

$$-2\operatorname{Re}\int_{\mathbb{R}^N}\nabla\psi\Delta u\overline{\nabla u}dx=\operatorname{Re}\int_{\mathbb{R}^N}\Delta\psi\nabla u\overline{\nabla u}dx.$$

Therefore, we have

$$2\operatorname{Im} \int_{\mathbb{R}^{N}} \nabla \psi \frac{\overline{\partial u}}{\partial t} \nabla u dx = -2\operatorname{Im} \int_{\mathbb{R}^{N}} \nabla \psi \frac{\partial u}{\partial t} \overline{\nabla u} dx$$

$$= -2\operatorname{Re} \int_{\mathbb{R}^{N}} \nabla \psi \Delta u \overline{\nabla u} dx - 2\operatorname{Re} \int_{\mathbb{R}^{N}} \nabla \psi f(u) \overline{\nabla u} dx$$

$$= \operatorname{Re} \int_{\mathbb{R}^{N}} \Delta \psi \nabla u \overline{\nabla u} dx - 2\operatorname{Re} \int_{\mathbb{R}^{N}} \nabla \psi f(u) \overline{\nabla u} dx$$

$$= \int_{\mathbb{R}^{N}} \Delta \psi |\nabla u|^{2} dx + \int_{\mathbb{R}^{N}} \Delta \psi F(u) dx.$$

Combining the above two estimates, we obtain

$$\begin{split} &\frac{d^2}{dt^2} \int_{\mathbb{R}^N} \psi(x) |u(t,x)|^2 dx \\ &= & 2 \left[2 \operatorname{Im} \int_{\mathbb{R}^N} \nabla \psi \frac{\overline{\partial u}}{\partial t} \nabla u dx - \operatorname{Im} \int_{\mathbb{R}^N} \Delta \psi \bar{u} \frac{\partial u}{\partial t} dx \right] \\ &= & 2 \left[\int_{\mathbb{R}^N} \Delta \psi |\nabla u|^2 dx + \int_{\mathbb{R}^N} \Delta \psi F(u) dx - \int_{\mathbb{R}^N} \Delta \psi f(u) \bar{u} dx + \int_{\mathbb{R}^N} \Delta \psi |\nabla u|^2 dx \right. \\ &\left. + \frac{1}{2} \int_{\mathbb{R}^N} \nabla |u|^2 \nabla (\Delta \psi) dx \right] \\ &= & 4 \int_{\mathbb{R}^N} \Delta \psi |\nabla u|^2 dx + \frac{1}{2} \int_{\mathbb{R}^N} \nabla |u|^2 \nabla (\Delta \psi) dx + \int_{\mathbb{R}^N} \Delta \psi [F(u) - f(u) \bar{u}] dx, \end{split}$$

which implies that the conclusion is valid.

Proof of Lemma 9.1. The proof splits in two cases, the finite time blow-up case for u and the infinite time of existence for u.

Case 1 $T_+(u_0) < \infty$. Note first that $\lambda(t) \to \infty$ as $t \to T_+(u_0)$. If not, there exists $t_i \uparrow T_+(u_0)$ such that $\lambda(t_i) \to \lambda_0 < +\infty$. Let

$$v_i(x) = \frac{1}{\left[\lambda\left(t_i\right)\right]^{\frac{N-2}{2}}} u\left(\frac{x - x\left(t_i\right)}{\lambda\left(t_i\right)}, t_i\right).$$

By the compactness of \overline{K} , there exists $v(x) \in H^1$ such that $v_i \to v$ in H^1 . Hence,

$$u\left(x - \frac{x\left(t_{i}\right)}{\lambda\left(t_{i}\right)}, t_{i}\right) = \left[\lambda\left(t_{i}\right)\right]^{\frac{N-2}{2}} v_{i}\left(\left[\lambda\left(t_{i}\right)\right]x\right) \to \lambda_{0}^{\frac{N-2}{2}} v\left(\lambda_{0}x\right) \text{ in } H^{1}$$

because of $\lambda(t_i) \geq A_0, \lambda_0 \geq A_0$. Let now h(x,t) be the solution of (1.1) with data $\lambda_0^{\frac{N-2}{2}}v(\lambda_0 x)$ at time $T_+(u_0)$ and

$$||h||_{L^{\frac{2(N+2)}{N-2}}_{(T_{+}(u_{0})-\delta,T_{+}(u_{0})+\delta)}W^{1,\frac{2N(N+2)}{N^{2}+4}}} < +\infty$$

in an interval $(T_+(u_0) - \delta, T_+(u_0) + \delta)$. Let $h_i(x,t)$ be the solution with data at $T_+(u_0)$ equal to $u\left(x - \frac{x(t_i)}{\lambda(t_i)}, t_i\right)$. Then Lemmas 3.1 and 3.2 guarantee that

$$\sup_{i} \|h_{i}\|_{L_{(T_{+}(u_{0})-\delta,T_{+}(u_{0})+\delta)}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^{2}+4}}} < +\infty.$$

But $u\left(x-\frac{x(t_i)}{\lambda(t_i)},t+t_i-T_+\left(u_0\right)\right)=h_i(x,t)$, contradicting finite blow-up since $T_+\left(u_0\right)<\infty$.

Let us prove now a decay result for u from the concentration properties in L^{2^*} of u at $T_+(u_0)$. Let us now fix $\varphi \in C_0^{\infty}(\mathbb{R}^N)$, φ radial, $\varphi \equiv 1$ for $|x| \leq 1$, $\varphi \equiv 0$ for $|x| \geq 2$ and set $\varphi_R(x) = \varphi(\frac{x}{R})$. Define

$$y_R(t) = \int_{\mathbb{R}^N} |u(x,t)|^2 \varphi_R(x) dx, \quad t \in [0, T_+(u_0)).$$

We then have

$$|y_R'| \le C_N \int_{\mathbb{D}^N} |\nabla W|^2 dx. \tag{9.1}$$

Indeed, it follows Lemma 9.2 and 9.3 that

$$|y_R'| \leq \frac{2}{R} \left| \operatorname{Im} \int_{\mathbb{R}^N} \bar{u} \nabla u \nabla \varphi(\frac{x}{R}) dx \right|$$

$$\leq C_N \left(\int_{\mathbb{R}^N} |\nabla u|^2 dx \right)^{\frac{1}{2}} \cdot \left(\int_{\mathbb{R}^N} \frac{|u|^2}{|x|^2} dx \right)^{\frac{1}{2}} \leq C_N \int_{\mathbb{R}^N} |\nabla W|^2 dx,$$

We also have:

For all
$$R > 0$$
, $\int_{|x| < R} |u(x, t)|^2 dx \to 0$ as $t \to T_+(u_0)$. (9.2)

In fact, $u(y,t) = [\lambda(t)]^{\frac{N-2}{2}} v(\lambda(t)y + x(t), t)$, so that

$$\int_{|x| < R} |u(x,t)|^2 dx$$
= $[\lambda(t)]^{-2} \int_{|y| < R\lambda(t)} |v(y+x(t),t)|^2 dy$
= $[\lambda(t)]^{-2} \int_{B(x(t),R\lambda(t))} |v(z,t)|^2 dz$

$$= [\lambda(t)]^{-2} \int_{B(x(t),R\lambda(t))\cap B(0,\varepsilon R\lambda(t))} |v(z,t)|^2 dz + [\lambda(t)]^{-2} \int_{B(x(t),R\lambda(t))\backslash B(0,\varepsilon R\lambda(t))} |v(z,t)|^2 dz$$

$$= A + B.$$

By Hölder inequality $(\frac{2}{2^*} + \frac{2}{N} = 1)$,

$$A \leq [\lambda(t)]^{-2} (\varepsilon R \lambda(t))^{N \cdot \frac{2}{N}} ||v||_{L^{2^*}}^2 \leq \varepsilon^2 R^2 C_3 \int_{\mathbb{R}^N} |\nabla W|^2 dx,$$

which is small with ε . Now, fixed $\varepsilon > 0$, note that $\lambda(t) \to +\infty$ as $t \to T_+(u_0)$, by Lemma 9.2 and Hölder inequality, we have

$$B \leq [\lambda(t)]^{-2} (R\lambda(t))^{N\frac{2}{N}} ||v||_{L^{2^*}(|x| \geq \varepsilon R\lambda(t))}^2 = R^2 ||v||_{L^{2^*}(|x| \geq \varepsilon R\lambda(t))}^2 \to 0 \text{ as } t \to T_+(u_0),$$

According to (9.1) and (9.2), we have

$$y_R(0) \le y_R(T_+(u_0)) + C_N T_+(u_0) \int_{\mathbb{R}^N} |\nabla W|^2 dx = C_N T_+(u_0) \int_{\mathbb{R}^N} |\nabla W|^2 dx.$$

Hence, letting $R \to +\infty$ we obtain $u_0 \in L^2(\mathbb{R}^N)$. Arguing as before,

$$|y_R(t) - y_R(T_+(u_0))| \le C_N(T_+(u_0) - t) \int_{\mathbb{R}^N} |\nabla W|^2 dx,$$

so that

$$y_R(t) \le C_N \left(T_+ \left(u_0 \right) - t \right) \int_{\mathbb{R}^N} |\nabla W|^2 dx.$$

Letting $R \to \infty$, we see that

$$||u(t)||_{L^2}^2 \le C_N (T_+(u_0) - t) \int_{\mathbb{R}^N} |\nabla W|^2 dx$$

and so by the conservation of the L^2 norm $||u(T_+(u_0))||_{L^2} = ||u_0||_{L^2} = 0$. But then $u \equiv 0$, contradicting $T_+(u_0) < \infty$.

Case 2 $T_+(u_0) = +\infty$. In this case we assume, in addition, that $|x(t)| \leq C_0$. We first note that for each $\varepsilon > 0$, there exists $R(\varepsilon) > 0$ such that, for all $t \in [0, \infty)$, it holds

$$\int_{|x|>R(\varepsilon)} \left(|\nabla u|^2 + F(u) + \frac{|u|^2}{|x|^2} \right) dx \le \varepsilon, \tag{9.3}$$

$$\int_{|x|>R(\varepsilon)} \left(|\nabla u|^2 + f(u)\overline{u} + \frac{|u|^2}{|x|^2} \right) dx \le \varepsilon, \tag{9.4}$$

In fact, $u(y,t) = [\lambda(t)]^{\frac{N-2}{2}} v(\lambda(t)y + x(t), t)$, so that

$$\int_{|y|>R(\varepsilon)} |\nabla u(y,t)|^2 dy = \int_{|y|>R(\varepsilon)} [\lambda(t)]^N |\nabla v(\lambda(t)y + x(t),t)|^2 dy$$

$$= \int_{|z|>R(\varepsilon)\lambda(t)} |\nabla v(z + x(t),t)|^2 dz$$

$$\leq \int_{|z| \geq R(\varepsilon)A_0} |\nabla v(z + x(t), t)|^2 dz$$

$$\leq \int_{|\alpha| > R(\varepsilon)A_0 - C_0} |\nabla v(\alpha, t)|^2 d\alpha$$

and the statement for this term now follows from Lemma 9.2 iv). The other terms are handled similarly. There exists $R_0 > 0$ such that, for all $t \in [0, +\infty)$, we have

$$8\int_{|x| \le R_0} |\nabla u|^2 dx - 8\int_{|x| \le R_0} F(u) dx \ge C_{\delta_0} \int_{\mathbb{R}^N} |\nabla u_0|^2 dx. \tag{9.5}$$

$$8 \int_{|x| \le R_0} |\nabla u|^2 dx - 8 \int_{|x| \le R_0} f(u) \overline{u} dx \ge C_{\delta_0} \int_{\mathbb{R}^N} |\nabla u_0|^2 dx.$$
 (9.6)

In fact, (6.8) combined with Lemma 9.2 i) yields

$$8 \int_{\mathbb{R}^N} |\nabla u|^2 dx - 8 \int_{\mathbb{R}^N} F(u) dx \ge \widetilde{C}_{\delta_0} \int_{\mathbb{R}^N} |\nabla u_0|^2 dx.$$

Now combine this with (9.3), with $\varepsilon = \varepsilon_0 \int_{\mathbb{R}^N} |\nabla u_0|^2 dx$ to obtain (9.5). Similarly, it can be proven that (9.6) is established.

To prove Case 2, we choose $\varphi \in C_0^{\infty}(\mathbb{R}^N)$, radial, with $\varphi(x) = |x|^2$ for $|x| \leq 1$, $\varphi(x) \equiv 0$ for $|x| \geq 2$. Define

$$z_R(t) = \int_{\mathbb{R}^N} |u(x,t)|^2 R^2 \varphi(\frac{x}{R}) dx,$$

then we have

$$|z_R'(t)| \le C_{N,\delta_0} \int_{\mathbb{R}^N} |\nabla u_0|^2 R^2 dx \text{ for } t > 0,$$

$$z_R'' \ge C_{N,\delta_0} \int_{\mathbb{R}^N} |\nabla u_0|^2 dx \text{ for } R \text{ large enough, } t > 0.$$

In fact, from Lemma 9.3 i),

$$|z_R'(t)| \leq 2R \left| \operatorname{Im} \int_{\mathbb{R}^N} \overline{u} \nabla u \nabla \varphi(\frac{x}{R}) dx \right| \leq C_N R \int_{0 \leq x \leq 2R} \frac{|x|}{|x|} |\nabla u| |u| dx$$

$$\leq C_N R^2 \left(\int_{\mathbb{R}^N} |\nabla u|^2 dx \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^N} \frac{|u|^2}{|x|^2} dx \right)^{\frac{1}{2}}$$

$$\leq C_N R^2 \int_{\mathbb{R}^N} |\nabla u_0|^2 dx$$

because of Lemma 9.2 i). In view of (9.3), (9.4), (9.5), (9.6) and Lemma 9.3, ii), we have

$$z_R''(t) = 4 \int_{\mathbb{R}^N} \Delta \varphi |\nabla u|^2 dx + \frac{1}{2} \int_{\mathbb{R}^N} \nabla |u|^2 \nabla (\Delta \varphi) dx + \int_{\mathbb{R}^N} \Delta \varphi [F(u) - f(u)\bar{u}] dx$$

$$\geq 4 \int_{\mathbb{R}^N} [|\nabla u|^2 - F(u)] dx + 4 \int_{\mathbb{R}^N} [|\nabla u|^2 - f(u)\bar{u}] dx$$

$$\geq 4 \int_{|x| \le R} [|\nabla u|^2 - F(u)] dx + 4 \int_{|x| \le R} [|\nabla u|^2 - f(u)\bar{u}] dx$$

$$-C_N \int_{R \le |x| \le 2R} \left[|\nabla u|^2 + \frac{|u|^2}{|x|^2} + |u|^{2^*} \right] dx$$

$$\geq C_{N,\delta_0} \int_{\mathbb{R}^N} |\nabla u_0|^2 dx$$

for R large. If we now integrate in t, we have $z_R'(t) - z_R'(0) \ge C_{N,\delta_0} t \int_{\mathbb{R}^N} |\nabla u_0|^2 dx$, but we also have $|z_R'(t) - z_R'(0)| \le 2C_N R^2 \int_{\mathbb{R}^N} |\nabla u_0|^2 dx$, a contradiction for t large, unless $\int_{\mathbb{R}^N} |\nabla u_0|^2 dx = 0$.

Proof of Theorem 9.1. Assume that $u_0 \not\equiv 0$ so that $\int_{\mathbb{R}^N} |\nabla u_0|^2 dx > 0$ and because of Lemma 9.2 i) (which is still valid here), $E(u_0) \geq C_{1,\delta_0} \int_{\mathbb{R}^N} |\nabla u_0|^2 dx$ and hence $E(u_0) > 0$. Because of Lemma 9.1, we only need to treat the case where there exists $\{t_n\}_{n=1}^{\infty}$, $t_n \geq 0$, $t_n \uparrow T_+(u_0)$, so that

$$\lambda\left(t_{n}\right) \to 0.$$

(If $t_n \to t_0 \in [0, T_+(u_0))$, we obtain for all R > 0, $\int_{|x| \ge R} |v(t_0)|^{2*} dx = 0$ but $\int_{\mathbb{R}^N} |\nabla v(t_0)|^2 dx > 0$). After possibly redefining $\{t_n\}_{n=1}^{\infty}$ we can assume that

$$\lambda(t_n) \le 2 \inf_{t \in [0, t_n]} \lambda(t)$$

and from our hypothesis

$$w_n(x) = \frac{1}{\lambda(t_n)^{\frac{N-2}{2}}} u\left(\frac{x}{\lambda(t_n)}, t_n\right) \to w_0 \text{ in } H^1.$$

By Theorem 6.1 we have

$$E(W) > E(w_0) = E(u_0) > 0, \int_{\mathbb{R}^N} |\nabla u(t)|^2 dx \le (1 - \delta_1) \int_{\mathbb{R}^N} |\nabla W|^2 dx$$

so that $\int_{\mathbb{R}^N} |\nabla w_0|^2 dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx$. Thus $w_0 \not\equiv 0$. Let us now consider solutions of (1.1), $w_n(x,\tau), w_0(x,\tau)$ with data $w_n(-,0), w_0(-,0)$ at $\tau=0$, defined in maximal intervals $\tau\in (-T_-(w_n),0]$, $\tau\in (-T_-(w_0),0]$ respectively. Since $w_n\to w_0$ in H^1 , $\lim_{n\to\infty} T_-(w_n)\geq T_-(w_0)$ and for each $\tau\in (-T_-(w_0),0]$, $w_n(x,\tau)\to w_0(x,\tau)$ in H^1 .

Note that by uniqueness in (1.1), for $0 \le t_n + \frac{\tau}{[\lambda(t_n)]^2}$, $w_n(x,\tau) = \frac{1}{[\lambda(t_n)]^{\frac{N-2}{2}}} u(\frac{x}{[\lambda(t_n)]}, t_n + \frac{\tau}{[\lambda(t_n)]^2})$. Remark that $\lim_{n \to \infty} \tau_n = \lim_{n \to \infty} t_n [\lambda(t_n)]^2 \ge T_-(w_0)$ and thus for all $\tau \in (-T_-(w_0), 0]$ for n large, $0 \le t_n + \frac{\tau}{\lambda(t_n)^2} \le t_n$. Indeed, if $\tau_n \to \tau_0 < T_-(w_0)$, then $w_n(x, -\tau_n) = \frac{1}{[\lambda(t_n)]^{\frac{N-2}{2}}} u_0\left(\frac{x}{\lambda(t_n)}\right) \to w_0(x, -\tau_0)$ in H^1 with $\lambda(t_n) \to 0$ which is a contradiction from $u_0 \not\equiv 0, w_0 \not\equiv 0$.

Fix now $\tau \in (-T_{-}(w_0), 0]$, for n sufficiently large $v\left(x, t_n + \frac{\tau}{\lambda(t_n)^2}\right)$, $\lambda\left(t_n + \frac{\tau}{\lambda(t_n)^2}\right)$ are defined and we have

$$v\left(x,t_n + \frac{\tau}{\lambda \left(t_n\right)^2}\right)$$

$$= \frac{1}{\lambda \left(t_n + \frac{\tau}{\lambda(t_n)^2}\right)^{\frac{N-2}{2}}} u \left(\frac{x}{\lambda \left(t_n + \frac{\tau}{\lambda(t_n)^2}\right)}, t_n + \frac{\tau}{\lambda \left(t_n\right)^2}\right)$$

$$= \frac{1}{\widetilde{\lambda}_n(\tau)^{\frac{N-2}{2}}} w_n \left(\frac{x}{\widetilde{\lambda}_n(\tau)}, \tau\right),$$

where

$$\widetilde{\lambda}_n(\tau) = \frac{\lambda \left(t_n + \frac{\tau}{\lambda(t_n)^2}\right)}{\lambda(t_n)} \ge \frac{1}{2}$$

because of the fact $\lambda(t_n) \leq 2 \inf_{t \in [0,t_n]} \lambda(t)$. One can assume after passing to a subsequence that $\widetilde{\lambda}_n \left(t_n + \frac{\tau}{\lambda(t_n)^2} \right) \to \widetilde{\lambda}_0(\tau)$ with $\frac{1}{2} \leq \widetilde{\lambda}_0(\tau) \leq +\infty$ and $v\left(x, t_n + \frac{\tau}{\lambda(t_n)^2} \right) \to v_0(x,\tau)$ in H^1 , as $n \to \infty$. Remark that $\widetilde{\lambda}_0(\tau) < +\infty$. If not, we will have

$$\frac{1}{\widetilde{\lambda}_n(\tau)^{\frac{N-2}{2}}} w_0\left(\frac{x}{\widetilde{\lambda}_n(\tau)}, \tau\right) \to v_0(x, \tau),$$

which implies $w_0(x,\tau) = 0$ which contradicts $E(w_0) = E(u_0) > 0$. Thus $\widetilde{\lambda}_0(\tau) < +\infty$ and $v_0(x,\tau) = \frac{1}{\widetilde{\lambda}_0(\tau)^{\frac{N-2}{2}}} w_0\left(\frac{x}{\widetilde{\lambda}_0(\tau)},\tau\right)$ where $v_0(\tau) \in \overline{K}$. We thus obtain a contradiction from Lemma 9.1.

Corollary 9.1. Assume that $E(u_0) < E(W)$, $\int_{\mathbb{R}^N} |\nabla u_0|^2 dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx$ and u_0 is radial. Then the solution u of the Cauchy problem (1.1) with data u_0 at t=0 has time interval of existence $I=(-\infty,+\infty)$, $||u||_{L_I^{2(N+2)}W^{1,\frac{2N(N+2)}{N^2+4}}} < +\infty$ and there exists $u_{0,+},u_{0,-}$ in H^1 such that

$$\lim_{t \to +\infty} \|u(t) - e^{it\Delta} u_{0,+}\|_{H^1} = 0, \ \lim_{t \to -\infty} \|u(t) - e^{it\Delta} u_{0,-}\|_{H^1} = 0.$$

Moreover, if we define δ_0 so that $E(u_0) \leq (1 - \delta_0)E(W)$, there exists a function $M(\delta_0)$ so that

$$||u||_{L_{I}^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^{2}+4}}} \le M(\delta_{0}).$$

Proof. By the integral equation in Lemma 3.1, we know that u(t) is radial for each $t \in I$. Using Remark 8.1 and Theorem 9.1 we obtain $I = (-\infty, +\infty), \|u\|_{L_I^{\frac{2(N+2)}{N-2}}W^{1,\frac{2N(N+2)}{N^2+4}}} < +\infty$. Now Remark 5.1 finishes the proof of the first statement.

For the last statement, let

$$D_{\delta_0} = \left\{ u_0 \in H^1 \text{ radial}, \int_{\mathbb{R}^N} |\nabla u_0|^2 dx < \int_{\mathbb{R}^N} |\nabla W|^2 dx \text{ and } E(u_0) \le (1 - \delta_0) E(W) \right\}$$
$$M(\delta_0) = \sup_{u \in D_{\delta_0}} ||u||_{L_I^{\frac{2(N+2)}{N-2}} W^{1,\frac{2N(N+2)}{N^2+4}}}.$$

We need to prove $M(\delta_0) < +\infty$. If not there is a sequence $u_{0,n}$ in D_{δ_0} and the corresponding solutions u_n such that $\|u_n\|_{L_I^{2(N+2)} N^{-2} W^{1,\frac{2N(N+2)}{N^2+4}}} \to +\infty$ as $n \to +\infty$. Note that we can assume that $\|e^{it\Delta}u_{0,n}\|_{L_I^{2(N+2)} N^{1,\frac{2N(N+2)}{N^2+4}}} \ge \rho$, with ρ as in Lemma 3.1. Arguing as in the proof of Lemma 8.1, we would conclude that first J=1 in the decomposition given in Lemma 7.1 and then since $\|U_1\|_{L_I^{2(N+2)} N^{-2} W^{1,\frac{2N(N+2)}{N^2+4}}} < +\infty$ we reach a contradiction. \square

References

- [1] H. Bahouri, G. Perelman, Global well-posedness for the derivative nonlinear Schrödinger equation. Invent. Math., 229(2)(2022), 639–688.
- [2] H. Berestycki, P. L. Lions, Nonlinear scalar field equations, I. existence of a ground state, Arch. Ration. Mech. Anal., 82(1983), 313–345.
- [3] H. Berestycki, P. L. Lions, Nonlinear scalar field equations, II. Existence of infinitely many solutions, Arch. Rational Mech. Anal, 82(4)(1983), 347–375.
- [4] J. Bergh, J. Lofstrom, Interpolation Spaces. An Introduction. Grundlehren der Mathematischen Wissenschaften, No. 223. Berlin, New York: Springer 1976.
- [5] T. Boulenger, D. Himmelsbach, E. Lenzmann, Blow up for fractional NLS, J. Funct. Anal, 271(2016), 2569–2603.
- [6] J. Bourgain, A. Bulut, Almost sure global well posedness for the radial nonlinear Schrödinger equation on the unit ball I: the 2D case, Annales de l'Institut Henri Poincaré C, Analyse non linéaire. No longer published by Elsevier, 2014, 31(6)(2014), 1267–1288.
- [7] J. Bourgain, Global well-posedness of defocusing critical nonlinear Schrödinger equation in the radial case, J. Am. Math. Soc., 12(1999), 145–171.
- [8] J. Bourgain, New global well-posedness results for nonlinear Schrödinger equations, AMS Colloquium Publications, 46, 1999.
- [9] T. Cazenave, Semilinear Schrödinger Equations. Courant Lecture Notes in Mathematics, vol. 10. New York: New York University Courant Institute of Mathematical Sciences 2003.
- [10] T. Cazenave, An Introduction to Nonlinear Schrödinger Equations, Text. Metod. Mat., vol. 26, Univ. Fed. Rio de Janeiro, 1993.
- [11] T. Cazenave, F. B. Weissler, The Cauchy problem for the critical nonlinear Schrödinger equation in H^s , Nonlinear Anal., Theory Methods Appl., 14(1990), 807–836.

- [12] V. D. Dinh, Well-posedness of nonlinear fractional Schrödinger and wave equations in Sobolev spaces, Int. J. Appl. Math., 31 (2018), 483–525.
- [13] J. Ginibre, G. Velo, On a class of nonlinear Schrödinger equations, I, The Cauchy problem, general case, J. Funct. Anal., 32(1)(1979), 1–32.
- [14] J. Ginibre and G. Velo, Scattering theory in the energy space for a class of nonlinear Schrödinger equations, J. Math. Pures Appl., 64(1984), 363–401.
- [15] D. Hennig, N. I. Karachalios, D. Mantzavinos, J. Cuevas-Maraver, I. G. Stratis, On the proximity between the wave dynamics of the integrable focusing nonlinear Schrödinger equation and its non-integrable generalizations, J. Differ. Equ., 397(2024), 106-165.
- [16] M. Keel, T. Tao, Endpoint Strichartz estimates, Am. J. Math., 120(1998), 955–980.
- [17] C. E. Kenig, F. Merle, Global well-posedness, scattering and blow-up for the energy-critical, focusing, non-linear Schrödinger equation in the radial case, Invent. Math., 166(3)(2006), 645–675.
- [18] S. Keraani, On the defect of compactness for the Strichartz estimates of the Schrödinger equations, J. Differ. Equations, 175(2001), 353–392.
- [19] S. Keraani, On the blow up phenomenon of the critical Schrödinger equation, J. Funct. Anal., 235(2006), 171–192.
- [20] F. Merle, On uniqueness and continuation properties after blow-up time of self-similar solutions of nonlinear schrödinger equation with critical exponent and critical mass, Comm. Pure Appl. Math, 45(2)(1992), 203-254.
- [21] T. Oh, Y. Wang, Global well-posedness of the one-dimensional cubic nonlinear Schrödinger equation in almost critical spaces, J. Differ. Equ., 269(1)(2020), 612–640.
- [22] T. Tao, M. Visan, X. Zhang, Global well-posedness and scattering for the defocusing mass-critical nonlinear Schrödinger equation for radial data in high dimensions, Duke Math. J., 140(1)(2007), 165–202.
- [23] K. Yajima, Existence of solutions for Schrödinger evolution equations, Comm. Math. Phys., 110(1987), 415–426.
- [24] X. Yu, H. Yue, Z. Zhao, Global Well-posedness and scattering for fourth-order Schrödinger equations on waveguide manifolds, SIAM J. Math. A., 56(1)(2024), 1427–1458.