

Holographic spectral function of fermion in instantonic plasma

Si-wen Li¹, Yi-peng Zhang², Hao-qian Li³,

*Department of Physics, School of Science,
Dalian Maritime University,
Dalian 116026, China*

Abstract

Using the gauge-gravity duality, we investigate the fermionic correlation function in the D(-1)-D3 brane system which describes the instantonic plasma in holography. In this system, the charge of the D(-1) brane as the D-instanton gives the gluon condensate. To simplify the holographic setup, we first reduce briefly the ten-dimensional supergravity background produced by D(-1)-D3-branes to an equivalently five-dimensional background. Then using the standard method for computing the Green function in the AdS/CFT dictionary, we derive the equations for the fermionic correlation functions and solve them numerically with the infalling boundary condition. Our numerical results illustrate that the fermionic correlation function as spectral function includes two branches of the dispersion curves whose behavior is very close to the results obtained from the method of hard thermal loop. And the effective mass generated by the medium effect of fermion splits into two values due to the spin-dependent interactions induced by instantons. Therefore, this work in holography demonstrates the instantonic configuration is very influential to the fermionic plasmino in plasma.

¹Email: siwenli@dlmu.edu.cn

²Email: ypmahler111@dlmu.edu.cn

³Email: lihaoqian@dlmu.edu.cn

Contents

1	Introduction	2
2	The holographic background for instantonic plasma	4
2.1	The D3-brane background with D-instanton	4
2.2	The effective 5d background	5
3	The fermionic correlation function	6
3.1	General setup for the fermionic correlation function	6
3.2	Covariant Dirac equation for a probe fermion	7
3.3	Construction for the correlation function	8
4	The numerical analysis	12
5	Summary	16

1 Introduction

In quantum chromodynamics (QCD), instanton is known as the nontrivially topological excitation of the vacuum which contributes to the thermodynamics of QCD and relates to the breaking of chiral symmetry [1, 2, 3]. Instanton is also known to consist of constituents which is called BPS (Bogomol’nyi-Prasad-Sommerfield) monopoles or dyons [4], so there have been a lot of continuous researches to connect confinement and the breaking of chiral symmetry to the instanton constituents [5, 6, 7, 8, 9, 10, 11, 12, 13]. In particular, the dynamics of fermion involving instanton e.g. the spin-dependent interaction, fermionic quantum tunneling of the instantonic vacuum, is very interesting and widely discussed in the textbooks of QCD [1, 2, 3].

However, due to the property of asymptotic freedom, it is very challenging to investigate QCD in low-energy region by using the perturbative method in quantum field theory (QFT) since QCD is strongly coupled in low-energy region. Fortunately, the gauge-gravity duality and AdS/CFT correspondence provide an alternative way to study the strongly coupled QFT [14, 15, 16] in holography. In the top-down approach, instanton is constructed simply as a geometric background with instantonic D-branes (D-instanton) [17, 18, 19, 20, 21], for example, the black D(-1)-D3 brane system can describe holographically the plasma with instanton in which the D(-1)-brane plays the role of instanton, and its dual field theory is $\mathcal{N} = 4$ super Yang-Mills theory with instanton or theta angle. So there have been many works based on the holographic instanton e.g. thermodynamics of Yang-Mills theory [22], hadronic spectrum [23, 24], real-time dynamics [25], Chern-Simons theory [26, 27], chiral symmetry breaking [28], quark potential [29, 30, 31], Schwinger effect [32, 33, 34], the hadronic interaction [35, 36]. Despite all this, the study of fermion with the instanton remains to be blank to the existing works in holography

although it attracts great interests as it is reviewed in [1, 2, 3] in QFT. Therefore, the goal of this work is to explore the fermionic correlation function with instanton as a start point for the further continuous investigations, since it can be evaluated by using the most fundamental principle in holography.

To our goal, the setup in this work starts with the D(-1)-D3 brane system since it describes simply the instantonic plasma in holography, then we reduce briefly the 10d IIB supergravity background produced by black D3-branes with D(-1)-branes [17, 25, 28] to an effectively five-dimensional (5d) geometric background. Afterwards, we probe a bulk fermion in the 5d geometric background and derive its covariant Dirac equation. In order to evaluate the fermionic correlation function in the dual theory, we employ the standard technique in the AdS/CFT dictionary so that obtain the differential equations for the fermionic correlators in the bulk. By imposing the infalling boundary condition at the horizon, we afterwards solve the fermionic correlation functions numerically. Our numerical calculation illustrates that, without the D-instanton, there is only one branch of the holographic dispersion curves for fermion lying on the light cone which is similar as the existing works about holographic fermions [35, 36, 37, 38, 39, 40]. In the presence of the D-instanton, the fermionic correlation function has a correction which in leading order splits the dispersion curves into two branches at small momentum, and this behavior is very close to the dispersion curves from the approach of hard thermal loop (HTL) in QFT. Furthermore, the two branches of the dispersion curves reveals that the effective bound mass of fermion in plasma has two values in the presence of the D-instanton. Remarkably it agrees qualitatively with effective potential for the spin-dependent interactions induced by instantons computed by the process of one-gluon exchange in QCD [1, 2]. Therefore, this work, as a start point, is a holographic reproduction of the properties of fermion and instanton and we believe the instantonic configuration in plasma is very influential to the fermionic plasmino.

The outline of this manuscript is given as follows. In Section 2, we collect the essentials of the 10d supergravity background produced by D3-branes with D(-1)-branes as D-instantons, and reduce it briefly to an effectively 5d gravity background. In Section 3, we demonstrate the principle in AdS/CFT dictionary to compute the holographic correlation function for spinor. Then we derive the covariant Dirac equation and the holographic counter term for spinor in order to obtain the differential equations for the fermionic correlators. In Section 4, we solve numerically the holographic fermionic correlation functions order by order with physical analysis and interpretation. The final section is the summary.

Notation

In this manuscript, the capital letters $L, M, N\dots$ refer to the indices of the bulk coordinate as x^M , the lowercase letters a, b, c refer to the corresponding indices of the coordinate in tangent space. The gamma matrices γ^a, Γ^M as the generators of the Clifford algebra with elfbein e_M^a are related as,

$$\{\gamma^a, \gamma^b\} = 2\eta^{ab}, \{\Gamma^M, \Gamma^N\} = 2g^{MN}, \Gamma^M = e_a^M \gamma^a, g_{MN} = e_M^a \eta_{ab} e_N^a, \quad (1.1)$$

where g_{MN} refers to the metric on a manifold and η_{ab} refers to the Minkowskian metric. The spin connection ω_{Mab} is defined as,

$$\omega_{Mab} = \eta_{cb} (e_a^N \partial_M e_N^c - e_a^N \Gamma_{MN}^K e_K^c), \quad (1.2)$$

where the affine connection Γ_{MN}^K is given by,

$$\Gamma_{MN}^K = g^{KL} (\partial_M g_{LN} + \partial_N g_{ML} - \partial_L g_{MN}). \quad (1.3)$$

The Greek letters as μ, ν, \dots refer to the indices of coordinate as x^μ at the holographic boundary. The lowercase letters i, j, k, l run over the spatial directions of the holographic boundary i.e. $i, j, k, l = 1, 2, 3$. The gamma matrices presented in this work are chosen as $\gamma^a = \{\gamma^\mu, \gamma\}$ as,

$$\gamma^\mu = i \begin{pmatrix} 0 & \sigma^\mu \\ \bar{\sigma}^\mu & 0 \end{pmatrix}, \gamma = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (1.4)$$

with $\sigma^\mu = (1, -\sigma^i)$, $\bar{\sigma}^\mu = (1, \sigma^i)$, $i = 1, 2, 3$, where σ^i s are the Pauli matrices.

2 The holographic background for instantonic plasma

In this section, let us briefly review the holographic background for instantonic plasma which corresponds to the system consisted of N_c D3-branes with N_D D-instantons i.e. the D(-1)-branes in the large- N_c limit in the type IIB string theory [17, 18, 25]. Then, we integrate out the S^5 part in order to obtain a five-dimensional (5d) background geometry for simplification.

2.1 The D3-brane background with D-instanton

The supergravity solution of N_c D3-branes with N_D D-instantons is in general described by a 10d deformed D3-brane solution with a non-trivial Ramond-Ramond (R-R) zero form C_0 . It is recognized as a marginal ‘‘bound state’’ of D3-branes with smeared N_D D(-1)-branes. In our notation, N_c D3-branes are identified as color branes, hence its number N_c denotes the color number. In the large N_c limit, the 10d bosonic type IIB supergravity action describes the low-energy dynamics of this system which is given in string frame as,

$$S_{\text{IIB}} = \frac{1}{2\kappa_{10}^2} \int d^{10}x \sqrt{-g} \left[e^{-2\Phi} (\mathcal{R} + 4\partial\Phi \cdot \partial\Phi) - \frac{1}{2} |F_1|^2 - \frac{1}{2} |F_5|^2 \right], \quad (2.1)$$

where $2\kappa_{10}^2 = (2\pi)^7 l_s^8$ is the 10d gravity coupling constant, l_s, g_s is respectively the string length and the string coupling constant. We use Φ to denote the dilaton field and $F_{1,5}$ is the field

strength of the R-R zero and four form $C_{0,4}$ respectively. The solution of N_c D3-branes with N_D D-instantons is the near-horizon solution of non-extremal D3-branes with a non-trivial C_0 , which in string frame reads [17, 18, 25],

$$\begin{aligned}
ds^2 &= e^{\frac{\phi}{2}} \left\{ \frac{r^2}{R^2} [-f(r) dt^2 + d\mathbf{x} \cdot d\mathbf{x}] + \frac{1}{f(r)} \frac{R^2}{r^2} dr^2 + R^2 d\Omega_5^2 \right\}, \\
e^\phi &= 1 + \frac{Q}{r_H^4} \ln \frac{1}{f(r)}, \quad f(r) = 1 - \frac{r_H^4}{r^4}, \quad F_5 = dC_4 = g_s^{-1} \mathcal{Q}_3 \epsilon_5, \\
F_1 &= dC_0, \quad C_0 = -ie^{-\phi} + i\mathcal{C}, \quad \phi = \Phi - \Phi_0, \quad e^{\Phi_0} = g_s,
\end{aligned} \tag{2.2}$$

where \mathcal{C} is a boundary constant for C_0 , ϵ_5 is the volume element of a unit S^5 . And the associated parameters are given as,

$$R^4 = 4\pi g_s N_c l_s^4, \quad \mathcal{Q}_3 = 4R^4, \quad Q = \frac{N_D (2\pi)^4 \alpha'^2}{N_c V_4} \mathcal{Q}_3. \tag{2.3}$$

Here, $x^\mu = \{t, \mathbf{x}\} = \{t, x^i\}$, $i = 1, 2, 3$ refers to the 4d spacetime where the D3-branes are extended along. And r is the holographic direction perpendicular to the D3-branes. The solution (2.2) is asymptotic $\text{AdS}_5 \times S^5$ at the holographic boundary $r \rightarrow \infty$ which describes geometrically that the D-instanton charge N_D is smeared over the worldvolume V_4 of the N_c coincident black D3-branes homogeneously with a horizon at $r = r_H$. And backreaction of the D-instantons has been taken into account in the background, thus it implies N_D/N_c must be fixed in the large- N_c limit. The dual theory of this background is conjectured as the 4d $\mathcal{N} = 4$ super Yang-Mills theory (SYM) in a self-dual gauge field (instantonic) background or with a dynamical axion at finite temperature characterized by the parameter Q [17]. Note that, in the language of hadron physics, C_0 is recognized as axion and the gluon condensate in this system can be evaluated as,

$$\langle \text{Tr} F_{\mu\nu} F^{\mu\nu} \rangle \simeq \frac{N_D}{16\pi^2 V_4} = \frac{Q}{(2\pi\alpha')^2 R^4} \frac{N_c}{(2\pi)^4}, \quad \mu, \nu = 0, 1, \dots, 3 \tag{2.4}$$

2.2 The effective 5d background

The S^5 part in the gravity solution (2.1) and (2.2) can be integrated out so that the background (2.2) can reduce to an 5d background equivalently. Since all the fields presented in (2.1) and (2.2) are expected to depend on the coordinates x^μ, r only, it is possible to integrate out the S^5 part in action with

$$ds_{(10d)}^2 = ds_{(5d)}^2 + ds_{S^5}^2. \tag{2.5}$$

So the action (2.1) becomes,

$$S_{\text{IIB}}^{(5\text{d})} = \frac{1}{2\kappa_5^2} \int d^5x \sqrt{-g^{(5\text{d})}} \left[\mathcal{R}^{5\text{d}} - \frac{1}{2} \partial\Phi \cdot \partial\Phi - \frac{1}{2} |dC_0|^2 - 2\Lambda \right], \quad (2.6)$$

where the cosmological constant Λ is obtained by

$$\Lambda = \frac{(2\pi l_s)^5}{2} \int_{S^5} dx^5 \sqrt{-g_{S^5}} |F_5|^2 = -\frac{6}{R^2}. \quad (2.7)$$

Accordingly, the 5d solution for the metric obtained from (2.2) satisfying the equations of motion by varying action (2.6) is

$$ds_{(5\text{d})}^2 = e^{\frac{\phi}{2}} \left\{ \frac{r^2}{R^2} [-f(r) dt^2 + d\mathbf{x} \cdot d\mathbf{x}] + \frac{1}{f(r)} \frac{R^2}{r^2} dr^2 \right\}, \quad (2.8)$$

and the solution for Φ and C_0 remain respectively to be the corresponding formulas given in (2.2). Note that the action (2.6) describes nothing but the gravity-dilaton-axion system, and we will use the 5d effective background (2.8) in z coordinate as,

$$\begin{aligned} ds_{(5\text{d})}^2 &= e^{\frac{\phi}{2}} \frac{R^2}{z^2} \left\{ -f(z) dt^2 + d\mathbf{x} \cdot d\mathbf{x} + \frac{dz^2}{f(z)} \right\}, \\ f(z) &= 1 - \frac{z^4}{z_H^4}, \quad e^{\phi} = 1 - q \ln f(z), \\ z &= \frac{R^2}{r}, \quad z_H = \frac{R^2}{r_H}, \quad q = \frac{z_H^4 Q}{R^8}. \end{aligned} \quad (2.9)$$

in which the holographic boundary is located at $z \rightarrow 0$ to continue our discussion.

3 The fermionic correlation function

In this section, we consider a probe fermion in the bulk geometry and evaluate its two-point Green function in holography by employing the standard method in the AdS/CFT dictionary. In particular, we will focus on the influence of the instanton charge denoted by q in the fermionic correlation function.

3.1 General setup for the fermionic correlation function

Let us start with the principle of the AdS/CFT which illustrates that the generating functional of the dual conformal field theory (CFT) Z_{CFT} is equal to its gravitational partition function Z_{gravity} in the bulk geometry. Namely, for a spinor field ψ in the 5d bulk, we have

$$Z_{\text{CFT}} [\bar{\psi}_0, \psi_0] = Z_{\text{gravity}} [\bar{\psi}, \psi] \Big|_{\bar{\psi}, \psi \rightarrow \bar{\psi}_0, \psi_0}, \quad (3.1)$$

with

$$\begin{aligned}
Z_{\text{CFT}} [\bar{\psi}_0, \psi_0] &= \left\langle \exp \left\{ \int_{\partial\mathcal{M}} (\bar{\eta}\psi_0 + \bar{\psi}_0\eta) d^Dx \right\} \right\rangle, \\
Z_{\text{gravity}} [\bar{\psi}, \psi] &= e^{-S_{\text{gravity}}^{\text{ren}}}, \\
S_{\text{gravity}}^{\text{ren}} &= \int_{\mathcal{M}} \mathcal{L}_{\text{gravity}}^{\text{ren}} [\bar{\psi}, \psi] \sqrt{-g} d^5x
\end{aligned} \tag{3.2}$$

Note that, we use \mathcal{M} to denote the bulk space and its holographic boundary is given at $\partial\mathcal{M} = \{z \rightarrow 0\}$. ψ_0 refers to the boundary value of ψ as a source of the boundary fermionic operator η , $\mathcal{L}_{\text{gravity}}^{\text{ren}}$ denotes the renormalized Lagrangian of the bulk field ψ . Then follow the standard steps in quantum field theory (QFT), the one-point function i.e. average value of $\bar{\eta}$ is given as

$$\langle \bar{\eta} \rangle = \frac{1}{Z_{\text{CFT}}} \frac{\delta Z_{\text{CFT}}}{\delta \psi_0} = \frac{1}{Z_{\text{gravity}}} \frac{\delta Z_{\text{gravity}}}{\delta \psi_0} = -\frac{\delta S_{\text{gravity}}^{\text{ren}}}{\delta \psi_0} \equiv \Pi_0, \tag{3.3}$$

where we have used the relation (3.1). Therefore, the two-point correlation function G_R of η is obtained as,

$$\langle \bar{\eta}(\omega, \vec{k}) \rangle = G_R(\omega, \vec{k}) \psi_0, \tag{3.4}$$

i.e.

$$\Pi_0 = G_R(\omega, \vec{k}) \psi_0, \tag{3.5}$$

where ω, \vec{k} refers to the frequency and 3-momentum of the associated Fourier modes. Altogether, it is possible to evaluate ψ_0, Π_0 in order to evaluate the two-point correlation function G_R of η by using the classical gravity action $S_{\text{gravity}}^{\text{ren}}$ in holography.

3.2 Covariant Dirac equation for a probe fermion

Keep the formulas in Section 3.1 in hand, let us investigate a probe fermion in the 5d holographic background (2.9). In general, the 5d homogeneous background metric can be written as,

$$ds_{(5\text{d})}^2 = g_{tt} dt^2 + g_{xx} d\mathbf{x} \cdot d\mathbf{x} + g_{zz} dz^2, \tag{3.6}$$

where g_{tt}, g_{xx}, g_{zz} depend on z only. For a probe fermion in bulk, its dynamic is described by the Dirac action given as,

$$S_{\text{bulk}} = i \int d^5x \sqrt{-g} \bar{\psi} (\Gamma^M \nabla_M - m) \psi, \tag{3.7}$$

so the Dirac equation reads

$$(\Gamma^M \nabla_M - m) \psi = 0, \quad (3.8)$$

where the covariant derivative operator is given with the spin connection as,

$$\nabla_M = \partial_M + \frac{1}{4} \omega_{Mab} \gamma^{ab}, \gamma^{ab} = \frac{1}{2} [\gamma^a, \gamma^b]. \quad (3.9)$$

Imposing the homogeneous metric (3.6), the Dirac operator is computed as,

$$\Gamma^M \nabla_M = \frac{1}{\sqrt{-g_{tt}}} \gamma^0 \partial_0 + \frac{1}{\sqrt{g_{xx}}} \gamma^i \partial_i + \frac{1}{\sqrt{g_{zz}}} \gamma \partial_z + \frac{1}{4g_{xx}g_{tt}\sqrt{g_{zz}}} [3g_{tt}g'_{xx} + g_{xx}g'_{tt}] \gamma. \quad (3.10)$$

To find a solution for the Dirac equation, we impose the ansatz for spinor ψ in Dirac representation as,

$$\begin{aligned} \psi &= \begin{pmatrix} \psi_R \\ \psi_L \end{pmatrix} = (-gg^{zz})^{-1/4} \int \frac{d^4 p}{(2\pi)^4} e^{ik \cdot x} \chi(z, k), \\ \chi(z, k) &= \begin{bmatrix} \chi_R(z, k) \\ \chi_L(z, k) \end{bmatrix}, \chi_{R,L} = \begin{pmatrix} \chi_{R,L}^{(1)} \\ \chi_{R,L}^{(2)} \end{pmatrix}, k_\mu = (k_0, k_i), \end{aligned} \quad (3.11)$$

into the Dirac equation (3.8), so it is simplified as,

$$\left[\sqrt{\frac{g_{xx}}{g_{zz}}} (\gamma \partial_z - m \sqrt{g_{zz}}) + i K_\mu \gamma^\mu \right] \chi(z, k) = 0, \quad (3.12)$$

where

$$K_\mu = \left(\sqrt{-\frac{g_{xx}}{g_{tt}}} k_0, k_i \right). \quad (3.13)$$

3.3 Construction for the correlation function

When the background metric (3.6) is inserted into (3.12), the Dirac equation reduces to two decoupled differential equations for $\chi_{R,L}$ as,

$$\begin{aligned} (\partial_z + m \sqrt{g_{zz}}) \left[(K \cdot \sigma)^{-1} \sqrt{\frac{g_{xx}}{g_{zz}}} (\partial_z - m \sqrt{g_{zz}}) \chi_R \right] &= -\sqrt{\frac{g_{zz}}{g_{xx}}} (K \cdot \bar{\sigma}) \chi_R, \\ (\partial_z - m \sqrt{g_{zz}}) \left[(K \cdot \bar{\sigma})^{-1} \sqrt{\frac{g_{xx}}{g_{zz}}} (\partial_z + m \sqrt{g_{zz}}) \chi_L \right] &= -\sqrt{\frac{g_{zz}}{g_{xx}}} (K \cdot \sigma) \chi_L, \end{aligned} \quad (3.14)$$

which can be solved analytically near the boundary $z \rightarrow 0$ as,

$$\begin{aligned}
\chi_R &= Cz^{1-mR} + Dz^{mR}, \\
\chi_L &= Az^{-mR} + Bz^{1+mR}.
\end{aligned} \tag{3.15}$$

where A, B, C, D are constant Weyl spinor dependent on 4-momentum k and the charge of the D-instanton q . Hence for $mR > 0$, the boundary value χ_0 of χ is given by A which is the most divergent term, defined as

$$\chi_0 = \lim_{z \rightarrow 0} z^{mR} \chi = \begin{pmatrix} 0 \\ A \end{pmatrix}. \tag{3.16}$$

Then recall the Dirac action (3.7) with (3.11), it leads to

$$\begin{aligned}
S_{\text{bulk}} &= i \int d^4x dz \sqrt{-g} \bar{\psi} (\Gamma^M \nabla_M - m) \psi \\
&= i \int d^4x (\bar{\chi} \gamma \chi) \Big|_{z_H}^0 - i \int d^4x dz \sqrt{\frac{g_{zz}}{g_{xx}}} \left[\sqrt{\frac{g_{xx}}{g_{zz}}} (\partial_z \bar{\chi} \gamma + m \sqrt{g_{zz}} \bar{\chi}) + i K_\mu \bar{\chi} \gamma^\mu \right] \chi.
\end{aligned} \tag{3.17}$$

Note that the last term in (3.17) vanishes since it is nothing but the conjugate equation of (3.12). Therefore, we can obtain a boundary action S_{bdry} from (3.17) as,

$$\begin{aligned}
S_{\text{bdry}} &= i \int d^4x (\bar{\chi} \gamma \chi) \Big|_{z \rightarrow 0} \\
&= i \int d^4x \left(D^\dagger A + C^\dagger A z^{1-2m} \right) \Big|_{z \rightarrow 0},
\end{aligned} \tag{3.18}$$

which implies the holographic counterterm S_{CT} at boundary is,

$$S_{\text{CT}} = -i \int d^4x C^\dagger A z^{1-2m} \Big|_{z \rightarrow 0}, \tag{3.19}$$

giving the holographic renormalized action as,

$$\begin{aligned}
S_{\text{gravity}}^{\text{ren}} &= S_{\text{bdry}} + S_{\text{CT}} \\
&= i \int d^4x D^\dagger A.
\end{aligned} \tag{3.20}$$

Picking up (3.20) with (3.3) and (3.5), the correlation function satisfies

$$D = -G_R A, \tag{3.21}$$




Figure 1: The real and imaginary parts of zero-th order correlation functions $G_{R,\alpha}^{(0)}$ as functions of ω, k . The parameter is chosen as $q = 0, m = 0.01, z_H = 1$.

in the Weyl representation. This is not surprised since the boundary value ψ_0 of ψ is in fact a chiral spinor. On the other hand, the correlation function in the 2×2 matrix must be able to be written by the combination of the Pauli matrices as,

$$G_R = \alpha + \beta k_i \sigma^i, \quad (3.22)$$

we can set that the momentum is along x^3 direction as $k_\mu = (-\omega, 0, 0, k)$ due to the rotation symmetry in $\mathbb{R}^3 = \{x^i\}$, so that the correlation function G_R is expected to be diagonal, as

$$G_R = \begin{pmatrix} G_R^{1,1} & 0 \\ 0 & G_R^{2,2} \end{pmatrix}. \quad (3.23)$$

To finalize this section, we introduce the ratio defined as

$$G_\alpha = (-1)^{\alpha+1} \frac{\chi_R^{(\alpha)}}{\chi_L^{(\alpha)}}, \quad \alpha = 1, 2. \quad (3.24)$$




Figure 2: Density plots of the zero-th order correlation functions $G_{R,\alpha}^{(0)}$ with the parameters as $q = 0, m = 0.01, z_H = 1$. The white regions refer to the peaks in the correlation functions.

According to (3.16) - (3.21), the correlation function is given by

$$G_R^{\alpha,\alpha} = (-1)^\alpha \lim_{z \rightarrow 0} z^{-2m} G_\alpha, \quad (3.25)$$

and the equations for the ratios can be obtained by recombining the components of the spinor χ with respect to the Dirac equation (3.12). As a result, one can find the equations for the ratios are given as,

$$\sqrt{\frac{g_{xx}}{g_{zz}}} G'_\alpha = (-1)^\alpha u - k + [k + (-1)^\alpha u] G_\alpha^2 + 2m\sqrt{g_{xx}} G_\alpha, \quad (3.26)$$

where “ $'$ ” refers to the derivative with respect to z , and

$$u = \omega \sqrt{-\frac{g_{xx}}{g_{tt}}} = \frac{\omega}{\sqrt{f}}. \quad (3.27)$$

Therefore the correlation function could be evaluated numerically by solving (3.26) with the infalling boundary condition $G_\alpha|_{z=z_H} = (-1)^\alpha i$ obtained by analyzing its asymptotic behavior near $z = z_H$ as [35, 36, 37, 38, 39]. Furthermore, when the D-instanton is taken into account, the correlation function given in (3.25) should also depend on the charge density of the D-instanton denoted as q . In order to evaluate exactly the q -dependence in the correlation function, we consider its leading order correction by expand G_α as,

$$G_\alpha = \mathcal{G}_\alpha + q\xi_\alpha, \quad (3.28)$$

where \mathcal{G}_α is the zero-th order solution satisfying the equations presented in (3.26) with $q = 0$ as (there is no sum for the repetitive indices),

$$\sqrt{f}\mathcal{G}'_\alpha = (-1)^\alpha u - k + [k + (-1)^\alpha u] \mathcal{G}_\alpha^2 + \frac{2mR}{z} \mathcal{G}_\alpha. \quad (3.29)$$

And ξ_α is the leading order correction satisfying the equations

$$\sqrt{f}\xi'_\alpha = [k + (-1)^\alpha u] 2\mathcal{G}_\alpha\xi_\alpha + \frac{2mR}{z} \left[\xi_\alpha - \frac{1}{4}\mathcal{G}_\alpha \ln f \right]. \quad (3.30)$$

By analyzing the asymptotic behavior of (3.29), we find the infalling boundary condition for \mathcal{G}_α could be $\mathcal{G}_\alpha|_{z=z_H} = (-1)^\alpha i$ which is exactly same as the infalling boundary condition for G_α . Therefore, the boundary condition for ξ_α is expected as $\xi_\alpha|_{z=z_H} = 0$. Altogether, we will solve (3.29) and (3.30) numerically in order to evaluate the associated correlation function G_R as the fermionic spectral function in holography.

4 The numerical analysis

We write down the correlation functions presented in (3.25) as the combinations of its zero-th order $G_{R,\alpha}^{(0)}$ and first order solution $G_{R,\alpha}^{(1)}$ as,

$$\begin{aligned} G_R^{\alpha,\alpha} &= G_{R,\alpha}^{(0)} + qG_{R,\alpha}^{(1)}, \\ G_{R,\alpha}^{(0)} &= (-1)^\alpha \lim_{z \rightarrow 0} z^{-2m} \mathcal{G}_\alpha, \\ G_{R,\alpha}^{(1)} &= (-1)^\alpha \lim_{z \rightarrow 0} z^{-2m} \xi_\alpha, \end{aligned} \quad (4.1)$$

where $\mathcal{G}_\alpha, \xi_\alpha$ are coefficients independent on q and they can be solved numerically through the equations (3.29) and (3.30). The corresponding numerical results for $\mathcal{G}_\alpha, \xi_\alpha$ are given in Figure 1 - Figure 5.

Let us compare our numerical results with the calculations by using the method of hard




Figure 3: The real and imaginary parts of leading order correlation functions $G_{R,\alpha}^{(1)}$ as functions of ω, k with $m = 0.01, z_H = 1$.

thermal loop (HTL) in thermal QCD. The approach of HTL with vanished chemical potential illustrates the dispersion curves of fermions (plasmino) is ⁴[41],

$$\begin{aligned} \omega(k) &\simeq m_f \pm \frac{1}{3}k + \frac{1}{3m_f}k^2, \quad k \ll 1; \\ \omega(k) &\simeq k, \quad k \gg 1, \end{aligned} \tag{4.2}$$

where m_f is the effective mass generated by the medium effect of fermion as,

$$m_f = \sqrt{\frac{C_F}{8}g_{YM}T}. \tag{4.3}$$

Here g_{YM} refers to the Yang-Mills coupling constant and C_F is a constant suggested to be $C_F = 4/3$ for fundamental quarks or $C_F = 1$ for electron. Accordingly, we can see the dispersion curves

⁴We pick up the modes of positive frequency with $\omega > 0$ and momentum $k > 0$, the case for the negative frequency can be obtained by replacing $\omega \rightarrow -\omega$.




Figure 4: Density plots of leading order correlation functions $G_{R,\alpha}^{(1)}$ as functions of ω, k with $m = 0.01, z_H = 1$. The white regions refer to the peaks in the correlation functions

presented in zero-th order correlation function given in Figure 1 and Figure 2 cover basically the behavior of dispersion curves from HTL at large momentum. For the small momentum, while the zero-th order dispersion curves may be close to the relation given in (5), it does not illustrate two branches of the dispersion curves as some related works in holography [35, 36, 37, 38, 39]. The reason may be that, according to the gravity/fluid correspondence [42, 43], the plasma described holographically by metric (2.8) with $q = 0$ corresponds to conformally ideal fluid of $\mathcal{N} = 4$ super Yang-Mills theory which is different from QED or QCD plasma. In particular, the interaction of the plasmino or the dissipation of the medium is not taken into account in the ideal fluid. While this issue may be expected to be figured out by taking into account the dissipation of the fluid somehow in the holographic background (2.8), we will focus on the influence on the instanton in this work.

For the case $q > 0$, the correlation function has a correction term $G_{R,\alpha}^{(1)}$ given by ξ_α as it is




Figure 5: The real and imaginary parts of leading order correlation functions $G_{R,\alpha}^{(1)}$ as functions of ω with various k and $m = 0.01, z_H = 1$. There are always two peaks for various k in the correlation function which correspond to the two branches of the dispersion curves.

illustrated in Figure 3 - Figure 5. For $\omega > 0$ and $k > 0$, it is clear to see that the dispersion curves have two branches at small momentum and they trend to merge for large momentum. By taking a look at the value of the correction terms, we can find the correction $G_{R,\alpha}^{(1)}$ is large/small for small/large momentum respectively which has an opposite behavior to the zero-th correlation function $G_{R,\alpha}^{(0)}$. Altogether, for $q > 0$, the first order correction $G_{R,\alpha}^{(1)}$ dominates the behavior of the dispersion curves at small momentum and it always illustrates two branches. At large momentum, zero-th order function $G_{R,\alpha}^{(0)}$ dominates the behavior of the dispersion curves which displays only one branch of the dispersion curves on the light cone. Therefore, the dispersion curves given by the total spectral function $G_R^{\alpha,\alpha}$ are very close to the relation for HTL (4.2).

Besides, our numerical results also illustrate that, i.e. for $\omega, k > 0$, the effective mass $m_f = \omega$ ($k = 0$) splits into two values as it is displayed in Figure 4. This may relate remarkably to the properties of the fermionic spin, since it is known that there is spin-dependent interactions among fermions induced by instantons in QCD [1, 2]. According to the calculations by one-gluon exchange, the massive quark potential dependent on the quantum number of spin in the presence of instanton, so the total mass of the plasmino including quark interaction splits into two values which leads to two branches in the dispersion. In this sense, our numerical evaluation might be a holographic reproduction of instanton-induced interaction with spin. However, for

the massless case, the spin-dependent interaction induced by instantons trend to be vanished, so it could be why the spectral function of massless fermion is not affected by the instanton charge. On the other hand, some existing works about D-instanton in gauge-gravity duality [24, 28, 29, 30, 31, 32, 33, 35, 36] also support that the property of mass for a massless hadron in plasma is not affected by the instanton, which could thus be a holographic confirmation.

5 Summary

In this work, we study the correlation function of fermion in the D(-1)-D3 brane system which describes holographically the hot plasma with instantonic configuration. To simplify the setup, we briefly reduce the 10d supergravity background to an 5d background of gravity-dilaton-axion system. By probing a spinor in the 5d bulk, we derive the equations for the holographic correlation function from the Dirac equation, then demonstrate the numerical calculations with the infalling boundary condition. In particular, we expand the correlation function as zero-th order solution plus first order solution. Our numerical results illustrate that the first order correction to the total correlation function including two branches of the fermionic dispersion curves is dominated for small momentum, while the zero-th order correlation function dominates the behavior of the fermionic dispersion curves at large momentum giving only one merged dispersion curves on light cone. This behavior is out of reach without the presence of the instanton and it is very close to the fermionic dispersion curves obtained by HTL. Furthermore, we also find the effective mass generated by the medium effect of fermion splits into two values due to the spin-dependent interactions among fermions induced by instantons which has been studied in the framework of QCD with instanton [1, 2]. Altogether, it means seemingly the instantonic configuration in the plasma is very influential for the spectral function of the plasmino.

Acknowledgements

We would like to thank Prof.Lin Shu for helpful discussion. This work is supported by the National Natural Science Foundation of China (NSFC) under Grant No. 12005033 and the Fundamental Research Funds for the Central Universities under Grant No. 3132024192.

References

- [1] T.Schäfer, E. Shuryak, “Instantons in QCD”, *Rev.Mod.Phys.* 70 (1998) 323-426, arXiv:hep-ph/9610451.
- [2] D. Gross, R. Pisarski, L. Yaffe, “QCD and Instantons at Finite Temperature”, *Rev.Mod.Phys.* 53 (1981) 43.

- [3] E. Vicari, H. Panagopoulos, “Theta dependence of SU(N) gauge theories in the presence of a topological term”, *Phys.Rept.* 470 (2009) 93-150, arXiv:0803.1593.
- [4] D. Tong, “TASI lectures on solitons: Instantons, monopoles, vortices and kinks”, arXiv:hep-th/0509216.
- [5] D. Diakonov, N. Gromov, V. Petrov, S. Slizovskiy, “Quantum weights of dyons and of instantons with nontrivial holonomy”, *Phys.Rev.D* 70 (2004) 036003, arXiv:hep-th/0404042.
- [6] D. Diakonov, V. Petrov, “Confining ensemble of dyons”, *Phys.Rev.D* 76 (2007) 056001, arXiv:0704.3181.
- [7] Y. Liu, E. Shuryak, I. Zahed, “Confining dyon-antidyon Coulomb liquid model. I.”, *Phys.Rev.D* 92 (2015) 8, 085006, arXiv:1503.03058.
- [8] Y. Liu, E. Shuryak, I. Zahed, “Light quarks in the screened dyon-antidyon Coulomb liquid model. II.”, *Phys.Rev.D* 92 (2015) 8, 085007, arXiv:1503.09148.
- [9] Y. Liu, E. Shuryak, I. Zahed, “The Instanton-Dyon Liquid Model III: Finite Chemical Potential”, *Phys.Rev.D* 94 (2016) 10, 105011, arXiv:1606.07009.
- [10] Y. Liu, E. Shuryak, I. Zahed, “Light Adjoint Quarks in the Instanton-Dyon Liquid Model IV”, *Phys.Rev.D* 94 (2016) 10, 105012, arXiv:1605.07584.
- [11] Y. Liu, E. Shuryak, I. Zahed, “The Instanton-Dyon Liquid Model V: Twisted Light Quarks”, *Phys.Rev.D* 94 (2016) 10, 105013, arXiv:1606.02996.
- [12] E. Poppitz, T. Schäfer, M. Unsal, “Continuity, Deconfinement, and (Super) Yang-Mills Theory”, *JHEP* 10 (2012) 115, arXiv:1205.0290.
- [13] E. Poppitz, T. Schäfer, M. Ünsal, “Universal mechanism of (semi-classical) deconfinement and theta-dependence for all simple groups”, *JHEP* 03 (2013) 087, arXiv:1212.1238.
- [14] J. Maldacena, “The Large N limit of superconformal field theories and supergravity”, *Adv.Theor.Math.Phys.* 2 (1998) 231-252, arXiv:hep-th/9711200.
- [15] O. Aharony, S. Gubser, J. Maldacena, H. Ooguri, Y. Oz, “Large N field theories, string theory and gravity”, *Phys.Rept.* 323 (2000) 183-386, arXiv:hep-th/9905111.
- [16] E. Witten, “Anti-de Sitter space and holography”, *Adv.Theor.Math.Phys.* 2 (1998) 253-291, arXiv:hep-th/9802150.
- [17] H. Liu, A. Tseytlin, “D3-brane D instanton configuration and N=4 superYM theory in constant selfdual background”, *Nucl.Phys.B* 553 (1999) 231-249, arXiv:hep-th/9903091.

- [18] J. Barbon, A. Pasquinucci, “Aspects of instanton dynamics in AdS / CFT duality”, *Phys.Lett.B* 458 (1999) 288-296, arXiv:hep-th/9904190.
- [19] L. Bartolini, Fr. Bigazzi, S. Bolognesi, A. Cotrone, A. Manenti, “Theta dependence in Holographic QCD”, *JHEP* 02 (2017) 029, arXiv:1611.00048.
- [20] S. Seki, S. Sin, “A New Model of Holographic QCD and Chiral Condensate in Dense Matter”, *JHEP* 10 (2013) 223, arXiv:1304.7097.
- [21] C. Wu, Z. Xiao, D. Zhou, “Sakai-Sugimoto model in D0-D4 background”, *Phys.Rev.D* 88 (2013) 2, 026016.
- [22] S. Li, “A holographic description of theta-dependent Yang-Mills theory at finite temperature”, *Chin.Phys.C* 44 (2020) 1, 013103, arXiv:1907.10277.
- [23] S. Li, T. Jia, “Matrix model and Holographic Baryons in the D0-D4 background”, *Phys.Rev.D* 92 (2015) 4, 046007, arXiv:1506.00068.
- [24] S. Li, “Holographic heavy-baryons in the Witten-Sakai-Sugimoto model with the D0-D4 background”, *Phys.Rev.D* 96 (2017) 10, 106018, arXiv:1707.06439.
- [25] S. Li, S. Lin, “D-instantons in Real Time Dynamics”, *Phys.Rev.D* 98 (2018) 6, 066002, arXiv:1711.06365.
- [26] S. Li, S. Luo, M. Tan, “Three-dimensional Yang-Mills-Chern-Simons theory from a D3-brane background with D-instantons”, *Phys.Rev.D* 104 (2021) 6, 066008, arXiv:2106.04038.
- [27] R. Argurio, A. Armoni, M. Bertolini, F. Mignosa, P. Niro, “Vacuum structure of large N QCD₃ from holography”, *JHEP* 07 (2020) 134, arXiv:2006.01755.
- [28] B. Gwak, M. Kim, B. Lee, Y. Seo, S. Sin, “Holographic D Instanton Liquid and chiral transition”, *Phys.Rev.D* 86 (2012) 026010, arXiv:1203.4883.
- [29] Z. Zhang, D. Hou, G. Chen, “Heavy quark potential and jet quenching parameter in a D-instanton background”, *Eur.Phys.J.A* 52 (2016) 12, 357, arXiv:1607.03985
- [30] Z. Zhang, D. Hou, G. Chen, “Imaginary potential of moving quarkonia in a D-instanton background”, *J.Phys.G* 44 (2017) 11, 115001, arXiv:1710.06579.
- [31] J. Chen, D. Hou, “Heavy quark potential and jet quenching parameter in a rotating D-instanton background”, *Eur.Phys.J.C* 84 (2024) 4, 447, arXiv:2202.00888.
- [32] S. Li, “Holographic Schwinger effect in the confining background with D-instanton”, *Eur.Phys.J.C* 81 (2021) 9, 797, arXiv:2005.11955.

- [33] L. Shahkarami, M. Dehghani, P. Dehghani, “Holographic Schwinger Effect in a D-Instanton Background”, *Phys.Rev.D* 97 (2018) 4, 046013, arXiv:1511.07986.
- [34] W. Cai, K. Li, S. Li, “Electromagnetic instability and Schwinger effect in the Witten–Sakai–Sugimoto model with D0–D4 background”, *Eur.Phys.J.C* 79 (2019) 11, 904, arXiv:1612.07087.
- [35] S. Li, H. Li, Y. Zhang, “The worldvolume fermion as baryon in holographic QCD with a theta angle”, arXiv:2402.01197.
- [36] S. Li, Y. Zhang, H. Li, “Correlation function of flavored fermion in holographic QCD”, *Phys.Rev.D* 109 (2024) 8, 086020, arXiv:2307.13357.
- [37] H. Liu, J. McGreevy, D. Vegh, “Non-Fermi liquids from holography”, *Phys.Rev.D* 83 (2011) 065029, arXiv:0903.2477.
- [38] Y. Seo, S. Sin, Y. Zhou, “Thermal Mass and Plasmino for Strongly Interacting Fermions”, *JHEP* 06 (2013) 076, arXiv:1305.1446.
- [39] Y. Seo, S. Sin, Y. Zhou, “Self-energy of Strongly Interacting Fermions in Medium: a Holographic Approach”, *Phys.Lett.B* 723 (2013) 207-212, arXiv:1205.3377.
- [40] L. Fang, X. Ge, Ji. Wu, H. Leng, “Anisotropic Fermi surface from holography”, *Phys.Rev.D* 91 (2015) 12, 126009, arXiv:1409.6062.
- [41] V. Klimov, “Spectrum of Elementary Fermi Excitations in Quark Gluon Plasma”, *Sov.J.Nucl.Phys.* 33 (1981) 934-935, *Yad.Fiz.* 33 (1981) 1734-1736.
- [42] S. Bhattacharyya, V. Hubeny, S. Minwalla, M. Rangamani, “Nonlinear Fluid Dynamics from Gravity”, *JHEP* 02 (2008) 045, arXiv:0712.2456.
- [43] S. Bhattacharyya, R. Loganayagam, S. Minwalla, S. Nampuri, S. Trivedi, Sp. Wadia, “Forced Fluid Dynamics from Gravity”, *JHEP* 02 (2009) 018, arXiv:0806.0006.