
BAYESIAN OUTCOME WEIGHTED LEARNING

A PREPRINT

✉ **Sophia Yazzourh**

Institut de Mathématiques de Toulouse
UMR5219 - Université de Toulouse
CNRS - UPS IMT
F-31062 Toulouse Cedex 9, France
sophia.yazzourh@math.univ-toulouse.fr

✉ **Nikki L. B. Freeman**

Department of Biostatistics and Bioinformatics
Duke Clinical Research Institute
Duke University
Durham, North Carolina 27701 USA
nikki.freeman@duke.edu

ABSTRACT

One of the primary goals of statistical precision medicine is to learn optimal individualized treatment rules (ITRs). The classification-based, or machine learning-based, approach to estimating optimal ITRs was first introduced in outcome-weighted learning (OWL). OWL recasts the optimal ITR learning problem into a weighted classification problem, which can be solved using machine learning methods, e.g., support vector machines. In this paper, we introduce a Bayesian formulation of OWL. Starting from the OWL objective function, we generate a pseudo-likelihood which can be expressed as a scale mixture of normal distributions. A Gibbs sampling algorithm is developed to sample the posterior distribution of the parameters. In addition to providing a strategy for learning an optimal ITR, Bayesian OWL provides a natural, probabilistic approach to estimate uncertainty in ITR treatment recommendations themselves. We demonstrate the performance of our method through several simulation studies.

Keywords Precision medicine · Individualized treatment rule · Bayesian machine learning

1 Introduction

The task of statistical precision medicine is to learn from data how to match patients to treatments with the aim of improving health outcomes [Kosorok and Laber, 2019]. One way to operationalize this goal is through individualized treatment regimes (ITRs), functions that map from patient characteristics to treatment recommendations. Ideally, we would like to learn ITRs that if followed in practice would lead to better outcomes on average in the target population than if another treatment strategy was used, e.g., a one-size-fits-all approach. These ITRs are called optimal ITRs [Kosorok and Laber, 2019, Murphy, 2003].

In the language of reinforcement learning, we focus on the "batch, off-policy" setting. By "batch" we mean that data have been previously collected and no new data will be received, and by "off policy" we mean that the strategy for assigning treatments in the observed data (e.g., through randomization as in a clinical trial) may not be the optimal strategy (or alternatively, regime or policy) [Sutton and Barto, 2018]. Within this setting, a large number of methods and approaches have been developed to learn such ITRs from data.

Some approaches estimate the expected value of the outcome we would expect under a particular ITR without any parametric assumptions. Then, the optimal ITR may be learned by searching over a class of ITRs. Examples of this general strategy include those proposed by Orellana et al. [2010], Robins et al. [2008], and Zhang et al. [2013].

Another class of approaches, sometimes referred to as indirect methods, model the mean of the outcome conditional on treatment and covariates. For a multi-stage ITR, sometimes called a dynamic treatment regime, this entails specifying a conditional mean model for each stage. Through these estimated conditional means, optimal ITRs can be deduced. One of the most popular regression-based frameworks for learning optimal ITRs is Q-learning [Watkins, 1989, Murphy, 2005]. Q-learning has been used to learn optimal ITRs in many settings, including clinical trial data [Schulte et al., 2014], observational data [Moodie et al., 2012], and in the presence of censoring for time-to-event data Goldberg and Kosorok [2012].

Machine learning or classification-based optimal ITR learning approaches convert the optimal ITR learning problem into the classification framework by which machine learning methods can be employed. Zhao et al. [2012] introduced outcome weighted learning (OWL) which leverages a simple value function estimator and the Radon-Nikodym theorem to rewrite the value function as a weighted classification problem. Consequently, learning the ITR that optimizes the value function can be solved as minimizing the classification loss function. Since its introduction, a number of extensions to OWL have been made including backwards outcome weighted learning (BOWL) and simultaneous outcome weighted learning (SOWL) for learning optimal multi-stage treatment regimes [Zhao et al., 2015], residual weighted learning (RWL) [Zhou et al., 2017] and augmented outcome-weighted learning (AOL) [Liu et al., 2018] which improve the finite sample properties of OWL, robust outcome weighted learning (ROWL) which uses an angle-based classification approach [Fu et al., 2019], and efficient augmentation and relaxation learning (EARL) which employs both a propensity model and outcome model and has the double robustness property Zhao et al. [2019].

Finally, a few Bayesian approaches for learning optimal ITRs have also been proposed. The Bayesian machine learning (BML) approach was introduced by Murray et al. [2018]. It employs Bayesian modeling within a framework that closely aligns with Q-learning by modeling the outcomes at each stage. Likelihood-based approaches, strategies that model both the distribution of the final outcome and the intermediate outcomes, have also been proposed within the Bayesian framework [Thall et al., 2002, 2007, Arjas and Saarela, 2010, Zajonc, 2012, Xu et al., 2016, Yu and Bondell, 2023].

The focus of this paper will be on the machine learning, or classification-based method, for learning optimal ITRs. Although classification-based approaches are powerful and avoid estimating models that are not the target of the analysis itself, there are limitations. For example, many machine learning methods for classification do not naturally quantify uncertainty, e.g., quantification of the uncertainty of a particular prediction. While this may be acceptable in some cases, being unable to quantify uncertainty is a serious gap when generating evidence for health care decision-making. Moreover, machine learning analyses are often evaluated in terms of predictive power which does not necessarily translate into inferential capability.

In this paper, we present a Bayesian approach to OWL. To our knowledge, this is the first Bayesian optimal ITR learning strategy to directly learn optimal ITRs. Using a construction similar to Polson and Scott [2011], we construct a pseudo-likelihood from the weighted classification loss function. Once transformed from an optimization-based framework to a probabilistic framework, our method generates an entire posterior distribution that can be used for inference and, most powerfully, for uncertainty quantification of the treatment recommendations themselves. Our main contributions are as follows:

1. We propose a Bayesian approach to learning optimal ITRs that leverages the classification-based framework and avoids modeling the outcome or nuisance conditional mean models.
2. We propose a simple Gibbs sampling algorithm for learning such an optimal ITR.
3. We demonstrate how to use our resulting pseudo-posterior distribution to quantify uncertainty in the treatment recommendations.

In Section 2, we set the notation and review OWL and Bayesian support vector machines. In Section 3 we construct the probabilistic formulation of the OWL classification problem, derive a Gibbs sampling algorithm for estimation, and detail our approach to uncertainty quantification. In Section 4 we demonstrate the performance of our approach through simulation studies. We conclude in Section 5 with a discussion of our results and future work.

2 Background

2.1 Setting

We let $A \in \mathcal{A} = \{-1, 1\}$ denote the action, or treatment, and assume that observed treatments are assigned randomly as in a clinical trial with $P(A = 1) = \rho$ known. Let $X_i = (X_{i,1}, \dots, X_{i,p})^\top \in \mathcal{X}$ denote the p -dimensional biomarker and prognostic information vector, and let R denote the outcome (bigger is better). We further assume that the reward can be rescaled so that $R > 0$. Then, the observed data is iid replicates of (A_i, X_i, R_i) for $i = 1, \dots, n$.

An ITR is a function d that maps from \mathcal{X} to a recommended treatment in \mathcal{A} . For a given ITR d , the value of d is $V(d) = \mathbb{E}[R(d)]$, where $R(d)$ is the reward we would observe if treatments were allocated according to rule d . An optimal ITR d^{opt} satisfies $V(d^{\text{opt}}) \geq V(d)$ for all $d \in \mathcal{D}$, where \mathcal{D} is a class of ITRs. Our goal is to learn an optimal ITR d^{opt} . Under the assumptions of causal consistency, the stable unit treatment value assumption, no unmeasured confounding, and positivity, $V(d)$ can be identified from the observed data and $V(d) = \mathbb{E}\{\max_{A \in \mathcal{A}} \mathbb{E}[R|A = d(\mathbf{x}), X = \mathbf{x}]\}$.

2.2 Outcome weighted learning

If we let P denote the distribution of (X, A, R) , and P^d denote the distribution of (X, A, R) when $A = d(X)$, then the reward we would expect if ITR $d(X)$ were followed is given by

$$\mathbb{E}^d(R) = \int R dP^d = \int R \frac{dP^d}{dP} dP = \mathbb{E} \left[\frac{\mathbb{1}(A = d(X))}{A\rho + (1-A)/2} R \right]. \quad (1)$$

Zhao et al. [2012] showed that maximizing Equation (1) is equivalent to a weighted classification problem and thereby solvable using techniques from machine learning. Specifically, they proposed OWL, a strategy for learning an optimal ITR using a convex surrogate loss function in place of the zero-one loss function and strategies from support vector machines. OWL minimizes the objective function

$$Q_n^{\text{OWL}}(\beta) = \frac{1}{n} \sum_{i=1}^n \frac{R_i}{A_i\rho + (1-A_i)/2} (1 - A_i h(X_i, \beta))_+ \quad (2)$$

where $(z)_+ = \max(z, 0)$ denotes the hinge loss function and $h(\cdot)$ is the ITR parameterized by β . Song et al. [2015] introduced a penalized variant of OWL that included a regularization term $p_\lambda(\beta)$ for the ITR parameters. POWL minimizes the objective function

$$Q_n^{\text{POWL}}(\beta) = \frac{1}{n} \sum_{i=1}^n \frac{R_i}{A_i\rho + (1-A_i)/2} (1 - A_i h(X_i, \beta))_+ + \sum_{j=1}^p p_\lambda(|\beta_j|) \quad (3)$$

where $p_\lambda(\beta)$ is a penalty function and λ is a tuning parameter.

2.3 Bayesian support vector machines

Although the pure machine learning framework is powerful, it is limited in its ability to capture and model uncertainty as in a statistical framework. Polson and Scott [2011] bridged this gap between pure machine learning and statistical modeling for SVMs by showing how to cast SVM into a Bayesian framework. They considered the L^α -norm regularized support vector classifier that chooses β to minimize

$$d_\alpha(\beta, \nu) = \sum_{i=1}^n \max(1 - r_i \mathbf{x}_i^\top \beta, 0) + \nu^{-\alpha} \sum_{j=1}^k |\beta_j / \sigma_j|^\alpha \quad (4)$$

where σ_j is the standard deviation of the j -th element of \mathbf{x} and ν is a tuning parameter. For this objective function, the learned classifier is a linear classifier. Polson and Scott [2011] shows that minimizing Equation (4) is equivalent to finding the mode of the pseudo-posterior distribution $p(\beta|\nu, \alpha, y)$

$$p(\beta|\nu, \alpha, r) \propto \exp(-d_\alpha(\beta, \nu)) \propto C_\alpha(\nu) L(r|\beta) p(\beta|\nu, \alpha) \quad (5)$$

where C_α is a pseudo-posterior normalization constant. Thus, the data dependent factor $L(y|\beta)$ is a pseudo-likelihood

$$L(r|\beta) = \prod_i L_i(r_i|\beta) = \exp \left\{ x - 2 \sum_{i=1}^n \max(1 - r_i \mathbf{x}_i^\top \beta, 0) \right\}. \quad (6)$$

The main theoretical result from Polson and Scott [2011] is that the pseudo-likelihood contribution $L_i(r_i|\beta)$ is a location-scale mixture of normals (Polson and Scott [2011], Theorem 1).

3 Our approach

We follow the strategy employed by Polson and Scott [2011] to cast the OWL objective function into a probabilistic Bayesian learning framework. The conversion is not one-to-one since Polson and Scott [2011] constructed a Bayesian model for a standard SVM whereas the objective function for OWL Equation (2) is a weighted SVM problem. We first employ a non-penalty prior to mimic original formulation of OWL as in Zhao et al. [2012], and later we demonstrate the inclusion of a penalty prior on the the ITR coefficients. Minimizing Equation (2) is equivalent to finding the mode of the pseudo-posterior which we can write as

$$\begin{aligned} p(\mathbf{x}|a_i, \nu, \alpha) &\propto \exp(-Q_n(\beta, \nu, \alpha)) \\ &\propto \exp \left\{ \sum_{n=1}^n \frac{r_i}{a_i\rho + (1-A_i)/2} (1 - a_i h(\mathbf{x}_i, \beta))_+ \right\} \prod_{j=1}^p p(\beta_j|\mu_0, \sigma_0^2) \\ &\propto C(\nu, \alpha) L(a|\beta) p(\beta|\mu_0, \sigma_0^2). \end{aligned} \quad (7)$$

Throughout, we will assume $R > 0$. When this is not the case, a distance-preserving transformation of R from \mathbb{R} to \mathbb{R}^+ can be used. Assuming that h is linear, i.e., $h(\mathbf{x}_i, \boldsymbol{\beta}) = \mathbf{x}_i^\top \boldsymbol{\beta}$ and following the strategy taken in Theorem 1 of Polson and Scott [2011], the contribution of a single observation to the pseudo-likelihood is given by

$$\begin{aligned} L_i(a_i|r_i, \mathbf{x}_i, \boldsymbol{\beta}) &= \exp \left\{ -2 \frac{r_i}{a_i \rho + (1 - a_i)/2} \max(1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta}, 0) \right\} \\ &= \mathbb{1}(a_i = 1) \int_0^\infty \frac{1}{\sqrt{2\pi\lambda_i}} \exp \left\{ -\frac{1}{2\lambda_i} \left(\frac{r_i}{\rho} + \lambda_i - \frac{r_i}{\rho} a_i \mathbf{x}_i^\top \boldsymbol{\beta} \right)^2 \right\} d\lambda_i \\ &\quad + \mathbb{1}(a_i = -1) \int_0^\infty \frac{1}{\sqrt{2\pi\lambda_i}} \exp \left\{ -\frac{1}{2\lambda_i} \left(\frac{r_i}{1-\rho} + \lambda_i - \frac{r_i}{1-\rho} a_i \mathbf{x}_i^\top \boldsymbol{\beta} \right)^2 \right\} d\lambda_i, \end{aligned} \quad (8)$$

or in other words that $L_i(a_i, \lambda_i|r_i, \mathbf{x}_i, \boldsymbol{\beta})$ is a scale mixture of Gaussians (full details are contained in Section 6.1).

3.1 Prior specification for the ITR parameters

In their formulation of Bayesian SVM, Polson and Scott [2011] use the exponential power prior for $\boldsymbol{\beta}$, a prior that can be show to be equivalent to L1-regularization of the regression parameters. Regularization of the OWL parameters have been explored as in Song et al. [2015]. In this paper, we first construct our method as an analogy to the original formulation of OWL without penalization. We make this choice because (1) our primary aim is to develop a Bayesian classification-based ITR learning approach, and because (2) L1-regularization does not necessarily yield sparse rules (see the discussion in Section 4.1 of Polson and Scott [2011]). However, regularization help avoid overfitting, a common problem in machine learning. Thus, we also explore penalty priors for $\boldsymbol{\beta}$, including the exponential power prior distribution and the spike-and-slab prior distribution.

3.1.1 Normal prior distribution for $\boldsymbol{\beta}$

We first consider the case with normal distribution priors on the treatment rule parameters $\beta_j \sim N(\mu_0, \sigma_0^2)$ for $j = 1, \dots, p$ where μ_0 and σ_0^2 are hyperparameters. With the pseudo-likelihood and a suitable prior for the ITR parameters defined, we can write the pseudo-posterior distribution as

$$\begin{aligned} p(\boldsymbol{\beta}, \boldsymbol{\lambda}|\mathbf{x}, \mathbf{r}, \mathbf{a}, \alpha, \nu) &\propto \prod_{\{i:a_i=1\}}^n \lambda_i^{-1/2} \cdot \exp \left\{ -\frac{1}{2} \sum_{i:a_i=1}^n \frac{\left(\frac{r_i}{\rho} + \lambda_i - \frac{r_i}{\rho} a_i \mathbf{x}_i^\top \boldsymbol{\beta} \right)^2}{\lambda_i} \right\} \\ &\quad \times \prod_{\{i:a_i=-1\}}^n \lambda_i^{-1/2} \cdot \exp \left\{ -\frac{1}{2} \sum_{i:a_i=-1}^n \frac{\left(\frac{r_i}{1-\rho} + \lambda_i - \frac{r_i}{1-\rho} a_i \mathbf{x}_i^\top \boldsymbol{\beta} \right)^2}{\lambda_i} \right\} \\ &\quad \times \prod_{j=1}^p \frac{1}{\sqrt{2\pi\sigma_0^2}} \exp \left\{ -\frac{1}{2} \frac{(\beta_j - \mu_{0,j})^2}{\sigma_0^2} \right\} \end{aligned} \quad (9)$$

where $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_n)^\top$, $\mathbf{r} = (r_1, \dots, r_n)^\top$, and $\mathbf{a} = (a_1, \dots, a_n)^\top$.

3.2 Exponential power prior distribution for $\boldsymbol{\beta}$

Rather than use normal priors for the coefficients of the rule, Polson and Scott [2011] employed an exponential power prior on $\boldsymbol{\beta}$. This prior contains the regularization penalty, and from Theorem 2 of Polson and Scott [2011], the double exponential prior regularization penalty can be written as

$$p(\beta_j|\nu, \alpha = 1) = \int_0^\infty \phi(\beta_j|0, \nu^2 \omega_j \sigma_j^2) \frac{1}{2} e^{-\frac{\omega_j}{2}} d\omega_j \quad (10)$$

where $p(\omega_j|\alpha) \propto \omega_j^{-\frac{3}{2}} St_{\alpha/2}^+(\omega_j^{-1})$ and $St_{\alpha/2}^+$ is the density function of a positive stable random variable of index $\alpha/2$. In particular, when $\alpha = 1$, $p(\omega_j|\alpha) \sim Exponential(2)$ (Corollary 1 of Polson and Scott [2011]).

Under this prior distribution specification, we can write the pseudo-posterior distribution as

$$\begin{aligned}
 p(\boldsymbol{\beta}, \boldsymbol{\lambda}, \boldsymbol{\omega} | \mathbf{x}, \mathbf{r}, \mathbf{a}, \alpha, \nu) &\propto \prod_{\{i:a_i=1\}}^n \lambda_i^{-1/2} \cdot \exp \left\{ -\frac{1}{2} \sum_{i:a_i=1}^n \frac{\left(\frac{r_i}{\rho} + \lambda_i - \frac{r_i}{\rho} a_i \mathbf{x}_i^\top \boldsymbol{\beta} \right)^2}{\lambda_i} \right\} \\
 &\times \prod_{\{i:a_i=-1\}}^n \lambda_i^{-1/2} \cdot \exp \left\{ -\frac{1}{2} \sum_{i:a_i=-1}^n \frac{\left(\frac{r_i}{1-\rho} + \lambda_i - \frac{r_i}{1-\rho} a_i \mathbf{x}_i^\top \boldsymbol{\beta} \right)^2}{\lambda_i} \right\} \\
 &\times \prod_{j=1}^p \omega_j^{-\frac{1}{2}} \cdot \exp \left\{ -\frac{1}{2\nu^2} \sum_{j=1}^p \frac{\beta_j^2}{\sigma_j^2 \omega_j} \right\} \cdot \prod_{j=1}^p p(\omega_j | \alpha). \tag{11}
 \end{aligned}$$

where $\boldsymbol{\omega} = (\omega_1, \dots, \omega_p)^\top$.

3.3 Spike-and-slab prior distribution for $\boldsymbol{\beta}$

Polson and Scott [2011] also explored the use of a spike-and-slab prior for $\boldsymbol{\beta}$. The spike-and-slab prior is a Bayesian approach used for variable selection. It combines a "spike" component, which is a Dirac delta function at zero, to induce sparsity by shrinking some coefficients exactly to zero, and a "slab" component that allows other coefficients to vary freely [Mitchell and Beauchamp, 1988, George and McCulloch, 1993]. Thus, the spike-and-slab prior on the j th coefficient β_j can be written as

$$p(\beta_j | \gamma_j, \nu^2) = \gamma_j N(0, \nu^2 \sigma_j^2) + (1 - \gamma_j) \delta_0(\beta_j) \tag{12}$$

where $\delta_0(\cdot)$ is the Dirac measure (point mass at 0). The prior on γ_j is given by

$$p(\gamma_j | \pi) = \pi^{\gamma_j} (1 - \pi)^{1 - \gamma_j}. \tag{13}$$

Letting \odot denote elementwise multiplication, i.e., where $(a_1, \dots, a_n) \odot (b_1, \dots, b_n) = (a_1 b_1, \dots, a_n b_n)$, the full pseudo-posterior when a spike-and-slab prior distribution is specified for $\boldsymbol{\beta}$ can be written as

$$\begin{aligned}
 p(\boldsymbol{\beta}, \boldsymbol{\lambda}, \boldsymbol{\gamma} | \mathbf{y}, \mathbf{X}, \pi, \nu) &= \prod_{i=1}^n p(y_i | \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\lambda}) \prod_{j=1}^p [p(\beta_j | \gamma_j, \nu^2) p(\gamma_j | \pi)] \\
 &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi\lambda_i}} \exp \left\{ \frac{1}{2} \frac{(1 + \lambda_i - y_i x_i^\top (\boldsymbol{\gamma} \odot \boldsymbol{\beta}))^2}{\lambda_i} \right\} \\
 &\times \prod_{j=1}^p [(\gamma_j N(0, \nu^2 \sigma_j^2) + (1 - \gamma_j) \delta_0(\beta_j)) \pi^{\gamma_j} (1 - \pi)^{1 - \gamma_j}]. \tag{14}
 \end{aligned}$$

3.4 Estimation

To draw from the pseudo-posterior distribution, Polson and Scott [2011] employed two algorithms, an expectation-minimization (EM) approach and a Gibbs sampling approach. The approach we take is the latter. Although sampling the pseudo-posterior is likely to be more time intensive than estimation via the EM algorithm, the rationale for a fully Bayesian approach is to enable uncertainty quantification (Section 3.5).

3.4.1 Normal prior distribution for $\boldsymbol{\beta}$

The full pseudo-posterior distribution under normal priors for $\boldsymbol{\beta}$ Equation (9) has two unknown parameters, $\boldsymbol{\beta}$ and $\boldsymbol{\lambda}$. To sample these parameters, we derive a Gibbs sampling algorithm, which entails sequentially sampling each parameter conditionally on the most up-to-date values of the other parameters. We give a high level summary of the derivation in this section and full details in Section 6.2. The conditional distribution of $\lambda_i | \boldsymbol{\beta}, \mathbf{x}_i, \mathbf{a}_i, r_i$ (up to a normalizing constant) can be written as:

$$p(\lambda_i | \boldsymbol{\beta}, \mathbf{x}_i, \mathbf{a}_i, r_i) \propto \mathbb{1}(a_i = -1) \lambda_i^{-1/2} \cdot \exp \left\{ -\frac{1}{2} \left(\lambda_i + \left(\frac{r_i}{1-\rho} \right)^2 (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta}) \lambda_i^{-1} \right) \right\}.$$

From Devroye [1986], page 479, a random variable has the generalized inverse Gaussian distribution $\mathcal{GIG}(\gamma, \psi, \chi)$ if its density function is $p(x|\gamma, \psi, \chi) = C(\gamma, \psi, \chi)x^{\gamma-1} \exp\{-\frac{1}{2}(\frac{\chi}{x} + \psi x)\}$, where $C(\gamma, \psi, \chi)$ is a normalization constant. Thus

$$p(\lambda_i|\boldsymbol{\beta}, \mathbf{x}_i, a_i, r_i) \sim \mathbb{1}(a_i = 1)\mathcal{GIG}\left(\frac{1}{2}, 1, \left(\frac{r_i}{\rho}\right)^2 (1 - a_i\mathbf{x}_i^\top\boldsymbol{\beta})^2\right) \\ + \mathbb{1}(a_i = -1)\mathcal{GIG}\left(\frac{1}{2}, 1, \left(\frac{r_i}{1-\rho}\right)^2 (1 - a_i\mathbf{x}_i^\top\boldsymbol{\beta})^2\right). \quad (15)$$

The conditional distribution of $\boldsymbol{\beta}|\boldsymbol{\lambda}, \boldsymbol{\omega}, \mathbf{r}, \mathbf{a}, \mathbf{x}$ follows from standard arguments for Bayesian linear models. The notable difference from such a standard model is that $\boldsymbol{\beta}|\boldsymbol{\lambda}, \boldsymbol{\omega}, \mathbf{r}, \mathbf{a}, \mathbf{x}$ is a mixture over two distributions, one for when the observed treatment in the data under analysis is 1 and one for when the observed treatment is -1 . The conditional distribution of $\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{r}, \mathbf{a}, \mathbf{x}$ has the form

$p(\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{r}, \mathbf{a}, \mathbf{x})$

$$\propto \exp\left\{-\frac{1}{2}\sum_{\{i:a_i=1\}}\left(-2\frac{r_i}{\rho}a_i\mathbf{x}_i^\top\boldsymbol{\beta}\left(1+\frac{r_i}{\rho\lambda_i}\right)+\left(\frac{r_i}{\rho}\right)^2\frac{1}{\lambda_i}(a_i\mathbf{x}_i^\top\boldsymbol{\beta})^2\right)\right\} \\ \cdot \exp\left\{-\frac{1}{2}\sum_{\{i:a_i=-1\}}\left(-2\frac{r_i}{1-\rho}a_i\mathbf{x}_i^\top\boldsymbol{\beta}\left(1+\frac{1}{(1-\rho)\lambda_i}\right)+\left(\frac{r_i}{1-\rho}\right)^2\frac{1}{\lambda_i}(a_i\mathbf{x}_i^\top\boldsymbol{\beta})^2\right)\right\} \exp\left\{-\frac{1}{2}\sum_{j=1}^p\frac{(\beta_j-\mu_{0,1})^2}{\sigma_0^2}\right\}$$

Let $n_1 = \sum_{i=1}^n \mathbb{1}(a_i = 1)$ and $n_{-1} = \sum_{i=1}^n \mathbb{1}(a_i = -1)$. Define \mathbf{X}_1 , \mathbf{W}_1 , \mathbf{R}_1 , and $\boldsymbol{\Lambda}_1$ as

$$\mathbf{X}_1 \equiv \begin{pmatrix} a_1x_{1,1} & \cdots & a_1x_{1,p} \\ \vdots & & \vdots \\ a_{n_1}x_{n_1,1} & \cdots & a_{n_1}x_{n_1,p} \end{pmatrix}_{(n_1 \times p)}, \quad \mathbf{W}_1 \equiv \begin{pmatrix} 1 + \frac{r_1}{\lambda_1} \\ \vdots \\ 1 + \frac{r_{n_1}}{\lambda_{n_1}} \end{pmatrix}_{(n_1 \times 1)}, \\ \mathbf{R}_1 \equiv \text{diag}(r_1/\rho, \dots, r_{n_1}/\rho)_{(n_1 \times n_1)}, \quad \text{and} \quad \boldsymbol{\Lambda}_1 = \text{diag}(\lambda_1, \dots, \lambda_{n_1}). \quad (16)$$

Define \mathbf{X}_{-1} , \mathbf{W}_{-1} , \mathbf{R}_{-1} , and $\boldsymbol{\Lambda}_{-1}$ analogously. Additionally define and $\boldsymbol{\Sigma}$ as $\text{diag}(\sigma_1, \dots, \sigma_p)$.

Then, we have that

$$p(\boldsymbol{\beta}|\boldsymbol{\lambda}, \boldsymbol{\omega}, \mathbf{r}, \mathbf{a}, \mathbf{x}) \\ \propto \exp\left\{-\frac{1}{2}\left[\boldsymbol{\beta}^\top \underbrace{(\mathbf{X}_1^\top \mathbf{R}_1^\top \boldsymbol{\Lambda}_1^{-1} \mathbf{R}_1 \mathbf{X}_1 + \mathbf{X}_{-1}^\top \mathbf{R}_{-1}^\top \boldsymbol{\Lambda}_{-1}^{-1} \mathbf{R}_{-1} \mathbf{X}_{-1} + \boldsymbol{\Sigma}^{-1})}_{\equiv B_1^{-1}} \boldsymbol{\beta} \right. \right. \\ \left. \left. - 2 \underbrace{(\mathbf{W}_1^\top \mathbf{R}_1 \mathbf{X}_1 + \mathbf{W}_{-1}^\top \mathbf{R}_{-1} \mathbf{X}_{-1} + \boldsymbol{\mu}_0^\top \boldsymbol{\Sigma}^{-1})}_{\equiv b_1} \boldsymbol{\beta} \right]\right\} \\ \propto \exp\left\{-\frac{1}{2}(\boldsymbol{\beta} - B_1 b_1)^\top B_1^{-1}(\boldsymbol{\beta} - B_1 b_1)\right\}.$$

In other words, the conditional distribution of $\boldsymbol{\beta}$ given $\boldsymbol{\lambda}, \boldsymbol{\omega}$, and is multivariate normal with mean $B_1 b_1$ and variance-covariance matrix B_1 . With the necessarily conditional distributions derived, the Gibbs sampling algorithm for sampling from the posterior distribution is given in Box 1.

Box 1. Gibbs sampling algorithm for normal distribution priors on $\boldsymbol{\beta}$

Initialize $\boldsymbol{\lambda}$ and $\boldsymbol{\beta}$; set the hyperparameters $\boldsymbol{\mu}_0$ and σ_0^2 .

Step 1: Draw $\boldsymbol{\beta}^{(g+1)}|\boldsymbol{\lambda}^{(g)}, \mathbf{r}, \mathbf{a}, \mathbf{x} \sim \mathcal{N}(B_1^{(g)} b_1^{(g)}, B_1^{(g)})$.

Step 2: Draw $\boldsymbol{\lambda}^{-1(g+1)}|\boldsymbol{\beta}^{(g)}, \mathbf{r}, \mathbf{a}, \mathbf{x}$ where

$$\lambda_i \sim \mathbb{1}(a_i = 1)\mathcal{GIG}\left(\frac{1}{2}, 1, \left(\frac{r_i}{\rho}\right)^2 (1 - a_i\mathbf{x}_i^\top\boldsymbol{\beta})^2\right) + \mathbb{1}(a_i = -1)\mathcal{GIG}\left(\frac{1}{2}, 1, \left(\frac{r_i}{1-\rho}\right)^2 (1 - a_i\mathbf{x}_i^\top\boldsymbol{\beta})^2\right).$$

Repeat Steps 1 and 2 until the chains converge.

3.4.2 Exponential power prior distribution for β

The full pseudo-posterior distribution when an exponential power prior distribution is specified for β (Equation (11)) has three unknown parameters, β , λ , and ω . To sample these parameters, we derive a Gibbs sampling algorithm, which entails sequentially sampling each parameter conditionally on the most up-to-date values of the other parameters. We give a high level summary of the derivation in this section and full details in Section 6.2. The conditional distribution of $\lambda_i | \beta, \mathbf{x}_i, \mathbf{a}_i, \mathbf{r}_i$ is the same as in the case with normal prior distributions for β given in Equation (15).

The conditional distribution of $\beta | \lambda, \omega, \mathbf{r}, \mathbf{a}, \mathbf{x}$ follows from standard arguments for Bayesian linear models. The notable difference from such a standard model is that $\beta | \lambda, \omega, \mathbf{r}, \mathbf{a}, \mathbf{x}$ is a mixture over two distributions, one for when the observed treatment in the data under analysis is 1 and one for when the observed treatment is -1 . The third term of the conditional distribution is the penalty. The conditional distribution of $\beta | \lambda, \omega, \mathbf{r}, \mathbf{a}, \mathbf{x}$ has the form

$p(\beta | \lambda, \omega, \mathbf{r}, \mathbf{a}, \mathbf{x})$

$$\begin{aligned} &\propto \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=1\}} \left(-2 \frac{r_i}{\rho} a_i \mathbf{x}_i^\top \beta \left(1 + \frac{r_i}{\rho \lambda_i} \right) + \left(\frac{r_i}{\rho} \right)^2 \frac{1}{\lambda_i} (a_i \mathbf{x}_i^\top \beta)^2 \right) \right\} \\ &\cdot \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=-1\}} \left(-2 \frac{r_i}{1-\rho} a_i \mathbf{x}_i^\top \beta \left(1 + \frac{1}{(1-\rho)\lambda_i} \right) + \left(\frac{r_i}{1-\rho} \right)^2 \frac{1}{\lambda_i} (a_i \mathbf{x}_i^\top \beta)^2 \right) \right\} \exp \left\{ -\frac{1}{2\nu^2} \sum_{j=1}^p \frac{\beta_j^2}{\sigma_j^2 \omega_j} \right\}. \end{aligned}$$

Letting $\Omega \equiv \text{diag}(\omega_1, \dots, \omega_p)_{(p \times p)}$, we have that

$p(\beta | \lambda, \omega, \mathbf{r}, \mathbf{a}, \mathbf{x})$

$$\begin{aligned} &\propto \exp \left\{ -\frac{1}{2} \left[\beta^\top \underbrace{(\mathbf{X}_1^\top \mathbf{R}_1^\top \Lambda_1^{-1} \mathbf{R}_1 \mathbf{X}_1 + \mathbf{X}_{-1}^\top \mathbf{R}_{-1}^\top \Lambda_{-1}^{-1} \mathbf{R}_{-1} \mathbf{X}_{-1} + \nu^{-2} \Omega^{-1} \Sigma^{-1})}_{\equiv B_2^{-1}} \beta - 2 \underbrace{(\mathbf{W}_1^\top \mathbf{R}_1 \mathbf{X}_1 + \mathbf{W}_{-1}^\top \mathbf{R}_{-1} \mathbf{X}_{-1})}_{\equiv b_2} \beta \right] \right\} \\ &\propto \exp \left\{ -\frac{1}{2} (\beta - B_2 b_2)^\top B_2^{-1} (\beta - B_2 b_2) \right\}. \end{aligned}$$

Thus, the conditional distribution of β given λ, ω , and is multivariate normal with mean $B_2 b_2$ and variance-covariance matrix B_2 . Full details of this derivation are given in Section 6.2.

Finally, the conditional distribution of $\omega | \beta, \nu$ is the same as that given in Corollary 3 of Polson and Scott [2011]. For $\alpha = 1$, the full conditional distribution of ω is $\omega_j^{-1} | \beta_j, \nu \sim \mathcal{IG}(\nu \sigma_j / |\beta_j|, 1)$. Together, with these three conditional distributions, we can summarize the Gibbs sampling algorithm as in Box 2.

Box 2. Gibbs sampling algorithm for exponential power distribution prior on β

Initialize λ, β and ω .

Step 1: Draw $\beta^{(g+1)} | \nu, \Lambda^{(g)}, \Omega^{(g)}, \mathbf{r}, \mathbf{a}, \mathbf{x} \sim \mathcal{N}(B_2^{(g)} b_2^{(g)}, B_2^{(g)})$.

Step 2: Draw $\lambda^{-1(g+1)} | \beta^{(g+1)}, \mathbf{r}, \mathbf{a}, \mathbf{x}$ where

$$\begin{aligned} \lambda_i^{(g+1)} | \beta^{(g+1)}, \nu, r_i, x_i \sim &\mathbb{1}(a_i = 1) \mathcal{GIG} \left(\frac{1}{2}, 1, \left(\frac{r_i}{\rho} \right)^2 (1 - a_i \mathbf{x}_i^\top \beta^{(g+1)})^2 \right) \\ &+ \mathbb{1}(a_i = -1) \mathcal{GIG} \left(\frac{1}{2}, 1, \left(\frac{r_i}{1-\rho} \right)^2 (1 - a_i \mathbf{x}_i^\top \beta^{(g+1)})^2 \right). \end{aligned}$$

Step 3: Draw $\omega_j^{-1(g+1)} | \beta_j^{(g+1)}, \nu \sim \mathcal{IG}(\nu \sigma_j / |\beta_j|^{-1}, 1)$
Repeat Steps 1, 2, and 3 until the chains converge.

3.4.3 Spike-and-slab prior distribution for β

The full pseudo-posterior distribution when a spike-and-slab prior is specified for β has three unknown parameters, β , λ , and γ . Similarly to the procedure outlined earlier, we employ a Gibbs sampling algorithm to obtain these parameters. The conditional distribution of $\lambda_i | \beta, \mathbf{x}_i, \mathbf{a}_i, \mathbf{r}_i$ is the same as in the case with normal prior distributions for β given in Equation (15).

The conditional distribution of β given λ , \mathbf{r} , \mathbf{a} , and \mathbf{x} mirrors the previous exponential power prior distribution, but without the third term of the penalty term. The adjustments are slight, with only the following modifications:

$$\begin{aligned} B_\gamma^{-1} &= X_1^T R_1^T \Lambda_1^{-1} R_1 X_1 + X_{-1}^T R_{-1}^T \Lambda_{-1}^{-1} R_{-1} X_{-1} \\ b_\gamma &= B_\gamma (W_1^T R_1 X_1 + W_{-1}^T R_{-1} X_{-1}) \end{aligned}$$

The spike-and-slab prior induces sparsity in coefficients through the parameters γ . The conditional distribution of γ given λ , \mathbf{r} , \mathbf{a} , and \mathbf{x} may be written as in Polson and Scott [2011] with b_γ and B_γ previously introduced :

$$p(\gamma | \lambda, \mathbf{r}, \mathbf{a}, \mathbf{x}, \nu) \propto p(\gamma) \frac{|\Sigma_\gamma^{-1} / \nu^2|^{1/2}}{|B_\gamma^{-1}|^{1/2}} \exp\left(-\frac{1}{2} \sum_{i=1}^n \frac{(1 + \lambda_i - a_i x_i^T b_\gamma)^2}{\lambda_i} - \frac{1}{2\nu^2} b_\gamma^T \Sigma_\gamma^{-1} b_\gamma\right)$$

By exploiting the quadratic term, the conditional distribution can be rewritten as in step 2 of Box 3. This form includes static terms that do not need to be recomputed in every iteration. To sample γ , a second Gibbs sampler nested within the first must be implemented.

Box 3. Gibbs sampling algorithm for spike-and-slab prior distribution on β

Initialize β and γ .

Step 1: Draw $\lambda^{-1(g+1)} | \beta^{(g+1)}, \mathbf{r}, \mathbf{a}, \mathbf{x}$ where

$$\begin{aligned} \lambda_i^{(g+1)} | \beta^{(g+1)}, \nu, r_i, x_i \sim & \mathbb{1}(a_i = 1) \mathcal{GIG}\left(\frac{1}{2}, 1, \left(\frac{r_i}{\rho}\right)^2 (1 - a_i \mathbf{x}_i^\top \beta^{(g+1)})^2\right) \\ & + \mathbb{1}(a_i = -1) \mathcal{GIG}\left(\frac{1}{2}, 1, \left(\frac{r_i}{1-\rho}\right)^2 (1 - a_i \mathbf{x}_i^\top \beta^{(g+1)})^2\right). \end{aligned}$$

Step 2: For $i = 1, \dots, k$ draw γ_i from $p(\gamma_i | \gamma_{-i})$ which is proportional to

$$p(\gamma | \lambda, \mathbf{r}, \mathbf{a}, \mathbf{x}, \nu) \propto p(\gamma) \frac{|\Sigma_\gamma^{-1} / \nu^2|^{1/2}}{|B_\gamma^{-1}|^{1/2}} \exp\left(-\frac{1}{2} [c(\lambda) + b_\gamma^T (X^T \Lambda^{-1} X)_\gamma b_\gamma - 2b_\gamma^T [X^T (1 + \lambda^{-1})]_\gamma] - \frac{1}{2\nu^2} b_\gamma^T \Sigma_\gamma^{-1} b_\gamma\right)$$

Step 3: When $\gamma_i = 1$, draw $\beta_\gamma^{(g+1)} | \nu, \Lambda^{(g)}, \mathbf{r}, \mathbf{a}, \mathbf{x} \sim \mathcal{N}(b_\gamma^{(g)}, B_\gamma^{(g)})$.
Repeat Steps 1, 2, and 3 until the chains converge.

3.5 Prediction and uncertainty quantification

Using the posterior predictive distribution, we can make treatment recommendations for a new patient and quantify our uncertainty in our recommendation. Let $\Theta = \{\beta, \lambda\}$ and \tilde{a} denote the recommended treatment for a new patient with features $\tilde{\mathbf{x}}$. Then

$$p(\tilde{a} = 1 | \tilde{\mathbf{x}}, \mathbf{X}, \mathbf{r}, \mathbf{a}) = \int_{\Theta} p(\tilde{a} = 1 | \tilde{\mathbf{x}}, \mathbf{X}, \mathbf{r}, \mathbf{a}, \beta, \lambda) p(\beta, \lambda | \mathbf{x}, \mathbf{r}, \mathbf{a}) d\theta.$$

For the class predictions, we can use the probit model which has the form

$$p(a = 1 | x) = \Phi(x^\top \beta)$$

where Φ is the cumulative distribution function of the standard normal distribution. Thus we can write the posterior predictive distribution as

$$p(\tilde{a} = 1 | \tilde{\mathbf{x}}, \mathbf{X}, \mathbf{r}, \mathbf{a}) = \int_{\Theta} \Phi(\tilde{\mathbf{x}}^\top \beta) p(\beta, \lambda | \mathbf{x}, \mathbf{r}, \mathbf{a}) d\theta. \quad (17)$$

4 Simulation study

4.1 Classification performance

We conducted simulation studies to assess the classification performance of the proposed method, following Zhao et al. [2012] and Song et al. [2015]. We compared the performance of OWL, Bayesian OWL with normal priors for β , Bayesian OWL with exponential power prior for β , and Bayesian OWL with spike-and-slab prior for β . For each simulated patient, we generated a 10-dimensional vector of patient features, X_1, \dots, X_{10} , drawn independently and uniformly distributed on $[-1, 1]$. Treatment A was drawn from $\{-1, 1\}$ independently of the prognostic variables with $\mathbb{P}(A = 1) = 1/2$. The outcome variable R was normally distributed with mean $Q_0 = 1 + 2X_1 + X_2 + 0.5X_3 + T_0(X, A)$ and standard deviation 1, where $T_0(X, A)$ was the interaction term between treatment and patient features. We examined two scenarios for the treatment-feature interaction term:

- Scenario 1 : $T_0(A, X) = (X_1 + X_2)A$
- Scenario 2 : $T_0(A, X) = 0.442(1 - X_1 - X_2)A$

Both scenarios 1 and 2 had linear decision boundaries determined by X_1 and X_2 . For scenario 1, the true optimal rule was given by $\mathbb{1}(X_1 + X_2 > 0)$, while for scenario 2, it was $\mathbb{1}(1 - X_1 - X_2 > 0)$. OWL was implemented with a linear kernel. For Bayesian OWL, Gibbs sampling was used to draw from the posterior distributions of the parameters 500 times. The first 150 draws were discarded as "burn-in" and point estimates of β were computed by taking the mean of the draws from the posterior distribution. Throughout, we set the hyperparameter $\nu = 0.8$.

For each scenario, we varied the training dataset from 100 to 200, 400 and 800 and tested on 1000 patients. For each training set size, we conducted 200 simulation runs. We evaluated classification performance using the misclassification rate, the ratio of the number of patients recommended a treatment counter to the true optimal rule divided by the total number of patients in the simulation run ($\frac{\text{Number of patients misclassified}}{\text{Total number of patients}}$). The simulation results are presented in Table 1 and 2.

n	OWL	Bayesian OWL Normal Prior	Bayesian OWL Exponential Power Prior	Bayesian OWL Spike and Slab
100	0.24	0.38	0.38	0.39
200	0.18	0.34	0.34	0.34
400	0.13	0.29	0.29	0.30
800	0.10	0.24	0.24	0.26

Table 1: Misclassification rates for different methods and sample sizes for scenario 1.

n	OWL	Bayesian OWL Normal Prior	Bayesian OWL Exponential Power Prior	Bayesian OWL Spike and Slab
100	0.22	0.38	0.38	0.39
200	0.15	0.34	0.34	0.34
400	0.13	0.31	0.31	0.30
800	0.10	0.25	0.25	0.22

Table 2: Misclassification rates for different methods and sample sizes for scenario 2.

As expected, the classification performance improved among all the ITR learning methods evaluated as the sample size increased. However, OWL consistently outperformed Bayesian OWL in all sample sizes and in both scenarios. We hypothesize that, with additional hyperparameter tuning, the performance of Bayesian OWL can be improved. Ordinarily, one would be hesitant to propose a method that is dominated by an existing method. However, the dominance of OWL is with respect to the misclassification rate. OWL, even with 800 samples in our simulation, has a 10% misclassification rate, and there is no way to determine which of the 10% of the simulated patients are likely misclassified (given a non-optimal treatment recommendation). In contrast, Bayesian OWL yields the entire posterior distribution of the estimated optimal ITR and thereby allows for immediate uncertainty quantification of individual-level treatment recommendations. In essence, Bayesian OWL can inform us of which treatment recommendations it is less certain about whereas OWL cannot. We demonstrate this in Section 4.2.

4.2 Treatment recommendation uncertainty quantification

To highlight the utility of quantifying the uncertainty of individual-level treatment recommendations, we generated a data set of 1000 patients under Scenario 1. We used these simulated data to train a Bayesian OWL model using the exponential power prior. Next, we used the same generative approach to simulate another 1000 patients. Specifically, we used a fine grid to generate X_1 and X_2 , the key variables in the true optimal rule (i.e., tailoring variables). The rationale for this approach was to generate a simulated set of patients whose characteristics covered the domain of the true optimal ITR so that we could estimate the uncertainty for combinations of X_1 and X_2 throughout the domain, $\mathcal{X}_1 \times \mathcal{X}_2 \in [-1, 1]^2$.

We evaluated the coefficients for the trained Bayesian OWL model. With an exponential power prior on β , which is analogous to $L1$ regularization, we would expect the coefficients of the features in the true optimal rule (tailoring variables) to be large and the coefficients of the features not in the rule to be driven close to 0. The magnitudes of the coefficients are displayed in Figure 1. As we would hope based on our knowledge of the true optimal rule, the magnitudes of the coefficients for X_1 and X_2 were larger than those for the other features, indicating that the estimated ITR using Bayesian OWL made decisions based on the correct patient features.




Figure 1: Feature importance

Figure 2 demonstrates Bayesian OWL’s ability to quantify uncertainty in its treatment recommendations. In Scenario 1, the true optimal ITR divides patients into two groups: patients in the upper-right half of the graph (where $X_1 + X_2 > 0$) should ideally get treatment $A = 1$, while those in the lower-left half should get treatment $A = -1$. Using the posterior predictive distribution as in Section 3.5, we computed the uncertainty associated with recommending Treatment 1 for patients whose features X_1 and X_2 lie in the upper-right half of the graph and the uncertainty associated with recommending Treatment -1 in the lower-left half. Notably, uncertainty was evaluated *individually* for each of the 1000 patients in our test cohort. This means that for each individual, we estimated how certain or uncertain we were about their specific treatment recommendation given their features X_1 and X_2 . Moreover, because our test set included simulated patients whose features spanned the domain of the ITR, we were able to compute uncertainty for every “type” of patient who could be recommended a treatment using the estimated ITR. We visualize the uncertainty across the domain of the ITR using a heat map (Figure 2). Certainties close to 1 (less uncertainty) are lighter in color and depicted with yellow and light green. In contrast, certainties close to 0 (more uncertain) are darker in color and depicted with purple and dark blue. As expected, Bayesian OWL is more certain about treatment recommendations for patients whose features are far from the decision boundary than for those that are close to the decision boundary.

Furthermore, in Figure 2, we have included misclassified individuals in our simulation. Those who were recommended treatment -1 but should have been recommended treatment 1 are indicated by red points, and those who were recommended treatment 1 but should have been recommended treatment -1 are indicated by orange points. We observe that the misclassified patients are located near the boundary, as we expect, and most noteworthy that they are located in regions where the model exhibits the greatest uncertainty (regions in which the background is shaded purple).




Figure 2: Heatmap of uncertainty quantification

5 Discussion

In this paper, we introduced a Bayesian formulation of OWL. To our knowledge, this is the first Bayesian strategy for *directly* learning an ITR. Moreover, we demonstrate how the Bayesian approach can enable us to quantify the uncertainty in our treatment recommendations. Both tasks, learning the optimal ITR and uncertainty quantification, can be implemented through a simple Gibbs sampling strategy for sampling the posterior distribution.

One may wonder why a Bayesian approach to OWL is needed since we already have OWL which can rely on methods from convex optimization. We believe that the answer to this question is two-fold. First, OWL is an appealing strategy for learning optimal ITRs because it models the decision rule directly rather than modeling conditional mean models and backing out the optimal ITR. Second, our Bayesian formulation of OWL yields a full pseudo-posterior distribution which means that we can quantify uncertainty in ITR’s treatment recommendations *at the individual level*. This is in contrast to the more common approach in the ITR literature which involves estimating the uncertainty in the value of the proposed ITR, i.e., the uncertainty in the expected value we would observe if everyone in the population were treated according to the rule. This may prove useful in the design and implementation of clinical studies by providing a strategy for identifying the types of patients for whom we feel confident in our ability to make treatment recommendations and the patient types that may require additional sampling and information to improve the recommendations. By casting the direct ITR learning approach into a probabilistic framework, we have widened the inferential possibilities for directly learned ITRs.

Our work has limitations. For example, we only examine linear rules. Although linear rules are simple to understand, there may be times when a nonlinear rule is desired or a nonlinear rule significantly outperforms a linear rule, i.e., clinically meaningful improvement from the nonlinear rule is worth the decrease in interpretability. Henao et al. [2014] proposed a strategy for Bayesian SVM for nonlinear decision boundaries, which is likely a good blueprint for extending this work to the nonlinear rule setting. Moreover, our approach does not attempt to do variable selection. This limits its applicability in high dimensional settings in which there is no information or weak information as to which tailoring covariates should be included in the treatment rule. Polson and Scott [2011] considers L1-regularization

on the decision boundary coefficients as well as a spike-and-slab prior to induce sparsity. These may be reasonable strategies to approximate the penalized version of OWL introduced by Song et al. [2015]. Finally, in simulation studies, OWL outperforms Bayesian OWL with respect to the misclassification rate, which is only ameliorated by the fact that Bayesian OWL can tell us when it is uncertain about its recommendations whereas OWL cannot.

While the direct learning, or classification, approach to learning optimal ITRs is incredibly powerful, leveraging tools from machine learning, inference and uncertainty quantification at the individual treatment recommendation level continues to be a challenge. Bayesian OWL overcomes this limitation by fully leveraging the benefits of direct-learning and the use of a probabilistic framework. Generating precision medicine evidence with wider inferential potential can improve our ability to build trust in these treatment algorithms and ultimately improve how we deploy precision medicine evidence in real-world health care decision making.

References

- Michael R Kosorok and Eric B Laber. Precision medicine. *Annu Rev Stat Appl*, 6:263–286, March 2019.
- S A Murphy. Optimal dynamic treatment regimes. *J. R. Stat. Soc. Series B Stat. Methodol.*, 65(2):331–355, May 2003.
- Richard S Sutton and Andrew G Barto. *Reinforcement Learning, second edition: An Introduction*. MIT Press, November 2018.
- Liliana Orellana, Andrea Rotnitzky, and James M Robins. Dynamic regime marginal structural mean models for estimation of optimal dynamic treatment regimes, part i: main content. *Int. J. Biostat.*, 6(2):Article 8, 2010.
- James Robins, Liliana Orellana, and Andrea Rotnitzky. Estimation and extrapolation of optimal treatment and testing strategies. *Stat. Med.*, 27(23):4678–4721, October 2008.
- Baquin Zhang, Anastasios A Tsiatis, Eric B Laber, and Marie Davidian. Robust estimation of optimal dynamic treatment regimes for sequential treatment decisions. *Biometrika*, 100(3), 2013.
- Christopher John Cornish Watkins. *Learning from Delayed Rewards*. PhD thesis, King’s College, 1989.
- Susan A Murphy. A generalization error for Q-Learning. *J. Mach. Learn. Res.*, 6:1073–1097, July 2005.
- Phillip J Schulte, Anastasios A Tsiatis, Eric B Laber, and Marie Davidian. Q- and a-learning methods for estimating optimal dynamic treatment regimes. *Stat. Sci.*, 29(4):640–661, November 2014.
- Erica E M Moodie, Bibhas Chakraborty, and Michael S Kramer. Q-learning for estimating optimal dynamic treatment rules from observational data. *Can. J. Stat.*, 40(4):629–645, December 2012.
- Yair Goldberg and Michael R Kosorok. Q-learning with censored data. *Ann. Stat.*, 40(1):529–560, February 2012.
- Yingqi Zhao, Donglin Zeng, A John Rush, and Michael R Kosorok. Estimating individualized treatment rules using outcome weighted learning. *J. Am. Stat. Assoc.*, 107(449):1106–1118, September 2012.
- Ying-Qi Zhao, Donglin Zeng, Eric B Laber, and Michael R Kosorok. New statistical learning methods for estimating optimal dynamic treatment regimes. *J. Am. Stat. Assoc.*, 110(510):583–598, 2015.
- Xin Zhou, Nicole Mayer-Hamblett, Umer Khan, and Michael R Kosorok. Residual weighted learning for estimating individualized treatment rules. *J. Am. Stat. Assoc.*, 112(517):169–187, May 2017.
- Ying Liu, Yuanjia Wang, Michael R Kosorok, Yingqi Zhao, and Donglin Zeng. Augmented outcome-weighted learning for estimating optimal dynamic treatment regimens. *Stat. Med.*, 37(26):3776–3788, November 2018.
- Sheng Fu, Qinying He, Sanguo Zhang, and Yufeng Liu. Robust outcome weighted learning for optimal individualized treatment rules. *J. Biopharm. Stat.*, 29(4):606–624, July 2019.
- Ying-Qi Zhao, Eric B Laber, Yang Ning, Sumona Saha, and Bruce E Sands. Efficient augmentation and relaxation learning for individualized treatment rules using observational data. *J. Mach. Learn. Res.*, 20, 2019.
- Thomas A Murray, Ying Yuan, and Peter F Thall. A bayesian machine learning approach for optimizing dynamic treatment regimes. *J. Am. Stat. Assoc.*, 113(523):1255–1267, October 2018.
- Peter F Thall, Hsi-Guang Sung, and Elihu H Estey. Selecting therapeutic strategies based on efficacy and death in multicourse clinical trials. *J. Am. Stat. Assoc.*, 97(457):29–39, March 2002.
- Peter F Thall, Leiko H Wooten, Christopher J Logothetis, Randall E Millikan, and Nizar M Tannir. Bayesian and frequentist two-stage treatment strategies based on sequential failure times subject to interval censoring. *Stat. Med.*, 26(26):4687–4702, November 2007.
- Elja Arjas and Olli Saarela. Optimal dynamic regimes: presenting a case for predictive inference. *Int. J. Biostat.*, 6(2): Article 10, March 2010.

- Tristan Zajonc. Bayesian inference for dynamic treatment regimes: Mobility, equity, and efficiency in student tracking. *J. Am. Stat. Assoc.*, 107(497):80–92, March 2012.
- Yanxun Xu, Peter Müller, Abdus S Wahed, and Peter F Thall. Bayesian nonparametric estimation for dynamic treatment regimes with sequential transition times. *J. Am. Stat. Assoc.*, 111(515):921–935, October 2016.
- Weichang Yu and Howard D Bondell. Bayesian likelihood-based regression for estimation of optimal dynamic treatment regimes. *J. R. Stat. Soc. Series B Stat. Methodol.*, 85(3):551–574, July 2023.
- Nicholas G Polson and Steven L Scott. Data augmentation for support vector machines. *ba*, 6(1):1–23, March 2011.
- Rui Song, Michael Kosorok, Donglin Zeng, Yingqi Zhao, Eric Laber, and Ming Yuan. On sparse representation for optimal individualized treatment selection with penalized outcome weighted learning. *Stat.*, 4(1):59–68, 2015.
- T J Mitchell and J J Beauchamp. Bayesian variable selection in linear regression. *J. Am. Stat. Assoc.*, 83(404):1023–1032, December 1988.
- Edward I George and Robert E McCulloch. Variable selection via gibbs sampling. *J. Am. Stat. Assoc.*, 88(423):881–889, September 1993.
- Luc Devroye. *Non-Uniform Random Variate Generation*. Springer New York, 1986.
- Ricardo Henao, Xin Yuan, and L Carin. Bayesian nonlinear support vector machines and discriminative factor modeling. *Adv. Neural Inf. Process. Syst.*, pages 1754–1762, December 2014.

6 Appendix

6.1 Representation of the pseudolikelihood function

Because we have assumed that the reward is strictly positive, the weight $\frac{r_i}{a_i\rho+(1-a_i)/2}$ is also positive and can be brought inside of the maximization operator so that

$$\begin{aligned}
 & L_i(a_i|r_i, \mathbf{x}_i, \boldsymbol{\beta}) \\
 &= \exp \left\{ -2 \frac{r_i}{a_i\rho + (1-a_i)/2} \max(1 - a_i\mathbf{x}_i^\top \boldsymbol{\beta}, 0) \right\} \\
 &= \exp \left\{ -2 \max \left(\frac{r_i}{a_i\rho + (1-a_i)/2} (1 - a_i\mathbf{x}_i^\top \boldsymbol{\beta}, 0) \right) \right\} \\
 &= \mathbb{1}(a_i = 1) \exp \left\{ -2 \max \left(\frac{r_i}{\rho} (1 - a_i\mathbf{x}_i^\top \boldsymbol{\beta}, 0) \right) \right\} + \mathbb{1}(a_i = -1) \exp \left\{ -2 \max \left(\frac{r_i}{1-\rho} (1 - a_i\mathbf{x}_i^\top \boldsymbol{\beta}, 0) \right) \right\}.
 \end{aligned}$$

The derivation of the pseudolikelihood representation follows Polson and Scott [2011]: Andrews and Mallows (1974) showed that $\int_0^\infty \frac{a}{\sqrt{2\pi\lambda}} e^{-\frac{1}{2}(a^2\lambda+b^2\lambda^{-1})} d\lambda = e^{-|ab|}$. Setting $a = 1$ and $b = u$, we have

$$\int_0^\infty \frac{1}{\sqrt{2\pi\lambda}} e^{-\frac{1}{2}(\lambda+u^2\lambda^{-1})} d\lambda = e^{-|u|}.$$

Multiplying through by e^{-u} and recalling the identity $\max(u, 0) = \frac{1}{2}(|u| + u)$, we have

$$\begin{aligned}
 & e^{-u} \int_0^\infty \frac{1}{\sqrt{2\pi\lambda}} e^{-\frac{1}{2}(\lambda+u^2\lambda^{-1})} d\lambda = e^{-|u|} e^{-u} \\
 \implies & \int_0^\infty \frac{1}{\sqrt{2\pi\lambda}} e^{-\frac{1}{2}(\lambda+u^2\lambda^{-1})-u} d\lambda = e^{-|u|-u} \\
 \implies & \int_0^\infty \frac{1}{\sqrt{2\pi\lambda}} e^{-\frac{1}{2\lambda}(\lambda^2+u^2+2u\lambda)} d\lambda = e^{-|u|-u} \\
 \implies & \int_0^\infty \frac{1}{\sqrt{2\pi\lambda}} e^{-\frac{1}{2\lambda}(u+\lambda)^2} d\lambda = e^{-2\max(u,0)}.
 \end{aligned}$$

Thus we can write the individual contribution of each observation to the marginal likelihood as

$$\begin{aligned}
& L_i(a_i|\lambda_i, r_i, \mathbf{x}_i, \boldsymbol{\beta}) \\
&= \mathbb{1}(a_i = 1) \exp \left\{ -2 \max \left(\frac{r_i}{\rho} (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta}, 0) \right) \right\} + \mathbb{1}(a_i = -1) \exp \left\{ -2 \max \left(\frac{r_i}{1-\rho} (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta}, 0) \right) \right\} \\
&= \mathbb{1}(a_i = 1) \int_0^\infty \frac{1}{\sqrt{2\pi\lambda_i}} \exp \left\{ -\frac{1}{2\lambda_i} \left(\frac{r_i}{\rho} (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta}) + \lambda_i \right)^2 \right\} d\lambda_i \\
&\quad + \mathbb{1}(a_i = -1) \int_0^\infty \frac{1}{\sqrt{2\pi\lambda_i}} \exp \left\{ -\frac{1}{2\lambda_i} \left(\frac{r_i}{1-\rho} (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta}) + \lambda_i \right)^2 \right\} d\lambda_i \\
&= \mathbb{1}(a_i = 1) \int_0^\infty \frac{1}{\sqrt{2\pi\lambda_i}} \exp \left\{ -\frac{1}{2\lambda_i} \left(\frac{r_i}{\rho} + \lambda_i - \frac{r_i}{\rho} a_i \mathbf{x}_i^\top \boldsymbol{\beta} \right)^2 \right\} d\lambda_i \\
&\quad + \mathbb{1}(a_i = -1) \int_0^\infty \frac{1}{\sqrt{2\pi\lambda_i}} \exp \left\{ -\frac{1}{2\lambda_i} \left(\frac{r_i}{1-\rho} + \lambda_i - \frac{r_i}{1-\rho} a_i \mathbf{x}_i^\top \boldsymbol{\beta} \right)^2 \right\} d\lambda_i,
\end{aligned}$$

and that $L_i(a_i, \lambda_i | r_i, \mathbf{x}_i, \boldsymbol{\beta})$ is a scale mixture of Gaussians.

6.2 Derivation of the Gibbs sampling algorithm

6.2.1 Conditional distribution of $\lambda_i | \boldsymbol{\beta}, \mathbf{x}_i, a_i, r_i$

$$\begin{aligned}
& p(\lambda_i | \boldsymbol{\beta}, \mathbf{x}_i, a_i, r_i) \\
&\propto \mathbb{1}(a = 1) \lambda_i^{-1/2} \cdot \exp \left\{ -\frac{1}{2} \frac{\left(\lambda_i - \frac{r_i}{\rho} (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta}) \right)^2}{\lambda_i} \right\} \\
&\quad \times \mathbb{1}(a = -1) \lambda_i^{-1/2} \cdot \exp \left\{ -\frac{1}{2} \frac{\left(\lambda_i - \frac{r_i}{1-\rho} (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta}) \right)^2}{\lambda_i} \right\} \\
&= \mathbb{1}(a = 1) \lambda_i^{-1/2} \cdot \exp \left\{ -\frac{1}{2} \frac{\left(\lambda_i^2 - 2\lambda_i \left(\frac{r_i}{\rho} \right) (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta}) + \left(\frac{r_i}{\rho} \right)^2 (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta})^2 \right)}{\lambda_i} \right\} \\
&\quad \times \mathbb{1}(a = -1) \lambda_i^{-1/2} \cdot \exp \left\{ -\frac{1}{2} \frac{\left(\lambda_i^2 - 2\lambda_i \left(\frac{r_i}{1-\rho} \right) (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta}) + \left(\frac{r_i}{1-\rho} \right)^2 (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta})^2 \right)}{\lambda_i} \right\} \\
&\propto \mathbb{1}(a = 1) \lambda_i^{-1/2} \cdot \exp \left\{ -\frac{1}{2} \left(\lambda_i + \left(\frac{r_i}{\rho} \right)^2 (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta}) \lambda_i^{-1} \right) \right\} \\
&\quad \times \mathbb{1}(a = -1) \lambda_i^{-1/2} \cdot \exp \left\{ -\frac{1}{2} \left(\lambda_i + \left(\frac{r_i}{1-\rho} \right)^2 (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta}) \lambda_i^{-1} \right) \right\}
\end{aligned}$$

From Devroye [1986], page 479, a random variable has the generalized inverse Gaussian distribution $\mathcal{GIG}(\gamma, \psi, \chi)$ if its density function is

$$p(x|\gamma, \psi, \chi) = C(\gamma, \psi, \chi) x^{\gamma-1} \exp \left\{ -\frac{1}{2} \left(\frac{\chi}{x} + \psi x \right) \right\},$$

where $C(\gamma, \psi, \chi)$ is a normalization constant. Thus

$$\begin{aligned}
 p(\lambda_i | \boldsymbol{\beta}, \mathbf{x}_i, a_i, r_i) \\
 \sim \mathbb{1}(a_i = 1) \mathcal{GIG} \left(\frac{1}{2}, 1, \left(\frac{r_i}{\rho} \right)^2 (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta})^2 \right) + \mathbb{1}(a_i = -1) \mathcal{GIG} \left(\frac{1}{2}, 1, \left(\frac{r_i}{1 - \rho} \right)^2 (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta})^2 \right).
 \end{aligned} \tag{18}$$

Recall that if $X \sim \mathcal{GIG}(\frac{1}{2}, \lambda, \chi)$, then $X^{-1} \sim \mathcal{IG}(\mu, \lambda)$ where $\chi = \lambda/\mu^2$ and \mathcal{IG} denotes the inverse Gaussian distribution. Consequently, we can write the

$$\begin{aligned}
 p(\lambda_i | \boldsymbol{\beta}, \mathbf{x}_i, a_i, r_i) \\
 \sim \mathbb{1}(a_i = 1) \mathcal{GIG} \left(\frac{1}{2}, 1, \left(\frac{r_i}{\rho} \right)^2 (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta})^2 \right) + \mathbb{1}(a_i = -1) \mathcal{GIG} \left(\frac{1}{2}, 1, \left(\frac{r_i}{1 - \rho} \right)^2 (1 - a_i \mathbf{x}_i^\top \boldsymbol{\beta})^2 \right).
 \end{aligned} \tag{19}$$

6.2.2 Conditional distribution of $\beta|\lambda, \mu_0, \sigma_0^2, \mathbf{r}, \mathbf{a}, \mathbf{x}$ (Normal prior)

$$\begin{aligned}
& p(\beta|\lambda, \mu_0, \sigma_0^2, \mathbf{r}, \mathbf{a}, \mathbf{x}) \\
& \propto \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=1\}} \frac{\left(\frac{r_i}{\rho} + \lambda_i - \left(\frac{r_i}{\rho} \right) a_i \mathbf{x}_i^\top \beta \right)^2}{\lambda_i} \right\} \cdot \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=-1\}} \frac{\left(\frac{r_i}{1-\rho} + \lambda_i - \left(\frac{r_i}{1-\rho} \right) a_i \mathbf{x}_i^\top \beta \right)^2}{\lambda_i} \right\} \\
& \quad \cdot \exp \left\{ -\frac{1}{2} \sum_{j=1}^p \frac{(\beta_j - \mu_{0,j})^2}{\sigma_0^2} \right\} \\
& \propto \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=1\}} \frac{\left(\lambda_i + \frac{r_i}{\rho} (1 - a_i \mathbf{x}_i^\top \beta) \right)^2}{\lambda_i} \right\} \cdot \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=-1\}} \frac{\left(\lambda_i + \frac{r_i}{1-\rho} (1 - a_i \mathbf{x}_i^\top \beta) \right)^2}{\lambda_i} \right\} \\
& \quad \cdot \exp \left\{ -\frac{1}{2} \sum_{j=1}^p \frac{\beta_j^2 - 2\beta_j \mu_{0,j} + \mu_{0,j}^2}{\sigma_0^2} \right\} \\
& = \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=1\}} \frac{\lambda_i^2 + 2\lambda_i \frac{r_i}{\rho} (1 - a_i \mathbf{x}_i^\top \beta) + \left(\frac{r_i}{\rho} \right)^2 (1 - a_i \mathbf{x}_i^\top \beta)^2}{\lambda_i} \right\} \\
& \quad \cdot \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=-1\}} \frac{\lambda_i^2 + 2\lambda_i \frac{r_i}{1-\rho} (1 - a_i \mathbf{x}_i^\top \beta) + \left(\frac{r_i}{1-\rho} \right)^2 (1 - a_i \mathbf{x}_i^\top \beta)^2}{\lambda_i} \right\} \exp \left\{ -\frac{1}{2} \sum_{j=1}^p \frac{\beta_j^2 - 2\beta_j \mu_{0,j} + \mu_{0,j}^2}{\sigma_0^2} \right\} \\
& \propto \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=1\}} \left(-2\frac{r_i}{\rho} a_i \mathbf{x}_i^\top \beta + \left(\frac{r_i}{\rho} \right)^2 \frac{1}{\lambda_i} (-2a_i \mathbf{x}_i^\top \beta + (a_i \mathbf{x}_i^\top \beta)^2) \right) \right\} \\
& \quad \cdot \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=-1\}} \left(-2\frac{r_i}{1-\rho} a_i \mathbf{x}_i^\top \beta + \left(\frac{r_i}{1-\rho} \right)^2 \frac{1}{\lambda_i} (-2a_i \mathbf{x}_i^\top \beta + (a_i \mathbf{x}_i^\top \beta)^2) \right) \right\} \\
& \quad \cdot \exp \left\{ -\frac{1}{2} \sum_{j=1}^p \frac{\beta_j^2 - 2\beta_j \mu_{0,j}}{\sigma_0^2} \right\} \\
& \propto \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=1\}} \left(-2\frac{r_i}{\rho} a_i \mathbf{x}_i^\top \beta \left(1 + \frac{r_i}{\rho \lambda_i} \right) + \left(\frac{r_i}{\rho} \right)^2 \frac{1}{\lambda_i} (a_i \mathbf{x}_i^\top \beta)^2 \right) \right\} \\
& \quad \cdot \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=-1\}} \left(-2\frac{r_i}{1-\rho} a_i \mathbf{x}_i^\top \beta \left(1 + \frac{1}{(1-\rho)\lambda_i} \right) + \left(\frac{r_i}{1-\rho} \right)^2 \frac{1}{\lambda_i} (a_i \mathbf{x}_i^\top \beta)^2 \right) \right\} \\
& \quad \cdot \exp \left\{ -\frac{1}{2} \sum_{j=1}^p \frac{\beta_j^2 - 2\beta_j \mu_{0,j}}{\sigma_0^2} \right\}
\end{aligned}$$

Consider the summation inside the exponential of the first term, we have

$$\sum_{\{i:a_i=1\}} \left(\underbrace{-2\frac{r_i}{\rho} a_i \mathbf{x}_i^\top \beta \left(1 + \frac{r_i}{\rho \lambda_i} \right)}_{\text{Term 1}} + \underbrace{\left(\frac{r_i}{\rho} \right)^2 \frac{1}{\lambda_i} (a_i \mathbf{x}_i^\top \beta)^2}_{\text{Term 2}} \right).$$

Let $n_1 = \sum_{i=1}^n \mathbb{1}(a_i = 1)$ Working with the summation over the first term

$$\begin{aligned} \text{Term 1} &= \sum_{\{a_i=1\}} \frac{r_i}{\rho} a_i \mathbf{x}_i^\top \boldsymbol{\beta} \left(1 + \frac{r_i}{\rho \lambda_i}\right) \\ &= \frac{r_1}{\rho} a_1 \mathbf{x}_1^\top \boldsymbol{\beta} \left(1 + \frac{r_1}{\rho} \lambda_1\right) + \cdots + \frac{r_{n_1}}{\rho} a_{n_1} \mathbf{x}_{n_1}^\top \boldsymbol{\beta} \left(1 + \frac{r_{n_1}}{\rho} \lambda_{n_1}\right) \\ &= \left(\frac{r_1}{\rho} \left(1 + \frac{r_1}{\rho \lambda_1}\right), \dots, \frac{r_{n_1}}{\rho} \left(1 + \frac{r_{n_1}}{\rho \lambda_{n_1}}\right)\right) \begin{pmatrix} a_1 \mathbf{x}_1^\top \boldsymbol{\beta} \\ \vdots \\ a_{n_1} \mathbf{x}_{n_1}^\top \boldsymbol{\beta} \end{pmatrix} \end{aligned}$$

Define \mathbf{X}_1 , \mathbf{W}_1 , and \mathbf{R}_1 as

$$\begin{aligned} \mathbf{X}_1 &\equiv \begin{pmatrix} a_1 x_{1,1} & \cdots & a_1 x_{1,p} \\ \vdots & & \vdots \\ a_{n_1} x_{n_1,1} & \cdots & a_{n_1} x_{n_1,p} \end{pmatrix}_{(n_1 \times p)}, & \mathbf{W}_1 &\equiv \begin{pmatrix} 1 + \frac{r_1}{\lambda_1} \\ \vdots \\ 1 + \frac{r_{n_1}}{\lambda_{n_1}} \end{pmatrix}_{(n_1 \times 1)}, \text{ and} \\ \mathbf{R}_1 &\equiv \text{diag}(r_1/\rho, \dots, r_{n_1}/\rho)_{(n_1 \times n_1)}. \end{aligned} \tag{20}$$

Then,

$$\text{Term 1} = \frac{1}{\rho} \mathbf{W}_1^\top \mathbf{R}_1 \mathbf{X}_1 \boldsymbol{\beta} \tag{21}$$

because

$$\begin{aligned} \mathbf{W}_1^\top \mathbf{R}_1 \mathbf{X}_1 \boldsymbol{\beta} &= \begin{pmatrix} 1 + \frac{r_1}{\lambda_1} & \cdots & 1 + \frac{r_{n_1}}{\lambda_{n_1}} \end{pmatrix} \begin{pmatrix} r_1/\rho & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & r_{n_1}/\rho \end{pmatrix} \begin{pmatrix} a_1 x_{1,1} & \cdots & a_1 x_{1,p} \\ \vdots & & \vdots \\ a_{n_1} x_{n_1,1} & \cdots & a_{n_1} x_{n_1,p} \end{pmatrix} \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_p \end{pmatrix} \\ &= \frac{1}{\rho} \begin{pmatrix} r_1 \left(1 + \frac{r_1}{\lambda_1}\right) & \cdots & r_{n_1} \left(1 + \frac{r_{n_1}}{\lambda_{n_1}}\right) \end{pmatrix} \begin{pmatrix} a_1 \mathbf{x}_1^\top \boldsymbol{\beta} \\ \vdots \\ a_{n_1} \mathbf{x}_{n_1}^\top \boldsymbol{\beta} \end{pmatrix}. \end{aligned}$$

Next, consider the summation over Term 2,

$$\begin{aligned} \text{Term 2} &= \sum_{\{i:a_i=-1\}} \left(\frac{r_i}{\rho}\right)^2 \frac{1}{\lambda_i} (a_i \mathbf{x}_i^\top \boldsymbol{\beta})^2 \\ &= \left(\frac{1}{\rho^2}\right) \left(\frac{r_1^2}{\lambda_1}\right) a_1^2 (\mathbf{x}_1^\top \boldsymbol{\beta})^2 + \cdots + \left(\frac{1}{\rho^2}\right) \left(\frac{r_{n_1}^2}{\lambda_{n_1}}\right) a_{n_1}^2 (\mathbf{x}_{n_1}^\top \boldsymbol{\beta})^2. \end{aligned}$$

Observe that

$$\begin{aligned}
 & a_i^2 \boldsymbol{\beta}^\top \mathbf{x}_i \mathbf{x}_i^\top \boldsymbol{\beta} \\
 &= \boldsymbol{\beta}^\top \begin{pmatrix} a_i x_{i,1} \\ \vdots \\ a_i x_{i,p} \end{pmatrix} (a_i x_{i,1} \ \cdots \ a_i x_{i,p}) \boldsymbol{\beta} \\
 &= \boldsymbol{\beta}^\top \begin{pmatrix} a_i^2 x_{i,1}^2 & a_i^2 x_{i,1} x_{i,2} & \cdots & a_i^2 x_{i,1} x_{i,p} \\ a_i^2 x_{i,1} x_{i,2} & a_i^2 x_{i,2}^2 & \cdots & a_i^2 x_{i,2} x_{i,p} \\ \vdots & \vdots & \ddots & \vdots \\ a_i^2 x_{i,1} x_{i,p} & a_i^2 x_{i,2} x_{i,p} & \cdots & a_i^2 x_{i,p}^2 \end{pmatrix} \boldsymbol{\beta} \\
 &= (\beta_1 \ \cdots \ \beta_p) \begin{pmatrix} a_i^2 x_{i,1}^2 & a_i^2 x_{i,1} x_{i,2} & \cdots & a_i^2 x_{i,1} x_{i,p} \\ a_i^2 x_{i,1} x_{i,2} & a_i^2 x_{i,2}^2 & \cdots & a_i^2 x_{i,2} x_{i,p} \\ \vdots & \vdots & \ddots & \vdots \\ a_i^2 x_{i,1} x_{i,p} & a_i^2 x_{i,2} x_{i,p} & \cdots & a_i^2 x_{i,p}^2 \end{pmatrix} \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_p \end{pmatrix} \\
 &= (\beta_1 a_i^2 x_{i,1}^2 + \beta_2 a_i^2 x_{i,1} x_{i,2} + \cdots + \beta_p a_i^2 x_{i,1} x_{i,p} \ \cdots \ \beta_1 a_i^2 x_{i,1} x_{i,p} + \beta_2 a_i^2 x_{i,2} x_{i,p} + \cdots + \beta_p a_i^2 x_{i,p}^2) \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_p \end{pmatrix} \\
 &= \beta_1^2 a_i^2 x_{i,1}^2 + \beta_1 \beta_2 a_i^2 x_{i,1} x_{i,2} + \cdots + \beta_1 \beta_p a_i^2 x_{i,1} x_{i,p} + \cdots + \beta_1 \beta_p a_i^2 x_{i,1} x_{i,p} + \beta_2 \beta_p a_i^2 x_{i,2} x_{i,p} + \cdots + \beta_p^2 a_i^2 x_{i,p}^2 \\
 &= \sum_{j=1}^p a_i^2 x_{i,j}^2 \beta_j^2 + 2 \sum_{j=1}^p \sum_{k \neq j}^p a_i^2 x_{i,j} x_{i,k} \beta_j \beta_k \\
 &= (a_i x_{i,1} \beta_1 + \cdots + a_i x_{i,p} \beta_p) \cdot a_i (x_{i,1} \beta_1 + \cdots + x_{i,p} \beta_p) \\
 &= (a_i \mathbf{x}_i^\top \boldsymbol{\beta}) \cdot (a_i \mathbf{x}_i^\top \boldsymbol{\beta}) \\
 &= (a_i \mathbf{x}_i^\top \boldsymbol{\beta})^2.
 \end{aligned}$$

Also observe that

$$\begin{aligned}
 \boldsymbol{\beta}^\top \mathbf{X}_1^\top \mathbf{X}_1 \boldsymbol{\beta} &= \boldsymbol{\beta}^\top \begin{pmatrix} a_1 x_{1,1} & \cdots & a_{n_1} x_{n_1,1} \\ \vdots & \cdots & \vdots \\ a_1 x_{1,p} & \cdots & a_{n_1} x_{n_1,p} \end{pmatrix} \begin{pmatrix} a_1 x_{1,1} & \cdots & a_1 x_{1,p} \\ \vdots & \cdots & \vdots \\ a_{n_1} x_{n_1,1} & \cdots & a_{n_1} x_{n_1,p} \end{pmatrix} \boldsymbol{\beta} \\
 &= (a_1 \boldsymbol{\beta}^\top \mathbf{x}_1 \ \cdots \ a_{n_1} \boldsymbol{\beta}^\top \mathbf{x}_{n_1}) \begin{pmatrix} a_1 \mathbf{x}_1^\top \boldsymbol{\beta} \\ \vdots \\ a_{n_1} \mathbf{x}_{n_1}^\top \boldsymbol{\beta} \end{pmatrix} \\
 &= \begin{pmatrix} a_1^2 \boldsymbol{\beta}^\top \mathbf{x}_1 \mathbf{x}_1^\top \boldsymbol{\beta} \\ \vdots \\ a_{n_1}^2 \boldsymbol{\beta}^\top \mathbf{x}_{n_1} \mathbf{x}_{n_1}^\top \boldsymbol{\beta} \end{pmatrix} \tag{22}
 \end{aligned}$$

Define $\boldsymbol{\Lambda}_1 = \text{diag}(\lambda_1, \dots, \lambda_{n_1})$. Then

$$\begin{aligned}
 \text{Term 2} &= \left(\frac{1}{\rho^2}\right) \left(\frac{r_1^2}{\lambda_1}\right) a_1^2 (\mathbf{x}_1^\top \boldsymbol{\beta})^2 + \cdots + \left(\frac{1}{\rho^2}\right) \left(\frac{r_{n_1}^2}{\lambda_{n_1}}\right) a_{n_1}^2 (\mathbf{x}_{n_1}^\top \boldsymbol{\beta})^2 \\
 &= \left(\frac{1}{\rho^2}\right) \left(\begin{pmatrix} r_1^2 \\ \lambda_1 \end{pmatrix} \ \cdots \ \begin{pmatrix} r_{n_1}^2 \\ \lambda_{n_1} \end{pmatrix} \right) \begin{pmatrix} a_1^2 \boldsymbol{\beta}^\top \mathbf{x}_1 \mathbf{x}_1^\top \boldsymbol{\beta} \\ \vdots \\ a_{n_1}^2 \boldsymbol{\beta}^\top \mathbf{x}_{n_1} \mathbf{x}_{n_1}^\top \boldsymbol{\beta} \end{pmatrix} \\
 &= \left(\frac{1}{\rho^2}\right) \left(\begin{pmatrix} r_1^2 \\ \lambda_1 \end{pmatrix} \ \cdots \ \begin{pmatrix} r_{n_1}^2 \\ \lambda_{n_1} \end{pmatrix} \right) \boldsymbol{\beta}^\top \mathbf{X}_1^\top \mathbf{X}_1 \boldsymbol{\beta} \\
 &= \boldsymbol{\beta}^\top \mathbf{X}_1^\top \mathbf{R}_1^\top \boldsymbol{\Lambda}_1^{-1} \mathbf{R}_1 \mathbf{X}_1 \boldsymbol{\beta}. \tag{23}
 \end{aligned}$$

Let $n_{-1} = \sum_{i=1}^n \mathbb{1}(a_i = -1)$. Similarly, define \mathbf{X}_{-1} , \mathbf{W}_{-1} , \mathbf{R}_{-1} , and $\mathbf{\Lambda}_{-1}$ as

$$\mathbf{X}_{-1} \equiv \begin{pmatrix} a_1 x_{1,1} & \cdots & a_1 x_{1,p} \\ \vdots & & \vdots \\ a_{n_{-1}} x_{n_{-1},1} & \cdots & a_{n_{-1}} x_{n_{-1},p} \end{pmatrix}_{(n_{-1} \times p)}, \quad \mathbf{W}_{-1} \equiv \begin{pmatrix} 1 + \frac{r_1}{\lambda_1} \\ \vdots \\ 1 + \frac{r_{n_{-1}}}{\lambda_{n_{-1}}} \end{pmatrix}_{(n_{-1} \times 1)},$$

$$\mathbf{R}_{-1} \equiv \text{diag}(r_1/(1-\rho), \dots, r_{n_{-1}}/(1-\rho))_{(n_{-1} \times n_{-1})}, \text{ and } \mathbf{\Lambda}_{-1} = \text{diag}(\lambda_1, \dots, \lambda_{n_{-1}}).$$

Additionally define $\mathbf{\Sigma} \equiv \text{diag}(\sigma_1, \dots, \sigma_p)$ so that we can write

$$\exp \left\{ -\frac{1}{2} \sum_{j=1}^p \frac{\beta_j^2 - 2\beta_j \mu_{0,j}}{\sigma_0^2} \right\} = \exp \left\{ -\frac{1}{2} (\boldsymbol{\beta}^\top \mathbf{\Sigma}^{-1} \boldsymbol{\beta} - 2\boldsymbol{\mu}_0^\top \mathbf{\Sigma}^{-1} \boldsymbol{\beta}) \right\}. \quad (24)$$

Thus, we have that

$$\begin{aligned} & p(\boldsymbol{\beta} | \boldsymbol{\lambda}, \boldsymbol{\omega}, \mathbf{r}, \mathbf{a}, \mathbf{x}) \\ & \propto \exp \left\{ -\frac{1}{2} \left(-2\mathbf{W}_1^\top \mathbf{R}_1 \mathbf{X}_1 \boldsymbol{\beta} + \boldsymbol{\beta}^\top \mathbf{X}_1^\top \mathbf{R}_1^\top \mathbf{\Lambda}_1^{-1} \mathbf{R}_1 \mathbf{X}_1 \boldsymbol{\beta} \right) \right\} \\ & \quad \cdot \exp \left\{ -\frac{1}{2} \left(-2\mathbf{W}_{-1}^\top \mathbf{R}_{-1} \mathbf{X}_{-1} \boldsymbol{\beta} + \boldsymbol{\beta}^\top \mathbf{X}_{-1}^\top \mathbf{R}_{-1}^\top \mathbf{\Lambda}_{-1}^{-1} \mathbf{R}_{-1} \mathbf{X}_{-1} \boldsymbol{\beta} \right) \right\} \\ & \quad \cdot \exp \left\{ -\frac{1}{2} (-2\boldsymbol{\mu}_0^\top \mathbf{\Sigma}^{-1} \boldsymbol{\beta} + \boldsymbol{\beta}^\top \mathbf{\Sigma}^{-1} \boldsymbol{\beta}) \right\} \\ & = \exp \left\{ -\frac{1}{2} \left[\boldsymbol{\beta}^\top \underbrace{(\mathbf{X}_1^\top \mathbf{R}_1^\top \mathbf{\Lambda}_1^{-1} \mathbf{R}_1 \mathbf{X}_1 + \mathbf{X}_{-1}^\top \mathbf{R}_{-1}^\top \mathbf{\Lambda}_{-1}^{-1} \mathbf{R}_{-1} \mathbf{X}_{-1} + \mathbf{\Sigma}^{-1})}_{\equiv B_1^{-1}} \boldsymbol{\beta} \right. \right. \\ & \quad \left. \left. - 2 \underbrace{(\mathbf{W}_1^\top \mathbf{R}_1 \mathbf{X}_1 + \mathbf{W}_{-1}^\top \mathbf{R}_{-1} \mathbf{X}_{-1} + \boldsymbol{\mu}_0^\top \mathbf{\Sigma}^{-1})}_{\equiv b_1} \boldsymbol{\beta} \right] \right\} \\ & = \exp \left\{ -\frac{1}{2} (\boldsymbol{\beta} - B_1 b_1)^\top B^{-1} (\boldsymbol{\beta} - B_1 b_1) - b_1^\top B_1 b_1 \right\} \\ & \propto \exp \left\{ -\frac{1}{2} (\boldsymbol{\beta} - B_1 b_1)^\top B_1^{-1} (\boldsymbol{\beta} - B_1 b_1) \right\}. \end{aligned}$$

The conditional distribution of $\boldsymbol{\beta}$ given $\boldsymbol{\lambda}$ is multivariate normal with mean $B_1 b_1$ and variance-covariance matrix B_1 .

6.2.3 Conditional distribution of $\boldsymbol{\beta} | \boldsymbol{\lambda}, \boldsymbol{\omega}, \mathbf{r}, \mathbf{a}, \mathbf{x}$ (Exponential power prior)

$$\begin{aligned} & p(\boldsymbol{\beta} | \boldsymbol{\lambda}, \boldsymbol{\omega}, \mathbf{r}, \mathbf{a}, \mathbf{x}) \\ & \propto \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=1\}} \frac{\left(\frac{r_i}{\rho} + \lambda_i - \left(\frac{r_i}{\rho} \right) a_i \mathbf{x}_i^\top \boldsymbol{\beta} \right)^2}{\lambda_i} \right\} \cdot \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=-1\}} \frac{\left(\frac{r_i}{1-\rho} + \lambda_i - \left(\frac{r_i}{1-\rho} \right) a_i \mathbf{x}_i^\top \boldsymbol{\beta} \right)^2}{\lambda_i} \right\} \\ & \quad \cdot \exp \left\{ -\frac{1}{2\nu^2} \sum_{j=1}^p \frac{\beta_j^2}{\sigma_j^2 \omega_j} \right\} \\ & \propto \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=1\}} \left(-2 \frac{r_i}{\rho} a_i \mathbf{x}_i^\top \boldsymbol{\beta} \left(1 + \frac{r_i}{\rho \lambda_i} \right) + \left(\frac{r_i}{\rho} \right)^2 \frac{1}{\lambda_i} (a_i \mathbf{x}_i^\top \boldsymbol{\beta})^2 \right) \right\} \\ & \quad \cdot \exp \left\{ -\frac{1}{2} \sum_{\{i:a_i=-1\}} \left(-2 \frac{r_i}{1-\rho} a_i \mathbf{x}_i^\top \boldsymbol{\beta} \left(1 + \frac{1}{(1-\rho)\lambda_i} \right) + \left(\frac{r_i}{1-\rho} \right)^2 \frac{1}{\lambda_i} (a_i \mathbf{x}_i^\top \boldsymbol{\beta})^2 \right) \right\} \\ & \quad \cdot \exp \left\{ -\frac{1}{2\nu^2} \sum_{j=1}^p \frac{\beta_j^2}{\sigma_j^2 \omega_j} \right\} \end{aligned}$$

The summation inside the first and second exponential terms are the same as in the derivation under the normal distribution prior for β (Section 6.2.2). Letting $\Omega \equiv \text{diag}(\omega_1, \dots, \omega_p)_{(p \times p)}$, we can write

$$\begin{aligned}
\sum_{j=1}^p \frac{\beta_j^2}{\sigma_j^2 \omega_j} &= (1/(\sigma_1^2 \omega_1) \quad \dots \quad 1/(\sigma_p^2 \omega_p)) \begin{pmatrix} \beta_1^2 \\ \vdots \\ \beta_p^2 \end{pmatrix} \\
&= (\beta_1^2 \quad \dots \quad \beta_p^2) \begin{pmatrix} 1/(\sigma_1^2 \omega_1) & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 1/(\sigma_p^2 \omega_p) \end{pmatrix} \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_p \end{pmatrix} \\
&= (\beta_1^2 \quad \dots \quad \beta_p^2) \begin{pmatrix} 1/\sigma_1^2 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 1/\sigma_p^2 \end{pmatrix} \begin{pmatrix} 1/\omega_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 1/\omega_p \end{pmatrix} \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_p \end{pmatrix} \\
&= \beta^\top \Omega^{-1} \Sigma^{-1} \beta
\end{aligned}$$

Thus, we have that

$$\begin{aligned}
&p(\beta | \lambda, \omega, \mathbf{r}, \mathbf{a}, \mathbf{x}) \\
&\propto \exp \left\{ -\frac{1}{2} \left(-2\mathbf{W}_1^\top \mathbf{R}_1 \mathbf{X}_1 \beta + \beta^\top \mathbf{X}_1^\top \mathbf{R}_1^\top \Lambda_1^{-1} \mathbf{R}_1 \mathbf{X}_1 \beta \right) \right\} \\
&\quad \cdot \exp \left\{ -\frac{1}{2} \left(-2\mathbf{W}_{-1}^\top \mathbf{R}_{-1} \mathbf{X}_{-1} \beta + \beta^\top \mathbf{X}_{-1}^\top \mathbf{R}_{-1}^\top \Lambda_{-1}^{-1} \mathbf{R}_{-1} \mathbf{X}_{-1} \beta \right) \right\} \\
&\quad \cdot \exp \left\{ -\frac{1}{2} \nu^{-2} \beta^\top \Omega^{-1} \Sigma^{-1} \beta \right\} \\
&= \exp \left\{ -\frac{1}{2} \left[\beta^\top \underbrace{(\mathbf{X}_1^\top \mathbf{R}_1^\top \Lambda_1^{-1} \mathbf{R}_1 \mathbf{X}_1 + \mathbf{X}_{-1}^\top \mathbf{R}_{-1}^\top \Lambda_{-1}^{-1} \mathbf{R}_{-1} \mathbf{X}_{-1} + \nu^{-2} \Omega^{-1} \Sigma^{-1})}_{\equiv B_2^{-1}} \beta \right. \right. \\
&\quad \left. \left. - 2 \underbrace{(\mathbf{W}_1^\top \mathbf{R}_1 \mathbf{X}_1 + \mathbf{W}_{-1}^\top \mathbf{R}_{-1} \mathbf{X}_{-1})}_{\equiv b_2} \beta \right] \right\} \\
&= \exp \left\{ -\frac{1}{2} (\beta - B_2 b_2)^\top B_2^{-1} (\beta - B_2 b_2) - b_2^\top B_2 b_2 \right\} \\
&\propto \exp \left\{ -\frac{1}{2} (\beta - B_2 b_2)^\top B_2^{-1} (\beta - B_2 b_2) \right\}.
\end{aligned}$$

The conditional distribution of β given λ, ω , and is multivariate normal with mean $B_2 b_2$ and variance-covariance matrix B_2 .