Categorified Open Topological Field Theories

Lukas Müller a and Lukas Woike b

^a Perimeter Institute N2L 2Y5 Waterloo Canada

^b Institut de Mathématiques de Bourgogne UMR 5584 CNRS & Université de Bourgogne F-21000 Dijon France

In this short note, we classify linear categorified open topological field theories in dimension two by pivotal Grothendieck-Verdier categories. In combination with recently developed string-net techniques, this leads to a new description of the spaces of conformal blocks of Drinfeld centers $Z(\mathcal{C})$ of pivotal finite tensor categories \mathcal{C} in terms of the modular envelope of the cyclic associative operad. If \mathcal{C} is unimodular, we prove that the space of conformal blocks inherits the structure of a module over the algebra of class functions of \mathcal{C} for every free boundary component. As a further application, we prove that the sewing along a boundary circle for the modular functor for $Z(\mathcal{C})$ can be decomposed into a sewing procedure along an interval and the application of the partial trace. Finally, we construct mapping class group representations from Grothendieck-Verdier categories that are not necessarily rigid and make precise how these generalize existing constructions.

CONTENTS

1	Introduction and summary	1
2	Classification of categorified open topological field theories	5
3	A topological characterization of rigidity	6
4	Comparison with the string-net construction	7
5	The action of the algebra of class functions	10
6	A gluing result involving the partial trace	11

1 INTRODUCTION AND SUMMARY

It is a classical result in quantum topology that any symmetric Frobenius algebra A in the category of vector spaces over a fixed field k extends uniquely to an open two-dimensional topological field theory; this follows from the results of [Cos07, LP08, WW16]. A symmetric Frobenius algebra A is exactly a cyclic associative algebra in vector spaces, where the notion of cyclic algebra over a cyclic operad [GK95] is set up such that the symmetry of the non-degenerate pairing $\kappa : A \otimes A \longrightarrow k$ is included. We can now rephrase the above extension result by saying that a cyclic associative algebra in vector spaces extends to a modular algebra in the sense of [GK98] over the open surface operad.

This note is concerned with a higher categorical analogue of this statement: In [MW23b] the framework of [Cos04] is adapted to define, for any category-valued cyclic operad \mathcal{O} , cyclic \mathcal{O} algebras with values in a symmetric monoidal bicategory \mathcal{S} , up to coherent isomorphism. In the cases of interest in quantum algebra, \mathcal{S} can be for example the symmetric monoidal bicategory Lex^t of finite categories over an algebraically closed field k [EO04] (linear abelian categories with finite-dimensional morphism spaces, enough projective objects, finite many simple objects up to isomorphism and finite length for every object). The 1-morphisms are left exact functors, the 2-morphisms are linear natural transformations, and the monoidal product is the Deligne product \boxtimes . By [MW23b, Theorem 4.12] cyclic associative algebras in Lex[†] are equivalent to *pivotal* Grothendieck-Verdier categories in Lex^f in the sense of [BD13]. A Grothendieck-Verdier category C in Lex^f is a monoidal category in Lex^f equipped with an object $K \in C$, the so-called dualizing ob*ject*, such that the hom functors $\mathcal{C}(K, X \otimes -)$ are representable with representing object $DX \in \mathcal{C}$, i.e. $\mathcal{C}(K, X \otimes -) \cong \mathcal{C}(DX, -)$, such that the functor $D: \mathcal{C}^{\mathrm{opp}} \longrightarrow \mathcal{C}$, called Grothendieck-Verdier duality, is an equivalence. One has canonical isomorphisms $DI \cong K$ and $D^2K \cong K$. A pivotal structure on a Grothendieck-Verdier category \mathcal{C} is a monoidal isomorphism $\omega : \mathrm{id}_{\mathcal{C}} \xrightarrow{\cong} D^2$ such that its component $\omega_K : K \longrightarrow D^2 K$ at the dualizing object coincides with the canonical isomorphism $K \cong D^2 K$. A pivotal finite tensor category in the sense of [EO04, EGNO15], which by definition has a rigid monoidal product (every object has left and right duals that in this case coincide), is an example of a pivotal Grothendieck-Verdier category, but the notion of a pivotal Grothendieck-Verdier category is more general. A rich source of Grothendieck-Verdier categories are vertex operator algebras [ALSW21]. We should mention that the conventions regarding Grothendieck-Verdier duality in [BD13] are dual to ours. The conventions from [BD13] are recovered if we consider cyclic associative algebras in Rex^f, the symmetric monoidal bicategory of finite categories, right exact functors and linear natural transformations. In this note, we treat both situations in parallel.

We show that a cyclic associative Lex^f-valued or Rex^f-valued algebra, i.e. a pivotal Grothendieck-Verdier category in Lex^f or Rex^f, extends uniquely (up to a contractible choice) to a modular algebra over the open surface operad. This is a higher categorical incarnation of the connection between open topological field theories and symmetric Frobenius algebras mentioned in the beginning.

Theorem 2.2 (Classification of open topological field theories in dimension two with values in Lex^t or Rex^f). The 2-groupoid of Lex^f -valued or Rex^f -valued open topological field theories in dimension two is equivalent to the 2-groupoid of pivotal Grothendieck-Verdier categories in Lex^f or Rex^f , respectively. In particular, for every pivotal Grothendieck-Verdier category in Lex^f or Rex^f , the extension to an open topological field theory of dimension two is unique up to equivalence.

The pivotal Grothendieck-Verdier categories appearing in this classification are not necessarily rigid, but we characterize the rigid case topologically in Theorem 3.2 in terms of adjunctions between the value of the corresponding open field theory on different surfaces. It turns out that, just like in [BJS21], it is more natural to only require duals and the pivotal structure to exist for the subcategory of projective objects.

To give further context, we should mention that a modular algebra over the open surface operad is exactly the 'open version' of a modular functor [Seg88, MS89, Tur94, Til98, BK01]. Given a pivotal Grothendieck-Verdier category \mathcal{C} in Lex^f or Rex^f, one obtains the extension to the open surface operad as follows: As cyclic associative algebra, \mathcal{C} extends to a modular algebra over the derived modular envelope of the associative operad, see [Cos04] for the definition of the derived modular envelope and [MW23b, Proposition 7.1] for the extension procedure for algebras. The resulting modular algebra $\widehat{\mathcal{C}}$ is called the *modular extension* of \mathcal{C} . We use a refinement of [Gia11, Theorem B] and [Cos04] to prove that $\widehat{\mathcal{C}}$ is an open topological field theory. The equivalence of Theorem 2.2 then becomes exactly the assignment $\mathcal{C} \mapsto \widehat{\mathcal{C}}$. Concretely, for a surface (for us always compact oriented) with at least one boundary component per connected component and n parametrized intervals embedded in its boundary $\widehat{\mathcal{C}}$ provides us with a left (or right) exact functor $\widehat{\mathcal{C}}(\Sigma; -) : \mathcal{C}^{\boxtimes n} \longrightarrow$ vect to the category of finite-dimensional k-vector spaces. The mapping class group Map(Σ) acts through natural automorphisms of $\widehat{\mathcal{C}}(\Sigma; -)$.

One of the main results of this note concerns exactly these mapping class group representations: Suppose that \mathcal{C} is a pivotal finite tensor category [EO04, EGNO15] (a finite linear category with rigid bilinear monoidal product, simple unit and a pivotal structure) whose two-sided duality we denote by $-^{\vee}$. Consider now the Drinfeld center $Z(\mathcal{C})$ of \mathcal{C} , i.e. the category of pairs of objects $X \in \mathcal{C}$ and half braidings $X \otimes - \cong - \otimes X$. There is a forgetful functor $U : Z(\mathcal{C}) \longrightarrow \mathcal{C}$ whose left and right adjoint we denote by $L : \mathcal{C} \longrightarrow Z(\mathcal{C})$ and $R : \mathcal{C} \longrightarrow Z(\mathcal{C})$, respectively, see [Shi17a] for a description of these adjoints. If \mathcal{C} is spherical in the sense of [DSPS20], then $Z(\mathcal{C})$ is a so-called modular category by [Shi23], a ribbon braided monoidal category with a non-degeneracy condition on the braiding. From a modular category \mathcal{A} , one may build a modular functor $\mathfrak{F}_{\mathcal{A}}$ using the construction from [Lyu95a, Lyu95b, Lyu96]. In particular, we obtain (possibly projective) mapping class group representations $\mathfrak{F}_{\mathcal{A}}(\Sigma; X_1, \ldots, X_n)$ for all surfaces Σ with labels $X_1, \ldots, X_n \in \mathcal{A}$ for the boundary components of Σ . One calls $\mathfrak{F}_{\mathcal{A}}(\Sigma; X_1, \ldots, X_n)$ the space of conformal blocks for \mathcal{A}, Σ and the labels X_1, \ldots, X_n . The modular functor for $Z(\mathcal{C})$ can be built even if \mathcal{C} is not spherical [BW22, Section 8.4] using [MW22]. The following result relates the mapping class group representations coming from $\widehat{\mathcal{C}}$ to the ones coming from $\mathfrak{F}_{Z(\mathcal{C})}$ using the left adjoint $L : \mathcal{C} \longrightarrow Z(\mathcal{C})$:

Theorem 4.3. Let C be a pivotal finite tensor category and Σ a surface with n boundary components, at least one boundary component per connected component, and exactly one interval embedded in each boundary component. Then there are for $X_1, \ldots, X_n \in C$ canonical natural $Map(\Sigma)$ -equivariant isomorphisms

$$\widehat{\mathcal{C}}(\varSigma; X_1, \dots, X_n) \xrightarrow{\cong} \mathfrak{F}_{Z(\mathcal{C})}(\varSigma; LX_1, \dots, LX_n)$$
(1.1)

between the value of the modular extension $\widehat{\mathcal{C}}$ of \mathcal{C} on Σ and the space conformal blocks for $Z(\mathcal{C})$ on Σ .

If \mathcal{C} is a modular category, then the right hand side (1.1) is the space of conformal blocks for the modular category $\overline{\mathcal{C}} \boxtimes \mathcal{C}$ (the bar indicates inversion of the braiding and the ribbon structure; passing to $\overline{\mathcal{C}} \boxtimes \mathcal{C}$ is often called the *combination of left and right movers*), but surprisingly the left hand side can be defined without rigidity. This allows us to prove the following statement:

Corollary 4.4. Let Σ be a surface with n boundary components, at least one per connected component, and let \mathcal{C} be a pivotal Grothendieck-Verdier in Lex^f or Rex^f (not necessarily a rigid one). Then the modular extension $\widehat{\mathcal{C}}(\Sigma; X_1, \ldots, X_n)$ of \mathcal{C} evaluated on Σ and boundary labels in \mathcal{C} carries a representation of the mapping class group of Σ that generalizes the space of conformal blocks after combination of left and right movers in the following sense: If \mathcal{C} is a modular category, then $\widehat{\mathcal{C}}(\Sigma; X_1, \ldots, X_n)$ is Map (Σ) -equivariantly isomorphic to the space of conformal blocks $\mathfrak{F}_{\overline{\mathcal{C}}\boxtimes\mathcal{C}}(\Sigma; \widetilde{X}_1, \ldots, \widetilde{X}_n)$ for $\overline{\mathcal{C}}\boxtimes\mathcal{C}$ on Σ with boundary labels $\widetilde{X}_i = \int^{Y \in \mathcal{A}} Y^{\vee} \boxtimes X_i \otimes Y$ for $1 \leq i \leq n$.

We give a formula for $\widehat{\mathcal{C}}(\Sigma; -)$ in (4.1). Corollary 4.4 is to the best of our knowledge the first construction of mapping class group representations from Grothendieck-Verdier categories that do not necessarily have the property of being rigid. It applies in particular to ribbon Grothendieck-Verdier categories in the sense of [BD13]. The construction of mapping class group representations from Grothendieck-Verdier categories was posed as an open problem in [ALSW21, Section 4]. Note that Corollary 4.4 does not directly generalize Lyubashenko's modular functor construction to Grothendieck-Verdier categories. What we find is instead a partial generalization of the modular functor obtained after combining left and right movers.

We discuss the following further applications of our main results: Suppose that \mathcal{C} is a pivotal finite tensor category that is unimodular, i.e. it has the additional property that its distinguished invertible object $\alpha \in \mathcal{C}$ [ENO04] controlling the quadruple dual via $-^{\vee\vee\vee\vee} \cong \alpha \otimes - \otimes \alpha^{-1}$ is isomorphic to the unit. Then we use factorization homology techniques similar to those in [BW22] to show that, for each free boundary component, $\widehat{\mathcal{C}}(\Sigma; X_1, \ldots, X_n)$ inherits the structure of a module over the algebra of class functions of \mathcal{C} :

Theorem 5.1. Let C be a unimodular pivotal finite tensor category and Σ a connected surface with at least one boundary component, $n \geq 0$ boundary intervals labeled by $X_1, \ldots, X_n \in C$ and $\ell \geq 0$ free boundary circles. Then the vector space $\widehat{C}(\Sigma; X_1, \ldots, X_n)$ associated by the modular extension \widehat{C} of C to Σ and the labels X_1, \ldots, X_n is naturally a module over $\mathsf{CF}(C)^{\otimes \ell}$, the ℓ -th tensor power of the algebra of class functions of C.

Finally, we prove a technical gluing result for the modular functor $\mathfrak{F}_{Z(\mathcal{C})}$ when \mathcal{C} is a pivotal finite tensor category. This gluing result is motivated as follows: Theorem 4.3 compares $\widehat{\mathcal{C}}$ with the restriction of the open-closed modular functor $\mathfrak{F}_{Z(\mathcal{C})}$ to its open part. This means that Theorem 4.3 does not contain a comparison for gluing operations along circles, and indeed this turns out to be a subtle point because open topological field theories do not 'know' about the gluing along boundary circles. As a remedy, we conclude first in Lemma 6.1 that for any surface $\Sigma \in O(n)$ and any disk D embedded in its interior, the embedding $\Sigma \setminus D \longrightarrow \Sigma$ induces a map $p : \widehat{\mathcal{C}}(\Sigma \setminus D; -) \longrightarrow \widehat{\mathcal{C}}(\Sigma; -)$ that we refer to as partial trace (we justify the name in Remark 6.2). The construction of p uses the string-net model for $\widehat{\mathcal{C}}$, and we insist on the point that our construction of p needs rigidity. We then prove:

Theorem 6.3. Let \mathcal{C} be a pivotal finite tensor category and Σ a surface with n + 2 boundary components, $n \geq 1$, at least one per connected component. Consider a sewing $\Sigma \longrightarrow \Sigma'$ that identifies two boundary components of Σ . Then for $X, Y_1, \ldots, Y_n \in \mathcal{C}$, the following diagram commutes:

$$\begin{split} \mathfrak{F}_{Z(\mathcal{C})}(\varSigma; LX^{\vee}, LX, LY_1, \dots, LY_n) & \stackrel{\cong}{\longrightarrow} \widehat{\mathcal{C}}(\varSigma; X^{\vee}, X, Y_1, \dots, Y_n) \\ & & \downarrow^{\widehat{\mathcal{C}} \text{ evaluated on sewing along circle}} \\ \mathfrak{F}_{Z(\mathcal{C})} \text{ evaluated on sewing along circle} \\ & & \widehat{\mathcal{C}}(\varSigma \setminus D; Y_1, \dots, Y_n) \\ & & \downarrow^{\text{partial trace map}} \\ \mathfrak{F}_{Z(\mathcal{C})}(\varSigma'; LY_1, \dots, LY_n) & \stackrel{\cong}{\longrightarrow} \widehat{\mathcal{C}}(\varSigma; Y_1, \dots, Y_n) \;, \end{split}$$

where the horizontal isomorphisms are the ones from Theorem 4.3.

ACKNOWLEDGMENTS. We thank Yang Yang for helpful discussions related to this project. LM gratefully acknowledges support of the Simons Collaboration on Global Categorical Symmetries. Research at Perimeter Institute is supported in part by the Government of Canada through the Department of Innovation, Science and Economic Development and by the Province of Ontario through the Ministry of Colleges and Universities. The Perimeter Institute is in the Haldimand Tract, land promised to the Six Nations. LW gratefully acknowledges support by the ANR project CPJ n°ANR-22-CPJ1-0001-01 at the Institut de Mathématiques de Bourgogne (IMB). The IMB receives support from the EIPHI Graduate School (contract ANR-17-EURE-0002).

2 Classification of categorified open topological field theories

Throughout this article, a surface refers to a compact oriented two-dimensional manifold Σ with an embedding of a disjoint union of standard circles and intervals into the boundary $\partial \Sigma$ of Σ . We call the complement of the embedding the free boundary, the image of the circles the closed boundary, and the image of the intervals the open boundary. This embedding of circles and intervals into the boundary of Σ is called the boundary parametrization. All diffeomorphisms between surfaces are required to preserve the orientation and the boundary parametrization.

We now recall very briefly the definition of the open surface operad O, a groupoid-valued operad. For details, we refer to the definition in [MSWY23, Section 3]; similar definitions of a slightly different flavor can be found in [Cos07, Gia11]. Its groupoid O(n) of arity n operations for $n \ge -1$ (to be read as n inputs and one output) is the groupoid whose objects are connected surfaces with at least one boundary component and a boundary parametrization that embeds n+1 intervals into the boundary, but no circles. The morphisms are mapping classes (isotopy classes of diffeomorphisms preserving the orientation and boundary parametrization), and the operadic composition glues surfaces along the parametrized intervals. The operad O is cyclic in the sense of [GK95], i.e. the inputs can be consistently exchanged with the output, and moreover modular in the sense of [GK98], i.e. it is equipped with self-compositions of operations.

Definition 2.1. A categorified open topological field theory with values in Lex^f or Rex^f is a modular algebra over O with values in Lex^f or Rex^f, respectively.

Explicitly, a categorified open topological field theory has an underlying finite linear category \mathcal{C} and for any $\Sigma \in O(n)$ a left or right exact functor $\mathcal{C}^{\boxtimes (n+1)} \longrightarrow \text{vect}$ with an action of the mapping class group $Map(\Sigma)$ of Σ ; moreover, this assignment is compatible with gluing. This is equivalent to the usual definition of open topological field theories as symmetric monoidal functors from the $(\infty, 1)$ -category $Bord_2^o$ of open bordisms to Lex^f or Rex^f .

We will now state and prove the classification result for categorified open topological field theories in dimension two:

Theorem 2.2 (Classification of open topological field theories in dimension two with values in Lex^t or Rex^f). The 2-groupoid of Lex^f -valued or Rex^f -valued open topological field theories in dimension two is equivalent to the 2-groupoid of pivotal Grothendieck-Verdier categories in Lex^f or Rex^f , respectively. In particular, for every pivotal Grothendieck-Verdier category in Lex^f or Rex^f , the extension to an open topological field theory of dimension two is unique up to equivalence.

Proof. A cyclic operad can be 'freely completed' to a modular one via the (derived) modular envelope of [Cos04]. The version of this construction adapted to Cat-valued operads [MW23b, Section 7.1], when applied to the cyclic associative operad As, leads to the modular operad U_f As that, after applying the nerve functor B and geometric realization |-|, yields a topological operad $|BU_f$ As|. It comes by [Gia11, Theorem B] with a map $|BU_f$ As| $\longrightarrow |BO|$ that induces a bijection on π_0 (a bijection on the set of path components) and a homotopy equivalence on all components except for the annulus without boundary intervals. For the modular operad $|BU_f$ As|, the component of the annulus without boundary intervals is the dihedral homology of the algebra As(1) of unary operations of the associative operad by [MW23a, Theorem 3.6]. But As(1) is just a point, which implies that its dihedral homology is the classifying space BO(2) of the two-dimensional orthogonal group. The topological group O(2) is the group of diffeomorphisms of the annulus, seen as operation of the operad O with no boundary intervals. Its mapping class group is the group $\pi_0(O(2)) \cong \mathbb{Z}_2$. The restriction of $|BU_f$ As| $\longrightarrow |BO|$ to the annulus component is now the projection map $BO(2) \longrightarrow B\pi_0(O(2))$. Clearly, this map induces an equivalence after applying the fundamental groupoid functor. Therefore, we obtain an equivalence

$$\Pi |B\mathsf{U}_{\mathsf{f}} \mathsf{As}| \xrightarrow{\simeq} \Pi |B\mathsf{O}| \simeq \mathsf{O}$$

$$\tag{2.1}$$

of groupoid-valued modular operads. By [MW23a, Theorem 4.2] we have moreover canonical inverse equivalences

$$\mathsf{CycAlg}(\mathsf{As},\mathcal{S}) \xrightarrow[]{\text{modular extension}}_{\text{restriction}} \mathsf{ModAlg}\left(\Pi|B\mathsf{U}_{\int}\mathsf{As}|,\mathcal{S}\right)$$
(2.2)

between the 2-groupoid of cyclic As-algebras with values in any symmetric monoidal bicategory S and modular $\Pi |BU_f As|$ -algebras in S. Once we specify S to be Lex^f or Rex^f, we find on the right hand side categorified open topological field theories thanks to (2.1) while on the left hand side we find pivotal Grothendieck-Verdier categories by [MW23b, Theorem 4.12].

3 A TOPOLOGICAL CHARACTERIZATION OF RIGIDITY

The pivotal Grothendieck-Verdier categories in Theorem 2.2 are not necessarily rigid. In this section, we will characterize those open field theories corresponding to rigid categories.

Let S be a symmetric monoidal bicategory with monoidal product \boxtimes . A rigid algebra [Dé23, Definition 2.1.1] in S is an associative algebra A in S, up to coherent isomorphism, such that the multiplication $\mu : A \boxtimes A \longrightarrow A$ and the unit $\eta : I \longrightarrow A$ have right adjoints $\delta : A \longrightarrow A \boxtimes A$ and $\varepsilon : A \longrightarrow I$, respectively, such that the canonical lax bimodule structure on δ is strong. This allows us to define the non-degenerate pairing $\kappa : A \boxtimes A \xrightarrow{\mu} A \xrightarrow{\varepsilon} I$ on A with copairing $\Delta : I \xrightarrow{\eta} A \xrightarrow{\delta} A \boxtimes A$. A symmetric rigid algebra in S is a cyclic associative algebra in S whose underlying algebra and non-degenerate pairing come from a rigid algebra. Explicitly, a symmetric rigid algebra is a rigid algebra equipped with a natural isomorphism $\Sigma : \kappa \longrightarrow \kappa \circ \sigma$ (here $\sigma : A \boxtimes A \longrightarrow A \boxtimes A$ is the symmetric braiding) squaring to the identity and satisfying [MW23b, Definition 4.1 (H1)].

Following [BJS21], we call a monoidal category $C \in \mathsf{Rex}^{\mathsf{f}} p$ -rigid if all its projective objects have both a left and right dual which is projective. A central result [BJS21, Definition-Proposition 4.1] shows that p-rigid categories are exactly the rigid algebras in $\mathsf{Rex}^{\mathsf{f}}$. In particular, the tensor product of a p-rigid monoidal category C is exact and hence preserves projective objects. Based on this observation, we can define a *p*-pivotal category as a p-rigid category for which in addition the subcategory of projective objects is equipped with a monoidal (non-unital) natural isomorphism $-^{\vee\vee} \longrightarrow \mathrm{id}$.

Proposition 3.1. Symmetric rigid algebras in Rex^f are p-pivotal categories.


Proof. From [BJS21, Definition-Proposition 4.1], we know that rigid algebras are exactly p-rigid categories. Hence, we only have to work out the additional structure coming from the symmetry isomorphism Σ . The natural isomorphism Σ of a rigid symmetric algebra is uniquely fixed by its value on the subcategory $\operatorname{Proj} \mathcal{C}$ of projective objects, which is a natural isomorphism between the two functors $(\operatorname{Proj} (\mathcal{C} \boxtimes \mathcal{C}))^{\operatorname{opp}} \longrightarrow \operatorname{vect}$ sending $P \boxtimes Q$ for $P, Q \in \operatorname{Proj} \mathcal{C}$ to $\mathcal{C}(P \otimes Q, I)$ and $\mathcal{C}(Q \otimes P, I)$, respectively. Thanks to duality, this is equivalent to a natural transformation between the functors $\mathcal{C}(P, Q^{\vee})$ and $\mathcal{C}(P, ^{\vee}Q)$ that via the Yoneda Lemma amount to a natural isomorphism $Q^{\vee} \longrightarrow {}^{\vee}Q$. The condition [MW23b, Definition 4.1 (H1)] on Σ implies that this is a pivotal structure.

Theorem 3.2. Let C be a cyclic associative algebra in S, and \hat{C} its associated open topological field theory. Then the following are equivalent:

- (i) There are adjunctions $\widehat{\mathcal{C}}(\diamondsuit) \dashv \widehat{\mathcal{C}}(\bigtriangledown)$ and $\widehat{\mathcal{C}}(\bigtriangledown) \dashv \widehat{\mathcal{C}}(\lhd)$ such that the natural isomorphisms making $\widehat{\mathcal{C}}(\backsim)$ a bimodule functor agree with those coming from the adjunction.
- (ii) C is a symmetric rigid algebra.

If $S = \text{Rex}^{f}$, then (ii) says exactly that C is p-pivotal by Proposition 3.1.

Proof. If \mathcal{C} is a symmetric rigid algebra, the monoidal product $\widehat{\mathcal{C}}(\diamondsuit)$ has a right adjoint $\delta: \mathcal{C} \longrightarrow \mathcal{C} \boxtimes \mathcal{C}$, and this adjoint is given by $\widehat{\mathcal{C}}(\triangleleft)$. Indeed, the following computation in the graphical calculus in symmetric monoidal bicategories (see e.g. [MW23b, Remark 2.6]) shows that $\widehat{\mathcal{C}}(\triangleleft)$ is given by



The second step uses the strong bimodule structure on δ . Therefore, (ii) implies (i).

Condition (i) implies that C is rigid: Both the multiplication and the unit have a right adjoint by definition, and the lax bimodule structure on the right adjoint of the multiplication is strong because it is required to be given by isomorphisms.

4 Comparison with the string-net construction

In this section, we compare the modular extension $\widehat{\mathcal{C}}$ of a pivotal finite tensor category \mathcal{C} with the string-net construction of [MSWY23]. This will be the main technical tool for the proof Theorem 4.3.

The string-net construction of Levin-Wen [LW05] can be used to construct categorified openclosed topological field theories. In the semisimple case, it was refined further in [Kir11] and more recently [Bar22]. A generalization of string-net techniques to pivotal bicategories is given in [FSY23]. In [MSWY23], string-nets are used to build an open-closed categorified topological field theory from a not necessarily semisimple pivotal finite tensor category C, and it is shown that its closed part is the Lyubashenko modular functor [Lyu95a, Lyu95b, Lyu96] for Z(C). We briefly review the construction of the categorified string-net topological field theory closely following the presentation in [MSWY23, Section 2].

A graph Γ in a surface Σ is a finite abstract graph Γ together with an embedding of its geometric realization $|\Gamma|$ into Σ mapping the endpoints of its leaves (half edges attached to only one vertex), and only those, to the open or closed boundary (see Figure 1 for an example).

Let \mathcal{C} be a pivotal finite tensor category. Informally, the \mathcal{C} -string-net space associated to a surface Σ consists of graphs in Σ whose edges and vertices are labeled with objects and morphisms of $\operatorname{Proj}\mathcal{C}$, respectively, modulo relations holding in \mathcal{C} . This idea is implemented via a colimit: We write \mathcal{C} -Graphs(Σ) for the category whose objects consist of graphs Γ in Σ together with a labeling of its edges by objects of $\operatorname{Proj}\mathcal{C}$ and an orientation; we denote the labeling by \underline{X} , so that the entire object is a pair (Γ, \underline{X}). Morphisms are generated by replacing graphs in disks by corollas. We would like to avoid reproducing here the full definition with all its technical details. The reader can find an example of such a replacement with a corolla within a disk in Figure 1. The details

are given in [MSWY23, Definition 2.1]. A boundary label for Σ is a finite collection of points in the open and closed boundary labeled each with an object $P \in \operatorname{Proj} \mathcal{C}$ and a sign + or - encoding whether edges ending at the point are incoming or outgoing; we agree that + corresponds to outwards pointing orientation. A \mathcal{C} -labeled graph in Σ has a corresponding boundary label. For a fixed boundary label B, we denote by \mathcal{C} -Graphs(Σ ; B) the full subcategory of \mathcal{C} -Graphs(Σ) of graphs restricting to B on $\partial \Sigma$. There is a functor

$$\mathbb{E}^{\Sigma,B}_{\mathcal{C}}: \mathcal{C}\text{-}\mathsf{Graphs}(\Sigma;B) \longrightarrow \mathsf{Vect}$$

to the category Vect of (not necessarily finite-dimensional) vector spaces over k sending a C-labeled graph to the vector space of compatible labels of the vertices with morphisms in C. The evaluation of the functor on morphisms is induced by the graphical calculus for C. We refer to [MSWY23, Definition 2.3] for more details. To see that the functor is well-defined, we need C to be pivotal.




Figure 1: An illustration of an object of C-Graphs(Σ) and a morphism corresponding to a disk replacement. The labels a, \ldots, o are elements of Proj C. We have not drawn is orientation associated to every edge. The free boundary of Σ is on the right and drawn in red.

Definition 4.1 ([MSWY23, Definition 2.3] following [FSY23, Section 3.2]). Let C be a pivotal finite tensor category, Σ a surface and B a C-boundary label for Σ . The string-net space $\operatorname{sn}_{\mathcal{C}}(\Sigma; B)$ is defined as the colimit

$$\mathsf{sn}_{\mathcal{C}}(\varSigma;B) := \operatornamewithlimits{colim}_{(\varGamma,\underline{X})\in\mathcal{C}\text{-}\mathsf{Graphs}(\varSigma;B)} \mathbb{E}_{\mathcal{C}}^{\varSigma,B}(\varGamma,\underline{X})$$

It is explained in [MSWY23, Remark 2.17] that $\operatorname{sn}_{\mathcal{C}}(\Sigma; B)$ is isomorphic to the two-dimensional admissible skein module for \mathcal{C} and Σ in the sense of [CGPM23]. The mapping class group of Σ geometrically acts on the string-net space.

Using string-net spaces, we can assign a k-linear category $\operatorname{sn}_{\mathcal{C}}(S)$ to every oriented 1-dimensional manifold S. An object of $\operatorname{sn}_{\mathcal{C}}(S)$ is a collection of points in S labeled with objects of $\operatorname{Proj}\mathcal{C}$ such that there is at least one point in every connected component of S. The space of morphisms from \underline{X} to \underline{Y} is the vector space $\operatorname{sn}_{\mathcal{C}}([0,1] \times S, \underline{X} \sqcup \underline{Y})$ with the points in \underline{X} labeled by – and those in \underline{Y} by +. Composition is defined by gluing of cylinders.

The string-net construction naturally extends to a categorified topological field theory with values in the bicategory $\mathsf{Bimod}^{\mathsf{f}}$ of k-linear pre-finite categories (see [MSWY23, Section 5]), bimodules (a

bimodule from \mathcal{C} to \mathcal{D} is a linear functor $M : \mathcal{C} \otimes \mathcal{D}^{\text{opp}} \longrightarrow \text{vect}$), and natural transformations. The composition of two bimodules $M : \mathcal{C} \otimes \mathcal{D}^{\text{opp}} \longrightarrow \text{vect}$ and $M' : \mathcal{D} \otimes \mathcal{E}^{\text{opp}} \longrightarrow \text{vect}$ is defined by the coend $\int^{d \in \mathcal{D}} M(-,d) \otimes M'(d,-)$. Explicitly, the open-closed topological field theory built from string-nets assigns to a surface Σ with incoming boundary S and outgoing boundary S'the bimodule $\operatorname{sn}_{\mathcal{C}}(\Sigma, -, -) : \operatorname{sn}_{\mathcal{C}}(S) \otimes \operatorname{sn}_{\mathcal{C}}(S')^{\operatorname{opp}} \longrightarrow \text{vect}$. This Bimod^f-valued field theory can be turned into a Rex^f-valued one by applying the finite free cocompletion, i.e. the equivalence $\operatorname{Bimod}^{f} \simeq \operatorname{Rex}^{f}$ of symmetric monoidal bicategories [MSWY23, Section 5] sending a pre-finite linear category \mathcal{A} to the category of linear functors $\mathcal{A}^{\operatorname{opp}} \longrightarrow \operatorname{vect}$, and a bimodule to the right exact functor it induces. We denote the Rex^{f} -valued open-closed topological field theory induced by $\operatorname{sn}_{\mathcal{C}}$ by $\operatorname{SN}_{\mathcal{C}}$.

Theorem 4.2. Let C be a pivotal finite tensor category. Then the following constructions agree, up to canonical equivalence, as open Rex^f-valued topological field theories:

- (i) The modular extension $\widehat{\mathcal{C}}$ of \mathcal{C} .
- (ii) The open part of the open-closed string-net topological field theory $SN_{\mathcal{C}}$.

Proof. By Theorem 2.2 it suffices to show that the two constructions each produce a $\mathsf{Rex}^{\mathsf{t}}$ -valued open topological field theory, i.e. a $\mathsf{Rex}^{\mathsf{f}}$ -valued modular O-algebra, whose restriction to disks with intervals embedded in its boundary is \mathcal{C} , seen as a cyclic associative algebra. For (i), this holds by (2.2). For $\mathsf{SN}_{\mathcal{C}}$ in (ii), it follows from [MSWY23, Theorem 5.9] that we obtain a $\mathsf{Rex}^{\mathsf{f}}$ -valued open-closed modular functor. After restriction to the open part, we obtain an open topological field theory whose restriction to disks is \mathcal{C} , as cyclic associative algebra, as explained in [MSWY23, Section 6]. This proves the equivalence of (i) and (ii).

Theorem 4.3. Let C be a pivotal finite tensor category and Σ a surface with n boundary components, at least one boundary component per connected component, and exactly one interval embedded in each boundary component. Then there are for $X_1, \ldots, X_n \in C$ canonical natural $Map(\Sigma)$ -equivariant isomorphisms

$$\widehat{\mathcal{C}}(\Sigma; X_1, \dots, X_n) \xrightarrow{\cong} \mathfrak{F}_{Z(\mathcal{C})}(\Sigma; LX_1, \dots, LX_n)$$

between the value of the modular extension $\widehat{\mathcal{C}}$ of \mathcal{C} on Σ and the space conformal blocks for $Z(\mathcal{C})$ on Σ .

Proof. We have natural $Map(\Sigma)$ -equivariant isomorphisms

$$\widehat{\mathcal{C}}(\varSigma; X_1, \dots, X_n) \xrightarrow{\cong} \mathsf{SN}_{\mathcal{C}}(\varSigma; X_1, \dots, X_n) \xrightarrow{\cong} \mathfrak{F}_{Z(\mathcal{C})}(\varSigma; LX_1, \dots, LX_n)$$

where the first isomorphism is a consequence of Theorem 4.2, and the second follows from [MSWY23, Theorem 7.1]. $\hfill \Box$

Corollary 4.4. Let Σ be a surface with n boundary components, at least one per connected component, and let \mathcal{C} be a pivotal Grothendieck-Verdier in Lex^f or Rex^f (not necessarily a rigid one). Then the modular extension $\widehat{\mathcal{C}}(\Sigma; X_1, \ldots, X_n)$ of \mathcal{C} evaluated on Σ and boundary labels in \mathcal{C} carries a representation of the mapping class group of Σ that generalizes the space of conformal blocks after combination of left and right movers in the following sense: If \mathcal{C} is a modular category, then $\widehat{\mathcal{C}}(\Sigma; X_1, \ldots, X_n)$ is $Map(\Sigma)$ -equivariantly isomorphic to the space of conformal blocks $\mathfrak{F}_{\overline{\mathcal{C}}\boxtimes\mathcal{C}}(\Sigma; \widetilde{X}_1, \ldots, \widetilde{X}_n)$ for $\overline{\mathcal{C}}\boxtimes\mathcal{C}$ on Σ with boundary labels $\widetilde{X}_i = \int^{Y \in \mathcal{A}} Y^{\vee} \boxtimes X_i \otimes Y$ for $1 \leq i \leq n$.

If Σ is connected of genus g with one boundary component (for simplicity), then (in the Lex^f-

version) one can show

$$\widehat{\mathcal{C}}(\Sigma;X) \cong \mathcal{C}\left(K, X \otimes \left(\otimes^{(4)} \left(\int^{Y,Y' \in \mathcal{C}} Y \boxtimes Y' \boxtimes DY \boxtimes DY'\right)\right)^{\otimes g}\right),\tag{4.1}$$

using a ribbon graph presentation of Σ and the techniques of [MW23b, Example 7.3]; here $\otimes^{(4)}$: $\mathcal{C}^{\boxtimes 4} \longrightarrow \mathcal{C}$ is the monoidal product seen as arity four operation.

Proof of Corollary 4.4. We obtain the mapping class group representations thanks to Theorem 2.2. In order to apply this result, we place exactly one interval on each boundary circle. Assume now that \mathcal{C} is modular: Then \mathcal{C} , by the results of [Shi19], is factorizable in the sense of [ENO04], i.e. the monoidal product induces a ribbon equivalence $\overline{\mathcal{C}} \boxtimes \mathcal{C} \xrightarrow{\simeq} Z(\mathcal{C})$. Now the statement follows from Theorem 4.3.

5 The action of the algebra of class functions

For a finite tensor category \mathcal{C} , one may define an algebra of class functions. We give here the definition from [Shi17b], where also the connection to more classical notions of class functions is made. We denote by $R : \mathcal{C} \longrightarrow Z(\mathcal{C})$ the right adjoint to the forgetful functor $U : Z(\mathcal{C}) \longrightarrow \mathcal{C}$ from the Drinfeld center $Z(\mathcal{C})$ of \mathcal{C} to \mathcal{C} . Then the algebra $\mathsf{CF}(\mathcal{C})$ of class functions of \mathcal{C} is the vector space $\mathcal{C}(\mathbb{A}, I)$, where $\mathbb{A} = \int_{X \in \mathcal{C}} X^{\vee} \otimes X = URI$ is the canonical end of \mathcal{C} . The algebra structure comes from the fact that $\mathcal{C}(\mathbb{A}, I) = \mathcal{C}(URI, I) \cong \mathsf{End}_{Z(\mathcal{C})}(RI)$ can be identified with the endomorphisms of RI.

Theorem 5.1. Let C be a unimodular pivotal finite tensor category and Σ a connected surface with at least one boundary component, $n \geq 0$ boundary intervals labeled by $X_1, \ldots, X_n \in C$ and $\ell \geq 0$ free boundary circles. Then the vector space $\widehat{C}(\Sigma; X_1, \ldots, X_n)$ associated by the modular extension \widehat{C} of C to Σ and the labels X_1, \ldots, X_n is naturally a module over $\mathsf{CF}(C)^{\otimes \ell}$, the ℓ -th tensor power of the algebra of class functions of C.

Proof. We begin with the observation of [BW22, Section 4] that the modular extension descends to factorization homology, but instead of using this fact for the modular extension of cyclic framed E_2 -algebras, we profit from this principle one dimension lower. Let us give the details: Denote by S the free boundary of Σ consisting of ℓ circles. For any finite set J and an embedding $\varphi : (0,1)^{\sqcup J} \longrightarrow S$, we denote by Σ^{φ} the surface Σ but with the intervals that are part of φ as additional parametrized intervals. The modular extension $\widehat{\mathcal{C}}$ of \mathcal{C} then provides us with a functor

$$\widehat{\mathcal{C}}(\Sigma^{\varphi}; -): \mathcal{C}^{\boxtimes J} \boxtimes \mathcal{C}^{\boxtimes n} \longrightarrow \text{vect} .$$
(5.1)

In other words, we can choose to embed a family of intervals into the free boundary and evaluate with $\hat{\mathcal{C}}$. The naturality in the embedding φ implies that the functors (5.1) descend to factorization homology and give us a functor

$$\varPhi_{\mathcal{C}}(\varSigma): \left(\int_{S} \mathcal{C}\right) \boxtimes \mathcal{C}^{\boxtimes n} \longrightarrow \mathsf{vect}$$

with a canonical isomorphism $\Phi_{\mathcal{C}}(\Sigma; \mathcal{O}_S, -) \cong \widehat{\mathcal{C}}(\Sigma; -) : \mathcal{C}^{\boxtimes n} \longrightarrow \text{vect}$ for the quantum character sheaf $\mathcal{O}_S \in \int_S \mathcal{C}$ from [BZBJ18, Section 5.1]. In particular,

$$\widehat{\mathcal{C}}(\Sigma; X_1, \dots, X_n) \cong \Phi_{\mathcal{C}}(\Sigma; \mathcal{O}_S, X_1, \dots, X_n)$$

as vector spaces. But now clearly $\widehat{\mathcal{C}}(\Sigma; X_1, \ldots, X_n)$ becomes a module over the endomorphism algebra $\operatorname{End}_{\int_S \mathcal{C}}(\mathcal{O}_S)$ of \mathcal{O}_S . Note that this is nothing but a one-dimensional version of the skein algebra [GJS23, Definition 2.6]; we mention this as a helpful connection for some readers, but it is by no means needed for the understanding.

We will now finish the proof by showing that

$$\operatorname{End}_{\int_{\mathcal{C}} \mathcal{C}}(\mathcal{O}_S) \cong \operatorname{CF}(\mathcal{C})^{\otimes \ell} \tag{5.2}$$

as algebras. Whenever $S = S' \sqcup S''$, we have an equivalence $\int_S \mathcal{C} \simeq \int_{S'} \mathcal{C} \boxtimes \int_{S''} \mathcal{C}$ under which \mathcal{O}_S corresponds to $\mathcal{O}_{S'} \boxtimes \mathcal{O}_{S''}$; this is just the monoidality of factorization homology with respect to disjoint union. Therefore, we can prove (5.2) for $S = \mathbb{S}^1$, i.e. $\ell = 1$. Up to this point, the proof would apply to any pivotal Grothendieck-Verdier category, but $\operatorname{End}_{\int_{\mathbb{S}^1} \mathcal{C}}(\mathcal{O}_{\mathbb{S}^1}) \cong \operatorname{CF}(\mathcal{C})$ will need that \mathcal{C} is actually a unimodular pivotal finite tensor category. Indeed, thanks to \mathcal{C} being a pivotal finite tensor category and the comparison result Theorem 4.2, $\widehat{\mathcal{C}}$ extends to an open-closed modular functor, namely the one built in [MSWY23] using string-nets, that associates to the circle $\int_{\mathbb{S}^1} \mathcal{C}$ (that can be understood as finite free cocompletion of the string-net category of \mathbb{S}^1) or, equivalently, the Drinfeld center $Z(\mathcal{C})$. The equivalence $\int_{\mathbb{S}^1} \mathcal{C} \simeq Z(\mathcal{C})$ as linear categories is induced by the left adjoint $L : \mathcal{C} \longrightarrow Z(\mathcal{C})$ to the forgetful functor $U : Z(\mathcal{C}) \longrightarrow \mathcal{C}$, see [MSWY23, Theorem 5.9]. In particular, under this equivalence $\mathcal{O}_{\mathbb{S}^1}$ corresponds to $LI \in Z(\mathcal{C})$, thereby proving that $\operatorname{End}_{\int_{\mathbb{S}^1} \mathcal{C}}(\mathcal{O}_{\mathbb{S}^1}) \cong \operatorname{End}_{Z(\mathcal{C})}(LI)$. Since \mathcal{C} is unimodular, L is also right adjoint to U [Shi17a]. This leaves us with the isomorphism of algebras $\operatorname{End}_{\int_{\mathbb{S}^1} \mathcal{C}}(\mathcal{O}_{\mathbb{S}^1}) \cong \operatorname{End}_{Z(\mathcal{C})}(RI) = \operatorname{CF}(\mathcal{C})$. This finishes the proof.

6 A Gluing result involving the partial trace

In this section, we use the results of the previous sections to prove a gluing result for the spaces of conformal blocks for a Drinfeld center. More precisely, we decompose the gluing along a circle into the gluing along an interval and the partial trace map.

Lemma 6.1. Let C be a pivotal finite tensor category and $\Sigma \in O(n)$. Then the extension of \widehat{C} to an open-closed modular functor induces for any disk D embedded in the interior of Σ a natural map

$$p:\widehat{\mathcal{C}}(\Sigma \setminus D; -) \longrightarrow \widehat{\mathcal{C}}(\Sigma; -)$$

coming from the embedding $\Sigma \setminus D \subset \Sigma$. We call p the partial trace map associated to \mathcal{C} , Σ and D.

Proof. By their very definition string-nets are functorial with respect to embeddings of surfaces. In particular, the inclusion $\Sigma \setminus D \subset \Sigma$ induces a map $\mathsf{SN}_{\mathcal{C}}(\Sigma \setminus D; -) \longrightarrow \mathsf{SN}_{\mathcal{C}}(\Sigma; -)$. Now the statement follows from Theorem 4.2.

Remark 6.2. Let us justify the use of the term 'partial trace' in Lemma 6.1 by showing that in a specific special case it actually reduces to the partial trace: Suppose that Σ is itself a disk, then $\Sigma \setminus D$ is an annulus A. With excision [MSWY23, Theorem 3.2 & Proposition 4.2] for stringnets we see (in the Lex^f-version) that $SN_{\mathcal{C}}(A; -) \cong \mathcal{C}(I, \mathbb{F} \otimes -)$ for the coend $\mathbb{F} = \int^{X \in \mathcal{C}} X^{\vee} \otimes X$ and $SN_{\mathcal{C}}(\Sigma; -) \cong \mathcal{C}(I, -)$ and that p in this case is induced by the map $\mathbb{F} \longrightarrow I$ coming from the evaluations $X^{\vee} \otimes X \longrightarrow I$. If we see a map $f: X \longrightarrow X \otimes Y$ as a map $I \longrightarrow X^{\vee} \otimes X \otimes Y \longrightarrow \mathbb{F} \otimes Y$ and hence as an element $f \in SN_{\mathcal{C}}(A; Y)$ (denoted by the same symbol by abuse of notation) then p(f) is the partial trace of f taken over X. Note that the partial trace coincides also the counit of the adjunction $\widehat{\mathcal{C}}(\Sigma) \dashv \widehat{\mathcal{C}}(\varsigma)$ that we have at our disposal since \mathcal{C} is rigid (Theorem 3.2). **Theorem 6.3.** Let \mathcal{C} be a pivotal finite tensor category and Σ a surface with n + 2 boundary components, $n \geq 1$, at least one per connected component. Consider a sewing $\Sigma \longrightarrow \Sigma'$ that identifies two boundary components of Σ . Then for $X, Y_1, \ldots, Y_n \in \mathcal{C}$, the following diagram commutes:

$$\begin{split} \mathfrak{F}_{Z(\mathcal{C})}(\varSigma; LX^{\vee}, LX, LY_1, \dots, LY_n) & \stackrel{\cong}{\longrightarrow} \widehat{\mathcal{C}}(\varSigma; X^{\vee}, X, Y_1, \dots, Y_n) \\ & \downarrow^{\widehat{\mathcal{C}} \text{ evaluated on sewing along circle}} \\ \mathfrak{F}_{Z(\mathcal{C})} \text{ evaluated on sewing along circle} \\ & \widehat{\mathcal{C}}(\varSigma \setminus D; Y_1, \dots, Y_n) \\ & \downarrow^{\text{partial trace map}} \\ \mathfrak{F}_{Z(\mathcal{C})}(\varSigma'; LY_1, \dots, LY_n) & \stackrel{\cong}{\longrightarrow} \widehat{\mathcal{C}}(\varSigma; Y_1, \dots, Y_n) , \end{split}$$

where the horizontal isomorphisms are the ones from Theorem 4.3.

Proof. Since $\mathfrak{F}_{Z(\mathcal{C})} \simeq \mathsf{SN}_{\mathcal{C}}$ as modular functors [MSWY23, Theorem 7.1], the gluing operation for the modular functor $\mathfrak{F}_{Z(\mathcal{C})}$ is exactly the gluing for string-nets which by the excision result [MSWY23, Theorem 3.2] for string-nets is the composition of gluing along the boundary intervals (here by assumption exactly one per boundary component) and the embedding of the string-nets; this is depicted schematically in Figure 2.




Figure 2: The gluing along a boundary circle can be decomposed into the gluing along a boundary interval and an embedding.

But the gluing of the string-nets along an interval can, when composed with the isomorphism $\mathsf{SN}_{\mathcal{C}}(\varSigma; -) \cong \widehat{\mathcal{C}}(\varSigma; -)$ from Theorem 4.2, agrees with the gluing along intervals for the open modular functor $\widehat{\mathcal{C}}$ while the embedding of the string-nets along $\varSigma \setminus D \longrightarrow \varSigma$ yields, when transferred to $\widehat{\mathcal{C}}$, exactly the partial trace.

References

- [ALSW21] R. Allen, S. Lentner, C. Schweigert, and S. Wood. Duality structures for module categories of vertex operator algebras and the Feigin Fuchs boson. arXiv:2107.05718 [math.QA], 2021.
- [Bar22] B. Bartlett. Three-dimensional TQFTs via string-nets and two-dimensional surgery. arXiv:2206.13262 [math.QA], 2022.
- [BD13] M. Boyarchenko and V. Drinfeld. A duality formalism in the spirit of Grothendieck and Verdier. *Quantum Top.*, 4(4):447–489, 2013.
- [BJS21] A. Brochier, D. Jordan, and N. Snyder. On dualizability of braided tensor categories. Compositio Math., 157(3):435–483, 2021.
- [BK01] B. Bakalov and A. Kirillov. Lectures on tensor categories and modular functors, volume 21 of University Lecture Series. Am. Math. Soc., 2001.
- [BW22] A. Brochier and L. Woike. A classification of modular functors via factorization homology. arXiv:2212.11259 [math.QA], 2022.
- [BZBJ18] D. Ben-Zvi, A. Brochier, and D. Jordan. Integrating quantum groups over surfaces. J. Top., 11(4):874–917, 2018.
- [CGPM23] F. Costantino, N. Geer, and B. Patureau-Mirand. Admissible skein modules. arXiv:2302.04493 [math.GT], 2023.
- [Cos04] K. Costello. The A-infinity operad and the moduli space of curves. arXiv:math/0402015 [math.AG], 2004.
- [Cos07] K. Costello. Topological conformal field theories and Calabi-Yau categories. Adv. Math., 210:165–214, 2007.
- [Dé23] T. Décoppet. Rigid and separable algebras in fusion 2-categories. Adv. Math., 419:108967, 2023.
- [DSPS20] C. L. Douglas, C. Schommer-Pries, and N. Snyder. Dualizable Tensor Categories. Mem. Amer. Math. Soc., 2020.
- [EGN015] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik. Tensor categories, volume 205 of Math. Surveys Monogr. Am. Math. Soc., 2015.
- [ENO04] P. Etingof, D. Nikshych, and V. Ostrik. An analogue of Radford's S⁴ formula for finite tensor categories. Int. Math. Res. Not., 54:2915–2933, 2004.
- [EO04] P. Etingof and V. Ostrik. Finite tensor categories. Mosc. Math. J., 4(3):627–654, 2004.
- [FSY23] J. Fuchs, C. Schweigert, and Y. Yang. String-net models for pivotal bicategories. arXiv:2302.01468 [math.QA], 2023.
- [Gia11] J. Giansiracusa. The framed little 2-discs operad and diffeomorphisms of handlebodies. J. Top., 4(4):919–941, 2011.
- [GJS23] S. Gunningham, D. Jordan, and P. Safronov. The finiteness conjecture for skein modules. Invent. Math., 232:301–363, 2023.
- [GK95] E. Getzler and M. Kapranov. Cyclic operads and cyclic homology. In R. Bott and S.-T. Yau, editors, Conference proceedings and lecture notes in geometry and topology, pages 167–201. Int. Press, 1995.
- [GK98] E. Getzler and M. Kapranov. Modular operads. Compositio Math., 110:65–126, 1998.
- [Kir11] A. Kirillov. String-net model of Turaev-Viro invariants. arXiv:1106.6033 [math.AT], 2011.
- [LP08] A. D. Lauda and H. Pfeiffer. Open-closed strings: Two-dimensional extended TQFTs and Frobenius algebras. Topology Appl., 155(7):623–666, 2008.
- [LW05] M. A. Levin and X.-G. Wen. String-net condensation: A physical mechanism for topological phases. Phys. Rev. B, 71:045110, 2005.
- [Lyu95a] V. V. Lyubashenko. Invariants of 3-manifolds and projective representations of mapping class groups via quantum groups at roots of unity. Comm. Math. Phys., 172:467–516, 1995.
- [Lyu95b] V. V. Lyubashenko. Modular transformations for tensor categories. J. Pure Appl. Alg., 98:279–

327, 1995.

- [Lyu96] V. V. Lyubashenko. Ribbon abelian categories as modular categories. J. Knot Theory and its Ramif., 5:311–403, 1996.
- [MS89] G. Moore and N. Seiberg. Classical and Quantum Conformal Field Theory. Comm. Math. Phys., 123:177–254, 1989.
- [MSWY23] L. Müller, C. Schweigert, L. Woike, and Y. Yang. The Lyubashenko modular functor for Drinfeld centers via non-semisimple string-nets. arXiv:2312.140109 [math.QA], 2023.
- [MW22] L. Müller and L. Woike. The distinguished invertible object as ribbon dualizing object in the Drinfeld center. arXiv:2212.07910 [math.QA], 2022.
- [MW23a] L. Müller and L. Woike. Classification of Consistent Systems of Handlebody Group Representations. Int. Math. Res. Not., rnad178, 2023.
- [MW23b] L. Müller and L. Woike. Cyclic framed little disks algebras, Grothendieck-Verdier duality and handlebody group representations. *Quart. J. Math.*, 74(1):163–245, 2023.
- [Seg88] G. Segal. Two-dimensional conformal field theories and modular functors. In IX International Conference on Mathematical Physics (IAMP), 1988.
- [Shi17a] K. Shimizu. On unimodular finite tensor categories. Int. Math. Res. Not., 2017(1):277–322, 2017.
- [Shi17b] K. Shimizu. The monoidal center and the character algebra. J. Pure Appl. Alg., 221(9):2338– 2371, 2017.
- [Shi19] K. Shimizu. Non-degeneracy conditions for braided finite tensor categories. Adv. Math., 355:106778, 2019.
- [Shi23] K. Shimizu. Ribbon structures of the Drinfeld center of a finite tensor category. Kodai Math. J., 46(1):75–114, 2023.
- [Til98] U. Tillmann. S-Structures for k-Linear Categories and the Definition of a Modular Functor. J. London Math. Soc., 58(1):208–228, 1998.
- [Tur94] V. G. Turaev. Quantum Invariants of Knots and 3-Manifolds, volume 18 of Studies in Math. De Gruyter, 1994.
- [WW16] N. Wahl and C. Westerland. Hochschild homology of structured algebras. Adv. Math., 288:240– 307, 2016.

Perimeter Institute, N2L 2Y5 Waterloo, Canada

Institut de Mathématiques de Bourgogne, UMR 5584, CNRS & Université de Bourgogne, F-21000 Dijon, France