# INDEPENDENT [k]-ROMAN DOMINATION ON GRAPHS

ATÍLIO G. LUIZ AND FRANCISCO ANDERSON SILVA VIEIRA

ABSTRACT. Given a function  $f: V(G) \to \mathbb{Z}_{\geq 0}$  on a graph  $G, AN(v)$  denotes the set of neighbors of  $v \in$  $V(G)$  that have positive labels under f. In 2021, Ahangar et al. introduced the notion of [k]-Roman Dominating Function ([k]-RDF) of a graph G, which is a function  $f: V(G) \to \{0, 1, \ldots, k+1\}$  such that  $\sum_{u \in N[v]} f(u) \geq k + |AN(v)|$  for all  $v \in V(G)$  with  $f(v) < k$ . The weight of  $f$  is  $\sum_{v \in V(G)} f(v)$ . The [k]-Roman domination number, denoted by  $\gamma_{[k]}(G)$ , is the minimum weight of a [k]-RDF of G. The notion of  $[k]$ -RDF for  $k = 1$  has been extensively investigated in the scientific literature since 2004, when introduced by Cockayne et al. as Roman Domination. An independent [k]-Roman dominating function ([k]-IRDF)  $f: V(G) \to \{0, 1, \ldots, k+1\}$  of a graph G is a [k]-RDF of G such that the set of vertices with positive labels is an independent set. The independent [k]-Roman domination number of G is the minimum weight of a [k]-IRDF of G and is denoted by  $i_{[k]R}(G)$ . In this paper, we propose the study of independent [k]-Roman domination on graphs for arbitrary  $k \geq 1$ . We prove that, for all  $k \geq 3$ , the decision problems associated with  $i_{[k]R]}(G)$  and  $\gamma_{[k]R]}(G)$  are NP-complete for planar bipartite graphs with maximum degree 3. We also present lower and upper bounds for  $i_{[k]R]}(G)$ . Moreover, we present lower and upper bounds for the parameter  $i_{[k]R]}(G)$  for two families of 3-regular graphs called generalized Blanuša snarks and Loupekine snarks.

#### 1. Introduction

Let  $G = (V(G), E(G))$  be a graph with vertex set  $V(G)$  and edge set  $E(G)$ . Two vertices  $u, v \in V(G)$ are adjacent or neighbors if  $uv \in E(G)$ . We say that G is trivial if  $|V(G)| = 1$ . For every vertex  $v \in V(G)$ , the open neighborhood of v is the set  $N(v) = \{u \in V(G) : uv \in E(G)\}\)$ , and the closed neighborhood of v is the set  $N[v] = \{v\} \cup N(v)$ . The degree of a vertex  $v \in V(G)$  is the number of neighbors of v and is denoted  $d_G(v)$ . A leaf vertex of G is a vertex  $v \in V(G)$  with  $d_G(v) = 1$ . Graph G is r-regular if  $d_G(v) = r$  for all  $v \in V(G)$ . The maximum degree of G is denoted by  $\Delta(G)$ . For any subset  $S \subseteq V(G)$ , the *induced subgraph* G[S] is the graph whose vertex set is S and whose edge set consists of all edges in  $E(G)$  that have both endpoints in S. As usual,  $P_n$  denotes a path on  $n \geq 1$ vertices and  $C_n$  denotes a cycle on  $n \geq 3$  vertices.

A set  $S \subseteq V(G)$  is called a *dominating set* of G if every vertex  $v \in V(G) \backslash S$  is adjacent to a vertex in S. The domination number of G, denoted  $\gamma(G)$ , is the minimum cardinality of a dominating set of G. A set of vertices  $S \subseteq V(G)$  is called *independent* if no two vertices in S are adjacent. An *independent* dominating set of G is an independent set  $S \subseteq V(G)$  that is also dominating. The *independent domination number* of  $G$ , denoted  $i(G)$ , is the minimum cardinality of an independent dominating set of G. The domination on graphs has been extensively studied in the scientific literature, giving rise to many variations [\[1\]](#page-17-0), a well-known of them being the Roman domination [\[2\]](#page-17-1).

Let G be a graph. For any function  $f: V(G) \to \mathbb{Z}_{\geq 0}$  and  $S \subseteq V(G)$ , define  $f(S) = \sum_{v \in S} f(v)$ . A Roman dominating function (RDF) on G is a function  $f: V(G) \to \{0,1,2\}$  such that every vertex  $u \in V(G)$  with label  $f(u) = 0$  is adjacent to at least one vertex  $v \in V(G)$  with label  $f(v) = 2$ . The weight of an RDF f is defined as  $\omega(f) = f(V(G)) = \sum_{v \in V(G)} f(v)$ . The Roman domination number of G is the minimum weight over all RDFs on G and is denoted by  $\gamma_R(G)$ .

The conception of Roman domination on graphs was motivated by defense strategies devised by the Roman Empire during the reign of Emperor Constantine, 272-337 AD [\[3,](#page-17-2) [4\]](#page-17-3). The idea behind Roman domination is that labels 1 or 2 represent either one or two Roman legions stationed at a given Roman province (vertex  $v$ ). A neighboring province (an adjacent vertex  $u$ ) is considered to be unsecured if no legions are stationed there (i.e.  $f(u) = 0$ ). An unsecured province u can be secured by sending a legion to u from an adjacent province v, by respecting the condition that a legion cannot be sent from a province v if doing so leaves that province without a legion. Thus, two legions must be stationed at a province  $(f(v) = 2)$  before one of the legions can be sent to an adjacent province.

Results on Roman domination and its variants have been collected in [\[5,](#page-17-4) [6,](#page-17-5) [7,](#page-17-6) [8,](#page-17-7) [9\]](#page-17-8), summing up to more than two hundred papers. Many of these variants aim to increase the effectiveness of the defensive strategy modeled by Roman domination. In 2021, Ahangar et al. [\[10\]](#page-17-9) introduced the notion of [k]-Roman domination, a generalization of Roman domination, which groups many of these Roman domination's variants under the same definition. The idea behind [k]-Roman domination is that any unsecured province could be defended by at least  $k$  legions without leaving any secure neighboring province without military forces.

Let G be a graph and  $f: V(G) \to \mathbb{Z}_{\geq 0}$  be a function. We say that a vertex v of G is active under f if  $f(v) \geq 1$ . For any vertex  $v \in V(G)$ , the *active neighborhood* of v under f, denoted by  $AN(v)$ , is the set of vertices  $w \in N(v)$  such that  $f(w) \geq 1$ . For any integer  $k \geq 1$ , a  $\lfloor k \rfloor$ -Roman dominating function on G, also called [k]-RDF, is a function  $f: V(G) \to \{0, 1, \ldots, k+1\}$  such that  $f(N[v]) \geq k + |AN(v)|$  for every vertex  $v \in V(G)$  with  $f(v) < k$ . The weight of a [k]-RDF f on G is defined as  $\omega(f) = f(V(G)) = \sum_{v \in V(G)} f(v)$ . The [k]-Roman domination number of G is the minimum weight that a [k]-RDF of G can have, and is denoted by  $\gamma_{[kR]}(G)$ . A [k]-RDF of G with weight  $\gamma_{[kR]}(G)$ is called a  $\gamma_{[k]}$ -function of G or  $\gamma_{[k]}(G)$ -function. Given a [k]-RDF  $f: V(G) \to \{0, 1, \ldots, k+1\}$  of a graph G, define  $V_i = \{u \in V(G) : f(u) = i\}$  for  $0 \le i \le k+1$ . We call  $(V_0, V_1, \ldots, V_{k+1})$  the ordered partition of  $V(G)$  under f. Since there exists a 1-1 correspondence between the functions  $f: V(G) \to \{0, 1, \ldots, k+1\}$  and the ordered partitions  $(V_0, V_1, \ldots, V_{k+1})$  of  $V(G)$ , it is common to use the notation  $f = (V_0, V_1, \ldots, V_{k+1})$  to refer to a [k]-RDF of G. By the definition of ordered partition, we can alternatively define the weight of a [k]-RDF f as  $\omega(f) = \sum_{p=0}^{k+1} p|V_p|$ . Figure [1](#page-1-0) shows some graphs endowed with  $[k]$ -RDFs.

<span id="page-1-0"></span>

(A) Path with a  $[k]$ -RDF with weight  $k + 1$ . (B) Tree with a  $[k]$ -RDF with weight  $2k + 2$ .

FIGURE 1. Two graphs endowed with  $[k]$ -Roman dominating functions.

For every  $k \geq 1$ , the [k]-Roman Domination Problem is to determine  $\gamma_{kR}(G)$  for an arbitrary graph G. Khalili et al. [\[11\]](#page-17-10) proved that the decision version of the  $k$ -Roman Domination Problem is  $N$ P-Complete even when restricted to bipartite and chordal graphs. Moreover, Valenzuela-Tripodoro et al. [\[12\]](#page-17-11) proved that the decision version of  $[k]$ -RDP is  $N \mathcal{P}$ -Complete even when restricted to star convex and comb convex bipartite graphs.

Note that [1]-Roman domination is equivalent to the original Roman domination definition. In addition, [2]-Roman domination has been previously studied [\[13\]](#page-17-12) under the name of Double Roman domination, as well as [3]-Roman domination has been investigated [\[10\]](#page-17-9) under the name of Triple Roman domination, and [4]-Roman domination has been recently studied under the name of Quadruple Roman domination [\[14\]](#page-17-13). Recently, Khalili et al. [\[11\]](#page-17-10) and Valenzuela-Tripodoro et al. [\[12\]](#page-17-11) presented sharp upper and lower bounds for the [k]-Roman domination number for all  $k \geq 1$ .

Given a [k]-Roman dominating function  $f = (V_0, V_1, \ldots, V_{k+1})$  on a graph G, we observe that the set of vertices  $S = V_1 \cup V_2 \cup \cdots \cup V_{k+1}$  is a dominating set of G since  $V(G) \setminus S = V_0$  and every vertex in  $V_0$ is adjacent to a vertex in S. This connection between dominating sets and the set of active vertices of a graph G under a [k]-Roman dominating function makes it possible to relate the parameters  $\gamma(G)$  and  $\gamma_{kR}(G)$  as well as to extend some restrictions traditionally imposed on dominating sets to [k]-Roman dominating functions. An example is the concept of independent dominating set: one may require the dominating set of active vertices of  $G$  to be also independent. Indeed, in their seminal paper, Cockayne et al. [\[2\]](#page-17-1) introduced the notion of Roman dominating functions  $f = (V_0, V_1, V_2)$  whose set of active vertices  $V_1 \cup V_2$  is an independent set, which are called *independent Roman dominating functions*. In 2019, Maimani et al. [\[15\]](#page-17-14) introduced the notion of independent double Roman dominating function, which is a [2]-Roman dominating function  $f = (V_0, V_1, V_2, V_3)$  of a graph G such that the set of active vertices  $V_1 \cup V_2 \cup V_3$  is an independent set. When studying independent Roman domination and independent double Roman domination, one can observe some differences but many similarities. Thus, based on the previous observations, we propose a generalization of independent Roman domination and independent double Roman domination, defined as follows.

A [k]-Roman dominating function  $f = (V_0, V_1, \ldots, V_{k+1})$  on a graph G is called an *independent* [k]-Roman dominating function, or [k]-IRDF for short, if the set of active vertices  $V_1 \cup V_2 \cup \cdots \cup V_{k+1}$  is an independent set. The *independent* [k]-Roman domination number  $i_{[k]}(G)$  is the minimum weight of a [k]-IRDF on G, and a [k]-IRDF of G with weight  $i_{[kR]}(G)$  is called an  $i_{[kR]}$ -function of G or  $i_{[kR]}(G)$ function. The Independent [k]-Roman Domination Problem consists in determining  $i_{[kR]}(G)$  for an arbitrary graph G. Since every [k]-IRDF is a [k]-RDF, we trivially obtain that  $\gamma_{kR}(G) \leq i_{[kR]}(G)$  for every graph G. As an example, Figure [2](#page-2-0) shows some graphs with  $i_{[k]}$ -functions.

<span id="page-2-0"></span>

FIGURE 2. Three graphs endowed with independent  $[k]$ -Roman dominating functions.

From the definition of independent [k]-Roman domination, we know that the active vertices  $v \in$  $V(G)$  with  $f(v) < k$  must have at least one active neighbor since the condition  $f(N[v]) \geq k + |AN(v)|$ must be satisfied. In addition to the previous condition, an independent  $[k]$ -Roman domination function also imposes that the set of active vertices must be independent. However, these two conditions considered simultaneously imply that an independent  $[k]$ -Roman dominating function does not assign labels from the set  $\{1, 2, \ldots, k-1\}$  to the vertices of a graph G. These initial observations concerning [k]-IRDFs are explicitly stated in the following propositions.

<span id="page-2-3"></span><span id="page-2-1"></span>**Proposition 1.** If G is a graph, then 
$$
\gamma_{[kR]}(G) \leq i_{[kR]}(G)
$$
.

**Proposition 2.** If  $f = (V_0, V_1, \ldots, V_{k+1})$  is a [k]-IRDF of a graph G, then  $V_i = \emptyset$  for all  $i \in$  $\{1, 2, \ldots, k-1\}.$ 

By Proposition [2,](#page-2-1) we can represent a [k]-IRDF  $f = (V_0, V_1, \ldots, V_{k+1})$  simply as  $f = (V_0, V_k, V_{k+1})$ . Moreover, note that the weight of a [k]-IRDF  $f = (V_0, V_k, V_{k+1})$  is also given by  $\omega(f) = k|V_k| + (k +$  $1)|V_{k+1}|.$ 

In this paper, we propose the study of independent  $[k]$ -Roman domination on graphs for arbitrary  $k \geq 1$ . The next sections of this paper are organized as follows. In Section [2,](#page-2-2) we prove that, for all  $k \geq 3$ , the decision versions of the Independent [k]-Roman Domination Problem and [k]-Roman Domination Problem are  $\mathcal{NP}$ -complete, even when restricted to planar bipartite graphs with maximum degree 3. In Section [3,](#page-5-0) we present some sharp lower and upper bounds for the independent  $[k]$ -Roman domination number of arbitrary graphs. In Sections [4](#page-7-0) and [5,](#page-13-0) we present specific lower bounds and upper bounds for the independent [k]-Roman domination number for two infinite families of 3-regular graphs called Generalized Blanuša Snarks and Loupekine snarks. Section [6](#page-17-15) presents our concluding remarks.

## 2. Complexity results

<span id="page-2-2"></span>In this section, we show that, for every integer  $k \geq 3$ , the decision versions of the [k]-Roman Domination Problem  $([k]$ -ROM-DOM) and the Independent  $[k]$ -Roman Domination Problem  $([k]$ -IROM-DOM) are  $N\mathcal{P}$ -complete when restricted to graphs with maximum degree 3. We remark that the  $\mathcal{NP}$ -completeness of [1]-ROM-DOM and [1]-IROM-DOM, when restricted to this same class of graphs, have already been established [\[16\]](#page-17-16). In the remaining of this section, we deal with  $k \geq 3$ . Consider the following decision problems.

## $[k]$ -ROM-DOM

**Instance:** A graph G and a positive integer  $\ell$ . **Question:** Does G have a [k]-Roman dominating function with weight at most  $\ell$ ?

### [k]-IROM-DOM

**Instance:** A graph G and a positive integer  $\ell$ .

**Question:** Does G have an independent [k]-Roman dominating function with weight at most  $\ell$ ?

For  $k \geq 3$ , we show that [k]-ROM-DOM and [k]-IROM-DOM are  $N\mathcal{P}$ -complete when restricted to graphs with maximum degree 3 through a reduction from the vertex cover problem. A vertex cover of a graph G is a set of vertices  $S \subseteq V(G)$  such that each edge of G is incident to some vertex in S. The vertex covering number of G, denoted  $\tau(G)$ , is the cardinality of a smallest vertex cover of G. Given a graph G and a positive integer  $\ell$ , the Vertex Cover Problem consists in deciding whether G has a vertex cover S with cardinality at most  $\ell$ . B. Mohar [\[17\]](#page-17-17) proved that the Vertex Cover Problem is  $N\mathcal{P}$ complete even when restricted to 2-connected planar 3-regular graphs. We use this result to construct a polynomial time reduction from the Vertex Cover Problem to [k]-ROM-DOM ([k]-IROM-DOM). The construction is described as follows: given a 2-connected planar 3-regular graph  $G$ , construct a new graph F from G by replacing each edge  $e = uv \in E(G)$  by a gadget  $G_e$  illustrated in Figure [3.](#page-3-0) Note that F can be constructed in polynomial time on  $|E(G)|$  and also that F is a planar bipartite graph with maximum degree 3.

<span id="page-3-0"></span>

FIGURE 3. Gadget  $G_e$  used in the reduction.

In order to prove the  $\mathcal{NP}$ -completeness result, given in Theorem [9,](#page-5-1) we need the following auxiliary results.

<span id="page-3-1"></span>**Lemma 3** (Khalili et al. [\[11\]](#page-17-10)). If  $k \geq 2$ , then in a  $\gamma_{kR}(G)$ -function of a graph G, no vertex needs to be assigned the label 1.

<span id="page-3-2"></span>**Proposition 4.** Let  $k \geq 1$  be an integer. If G is a connected graph with at least 3 vertices, then in a  $[k]$ -Roman dominating function of G with weight  $\gamma_{[k]}(G)$ , no leaf vertex of G needs to be assigned the label  $k + 1$ .

*Proof.* Let G be a connected graph on at least 3 vertices and f a  $\gamma_{[k]}$ -function of G. Let  $v \in V(G)$  be a leaf vertex and let  $w \in V(G)$  be the neighbor of v. For the purpose of contradiction, suppose that f needed to assign  $k+1$  to v, that is,  $f(v) = k+1$ . Since f is a  $\gamma_{[k]}$ -function,  $f(w) \leq k$  (otherwise, by assigning 0 to v we obtain a [k]-RDF with weight smaller then  $\omega(f)$ ). We modify the labeling f by exchange the labels of the vertices  $v$  and  $w$  and maintaining the labels of all the other vertices the same. Note that f continues to be a  $[k]$ -Roman dominating function with the same weight as before and the vertex v does not have the weight  $k+1$  anymore.

<span id="page-3-3"></span>**Proposition 5.** Let  $k \geq 1$  be an integer and G be a connected graph with at least 3 vertices. In any  $\gamma_{[kR]}$ -function f of G, no leaf vertex needs to be assigned a label different from 0 or k.

*Proof.* Let G and f be as in the hypothesis and let  $v \in V(G)$  be a leaf vertex with neighbor  $w \in$  $V(G)$ . By Lemma [3](#page-3-1) and Proposition [4,](#page-3-2)  $f(v) \notin \{1, k+1\}$ . If  $f(v) \in \{0, k\}$ , then f is the desired function. Thus, suppose that f needs to assign a label in the set  $\{2, 3, \ldots, k-1\}$  to vertex v, that is,  $f(v) \in \{2, 3, \ldots, k-1\}$ . In this case, the neighbor w of v has  $f(w) \neq 0$  and is, thus, an active neighbor of v. By the definition of  $[k]$ -RDF,  $f(N[v]) = f(v) + f(w) \geq k + |AN(v)| = k + 1$ . Thus,  $f(v) + f(w) \geq k+1$ . We modify the labeling f by assigning label  $f(v) + f(w)$  to vertex w, by assigning label 0 to vertex  $v$ , and maintaining the labels of all the remaining vertices of  $G$  the same. Note that f continues to be a  $[k]$ -RDF with the same weight as before and the new label of v does not belong to the set  $\{2, 3, \ldots, k-1\}$ , which is a contradiction.

<span id="page-3-4"></span>**Lemma 6.** Let  $k \geq 2$  be an integer. Given a 2-connected planar 3-regular graph G, let F be the graph constructed from G by replacing each edge  $e = uv$  in G by a gadget  $G_e$  shown in Figure [3.](#page-3-0) Then, any  $\gamma_{[k]}$ -function f of F satisfies  $(f(x_6^e), f(x_7^e)) \in \{(0, k+1), (k, 0)\}$  and  $(f(x_9^e), f(x_{10}^e)) \in$  $\{(k+1,0),(0,k)\}.$ 

*Proof.* Let f be a  $\gamma_{[k]}$ -function of F. We only analyze the values  $f(x_6)$  and  $f(x_7)$  since the analysis for  $f(x_9^e)$  and  $f(x_{10}^e)$  is analogous and follows from the symmetry of F along the vertical axis.

Note that  $x_6^e$  is a leaf vertex and  $N(x_6^e) = \{x_7^e\}$ . By Proposition [5,](#page-3-3) either  $f(x_6^e) = 0$  or  $f(x_6^e) = k$ . If  $f(x_6^e) = 0$ , then  $f(x_7^e) = k + 1$  by the definition of [k]-RDF, and the result follows. Thus, suppose that  $f(x_6^e) = k$ . By Lemma [3,](#page-3-1)  $f(x_7^e) \neq 1$ . If  $f(x_7^e) \geq 2$ , then  $f(x_6^e) + f(x_7^e) \geq k + 2$ . Hence, it would be possible to obtain a [k]-RDF with smaller weight by assigning label  $f(x_6^e) + f(x_7^e) - 1$  to  $x_7^e$  and 0 to v, thus contradicting the choice of f. Therefore, we obtain that  $f(x_7^e) = 0$ , and the result follows.  $\Box$ 

<span id="page-4-0"></span>**Lemma 7.** Let  $k \geq 3$  be an integer. Given a 2-connected planar 3-regular graph G, let F be a graph constructed from G by replacing each edge  $e = uv$  in G by a gadget  $G_e$  illustrated in Figure [3.](#page-3-0) Let  $U_e = \{x_2^e, x_3^e, x_4^e, x_5^e, x_7^e, x_8^e, x_9^e, x_{10}^e\} \subset V(G_e)$ . Then, in any  $\gamma_{[kR]}$ -function f of the graph F, we have that the function f restricted to  $U_e$  is a  $[k]$ -RDF of  $F[U_e]$  with weight  $f(U_e) = 3k + 2$ . Moreover,  $(f(x_2^e), f(x_4^e)) \in \{(0,0), (k+1,0), (0, k+1)\}.$ 

*Proof.* Let  $k \geq 3$  be an integer. Let G and F be as in the hypothesis. Let  $f: V(F) \to \{0, 1, \ldots, k+1\}$ be a  $\gamma_{[k]}$ -function of F. For each gadget  $G_e \subset F$ , define  $U_e = \{x_2^e, x_3^e, x_4^e, x_6^e, x_7^e, x_8^e, x_9^e, x_{10}^e\} \subset V(G_e)$ . By Lemma [6,](#page-3-4)  $(f(x_6^e), f(x_7^e)) \in \{(0, k+1), (k, 0)\}\$  and  $(f(x_9^e), f(x_{10}^e)) \in \{(k+1, 0), (0, k)\}\$ . Thus, there are four cases to analyze, depending on the values of the labels  $f(x_6^e)$ ,  $f(x_7^e)$ ,  $f(x_9^e)$  and  $f(x_{10}^e)$ .

**Case 1:**  $(f(x_6^e), f(x_7^e)) = (k, 0)$  and  $(f(x_9^e), f(x_{10}^e)) = (0, k)$ . We claim that this case cannot occur. For the purpose of contradiction, suppose it occurs. Since f is a  $\gamma_{[k]}$ -function and  $x_8^e$  has no active neighbor, we have that  $f(x_8^e) = k$ . Note that  $\sum_{i=6}^{10} f(x_i^e) = 3k$ . Thus, we can redefine the labels of some vertices of F so as to obtain another  $[k]$ -RDF f' of F with smaller weight than f such that  $\sum_{i=6}^{10} f'(x_i^e) = 2k + 2 < 3k$ , as follows: define  $(f'(x_6^e), f'(x_7^e), f'(x_8^e), f'(x_9^e), f'(x_{10}^e)) = (0, k + 1, 0, k + 1)$ 1,0) and make  $f'(x) = f(x)$  for every remaining vertex x of F. This contradicts the choice of f as a  $\gamma_{[k]}$ -function.

**Case 2:**  $(f(x_6^e), f(x_7^e)) = (0, k+1)$  and  $(f(x_9^e), f(x_{10}^e)) = (k+1, 0)$ . Since f is a  $\gamma_{[k]}$ -function, we have that  $f(x_8^e) = 0$ . By the definition of [k]-RDF, we have that  $f(N[x_3^e]) = f(x_2^e) + f(x_3^e) + f(x_4^e) \ge$  $k + |AN(x_3^e)| \ge k$ . All these facts imply that  $f(U_e) = \sum_{w \in U_e} f(w) \ge 3k + 2$ . Moreover, a [k]-RDF of  $F[U_e]$  with weight  $3k+2$  is obtained by assigning labels  $\tilde{f}(x_2^e) = 0$ ,  $f(x_3^e) = k$  and  $f(x_4^e) = 0$ . Therefore,  $f(U_e) = 3k + 2$ ,  $(f(x_2^e), f(x_4^e)) = (0, 0)$ , and the result follows.

**Case 3:**  $(f(x_6^e), f(x_7^e)) = (0, k+1)$  and  $(f(x_9^e), f(x_{10}^e)) = (0, k)$ . Since f is a  $\gamma_{[kR]}$ -function, we have that  $f(x_8^e) = 0$ . Moreover, since  $f(x_9^e) = 0$  and  $f(x_8^e) + f(x_{10}^e) < k+1$  we obtain that  $f(x_4^e) \neq 0$ . By the definition of [k]-RDF and since  $f(x_4^e) \neq 0$ , we have that  $f(N[x_3^e]) = f(x_2^e) + f(x_3^e) + f(x_4^e) \ge$  $k + |AN(x_3^e)| \ge k + 1$ . All these facts imply that  $f(U_e) = \sum_{w \in U_e} f(w) \ge 3k + 2$ . From the previous facts, we have that  $f(U_e) = 3k + 2$  only if  $f(N[x_3^e]) = k + |\tilde{A}N(x_3^e)| = k + 1$ , which implies that  $f(x_2^e) = 0$ . Thus,  $f(x_2^e) + f(x_3^e) + f(x_4^e) = 0 + f(x_3^e) + f(x_4^e) = k + 1$ , i.e.,  $f(x_3^e) + f(x_4^e) = k + 1$ . Since f is a  $\gamma_{[k]}$ -function, we obtain that  $f(x_3^e) = 0$  and  $f(x_4^e) = k+1$ . Therefore,  $(f(x_2^e), f(x_4^e)) = (0, k+1)$ and the result follows.

**Case 4:**  $(f(x_6^e), f(x_7^e)) = (k, 0)$  and  $(f(x_9^e), f(x_{10}^e)) = (k + 1, 0)$ . The proof for this case is analogous to the proof of the previous case and follows from the symmetry of  $G_e$  along the vertical axis.

Therefore, in any  $\gamma_{[k]}$ -function f of the graph F, we have that the function f restricted to  $U_e$  is a [k]-RDF of  $F[U_e]$  with weight  $f(U_e) = 3k + 2$ . Moreover,  $(f(x_2^e), f(x_4^e)) \in \{(0,0), (k+1,0), (0, k+1)\}$ .

<span id="page-4-1"></span>**Theorem 8.** Let  $k \geq 3$  be an integer. Given a 2-connected planar 3-regular graph G, let F be a planar bipartite graph with  $\Delta(F) = 3$  constructed from G by replacing each edge e = uv in G by a gadget  $G_e$ illustrated in Figure [3.](#page-3-0) Then,

$$
\gamma_{[kR]}(F) = i_{[kR]}(F) = \tau(G) + k|V(G)| + (3k+2)|E(G)|.
$$

*Proof.* Let G and F be as in the statement of the theorem. Let C be a vertex cover of G with  $|C| = \tau(G).$ 

We initially prove that  $i_{[kR]}(F) \leq \tau(G) + k|V(G)| + (3k+2)|E(G)|$ . In order to do this, we construct an appropriate  $[k]$ -IRDF  $f = (V_0, V_k, V_{k+1})$  of F as follows. First, define two empty sets  $D_k$  and  $D_{k+1}$ . For each gadget  $G_e \subset F$ , associated with an edge  $e = uv \in E(G)$ , do the following: if  $v \in C$ , then, add the vertex  $x_6^e$  to  $D_k$  and add the vertices  $x_2^e$  and  $x_9^e$  to  $D_{k+1}$ ; otherwise, add the vertex  $x_{10}^e$ to  $D_k$  and add the vertices  $x_4^e$  and  $x_7^e$  to  $D_{k+1}$ . Note that  $|D_k| = |E(G)|$  and  $|D_{k+1}| = 2|E(G)|$ . Define the function  $f = (V_0, V_k, V_{k+1})$  such that  $V_0 = V(F) \setminus (V(G) \cup D_k \cup D_{k+1}), V_k = D_k \cup V(G) \setminus C$ and  $V_{k+1} = D_{k+1} \cup C$ . From the definition of f, we have that f is a [k]-IRDF of F with weight  $\omega(f) = k|D_k \cup V(G) \setminus C| + (k+1)|D_{k+1} \cup C| = k(|E(G)| + |V(G)| - \tau(G)) + (k+1)(2|E(G)| + \tau(G)) =$  $\tau(G) + k|V(G)| + (3k+2)|E(G)|$ . Therefore,  $i_{[kR]}(F) \leq \omega(f) = \tau(G) + k|V(G)| + (3k+2)|E(G)|$ .

Next, we show that  $\gamma_{kR}(F) \geq \tau(G) + k|V(G)| + (3k+2)|E(G)|$ . Let  $f = (V_0, \emptyset, V_2, \ldots, V_{k+1})$ be a  $\gamma_{[k]}$ -function of F. Let  $G_e$  be a gadget of F, for any edge  $e = uv \in E(G)$ . Define the set  $U_e = \{x_2^e, x_3^e, x_4^e, x_6^e, x_7^e, x_8^e, x_9^e, x_{10}^e\} \subset V(G_e) \subset V(F)$ . By Lemma [7,](#page-4-0) the function f restricted to  $U_e$  has weight  $3k+2$  and is a [k]-RDF of the subgraph induced by  $U_e$ . Moreover,  $(f(x_2^e), f(x_4^e)) \in$  $\{(0, k + 1), (k + 1, 0), (0, 0)\}.$  Let  $S = \{x \in U_e : f(x) \neq 0, e \in E(G)\}.$  Let  $V' \subset V(F)$  be the set of vertices that are not adjacent to some vertex in S and are not in S, that is,  $V' = V(F) \setminus N[S]$ . Let F' be the induced subgraph  $F[V']$ . For each  $e \in E(G)$ , all the vertices in  $U_e$  and at most one of the vertices  $x_1^e$  and  $x_5^e$  are not in V'. This implies that F' is a forest of trees with  $|V(G)|$ components such that each component is a star whose central vertex is a vertex  $z \in V(G)$ . Let T be a component of F'. If T is a single vertex (i.e.  $V(T) = \{z\}$ ), then  $f(z) = k$ . On the other hand, if T is not a single vertex, then z is the central vertex of the star T and  $f(z) = k + 1$ . Let  $D = V(G) \cap V_{k+1}$ . From the above discussion, we conclude that D is a vertex cover of G. Since each subset  $U_e$  contributes with  $3k+2$  to the weight of f and there are  $|E(G)|$  of these subsets, then they contribute to a total of  $(3k+2)|E(G)|$  to the weight of f. From these facts we obtain that  $\omega(f) = (k+1)|D| + k(|V(G)| - |D|) + (3k+2)|E(G)| = |D| + k|V(G)| + (3k+2)|E(G)|$ . Thus,  $\tau(G) \leq |D| = \omega(f) - k|V(G)| - (3k+2)|E(G)| = \gamma_{kR}(G) - k|V(G)| - (3k+2)|E(G)|$ . Therefore,  $\gamma_{[kR]}(G) \geq \tau(G) + k|V(G)| + (3k+2)|E(G)|.$ 

Since  $\gamma_{[kR]}(G) \leq i_{[kR]}(G)$  (see Proposition [1\)](#page-2-3), we have that  $\tau(G) + k|V(G)| + (3k+2)|E(G)| \leq$  $\gamma_{[kR]}(G) \leq i_{[kR]}(G) \leq \tau(G) + k|V(G)| + (3k+2)|E(G)|$ , and the result follows.

<span id="page-5-1"></span>**Theorem 9.** Let  $k \geq 3$  be an integer. Then, [k]-ROM-DOM (resp. [k]-IROM-DOM) is NP-complete even when restricted to planar bipartite graphs G with  $\Delta(G) = 3$ .

*Proof.* We first show that [k]-ROM-DOM (resp. [k]-IROM-DOM) is a member of  $N\mathcal{P}$ . Given any instance  $(G, \ell)$  of [k]-ROM-DOM (resp. [k]-IROM-DOM) and a certificate function  $f: V(G) \to$  $\{0,1,\ldots,k+1\}$ , we can verify (in polynomial time) if  $\sum_{v\in V(G)} f(v) \leq \ell$  and if  $\sum_{u\in N[v]} f(u) \geq$  $|AN(v)| + k$  for every  $v \in V(G)$  (resp. in the case of [k]-IROM-DOM, it is also necessary to check if  $V_k \cup V_{k+1}$  is an independent set). Next, we show that [k]-ROM-DOM (resp. [k]-IROM-DOM) is  $\mathcal{NP}$ -hard. Recall that we showed how to construct a planar bipartite graph F with  $\Delta(F) = 3$  from a given 2-connected planar 3-regular graph G in polynomial time on  $|E(G)|$ . From Theorem [8,](#page-4-1) we deduce that there exists a polynomial time algorithm that calculates  $\tau(G)$  if and only if there exists a polynomial time algorithm that calculates  $\gamma_{kR}(F)$  (resp.  $i_{[kR]}(F)$ ). However, since VCP is NPcomplete even when restricted to 2-connected planar 3-regular graphs, we obtain, from this reduction, that  $[k]$ -ROM-DOM (resp.  $[k]$ -IROM-DOM) is  $N\mathcal{P}$ -complete even when restricted to planar bipartite graphs with maximum degree 3.  $\Box$ 

## 3. BOUNDS FOR THE INDEPENDENT  $[k]$ -ROMAN DOMINATION NUMBER

<span id="page-5-0"></span>In this section, we present some lower and upper bounds for the independent  $[k]$ -Roman domination number of arbitrary graphs. Since the set  $V_k \cup V_{k+1}$  is an independent dominating set in every [k]-IRDF  $f = (V_0, V_k, V_{k+1})$  of a graph G, it seems reasonable that  $i_{[kR]}(G)$  and  $i(G)$  are related, such as shown in the following proposition.

<span id="page-5-2"></span>**Proposition 10.** Let  $k \geq 1$  be an integer. If G is a graph, then  $k \cdot i(G) \leq i_{kR}(G) \leq (k+1) \cdot i(G)$ .

*Proof.* Given a minimum independent dominating set S of a graph G, we define a  $[k]$ -IRDF  $f =$  $(V_0, V_k, V_{k+1})$  of G with weight  $(k+1)i(G)$  by making  $V_{k+1} = S$ ,  $V_k = \emptyset$  and  $V_0 = V(G) \setminus S$ . Hence,  $i_{[kR]}(G) \le \omega(f) = (k+1)i(G).$ 

Now, let  $f = (V_0, V_k, V_{k+1})$  be an  $i_{[k]}$ -function of a graph G. Since  $i(G) \leq |V_k| + |V_{k+1}|$ , we have that  $k \cdot i(G) \leq k(|V_k| + |V_{k+1}|) \leq k|V_k| + (k+1)|V_{k+1}| = i_{kR}(G)$ , and the result follows.

The lower bound presented in Proposition [10](#page-5-2) is tight since it is attained by empty graphs. Moreover, graphs whose independent [k]-Roman domination number equals the upper bound given in Proposition [10](#page-5-2) receive a specific name. We say that a graph G is independent  $|k\rangle$ -Roman when  $i_{[k]R}(G) = (k+1)i(G)$ . The next lemma is a generalization of a result of Shao et al. [\[18\]](#page-17-18) and presents a characterization of independent [k]-Roman graphs.

<span id="page-5-3"></span>**Lemma 11.** Let G be a graph. Then, G is independent [k]-Roman if and only if G has an  $i_{[kR]}$ -function  $f = (V_0, V_k, V_{k+1})$  such that  $V_k = \emptyset$ .

*Proof.* Let G be a graph. First, suppose that G has an  $i_{[k]}(G)$ -function  $f = (V_0, V_k, V_{k+1})$  such that  $V_k = \emptyset$ . This implies that  $i(G) \leq |V_k| + |V_{k+1}| = |V_{k+1}|$  and that  $i_{[k]}(G) = (k+1)|V_{k+1}|$ . Thus,  $(k+1)i(G) \leq (k+1)|V_{k+1}| = i_{kR}(G)$ . By Proposition [10,](#page-5-2)  $i_{kR}(G) \leq (k+1)i(G)$ . Therefore,  $i_{[kR]}(G) = (k+1)i(G)$  and G is independent [k]-Roman.

Now, consider G independent [k]-Roman. For the purpose of contradiction, suppose that every  $i_{[kR]}(G)$ -function  $f = (V_0, V_k, V_{k+1})$  has  $V_k \neq \emptyset$ . Let  $f = (V_0, V_k, V_{k+1})$  be an  $i_{[kR]}$ -function of G with  $|V_k|$  as minimum as possible. From the definition of [k]-IRDF, we know that  $i(G) \leq |V_k \cup V_{k+1}|$  $|V_k| + |V_{k+1}|.$ 

In fact, we claim that  $i(G) = |V_k| + |V_{k+1}|$ . In order to prove this claim, suppose that there exists a minimum independent dominating set S of G such that  $|S|$  <  $|V_k|$  +  $|V_{k+1}|$ . Let  $g = (V_0^g, V_k^g, V_{k+1}^g)$  be a function with  $V_k^g = \emptyset$ ,  $V_{k+1}^g = S$  and  $V_0^g = V(G) \setminus S$ . Thus, g is a [k]-IRDF of G with  $|V_{k+1}^g| = i(G)$ . Since G is independent [k]-Roman, we have that  $(k+1)i(G) = i_{[k]R]}(G)$ . Moreover, by the definition of [k]-IRDF, we know that  $i_{[k]}(G) = k|V_k| + (k+1)|V_{k+1}|$ . Then, we have that  $\omega(g) = (k+1)|V_{k+1}^g| =$  $(k+1)i(G) = i_{[k]}(G) = k|V_k| + (k+1)|V_{k+1}| = \omega(f)$ . Thus, we have  $\omega(g) = \omega(f) = i_{[k]}(G)$ , but  $V_k^g = \emptyset$ . In other words, we found an  $i_{[kR]}(G)$ -function  $g = (V_0^g, V_k^g, V_{k+1}^g)$  with  $V_k^g = \emptyset$ , which is a contradiction. Therefore,  $i(G) = |V_k| + |V_{k+1}|$  as claimed.

Since G is independent [k]-Roman, we have that  $(k+1)i(G) = i_{[k]}(G)$  and, thus,  $(k+1)(|V_k| +$  $|V_{k+1}| = (k+1)i(G) = i_{[k]}(G) = k|V_k| + (k+1)|V_{k+1}|$ , implying that  $|V_k| = 0$ , which is a contradiction. Therefore, we conclude that G has an  $i_{[k]}(G)$ -function  $f = (V_0, V_k, V_{k+1})$  such that  $V_k = \emptyset$ .

Another useful kind of lower bound connects the parameter with the maximum degree and number of vertices of the graph. As an example, in what concerns the [k]-Roman domination number, Valenzuela-Tripodoro et al. [\[12\]](#page-17-11) presented the following lower bound for the  $[k]$ -Roman domination number of nontrivial connected graphs.

<span id="page-6-0"></span>**Theorem 12** (Valenzuela-Tripodoro et al. [\[12\]](#page-17-11)). Let  $k \geq 1$  be an integer. Let G be a nontrivial connected graph with maximum degree  $\Delta(G) \geq k$ . Then  $\gamma_{[k]}(G) \geq \frac{|V(G)|(k+1)}{\Delta(G)+1}$ .

<span id="page-6-1"></span>Since  $i_{[kR]}(G) \geq \gamma_{[kR]}(G)$ , we immediatelly obtain the following corollary from Theorem [12.](#page-6-0)

**Corollary 13.** Let  $k \geq 1$  be an integer. Let G be a nontrivial connected graph with maximum degree  $\Delta(G) \geq k$ . Then  $i_{[kR]}(G) \geq \frac{|V(G)|(k+1)}{\Delta(G)+1}$ .

We remark that Corollary [13](#page-6-1) only applies for the cases where  $k \leq \Delta(G)$ . In the next theorem we present a new lower bound for the independent  $[k]$ -Roman domination number of connected graphs  $G$ with  $k \geq \Delta(G)$  and  $k \geq 4$ .

<span id="page-6-2"></span>**Theorem 14.** Let  $k \geq 4$  be an integer. If G is a nontrivial connected graph with  $k \geq \Delta(G) \geq 1$ , then

$$
i_{[kR]}(G) \ge \frac{|V(G)|(k+1)}{\Delta(G)+1}.
$$

Moreover, if  $i_{[k]}(G) = \frac{|V(G)|(k+1)}{\Delta(G)+1}$ , then G is independent [k]-Roman.

*Proof.* Let  $k \geq 4$  be an integer and G be a nontrivial connected graph with maximum degree  $\Delta \geq 1$ . Suppose that  $k \geq \Delta$ . Let  $f: V(G) \to \{0, k, k+1\}$  be an  $i_{[k]}(G)$ -function. Recall that  $V_i = \{w \in$  $V(G)$ :  $f(w) = i$  for  $i \in \{0, k, k+1\}$ . In this proof, we use a discharging procedure similar to the approach used by Shao et al. [\[18\]](#page-17-18). Our discharging procedure is described as follows. Firstly, each vertex  $v \in V(G)$  is assigned the initial charge  $s(v) = f(v)$ . Next, we apply the discharging procedure defined by means of the following two rules:

**Rule 1:** every vertex  $v \in V(G)$  with  $s(v) = k+1$  sends a charge of  $\frac{k+1}{\Delta+1}$  to each vertex in  $N(v) \cap V_0$ ; **Rule 2:** every vertex  $v \in V(G)$  with  $s(v) = k$  sends a charge of  $\frac{(k-2)(k+1)}{k(\Delta+1)}$  to each vertex in  $N(v) \cap V_0$ .

Denote by  $s'(v)$  the final charge of vertex v after applying the discharging procedure. Note that:

I. for each vertex  $v \in V(G)$  with  $f(v) = k + 1$ , since it sends charge to at most  $d_G(v)$  vertices, by Rule 1 we obtain that the final charge of v is  $s'(v) \geq s(v) - d_G(v) \frac{k+1}{\Delta+1} \geq (k+1) - \frac{\Delta(k+1)}{\Delta+1} = \frac{k+1}{\Delta+1}$ , that is,  $s'(v) \geq \frac{k+1}{\Delta+1}$ ;

II. for each vertex  $v \in V(G)$  with  $f(v) = k$ , since it sends charge to at most  $d_G(v)$  vertices, by Rule 2 we obtain that the final charge of v is  $s'(v) \geq s(v) - d_G(v) \frac{(k-2)(k+1)}{k(\Delta+1)} \geq k - \frac{\Delta(k-2)(k+1)}{k(\Delta+1)} =$ 

$$
\frac{k^2 + \Delta k + 2\Delta}{k(\Delta + 1)} > \frac{k^2 + \Delta k}{k(\Delta + 1)} = \frac{k + \Delta}{\Delta + 1} \ge \frac{k + 1}{\Delta + 1}, \text{ that is, } s'(v) > \frac{k + 1}{\Delta + 1}.
$$

From the previous analysis, we obtain that  $s'(v) \geq \frac{k+1}{\Delta+1}$  for all  $v \in V(G)$  with  $f(v) > 0$ . Now, let us analyze an arbitrary vertex  $v \in V(G)$  with  $f(v) = 0$ . Since f is a [k]-IRDF, we have that  $f(N[v]) = f(N(v)) \ge |AN(v)| + k$ . So, either v has at least one neighbor  $w \in V_{k+1}$  or v has at least two neighbors  $u_1, u_2 \in V_k$ . If v has at least one neighbor  $w \in V_{k+1}$ , then  $s'(v) \geq \frac{k+1}{\Delta+1}$  since w sent a charge of  $\frac{k+1}{\Delta+1}$  to v. On the other hand, if v has at least two neighbors  $u_1, u_2 \in V_k$ , each of these neighbors sent a charge of  $\frac{(k-2)(k+1)}{k(\Delta+1)}$  to v and, thus,  $s'(v) \geq 2 \cdot \frac{(k-2)(k+1)}{k(\Delta+1)} \geq \frac{k+1}{\Delta+1}$  for  $k \geq 4$ . Hence, we obtain that  $s'(v) \geq \frac{k+1}{\Delta+1}$  for all  $v \in V(G)$ . Moreover, since the discharging procedure does not change the total value of charge in  $G$ , we obtain that

<span id="page-7-1"></span>
$$
(1) \quad i_{[k]}(G) = \omega(f) = \sum_{v \in V(G)} f(v) = \sum_{v \in V(G)} s(v) = \sum_{v \in V(G)} s'(v) \ge \sum_{v \in V(G)} \frac{k+1}{\Delta+1} = |V(G)| \cdot \frac{(k+1)}{\Delta+1}.
$$

Therefore,  $i_{[k]}(G) = \omega(f) \ge \frac{|V(G)|(k+1)}{\Delta+1}$ . From now on, suppose that  $\omega(f) = \frac{|V(G)|(k+1)}{\Delta+1}$ . In this case, by the inequality chain [\(1\)](#page-7-1), we have that  $s'(v) = \frac{k+1}{\Delta+1}$  for all  $v \in V(G)$ . This implies that no vertex of G was assigned label k since  $s'(w) > \frac{k+1}{\Delta+1}$  for every vertex  $w \in V(G)$  with  $f(w) = k$ . Hence, by Lemma [11,](#page-5-3) G is independent  $[k]$ -Roman.

<span id="page-7-2"></span>The next result follows immediately from Corollary [13](#page-6-1) and Theorem [14.](#page-6-2)

**Theorem 15.** Let  $k \geq 1$  be an integer. If G is a nontrivial connected graph with  $\Delta(G) \geq 3$ , then  $i_{[kR]}(G) \geq \frac{|V(G)|(k+1)}{\Delta(G)+1}.$ 

The lower bound presented in Theorem [15](#page-7-2) is tight, which can be seen by analyzing Cartesian products of some paths and cycles. Given arbitrary graphs  $G$  and  $H$ , the Cartesian product of  $G$ and H is the graph  $G \Box H$  with vertex set  $V(G \Box H) = \{(u, v) : u \in V(G), v \in V(H)\}\)$ . Two vertices  $(u_1, v_1)$  and  $(u_2, v_2)$  of  $G\Box H$  are adjacent if and only if either  $u_1 = u_2$  and  $v_1v_2 \in E(H)$ ; or  $v_1 = v_2$ and  $u_1u_2 \in E(G)$ . Let  $P_2 = (w_1, w_2)$  be a path with two vertices and  $C_{4p} = (v_1, v_2, \ldots, v_{4p})$  be a cycle with [4](#page-7-3)p vertices,  $p \geq 1$ . As an example, Figure 4 shows the graph  $P_2 \square C_8$ . By Theorem [15,](#page-7-2)  $i_{[k]R]}(P_2 \Box C_{4p}) \geq 2p(k+1)$ . In addition, a [k]-IRDF of  $P_2 \Box C_{4p}$  with weight  $2p(k+1)$  is easily obtained by assigning label  $k + 1$  to the set of vertices  $\{(w_1v_i): i = 2t, t \equiv 1 \pmod{2}\} \cup \{(w_2v_j): j = 4t, t \ge 1\}.$ Therefore,  $i_{[k]}(P_2 \Box C_{4p}) = 2p(k+1)$  and, by Theorem [14,](#page-6-2)  $P_2 \Box C_{4p}$  is independent [k]-Roman for all  $k \geq 4$ .

<span id="page-7-3"></span>

FIGURE 4. Cartesian product  $P_2 \Box C_8$  with an  $i_{[kR]}$ -function.

# 4. THE INFINITE FAMILY OF GENERALIZED BLANUŠA SNARKS

<span id="page-7-0"></span>A cut-edge of a graph G is an edge whose deletion increases the number of connected components of G. A snark is a connected 3-regular graph G without cut-edges that does not admit an assignment of labels  $f: E(G) \to \{1,2,3\}$  to its edges so that any two adjacent edges receive distinct labels. The origin of snarks is connected with the Four-Color Problem [\[19\]](#page-17-19) and their study began in 1898 when the first snark was constructed by Petersen [\[20\]](#page-17-20). In 1946, Blanuša constructed two snarks, called Blanuša snarks [\[21\]](#page-18-0). From Blanuša snarks, Watkins constructed two infinite families of snarks, called Generalized Blanuša Snarks [\[22\]](#page-18-1), which are considered in this section.

Luiz [\[16\]](#page-17-16) determined the exact value of the parameter  $i_{1R}(G)$  for every generalized Blanuša snark G. Therefore, in this section, we only analyze values of  $i_{[kR]}$  for the generalized Blanuša snarks for values of  $k \geq 2$ .

<span id="page-8-0"></span>The members of the family of *generalized Blanuša snarks* are graphs formed from subgraphs called construction blocks, denoted  $B_0^1$ ,  $B_0^2$  and L (see Figure [5\)](#page-8-0). A generalized Blanuša snark contains as subgraphs one of the graphs  $B_0^1, B_0^2$  and i copies of the graph L, called  $L_1, L_2, \ldots, L_i$ . Vertices a, b, c and d, belonging to both  $B_0^1$  and  $B_0^2$ , and the vertices  $x_j$ ,  $y_j$ ,  $w_j$  and  $z_j$ , belonging to L, are called border vertices.



FIGURE 5. Construction blocks  $B_0^1$ ,  $B_0^2$  and L of the generalized Blanuša snarks.

In the next paragraphs, we define these families of graphs based on a recursive construction. Let  $\mathfrak{B}^1 = \{B_1^1, B_2^1, B_3^1, \ldots\}$  and  $\mathfrak{B}^2 = \{B_1^2, B_2^2, B_3^2, \ldots\}$  be the first and the second families of generalized Blanuša snarks, respectively. The first member of  $\mathfrak{B}^1$ , the snark  $B_1^1$ , has vertex set  $V(B_1^1) = V(B_0^1) \cup$  $V(L_1)$  and edge set  $E(B_1^1) = E(B_0^1) \cup E(L_1) \cup \{cy_1, dx_1, az_1, bw_1\}$  (see Figure [6\(a\)\)](#page-8-1). The second snark  $\text{Im }\mathfrak{B}^1\text{, snark }B_2^1\text{, has vertex set }V(B_2^1)=V(B_0^1)\cup V(L_1)\cup V(L_2)\text{ and edge set }E(B_2^1)=E(B_0^1)\cup E(L_1)\cup E(L_2)\text{ and }E(L_1^1)=E(L_1^1)\cup E(L_1)\text{ and }E(L_2^1)=E(L_1^1)\cup E(L_2)\text{ and }E(L_1^1)=E(L_1^1)\cup E(L_1)\text{ and }E(L_2^1)=E(L_1^1)\cup E(L_2)\text{ and }E(L_1^1)=E(L_1^1)\cup E(L_1)\text{ and }E(L_2^$  $E(L_2) \cup \{cy_1, dx_1, w_1y_2, z_1x_2, az_2, bw_2\}$  (see Figure [6\(b\)\)](#page-8-1). The smallest snark of family  $\mathfrak{B}^2$ , graph  $B_1^2$ , has vertex set  $V(B_1^2) = V(B_0^2) \cup V(L_1)$  and edge set  $E(B_1^2) = E(B_0^2) \cup E(L_1) \cup \{bw_1, az_1, cy_1, dx_1\}$ (see Figure [7\(a\)\)](#page-8-2). The second snark in  $\mathfrak{B}^2$ ,  $B_2^2$ , has vertex set  $V(B_2^2) = V(B_0^2) \cup V(L_1) \cup V(L_2)$  and edge set  $E(B_2^2) = E(B_0^2) \cup E(L_1) \cup E(L_2) \cup \{bw_2, az_2, cy_1, dx_1, z_1x_2, w_1y_2\}$  (see Figure [7\(b\)\)](#page-8-2).

<span id="page-8-1"></span>

FIGURE 6. The first two smallest members of the family  $\mathfrak{B}^1$ .

<span id="page-8-2"></span>

FIGURE 7. The first two smallest members of the family  $\mathfrak{B}^2$ .

In order to construct larger generalized Blanuša snarks, we use a subgraph  $LG_i$ , called *link graph*, with vertex set  $V(LG_i) = V(L_{i-1}) \cup V(L_i)$  and edge set  $E(LG_i) = E(L_{i-1}) \cup E(L_i) \cup \{w_{i-1}y_i, z_{i-1}x_i\}$ (see Figure [8\)](#page-9-0). Let  $t \in \{1,2\}$ . For each integer i, with  $i \geq 3$ , the snark  $B_i^t$  is obtained recursively from the snark  $B_{i-2}^t$  and the link graph  $LG_i$  according to the following rules:

- (i)  $V(B_i^t) = V(B_{i-2}^t) \cup V(LG_i);$
- <span id="page-9-0"></span>(ii)  $E(B_i^t) = (E(B_{i-2}^t) \setminus E_{i-2}^{out}) \cup E(LG_i) \cup E_i^{in}$ , where
	- $E_{i-2}^{out} = \{az_{i-2}, bw_{i-2}\};$  and
	- $E_i^{in} = \{w_{i-2}y_{i-1}, z_{i-2}x_{i-1}, az_i, bw_i\}.$



Figure 8. The link graph  $LG_i$ .

Theorem [16](#page-9-1) establishes an upper bound for the independent [k]-Roman domination number of generalized Blanuša snarks.

<span id="page-9-1"></span>**Theorem 16.** Let  $k \geq 2$  be an integer. If  $B_i^t$  is a generalized Blanuša snark, with  $t \in \{1,2\}$  and  $i \geq 1$ , then,

$$
i_{[kR]}(B_i^t) \le \begin{cases} (k+1)(2i+2) + 2k & \text{if } t = 1 \text{ and } i \ge 3 \text{ with } i \text{ odd;}\\ (k+1)(2i+3) & \text{otherwise.} \end{cases}
$$

*Proof.* Initially, we separately show that the snark  $B_i^t$ , with  $i = 1$  and  $t \in \{1, 2\}$  has a [k]-IRDF with weight equal to  $5(k + 1) = (k + 1)(2i + 3)$ . This special case is shown in Figure [9,](#page-9-2) with  $B_1^1$  and  $B_1^2$ endowed with their respective  $[k]$ -IRDFs.

<span id="page-9-2"></span>

(A) Snark  $B_1^1$  with a [k]-IRDF with weight  $5(k+1)$ .



(B) Snark  $B_1^2$  with a [k]-IRDF with weight  $5(k+1)$ .

FIGURE 9. Independent [k]-Roman domination functions of snarks  $B_1^1$  and  $B_1^2$ with weight  $5(k+1)$ .

Next, we prove by strong induction on i that every snark  $B_i^t$ , with  $t \in \{1,2\}$  and  $i \geq 2$ , has a [k]-IRDF  $f_i$  with the following properties: (i)  $f_i(a) = k + 1$ ,  $f_i(b) = 0$ ,  $f_i(w_i) = k + 1$  and  $f_i(z_i) = 0$ ; (ii)  $\omega(f_i) = (k+1)(2i+2) + 2k$  if  $t = 1, i \geq 3$  and i odd; or  $\omega(f_i) = (k+1)(2i+3)$  otherwise. We call special a [k]-IRDF  $f_i$  of  $B_i^t$  that satisfies the previous two properties. The induction is based on the recursive construction of the families  $\mathfrak{B}^1$  and  $\mathfrak{B}^2$ .

For the base case, consider the snarks  $B_i^t$  with  $i \in \{2,3\}$  and  $t \in \{1,2\}$ . For  $i = 2$ , Figures [10\(a\)](#page-10-0) and [11\(a\)](#page-10-1) exhibit the snarks  $B_2^1$  and  $B_2^2$ , respectively, with their special [k]-IRDFs with weight  $7(k +$  $1) = (k+1)(2i+3)$ . For  $i = 3$ , the snark  $B_3^1$  is illustrated in Figure [10\(b\)](#page-10-0) with a special [k]-IRDF  $f_3$ with weight  $8(k+1) + 2k$ ; and the snark  $B_3^2$  is illustrated in Figure [11\(b\)](#page-10-1) with a special [k]-IRDF  $f_3$ with weight  $9(k + 1)$ .

For the inductive step, consider a snark  $B_i^t$  with  $i \geq 4$  and  $t \in \{1, 2\}$ . By the recursive construction of generalized Blanuša snarks, we know that  $B_i^t$  can be constructed from the link graph  $LG_i$  and the snark  $B_{i-2}^t$ . Figure [12](#page-11-0) shows the link graph  $LG_i$  with a vertex labeling  $\varphi: V(LG_i) \to \{0, k+1\}$ with weight 4(k + 1). Also, by induction hypothesis, the snark  $B_{i-2}^t$  has a special [k]-IRDF  $f_{i-2}$ with weight  $\omega(f_{i-2}) = (k+1)(2(i-2)+2) + 2k$  when  $t = 1$  and  $i \geq 3$ , i odd; or with weight  $\omega(f_{i-2}) = (k+1)(2(i-2)+3)$  otherwise. Since  $f_{i-2}$  is special, we also have that  $f_{i-2}(a) = k+1$ ,

<span id="page-10-0"></span>

(A) Snark  $B_2^1$  with a special [k]-IRDF with weight  $7(k+1)$ .



(B) Snark  $B_3^1$  with a special [k]-IRDF with weight  $8(k + 1) + 2k$ .

FIGURE 10. Special independent  $[k]$ -Roman dominating functions for the snarks  $B_2^1$  and  $B_3^1$ .

<span id="page-10-1"></span> $f_{i-2}(b) = 0, f_{i-2}(w_{i-2}) = k+1, f_{i-2}(z_{i-2}) = 0$ , for  $a, b, w_{i-2}, z_{i-2} \in V(B^t_{i-2})$ . Thus, we define a vertex labeling  $f_i$  for  $B_i^t$  as follows. For every vertex  $v \in V(B_i^t)$ ,

$$
f_i(v) = \begin{cases} f_{i-2}(v) & \text{if } v \in V(B_{i-2}^t) \cap V(B_i^t); \\ \varphi(v) & \text{if } v \in V(LG_i) \cap V(B_i^t). \end{cases}
$$



(A) Snark  $B_2^2$  with a special [k]-IRDF with weight  $7(k + 1).$ 



(B) Snark  $B_3^2$  with a special [k]-IRDF with weight  $9(k + 1)$ .

FIGURE 11. Special independent [k]-Roman dominating functions for  $B_2^2$  and  $B_3^2$ .

Next, we prove that  $f_i$  is an [k]-IRDF of  $B_i^t$ . By induction hypothesis, the [k]-IRDF  $f_{i-2}$  of  $B_{i-2}^t$ is such that  $f_{i-2}(a) = k+1$ ,  $f_{i-2}(b) = 0$ ,  $f_{i-2}(w_{i-2}) = k+1$ ,  $f_{i-2}(z_{i-2}) = 0$ . This implies that the labeling  $f_i$  restricted to subgraph  $B_{i-2}^t - E_{i-2}^{out} \subset B_i^t$  is almost a [k]-IRDF of  $B_{i-2}^t - E_{i-2}^{out}$  since



<span id="page-11-0"></span>FIGURE 12. Link graph  $LG_i$  with a vertex labeling  $\varphi$ . Note that the vertices  $y_{i-1}, z_i$ and its neighbors have label 0.

 $z_{i-2}$  and b are the only vertices with label 0 in  $B_{i-2}^t - E_{i-2}^{out}$  such that  $f(N[z_{i-2}]) < |AN(z_{i-2})| + k$ and  $f(N[b]) < |AN(b)| + k$ . Also, by construction, the labeling  $f_i$  restricted to subgraph  $LG_i \subset B_i^t$ assigns label 0 to vertices  $y_{i-1}$  and  $z_i$  an these are the only vertices with label 0 in  $LG_i$  that have  $f(N[y_{i-1}]) < |AN(y_{i-1})| + k$  and  $f(N[z_i]) < |AN(z_i)| + k$ . Additionally, no two vertices with label  $k+1$  in  $LG_i$  are adjacent. Thus,  $f_i$  restricted to  $LG_i$  is almost a [k]-IRDF of  $LG_i$  since  $y_{i-1}$  and  $z_i$ are the only vertices of  $LG_i$  that have label 0 and  $f(N[y_{i-1}]) = f(N[z_i]) = 0$ . Therefore, in order to prove that  $f_i$  is a [k]-IRDF of  $B_i^t$ , it suffices to show that the vertices  $y_{i-1}, z_{i-2}, z_i, b$  have a neighbor in  $B_i^t$  with label  $k+1$ . This comes down to analyzing the labels of the endpoints of the edges in the set  $E_i^{in} = \{w_{i-2}y_{i-1}, z_{i-2}x_{i-1}, az_i, bw_i\}$  and verify if the vertices  $w_{i-2}, x_{i-1}, a, w_i$  have label  $k+1$ . From the definition of  $f_i$ , we have that  $f_i(w_{i-2}) = f_{i-2}(w_{i-2}) = k+1$ ,  $f_i(x_{i-1}) = \varphi(x_{i-1}) = k+1$ ,  $f_i(a) = f_{i-2}(a) = k+1$  and  $f_i(w_i) = \varphi(w_i) = k+1$ . Thus, the vertices  $y_{i-1}, z_{i-2}, z_i, b$  (that have label 0) are adjacent to vertices with label  $k+1$  in  $B_i^t$ , that is, the function  $f_i$  is a [k]-IRDF of  $B_i^t$ .

Now, we prove that  $f_i$  is special. The weight of  $f_i$  is given by the sum of the weights of the functions  $f_{i-2}$  and  $\varphi$ . Thus, if  $t = 1$ ,  $i \geq 5$  and i odd, then  $\omega(f_i) = \omega(f_{i-2}) + \omega(\varphi) = (k+1)(2(i-1))$  $2) + 2$  + 2k + 4(k + 1) = (k + 1)(2i + 2) + 2k; otherwise, we have that  $\omega(f_i) = \omega(f_{i-2}) + \omega(\varphi)$  =  $(k+1)(2(i-2)+3)+4(k+1)=(k+1)(2i+3)$ . Note that  $f_i(a)=k+1$ ,  $f_i(b)=0$ ,  $f_i(w_i)=k+1$ and  $f_i(z_i) = 0$  since these are the labels of each of these vertices in the subgraphs  $B_{i-2}^t$  and  $LG_i$ . Therefore,  $f_i$  is a special [k]-IRDF of  $B_i^t$ , and the result follows.

By Theorem [14,](#page-6-2)  $i_{[k]}(B_i^t) \ge (k+1)(2i+2.5)$  for  $k \ge 4$ . However, for increasingly larger values of  $k$ , this lower bound moves away from the upper bounds given in Theorem [16.](#page-9-1) Therefore, better lower bounds are needed. Theorems [20](#page-12-0) and [21](#page-12-1) establish better lower bounds for the parameter  $i_{[kR]}(B_i^t)$ . In order to prove these results, we first present some additional definitions and auxiliary lemmas and theorems.

Given a graph G and two disjoint sets  $S_1 \subset V(G)$  and  $S_2 \subset V(G)$ , we denote by  $E(S_1, S_2)$  the set of edges  $uv \in E(G)$  such that  $u \in S_1$  and  $v \in S_2$ . Also, given  $S \subseteq V(G)$ , we denote by  $N(S)$  the set of vertices  $\{w \in V(G) \backslash S : uw \in E(G) \text{ and } u \in S\}$ . We also define  $N[S] = S \cup N(S)$ .

<span id="page-11-3"></span>**Lemma 17.** Let 
$$
k \geq 2
$$
 be an integer. If G is a 3-regular graph with n vertices and  $f = (V_0, V_k, V_{k+1})$  is an  $i_{[k]}$ -function of G, then  $|V_k| \leq \frac{8i_{[k]}(G) - 2(k+1)n}{3k-5}$  and  $|V_{k+1}| \geq \frac{2kn - 5i_{[k]}(G)}{3k-5}$ .

*Proof.* Let G be a 3-regular graph with n vertices and  $f = (V_0, V_k, V_{k+1})$  be an  $i_{[kR]}$ -function of G. Thus,  $i_{[kR]}(G) = \omega(f) = k|V_k| + (k+1)|V_{k+1}|$ . This fact implies that

.

<span id="page-11-1"></span>(2) 
$$
|V_{k+1}| = \frac{i_{[k]}(G) - k|V_k|}{k+1} \quad \text{and} \quad |V_k| = \frac{i_{[k]}(G) - (k+1)|V_{k+1}|}{k}
$$

Since  $k > 2$ , each vertex  $v \in V(G)$  with  $f(v) = 0$  has at least one neighbor with label  $k + 1$  or at least two neighbors with label k. Let  $S = V_0 \cap N(V_{k+1})$  and  $T = V_0 \setminus S$ . Since G is 3-regular, each vertex in  $V_{k+1}$  is adjacent to at most 3 vertices in S. Thus,  $|S| \leq 3|V_{k+1}|$ . Similarly, since each vertex in  $V_k$  is adjacent to at most 3 vertices in T and since each vertex in T has at least two neighbors in  $V_k$ , we obtain that  $2|T| \leq |E(V_k, T)| \leq 3|V_k|$ , which imples that  $|T| \leq \frac{3|V_k|}{2}$ . Therefore,  $|V_0| = |S| + |T| \leq 3|V_{k+1}| + \frac{3|V_k|}{2}$  $\frac{v_{k\perp}}{2}$ .

From the definition of [k]-IRDF, it follows that  $n = |V_0| + |V_k| + |V_{k+1}|$ . Hence,  $n = |V_0| + |V_k| +$  $|V_{k+1}| \leq 3|V_{k+1}| + \frac{3|V_k|}{2} + |V_k| + |V_{k+1}| = 4|V_{k+1}| + \frac{5|V_k|}{2}$  $\frac{v_{k}}{2}$ , that is,

<span id="page-11-2"></span>(3) 
$$
n \le 4|V_{k+1}| + \frac{5|V_k|}{2}
$$

From Equation [\(2\)](#page-11-1) and Inequality [\(3\)](#page-11-2), we have that  $n \leq 4 \cdot \frac{i_{[k]}(G) - k|V_k|}{k+1} + \frac{5|V_k|}{2} = \frac{8i_{[k]}(G) - (3k-5)|V_k|}{2(k+1)}$ . From this last inequality, we conclude that  $|V_k| \leq \frac{8i_{[kR]}(G) - 2(k+1)n}{3k-5}$ . Also, from Equation [\(2\)](#page-11-1) and Inequality [\(3\)](#page-11-2), we have that  $n \leq 4|V_{k+1}| + \frac{5i_{[k]}(G) - 5(k+1)|V_{k+1}|}{2k} = \frac{8k|V_{k+1}| + 5i_{[k]}(G) - 5(k+1)|V_{k+1}|}{2k} =$  $5i_{[kR]}(G)+(3k-5)|V_{k+1}|$  $\frac{(3k-5)|V_{k+1}|}{2k}$ . From this last inequality, we conclude that  $|V_{k+1}| \ge \frac{2kn-5i_{[kR]}(G)}{3k-5}$ . □

<span id="page-12-3"></span>**Lemma 18.** Let G be a graph and  $k \ge 1$  be an integer. For any  $i_{[k]}$ -function  $f = (V_0, V_k, V_{k+1})$  of G, we have that  $|V_{k+1}| \leq i_{kR}(G) - k \cdot i(G)$  and  $|V_k| \geq (k+1)i(G) - i_{kR}(G)$ .

*Proof.* Let G be a graph with an  $i_{[k]R]}$ -function  $f = (V_0, V_k, V_{k+1})$ . Since  $V_k \cup V_{k+1}$  is an independent dominating set of G, we have  $i(G) \leq |V_k| + |V_{k+1}|$ . Hence,  $k \cdot i(G) \leq k|V_k| + k|V_{k+1}| = k|V_k| +$  $(k+1)|V_{k+1}| - |V_{k+1}| = i_{[k]}(G) - |V_{k+1}|$ . This implies that  $|V_{k+1}| \leq i_{[k]}(G) - k \cdot i(G)$ . In addition,  $(k+1)i(G) \leq (k+1)|V_k| + (k+1)|V_{k+1}| = i_{[k]}(G) + |V_k|$ . This implies that  $|V_k| \geq (k+1)i(G) - i_{[k]}(G)$ , and the result follows.

The next result is used in our proofs and determines the domination number and independent domination number for generalized Blanuša snarks.

<span id="page-12-2"></span>**Theorem 19** (A. Pereira [\[23\]](#page-18-2)). Let  $B_i^t$  be a generalized Blanuša snark with  $t \in \{1,2\}$  and  $i \geq 1$ . Then,

$$
i(B_i^t) = \gamma(B_i^t) = \begin{cases} 2i + 4 & \text{if } t = 1 \text{ and } i \ge 3 \text{ with } i \text{ odd;}\\ 2i + 3 & \text{otherwise.} \end{cases}
$$

<span id="page-12-0"></span>**Theorem 20.** Let  $k \geq 2$  be an integer. Let  $B_i^t$  be a generalized Blanuša snark such that  $t = 1$  and  $i \geq 3$  with i odd. Then,

$$
i_{[kR]}(B_i^t) \ge (k+1)(2i+2) + 2k - 2.
$$

*Proof.* By the definition of  $B_i^t$ , we have that  $|V(B_i^t)| = 8i + 10$ . Define  $n = 8i + 10$ . Let  $f =$  $(V_0, V_k, V_{k+1})$  be an  $i_{[k]}$ -function of  $B_i^t$ . For the purpose of contradiction, suppose that  $i_{[k]}(B_i^t) \leq$  $(k+1)(2i+2)+2k-3$ . Next, we find a lower bound for  $|V_k|$ . By Theorem [19](#page-12-2) and Theorem [18,](#page-12-3)  $|V_k| \ge (k+1)i(G)-i_{[k]}(G) \ge (k+1)(2i+4)-[(k+1)(2i+2)+2k-3]=5.$  Thus,  $|V_k| \ge 5.$  Next, we find an upper bound for  $|V_k|$ . By Lemma [17,](#page-11-3)  $|V_k| \le \frac{8i_{[k]}(B_i^t) - 2(k+1)n}{3k-5} \le \frac{8((k+1)(2i+2)+2k-3)-2(k+1)(8i+10)}{3k-5} =$  $\frac{12k-28}{3k-5}$  < 4 for all  $k \ge 2$ . That is,  $|V_k|$  < 4. However, these facts imply that  $5 \le |V_k|$  < 4, which is a  $\Box$ contradiction.

<span id="page-12-1"></span>**Theorem 21.** Let  $k \geq 4$  be an integer. Let  $B_i^t$  be a generalized Blanuša snark such that  $t = 2$ , or  $t = 1$  with  $i = 1$ , or  $t = 1$  with i even. Then,

$$
i_{[kR]}(B_i^t) = (k+1)(2i+3).
$$

*Proof.* Let  $k \geq 4$  be an integer. By Theorem [16,](#page-9-1)  $i_{[kR]}(B_i^t) \leq (k+1)(2i+3)$ . So, in order to conclude the proof, it suffices to prove that  $i_{[k]}(B_i^t) \ge (k+1)(2i+3)$ . By the definition of  $B_i^t$ , we have that  $|V(B_i^t)| = 8i + 10$ . Define  $n = 8i + 10$ . Let  $f = (V_0, V_k, V_{k+1})$  be an  $i_{[k]}$ -function of  $B_i^t$ . For the purpose of contradiction, suppose that  $i_{[kR]}(B_i^t) \leq (k+1)(2i+3) - 1$ .

By Lemma [17,](#page-11-3)  $|V_k| \le \frac{8i_{[kR]}(B_i^k) - 2(k+1)n}{3k-5} \le \frac{8((k+1)(2i+3)-1) - 2(k+1)(8i+10)}{3k-5} = \frac{(k+1)[8(2i+3)-2(8i+10)] - 8}{3k-5} = \frac{4k-4}{3k-5}$ . That is,  $|V_k| \le \frac{4k-4}{3k-5}$ . For  $k \ge 4$ , we have that  $\frac{4k-4}{3k-5} < 2$ . This imp  $k \geq 4$ . On the other hand, by Lemma [18](#page-12-3) and Theorem [19,](#page-12-2)  $|V_k| \geq (k+1)i(B_i^t) - i_{[kR]}(B_i^t) \geq$  $(k+1)(2i+3) - (k+1)(2i+3) + 1 = 1$ . These facts imply that  $|V_k| = 1$ .

By the definition of [k]-RDF,  $i_{[k]R}(B_i^t) = k|V_k| + (k+1)|V_{k+1}| = k + (k+1)|V_{k+1}|$ . Moreover, since  $i(B_i^t) = 2i + 3$ , we have that  $2i + 3 = i(B_i^t) \leq |V_k| + |V_{k+1}| = 1 + |V_{k+1}|$ , which implies that  $|V_{k+1}| \geq 2i+2$ . Hence,  $i_{[k]}(B_i^t) = (k+1)|V_{k+1}|+k \geq (k+1)(2i+2)+k$ . From these facts, we have that  $(k+1)(2i+2)+k \leq i_{[k]}(B_i^t) \leq (k+1)(2i+3)-1.$  However, since  $(k+1)(2i+2)+k = (k+1)(2i+3)-1,$ we obtain that  $i_{[k]}(B_i^t) = (k+1)(2i+2) + k$ . Since  $i_{[k]}(B_i^t) = (k+1)(2i+2) + k$  and  $|V_k| = 1$  we obtain that  $|V_{k+1}| = 2i + 2$ .

Since  $B_i^t$  is 3-regular, each vertex in  $V_{k+1}$  dominates at most 3 vertices in  $V_0$ . Thus,  $|N(V_{k+1})| \le$  $3|V_{k+1}|$ . This implies that  $|N[V_{k+1}]| = |V_{k+1}| + |N(V_{k+1})| \le (2i+2) + 3(2i+2) = 8i+8$ . In other words, there are at most  $8i + 8$  vertices that are either in  $V_{k+1}$  or are dominated by vertices in  $V_{k+1}$ . Since  $|V(B_i^t)| = 8i + 10$ , it remains  $|V(B_i^t)| - (8i + 8) = 2$  vertices in the set  $V_0 \cup V_k$  that are not

dominated by vertices with label  $k + 1$ . One of these vertices belong to the set  $V_k$ , since  $|V_k| = 1$ , and the other vertex, say w, belongs to the set  $V_0$ . Since  $f(w) = 0$  and w has no neighbor in the set  $V_{k+1}$ , we conclude that  $f(N[w]) < k + |AN(w)|$ , which is a contradiction.

<span id="page-13-3"></span>**Corollary 22.** Let  $k \geq 4$  be an integer. If  $B_i^t$  is a generalized Blanuša snark, with  $t = 2$ , or  $t = 1$ with  $i = 1$ , or  $t = 1$  with i even, then  $B_i^t$  is an independent [k]-Roman graph.

<span id="page-13-0"></span>*Proof.* By Theorem [21](#page-12-1) and Theorem [19,](#page-12-2) we have that  $i_{[k]}(B_i^t) = (k+1)(2i+3) = (k+1)i(B_i^t)$ . Therefore,  $B_i^t$  is an independent [k]-Roman graph.

## 5. The infinite family of Loupekine Snarks

Around 1975, F. Loupekine proposed a method of construction of infinite families of snarks using subgraphs of other known snarks. Loupekine's method was originally presented by Isaacs [\[24\]](#page-18-3) in 1976. In this section, we consider two subfamilies of Loupekine snarks, called  $LP_1$ -snarks and  $LP_0$ -snarks, which are both obtained from fixed subgraphs called basic blocks. A basic block  $B_i$  is illustrated in Figure [13.](#page-13-1) Note that  $B_i$  has five different degree-2 vertices, namely  $r_i, s_i, t_i, u_i, v_i$ , called border vertices. The construction of the two families is described in the next paragraphs.

<span id="page-13-1"></span>Let  $\ell \geq 3$  be an odd integer. An  $\ell$ -LP<sub>1</sub>-snark  $G_L$  is constructed from  $\ell$  basic blocks  $B_0, B_1, \ldots, B_{\ell-1}$ . For each  $i \in \{0, \ldots, \ell-1\}$ , we connect the border vertices  $s_i$  and  $v_i$  of block  $B_i$  to the border vertices  $r_{i+1}$  and  $u_{i+1}$  of block  $B_{i+1}$  (indexes taken modulo  $\ell$ ) with a pair of edges from the set  $E_{i,i+1}$  that comprises either a pair of laminar edges  $\{s_i r_{i+1}, v_i u_{i+1}\}$  or a pair of intersecting edges  $\{s_i u_{i+1}, v_i r_{i+1}\}$ , but not both. These edges connecting two consecutive basic blocks are called *plug-edges*.



FIGURE 13. Basic Block  $B_i$ .

Note that, after the addition of the plug-edges, the border vertices  $t_i$ , with  $0 \le i \le \ell - 1$ , still have degree 2. Thus, in the next step of the construction, any three distinct border vertices  $t_i, t_j, t_s$ , all of them with degree two, are linked to a new vertex  $z_{i,j,s}$ , called *link-vertex*, by adding vertex  $z_{i,j,s}$  and three new edges  $t_i z_{i,j,s}$ ,  $t_j z_{i,j,s}$  and  $t_s z_{i,j,s}$  to  $G_L$ . The previous operation can be done an odd number q of times, with  $1 \le q \le \lfloor \ell/3 \rfloor$ . Since  $\ell$  is odd, an even number  $\ell - 3q$  of border vertices with degree two remain. If  $\ell - 3q > 0$ , the remaining border vertices with degree two are paired up and each pair  $t_i$  and  $t_j$  is linked by a new edge  $t_it_j$ , called a repairing edge, thus concluding the construction of an  $\ell$ -LP<sub>1</sub>-snark  $G_L$ . Figure [14](#page-13-2) shows a 5-LP<sub>1</sub>-snark.

<span id="page-13-2"></span>

FIGURE 14. An  $LP_1$ -snark with 5 basic blocks and one linkvertex.

In the previous construction, if each link-vertex of  $G_L$  is connected to three consecutive basic blocks  $B_i, B_{i+1}, B_{i+2}$ , and if all repairing edges also connect consecutive basic blocks (are of the form  $t_i t_{i+1}$ ),

<span id="page-14-0"></span>then  $G_L$  is said to be an  $\ell$ -LP<sub>0</sub>-snark. Figure [15](#page-14-0) shows a 5-LP<sub>0</sub>-snark. Every LP<sub>0</sub>-snark is also an  $LP_1$ -snark but the converse is not true.



FIGURE 15. An  $LP_0$  with 5 basic blocks and one link-vertex.

Theorem [23](#page-14-1) establishes an upper bound for the independent [k]-Roman domination number of  $LP_1$ -snarks.

<span id="page-14-1"></span>**Theorem 23.** Let  $k \geq 1$  be an integer. Let  $G_L$  be an  $\ell$ -LP<sub>1</sub>-snark with  $\sigma$  link-vertices, with  $\ell \geq 3$ ,  $\ell$ odd and  $\sigma \geq 1$ . Then,  $i_{[kR]}(G_L) \leq 2(k+1)\ell + k\sigma$ .

*Proof.* For each  $i \in \{0, 1, \ldots, \ell - 1\}$ , a function  $g_i: V(B_i) \to \{0, k + 1\}$  for basic block  $B_i$  is defined in Figure [16.](#page-14-2) Note that  $g_i$  has weight  $\omega(g_i) = 2(k+1)$ . Also, note that, under  $g_i$ , every vertex of  $B_i$  with label 0 is adjacent to a vertex of  $B_i$  with label  $k+1$  and no two vertices with label  $k+1$  are adjacent. Hence,  $g_i$  is an [k]-IRDF of  $B_i$ .

Define a function  $f: V(G_L) \to \{0, k, k+1\}$  for  $G_L$  as follows. For each vertex  $v \in V(G_L)$ ,

$$
f(v) = \begin{cases} g_i(v) & \text{if } v \in V(B_i), \text{ for } 0 \le i \le \ell - 1; \\ k & \text{if } v \text{ is a link-vertex.} \end{cases}
$$

<span id="page-14-2"></span>The weight of f is given by  $\omega(f) = \sum_{i=0}^{\ell-1} \omega(g_i) + k\sigma = 2(k+1)\ell + k\sigma$ . It remains to show that f is a [k]-IRDF. Note that the vertices that receive label  $k+1$  are the vertices  $p_i$  and  $q_i$ , for  $0 \le i \le \ell-1$ , and these vertices form an independent set of  $G<sub>L</sub>$ . Moreover, the set of link-vertices also form an independent set of  $G_L$ . Every link-vertex has label k and is adjacent to three vertices that have label 0. Therefore, no two vertices with labels  $k$  or  $k+1$  are adjacent and, as previously argued, every vertex of  $B_i$  with label 0 is adjacent to a vertex with label  $k+1$  that also belongs to  $B_i$ . Therefore, f is a [k]-IRDF of  $G_L$  with weight  $2(k+1)\ell + k\sigma$ .



FIGURE 16. Basic block  $B_i$  with a [k]-IRDF  $g_i$  with weight  $2(k+1)$ .

Theorem [24](#page-15-0) shows a better upper bound for  $i_{[kR]}$  than that shown in Theorem [23](#page-14-1) when restricted to the subfamily of  $LP_0$ -snarks. Next, we define two subgraphs of an  $LP_0$ -snark that are used in our proof.

Let  $G_L$  be an  $\ell$ -LP<sub>0</sub>-snark. Given a repairing edge  $t_i t_{i+1}$  of  $G_L$ , a *double gadget*  $G_d$  of  $G_L$  is a subgraph of  $G_L$  induced by the set of vertices  $V(B_i) \cup V(B_{i+1})$ . On the other hand, given a linkvertex  $z_{i,i+1,i+2}$  of  $G_L$ , a triple gadget  $G_t$  of  $G_L$  is a subgraph of  $G_L$  induced by the set of vertices  $V(B_i) \cup V(B_{i+1}) \cup V(B_{i+2}) \cup \{z_{i,i+1,i+2}\}.$  Figure [17](#page-15-1) shows a scheme of a double gadget and a triple gadget.

<span id="page-15-1"></span>

FIGURE 17. Gadgets of an  $LP_0$ -snark. Dashed edges represent the possible configurations for plug-edges connecting two consecutive blocks: either laminar edges or intersecting edges.

<span id="page-15-0"></span>**Theorem 24.** Let  $k \geq 1$  be an integer. Let  $G_L$  be an  $\ell$ -LP<sub>0</sub>-snark with  $\sigma$  link-vertices, with  $\sigma \geq 1$ and odd  $\ell \geq 3$ . Then,  $i_{[kR]}(G_L) \leq 2(k+1)\ell + (k-1)\sigma$ .

*Proof.* Let  $G_L$  be an  $\ell$ -LP<sub>0</sub>-snark. In order to prove the theorem, we construct a [k]-IRDF  $f: V(G_L) \to$  $\{0, k, k+1\}$  of  $G_L$  with weight  $2(k+1)\ell + (k-1)\sigma$ . Let  $G_1, G_2, \ldots, G_r$  be gadgets of  $G_L$  given in cyclic sequence, with  $r \geq 1$ . Figure [18\(a\)](#page-15-2) exhibits an [k]-IRDF  $\pi_d : V(G_d) \to \{0, k+1\}$  with weight  $4(k + 1)$  for double gadgets and Figure [18\(b\)](#page-15-2) exhibits an [k]-IRDF  $\pi_t: V(G_t) \to \{0, k, k+1\}$  with weight  $5(k+1) + 2k$  for triple gadgets. Note that, in both labelings  $\pi_d$  and  $\pi_t$ , the set of vertices with label k or  $k + 1$  form an independent set and every vertex with label 0 is adjacent to a vertex with label  $k + 1$ . Hence,  $\pi_d$  and  $\pi_t$  are [k]-IRDFs.

For every vertex  $v \in V(G_L)$ , define:  $f(v) = \pi_d(v)$  if v belongs to a double gadget  $G_d \subset G_L$ ; or  $f(v) = \pi_t(v)$ , otherwise. Next, we prove that the labeling f is a [k]-IRDF of  $G_L$ . By the definition of f, we have that f restricted to double gadgets or triple gadgets is a  $[k]$ -IRDF. Thus, it remains to show that  $V_k \cup V_{k+1}$  is an independent set of  $G_L$  (Recall that  $V_i \subseteq V(G_L)$  is the set of vertices with label i). In order to see this, recall that the set of vertices  $V_k \cup V_{k+1}$  form an independent set when restricted to double or triple gadgets. This property still holds when considering the whole graph since the plug-edges that connect two consecutive gadgets  $G_i$  and  $G_{i+1}$  connect border vertices that have label 0. Therefore, f is an  $[k]$ -IRDF of  $G_L$ .

The weight of f is the sum of the weight of its gadgets. Note that  $\ell = 2p_2 + 3p_3$ , where  $p_2$  is the number of double gadgets and  $p_3 = \sigma$  is the number of triple gadgets. Thus,  $\omega(f) = \omega(\pi_d) \cdot p_2 + \sigma$  $\omega(\pi_t) \cdot p_3 = 4(k+1)p_2 + (5(k+1)+2k)p_3 = 2(k+1)(2p_2+3p_3) + (k-1)p_3 = 2(k+1)\ell + (k-1)p_3 = 2(k+1)\ell + (k-1)p_4$  $2(k+1)\ell + (k-1)\sigma$ .

<span id="page-15-2"></span>

(A) [k]-IRDF  $\pi_d$  of  $G_d$  with weight  $4(k+1)$ .

(b) [k]-IRDF  $\pi_t$  of  $G_t$  with weight  $5(k+1) + 2k$ .

FIGURE 18. Double gadget  $G_d$  with a [k]-IRDF  $\pi_d : V(G_d) \to \{0, k+1\}$  and triple gadget  $G_t$  with a [k]-IRDF  $\pi_t: V(G_t) \to \{0, k, k+1\}.$ 

<span id="page-15-3"></span>Theorem [25](#page-15-3) establishes an upper bound for the  $[k]$ -Roman domination number of  $LP_0$ -snarks.

**Theorem 25.** Let  $k \geq 1$  be an integer. Let  $G_L$  be an  $\ell$ -LP<sub>0</sub>-snark with  $\ell \geq 3$ ,  $\ell$  odd. Then,  $\gamma_{[kR]}(G_L) \leq$  $2(k + 1)\ell$ .

*Proof.* Let  $G_L$  be an  $\ell$ -LP<sub>0</sub>-snark. We construct a [k]-RDF  $f: V(G_L) \to \{0, k, k+1\}$  for  $G_L$  with weight  $2(k+1)\ell$ . Figure [19](#page-16-0) exhibits a  $[k]$ -RDF  $\pi_t: V(G_t) \to \{0, k+1\}$  with weight  $6(k+1)$  for triple gadgets  $G_t \subset G_L$ ; and double gadgets  $G_d \subset G_L$  are assigned the [k]-IRDF  $\pi_d : V(G_d) \to \{0, k+1\}$ with weight  $4(k + 1)$  shown in Figure [18\(a\).](#page-15-2) Note that, in both labelings  $\pi_d$  and  $\pi_t$ , every vertex with label 0 is adjacent to a vertex with label  $k + 1$ . Hence,  $\pi_d$  e  $\pi_t$  are [k]-RDFs. For every vertex  $v \in V(G_L)$ , define:  $f(v) = \pi_d(v)$  if v belongs to a double gadget  $G_d \subset G_L$ ; or  $f(v) = \pi_t(v)$ , otherwise. By the definition of  $f$ , we have that  $f$  restricted to each gadget is a  $[k]$ -RDF (in fact, by the definition of  $\pi_d$  and  $\pi_d$ , any vertex with label 0 is already adjacent to a vertex with label  $k + 1$ ). Hence, f is a  $[k]$ -RDF of  $G_L$ . Furthermore, the weight of f is the sum of the weight of its gadgets. Note that  $\ell = 2p_2 + 3p_3$ , where  $p_2$  is the number of double gadgets and  $p_3$  is the number of triple gadgets. Thus,  $\omega(f) = \omega(\pi_d) \cdot p_2 + \omega(\pi_t) \cdot p_3 = 4(k+1)p_2 + 6(k+1)p_3 = 2(k+1)(2p_2 + 3p_3) = 2(k+1)\ell.$ 

<span id="page-16-0"></span>

FIGURE 19. Triple gadget  $G_t$  with a  $[k]$ -RDF  $\pi_t: V(G_t) \to \{0, k+1\}$  with weight  $6(k+1)$ .

By Theorem [15,](#page-7-2) a k-LP<sub>0</sub>-snark G with n vertices has  $i_{[kR]}(G) \geq \left\lceil \frac{(k+1)n}{4} \right\rceil$  $\frac{(-1)n}{4}$ . However, this lower bound is improved, for all  $k \geq 4$ , in Theorem [27.](#page-16-1) In order to prove it, we need the following lower bound on the domination number of  $LP_0$ -snarks [\[23\]](#page-18-2).

<span id="page-16-2"></span>**Theorem 26** (A. Pereira [\[23\]](#page-18-2)). Let  $G_L$  be an  $\ell$ -LP<sub>0</sub>-snark with  $\ell \geq 3$  and  $\sigma$  link-vertices. Then,  $\gamma(G_L) \geq \lfloor \frac{n}{4} \rfloor + 1$ , where  $n = 7\ell + \sigma$ .

<span id="page-16-1"></span>**Theorem 27.** Let  $k \geq 4$  be an integer. Let  $G_L$  be an  $\ell$ -LP<sub>0</sub>-snark with  $\ell \geq 3$  and  $\sigma$  link-vertices. Then,  $i_{[kR]}(G_L) \geq \left\lceil \frac{(k+1)n}{4} \right\rceil$  $\frac{(-1)n}{4} + 1$ , where  $n = |V(G_L)| = 7\ell + \sigma$ .

*Proof.* Let  $G_L$  and k be as stated in the hypothesis. For the purpose of contradiction, suppose that  $i_{[kR]}(G_L) \leq \left\lceil \frac{(k+1)n}{4} \right\rceil$  $\frac{(-1)n}{4}$ . By Theorem [15,](#page-7-2) we have that  $i_{[kR]}(G_L) \ge \left\lceil \frac{(k+1)n}{4} \right\rceil$  $\frac{(-1)n}{4}$ . From these two inequalities we obtain that  $i_{[k]}(G_L) = \left[\frac{(k+1)n}{4}\right]$  $\frac{(-1)n}{4}$ . This fact along with Theorem [14](#page-6-2) and the fact that  $k \geq 4$ imply that  $G_L$  is independent [k]-Roman, that is,  $i_{[kR]}(G_L) = (k+1)i(G_L)$ . By Theorem [26,](#page-16-2) we obtain that  $i_{[kR]}(G_L) = (k+1)i(G_L) \ge (k+1)\gamma(G_L) \ge (k+1)\left\lfloor \frac{n}{4} \right\rfloor + (k+1)$ . Thus, we have that  $(k+1)n$  $\left\lfloor \frac{n+1}{4} \right\rfloor = i_{[k]}(G_L) = (k+1)i(G_L) \ge (k+1) \left\lfloor \frac{n}{4} \right\rfloor + (k+1)$ , that is,

<span id="page-16-3"></span>(4) 
$$
\left\lceil \frac{(k+1)n}{4} \right\rceil \ge (k+1) \left\lfloor \frac{n}{4} \right\rfloor + (k+1).
$$

Since  $n = 7\ell + \sigma$  and  $\ell$  and  $\sigma$  are odd numbers, we obtain that n is an even number. We split the proof into two cases depending of the value of n modulo 4.

**Case 1.**  $n \equiv 0 \pmod{4}$ . Define  $n = 4p, p \in \mathbb{Z}$ . From inequality [\(4\)](#page-16-3) we have that  $\frac{(k+1)4p}{4}$  $\frac{(-1)4p}{4}$   $\geq$  $(k+1)\left\lfloor \frac{4p}{4} \right\rfloor + (k+1)$ , which implies that  $(k+1)p \ge (k+1)p + (k+1)$ , which is a contradiction.

**Case 2.**  $n \equiv 2 \pmod{4}$ . Define  $n = 4p + 2$ ,  $p \in \mathbb{Z}$ . From inequality [\(4\)](#page-16-3) we have that  $\frac{(k+1)(4p+2)}{4}$  $\frac{(4p+2)}{4}$   $\geq$  $(k+1)\left\lfloor\frac{4p+2}{4}\right\rfloor+(k+1)$ , which implies that  $(k+1)p+\frac{k+1}{2}\geq (k+1)p+(k+1)$ , which is a contradiction. Since both cases led us to a contradiction, we conclude that  $i_{[kR]}(G_L) \geq \left[\frac{(k+1)n}{4}\right]$  $\frac{-1}{4}$  + 1.

### 6. Closing Remarks

<span id="page-17-15"></span>In this work, we prove that, for all  $k \geq 3$ , the independent [k]-Roman domination problem and the  $[k]$ -Roman domination problem are  $\mathcal{NP}$ -complete even when restricted to planar bipartite graphs with maximum degree 3 and also present lower and upper bounds for the parameter  $i_{[kR]}(G)$ . Moreover, we investigate  $i_{[kR]}(G)$  for two families of 3-regular graphs called generalized Blanuša snarks and Loupekine snarks.

In Corollary [22,](#page-13-3) we present an infinite family of independent [k]-Roman graphs, which are graphs that have  $i_{kR}(G) = (k+1)i(G)$ . An interesting open problem is finding other classes of independent  $[k]$ -Roman graphs.

Adabi et al. [\[25\]](#page-18-4) proved that any graph G with  $\Delta(G) \leq 3$ , has  $\gamma_{[k]}(G) = i_{[k]}(G)$  for  $k = 1$ . We remark that the family of planar bipartite graphs with maximum degree 3 constructed in the reduction shown in Section [2](#page-2-2) is an example of infinite family of graphs with  $\Delta(G) = 3$  for which  $\gamma_{kR}(G) = i_{kR}(G)$  for all  $k \geq 1$ . Thus, another interesting line of research is finding other classes of graphs with  $\Delta(G) \leq 3$  for which  $\gamma_{[k]}(G) = i_{[k]}(G)$  for  $k \geq 2$ . In fact, we conjecture that this property holds for all generalized Blanuša snarks.

## 7. Acknowledgments

This work was partially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq.

#### **REFERENCES**

- <span id="page-17-0"></span>[1] T. Haynes, S. Hedetniemi, M. Henning, Domination in Graphs: Core Concepts, Springer Monographs in Mathematics, Springer International Publishing, 2023.
- <span id="page-17-1"></span>[2] E. J. Cockayne, P. A. Dreyer Jr, S. M. Hedetniemi, S. T. Hedetniemi, Roman domination in graphs, Discrete mathematics 278 (1-3) (2004) 11–22.
- <span id="page-17-3"></span><span id="page-17-2"></span>[3] I. Stewart, Defenden the Roman empire!, Scientific American 281 (6) (1999) 136–139.
- [4] C. S. ReVelle, K. E. Rosing, Defendens Imperium Romanum: A Classical Problem in Military Strategy, The American Mathematical Monthly 107 (7) (2000) 585–594.
- <span id="page-17-4"></span>[5] M. Chellali, N. J. Rad, S. Sheikholeslami, L. Volkmann, A survey on Roman domination parameters in directed graphs, J. Combin. Math. Combin. Comput. (115) (2020) 141–171.
- <span id="page-17-5"></span>[6] M. Chellali, N. J. Rad, S. M. Sheikholeslami, L. Volkmann, Varieties of Roman domination II, AKCE International Journal of Graphs and Combinatorics 17 (3) (2020) 966–984.
- <span id="page-17-6"></span>[7] M. Chellali, N. Jafari Rad, S. M. Sheikholeslami, L. Volkmann, Roman Domination in Graphs, Springer International Publishing, Cham, 2020, pp. 365–409.
- <span id="page-17-7"></span>[8] M. Chellali, N. J. Rad, S. M. Sheikholeslami, L. Volkmann, Varieties of Roman Domination, Springer International Publishing, Cham, Switzerland, 2021, pp. 273–307.
- <span id="page-17-8"></span>[9] M. Chellali, N. J. Rad, S. M. Sheikholeslami, L. Volkmann, The Roman Domatic Problem in Graphs and Digraphs: A Survey, Discussiones Mathematicae Graph Theory 42 (3) (2022) 861–891.
- <span id="page-17-9"></span>[10] H. Abdollahzadeh Ahangar, M. Álvarez, M. Chellali, S. Sheikholeslami, J. Valenzuela-Tripodoro, Triple roman domination in graphs, Applied Mathematics and Computation 391 (2021) 125444. [doi:https://doi.org/10.1016/j.amc.2020.125444](https://doi.org/https://doi.org/10.1016/j.amc.2020.125444).
- <span id="page-17-10"></span>[11] M. C. N. Khalili, J. Amjadi, S. M. Sheikholeslami, On [k]-roman domination in graphs, AKCE International Journal of Graphs and Combinatorics 20 (3) (2023) 291–299. [doi:10.1080/09728600.2023.2241531](https://doi.org/10.1080/09728600.2023.2241531).
- <span id="page-17-11"></span>[12] M. C. L. J. C. Valenzuela-Tripodoro, M. A. Mateos-Camacho, M. P. Alvarez-Ruiz, Further results on the [k]-roman domination in graphs, Bulletin of the Iranian Mathematical Society 50 (2) (2024) 1–16. [doi:10.1007/s41980-024-00872-1](https://doi.org/10.1007/s41980-024-00872-1).
- <span id="page-17-12"></span>[13] R. A. Beeler, T. W. Haynes, S. T. Hedetniemi, Double Roman domination, Discrete Applied Mathematics 211 (2016) 23–29. [doi:https://doi.org/10.1016/j.dam.2016.03.017](https://doi.org/https://doi.org/10.1016/j.dam.2016.03.017).
- <span id="page-17-13"></span>[14] J. Amjadi, N. Khalili, Quadruple roman domination in graphs, Discrete Mathematics, Algorithms and Applications 14 (03) (2022) 2150130. [doi:10.1142/S1793830921501305](https://doi.org/10.1142/S1793830921501305).
- <span id="page-17-14"></span>[15] H. Maimani, M. Momeni, S. N. Moghaddam, F. R. Mahid, S. M. Sheikholeslami, Independent double Roman domination in graphs, Bulletin of the Iranian Mathematical Society 46 (2) (2019) 543–555. [doi:10.1007/s41980-019-00274-8](https://doi.org/10.1007/s41980-019-00274-8).
- <span id="page-17-16"></span>[16] A. G. Luiz, Roman domination and independent roman domination on graphs with maximum degree three, Discrete Applied Mathematics 348 (2024) 260–278. [doi:https://doi.org/10.1016/j.dam.2024.02.006](https://doi.org/https://doi.org/10.1016/j.dam.2024.02.006).
- <span id="page-17-17"></span>[17] B. Mohar, Face covers and the genus problem for apex graphs, Journal of Combinatorial Theory, Series B 82 (1) (2001) 102–117. [doi:https://doi.org/10.1006/jctb.2000.2026](https://doi.org/https://doi.org/10.1006/jctb.2000.2026).
- <span id="page-17-18"></span>[18] Z. Shao, P. Wu, H. Jiang, Z. Li, J. Žerovnik, X. Zhang, Discharging approach for double Roman domination in graphs, IEEE Access 6 (2018) 63345–63351. [doi:10.1109/ACCESS.2018.2876460](https://doi.org/10.1109/ACCESS.2018.2876460).
- <span id="page-17-20"></span><span id="page-17-19"></span>[19] P. G. Tait, Remarks on the colouring of maps, Proceedings of the Royal Society of Edinburgh 10 (4) (1880) 501–503.
- [20] J. Petersen, Sur le théorème de tait, L'Intermédiaire des Mathématiciens 5 (1898) 225–227.
- <span id="page-18-1"></span><span id="page-18-0"></span>[21] D. Blanuša, Problem cetiriju boja (croatian), hrvatsko priordoslorno drusštvo glasnik mat-fiz, Astr Ser II 1 (1946) 31–42.
- <span id="page-18-2"></span>[22] J. J. Watkins, On the construction of snarks, Ars Combinatoria 16 (1983) 111–124.
- [23] A. A. Pereira, Dominating sets in cubic graphs, Master thesis (in portuguese), University of Campinas, Institute of Computing, Campinas, S˜ao Paulo, Brazil, Available at: https://hdl.handle.net/20.500.12733/1640296 (2020).
- <span id="page-18-3"></span>[24] R. Isaacs, Loupekhine's snarks: a bifamily of non-Tait-colorable graphs, Journal of Combinatorial Theory, Series B (1976).
- <span id="page-18-4"></span>[25] N. J. R. M. Adabi, E. Ebrahimi Targhi, M. S. Moradi, Properties of independent Roman domination in graphs, Australasian Journal of Combinatorics 52 (2012) 11–18.

FEDERAL UNIVERSITY OF CEARÁ, CAMPUS QUIXADÁ, QUIXADÁ, CEARÁ, BRAZIL Email address: gomes.atilio@ufc.br

FEDERAL UNIVERSITY OF CEARÁ, CAMPUS QUIXADÁ, QUIXADÁ, CEARÁ, BRAZIL

Email address: andersonsilva@alu.ufc.br