

New construction of a vacuum doubly rotating black ring by the Ehlers transformation

Ryotaku Suzuki* and Shinya Tomizawa†

Mathematical Physics Laboratory, Toyota Technological Institute

Hisakata 2-12-1, Tempaku-ku, Nagoya, Japan 468-8511

(Dated: June 18, 2024)

Using the Ehlers transformation, we derive an exact solution for a doubly rotating black ring in five-dimensional vacuum Einstein theory. It is well-known that the vacuum Einstein theory with three commuting Killing vector fields can be reduced to a non-linear sigma model with $SL(3, \mathbb{R})$ target space symmetry. As shown previously by Giusto and Saxena, the $SO(2, 1)$ subgroup in the $SL(3, \mathbb{R})$ can generate a rotating solution from a static solution while preserving asymptotic flatness. This so-called Ehlers transformation actually transforms the five-dimensional Schwarzschild black hole into the five-dimensional Myers-Perry black hole. However, unlike the case with the black hole, applying this method directly to the static black ring or the Emparan-Reall black ring, does not yield a regular rotating black ring due to the emergence of a Dirac-Misner string singularity. To solve this undesirable issue, we use a singular vacuum solution of a rotating black ring/lens that already possesses a Dirac-Misner string singularity as the seed solution for the Ehlers transformation. The resulting solution is regular, indicating the absence of curvature singularities, conical singularities, orbifold singularities, Dirac-Misner string singularities, and closed timelike curves both on and outside the horizon. We show that this solution obtained by the Ehlers transformation coincides precisely with the Pomeransky-Sen'kov solution. We expect that applying this method to other theories may lead to the finding of new exact solutions, such as solutions for black lenses and capped black holes, as well as black ring configurations.

I. INTRODUCTION

The study of black hole solutions in Einstein gravity has proven to be an invaluable platform for examining both classical and quantum gravitational theories. Specifically, the exploration of higher-dimensional black holes has captured significant scientific interest over the past twenty years. This attention is highlighted by achievements such as the microscopic derivation of Bekenstein-Hawking entropy [1] and the theoretical possibility of creating black holes in accelerators due to the presence of large extra dimensions [2]. Despite significant progress in understanding higher-dimensional black holes, our comprehension of higher-dimensional gravity remains superficial, as it encompasses a broader spectrum of complexities and degrees of freedom. According to the topology theorem for stationary black holes in five dimensions [3–5], the topology of the event horizon's cross-section is restricted to a sphere S^3 , a ring $S^1 \times S^2$, or lens spaces $L(p, q)$, provided the spacetime is asymptotically flat and supports two commuting axial Killing vector fields. In the first two cases, we possess established exact solutions to the vacuum Einstein equations [6–9]. By contrast, securing a vacuum black hole solution with the lens space topology, denoted as $L(p, q)$, remains challenging and yet unresolved. Various types of black hole solutions theory have so far been found, with the help of, in part, recent development of solution generating techniques. However, the classification of those black hole solutions has not yet been achieved, even in five-dimensional vacuum Einstein theory, since the development of the solution-generating methods is still insufficient.

Emparan and Reall [8] first found the exact solution for an S^1 -rotating black ring, demonstrating that five-dimensional vacuum Einstein theory admits the existence of an S^1 -rotating spherical black hole and two rotating black rings with identical conserved charges, thereby clearly showcasing the non-uniqueness feature in higher dimensions. In general, since in five dimensions, black holes can rotate around two independent axes, it is natural to consider whether there exists a more general black ring solution with two independent rotations, S^1 rotation and S^2 rotation. The first derivation of a black ring with S^2 -rotation was independently achieved by Mishima and Iguchi [10] and Figueras [11] by different methods, though their solutions, which are the same, contain conical singularities.

*Electronic address: sryotaku@toyota-ti.ac.jp

†Electronic address: tomizawa@toyota-ti.ac.jp

The inverse scattering method (ISM), initially developed by Belinski and Zakharov [12], has proven indispensable for constructing the most general black ring solutions that rotate along both the S^1 and S^2 axes. However, direct application of the ISM to higher dimensions tends to yield singular solutions. Pomeransky adeptly modified the method, successfully applying it to five-dimensional black holes by deriving the five-dimensional Myers-Perry solution from the five-dimensional Schwarzschild solution [13]. The ISM was also shown to facilitate the generation of S^2 -rotating black rings [14]. Nevertheless, crafting the S^1 -rotating black ring using the ISM has proven significantly more challenging due to the inevitable emergence of naked curvature singularities from a regular seed solution. A significant advancement in this direction was made in Refs. [15, 16], which found the appropriate seed, a certain singular solution, to generate the S^1 -rotating black ring. This progress enabled the creation of the doubly rotating black ring using the ISM, and consequently, Pomeransky and Sen'kov successfully obtained the balanced doubly rotating black ring solution [9]. Though they gave only the balanced solution in their paper [9], the unbalanced generalization of the black ring was later presented explicitly in Ref. [17]. Furthermore, a more compact form of this solution was presented by Ref. [18].

Meanwhile, the sigma model approach has been established as one of the most effective methods for generating exact solutions to Einstein's equations [19]. In vacuum Einstein theory with a single Killing vector, this theory can be reduced to a non-linear sigma model coupled with three-dimensional gravity through dimensional reduction [20]. Here, the target space of this sigma model is represented by the coset $SL(2, \mathbb{R})/SO(2)$, which facilitates the generation of new solutions by applying group transformations to the coset representative of a seed solution. Similarly, for the four-dimensional Einstein-Maxwell theory with a single Killing vector, the corresponding coset is $SU(2, 1)/S[U(2) \times U(1)]$ [21]. The $SU(2, 1)$ symmetry provides two significant transformations: the Ehlers transformation [22], which adds a nut charge, and the Harrison transformation [23], which adds an electric or magnetic charge to a given seed solution. However, the $SU(2, 1)$ symmetry cannot simply add rotation to a non-rotating solution while preserving asymptotic flatness. For a stationary and axisymmetric solution, this can be achieved by combining transformations associated with an arbitrarily chosen direction in the two Killing vector space with transitions to other linear combinations of the two Killing vectors. For example, as pointed out by Clement [24, 25], the Kerr solution can be obtained from the Schwarzschild solution using this transformation. As Maison showed [26], in d -dimensional ($d > 4$) vacuum Einstein theory with $(d - 3)$ commuting Killing vectors, the coset is represented by $SL(d - 2, \mathbb{R})/SO(2, d - 4)$. Specifically, for $d = 5$, Giusto and Saxena [27] demonstrated that by using the $SO(2, 1)$ subgroup of the $SL(3, \mathbb{R})$ symmetry, which preserves asymptotic flatness, one can transform a static solution into a one-parameter family of stationary solutions with angular momentum. They illustrated this by deriving the Myers-Perry black hole from the five-dimensional Schwarzschild solution.

Additionally, for five-dimensional minimal supergravity, which is the Einstein-Maxwell-Chern-Simons theory with a specific coupling constant, two Killing vectors with the coset is given by $G_{2(2)}/[SL(2, \mathbb{R}) \times SL(2, \mathbb{R})]$ or $G_{2(2)}/SO(4)$, depending on whether the Killing vectors chosen are a timelike one and a spacelike one, or two spacelike ones, respectively [28, 29]. Utilizing the $G_{2(2)}$ symmetry, Bouchareb et al. [30] developed a method for generating solutions, including an electric Harrison transformation that transforms a five-dimensional vacuum solution into an electrically charged one in five-dimensional minimal supergravity. Applying this transformation to the five-dimensional vacuum rotating black hole (the Myers-Perry solution [7]) results in the five-dimensional charged rotating black hole (the Cvetič-Youm solution [31]). However, as noted by the authors, applying this Harrison transformation to the vacuum doubly rotating black ring (the Pomeransky-Sen'kov solution [9]) does not produce a regular charged doubly spinning black ring solution, as the resulting solution suffers from a Dirac-Misner string singularity. In our previous work [32], we addressed this serious problem and constructed the exact solution for a charged rotating black ring. The key idea was to use the electric Harrison transformation on a vacuum solution of a rotating black ring that possesses a Dirac-Misner string singularity. This transformation also provided the first example of a non-BPS exact solution representing an asymptotically flat, stationary spherical black hole, referred to as a capped black hole, with a domain of outer communication (DOC) having a nontrivial topology, in the sense that the DOC on a timeslice has the topology of $[R^4 \# CP^2] \setminus B^4$ [33, 34].

The aim of this paper is to present a new construction of a vacuum solution for a doubly rotating black ring in five-dimensional Einstein theory using the $SL(3, \mathbb{R})$ transformation in a non-linear sigma model approach. The basic procedure resembles the construction of the charged dipole black ring or the capped black hole solution in our previous works [33, 34], in which we used the black lens solution which possesses a Dirac-Misner string singularity as a seed solution for the Harrison transformation. We also use the same seed solution as for the the Harrison transformation—the singular solution with a Dirac-Misner string singularity—for the $SL(3, \mathbb{R})$ transformation. This singularity will be removed by appropriately adjusting the solution's parameters not before but after the $SL(3, \mathbb{R})$ transformation. By applying the $SL(3, \mathbb{R})$ transformation to this vacuum solution, we add rotation to the vacuum seed solution. After applying the transformation to the seed, we select solution's parameters appropriately to ensure that the resulting solution is regular, free from Dirac-Misner string singularity, and devoid of conical singularities within the black ring.

Consequently, the final solution represents a doubly rotating black ring characterized by its regularity—it does not have curvature singularities, conical singularities, Dirac-Misner string singularities, and orbifold singularities on the rods (which include the rotational axes and the horizon). According to the uniqueness theorem for vacuum black ring solutions [5, 17, 35], our derived solution is expected to be isometric to the well-known Pomeransky-Sen'kov solution [9], provided that the spacetime described by our solution is regular and free from closed timelike curves on and outside the event horizon. However, demonstrating the equivalence of the two solutions is challenging due to the apparent differences arising from the complex and lengthy expressions of our solution, which stem from significant differences in the parameters used in two different solution-generating methods. By analyzing the rod data and asymptotic charges of both solutions, we identify the relationship between the two sets of parameters. Consequently, after correlating the differing parameter sets of the two solutions, we establish that our solution precisely coincides with the Pomeransky-Sen'kov solution.

The remainder of this paper is dedicated to constructing the aforementioned black ring solution. In the next section (Sec. II), we will briefly review the solution-generation method that employs the $SL(3, \mathbb{R})$ invariance to add angular momentum to a vacuum seed solution within the non-linear sigma model framework of five-dimensional Einstein theory. In Sec. III, we will introduce the vacuum solution of the rotating black lens, which includes a Dirac-Misner string singularity. This solution will serve as the seed for the $SL(3, \mathbb{R})$ transformation. Previously, this same vacuum solution was utilized in Refs. [33, 34] to construct a charged black ring solution and a capped black hole, employing it as a seed for the Harrison transformation. In alignment with these earlier works [33, 34], we will retain the Dirac-Misner string singularity, choosing not to eliminate it." In Sec. IV, we will apply the $SL(3, \mathbb{R})$ -transformation to the vacuum solution, leading to a corresponding rotating vacuum solution. By setting appropriate boundary conditions on the parameters, we will ensure the absence of any singularities—including curvature, conical, and Dirac-Misner string singularities—on the rotational axes and the horizon. This will allow us to derive a doubly rotating black ring solution in the C-metric form. Initially, this solution appears to differ from the well-known Pomeransky-Sen'kov solution. However, in Sec.V, we will show that the resulting solution exactly aligns with the Pomeransky-Sen'kov solution. Finally, in Sec. VI, we will summarize our findings and outline future research directions

II. PRELIMINARY

Under the assumption of space-time symmetry, the Einstein equations simplify to equations for scalar fields coupled with a three-dimensional gravity. We then review that this system of scalar fields is governed by a nonlinear sigma model.

A. Vacuum Einstein gravity with two commuting Killing vectors

First, we consider a spacetime with two mutually commuting Killing vector fields ξ_a ($a = 0, 1$) such that $[\xi_a, \xi_b] = 0$. Then, by introducing the coordinates x^a as $\xi_a = \partial/\partial x^a$, the metric can be written as

$$ds^2 = \lambda_{ab}(dx^a + a^a_i dx^i)(dx^b + a^b_j dx^j) + \tau^{-1} h_{ij} dx^i dx^j, \quad (1)$$

where the functions, $\lambda_{ab} = (\xi_a | \xi_b)$, $\tau := -\det(\lambda_{ab})$, a^a_i , the three-dimensional metric h_{ij} ($i = 2, 3, 4$) are independent of the coordinates x^a . Let us define the twist one-forms by

$$V_a = *(\xi_1 \wedge \xi_2 \wedge d\xi_a), \quad (2)$$

whose exterior derivative can be written as

$$dV_a = 2 * (\xi_1 \wedge \xi_2 \wedge R(\xi_a)), \quad (3)$$

where $R(\xi_a)$ is the Ricci one-form. Therefore, by the vacuum Einstein equation, since the right-hand side in Eq. (3) vanishes, there locally exists the twist potentials ω_a such that

$$d\omega_a = V_a, \quad (4)$$

which can be written as

$$\partial_k \omega_a = \tau \sqrt{|h|} \lambda_{ab} \varepsilon_{ijk} h^{im} h^{jn} \partial_m a^b_n. \quad (5)$$

Then, the vacuum Einstein equations can be expressed as the field equations for the five scalar fields, $\{\lambda_{ab}, \omega_a\}$,

$$\Delta_h \lambda_{ab} = \lambda^{cd} h^{ij} \frac{\partial \lambda_{ac}}{\partial x^i} \frac{\partial \lambda_{bd}}{\partial x^j} + \tau^{-1} h^{ij} \frac{\partial \omega_a}{\partial x^i} \frac{\partial \omega_b}{\partial x^j}, \quad (6)$$

$$\Delta_h \omega_a = \tau^{-1} h^{ij} \frac{\partial \tau}{\partial x^i} \frac{\partial \omega_a}{\partial x^j} + \lambda^{bc} h^{ij} \frac{\partial \lambda_{ab}}{\partial x^i} \frac{\partial \omega_c}{\partial x^j}, \quad (7)$$

and the Einstein equations for the three-dimensional metric h_{ij} , which is coupled with the five scalar fields,

$$R_{ij}^h = \frac{1}{4} \lambda^{ab} \lambda^{cd} \frac{\partial \lambda_{ac}}{\partial x^i} \frac{\partial \lambda_{bd}}{\partial x^j} + \frac{1}{4} \tau^{-2} \frac{\partial \tau}{\partial x^i} \frac{\partial \tau}{\partial x^j} - \frac{1}{2} \tau^{-1} \lambda^{ab} \frac{\partial \omega_a}{\partial x^i} \frac{\partial \omega_b}{\partial x^j}, \quad (8)$$

where Δ_h is the Laplacian and R_{ij}^h denotes the Ricci tensor with respect to h_{ij} , respectively.

1. $SL(3, \mathbb{R})$ nonlinear sigma model

Thus, as a consequence of the existence of two isometries ξ_a , we have five scalar fields λ_{ab}, ω_a ($a = 0, 1$), which we denote collectively by coordinates $\Phi^A = (\lambda_{ab}, \omega_a)$. Then, we can find that the equations of motion, eqs. (6), (7) and (8) are derived from the following action for sigma-model Φ^A coupled with three-dimensional gravity with respect to the metric h_{ij} ,

$$S = \int \left(R^h - G_{AB} \frac{\partial \Phi^A}{\partial x^i} \frac{\partial \Phi^B}{\partial x^j} h^{ij} \right) \sqrt{|h|} d^3 x, \quad (9)$$

where R^h denotes the Ricci scalar with respect to h_{ij} , and the target space metric, G_{AB} , is given by

$$G_{AB} d\Phi^A d\Phi^B = \frac{1}{4} \text{Tr}(\lambda^{-1} d\lambda \lambda^{-1} d\lambda) + \frac{1}{4} \tau^{-2} d\tau^2 - \frac{1}{2} \tau^{-1} V^T \lambda^{-1} V, \quad (10)$$

with $\lambda = (\lambda_{ab})$, $\omega = (\omega_1, \omega_2)^T$ and $V = d\omega$. It can be actually shown that varying the action by h_{ij} derives the equation (8),

$$R_{ij}^h = G_{AB} \frac{\partial \Phi^A}{\partial x^i} \frac{\partial \Phi^B}{\partial x^j}, \quad (11)$$

and also varying the action by Φ^A can derive the equations (6) and (7),

$$\Delta_h \Phi^A + h^{ij} \Gamma_{BC}^A \frac{\partial \Phi^B}{\partial x^i} \frac{\partial \Phi^C}{\partial x^j} = 0, \quad (12)$$

where Γ_{BC}^A is the Christoffel symbol with respect to G_{AB} .

2. Coset matrix

Maison [26] showed that the action for the scalar fields Φ^A is invariant under the global $SL(3, \mathbb{R})$ transformation, introducing the $SL(3, \mathbb{R})$ matrix χ , defined by the 3×3 matrix

$$\chi = \begin{pmatrix} \lambda - \frac{\omega \omega^T}{\tau} & \frac{\omega}{\tau} \\ \frac{\omega^T}{\tau} & -\frac{1}{\tau} \end{pmatrix}, \quad (13)$$

where this matrix χ is symmetric, $\chi^T = \chi$, and unimodular, $\det(\chi) = 1$. Then, the action (24) can be written in terms of χ as follows

$$S = \int \left(R^h - \frac{1}{4} h^{ij} \text{tr}(\chi^{-1} \partial_i \chi \chi^{-1} \partial_j \chi) \right) \sqrt{|h|} d^3 x, \quad (14)$$

which is invariant under the transformation

$$\chi \rightarrow \chi' = g\chi g^T, \quad h \rightarrow h \quad (15)$$

with $g \in SL(3, \mathbb{R})$. The equations of motion (8), (6) and (7) can be written, in terms of χ , as

$$d \star_3 (\chi^{-1} d\chi) = 0, \quad (16)$$

$$R_{ij}^h = \frac{1}{4} \text{tr}(\chi^{-1} \partial_i \chi \chi^{-1} \partial_j \chi). \quad (17)$$

Thus, the presence of two commuting Killing vector fields reduces the five-dimensional vacuum Einstein theory to a three-dimensional non-linear sigma model with a $SL(3, \mathbb{R})$ target space symmetry. If both two Killing vectors are spacelike, it is described by the $SL(3, \mathbb{R})/SO(3)$ sigma model coupled to gravity, while if one of the two Killing vectors is timelike, the symmetry is replaced with $SL(3, \mathbb{R})/SO(2, 1)$ [26].

B. Einstein gravity with three commuting Killing vectors

It was seen from the previous subsection that assuming the existence of one timelike Killing vector $\xi_0 = \partial/\partial t$ and one spacelike axial Killing vector $\xi_1 = \partial/\partial \psi$, this theory reduces to the $SL(3, \mathbb{R})/SO(2, 1)$ non-linear sigma models coupled to three-dimensional gravity [26]. Furthermore, if we assume the presence of another spacelike Killing vector $\xi_2 = \partial/\partial x_2$ such that three Killing vectors mutually commute, $[\xi_I, \xi_J]$ ($I, J = 0, 1, 2$), it can be shown that the Einstein gravity can be completely reduced to the non-linear sigma model as follows: Under a further assumption of the presence of the third spacelike axial Killing vector $\xi_2 = \partial/\partial x^2$, the two-dimensional surface orthogonal to three ξ_I ($I, J = 0, 1, 2$) is integrable, as shown in Ref. [36, 37]. Under the symmetry assumptions, the metric can be written in the Weyl-Papapetrou form:

$$ds^2 = \lambda_{ab}(dx^a + a^a{}_2 dx^2)(dx^b + a^b{}_2 dx^2) + \tau^{-1} \rho^2 (dx^2)^2 + \tau^{-1} e^{2\sigma} (d\rho^2 + dz^2), \quad (18)$$

where the coordinates x^a ($a = 0, 1$) and x^2 denote the Killing coordinates and all functions λ_{ab} , $\tau := -\det(\lambda_{ab})$, $a^a{}_2$, σ are independent of x^a and x^2 . The coordinates (ρ, z) that span a two-dimensional base space, $\Sigma = \{(\rho, z) | \rho \geq 0, -\infty < z < \infty\}$, are globally well-defined, harmonic, and mutually conjugate on Σ .

In this coordinate system, the equation of motion (12) for Φ^A reduces to

$$\Delta_\gamma \Phi^A + \Gamma_{BC}^A [\Phi_{,\rho}^B \Phi_{,\rho}^C + \Phi_{,z}^C \Phi_{,z}^C] = 0, \quad (19)$$

where Δ_γ is the Laplacian with respect to the abstract three-dimensional metric $\gamma = d\rho^2 + dz^2 + \rho^2 d\varphi^2$. On the other hand, the function σ is determined by

$$\begin{aligned} \frac{2}{\rho} \sigma_{,\rho} &= R_{\rho\rho}^h - R_{zz}^h \\ &= G_{AB} [\Phi_{,\rho}^A \Phi_{,\rho}^B - \Phi_{,z}^A \Phi_{,z}^B] \\ &= \frac{1}{4} \tau^{-2} [(\tau_{,\rho})^2 - (\tau_{,z})^2] + \frac{1}{4} \lambda^{ab} \lambda^{cd} (\lambda_{ac,\rho} \lambda_{bd,\rho} - \lambda_{ac,z} \lambda_{bd,z}) - \frac{1}{2} \tau^{-1} \lambda^{ab} (\omega_{a,\rho} \omega_{b,\rho} - \omega_{a,z} \omega_{b,z}), \end{aligned} \quad (20)$$

$$\begin{aligned} \frac{1}{\rho} \sigma_{,z} &= R_{\rho z}^h \\ &= G_{AB} \Phi_{,\rho}^A \Phi_{,z}^B \\ &= \frac{1}{4} \tau^{-2} \tau_{,\rho} \tau_{,z} + \frac{1}{4} \lambda^{ab} \lambda^{cd} \lambda_{ac,\rho} \lambda_{bd,z} - \frac{1}{2} \tau^{-1} \lambda^{ab} \omega_{a,\rho} \omega_{b,z}, \end{aligned} \quad (21)$$

where the integrability of the function σ , $\sigma_{,\rho z} = \sigma_{,z\rho}$, is assured by Eq. (19). From Eq. (4) or (5), the metric functions $a^a{}_2$ ($a = 0, 1$) are determined by

$$\partial_\rho a^a{}_2 = -\frac{\rho}{\tau} \lambda^{ab} \partial_z \omega_b, \quad \partial_z a^a{}_2 = \frac{\rho}{\tau} \lambda^{ab} \partial_\rho \omega_b. \quad (22)$$

Thus, once one determines the five scalar fields $\Phi^A = (\lambda_{ab}, \omega_a)$ through Eq. (19), one can obtain the solutions to the five-dimensional vacuum Einstein equations with three Killing isometries. It turns out that Eq. (19) are derived from

the following action

$$\begin{aligned} S &= \int d\rho dz \rho [G_{AB}(\partial\Phi^A)(\partial\Phi^B)] \\ &= \int d\rho dz \rho \left[\frac{1}{4} \text{Tr}(\lambda^{-1} \partial\lambda \lambda^{-1} \partial\lambda) + \frac{1}{4} \tau^{-2} \partial\tau^2 - \frac{1}{2} \tau^{-1} v^T \lambda^{-1} v \right] \end{aligned} \quad (23)$$

$$= \frac{1}{4} \int d\rho dz \rho \text{tr} (\chi^{-1} \partial_i \chi \chi^{-1} \partial^i \chi) . \quad (24)$$

In terms of the coset matrix χ , Eqs. (19) and (20), (21) can be written, respectively, as

$$\partial_\rho(\rho \partial_\rho \chi \chi^{-1}) + \partial_z(\rho \partial_z \chi \chi^{-1}) = 0, \quad (25)$$

and

$$\frac{2}{\rho} \sigma_{,\rho} = -\frac{1}{\rho^2} + \frac{1}{4\rho^2} \text{tr}[(\rho \partial_\rho \chi \chi^{-1})^2 - (\rho \partial_z \chi \chi^{-1})^2], \quad (26)$$

$$\frac{1}{\rho} \sigma_{,z} = \frac{1}{4\rho^2} \text{tr}[(\rho \partial_\rho \chi \chi^{-1})(\rho \partial_z \chi \chi^{-1})]. \quad (27)$$

C. Ehlers transformation

When one constructs a new asymptotically flat vacuum solution from a certain vacuum seed by using the $SL(3, \mathbb{R})$ transformation, it is convenient to use a special $SL(3, \mathbb{R})$ transformation that preserves the asymptotic metric at infinity. This ensures that the five-dimensional Minkowski metric, which can be expressed in terms of the angular coordinates (ψ, ϕ) with the periodicity 2π ,

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2 \theta d\psi^2 + \cos^2 \theta d\phi^2), \quad (28)$$

is maintained, or, alternatively, in terms of the Euler angles $\phi_\pm := \psi \pm \phi$ and $\Theta := 2\theta$, it can be also written as

$$ds^2 = -dt^2 + dr^2 + \frac{r^2}{4} [(d\phi_+ + \cos \Theta d\phi_-)^2 + d\Theta^2 + \sin^2 \Theta d\phi_-^2], \quad (29)$$

Giusto and Saxena showed in Ref.[27] that the $SL(3, \mathbb{R})$ -transformation preserving asymptotic flatness is given by the one that leaves the limit $r \rightarrow \infty$ of the coset matrix χ corresponding to the Minkowski metric (29):

$$\lim_{r \rightarrow \infty} \chi = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{bmatrix}, \quad (30)$$

rather than the one with the orthogonal angles (28), invariant under the transformation (15)¹. This can be expressed by the subgroup $SO(2, 1)$ within the $SL(3, \mathbb{R})$. Moreover, among the subgroup with three parameters, only one parameter family results in a nontrivial transformation. More precisely, choosing the two Killing vectors $(\xi_0, \xi_1) = (\partial/\partial t, \partial/\partial \psi)$ or $(\xi_0, \xi_1) = (\partial/\partial t, \partial/\partial \phi)$ results in only trivial transformations but selecting other Killing vectors $(\xi_0, \xi_1) = (\partial/\partial t, \partial/\partial \phi_+)$ or $(\xi_0, \xi_1) = (\partial/\partial t, \partial/\partial \phi_-)$ yields a physical transformation that adds angular momentum to a static seed solution, which is expressed as

$$\chi \rightarrow \chi' = g_r^T \chi g_r, \quad g_r = \begin{bmatrix} 1 & 0 & \tilde{\beta} \\ -\tilde{\beta} & 1 & -\frac{\tilde{\beta}^2}{2} \\ 0 & 0 & 1 \end{bmatrix} \in SL(3, \mathbb{R}). \quad (31)$$

¹ It should be noted that χ at $r \rightarrow \infty$ in Ref.[27] has the opposite signature to the one we used, due to the difference in the signature of the twist potentials (4).

Under this transformation, the set of the scalar fields $\Phi^A = \{\lambda_{ab}, \omega_a\}$ is changed to the set of different ones $\Phi'^A = \{\lambda'_{ab}, \omega'_a\}$ as follows

$$\begin{aligned}
\lambda'_{00} &= D^{-1} \left[\lambda_{00} - 2\tilde{\beta}\lambda_{01} + \tilde{\beta}^2 (\lambda_{11} + \epsilon^{ij}\omega_i\lambda_{j0}) - \tilde{\beta}^3 \epsilon^{ij}\omega_i\lambda_{j1} + \frac{\tilde{\beta}^4 (\tau (-\omega_0^2 + \tau\lambda_{00}) + (\epsilon^{ij}\omega_i\lambda_{j0})^2)}{4\lambda_{00}} \right], \\
\lambda'_{01} &= D^{-1} \left[\lambda_{01} - \tilde{\beta} (\lambda_{11} + \epsilon^{ij}\omega_i\lambda_{j0}) + \frac{3}{2}\tilde{\beta}^2 \epsilon^{ij}\omega_i\lambda_{j1} - \frac{\tilde{\beta}^3 (\tau (-\omega_0^2 + \tau\lambda_{00}) + (\epsilon^{ij}\omega_i\lambda_{j0})^2)}{2\lambda_{00}} \right], \\
\lambda'_{11} &= D^{-1} \left[\lambda_{11} - 2\tilde{\beta}\epsilon^{ij}\omega_i\lambda_{j1} + \frac{\tilde{\beta}^2 (\tau (-\omega_0^2 + \tau\lambda_{00}) + (\epsilon^{ij}\omega_i\lambda_{j0})^2)}{\lambda_{00}} \right], \\
\omega'_0 &= D^{-1} \left[\omega_0 + \tilde{\beta} (-\omega_1 - \omega_0^2 + \tau\lambda_{00}) + \frac{3}{2}\tilde{\beta}^2 (\omega_0\omega_1 - \tau\lambda_{01}) + \frac{1}{2}\tilde{\beta}^3 (-\omega_1^2 + \tau\lambda_{11}) \right], \\
\omega'_1 &= D^{-1} \left[\omega_1 + \tilde{\beta} (-\omega_0\omega_1 + \tau\lambda_{01}) + \frac{1}{2}\tilde{\beta}^2 (\omega_1^2 - \tau\lambda_{11}) \right], \\
\tau' &= \tau/D,
\end{aligned} \tag{32}$$

with

$$D = 1 - 2\tilde{\beta}\omega_0 + \tilde{\beta}^2 (\omega_1 + \omega_0^2 - \tau\lambda_{00}) - \tilde{\beta}^3 (\omega_0\omega_1 - \tau\lambda_{01}) + \frac{1}{4}\tilde{\beta}^4 (\omega_1^2 - \tau\lambda_{11}), \tag{33}$$

where the new parameter $\tilde{\beta}$ is physically related to the angular momentum with respect to $\partial/\partial\phi_+$ or $\partial/\partial\phi_-$, depending on the choice of the Killing vector $\xi_1 = \partial/\partial\phi_+$ or $\partial/\partial\phi_-$. The one-forms $a'^a_i dx^i$ ($a = 0, 1$) for the transformed solution are determined by the five scalar functions λ'_{ab}, ω'_a from Eqs. (5), after replacing λ_{ab}, ω_a with λ'_{ab}, ω'_a and substituting Eq. (32). Thus, a'^a_i can be determined from a^a_i and $\{\lambda_{ab}, \omega_a\}$ by

$$\begin{aligned}
\partial_m(a'^0_n - a^0_n) - \partial_n(a'^0_m - a^0_m) &= \sqrt{h}\epsilon_{mnl}h^{\ell k} \left[\tilde{\beta}(2(\tilde{\mathbf{X}}^0_0)_k + (\tilde{\mathbf{X}}^1_1)_k) + \frac{\tilde{\beta}^2}{2}(2(\tilde{\mathbf{Y}}^1)_k - (\tilde{\mathbf{X}}^0_1)_k) + \frac{\tilde{\beta}^3}{2}(\tilde{\mathbf{Y}}^0)_k \right], \\
\partial_m(a'^1_n - a^1_n) - \partial_n(a'^1_m - a^1_m) &= \sqrt{h}\epsilon_{mnl}h^{\ell k} \left[\tilde{\beta}(\tilde{\mathbf{X}}^1_0)_k + \frac{3}{2}\tilde{\beta}^2(\tilde{\mathbf{X}}^0_0)_k + \frac{\tilde{\beta}^3}{2}((\tilde{\mathbf{Y}}^1)_k - (\tilde{\mathbf{X}}^0_1)_k) + \frac{\tilde{\beta}^4}{4}(\tilde{\mathbf{Y}}^0)_k \right],
\end{aligned} \tag{34}$$

where the one-forms $(\tilde{\mathbf{X}}^a_b)_k$ and $(\tilde{\mathbf{Y}}^a)_k$ can be expressed in terms of only the scalar fields Φ^A as :

$$\begin{aligned}
\tilde{\mathbf{X}}^a_b &= \lambda^{ac}d\lambda_{cb} - \tau^{-1}\lambda^{ac}d\omega_c\omega_b + \tau^{-1}\delta_1^a\delta_b^0\lambda^{0c}d\omega_c, \\
\tilde{\mathbf{Y}}^a &= -\epsilon^{ab}d\omega_b - \tau^{-1}\epsilon^{ab}\omega_b\omega_c\lambda^{cd}d\omega_d + \epsilon^{ab}d\lambda_{bc}\lambda^{cd}\omega_d + \tau^{-1}\epsilon^{ab}\omega_b d\tau.
\end{aligned} \tag{35}$$

The assumption of a further Killing vector $\xi_2 = \partial/\partial\phi_-$ or $\xi_2 = \partial/\partial\phi_+$, depending on the choice of $\xi_1 = \partial/\partial\phi_+$ or $\xi_1 = \partial/\partial\phi_-$, simplifies Eqs. (34) to

$$\partial_\rho(a'^0_2 - a^0_2) = -\rho \left[\tilde{\beta}(2(\tilde{\mathbf{X}}^0_0)_z + (\tilde{\mathbf{X}}^1_1)_z) + \frac{\tilde{\beta}^2}{2}(2(\tilde{\mathbf{Y}}^1)_z - (\tilde{\mathbf{X}}^0_1)_z) + \frac{\tilde{\beta}^3}{2}(\tilde{\mathbf{Y}}^0)_z \right], \tag{36}$$

$$\partial_z(a'^0_2 - a^0_2) = \rho \left[\tilde{\beta}(2(\tilde{\mathbf{X}}^0_0)_\rho + (\tilde{\mathbf{X}}^1_1)_\rho) + \frac{\tilde{\beta}^2}{2}(2(\tilde{\mathbf{Y}}^1)_\rho - (\tilde{\mathbf{X}}^0_1)_\rho) + \frac{\tilde{\beta}^3}{2}(\tilde{\mathbf{Y}}^0)_\rho \right], \tag{37}$$

$$\partial_\rho(a'^1_2 - a^1_2) = -\rho \left[\tilde{\beta}(\tilde{\mathbf{X}}^1_0)_z + \frac{3}{2}\tilde{\beta}^2(\tilde{\mathbf{X}}^0_0)_z + \frac{\tilde{\beta}^3}{2}((\tilde{\mathbf{Y}}^1)_z - (\tilde{\mathbf{X}}^0_1)_z) + \frac{\tilde{\beta}^4}{4}(\tilde{\mathbf{Y}}^0)_z \right], \tag{38}$$

$$\partial_z(a'^1_2 - a^1_2) = \rho \left[\tilde{\beta}(\tilde{\mathbf{X}}^1_0)_\rho + \frac{3}{2}\tilde{\beta}^2(\tilde{\mathbf{X}}^0_0)_\rho + \frac{\tilde{\beta}^3}{2}((\tilde{\mathbf{Y}}^1)_\rho - (\tilde{\mathbf{X}}^0_1)_\rho) + \frac{\tilde{\beta}^4}{4}(\tilde{\mathbf{Y}}^0)_\rho \right], \tag{39}$$

The integration can be written as

$$\begin{aligned}
a'^0_2 &= a^0_2 + \tilde{\beta}(2\mathbf{X}^0_0 + \mathbf{X}^1_1) + \frac{\tilde{\beta}^2}{2}(2\mathbf{Y}^1 - \mathbf{X}^0_1) + \frac{\tilde{\beta}^3}{2}\mathbf{Y}^0, \\
a'^1_2 &= a^1_2 + \tilde{\beta}\mathbf{X}^1_0 + \frac{3}{2}\tilde{\beta}^2\mathbf{X}^0_0 + \frac{\tilde{\beta}^3}{2}(\mathbf{Y}^1 - \mathbf{X}^0_1) + \frac{\tilde{\beta}^4}{4}\mathbf{Y}^0,
\end{aligned} \tag{40}$$

where \mathbf{X}^a_b and \mathbf{Y}^a are the solutions of

$$\begin{aligned}\partial_\rho \mathbf{X}^a_b &= -\rho (\tilde{\mathbf{X}}^a_b)_z, & \partial_z \mathbf{X}^a_b &= \rho (\tilde{\mathbf{X}}^a_b)_\rho, \\ \partial_\rho \mathbf{Y}^a &= -\rho (\tilde{\mathbf{Y}}^a)_z, & \partial_z \mathbf{Y}^a &= \rho (\tilde{\mathbf{Y}}^a)_\rho.\end{aligned}\quad (41)$$

Thus, one can obtain the new metric describing the rotating solution to the vacuum Einstein equations with the same Killing isometries. As shown previously in Ref. [30], applying this transformation to the five-dimensional static black hole solution (the five-dimensional Schwarzschild-Tangherlini solution [38]) generates the five-dimensional rotating black hole solution (the five-dimensional Myers-Perry solution [7]). As will be seen in the following section, however, when applied to vacuum black rings, such as the static black ring and the singly rotating black ring, a Dirac-Misner string singularity inevitably appears on the disk inside the ring, similar to the Harrison transformation studied in Ref. [30]. In the following section, to solve this undesirable problem, we will choose the vacuum rotating black ring having a Dirac-Misner string singularity on the disk inside the ring as the seed of the transformation, as performed in our previous work [32] for the Harrison transformation.

III. SEED SOLUTION FOR THE EHLERS TRANSFORMATION

In our previous works [33, 34], as a vacuum seed for the Harrison transformation, we chose the vacuum rotating black lens with a horizon cross-section of lens space $L(n; 1)$ ($n = 0, 1, 2, \dots$) [39], which initially has a Dirac-Misner string singularity. It should be noted that in Ref. [39], they removed this singularity, but we did not, since we eliminated it by appropriately adjusting the parameters of the solution after the Harrison transformation. Now, we follow the same procedure as in Ref. [33, 34]. Furthermore, we use the same seed solution as the seed used when constructing the capped black hole or charged dipole black ring [33, 34].

A. Black lens with a Dirac-Misner string singularity as a seed

The metric for the vacuum seed solution (see Refs. [33, 34] for details on its construction) that we use for the Ehlers transformation is written in the C-metric form as follows:

$$\begin{aligned}ds^2 &= -\frac{\bar{H}(y, x)}{\bar{H}(x, y)}(d\bar{t} + \bar{\Omega}_\psi(x, y)d\bar{\psi} + \bar{\Omega}_\phi(x, y)d\bar{\phi})^2 + \frac{\bar{F}(y, x)}{\bar{H}(y, x)}d\bar{\psi}^2 - \frac{2\bar{J}(x, y)}{\bar{H}(y, x)}d\bar{\psi}d\bar{\phi} - \frac{\bar{F}(x, y)}{\bar{H}(y, x)}d\bar{\phi}^2 \\ &+ \frac{\ell^2 \bar{H}(x, y)}{4(1-\gamma)^3(1-\nu)^2(1-a^2)(x-y)^2} \left(\frac{dx^2}{G(x)} - \frac{dy^2}{G(y)} \right),\end{aligned}\quad (42)$$

where

$$G(u) = (1-u^2)(1+\nu u), \quad (43)$$

$$\begin{aligned}\bar{H}(x, y) &= \left[\nu b^2 c_1^2 (1+\nu)^2 (\gamma-\nu)(1-\gamma)(1-x^2) + \nu(1-\gamma)(\gamma-\nu) (b(1-\nu)^2(1-x) - 2c_1(1+\nu x))^2 \right. \\ &+ b^2 c_1^2 (1+\nu)^3 (x+1)(\gamma-\nu)^2 \left. \right] (1+y)^2 - [d_5(1-x)^2 + d_6(1-x^2) + 2(1-\nu)^{-1}d_6(1+x)(1+\nu x)](1+y) \\ &+ 4(1-a^2)(1-\gamma)^3(1-\nu)^4(1-x) + 2c_2^2(1-\gamma)(1-\nu)^2(1-x^2) - 4d_1(1-\gamma)(1-\nu)^2(1+\nu)(1+x),\end{aligned}\quad (44)$$

$$\begin{aligned}\bar{F}(x, y) &= \frac{2\ell^2}{(1-a^2)(x-y)^2} \left[4 \left[(1-a^2)^2 (y-1)(1-\gamma)^3(1-\nu)^3 - d_1^2(1+y) \right] (1+y\nu)G(x) \right. \\ &+ 4[(1-\nu)c_2 - (1-ab)(\gamma-\nu)(1+\nu)c_1]^2 (1+x\nu)(1+x)G(y) \\ &+ \nu^{-1}(1-\nu)^3(\gamma-\nu)(d_3^2(1-x^2)G(y) - c_3^2(1-y^2)G(x)) \\ &+ \left. \frac{G(x)G(y) \left[(1-a^2)(1-\gamma)d_4 - (a-b)^2y(1-\gamma)^2(\gamma-\nu)\nu c_2^2 + x(\gamma-\nu)\nu (bd_1 - c_1c_3)^2 \right]}{\nu(1-\gamma)} \right],\end{aligned}\quad (45)$$

$$\begin{aligned}\bar{J}(x, y) &= \frac{2\ell^2(1+x)(1+y)}{(1-a^2)(x-y)} \left[4d_1((a-b)(1-\gamma)(\gamma-\nu)(1+\nu) - ad_2)(1+\nu x)(1+\nu y) \right. \\ &+ \left. d_3c_3(1-\nu)^3(\gamma-\nu)(1-x)(1-y) - (a-b)(\gamma-\nu)c_2(c_1c_3 - bd_1)(1-x)(1-y)(1+x\nu)(1+y\nu) \right],\end{aligned}\quad (46)$$

$$\bar{\Omega}_\psi(x, y) = \frac{v_0 \ell (1+y)(1-\nu)}{\nu \bar{H}(y, x)} \left[c_2 (c_1 c_3 - b d_1) (1-x)(1+x\nu)(1+y\nu) \right. \\ \left. - (1-\nu)^2 d_2 c_3 (1-x) + (1+x\nu) d_1 (2\nu(1-ab)(1-\gamma)(1+\nu)(1+x) + (1-3\nu-x(1+\nu)) c_3) \right], \quad (47)$$

$$\bar{\Omega}_\phi(x, y) = \frac{v_0 \ell (1+x)}{\bar{H}(y, x)} \left[b(1+x) d_1 (d_2(1+y)(1+y\nu) + \nu c_3 (1-y^2) (1-\nu)) \right. \\ \left. + \frac{2(a-b)(1-\gamma)^2 (2d_1(1+x\nu)(1+y\nu)^2 - (1-\nu)^2 \nu c_3 (1-y)(x+y+\nu+xy\nu))}{1+\nu} \right], \quad (48)$$

and the coefficients are given by

$$\begin{aligned} v_0 &:= \sqrt{\frac{2(\gamma^2 - \nu^2)}{(1-a^2)(1-\gamma)}}, \\ c_1 &:= (1-\gamma)a + (\gamma-\nu)b, \\ c_2 &:= 2a(1-\gamma)\nu + b(\gamma-\nu)(1+\nu), \\ c_3 &:= 2(1-\gamma)\nu + b^2(\gamma-\nu)(1+\nu), \\ d_1 &:= (\nu+1)c_1^2 - (1-\gamma)(1-\nu)^2, \\ d_2 &:= b(\nu+1)c_1(\gamma-\nu) + 2\nu(1-\gamma)(1-\nu), \\ d_3 &:= (a^2-1)b(\gamma-1)(\nu+1) - ac_3, \\ d_4 &:= b^2(\gamma-\nu) [(\nu+1)^2 c_1^2 (-3(1-\gamma)\nu - \nu^2 + 1) - (1-\gamma)(1-\nu)^4 (2\nu+1)] \\ &\quad + (1-\gamma) \left[((1-\nu)c_2 - 2\nu^2 c_1)^2 - 4\nu^2 c_1^2 (-\gamma(\nu+2) + 3\nu^2 + 1) \right], \\ d_5 &:= (1-\gamma)(1-\nu)^3 [(\gamma-3\nu)(b^2(\nu-1)(\gamma-\nu) - c_1^2) - 2bc_1(3\nu-1)(\gamma-\nu)], \\ d_6 &:= c_1(1-\gamma)(1-\nu)(1-\nu^2)(c_2 - (1-\gamma)(a-b)(\gamma-\nu)). \end{aligned} \quad (49)$$

We assume the ranges of the coordinates as

$$-\infty < \bar{t} < \infty, \quad -1 \leq x \leq 1, \quad -1/\nu \leq y \leq -1, \quad (50)$$

where, unlike the Harrison transformation studied in our previous works [32–34], we do not impose the periodicities at this stage:

$$0 \leq \bar{\psi} \leq 2\pi, \quad 0 \leq \bar{\phi} \leq 2\pi, \quad (51)$$

which correspond to the conditions for the absence of conical singularities on $y = -1$ and $x = -1$, respectively. After performing the Ehlers transformation on this seed solution, we will impose the periodicities on the angular coordinates ψ and ϕ . This solution has five parameters $(\ell, \nu, \gamma, a, b)$, which have the ranges

$$\ell > 0, \quad 0 < \nu < \gamma < 1, \quad -1 < a < 1. \quad (52)$$

The boundaries of the C-metric coordinate (x, y) for the seed solution can be described as follows:

(i) $\bar{\phi}$ -rotational axis: $\partial\Sigma_{\bar{\phi}} = \{(x, y) | x = -1, -1 < y < -1/\nu\}$ with the rod vector $v_{\bar{\phi}} := (0, 0, 1) = \partial/\partial\bar{\phi}$, where it should be noted that in the assumption of the periodicity $\bar{\phi} \sim \bar{\phi} + 2\pi$ from the coordinate ranges (51) ensures the absence of the conical singularities, but we do not impose this.

(ii) Horizon: $\partial\Sigma_{\mathcal{H}} = \{(x, y) | -1 < x < 1, y = -1/\nu\}$ with the rod vector $v_{\mathcal{H}} := (1, \bar{\omega}_\psi, \bar{\omega}_\phi)$, where

$$\begin{aligned} \bar{\omega}_\psi &= \frac{v_0 (1-a^2) (1-\gamma)}{2\ell(\gamma+\nu) (1-a^2 + a(a-b)\gamma - (1-ab)\nu)}, \\ \bar{\omega}_\phi &= \bar{\omega}_\psi \frac{2a(1-\gamma)\nu - (1-a^2)b(1-\gamma)(1+\nu) + ab^2(\gamma-\nu)(1+\nu)}{2(1-\gamma)\nu + b^2(\gamma-\nu)(1+\nu)}. \end{aligned} \quad (53)$$

(iii) Inner axis: $\partial\Sigma_{\text{in}} = \{(x, y) | x = 1, -1 < y < -1/\nu\}$ with the rod vector

$$v_{\text{in}} := (v_{\text{in}}^{\bar{t}}, v_{\text{in}}^{\bar{\psi}}, 1), \quad (54)$$

with

$$v_{\text{in}}^{\bar{t}} := v_0 \ell (a - b), \quad v_{\text{in}}^{\bar{\psi}} := \frac{ad_1 + (1 - \gamma)(1 + \nu)(1 - a^2)c_1}{d_1}, \quad (55)$$

where we note that the presence of the nonzero t -component $v_{\text{in}}^{\bar{t}}$ denotes the existence of a the Dirac-Misner string singularity.

(iv) $\bar{\psi}$ -rotational axis: $\partial\Sigma_{\bar{\psi}} = \{(x, y) | -1 < x < 1, y = -1\}$ with the rod vector $v_{\bar{\psi}} := (0, 1, 0) = \partial/\partial\bar{\psi}$, where the periodicity $\bar{\psi} \sim \bar{\psi} + 2\pi$ from the coordinate ranges (51) also ensures the absence of the conical singularities, but as mentioned above, we do not impose this.

(v) Infinity: $\partial\Sigma_{\infty} = \{(x, y) | x \rightarrow y \rightarrow -1\}$. In the limit $x \rightarrow y \rightarrow -1$ ($r \rightarrow \infty$), the metric (locally) asymptotes to the five-dimensional Minkowski metric

$$ds^2 \simeq -d\bar{t}^2 + dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\bar{\psi}^2 + r^2 \cos^2 \theta d\bar{\phi}^2, \quad (56)$$

where the coordinates (r, θ) are defined by

$$x = -1 + \frac{4\ell(1 - \nu) \cos^2 \theta}{r^2}, \quad y = -1 - \frac{4\ell(1 - \nu) \sin^2 \theta}{r^2}. \quad (57)$$

As previously discussed in Ref. [34], by setting $a = b$ where the metric precisely describes the rotating black lens in Ref. [39], the Dirac-Misner string singularity [40] on the inner axis $\partial\Sigma_{\text{in}}$ can be removed. However, similar to the approach taken for the charged dipole black ring in Ref. [32] and the capped black hole [34], we choose to use the vacuum black lens with a Dirac-Misner string singularity as the seed for the Ehlers transformation. Therefore, we do not assume its absence before the transformation. In the following section, we will eliminate this singularity after the transformation by appropriately adjusting the parameters of the solution.

B. The seed solution in the Euler angles

The Ehlers transformation (31) requires the choice of either $(\xi_0, \xi_1) = (\partial/\partial t, \partial/\partial\phi_+)$ or $(\xi_0, \xi_1) = (\partial/\partial t, \partial/\partial\phi_-)$, where ϕ_{\pm} are the Euler angles introduced in Sec. II C. In the choice of the Killing vectors $(\xi_0, \xi_1) = (\partial/\partial t, \partial/\partial\phi_+)$ and $\xi_2 = \partial/\partial\phi_-$ ($(x^0, x^1, x^2) = (t, \phi_+, \phi_-)$), the scalar fields $\Phi^A = \{\lambda_{ab}, \omega_a\}$ and the functions τ, a^a_2 can be expressed as

$$\begin{aligned} \lambda_{00} &= -\frac{\bar{H}(y, x)}{\bar{H}(x, y)}, \quad \lambda_{01} = -\frac{\bar{H}(y, x)}{2\bar{H}(x, y)} (\bar{\Omega}_{\psi}(x, y) \pm \bar{\Omega}_{\phi}(x, y)), \\ \lambda_{11} &= -\frac{\bar{F}(x, y) - \bar{F}(y, x) \pm 2\bar{J}(x, y)}{4\bar{H}(y, x)} - \frac{\bar{H}(y, x)(\bar{\Omega}_{\phi}(x, y) \pm \bar{\Omega}_{\psi}(x, y))^2}{4\bar{H}(x, y)}, \\ a_2^0 &= \frac{\bar{F}(x, y)\bar{\Omega}_{\psi}(x, y) \pm \bar{F}(y, x)\bar{\Omega}_{\phi}(x, y) \pm \bar{J}(x, y)(\bar{\Omega}_{\psi}(x, y) \mp \bar{\Omega}_{\phi}(x, y))}{\bar{F}(x, y) - \bar{F}(y, x) \pm 2\bar{J}(x, y)}, \\ a_2^1 &= -\frac{\bar{F}(x, y) + \bar{F}(y, x)}{\bar{F}(x, y) - \bar{F}(y, x) \pm 2\bar{J}(x, y)}, \\ \tau &= -\frac{\bar{F}(x, y) - \bar{F}(y, x) \pm 2\bar{J}(x, y)}{4\bar{H}(x, y)}, \end{aligned} \quad (58)$$

and one can find the twist potential is given by²

$$\begin{aligned} \omega_0 &= \frac{1}{2} (\bar{\Omega}_{\psi}(y, x) \pm \bar{\Omega}_{\phi}(y, x)), \\ \omega_1 &= -\tau + \frac{2\bar{K}_{\pm}(x, y)}{\bar{H}(x, y)}, \end{aligned} \quad (59)$$

² The Jacobian for the transformation $(\psi, \phi) \rightarrow (\phi_+, \phi_-)$ becomes 1/2. Hence the duality conditions for Eqs. (59) and (61) must be given in the rescaled canonical coordinate $(\bar{\rho}, \bar{z}) = (\rho/2, z/2)$.

where $\bar{K}_\pm(x, y)$ is a polynomial of x and y given by

$$\begin{aligned} \bar{K}_\pm(x, y) = & \frac{\ell^2}{1 \pm a} \left[-2k_1^\pm(x+1)(y^2-1) + k_2^\pm(x+y-2)(\nu(-xy+x+y+1)+2) + k_3^\pm(x+1)(y+1) \right. \\ & + k_4^\pm(\nu^2(x^2(y-1)+x(y^2+7y+2)-y^2+2y+1) + \nu(-(x^2(y-1))-x(y^2-5)+y^2+5y+6) + (x+1)(y+1)) \\ & + k_5^\pm(\nu^2(x^2(y-1)+x(y^2+3)-y^2+3y+2) + \nu(x^2(-y)+x^2-xy^2+8xy+x+y^2+y+6) + 4(x+y)) \\ & \left. + k_6^\pm(x+1)(y+1)(x-y) + k_7^\pm(x-1)(y-1)(x-y) + k_8^\pm(\nu x+1)(x-y)(\nu y+1) + k_9^\pm \bar{H}(x, y) \right], \end{aligned} \quad (60)$$

where $\{k_i^\pm\}_{i=1,\dots,9}$ are given in the Appendix and k_9^+ is the integration constant determined so that the coset matrix has the desirable behavior at the infinity (30).

IV. TRANSFORMED SOLUTION

In what follows, for a simpler expression, we rescale the transformation parameter as $\beta := \ell v_0 \tilde{\beta}/4$. The new scalar fields $\Phi'^A = \{\lambda'_{ab}, \omega'_a\}$ and the new functions τ' , a'^a after the transformation (31) can be obtained through Eqs. (32) and (40). The functions a'^a ($a = 0, 1$) can be expressed into the following lengthy form

$$\begin{aligned} a'^a{}_2 = & \mathcal{A}_0^a + \frac{1}{(x-y)^2(\bar{F}(x, y) - \bar{F}(y, x) + 2\bar{J}(x, y))} \left[\mathcal{A}_1^a(\nu y(x^2 - y^2) + \nu^2 x(x^2 - 1)y - y^2 + 1) \right. \\ & + \mathcal{A}_2^a(-\nu x^3 y^3 - x^2 y^3 - xy^2 + \nu xy + x + y^3) + \mathcal{A}_3^a(x-y)(\nu x^2 y^2 + \nu xy + x + y) - \mathcal{A}_4^a x^3(y^2 - 1)(\nu y + 1) \\ & \left. - \frac{\mathcal{A}_5^a(x^2 - 1)y^3(\nu x + 1)}{\nu} + \mathcal{A}_6^a(x^2 - 1)y^2(\nu x + 1) - \mathcal{A}_7^a(x^2 - 1)y(\nu x + 1) + \frac{\mathcal{A}_8^a(x^2 - 1)y^4(\nu x + 1)}{\nu} \right], \end{aligned} \quad (61)$$

where due to the lengthy expressions, we do not show the coefficients \mathcal{A}_k^a ($k = 0, \dots, 8$) explicitly. Note that the integration constants \mathcal{A}_0^a are set so that $a'^a{}_2$ asymptotes to $0 + C \cos(2\theta)$ with a constant C as $r \rightarrow \infty$ in Eq. (77). The transformed metric is not at the rest frame but has a global rotation at infinity, which can be removed by redefining the coordinates $(\bar{t}, \phi_+, \phi_-) \rightarrow (t, \psi, \phi)$.

$$\bar{t} = t + \mathcal{C}(\psi + \phi), \quad \phi_+ = \sqrt{D_0}(\psi + \phi), \quad \phi_- = \frac{1}{\sqrt{D_0}}(\psi - \phi), \quad (62)$$

where $D_0 := D|_{(x,y)=(-1,-1)}$ is given by

$$D_0 := 1 + D_{0,2}\beta^2 + D_{0,3}\beta^3 + D_{0,4}\beta^4, \quad (63)$$

and

$$\mathcal{C} := \frac{\ell v_0}{8\sqrt{D_0}}(2D_{0,2}\beta + 3D_{0,3}\beta^2 + 4D_{0,4}\beta^3), \quad (64)$$

with the coefficients

$$\begin{aligned}
D_{0,2} &= -\frac{6\left((1-a^2)(1-\nu) + (b-a)^2(\gamma-\nu)\right)}{\gamma-\nu}, \\
D_{0,3} &= \frac{1}{\gamma^2-\nu^2} \left[-8(1-a)^2(1+a)(1+b) - 8(a-b)^3\gamma^2 + 8(1-a)(1-\gamma)(-3a + (3-a+a^2)b + (1-a)b^2)\nu \right. \\
&\quad \left. + 8(1-3ab + (1+a+a^2)b^2 - b^3)\nu^2 - 8(1-a)\gamma(a(1+2a) - (1+3a-a^2)b + (1-a)b^2) \right], \\
D_{0,4} &= \frac{1}{(\gamma^2-\nu^2)^2} \left[-3(a-b)^4\gamma^4 + 4(1-a)^3(1+a)(3+2b)\nu - (1-a)^2\{3-26a+7a^2+8(2-3a+a^2)b+16b^2\}\nu^2 \right. \\
&\quad - 2(1-a)\{3-9a+2(6-5a+5a^2)b - (1+5a)b^2 - 4(1-a)b^3\}\nu^3 \\
&\quad + (1-b)\{-3-3(1-4a)b - (5-4a+8a^2)b^2 + 3b^3\}\nu^4 \\
&\quad + \gamma^3\{-2(1-a)(a-b)^2(1+5a+4(-1+a)b) + 2(-1+a)(-a+b)^2(3(-3+a) + 4(-1+a)b)\nu\} \\
&\quad + \gamma^2\{(1-a)^2(1-6a-11a^2+8(1+a-2a^2)b+16ab^2) \\
&\quad + 2(1-a)(3(1+a-8a^2+4a^3) + 2(-4+21a-17a^2+6a^3)b + (-17+27a-16a^2)b^2 - 4(1-a)b^3)\nu \\
&\quad + \nu^2(-7+40a-60a^2+40a^3-7a^4-4(6-17a+24a^2-9a^3+2a^4)b \\
&\quad + 2(-7+28a-11a^2+8a^3)b^2 - 8(1+a+a^2)b^3 + 6b^4)\} \\
&\quad + \gamma\{-4(1-a)^3(1+a)(1+2b) - 2(1-a)^3(9+15a+4(-1+3a)b-8b^2)\nu \\
&\quad + 2(1-a)(8-28a+33a^2-7a^3 + (20-54a+30a^2-8a^3)b + (17-19a+8a^2)b^2 - 4(1-a)b^3)\nu^2 \\
&\quad \left. + 2(1-a)(3-9a+2(6-5a+5a^2)b - (1+5a)b^2 - 4(1-a)b^3)\nu^3\right].
\end{aligned}$$

In the new coordinates (62), one can write the transformed metric again in the C-metric form

$$\begin{aligned}
ds^2 &= -\frac{H(y,x)}{H(x,y)}(dt + \Omega_\psi(x,y)d\psi + \Omega_\phi(x,y)d\phi)^2 + \frac{F(y,x)}{H(y,x)}d\psi^2 - \frac{2J(x,y)}{H(y,x)}d\psi d\phi - \frac{F(x,y)}{H(y,x)}d\phi^2 \\
&\quad + \frac{\ell^2 H(x,y)}{4(1-\gamma)^3 D_0(1-\nu)^2(1-a^2)(x-y)^2} \left(\frac{dx^2}{G(x)} - \frac{dy^2}{G(y)} \right), \tag{65}
\end{aligned}$$

where each functions are given by

$$\begin{aligned}
H(x, y) = & (1+y)^2 \left\{ \nu d_1 f_7^2 (1-\nu) (1-x^2) (2(\gamma-1)\nu + c_3) + 2c_3 c_1^2 f_6^2 (\gamma-\nu)(\nu x+1)^2 \right. \\
& - (\gamma-\nu)(1-\nu) (c_1^2 c_3 f_6^2 - 2(1-\gamma)\nu (c_1 - b(1-\nu))^2 f_8^2) (1-x)(1+x\nu) \\
& \left. - d_1 (\gamma-\nu) (2(\gamma-1)\nu + c_3) - 4(\gamma-1)d_2 \nu (\nu-1) \right\} \\
& - (1+y) \left\{ d_5 g_6 (1-x)^2 + d_6 g_7 (1-x^2) + 2(1-\nu)^{-1} d_6 g_8 (1+x)(1+x\nu) \right\} \\
& + 4(1-a^2)(1-x)(1-\gamma)^3 (1-\nu)^4 D_0 - 4(1+x)(1-\gamma)(1-\nu)^2 (1+\nu) d_1 f_2^2 + 2(1-x^2)(1-\gamma)(1-\nu)^2 c_2^2 f_5^2,
\end{aligned} \tag{66}$$

$$\begin{aligned}
F(x, y) = & \frac{2\ell^2}{D_0(1-a^2)(x-y)^2} \left[4(1+y\nu)G(x) \left\{ (1-a^2)^2 (-1+y)(1-\gamma)^3 (1-\nu)^3 D_0^2 - (1+y)d_1^2 f_1^2 f_2^2 \right\} \right. \\
& + 4(1+x)(1+x\nu)G(y) \left((1-ab)(\gamma-\nu)(1+\nu)c_1 - (1-\nu)c_2 \right)^2 g_{13}^2 \\
& + \nu^{-1}(1-\nu)^3 (\gamma-\nu) \left\{ (y^2-1)G(x)c_3^2 f_3^2 f_4^2 + (1-x^2)G(y)d_3^2 g_2^2 \right\} \\
& \left. + G(x)G(y) \left\{ \frac{x(\gamma-\nu)g_5^2(c_1 c_3 - b d_1)^2}{1-\gamma} - (a-b)^2 y(1-\gamma)(\gamma-\nu)c_2^2 f_5^2 f_8^2 + \frac{(1-a^2)d_4 D_0 g_4}{\nu} \right\} \right],
\end{aligned} \tag{67}$$

$$\begin{aligned}
J(x, y) = & \frac{2\ell^2(1+x)(1+y)}{D_0(1-a^2)(x-y)} \left[4d_1 f_1 f_2 g_{13} \left\{ (1-ab)(\gamma-\nu)(1+\nu)c_1 - (1-\nu)c_2 \right\} (1+x\nu)(1+y\nu) \right. \\
& \left. - (1-x)(1-y)(\gamma-\nu) \left\{ (a-b)(1+x\nu)(1+y\nu)c_2 f_5 f_8 g_5 (c_1 c_3 - b d_1) + (1-\nu)^3 c_3 d_3 f_3 f_4 g_2 \right\} \right],
\end{aligned} \tag{68}$$

$$\begin{aligned}
\Omega_\psi(x, y) = & \frac{\ell v_0(1+y)}{\sqrt{D_0}H(y, x)} \left[\nu^{-1} c_2 (c_1 c_3 - b d_1) f_5 g_9 (1-x)(1-\nu)(1+x\nu)(1+y\nu) \right. \\
& + (1+y\nu) \left\{ (1+x)(1+x\nu)c_1 c_3 f_2 f_9 f_{10} - \frac{1}{2} b (1-x^2) (\gamma-\nu)(1-\nu)c_3 f_4 f_{10} f_{12} \right\} \\
& - 2(1+x)(1-\nu)(1+x\nu)d_1 f_2 f_{11} (c_3 f_3 - (1-ab)(1-\gamma)(1+\nu)f_{17}) \\
& \left. + \nu^{-1} c_3 (d_1 - d_2) f_4 g_3 (1-x)(1-\nu)^2 (1+x\nu) + c_3 d_2 f_4 g_{10} (1-\nu)^2 (1-x^2) \right],
\end{aligned} \tag{69}$$

$$\begin{aligned}
\Omega_\phi(x, y) = & \frac{\ell v_0(1+x)}{\sqrt{D_0}H(y, x)} \left[b(1+x)(1+y)(1+y\nu)d_1 d_2 f_1 f_{13} f_{14} \right. \\
& + \frac{(1+x)(1-y^2)(1-\nu)\nu c_3 (b(1+\nu)d_1 - 2(a-b)(1-\gamma)^2(1-\nu)\nu) f_3 g_{11}}{1+\nu} \\
& + \frac{(y-1)(1-\gamma)(1-\nu)\nu(1+y\nu + x(1+y(4-3\nu)))c_3 f_1 f_3 f_{10}}{1+\nu} \\
& \left. + \frac{2(a-b)(1-\gamma)^2 ((y-1)(x+y)(1-\nu)^3 \nu c_3 f_3 g_{12} + 2(1+x\nu)(1+y\nu)^2 d_1 f_1 g_1)}{1+\nu} \right],
\end{aligned} \tag{70}$$

where $\{f_i\}_{i=1,\dots,17}$ and $\{g_i\}_{i=1,\dots,13}$ are the polynomials of β up to $\mathcal{O}(\beta^2)$ and $\mathcal{O}(\beta^4)$, respectively, whose coefficients are functions of (a, b, γ, ν) . Both f_i and g_i are defined so that $f_i(\beta=0) = 1$ and $g_i(\beta=0) = 1$ except f_{10} . Note that D_0 defined in Eq. (63) has a simpler expression in terms of f_i

$$D_0 = f_8^2 - \frac{d_1 (f_1^2 - f_8^2)}{(1-a^2)(1-\gamma)(1-\nu^2)} + \frac{c_3 (f_3^2 - f_8^2)}{(1-a^2)(1-\gamma)(1+\nu)}. \tag{71}$$

A. Asymptotic infinity and the rod structure

The boundaries for the transformed solution in the C-metric coordinates, including the infinity, can be summarized as follows:

(i) ϕ -rotational axis : $\partial\Sigma_\phi = \{(x, y)|x = -1, -1 < y < -1/\nu\}$ with the rod vector $v_\phi := (0, 0, 1) = \partial/\partial\phi$, where we impose the periodicity of the coordinate ϕ as $\phi \sim \phi + 2\pi$ to ensures the absence of the conical singularities on $\partial\Sigma_\phi$,

(ii) Horizon: $\partial\Sigma_{\mathcal{H}} = \{(x, y)|-1 < x < 1, y = -1/\nu\}$ with the rod vector $v_{\mathcal{H}} := (1, \omega_\psi, \omega_\phi)$, where

$$\omega_\psi = \bar{\omega}_\psi \frac{\sqrt{D_0} f_4}{g_3}, \quad \omega_\phi = \bar{\omega}_\phi \frac{\sqrt{D_0} g_2}{f_3 g_3}, \quad (72)$$

(iii) Inner axis: $\partial\Sigma_{\text{in}} = \{(x, y)|x = 1, -1 < y < -1/\nu\}$ with the rod vector

$$v_{\text{in}} = (v_{\text{in}}^t, v_{\text{in}}^\psi, 1) \quad (73)$$

where $v_{\text{in}}^\psi = v_{\text{in}}^{\bar{\psi}} + \delta v_{\text{in}}^\psi$, with

$$v_{\text{in}}^t = \frac{\ell v_0 (a - b + \tilde{g}_1)}{\sqrt{D_0} f_1}, \quad (74)$$

$$\delta v_{\text{in}}^\psi = \frac{(1 - \nu) c_2 (f_5 f_{16} - f_1 f_2) - c_1 (f_6 f_{15} - f_1 f_2) (1 - ab) (\gamma - \nu) (1 + \nu)}{d_1 f_1 f_2}, \quad (75)$$

and \tilde{g}_1 is given by

$$1 + \frac{\tilde{g}_1}{a - b} := g_1 + \frac{c_3 f_3 f_{10} \nu}{(1 - \gamma) d_1 (\nu + 1) (a - b)} = f_{13} f_{14} + \frac{c_3 f_3 f_{10}}{2(a - b)(1 - \gamma) d_1} - \frac{c_1 c_3 (f_{13} f_{14} - f_{18} f_{19})}{2(a - b)(1 - \gamma)^2 \nu}. \quad (76)$$

Here, we should note that the presence of the t -component denotes the existence of a Dirac-Misner string singularity.

(iv) ψ -rotational axis: $\partial\Sigma_\psi = \{(x, y)|-1 < x < 1, y = -1\}$ with the rod vector $v_\psi := (0, 1, 0) = \partial/\partial\psi$, where we impose the periodicity of the coordinate ψ as $\psi \sim \psi + 2\pi$ to ensures the absence of the conical singularities on $\partial\Sigma_\psi$,

(v) Infinity: $\partial\Sigma_\infty = \{(x, y)|x \rightarrow y \rightarrow -1\}$: Introducing the coordinates (r, θ) by

$$x = -1 + \frac{4\ell(1 - \nu) \cos^2 \theta}{r^2}, \quad y = -1 - \frac{4\ell(1 - \nu) \sin^2 \theta}{r^2}, \quad (77)$$

we find that the limit $x \rightarrow y \rightarrow -1$ ($r \rightarrow \infty$) corresponds to the asymptotic infinity, since in this limit, the metric behaves as the five-dimensional Minkowski metric

$$ds^2 = - \left(1 - \frac{8G_5 M}{3\pi r^2} \right) dt^2 - \frac{8G_5 J_\psi \sin^2 \theta}{\pi r^2} dt d\psi - \frac{8G_5 J_\phi \cos^2 \theta}{\pi r^2} dt d\phi + dr^2 + r^2 \sin^2 \theta d\psi^2 + r^2 \cos^2 \theta d\phi^2 + r^2 d\theta^2, \quad (78)$$

where the mass M and two angular momenta J_ψ, J_ϕ can be written as

$$M = \frac{3\ell^2 \pi (\gamma + \nu) (d_1 - (1 - \nu) c_3)}{4(1 - a^2)(1 - \gamma)^2 (1 + \nu) D_0} \left[f_8^2 + \frac{(1 + \nu) c_2^2 (f_8^2 - f_5^2)}{(1 - \nu)(\gamma + \nu) (d_1 - (1 - \nu) c_3)} + \frac{(1 + \nu) d_1 ((1 - \gamma) (f_8^2 - f_1^2) - (1 + \nu) (f_8^2 - f_2^2))}{(1 - \nu)(\gamma + \nu) (d_1 - (1 - \nu) c_3)} \right], \quad (79)$$

$$J_\psi = - \frac{\ell^3 \pi v_0 (c_3 (d_1 - d_2) f_4 g_3 + c_2 (c_1 c_3 - b d_1) f_5 g_9)}{4D_0^{3/2} \nu (1 - a^2) (1 - \gamma)^3}, \quad (80)$$

$$J_\phi = - \frac{\ell^3 \pi v_0 (2\nu c_3 f_3 f_1 f_{10} - (2\nu c_3 f_3 g_{12} + d_1 f_1 g_1) (a - b) (1 - \gamma) (1 - \nu))}{2D_0^{3/2} (1 - a^2) (1 - \gamma)^2 (1 - \nu^2)}, \quad (81)$$

where we used the expression Eq. (71) in the mass formula.

Finally, we summarize some observations on how the Ehlers transformation changes the rod structure. This transformation does not change the positions of both the rotational axes at $x = -1$ and at $y = -1$. Additionally, it does not alter the directions of the rod vectors after the removal of the global rotation by the coordinate transformation. However, it does not preserve the regularity, specifically the absence of conical singularities on the two axes. Therefore, to ensure the regularity of the transformed solution, we did not impose regularity on the seed solution. Additionally, this transformation also does not change the positions of the horizon $\partial\Sigma_{\mathcal{H}}$ at $y = -1/\nu$ and the inner axis $\partial\Sigma_{\text{in}}$ at $x = 1$, but it changes the directions of the rod vectors. Specifically, $v_{\mathcal{H}}$ and v_{in} change from $(1, \bar{\omega}_\psi, \bar{\omega}_\phi)$ to $(1, \omega_\psi, \omega_\phi)$ and from $(v_{\text{in}}^{\bar{t}}, v_{\text{in}}^{\bar{\psi}}, 1)$ to $(v_{\text{in}}^t, v_{\text{in}}^\psi, 1) = ((v_{\text{in}}^{\bar{t}} + \tilde{g}_1)D_0^{-\frac{1}{2}}f_1^{-1}, v_{\text{in}}^{\bar{\psi}} + \delta v_{\text{in}}^\psi, 1)$, respectively.

Since the presence of the nonzero t -component v_{in}^t in v_{in} still yields a Dirac-Misner string singularity, one can remove it by setting

$$a - b + \tilde{g}_1 = 0, \quad (82)$$

where note that $\tilde{g}_1 \rightarrow 0$ as $\beta \rightarrow 0$. Therefore, if we choose the seed solution with $\beta = 0$ that does not have the Dirac-Misner string singularity (corresponding to $a = b$), as in Ref. [30], the Dirac-Misner string singularity inevitably appears due to $\tilde{g}_1 \neq 0$ for $\beta \neq 0$. Thus, to obtain a regular solution without a Dirac-Misner string singularity by the transformation, we must choose a singular seed solution having a Dirac-Misner string singularity. Moreover, since $v_{\text{in}}^\psi \neq v_{\text{in}}^{\bar{\psi}}$ for $\beta \neq 0$, this transformation does not preserve the horizon topology, which implies that even if the horizon cross-section of the seed solution has a ring topology $S^2 \times S^1$ ($v_{\text{in}}^{\bar{\psi}} = 0$), the transformed solution does not ($v_{\text{in}}^\psi \neq 0$).

B. Regularity conditions for black rings

Now, we consider the condition that the transformed solution describes a regular black ring, meaning that the spacetime has neither curvature singularities, conical singularities, orbifold singularities, nor Dirac-Misner string singularities either inside or outside the horizons.

As shown in Ref. [5], the topology of the horizon cross-section is assured to be $S^2 \times S^1$ when the rod vector on $\partial\Sigma_{\text{in}}$ can be expressed in the form $v_{\text{in}} = (0, 1, 0)$. This requires that

$$c = \frac{\ell v_0(a - b + \tilde{g}_1)}{\sqrt{D_0}f_1} = 0, \quad (83)$$

$$n = \frac{-c_1 f_6 f_{15}(1 - ab)(\gamma - \nu)(1 + \nu) + (1 - \nu)c_2 f_5 f_{16}}{d_1 f_1 f_2} = 0, \quad (84)$$

where Eq. (83) is the condition for the absence of a Dirac-Misner string singularity on $\partial\Sigma_{\text{in}}$ and Eq. (84) assures the ring topology of the horizon cross-section. Moreover, the absence of conical singularities on $\partial\Sigma_{\text{in}}$ requires the condition:

$$\lim_{\rho \rightarrow 0} \sqrt{\frac{\rho^2 g_{\rho\rho}}{g_{ij} v_{\text{in}}^i v_{\text{in}}^j}} = \frac{\Delta\phi}{2\pi} \iff \frac{d_1^2 f_1^2 f_2^2}{(1 - a^2)^2 (1 - \gamma)^3 (1 - \nu)^2 (1 + \nu) D_0^2} = 1. \quad (85)$$

From these three conditions, the number of the independent parameters are reduced from six ($\ell, a, b, \gamma, \nu, \beta$) to three. The two conditions (84) and (85) can be solved with respect to a and b as functions of (γ, ν, β) . Whether regular or singular, these equations have many branches, one of which corresponds to the Pomerasky-Sen'kov solution:

$$a = \frac{(\gamma - \nu)\sqrt{1 + \nu} - \sqrt{2(1 - \nu)\nu(\gamma - \nu)}}{\beta[(1 - \gamma)\sqrt{1 + \nu} \pm (1 - \nu)\sqrt{1 - \gamma}]} + \frac{\sqrt{\nu + 1}(\gamma - \nu) \pm 2\sqrt{1 - \gamma}(1 - \nu) + (\gamma - \nu)^{-1}\sqrt{2(1 - \nu)\nu(\gamma - \nu)} \left(1 - 2\gamma + \nu \mp (1 - \nu)\sqrt{(1 - \gamma)/(1 + \nu)}\right)}{(1 - \gamma)\sqrt{\nu + 1} \pm \sqrt{1 - \gamma}(1 - \nu)}, \quad (86)$$

$$b = \frac{\sqrt{1 + \nu}(1 - \gamma) \mp \sqrt{1 - \gamma}(1 - \nu)}{(\gamma - \nu)\sqrt{1 + \nu}}. \quad (87)$$

Together with Eqs. (86) and (87), further imposing Eq. (83), also determined β as the function of γ and ν :

$$\beta = \frac{1}{\mathcal{B}_1 \pm \mathcal{B}_2 \sqrt{(1-\gamma)(1+\nu)}} \left[(1-\nu^2) (\gamma^3 (-1+4\nu-3\nu^2) + \gamma^2 (8-3\nu+4\nu^2+7\nu^3)) \right. \\ \left. + \gamma (-8-8\nu+\nu^2-12\nu^3-5\nu^4) + \nu (8+3\nu^2+4\nu^3+\nu^4) \right. \\ \left. \pm 2\sqrt{1+\nu}\sqrt{1-\gamma}(1-\nu) (\gamma^2 (-2+\nu+2\nu^2+3\nu^3) - 4\gamma (-1-\nu+\nu^3+\nu^4) + \nu (-4-2\nu-\nu^2+2\nu^3+\nu^4)) \right. \\ \left. + 2(\gamma-\nu)\sqrt{2(-1+\nu)\nu(-\gamma+\nu)} (\sqrt{1+\nu}(1-\nu) (4+5\nu+3\nu^2-\gamma(3+3\nu+2\nu^2)) \right. \\ \left. \pm \sqrt{1-\gamma}(1+\nu) (-4+\gamma(-1+\nu)^2+\nu-4\nu^2-\nu^3)) \right], \quad (88)$$

where \mathcal{B}_1 and \mathcal{B}_2 are given as

$$\mathcal{B}_1 = (1-\nu)^2 (-16-12\nu-37\nu^2-29\nu^3-3\nu^4+\nu^5-\gamma^2(5+17\nu+11\nu^2-\nu^3)+2\gamma(10+19\nu+25\nu^2+9\nu^3+\nu^4)), \\ \mathcal{B}_2 = (1-\nu)(16-12\nu+33\nu^2+5\nu^3+23\nu^4-\nu^5-\gamma^2(-1+\nu)(1+\nu)^2-2\gamma(6+7\nu+13\nu^2+5\nu^3+\nu^4)). \quad (89)$$

Finally, substituting this to Eq. (86), we can obtain

$$a = \frac{(\gamma-3)(1-\nu)\sqrt{\nu+1} \pm \sqrt{1-\gamma}(\nu^2+3)}{(\gamma-1)(\nu-1)\sqrt{\nu+1} \mp \sqrt{1-\gamma}(\nu+1)^2}. \quad (90)$$

V. EQUIVALENCE WITH THE POMERANSKY-SEN'KOV SOLUTION

To summarize, it can be seen from Eqs. (87), (88) and (90) that the transformed solution can be expressed only in terms of the three parameters (ℓ, ν, γ) . Now, we demonstrate that either of the above (\pm) -branches is identical to the Pomeransky-Sen'kov solution [9], where the metric is given by Eq. (B1) and is three parameter family of (k, α_1, α_2) . This solution describes a doubly rotating black ring with $J_\psi > 0$ and $J_\phi > 0$ for the parameter range of $k > 0$ and $0 \leq \alpha_2 \leq \alpha_1 < 1$. The parameter region is bounded by the singly rotating solution, i.e., Emparan-Reall black ring solution, $\{(\alpha_1, \alpha_2) | 0 < \alpha_1 < 1, \alpha_2 = 0\}$ and the extremal solution $\{(\alpha_1, \alpha_2) | 0 < \alpha_1 = \alpha_2 < 1\}$. Since the $(-)$ -branch yields the same solution as the $(+)$ -branch and these differs only in the signature of the angular momentum around the ψ -rotational axis ($J_\psi \rightarrow -J_\psi$), we may consider only the $(+)$ -branch without loss of generality.

According to the uniqueness theorem for a vacuum black ring solution [5, 17, 35], under the assumptions of the presence of three mutually commuting Killing vector fields, a timelike Killing vector field and two axial Killing vector fields, besides the trivial topology of $(\mathbb{R} \times \{\mathbb{R}^4 \setminus D^2 \times S^2\})$ of the domain of outer communication, an asymptotically flat, stationary rotating black ring with non-degenerate connected event horizon of the cross-section topology $S^2 \times S^1$ is characterized uniquely by the mass, two independent angular momenta, and additional information on the rod structure such as the ratio of the S^2 radius to the S^1 . Therefore, if our transformed solution exactly coincides with the Pomeransky-Sen'kov solution, using this theorem, we can extract the relation between two sets of the parameters, (ℓ, ν, γ) in our transformed solution and (k, α_1, α_2) in the Pomeransky-Sen'kov solution by comparing the rod length and asymptotic charges between two solutions. To relate these two three-parameter families of solutions, it is sufficient to consider only the rod data (the rod lengths of the horizon and inner axis) and the mass for the two solutions. In the following discussion, we examine the correspondence between our solution obtained in the previous section and the non-extremal Pomeransky-Sen'kov solution, whose parameter region is defined by $(\alpha_1, \alpha_2) | 0 \leq \alpha_2 < \alpha_1 < 1$, as the uniqueness theorem applies only to non-degenerate solutions.

In the Weyl-Papapetrou coordinates (ρ, z) , the rod structure, for both solutions, our transformed solution (65) and Pomeransky-Sen'kov metric (B1), can be summarized as follows:

Solutions	Rods
Our transformed solution	$(-\infty, -\nu\ell^2), (-\nu\ell^2, \nu\ell^2), (\nu\ell^2, \ell^2), (\ell^2, \infty)$
Pomeransky-Sen'kov solution	$(-\infty, -\frac{\alpha_1-\alpha_2}{1-\alpha_1\alpha_2}k^2), (-\frac{\alpha_1-\alpha_2}{1-\alpha_1\alpha_2}k^2, \frac{\alpha_1-\alpha_2}{1-\alpha_1\alpha_2}k^2), (\frac{\alpha_1-\alpha_2}{1-\alpha_1\alpha_2}k^2, k^2), (k^2, \infty)$

Then, two metrics have the rods of the same length if

$$\ell = k, \quad \nu = \frac{\alpha_1 - \alpha_2}{1 - \alpha_1\alpha_2}. \quad (91)$$




FIG. 1: $\nu = \text{const.}$ curves (blue curves) in the (α_1, α_2) plane mapped by Eq. (93). Each curves correspond to $\nu = 1/10, 2/10, \dots, 9/10$ and $99/100$. γ runs from $\nu(3 - \nu)/(1 + \nu)$ to ν , from the bottom to the top. The $\nu = \text{const.}$ curve approaches the extremal limit $\alpha_1 = \alpha_2$ (gray line) at $\nu \rightarrow 0$.

Moreover, as the third condition, we compare the mass (79) for our transformed solution with the mass (B6) for the Pomeransky-Sen'kov solution. Using Eqs. (87), (88) and (90), we can express the mass (79) as a simpler form,

$$M = \frac{3\ell^2\pi(\gamma + \nu) \left(\nu(\gamma - \nu) + \sqrt{2\nu(1 - \nu^2)(\gamma - \nu)} \right)}{2(\gamma - \nu)(2 - \nu^2 - \gamma\nu)}. \quad (92)$$

Equating Eq. (92) with Eq. (B6), along with Eq. (91), one can express (α_1, α_2) as functions of (ν, γ)

$$\alpha_1 = U_+ - \sqrt{U_+^2 - 1}, \quad \alpha_2 = U_- - \sqrt{U_-^2 - 1}, \quad (93)$$

where

$$U_{\pm} = \frac{\nu(-\gamma^2(1 \mp \nu)^2 + 2\gamma(\nu^3 \pm 2\nu^2 + \nu \mp 4) + \nu(3\nu^3 \pm 2\nu^2 + 3\nu \pm 8)) + 2\sqrt{2}(\gamma + \nu)\sqrt{\nu(1 - \nu^2)^3(\gamma - \nu)}}{\nu(\gamma(1 \mp \nu) - \nu(3 \pm \nu))^2}. \quad (94)$$

Note that the \pm symbol in U_{\pm} are not related to the previous (\pm) -branches.

It should be noted that Eq. (93) does not provide a one-to-one mapping from the range of $\{(\nu, \gamma) \mid 0 < \nu < \gamma < 1\}$ to the range of $\{(\alpha_1, \alpha_2) \mid 0 \leq \alpha_2 < \alpha_1 < 1\}$, because as shown in Fig.1 the parameter subregion of the (+)-branch for our solution: $\{(\nu, \gamma) \mid \nu < \gamma \leq \nu(3 - \nu)/(1 + \nu), 0 < \nu < 1\}$ has a one-to-one correspondence to the entire parameter region of the Pomeransky-Sen'kov solution: $\{(\alpha_1, \alpha_2) \mid 0 \leq \alpha_2 < \alpha_1 < 1\}$. Moreover, using Eqs. (87), (88), (90) and (93), one can easily show that the two angular momenta $|J_{\psi}|$ and $|J_{\phi}|$ for our solution, given by Eqs. (80) and (81) exactly coincide with those for the Pomeransky-Sen'kov solution, given by Eqs. (B7) and (B8). Therefore, it can be expected from the uniqueness theorem [5, 17] that the two metrics, the (+)-branch metric for our solution and the Pomeransky-Sen'kov solution, are isometric, as long as the metric for the (+)-branch is regular and free from closed timelike curves on and outside the horizon.

Actually, one can directly show that the (+)-branch metric (65) coincides with the Pomeransky-Sen'kov metric (B1) using Eqs. (87), (88), (90) and (93) as follows. As given in the Appendix B, the metric of the Pomeransky-Sen'kov solution is written in the C-metric coordinates (x, y) as our transformed metric, but it should be noted that the choice of the C-metric coordinates (x, y) is not unique. For the latter, the function $G(u)$ in Eq. (43) can be written as a cubic function with respect to u , whereas for the former, the corresponding function (B2) can be expressed in a quartic function as

$$G_4(u) = (1 - u^2)(1 + \alpha_1 u)(1 + \alpha_2 u), \quad (95)$$

where the outer and inner horizons correspond to $y = -1/\alpha_1$ and $y = -1/\alpha_2$. These two metrics apparently differ, but are related by the coordinate transformation $(x, y) \rightarrow (x_{\text{new}}, y_{\text{new}})$, referred to as the Möbius transformation [18],

which is defined by

$$x = \frac{x_{\text{new}} + \alpha_2}{1 + \alpha_2 x_{\text{new}}}, \quad y = \frac{y_{\text{new}} + \alpha_2}{1 + \alpha_2 y_{\text{new}}}, \quad \nu = \frac{\alpha_1 - \alpha_2}{1 - \alpha_1 \alpha_2}, \quad (96)$$

which transforms the C-metric written in the cubic function to the C-metric written in the quartic function.

$$\frac{dx^2}{G(x)} - \frac{dy^2}{G(y)} = (1 - \alpha_1 \alpha_2) \left[\frac{dx_{\text{new}}^2}{G_4(x_{\text{new}})} - \frac{dy_{\text{new}}^2}{G_4(y_{\text{new}})} \right]. \quad (97)$$

Thus, in the new coordinates $(x_{\text{new}}, y_{\text{new}})$, using Eqs. (87), (88), (90) and (93), it is straightforward to show each of (+) or (-)-branch for our metric (65) obtained by the Ehlers transformation coincides with those in the Pomeransky-Sen'kov metric (B1).

VI. SUMMARY AND DISCUSSION

In this paper, by employing the sigma model approach, we have obtained an exact solution for a doubly rotating black ring to the five-dimensional vacuum Einstein equation. Giusto and Saxena have previously shown that the $SO(2, 1)$ subgroup of the $SL(3, \mathbb{R})$ symmetry can produce a one-parameter family of stationary solutions with angular momenta. However, unlike black holes, merely applying this technique to the static black ring or the Emparan-Reall black ring does not result in a regular singly rotating black ring or a regular doubly rotating black ring, due to the emergence of a Dirac-Misner string singularity. To tackle this problem, we have utilized a vacuum solution of a rotating black ring, which already includes a Dirac-Misner string singularity, as the seed solution for the Ehlers transformation. The fundamental idea is similar to the construction of a charged black ring performed in our previous work, where as a vacuum seed for the electric Harrison transformation, we chose the black ring solution with a Dirac-Misner string singularity. By appropriately adjusting the solution's parameters after the transformation, the resultant solution has proven to be regular, showing no signs of curvature singularities, conical singularities, orbifold singularities, Dirac-Misner string singularities, or closed timelike curves, both on and outside the horizon. As expected from the uniqueness theorem for a stationary and bi-axisymmetric black ring [5], this solution exactly coincides with the Pomeransky-Sen'kov solution.

According to the uniqueness theorem for charged black rings in the five-dimensional supergravity [35, 41], under the assumptions of the trivial topology $(\mathbb{R} \times \{\mathbb{R}^4 \setminus D^2 \times S^2\})$ of the domain of outer communication, an asymptotically flat, stationary and bi-axisymmetric charged rotating black ring with non-degenerate connected event horizon is characterized uniquely by the mass, electric charge, two independent angular momenta, dipole charge, and additional information on the rod structure such as the ratio of the S^2 radius to the S^1 radius. Elvang *et al.* [42] constructed an example of such a dipole ring, which is originating from a seven-parameter family of non-supersymmetric black ring solutions, furthermore, we gave another example, which was constructed by the electric Harrison transformation. These both solutions carry the five physical quantities, but three only are independent among them. As conjectured by the authors in Ref. [42], it is natural that a more general non-Bogomol'nyi-Prasad-Sommerfield (BPS) black ring solution exists, characterized by its mass, two independent angular momenta, electric charge, and a dipole charge, which are independent of the other asymptotic conserved charges, which finally may have been constructed in Ref. [43]. However, this solution has its considerably complex and formal expression, at this point, we do not know whether this solution truly describes a regular black ring solution with the five independent parameters, including the regularity, the horizon topology, asymptotic flatness, as well as physical properties. Therefore, to investigate them, we need a more compact form of the solution, such as the C-metric form. We anticipate that applying our method to the five-dimensional minimal supergravity may enable us to derive such an exact solution.

Acknowledgement

R.S. was supported by JSPS KAKENHI Grant Number JP24K07028. S.T. was supported by JSPS KAKENHI Grant Number 21K03560.

Appendix A: Coefficients k_1^\pm, \dots, k_9^\pm

$$\begin{aligned}
k_1^\pm &= \frac{c_1 c_3 (\nu - 1)(\gamma + \nu) (2(\gamma - 1)c_1 \nu (\nu + 1) \pm 2(1 - \gamma)\nu (\nu^2 - 1) \pm d_1(1 - \nu) \pm 2d_2\nu)}{8(\gamma - 1)}, \\
k_2^\pm &= -\frac{(1 \pm a)(1 - \gamma)d_1(1 - \nu)^3(\gamma + \nu)}{4(\nu + 1)}, \\
k_3^\pm &= \frac{c_3 d_1 (1 - \nu^2)(\gamma + \nu) ((1 - \gamma)(1 - \nu) \pm 4(\gamma - \nu)\nu b)}{8(\gamma - 1)\nu}, \\
k_4^\pm &= \frac{c_3 d_1 (1 - \nu)(\gamma + \nu)}{8\nu}, \\
k_5^\pm &= \frac{(1 \pm a)(\gamma - 1)c_3 (\nu - 1)^3(\gamma + \nu)}{4(\nu + 1)}, \\
k_6^\pm &= -\frac{\nu}{4} \left[2c_3^2 (\nu - 1)^2 (\nu - \gamma) - 2(1 \pm a)(1 - \gamma)c_3 (\nu - 1)^2 (2(1 - \gamma)\nu \pm b(\nu + 1)(\gamma - \nu)) \right. \\
&\quad \left. - (1 \mp a)(1 - \gamma)d_1(1 - \nu) (3\gamma\nu + \gamma - \nu^2 - 3\nu \pm 2b(\nu + 1)(\gamma - \nu)) + c_3 d_1 (\nu - 1)(3\nu - \gamma \mp 2a(1 - \gamma)) \right], \\
k_7^\pm &= \frac{c_3 (1 - \nu)^3}{8(\nu + 1)} \left[4a(1 - \gamma)c_1 (\nu + 1)\nu \pm 2a(\gamma - 1)(\nu - 1)\nu(\gamma + \nu) \pm 4(\gamma - 1)c_2\nu \right. \\
&\quad \left. + 2(\gamma - 1)(\nu - 1)\nu(\gamma - \nu - 2) - d_1(\gamma + \nu) \right], \\
k_8^\pm &= \frac{d_1 (\nu - 1)}{2(\nu + 1)} \left[2ab(\gamma - 1)(\nu + 1)(\gamma - \nu) \pm (1 - \gamma) (a(\nu - 1)(\gamma + \nu) + 2c_2) \right. \\
&\quad \left. + b^2(\nu + 1) (\nu^2 - \gamma^2) + (1 - \gamma)(\nu + 1)(\gamma - 3\nu) \right] \\
k_9^\pm &= \frac{(\gamma + \nu) (1 - a^2 + \gamma(b - a)^2 - \nu (1 - 2ab + b^2))}{8(1 \mp a)(1 - \gamma)}.
\end{aligned}$$

Appendix B: Pomeransky-Sen'kov's doubly-spinning black rings

Here we recapitulate the metric for the doubly-spinning black rings [9]³

$$\begin{aligned}
ds^2 &= -\frac{H_{\text{PS}}(y, x)}{H_{\text{PS}}(x, y)} (dt + \Omega_{\text{PS}})^2 + \frac{1}{H_{\text{PS}}(y, x)} [F_{\text{PS}}(x, y)d\psi^2 - 2J_{\text{PS}}(x, y)d\psi d\phi - F_{\text{PS}}(y, x)d\phi^2] \\
&\quad + \frac{2k^2 H_{\text{PS}}(x, y)}{(x - y)^2 (1 - \alpha_1 \alpha_2)^2} \left(\frac{dx^2}{G_4(x)} - \frac{dy^2}{G_4(y)} \right), \tag{B1}
\end{aligned}$$

³ We have exchanged ψ and ϕ so that ψ generates the S^1 -rotation.

where

$$\begin{aligned}
G_4(u) &= (1-u^2)(1+\alpha_1 u)(1+\alpha_2 u), \\
H_{\text{PS}}(x, y) &= 1 - \alpha_1^2 \alpha_2^2 + 2(1-x^2)y\alpha_1\alpha_2(\alpha_1+\alpha_2) + (\alpha_1+\alpha_2)^2 + 2x(\alpha_1+\alpha_2)(1-y^2\alpha_1^2\alpha_2^2) \\
&\quad + x^2 y^2 \alpha_1 \alpha_2 (1 - \alpha_1^2 \alpha_2^2 - (\alpha_1 + \alpha_2)^2), \\
F_{\text{PS}}(x, y) &= \frac{2k^2}{(x-y)^2(1-\alpha_1\alpha_2)^2} \left[G(y) \left\{ 2(\alpha_1+\alpha_2)^2 + x(1+\alpha_1^2)(\alpha_1+\alpha_2)(1+\alpha_2^2) \right. \right. \\
&\quad \left. \left. + x^3(\alpha_1+\alpha_2)(1-3\alpha_1^2\alpha_2^2+2\alpha_1^3\alpha_2^3 - (\alpha_1+\alpha_2)^2) - x^4\alpha_1\alpha_2(1-\alpha_1\alpha_2)(-1+\alpha_1^2\alpha_2^2+(\alpha_1+\alpha_2)^2) \right. \right. \\
&\quad \left. \left. + x^2(1+\alpha_1\alpha_2)(-(\alpha_1+\alpha_2)^2+(-1+\alpha_1\alpha_2)^2) \right\} \right. \\
&\quad \left. + (1-y^2)G(x)((1+\alpha_1\alpha_2)(-(\alpha_1+\alpha_2)^2+(-1+\alpha_1\alpha_2)^2) - y(\alpha_1+\alpha_2)(-1+\alpha_2^2+\alpha_1^2(1+3\alpha_2^2))) \right], \\
J_{\text{PS}}(x, y) &= \frac{2k^2(1-x^2)(1-y^2)(\alpha_1+\alpha_2)\sqrt{\alpha_1\alpha_2}}{(x-y)(1-\alpha_1\alpha_2)^2} \left[1 + (\alpha_1+\alpha_2)^2 - \alpha_1^2\alpha_2^2 + 2\alpha_1\alpha_2(\alpha_1+\alpha_2)(x+y) \right. \\
&\quad \left. + \alpha_1\alpha_2((\alpha_1+\alpha_2)^2 + \alpha_1^2\alpha_2^2 - 1)xy \right], \\
\Omega_{\text{PS}}(x, y) &= -\frac{2k(\alpha_1+\alpha_2)\sqrt{\alpha_1\alpha_2(1-\alpha_1^2)(1-\alpha_2^2)}}{H(y, x)} \left[(1-x^2)y d\phi \right. \\
&\quad \left. - \frac{(1+y)(1+\alpha_1+\alpha_2-\alpha_1\alpha_2+2x(1-y)\alpha_1\alpha_2-x^2y\alpha_1\alpha_2(-1+\alpha_1+\alpha_2+\alpha_1\alpha_2))}{(1-\alpha_1)(1-\alpha_2)} d\psi \right].
\end{aligned} \tag{B2}$$

For the convenience, we have replaced the original parameters (λ, ν) in Ref. [9] with the following (α_1, α_2)

$$\lambda = \alpha_1 + \alpha_2, \quad \nu = \alpha_1\alpha_2, \quad 0 \leq \alpha_2 \leq \alpha_1 < 1. \tag{B3}$$

On the horizon at $y = -1/\alpha_1$, we obtain the horizon velocity

$$\omega_{\psi}^{\text{PS}} = \frac{\sqrt{(1-\alpha_1^2)(1-\alpha_2^2)}}{2k(1+\alpha_1)(1+\alpha_2)}, \quad \omega_{\phi}^{\text{PS}} = \frac{(1-\alpha_1)(1+\alpha_1^2)(1-\alpha_2)\alpha_2}{2k\sqrt{\alpha_1\alpha_2}(\alpha_1+\alpha_2)\sqrt{(1-\alpha_1^2)(1-\alpha_2^2)}}. \tag{B4}$$

and the surface gravity

$$\kappa_{\text{PS}} = \frac{(1-\alpha_1)(\alpha_1-\alpha_2)(1-\alpha_1\alpha_2)}{4k\alpha_1(1+\alpha_2)(\alpha_1+\alpha_2)}. \tag{B5}$$

The asymptotic charges are given by

$$M_{\text{PS}} = \frac{3k^2\pi(\alpha_1+\alpha_2)}{G_5(1-\alpha_1)(1-\alpha_2)}, \tag{B6}$$

$$J_{\phi}^{\text{PS}} = \frac{4k^3\pi\sqrt{\alpha_1\alpha_2}(\alpha_1+\alpha_2)\sqrt{(1-\alpha_1^2)(1-\alpha_2^2)}}{G_5(1-\alpha_1)(1-\alpha_2)(1-\alpha_1\alpha_2)^2}, \tag{B7}$$

$$J_{\psi}^{\text{PS}} = \frac{2k^3\pi(\alpha_1+\alpha_2)\sqrt{(1-\alpha_1^2)(1-\alpha_2^2)}(1+\alpha_2+\alpha_1^2\alpha_2(1+\alpha_2)+\alpha_1(1-6\alpha_2+\alpha_2^2))}{G_5(1-\alpha_1)^2(1-\alpha_2)^2(1-\alpha_1\alpha_2)^2}. \tag{B8}$$

Appendix C: Coefficients f_i and g_i

The explicit forms of the coefficients f_i and g_i are shown below.

1. f_1, \dots, f_{22}

f_i are quadratic functions of β . f_i for $14 \leq i \leq 22$ does not appear in the metric functions but does only in the expressions of g_i .

$$f_1 = 1 + 2(a-b)\beta + \frac{\beta^2}{\gamma^2 - \nu^2} \left[(a-b)^2\gamma^2 - (1-a)^2\gamma(1+2b+(3+2b)\nu) \right. \\ \left. + \nu((1-a)^2(3+2b) + (1-2a+2(1-a+a^2)b-b^2)\nu) \right], \quad (C1)$$

$$f_2 = 1 + 2(1-ab)\beta + \frac{\beta^2}{\gamma^2 - \nu^2} \left[((2-a)a - 2(1-a+a^2)b - (1-2a)b^2)\gamma^2 \right. \\ \left. + (1-a)^2\gamma(1+2b+(3+2b)\nu) - \nu((1-a)^2(3+2b) + (1-b)(1+(1-2a)b)\nu) \right], \quad (C2)$$

$$f_3 = 1 + 2(a-b)\beta + \frac{\beta^2}{\gamma^2 - \nu^2} \left[(a-b)^2\gamma^2 - 2(1-a)^2(1+a) + (1-a)^2(1-2a+2b)\nu \right. \\ \left. + (1-2a+2(1-a+a^2)b-b^2)\nu^2 + (1-a)^2\gamma(1+2a-2b-(1-2a+2b)\nu) \right], \quad (C3)$$

$$f_4 = 1 - \frac{2\beta}{\gamma - \nu} (a^2 - 1 - a(a-b)\gamma + \nu(1-ab)) + \frac{\beta^2}{\gamma^2 - \nu^2} \left[2(1-a)^2(1+a) - (1-2a)(a-b)^2\gamma^2 \right. \\ \left. - (1-a)(1-5a+4b)\nu - (1-2ab-b^2+2ab^2)\nu^2 - (1-a)\gamma(1-a-4a^2+4ab-(1-5a+4b)\nu) \right], \quad (C4)$$

$$f_5 = 1 - \frac{2\beta}{c_2} (2a(-1+\gamma)\nu + ab^2(\gamma-\nu)(1+\nu) + b(-1+a^2-a^2\gamma-(2+a^2)(-1+\gamma)\nu+\nu^2)) \\ + \frac{\beta^2}{(\gamma^2 - \nu^2)c_2} \left[b\gamma(2(1-a)^2(1+a) + (1-a)(-1+a+4a^2-4ab)\gamma + (-1+2a)(-a+b)^2\gamma^2) \right. \\ \left. + \nu \left\{ (2(-2+a+6a^2-4a^3) + (4-20a+17a^2-6a^3)b + 4(2-3a+2a^2)b^2 + (1-2a)b^3)\gamma^2 \right. \right. \\ \left. \left. + (-a+b)^2(-4+2a-b+2ab)\gamma^3 - 2(1-a)^2(1+a)(2+b) + 2(1-a)^2(4+5a+(-1+3a)b-2b^2)\gamma \right\} \right. \\ \left. + \nu^2 \left\{ (1-a)(2(2-3a+a^2) + (3-9a+2a^2)b+4b^2) \right. \right. \\ \left. \left. + (4(a-2)(1-a)^2 + (-10+26a-21a^2+4a^3)b - 2(4-7a+2a^2)b^2 + (1-2a)b^3)\gamma \right. \right. \\ \left. \left. + (2(2-a)(1-a)^2 + (5-14a+10a^2-2a^3)b + (4-6a+4a^2)b^2 + (1-2a)b^3)\gamma^2 \right\} \right. \\ \left. - (2a - (6-6a+5a^2)b + 4ab^2 + (1-2a)b^3)(1-\gamma)\nu^3 + (1-b)b(1+(1-2a)b)\nu^4 \right], \quad (C5)$$

$$f_6 = 1 - \frac{2\beta}{(\gamma - \nu)(1 + \nu)c_1} \left[a^2 - 1 + (2 - 2a + a^2 + b - 2ab)\nu - (1-b)(1-2a+b)\nu^2 - (1-b)b\nu^3 \right. \\ \left. + (a-b)^2\gamma^2(1+\nu) + \gamma(1-2a^2-b+2ab-2(1-a+a^2-2ab+b^2)\nu + (1-b)(1-2a+2b)\nu^2) \right] \\ + \frac{\beta^2}{(\gamma^2 - \nu^2)(1 + \nu)c_1} \left[2(1-a)^2(1+a) - (1-a)(2-5a-a^2+4b)\nu \right. \\ \left. + (-2-2a+6a^2-3a^3 + (3-4a+3a^2)b + (2-3a)b^2)\nu^2 + (2-3a+3a^2b-3ab^2+b^3)\nu^3 + (1-b)^2b\nu^4 \right. \\ \left. + (a-b)\gamma^2(4a^2-1-2a+(2-3a)b + ((2-a)^2-2ab+b^2)\nu - (3(1-a)^2 + (2-a)b-b^2)\nu^2) \right. \\ \left. - (a-b)^3\gamma^3(1+\nu) + (1-a)\gamma(a-2+5a^2-4ab+(2-9a-a^2+8b)\nu) \right. \\ \left. + (2+5a-12a^2+6a^3 - (8-12a+7a^2)b + 3ab^2-b^3)\gamma\nu^2 - (2-3a+3a^2b-3ab^2+b^3)\gamma\nu^3 \right], \quad (C6)$$

$$\begin{aligned}
f_7 = & 1 - \frac{2\beta}{b(\gamma - \nu)(1 + \nu)} \left[(b-1)b\gamma + (2-2a+b-b^2)(1-\gamma)\nu + (1-b)b\nu^2 \right] \\
& + \frac{\beta^2}{b(\gamma^2 - \nu^2)(1 + \nu)} \left[b\gamma((1-a)^2 + ((2-a)a - 2b + b^2)\gamma) - (1-b)^2b\nu^3 \right. \\
& + (-(1-a)^2(4+3b) + 4(1-a)(1-2a + (2-a)b)\gamma + (4(1-a)a - (4-6a+a^2)b - 2b^2 + b^3)\gamma^2)\nu \\
& \left. + (4(-1+a) + (2-3a)ab + 2b^2 - b^3 + (1-a)(4-b-3ab)\gamma)\nu^2 \right], \tag{C7}
\end{aligned}$$

$$\begin{aligned}
f_8 = & 1 + \frac{2\beta}{(a-b)(\gamma - \nu)} \left[1 - a^2 + (a-b)^2\gamma - (1-2ab+b^2)\nu \right] \\
& - \frac{\beta^2}{(a-b)(\gamma^2 - \nu^2)} \left[2(-1+a+a^2-a^3) - a\gamma(1+2a-3a^2) - (a^3-b^3)\gamma^2 - b^2\gamma(2+2a^2+a(-4+3\gamma)) \right. \\
& + b(-2(1-a)^2(1+a) + (1+4a-7a^2+2a^3)\gamma + 3a^2\gamma^2) \\
& - \nu(1-a)(\gamma-1)((a-5)a + (5-3a+2a^2)b + 2(1-a)b^2) \\
& \left. - (-2+a+(-1+6a-2a^2)b + (-2+a-2a^2)b^2 + b^3)\nu^2 \right], \tag{C8}
\end{aligned}$$

$$\begin{aligned}
f_9 = & 1 + 2(a-b)\beta + \frac{\beta^2}{(\gamma^2 - \nu^2)c_1} \left[-2(1-a)^2(1+a) - (a-b)^3\gamma^3 + (1-a)^2(4+a+2b)\nu - (1-b)b(1-2a+b)\nu^3 \right. \\
& - (2-5a+4a^2 + (1-2a-a^2)b + ab^2)\nu^2 - (a-b)\gamma^2(1-a(2-b) - ((1-a)^2 - ab + b^2)\nu) \\
& \left. - \gamma((1-a)^2(-2-3a+2b) + 2(2-3a+a^3)\nu + (-2+5a-4a^2 + (-2+4a+a^2)b - 3ab^2 + b^3)\nu^2) \right], \tag{C9}
\end{aligned}$$

$$f_{10} = \frac{4(1-a)(1-\gamma)d_1}{\gamma^2 - \nu^2} \beta(1 + (a-b)\beta), \tag{C10}$$

$$\begin{aligned}
f_{11} = & 1 - \frac{2\beta(-1+a-a\gamma + b(-1+a+\gamma) + (-1+ab)\nu)}{\gamma + \nu} \\
& + \frac{\beta^2}{\gamma^2 - \nu^2} \left[(a-b)^2\gamma^2 + \nu((1-a)(a-3+2b^2) - (1-b)(1+b-2ab)\nu) + (1-a)\gamma(1+a-2b^2 + (3-a-2b^2)\nu) \right], \tag{C11}
\end{aligned}$$

$$f_{12} = f_1 + \frac{c_3(f_1 - f_3)}{(a+1)b(1-\gamma)(\nu+1)}, \tag{C12}$$

$$f_{13} = f_7 + \frac{(b(\gamma - \nu^2) - (2+b)(1-\gamma)\nu)f_{10}}{2b(1+\nu)d_1}, \tag{C13}$$

$$\begin{aligned}
f_{14} = & 1 - \frac{2\beta}{d_2} \left[b(-1 + a^2 + \gamma - 2a^2\gamma - (1 - 2a)b\gamma + (a - b)^2\gamma^2) \right. \\
& + (-2 + (2 + a^2)b + (1 - 2a)b^2 - 2(-1 + b + a^2b - 2ab^2 + b^3)\gamma + b(-a + b)^2\gamma^2)\nu \\
& \left. + (2 - b - 2ab^2 + b^3 + (-2 + b + (1 + 2a)b^2 - 2b^3)\gamma)\nu^2 + (-1 + b)b^2\nu^3 \right] \\
& + \frac{\beta^2}{(\gamma^2 - \nu^2)d_2} \left[b\gamma(2(1 - a)^2(1 + a) + (1 - a)(-2 + a + 5a^2 - 4ab)\gamma \right. \\
& + (a(-1 - 2a + 4a^2) + (1 + 4a - 7a^2)b + (-2 + 3a)b^2)\gamma^2 - (a - b)^3\gamma^3) \\
& + (-2(1 - a)^2(1 + a)(2 + b) + 2(1 - a)^2(5 + 6a + 4ab - 2b^2)\gamma \\
& + (-6 + 4a + 16a^2 - 12a^3 + (2 - 22a + 26a^2 - 11a^3)b + (9 - 16a + 11a^2)b^2 + (2 - 3a)b^3)\gamma^2 \\
& - 2((3 - 2a)a^2 - 3a(2 - 2a + a^2)b + (3 - 4a + 5a^2)b^2 - 3ab^3 + b^4)\gamma^3 + b(-a + b)^3\gamma^4)\nu \\
& + ((1 - a)(2 - 6a + 4a^2 + (2 - 9a + 3a^2)b + 4b^2) \\
& + (6(1 - a)^2(-1 + 2a) + (-12 + 37a - 34a^2 + 8a^3)b + (-13 + 20a - 5a^2)b^2 + (2 - 3a)b^3)\gamma \\
& + (4 - 24a + 30a^2 - 12a^3 + (18 - 39a + 32a^2 - 7a^3)b + 2(5 - 10a + 4a^2)b^2)\gamma^2 \\
& + (2a(4 - 5a + 2a^2) + (-8 + 13a - 10a^2 + 2a^3)b + (-3 + 4a - 3a^2)b^2 + (2 + 3a)b^3 - 2b^4)\gamma^3)\nu^2 \\
& + (2(a - 2)a - (-10 + 14a - 10a^2 + a^3)b + (3 - 8a + a^2)b^2 + (-2 + 3a)b^3 \\
& + 2(-1 + 4a - 2a^2 + (-8 + 14a - 10a^2 + a^3)b - (3 - 8a + a^2)b^2 - 3ab^3 + b^4)\gamma \\
& + (2(-1 + a)^2 - (-6 + 14a - 10a^2 + a^3)b + (5 - 8a + a^2)b^2 + (-2 + 3a)b^3)\gamma^2)\nu^3 - (1 - b)^2b^2\nu^5 \\
& \left. + (2 - (2 + a)b + (2 - 4a + a^2)b^2 + 3ab^3 - b^4 + (-2 + (2 + a)b - (1 - 4a + a^2)b^2 - (2 + 3a)b^3 + 2b^4)\gamma)\nu^4 \right], \tag{C14}
\end{aligned}$$

$$\begin{aligned}
f_{15} = & 1 + \frac{2\beta(1 - a^2 + (a - b)^2\gamma - (1 - 2ab + b^2)\nu)}{(1 - ab)(\gamma - \nu)} \\
& - \frac{\beta^2}{(1 - ab)(\gamma^2 - \nu^2)} \left[-2(1 - a)^2(1 + a)(1 + b) - (a - b)^2(-1 + 2a + (-2 + a)b)\gamma^2 \right. \\
& + (1 - a)(1 - 5a + (6 - 3a + a^2)b - 2(-1 + a)b^2)\nu - (1 - b)(-1 + (-1 + 3a)b + (-2 + a)b^2)\nu^2 \\
& \left. - (1 - a)\gamma(-1 + a + 4a^2 + (-2 - 5a + 3a^2)b + 2(1 - a)b^2 + (1 - 5a + (6 - 3a + a^2)b - 2(-1 + a)b^2)\nu) \right], \tag{C15}
\end{aligned}$$

$$\begin{aligned}
f_{16} = & 1 + 2(1 - ab)\beta - \frac{\beta^2}{(1 - \nu)(\gamma^2 - \nu^2)} \left[2(1 - a)^2(1 + a) + (1 - a)^2(-1 + 2a(1 + b))\nu \right. \\
& - ((-2 + a)a - 2(-2 + 4a - 4a^2 + a^3)b + (3 - 6a + 2a^2)b^2)\nu^2 - (1 - 2ab + (1 - 2a + 2a^2)b^2)\nu^3 \\
& + (a - b)\gamma^2(a(-3 + 2a) + (1 - 2a + 2a^2)b + (4 - 5a + 2a^2 + (3 - 6a + 2a^2)b)\nu) \\
& \left. + (1 - a)^2\gamma(-3 - 2a(2 + b) + (2 - 4a + 4(1 - a)b + 4b^2)\nu + (1 - 2(-2 + a)b + 4b^2)\nu^2) \right], \tag{C16}
\end{aligned}$$

$$\begin{aligned}
f_{17} = & 1 - \frac{2\beta}{(1-ab)(1-\gamma)(\gamma-\nu)(1+\nu)} \left[-1 + a^2 + (2 + a^2 + a(-2 - 2b + b^2))\nu + (-1 + b)(1 - 2a + b)\nu^2 \right. \\
& + (1-a)b^2\nu^3 + (a-b)^2\gamma^2(1+\nu) \\
& \left. + \gamma(1 - 2a^2 + 2ab - ab^2 - 2(1-a+a^2-2ab+b^2)\nu + (1-2a+2ab+(-2+a)b^2)\nu^2) \right] \\
& - \frac{\beta^2}{(1-ab)(1-\gamma)(1+\nu)(\gamma^2-\nu^2)} \left[(1-a)(3-5a-2a^2+(4-3a+3a^2)b-2(1-a)ab^2)\nu \right. \\
& + (2+2a-7a^2+4a^3-(2+a^3)b+(1-4a+10a^2-4a^3)b^2+(2-5a+2a^2)b^3)\nu^2 - 2(1-a)^2(1+a)(1+b) \\
& + (-3+4a+(a-4a^2)b+(3-4a+6a^2-2a^3)b^2+(2-7a+4a^2)b^3)\nu^3 \\
& + 2(1-a)b^2(1-ab)\nu^4 + (a-b)^2(-1+2a+(-2+a)b)\gamma^3(1+\nu) \\
& + \gamma^2 \left\{ -1 + 2a + 4a^2 - 6a^3 - (2 + 5a - 14a^2 + 4a^3)b + (3 - 10a + 6a^2 - 2a^3)b^2 + (2 - 3a + 2a^2)b^3 \right. \\
& + (a(-4 + 5a - 2a^2) + (4 - 10a + 14a^2 - 5a^3)b + (5 - 18a + 14a^2 - 4a^3)b^2 + (6 - 9a + 4a^2)b^3)\nu \\
& \left. + (1-a)(1+3a-4a^2+(-2+a+a^2)b+2(1-3a+a^2)b^2-2(-2+a)b^3)\nu^2 \right\} \\
& - \gamma \left\{ (1-a)(-3+a+6a^2+(-4-5a+5a^2)b+2(-1+a)ab^2) \right. \\
& - (1-a)(-3+9a+2a^2+(-8+7a-7a^2)b-2(2-5a+3a^2)b^2+4(-1+a)b^3)\nu \\
& + (3+4a-14a^2+8a^3+(-4+3a-2a^3)b+(7-16a+18a^2-6a^3)b^2+(6-15a+8a^2)b^3)\nu^2 \\
& \left. + (-3+4a+(a-4a^2)b+(3-4a+6a^2-2a^3)b^2+(2-7a+4a^2)b^3)\nu^3 \right\} \Big], \tag{C17}
\end{aligned}$$

$$\begin{aligned}
f_{18} = & 1 - \frac{2\beta(-1+a^2+(a-b)^2\gamma^2+(1-2a+b)\nu+(1-b)b\nu^2-\gamma(-1+2a^2+b-2ab+(1-2a+b)\nu))}{(\gamma+\nu)c_1} \\
& + \frac{\beta^2}{(\gamma^2-\nu^2)c_1} \left[2(1-a)^2(1+a) - (a-b)^3\gamma^3 - (1-a)(4+a^2-a(1+4b))\nu \right. \\
& + (2-3a-(1-4a+a^2)b+(-2+a)b^2)\nu^2 + (1-b)^2b\nu^3 \\
& + (a-b)\gamma^2(-1-2a+4a^2+(2-3a)b+((-1+a)^2+(2-3a)b+b^2)\nu) \\
& + \gamma((1-a)(-2+a+5a^2-4ab)+2(1-a)(2-a+a^2-4ab+2b^2)\nu) \\
& \left. - (2-3a+(4-a)ab-(4-a)b^2+b^3)\gamma\nu^2 \right], \tag{C18}
\end{aligned}$$

$$f_{19} = 1 + \frac{2\beta(1-a+(a-b)\gamma-(1-b)\nu)}{\gamma-\nu} + \frac{\beta^2(1-a^2-2b(1-a(1-\gamma))+a^2\gamma+b^2\gamma-(1-b)^2\nu)}{\gamma-\nu}, \tag{C19}$$

$$\begin{aligned}
f_{20} = & 1 - \frac{2\beta}{(\gamma - \nu)(c_3 - (1 - ab)(1 - \gamma)(1 + \nu))} \left[1 - a^2 - (2 + a^2 + a(-2 - 2b + b^2))\nu \right. \\
& + (1 - 2(1 - a)b - (1 + a)b^2 + b^3)\nu^2 - (1 - b)b^2\nu^3 - (a - b)\gamma^2(a - b + b^2 + (-2 + a - b + b^2)\nu) \\
& \left. + \gamma(-1 + 2a^2 - 2ab + ab^2 + 2((-1 + a)^2 + b - 2ab + (1 + a)b^2 - b^3)\nu + (-1 - 2(-1 + a)b + (2 + a)b^2 - 2b^3)\nu^2) \right] \\
& + \frac{\beta^2}{(\gamma^2 - \nu^2)(c_3 - (1 - ab)(1 - \gamma)(1 + \nu))} \left[-2(1 - a)^2(1 + a)(1 + b) \right. \\
& - (1 - a)(1 + 5a - 2a^2 + (-4 + 3a - 3a^2)b - 2(1 - a)b^2)\nu \\
& + (4 - 6a + 3a^2 - (-2 + 8a - 4a^2 + a^3)b + (2 - 2a + 3a^2)b^2 - ab^3)\nu^2 \\
& + (-1 + (4 - 3a)b + (-1 + 2a + a^2)b^2 - (2 + a)b^3 + b^4)\nu^3 + (-1 + b)^2b^2\nu^4 \\
& + (a - b)^2\gamma^3(-1 + 2a + (-2 + a)b + b^2 + (-3 + 2a + (-2 + a)b + b^2)\nu) \\
& + \gamma^2(-1 + 2a + 4a^2 - 6a^3 - (2 + 5a - 14a^2 + 4a^3)b + (4 - 10a + 3a^2)b^2 + ab^3 \\
& \quad + (-2 - 4a + 13a^2 - 6a^3 + (8 - 22a + 18a^2 - 5a^3)b + (7 - 14a + 5a^2)b^2 + (2 + a)b^3 - b^4)\nu \\
& \quad + (3(-1 + a)^2 - (-2 + a)(-1 + a)^2b + (1 - 4a + 2a^2)b^2 + 2b^3 - b^4)\nu^2) \\
& + \gamma((-1 + a)(-3 + a + 6a^2 + (-4 - 5a + 5a^2)b - 2(-1 + a)b^2) \\
& \quad + (-1 + a)(-3 - 11a + 6a^2 + (12 - 11a + 7a^2)b - 6(-1 + a)b^2)\nu \\
& \quad - (7 - 12a + 6a^2 + (4 - 13a + 8a^2 - 2a^3)b + (5 - 6a + 5a^2)b^2 - (2 + a)b^3 + b^4)\nu^2 \\
& \quad \left. + (1 + (-4 + 3a)b - (-1 + 2a + a^2)b^2 + (2 + a)b^3 - b^4)\nu^3) \right], \tag{C20}
\end{aligned}$$

$$\begin{aligned}
f_{21} = & 1 + \frac{2\beta(1 - a^2 + a(a - b)\gamma + (2 - a)(a - b)(1 - \gamma)\nu - (1 - ab)\nu^2)}{(1 - \nu)(\gamma + \nu)} \\
& + \frac{\beta^2}{(\gamma^2 - \nu^2)(1 - \nu)} \left[2(1 - a)^2(1 + a) - (1 - a)(3 + 2a^2 - a(1 + 4b))\nu + ((a - 2)a + 2(3 - 2a)ab - (3 - 2a)b^2)\nu^2 \right. \\
& + (1 - b)(1 + (1 - 2a)b)\nu^3 + (a - b)^2\gamma^2(-1 + 2a + (-3 + 2a)\nu) \\
& \left. + (1 - a)\gamma(-1 + a + 4a^2 - 4ab + (2 - 2a + 4a^2 - 8ab + 4b^2)\nu + (-1 + a - 4ab + 4b^2)\nu^2) \right], \tag{C21}
\end{aligned}$$

$$\begin{aligned}
f_{22} = & 1 + \frac{2\beta(1 - a + (a - b)\gamma + (1 - b)\nu)}{\gamma + \nu} \\
& + \frac{\beta^2((a - b)^2\gamma^2 + \nu((1 - a)(1 - 3a + 2b) - (1 - b)^2\nu) + (1 - a)\gamma(1 + a - 2b - \nu + 3a\nu - 2b\nu))}{\gamma^2 - \nu^2}, \tag{C22}
\end{aligned}$$

2. g_1, \dots, g_{13}

$$g_1 = f_{13}f_{14} - \frac{c_1c_3(f_{13}f_{14} - f_{18}f_{19})}{2(a-b)(1-\gamma)^2\nu} + \frac{(1-\nu)c_3f_3f_{10}}{2(a-b)(1-\gamma)(1+\nu)d_1}, \quad (\text{C23})$$

$$g_2 = D_0 + \frac{c_3(D_0 - f_3f_4)}{d_3}, \quad (\text{C24})$$

$$g_3 = f_{14}f_{21} - \frac{d_1(f_{14}f_{21} - f_2f_{22})}{(1-\gamma)(1+\nu)(ac_1 - 1 + \nu)} - \frac{b\nu(\gamma - \nu)f_{10}f_{12}}{(1-\gamma)(1-\nu^2)(ac_1 - 1 + \nu)}, \quad (\text{C25})$$

$$g_4 = 1 - \frac{4\beta}{d_4}\mathcal{G}_4^{(1)} + \frac{4\beta^3}{(\gamma^2 - \nu^2)d_4} \sum_{i=1}^2 \mathcal{G}_4^{(2,i)} + \frac{6\beta^2}{(\gamma - \nu)d_4} \sum_{i=1}^2 \mathcal{G}_4^{(3,i)} + \frac{\beta^4}{(\gamma^2 - \nu^2)^2d_4} \sum_{i=1}^6 \mathcal{G}_4^{(4,i)}, \quad (\text{C26})$$

$$g_5 = \frac{c_1c_3f_3f_6 - bd_1f_1f_7}{c_1c_3 - bd_1}, \quad (\text{C27})$$

$$g_6 = f_8^2 + \frac{2(1-\gamma)^2(1-\nu)^2\nu d_1(f_1^2 - f_8^2)}{(1+\nu)d_5} + \frac{(1-\gamma)^2(1-\nu)^4c_3(f_3^2 - f_8^2)}{(1+\nu)d_5}, \quad (\text{C28})$$

$$g_7 = g_8 + \frac{16(1-a)^2(1-\gamma)^2(1-\nu)^2\nu c_3d_1}{(\gamma^2 - \nu^2)(1+\nu)d_6} \beta^2(1 + (a-b)\beta)^2, \quad (\text{C29})$$

$$g_8 = f_{14}^2 - \frac{2(1-\gamma)(1-\nu)^2\nu d_1(f_{14}^2 - f_2^2)}{d_6} + \frac{(\gamma - \nu)(1-\nu)^2(1+\nu)c_1^2c_3(f_{14}^2 - f_6^2)}{2d_6\nu}, \quad (\text{C30})$$

$$g_9 = \frac{c_1c_3f_4f_{18} - bd_1f_2f_{13}}{c_1c_3 - bd_1}, \quad (\text{C31})$$

$$g_{10} = f_{14}f_{21} - \frac{bf_{10}f_{12}(\nu + 1)(\gamma - \nu)}{2d_2(\nu - 1)}, \quad (\text{C32})$$

$$g_{11} = f_{18}f_{20} - \frac{(1-\gamma)(4 + \nu - \nu^2)f_1f_{10}}{(1+\nu)(2(a-b)(1-\gamma)^2(1-\nu)\nu - b(1+\nu)d_1)} - \frac{(a-b)(1-\gamma)^2(1-\nu)^2(f_{21}f_8 - f_{18}f_{20})}{2(a-b)(1-\gamma)^2(1-\nu)\nu - b(1+\nu)d_1}, \quad (\text{C33})$$

$$g_{12} = f_{21}f_8 + \frac{(2+\nu)f_1f_{10}}{(a-b)(1-\gamma)(1-\nu^2)}, \quad (\text{C34})$$

$$g_{13} = \frac{(1-ab)(\gamma - \nu)(1+\nu)c_1f_6f_{15} - (1-\nu)c_2f_5f_{16}}{(1-ab)(\gamma - \nu)(1+\nu)c_1 - (1-\nu)c_2}, \quad (\text{C35})$$

where lengthy coefficients in g_4 are shown below

$$\begin{aligned}
\mathcal{G}_4^{(1)} = & b^3\gamma(-1+a^2 - (-1+a+2a^2-2ab)\gamma + (a+a^2-b-2ab+b^2)\gamma^2) \\
& + b(- (1-a^2)(2-b^2) + (2(2-a-3a^2) + (1+3a-3a^2+3a^3)b + (1+a)^2b^2 - 4ab^3)\gamma \\
& - (2-4a-6a^2 + (3+6a-6a^2+9a^3)b + (1+7a-10a^2)b^2 - (3+2a)b^3 + 3b^4)\gamma^2 \\
& - (2a(1+a) - (2+3a-3a^2+9a^3)b + (1-5a+19a^2)b^2 + (2-11a)b^3 + b^4)\gamma^3 + 3b(-a+b)^3\gamma^4)\nu \\
& + (4a(1+a-2a^2(1-\gamma))(-1+\gamma)^3 - 2b(1-\gamma)^2(-4-2a+3a^2+a^3 - (-2-2a+15a^2+a^3)\gamma + 12a^2\gamma^2) \\
& + b^2(1-\gamma)(-1-3a+3a^2-3a^3+2(1-5a-4a^2+6a^3)\gamma + a(37+5a-15a^2)\gamma^2 + 6a(-4+a^2)\gamma^3) \\
& + 3b^5\gamma(1+\gamma-5\gamma^2+2\gamma^3) + b^3(-1+\gamma)(2+a-a^2+2a(-5+11a)\gamma + (7+3a-39a^2)\gamma^2 + 2(-4+9a^2)\gamma^3) \\
& + b^4(2a+(-3+2a)\gamma + (6-37a)\gamma^2 + 51a\gamma^3 - 18a\gamma^4))\nu^2 \\
& + (4a(2+a+a^2(1-\gamma))(1-\gamma)^3 - 2b(1-\gamma)^2(4+a+11a^2 - (4+3a+17a^2)\gamma + 6a^2\gamma^2) \\
& + b^2(1-\gamma)(a(13+5a-6a^2) + a(-52-2a+15a^2)\gamma + (2+51a-3a^2-12a^3)\gamma^2 + 3a(-4+a^2)\gamma^3) \\
& - b^3(1-\gamma)(1+5a-12a^2 + (-6-6a+47a^2)\gamma + (17-5a-44a^2)\gamma^2 + (-4+9a^2)\gamma^3) \\
& + b^5(-1-3\gamma+27\gamma^2-23\gamma^3+3\gamma^4) + b^4(1-2a+(-6+41a)\gamma - 91a\gamma^2 + (2+61a)\gamma^3 - 9a\gamma^4))\nu^3 \\
& + (-4a+2(-5+a)a^2b - 3a(-5+a+a^2)b^2 + (1-3a+17a^2)b^3 + (2-15a)b^4 + b^5 \\
& + (12a+(-4+30a^2-6a^3)b + (-2-53a+11a^2+9a^3)b^2 + (16-7a-50a^2)b^3 + 65ab^4 - 21b^5)\gamma \\
& + (-12a+(8-30a^2+6a^3)b + (2+61a-13a^2-9a^3)b^2 + (-30+14a+49a^2)b^3 - (6+73a)b^4 + 33b^5)\gamma^2 \\
& + (4a-2(2-5a^2+a^3)b + a(-23+5a+3a^2)b^2 + (13-4a-16a^2)b^3 + (1+23a)b^4 - 10b^5)\gamma^3)\nu^4 \\
& + b(2-2a+(11-5a)ab + (-8+5a+7a^2)b^2 - 16ab^3 + 6b^4 \\
& - (4-4a+(1+22a-10a^2)b + (-20+13a+14a^2)b^2 - (6+35a)b^3 + 21b^4)\gamma \\
& + (2-2a+(1+11a-5a^2)b + (-12+8a+7a^2)b^2 - (3+19a)b^3 + 12b^4)\gamma^2)\nu^5 \\
& - b^2(-1+(3-4a)b + (2+5a)b^2 - 5b^3 + (1+(-3+4a)b - (3+5a)b^2 + 6b^3)\gamma)\nu^6 - (1-b)b^4\nu^7, \tag{C36}
\end{aligned}$$

$$\begin{aligned}
\mathcal{G}_4^{(2,1)} = & b^2\gamma (2(1-a)^3(1+a)^2 + (1-a)^2(1+a)(-4+2a+8a^2+b-7ab)\gamma \\
& + (1-a)(2(1-2a-6a^2+3a^3+6a^4) + (1+16a-2a^2-21a^3)b + (-4-4a+8a^2)b^2 + (1+a)b^3)\gamma^2 \\
& - (-2a(1+a-5a^2-a^3+4a^4) + (2+7a-18a^2-14a^3+21a^4)b \\
& + (-5+6a+21a^2-16a^3)b^2 + (2-9a+a^2)b^3 + 2ab^4)\gamma^3 + (a-b)^3(2-2a^2+(-3+a)b+b^2)\gamma^4) \\
- & b(2(1-a)^3(1+a)^2(2+b) - 2(1-a)^3(1+a)(9+8a+3(-1+a)b-4b^2)\gamma \\
& + (1-a)(24-10a-58a^2+8a^3+24a^4 + (-13+56a+16a^2-38a^3+7a^4)b \\
& - (23+4a-32a^2+23a^3)b^2 + 12(-1+a)ab^3 + (1+a)b^4)\gamma^2 \\
& - (10-24a-32a^2+62a^3+4a^4-16a^5 + (1+76a-78a^2-50a^3+47a^4-8a^5)b \\
& + (-34+17a+66a^2-77a^3+36a^4)b^2 + (7-25a+63a^2-37a^3)b^3 + (1-18a+5a^2)b^4 + 4ab^5)\gamma^3 \\
& - (2a(3+4a-15a^2+2a^3+2a^4) + (-6-27a+64a^2+6a^3-21a^4+12a^5)b + (3+39a-31a^2)b^4 \\
& + (19-32a-24a^2+53a^3-50a^4)b^2 + (-2+11a-66a^2+68a^3)b^3 - (9+2a)b^5 + 3b^6)\gamma^4 \\
& + (a-b)^3(2(a-3) + a(10a-3)b + (6-14a)b^2 + b^3)\gamma^5 - 3b(a-b)^5\gamma^6)\nu \\
- & (b^7\gamma^3(2+3\gamma-15\gamma^2+6\gamma^3) + b^6\gamma^3(-6+4a+(18-55a)\gamma+81a\gamma^2-30a\gamma^3) \\
& + 4a(-1+\gamma)^3(2(1-a)^3(1+a) + (1-a)^2(1+5a+6a^2)\gamma + 3a^2(1+a-2a^2)\gamma^2 + 2a^4\gamma^3) \\
& - 2b(1-\gamma)^2((1-a)^3(7+13a+6a^2) + (-1+a)^2(-15-19a-5a^2+15a^3)\gamma \\
& + (2+a-24a^2-44a^3+75a^4-12a^5)\gamma^2 + a^2(18+24a-65a^2+3a^3)\gamma^3 + 20a^4\gamma^4) \\
& - b^3(1-\gamma)((1-a)^2(7+6a-a^2) - 3(-2+13a-20a^2+7a^3+2a^4)\gamma \\
& + (-49+36a+12a^2-26a^3+15a^4)\gamma^2 + 2(24+27a-94a^2+24a^3+11a^4)\gamma^3 \\
& - (12+48a-202a^2+9a^3+60a^4)\gamma^4 + 10a^2(-8+3a^2)\gamma^5) \\
& + b^5\gamma((1-a^2)(1-3\gamma) - (2-42a+54a^2)\gamma^2 - (12+45a-163a^2)\gamma^3 + 3(4+a-57a^2)\gamma^4 - (8-60a^2)\gamma^5) \\
& + b^4\gamma(-12(-1+a)^2 + (21-49a+33a^2-9a^3)\gamma + (7-23a-33a^2+75a^3)\gamma^2 \\
& + (-24+110a+15a^2-183a^3)\gamma^3 + 3(4-34a-a^2+59a^3)\gamma^4 + (40a-60a^3)\gamma^5) \\
& - b^2(-1+\gamma)(2(-1+a)^2(-5+2a+8a^2+a^3) - 2(-1+7a^2-6a^3-6a^4+6a^5)\gamma \\
& + (-1-84a-38a^2+168a^3-75a^4+18a^5)\gamma^2 - 2(-5-48a-63a^2+174a^3-33a^4+a^5)\gamma^3 \\
& - a(36+72a-274a^2+15a^3+12a^4)\gamma^4 + (-80a^3+6a^5)\gamma^5)\nu^2 \\
- & (-4a(1-\gamma)^3((1-a)^2(-1+9a+4a^2) - (1-a)^2(5+18a+7a^2)\gamma + a^2(-9+3a+2a^2)\gamma^2 + a^4\gamma^3) \\
& + b^7\gamma^2(2-2\gamma+24\gamma^2-23\gamma^3+3\gamma^4) \\
& - 2b(1-\gamma)^2(-15+8a+30a^2-33a^3+13a^4-3a^5 + (22+12a-90a^2+59a^3-11a^4+6a^5)\gamma \\
& - (10+21a-114a^2+41a^3+7a^4+3a^5)\gamma^2 + a^2(-54+15a-5a^2)\gamma^3 + 10a^4\gamma^4) \\
& + b^6\gamma(4a-2(3+2a)\gamma+6(-2+7a)\gamma^2-130a\gamma^3+(6+103a)\gamma^4-15a\gamma^5) \\
& + b^2(1-\gamma)(3-51a+44a^2+34a^3-39a^4+9a^5 + (-14+39a+58a^2-196a^3+126a^4-25a^5)\gamma \\
& + 2(2+75a-121a^2+158a^3-69a^4+9a^5)\gamma^2 + 2(5-123a+79a^2-110a^3+27a^4+3a^5)\gamma^3 \\
& + a(108-18a+106a^2-3a^3-11a^4)\gamma^4 + a^3(-40+3a^2)\gamma^5) \\
& - b^3(1-\gamma)(-23+35a-12a^2-7a^3+7a^4 + (35-30a-78a^2+118a^3-53a^4)\gamma \\
& + 2(33-81a+152a^2-93a^3+27a^4)\gamma^2 + 2(-39+62a-156a^2+45a^3+19a^4)\gamma^3 \\
& + (36+6a+138a^2-15a^3-61a^4)\gamma^4 + 5a^2(-8+3a^2)\gamma^5) \\
& - b^4(-4(2-3a+a^3) + (1+a-39a^2+29a^3)\gamma + (27-105a+129a^2-61a^3)\gamma^2 \\
& - 2(27-113a+66a^2+28a^3)\gamma^3 + (32-212a+69a^2+229a^3)\gamma^4 \\
& + (6+98a-27a^2-167a^3)\gamma^5 + 10a(-2+3a^2)\gamma^6) \\
& + b^5(-1+a^2+(1-18a+5a^2)\gamma+(4+18a-4a^2)\gamma^2+(14+6a-100a^2)\gamma^3+(-22+15a+253a^2)\gamma^4 \\
& + (20-21a-185a^2)\gamma^5+(-4+30a^2)\gamma^6)\nu^3,
\end{aligned} \tag{C37}$$

$$\begin{aligned}
\mathcal{G}_4^{(2;2)} = & -(-4a(1-\gamma)^3(-7-4a+21a^2-11a^3+a^4+(7+3a-27a^2+17a^3-2a^4)\gamma+a^2(6-6a+a^2)\gamma^2) \\
& +2b(1-\gamma)^2(-16+a-6a^2-2a^3+27a^4-6a^5+(18+18a-48a^2+42a^3-65a^4+15a^5)\gamma \\
& - (14+11a-90a^2+82a^3-49a^4+12a^5)\gamma^2+a^2(-36+42a-11a^2+3a^3)\gamma^3) \\
& -b^7\gamma(3+2\gamma+6\gamma^2-27\gamma^3+10\gamma^4)+b^3(1-\gamma)(-33+20a-30a^2+57a^3-24a^4 \\
& +2(48-69a+50a^2-72a^3+35a^4)\gamma+(-47+72a+26a^2+56a^3-45a^4)\gamma^2 \\
& +(-22+85a-88a^2+37a^3-24a^4)\gamma^3+(24-60a-8a^2-6a^3+23a^4)\gamma^4) \\
& +b^6(-2a+(9-2a)\gamma+2(-6+17a)\gamma^2+26a\gamma^3-(18+97a)\gamma^4+(3+41a)\gamma^5) \\
& +b^5(1+9a-4a^2+a(-51+56a)\gamma+(10+42a-116a^2)\gamma^2+(11-42a-5a^2)\gamma^3 \\
& +2(-9+27a+68a^2)\gamma^4+(14-12a-67a^2)\gamma^5) \\
& +b^4(-8+19a-30a^2+17a^3+(25-67a+138a^2-115a^3)\gamma+2(-21+41a-90a^2+94a^3)\gamma^2 \\
& +(2-26a+96a^2-44a^3)\gamma^3+(29+24a-36a^2-101a^3)\gamma^4+(-12-32a+12a^2+55a^3)\gamma^5) \\
& +b^2(21+72a-66a^2+14a^3-33a^4+4a^5+(-13-290a+208a^2+82a^3+65a^4-12a^5)\gamma \\
& +2(-15+159a-63a^2-179a^3+9a^4+4a^5)\gamma^2+2(6+9a-108a^2+221a^3-57a^4+4a^5)\gamma^3 \\
& +(10-190a+308a^2-208a^3+79a^4-12a^5)\gamma^4+a(72-108a+28a^2-15a^3+4a^4)\gamma^5))\nu^4 \\
& - (4a(1-\gamma)^3(-9+5a+6a^2-6a^3+(3-2a-6a^2+6a^3)\gamma) \\
& -2b(1-\gamma)^2(-2-19a-28a^2+35a^3-19a^4+a^5+(5+18a+32a^2-47a^3+38a^4-2a^5)\gamma \\
& +(-6+6a-4a^2+12a^3-19a^4+a^5)\gamma^2) \\
& +b^7(1+3\gamma-21\gamma^2+2\gamma^3+9\gamma^4)+b^6(-3+2a+(18-59a)\gamma+117a\gamma^2+(12-38a)\gamma^3-(9+22a)\gamma^4) \\
& +b^5(-1+21a-27a^2-(14+27a-155a^2)\gamma+(9+6a-249a^2)\gamma^2-(5+24a-105a^2)\gamma^3-(7-24a-16a^2)\gamma^4) \\
& +b^4(-7+2a-33a^2+39a^3-3(4-28a-13a^2+59a^3)\gamma+(42-206a+60a^2+234a^3)\gamma^2 \\
& -(47-161a+69a^2+93a^3)\gamma^3+(30-41a+3a^2-3a^3)\gamma^4) \\
& +b^3(18+12a+22a^2+21a^3-23a^4+(-56-238a^2+53a^3+68a^4)\gamma+(119-177a+564a^2-221a^3-66a^4)\gamma^2 \\
& +(-106+279a-502a^2+199a^3+20a^4)\gamma^3+(25-114a+154a^2-52a^3+a^4)\gamma^4) \\
& -b^2(22+28a+8a^2+70a^3-37a^4+a^5-(44+198a-126a^2+354a^3-147a^4+4a^5)\gamma \\
& +3(7+123a-136a^2+214a^3-73a^4+2a^5)\gamma^2+(5-256a+406a^2-502a^3+145a^4-4a^5)\gamma^3 \\
& +(-4+57a-132a^2+144a^3-36a^4+a^5)\gamma^4))\nu^5 \\
& - (4a(3-2a)(1-\gamma)^3-2b(-1+\gamma)^2(-7+21a-14a^3+\gamma-17a\gamma+14a^3\gamma) \\
& +b^7(-1+21\gamma-28\gamma^2+4\gamma^3)+b^6(-6+21a-107a\gamma+4(3+29a)\gamma^2+(6-30a)\gamma^3) \\
& +b^5(2+3a-47a^2+(-23+39a+176a^2)\gamma+(25-42a-187a^2)\gamma^2+2(-8+29a^2)\gamma^3) \\
& +b^4(18-48a+21a^2+43a^3-(22-152a+117a^2+125a^3)\gamma+(21-193a+147a^2+121a^3)\gamma^2 \\
& -(13-89a+51a^2+39a^3)\gamma^3) \\
& +b^2(-10+12a+32a^2-76a^3+21a^4+(2-12a-92a^2+228a^3-63a^4)\gamma \\
& +(29-12a+88a^2-228a^3+63a^4)\gamma^2+(-21+12a-28a^2+76a^3-21a^4)\gamma^3) \\
& +b^3(-5-8a+80a^2-49a^3-7a^4+(-13+83a-312a^2+172a^3+21a^4)\gamma \\
& +(17-119a+384a^2-197a^3-21a^4)\gamma^2+(1+44a-152a^2+74a^3+7a^4)\gamma^3))\nu^6 \\
& - b(6(a-1)+(17-27a+4a^2)b-(5-11a-30a^2+28a^3)b^2-(2+30a-33a^2-17a^3)b^3+(13-21a-34a^2)b^4 \\
& +28ab^5-6b^6+(12(1-a)+(-33+54a-8a^2)b+(-2-13a-60a^2+56a^3)b^2+(13+57a-78a^2-34a^3)b^3 \\
& +5(-3+9a+16a^2)b^4-(18+65a)b^5+21b^6)\gamma \\
& +(-6(1-a)+(16-27a+4a^2)b+(7+2a+30a^2-28a^3)b^2+(-15-27a+45a^2+17a^3)b^3 \\
& -2(-7+12a+23a^2)b^4+(6+37a)b^5-11b^6)\gamma^2)\nu^7 \\
& + (1-b)b^2(1-8(1-a)b+(2-a-9a^2)b^2+(1+11a)b^3-5b^4 \\
& -(1-8(1-a)b+(1-a-9a^2)b^2+(3+11a)b^3-6b^4)\gamma)\nu^8-(1-b)^3b^4\nu^9, \tag{C38}
\end{aligned}$$

$$\begin{aligned}
\mathcal{G}_4^{(3,1)} &= (a-b)^4 \gamma^5 \nu (3b^2 + (-8+6b^2)\nu + (-4+3b^2)\nu^2) \\
&- (a-b)^2 \gamma^4 (b^2(-1+a^2+2b-b^2) + b(-4+(1-4a+10a^2)b + (4-12a)b^2 + b^3)\nu \\
&\quad + (8(1+a-5a^2) + (-4+52a)b + (-13-4a+23a^2)b^2 - 42ab^3 + 18b^4)\nu^2 \\
&\quad + (-4(5+4a^2) + (4+56a)b + (-29+4a+20a^2)b^2 - 4(1+12a)b^3 + 29b^4)\nu^3 \\
&\quad + (-4(-3+2a+a^2) + 4(1+3a)b + 2(-9+2a+3a^2)b^2 - 2(1+9a)b^3 + 13b^4)\nu^4) \\
&+ \gamma^3 (b^2(1-4a^2+3a^4 + (-2+4a+4a^2-4a^3)b - (-1+4a+a^2)b^2 + 2ab^3) \\
&\quad + b(4-12a^2 + (-9+20a+8a^2-12a^3+13a^4)b - 2(1+3a-14a^2+16a^3)b^2 \\
&\quad \quad + (-3-24a+19a^2)b^3 + 4(2+a)b^4 - 4b^5)\nu \\
&\quad + 2(-4a(1-4a-4a^2+10a^3) + 2(-2-11a-16a^2+51a^3)b + (19+20a-90a^2-8a^3+17a^4)b^2 \\
&\quad \quad + (-6+28a+20a^2-60a^3)b^3 + (-3-20a+71a^2)b^4 + (8-30a)b^5 + 2b^6)\nu^2 \\
&\quad + 2(-6a(-2+7a+2a^3) + 2a(25-2a+51a^2)b + (-25+8a-176a^2+4a^3+22a^4)b^2 \\
&\quad \quad - 2(3-55a+2a^2+46a^3)b^3 + (-17+141a^2)b^4 - 92ab^5 + 21b^6)\nu^3 \\
&\quad + (-8a(3-7a+4a^2+2a^3) + 4(2-9a+16a^2+17a^3)b + (1-40a-176a^2+16a^3+23a^4)b^2 \\
&\quad \quad - 2(-7-82a+22a^2+58a^3)b^3 + (-51+44a+219a^2)b^4 - 2(8+91a)b^5 + 56b^6)\nu^4 \\
&\quad + (-4(-2+a)a + 4(-1-5a+5a^2+a^3)b + (19-36a-16a^2+4a^3+3a^4)b^2 + (14+42a-20a^2-24a^3)b^3 \\
&\quad \quad + (-27+24a+59a^2)b^4 - 4(2+15a)b^5 + 22b^6)\nu^5) \\
&+ \gamma^2 ((-1+a^2)b^2(2-3a^2+2(-1+a)b+b^2) \\
&\quad + b(-8+12a^2+(16-16a-11a^2+12a^3-9a^4)b+2(1-10a^2+8a^3)b^2+12ab^3-6ab^4)\nu \\
&\quad + (8a(3-6a-6a^2+10a^3) + (12+40a+72a^2-140a^3)b + (-42-36a+74a^2+24a^3-24a^4)b^2 \\
&\quad \quad + (4+4a-58a^2+74a^3)b^3 + (3+48a-57a^2)b^4 - 12b^5 + 6b^6)\nu^2 \\
&\quad - 2(-2a(-18+33a+4a^3) + 2(-2+17a-12a^2+61a^3)b + (-10+28a-170a^2+21a^4)b^2 \\
&\quad \quad - 2(5-37a-8a^2+40a^3)b^3 + (-7-24a+105a^2)b^4 + (12-48a)b^5 + 3b^6)\nu^3 \\
&\quad + (24a(3-4a+2a^2+a^3) - 4(4+9a+18a^2+21a^3)b + (50+16a+263a^2-24a^3-33a^4)b^2 \\
&\quad \quad + 2(5-111a+25a^2+79a^3)b^3 - 3(-13+8a+93a^2)b^4 + 204ab^5 - 48b^6)\nu^4 \\
&\quad - (-12(-2+a)a + 4a(-17+15a+3a^2)b + (36-72a-55a^2+12a^3+9a^4)b^2 + (22+108a-52a^2-64a^3)b^3 \\
&\quad \quad + 10(-5+6a+15a^2)b^4 - 6(4+25a)b^5 + 54b^6)\nu^5 \\
&\quad + b(4-4a+(-6+20a-6a^2)b+2(-8-2a+5a^2+3a^3)b^2+(15-24a-25a^2)b^3+12(1+3a)b^4-18b^5)\nu^6), \\
\end{aligned} \tag{C39}$$

$$\begin{aligned}
\mathcal{G}_4^{(3,2)} = & -\nu \left((1-a^2)^2 b^2 + (1-a^2) (8(a-1)a + 4b - (8-4a+a^2) b^2 - 2(1-a)b^3 + b^4) \nu \right. \\
& + (-24(-1+a)a + 4(-2-3a-5a^2+9a^3) b + (21-11a^2-4a^3+2a^4) b^2 \\
& \quad + (2-8a+12a^2-8a^3) b^3 + 2(-1-2a+a^2) b^4 + 2ab^5) \nu^2 \\
& + (-4a(6-5a+2a^2+a^3) + 4a(9+a+2a^2) b + (-20+16a-49a^2+4a^3+5a^4) b^2 + 2a(7+4a-8a^2) b^3 \\
& \quad + a(-12+19a) b^4 - 2(-1+a) b^5 - b^6) \nu^3 \\
& + (-4(-2+a)a + 4(2-7a+5a^2+a^3) b + (-1-23a^2+4a^3+3a^4) b^2 - 2(4-15a+6a^2+8a^3) b^3 \\
& \quad + (1-4a+31a^2) b^4 + (4-18a) b^5 + b^6) \nu^4 \\
& + 2b(2(-1+a) + (4-10a+3a^2) b + (1+8a-5a^2-3a^3) b^2 + (-5+6a+9a^2) b^3 - 11ab^4 + 3b^5) \nu^5 \\
& \left. + b^2(-1+6(1-a)b + (-3+8a+3a^2) b^2 - 4(1+2a)b^3 + 5b^4) \nu^6 + (-1+b)^2 b^4 \nu^7 \right) \\
+ \gamma \left(& (1-a^2)^2 b^2 + 2(1-a^2) b(2+(-3+2a-2a^2) b + 2(-1+a) b^2 + b^3) \nu \right. \\
& + (-8a(3-4a-4a^2+5a^3) + 4a(-3-8a+9a^2) b + 4(1+4a+a^2-4a^3+2a^4) b^2 \\
& \quad + (2-12a+28a^2-20a^3) b^3 + 3(-1-4a+a^2) b^4 + 6ab^5) \nu^2 \\
& - (4a(-18+23a+a^3) + 4(4+a+14a^2-37a^3) b + (-31-32a+132a^2+8a^3-17a^4) b^2 \\
& \quad + 2(1-9a-20a^2+30a^3) b^3 + (1+40a-57a^2) b^4 + 4(-2+a) b^5 + 4b^6) \nu^3 \\
& + (-8a(9-9a+4a^2+2a^3) + 4(2+21a+8a^2+11a^3) b + (-59+24a-182a^2+16a^3+21a^4) b^2 \\
& \quad - 4(5-30a+2a^2+22a^3) b^3 + 2(-3-12a+67a^2) b^4 + (16-68a) b^5 + 4b^6) \nu^4 \\
& + (-12(-2+a)a + 4(3-19a+15a^2+3a^3) b + (16-36a-62a^2+12a^3+9a^4) b^2 \\
& \quad - 4a(-24+11a+14a^2) b^3 + 2(-14+16a+61a^2) b^4 - 108ab^5 + 27b^6) \nu^5 \\
& + b(8(a-1) + 2(7-20a+6a^2) b + 2(9+10a-10a^2-6a^3) b^2 + (-23+36a+43a^2) b^3 - 2(8+29a) b^4 + 26b^5) \nu^6 \\
& \left. + b^2(-1-6(-1+a)b + (-1+8a+3a^2) b^2 - 8(1+a) b^3 + 7b^4) \nu^7 \right), \tag{C40}
\end{aligned}$$

$$\begin{aligned}
\mathcal{G}_4^{(4,1)} &= (a-b)^6 \gamma^8 \nu (3b^2 + (-8+6b^2)\nu + (-4+3b^2)\nu^2) \\
&- (a-b)^4 \gamma^7 (-b^2(3-3a^2+2(-2+a)b+b^2) + b(4(-2+a) + (-2+9a^2)b + (8-16a)b^2 + b^3)\nu \\
&\quad + (8(2+2a-7a^2) - 8(3-11a+2a^2)b + 3(-4+5a^2)b^2 - 36ab^3 + 15b^4)\nu^2 \\
&\quad + (-4(14-8a+a^2) + 4(2+9a)b + 3(-6+5a^2)b^2 - 8(1+4a)b^3 + 23b^4)\nu^3 \\
&\quad + (40-48a+12a^2 + 8(3-4a+2a^2)b + (-13+6a^2)b^2 - 2(2+5a)b^3 + 10b^4)\nu^4) \\
&+ (-a+b)^2 \gamma^6 (2b^2(1+4a-9a^2-4a^3+8a^4 + (-4+4a+14a^2-13a^3)b + (2-8a+4a^2)b^2 + ab^3) \\
&\quad + b(8a(5-8a+2a^3) + (-27+32a+36a^2-32a^3+11a^4)b + (8-20a+40a^2-42a^3)b^2 \\
&\quad + (-4-40a+41a^2)b^3 - 4(-3+a)b^4 - 3b^5)\nu \\
&\quad + (-8(3-4a-18a^3+22a^4) - 4a(26+66a-147a^2+32a^3)b + (68+208a-554a^2+208a^3-29a^4)b^2 \\
&\quad - 2(28-96a+56a^2+17a^3)b^3 + 2(-11-8a+55a^2)b^4 + (24-58a)b^5 + 3b^6)\nu^2 \\
&\quad + (4(19-96a^2+56a^3+6a^4) - 8(8-69a+59a^2-29a^3+6a^4)b + (-218+336a-526a^2+192a^3-7a^4)b^2 \\
&\quad - 2(36-138a+60a^2+25a^3)b^3 + 2(-13+16a+58a^2)b^4 - 94ab^5 + 21b^6)\nu^3 \\
&\quad + (4(-28+24a+78a^2-92a^3+25a^4) + 4a(-58+114a-103a^2+32a^3)b + (98-312a+404a^2-200a^3+37a^4)b^2 \\
&\quad - 4(-16+30a-21a^2+9a^3)b^3 + (-2+32a+26a^2)b^4 - 8(3+5a)b^5 + 21b^6)\nu^4 \\
&\quad + (4(15-32a+18a^2+a^4) + 8(8-32a+43a^2-27a^3+4a^4)b + (77-272a+370a^2-160a^3+20a^4)b^2 \\
&\quad - 4(-16+36a-20a^2+a^3)b^3 + (2+8a-13a^2)b^4 + 2(-6+a)b^5 + 6b^6)\nu^5) \\
&+ \gamma^5 ((-1+a)b^2(1-7a-2a^2+38a^3-2a^4-34a^5+2(2+5a-35a^2-9a^3+45a^4)b \\
&\quad + (-5+35a+35a^2-77a^3)b^2 + 4(-1-4a+5a^2)b^3 + (1+a)b^4) \\
&\quad - b(4(2-11a+54a^3-48a^4-18a^5+20a^6) + (14+88a-465a^2+296a^3+341a^4-288a^5+30a^6)b \\
&\quad - 4(12-66a+8a^2+134a^3-110a^4+27a^5)b^2 + (-35-96a+348a^2-368a^3+151a^4)b^3 \\
&\quad - 4(-6+21a-42a^2+22a^3)b^4 + (5-32a+11a^2)b^5 + 4ab^6)\nu \\
&\quad + (8(-2+6a+3a^2-16a^3+30a^4-60a^5+40a^6) + 8(3-36a+46a^2-20a^3+162a^4-208a^5+48a^6)b \\
&\quad + (145-112a-408a^2-1392a^3+2810a^4-1120a^5+149a^6)b^2 \\
&\quad + (-52+470a+688a^2-2088a^3+1240a^4-300a^5)b^3 + (-136-168a+726a^2-576a^3+123a^4)b^4 \\
&\quad + 4(7-22a+7a^2+22a^3)b^5 + (-7+48a-67a^2)b^6 + (-4+6a)b^7 + b^8)\nu^2 \\
&\quad - (4(-16+40a+33a^2+32a^3-292a^4+192a^5-2a^6) - 4(-12+170a+56a^2-888a^3+818a^4-285a^5+72a^6)b \\
&\quad + (352+240a-3782a^2+5136a^3-3860a^4+1504a^5-179a^6)b^2 \\
&\quad + 4(-40+480a-988a^2+1208a^3-626a^4+121a^5)b^3 + (-418+1440a-2632a^2+1760a^3-439a^4)b^4 \\
&\quad + 4(-38+137a-128a^2+26a^3)b^5 + (-24+32a+57a^2)b^6 + (8-28a)b^7 + b^8)\nu^3 \\
&\quad + (-4(24-64a-85a^2+144a^3+176a^4-268a^5+88a^6) - 2(-122+421a-128a^2-706a^3+250a^4+5a^5)b^3 \\
&\quad - 8(-2+124a-152a^2-122a^3+273a^4-205a^5+46a^6)b + (137-104a-396a^2-80a^3+302a^4)b^4 \\
&\quad + (493-1048a+599a^2+984a^3-2402a^4+832a^5-65a^6)b^2 - 4(3-15a-45a^2+101a^3)b^5 \\
&\quad + (7-64a+232a^2)b^6 - 64ab^7 + 9b^8)\nu^4 \\
&\quad + (-4(-16+56a+31a^2-192a^3+182a^4-56a^5+9a^6) - 4(-14-91a+312a^2-598a^3+578a^4-317a^5+52a^6)b \\
&\quad + (-178+840a-3335a^2+4728a^3-3827a^4+1216a^5-125a^6)b^2 \\
&\quad + 8(-26+245a-482a^2+575a^3-258a^4+29a^5)b^3 + (-397+1312a-2480a^2+1392a^3+72a^4)b^4 \\
&\quad - 8(16-67a+43a^2+58a^3)b^5 + (-31+428a^2)b^6 + (8-168a)b^7 + 25b^8)\nu^5 \\
&\quad + (4(-4+20a-27a^2+16a^3+6a^4-12a^5+3a^6) - 8(5-24a+70a^2-70a^3+33a^4-14a^5+2a^6)b \\
&\quad + (-93+640a-1098a^2+1152a^3-860a^4+256a^5-26a^6)b^2 - 4(5-58a+43a^2+46a^3)b^5 \\
&\quad + 2(-94+301a-576a^2+692a^3-316a^4+38a^5)b^3 + 2(-31+172a-427a^2+272a^3+6a^4)b^4 \\
&\quad + (-35+16a+194a^2)b^6 + (4-86a)b^7 + 14b^8)\nu^6), \tag{C41}
\end{aligned}$$

$$\begin{aligned}
\mathcal{G}_4^{(4,2)} = & \gamma^4 (2(1-a)^2 b^2 (3(1-2a-5a^2+4a^3+6a^4) + (2+17a-12a^2-31a^3)b + (-3+2a+13a^2)b^2) \\
& + (1-a)b(-8(-6+15a+17a^2-42a^3-6a^4+20a^5) + (26+258a-496a^2-184a^3+427a^4-85a^5)b \\
& + 2(117a-56+121a^2-245a^3+95a^4)b^2 - (17+129a-263a^2+145a^3)b^3 \\
& + 12(1-4a+3a^2)b^4 + (1+a)b^5) \nu \\
& + (8(8-26a+19a^2+24a^3-80a^4+100a^5-45a^6) - 4(26-155a+176a^2-100a^3+360a^4-446a^5+140a^6)b \\
& + (-322+368a+311a^2+680a^3-1913a^4+1096a^5-204a^6)b^2 \\
& + 2(50-279a+216a^2+148a^3-292a^4+144a^5)b^3 \\
& + (149-408a+426a^2-32a^3-111a^4)b^4 + 4(16-33a+9a^2+4a^3)b^5 - (7-16a+a^2)b^6 - 2ab^7) \nu^2 \\
& + (-4(68-168a+35a^2-96a^3+500a^4-400a^5+65a^6) + 4(-29+88a-136a^2+75a^3)b^5 \\
& - 8(-17+141a+52a^2-681a^3+778a^4-375a^5+90a^6)b + (10+96a-56a^2)b^6 \\
& + (702-272a-4072a^2+7920a^3-7154a^4+3104a^5-456a^6)b^2 - 2(10+3a)b^7 + 5b^8 \\
& + 2(-24+652a-2248a^2+3360a^3-2124a^4+517a^5)b^3 + (-247+1376a-2788a^2+2336a^3-816a^4)b^4) \nu^3 \\
& + (4(112-288a-85a^2+456a^3+28a^4-380a^5+170a^6) \\
& + 4(4+406a-776a^2+482a^3+134a^4-441a^5+120a^6)b \\
& + (-794+2600a-5662a^2+4760a^3-382a^4+96a^5-25a^6)b^2 \\
& + 2(-362+2067a-2868a^2+1598a^3-954a^4+255a^5)b^3 \\
& + (-728+1840a-1960a^2+1776a^3-891a^4)b^4 \\
& + 4(-26+41a-122a^2+182a^3)b^5 + (36+128a-393a^2)b^6 + 8(-5+13a)b^7 - 5b^8) \nu^4 \\
& - (8(44-148a+58a^2+232a^3-326a^4+140a^5-15a^6) \\
& + 8(-28-2a-194a^2+670a^3-717a^4+365a^5-70a^6)b \\
& + (122+56a-4686a^2+7592a^3-6629a^4+2496a^5-302a^6)b^2 \\
& + 2(8+1051a-2092a^2+3086a^3-1616a^4+209a^5)b^3 + (-473+1328a-3012a^2+1656a^3+288a^4)b^4 \\
& - 4(43-171a+93a^2+217a^3)b^5 + (-35+32a+730a^2)b^6 - 292ab^7 + 45b^8) \nu^5 \\
& + (128-560a+948a^2-736a^3+8a^4+240a^5-60a^6+4(22-221a+632a^2-662a^3+386a^4-155a^5+20a^6)b \\
& + (430-2304a+4391a^2-5192a^3+3677a^4-1144a^5+133a^6)b^2 \\
& + (620-2482a+4752a^2-5032a^3+2392a^4-376a^5)b^3 + (305-1288a+2502a^2-1696a^3+76a^4)b^4 \\
& + 4(10-98a+113a^2+114a^3)b^5 + (31-144a-506a^2)b^6 + (40+258a)b^7 - 55b^8) \nu^6 \\
& + (-4(4-16a+39a^2-48a^3+22a^4) + 8(5-10a+20a^2-30a^3+17a^4+a^5)b + (20+74a)b^7 - 20b^8 \\
& + (74-544a+1100a^2-960a^3+374a^4-96a^5+9a^6)b^2 + (176-668a+944a^2-696a^3+336a^4-66a^5)b^3 \\
& + (71-96a+204a^2-272a^3+59a^4)b^4 + 4(-17+8a+22a^2+17a^3)b^5 + (14-64a-113a^2)b^6) \nu^7), \quad (\text{C42})
\end{aligned}$$

$$\begin{aligned}
\mathcal{G}_4^{(4,3)} = & \gamma^3 (-(1-a)^3(1+a)b^2(9-19a^2+2a(-3+8b)) \\
& +(1-a)^3b(4(-22-7a+57a^2+40a^3)+(-27-249a-109a^2+113a^3)b-32(-3+5a^2)b^2+8(1+7a)b^3)\nu \\
& +(-1+a)(8(12-30a+15a^2+31a^3-59a^4+31a^5)+8(-17+54a-16a^2+12a^3-83a^4+50a^5)b \\
& +(-371+389a+186a^2-158a^3-129a^4+95a^5)b^2+4(-6-49a+183a^2-139a^3+3a^4)b^3 \\
& +(106-454a+410a^2-38a^3)b^4-8(-5+4a+a^2)b^5-2(1+a)b^6)\nu^2 \\
& +(4(1-a)^2(112-40a-107a^2-238a^3+141a^4)+4(-26+187a+132a^2-1178a^3+1594a^4-951a^5+240a^6)b \\
& -(619-840a-1579a^2+5448a^3-6285a^4+3136a^5-531a^6)b^2 \\
& +8(2-9a+30a^2-18a^3)b^5+2(5-32a+11a^2)b^6 \\
& -8(36-94a-128a^2+470a^3-382a^4+103a^5)b^3+2(-63-64a+420a^2-496a^3+203a^4)b^4+8ab^7)\nu^3 \\
& +(-4(208-516a-49a^2+864a^3-438a^4-260a^5+195a^6)-4(-18+43a-100a^2+91a^3)b^5 \\
& -8(17+130a-486a^2+738a^3-427a^4-12a^5+30a^6)b+(5-144a+149a^2)b^6 \\
& +(297-2136a+8111a^2-10120a^3+5308a^4-1824a^5+220a^6)b^2+(20-6a)b^7-5b^8 \\
& +(664-4780a+8224a^2-7148a^3+4012a^4-882a^5)b^3+(880-2416a+3102a^2-2288a^3+757a^4)b^4)\nu^4 \\
& +(-8(-96+320a-242a^2-288a^3+574a^4-280a^5+25a^6)+4(200-344a+356a^2+360a^3-374a^4+67a^5)b^3 \\
& -4(-86+327a-68a^2-1314a^3+1676a^4-870a^5+200a^6)b+(-44-192a-596a^2+256a^3+271a^4)b^4 \\
& +(879-2520a-269a^2+3320a^3-4465a^4+2304a^5-358a^6)b^2 \\
& -4(34-139a+76a^2+98a^3)b^5+(-54-32a+351a^2)b^6+(40-116a)b^7+5b^8)\nu^5 \\
& +(8(-44+182a-341a^2+288a^3-34a^4-60a^5+15a^6)-8(1-201a+566a^2-636a^3+442a^4-170a^5+20a^6)b \\
& +(-591+2680a-6219a^2+8488a^3-5887a^4+2016a^5-272a^6)b^2 \\
& +4(-158+781a-1532a^2+1550a^3-842a^4+173a^5)b^3+(-450+1664a-2664a^2+1920a^3-305a^4)b^4 \\
& -8(10-48a+63a^2+21a^3)b^5+(-50+128a+283a^2)b^6-160ab^7+27b^8)\nu^6 \\
& +(4(16-56a+151a^2-192a^3+88a^4)-4(38-81a+212a^2-274a^3+136a^4+8a^5)b \\
& +(-233+1848a-3547a^2+3368a^3-1463a^4+384a^5-36a^6)b^2 \\
& +4(-152+508a-804a^2+584a^3-270a^4+63a^5)b^3+(-118+448a-560a^2+576a^3-261a^4)b^4 \\
& +4(30-67a+16a^2+14a^3)b^5+(8+96a-13a^2)b^6-4(10+9a)b^7+19b^8)\nu^7 \\
& +(4a(-4+3a)+(8+80a^2-80a^3)b+(-68+192a-422a^2+384a^3-73a^4)b^2 \\
& +2(-28+190a-280a^2+110a^3-22a^4+9a^5)b^3-2(22-16a-83a^2+40a^3+27a^4)b^4 \\
& +4(12-53a+32a^2+15a^3)b^5+(31+16a-70a^2)b^6+(-20+22a)b^7+2b^8)\nu^8), \tag{C43}
\end{aligned}$$

$$\begin{aligned}
& \mathcal{G}_4^{(4,4)} + \nu^2 (4(1-a)^4(1+a)^2b(4+3b) + (1-a)^3(1+a) (16(-1-2a+3a^2) - 16(2-a-3a^2))b \\
& \quad - (63-42a-5a^2)b^2 - 16(2-a)b^3) \nu \\
& + (1-a)^2 (16(4-a-12a^2+7a^3+2a^4) - 4(-10+19a+36a^2-53a^3+8a^4))b \\
& \quad + (89-62a-42a^2+74a^3-35a^4)b^2 + 2(34-47a-12a^2+a^3)b^3 + 2(17-6a+a^2)b^4) \nu^2 \\
& - (1-a) (16(6-16a+9a^2+13a^3-13a^4+a^5) + 8(1-35a+45a^2-14a^3-5a^4+6a^5))b \\
& \quad + (63-49a+298a^2-390a^3+130a^4+2a^5)b^2 + 2a(-77+79a-51a^2+17a^3)b^3 + (49-63a+25a^2+17a^3)b^4 \\
& \quad - 4(-5+4a+a^2)b^5 - (1+a)b^6) \nu^3 \\
& + (4(16-64a+133a^2-120a^3+26a^4+12a^5-3a^6) + 4(-16-18a+76a^2-126a^3+122a^4-43a^5+4a^6))b \\
& \quad + (21+296a-402a^2+176a^3+16a^4-120a^5+29a^6)b^2 - 2(62-101a+192a^2-160a^3+8a^4+10a^5)b^3 \\
& \quad + (1+88a-66a^2-64a^3+49a^4)b^4 - 4(-4+9a-15a^2+8a^3)b^5 + (1-16a+7a^2)b^6 + 2ab^7) \nu^4 \\
& + (-8(2-4a+17a^2-24a^3+11a^4) + 8(6-13a+46a^2-47a^3+17a^4+a^5))b - (13+64a-44a^2+80a^3-64a^4)b^4 \\
& \quad + (-37-120a+235a^2-488a^3+341a^4-96a^5+9a^6)b^2 + (32+52a+64a^2-24a^3+72a^4-54a^5)b^3 \\
& \quad - 4(-7+14a-26a^2+21a^3)b^5 + (2-32a+36a^2)b^6 - 2(-2+a)b^7 - b^8) \nu^5 \\
& + (4(4-3a)a + 4(2-3a-20a^2+20a^3))b + (15-160a+435a^2-384a^3+73a^4)b^2 \\
& \quad - 2(-22+93a-140a^2+66a^3-22a^4+9a^5)b^3 + (40-48a-88a^2+80a^3+39a^4)b^4 \\
& \quad - 4(6-15a+6a^2+22a^3)b^5 + (-8+65a^2)b^6 + (8-24a)b^7 + b^8) \nu^6 \\
& + b(8(-1+a) - (-27+32a+a^2))b - 2a(31-76a+38a^2)b^2 \\
& \quad + (5-16a-60a^2+72a^3+9a^4)b^3 - 4(1-17a+23a^2+8a^3)b^4 + (-17+32a+55a^2)b^5 - 34ab^6 + 6b^7) \nu^7 \\
& + (-1+b)^2b^2(-1-10(-1+a)b+2(1-3a)^2b^2+(2-14a)b^3+5b^4) \nu^8 + (1-b)^4b^4\nu^9, \tag{C44}
\end{aligned}$$

$$\begin{aligned}
\mathcal{G}_4^{(4,5)} = & -\gamma\nu (16(1-a)^4(1+a)^2b(1+b) \\
& +(1-a)^3(1+a)(16(1-a)^2+32(1-a^2)b-(41-102a+45a^2)b^2-(64-48a)b^3)\nu \\
& -(1-a)^3(-64(-2-3a+a^2+4a^3)-4(-22-7a+25a^2+72a^3)b+(155+217a-19a^2-81a^3)b^2 \\
& \quad -32(1-4a+a^2)b^3+8(-9+a)b^4)\nu^2 \\
& +(1-a)(-8(-44+58a+93a^2-171a^3+39a^4+25a^5)+8(17-38a-96a^2+228a^3-125a^4+14a^5)b \\
& \quad +(434-350a-888a^2+1400a^3-765a^4+163a^5)b^2-2(-106+163a+139a^2-287a^3+83a^4)b^3 \\
& \quad +(27+35a-61a^2-13a^3)b^4-12(3-4a+a^2)b^5+(1+a)b^6)\nu^3 \\
& +(4(1-a)^2(-112+168a+29a^2-238a^3+21a^4)+4(-20+310a-548a^2+208a^3+162a^4-181a^5+68a^6)b \\
& \quad +(-473+1360a-2436a^2+2896a^3-1576a^4+192a^5+53a^6)b^2 \\
& \quad +4(-76+394a-664a^2+506a^3-194a^4+29a^5)b^3-(283-736a+660a^2-144a^3-63a^4)b^4 \\
& \quad +(-8+12a+72a^2-56a^3)b^5+(5-32a+11a^2)b^6+4ab^7)\nu^4 \\
& +(4(68-272a+555a^2-496a^3+98a^4+60a^5-15a^6)-8(22+70a-220a^2+310a^3-277a^4+100a^5-10a^6)b \\
& \quad +(58+600a-73a^2-1208a^3+1145a^4-736a^5+142a^6)b^2+(-348+490a-688a^2+352a^3+448a^4-208a^5)b^3 \\
& \quad +(104-24a+6a^2-288a^3+215a^4)b^4-4(-25+68a-101a^2+64a^3)b^5 \\
& \quad +(1-80a+77a^2)b^6+2(6-a)b^7-3b^8)\nu^5 \\
& +(-4(16-40a+141a^2-192a^3+88a^4)+4(42-93a+316a^2-342a^3+136a^4+8a^5)b \\
& \quad +(15-1000a+1817a^2-2424a^3+1397a^4-384a^5+36a^6)b^2 \\
& \quad -4(-64+136a-364a^2+248a^3-138a^4+57a^5)b^3 \\
& \quad +(54-384a+144a^2-192a^3+271a^4)b^4-4(-6+a-40a^2+90a^3)b^5 \\
& \quad +(-28-32a+215a^2)b^6+(24-68a)b^7+3b^8)\nu^6 \\
& +(12(4-3a)a+8(1-3a-30a^2+30a^3)b+(94-512a+1292a^2-1152a^3+219a^4)b^2 \\
& \quad -2(-94+395a-560a^2+242a^3-66a^4+27a^5)b^3+(55-88a-276a^2+240a^3+132a^4)b^4 \\
& \quad -4(17-77a+77a^2+59a^3)b^5+(-31+64a+266a^2)b^6-128ab^7+21b^8)\nu^7 \\
& +b(16(-1+a)+(53-64a-2a^2)b-8(-2+17a-38a^2+19a^3)b^2+(-5+64a+134a^2)b^5 \\
& \quad +(-11-32a-96a^2+144a^3+18a^4)b^3-8(2-20a+29a^2+8a^3)b^4-8(3+10a)b^6+21b^7)\nu^8 \\
& +(-1+b)^2b^2(-1-10(-1+a)b+3(1-4a+6a^2)b^2-14ab^3+6b^4)\nu^9, \tag{C45}
\end{aligned}$$

$$\begin{aligned}
\mathcal{G}_4^{(4,6)} = & \gamma^2 (4(-1+a)^4(1+a)^2b^2 - (-1+a)^3(1+a)b(-16(-4+a+5a^2) + (31+54a-69a^2)b + 16(-2+3a)b^2) \nu \\
& + (1-a)^2(-16(-4+7a-9a^3+6a^4) + (-40+76a+16a^2+44a^3-96a^4)b \\
& + (-223+234a+168a^2-258a^3+31a^4)b^2 - 4(34+a-92a^2+33a^3)b^3 + 12(3-10a+3a^2)b^4) \nu^2 \\
& + (1-a)(32(-11+12a+12a^2+4a^3-34a^4+17a^5) + 8(-5-4a-50a^2+184a^3-219a^4+90a^5)b \\
& + (101-275a+90a^2+1074a^3-1193a^4+311a^5)b^2 - 4(-76+225a-75a^2-97a^3+55a^4)b^3 \\
& + 2(97-159a+73a^2+17a^3)b^4 + (8+32a-40a^2)b^5 - 2(1+a)b^6) \nu^3 \\
& - (-4(192-460a-67a^2+936a^3-682a^4-52a^5+133a^6) - 4(-18+42a-39a^2+14a^3)b^5 \\
& + 4(-50+25a+580a^2-1466a^3+1242a^4-349a^5+16a^6)b \\
& + (-461+56a+4391a^2-7776a^3+5605a^4-2072a^5+289a^6)b^2 + 2(-1-8a+5a^2)b^6 + 2ab^7 \\
& + (40-1448a+3736a^2-4588a^3+2836a^4-622a^5)b^3 + 2(172-532a+741a^2-472a^3+103a^4)b^4) \nu^4 \\
& - (-4(-208+704a-671a^2-384a^3+1082a^4-560a^5+45a^6) \\
& - 8(-31+250a-306a^2-191a^3+457a^4-280a^5+80a^6)b \\
& + (1049-3240a+3832a^2-2712a^3+199a^4+736a^5-212a^6)b^2 \\
& + (944-3350a+5048a^2-2996a^3+744a^4-44a^5)b^3 + (353-1232a+1168a^2-456a^3+24a^4)b^4 \\
& + 4(-20+71a-77a^2+22a^3)b^5 + (8+26a^2)b^6 - 2(2+3a)b^7 + b^8) \nu^5 \\
& + (-8(-56+226a-447a^2+392a^3-66a^4-60a^5+15a^6) \\
& + 4(-38-351a+1004a^2-1244a^3+996a^4-370a^5+40a^6)b \\
& + (291-712a+3255a^2-5832a^3+4199a^4-1744a^5+278a^6)b^2 \\
& - 4(6+239a-556a^2+630a^3-518a^4+145a^5)b^3 + (300-832a+1088a^2-992a^3+383a^4)b^4 \\
& - 4(-46+115a-142a^2+82a^3)b^5 + (-6-64a+99a^2)b^6 + (8-16a)b^7 + b^8) \nu^6 \\
& + (-12(8-24a+73a^2-96a^3+44a^4) + 8(29-64a+198a^2-231a^3+102a^4+6a^5)b \\
& + (211-2184a+4029a^2-4344a^3+2145a^4-576a^5+54a^6)b^2 \\
& - 8(-82+245a-458a^2+326a^3-153a^4+45a^5)b^3 + (124-672a+456a^2-416a^3+409a^4)b^4 \\
& - 16(6-18a+6a^2+25a^3)b^5 + (8-32a+305a^2)b^6 - 104ab^7 + 15b^8) \nu^7 \\
& + (12(4-3a)a + 4(-2-3a-60a^2+60a^3)b + (147-544a+1279a^2-1152a^3+219a^4)b^2 \\
& - 2(-100+492a-700a^2+286a^3-66a^4+27a^5)b^3 + (64-72a-354a^2+240a^3+147a^4)b^4 \\
& - 4(28-115a+103a^2+52a^3)b^5 + (-24+48a+271a^2)b^6 - 2(4+63a)b^7 + 23b^8) \nu^8 \\
& + b(8(-1+a) - (-26+32a+a^2)b - 2(-8+37a-76a^2+38a^3)b^2 + (-21-16a-36a^2+72a^3+9a^4)b^3 \\
& - 4(-2-23a+35a^2+8a^3)b^4 + (-18+32a+79a^2)b^5 - 2(2+23a)b^6 + 10b^7) \nu^9). \tag{C46}
\end{aligned}$$

-
- [1] A. Strominger and C. Vafa, “Microscopic origin of the Bekenstein-Hawking entropy,” *Phys. Lett. B* **379**, 99 (1996) [hep-th/9601029].
- [2] P. C. Argyres, S. Dimopoulos and J. March-Russell, “Black holes and submillimeter dimensions,” *Phys. Lett. B* **441**, 96 (1998) [hep-th/9808138].
- [3] M. I. Cai and G. J. Galloway, “On the Topology and area of higher dimensional black holes,” *Class. Quant. Grav.* **18**, 2707 (2001) [hep-th/0102149].
- [4] G. J. Galloway and R. Schoen, “A Generalization of Hawking’s black hole topology theorem to higher dimensions,” *Commun. Math. Phys.* **266**, 571 (2006) [gr-qc/0509107].
- [5] S. Hollands and S. Yazadjiev, “Uniqueness theorem for 5-dimensional black holes with two axial Killing fields,” *Commun. Math. Phys.* **283**, 749-768 (2008) [arXiv:0707.2775 [gr-qc]].
- [6] F. R. Tangherlini, “Schwarzschild field in n dimensions and the dimensionality of space problem,” *Nuovo Cim.* **27**, 636 (1963).
- [7] R. C. Myers and M. J. Perry, “Black Holes in Higher Dimensional Space-Times,” *Annals Phys.* **172**, 304 (1986)
- [8] R. Emparan and H. S. Reall, “A Rotating black ring solution in five-dimensions,” *Phys. Rev. Lett.* **88**, 101101 (2002) [arXiv:hep-th/0110260 [hep-th]].

- [9] A. A. Pomeransky and R. A. Sen'kov, "Black ring with two angular momenta," [arXiv:hep-th/0612005 [hep-th]].
- [10] T. Mishima and H. Iguchi, "New axisymmetric stationary solutions of five-dimensional vacuum Einstein equations with asymptotic flatness," Phys. Rev. D **73**, 044030 (2006) [arXiv:hep-th/0504018 [hep-th]].
- [11] P. Figueras, "A Black ring with a rotating 2-sphere," JHEP **07**, 039 (2005) [arXiv:hep-th/0505244 [hep-th]].
- [12] V. A. Belinsky and V. E. Zakharov, "Stationary Gravitational Solitons with Axial Symmetry," Sov. Phys. JETP **50**, 1 (1979).
- [13] A. A. Pomeransky, "Complete integrability of higher-dimensional Einstein equations with additional symmetry, and rotating black holes," Phys. Rev. D **73** (2006) 044004 [hep-th/0507250].
- [14] S. Tomizawa, Y. Morisawa and Y. Yasui, "Vacuum solutions of five dimensional Einstein equations generated by inverse scattering method," Phys. Rev. D **73**, 064009 (2006) [arXiv:hep-th/0512252 [hep-th]].
- [15] H. Iguchi and T. Mishima, "Solitonic generation of five-dimensional black ring solution," Phys. Rev. D **73**, 121501 (2006) [arXiv:hep-th/0604050 [hep-th]].
- [16] S. Tomizawa and M. Nozawa, "Vacuum solutions of five-dimensional Einstein equations generated by inverse scattering method. II. Production of black ring solution," Phys. Rev. D **73**, 124034 (2006) [arXiv:hep-th/0604067 [hep-th]].
- [17] Y. Morisawa, S. Tomizawa and Y. Yasui, "Boundary Value Problem for Black Rings," Phys. Rev. D **77**, 064019 (2008) [arXiv:0710.4600 [hep-th]].
- [18] Y. Chen, K. Hong and E. Teo, Phys. Rev. D **84**, 084030 (2011) doi:10.1103/PhysRevD.84.084030 [arXiv:1108.1849 [hep-th]].
- [19] H. Stephani, D. Kramer, M. A. H. MacCallum, C. Hoenselaers, and E. Herlt, *Exact Solutions of Einstein's Field Equations*, 2nd ed. (Cambridge University Press, Cambridge, England, 2003).
- [20] F. J. Ernst, "New formulation of the axially symmetric gravitational field problem," Phys. Rev. **167**, 1175-1179 (1968).
- [21] F. J. Ernst, "New Formulation of the Axially Symmetric Gravitational Field Problem. II," Phys. Rev. **168**, 1415-1417 (1968).
- [22] J. Ehlers. Konstruktionen und Charakterisierung von Lösungen der Einsteinschen Gravitationsfeldgleichungen. Phd thesis, Hamburg University, 1957.
- [23] B. K. Harrison. New Solutions of the Einstein-Maxwell Equations from Old. Journal of Mathematical Physics, **9** (11), 1744-1752, (1968).
- [24] G. Clement, "From Schwarzschild to Kerr: Generating spinning Einstein-Maxwell fields from static fields," Phys. Rev. D **57**, 4885-4889 (1998) [arXiv:gr-qc/9710109 [gr-qc]].
- [25] G. Clement, "Generating rotating fields in general relativity," Grav. Cosmol. **5**, 281-284 (1999) [arXiv:gr-qc/9910096 [gr-qc]].
- [26] D. Maison, "EHLERS-HARRISON TYPE TRANSFORMATIONS FOR JORDAN'S EXTENDED THEORY OF GRAVITATION," Gen. Rel. Grav. **10**, 717-723 (1979)
- [27] S. Giusto and A. Saxena, "Stationary axisymmetric solutions of five dimensional gravity," Class. Quant. Grav. **24**, 4269-4294 (2007) [arXiv:0705.4484 [hep-th]].
- [28] S. Mizoguchi and N. Ohta, "More on the similarity between $D = 5$ simple supergravity and M theory," Phys. Lett. B **441**, 123-132 (1998) [arXiv:hep-th/9807111 [hep-th]].
- [29] S. Mizoguchi and G. Schroder, "On discrete U duality in M theory," Class. Quant. Grav. **17**, 835-870 (2000) [arXiv:hep-th/9909150 [hep-th]].
- [30] A. Bouchareb, G. Clement, C. M. Chen, D. V. Gal'tsov, N. G. Scherbukh and T. Wolf, "G(2) generating technique for minimal $D=5$ supergravity and black rings," Phys. Rev. D **76**, 104032 (2007) [erratum: Phys. Rev. D **78**, 029901 (2008)] [arXiv:0708.2361 [hep-th]].
- [31] M. Cvetič and D. Youm, "General rotating five-dimensional black holes of toroidally compactified heterotic string," Nucl. Phys. B **476**, 118-132 (1996) [arXiv:hep-th/9603100 [hep-th]].
- [32] R. Suzuki and S. Tomizawa, "New construction of a charged dipole black ring by the Harrison transformation," Phys. Rev. D **109**, no.8, 084020 (2024) [arXiv:2402.07589 [hep-th]].
- [33] R. Suzuki and S. Tomizawa, Phys. Rev. D **109**, no.12, L121503 (2024) [arXiv:2311.11653 [hep-th]].
- [34] R. Suzuki and S. Tomizawa, "Solution Generation of a Capped Black Hole," [arXiv:2403.17796 [hep-th]].
- [35] S. Tomizawa, Y. Yasui and A. Ishibashi, "Uniqueness theorem for charged dipole rings in five-dimensional minimal supergravity," Phys. Rev. D **81**, 084037 (2010) [arXiv:0911.4309 [hep-th]].
- [36] R. Emparan and H. S. Reall, Phys. Rev. D **65**, 084025 (2002).
- [37] T. Harmark, Phys. Rev. D **70**, 124002 (2004); T. Harmark and P. Olesen, Phys. Rev. D **72**, 124017 (2005).
- [38] F. R. Tangherlini, Nuovo. Cimento, **27**, 636 (1963).
- [39] Y. Chen and E. Teo, "A Rotating black lens solution in five dimensions", Phys. Rev. D **78**, 064062 (2008).
- [40] C. W. Misner, "The Flatter regions of Newman, Unti and Tamburino's generalized Schwarzschild space," J. Math. Phys. **4**, 924-938 (1963)
- [41] S. Tomizawa, Y. Yasui and A. Ishibashi, "Uniqueness theorem for charged rotating black holes in five-dimensional minimal supergravity," Phys. Rev. D **79**, 124023 (2009) [arXiv:0901.4724 [hep-th]].
- [42] H. Elvang, R. Emparan and P. Figueras, "Non-supersymmetric black rings as thermally excited supertubes," JHEP **02**, 031 (2005) [arXiv:hep-th/0412130 [hep-th]].
- [43] A. Feldman and A. A. Pomeransky, "General triple charged black ring solution in supergravity," JHEP **08**, 059 (2014) [arXiv:1406.4617 [hep-th]].