

THE SPECIAL UNITARY GROUPS $SU(2n)$ AS FRAMED MANIFOLDS

HARUO MINAMI

ABSTRACT. Let $[SU(2n), \mathcal{L}]$ denote the bordism class of $SU(2n)$ ($n \geq 2$) equipped with the left invariant framing \mathcal{L} . Then it is well known that $e_{\mathbb{C}}([SU(2n), \mathcal{L}]) = 0$ in \mathbb{O}/\mathbb{Z} where $e_{\mathbb{C}}$ denotes the complex Adams e -invariant. In this note we show that replacing \mathcal{L} by the twisted framing by a specific map it can be transformed into a generator of $\text{Im } e_{\mathbb{C}}$. In addition to that we also show that the same procedure affords an analogous result for a quotient of $SU(2n+1)$ by a circle subgroup which inherits a canonical framing from $SU(2n+1)$ in the usual way.

1. INTRODUCTION

Let G be a simply-connected compact Lie group of dimension $4l-1$ and of rank ≥ 2 . Let us consider it to be the framed manifold equipped with the left invariant framing \mathcal{L} and write $[G, \mathcal{L}]$ for its bordism class in π_{4l-1}^S . Then it is well known [3] that

$$e_{\mathbb{C}}([G, \mathcal{L}]) = 0$$

where $e_{\mathbb{C}} : \pi_{4l-1}^S \rightarrow \mathbb{O}/\mathbb{Z}$ denotes the complex Adams e -invariant. In view of this result we wish to raise the problem of the existence of a map $f : G \rightarrow GL(s, \mathbb{R})$ such that $e_{\mathbb{C}}([G, \mathcal{L}^f])$ is nonzero and if possible, is equal to $(-1)^{\ell-1} B_{\ell}/2\ell$ [2] where \mathcal{L}^f denotes the framing obtained by twisting \mathcal{L} by f and B_l the l th Bernoulli number.

In this note we consider the case $G = SU(2n)$ ($n \geq 2$) where $\dim G = 4n^2 - 1$. The main result is the following theorem.

Theorem. *Let $\rho : SU(2n) \rightarrow GL(4n, \mathbb{R})$ ($n \geq 2$) be the standard real representation of $SU(2n)$. Then we have*

$$e_{\mathbb{C}}([SU(2n), \mathcal{L}^{(n-1)\rho}]) = (-1)^{n-1} B_{n^2}/2n^2$$

The proof is based on Proposition 2.1 of [5]. But also in the case $G = SU(2n+1)$ it can be seen that the procedure used there is adaptable to its quotient by a circle subgroup C , considered as a framed manifold equipped with the induced framing \mathcal{L}_C . Now this framing can be obtained by applying the fact that \mathcal{L} is a C -equivariant framing [4] to the equation (*) of [5, p. 16]. Here let $S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$ and put

$$C = \{\text{diag}(1, \dots, 1, z, \bar{z}) \mid z \in S^1\}$$

where $\text{diag}(a_1, \dots, a_{2n+1})$ denotes the diagonal matrix with a_1, \dots, a_{2n+1} on the diagonal. Then in fact we have

Proposition. $e_{\mathbb{C}}([SU(2n+1)/C, \mathcal{L}_C]) = -B_{n^2+n}/2(n^2+n)$ ($n \geq 1$).

2. TENSOR PRODUCT DECOMPOSITION OF E

Let $G = SU(m)$ ($m \geq 2$) and S be the circle subgroup of G generated by $d(z) = \text{diag}(\bar{z}^{m-1}, z, \dots, z)$ with $z \in S^1$. Regard G as an S -space endowed with $d(z)$ acting on $g \in G$ by the rule $(g, z) \rightarrow gd(z)$ and consider it to be the principal bundle along with the natural projection $p : G \rightarrow G/S$. Let $E = G \times_S \mathbb{C}$ denote the canonical complex line bundle over G/S associated to p where S acts on \mathbb{C} as S^1 . Then its unit sphere bundle $\pi : S(E) \rightarrow G/S$ is naturally isomorphic to $p : G \rightarrow G/S$ as a principal S -bundle over G/S , which is expressed as $(S(E), \pi, G/S) \cong (G, p, G/S)$ as usual.

In order to introduce some further notations we first recall the case $m = 2$. Write

$$R(rz, v) = \begin{pmatrix} rz & v \\ -\bar{v} & r\bar{z} \end{pmatrix}, \quad r \geq 0, \quad z \in S^1, \quad v \in \mathbb{C},$$

for the elements of $SU(2)$. Then $R(rz, v)d(z) = R(r, zv)$ and so letting it correspond to the element $(1 - 2r^2, 2rzv) \in S^2$ we have a homeomorphism between $SU(2)/S$ and S^2 . Thinking of p as a principal S -bundle over S^2 via this homeomorphism we have $(S(L), \pi, S^2) \cong (SU(2), p, S^2)$ where L is used instead of E . From the formula of [5, p. 40] we have $e([S(L), \Phi_L]) = B_1/4$ where Φ_L denotes the trivialization of the stable tangent bundle of $S(L)$. In this case Φ_L coincides with \mathcal{L} via the above isomorphism and so we have $e([SU(2), \mathcal{L}]) = 1/24$.

For brevity we write $(r, zv)_R$ for $(1 - 2r^2, 2rzv) \in S^2$ and identify

$$(r, zv)_R = p(R(rz, v)) = R(r, zv).$$

Here in particular $(r, zv)_R$ represents $(0, 1)_R$ if $r = 0$ and $(1, 0)_R$ if $r = 1$. The latter is assumed to be specified as the base point of S^2 for the reason that it is the image of the identity matrix of $SU(2)$ by p .

Suppose $m \geq 3$ and let $0 \leq j \leq m-2$ and $1 \leq i \leq m_j$ where $m_j = m - j - 1$. Further we assume that the factors used in the product constructions are arranged in ascending order of their suffixes derived by i, j . For fixed j we put

$$R_{i;j}(r_{i;j}z, v_{i;j}) = \begin{pmatrix} I_j & 0 & 0 & 0 & 0 \\ 0 & r_{i;j}z & 0 & v_{i;j} & 0 \\ 0 & 0 & I_{i-1} & 0 & 0 \\ 0 & -\bar{v}_{i;j} & 0 & r_{i;j}\bar{z} & 0 \\ 0 & 0 & 0 & 0 & I_{m_j-i} \end{pmatrix} \quad \text{with } R(r_{i;j}z, v_{i;j}) \in SU(2)$$

where I_s denotes the identity matrix of order s . Then we have

$$(1) \quad \prod_{i=1}^{m_j} R_{i;j}(r_{i;j}z, v_{i;j}) = \begin{pmatrix} I_j & 0 & 0 & 0 & \cdots & 0 \\ 0 & a_{0;j} & b_{1;j} & b_{2;j} & \cdots & b_{m_j;j} \\ 0 & a_{1;j} & c_{1,1;j} & c_{1,2;j} & \cdots & c_{1,m_j;j} \\ 0 & a_{2;j} & 0 & c_{2,2;j} & \cdots & c_{2,m_j;j} \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & a_{m_j;j} & 0 & \cdots & 0 & c_{m_j,m_j;j} \end{pmatrix}$$

where

$$\begin{aligned} a_{0;j} &= r_{1;j} \cdots r_{m_j;j} z^{m_j}, \\ a_{s;j} &= -r_{s+1;j} \cdots r_{m_j;j} z^{m_j-s} \bar{v}_{s;j} \quad (1 \leq s \leq m_j - 1), \\ a_{m_j;j} &= -\bar{v}_{m_j;j}, \end{aligned}$$

$$\begin{aligned}
b_{1;j} &= v_{1;j}, \\
b_{s;j} &= r_{1;j} \cdots r_{s-1;j} z^{s-1} v_{s;j} \quad (2 \leq s \leq m_j), \\
c_{s,t;j} &= -r_{s+1;j} \cdots r_{t-1;j} z^{t-s-1} \bar{v}_{s;j} v_{t;j} \quad (1 \leq s \leq t-2, 3 \leq t \leq m_j), \\
c_{s,s+1;j} &= -\bar{v}_{s;j} v_{s+1;j} \quad (1 \leq s \leq m_j - 1), \\
c_{s,s;j} &= r_{s;j} \bar{z} \quad (1 \leq s \leq m_j), \\
c_{s,t;j} &= 0 \quad (s > t).
\end{aligned}$$

Let $d_j(z) = \text{diag}(1, \cdot^{\cdot^{\cdot}}, 1, \bar{z}^{m_j}, z, \dots, z)$. By referring to the above calculation we have

$$(2) \quad \left(\prod_{i=1}^{m_j} R_{i;j}(r_{i;j}z, v_{i;j}) \right) d_j(z) = \prod_{i=1}^{m_j} R_{i;j}(r_{i;j}, z^i v_{i;j}).$$

The matrix representation of the right-hand side of this equation has a similar form to that of (1). In fact, restricting ourselves to the part we need for later use we have

$$\begin{aligned}
\underline{a}_{0;j} &= r_{1;j} \cdots r_{m_j;j}, \\
(3) \quad \underline{a}_{s;j} &= -r_{s+1;j} \cdots r_{m_j;j} \bar{w}_{s;j} \quad (1 \leq s \leq m_j - 1), \\
\underline{a}_{m_j;j} &= -\bar{w}_{m_j;j},
\end{aligned}$$

where $\underline{a}_{s;j}$ denotes its matrix element corresponding to $a_{s;j}$ and $w_{s;j}$ stands for $z^s v_{s;j}$.

Let $D_0(z) = I_m$ and $D_j(z) = \text{diag}(z^{m-1}, \bar{z}, \cdot^{\cdot^{\cdot}}, \bar{z}, \bar{z}^{m-j}, 1, \dots, 1)$ ($1 \leq j \leq m-2$). Clearly $D_j(z)d(z) (= d(z)D_j(z)) = d_j(z)$. Define $R_j^{\{i\}}(r_{i;j}z, v_{i;j})$ to be the product

$$\left(\prod_{s=1}^{m_j} R_{s;j}(r_{s;j}z, v_{s;j}) \right) D_j(z) \text{ with } r_{s;j} = 1 \text{ for all } s, j \text{ except } s = i.$$

Then by (2) we have

$$(4) \quad R_j^{\{i\}}(r_{i;j}z, v_{i;j})d(z) = R_{i;j}(r_{i;j}, z^i v_{i;j}), \quad z \in S^1$$

since $R_{s;j}(r_{s;j}, z^s v_{s;j}) = I_m$ if $r_{s;j} = 1$. Hence putting

$$R_j(r_{i;j}z_i, v_{i;j}) = \prod_{i=1}^{m_j} R_j^{\{i\}}(r_{i;j}z_i, v_{i;j}), \quad z_i \in S^1.$$

we have due to the commutativity of $d(z')$ and $R_j^{\{i\}}(r_{i;j}z, v_{i;j})$ ($z, z' \in S^1$)

$$(5) \quad R_j(r_{i;j}z_i, v_{i;j})d(z_1) \cdots d(z_{m_j}) = \prod_{i=1}^{m_j} R_{i;j}(r_{i;j}, z_i^i v_{i;j}) \quad (j \geq 1).$$

But in the case of $j = 0$, differing from this, we have

$$(5b) \quad R_0(r_{i;0}z_i, v_{i;0})d(z_1) \cdots d(z_{m-1}) = R_{1;0}(r_{1;0}, z_1 v_{1;0}) \prod_{i=2}^{m-1} R_{i;0}(r_{i;0}, z_{i-1}^m z_i^i v_{i;0}).$$

This can be shown using induction. In fact, as in (5), we first have

$$R_0(r_{i;0}z_i, v_{i;0})d(z_{m-1}) = \left(\prod_{s=1}^{m-2} R_{s;0}(r_{s;0}z_s, v_{s;0}) \right) R_{m-1;0}(r_{m-1;0}, z_{m-1}^{m-1} v_{m-1;0}).$$

By letting $d(z_{m-2})$ act on both sides of this it can be derived that

$$\begin{aligned}
R_0(r_{i;0}z_i, v_{i;0})d(z_{m-1})d(z_{m-2}) &= \left(\prod_{i=1}^{m-3} R_{i;0}(r_{i;0}z_i, v_{i;0}) \right) \\
&\quad R_{m-2;0}(r_{m-2;0}, z_{m-2}^{m-2} v_{m-2;0}) R_{m-1;0}(r_{m-1;0}, z_{m-2}^m z_{m-1}^{m-1} v_{m-1;0}).
\end{aligned}$$

Repeating this procedure we can achieve the equation (5b). Further these equations have the following extensions.

$$(6) \quad \begin{aligned} R_j^{\{i\}}(r_{i;j}z z', v_{i;j})d(z) &= R_j^{\{i\}}(r_{i;j}z', z^i v_{i;j}) \quad (j \geq 0), \\ R_j(r_{i;j}z_i z'_i, v_{i;j})d(z_1) \cdots d(z_{m_j}) &= \begin{cases} R_j(r_{i;j}z'_s, z'_i v_{i;j}) & (j \geq 1), \\ R_j(r_{i;j}z'_i, z_{i-1}^n z'_i v_{i;j}) & (j = 0) \end{cases} \quad (\text{where } z_{-1} = 1). \end{aligned}$$

From (4) we see that $p(R_j^{\{i\}}(r_{i;j}z, v_{i;j})) = p(R_{i;j}(r_{i;j}, z^i v_{i;j}))$ holds for $p : G \rightarrow G/S$. This permits us to embed $S_{i;j}^2$ in G/S by means of the injective map $(r_{i;j}, w_{i;j})_R \rightarrow p(R_j^{\{i\}}(r_{i;j}z, v_{i;j}))$ where $S_{i;j}^2 = S^2$ denotes the 2-sphere consisting of $(r_{i;j}, w_{i;j})_R$. From (6) we also see that

$$P_j^{\{i\}} = \{R_j^{\{i\}}(r_{i;j}z, v_{i;j}) | (r_{i;j}, w_{i;j})_R \in S^2, z \in S^1\} \subset G$$

is closed under the action of S . Combining these two facts we have that $q_{i;j} = p|P_j^{\{i\}} : P_j^{\{i\}} \rightarrow S_{i;j}^2 \subset G/S$ provides the projection map of a principal S -bundle over $S_{i;j}^2$. Then $q_{i;j}$ factors through the map $P_j^{\{i\}} \rightarrow SU(2)$ given by $R_j^{\{i\}}(r_{i;j}z, v_{i;j}) \rightarrow R(r_{i;j}z, z^{i-1}v_{i;j})$ which yields an isomorphism $(P_j^{\{i\}}, q_{i;j}, S_{i;j}^2) \cong (SU(2), p, S^2)$. Let us write $L_{i;j}$ for the canonical complex line bundle over $S_{i;j}^2$ associated to $q_{i;j}$. Then since $(S(L_{i;j}), \pi, S_{i;j}^2) \cong (P_j^{\{i\}}, q_{i;j}, S_{i;j}^2)$ we have

$$(7) \quad (S(L_{i;j}), \pi, S_{i;j}^2) \cong (SU(2), p, S^2) \quad (\text{identifying } S_{i;j}^2 = S^2).$$

Let us write $(S_j^2)^{m_j} = S_{1;j}^2 \times \cdots \times S_{m_j;j}^2$ and let $\phi_j : (S_j^2)^{m_j} \rightarrow G/S$ be the map given by $x_j = (x_{1;j}, \cdots, x_{m_j;j}) \rightarrow p(R_j(r_{i;j}, w_{i;j}))$ where $x_{i;j} = (r_{i;j}, w_{i;j})_R \in S_{i;j}^2$. Let us also write $(S^2)^{(m^2-m)/2} = (S_0^2)^{m_0} \times \cdots \times (S_{m-2}^2)^{m_{m-2}}$ and let $\phi : (S^2)^{(m^2-m)/2} \rightarrow G/S$ be the map defined by $\phi(x) = \phi_0(x_0) \cdots \phi_{m-2}(x_{m-2})$ where $x = (x_0, \cdots, x_{m-2})$ with $x_j \in (S_j^2)^{m_j}$. Let us put $R(r_{i;j}z_i, v_{i;j}) = \prod_{j=0}^{m-2} R_j(r_{i;j}z_i, v_{i;j})$. Then, as in the case of $P_j^{\{i\}}$, using (5), (5b), (6),

$$P = \{R(r_{i;j}z_i, v_{i;j}) | (r_{i;j}, w_{i;j})_R \in S_{i;j}^2, z_i \in S^1\} \subset G$$

forms the total space of a principal S -bundle endowed with the projection map $q : P \rightarrow (S^2)^{(m^2-m)/2}$ such that $\phi \circ q = p|P$. Let us write $L_j^{m_j} = L_{1;j} \boxtimes \cdots \boxtimes L_{m_j;j}$ and also put $L^{(m^2-m)/2} = L_0^{m_0} \boxtimes \cdots \boxtimes L_{m-2}^{m_{m-2}}$. Then $(S(L^{(m^2-m)/2}), \pi, (S^2)^{(m^2-m)/2}) \cong (P, q, (S^2)^{(m^2-m)/2})$ for the reason similar to (7) and therefore we have

$$(8) \quad (S(L^{(m^2-m)/2}), \pi, (S^2)^{(m^2-m)/2}) \cong \phi^*(G, p, G/S)$$

where the right hand side denotes the induced bundle of $(G, p, G/S)$ by ϕ .

Lemma 1. $\phi : (S^2)^{(m^2-m)/2} \rightarrow G/S$ is injective.

Proof. By assumption $\phi(x) = \phi_0(x_0) \cdots \phi_{m-2}(x_{m-2})$ and $\phi_j(x) = p(R_j(r_{i;j}, w_{i;j}))$. Here $R_j(r_{i;j}, w_{i;j})$ can be written in the form $R_j(r_{i;j}, w_{i;j}) = \text{diag}(I_j, M_j)$ where the first column of M_j consists of $\underline{a}_{0;j}, \underline{a}_{1;j}, \cdots, \underline{a}_{m-j-1;j}$ given in (3).

Suppose $\phi(x) = \phi(x')$, that is, $\phi_0(x_0) \cdots \phi_{m-2}(x_{m-2}) = \phi_0(x'_0) \cdots \phi_{m-2}(x'_{m-2})$. Then due to the definition this can be interpreted as meaning that

$$R_0(r_{i;0}, w_{i;0j}) \cdots R_{m-2}(r_{i;m-2}, w_{i;m-2}) = R_0(r'_{i;0}, w'_{i;0j}) \cdots R_{m-2}(r'_{i;m-2}, w'_{i;m-2})$$

where we denote by attaching $'$ to an element accompanied by x its corresponding element accompanied by x' . In view of the form of $R_j(r_{i;j}, w_{i;j})$ we therefore have

$$\underline{a}_{0;0} = \underline{a}'_{0;0}, \dots, \underline{a}_{m-1;0} = \underline{a}'_{m-1;0}.$$

Substituting the result of (3) into these equations we have $r_{1;0} \cdots r_{m-1;0} = r'_{1;0} \cdots r'_{m-1;0}$, $r_{s+1;0} \cdots r_{m-1;0} \bar{w}_{s;0} = r'_{s+1;0} \cdots r'_{m-1;0} \bar{w}'_{s;0}$ ($1 \leq s \leq m-1$) and $\bar{w}_{m-1;0} = \bar{w}'_{m-1;0}$ where $r_{s;0}^2 + |w_{s;0}|^2 = 1$ and $r_{s;0} \geq 0$. For convenience we assume that $r_{i;0} > 0$ and moreover proceed by induction on i in reverse order. First from $\underline{a}_{m-1;0} = \underline{a}'_{m-1;0}$ we have $w_{m-1;0} = w'_{m-1;0}$ and so it follows that $r_{m-1;0} = r'_{m-1;0}$. Next from the second equation $\underline{a}_{m-2;0} = \underline{a}'_{m-2;0}$ we see that $r_{m-1;0} w_{m-2;0} = r'_{m-1;0} w'_{m-2;0}$. But since $r_{m-1;0} = r'_{m-1;0} > 0$ we have $w_{m-2;0} = w'_{m-2;0}$ and so similarly it follows that $r_{m-2;0} = r'_{m-2;0}$. Repeating this procedure we obtain subsequently $w_{m-3;0} = w'_{m-3;0}$, $r_{m-3;0} = r'_{m-3;0}$, $w_{m-4;0} = w'_{m-4;0}$, $r_{m-4;0} = r'_{m-4;0}$, \dots , $r_{2;0} = r'_{2;0}$ and finally we obtain $w_{1;0} = w'_{1;0}$, $r_{1;0} = r'_{1;0}$. At this point we conclude that $x_0 = x'_0$ and so the assumption above can be rewritten as

$$\phi_1(x_1) \cdots \phi_{m-2}(x_{m-2}) = \phi_1(x'_1) \cdots \phi_{m-2}(x'_{m-2}).$$

From this by looking at the form of $R_j(r_{i;j}, w_{i;j})$ again we have $\underline{a}_{0;1} = \underline{a}'_{0;1}, \dots, \underline{a}_{m-2;1} = \underline{a}'_{m-2;1}$. Replacing the previous equations by these ones and the condition $r_{i;0} > 0$ by $r_{i;1} > 0$ in the argument there we can get $x_1 = x'_1$, hence the rewritten assumption above is also rewritten as

$$\phi_2(x_2) \cdots \phi_{m-2}(x_{m-2}) = \phi_2(x'_2) \cdots \phi_{m-2}(x'_{m-2}).$$

Repeating this step we obtain $x_2 = x'_2, \dots, x_{m-2} = x'_{m-2}$ via inductive reduction, which implies $x = x'$. This proves the lemma.

In the above, if $r_{m_j;j} = 0$, then we think of it as not being zero by exchanging the m_j th and other proper rows and columns depending on the case. In the case that there are no such row and column, this approach leads us to obtain directly $r_{i;j} = r'_{i;j} = 0$ ($1 \leq i \leq m_j$) and so $x_j = x'_j$. \square

3. PROOFS OF THEOREM AND PROPOSITION

For any $k \in \mathbb{Z}$ such that $1 \leq k \leq m/2 - 1$ we put

$$D(z_{2k-1}, z z_{2k}) = \text{diag}(1, \binom{2k-1}{\dots}, 1, z_{2k-1}, \bar{z}_{2k-1} z z_{2k}, \bar{z} \bar{z}_{2k}, 1, \dots, 1)$$

for $z_{2k-1}, z_{2k}, z \in S^1$. Let Q_k be the subspace of G consisting of $D(z_{2k-1}, z z_{2k}) d(\bar{z})$. Then it forms the total space of a principal S -bundle over $T_k = S^1 \times S^1$ along with the projective map $p_k : Q_k \rightarrow T_k$ given by $D(z_{2k-1}, z z_{2k}) d(\bar{z}) \rightarrow (z_{2k-1}, z z_{2k})$ where T_k is considered as an subspace of G/S under $\psi_k : (z_{2k-1}, z z_{2k}) \rightarrow p(D(z_{2k-1}, z z_{2k}))$.

Let $\mu : T_k \rightarrow S^2$ be the map defined by $\mu(e^{\eta i}, z e^{\theta i}) = (-\cos \mu, z e^{\theta i} \sin \mu)$ where $z_{2k-1} = e^{\eta i}$ ($0 \leq \eta \leq \pi$) and $z_{2k} = e^{\theta i}$ ($0 \leq \theta < 2\pi$). Taking into account the fact that a principal circle bundle over S^1 is trivial we see that the classifying map of p_k factors through S^2 where the restriction of p_k to $\{1\} \times S^1 \subset T_k$ is viewed as being trivial. Then we have

Lemma 2. $(Q_k, p_k, T_k) \cong \mu^*(SU(2), p, S^2)$ ($1 \leq k \leq m/2 - 1$).

Proof. In order to prove this it suffices to show that there is a bundle map covering μ . From the fact that μ can be rewritten as $\mu(e^{\eta i}, z e^{\theta i}) = (\cos(\eta/2), z e^{\theta i} \sin(\eta/2))_R$ we see

that the map $\tilde{\mu} : Q_k \rightarrow SU(2)$ given by $D(e^{\eta i}, ze^{\theta i})d(\bar{z}) \rightarrow (\bar{z} \cos(\eta/2), e^{\theta i} \sin(\eta/2))$ in fact provides the required bundle map. \square

Let $m = 2n$ or $2n+1$ and put $T^{n-1} = T_1 \times \cdots \times T_{n-1}$. Let $\phi_j : x_j \rightarrow p(R_j(r_{i;j}, w_{i;j}))$ and $\psi_k : y_k \rightarrow p(D(z_{2k-1}, \zeta_{2k}))$ be the injective maps of $(S^2)^{m_j}$ and T_k into G/S , respectively, described above where $x_j = (x_{1;j}, \dots, x_{m_j;j})$ with $x_{i;j} = (r_{i;j}, w_{i;j})_R \in S^2_{i;j}$ and $y_k = (z_{2k-1}, \zeta_{2k}) \in T_k$. Then by putting for $x = (x_0, \dots, x_{m-2})$ and $y = (y_1, \dots, y_{n-1})$

$$\psi(x, y) = \phi_0(x_0) \cdots \phi_{m-2}(x_{m-2}) \psi_1(y_1) \cdots \psi_{n-1}(y_{n-1})$$

we have a map of $(S^2)^{(m^2-m)/2} \times T^{n-1}$ into G/S . Let us put

$$\tilde{R}(r_{i;j}z_i, v_{i;j}; z_{2k-1}, z'_k z_{2k}) = \prod_{j=0}^{m-2} R_j(r_{i;j}z_i, v_{i;j}) \prod_{k=1}^{n-1} D(z_{2k-1}, z'_k z_{2k})d(\bar{z}'_k).$$

Then as in the case of P

$$\tilde{P} = \{\tilde{R}(r_{i;j}z_i, v_{i;j}; z_{2k-1}, z'_k z_{2k}) | (r_{i;j}, w_{i;j})_R \in S^2_{i;j}, z'_k, z_{2k-1}, z_{2k} \in S^1\} \subset G$$

forms the total space of a principal S -bundle endowed with the projection map $\tilde{q} : \tilde{P} \rightarrow (S^2)^{(m^2-m)/2} \times T^{n-1}$ such that $\psi \circ \tilde{q} = p|_P$. Let \tilde{L}_k be the complex line bundle over T_k associated to p_k . Write $\tilde{L}^{n-1} = \tilde{L}_1 \boxtimes \cdots \boxtimes \tilde{L}_{n-1}$ and put $E = L^{(m^2-m)/2} \boxtimes \tilde{L}^{n-1}$. Then we have $(S(E), \pi, (S^2)^{(m^2-m)/2} \times T^{n-1}) \cong (\tilde{P}, q, (S^2)^{(m^2-m)/2} \times T^{n-1})$ and therefore

$$(9) \quad (S(E), \pi, (S^2)^{(m^2-m)/2} \times T^{n-1}) \cong \psi^*(G, p, G/S).$$

Lemma 3. $\psi : (S^2)^{(m^2-m)/2} \times T^{n-1} \rightarrow G/S$ is injective.

Proof. In terms of the notation of the proof of Lemma 1 we prove that if we suppose that $\psi(x, y) = \psi(x', y')$, then it follows that $(r_{i;j}, w_{i;j})_R = (r'_{i;j}, w'_{i;j})_R$ ($0 \leq j \leq m-2$) and $(z_{2k-1}, \zeta_{2k}) = (z'_{2k-1}, \zeta'_{2k})$ ($1 \leq k \leq n-1$), the latter of which is the part added to the proof of injectivity of ϕ . The proof can be proceeded along the same lines as in the proof of the injectivity of ϕ above. But in fact there are the following changes in the use of the elements $\bar{w}_{m_j;j}$ each of which is a starting point of the proof for $\phi_j: \bar{w}_{m-1;0} \rightarrow \bar{w}_{m-1;0}$ (i.e. no change), $\bar{w}_{m-2;1} \rightarrow z_1 \bar{w}_{m-2;1}$, $\bar{w}_{m-3;2} \rightarrow (\bar{z}_1 \zeta_2) \bar{w}_{m-3;2}$ and in general $\bar{w}_{m-2k-2;2k+1} \rightarrow (z_{2k+1} \bar{\zeta}_{2k}) \bar{w}_{m-2k-2;2k+1}$, $\bar{w}_{m-2k-3;2k+2} \rightarrow (\bar{z}_{2k+1} \zeta_{2k+2}) \bar{w}_{m-2k-3;2k+2}$ for $k = 1, \dots, n-2$. Similarly substituting these elements into the equations $\underline{a}_{0;j} = \underline{a}'_{0;j}$, $\underline{a}_{1;j} = \underline{a}'_{1;j}$, \dots , $\underline{a}_{m-j-1;j} = \underline{a}'_{m-j-1;j}$ we can obtain $(x, y) = (x', y')$.

First from $\bar{w}_{m-1;0} = \bar{w}'_{m-1;0}$ we have $x_0 = x'_0$, which shows that $z_1 \bar{w}_{m-2;1} = z'_1 \bar{w}'_{m-2;1}$, so we have $z_1 r_{m-2;1} = z'_1 r'_{m-2;1}$. If $r_{m-2;1} > 0$, then from the second equation it follows that $r_{m-2;1} = r'_{m-2;1}$ and so $z_1 = z'_1$. If $r_{m-2;1} = 0$, then since $x_0 = x'_0 = (0, 1)_R$ are satisfied we get $z_1 = z'_1$. However in this case in order to move the next step we need to use the approach of exchanging rows and columns. In any case it can be concluded that $x_1 = x'_1$, $z_1 = z'_1$. Next from $(\bar{z}_1 \zeta_2) \bar{w}_{m-3;2} = (\bar{z}'_1 \zeta'_2) \bar{w}'_{m-3;2}$ we have $\zeta_2 \bar{w}_{m-3;2} = \zeta'_2 \bar{w}'_{m-3;2}$ and so it follows that $x_2 = x'_2$, $\zeta_2 = \zeta'_2$. Therefore by $(z_3 \bar{\zeta}_2) \bar{w}_{m-4;3} = (z'_3 \bar{\zeta}'_2) \bar{w}'_{m-4;3}$ we have $z_3 \bar{w}_{m-4;3} = z'_3 \bar{w}'_{m-4;3}$ and so in a similar way to the previous case we have $x_3 = x'_3$, $z_3 = z'_3$. Subsequently also by $(\bar{z}_3 \zeta_4) \bar{w}_{m-5;4} = (\bar{z}'_3 \zeta'_4) \bar{w}'_{m-5;4}$ we have $\zeta_4 \bar{w}_{m-5;4} = \zeta'_4 \bar{w}'_{m-5;4}$ and so it follows that $x_4 = x'_4$, $\zeta_4 = \zeta'_4$. At this point we can conclude that $x_1 = x'_1$, $x_2 = x'_2$, $x_3 = x'_3$, $x_4 = x'_4$ and $y_1 = y'_1$, $y_2 = y'_2$. These tell us that through repetition of these procedures we can arrive at the desired result. \square

Proof of Theorem. First we note that $\dim((S^2)^{(m^2-m)/2} \times T^{n-1}) = \dim G/S$. In general $SU(m)$ is embedded in $\mathbb{C}^{m(m-1)}$ as the subspace of $(m-1)$ -tuples (x_1, \dots, x_{m-1}) with

$x_i \in \mathbb{C}^m$, $|x_i| = 1$ and $x_i \perp x_j$ if $i \neq j$ (cf. [1] and [6]). In view of this, if we think of ψ as a map between manifolds of the same dimension $m^2 - 2$, then by taking into account its differentiability we can see from Lemma 3 that ψ is a homeomorphism and so by construction it can also be taken to be a diffeomorphism.

Identify G/S with $(S^2)^{(m^2-m)/2} \times T^{n-1}$ via ψ and write E for $L^{(m^2-m)/2} \boxtimes \tilde{L}^{n-1}$ in (9). Then combining the equations of (7) and Lemma 2 we have from Proposition 2.1 of [5]

$$e_{\mathbb{C}}([S(E), \Phi_E]) = (-1)^{n-1} B_{n^2}/2n^2$$

where Φ_E denotes the trivialization of the stable tangent space of $S(E)$ derived by the induced framing \mathcal{L}_S on G/S .

Now $D(z_{2k-1}, zz_{2k}) \in Q_k$ equals $D(z_{2k-1}, zz_{2k}) = R_{1;2k-1}(z_{2k-1}, 0)R_{1;2k}(zz_{2k}, 0)$ where $R_{1;2k-1}(z_{2k-1}, 0)$ and $R_{1;2k}(zz_{2k}, 0)$, respectively, constitute the product components of $P \subset G$ which belong to $q_{0,2k-1}^{-1}((1, 0)_R)$ and $q_{0,2k}^{-1}((1, 0)_R)$. Putting \mathcal{L} on Φ_E , from this we see that the restrictions of \mathcal{L} to Q_k and P overlap and therefore the restriction of \mathcal{L} to $T_k \subset G$ must be stably trivial. This tells us that if we let ρ denote the realification of the standard complex representation $G \hookrightarrow GL(m, \mathbb{C})$, then Φ_E must be stably equivalent to $\mathcal{L}^{(n-1)\rho}$. Substituting this into the equation above we have the desired result. \square

Proof of Proposition. Let $m = 2n + 1$. Let C be as above. Then clearly $S \cap C = \{I_{2n+1}\}$, so $S \times C$ becomes a subgroup of G . Letting $r : G \rightarrow G/C$ and $\tilde{r} : G/C \rightarrow G/(S \times C)$ be the quotient maps, the pair (r, \tilde{r}) gives a bundle map between principal S -bundles $G \rightarrow G/S$ and $G/C \rightarrow G/(S \times C)$. Consider the composition $r \circ \iota : \tilde{P} \hookrightarrow G \rightarrow G/C$ and $\tilde{r} \circ \psi : (S^2)^{(m^2-m)/2} \times T^{n-1} \rightarrow G/S \rightarrow G/(S \times C)$ where ι denotes the inclusion of \tilde{P} into G . Then clearly $(\tilde{r} \circ \psi) \circ \tilde{q} = \tilde{r} \circ (r \circ \iota)$ holds and $\dim((S^2)^{(m^2-m)/2} \times T^{n-1}) = \dim G/(S \times C)$. Following the proof of Lemma 3 we can see that $\tilde{r} \circ \psi$ is injective and so it becomes a homeomorphism. Therefore by replacing ψ by $\tilde{r} \circ \psi$ in the proof of the theorem above we can get a modified form in the case of $m = 2n + 1$. This proves the proposition. \square

REFERENCES

- [1] L. Astey, M. A. Guest and G. Pastor, *Lie groups as framed boundaries*, Osaka J. Math. **25** (1988), 891–907.
- [2] J. F. Adams, *On the groups $J(X)$ –IV*, Topology **5** (1966), 21–71.
- [3] M. F. Atiyah and L. Smith, *Compact Lie groups and the stable homotopy of spheres*, Topology **13** (1974), 135–142.
- [4] K. Knapp, *Rank and Adams filtration of a Lie group*, Topology **17** (1978), 41–52.
- [5] P. Löffler and L. Smith, *Line bundles over framed manifolds*, Math. Z. **138** (1974), 35–52.
- [6] B. Steer, *Orbits and the homotopy class of a compactification of a classical map*, Topology **15** (1976), 383–393.

H. MINAMI: PROFESSOR EMERITUS, NARA UNIVERSITY OF EDUCATION
Email address: hminami@camel.plala.or.jp