A Trifecta of Modelling Tools: Navigating COMPAS, Shark and Bayesian Inference for Binary Black Hole Model Selection

Liana Rauf^{1,2*}, Cullan Howlett^{1,2}, Simon Stevenson^{2,3}, Jeff Riley^{2,4}

¹*School of Mathematics and Physics, The University of Queensland, Brisbane, QLD 4072, Australia*

³*Centre for Astrophysics and Supercomputing, Swinburne University of Technology, Mail H39, PO Box 218, VIC 3122, Australia*

⁴*School of Physics and Astronomy, Monash University, Clayton VIC 3800, Australia*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

Gravitational waves (GWs) have revealed surprising properties of binary black hole (BBH) populations, but there is still mystery surrounding how these compact objects evolve. We apply Bayesian inference and an efficient method to calculate the BBH merger rates in the Shark host galaxies, to determine the combination of COMPAS parameters that outputs a population most like the GW sources from the LVK transient catalogue. For our COMPAS models, we calculate the likelihood with and without the dependence on the predicted number of BBH merger events. We find strong correlations between hyper-parameters governing the specific angular momentum (AM) of mass lost during mass transfer, the mass-loss rates of Wolf–Rayet stars via winds and the chemically homogeneous evolution (CHE) formation channel. For the AM loss, we also find opposing trends in the likelihood with and without the dependence on the number of GW events. We conclude that analysing the marginalised *and* unmarginalised likelihood is a good indicator of whether the population parameters distribution and number of observed events reflect the LVK data. In doing so, we see that the majority of the models preferred in terms of the population-level parameters of the BBHs greatly overpredict the number of events we should have observed to date. Looking at the smaller number of models which perform well with both likelihoods, we find that those with no CHE, AM loss occurring closer to the donor during the first mass-transfer event, and/or higher rates of mass-loss from Wolf–Rayet winds are generally preferred by current data. We find these conclusions to be robust to our choice of selection criteria. However, we do identify several notable events that all of our models struggle to reproduce, and that, even when incorporating selection effects into our models to match our selection of LVK data, COMPAS tends to underpredict the proportion of events with chirp mass > $50M_{\odot}$ and $z > 0.5$, leaving avenues for future investigations.

Key words: gravitational waves – black hole physics – stars: evolution – galaxies: star formation – methods: analytical – methods: numerical – methods: statistical

1 INTRODUCTION

The detection of gravitational waves (GWs) have prompted the astrophysical community to revise the models for binary black hole (BBH) formation and evolution. There are currently two main formation channels proposed:

(i) Isolated: The formation of merging BBHs through the evolution of isolated binaries. These stars are gravitationally bound since birth and are not perturbed by other stars or compact objects.

(ii) Dynamical: BBH mergers occurring in dense environments via three-body encounters followed by energy exchange and hardening of the binary. Some examples of dense environments are young star, nuclear or globular clusters, or in accretion disks of the central supermassive BHs in galaxies.

GWs can be used to gain insight into the poorly constrained aspects of massive stellar binary evolution and help in distinguishing between binary formation channels. The detailed origins and formation channels for binary systems are still unknown, particularly what fraction of these systems come from isolated and dynamical channels.

There are also aspects of stellar evolution that are poorly understood, such as the common envelope phase, mass transfer efficiency and angular momentum loss. This information is critical for understanding the population of BBHs observed by LIGO, Virgo and KAGRA (LVK), including their mass and redshift distributions, and to inform us on what is required for future detectors [\(Mapelli](#page-21-0) [2019,](#page-21-0) [2020;](#page-21-1) [Mandel & Broekgaarden](#page-21-2) [2022;](#page-21-2) [Arca Sedda et al.](#page-20-0) [2023\)](#page-20-0). Com-bining the data from GWTC-1 [\(Abbott et al.](#page-20-1) [2019a\)](#page-20-1), GWTC-2 [\(Ab](#page-20-2)[bott et al.](#page-20-2) [2021a\)](#page-20-2) and GWTC-3 [\(Abbott et al.](#page-20-3) [2023b\)](#page-20-3), there is a total of 90 confident ($p_{astro} > 0.5$) GW event candidates observed to date, including 85 BBHs, 2 binary neutron stars (BNSs), and 3 black-holeneutron-stars (BHNSs). This growing population of GWs, including those to be released from the ongoing fourth LVK observing run (O4), provides key information on the source masses, spins and local merger rates [\(Abbott et al.](#page-20-4) [2023a\)](#page-20-4) of binary compact objects, which can be used to answer some of these questions.

²*OzGrav: The ARC Centre of Excellence for Gravitational Wave Discovery, Australia*

[★] E-mail: l.rauf@uq.net.au

However, there are many formation pathways that could lead to the same individual source properties. Simulations of ensembles of events, produced by varying the physical assumptions, can provide better insight into how these binary systems evolve and merge over time. When combined with the full set of data observed to date, they can potentially break the degeneracy between uncertainties in stellar evolution parameters and BBH populations. They also enable us to produce merger rates that can be compared to observations. [Mandel](#page-21-3) [& Broekgaarden](#page-21-3) [\(2021\)](#page-21-3) provide a comprehensive review of the local merger rates for the isolated and dynamical formation channels. The studies in the review combine population synthesis codes with different assumptions for the metallicity dependent star formation rate (SFR) density, which plays a crucial role in the evolution of massive binaries [\(Neijssel et al.](#page-21-4) [2019;](#page-21-4) [Chruślińska](#page-21-5) [2024\)](#page-21-5). [Broekgaarden et al.](#page-21-6) [\(2022a\)](#page-21-6) utilises COMPAS [\(Stevenson et al.](#page-22-0) [2017;](#page-22-0) [Vigna-Gómez et al.](#page-22-1) [2018;](#page-22-1) [Team COMPAS: Riley, J. et al.](#page-22-2) [2022\)](#page-22-2) to explore the impact of stellar evolution parameters and metallicity specific star formation, $S(z, Z)$, on the merger rate, chirp mass and coalescence time distributions. They found that uncertainties in the binary evolution models only led to variations in the BBH merger rate of up to a factor 2, with the BBH merger rates most sensitive to $S(z, Z)$. [Santoliquido et al.](#page-22-3) [\(2021\)](#page-22-3) and [Neijssel et al.](#page-21-4) [\(2019\)](#page-21-4) also come to similar conclusions. Many of these studies choose to explore the variations in binary evolution models over a grid of metallicities, and generate populations for each grid point. Some studies have shown that population III stars can also contribute to the total BBH merger rate density [\(San](#page-22-4)[toliquido et al.](#page-22-4) [2024;](#page-22-4) [Tanikawa](#page-22-5) [2024\)](#page-22-5). These studies emphasise the need to further study the assumptions in binary evolution models to understand the impact on the merger rate, specifically the *combined* impact of model variations [\(Stevenson & Clarke](#page-22-6) [2022\)](#page-22-6).

There have also been various studies proposing methods for constraining different binary evolution models. [Wong et al.](#page-22-7) [\(2023\)](#page-22-7) applies "backward" modelling of each GW event to the progenitors to avoid fixed astrophysical assumptions, by allowing variations in the hyper-parameters. However, they note that their methods become less efficient as the dimensionality of the parameter space increases. [Bar](#page-20-5)[rett et al.](#page-20-5) [\(2018\)](#page-20-5) uses Fisher information matrices to quantitatively analyse how the observed BBH population will constrain the COMPAS models, given the chirp mass distribution and accounting for the metallicity specific cosmic SFR, selection effects and measurement uncertainty. [Riley & Mandel](#page-22-8) [\(2023\)](#page-22-8) uses artificial neural networks to generate an interpolant to calculate the merger detection rates for double compact objects for varying prescriptions of the metallicity specific SFR. This is used to create GW populations for each model, where the chirp mass and redshift distribution are compared using Bayesian analysis to the LVK data. [Delfavero et al.](#page-21-7) [\(2023\)](#page-21-7) calculates a joint likelihood for binary evolution models. They fit normal distribution fits to the likelihood grids and implement Gaussian process regression to interpolate between models, the detected number of events and the likelihood. This is also used to quantify the systematic uncertainty in the parameters. However, they emphasise that the inference on a limited subspace of models could be biased. These studies highlight the high computational costs for multi-dimensional parameter studies. However, as more observations are made, the uncertain astrophysics will produce correlated impacts in the binary population, so there is a crucial need for more parameters covering a large range in model inference. Realistically, all formation channels (within the isolated and dynamical channels) must be considered to obtain a parameter space representative of the true BBH population [\(Zevin et al.](#page-22-9) [2021;](#page-22-9) [Cheng et al.](#page-21-8) [2023a\)](#page-21-8)

To build on this body of work, we develop here a new method of combining a more accurate, non-parametric star-formation and

MNRAS **000**, [1](#page-0-0)[–23](#page-26-0) (0000)

metallicity history of the Universe with a wider range of different massive binary evolution models. We then use these models alongside data from GWTC-1 [\(Abbott et al.](#page-20-1) [2019a\)](#page-20-1), GWTC-2 [\(Abbott](#page-20-2) [et al.](#page-20-2) [2021a\)](#page-20-2) and GWTC-3 [\(Abbott et al.](#page-20-3) [2023b\)](#page-20-3) to identify the most likely avenues for generating binary compact objects, and to rule out those that are unable to produce the distributions of objects seen to date. We do this by improving on [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10), where we combined runs from the COMPAS rapid population synthesis code, for formation and evolution of BBH formed through isolated binary evolution, with a star formation rate and metallicity history for the observed host galaxies predicted by the semi-analytic galaxy evolution code Shark [\(Lagos et al.](#page-21-9) [2018,](#page-21-9) [2019\)](#page-21-10). In our original work, we predicted (among other things) the BBH merger rate as a function of redshift for four COMPAS runs, where we varied the remnant mass prescription. We found the merger rate ranged over several magnitudes and we concluded that the [Mandel & Müller](#page-21-11) (2020) remnant mass prescription was the best fit out of the four models to the local merger rate currently predicted using the BBH mergers from GWTC-3. However, we also found that cross-matching between the SHARK galaxies and the COMPAS population was prohibitively time-consuming for exploring larger portions of our parameter space. Since we explored only rates, we were also not able to confidently rule out any models that do not align with the GW population-level parameters, and did not test the impact of varying sub-parameters of these models such as those covering other aspects of binary evolution rather than just the mass of the remnant compact objects.

So, in this paper, we substantially extend on our work in [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) and explore an alternative and efficient method of predicting the BBH merger rate and population level parameters such as the binary masses and redshift distributions. We do so by rescaling a high-fidelity single run of our original simulation, varying some of the most relevant COMPAS input parameters. This allows us to test in this work > 300 different models for population synthesis, varying ingredients/model assumptions such as the common envelope efficiency, mass transfer accretion efficiency, mass-loss rates of Wolf–Rayet stars, chemically homogeneous evolution model and specific angular momentum loss. For each of these we evaluate the ability of the model to reproduce the observed distribution of black hole masses and redshifts, using a Bayesian framework to account for selection effects. We also explore the impact on our findings with and without including the merger rate as a fitting parameter (i.e., including a comparison of the model-predicted and observed merger rates in evaluating our model posteriors). Compared to previous studies this work also has the improvement of incorporating a basic estimate of sampling error in the simulated model merger rates and posteriors. This allows us to set a threshold for what we consider to be tangible differences in model performance, and also demonstrates more generally the importance of further investigation into the uncertainties in the typical 'simulation-based inference' used in GW population studies.

This paper is structured as follows. In Section [2,](#page-2-0) we summarise recent progress in our understanding of various massive binary evolution models and parameters, motivating the range of models we explore in this work. In Section [3,](#page-6-0) we show that starting with a fiducial model, where the merger rate in a set of simulated galaxies has been evaluated by carefully cross-matching between COMPAS and SHARK, we can rescale the simulation to produce the time and metallicity dependent GW merger rate for another model where the COMPAS parameters have been altered. In Section [4,](#page-8-0) we implement Bayesian inference to determine which of these COMPAS runs output a BBH population that best matches the current observed population. We do so by comparing the chirp mass, mass ratio and redshift distributions

from each model with the LVK posterior sample distributions. We then validate the total number of theoretical and detected GW events expected for the models and the models posteriors by comparing to the values obtained using the four 'brute-force' evaluations from [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10). We repeat the method for a wide range of new COMPAS runs in Section [5,](#page-12-0) identifying the best and worst fit models to the data and studying the impact of different selection effects and analysis choices on the conclusions we can draw about stellar evolution. We conclude in Section 6 . In this work, we use the SHARK cosmology H_0 , $\Omega_m = \{67.8, 0.308\}$ where necessary.

2 COMPAS**: OVERVIEW OF MODELS**

In the following section, we detail the physics and the implementation in COMPAS of the common envelope, mass transfer, mass loss via winds, angular momentum loss and formation channels such as the isolated and chemically homogeneous evolution. We review the literature on the assumptions and parameters associated with these processes that can impact the evolution of the binary stars and, hence, the number and source properties of GWs for each COMPAS model. We conclude with a summary of the big questions surrounding these prescriptions and their limitations in the context of BBH population modelling.

2.1 Isolated formation channels

In this paper, we focus on the isolated formation channel. This is the most commonly studied channel. It involves massive stars formed on the zero-age-main-sequence (ZAMS). We describe, one of the formation channels, the classical isolated binary evolution via a common envelope phase as follows:

(i) The primary star in the binary reaches the end of main sequence (MS) evolution. Hydrogen burning in the core causes the outer stellar envelope to expand past the Roche lobe and transfer mass to the secondary star.

(ii) Mass transfer from the primary stops and leaves a Wolf-Rayet star. This then reaches the end of its life, undergoes a supernova and forms a compact object. The system is now an X-ray binary.

(iii) The secondary evolves and transfers mass back to the primary (compact) object. This process is dynamically unstable and forms a common envelope, which encompasses the compact object.

(iv) Orbital energy released due to the compact object rapidly in-spiralling can remove the common envelope.

(v) The secondary star evolves and eventually also forms a compact object.

(vi) The double compact object (DCO) system slowly in-spirals due to orbital energy loss via GWs.

However, there are many stages within this formation channel where the binary may not survive, or the events can subtly differ from the above prescription leading to dramatically different outcomes. The system may not survive the first supernova explosion or the common envelope may be not be ejected resulting in the compact object merging with the secondary core before a DCO is formed. Alternatively, the second mass transfer could be stable and avoid the common envelope (e.g., [van den Heuvel et al.](#page-22-11) [2017;](#page-22-11) [Neijssel et al.](#page-21-4) [2019;](#page-21-4) [van Son et al.](#page-22-12) [2022a;](#page-22-12) [Gallegos-Garcia et al.](#page-21-12) [2021;](#page-21-12) [Marchant](#page-21-13) [et al.](#page-21-13) [2021\)](#page-21-13). The wind mass-loss rates and remnant prescription can also impact the orbital separation, as well as the final masses. There are various circumstances and factors that can affect the separation of DCO systems, which is strongly correlated with the delay time and

the merger rate. On one hand, this makes GW sources unique probes of cosmic history [\(Neijssel et al.](#page-21-4) [2019;](#page-21-4) [Mapelli](#page-21-14) [2021;](#page-21-14) [Mandel &](#page-21-15) [Farmer](#page-21-15) [2022;](#page-21-15) [Costa et al.](#page-21-16) [2023\)](#page-21-16), but on the other, leads to difficulties in interpreting or identifying the most important/critical points in the evolution of the system.

As an example, [Stevenson et al.](#page-22-0) [\(2017\)](#page-22-0) used COMPAS to show that the first three GW events detected by LIGO: GW150914, GW151226 and GW151012, can be explained by the classical isolated channel with mass transfer and common envelope phase, and characterised their progenitor metallicity and masses. However, [Dorozsmai & Too](#page-21-17)[nen](#page-21-17) [\(2022\)](#page-21-17) found that the stable channel, consisting of two stable mass transfer phases, dominates the GW sources — formation of GW sources via a radiative common envelope channel is not possible for low mass transfer efficiency ($f_{\text{MT}} \leq 0.3$) due to the mass ratio being below the critical mass ratio ($q_{\text{crit}} = M_d / M_a$, where M_a is the acretor mass and M_d is the donor mass). [Picco et al.](#page-21-18) [\(2023\)](#page-21-18) similarly found that stable mass transfer, where mass and angular momentum is lost through the L2 point, is a possible formation channel for BBHs. Comparing the two channels, [van Son et al.](#page-22-12) [\(2022a\)](#page-22-12) found that the common envelope channel tends to dominate BBH mergers with primary masses < $30M_{\odot}$ binaries with short delay times (< 1Gyr). They also found that the common envelope phase is rare for massive stars due to heavy mass loss preventing them from filling their Roche lobe. As such, the stable mass transfer channel, which widens the binary orbit as mass is lost from the system during secondary mass transfer, forms primary BH masses > $30M_{\odot}$ and systems with long delay times $($ > 1Gyr).

2.1.1 Common envelope evolution (α_{CE}) and Mass transfer $efficiency(f_{MT})$

Given the above, in the formation of isolated binary stars, uncertainties remain in the common envelope (CE) phase and mass transfer process, which play a key role in the final component and orbital properties.

The mass transfer efficiency is the fraction of mass lost by the donor star and accreted by the companion star. [Team COMPAS: Riley, J.](#page-22-2) [et al.](#page-22-2) [\(2022\)](#page-22-2) uses a simplified approach for the mass transfer. They parameterise the fraction of mass lost by the donor to the accretor with f_{MT} ,

$$
\dot{M}_a = -f_{\rm MT} \dot{M}_d,\tag{1}
$$

with $0 \le f_{\text{MT}} \le 1$. This fraction depends on the properties of the binary prior to mass transfer and varies over time. We use a fixed fraction prescription in COMPAS, so the fraction of mass accreted is assumed to be constant over the mass transfer phase. $f_{\text{MT}} = 1$ corresponds to fully conservative mass transfer, so all the mass lost from the donor is accreted. $f_{\text{MT}} < 1$ is non-conservative and $f_{\text{MT}} = 0$ is fully non-conservative. In the default setting, the mass transfer efficiency is determined by the ratio of the Kelvin–Helmholtz thermal timescale of the two stars,

$$
\tau_{\rm KH} = \frac{E_{\rm int}}{L}.\tag{2}
$$

This is the time taken for the star's internal energy, E_{int} , to be radiated at it's current luminosity, L . The maximum accretion rate onto a stellar accretor is,

$$
\dot{M}_a = \frac{CM_a}{\tau_{\text{KH,a}}},\tag{3}
$$

where the default factor $C = 10$ is assumed to account for the expansion of the star due to mass loss [\(Paczyński & Sienkiewicz](#page-21-19) [1972;](#page-21-19) [Neo et al.](#page-21-20) [1977;](#page-21-20) [Hurley et al.](#page-21-21) [2002;](#page-21-21) [Schneider et al.](#page-22-13) [2015\)](#page-22-13). This is calculated for every mass transfer event and is non-constant. Accretion onto the compact objects, such as black holes and neutron stars, is assumed to be Eddington limited such that,

$$
\dot{M}_a = \dot{M}_{\text{edd}} = 1.5 \times 10^{-8} \frac{R_*}{10 \text{km}} \frac{M_\odot}{\text{yr}},\tag{4}
$$

where the black hole radius, R_{\ast} , is the Schwarzschild radius. In the unstable mass transfer scenario, for simplicity, the CE phase is parameterised by COMPAS in the context of energy balance with the α_{CE} formalism [\(Xu & Li](#page-22-14) [2010\)](#page-22-14). The CE efficiency, α_{CE} , describes the efficiency of converting the orbital energy into kinetic energy to unbind the common envelope. This unbinding energy is defined as,

$$
E_{\text{bind}} = \alpha_{\text{CE}} \left(-\frac{GM_d M_2}{2a_i} + \frac{GM_{\text{core}} M_2}{2a_f} \right),\tag{5}
$$

where G is the gravitational constant, M_d is the mass of the pre-CE donor, M_2 is the mass of the secondary, M_{core} is the mass of the donor's core, a_i and a_f are the pre- and post-CE orbital separations. Increasing α_{CE} increases the unbinding energy, so the CE can be ejected more efficiently at wider binary separations. Too small values of α_{CE} would require the binary to shrink in order to obtain enough unbinding energy [\(Tutukov & Yungelson](#page-22-15) [1979;](#page-22-15) [Webbink](#page-22-16) [1984;](#page-22-16) [Ivanova et al.](#page-21-22) [2013;](#page-21-22) [Hirai & Mandel](#page-21-23) [2022;](#page-21-23) [Stevenson & Clarke](#page-22-6) [2022\)](#page-22-6). [de Kool](#page-22-17) [\(1990\)](#page-22-17) proposed an alternative version of the binding energy, introducing a new parameter, λ , to characterise the central concentration of the donor's envelope,

$$
E_{\text{bind}} = -\frac{GM_1 M_{\text{env}}}{\lambda a_i r_L},\tag{6}
$$

where $M_{\text{env}} = M_1 - M_{\text{core}}$ is the mass of the envelope and $r_L = R_L/a_i$ is the ratio of the Roche lobe radius and pre-CE orbital separation. The post-CE orbital separation is determined by substituting Eq [6](#page-3-0) into Eq [5,](#page-3-1)

$$
\frac{a_f}{a_i} = \frac{M_{\text{core}}M_2}{M_1} \frac{1}{M_2 + 2M_{\text{env}}/\alpha_{\text{CE}}\lambda r_L}.
$$
\n(7)

Due to lack of understanding and convenience in population synthesis calculations, α_{CE} and λ are set to constants, but *should* be varied depending the stellar and binary parameters ($Xu \& Li 2010$).

Progenitors of GW sources provide an alternative path for comparison between models and observations. Zuo $\&$ Li [\(2014\)](#page-22-18) uses the binary population synthesis code by [Hurley et al.](#page-21-24) [\(2000\)](#page-21-24) and [Hurley](#page-21-21) [et al.](#page-21-21) [\(2002\)](#page-21-21) to study high-mass X-ray binaries (HMXB) and the correlation between their apparent luminosity and displacement from star clusters. A high value of $\alpha_{\text{CE}} = 0.8 - 1.0$ is preferred when comparing to observations by [Kaaret et al.](#page-21-25) [\(2004\)](#page-21-25). However, the ambiguity in the definition of the donor core mass boundary introduces a large uncertainty (almost two orders of magnitude), which translates directly to the expected value of α_{CE} . [Romero-Shaw et al.](#page-22-19) [\(2023\)](#page-22-19) uses COMPAS to study black hole high-mass X-ray binaries (BH-HMXB). They found for the case where $f_{\text{MT}} \leq 0.1$ (and $f_{\text{WR}} = 0.1$, explained in Section [2.1.2\)](#page-3-2), that 8% of BH-HMXBs end up as BBHs or BHNSs. [de Mink et al.](#page-22-20) [\(2007\)](#page-22-20) attempts to fit the mass transfer efficiency fraction for massive close binaries. They find a wide spread in $f_{\rm MT}$ and conclude that a single value is not valid. Therefore, more electromagnetic observations are required to constrain modelling degeneracies of massive binaries.

There have been many studies on the mass transfer and common envelope phase. Many of these have explored how varying these parameters affect the BBH merger rate, the BH mass distribution range and which binary formation channels dominate these BBH mergers [\(Santoliquido et al.](#page-22-3) [2021;](#page-22-3) [Grichener](#page-21-26) [2023;](#page-21-26) [van Son et al.](#page-22-21)

[2020,](#page-22-21) [2022a](#page-22-12)[,b;](#page-22-22) [Marchant et al.](#page-21-13) [2021\)](#page-21-13). To infer the formation channels of BBH mergers, many studies further explore the connection between the models and data [\(Cheng et al.](#page-21-27) [2023b\)](#page-21-27). [Bouffanais et al.](#page-21-28) [\(2021\)](#page-21-28) find the preferred parameter range for the common envelope and mass transfer efficiency is [4, 7] and [0.7, 0.8] respectively. They also showed that increasing the common envelope efficiency increases the uncertainty on the local BBH merger rate density. However studies such as [Bavera et al.](#page-20-6) [\(2021\)](#page-20-6)[,Klencki et al.](#page-21-29) [\(2021\)](#page-21-29) and [Cui](#page-21-30) [& Li](#page-21-30) [\(2023\)](#page-21-30) find there is a preference for lower α_{CE} (0.2 < α_{CE} < 1, $\alpha_{\text{CE}} \leq 0.7$ and $\alpha_{\text{CE}} = 0.5$, 1 respectively). [Klencki et al.](#page-21-29) [\(2021\)](#page-21-29) further questions the validity of the CE channel for producing massive BHs. They find that successful ejection of the CE is possible if the donor is a massive convective-envelope giant.

We briefly mention the various studies that explore the uncertainties in parameters. But another uncertainty to note is in how these parameters are implemented in population synthesis codes. [Gallegos-Garcia et al.](#page-21-12) [\(2021\)](#page-21-12) highlights the uncertainty in stellar evolution tracks by comparing the mass transfer and common envelope treatments in the population synthesis codes MESA and COSMIC. The COSMIC models primarily form BBH mergers through the common envelope phase, while the stable mass transfer formation channel is preferred with MESA. Increasing α_{CE} also increases the variation in the BBH merger rates between the models, due to variations in merger times. [Agrawal et al.](#page-20-7) [\(2022a\)](#page-20-7) and [Romagnolo et al.](#page-22-23) [\(2023\)](#page-22-23) have compared population synthesis codes to explore the uncertainty in radial expansion of massive stars and the variation in the remnant mass.

2.1.2 Wolf–Rayet Winds (f_{WR})

Stellar winds are outflows of gas from the atmosphere of a star. Wolf–Rayet stars are evolved stars with high mass loss rates, where the hydrogen envelope of the stars are removed. To account for uncertainties in the mass-loss rates of Wolf–Rayet stars, in COMPAS, the Wolf–Rayet multiplier, f_{WR} , is used to parameterise the mass loss rate [\(Belczynski et al.](#page-21-31) [2010\)](#page-21-31),

$$
\frac{dM}{dt} = f_{\rm WR} \times 10^{-13} L^{1.5} \left(\frac{Z}{Z_{\odot}}\right)^{m} M_{\odot} \text{yr}^{-1},\tag{8}
$$

where M is the mass of the helium star, L is the luminosity, Z is the metallicity, $Z_{\odot} = 0.014$ [\(Asplund et al.](#page-20-8) [2009\)](#page-20-8) and $m = 0.86$ [\(Vink](#page-22-24) [& de Koter](#page-22-24) [2005\)](#page-22-24). Wolf–Rayet stars have been studied to understand their explosion mechanisms and determine if they are progenitors of gamma ray bursts [\(Detmers et al.](#page-21-32) [2008\)](#page-21-32). More recently, Wolf–Rayet stars have been used to study GW sources [\(Vink & Harries](#page-22-25) [2017;](#page-22-25) [Bogomazov et al.](#page-21-33) [2018\)](#page-21-33). Decreasing f_{WR} leads to an increase in the BH masses [\(Dorozsmai & Toonen](#page-21-17) [2022;](#page-21-17) [Romero-Shaw et al.](#page-22-19) [2023\)](#page-22-19).

HMXBs are usually ideal candidates to study and constrain the effects of stellar mass loss in interacting binaries, as radiation driven winds are expected for giant stars. [Schrøder et al.](#page-22-26) [\(2021\)](#page-22-26) finds that the orbital evolution of HMXBs depends on the wind velocity and mass ratio. Slow winds and high mass objects results in the binary shrinking, due to drag forces. In the case where there is no drag, the binary separation widens. GWs can provide an indirect approach to study the separation of binaries, as their progenitors include HMXBs. However, [Liotine et al.](#page-21-34) [\(2023\)](#page-21-34) found that no currently observed HMXB is predicted to form a merging BBH with high probability. They use COSMIC [\(Breivik et al.](#page-21-35) [2020\)](#page-21-35) to conclude that the BHs in the detectable HMXBs are not likely to have masses > $35M_{\odot}$, the detectable HMXBs will not merge as BBHs within the Hubble time and the distribution of redshifts, masses and metallicities of detectable HMXBs and BBHs are different. [Fishbach & Kalogera](#page-21-36) [\(2022\)](#page-21-36) argue

that discrepancies in the BH masses in BBHs and HMXBs are due to GW observational selection effects.

[Neijssel et al.](#page-21-37) [\(2021\)](#page-21-37) and [Romero-Shaw et al.](#page-22-19) [\(2023\)](#page-22-19) find a preference for low f_{WR} (< 0.2) for HMXB systems such as Cygnus X-1 [\(Miller-Jones et al.](#page-21-38) [2021\)](#page-21-38). [Stevenson & Clarke](#page-22-6) [\(2022\)](#page-22-6) explores the correlated impact of model parameters on the merger rate. They identified a correlation between the Wolf-Rayet factor and metallicity, as increasing both parameters increases mass loss via stellar winds. They found most models that match the current observations have $f_{\text{WR}} > 1$. This contradicts the previously mentioned studies due to COMPAS over-predicting the rates and masses of BBHs formed through chemically homogeneous evolution (see Section [2.2\)](#page-4-0).

[Riley et al.](#page-22-27) [\(2021\)](#page-22-27) studies the CHE binaries as a formation channel for various f_{WR} with COMPAS. They found a correlation between the delay time and f_{WR} , as increasing mass loss leads to wider binaries. This results in the merger rate decreasing with increasing f_{WR} . For $f_{\text{WR}} = 1$, the CHE channel contributes 20 Gpc⁻³yr⁻¹ to the total rate of 50 Gpc⁻³yr⁻¹, which is an overestimation compared to GWTC-3. Due to the short delay times, the BBH merger rate peaks at higher redshifts ($z \approx 3 - 4$) compared to the peak SFR.

2.1.3 Specific angular momentum loss (f_γ)

Mass transfer from a massive donor to a lower-mass accretor adds specific angular momentum to the transferred mass, so conservation of angular momentum requires that binary's orbit shrinks. If mass transfer is non-conservative, angular momentum, $J_{\rm orb}$, is carried away by mass lost from the system. The change in angular momentum is parameterised by γ ,

$$
\frac{j_{\text{orb}}}{\dot{M}_d} = \gamma \frac{J_{\text{orb}}}{M_d}.
$$
\n(9)

The orbital evolution of the binary is given by,

$$
\frac{\dot{a}}{a} = -2\frac{\dot{M}_d}{M_d} \left[1 - f_{\text{MT}} \frac{M_d}{M_a} - (1 - f_{\text{MT}}) \left(\gamma + \frac{1}{2} \right) \frac{M_d}{M_d + M_a} \right]
$$
\n
$$
= -2\frac{\dot{M}_d}{M_d} \left[1 - \frac{f_{\text{MT}}}{q} - (1 - f_{\text{MT}}) \left(\gamma + \frac{1}{2} \right) \frac{1}{1 + q} \right].
$$
\n(10)

[Willcox et al.](#page-22-28) [\(2023\)](#page-22-28) introduces a new parametrisation, a_{γ} ,

$$
a_{\gamma} = a_{\text{acc}} + f_{\gamma}(a_{\text{L2}} - a_{\text{acc}}),\tag{11}
$$

which is the effective decoupling radius between the accretor, a_{acc} , and the L2 point, a_{L2} . COMPAS default model assumes isotropic re-emission of matter during non-conservative stable mass transfer $(f_{\gamma} = 0)$. In this model, mass lost from the donor is transported near the accretor and ejected via fast isotropic winds. Hydrodynamical simulations indicate that this mass is lost between the accretor and L2 point [\(MacLeod et al.](#page-21-39) [2018a](#page-21-39)[,b\)](#page-21-40). f_{γ} determines the point where this angular momentum is lost. If the donor star is growing rapidly with respect to its Roche Lobe, the system will lose mass through the outer Lagrangian point (L2), which causes the orbit to shrink. This corresponds to $f_{\gamma} = 1$. Angular momentum loss directly from the accretor corresponds to $f_{\gamma} = 0$ [\(Costa et al.](#page-21-16) [2023;](#page-21-16) [Romero-Shaw](#page-22-19) [et al.](#page-22-19) [2023\)](#page-22-19).

[Lu et al.](#page-21-41) [\(2023\)](#page-21-41) and [Marchant et al.](#page-21-13) [\(2021\)](#page-21-13) found that their models predicted high mass transfer rates, which lead to mass loss through the L2 point. [MacLeod et al.](#page-21-39) [\(2018a\)](#page-21-39); [MacLeod & Loeb](#page-21-42) [\(2020\)](#page-21-42) argue that the specific angular momentum is ejected between the accretor and the L2 point. [Willcox et al.](#page-22-28) [\(2023\)](#page-22-28) uses COMPAS to study the stable mass transfer channel. They found that increasing f_{γ} leads to a reduction in the number of systems that undergo stable mass transfer. Increasing $f_{\rm MT}$ while decreasing f_{γ} increases the likelihood of stable mass transfer. For $f_{\gamma} \geq 0.7$, nearly all systems experience common envelope evolution.

[van Son et al.](#page-22-22) [\(2022b\)](#page-22-22) studies the impact of f_{γ} on the local BBH merger rate and the primary BH mass distribution. for $f_{\gamma} = 0.75$ and $f_y = 1$, there are no BBHs formed through the stable mass transfer channel. The merger rate drops from $141 \text{Gpc}^{-3} \text{yr}^{-1}$ with $f_{\gamma} = 0.25$ to 16Gpc⁻³yr⁻¹ with $f_{\gamma} = 0.5$, which implies the stable mass transfer contributes strongly to the merger rate for f_{γ} < 0.5. As the low mass end of the primary BH mass distribution is dominated by the stable mass transfer channel, as f_{γ} increases, the minimum primary BH mass increases.

2.2 Chemically homogeneous evolution

In the chemically homogeneous evolution (CHE) channel, the stars do not expand. They are metal poor and rapidly rotating in tight binaries. The rotation efficiently transports hydrogen to the core and helium to the envelope. This evolves and forms a Wolf–Rayet star, which contracts rather than expands. Wind-driven mass loss does not widen the binary significantly for low metallicity systems and there is no mass transfer, with the exception of massive over-contact binaries [\(Marchant et al.](#page-21-43) [2016\)](#page-21-43). Highly spinning DCOs would produce GWs in the age of the Universe [\(Nathaniel](#page-21-44) [2017;](#page-21-44) [Mandel & Farmer](#page-21-15) [2022;](#page-21-15) [Costa et al.](#page-21-16) [2023\)](#page-21-16). [de Mink et al.](#page-22-29) [\(2009,](#page-22-29) [2010\)](#page-22-30) introduces the concept of CHE to justify short period massive binaries such as the HMXB M33 X-7 [\(Orosz et al.](#page-21-45) [2007\)](#page-21-45).

Mandel $\&$ de Mink [\(2016\)](#page-21-46) found that the CHE channel significantly contributes to the BBH merger rate (10.5 ± 0.5 Gpc⁻³yr⁻¹ at redshift 0). These systems typically have a total mass ranging between 50 – 110 M_{\odot} . [Marchant et al.](#page-21-43) [\(2016\)](#page-21-43) explores the parameter space for forming CHE binaries. They found that at low metallicity, BBHs will merge within the age of the Universe due to weaker winds ($t_{\text{delay,min}} = 0.4 \text{ Gyr}$ for $Z_{\odot}/50$). There is strong evidence for increasing chirp masses with decreasing metallicity, with masses above the pair instability SN gap. If these mergers were detected, they could probe the evolution of massive stars in the early Universe. In addition to calculating the merger rate and mass distribution for the CHE channel, [du Buisson et al.](#page-22-31) [\(2020\)](#page-22-31) finds that the cosmological merger rate (5.8 Gpc⁻³yr⁻¹ at redshift 0) can vary up to 40% for extreme deviations in the metallicity specific SFR.

In COMPAS, there are two CHE modes, 'optimistic' and 'pes-simistic', which refer to how a CH star evolves on the MS [\(Riley](#page-22-27) [et al.](#page-22-27) [2021\)](#page-22-27). In the 'optimistic' mode, if the star begins the main sequence with fast enough rotation to evolve as a CHE star, it is assumed the star continues rotating fast enough to remain CH throughout the MS and the spin is not checked. For the CHE mode 'pessimistic' the spin is checked at every timestep, and if it has slowed below the CH metallicity-dependent rotational frequency threshold, it is immediately converted to a regular MS star and evolve to the end of the MS [\(Team COMPAS: Riley, J. et al.](#page-22-2) [2022\)](#page-22-2).

2.3 Remnant mass prescription

GWs can be used to infer the underlying mass distributions of BBHs. This remnant mass distributions depend on the metallicity and the type of explosion mechanism. Their formation can be described in three stages; stellar collapse and bounce, convective engine, and post-explosion fallback. Massive stars undergo a series of successful burning stages, building an iron core until electron degeneracy pressure can no longer support this. Electron capture and dissociation of the core elements into alpha particles accelerates the core collapse. This collapse is halted when the core reaches nuclear densities and nuclear forces cause the core to bounce and emit a shock outwards. This results in a mixing of the hot and cold layers in the star, known as convection. The convective engine provides the means to convert potential energy released in the collapse into explosion energy in the form of a supernovae [\(Fryer et al.](#page-21-47) [2012;](#page-21-47) [Fryer et al.](#page-21-48) [2021\)](#page-21-48). The commonly accepted supernova explosion mechanism is neutrino driven, where some of the neutrinos emitted during the core collapse are reabsorbed, powering an explosion [\(Janka et al.](#page-21-49) [2007;](#page-21-49) [Marek & Janka](#page-21-50) [2009;](#page-21-50) [Janka](#page-21-51) [2012,](#page-21-51) [2017\)](#page-21-52).

[Fryer et al.](#page-21-47) [\(2012\)](#page-21-47) provide analytic prescriptions for estimating the remnant mass in population synthesis codes, based on at the time recent understanding of supernova and gamma-ray burst explosions. In their model/fit, the remnant mass depends on the core, fallback mass and explosion mechanism. They explore two neutrino-driven explosion mechanisms– the rapid convection, which leads to explosion in the 250 ms after bounce, and the delayed convection, which can occur over a longer timescale. [Fryer et al.](#page-21-53) [\(2022\)](#page-21-53) altered the remnant mass prescription from [Fryer et al.](#page-21-47) [\(2012\)](#page-21-47) to study the growth rate of convection. They introduce a new parameter in their models denoted f_{mix} , the mixing growth time for convective instabilities. A fast growth time, $f_{\text{mix}} = 4.0$, and a slow growth time, $f_{\text{mix}} = 0.5$, correspond to the rapid and delayed models respectively from [Fryer](#page-21-47) [et al.](#page-21-47) [\(2012\)](#page-21-47). Whether a star undergoes a successful or failed supernovae provides clues on the final remnant object. However, there are uncertainties surrounding these mechanisms [\(Mapelli](#page-21-14) [2021\)](#page-21-14). GW detection from a core collapse SN can be distinguish between explosion mechanisms [\(Powell et al.](#page-22-32) [2016,](#page-22-32) [2023\)](#page-22-33)

[Mandel & Müller](#page-21-11) [\(2020\)](#page-21-11) develop probabilistic prescriptions for the remnant mass based on results from 3D supernovae simulations in [Müller et al.](#page-21-54) [\(2016\)](#page-21-54). Instead of a single value prescription, the BH mass branches over a range of core masses, depending on the fallback strength. If the BH is not formed by complete fallback, the mass follows as normal distribution. These models are more realistic as they capture the stochasticity of stellar evolution.

[Schneider et al.](#page-22-34) [\(2021\)](#page-22-34) also applies the [Mandel & Müller](#page-21-11) [\(2020\)](#page-21-11) semi-analytic SN prescription, but the resulting remnant mass prescription differs. The model focuses more on the removal of the hydrogen-rich common envelope and the evolution steps before core collapse. They also explore the amount of fallback. Their models overestimate the BH masses by 10% if the BHs are not formed by direct collapse. They do not account for any convective boundary mixing beyond the core helium burning.

[Schneider et al.](#page-22-34) [\(2023\)](#page-22-35) uses the models in Schneider et al. [\(2021\)](#page-22-34) and the stellar evolution code MESA [\(Paxton et al.](#page-21-55) [2011\)](#page-21-55) to explore the black hole mass distribution and chirp mass distribution of BBH mergers. The ability of stars to explode in the delayed neutrino-driven SN mechanism is characterised by the compactness of the progenitor cores [\(O'Connor & Ott](#page-21-56) [2011;](#page-21-56) [Ertl et al.](#page-21-57) [2016\)](#page-21-57). This depends on the mass and radius, and a high compactness indicates unsuccessful explosions and collapse into a BH. Thus, the chirp mass distribution is bimodal and the model predicts a lack of chirp masses in the range 9 – 13 M_{\odot} . Mass accretion can influence the final outcome of core collapse and final remnant mass. [Schneider et al.](#page-22-36) [\(2024\)](#page-22-36) studies models with MESA that allow for multiple mergers. They find BH masses in the pair instability SN gap, which can explain events such as GW190521. They note that for a more complete picture of stellar mergers, further detailed multi-dimensional, hydrodynamic simulations and a systematic exploration of the relevant parameter space is needed.

2.4 Summary

We outline the parameters and model assumptions varied in COMPAS in Table [1.](#page-6-1) We note that the initial conditions and distributions for the binary systems are set to default (except for the metallicity where we use Shark), as shown in Table [A1.](#page-22-37) Future studies may choose to vary these parameters. As discussed, we are aware of the limitations and approximations made in COMPAS and other population synthesis codes, that may cause discrepancies in the outputs. We summarize these limitations and the key questions below:

(i) The binding energy formalism utilises a fixed α_{CE} , but is this a valid approach? [Hirai & Mandel](#page-21-23) [\(2022\)](#page-21-23) proposes a two-step CE process, which depends on the evolutionary stage of the donor and the companion's mass. They find that the predicted post-CE separation is wider compared to the α_{CE} formalism. This may pose a problem for BBHs merging within Hubble time. [Picker et al.](#page-21-58) [\(2024\)](#page-21-58) has recently implemented fits for the convective envelope mass and associated values of λ to be used in conjunction with the two-step CE formalism. They find the effective combined value of $\alpha_{\text{CE}}\lambda$ can range from $\sim 10^{-3}$ to 10³ when converted to the α-formalism.

(ii) Are the masses in COMPAS overestimated and how does this impact the CHE binaries? CHE stars evolve from the hydrogen-ZAMS directly to the helium-ZAMS. When a CHE star is close to the end of the MS, it is already rich in helium, and may start to resemble a helium/ Wolf–Rayet star. However, COMPAS implements the same wind loss prescription throughout the MS, instead of a specific Wolf–Rayet wind prescription with higher mass loss rates. This change is applied to the MESA CHE modelling [\(Marchant et al.](#page-21-43) [2016;](#page-21-43) [du Buisson et al.](#page-22-31) [2020;](#page-22-31) [Marchant et al.](#page-21-59) [2023\)](#page-21-59).

(iii) Mass loss via radiation-driven winds is still an uncertainty in massive star progenitor models and accurate predictions are necessary for predicting their evolutionary paths. [Björklund et al.](#page-21-60) [\(2023\)](#page-21-60) derive radiation-driven mass-loss from hot, massive stars depending on their fundamental stellar parameters and implement this in MESA. They find that lower mass-loss rates potentially allow for the creation of high-mass black holes even at higher metallicities compared to previous models.

(iv) At masses > $50M_{\odot}$, [Hurley et al.](#page-21-24) [\(2000\)](#page-21-24) smoothly extrapolate the Pols et al. (1998) model grid up to 150 M_{\odot} , which is not well tested outside of the prescribed mass range. Significant improvement on this extrapolation method is far from trivial, as the evolution of high mass binaries is still very uncertain. This is reflected by the discrepancies between population synthesis codes in their stellar tracks [\(Agrawal](#page-20-9) [et al.](#page-20-9) [2020,](#page-20-9) [2022a](#page-20-7)[,b\)](#page-20-10).

(v) There are many uncertainties in the mass transfer efficiency and angular momentum loss, which influence the Roche lobe response to mass loss. Uncertainties in how a donor star responds to mass loss, combined with the Roche lobe response uncertainties contributes to the lack of understanding around the mass transfer stability boundary as a function of donor mass and radius [\(Ge et al.](#page-21-61) [2010,](#page-21-61) [2015,](#page-21-62) [2020;](#page-21-63) [Temmink et al.](#page-22-39) [2023\)](#page-22-39).

(vi) Is the angular momentum loss occurring closer to the accretor or the L2 point? And should the implementation of f_{γ} vary for degenerate and non-degenerate accretors? When performing the analysis in this paper, split functionality for AM loss from degenerate and non-degenerate accretors was yet to be implemented in COMPAS. Physically, it is more accurate for f_{γ} to be implemented differently for the primary and secondary star, particular when the accretor is a compact object. It is not clear how this would impact our current results, where we have only considered the angular momentum loss near the non-degenerate object. [Gallegos-Garcia et al.](#page-21-64) [\(2023\)](#page-21-64) derive an analytic model for mass loss from the binary that is informed

Table 1. COMPAS hyper-parameters explored in this paper. Note that some prescriptions are only used in conjunction with others. The mass transfer fraction is only applied if the mass transfer accretion efficiency prescription is set to FIXED_FRACTION. The default is THERMAL. Similarly, the angular momentum loss linear fraction is only applied when the mass transfer angular momentum loss prescription is MACLEOD_LINEAR. The default is ISOTROPIC ($f_{\gamma} = 0$). Refer to Tables [C1](#page-24-0)[,C2](#page-25-0) and [C3](#page-26-0) in Appendix [C](#page-22-40) for all relevant prescriptions used for each model.

by observations of X-ray binary accretion disks and hydrodynamical simulations, which considers angular momentum loss from a disk wind around a compact object accretor. They find their model predicts more angular momentum loss compared to the standard case of isotropic re-emission and the orbital separation of a binary can be considerably affected by mass loss via a disk wind.

(vii) COMPAS has two modes; single stellar evolution (SSE) and binary stellar evolution (BSE). The [Hurley et al.](#page-21-24) [\(2000,](#page-21-24) [2002\)](#page-21-21) SSE algorithms are implemented in the BSE mode at all evolutionary phases and provides the stellar attributes for both components of the binary. However, the fitting formulae are not well behaved outside of the metallicity range $Z \in [10^{-4}, 0.03]$.

(viii) The SSE tracks in COMPAS are based on non-rotating stars and do not account for effects on stellar evolution due to mild rotation. Thus, the modelling of rotational mixing, core overshooting (convection beyond the Schwarzschild criterion) and donor stripping is not accurate. Many studies had investigated the impact of these parameters on stellar structures using MESA and PARSEC [\(Claret &](#page-21-65) [Torres](#page-21-65) [2018;](#page-21-65) [Costa et al.](#page-21-66) [2019;](#page-21-66) [Laplace et al.](#page-21-67) [2020;](#page-21-67) [Temaj et al.](#page-22-41) [2024\)](#page-22-41).

3 PREDICTING THE GW MERGER RATE FOR DIFFERENT COMPAS MODELS

In this section we demonstrate our method for rescaling the merger rate from a fiducial, carefully cross-matched, simulation of galaxy formation (via Shark) and binary population synthesis (from COMPAS). This allows us to produce predictions for the rate and properties of an ensemble of GW events for any model without the large computational cost of redoing the cross-matching from scratch. From [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10), the number of GW events per SHARK galaxy that occur at time t_i from stars born at time t_i using COMPAS model α was written as

$$
\frac{dN_{\text{GW}}^{\alpha}}{dN_{\text{gal}}}(Z, t_i, t_j) = \frac{\psi(Z, t_i)}{M_{\text{tot}}^{\alpha}} \Delta t_i \sum_{n=1}^{N_{\text{binary}}} \vartheta(t_i + \tau_n^{\alpha}(Z), t_j, \Delta t_j), \quad (12)
$$

where

$$
\Delta t_j = t_j - t_{j-1},\tag{13}
$$

$$
\vartheta(t, t_j, \Delta t_j) = \begin{cases} 1 & \text{if } t_j - \frac{\Delta t_j}{2} < t < t_j + \frac{\Delta t_j}{2}, \\ 0 & \text{otherwise.} \end{cases} \tag{14}
$$

and $\psi(Z, t_i)$ is the metallicity (Z) dependent star formation rate in the galaxy when the progenitor stars were born. M_{tot}^{α} is a normalisation constant for a given model α (see Eq 10 in [Rauf et al.](#page-22-10) [2023\)](#page-22-10), which normalises the star formation rate based on the chosen stellar initial mass function, in such a way that it cancels out the dependence of the number of GW events on the number of binaries that we simulate N_{binary} , and any lower or upper mass cuts we apply when running COMPAS. $\vartheta(t, t_i, \Delta t_i)$ is a counter for the number of BBH mergers in SHARK snapshot, j. $\tau_n^{\alpha}(Z)$ is the coalescence time of each binary outputted from the model α . If the merger time, $t = t_i + \tau_n^{\alpha}(Z)$, falls in the i^{th} bin, a count is added to that bin, otherwise it does not contribute. Overall, this equation encapsulates that the rate consists of two contributions, one from the Shark host galaxy and the other from population synthesis (in our case from COMPAS).

3.1 The effective factor for different models

Cross-matching the star formation history of every galaxy in a simulation with a COMPAS run is computationally expensive. To simplify this, we now demonstrate how we can rescale the rate. If instead of considering GW events on a per galaxy basis we consider them in terms of number per metallicity and time t_i and t_j bins, we can write

$$
\frac{d^3 N_{\text{GW}}^{\alpha}}{dZ dt_i dt_j} = \frac{d^3 N_{\text{gal}}}{dZ dt_i dt_j} \frac{dN_{\text{GW}}^{\alpha}}{dN_{\text{gal}}}
$$
\n
$$
= \frac{d^3 N_{\text{gal}}}{dZ dt_i dt_j} \frac{\overline{\psi}(Z, t_i)}{M_{\text{tot}}^{\alpha}} \Delta t_i \sum_n \theta(t_i + \tau_n^{\alpha}(Z, t_i), t_j, \Delta t_j).
$$
\n(15)

In the second line we have substituted in Eq. [12](#page-6-2) and use $\overline{\psi}$, $\overline{\theta}$ to acknowledge that this is the average of the star-formation history and Heaviside function over all galaxies that fall into the the bin $dZdt_i dt_j$ (i.e., we assume that all galaxies in this bin produce the same number of gravitational waves, which is strictly true only for infinitesimally small bins). We now can introduce a new population synthesis model, β , where the merger rate is unknown. If we have the merger rate from model α , we can take a ratio of the two models and rearrange,

$$
\frac{d^3 N_{\text{GW},k}^{\beta}}{dZ dt_i dt_j} = \frac{M_{\text{tot}}^{\alpha}}{M_{\text{tot}}^{\beta}} \frac{d^3 N_{\text{GW}}^{\alpha}}{dZ dt_i dt_j} \frac{\sum_n \theta(t_i + \tau_n^{\beta}(Z, t_i), t_j, \Delta t_j)}{\sum_n \theta(t_i + \tau_n^{\alpha}(Z, t_i), t_j, \Delta t_j)} = \frac{M_{\text{tot}}^{\alpha}}{M_{\text{tot}}^{\beta}} \frac{d^3 N_{\text{GW}}^{\alpha}}{dZ dt_i dt_j} f_{\text{eff}}^{\alpha \beta}(Z, t_i, t_j, \Delta t_j). \tag{16}
$$

We hence find that retaining the information from SHARK is not required as the star formation history, $\overline{\psi}(Z, t_i)$, and the galaxy distribution are independent of the population synthesis model. In practice we also expect that for most use cases $\frac{M_{tot}^{\alpha}}{M_{tot}^{\beta}} \approx 1$ since changes to the initial mass function between different literature values primarily affect stars that are too small to become binary black holes or neutron stars, although we retain it here for completeness.^{[1](#page-6-3)} The only caveat

 1 Note this factor also needs to be included if one were to rescale between models where different lower or upper mass limits have been used to draw

Figure 1. An example of the effective factor between two COMPAS models with different remnant mass prescriptions α = [Fryer et al.](#page-21-47) [\(2012\)](#page-21-47) and β = Mandel $\&$ Müller [\(2020\)](#page-21-11) in bins of merger time and metallicity for binaries forming at redshift 2.5 (corresponding to $t_i \approx 2.62 \text{Gyr}$). In this plot and the rest of our work, an un-equal metallicity binning is chosen such that the number of Shark galaxies in each metallicity bin in the same. Unshaded bins indicate regions of the parameter space where there are no gravitational waves forming in either one or both of the models.

then is on the free choice of binning used to evaluate what we call the 'effective factor' $f_{\text{eff}}^{\alpha\beta}$ between the two models — although the dependence on galaxy simulation has been absorbed into our fiducial $\tilde{d}^3 N_{\rm GW}^{\alpha}$ $\frac{d^2 N_{\text{GW}}}{dZ dt_i dt_j}$, there is still the implicit assumption that the rate in each galaxy in the bin is the same. We find that this approximation remains valid over a range of metallicity binning choices, even for the quite wide bins we adopt as our default (100 un-evenly spaced bins for log Z ∈ [−7.0, −0.3]). We show an example distribution of f_{eff} in these metallicity and merger time bins in Fig. [1](#page-7-0) for a single choice of binary formation redshift of 2.5 (corresponding to $t_i \approx 2.62 \text{Gyr}$).

3.2 Generating the merger rate

Here we give a qualitative overview of our algorithm for predicting the GW merger rate for our different COMPAS models.

(i) The starting point of our algorithm is a high-fidelity set of GW events simulated from a combination of Shark and COMPAS model α . The detailed steps used for generating these are covered in [Rauf](#page-22-10) [et al.](#page-22-10) [\(2023\)](#page-22-10) and so will not be repeated here. Briefly, this is done by cross-matching the galaxy metallicities to the binary population metallicities outputted from COMPAS. We inject this population into the Shark galaxies and track their evolution until they merge using Eq [14.](#page-6-4)

(ii) From this, $\frac{d^3 N_{\text{GW}}^{\alpha}}{dZ dt_i dt_i}$ is calculated by taking the fiducial merger rate outputs and binning in terms of metallicity and birth and merger time. For the metallicity we use quantiles to determine the bin edges such that there is an even number of galaxies in each bin. For the birth and merger time, we simply use the 170 snapshots from SHARK to define the 170 bins.

stars from the Initial Mass Function (IMF), where a different mass ratio distribution for the initial binaries is assumed, or the overall binary fraction is changed

Figure 2. Volumetric BBH merger rate with remnant mass prescription from [Mandel & Müller](#page-21-11) [\(2020\)](#page-21-11) as a function of redshift. The blue line is the true merger rate from running COMPAS and Shark. The red dashed line is the merger rate rescaled with the effective factor calculated using the full COMPAS run containing 170×10^6 binaries. The grey lines are the rescaled merger rate where the effective factor has been calculated using a unique sub-sample of 10⁶ binaries from the full COMPAS run. The green line is the average over 170 of these subsamples.

(iii) Having established our metallicity, birth time and merger time bins, we take our full list COMPAS binaries, read in the metallicity and evolution time, which is added to the birth time $(t_{\text{birth}} + t_{\text{BBH}} +$ $t_{\text{coalescence}}$), and add these to the corresponding bins. We apply this to both the fiducial and new model, take the ratio and multiply by the merger rate in (ii) to estimate the new merger rate, shown in Eq [16.](#page-6-5)

Once we have our estimate for the number of gravitational waves in our new model β we evaluate the volumetric merger rate at time t_i by simply summing over the metallicity bins and birth time bins,

$$
R_{\rm GW}^{\beta}(t_{\rm j}) = \frac{1}{V_{\rm sim}} \sum_{i=0}^{170} \sum_{k=0}^{100} \frac{d^3 N_{\rm GW}^{\beta}}{dZ_k dt_i dt_j},\tag{17}
$$

where V_{sim} is the volume of our original SHARK simulation.

For efficiency, we choose a sample $\tilde{N}_{\beta}^{\text{GW}}$ of 10⁶ binaries from our β COMPAS run to estimate the effective factor, while our fiducial model α has 170 × 10⁶.^{[2](#page-7-1)} The merger rate is reduced by a factor of 170 but this is compensated for by the normalisation $\frac{M_{tot}^{\alpha}}{M_{tot}^{\beta}} \sim 170$. Thus, this formalism removes the dependence on the sample size. We test the validity of our rescaling procedure and using only 10^6 binaries for model β by splitting one of our fiducial COMPAS runs from [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) (using the remnant mass model of [Mandel](#page-21-11) [& Müller](#page-21-11) [2020\)](#page-21-11) into 170 by 10^6 sub-samples. We then use these to estimate the rescaled merger rate for this model given α = [Fryer](#page-21-47) [et al.](#page-21-47) [\(2012\)](#page-21-47), and look at the spread in the merger rate. We show in Fig. [2](#page-7-2) that overall the rescaling method using equal numbers of binaries for models α and β recovers the true merger rate from our high-fidelity cross-matching, but also that using only 10⁶ binaries for

² Note that we draw only ~ 10⁶ unique masses for this process from the IMF, but each one is randomly assigned a different metallicity from SHARK, resulting in ∼ 170 × 10⁶ unique *combinations* of initial masses and metallicities, and so ~ 170×10^6 unique binaries.

Figure 3. Volumetric rate plot as a function of redshift from [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) recreated with the effective factor scaling method. The error on the plots are the range of merger rates from sampling from each COMPAS run 170 times.

model β returns, on average, an unbiased estimate of the merger rate. The use of a smaller number of binaries in the rescaling procedure does introduce additional Poisson scatter in the merger rate, which we propagate through into our model comparison/evaluation (see Section [4.](#page-8-0)

In Fig. [3,](#page-8-1) we show the volumetric BBH merger rates in the SHARK simulations using the effective factor approximation for four different models. In all cases we find that we are able to recover rates from [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) very well. We note that the plot is noisier than in [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) due to the smaller sample of BBHs taken from each COMPAS run, but again this noise arises only from Poisson sampling error.

4 BAYESIAN INFERENCE AND MODEL COMPARISON

In this section we outline our procedure for comparing different COMPAS models to data from GWTC-1 [\(Abbott et al.](#page-20-1) [2019a\)](#page-20-1), GWTC-2 [\(Abbott et al.](#page-20-2) [2021a\)](#page-20-2) and GWTC-3 [\(Abbott et al.](#page-20-3) [2023b\)](#page-20-3) including our rescaling methodology. We validate this by method by comparing the likelihoods given the data for the four models from [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) for which we have both 'brute-force' measurements of the GW rate and our rescaled calculations.

4.1 Selection criteria

The Bayesian analysis will depend strongly on N_{obs} , which is the number of *confident* GW detections. How confident we are in the GW signals and the validity of the sources is determined by the following ranking statistics:

(i) Signal-to-noise ratio (SNR): This is proportional to the signal amplitude divided by the noise amplitude. The higher the SNR, the stronger the signal compared to the noise, and more likely that the signal will be detected. This is the main ranking statistic when the signal shape is known from theory, such as the signal for a BBH merger. We use the network SNR, which is the square root of the sum of the squares of SNRs from the triggers which form the GW event (only from the detectors that produce triggers contributing to the event; [Cabourn Davies & Harry](#page-21-68) [2022\)](#page-21-68). The SNR, ρ , in a single detector is defined as,

$$
\rho^2 = \langle h|h \rangle = 4\mathcal{R} \int_{f_{\text{min}}}^{f_{\text{max}}} \frac{h^*(f)h(f)}{S(f)} df,
$$
\n(18)

where $h(f)$ is the waveform measured by the detector, $S(f)$ is the power spectrum density and f_{min} and f_{max} are the minimum and maximum limits for the considered frequency range [\(Farr et al.](#page-21-69) [2015;](#page-21-69) [Barrett et al.](#page-20-5) [2018\)](#page-20-5).

(ii) False alarm rate (FAR): Once the threshold for the SNR is set, the FAR can be determined, providing an alternative way to estimate the significance of the event [\(Zheng et al.](#page-22-42) [2021\)](#page-22-42). The FAR is the number of noise events with a signal equal to or greater than the GW event, divided by the total duration of the background data. The false alarm rate of single-detector candidates given ρ is,

$$
FAR(\rho) = \int_{\rho}^{\rho_{\text{max}}} \Lambda_n p_n(\rho') d\rho', \qquad (19)
$$

where Λ_n is the mean Poisson rate of noise triggers and $p_n(\rho')$ is the probability densities describing the distribution of ρ [\(Callister et al.](#page-21-70) [2017\)](#page-21-70).

(iii) p_{astro} : The probability that a particular event is of astrophysical origin, as opposed to noise fluctuations or terrestrial, is defined by p_{astro} . It is measured by comparing the GW event rate to the background event rate for a fixed SNR and FAR. The probability of a trigger, x , being astrophysical is,

$$
p_{astro}(x) = \frac{p(x|S)\pi(S)}{p(x|S)\pi(S) + p(x|N)\pi(N)},
$$
\n(20)

where S and N describe the signal and noise model, as well as the distribution of signal and noise sources respectively [\(Callister et al.](#page-21-70) [2017;](#page-21-70) [Broekgaarden et al.](#page-21-71) [2023\)](#page-21-71).

For our most rigorous selection, we combine FAR ≤ 0.25 yr⁻¹ and $p_{astro} \geq 0.9$, as the FAR alone will not provide information on the sources being astrophysical, but rather the behaviour of the signal background. The p_{astro} reveals more events in the population rich areas of the parameter space, and is a more realistic metric for setting threshold for electromagnetic follow-up [\(Lynch et al.](#page-21-72) [2018;](#page-21-72) [Andres et al.](#page-20-11) 2022). The caveat of p_{astro} is that that independent pipelines will obtain different p_{astro} due to a difference in data re-duction techniques. [Banagiri et al.](#page-20-12) [\(2023\)](#page-20-12) proposes an unified p_{astro} by combining triggers from all pipelines, which incorporates extra information about a signal. [Galaudage et al.](#page-21-73) [\(2020\)](#page-21-73) also emphasises the need for a pipeline independent p_{astro} , as they show that the posterior distribution of the population hyper-parameters can change with an astrophysically motivated prior. Given these caveats, we also investigate a simpler, broader selection using only an $SNR \geq 8$ cut, which matches that used in evaluating our detection probability function $p_{\text{det}}(\theta)$ for our models and provides us with a larger samples of events. However, the downside of this is that it potentially allows events into our sample that may not be astrophysical. In these cases, we allow our Bayesian framework to decide if these events should be ruled out.

Our supersample of GW events for querying the PDF for each COMPAS model consists of confirmed BBHs and BH-mass-gap events such as GW190521 [\(Abbott et al.](#page-20-13) [2020a\)](#page-20-13) and GW190814 [\(Abbott](#page-20-14) [et al.](#page-20-14) [2020c\)](#page-20-14). This gives a total of 93 catalog GW detections. Of these, the following GW events are *not* included in our analysis for the following reasons:

- GW170817: Confirmed BNS [\(Abbott et al.](#page-20-15) [2017\)](#page-20-15)
- GW190425: Confirmed BNS [\(Abbott et al.](#page-20-16) [2020b\)](#page-20-16)
- GW190426_152155: Not astrophysical [\(Abbott et al.](#page-20-2) [2021a\)](#page-20-2)

10 *Rauf et al.*

- GW190909_114149: Not astrophysical [\(Abbott et al.](#page-20-2) [2021a\)](#page-20-2)
- GW191219_163120: Confirmed NSBH [\(Abbott et al.](#page-20-3) [2023b\)](#page-20-3)
- GW200115 042309: Confirmed NSBH [\(Abbott et al.](#page-20-17) [2021b\)](#page-20-17)

We then apply our two sets of selection criteria. We have 79 GW events with SNR $\geq 8^3$ $\geq 8^3$ and 60 GW events with $p_{astro} \geq 0.9$ and FAR ≤ 0.25 yr⁻¹. Tab [B1](#page-23-0) lists the events used with each selection criteria. Out of the 87 BBH GW events we could have included (93 less the 6 listed above) the following GW events do not fit either criteria:

- GW190403_051519 • GW200220_061928
- GW190514_065416 • GW200306_093714
- GW191113_071753
- GW200208 222617

• GW200308_173609

• GW200322_091133

4.2 Fitting LVK posterior for various COMPAS **models**

Our basic fitting methodology is based on the widely-used Bayesian framework (e.g., [Stevenson et al.](#page-22-43) [2015;](#page-22-43) [Bouffanais et al.](#page-21-74) [2019;](#page-21-74) [Man](#page-21-75)[del et al.](#page-21-75) [2019;](#page-21-75) [Vitale et al.](#page-22-44) [2022;](#page-22-44) [Périgois et al.](#page-21-76) [2023\)](#page-21-76). Given the BBH GW data from some observing runs, H , the posterior for a model specified via hyper-parameters λ (for instance a certain mass transfer efficiency f_{MT} or specific angular momentum loss f_{γ}), takes the form of an inhomogeneous Poisson distribution,

$$
\mathcal{L}(\lambda, N_{\lambda}|\mathcal{H}) = \pi(\lambda)e^{-\mu_{\lambda}}N_{\lambda}^{N_{\text{obs}}}\prod_{k}^{N_{\text{obs}}} \mathcal{I}^{k}.
$$
 (21)

 N_{λ} is the number of GW events predicted by the model in a given time-frame, while μ_{λ} is the number the model suggests we would actually detect. For convenience we can use these to define some detection efficiency $\beta(\lambda) = \frac{\mu_{\lambda}}{N_{\lambda}}$. N_{obs} is the actual number of detected events in our dataset. We assume that *a priori* all hyper-parameter combinations are equally likely, so $\pi(\lambda) = 1$. Finally, \hat{I}^k denotes the likelihood of observing GW event h^k given the hyper-parameters λ , written as a marginalisation over the binary properties predicted by that model,

$$
\mathcal{I}^{k} = \int \mathcal{L}(h^{k}|\theta) p_{\text{sim}}(\theta|\lambda) d\theta = \frac{1}{N_{s}^{k}} \sum_{i=1}^{N_{s}^{k}} \frac{p_{\text{sim}}(\theta_{i}^{k}|\lambda)}{\pi^{k}(\theta_{i}^{k})},
$$
(22)

where $\theta = \{M_c, q, z\}$ are the GW event parameters chirp mass, mass ratio and redshift respectively and $p_{\text{sim}}(\theta|\lambda)$ is the probability distribution function for the event-level parameters generated by COMPAS using the Shark metallicity as input. The latter sum arises from the use of the N_s^k publicly available posterior samples to represent the likelihood of event k, $\mathcal{L}(h^k|\theta)$. Formally $p_{\text{sim}}(\theta|\lambda)$ is obtained by marginalising over all the metallicities and birth times used that can give rise to an event with parameters θ :

$$
p_{\text{sim}}(\theta|\lambda) = \int_0^\infty \int_0^{t_j} p_{\text{sim}}(\mathcal{M}_c, q, z(t_j)|Z, t_i, \lambda) dt_i dZ.
$$
 (23)

 N_{λ} is the number of GW events in the Universe within an observing time that is expected from each COMPAS model. We can evaluate this given our rescaled merger rate density (expressed in individual metallicity and birth time bins, and in units of Mpc⁻³yr⁻¹) and p_{sim} .

$$
N_{\lambda} = 4\pi T_{\text{obs}} \int_{\theta \in V_{\theta}} \int_0^{\infty} \int_0^{t_j(z)} p_{\text{sim}}(\theta | Z, t_i, \lambda)
$$

$$
R_{\text{GW}}^{\lambda}(Z, t_i, t_j(z)) \chi^2(z) \frac{c}{H(z)} \frac{1}{1+z} dt_i dZ d\theta, \qquad (24)
$$

where $\chi(z)$ is the comoving distance to the redshift z, c is the speed of light, $H(z)$ is the Hubble parameter, and T_{obs} is the total LVK observing time (duty cycle). This is 118.0, 138.8 and 142.0 days for O1, O2 and O3 respectively, which we converted to years.

We can extend this to the predicted *detected* number of GW events using a detection probability for the event-level parameters $p_{\text{det}}(\mathcal{M}_c, q, z)$ given a particular detector configuration and signalto-noise threshold.

$$
\mu_{\lambda} = 4\pi T_{\text{obs}} \int_{\theta \in V_{\theta}} \int_0^{\infty} \int_0^{t_j(z)} p_{\text{sim}}(\theta | Z, t_i, \lambda) p_{\text{det}}(\theta)
$$

$$
R_{\text{GW}}^{\lambda}(Z, t_i, t_j(z)) \chi^2(z) \frac{c}{H(z)} \frac{1}{1+z} dt_i dZ d\theta. \tag{25}
$$

For Eq. [21,](#page-9-1) we are assuming that N_{λ} events are all formed via the isolated formation channel. There is some evidence for some events in GWTC-3 [\(Abbott et al.](#page-20-3) [2023b\)](#page-20-3) forming via the dynamical channel [\(Arca Sedda](#page-20-18) [2021;](#page-20-18) [Romero-Shaw et al.](#page-22-45) [2022\)](#page-22-45). To avoid any assumptions on the formation channel and discount information on the predicted vs. observed GW rate, we can marginalise over N_{λ} to remove its dependence by multiplying with a prior $\pi(N_\lambda) = \frac{1}{N_\lambda}$ and integrating [\(Fishbach et al.](#page-21-77) [2018\)](#page-21-77).

$$
\mathcal{L}(\lambda|\mathcal{H}) = \int \frac{1}{N_{\lambda}} e^{-N_{\lambda}\beta(\lambda)} N_{\lambda}^{N_{\text{obs}}} \prod_{k}^{N_{\text{obs}}} \mathcal{I}^{k} dN_{\lambda}
$$

$$
= (N_{\text{obs}} - 1)! \prod_{k}^{N_{\text{obs}}} \frac{\mathcal{I}^{k}}{\beta(\lambda)},
$$
(26)

We make model inferences using both of these posteriors in our work and compare how our conclusions change when N_{λ} is included as information.

In order to evaluate our posteriors, we first calculate the model volumetric rate using a variant of Eq. [17](#page-7-3) but kept in separate metallicity and birth time bins. Given $T_{\rm obs}$, the merger rate and the appropriate volumetric factors, we then find the number of GW events that would fall in each metallicity and time bin, $N_{\lambda,i,j}$, and draw that many events (characterised by a chirp mass and mass ratio) from our full list of 10^6 COMPAS binaries.^{[4](#page-9-2)} We store the chirp mass, mass ratio and merger redshift of the events across all metallicity and birth time bins before then building the PDF $p_{\text{SIM}}(\theta|\lambda)$ using Kernel Density Estimation, in particular using the fast KDEpy.FFTKDE package. The bandwidth for our KDE is calculated using Silverman's rule of thumb [\(Silverman](#page-22-46) [1986\)](#page-22-46) for each hyper-parameter distribution and averaged to obtain the single band-width for the full 3D distribution. As a by-product of method we can also estimate N_{λ} and μ_{λ} at the same time by simply keeping track of how many COMPAS binaries we are drawing.

For $p(\theta_i^k|\lambda)$, we query the KDE-built PDF using the chirp mass, mass ratio and redshift from the public LVK posterior samples. We construct $\pi^{k}(\theta_i^k)$ by building another KDE with the public LVK

³ We use the network SNR provided in [https://gwosc.org/eventapi/](https://gwosc.org/eventapi/html/GWTC/?page=1&pagesize=100) [html/GWTC/?page=1&pagesize=100](https://gwosc.org/eventapi/html/GWTC/?page=1&pagesize=100).

⁴ We find $N_{COMPAS,i,i}$ in each metallicity and snapshot bins from the COMPAS run. We calculate a weight defined as $N_{\lambda,i,j}/N_{\text{COMPAS},i,j}$ to avoid repeatedly drawing from the COMPAS run. We apply this weight to the PDF. When calculating the detected PDF, we multiply this weight by p_{det} .

prior samples and querying this again with the posterior samples at each point θ_i^k . In the case where $p_{\text{sim}}(\theta|\lambda) = 0$ for all posterior samples, this implies that the COMPAS model cannot reproduce the BBHs in the posterior samples, as they fall outside of the KDE. We flag these events and ultimately remove them from the analysis for that specific model comparison. For each of the $N_{\lambda,i,j}$, we calculate the detection probability, p_{det} , based on the masses, redshift and luminosity distance. The detection probability comes from a selection effect function in COMPAS, which takes the BH masses, redshift and luminosity distance as inputs [\(Barrett et al.](#page-20-5) [2018;](#page-20-5) [Team COMPAS:](#page-22-2) [Riley, J. et al.](#page-22-2) [2022\)](#page-22-2). The SNR is calculated by computing the source waveform using the waveform model from the LAL software suite [\(LIGO Scientific Collaboration et al.](#page-21-78) [2018\)](#page-21-78).

We apply the same procedure also to estimate marginalised, observed 1D distributions $p_{\text{SIM_DET}}(\mathcal{M}_c|\lambda)$, $p_{\text{SIM_DET}}(q|\lambda)$ and $p_{\text{SIM DET}}(z|\lambda)$.^{[5](#page-10-0)} We show these 1D distributions for our four fiducial models in Fig. [4](#page-11-0) alongside the GWTC data, with our main observations being:

• The models follow the chirp mass distribution reasonably well except for the chirp mass range 100–130 M⊙. These models predict high mass BHs that either may have not been detected due to the rarity of finding these types of events with current detectors, as they may be associated with noise in the low frequency band, or as a result of inaccuracies in the models resulting in an incorrect representation of the observed population.

• The mass ratio distribution is quite noisy but overall relatively flat, similar to the observed distribution. However for [Mandel &](#page-21-11) [Müller](#page-21-11) [\(2020\)](#page-21-11) and [Schneider et al.](#page-22-34) [\(2021\)](#page-22-34) there is a peak in the distribution after $q \approx 0.8$ indicating a preference for equal mass binaries. [Mandel & Müller](#page-21-11) [\(2020\)](#page-21-11) with $f_{\text{WR}} = 0.2$ has a preference for $q \sim 0.6$ and [Fryer et al.](#page-21-47) [\(2012\)](#page-21-47) for $q \sim 0.7 - 0.8$.

• The shape of the redshift distributions for the models fit the observed distribution quite well. Overall, the [Fryer et al.\(2012\)](#page-21-47) model is the best fit, while the other models have a wider distribution. This indicates BBHs from these models are more likely to be detected at higher redshifts.

4.3 Quantifying the uncertainty in N_A , $p(\lambda, N_A|\mathcal{H})$ and $p(\lambda|\mathcal{H})$

For our four models in [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) we were able to split COMPAS outputs into 170 sub-samples to compute 170 effective factors and produce 170 theoretical, N_{λ} , and detected, μ_{λ} , numbers of GW events. We show the distribution of the detected number of events in Fig. [5.](#page-11-1) For the error on the total number of events, we can use the standard deviation across our 170 subsamples. However, if we have only one sample of N_{sample} binaries for a COMPAS, we require an alternative approach to assign an error for these models.^{[6](#page-10-1)} We find that the ratio of the standard deviation to the mean number of events as a function of $\log N_{\lambda}$ for the four models has a linear trend such that the relative error increases as the total number of GW events

generated by the model decreases, in a roughly Poissonion fashion. In Fig. [6](#page-11-2) we show the linear fits for the standard deviation in N_{λ} and μ_{λ} , which we show in the following equations:

$$
\frac{\Delta N_{\lambda}}{N_{\lambda}} \approx -0.04 \log N_{\lambda} + 0.55
$$
 (27)

$$
\frac{\Delta \mu_{\lambda}}{\mu_{\lambda}} \approx -0.03 \log N_{\lambda} + 0.43 \tag{28}
$$

We can now apply this to single samples, assuming similar Gaussian distributions for other COMPAS models. To justify this, we find that the mean μ_{λ} over the 170 sub-samples in Fig. [5](#page-11-1) is equivalent to the μ_{λ} we obtain using the true merger rate from the full COMPAS output (one sample) for all four models that we test.

Now to address the uncertainty in \mathcal{I}^k ; We find another linear trend for the standard deviation in $\sum_{k}^{N_{\text{obs}}} \log T^{k}$ over the 170 sub-samples as a function of the mean $\log N_\lambda$ for both our selection criteria, as shown in Fig. [7.](#page-12-1)

The linear fits for the standard deviation in the likelihood sums are given by:

$$
\frac{1}{N_{\text{obs}}} \Delta \sum_{k}^{N_{\text{obs}}} \log \mathcal{I}^k \approx -0.05 \log N_{\lambda} + 0.70 \tag{29}
$$

For new COMPAS models shown in Section [5,](#page-12-0) we are able to use Eq. [29](#page-10-2) and to calculate the error on $\sum_{k}^{N_{\text{obs}}} \log T^{k}$ given N_{λ} without needing to resample. We can now propagate the error in N_{λ} , μ_{λ} and $\sum_{k}^{N_{\text{obs}}} \log T^{k}$ into the error on the log-likelihood.

$$
\Delta \log \mathcal{L}(\lambda|\mathcal{H}) = \sqrt{\left(N_{\text{obs}}\frac{\Delta \mu_{\lambda}}{\mu_{\lambda}}\right)^{2} + \left(N_{\text{obs}}\frac{\Delta N_{\lambda}}{N_{\lambda}}\right)^{2} + \left(\Delta \sum_{k}^{N_{\text{obs}}} \log T^{k}\right)^{2}}
$$

$$
\Delta \log \mathcal{L}(\lambda, N_{\lambda}|\mathcal{H}) = \sqrt{(\Delta \mu_{\lambda})^{2} + \left(N_{\text{obs}}\frac{\Delta N_{\lambda}}{N_{\lambda}}\right)^{2} + \left(\Delta \sum_{k}^{N_{\text{obs}}} \log T^{k}\right)^{2}}
$$
(30)

Although only rudimentary given the small number of models available to us to generate fitting formulae, even adopting this simple uncertainty on the likelihood allows us to consider the statistical variations in model evaluation using population synthesis. As such it enables us to draw more confident conclusions on which hyperparameters cause major changes in the likelihood, and whether these changes are physically significant.

4.4 Comparing posteriors with and without rescaling merger rate

Fig. [8](#page-13-0) shows the I^k term for individual GW events. This is a practical approach to determine the probability of individual GW events occurring in the populations generated by the COMPAS models. The [Schneider et al.](#page-22-34) [\(2021\)](#page-22-34) model generally returns high probabilities when querying the PDF, while the [Fryer et al.](#page-21-47) [\(2012\)](#page-21-47) model is disfavoured. We find some events are outliers and are not well reproduced by some or all of the models relative to the other events. As an example, [Mandel & Müller](#page-21-11) [\(2020\)](#page-21-11) with $f_{WR} = 0.2$ and [Schneider](#page-22-34) [et al.](#page-22-34) [\(2021\)](#page-22-34) are less likely to output the progenitors of GW190814, GW190917_114630 and GW200210_092254 compared to [Fryer](#page-21-47) [et al.](#page-21-47) [\(2012\)](#page-21-47) and [Mandel & Müller](#page-21-11) [\(2020\)](#page-21-11). For GW190426_190642, GW190602_175927 and GW191109_010717 [Schneider et al.](#page-22-34) [\(2021\)](#page-22-34) is distinctly preferred over the other models, although in these cases, and especially that of GW190521, the models are generally less able

 5 $p_{\text{SIM_DET}}$ refers to the detected PDF, which is marginalised over the other two fitting parameters *after* the detection probability is applied.

 $6\,$ Given the number of models we test in this work, the alternatives of simply using more samples or bootstrapping using the $N_{\text{sample}} = 10^6$ samples would be computationally challenging. Furthermore, subsampling or bootstrapping to generate an error would fail to capture the impact of progenitors of rare events that are not represented in the $N_{\text{sample}} = 10^6$, but which are present in the distributions obtained from our four models with $N_{\text{sample}} = 170 \times 10^6$, and from which our errors are extrapolated.

Figure 4. Distribution of M_c , q and z for different models. Difference colours refer to the different remnant mass prescriptions investigated in [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10). The grey shaded regions are the inferred distribution from the LVK collaborations, calculated using the posterior samples for the $N_{\text{obs}} = 79$ GW events based on the $SNR \geq 8$ cut.

Figure 5. Normalised histograms of the detected number of BBH merger events calculated using Eq. [25](#page-9-3) over the LVK observing time for the last three observing runs. The coloured distributions are obtained from the predictions from the model for 170 sub-samples. The grey band represents the uncertainty in the observed number of BBH mergers depending on the selection criteria adopted (see Section [4.1\)](#page-8-2). The dotted lines are the number of events predicted using the true merger rate from [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10).

to produce the required event-level properties. This is expected for GW190521 as both masses, $\sim 85M_{\odot}$ and $\sim 66M_{\odot}$, most likely lie in the pair-instability supernovae mass gap, making it atypical of bina-

Figure 6. Relative error on the numbers of generated and detected events versus the average log N_{λ} for the four [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) models from 170 sub-samples. The crosses correspond to the relative error from the models and the dashed lines are linear fits to the points. The black points and fit are for N_λ and the red points and fits are for μ_λ .

ries formed via the isolated formation channel [\(Abbott et al.](#page-20-13) [2020a\)](#page-20-13).

For a more comprehensive assessment of the models, we can analyse the likelihoods. In Fig. [9](#page-14-0) we now incorporate the detection probability and show the likelihoods with and without N_{λ} dependence. The

Figure 7. The standard deviation in the sum of the log-likelihood per event, $\log \frac{I}{N_{\text{obs}}}$, versus the mean $\log N_{\lambda}$ over the 170 sub-samples for the four [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) models. We show the linear fit (green dashed line) that applies to both sets of selection criteria; FAR ≤ 0.25 yr⁻¹ and $p_{astro} \geq 0.9$ (black) and $SNR \geq 8$ (red).

[Mandel & Müller](#page-21-11) [\(2020\)](#page-21-11) appears to the best fit to the observed population. This can be justified with Fig. [4.](#page-11-0) The [Mandel & Müller](#page-21-11) [\(2020\)](#page-21-11) model has two peaks in the chirp mass distribution and a relatively flat mass ratio distribution, similar to the LVK population distribution. If we assume that the models output accurate populations, then incorporating the number of GW events makes the likelihood worse – [Fryer et al.](#page-21-47) [\(2012\)](#page-21-47) greatly overestimates the number of detected events, while [Schneider et al.](#page-22-34) [\(2021\)](#page-22-34) underestimates it. In general the latter is preferred as it leaves 'room' for some of the LVK observed events to arise from the dynamical formation channel. The exception is the [Mandel & Müller](#page-21-11) [\(2020\)](#page-21-11) model, which performs well since the model predicts the same number of GW events within the observed range. The true number of events predicted is ∼ 60, comparable to N_{obs} for the $p_{\text{astro}} \ge 0.9$ and FAR ≤ 0.25yr^{-1} selection criteria. For SNR \geq 8, we see an increase in the likelihood for all models, but the distributions remain the same. The biggest difference in the unmarginalised likelihood for our two selections occurs for the [Fryer](#page-21-47) [et al.](#page-21-47) [\(2012\)](#page-21-47) model, which produces the largest number of detected events out of the four models. This is due to N_{obs} increasing to 79, such that the difference between the predicted μ_{λ} and the number of observed LVK events is slightly reduced.

5 RESULTS

Having validated our methodology using the four models analysed in [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10), this section discusses the outputs from our new set of COMPAS models where we vary the range of hyper-parameters described previously in Section [2](#page-2-0) (see also Table [1.](#page-6-1) We start with describing the trends in the number of detected events when varying the COMPAS hyper-parameters. We calculate the Bayes factor to identify groups of models that are more preferred. We then show our likelihood analysis, where we discuss the best and worst models when compared to the observed population parameters. We follow up this analysis and look at specific events and their \mathcal{I}^k value to determine which models are more and less likely to generate the progenitors of these events. See Tables [C1,](#page-24-0) [C2](#page-25-0) and [C3](#page-26-0) in Appendix [C](#page-22-40) for a detailed description of the hyper-parameters for all 304 COMPAS models in this analysis.

5.1 μ _{GW} versus COMPAS parameters

Since we explore the number of GW events as a parameter in our posterior analysis, in this section we comment briefly on the trends between the detected number of GW events, μ_{λ} , and the COMPAS hyper-parameters.

In the left plot of Fig. [10,](#page-15-0) we see μ _{GW} increase with mass transfer fraction, f_{MT} , when the corresponding angular momentum is lost directly from the accretor ($f_{\gamma} = 0$). Increasing the mass transfer can lead to more equal mass objects, which we find most of our COMPAS models producing. This trend disappears for higher f_{γ} . We credit this to non-trivial evolutionary effects such as the stability prescriptions, that determine any future interactions within the binary. This preference for intermediate values of f_{γ} agrees with the findings by [Santoliquido et al.](#page-22-3) [\(2021\)](#page-22-3). Irrespective of f_{MT} , μ_{GW} peaks for moderate values of $f_{\gamma} = 0.5$ but drops again for high f_{γ} when angular momentum loss occurs closer to the L2 point. At this point, mass is lost at higher velocities so more angular momentum is lost from the system, which in turn can induce CE evolution and merger during the mass transfer stage, stopping the formation of a BBH. These results are in agreement with [Lu et al.](#page-21-41) [\(2023\)](#page-21-41); [van Son et al.](#page-22-22) [\(2022b\)](#page-22-22) that the number of stable mass channel objects increases for decreasing f_{γ} , as opposed to the objects formed by CE evolution. [Marchant](#page-21-13) [et al.](#page-21-13) [\(2021\)](#page-21-13) also found that the stable channel dominates the BBH merger rate. We note that when $f_{\text{MT}} = 1$, f_{γ} should have no impact. Any variations in the number of events at $f_{\text{MT}} = 1$ is hence due to Eddington-limited accretion occurring after the first mass-transfer event(s), once the accretor has become a BH. \prime

We show the models that produce the closest μ_{λ} to the observed LVK events in the right plot of Fig. [10.](#page-15-0) We find that for μ _{GW} to be below 100 events (closer to the number of detected events from all observing runs), a CE efficiency $\alpha_{\text{CE}} = 10, 100$ and $f_{\gamma} = 0.75, 1.0$ with no CHE is required. This implies that the CE is ejected earlier in the evolution and the orbit tightens at a wider separation, which means longer merger times and less events in the same observed time. This result validates the studies by [Santoliquido et al.](#page-22-3) [\(2021\)](#page-22-3), [Bouffanais et al.](#page-21-28) [\(2021\)](#page-21-28) and [Grichener](#page-21-26) [\(2023\)](#page-21-26). With the exception of these models, the CE efficiency has no significant impact on the populations, as the range for the number of events does not vary significantly with α_{CE} .

In Figure [11,](#page-15-1) μ _{GW} decreases with increasing f_{WR} . We find this for any CHE mode and remnant mass prescription, but we show this for [Schneider et al.](#page-22-34) [\(2021\)](#page-22-34). This is due to increasing the mass loss rates leading to smaller BHs and wider separations, so less mergers occur in the same observing time. These predictions agree with the work by [Riley et al.](#page-22-27) [\(2021\)](#page-22-27) and [Stevenson & Clarke](#page-22-6) [\(2022\)](#page-22-6) using COMPAS. We find that for no CHE, the rate of change in μ_{λ} across f_{WR} is smaller compared to the pessimistic and optimistic cases.

The pessimistic and optimistic CHE modes produced > 1000 events due to their tight orbits. This suggests the CHE channel is not

 7 As a result of this finding, dual functionality in the AM loss was later implemented in COMPAS, as discussed in Section [2.4.](#page-5-0) If specific AM loss is set at the compact object as well as prior to the formation of the first compact object we expect the number of events will no longer vary for $f_{\text{MT}} = 1$, although at time of writing this functionality is yet to be comprehensively tested.

Figure 8. I^k term for each GW event with SNR ≥ 8 . Each colour represents a different remnant mass prescription from COMPAS and the error bar represents the sampling error over 170 samples when building the simulated PDF of M_c , q and z given the model.

Figure 9. Normalised histograms of the log likelihoods obtained using 170 sub-samples for the models explored in [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) with and without the merger rate dependence. The top two panels are the likelihoods comparing the models to the GW events that fit the FAR ≤ 0.25 yr⁻¹ and $p_{astro} \geq 0.9$ criteria, while the bottom two panels compares to the SNR ≥ 8 criteria. The dotted lines are the likelihoods using the true merger rates calculated from [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10). In all cases we see that the true likelihood is well represented within our distribution of 170 subsamples, indicating our use of a relatively smaller number of 10⁶ binaries is not expected to bias our results.

ideal for generating the current LVK detected population. We verify this with the likelihoods in Section [5.2.](#page-14-1)

5.2 Bayes Factor and Likelihood Analysis

To identify groups of models that are more preferred than others, we use the Bayes factor, β . Since the priors for all models are the same, the Bayes factor is the ratio of the likelihoods between two models, here denoted α and κ :

$$
\mathcal{B}(\lambda|\mathcal{H}) = \frac{\mathcal{L}(\lambda_{\alpha}|\mathcal{H})}{\mathcal{L}(\lambda_{\kappa}|\mathcal{H})}; \qquad \mathcal{B}(\lambda, N_{\lambda}|\mathcal{H}) = \frac{\mathcal{L}(\lambda_{\alpha}, N_{\lambda_{\alpha}}|\mathcal{H})}{\mathcal{L}(\lambda_{\kappa}, N_{\lambda_{\kappa}}|\mathcal{H})}. \quad (31)
$$

In Fig. [12,](#page-16-0) we show the Bayes factor with the likelihoods using the GW events with the SNR ≥ 8 cut.

We note that the Bayes factor indicates there is strong evidence that some groups of models are preferred more than others, which we will further discuss in the likelihood analysis. We emphasise these with highlighted boxes in Fig. [12](#page-16-0) labeled A-G. We can understand the range of Bayes factor by considering the different components that enter Eq [31.](#page-14-2) The difference in $\log \beta(\lambda)$ between different models ranges from ~ 0.05 – 3.44. However, we find that the I^k terms varies over 50 orders of magnitude (see bottom plot in Fig. [18\)](#page-19-1) and therefore dominates the Bayes factor. This confirms that some of the models are distinguishable and strongly preferred (more likely to produce progenitors of particular GW events) over others purely based on the masses and redshift distributions they predict. When including N_{λ} as a parameter in the likelihood, the $-\mu_{\lambda} + N_{\text{obs}} \log(\mu_{\lambda})$ terms this introduces into the log-likelihood has significant weight, and comes to dominate the Bayes factor, meaning that a comparison of two models using this likelihood strongly favours those that produce a number of events close to the observed number of events in our data sample, even if the event-level distributions are less well matched. Hence, we carefully consider the results using both of our likelihoods.

Having established groups of models that are strongly preferred over other models, we now discuss the likelihoods for individual cases and the trends within these groups. The following results discussed use the $SNR > 8$ selection cut. We see an increase in the likelihood across all models, but the trends don't change substantially with the selection criteria and we find the differences are made more distinct when using the SNR \geq 8 sample. We show this in Tables [C1,](#page-24-0) [C2](#page-25-0) and C₃ with Bayes factors between our various models.

We see an increase in the likelihood from $f_{\gamma} = 0$ to $f_{\gamma} = 0.5$ and we find that the likelihoods are similar with no CHE and pessimistic and optimistic CHE. In Fig. [13,](#page-16-1) we show the likelihoods for fixed f_{γ} . It is clear that $f_{\gamma} = 0.5$ is preferred over $f_{\gamma} = 1.0$. Less BBH mergers are produced with $f_{\gamma} = 1.0$. Models with $f_{\gamma} = 0.5$ produce more equal mass binaries, which is more consistent with the LVK population [\(Abbott et al.](#page-20-19) [2019b\)](#page-20-19). However, $f_{\gamma} = 0.75$ is unique in that for low mass transfer $f_{\text{MT}} < 0.5$ the population is disfavoured similar to $f_{\gamma} = 1.0$, but for high mass transfer $f_{\text{MT}} > 0.5$ the model is more favourable. In all cases, we see little impact from the choice of common envelope efficiency. This highlights the degeneracy between stellar modelling assumptions and populations. Ultimately, we find that in order to reproduce the observed events from LVK, we need either momentum loss to occur close to the accretor, or, if occurring further away, for the mass transfer fraction to be high. These trends are also highlighted by boxes A and B in Fig. [12,](#page-16-0) where moderate values of f_{γ} are strongly favoured when comparing population parameters.

Fig. [14](#page-16-2) then explores different combinations of the remnant mass prescription, CHE mode and Wolf–Rayet multiplier. Overall, we see weaker trends here than with the mass transfer fraction. However, the remnant mass prescription can be seen to impact the likelihood, with the [Fryer et al.](#page-21-47) [\(2012\)](#page-21-47) or [Mandel & Müller](#page-21-11) [\(2020\)](#page-21-11) prescriptions preferred. It is worth reiterating that, of these two, the [Fryer et al.](#page-21-47) [\(2012\)](#page-21-47) was seen to greatly over-predict the number of observed events to date. We find a weak trend of increasing likelihood in the absence of CHE, most evident for the [Schneider et al.](#page-22-34) [\(2021\)](#page-22-34) model, but overall less significant an impact than was seen for changes in the mass transfer and AM loss parameters. We also see a weak preference for larger f_{WR} depending on the remnant mass model.

Figure 10. Detected number of BBH merger events versus the mass transfer fraction for no CHE at fixed f_y . Left: We fix the common envelope evolution parameter $\alpha_{\text{CE}} = 0.1$ and show results for different specific angular momentum loss parameters. *Right*: We show the models for any f_{γ} and α_{CE} which output μ_{λ} close to the number of observed LVK events given by the grey band.

Figure 11. Detected number of BBH merger events versus the Wolf–Rayet multiplier for fixed CHE modes and the [Schneider et al.](#page-22-34) [\(2021\)](#page-22-34) remnant mass prescription. The grey band is the observed number of LVK events.

5.2.1 Likelihood including

When including the dependence on the number of GW events, we are able to identify more distinct groups of trends, particularly with the CHE modes as shown in Fig. [15.](#page-17-0) This is in contrast to the likelihoods without the number of events, indicating that the CHE mode more strongly affects just the number of BBH mergers, rather than the underlying distributions of mass or redshift. The likelihood generally reduces when switching from no CHE to Pessimistic or Optimistic, which is not as notable for the likelihood without N_{λ} dependence. We also see this impact in Fig. [12,](#page-16-0) where box C and D correspond

to no CHE being strongly preferred over Pessimistic and Optimistic CHE and box E implies Pessimistic is strongly preferred over Optimistic CHE. Allowing for CHE would mean tighter orbits and more mergers within Hubble time, which would over-predict the number of observed events when compared to N_{obs} . In addition, the likelihood now decreases from $f_{\gamma} = 0.25$ to $f_{\gamma} = 0.5$. This is also due to the increased number of GW events, shifting them further from the observed number of LVK events. It then increases again with further increasing $f_{\gamma} = 0.75, 1.0$ due to the reduced numbers of events. Within each choice of CHE, we see a preference for low mass transfer efficiency for $f_{\gamma} = 0$. This is required to keep the predicted number of events low or comparable to the actual number of LVK events, as these produce unequal mass wide binaries with long merger times.

Although no CHE is preferred, this can depend on f_{γ} . We show a subset of no CHE with varying f_{γ} in Fig. [15.](#page-17-0) Again we note the degeneracy in the hyper-parameters and populations. Low values of f_{MT} and $f_{\gamma} = 0.25$ produce similar populations to $f_{\gamma} = 0$, while models with higher values of f_{MT} at $f_{\gamma} = 0.25$ appear similar to Pessimistic CHE with $f_{\gamma} = 0$ populations. Conversely, further increases to $f_{\gamma} = 0.5$ and *high* f_{MT} produce high-likelihood populations similar to $f_{\gamma} = 0$, with now a decreasing likelihood as f_{MT} decreases. The likelihood further increases with $f_{\gamma} = 0.75$ particularly again for high f_{MT} . There is hence a flip in the behaviour of increasing mass transfer fraction depending on whether the momentum loss is occurring closer to the accretor or L2 point. These changes in likelihood closely reflect the trends seen in the number of predicted events seen in Fig. [10,](#page-15-0) but further understanding of this complex behaviour warrants further study.

Finally, in Fig. [16,](#page-18-0) we find a strong case for likelihood increasing with f_{WR} , with the exception of [Schneider et al.](#page-22-34) [\(2021\)](#page-22-34) with no CHE. This is also shown in Fig. [12](#page-16-0) by boxes F and G, where $f_{\gamma} = 0$ and thermal timescale mass transfer rates are strongly preferred over moderate values of f_{γ} and fixed f_{MT} . For fixed CHE and remnant mass prescription, there is a preference for models with $f_{WR} = 10$. This is because mass loss via strong winds can widen binaries. This would increase the merger time and decrease the number of GW

Figure 12. Comparison of models with SNR \geq 8 using the Bayes factor. Each grid is coloured by $\log_{10}(\mathcal{B})$. For $\log_{10}(\mathcal{B}) > 0$, the model in the row (y-axis) is favoured over than the model in the column (x-axis), while for $\log_{10}(\mathcal{B}) < 0$ the model in the row is disfavoured over the model in the column. Left plot is the Bayes factor for the marginalised likelihood and right plot is the Bayes factor for the unmarginalised likelihood.

Figure 13. Likelihoods for models with no CHE. Each band corresponds to a fixed f_{γ} . Within each band, excluding the Rauf et al. (2023) models in blue, the mass transfer fraction and common envelope efficiency are increasing with increasing model number, within the ranges in Table [1,](#page-6-1) with f_{MT} varying more rapidly.

events closer to the LVK detected number of events. Overall, again no CHE is preferred over pessimistic and optimistic CHE — in the presence of CHE, one needs a rather finely-tuned set of: high Wolf– Rayet factor, high f_{γ} and low mass transfer efficiency to recover a reasonable number of mergers.

Figure 14. Marginalised likelihood for varying CHE and remnant mass prescription. In each band, the Wolf–Rayet multiplier is increased with increasing model number, across the values given in Table [1.](#page-6-1)

Given the main assumption in this paper being all GW events originate from the isolated binary formation channel, the observed number of events in the LVK data places an upper bound on μ_{λ} . Any models that output $\leq N_{\text{obs}}$ cannot be ruled out. However, even in the case where the number of detected events is close to N_{obs} , this is not a good indicator of if the model population will match

Figure 15. Model versus unmarginalised likelihood, where each coloured band corresponds to the a fixed CHE mode and f_{γ} . Within each band, excluding the [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) models, the mass transfer fraction and common envelope efficiency are increasing with increasing model number, within the ranges in Table [1,](#page-6-1) with f_{MT} varying more rapidly.

the observed population. We propose checking the marginalised and unmarginalised likelihood simultaneously to find if there is overlap in models. This will determine the models with comparable populations *and* detected number of events to the LVK data. We check the 30 largest likelihoods out of the 300 new models for both these cases separately, and identify the common models and their hyperparameters. We again find a clear preference for no CHE for $f_{\text{WR}} \geq 1$ and $f_y = 0$, in that these models often return representative numbers of detected events, *and* populations of masses and redshifts, but the choice of remnant mass prescription varies between Mandel $\&$ [Müller](#page-21-11) [\(2020\)](#page-21-11), [Fryer et al.](#page-21-47) [\(2012\)](#page-21-47) and [Fryer et al.](#page-21-53) [\(2022\)](#page-21-53).

5.2.2 Comparisons of the best individual models to data

We now compare the visual population parameter distributions to the GW events with $SNR \geq 8$ posteriors for the 300 new models. Model 44 has the best likelihood $(\mathcal{L}(\lambda|\mathcal{H}))$ which implies that the model population best matches the observed LVK population. As shown in Fig. [17,](#page-18-1) we find that the chirp mass and mass ratio distribution falls within the 2σ contour of the GW events. The LVK chirp mass distribution has three distinct peaks. The best model does not replicate the

first peak but it does for the second and third and generally matches the overall chirp mass distribution. That said, the mass ratio distribution is uniform for the LVK data but the model does prefer high mass ratios ($q > 0.6$). Model 79 outputs the worst likelihood. The chirp mass and mass ratio distributions have very narrow peaks corresponding to the second peak in the LVK mass distribution around $M_c = 20M_{\odot}$ and at $q = 0.5$. The redshift distribution range is also smaller compared to Model 44 and the data. However, the number of detected events, μ_{λ} is closer to $N_{\text{obs}} = 79$. This demonstrates the intrinsic difficulty in population synthesis of returning the correct event-level properties and overall GW rate.

The best likelihood with a dependence on the number of events $(L(\lambda, N_{\lambda}|\mathcal{H}))$ is Model 274. This model has a peak in the chirp mass distribution corresponding to the first peak in the LVK mass distribution and recovers the higher mass peaks reasonably well too, albeit with absence in the very high mass tail. The mass ratio distribution is not uniform, there is a preference for mass ratios > 0.5 , but < 0.5 are also possible such that the model and data distributions agree reasonably well. This model outputs $\mu_{\lambda} \approx 83$ events. The worst likelihood is model 199. This model has has a rather high peak chirp mass around $40M_{\odot}$ and a strong preference for high mass ratios $(q > 0.8)$. It is likely that as a result of producing higher mass BHs with smaller separations, this models outputs the largest detected number of events, ∼ 3805, which negatively impacts the likelihood.

In all cases, our models seem to underpredict the high chirp mass and high mass ratio events with $z \geq 0.5$ when compared to the LVK distribution. It is possible that these events are not produced frequently by COMPAS. This may be a limitation in the SSE implementation, as discussed in Section [2.4.](#page-5-0) It is likely that these systems are not produced efficiently through the isolated channel, which could indicate that some of our LVK events progenitors form dynamically. And finally, these events may just be rare or 'exceptional' in any formation channel/ environment. In this case, approaches more like [Passenger et al.](#page-21-79) [\(2024\)](#page-21-79) — which uses the "normalised evidence" within a statistical framework to determine if hierarchical mergers in active galactic nuclei or globular clusters can justify exceptionally large events such as GW190521 — may be more appropriate.

5.3 Which data are the most informative?

Given our range of models and the form with which the likelihood is evaluated by summing over each GW event, we can also ask the question of which events allow for the greatest distinction between our various COMPAS models. For each GW event, we evaluate the distribution of $\log(\mathcal{I}^k)$ terms for all 300 models. We use the range and mean of the distributions to determine if the progenitors of the GW events are distinguishable between models. We also look for skewed distributions, which indicate that some models are more or less likely to produce progenitors of particular GW events. We show an example of 20 of these — the top with the least variance between models, and the ten with the most — in Fig. [18.](#page-19-1)

The top plot of Fig. [18](#page-19-1) shows the GW events where their \mathcal{I}^k terms are tightly constrained to high probabilities. This implies that all the models are very likely to produce the progenitors of these events. The distributions appear Gaussian. However, upon focusing on the $\log(I^k)$ < -5 region they appear to have a right skewed distribution, implying there a few models in this tail that are less preferred to form the progenitors of these events.

The bottom plot shows the GW events where the progenitors populations are more clearly distinguishable between models. In particular, GW190814 and GW190521 have wide bimodal distributions, apparently separating our COMPAS models into those that can, and

Figure 16. Unmarginalised likelihood for No CHE (*Left*) and pessimistic CHE (*Right*) for fixed remnant mass prescriptions. Within each band we vary the Wolf–Rayet multiplier with values given in Table [1,](#page-6-1) increasing the value with increasing model number.

Figure 17. M_c , q and z 3 σ distributions for the models with the best (green contour) and worst (red contour) likelihood. On the left we show the likelihood without the N_A dependence and the right plot shows the likelihood with the N_A dependence. The blue contour combines the posteriors for the $N_{obs} = 79$ GW events based on the $SNR \geq 8$ cut.

those that cannot, reproduce the characteristics of these events. In Figure [19](#page-19-2) we show these events with low probability of their progenitors forming in some models. They appear to be close to, or in, the pair-instability mass gap \sim 50 − 130 M_{\odot} [\(Woosley & Heger](#page-22-47) [2021\)](#page-22-47). GW190426_190642 has a total mass that supersedes the total mass of GW190521, and it could be argued that they are from

the same formation channel. However, GW190426_190642 is less certain to be a true event, having $p_{astro} = 0.75$ and $FAR = 4.1yr^{-1}$ [\(Banerjee](#page-20-20) [2022\)](#page-20-20). GW190602_175927 and GW191109_010717 have at least one component in the mass gap. The spin-orbit misalignment of GW191109_010717 provides further evidence it has a dynamical origin [\(Zhang et al.](#page-22-48) [2023\)](#page-22-48). However, GW191109_010717 has been

Figure 18. Distribution of $\log(I^k)$ terms for the 300 new COMPAS models. The top plot shows the GW events where the distribution are constrained to high log I^k values and the bottom plot shows GW events with broad distributions. Out of the distinguishable events, the pink distributions correspond to events with one component in the lower mass gap and the green distributions correspond to events with at least one component in the upper mass gap.

flagged as problematic due to data quality issues, which may lead to unreliable inference of the source parameters such as the precession [\(Tong et al.](#page-22-49) [2022\)](#page-22-49). GW190917_114630 and GW200210_092254 have a secondary component in the lower mass gap between the most massive neutron star and smallest BH, like GW190814. The low probability of generating these events by some models indicate that the isolated formation channel is not ideal for forming unequal mass binaries. However, the ability of some models to form the progenitors of these events suggests that this mass gap might be narrower or non-existent. More observations of similar events is paramount in confirming this [\(Zevin et al.](#page-22-50) [2020\)](#page-22-50).

Overall, looking at the models that perform well for these events, they tend to produce a dearth of higher mass BHs, with chirp mass distributions that peak around 40 M_{\odot} and mass ratio distributions that peak around $q \approx 0.9$. However, this often comes at the expense of producing enough lower mass or more extreme mass ratio binaries, to explain the mass and redshift distribution of the other GW events.

This is particularly emphasised for GW190521 where we found that the posterior samples often did not fall at all within the chirp mass or mass ratio range output from COMPAS. In these cases we assign a probability of zero to these events such that they make no difference to the sum in Eq [21](#page-9-1) and [26,](#page-9-3) but are also hence not represented in

Figure 19. Posterior distributions for 'outlier' events, (identified by the green distributions in Fig. [18\)](#page-19-1) that are not well reproduced by the majority of our COMPAS models. The blue contour is the full posterior distribution for LVK data with $SNR \geq 8$, which clearly indicates the majority of the data are at both lower mass and lower redshift. We show GW190426_190642 as an unfilled contour, as it does not fit the $p_{astro} \ge 0.9$ and FAR ≤ 0.25 yr⁻¹ criteria.

Fig. [18.](#page-19-1) Model choices that fail to produce any progenitor close to GW190521 include:

• No CHE (i.e., models 1-80 and 241-250), implying that formation channels with mass transfer and CE evolution produce smaller BH masses.

• The remnant mass prescriptions [Mandel & Müller](#page-21-11) [\(2020\)](#page-21-11) and [Schneider et al.](#page-22-34) [\(2021\)](#page-22-34), which tend to produce smaller BH masses. The exception being model 271-280, which use [Fryer et al.](#page-21-47) [\(2012\)](#page-21-47) and [Fryer et al.](#page-21-48) [\(2021\)](#page-21-48).

• Models 221 and 270, where CHE is set to Optimistic, but f_{γ} = 0.75 (Model 221) and/or $f_{WR} = 10$ (Model 270), which may be contributing to extreme mass loss from the progenitors again leading to a population of particularly low-mass BHs.

Overall, the difficulty in simultaneously producing both these larger events and the bulk of LVK data even across the large range of models we explore in this work lends credence to the idea of multiple formation pathways for these events.

6 CONCLUSION

In this work we present a thorough analysis of the ability of over 300 binary population synthesis models to reproduce the GW events observed so far within Gravitational Wave Transients Catalogues 1-3. To enable this, we developed an efficient method for predicting the BBH merger rate in the isolated channel for various COMPAS models (Section [3\)](#page-6-0), validated based on previous work from [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10). We then used this within a Bayesian inference method to calculate the likelihood of these models with and without the dependence on the number of GW events predicted by each model, N_{λ} (Section [4\)](#page-8-0).

The COMPAS population synthesis code contains a number of free

parameters — we introduce some of the most salient of these in Section [2.](#page-2-0) Our results fitting these models to data in Section [5](#page-12-0) demonstrate there is a preference for subsets of models dictated by the type of isolated formation channel and specific angular momentum loss. However, we frequently find opposing trends in the likelihoods with and without N_{λ} dependence — models that produce realistic event-level masses and redshifts frequently over-predict the number of events we should have observed to date. Looking at models which perform well with both likelihoods, we find that models with no Chemically Homogeneous Evolution, low to moderate values of f_{γ} (such that angular momentum loss occurs closer to the donor) and higher rates of mass-loss from Wolf–Rayet winds (larger f_{WR}) are generally preferred by current data.

We find that the selection effects impact our likelihoods, as the $SNR \geq 8$ cut on the GW events increases the likelihood for *all* models. This may vary with future GW detections, particularly if the BH masses are in the mass gaps. We also use the individual event likelihoods I^k across all our models term to distinguish individual events that are easy or more difficult to reproduce by our models. We identify the usual suspects GW190521 and GW190814 as typically having low probability of existing in our new models, as well as some outlier events in the the lower and upper mass gap. This motivates future studies to consider joint population inference and model selection studies across multiple formation channels [\(Gompertz et al.](#page-21-80) [2022;](#page-21-80) [Gayathri et al.](#page-21-81) [2023\)](#page-21-81).

Our method for efficiently calculating the merger rate can also be applied to other population synthesis codes. However, we note that it is only applicable in the case where the metallicity dependent SFR is the same for the fiducial and new model. We argue that using a different SFR may not impact the PDF significantly. [van Son et al.](#page-22-51) [\(2023\)](#page-22-51) finds that variation in the SFR do not impact the features in the primary BH mass distribution and have minimal impact on the low mass end of the distribution.

We acknowledge this work relies on various assumptions and simplifications. For future work, we consider including the effective spin as a population parameter in the analysis. The covariance between the mass ratio and effective spin can arise due to common envelope physics, mass transfer efficiency and Eddington limited accretion [\(Mandel & Fragos](#page-21-82) [2020;](#page-21-82) [Bavera et al.](#page-20-6) [2021;](#page-20-6) [Zevin & Bavera](#page-22-52) [2022;](#page-22-52) [Broekgaarden et al.](#page-21-83) [2022b;](#page-21-83) [Adamcewicz et al.](#page-20-21) [2023;](#page-20-21) [Olejak et al.](#page-21-84) [2024\)](#page-21-84). This could be used to further distinguish between COMPAS models. Alternatively, we can build the PDF only with M_c , mass ratio or redshift to test the validity of the model outputs. We could also consider including natal kick velocity as one of the hyper-parameters to vary in COMPAS in the future, as this is not well constrained for BHs [\(Callister et al.](#page-21-85) [2021;](#page-21-85) [Cui & Li](#page-21-30) [2023;](#page-21-30) [Vigna-Gómez et al.](#page-22-53) [2024\)](#page-22-53), or more fundamental parameters such as the binary fraction. Varying the selection function cuts applied to current data to restrict the analysis to only the strongest detections may further inform us on trends but given we found consistent results with our two different selection functions this may not necessarily provide improved constraints on the hyper-parameters.

Future GW detectors will observe GW events at a wider range of frequencies and redshifts, and will provide data with increasing precision [\(Broekgaarden et al.](#page-21-71) [2023\)](#page-21-71). This will answer a variety of astrophysical questions and provide a clearer picture of the BBH population distribution. In the absence of electromagnetic counterparts, BBH merger populations with the *correct* modelling are crucial for an unbiased inference of cosmological parameters [\(Mastrogiovanni](#page-21-86) [et al.](#page-21-86) [2021;](#page-21-86) [Gray et al.](#page-21-87) [2023\)](#page-21-87). With deeper catalogues and higher SNR detections, our selection criteria will be more refined, which will hopefully allow us to further constrain the uncertainties in massive binary evolution models. However, as always, with more GW events comes greater change of observing rare exceptional events to challenge these models. Until then, in the case of limited number of GW events we must rely on observations of massive binary stars to constrain the different assumptions for massive binary evolution. Observations of X-ray binaries, red supergiants, luminous blue variables and Wolf–Rayet stars can help constrain binaries at different evolutionary stages [\(Eldridge & Stanway](#page-21-88) [2022;](#page-21-88) [Marchant & Bodensteiner](#page-21-89) [2023;](#page-21-89) [Li et al.](#page-21-90) [2024\)](#page-21-90).

ACKNOWLEDGEMENTS

We thank Carole Perigois and Michella Mapelli for useful discussion on their paper that motivated this work. We thank Ilya Mandel, Eric Thrane and Paul Lasky for useful discussions on the methodology and results. We also thank Reinhold Wilcox and Floor Broekgaarden for useful discussions on the COMPAS hyper-parameters and relevant literature.

The authors acknowledge support from the Australian Research Council (ARC) Centre of Excellence for Gravitational Wave Discovery (OzGrav), through project numbers CE170100004 and CE230100016. LR and CH acknowledge support through ARC Discovery Project DP20220101395. SS is a recipient of an ARC Discovery Early Career Research Award (DE220100241).

DATA AVAILABILITY

Prior and posterior samples of GW events were provided by Carole Perigois. Simulations in this paper made use of the COMPAS rapid binary population synthesis code (version v02.38.08), which is freely available at [this repository.](http://github.com/TeamCOMPAS/COMPAS)

REFERENCES

- Abbott B. P., et al., 2017, [Phys. Rev. Lett.,](http://dx.doi.org/10.1103/PhysRevLett.119.161101) [119, 161101](https://ui.adsabs.harvard.edu/abs/2017PhRvL.119p1101A)
- Abbott B. P., et al., 2019a, [Physical Review X,](http://dx.doi.org/10.1103/PhysRevX.9.031040) [9, 031040](https://ui.adsabs.harvard.edu/abs/2019PhRvX...9c1040A)
- Abbott B. P., et al., 2019b, [ApJ,](http://dx.doi.org/10.3847/2041-8213/ab3800) [882, L24](https://ui.adsabs.harvard.edu/abs/2019ApJ...882L..24A)
- Abbott R., et al., 2020a, [Phys. Rev. Lett.,](http://dx.doi.org/10.1103/PhysRevLett.125.101102) [125, 101102](https://ui.adsabs.harvard.edu/abs/2020PhRvL.125j1102A)
- Abbott B. P., et al., 2020b, [ApJ,](http://dx.doi.org/10.3847/2041-8213/ab75f5) [892, L3](https://ui.adsabs.harvard.edu/abs/2020ApJ...892L...3A)
- Abbott R., et al., 2020c, [ApJ,](http://dx.doi.org/10.3847/2041-8213/ab960f) [896, L44](https://ui.adsabs.harvard.edu/abs/2020ApJ...896L..44A)
- Abbott R., et al., 2021a, [Physical Review X,](http://dx.doi.org/10.1103/PhysRevX.11.021053) [11, 021053](https://ui.adsabs.harvard.edu/abs/2021PhRvX..11b1053A)
- Abbott R., et al., 2021b, [ApJ,](http://dx.doi.org/10.3847/2041-8213/ac082e) [915, L5](https://ui.adsabs.harvard.edu/abs/2021ApJ...915L...5A)
- Abbott R., et al., 2023a, [Physical Review X,](http://dx.doi.org/10.1103/PhysRevX.13.011048) [13, 011048](https://ui.adsabs.harvard.edu/abs/2023PhRvX..13a1048A)
- Abbott R., et al., 2023b, [Physical Review X,](http://dx.doi.org/10.1103/PhysRevX.13.041039) [13, 041039](https://ui.adsabs.harvard.edu/abs/2023PhRvX..13d1039A)
- Adamcewicz C., Lasky P. D., Thrane E., 2023, [ApJ,](http://dx.doi.org/10.3847/1538-4357/acf763) [958, 13](https://ui.adsabs.harvard.edu/abs/2023ApJ...958...13A)
- Agrawal P., Hurley J., Stevenson S., Szécsi D., Flynn C., 2020, [MNRAS,](http://dx.doi.org/10.1093/mnras/staa2264) [497,](https://ui.adsabs.harvard.edu/abs/2020MNRAS.497.4549A) [4549](https://ui.adsabs.harvard.edu/abs/2020MNRAS.497.4549A)
- Agrawal P., Szécsi D., Stevenson S., Eldridge J. J., Hurley J., 2022a, [MNRAS,](http://dx.doi.org/10.1093/mnras/stac930) [512, 5717](https://ui.adsabs.harvard.edu/abs/2022MNRAS.512.5717A)
- Agrawal P., Stevenson S., Szécsi D., Hurley J., 2022b, [A&A,](http://dx.doi.org/10.1051/0004-6361/202244044) [668, A90](https://ui.adsabs.harvard.edu/abs/2022A&A...668A..90A)
- Andres N., et al., 2022, [Classical and Quantum Gravity,](http://dx.doi.org/10.1088/1361-6382/ac482a) [39, 055002](https://ui.adsabs.harvard.edu/abs/2022CQGra..39e5002A)
- Arca Sedda M., 2021, [ApJ,](http://dx.doi.org/10.3847/2041-8213/abdfcd) [908, L38](https://ui.adsabs.harvard.edu/abs/2021ApJ...908L..38A)
- Arca Sedda M., Mapelli M., Benacquista M., Spera M., 2023, [MNRAS,](http://dx.doi.org/10.1093/mnras/stad331) [520,](https://ui.adsabs.harvard.edu/abs/2023MNRAS.520.5259A) [5259](https://ui.adsabs.harvard.edu/abs/2023MNRAS.520.5259A)
- Asplund M., Grevesse N., Sauval A. J., Scott P., 2009, [ARA&A,](http://dx.doi.org/10.1146/annurev.astro.46.060407.145222) [47, 481](https://ui.adsabs.harvard.edu/abs/2009ARA&A..47..481A)
- Banagiri S., Berry C. P. L., Davies G. S. C., Tsukada L., Doctor Z., 2023, [Phys. Rev. D,](http://dx.doi.org/10.1103/PhysRevD.108.083043) [108, 083043](https://ui.adsabs.harvard.edu/abs/2023PhRvD.108h3043B)
- Banerjee S., 2022, [A&A,](http://dx.doi.org/10.1051/0004-6361/202142331) [665, A20](https://ui.adsabs.harvard.edu/abs/2022A&A...665A..20B)
- Barrett J. W., Gaebel S. M., Neijssel C. J., Vigna-Gómez A., Stevenson S., Berry C. P. L., Farr W. M., Mandel I., 2018, [MNRAS,](http://dx.doi.org/10.1093/mnras/sty908) [477, 4685](https://ui.adsabs.harvard.edu/abs/2018MNRAS.477.4685B)
- Bavera S. S., et al., 2021, [A&A,](http://dx.doi.org/10.1051/0004-6361/202039804) [647, A153](https://ui.adsabs.harvard.edu/abs/2021A&A...647A.153B)
- Belczynski K., Bulik T., Fryer C. L., Ruiter A., Valsecchi F., Vink J. S., Hurley J. R., 2010, [ApJ,](http://dx.doi.org/10.1088/0004-637X/714/2/1217) [714, 1217](https://ui.adsabs.harvard.edu/abs/2010ApJ...714.1217B)
- Björklund R., Sundqvist J. O., Singh S. M., Puls J., Najarro F., 2023, [A&A,](http://dx.doi.org/10.1051/0004-6361/202141948) [676, A109](https://ui.adsabs.harvard.edu/abs/2023A&A...676A.109B)
- Bogomazov A. I., Cherepashchuk A. M., Lipunov V. M., Tutukov A. V., 2018, [New Astron.,](http://dx.doi.org/10.1016/j.newast.2017.07.004) [58, 33](https://ui.adsabs.harvard.edu/abs/2018NewA...58...33B)
- Bouffanais Y., Mapelli M., Gerosa D., Di Carlo U. N., Giacobbo N., Berti E., Baibhav V., 2019, [ApJ,](http://dx.doi.org/10.3847/1538-4357/ab4a79) [886, 25](https://ui.adsabs.harvard.edu/abs/2019ApJ...886...25B)
- Bouffanais Y., Mapelli M., Santoliquido F., Giacobbo N., Iorio G., Costa G., 2021, [MNRAS,](http://dx.doi.org/10.1093/mnras/stab1589) [505, 3873](https://ui.adsabs.harvard.edu/abs/2021MNRAS.505.3873B)
- Breivik K., et al., 2020, [ApJ,](http://dx.doi.org/10.3847/1538-4357/ab9d85) [898, 71](https://ui.adsabs.harvard.edu/abs/2020ApJ...898...71B)
- Broekgaarden F. S., et al., 2022a, [MNRAS,](http://dx.doi.org/10.1093/mnras/stac1677) [516, 5737](https://ui.adsabs.harvard.edu/abs/2022MNRAS.516.5737B)
- Broekgaarden F. S., Stevenson S., Thrane E., 2022b, [ApJ,](http://dx.doi.org/10.3847/1538-4357/ac8879) [938, 45](https://ui.adsabs.harvard.edu/abs/2022ApJ...938...45B)
- Broekgaarden F. S., Banagiri S., Payne E., 2023, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2303.17628) [p.](https://ui.adsabs.harvard.edu/abs/2023arXiv230317628B) [arXiv:2303.17628](https://ui.adsabs.harvard.edu/abs/2023arXiv230317628B)
- Cabourn Davies G. S., Harry I. W., 2022, [Classical and Quantum Gravity,](http://dx.doi.org/10.1088/1361-6382/ac8862) [39, 215012](https://ui.adsabs.harvard.edu/abs/2022CQGra..39u5012C)
- Callister T. A., Kanner J. B., Massinger T. J., Dhurandhar S., Weinstein A. J., 2017, [Classical and Quantum Gravity,](http://dx.doi.org/10.1088/1361-6382/aa7a76) [34, 155007](https://ui.adsabs.harvard.edu/abs/2017CQGra..34o5007C)
- Callister T. A., Farr W. M., Renzo M., 2021, [ApJ,](http://dx.doi.org/10.3847/1538-4357/ac1347) [920, 157](https://ui.adsabs.harvard.edu/abs/2021ApJ...920..157C)
- Cheng A. Q., Zevin M., Vitale S., 2023a, [ApJ,](http://dx.doi.org/10.3847/1538-4357/aced98) [955, 127](https://ui.adsabs.harvard.edu/abs/2023ApJ...955..127C)
- Cheng A. Q., Zevin M., Vitale S., 2023b, [Astrophys. J.,](http://dx.doi.org/10.3847/1538-4357/aced98) 955, 127
- Chruślińska M., 2024, [Annalen der Physik,](http://dx.doi.org/10.1002/andp.202200170) [536, 2200170](https://ui.adsabs.harvard.edu/abs/2024AnP...53600170C)
- Claret A., Torres G., 2018, [ApJ,](http://dx.doi.org/10.3847/1538-4357/aabd35) [859, 100](https://ui.adsabs.harvard.edu/abs/2018ApJ...859..100C)
- Costa G., Girardi L., Bressan A., Marigo P., Rodrigues T. S., Chen Y., Lanza A., Goudfrooij P., 2019, [MNRAS,](http://dx.doi.org/10.1093/mnras/stz728) [485, 4641](https://ui.adsabs.harvard.edu/abs/2019MNRAS.485.4641C)
- Costa G., Chruślińska M., Klencki J., Broekgaarden F. S., Rodriguez C. L., Joseph T. D., Saracino S., 2023, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2311.15778) [p. arXiv:2311.15778](https://ui.adsabs.harvard.edu/abs/2023arXiv231115778C)
- Cui Z., Li X.-D., 2023, [MNRAS,](http://dx.doi.org/10.1093/mnras/stad1800) [523, 5565](https://ui.adsabs.harvard.edu/abs/2023MNRAS.523.5565C)
- Delfavero V., O'Shaughnessy R., Belczynski K., Drozda P., Wysocki D., 2023, [Phys. Rev. D,](http://dx.doi.org/10.1103/PhysRevD.108.043023) [108, 043023](https://ui.adsabs.harvard.edu/abs/2023PhRvD.108d3023D)
- Detmers R. G., Langer N., Podsiadlowski P., Izzard R. G., 2008, [A&A,](http://dx.doi.org/10.1051/0004-6361:200809371) [484,](https://ui.adsabs.harvard.edu/abs/2008A&A...484..831D) [831](https://ui.adsabs.harvard.edu/abs/2008A&A...484..831D)
- Dorozsmai A., Toonen S., 2022, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2207.08837) [p. arXiv:2207.08837](https://ui.adsabs.harvard.edu/abs/2022arXiv220708837D)
- Eldridge J. J., Stanway E. R., 2022, [ARA&A,](http://dx.doi.org/10.1146/annurev-astro-052920-100646) [60, 455](https://ui.adsabs.harvard.edu/abs/2022ARA&A..60..455E)
- Ertl T., Janka H. T., Woosley S. E., Sukhbold T., Ugliano M., 2016, [ApJ,](http://dx.doi.org/10.3847/0004-637X/818/2/124) [818,](https://ui.adsabs.harvard.edu/abs/2016ApJ...818..124E) [124](https://ui.adsabs.harvard.edu/abs/2016ApJ...818..124E)
- Farr W. M., Gair J. R., Mandel I., Cutler C., 2015, [Phys. Rev. D,](http://dx.doi.org/10.1103/PhysRevD.91.023005) [91, 023005](https://ui.adsabs.harvard.edu/abs/2015PhRvD..91b3005F)
- Fishbach M., Kalogera V., 2022, [ApJ,](http://dx.doi.org/10.3847/2041-8213/ac64a5) [929, L26](https://ui.adsabs.harvard.edu/abs/2022ApJ...929L..26F)
- Fishbach M., Holz D. E., Farr W. M., 2018, [ApJ,](http://dx.doi.org/10.3847/2041-8213/aad800) [863, L41](https://ui.adsabs.harvard.edu/abs/2018ApJ...863L..41F)
- Fryer C. L., Belczynski K., Wiktorowicz G., Dominik M., Kalogera V., Holz D. E., 2012, The Astrophysical Journal, 749, 91
- Fryer C. L., Karpov P., Livescu D., 2021, [Astronomy Reports,](http://dx.doi.org/10.1134/S1063772921100103) [65, 937](https://ui.adsabs.harvard.edu/abs/2021ARep...65..937F)
- Fryer C. L., Olejak A., Belczynski K., 2022, [ApJ,](http://dx.doi.org/10.3847/1538-4357/ac6ac9) [931, 94](https://ui.adsabs.harvard.edu/abs/2022ApJ...931...94F)
- Galaudage S., Talbot C., Thrane E., 2020, [Phys. Rev. D,](http://dx.doi.org/10.1103/PhysRevD.102.083026) [102, 083026](https://ui.adsabs.harvard.edu/abs/2020PhRvD.102h3026G)
- Gallegos-Garcia M., Berry C. P. L., Marchant P., Kalogera V., 2021, [ApJ,](http://dx.doi.org/10.3847/1538-4357/ac2610) [922, 110](https://ui.adsabs.harvard.edu/abs/2021ApJ...922..110G)
- Gallegos-Garcia M., Jacquemin-Ide J., Kalogera V., 2023, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2308.13146) [p.](https://ui.adsabs.harvard.edu/abs/2023arXiv230813146G) [arXiv:2308.13146](https://ui.adsabs.harvard.edu/abs/2023arXiv230813146G)
- Gayathri V., Bartos I., Rosswog S., Miller M. C., Veske D., Lu W., Marka S., 2023, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2307.09097) [p. arXiv:2307.09097](https://ui.adsabs.harvard.edu/abs/2023arXiv230709097G)
- Ge H., Hjellming M. S., Webbink R. F., Chen X., Han Z., 2010, [ApJ,](http://dx.doi.org/10.1088/0004-637X/717/2/724) [717,](https://ui.adsabs.harvard.edu/abs/2010ApJ...717..724G) [724](https://ui.adsabs.harvard.edu/abs/2010ApJ...717..724G)
- Ge H., Webbink R. F., Chen X., Han Z., 2015, [ApJ,](http://dx.doi.org/10.1088/0004-637X/812/1/40) [812, 40](https://ui.adsabs.harvard.edu/abs/2015ApJ...812...40G)
- Ge H., Webbink R. F., Chen X., Han Z., 2020, [ApJ,](http://dx.doi.org/10.3847/1538-4357/aba7b7) [899, 132](https://ui.adsabs.harvard.edu/abs/2020ApJ...899..132G)
- Gompertz B. P., Nicholl M., Schmidt P., Pratten G., Vecchio A., 2022, [MN-](http://dx.doi.org/10.1093/mnras/stac029)[RAS,](http://dx.doi.org/10.1093/mnras/stac029) [511, 1454](https://ui.adsabs.harvard.edu/abs/2022MNRAS.511.1454G)
- Gray R., et al., 2023, [J. Cosmology Astropart. Phys.,](http://dx.doi.org/10.1088/1475-7516/2023/12/023) [2023, 023](https://ui.adsabs.harvard.edu/abs/2023JCAP...12..023G)
- Grichener A., 2023, [MNRAS,](http://dx.doi.org/10.1093/mnras/stad1449) [523, 221](https://ui.adsabs.harvard.edu/abs/2023MNRAS.523..221G)
- Hirai R., Mandel I., 2022, [ApJ,](http://dx.doi.org/10.3847/2041-8213/ac9519) [937, L42](https://ui.adsabs.harvard.edu/abs/2022ApJ...937L..42H)
- Hurley J. R., Pols O. R., Tout C. A., 2000, [MNRAS,](http://dx.doi.org/10.1046/j.1365-8711.2000.03426.x) [315, 543](https://ui.adsabs.harvard.edu/abs/2000MNRAS.315..543H)
- Hurley J. R., Tout C. A., Pols O. R., 2002, [MNRAS,](http://dx.doi.org/10.1046/j.1365-8711.2002.05038.x) [329, 897](https://ui.adsabs.harvard.edu/abs/2002MNRAS.329..897H)
- Ivanova N., et al., 2013, [A&ARv,](http://dx.doi.org/10.1007/s00159-013-0059-2) [21, 59](https://ui.adsabs.harvard.edu/abs/2013A&ARv..21...59I)
- Janka H.-T., 2012, [Annual Review of Nuclear and Particle Science,](http://dx.doi.org/10.1146/annurev-nucl-102711-094901) [62, 407](https://ui.adsabs.harvard.edu/abs/2012ARNPS..62..407J)
- Janka H.-T., 2017, in Alsabti A.W., Murdin P., eds, , Handbook of Supernovae. p. 1095, [doi:10.1007/978-3-319-21846-5_109](http://dx.doi.org/10.1007/978-3-319-21846-5_109)
- Janka H. T., Langanke K., Marek A., Martínez-Pinedo G., Müller B., 2007, [Phys. Rep.,](http://dx.doi.org/10.1016/j.physrep.2007.02.002) [442, 38](https://ui.adsabs.harvard.edu/abs/2007PhR...442...38J)
- Kaaret P., Alonso-Herrero A., Gallagher J. S., Fabbiano G., Zezas A., Rieke M. J., 2004, [MNRAS,](http://dx.doi.org/10.1111/j.1365-2966.2004.07516.x) [348, L28](https://ui.adsabs.harvard.edu/abs/2004MNRAS.348L..28K)
- Klencki J., Nelemans G., Istrate A. G., Chruslinska M., 2021, [A&A,](http://dx.doi.org/10.1051/0004-6361/202038707) [645,](https://ui.adsabs.harvard.edu/abs/2021A&A...645A..54K) [A54](https://ui.adsabs.harvard.edu/abs/2021A&A...645A..54K)
- Kroupa P., 2002, Science, 295, 82
- LIGO Scientific Collaboration Virgo Collaboration KAGRA Collaboration 2018, LVK Algorithm Library - LALSuite, Free software (GPL), [doi:10.7935/GT1W-FZ16](http://dx.doi.org/10.7935/GT1W-FZ16)
- Lagos C. d. P., Tobar R. J., Robotham A. S. G., Obreschkow D., Mitchell P. D., Power C., Elahi P. J., 2018, [MNRAS,](http://dx.doi.org/10.1093/mnras/sty2440) [481, 3573](https://ui.adsabs.harvard.edu/abs/2018MNRAS.481.3573L)
- Lagos C. d. P., et al., 2019, [MNRAS,](http://dx.doi.org/10.1093/mnras/stz2427) [489, 4196](https://ui.adsabs.harvard.edu/abs/2019MNRAS.489.4196L)
- Laplace E., Götberg Y., de Mink S. E., Justham S., Farmer R., 2020, [A&A,](http://dx.doi.org/10.1051/0004-6361/201937300) [637, A6](https://ui.adsabs.harvard.edu/abs/2020A&A...637A...6L)
- Li Z., Zhu C., Lü G., Li L., Liu H., Guo S., Yu J., Lu X., 2024, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2405.11571) [p. arXiv:2405.11571](https://ui.adsabs.harvard.edu/abs/2024arXiv240511571L)
- Liotine C., Zevin M., Berry C. P. L., Doctor Z., Kalogera V., 2023, [ApJ,](http://dx.doi.org/10.3847/1538-4357/acb8b2) [946,](https://ui.adsabs.harvard.edu/abs/2023ApJ...946....4L) [4](https://ui.adsabs.harvard.edu/abs/2023ApJ...946....4L)
- Lu W., Fuller J., Quataert E., Bonnerot C., 2023, [MNRAS,](http://dx.doi.org/10.1093/mnras/stac3621) [519, 1409](https://ui.adsabs.harvard.edu/abs/2023MNRAS.519.1409L)
- Lynch R., Coughlin M., Vitale S., Stubbs C. W., Katsavounidis E., 2018, [ApJ,](http://dx.doi.org/10.3847/2041-8213/aacf9f) [861, L24](https://ui.adsabs.harvard.edu/abs/2018ApJ...861L..24L)
- MacLeod M., Loeb A., 2020, [ApJ,](http://dx.doi.org/10.3847/1538-4357/ab89b6) [895, 29](https://ui.adsabs.harvard.edu/abs/2020ApJ...895...29M)
- MacLeod M., Ostriker E. C., Stone J. M., 2018a, [ApJ,](http://dx.doi.org/10.3847/1538-4357/aacf08) [863, 5](https://ui.adsabs.harvard.edu/abs/2018ApJ...863....5M)
- MacLeod M., Ostriker E. C., Stone J. M., 2018b, [ApJ,](http://dx.doi.org/10.3847/1538-4357/aae9eb) [868, 136](https://ui.adsabs.harvard.edu/abs/2018ApJ...868..136M)
- Mandel I., Broekgaarden F. S., 2021, arXiv preprint arXiv:2107.14239
- Mandel I., Broekgaarden F. S., 2022, [Living Reviews in Relativity,](http://dx.doi.org/10.1007/s41114-021-00034-3) [25, 1](https://ui.adsabs.harvard.edu/abs/2022LRR....25....1M)
- Mandel I., Farmer A., 2022, [Phys. Rep.,](http://dx.doi.org/10.1016/j.physrep.2022.01.003) [955, 1](https://ui.adsabs.harvard.edu/abs/2022PhR...955....1M)
- Mandel I., Fragos T., 2020, [ApJ,](http://dx.doi.org/10.3847/2041-8213/ab8e41) [895, L28](https://ui.adsabs.harvard.edu/abs/2020ApJ...895L..28M)
- Mandel I., Müller B., 2020, Monthly Notices of the Royal Astronomical Society, 499, 3214
- Mandel I., de Mink S. E., 2016, [MNRAS,](http://dx.doi.org/10.1093/mnras/stw379) [458, 2634](https://ui.adsabs.harvard.edu/abs/2016MNRAS.458.2634M)
- Mandel I., Farr W. M., Gair J. R., 2019, [MNRAS,](http://dx.doi.org/10.1093/mnras/stz896) [486, 1086](https://ui.adsabs.harvard.edu/abs/2019MNRAS.486.1086M)
- Mapelli M., 2019, in Oskinova L. M., Bozzo E., Bulik T., Gies D. R., eds, , Vol. 346, High-mass X-ray Binaries: Illuminating the Passage from Massive Binaries to Merging Compact Objects. pp 397–416, [doi:10.1017/S1743921319001868](http://dx.doi.org/10.1017/S1743921319001868)
- Mapelli M., 2020, [Frontiers in Astronomy and Space Sciences,](http://dx.doi.org/10.3389/fspas.2020.00038) [7, 38](https://ui.adsabs.harvard.edu/abs/2020FrASS...7...38M)
- Mapelli M., 2021, in , Handbook of Gravitational Wave Astronomy. p. 16, [doi:10.1007/978-981-15-4702-7_16-1](http://dx.doi.org/10.1007/978-981-15-4702-7_16-1)
- Marchant P., Bodensteiner J., 2023, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2311.01865) [p. arXiv:2311.01865](https://ui.adsabs.harvard.edu/abs/2023arXiv231101865M)
- Marchant P., Langer N., Podsiadlowski P., Tauris T. M., Moriya T. J., 2016, [A&A,](http://dx.doi.org/10.1051/0004-6361/201628133) [588, A50](https://ui.adsabs.harvard.edu/abs/2016A&A...588A..50M)
- Marchant P., Pappas K. M. W., Gallegos-Garcia M., Berry C. P. L., Taam R. E., Kalogera V., Podsiadlowski P., 2021, [A&A,](http://dx.doi.org/10.1051/0004-6361/202039992) [650, A107](https://ui.adsabs.harvard.edu/abs/2021A&A...650A.107M)
- Marchant P., Podsiadlowski P., Mandel I., 2023, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2311.14041) [p.](https://ui.adsabs.harvard.edu/abs/2023arXiv231114041M) [arXiv:2311.14041](https://ui.adsabs.harvard.edu/abs/2023arXiv231114041M)
- Marek A., Janka H. T., 2009, [ApJ,](http://dx.doi.org/10.1088/0004-637X/694/1/664) [694, 664](https://ui.adsabs.harvard.edu/abs/2009ApJ...694..664M)
- Mastrogiovanni S., et al., 2021, [Phys. Rev. D,](http://dx.doi.org/10.1103/PhysRevD.104.062009) [104, 062009](https://ui.adsabs.harvard.edu/abs/2021PhRvD.104f2009M)
- Miller-Jones J. C. A., et al., 2021, [Science,](http://dx.doi.org/10.1126/science.abb3363) [371, 1046](https://ui.adsabs.harvard.edu/abs/2021Sci...371.1046M)
- Müller B., Heger A., Liptai D., Cameron J. B., 2016, [MNRAS,](http://dx.doi.org/10.1093/mnras/stw1083) [460, 742](https://ui.adsabs.harvard.edu/abs/2016MNRAS.460..742M)
- Nathaniel K., 2017
- Neijssel C. J., et al., 2019, [MNRAS,](http://dx.doi.org/10.1093/mnras/stz2840) [490, 3740](https://ui.adsabs.harvard.edu/abs/2019MNRAS.490.3740N)
- Neijssel C. J., Vinciguerra S., Vigna-Gómez A., Hirai R., Miller-Jones J. C. A., Bahramian A., Maccarone T. J., Mandel I., 2021, [ApJ,](http://dx.doi.org/10.3847/1538-4357/abde4a) [908, 118](https://ui.adsabs.harvard.edu/abs/2021ApJ...908..118N)
- Neo S., Miyaji S., Nomoto K., Sugimoto D., 1977, PASJ, [29, 249](https://ui.adsabs.harvard.edu/abs/1977PASJ...29..249N)
- O'Connor E., Ott C. D., 2011, [ApJ,](http://dx.doi.org/10.1088/0004-637X/730/2/70) [730, 70](https://ui.adsabs.harvard.edu/abs/2011ApJ...730...70O)
- Olejak A., Klencki J., Xu X.-T., Wang C., Belczynski K., Lasota J.-P., 2024, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2404.12426) [p. arXiv:2404.12426](https://ui.adsabs.harvard.edu/abs/2024arXiv240412426O)

Orosz J. A., et al., 2007, [Nature,](http://dx.doi.org/10.1038/nature06218) [449, 872](https://ui.adsabs.harvard.edu/abs/2007Natur.449..872O)

- Paczyński B., Sienkiewicz R., 1972, Acta Astron., [22, 73](https://ui.adsabs.harvard.edu/abs/1972AcA....22...73P)
- Passenger L., Thrane E., Lasky P. D., Payne E., Stevenson S., Farr B., 2024, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2405.09739) [p. arXiv:2405.09739](https://ui.adsabs.harvard.edu/abs/2024arXiv240509739P)
- Paxton B., Bildsten L., Dotter A., Herwig F., Lesaffre P., Timmes F., 2011, [ApJS,](http://dx.doi.org/10.1088/0067-0049/192/1/3) [192, 3](https://ui.adsabs.harvard.edu/abs/2011ApJS..192....3P)
- Périgois C., Mapelli M., Santoliquido F., Bouffanais Y., Rufolo R., 2023, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2301.01312) [p. arXiv:2301.01312](https://ui.adsabs.harvard.edu/abs/2023arXiv230101312P)
- Picco A., Marchant P., Sana H., Nelemans G., 2023, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2309.05736) [p.](https://ui.adsabs.harvard.edu/abs/2023arXiv230905736P) [arXiv:2309.05736](https://ui.adsabs.harvard.edu/abs/2023arXiv230905736P)
- Picker L., Hirai R., Mandel I., 2024, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2402.13180) [p. arXiv:2402.13180](https://ui.adsabs.harvard.edu/abs/2024arXiv240213180P)
- Pols O. R., Schröder K.-P., Hurley J. R., Tout C. A., Eggleton P. P., 1998, [MNRAS,](http://dx.doi.org/10.1046/j.1365-8711.1998.01658.x) [298, 525](https://ui.adsabs.harvard.edu/abs/1998MNRAS.298..525P)
- Powell J., Gossan S. E., Logue J., Heng I. S., 2016, [Phys. Rev. D,](http://dx.doi.org/10.1103/PhysRevD.94.123012) [94, 123012](https://ui.adsabs.harvard.edu/abs/2016PhRvD..94l3012P)
- Powell J., Iess A., Llorens-Monteagudo M., Obergaulinger M., Müller B., Torres-Forné A., Cuoco E., Font J. A., 2023, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2311.18221) [p.](https://ui.adsabs.harvard.edu/abs/2023arXiv231118221P) [arXiv:2311.18221](https://ui.adsabs.harvard.edu/abs/2023arXiv231118221P)
- Rauf L., Howlett C., Davis T. M., Lagos C. D. P., 2023, [Monthly Notices of](http://dx.doi.org/10.1093/mnras/stad1757) [the Royal Astronomical Society,](http://dx.doi.org/10.1093/mnras/stad1757) 523, 5719
- Riley J., Mandel I., 2023, [ApJ,](http://dx.doi.org/10.3847/1538-4357/accf90) [950, 80](https://ui.adsabs.harvard.edu/abs/2023ApJ...950...80R)
- Riley J., Mandel I., Marchant P., Butler E., Nathaniel K., Neijssel C., Shortt S., Vigna-Gómez A., 2021, [MNRAS,](http://dx.doi.org/10.1093/mnras/stab1291) [505, 663](https://ui.adsabs.harvard.edu/abs/2021MNRAS.505..663R)
- Romagnolo A., Belczynski K., Klencki J., Agrawal P., Shenar T., Szécsi D., 2023, [MNRAS,](http://dx.doi.org/10.1093/mnras/stad2366) [525, 706](https://ui.adsabs.harvard.edu/abs/2023MNRAS.525..706R)
- Romero-Shaw I., Lasky P. D., Thrane E., 2022, [ApJ,](http://dx.doi.org/10.3847/1538-4357/ac9798) [940, 171](https://ui.adsabs.harvard.edu/abs/2022ApJ...940..171R)
- Romero-Shaw I., Hirai R., Bahramian A., Willcox R., Mandel I., 2023, [MN-](http://dx.doi.org/10.1093/mnras/stad1732)[RAS,](http://dx.doi.org/10.1093/mnras/stad1732) [524, 245](https://ui.adsabs.harvard.edu/abs/2023MNRAS.524..245R)
- Sana H., et al., 2012, Science, 337, 444
- Santoliquido F., Mapelli M., Giacobbo N., Bouffanais Y., Artale M. C., 2021, [MNRAS,](http://dx.doi.org/10.1093/mnras/stab280) [502, 4877](https://ui.adsabs.harvard.edu/abs/2021MNRAS.502.4877S)
- Santoliquido F., Mapelli M., Iorio G., Costa G., Glover S. C. O., Hartwig T., Klessen R. S., Merli L., 2024, [MNRAS,](http://dx.doi.org/10.1093/mnras/stad3969) [528, 954](https://ui.adsabs.harvard.edu/abs/2024MNRAS.528..954S)
- Schneider F. R. N., Izzard R. G., Langer N., de Mink S. E., 2015, [ApJ,](http://dx.doi.org/10.1088/0004-637X/805/1/20) [805,](https://ui.adsabs.harvard.edu/abs/2015ApJ...805...20S) 20
- Schneider F., Podsiadlowski P., Müller B., 2021, Astronomy & Astrophysics, 645, A5
- Schneider F. R. N., Podsiadlowski P., Laplace E., 2023, [ApJ,](http://dx.doi.org/10.3847/2041-8213/acd77a) [950, L9](https://ui.adsabs.harvard.edu/abs/2023ApJ...950L...9S)
- Schneider F. R. N., Podsiadlowski P., Laplace E., 2024, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2403.03984) [p.](https://ui.adsabs.harvard.edu/abs/2024arXiv240303984S) [arXiv:2403.03984](https://ui.adsabs.harvard.edu/abs/2024arXiv240303984S)
- Schrøder S. L., MacLeod M., Ramirez-Ruiz E., Mandel I., Fragos T., Loeb A., Everson R. W., 2021, [arXiv e-prints,](http://dx.doi.org/10.48550/arXiv.2107.09675) [p. arXiv:2107.09675](https://ui.adsabs.harvard.edu/abs/2021arXiv210709675S)
- Silverman B. W., 1986, Density estimation for statistics and data analysis
- Stevenson S., Clarke T. A., 2022, [MNRAS,](http://dx.doi.org/10.1093/mnras/stac2936) [517, 4034](https://ui.adsabs.harvard.edu/abs/2022MNRAS.517.4034S)
- Stevenson S., Ohme F., Fairhurst S., 2015, [ApJ,](http://dx.doi.org/10.1088/0004-637X/810/1/58) [810, 58](https://ui.adsabs.harvard.edu/abs/2015ApJ...810...58S)
- Stevenson S., Vigna-Gómez A., Mandel I., Barrett J.W., Neijssel C. J., Perkins D., de Mink S. E., 2017, [Nature Communications,](http://dx.doi.org/10.1038/ncomms14906) [8, 14906](https://ui.adsabs.harvard.edu/abs/2017NatCo...814906S)
- Tanikawa A., 2024, arXiv e-prints, [p. arXiv:2403.04389](https://ui.adsabs.harvard.edu/abs/2024arXiv240304389T)
- Team COMPAS: Riley, J. et al., 2022, [ApJS,](http://dx.doi.org/10.3847/1538-4365/ac416c) [258, 34](https://ui.adsabs.harvard.edu/abs/2022ApJS..258...34R)
- Temaj D., Schneider F. R. N., Laplace E., Wei D., Podsiadlowski P., 2024, [A&A,](http://dx.doi.org/10.1051/0004-6361/202347434) [682, A123](https://ui.adsabs.harvard.edu/abs/2024A&A...682A.123T)
- Temmink K. D., Pols O. R., Justham S., Istrate A. G., Toonen S., 2023, [A&A,](http://dx.doi.org/10.1051/0004-6361/202244137) [669, A45](https://ui.adsabs.harvard.edu/abs/2023A&A...669A..45T)
- Tong H., Galaudage S., Thrane E., 2022, [Phys. Rev. D,](http://dx.doi.org/10.1103/PhysRevD.106.103019) [106, 103019](https://ui.adsabs.harvard.edu/abs/2022PhRvD.106j3019T)
- Tutukov A., Yungelson L., 1979, in Conti P. S., De Loore C. W. H., eds, , Vol. 83, Mass Loss and Evolution of O-Type Stars. pp 401–406
- Vigna-Gómez A., et al., 2018, [MNRAS,](http://dx.doi.org/10.1093/mnras/sty2463) [481, 4009](https://ui.adsabs.harvard.edu/abs/2018MNRAS.481.4009V)
- Vigna-Gómez A., et al., 2024, [Phys. Rev. Lett.,](http://dx.doi.org/10.1103/PhysRevLett.132.191403) [132, 191403](https://ui.adsabs.harvard.edu/abs/2024PhRvL.132s1403V)
- Vink J. S., Harries T. J., 2017, [A&A,](http://dx.doi.org/10.1051/0004-6361/201730503) [603, A120](https://ui.adsabs.harvard.edu/abs/2017A&A...603A.120V)
- Vink J. S., de Koter A., 2005, [A&A,](http://dx.doi.org/10.1051/0004-6361:20052862) [442, 587](https://ui.adsabs.harvard.edu/abs/2005A&A...442..587V)
- Vitale S., Gerosa D., Farr W. M., Taylor S. R., 2022, in , Handbook of Gravitational Wave Astronomy. p. 45, [doi:10.1007/978-981-15-4702-7_45-1](http://dx.doi.org/10.1007/978-981-15-4702-7_45-1) Webbink R. F., 1984, [ApJ,](http://dx.doi.org/10.1086/161701) [277, 355](https://ui.adsabs.harvard.edu/abs/1984ApJ...277..355W)
- Willcox R., MacLeod M., Mandel I., Hirai R., 2023, [ApJ,](http://dx.doi.org/10.3847/1538-4357/acffb1) [958, 138](https://ui.adsabs.harvard.edu/abs/2023ApJ...958..138W)
- Wong K. W. K., Breivik K., Farr W. M., Luger R., 2023, [ApJ,](http://dx.doi.org/10.3847/1538-4357/acc863) [950, 181](https://ui.adsabs.harvard.edu/abs/2023ApJ...950..181W)
- Woosley S. E., Heger A., 2021, [ApJ,](http://dx.doi.org/10.3847/2041-8213/abf2c4) [912, L31](https://ui.adsabs.harvard.edu/abs/2021ApJ...912L..31W)
- Xu X.-J., Li X.-D., 2010, [ApJ,](http://dx.doi.org/10.1088/0004-637X/716/1/114) [716, 114](https://ui.adsabs.harvard.edu/abs/2010ApJ...716..114X)
- Zevin M., Bavera S. S., 2022, [ApJ,](http://dx.doi.org/10.3847/1538-4357/ac6f5d) [933, 86](https://ui.adsabs.harvard.edu/abs/2022ApJ...933...86Z)
- Zevin M., Spera M., Berry C. P. L., Kalogera V., 2020, [ApJ,](http://dx.doi.org/10.3847/2041-8213/aba74e) [899, L1](https://ui.adsabs.harvard.edu/abs/2020ApJ...899L...1Z)
- Zevin M., et al., 2021, [ApJ,](http://dx.doi.org/10.3847/1538-4357/abe40e) [910, 152](https://ui.adsabs.harvard.edu/abs/2021ApJ...910..152Z)
- Zhang R. C., Fragione G., Kimball C., Kalogera V., 2023, [ApJ,](http://dx.doi.org/10.3847/1538-4357/ace4c1) [954, 23](https://ui.adsabs.harvard.edu/abs/2023ApJ...954...23Z)
- Zheng Y., Cavaglià M., Quitzow-James R., Mogushi K., 2021, Significance, 18
- Zuo Z.-Y., Li X.-D., 2014, [MNRAS,](http://dx.doi.org/10.1093/mnras/stu993) [442, 1980](https://ui.adsabs.harvard.edu/abs/2014MNRAS.442.1980Z)
- de Kool M., 1990, [ApJ,](http://dx.doi.org/10.1086/168974) [358, 189](https://ui.adsabs.harvard.edu/abs/1990ApJ...358..189D)
- de Mink S. E., Pols O. R., Hilditch R. W., 2007, [A&A,](http://dx.doi.org/10.1051/0004-6361:20067007) [467, 1181](https://ui.adsabs.harvard.edu/abs/2007A&A...467.1181D)
- de Mink S. E., Cantiello M., Langer N., Pols O. R., Brott I., Yoon S. C., 2009, [A&A,](http://dx.doi.org/10.1051/0004-6361/200811439) [497, 243](https://ui.adsabs.harvard.edu/abs/2009A&A...497..243D)
- de Mink S. E., Cantiello M., Langer N., Pols O. R., Yoon S. C., 2010, in Prša A., Zejda M., eds, Astronomical Society of the Pacific Conference

Distribution	Model
IMF	Kroupa (2002)
Initial mass range	$5 - 150 M_{\odot}$
	$5 - 200M_{\odot}$ (Rauf et al. 2023)
Semi-major axis	Flat-in-log
Orbital period	Flat-in-log
Mass ratio	Sana et al. (2012)
Eccentricity	ZERO (circular)
	Sana et al. (2012) (Rauf et al. 2023)
Metallicity	SHARK (Lagos et al. 2018)

Table A1. Summary of COMPAS default initial conditions.

Series Vol. 435, Binaries - Key to Comprehension of the Universe. p. 179 ([arXiv:0910.3694](http://arxiv.org/abs/0910.3694)), [doi:10.48550/arXiv.0910.3694](http://dx.doi.org/10.48550/arXiv.0910.3694)

- du Buisson L., et al., 2020, [MNRAS,](http://dx.doi.org/10.1093/mnras/staa3225) [499, 5941](https://ui.adsabs.harvard.edu/abs/2020MNRAS.499.5941D)
- van Son L. A. C., et al., 2020, [ApJ,](http://dx.doi.org/10.3847/1538-4357/ab9809) [897, 100](https://ui.adsabs.harvard.edu/abs/2020ApJ...897..100V)
- van Son L. A. C., et al., 2022a, [ApJ,](http://dx.doi.org/10.3847/1538-4357/ac64a3) [931, 17](https://ui.adsabs.harvard.edu/abs/2022ApJ...931...17V)
- van Son L. A. C., et al., 2022b, [ApJ,](http://dx.doi.org/10.3847/1538-4357/ac9b0a) [940, 184](https://ui.adsabs.harvard.edu/abs/2022ApJ...940..184V)
- van Son L. A. C., de Mink S. E., Chruślińska M., Conroy C., Pakmor R., Hernquist L., 2023, [ApJ,](http://dx.doi.org/10.3847/1538-4357/acbf51) [948, 105](https://ui.adsabs.harvard.edu/abs/2023ApJ...948..105V)
- van den Heuvel E. P. J., Portegies Zwart S. F., de Mink S. E., 2017, [MNRAS,](http://dx.doi.org/10.1093/mnras/stx1430) [471, 4256](https://ui.adsabs.harvard.edu/abs/2017MNRAS.471.4256V)

APPENDIX A: MODEL INITIAL CONDITIONS AND FIXED PARAMETERS

In Table [A1,](#page-22-37) we list the distributions and ranges for the input parameters used to define the initial conditions of our COMPAS binaries. These are fixed for all models explored in this paper and [Rauf et al.](#page-22-10) (2023)

APPENDIX B: GW DATA

In Table [B1,](#page-23-0) we list the GW events used as data for each selection criteria, as described in Section [4.1.](#page-8-2)

APPENDIX C: MODEL HYPER-PARAMETERS AND BAYES FACTOR DATA

Extending on Table [1,](#page-6-1) for each model in Tables [C1,](#page-24-0) [C2](#page-25-0) and [C3](#page-26-0) we detail the hyper-parameters and Bayes factor for each selection criteria. [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) (a)-(d) refer to the models explored in [Rauf](#page-22-10) [et al.](#page-22-10) [\(2023\)](#page-22-10), where we vary the remnant mass prescription (with the exception of [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) (c) where we also vary f_{WR}).

This paper has been typeset from a TEX/LATEX file prepared by the author.

Table B1. List of GW events included in the analysis for each selection criteria.

Table C1. Hyper-parameters for the [Rauf et al.](#page-22-10) [\(2023\)](#page-22-10) models and new Models 1-100 and the corresponding Bayes factor relative to the maximum likelihoods, for which the model is in bold. In green, we highlight the highest likelihood models, where the Bayes factor is > -10 for either the marginalised or unmarginalised likelihoods. MNRAS **⁰⁰⁰**, [1](#page-0-0)[–23](#page-26-0) (0000)

Table C2. Hyper-parameters for Models 101-200 and the corresponding Bayes factor relative to the maximum likelihoods, for which the model is in bold. In green, we highlight the highest likelihood models, where the Bayes factor is > −10 for either the marginalised or unmarginalised likelihoods. MNRAS **000**, [1](#page-0-0)[–23](#page-26-0) (0000)

Rauf et al.

Navigating COMPAS*, Shark and Bayesian Inference for BBH Model Selection* 27

Table C3. Hyper-parameters for Models 201-300 and the corresponding Bayes factor relative to the maximum likelihoods, for which the model is in bold. In green, we highlight the highest likelihood models, where the Bayes factor is > -10 for either the marginalised or unmarginalised likelihoods.