Simultaneous comparisons of the variances of k treatments with that of a control: a Levene-Dunnett type procedure

Ludwig A. Hothorn,

Im Grund 12, D-31867 Lauenau, Germany (e-mail:ludwig@hothorn.de)

(retired from Leibniz University Hannover

June 19, 2024

Abstract

There are some global tests for heterogeneity of variance in k-sample one-way layouts, but few consider pairwise comparisons between treatment levels. For experimental designs with a control, comparisons of the variances between the treatment levels and the control are of interest - in analogy to the location parameter with the Dunnett (1955) procedure. Such a many-to-one approach for variances is proposed using the Levene transformation, a kind of residuals. Its properties are characterized with simulation studies and corresponding data examples are evaluated with R code.

1 A Levene-Dunnett type test

Several tests (or pre-tests) on variance homogeneity in completely randomized one-way layouts exists for independent pairwise ratios, for global tests [4] or for ordered alternatives [6]. In clinical dose-finding studies and toxicological bioassays, simultaneous comparisons of the variances of k treatments with that of a control are of interest in a design with treatment levels T_j , j = 0, ..., k, i.e. $[T_j : T_0, T_1, T_2, ..., T_k]$ is commonly used, where T_0 represents a placebo or a negative control group. In principle the closed testing procedure can be used to achieve simultaneous, pairwise comparisons against control [1]. Whereas adjusted p-values are easy to estimated, simultaneous confidence intervals are complicated [3]. A straightforward approach is the use of Levene transformed variables $Z_{ji} = Y_{ji} - Median(Y_j)$ directly within the standard Dunnett procedure [2]. The common Dunnett test [2] can be formulated as a multiple contrast test: $maxT = max(t_1, ..., t_k)$ where t_{ξ} are standardised contrast tests: $t_{\xi} = \sum_{i=0}^k c_i \bar{y}_i / S \sqrt{\sum_i^k c_i^2/n_i}$ where c_i^{ξ} are the contrast coefficients. For a balanced design with k = 2 the contrast matrix is simple, e.g. for the one-tailed Dunnett test:

Table 1: Dunnett-type contrast matrix for k = 1 + 3 comparisons

The quantiles of the max-T test $t_{k,df,R,1-\alpha}$ are monotonic with k which provides elementary inference, i.e. adjusted p-values and/or simultaneous confidence intervals. As in the Dunnett original procedure, one- or two-sided hypothesis formulations are easily possible. The empirical behavior of familywise error rate (FWER) and power of this new Levene-Dunnett test (Lev-Dun) is characterized by means of a simulation study, whereby the behavior with the usual small n_i is of interest. The estimation of both adjusted p-values and simultaneous confidence intervals is demonstrated using an example.

2 Simulations

2.1 Simulations under global *H*₀

First, a balanced design k = 3 + 1 is considered.

n_i	$\hat{\alpha}$
3	0.001
5	0.010
10	0.042
20	0.044
30	0.046
50	0.045

Table 2: Empirical FWER $\hat{\alpha}$ in balanced design ($k = 3 + 1, n_i = const$, one-sided)

This test, based on an asymptotic transformation, controls the FWER from about $n_i = 10$ on. Second, selected unbalanced designs for k = 3 + 1 are considered.

n_i	$\hat{\alpha}$
10,10,10,10	0.042
16,8,8,8,8	0.033
8,8,8,12	0.041

Table 3: Empirical FWER $\hat{\alpha}$ in unbalanced designs (k = 3 + 1, N = 40, one-sided)

For selected unbalanced designs, the test becomes slightly conservative

2.2 Simulations under partial *H*₁

Different patterns of variance heterogeneity are considered in a balanced design.

σ_0	σ_1	σ_2	σ_3	$\hat{\pi}$	$\pi/\hat{\alpha}_{01}$	π/\hat{lpha}_{02}	$\pi/\hat{\alpha}_{03}$
const	const	const	↑	0.806	0.002	0.000	0.806
const	const	↑	↑	0.847	0.000	0.617	0.598
const	↑	↑	↑	0.789	0.417	0.403	0.413
const	↑	↑	const	0.849	0.598	0.597	0.000
const	↑	const	const	0.822	0.820	0.004	0.001
const	const	↑	const	0.810	0.001	0.810	0.003
const	↑	const	↑	0.836	0.593	0.000	0.584
↑	const	const	↑	0.049	0.000	0.000	0.049

Table 4: Global and elementary powers (size) for various variance patterns ($n_i = 10$, one-sided)

The power estimates depend on the particular variance pattern, the elementary power estimates are correct with respect to the underlying pattern, as are the elementary α_i . The one-sided version provides a correct directional testing.

Patterns of variance heterogeneity are also considered for selected unbalanced designs.

						$\pi/\hat{\alpha}_{02}$	
10	10	10	10	0.806	0.002	0.000	0.806
16	8	8	8	0.799	0.002	0.799	0.001
16	10	4	10	0.607	0.010	0.599	0.006

Table 5: Global and elementary powers (size) for a selected variance pattern ($\sigma_0 = \sigma_1 < \sigma_2 \sigma_3$) in unbalanced design (N = 40, one-sided)

Hardly any different power for a design with a higher sample sizes in the control n_0 (i.e., an optimal design for location effects), but, as expected, a significant drop in power when the group with the increased variance uses a significantly lower sample size (still for a constant total sample size N).

3 A modified Levene transformation

For designs with small sample sizes n_i , in treatment groups with odd n_i the value $Z_{ji} = 0$ occur which cause conservativeness [1]. A one-sided trimmed test version, omitting this single value zero, (modLev-Dun) avoid this.

n_i	LevDun	modLevDun
15	0.036	0.051
11	0.029	0.048
7	0.019	0.050
5	0.010	0.040
3	0.001	0.047

Table 6: Small sample size behavior: comparison of FWER estimates of Levene and modified Levene-Dunnett test

σ_1	σ_2	σ_3	σ_4	LevDun	$LevDun_{01}$	$LevDun_{02}$	$LevDun_{03}$	modLevDun	$modLevDun_{01}$	$modLevDun_{02}$	$modLevDun_{03}$
const	const	const	↑	0.845	0.001	0.000	0.845	0.869	0.001	0.000	0.869
const	const	↑	const	0.867	0.000	0.866	0.001	0.883	0.000	0.882	0.001
const	↑	const	const	0.848	0.848	0.000	0.000	0.878	0.878	0.001	0.000
↑	const	const	const	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
const	const	↑	↑	0.879	0.000	0.620	0.637	0.911	0.000	0.688	0.698
const	↑	↑	↑	0.861	0.444	0.452	0.451	0.910	0.521	0.531	0.541

Table 7: Comparison of elementary power estimates of Levene and modified Levene-Dunnett test for balanced small sample size $n_i = 11$

The control of the FWER even at small n_i and thus the higher power are impressive, although this modification uses a biased test statistic.

4 A global test for any heterogeneity

Sometimes global test results are of interest in unstructured k-sample designs. Instead the Dunnett manyto-one comparison contrasts, the multiple contrasts against the grand mean can be used [5]. While conventional F-tests use the sum of the quadratic deviation from the overall mean, the maximum test using linear deviations is used here. Such an approach allows both global claims (using the $min(p_x)$ approach) and adjusted p-values or simultaneous confidence intervals for the pairwise comparisons against the grand mean. This approach is demonstrated by an example below.

5 Examples

The first example uses the litter weight data in a toxicological bioassay (data available in library(multcomp)) [7]. The R-code is simple to estimate adjusted p-value or simultaneous confidence intervals

```
library(multcomp)
data(litter)
medrs<-tapply(litter$weight, litter$dose, median) # group-specific medians
medrsV<-medrs[as.integer(litter$dose)] # median for each subject
litter$tresponse<- abs(litter$weight-medrsV) # Levene transformation
mod2<-lm(tresponse~dose, data=litter) # linear model
summary(glht(mod2, linfct = mcp(dose ="Dunnett"), alternative="greater"))
plot(glht(mod2, linfct = mcp(dose ="Dunnett"), alternative="greater"))</pre>
```





Figure 1: Boxplot of litter weight data

Comparisons	p-value LevDun
5 - 0	0.022
50 - 0	0.491
500 - 0	0.022

Table 8: p-values for one-sided Levene-Dunnett test for the litter weight data

The second example uses the survival time data in [1] as an example for unstructured k-sample design:

```
boos<-data.frame(survival, group=as.factor(grp))
library(multcomp)
medrs<-tapply(boos$survival, boos$group, median) # group-specific medians
medrsV<-medrs[(boos$group)] # median for each subject
boos$tresponse<- abs(boos$survival-medrsV) # Levene transformation
mod3<-lm(tresponse~group, data=boos)
summary(glht(mod3, linfct = mcp(group="GrandMean")))
library(GLDEX)
no0<-fun.zero.omit(boos$tresponse)
boos0<-data.frame(no0, group0=as.factor(grp0))
mod4<-lm(no0~group0, data=boos0)
summary(glht(mod4, linfct = mcp(group0="GrandMean")))</pre>
```






Figure 2: Simultaneous confidence intervals for Levene transformations




Figure 3: Boxplot for Boos et al. 2023 example

The advantage of the new heterogeneity test compared to the F-test variant (in library(misty)) is obvious: not only is a global p-value is available (i.e. 'variance heterogeneity exists'), but the statement 'the variance at the tumor site='breast' is increased with respect to the overall variance' is also available.

6 Summary

The proposed Levene-Dunnett test can be recommended for randomized designs with a control or placebo group for the analysis of possible variance heterogeneity, whereby the modified version should be used for small sample sizes. An analogous heterogeneity test for unstructured designs is also possible. Two data examples were used to illustrate the simple R code and the problem-adapted interpretation.

Cancer site	LevDun	modLevDun
breast	0.002	0.002
bronchus	0.999	0.13
colon	0.999	0.78
ovary	0.31	0.54
stomach	0.99	0.79
global Levene min(p)-test	0.002	0.002
global Levene F-test	0.027	-

Table 9: p-values for 2-sided Levene-Dunnett and modLevDun test for survival data

References

- Dennis DB, Kaiyuan D., and Xiaoni L. Pairwise comparisons for levene-style variability parameters. Comm. Stat- B, 52(4):1562–1576, APR 3 2023.
- [2] C. W. Dunnett. A multiple comparison procedure for comparing several treatments with a control. J. Amer. Statist. Assoc., 50(272):1096–1121, 1955.
- [3] L.A. Hothorn. Comparisons of multiple treatment groups with a negative control or placebo group: Dunnett test vs. closed test procedure. *arXiv:2012.04277 (2020)*, 2020.
- [4] Katsileros A., Antonetsis N. A comparison of tests for homoscedasticity using simulation and empirical data. *Comm. Stats-A*, 31(1), JAN 2024.
- [5] F. Konietschke, S. Bosiger, E. Brunner, and L. A. Hothorn. Are multiple contrast tests superior to the anova? Intern. J. Biostat., 9(1):63–73, May 2013.
- [6] Pallmann P., Hothorn LA., and Djira GD. A Levene-type test of homogeneity of variances against ordered alternatives. Comput Stat., 29(6):1593–1608, DEC 2014.
- [7] P. H. Westfall. Multiple testing of general contrasts using logical constraints and correlations. J. Amer. Statist. Assoc., 92(437):299–306, March 1997.