Model-Based Inference and Experimental Design for Interference Using Partial Network Data

Steven Wilkins Reeves Department of Statistics, University of Washington

> Shane Lubold US Census Bureau

Arun G. Chandrasekhar Department of Economics, Stanford University, J-PAL, NBER

Tyler H. McCormick^{*}

Departments of Sociology and Statistics, University of Washington

Abstract

The stable unit treatment value assumption states that the outcome of an individual is not affected by the treatment statuses of others, however in many real world applications, treatments can have an effect on many others beyond the immediately treated. Interference can generically be thought of as mediated through some network structure. In many empirically relevant situations however, complete network data (required to adjust for these spillover effects) are too costly or logistically infeasible to collect. Partially or indirectly observed network data (e.g., subsamples, aggregated relational data (ARD), egocentric sampling, or respondent-driven sampling) reduce the logistical and financial burden of collecting network data, but the statistical properties of treatment effect adjustments from these design strategies are only beginning to be explored. In this paper, we present a framework for the estimation and inference of treatment effect adjustments using partial network data through the lens of structural causal models. We also illustrate procedures to assign treatments using only partial network data, with the goal of either minimizing estimator variance or optimally seeding. We derive single network asymptotic results applicable to a variety of choices for an underlying graph model. We validate our approach using simulated experiments on observed graphs with applications to information diffusion in India and Malawi.

^{*}Correspondence: tylermc@uw.edu. The authors greatly thank Lori Beaman, Stephane Bonhomme, Vincent Boucher, Carlos Cinelli, Paul Goldsmith-Pinkham, Kosuke Imai, Ben Letham, Fabrizia Mealli, Alex Philip and Alex Volfovsky for helpful comments and discussion.

1 Introduction

Interference occurs when one individual's treatment status impacts others' outcomes. Interference, also known as "spillover effects," appears in multiple scientific domains, including the study of infectious diseases (Hudgens and Halloran, 2008; Tchetgen and VanderWeele, 2012), studying peer influence (Manski, 1993; Bramoullé et al., 2009; De Giorgi et al., 2010; Epple and Romano, 2011; Goldsmith-Pinkham and Imbens, 2013), public policy (Malani et al., 2021; Imai et al., 2021), information diffusion (Banerjee et al., 2013, 2019), technology adoption (Beaman et al., 2021), online platforms (Saveski et al., 2017; Pouget-Abadie et al., 2018, 2019) and online marketplaces (Ha-Thuc et al., 2020; Johari et al., 2022), among other domains.

Interference violates the stable unit treatment value assumption (SUTVA), which states that an individual's outcome is not impacted by the treatment status of their peers. When SUTVA is violated, each potential outcome, the counterfactual outcome under a given treatment assignment, could depend on all treatment assignments within the population. Valid inference for treatment effects under SUTVA violations is an active area of research, with solutions typically depending on a combination of exposure maps and structural causal models. Exposure maps categorize respondents according to their network characteristics and the vector of treatment statuses (Aronow and Samii, 2017; Chandrasekhar et al., 2023), while structural causal models identify specific pathways for influence between individuals (van der Laan, 2012; Ogburn et al., 2022).

Estimating causal effects under interference typically requires complete network data, which is expensive and onerous to collect or may not be available due to privacy constraints. Partially observed network data takes many forms: subgraph samples where a researcher observes the presence/absence for only a subset of possible connections, egocentric sampling using either specific links or aggregates, or network-based sampling methods such as snowball sampling or respondent-driven sampling. In each case, incomplete network information introduces miss-measurement in the exposure map. A person may have treated peers, for example, but if links to those peers are not observed, the researcher will think their outcome is totally orthogonal to the treatment.

This paper introduces a framework for estimation and inference of causal effects under *partial* network data arising from a single graph. Partial here means that we may observe some or no links or aggregate summaries of links, which we will formalize later. With such data, we recover multiple estimands including various conditional or average treatment effects. To do this, we define a broad class of structural causal models that are amenable to estimation using partial data. This class covers many empirically relevant schemas for interference, such as diffusion and its generalizations. Estimation leverages a dual approach: first, by using an iterated expectation method for de-biased estimation of model parameters with partial network data, and second, by managing the dependence of exogenous noise in the outcomes. Chandrasekhar and Lewis (2011) also introduced an iterated expectations strategy for cases where multiple networks are available and data are independent across networks. We tackle the more challenging inference task of single network asymptotics. Our method applies when the underlying graph has features captured by the class of nodeexchangeable formation models, which we commonly see in practice and connect this to the problem of estimating effects of experiments. Previous methods (Breza et al., 2020, 2023) developed a related method to estimate network model features using a specific type of aggregated network data. Along with expanding to a wide range of partial data types, we extend this existing methodology to relax the requirement in previously published studies that traits be mutually distinct, a challenge to its usability in practice until now.

We also tackle the problem of experimental design associated with network exposure in scenarios where obtaining pristine network data ahead of randomized controlled trials (RCTs) is challenging or impossible. By collecting partial network data and employing a Bayesian optimization algorithm, we propose experimental designs that efficiently maximize treatment saturation tailored to specific estimands of interest. Our results demonstrate that this methodology not only surpasses traditional methods like inverse probability weighted (IPW) estimators in estimating global average treatment effects but also facilitates innovative seeding strategies that leverage the unique characteristics of partial network data. We demonstrate that these techniques can be used to assign treatment in such a way as to minimize estimator variance or to optimally seed for diffusion.

The remainder of the paper is structured as follows. We begin with a review of related work (section 1.1. section 2 defines the necessary background, then section 3 describes the procedure for estimation and inference. section 4 describes experimental design using partial network data and section 5 provides empirical examples. We conclude in section 6. Code to replicate the results in the paper is available at https://github.com/SteveJWR/ardexp, and an R package is available from https://github.com/SteveJWR/SBMNetReg.

1.1 Related Work

We first provide a brief review of literature related to inference with partial network data, then move to an overview of causal inference under interference. Complete network data collection can be prohibitively expensive and restricted by privacy concerns (Breza et al., 2020). Researchers typically work with partial network data derived from various sources such as survey samples, coarse geographic data, kinship information from censuses, or aggregated financial transactions. Comprehensive reviews of methods for handling network data can be found in De Paula (2017) or Graham (2020), and discussions on identification in network and related models are provided in Manski (2009). A direct approach is node subsampling, selecting a portion of nodes from the population and mapping the entire graph among them. If random sampling of nodes is infeasible, or if populations are sensitive or stigmatized, techniques like snowball or respondent-driven sampling offer a limited but focused view of the graph (Heckathorn, 1997; Goel and Salganik, 2009, 2010; Baraff et al., 2016; Green et al., 2020)

When complete edge enumeration among node subsets is impractical, researchers adopt standard survey methods such as Aggregated Relational Data (ARD) collection. The main intuition is that each of the partial network designs mentioned above can be used to estimate a breakdown of each respondent's network in terms of observable characteristics. In ARD surveys, respondents are asked, "How many people do you know with trait X?" for various traits. Additional conditions may be added in addition to collect the type of connection that is relevant (Feehan et al., 2016). Originally designed to estimate hard-toreach populations like HIV-positive men in the US (Killworth et al., 1998; Scutelniciuc, 2012; Jing et al., 2014), has been extended to a variety of other settings such as financial contagion models (Acemoglu et al., 2015) as well as more general network scale up methods utilized (NSUM) (Killworth et al., 1998; Kadushin et al., 2006; Feehan and Salganik, 2016; McCormick, 2020) and is notably 70 to 80% less costly than full network data collection (Breza et al., 2020). Another standard survey method, egocentric sampling, asks respondents to consider specific individuals in their networks and provide detailed information about them, unlike the aggregate focus of ARD and is commonly used in applications such as contact tracing (Potter et al., 2011), violence perpetration (Bond and Bushman, 2017) or adolescent substance measurement (Huang et al., 2014).

The first task in causal inference problems, particularly in the presence of interference, is defining the target estimand. The global average treatment effect (GATE), for example, assesses the impact of treating everyone versus treating no one, considering peer effects (Ugander et al., 2013). Other interests might include the effect of specific treatment allocations, like identifying influential individuals (Kempe et al., 2003; Banerjee et al., 2019), often limited by policy constraints (e.g., subsidies for the ultra-poor as in Anderson and Feder (2007)) or due to non-monotone peer effects dynamics (Banerjee et al., 2018): treating everyone may change interaction dynamics in equilibrium. More generally Aronow and Samii (2017) compare average treatment effects between two exposure configurations. A distinct but related line of work seeks to detect whether interference is present at all (Athey et al., 2018).

Models for peer influence like contagion (Jackson et al., 2008; Banerjee et al., 2013; Beaman et al., 2021; He and Song, 2023) or hearing models (Banerjee et al., 2019) structure interference analysis by identifying specific mechanisms that describe how connections between peers impact outcomes. Auerbach and Tabord-Meehan (2021) explore these effects through structural causal models focusing on nonparametric estimation, while our work emphasizes estimation, inference, and design using partial network data. Much of this literature assumes a fully observed graph, though a recent line of literature address imperfect or incompletely sampled graphs under certain conditions and for specific average causal effects (Hardy et al., 2019; Yu et al., 2022; Cortez et al., 2022). A related line of work examines sensitivity analysis for standard causal estimators under hidden treatment diffusion Tortú et al. (2021).

2 Environment

Let $i \in \{1, 2, ..., n\} = \mathscr{V}$ denote a population of interacting individuals and let $\mathscr{G} = \mathscr{V} \times \mathscr{E}$ be the network by which interference is propagated; where \mathscr{V} is the set of node vertices and $\mathscr{E} \subset \mathscr{V} \times \mathscr{V}$ is a set of edges (either directed or undirected). We can also extend this to weighted graphs, however binary networks are presented for simplicity. We can represent this graph by the adjacency matrix $G \in \{0, 1\}^{n \times n}$. We consider binary treatments denoted by a treatment vector $\mathbf{a} \in \{0, 1\}^n$ and let denote the potential outcome $Y_i(\mathbf{a}) \in \mathbb{R}$, under a treatment assignment \mathbf{a} , and Y_i denote the actual observed outcome. Lastly, we assume that we have access to pre-treatment node-level covariates $X_i \in \mathbb{R}^m$. In the remainder of the paper let O and o denote the usual big and little oh notation and O_P and o_P denote the stochastically bounded and convergence to 0 in probability for sequences of random variables. We use \widetilde{O} if we are suppressing logarithmic factors in the rate. Let $||\cdot||_p$ denote and p-norm, and let $||\cdot||_F$ denote the Frobenius norm.

2.1 A structural causal model

We use the framework of structural causal models, a nonparametric extension of structural equation models (Pearl, 2009). Similar approaches have been studied by Ogburn et al. (2022) and Auerbach and Tabord-Meehan (2021) in the case of fully observed networks. We derive a model that is amenable to estimation with partial data.

Let $Y_i(\mathbf{a})$ denote the potential outcome of Y_i under a treatment allocation \mathbf{a} . The exposure mapping V_i is represented as a function f_V such that $V_i = f_V(\mathbf{a}, \varphi_i(G)) \in \mathbb{R}^{p_V}$ where φ_i is the relevant graph information for individual *i* relative to their position with respect to treated individuals. We also allow for the potential outcome to be modulated by some additional confounder $S_i = f_S(\mathbf{X}, \vartheta_i(G)) \in \mathbb{R}^{p_S}$. We model the potential outcomes Y_i as a function of the exposure, type-value S_i and some additional noise ε_Y

$$Y_i = f_Y(S_i, V_i, \varepsilon_Y) \tag{1}$$

The benefits of structural causal models are that they allow for the characterization of all causal effects in a system, as well as the distributions of counterfactuals. However, they require correct specification of the causal process, i.e. correct specification of the exposure map and the relevant confounders. Even if one can propose a model for interference, estimation is not straightforward due to the fact that we only observe partial graph information in G^* . Many common models of interference can be expressed as structural causal models, and can be thought of as parameterizations of $f_Y(S_i, V_i, \varepsilon_Y) = f_Y(S_i, V_i, \varepsilon_Y; \beta_0)$. This then reduces the challenge to estimating β_0 using partially observed data. The exogenous noise, ϵ_Y , within our model is likely influenced by the graph structure, as interactions and peer effects can induce correlations in outcomes that extend beyond individual exposures. This complexity suggests that the noise, even if initially considered as external to the model, is intertwined with the network dynamics, reflecting the propagation and interference effects inherent in our structural causal framework.

We distinguish two types of target parameters. The first are the **outcome model** parameters, which parameterize the distribution of the outcome, exposure, and confounder (Y, S, V). Specifically, $f_Y(S_i, V_i, \epsilon_Y) = f_Y(S_i, V_i, \epsilon_Y; \beta_0)$ under parameterization $\beta \in \mathbb{R}^p$. The true model parameters are $\beta_0 \in \mathbb{R}^p$, identifiable through a moment equation m, $\mathbb{E}[m(Y_i, S_i, V_i, \beta_0)] = 0$, or through regression. In a simple diffusion model, this is the probability of infecting a neighboring node $q \in [0, 1]$.

The second set of parameters we consider are the **causal** parameters, those involving the distributions of the counterfactuals. The main causal parameter we will consider is the expected average potential outcome on the complete network G, $\Psi(\mathbf{a}|G) = \frac{1}{n} \sum_{i=1}^{n} \mathbb{E}[Y_i(\mathbf{a})]$, though these can also be made conditional on a covariate x: $\Psi(\mathbf{a}|x, G)$. Leveraging the structural causal model, we can define the these causal effects in terms of the structural causal model. We illustrate conditions for identification of these causal effects in section 2.3. While we focus on defining causal quantities through conditional means, the nonparametric identification can also apply to other functionals like quantiles.

Inference for learning the causal parameters under the above assumptions now amounts to learning the distributional relationship between Y_i and S_i, V_i . We consider settings where the assignment of treatments can be manipulated by an experimenter, which we discuss in section 4. If one leverages this model, either through assumption or estimation, then we can use a structural causal model to generate expected potential outcomes under different treatment assignments $f_Y(S_i, V_i, \varepsilon_Y)$, which is precisely what is done in the case of seeding. A contrast of these frameworks is included in the Appendix in section A.1. The applicability of a model to a new population parallels challenges in distribution shift, as explored in Shimodaira (2000) or Wilkins-Reeves et al. (2024).

Adding structure to the potential outcomes model is standard in fields like economics, where researchers often propose models to explain how information or behaviors spread across networks. Many of these models include a temporal element. In our setting, we consider outcomes at a fixed time T, i.e., $Y_i(\mathbf{a}) = Y_{i,T}(\mathbf{a})$. For instance, Banerjee et al. (2013) explore a latent diffusion process in micro-lending, Banerjee et al. (2019) study a hearing model for information diffusion, and Beaman et al. (2021) analyze behavior adoption in agriculture through complex contagion. Additionally, Centola and Macy (2007) differentiate the spread of information, often through single links, from behaviors that require multiple neighbors for network propagation.

2.1.1 Example: Contagion as a structural causal model

A foundational model of information diffusion is based on simple contagion, and generalizations of SIR (Susceptible-Infected-Recovered) models on networks (Kermack and McKendrick, 1927; Giles, 1977). These models have been further studied and extended in various settings (Jackson and Yariv, 2006; Aral et al., 2009; Romero et al., 2011; Chierichetti et al., 2011; Banerjee et al., 2013, 2019). Here we illustrate how the base model, under which many extensions are built, can be interpreted as a structural causal model. This interpretation can also be applied to complex contagion settings Centola and Macy (2007); Beaman et al. (2021); Cencetti et al. (2023).



(d) Contagion process at (e) Time period T = 1. (f) Time period T = 2. time period T = 0.

Figure 1: Contagion process where a single node is seeded in time T = 0 in blue, and infected nodes displayed in orange at times T = 1 and T = 2.

Consider a scenario where initially infected (treated) seeds **a** transmit the infection to each neighbor with probability q at each time-step $t \in 1, 2, ..., T$, after which they are no longer infectious. An infection status at time t is denoted as $Y_{it} = 1$. The overall outcome $Y_i = 1$ indicates whether a node was infected at any time up to T. For a simple case with T = 2, we model the transmission using Bernoulli random variables $\epsilon_{ij} \sim \text{Bernoulli}(q)$, representing potential infection from node i to node j. Let $\mathbf{E}_{ij} = \epsilon_{ij}$ and $\mathbf{D} = \mathbf{E} \odot G$. This setup is depicted in Figure 1. Given a random sample of the directed graph \mathbf{D} , one can characterize what would have happened if a node were treated, which is precisely the counterfactual. For instance in Figure 1 we seed the left most which proceeds to propagate in steps 1 and 2. Additionally, one can construct the relevant exposure map for any fixed number of time steps T.

2.1.2 Examples of Exposure Maps

We consider several examples of exposure maps, though this list is not exhaustive.

Example 2.1 (Local Interaction Effects). Simple examples of local network effects may include the total number of treated neighbors, $V_i = \sum_j G_{ij}a_j$, or the treated fraction of one's neighbors $V_i = \sum_j \frac{G_{ij}}{d_i}a_j$, where $d_i = \sum_j G_{ij}$ is degree.

Example 2.2 (Risk-Sharing Networks (Ambrus et al., 2014)). Equilibrium risk sharing is that the graph consists of C mutually exclusive communities such that any endowment vector within the community is aggregated and shared evenly. Let treatment **a** be an "endowment" and let $\overline{\mathbf{a}}_c = \sum_{j \in c} a_j$ be the sum of the endowment vector for community c, with |c| denoting its size. Then, $V_i = f_V(\mathbf{a}, \varphi_i) = \overline{\mathbf{a}}_c \cdot |c|^{-1}$. That is, the exposure is just a function of the total endowment of the community and nothing more.

Example 2.3 (Hearing Information (Banerjee et al., 2019)). Many phenomena, like the spread of diseases, information, or social behaviors, can be effectively modeled as contagion processes. These models show how such phenomena spread through networks (Keeling and

Rohani, 2008; Centola and Macy, 2007; Barrat et al., 2008; Pastor-Satorras et al., 2015; Cencetti et al., 2023).

Banerjee et al. (2019) introduces a message-passing model based on such a contagion process. The treatments, denoted by \mathbf{a} , represent a seed piece of information disseminated over a series of time steps, from 1 to T. After T time steps, no further message spreading occurs. We define a "hearing matrix" \mathbf{H}_0 , which calculates the expected number of times person j hears information from person i after T time steps, based on transmission probabilities.

The expected total number of messages that person j hears by time T is represented by V_i (the exposure) which affects their response Y_i through a link function Λ :

$$\mathbb{E}[Y_i|V_i] = \Lambda(\beta_0 + \beta_1 V_i).$$

A common assumption is propose a single transmission probability for each individual q, thus giving structure to the exposure map:

$$V_i = (\mathbf{H}\mathbf{a})_i$$
 the *i*th element of this vector
where $\mathbf{H} = \sum_{t=1}^T q^t G^t$

It is straightforward to generalize this to include heterogeneity in the diffusion time steps β_t and illustrate this model in equation (2):

$$\mathbb{E}[Y_i|V_i] = \Lambda(\beta_0 + \sum_{t=1}^T \beta_t \mathbf{e}_i^T G^t \mathbf{a}) = h(S_i, V_i; \beta).$$
(2)

Furthermore, we can relax this model to allow for additional heterogeneity through graphlevel statistics S_i , which may include node-level covariates X_i or individual graph-level information such as the degree d_i .

2.2 Examples of Partially Measured Network Data

Now that we've established our framework for interference, we next return to the data used for estimation. In our setting, we do not have access to the full graph G, but rather, have access to some summarizing function the graph $G^* = \zeta(G)$. Tsiatis (2006) uses the term coarsened data to refer to such partial measurements of missing data in general, not necessarily in the network setting. "Coarsened" is apt in our setting because our method works on partial graph structure that give (either directly or by construction) an estimate of linking rates across population members with different combinations of observable traits. A non-exhaustive set of examples of partially measured network data include induced subgraphs or egocentric sampling (Freeman, 1982; Almquist, 2012), respondent driven sampling (Heckathorn, 1997), aggregated relational data (Killworth et al., 1998), respondent driven sampling (Heckathorn, 1997; Goel and Salganik, 2009, 2010; Green et al., 2020) and more.

Example 2.4 (Induced subgraph). We sample $m \leq n$ of nodes in the graph randomly, with at least one node from each of the K communities. Let $G^* = G_{I_m,I_m}$ be the sub-graph induced by these m nodes where $I_m \subset \{1, 2, ..., n\}$ are the set of nodes that are sub-sampled from the whole population.

Example 2.5 (Respondent Driven Sampling). Let $i \in I_m \subset \{1, 2, ..., n\}$ denote the indices of a sample of individuals obtained through respondent driven sampling. An initial number of individuals are recruited as seeds, and subsequent individuals are recruited via referrals from the others in a population. Under this process we receive a subgraph of connected individuals G_{I_m,I_m} as well as the list of connections to additional nodes $I_{n\setminus m} = \{1, 2, ..., n\} \setminus$ $I_m \ G^* = G_{I_m,I_m}, G_{I_m,I_n\setminus m}.$

Example 2.6 (Aggregated Relational Data). Aggregated relational data consists of aggregated sums of connections to nodes of a given trait. Typically this is collected from a survey consisting of questions such as "How many many people do you know with [X] trait?". For a set of T traits, this consists of $X_{it}^* = \sum_{i=1}^n G_{ij}I(t_j = t)$.

In order to infer about the distribution of the missing part of the graph, we propose that $G \sim \theta_0$ where we assume that $\theta_0 \in \Theta$ denotes the parameters of a random graph model. In this case, for each *i*, there is an a latent ξ_i parameter such that $P(G_{ij} = 1|\xi_i, \xi_j) = \tilde{g}(\xi_i, \xi_j)$ for some function symmetric, measurable, \tilde{g} , known as a graphon (Lovász and Szegedy, 2006; Orbanz and Roy, 2015). Many common graph models, such as latent space models (Hoff et al., 2002; Handcock et al., 2007; Lubold et al., 2023; Wilkins-Reeves and McCormick, 2022), are included in this category. Graphons are appealing in this context because, following Airoldi et al. (2013) and Gao et al. (2015), they can be approximated arbitrarily well using latent types assigned to each node. Said another way, graphons introduce complex dependence in the network-generating mechanism through clustering induced by latent types associated with each node. In our inferential procedures in section 3, the general procedures involve estimation from a missing data perspective. This will involve estimating the graph model $\hat{\theta} := \hat{\theta}(G^*)$ then inferring about the distribution $G|G^*, \hat{\theta}$.

2.3 Nonparametric Identification of Causal Effects

We first illustrate an identification procedure for the causal effect without a-priori imposing any model structure. These are analogous to standard causal identification assumptions, adapted to our framework.

Definition 2.1 (Exposure Weak Ignorability). We say that an exposure assignment is weakly ignorable if the following holds:

$$Y_i(v) \perp \{V_i = v\} | S_i$$

Conditioning the graph confounder S_i captures the heterogeneity of the outcomes when observed with a given exposure. In simple contagion models, nodes are equivalent, and this independence occurs naturally without conditioning. section 5.1 discusses an example from Ugander and Yin (2023) where conditioning on node degree suffices for any randomization.

Definition 2.2 (Exposure Consistency). Exposure consistency holds if

$$V_i = v \implies Y_i = Y_i(v) = Y_i(\mathbf{a})$$

where $Y_i(v)$ is the potential outcome of individual i for the exposure v.

This assumption can be simply understood as the exposure is correctly specified.

Definition 2.3 (Conditional Independence of the Graph and Outcome). We assume that the outcome is conditionally independent of the outcome conditional on the exposure and the graph generative parameters

$$Y_i(\mathbf{a}) \perp \!\!\!\perp G | V_i, S_i.$$

This assumption states that once we have adjusted for V_i and S_i , then the potential outcomes are independent of the network G. These assumptions allow us to express the causal estimand through observational data.

 $P(Y_i(\mathbf{a}) = y | S_i = s, G) = P(Y_i(v) = y | S_i = s, G) \text{ By the exposure mapping}$ $= P(Y_i(v) = y | V_i = v, S_i = s, G) \text{ By weak ignorability}$ $= P(Y_i = y | V_i = v, S_i = s, G) \text{ By consistency}$ $= P(Y_i = y | V_i = v, S_i = s) \text{ Graph conditional independence}$ $\implies \Psi(\mathbf{a}|G) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}[Y_i | V_i = f_V(\mathbf{a}, \varphi_i), S_i = f_S(\mathbf{X}, \vartheta_i)]$

For brevity, we denote the true conditional mean $\mathbb{E}[Y_i|V_i = v, S_i = s]$ as $h_0(s, v)$ and denote $h(s, v; \beta)$ a model to estimate $h_0(s, v)$. Given a network model θ , observed graph data G^* ,

and a conditional model $h(s, v; \beta)$ we can also define the expected average treatment effect

$$\Psi(\mathbf{a}|\beta, G^*, \theta) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}[h(f_V(\mathbf{a}, \mathbf{X}; \varphi_i), f_S(\mathbf{X}; \vartheta_i); \beta) | \mathbf{a}, \mathbf{X}, G^*, \theta]$$
(3)

where under the correct model conditional model and graph model $\mathbb{E}[\Psi(\mathbf{a}|G)|\mathbf{a}, \mathbf{X}, \theta_0] = \Psi(\mathbf{a}|\beta_0, G^*, \theta_0)$. In Appendix A.2.5, we illustrate when this population average effect under any draw of the network $\Psi(\mathbf{a}|G)$ will be close to the average over the model class $\Psi(\mathbf{a}|\beta_0, G^*, \theta_0)$; and study plug-in estimators $\widehat{\Psi}(\mathbf{a}|G) = \Psi(\mathbf{a}|\widehat{\beta}, G^*, \widehat{\theta})$.

3 Inference

We outline our method for estimating parameters with partial network data. Developing these results requires two theoretical tools: a fast estimation rate for network model parameters θ_0 , and a suitable central limit theorem for scenarios with correlated outcomes. Outcome Model Parameters and Estimators Next we consider estimating the outcome model parameters β_0 . We present two methods for estimating such parameters, instrumentation in a linear model, and Z estimators. The iterated expectation procedure for estimating such parameters was introduced in Chandrasekhar and Lewis (2011), however, we extend inference to the single network setting. Similar approaches exist for peer effects models (Boucher and Houndetoungan, 2020).

3.0.1 Estimation in Linear Models

We first illustrate identification of the conditional model under a linear model assumption.

$$Y_i = \beta_0^T \widetilde{h}(S_i, V_i) + \varepsilon_i$$

where $\mathbb{E}[\varepsilon_i] = 0$ and there can be general correlation $\operatorname{Var}[\varepsilon] = \Sigma$. Without access to the network data, one can recover the model parameters through conditional expectation

$$\mathbb{E}[\mathbb{E}[Y|S(G), V(G), G, \mathbf{a}, \mathbf{X}] | \mathbf{a}, \mathbf{X}, G^*, \theta_0] = \beta_0^T \mathbb{E}[\widetilde{h}(S(G), V(\mathbf{a}, G)) | \mathbf{a}, \mathbf{X}, G^*, \theta_0]$$

where we create a new set of features $\widetilde{H}_i = \mathbb{E}[\widetilde{h}(S_i(G), V_i(\mathbf{a}, G))|\mathbf{a}, \mathbf{X}, G^*, \theta_0]$ by averaging over the network model. Here identification comes from the variation of these averaged features \widetilde{H}_i over the population. More clearly, letting $\widetilde{\mathbf{H}} \in \mathbb{R}^{n \times p}$ denote the design matrix of this model, identification comes from the linear independence of the columns of $\widetilde{\mathbf{H}}$.

3.0.2 Z estimators

In other cases, parameters may be defined through a moment equation, and can be used to construct a Z-estimator, for example, generalized linear models (GLMs). These parameters can be identified using an estimating equation approach where given a moment function $\widetilde{m}(Y_i, S_i, V_i; \beta)$ such that $\mathbb{E}[\widetilde{m}(Y_i, S_i, V_i; \beta) | \mathbf{a}, \mathbf{X}, G] = 0$ if and only if $\beta = \beta_0$. Through the use of iterated expectations, we can define a new estimating equation, by marginalizing over the draws of the graph model then applying iterated expectations

$$m_i(Y_i;\beta,\mathbf{a},\mathbf{X},G^*,\theta) := \mathbb{E}[\widetilde{m}(Y_i,S_i,V_i;\beta)|Y_i,\mathbf{a},\mathbf{X},G^*,\theta]$$
(4)

where $\mathbb{E}[m_i(Y_i, S_i, V_i; \beta_0, \theta_0) | G^*, \mathbf{a}, \mathbf{X}, \theta_0] = \mathbb{E}[\mathbb{E}[\widetilde{m}(Y_i, S_i, V_i; \beta_0) | \mathbf{a}, \mathbf{X}, G] | G^*, \mathbf{a}, \mathbf{X}, \theta_0] = 0.$

Identification arises from the variation of exposure and confounders, such that $\beta = \beta_0$ if and only if $\mathbb{E}\left[\frac{1}{n}\sum_{i=1}^n m_i(Y_i, S_i, V_i; \beta, \theta_0) \middle| G^*, \mathbf{a}, \mathbf{X}, \theta_0\right] = 0$. Exact conditions vary by parameter, but GLMs can use a similar strategy as linear models.

3.1 Inference with partially measured data.

We introduce a general procedure for estimating the outcome model parameters. We also illustrate inference for estimation of a causal target parameter on a particular graph G.

We present a pseudo-code approach to the procedure in Algorithm 1. Let $\tilde{Z}_i = (Y_i, S_i, V_i)$ denote the full (including unobserved) data, and let $\mathbf{Z} = (\mathbf{Y}, \mathbf{a}, \mathbf{X}, G^*)$ denote the observed data.

Algorithm 1 Z-estimation overview

- 1: Define an model for the relationship of \mathbf{Y} given the exposures \mathbf{V} and confounders \mathbf{S} (for instance, a regression model $\mathbb{E}[Y|V,S] = h(v,s;\beta_0), \beta \in \mathcal{B} \subset \mathbb{R}^p$ with parameters which can be estimation via the estimating function $\widetilde{m}_n(\widetilde{\mathbf{Z}};\beta)$. Let $\widetilde{m}_n(\widetilde{\mathbf{Z}};\beta) = \frac{1}{n} \sum_{i=1}^n m(\widetilde{Z}_i;\beta)$ denote the empirical estimating function.
- 2: Estimate a model of the network, using the node-level covariates $\hat{\theta} := \hat{\theta}(G^*)$.
- 3: Estimate $\hat{\beta}$ by solving the estimating equation $m_n(\mathbf{Y}; \hat{\beta}, G^*, \hat{\theta}) = 0$, where $m_n(\mathbf{Y}; \hat{\beta}, G^*, \hat{\theta}) = \frac{1}{n} \sum_{i=1}^n m_i(Y_i; \beta, \mathbf{a}, \mathbf{X}, G^*, \theta)$ where m_i is defined in equation (4).
- 4: (optional) Plug in $\hat{\beta}$ to $\Psi(\mathbf{a}|\hat{\beta}, G^*, \hat{\theta})$.

Step 1 asks the practitioner to propose a response model given the treatment, i.e. the causal model in section 2.1. Step 2 estimates the generative model given the partial network data and the node covariates observed. We give theoretical results where the formation model is a stochastic blockmodel, then give rate estimation relative to the more general graphon approach. Step 3 estimates the parameter by marginalizing the estimating function over the graph model. Lastly, Step 4 is optional if the target parameter is a plug-in estimator of the causal parameter using the regression model. We discuss inference for the plug-in estimate of causal parameters using a delta method argument in the Appendix A.2.5. We next give our asymptotic results, then provide an example of this algorithm in section 5.1.

3.2 Asymptotic Results

The asymptotic results for both the Z-estimator and the linear model will depend on being able to establish a central limit theorem based on the exogenous noise. To establish asymptotic properties for our outcomes on a network, we extend the application of the central limit theorem (CLT) to structures not commonly associated with traditional time series or spatial dependencies. Nonetheless when the exogenous noise is correlated, we will need a method of handling the central limit theorem. Specifically, we utilize a general version of the CLT for dependent data from Chandrasekhar et al. (2023). For brevity in presentation, we leave the full detail of this central limit theorem to the appendix.

We denote $g_i(\mathbf{Z}; \beta) = m_i(\mathbf{Y}; \mathbf{a}, \mathbf{X}, \beta, G^*, \theta_0)$ to be the moment function evaluated using the true generative model and correspondingly $g_n(\mathbf{Z}; \beta) = \frac{1}{n} \sum_{i=1}^n g_i(\mathbf{Z}; \beta)$. Further, define the (normalized) random vector of the estimating function evaluated at the correct model parameters $\mathcal{E}_i = \frac{1}{n} g_i(\mathbf{Z}; \beta_0)$. And lastly let $D_n(\mathbf{Z}; \beta_0) = \nabla_\beta g_n(\mathbf{Z}; \beta_0) \in \mathbb{R}^{p \times p}$ denote the gradient of the estimating equation $g_n(\mathbf{Z}; \beta)$. To develop valid inference, we must estimate the graph model quickly enough to disregard the graph estimation component during inference. We will next present the theorem and discuss the assumptions further.

Assumption 3.1 (Regularity Conditions for Z-Estimation). Suppose the following conditions hold for all n.

Consistency for a Z estimator

A1.
$$\mathbb{E}[g_n(\mathbf{Z};\beta)] = 0$$
 for $\beta = \beta_0$ and for all $\epsilon > 0$, $inf_{||\beta-\beta_0||>\epsilon}\mathbb{E}[g_n(\mathbf{Z};\beta)] > 0$
A2. $\sup_{\beta\in\mathcal{B}} \left| \left(\frac{\partial}{\partial\beta}\right)^l g_n(\mathbf{Z};\beta) - \left(\frac{\partial}{\partial\beta}\right)^l \mathbb{E}[g_n(\mathbf{Z};\beta)] \right| = o_P(1)$ for $l \in \{0,1,2\}$

Graph Model Regularity conditions

B1. $\hat{\theta}$ is an s(n)-consistent estimate of the graph parameters $||\hat{\theta} - \theta_0|| = o_P(s(n))$

B2. $\sup_{\beta \in \mathcal{B}} |m_n(\mathbf{Z}; \beta, \theta) - m_n(\mathbf{Z}; \beta, \theta')| \leq b_n(\mathbf{Z}) ||\theta - \theta'||$ where $b_n(\mathbf{Z}) = O_P(1)$ (that is, $b_n(\mathbf{Z})$ is stochastically bounded).

Central Limit Theorem (CLT)

C1. The random vectors $\mathcal{E}_{1:n}$ satisfy the affinity set conditions of Chandrasekhar et al. (2023) (restated as Theorem A.2.2 in the appendix) with corresponding covariance matrix $\Gamma_n = Var[\sum_{i=1}^n \mathcal{E}_i]$. Where $r(n) := \sqrt{\lambda_{min}(\Gamma_n)}$.

Theorem 3.2 (Single Network Z-estimator Asymptotics). Suppose that $\hat{\beta}$ is computed as per Algorithm 1, Assumptions 3.1 hold, and s(n) = o(r(n)). Then:

$$\Gamma_n^{-1/2} D(\beta_0)(\widehat{\beta} - \beta) \to_d N(0, I_p)$$
(5)

Where $\mathbb{E}[\nabla_{\beta}g_n(\mathbf{Z};\beta)|\mathbf{a},\mathbf{X},G^*,\theta_0]\Big|_{\beta=\beta_0} = D(\beta_0)$

The first set of assumptions ensures the consistency of Z-estimators, typically derived from uniform laws of large numbers as discussed in Andrews (1987); Newey and McFadden (1994). The second set involves conditions that make the graph model's estimation negligible, requiring the estimating functions to be smooth with respect to the graph parameters.

The final set of assumptions, stated in C1, are utilized so that \mathcal{E}_i satisfy a central limit theorem Chandrasekhar et al. (2023). This assumption is required if the data exhibit further dependence after controlling for graph parameters (if, for example, there are latent factors that impact both outcomes and the propensity to form ties). The main idea of Chandrasekhar et al. (2023) is to represent dependence in terms of "affinity sets" where the majority of dependence structure is captured within sets, leaving little between sets. In the modelling of social behaviours beyond just considering outcomes as a function of the exposure observed, outcomes may be further correlated, beyond examples of spatial dependence or heteroskedasticity. In practice we can include these dependencies through correlation terms matching the generative graph model, such as between blocks of a stochastic blockmodel or via latent positions in a latent space model.

Here, r(n) describes the effective rate at which the variance converges. For the estimation of the graph model θ_0 to be considered negligible, it must occur more rapidly than r(n). In cases of independent or minimally dependent noise, it is typical for $r(n) \approx n^{-1/2}$. Alternatively, in different scenarios, \mathcal{E}_i might exhibit correlation within densely connected blocks of the network, such as during a diffusion process in a stochastic blockmodel with k_n densely linked blocks (refer to Chandrasekhar et al. (2023), section 4.4). In such cases, r(n)is generally on the order of $k_n^{-1/2}$. If both r(n) and s(n) approach zero, but $\frac{s(n)}{r(n)}$ diverges or stabilizes at a nonzero constant, a consistent estimator for the outcome model parameters can still be obtained. However, its asymptotic distribution may be influenced by the graph model estimation, necessitating a tailored inference approach.

An analogous argument follows when conducting inference using a linear model. For the sake of brevity and avoiding repetition, we include it in the Appendix in section A.2.2. In Theorem 3.3 we present a summary.

Theorem 3.3. Let $\widetilde{H}_i(\theta) = \mathbb{E}[\widetilde{h}(S_i(G), V_i(G)) | \mathbf{a}, \mathbf{X}, G^*; \theta]$. The OLS estimator uses the model averaged coefficients $\widetilde{H}_i(\theta)$ in place of the true unobserved coefficients \widetilde{h}_i . Let $\mathsf{H}_n(\theta) = \frac{1}{n} \sum_{i=1}^n \widetilde{H}_i(\theta) \widetilde{H}_i^T(\theta)$. Given an estimate of the model parameters $\widehat{\theta}$, we define the

$$\widehat{\beta}_{lm} = \mathsf{H}_n^{-1}(\widehat{\theta}) \frac{1}{n} \sum_{i=1}^n \widetilde{H}_i(\widehat{\theta}) Y_i$$

Let $u_i = (\widetilde{h}(S_i(G), V_i(G)) - \widetilde{H}_i(\theta_0))\beta_0 + \epsilon_i$ and let $\Gamma_n = Var\left[\frac{1}{n}\sum_{i=1}^n u_i\right]$ Suppose the conditions of Theorem A.2.3 in the Appendix hold. Then

$$\Gamma_n^{-1/2} \mathsf{H}_n(\widehat{\theta})(\widehat{\beta}_{lm} - \beta_0) \to_d N(0, I_p)$$

3.3 Network Model Estimation

We next discuss the estimation of the generative model for the network using a variety of data types. We first demonstrate results for estimating parameters in a stochastic blockmodel. We then extend the result to view the blockmodel as an approximation of a graphon. Breza et al. (2020) and Breza et al. (2023) consistently estimate a generative model for ARD with mutually exclusive traits. We extend this work by introducing a novel method for estimating the stochastic blockmodel with non-mutually exclusive traits using constrained least squares approach. This innovation has major implications for practice since it can dramatically reduce survey length (since asking about multiple categories requires constructing separate questions for each intersection to make the traits mututally exclusive). Our approach applies to a range of partial network data, not just ARD, but we summarize the resulting rates using ARD for a variety of model classes in the appendix in Table A.2.3 for comparison with existing literature.

In the main text, we concentrate on estimating the stochastic blockmodel using ARD. In the Appendix in section A.2.3.1 we estimate generative models using partial network data such as subgraph sampling and develop similar rates for the stochastic blockmodel for subgraph sampling and reference a similar result for respondent driven sampling.

3.3.1 SBM Estimation with ARD

Recall that X_{it}^* represents a set of ARD response vectors. Breza et al. (2023) show that we can consistently estimate the connection probabilities between latent types, however, we present an improved version of the SBM estimator which allows for an non-mutually exclusive traits. Let n_t denote the total number of individuals of trait type t. Let N'_k denote the nodes in our sample in group k, and let n_k denote the number of nodes in the graph in group k. We cluster the node memberships according to Algorithm 2.

After we obtain a clustering, we can estimate the stochastic blockmodel. Let $\widehat{\Omega}_{kt} = \widehat{N}_{kt}/N_t$ where N_{kt} are the number of traits in the estimated group k and with trait t, and N_t are the number of individuals with trait t, and $\Omega_{kt} = N_{kt}/N_t$, the analogous population quantity. We next define the probability matrix of observing a connection of group k with

Algorithm 2 ARD SBM clustering procedure

- 1: Count the number of individuals with each trait n_t
- 2: Denote the normalized ARD responses $X_{it}^{\dagger} = X_{it}^*/n_t$.
- 3: Cluster the normalized ARD response vectors $\{X_i^{\dagger}\}_{i=1}^T$ into K groups using hierarchical agglomerative clustering into a set of clusters $\hat{k}_i \in \{1, 2, \dots, K\}$

a trait t. $\widetilde{\mathbf{P}}_{kt} = \sum_{k'} \mathbf{P}_{kk'} \omega_{k't}$, where $\widetilde{\mathbf{P}}_{kt} = P(G_{ij} = 1 | k_j = k, t_i = t)$. This relationship can be expressed in a linear system $\widetilde{\mathbf{P}} = \Omega \mathbf{P}$ where $\Omega \in \mathbb{R}^{T \times K}$ and $\Omega_{kt} = \omega_{kt}$. If Ω is of full column rank, then a unique solution will exist as:

$$\widehat{\mathbf{P}}_{kk'} = \left(\widehat{\Omega}^{\mathsf{T}}\widehat{\Omega}\right)^{-1}\widehat{\Omega}^{\mathsf{T}}\widehat{\widetilde{\mathbf{P}}} \quad \text{where } \widehat{\widetilde{\mathbf{P}}}_{kt} = \frac{1}{n_k n_t} \sum_{i \in \widehat{N}_k} X_{it}^*.$$

In general, one can symmetrize $\widehat{\mathbf{P}}_{kk'}$ after the estimate to ensure the constraints of an undirected stochastic blockmodel are satisfied. Alternatively, once can also minimize the constrained least squares objective which can be implemented using standard convex solvers such as CVX (Fu et al., 2020)

$$\widehat{\mathbf{P}} = \underset{0 \le \mathbf{P} \le 1: \mathbf{P} = \mathbf{P}^{\mathsf{T}}}{\operatorname{arg\,min}} \sum_{i=1}^{n} \sum_{t=1}^{T} (\widetilde{X}_{it} - \sum_{k'} \widehat{\Omega}_{k't} P_{k',k_i})^2.$$

Breza et al. (2023) develop a method for consistently estimating the stochastic blockmodel. We extend their result by obtaining a rate for estimating model parameters (Lemma 3.4) and relax the assumption that of mutually exclusive traits. We differentiate between the estimated cross-group probabilities $\mathbf{P}^{(\hat{\mathbf{k}})}$ and those under known membership $\mathbf{P}^{(\mathbf{k})}$.

Lemma 3.4. Suppose that we use the clustering strategy outlined in section 3.3.1 to cluster the observations based on aggregated relational data. Let $Z_k = (\widetilde{\mathbf{P}}_{k1}, \dots, \widetilde{\mathbf{P}}_{kT})$ and $\widetilde{\mathbf{P}}_{kt} = P(G_{ij} = 1 | k_i = k, t_j = t)$. Assume also that $\inf_{k,k'} ||Z_k - Z_k||_2 > 0$ and that $T \ge K$ where T is the number of discrete traits asked about and K is the true number of clusters.

Let $\hat{\mathbf{k}}$ denote the estimated cluster memberships and let $\widehat{\mathbf{P}}^{(\hat{\mathbf{k}})}$ be the corresponding estimate of the cross block probabilities. Let $\Omega_{kt} = N_{kt}/N_t$ denote the matrix which involves the fraction of the individuals in cluster k who also have trait t, and $\widehat{\Omega}$ the estimated counterpart based on membership clusters. Let $C_{\Omega} = \lambda_{max}((\Omega^T \Omega)^{-1})$ and $\lambda_{max}(\cdot)$ denotes the largest eigenvalue of a matrix and $C_{\Omega} > 0$. Then with probability at least $1 - \delta - \frac{1}{n}$

$$||\widehat{\mathbf{P}}^{(\widehat{\mathbf{k}})} - \mathbf{P}^{(\mathbf{k})}||_1 \le C_\Omega \frac{KT}{n} \sqrt{\frac{\log(2/\delta)\log(KT)}{2}}$$

We contrast our results to the optimal estimation rate for a stochastic blockmodel from Gao et al. (2015), $\tilde{O}_P(n^{-1/2})$. Our rate appears faster due to the complexity difference in clustering problems. Our clustering benefits from node-level traits, which provide extra information. As the network grows, the normalized ARD vector converges to its mean, simplifying clustering and resulting in a faster rate.

3.3.2 Misspecification of the Graph Model

We use a stochastic blockmodel as it effectively approximates a general graphon class. Even if θ_0 belongs to a smooth graphon class rather than a stochastic blockmodel, we can still bound the bias in estimating the relevant model parameters. Consider a scenario where edges are generated under a true graphon model \tilde{g} where $\eta_{ij} = \tilde{g}(\xi_i, \xi_j) = P(G_{ij} =$ $1|\xi)$ where $\xi \sim_{iid} P_{\xi} \in [0, 1]$. Let $\mathcal{H}_{\alpha}(M)$ denote a smooth graphon class defined via the α -M-Hölder class as follows. Let $\mathcal{D} = [0, 1]^2 \cap x \leq y$ denote the domain of (x, y). We define the norm $||\tilde{g}||_{\mathcal{H}_{\alpha}}$ as:

$$||\widetilde{g}||_{\mathcal{H}_{\alpha}} = \max_{j+k \leq \lfloor \alpha \rfloor} \sup_{x,y \in \mathcal{D}} |\nabla_{jk} \widetilde{g}(x,y)| + \max_{j+k=\lfloor \alpha \rfloor} \sup_{(x,y) \neq (x'y') \in \mathcal{D}} \frac{\nabla_{jk} \widetilde{g}(x,y) - \nabla_{jk} \widetilde{g}(x',y')}{(|x-x'|+|y-y'|)^{\alpha-\lfloor \alpha \rfloor}}$$

and the Hölder class corresponding to this norm as

$$\mathcal{H}_{\alpha}(M) = \{ ||\widetilde{g}||_{\mathcal{H}_{\alpha}} \le M : \widetilde{g}(x,y) = \widetilde{g}(y,x); 0 \le \widetilde{g}(x,y) \le 1 \}.$$

Prior work has focused on the approximability of a stochastic blockmodel to any element of a smooth graphon class. In particular there will always be some assignment of block memberships such that we can bound the 2-norm probability deviation from the true model.

Lemma 3.5. Suppose that θ_* corresponds to a true graphon model and θ_0 a corresponding approximating stochastic blockmodel. Denote the population estimating function, as a function of the model parameters

$$L_n(\beta, \theta) = \mathbb{E}[\widetilde{m}_n(\mathbf{Z}; \beta) | \mathbf{a}, \mathbf{X}, \theta]$$

where $L_n(\beta_0, \eta_0) = 0$ defines the population parameter β_0 under the misspecified model θ_0 , and let $L_n(\beta_*, \theta_*) = 0$ define the population solution β_* to the correctly specified graph model θ_* . Let η_0 and η_* be the pairwise edge probabilities corresponding to the models θ_0, θ_* respectively. Finally assume that:

D1. \mathcal{B} is compact

D2.
$$\sup_{\beta \in \mathcal{B}} |L_n(\beta, \eta) - L_n(\beta, \eta_*)| \le L ||\eta - \eta_*||_2 / n$$

D3. $\min_j \frac{\partial}{\partial \beta_j} L_n(\beta, \eta_*) \Big|_{\beta = \beta_*} = \lambda > 0$

Then the approximation error under the graph misspecification is bounded by the rate:

$$||\beta_0 - \beta_*|| = O(\lambda^{-1} K^{-\alpha \wedge 1}) \quad where \ a \wedge b = \min(a, b).$$
(6)

In practice, we don't directly select clusters; misspecified clusters relate to observed traits, thus this bound holds only under good alignment of clusters. This bound is a worstcase scenario and may be overly conservative regarding observed bias. Future work could involve sensitivity analysis of the response function and the latent graph model.

4 Experimental Design

So far, our focus has been on estimating model parameters given a treatment assignment **a**. We now explore experimental design methods that leverage partial network data to choose **a** to minimize the variance of our estimands. Leveraging partial network data for this purpose is particularly appealing in practice, since it requires substantially less investment than collecting full network data and could be collected as part of creating a sampling frame in settings where researchers collect data to construct the frame.

We consider saturation randomization experiments, which divide the dataset into J clusters of size n_j . A proportion τ_j of each cluster is assigned the treatment, totaling $n_t = \sum_{j=1}^{J} \tau_j n_j$, and generally will not be the same "blocks" as those in a graph model if the graph model uses discrete factors (e.g. stochastic blockmodel). Practically, due to budget constraints, the set of possible saturation levels τ is limited to $\mathcal{T} \subset [0, 1]^J$. For example, this could be due to limited resources like a finite vouchers in a vaccine trial.

4.1 Bayesian Optimization of Asymptotic Regression Estimators

Our goal is to optimize the asymptotic variance of a function of the model parameter $\hat{\beta}$ in section 3. We highlight this by optimizing the variance of the estimates of linear contrasts of the parameters $\phi^T \beta$. When using the stochastic block model for the network model these treatment blocks could align with the model blocks, however this need not (and likely won't) be the case. They could, instead, be based on observed characteristics (e.g. geography, classrooms).

Denote the variance of the target contrast parameter conditional on the treatment assignment as **a**: $v^{\phi}(\mathbf{a}; \theta) = \operatorname{Var}(\phi^T \hat{\beta} | \mathbf{a}, \theta)$. Ideally, the goal is to find a treatment assignment a^* that minimizes the variance of the contrast: $a^* = \arg \min_{\mathbf{a} \in \{0,1\}^n} v^{\phi}(\mathbf{a}; \theta)$. Without added structure, optimizing treatment assignments is NP-hard, requiring a search over 2^n possible assignments. By changing the objective to one where we optimize over a set of saturation levels over a set of groups $\tau \in [0, 1]^J$, we simplify the problem so that it is no longer NP- hard (i.e. since $J \ll n$ typically) and is therefore tractable. The distribution of treatment assignments, **a**, under τ is denoted by P_{τ} , and we aim to minimize:

$$\mathcal{V}(\tau;\theta) = \mathbb{E}_{\mathbf{a} \sim P_{\tau}}[v^{\phi}(\mathbf{a};\theta_0)].$$

In Algorithm 3, we present a method for evaluating the variance of a linear model using a generic feature map \tilde{h} for a given treatment assignment **a** and a graph model θ . A general approach for Z-estimators is detailed in the Appendix. Algorithm 3 operates under specific assumptions about the covariance matrix Σ , which may include correlations within densely connected network components. We will present our algorithm for minimizing this variance using Bayesian optimization, which accounts for the uncertainty in the outcome, given a graph model. In the appendix we give an extension which also which incorporates network model uncertainty $\hat{\theta}$ (section A.2.7).

Algorithm	3 Saturation Randomized Design Variance.
1: Inputs:	Variance structure $\operatorname{Var}[\mathbf{u}] = \Sigma$, Model estimate $\widehat{\theta}$.

2: Sample L draws from the graph model $\{\widehat{G}^{(l)}\}_{l=1}^L\sim \widehat{\theta}|G^*$

- 3: Sample R treatments $\{\mathbf{a}_r\}_{r=1}^R$ according to the block saturation levels τ .
- 4: for $r \leftarrow 1$ to R do
- 5: Compute the averaged features over draws from the graph model $\{\widehat{G}^{(l)}\}_{l=1}^{L}$

$$\widehat{H}_{ir}(\mathbf{a}) = \frac{1}{L} \sum_{l=1}^{L} \widetilde{h}(S_i(\widehat{G}^{(l)}) V_i(\mathbf{a}_r; \widehat{G}^{(l)}))$$

- 6: Compute the Hessian $\widehat{\mathsf{H}}_n(\mathbf{a}_r) = \frac{1}{n} \sum_{i=1}^n \widehat{H}_{ir}(\mathbf{a}) \widehat{H}_{ir}^T(\mathbf{a}).$
- 7: Compute the design matrix $\widehat{H}_r^T(\mathbf{a}) \in \mathbb{R}^{n \times p}$ where each row is $\widehat{H}_{ir}(\mathbf{a})$.
- 8: Compute the variance for a single draw of the treatment vector \mathbf{a}_r :

$$v^{\phi}(\mathbf{a}_r; \widehat{\theta}) = \phi^T \widehat{\mathbf{H}}_n^{-1}(\mathbf{a}_r) \widehat{H}_r^T(\mathbf{a}) \Sigma \widehat{H}_r(\mathbf{a}) \widehat{\mathbf{H}}_n^{-1}(\mathbf{a}_r) \phi$$

9: end for

10: Average over each of the draws $\mathcal{V}(\tau; \hat{\theta}) = \sum_{r=1}^{R} v^{\phi}(\mathbf{a}_r; \hat{\theta})$

Bayesian Optimization. Calculating the average variance $\mathcal{V}(\tau; \hat{\theta})$ in Algorithm 3 is computationally intensive to evaluate and often non-convex. Since the number of cluster saturation tends to be relatively small, this suggests that Bayesian optimization is an appropriate method for minimizing this saturation variance. Let $\mathcal{V}(\tau) := \mathcal{V}(\tau; \hat{\theta})$ denote our objective function of the variance evaluated using an estimate of the network model $\hat{\theta}$. Given a set of pilot points $\tau_1, \tau_2, \ldots, \tau_{n_0}$ (i.e. uniformly sampled on \mathcal{T}) we propose a Gaussian process prior satisfying

$$\mathcal{V}(\tau_{1:n_0}) \sim N(\mu_0(\tau_{1:n_0}, \Sigma_0(\tau_{1:n_0}, \tau_{1:n_0})))$$

where $\operatorname{Cov}[\mathcal{V}(\tau_i), \mathcal{V}(\tau_j)] = \Sigma_0(\tau_i, \tau_j)$ where Σ_0 is a positive semidefinite kernel function. As a default, we use the Gaussian kernel $\Sigma_0(x, x') = \alpha_0 \exp(-||x - x'||^2)$. We can then use this prior to define a posterior over remainder of the design space \mathcal{T}

$$\mathcal{V}(\tau)|\mathcal{V}(\tau_{1:n_0}) \sim N(\mu_n(\tau), \sigma_n^2(\tau))$$

$$\mu_n(\tau) = \Sigma_0(\tau, \tau_{1:n_0})\Sigma_0(\tau_{1:n_0}, \tau_{1:n_0})^{-1}(\mathcal{V}(\tau) - \mu_0(\tau_{1:n_0})) + \mu_0(\tau)$$

$$\sigma_n^2(\tau) = \Sigma_0(\tau, \tau) - \Sigma_0(\tau, \tau_{1:n_0})\Sigma_0(\tau_{1:n_0}, \tau_{1:n_0})^{-1}\Sigma_0(\tau_{1:n_0}, \tau).$$

From this posterior, we define an acquisition function $A(\tau)$. As a default, we choose the upper confidence bound (UCB) acquisition function $A(\tau) = \mu_n(\tau) - \kappa \sigma_n(\tau)$ for a chosen κ (where we set $\kappa = 2$). This method is implemented in the R package rBayesianOptimization, which uses GPfit (Yan, 2021; MacDonald et al., 2015). For a detailed review of Bayesian optimization techniques, refer to Frazier (2018). We evaluate the complete Bayesian optimization procedure in Algorithm 4, where we apply the procedure for N_0 iterations.

The quality of optimization over N_0 iterations depends on the smoothness of $\mathcal{V}(\tau)$. Since variance might diverge under some settings (e.g., as $\tau \to 0$), a simple alternative is to maximize $\exp(-\mathcal{V}(\tau))$ instead. The closeness of the maximizer after N_0 iterations hinges on the

Algorithm 4 Bayesian Optimization Procedure

- 1: Inputs: Graph model $\hat{\theta}$ and partial graph information G^* . Kernel function Σ_0 .
- 2: Sample $\tau_{1:n_0}$ uniformly from \mathcal{T} , as a pilot sample of the design points.
- 3: Update the posterior on $\mathcal{V}(\tau)$.
- 4: for $i \leftarrow n_0 + 1$ to $n_0 + N_0$ do
- 5: Update the posterior on $\mathcal{V}(\tau)|\mathcal{V}(\tau_{1:(i-1)})$.
- 6: Let τ_i be the minimizer of the acquisition function $A(\tau)$ (UCB).
- 7: Evaluate $\mathcal{V}(\tau_i)$ using Algorithm 3.
- 8: end for
- 9: Return the point $\tau_{1:(n_0+N_0)}$ with the smallest $\mathcal{V}(\tau)$

smoothness of $\exp(-\mathcal{V}(\tau))$, which we assume belongs to a reproducing kernel Hilbert space, \mathcal{H} , with a bounded kernel $\Sigma_0(x, x') \leq B$. This function's smoothness affects the approximation rate, detailed in Srinivas et al. (2009). For instance, with Gaussian kernel Σ_0 , the approximation error is $\exp(-\mathcal{V}(\tau^*)) \geq \frac{1}{N_0} \sum_{m=1}^{N_0} \exp(-\mathcal{V}(\tau_m)) + O_P(\frac{B\sqrt{\log(N_0)^{K+1}} + \log(N_0)^{K+1}}{\sqrt{N_0}}).$ Similar findings apply to Matern and linear kernels per Srinivas et al. (2009).

4.2 Designs for Optimal Seeding

Given a model of the potential outcomes, we may also leverage this model for optimal seeding, a task that is NP-hard (Kempe et al., 2003) in general. Many contagion models are exchangeable given an exposure, and with only block information available, then we can reduce our search space to that over block saturation. In our case, where exact network structures are unknown, we determine the optimal blocks for seeding. When $K \ll n$, this structure significantly reduces computational efforts, and we only need to decide how many seeds to allocate to each of the K clusters.

The model leveraged for the outcome $f_Y(V_i, S_i, \varepsilon_Y)$ could be a predefined model based on domain knowledge, such as complex contagion used by Beaman et al. (2021). In other scenarios, this might be estimated (e.g., simulation using $f_Y(V_i, S_i, \varepsilon_Y; \hat{\beta})$ in place of $f_Y(V_i, S_i, \varepsilon_Y)$). This is demonstrated in Algorithm 5 (line 5).

Algorithm 5 Optimal Seeding With Partial Network Data			
1: Inputs: Number of seeds N, Model estimate $\hat{\theta}$, number of graph draws L.			
2: Sample L draws from the graph model $\{\widehat{G}^{(l)}\}_{l=1}^{L} \sim \widehat{\theta} G^*$			
3: for $\tau \in \mathcal{T}$ do			
4: Sample <i>L</i> treatments $\{\mathbf{a}_l\}_{l=1}^L$ according to the block saturation levels τ .			
5: Compute the outcomes $Y_i^{(l,\mathbf{a}_l)}$ according to the outcome model $f_Y(V_i, S_i, \varepsilon_Y)$.			
6: Compute the average (and standard error) over draws of the network $\overline{Y}^{(\tau)} = \frac{1}{L} \sum_{l=1}^{L} Y_i^{(l,\mathbf{a}_l)}$			
7: end for			
8: Return saturation level τ with the largest value of $\overline{Y}^{(\tau)}$.			

When the total number of seeds (see Algorithm 5, line 3) is small, it is computationally feasible to implement exactly. Alternatively, we could use Bayesian optimization to control treatment saturation levels.

5 Data Analysis

In this section, we present three empirical examples to illustrate our framework's utility in estimating causal effects, designing experiments, and implementing seeding strategies. We adopt a semi-synthetic approach in our examples, where the outcomes are simulated based on processes derived from real networks. The networks analyzed pertain to observational and experimental studies focused on information diffusion in rural villages in India and Malawi, as discussed in Banerjee et al. (2013), Banerjee et al. (2019), and Beaman et al. (2021). These networks consist of 30-400 households per village. To ensure continuity across the examples, we generate ARD as the partial data type and model the networks using stochastic blockmodels for each case, however the use of other network generative

models and partial network datatype are applicable in these cases.

When covariates are available for all nodes, we use them to construct ARD. If covariates are missing, we apply the Leiden algorithm (Traag et al., 2019) in igraph (Csardi and Nepusz, 2006) to cluster the network and treat these clusters as traits. Table 1 details which datasets used actual traits versus clustering to manage trait numbers in our simulations. In Section 5.1, we use networks from Banerjee et al. (2013), which include various social

Network Dataset	Traits
Banerjee et al. (2013)	Leiden Cluster $K \in [4, 16]$
Banerjee et al. (2019)	Observed Traits (section A.4.5.2)
Beaman et al. (2021)	Leiden Cluster $K = 8$

Table 1: Summary of synthetic traits vs. real traits in the semi-synthetic simulations.

relations from 70 villages, each with 80 to 350 individuals per household. In Section 5.2, we use networks from Banerjee et al. (2019), consisting of 68 similar-sized villages, and repeat the simulations 500 times per village. In Section 5.3, we include networks from Beaman et al. (2021), excluding those with insufficient connections for diffusion. This leaves 114 villages with 30 to 350 households per village, and we repeat the simulations 2000 times per village.

5.1 Causal Effect Estimation

In our first example, the aim is to estimate the global average treatment. We consider the example from Ugander and Yin (2023) and generate a set of potential outcomes according to the following model

$$Y_i(\mathbf{0}) = \frac{d_i}{\overline{d}} \cdot (\alpha + bX_i + \sigma\epsilon_i), \quad Y_i(\mathbf{a}) = Y_i(\mathbf{0}) \cdot \left(1 + \delta a_i + \gamma \frac{\sum_{j \in [n]} G_{ij} a_j}{d_i}\right)$$

where $\epsilon_i \sim_{iid} N(0, 1)$ is some independent noise, and X_i is a covariate that varies throughout the network, d_i is the degree of individual i and \overline{d} is the average degree across the network. We set $\alpha = 1$, b = 1, $\delta = 1$, $\sigma = 0.5$ and $\gamma = -0.5$. The global average treatment effect in this model is $\frac{1}{n} \sum_{i=1}^{n} Y_i(\mathbf{0})(\delta + \gamma) = \Psi(\mathbf{a} = 1|G) - \Psi(\mathbf{a} = 0|G)$. The exposure is the individual treatment in conjunction with the average treatment of neighbors, and the graph confounder include the degree ratio and node level covariates

$$f_V(\mathbf{a};\varphi_i(G)) = \left(a_i, \frac{\sum_{j\in[n]} G_{ij}a_j}{\overline{d}}\right), \quad f_S(\mathbf{X};\vartheta_i(G)) = \left(\frac{d_i}{\overline{d}}, X_i\right).$$

We evaluate the effectiveness of graph cluster randomization by comparing a Horvitz-Thompson estimator Ugander et al. (2013) to a difference in means estimator under a cluster randomized design. In this design, half of the clusters receive no treatment (saturation of 0) and the other half receive full treatment (saturation of 1). We vary the number of clusters from 4 to 16 but display results only for 4 and 10 clusters in Figure 2 for clarity.



(a) Bias

(b) RMSE

Figure 2: Comparison of GATE estimators. ARD denotes our method using aggregated relational data. The "Full Network" method uses a regression approach with the full data available. DM is the difference in means and HT is the Horvitz-Thompson estimator.

Figure 2 shows that the full data regression model performs the best, as it leverages more information than the ARD approaches. However, the ARD version still effectively minimizes bias (Figure 2(a)) and RMSE (Figure 2(b)). In our simulations of dense graphs with few clusters, the Horvitz-Thompson Estimator faces challenges as the network grows—almost all nodes have at least one neighbor with a treatment different than their own. The difference in means estimator shows consistent bias, due to not using heterogeneous covariate information. While regression with complete data is most effective, using partial network data still yields comparably good results.

5.2 Experimental Design

We next highlight aspects of experimental design using an information diffusion example based on the hearing model referenced in section 2.1.2. At each time step the previously infected nodes are susceptible again the nodes infected in the last round will infect their neighbors with probability q_{t+1} . We repeat this for T = 3 rounds. Let N_i denote the total number of infections after the process. We then sample some binary response $P(Y_i = 1|N_i) = \text{logit}(\alpha_0 + \alpha_1 N_i)$ where α_0 and α_1 .

In this case, $V_i = \mathbb{E}[N_i|\mathbf{a}] = \sum_{t=0}^{3} \beta_t \mathbf{a}(G^t)_i$ where $\beta_t = \prod_{j=1}^{t} q_j$. We estimate the coefficients in each of these cases letting $V_i = \mathbb{E}[N_i|\mathbf{a}]$ be the exposure mapping. We then generate the outcomes according to the exposure received

$$\mathbb{E}[Y_i|S_i, V_i] = \Lambda(\alpha_0 + \alpha_1(\sum_{t=0}^3 \beta_t(G^t)_i \mathbf{a}))$$

where $\Lambda(\cdot)$ is the logistic function. For our experiments, we set $\beta = (0, 0.5, 0.05, 0.005)$.

In the dataset, seeds are assigned uniformly with either 3 or 5 seeds per network. Following our procedure in section 4, we compute the optimal seed allocations, ensuring no cluster receives more seeds than available in the actual experiment (either 3 or 5). In practice our Bayesian optimization procedure starts by randomly sampling the target space 20 times, followed by 20 iterations to refine saturation. We then compare the estimates for α_1 and all model parameters as shown in Figure 3. The results indicate that a more strategically designed experiment generally yields more significant gains than directly using the graph parameters. On average, using optimized designs rather than uniform random designs when collecting network data significantly reduced RMSE. Specifically, for estimating α_1 , the optimized design decreased RMSE by 38% (±12%) compared to 11% (±2%) with complete data (where the brackets refer to the 95% confidence interval of the mean estimate across simulations). For all parameters, the optimized design resulted in a 45% (±10%) reduction in RMSE, versus an 18% (±2%) reduction with complete data.



(a) Estimation of α_1 .

(b) Estimation of all model parameters.

Figure 3: Estimation of parameter α_1 and all model parameters β using the naive and optimized seeding. We observe that the potential gain found using a more efficient design is much greater than simply collecting complete network data.

5.3 Optimal Seeding

We apply our methodology to the seeding problem described in Beaman et al. (2021), where the diffusion of pit-planting technology among Malawian farmers follows a complex contagion process. The outcome model is defined as $Y_i = f_Y(S_i, V_i, \epsilon_Y)$, with individuals having a threshold $\varsigma_i \sim N_{[0,\infty)}(\lambda, 0.1)$ for spreading infection based on neighbor infections from the previous time (where $N_{(a,b)}(\mu, \sigma)$ refers to the μ, σ normal distribution truncated on the interval (a, b)). This process is simulated over three time periods to align with their experimental design, setting $\lambda = 2$ and repeating 2000 times for K = 8 clusters to determine optimal seeding groups.

We explore two seeding strategies: randomly assigning seeds to the top two members of optimal clusters, and seeding the nodes with the highest degrees within these clusters. We compare these strategies to common degree targeting, noting that our max degree method typically yields the highest adoption rates, especially in larger, sparser villages, as illustrated in Figure 4. However, in very small or dense networks, the performance differences between strategies are negligible. Across all graphs we find the optimal seeding strategy to increase adoption by 1.50 (± 0.16) times relative to degree seeding, while the optimal blocks was 1.13 (± 0.12) times and optimal degree within blocks increased adoption by 1.28 $\pm (0.13)$ times.

6 Conclusions

We introduce a framework that identifies causal effects under interference using a structural causal model, facilitating inference with partial network data. The framework is general and can be applied using broad class of outcome models and graph models. Our outcome modelling approach leveraging node-level heterogeneity and exposure mappings allow for



(a) Adoption ratio as a function of edge density. (b) Adoption ratio as a function of village size.
Figure 4: Comparison of different seeding methods under complex contagion. Model-based targeting of optimal blocks generally outperforms degree seeding, especially when targeting the highest degree nodes within those blocks.

the estimation of all causal effects, rather that other methods which tend to focus on a single causal effect like the GATE. Demonstrations through semi-synthetic problems highlight its effectiveness, matching or surpassing fully observed data methods in certain scenarios.

Our method highlights that directly modeling interference mechanisms offers several advantages, including leveraging transportability of outcome models for seeding and inference for experimental designs when estimating effects under interference.

Future studies might consider semiparametric approaches to estimation with partial data like those in Auerbach (2022). Additional structured assumptions on potential outcomes as suggested in Belloni et al. (2022) could also be explored. Currently, our focus has been on analyzing problems at a single time point. However, future research could extend to designing experiments with panel data and staggered rollouts. It would also be worth-while to develop classes of outcome models that more explicitly incorporate this temporal
structure.

References

- Acemoglu, D., A. Ozdaglar, and A. Tahbaz-Salehi (2015). Systemic risk and stability in financial networks. The American Economic Review 105(2), 564–608.
- Airoldi, E. M., T. B. Costa, and S. H. Chan (2013). Stochastic blockmodel approximation of a graphon: Theory and consistent estimation. <u>Advances in Neural Information Processing</u> Systems 26.
- Alidaee, H., E. Auerbach, and M. P. Leung (2020). Recovering network structure from aggregated relational data using penalized regression. arXiv preprint arXiv:2001.06052.
- Almquist, Z. W. (2012). Random errors in egocentric networks. <u>Social networks</u> <u>34</u>(4), 493–505.
- Ambrus, A., M. Mobius, and A. Szeidl (2014). Consumption risk-sharing in social networks. American Economic Review 104(1), 149–182.
- Anderson, J. R. and G. Feder (2007). Agricultural extension. <u>Handbook of agricultural</u> economics 3, 2343–2378.
- Andrews, D. W. (1987). Consistency in nonlinear econometric models: A generic uniform law of large numbers. Econometrica: Journal of the Econometric Society, 1465–1471.
- Aral, S., L. Muchnik, and A. Sundararajan (2009). Distinguishing influence-based contagion from homophily-driven diffusion in dynamic networks. <u>Proceedings of the National</u> Academy of Sciences 106(51), 21544–21549.

- Aronow, P. M. and C. Samii (2017). Estimating average causal effects under general interference, with application to a social network experiment. <u>Annals of Applied Statistics 11(4)</u>, 1912–1947.
- Athey, S., D. Eckles, and G. W. Imbens (2018). Exact p-values for network interference. Journal of the American Statistical Association 113(521), 230–240.
- Auerbach, E. (2022). Identification and estimation of a partially linear regression model using network data. Econometrica 90(1), 347–365.
- Auerbach, E. and M. Tabord-Meehan (2021). The local approach to causal inference under network interference. arXiv preprint arXiv:2105.03810.
- Banerjee, A., E. Breza, A. G. Chandrasekhar, and B. Golub (2018). When less is more: Experimental evidence on information delivery during india's demonetization. Technical report, National Bureau of Economic Research.
- Banerjee, A., A. G. Chandrasekhar, E. Duflo, and M. O. Jackson (2013). The diffusion of microfinance. Science 341(6144), 1236498.
- Banerjee, A., A. G. Chandrasekhar, E. Duflo, and M. O. Jackson (2019). Using gossips to spread information: Theory and evidence from two randomized controlled trials. <u>The</u> Review of Economic Studies.
- Baraff, A. J., T. H. McCormick, and A. E. Raftery (2016). Estimating uncertainty in respondent-driven sampling using a tree bootstrap method. <u>Proceedings of the National</u> Academy of Sciences 113(51), 14668–14673.
- Barrat, A., M. Barthelemy, and A. Vespignani (2008). <u>Dynamical processes on complex</u> networks. Cambridge university press.

- Beaman, L., A. BenYishay, J. Magruder, and A. M. Mobarak (2021). Can network theorybased targeting increase technology adoption? <u>American Economic Review</u> <u>111</u>(6), 1918–43.
- Belloni, A., F. Fang, and A. Volfovsky (2022). Neighborhood adaptive estimators for causal inference under network interference. arXiv preprint arXiv:2212.03683.
- Bond, R. M. and B. J. Bushman (2017). The contagious spread of violence among us adolescents through social networks. American journal of public health 107(2), 288–294.
- Boucher, V. and A. Houndetoungan (2020). Estimating peer effects using partial network data. Centre de recherche sur les risques les enjeux économiques et les politiques
- Bradley, R. C. (2005). Basic properties of strong mixing conditions. a survey and some open questions.
- Bramoullé, Y., H. Djebbari, and B. Fortin (2009). Identification of peer effects through social networks. Journal of econometrics 150(1), 41–55.
- Brennan, J., V. Mirrokni, and J. Pouget-Abadie (2022). Cluster randomized designs for one-sided bipartite experiments. arXiv preprint arXiv:2210.16415.
- Breza, E., A. G. Chandrasekhar, S. Lubold, T. H. McCormick, and M. Pan (2023). Consistently estimating network statistics using aggregated relational data. <u>Proceedings of</u> the National Academy of Sciences 120(21), e2207185120.
- Breza, E., A. G. Chandrasekhar, T. H. McCormick, and M. Pan (2020). Using aggregated relational data to feasibly identify network structure without network data. <u>American</u> Economic Review.

- Cencetti, G., D. A. Contreras, M. Mancastroppa, and A. Barrat (2023). Distinguishing simple and complex contagion processes on networks. <u>Physical Review Letters</u> <u>130</u>(24), 247401.
- Centola, D. and M. Macy (2007). Complex contagions and the weakness of long ties. American journal of Sociology 113(3), 702–734.
- Chandrasekhar, A. and R. Lewis (2011). Econometrics of sampled networks. <u>Unpublished</u> manuscript, MIT.[422].
- Chandrasekhar, A. G., M. O. Jackson, T. H. McCormick, and V. Thiyageswaran (2023). General covariance-based conditions for central limit theorems with dependent triangular arrays. <u>arXiv preprint arXiv:2308.12506</u>.
- Chatterjee, S. and P. Diaconis (2011). Estimating and understanding exponential random graph models. Arxiv preprint arXiv:1102.2650.
- Chatterjee, S., P. Diaconis, and A. Sly (2011). Random graphs with a given degree sequence. The Annals of Applied Probability, 1400–1435.
- Chierichetti, F., D. Liben-Nowell, and J. Kleinberg (2011). Reconstructing patterns of information diffusion from incomplete observations. <u>Advances in neural information</u> processing systems 24.
- Cortez, M., M. Eichhorn, and C. Yu (2022). Staggered rollout designs enable causal inference under interference without network knowledge. In <u>Advances in Neural Information</u> Processing Systems.
- Csardi, G. and T. Nepusz (2006). The igraph software package for complex network research. InterJournal, Complex Systems 1695(5), 1–9.

- De Giorgi, G., M. Pellizzari, and S. Redaelli (2010). Identification of social interactions through partially overlapping peer groups. <u>American Economic Journal: Applied</u> Economics 2(2), 241–275.
- De Paula, A. (2017). Econometrics of network models. In <u>Advances in economics and</u> <u>econometrics: Theory and applications, eleventh world congress</u>, pp. 268–323. Cambridge University Press Cambridge.
- Dempster, A. P., N. M. Laird, and D. B. Rubin (1977). Maximum likelihood from incomplete data via the em algorithm. <u>Journal of the royal statistical society: series B</u> (methodological) <u>39(1)</u>, 1–22.
- Epple, D. and R. E. Romano (2011). Peer effects in education: A survey of the theory and evidence. In Handbook of social economics, Volume 1, pp. 1053–1163. Elsevier.
- Feehan, D. M. and M. J. Salganik (2016). Generalizing the network scale-up method: a new estimator for the size of hidden populations. Sociological methodology 46(1), 153–186.
- Feehan, D. M., A. Umubyeyi, M. Mahy, W. Hladik, and M. J. Salganik (2016). Quantity versus quality: A survey experiment to improve the network scale-up method. <u>American</u> journal of epidemiology 183(8), 747–757.
- Frazier, P. I. (2018). A tutorial on bayesian optimization. arXiv preprint arXiv:1807.02811.
- Freeman, L. C. (1982). Centered graphs and the structure of ego networks. <u>Mathematical</u> Social Sciences 3(3), 291–304.
- Fu, A., B. Narasimhan, and S. Boyd (2020, 11). CVXR: An R package for disciplined convex optimization. Journal of Statistical Software 94(14), 1–34.

- Gao, C., Y. Lu, and H. H. Zhou (2015). Rate-optimal graphon estimation. <u>The Annals of</u> Statistics, 2624–2652.
- Giles, P. (1977). The mathematical theory of infectious diseases and its applications. Journal of the Operational Research Society 28(2), 479–480.
- Goel, S. and M. J. Salganik (2009). Respondent-driven sampling as markov chain monte carlo. Statistics in medicine 28(17), 2202–2229.
- Goel, S. and M. J. Salganik (2010). Assessing respondent-driven sampling. Proceedings of the National Academy of Sciences 107(15), 6743–6747.
- Goldsmith-Pinkham, P. and G. Imbens (2013). Social networks and the identification of peer effects. Journal of Business and Economic Statistics 31:3, 253–264.
- Graham, B. S. (2020). Network data. In <u>Handbook of econometrics</u>, Volume 7, pp. 111–218. Elsevier.
- Green, A., T. McCormick, and A. Raftery (2020). Consistency for the tree bootstrap in respondent-driven sampling. Biometrika 107(2), 497–504.
- Ha-Thuc, V., A. Dutta, R. Mao, M. Wood, and Y. Liu (2020). A counterfactual framework for seller-side a/b testing on marketplaces. In <u>Proceedings of the 43rd International</u> <u>ACM SIGIR Conference on Research and Development in Information Retrieval</u>, pp. 2288–2296.
- Handcock, M. S., A. E. Raftery, and J. M. Tantrum (2007). Model-based clustering for social networks. <u>Journal of the Royal Statistical Society Series A: Statistics in</u> Society 170(2), 301–354.

- Hardy, M., R. M. Heath, W. Lee, and T. H. McCormick (2019, 3). Estimating spillovers using imprecisely measured networks. arXiv.
- He, X. and K. Song (2023, 12). Measuring Diffusion Over a Large Network. <u>The Review</u> of Economic Studies, rdad115.
- Heckathorn, D. D. (1997). Respondent-driven sampling: a new approach to the study of hidden populations. Social problems 44(2), 174–199.
- Hoff, P. D., A. E. Raftery, and M. S. Handcock (2002, 12). Latent Space Approaches to Social Network Analysis. <u>https://doi.org/10.1198/016214502388618906</u> <u>97(460)</u>, 1090– 1098.
- Huang, G. C., J. B. Unger, D. Soto, K. Fujimoto, M. A. Pentz, M. Jordan-Marsh, and T. W. Valente (2014). Peer influences: the impact of online and offline friendship networks on adolescent smoking and alcohol use. Journal of Adolescent Health 54(5), 508–514.
- Hudgens, M. G. and M. E. Halloran (2008). Toward causal inference with interference. Journal of the American Statistical Association 103(482), 832–842.
- Imai, K., Z. Jiang, and A. Malani (2021). Causal inference with interference and noncompliance in two-stage randomized experiments. <u>Journal of the American Statistical</u> Association 116(534), 632–644.
- Jackson, M. O. et al. (2008). <u>Social and economic networks</u>, Volume 3. Princeton university press Princeton.
- Jackson, M. O. and L. Yariv (2006). Diffusion on social networks. <u>Economie</u> publique/Public economics (16).

- Jing, L., C. Qu, H. Yu, T. Wang, and Y. Cui (2014). Estimating the sizes of populations at high risk for HIV: a comparison study. PloS ONE 9(4), e95601.
- Johari, R., H. Li, I. Liskovich, and G. Y. Weintraub (2022). Experimental design in twosided platforms: An analysis of bias. Management Science 68(10), 7069–7089.
- Kadushin, C., P. D. Killworth, H. R. Bernard, and A. A. Beveridge (2006). Scale-up methods as applied to estimates of heroin use. Journal of Drug Issues 36(2), 417–440.
- Keeling, M. J. and P. Rohani (2008). <u>Modeling Infectious Diseases in Humans and Animals</u>. Princeton University Press.
- Kempe, D., J. Kleinberg, and É. Tardos (2003). Maximizing the spread of influence through a social network. In <u>Proceedings of the ninth ACM SIGKDD international conference</u> on Knowledge discovery and data mining, pp. 137–146.
- Kermack, W. O. and A. G. McKendrick (1927). A contribution to the mathematical theory of epidemics. <u>Proceedings of the royal society of london</u>. Series A, Containing papers of a mathematical and physical character 115(772), 700–721.
- Killworth, P. D., E. C. Johnsen, C. McCarty, G. A. Shelley, and H. R. Bernard (1998). A social network approach to estimating seroprevalence in the United States. <u>Social</u> Networks 20(1), 23–50.
- Killworth, P. D., C. McCarty, H. R. Bernard, G. A. Shelley, and E. C. Johnsen (1998). Estimation of seroprevalence, rape, and homelessness in the United States using a social network approach. Evaluation Review 22(2), 289–308.
- Lovász, L. and B. Szegedy (2006). Limits of dense graph sequences. <u>Journal of</u> Combinatorial Theory, Series B 96(6), 933–957.

- Lubold, S., A. G. Chandrasekhar, and T. H. McCormick (2023). Identifying the latent space geometry of network models through analysis of curvature. <u>Journal of the Royal</u> Statistical Society Series B: Statistical Methodology 85(2), 240–292.
- MacDonald, B., P. Ranjan, and H. Chipman (2015). Gpfit: An r package for fitting a gaussian process model to deterministic simulator outputs. Journal of Statistical Software <u>64</u>, 1–23.
- Malani, A., P. Holtzman, K. Imai, C. Kinnan, M. Miller, S. Swaminathan, A. Voena,B. Woda, and G. Conti (2021). Effect of health insurance in india: a randomized controlled trial. Technical report, National Bureau of Economic Research.
- Manski, C. F. (1993). Identification of endogenous social effects: The reflection problem. The review of economic studies 60(3), 531–542.

Manski, C. F. (2009). Identification for prediction and decision. Harvard University Press.

- McCormick, T. H. (2020). The network scale-up method. <u>The Oxford Handbook of Social</u> Networks, 153.
- Newey, W. K. and D. McFadden (1994). Large sample estimation and hypothesis testing. Handbook of econometrics 4, 2111–2245.
- Ogburn, E. L., O. Sofrygin, I. Diaz, and M. J. Van der Laan (2022). Causal inference for social network data. Journal of the American Statistical Association, 1–15.
- Orbanz, P. and D. M. Roy (2015). Bayesian models of graphs, arrays and other exchangeable random structures. <u>IEEE Transactions on Pattern Analysis and Machine</u> Intelligence 37(2), 437–461.

Pastor-Satorras, R., C. Castellano, P. Van Mieghem, and A. Vespignani (2015). Epidemic processes in complex networks. Reviews of modern physics 87(3), 925.

Pearl, J. (2009). Causality. Cambridge university press.

- Potter, G. E., M. S. Handcock, I. M. Longini Jr, and M. E. Halloran (2011). Estimating within-household contact networks from egocentric data. <u>The annals of applied</u> statistics 5(3), 1816.
- Pouget-Abadie, J., V. Mirrokni, D. C. Parkes, and E. M. Airoldi (2018). Optimizing cluster-based randomized experiments under monotonicity. In <u>Proceedings of the 24th</u> <u>ACM SIGKDD International Conference on Knowledge Discovery & Data Mining</u>, pp. 2090–2099.
- Pouget-Abadie, J., G. Saint-Jacques, M. Saveski, W. Duan, S. Ghosh, Y. Xu, and E. M. Airoldi (2019). Testing for arbitrary interference on experimentation platforms. Biometrika 106(4), 929–940.
- Roch, S. and K. Rohe (2018). Generalized least squares can overcome the critical threshold in respondent-driven sampling. <u>Proceedings of the National Academy of Sciences 115</u>(41), 10299–10304.
- Romero, D. M., B. Meeder, and J. Kleinberg (2011). Differences in the mechanics of information diffusion across topics: idioms, political hashtags, and complex contagion on twitter. In <u>Proceedings of the 20th international conference on World wide web</u>, pp. 695–704.
- Saveski, M., J. Pouget-Abadie, G. Saint-Jacques, W. Duan, S. Ghosh, Y. Xu, and E. M. Airoldi (2017). Detecting network effects: Randomizing over randomized experiments. In

Proceedings of the 23rd ACM SIGKDD international conference on knowledge discovery and data mining, pp. 1027–1035.

- Scutelniciuc, O. (2012). Network scale-up method experiences: Republic of kazakhstan. <u>Consultation on estimating population sizes through household surveys: Successes and</u> challenges (New York, NY).
- Shimodaira, H. (2000). Improving predictive inference under covariate shift by weighting the log-likelihood function. Journal of statistical planning and inference 90(2), 227–244.
- Srinivas, N., A. Krause, S. M. Kakade, and M. Seeger (2009). Gaussian process optimization in the bandit setting: No regret and experimental design. arXiv preprint arXiv:0912.3995.
- Tchetgen, E. J. T. and T. J. VanderWeele (2012). On causal inference in the presence of interference. Statistical methods in medical research 21(1), 55–75.
- Tortú, C., I. Crimaldi, F. Mealli, and L. Forastiere (2021). Causal effects with hidden treatment diffusion on observed or partially observed networks. <u>arXiv preprint</u> arXiv:2109.07502.
- Traag, V. A., L. Waltman, and N. J. Van Eck (2019). From louvain to leiden: guaranteeing well-connected communities. Scientific reports 9(1), 5233.
- Tran, V. C. and T. P. T. Vo (2021). Estimation of dense stochastic block models visited by random walks. Electronic Journal of Statistics 15(2), 5855–5887.
- Tsiatis, A. A. (2006). Semiparametric theory and missing data, Volume 4. Springer.
- Ugander, J., B. Karrer, L. Backstrom, and J. Kleinberg (2013, 8). Graph cluster randomization: Network exposure to multiple universes. Proceedings of the ACM SIGKDD

International Conference on Knowledge Discovery and Data Mining Part F128815, 329–337.

Ugander, J. and H. Yin (2023). Randomized graph cluster randomization. <u>Journal of</u> Causal Inference 11(1), 20220014.

Vaart, A. W. v. d. (1998, 10). Asymptotic Statistics. Cambridge University Press.

- van der Laan, M. J. (2012). Causal inference for networks.
- Viviano, D. (2020). Experimental design under network interference. <u>arXiv preprint</u> arXiv:2003.08421.
- Wilkins-Reeves, S., X. Chen, Q. Ma, C. Agarwal, and A. Hofleitner (2024). Multiply robust estimation for local distribution shifts with multiple domains. <u>arXiv preprint</u> arXiv:2402.14145.
- Wilkins-Reeves, S. and T. McCormick (2022). Asymptotically normal estimation of local latent network curvature. arXiv preprint arXiv:2211.11673.
- Wu, C. J. (1983). On the convergence properties of the em algorithm. <u>The Annals of</u> statistics, 95–103.
- Yan, Y. (2021). rBayesianOptimization.
- Yu, C. L., E. M. Airoldi, C. Borgs, and J. T. Chayes (2022). Estimating the total treatment effect in randomized experiments with unknown network structure. <u>Proceedings of the</u> National Academy of Sciences 119(44), e2208975119.

SUPPLEMENTARY MATERIAL

A.1 Comparing Frameworks of Interference

We contrast the approaches of a fixed outcome approach as in Aronow and Samii (2017) to a structural causal model approach. In the former approach, each individual has a distinct outcome under an exposure v, $Y_i(v)$. Though such an approach is robust for learning parameters such as average treatment effects $\frac{1}{n} \sum_{i=1}^{n} Y_i(v)$, the information in an individual *i*'s potential outcome is completely distinct from individual *j*. This important details has important downstream implications.

Consider the simple contagion model from the example in section 2.1.1 which takes place in a single time period (T = 1). Consider the nodes i, j in Figure 5 with seeded nodes in blue. Suppose that at time T = 1, that each neighbour of a treated node is infected with probability q. Since each one has only a single treated neighbor the distribution of the infection probability $P(Y_i = 1 | \mathbf{a}, G)$ i and j are equivalent as their exposures are identical (i.e. they are each connected to a single seed node). However, in the finite sample framework the potential outcomes of any two nodes with a single treated neighbor can be arbitrarily different $(Y_i(v) \neq Y_j(v))$.

This nonparametric structure imposed on the potential outcomes later imposes restrictions on the degree of influence of others a node can have for estimation, thereby limiting this framework to examples with local dependencies (a phenomena also seen in Ogburn et al. (2022)).



Figure 5: Equivalence of distribution of potential outcomes of nodes i and j are equivalent under this given treatment assignment as all of the rooted networks are equivalent.

A.1.1 Why not IPW estimators?

In many nonparametric approaches to estimating causal quantities under interference, inverse probability weighted (IPW) estimates can be developed given a randomization scheme, i.e. distribution of the assignments $P(\mathbf{a})$ (Aronow and Samii, 2017). This is useful as it can be used to develop estimators for causal effects any exchangeability assumptions on the potential outcomes. However when V_i is not observed directly, we must leverage additional structure in order to estimate any causal effects.

Our objective is to understand the model's structure and often apply it to tasks such as seeding. Thus, we rely on a correct model specification. The challenge with developing an IPW estimator arises when exposure is not observed. In such cases, it becomes impossible to determine which potential outcome was observed, violating the causal consistency assumption. Specifically, we don't know which potential outcome Y_i represents (i.e., which exposure $v, Y_i = Y_i(v)$).

A.2 Additional Methodological Details

In this section we discuss extensions to several aspects of the paper with respect to the paper. Before proceeding we also introduce the full statement of the generalized central limit theorem result which we use to derive our asymptotic results.

A.2.1 A central limit theorem for dependent data.

Although network models do not neatly fit into conventional time series or spatial dependency categories, we provide a general framework by satisfying the necessary conditions through common dependence assumptions such as *M*-dependence. This includes scenarios characterized by α -, ϕ -, or ρ -mixing (Bradley, 2005). Our approach begins by defining affinity sets, (sets for which there is high correlation with an outcome) that form the foundational framework for applying the CLT, setting the stage for demonstrating its relevance and utility in analyzing network data.

Definition A.2.1 (Affinity sets). Denote a triangular array of mean 0 random vectors $W_{1:n}^{(n)}$ with dimension p. Let $\mathcal{A}_{(i,d)}^{(n)}$ denote an affinity set which contains all of the variables in the triangular array which are highly correlated with $W_{i,d}^{(n)}$, the dth dimension of the ith random variable.

The affinity sets can be used to construct a matrix which contains the bulk of the covariance across observations and dimensions. The regularity conditions can be understood as control of the covariance within affinity sets (A.2.1), control of the covariance across affinity sets (A.2.2) and control of the covariance outside of the affinity sets (A.2.3). We collectively refer to these as the affinity set conditions. The affinity sets can be used to construct a covariance matrix $\Gamma_{n,dd'} = \sum_{i=1}^{n} \sum_{(j,d') \in \mathcal{A}_{(i,d)}^{(n)}} \operatorname{cov}(W_{i,d}^{(n)}, W_{j,d'}^{(n)}).$

$$\sum_{(i,d):(j,d'),(k,d'')} \mathbb{E}[W_{i,d}|W_{j,d'}W_{k,d''}] = o(||\Gamma_n||_F^{3/2}),$$
(A.2.1)

$$\sum_{(i,d),(j,d');(k,d''),(l,\widehat{d})} \operatorname{cov}(W_{i,d}W_{k,d''}, W_{j,d'}W_{l,\widehat{d}}) = o(||\Gamma_n||_F^2),$$
(A.2.2)

$$\sum_{(i,d)} \mathbb{E}[|\mathbf{W}_{-i,d}\mathbb{E}[W_{i,d}\mathbf{W}_{-i,d}]|] = o(||\Gamma_n||_F).$$
(A.2.3)

Theorem A.2.2 (Theorem 1 from Chandrasekhar et al. (2023)). Denote a mean 0 triangular array of random vectors $W_{1:n}^{(n)}$. If a collection of affinity sets $\mathcal{A}_{(i,d)}^{(n)}$ satisfy the conditions of equations (A.2.1), (A.2.2) and (A.2.3). Then

$$\Gamma_n^{-1/2} S_n \to_d N(0, I_p)$$

The authors illustrate several examples under which these conditions are sufficient for the this central limit theorem to hold. This theorem will be useful for proving our asymptotic results.

A.2.2 Inference for the OLS Estimator

Here we first give the full theorem and regularity conditions with respect to the linear model.

Theorem A.2.3. Let $\widetilde{H}_i(\theta) = \mathbb{E}[\widetilde{h}(S_i(G), V_i(G))|\mathbf{a}, \mathbf{X}, G^*; \theta]$. The OLS estimator uses the model averaged coefficients $\widetilde{H}_i(\theta)$ in place of the true unobserved coefficients \widetilde{h}_i . Let $\mathsf{H}_n(\theta) = \frac{1}{n} \sum_{i=1}^n \widetilde{H}_i(\theta) \widetilde{H}_i^T(\theta)$. Given an estimate of the model parameters $\widehat{\theta}$, we define the

$$\widehat{\beta}_{lm} = \mathsf{H}_n^{-1}(\widehat{\theta}) \frac{1}{n} \sum_{i=1}^n \widetilde{H}_i(\widehat{\theta}) Y_i$$

Let $u_i = (\tilde{h}(S_i(G), V_i(G)) - \tilde{H}_i(\theta_0))\beta_0 + \epsilon_i$. Suppose the following conditions hold for all n. Model Regularity conditions

- E1. $\hat{\theta}$ is a s(n)-consistent estimate of the graph parameters $||\hat{\theta} \theta_0|| = o_P(s(n))$
- E2. $|\mathsf{H}_n(\theta) \mathsf{H}_n(\theta')| \le b_n(\mathbf{Z}) ||\theta \theta'||$ where $b_n(\mathbf{Z}) = O_P(1)$ (that is, $b_n(\mathbf{Z})$ is stochastically bounded).
- *E3.* $\max_i ||\widetilde{H}_i(\theta) \widetilde{H}_i(\theta')|| \le b_n(\mathbf{Z})||\theta \theta'||$
- *E4.* $||H_i(\theta)|| \leq M < \infty$
- E5. $\left|\frac{1}{n}\sum_{i=1}^{n}|u_i|-\frac{1}{n}\sum_{i=1}^{n}\mathbb{E}[|u_i|]\right|=o_P(1)$

Lastly, let Γ_n denote a matrix that satisfies the following central limit theorem for the estimating function

Central Limit Theorem

F1. For the array of random variables $\mathcal{G}_i = \frac{1}{n} H_i(\theta_0) u_i$, there exists a set of affinity sets $\mathcal{A}_{(i,d)}^n$ such that (A.2.1), (A.2.2) and (A.2.3) are satisfied with a corresponding matrix Γ_n , where $\sqrt{\lambda_{min}(\Gamma_n)} = r(n)$.

Then if r(n) = o(s(n))

$$\Gamma_n^{-1/2} \mathsf{H}_n(\widehat{\theta})(\widehat{\beta}_{lm} - \beta_0) \to_d N(0, I_p)$$

A.2.3 Estimation of Network Models

We next discuss the estimation of generative models of network formation using several datatypes. We summarize the information for using ARD in Table A.2.3 as discuss similar rates for other datatypes.

A.2.3.1 Estimation of the Stochastic Blockmodel Using Sampled Data

We illustrate that it is possible to estimate the stochastic blockmodel using a diverse set of partial and sampled network data types. In each case, $\mathbf{P}_{kk'}$ refer to the cross-block

Network Model	Norm	ARD Rate
SBM	$\sum_{k,k'} \left \widehat{P}_{kk'} - P_{kk'} \right $	$\widetilde{O}_P(K/n)$
Latent Space	$\sup_{i\in\{1,2,\dots,n\}} \widehat{\theta}_i-\theta_i $	$O_P(\sqrt{\log(n)/n})$
Beta Model	$\sup_{i\in\{1,2,\dots,n\}} \widehat{\theta}_i-\theta_i $	$O_P(\sqrt{\log(n)/n})$
Low-Rank Graphon	$\tfrac{1}{n^2} \widehat{\eta}-\eta_0 _2$	$\widetilde{O}_P(1/T)$

Table 2: Summary of estimation rates with respect to model classes. The norms used for the latent space and beta models are with respect to their individual parameters θ_i . We let $\eta_{0,ij} = P(G_{ij} = 1|\theta_0)$ denote the probability of two nodes connecting in the graphon model. Rates for the latent space and beta models are derived in Breza et al. (2023) and the low-rank graphon in Alidaee et al. (2020).

probabilities, while $k_i \in \{1, 2, ..., K\}$ denote the node memberships. We consider *partial* network data to be any subset of the network data which can be used to generate an estimate of the generative model $\hat{\theta}$.

Example A.2.1 (Induced subgraph). We sample $m \le n$ of nodes in the graph randomly, with at least one node from each of the K communities. Let G' be the sub-graph induced by these m nodes. Let N'_k denote the set of sampled nodes in community k, assumed to be positive for each k. Let

$$\widehat{\mathbf{P}}_{kk'} = \frac{1}{|N'_k| |N'_{k'}|} \sum_{i \in N'_k} \sum_{j \in G'_{k'}} G'_{ij}$$

Example A.2.2 (Edges missing). Suppose that edges are missing according to some distribution. Let G' be the observed graph, and suppose that $P(G'_{ij} = 1 | X_{ij} = x)$ is the probability of observing the edge G'_{ij} , given dyad-level covariates X and the edge G_{ij} . Suppose that we have a consistent estimator of this conditional response. Then,

$$\widehat{\mathbf{P}}_{kk'} = \frac{1}{|N'_k| |N'_{k'}|} \sum_{i \in N'_k} \sum_{j \in G'_{k'}} \frac{G'_{ij}}{\widehat{P}(G'_{ij} = 1|X_{ij})}$$

Lemma A.2.4 (Rates for induced subgraph and Edges Missing). Consider an estimate for a stochastic blockmodel cross probabilities based on either the induced subgraph or the edges missing example of $m \leq n$. Let $m_k = |N_k| = \rho_k m$ for some $\rho_k \in (0, 1)$. Then with probability at least $1 - \delta$

$$|\widehat{\mathbf{P}}_{kk'} - \mathbf{P}_{kk'}| \le \frac{1}{\rho_k \rho_{k'} m} \sqrt{\frac{\log(2/\delta)}{2}}$$
(A.2.4)

Further, suppose that $\sup_{x} |\widehat{P}(G_{ij} = 1|X_{ij} = x) - P(G_{ij} = 1|X_{ij} = x)| = o_P(m^{-1})$ with $P(G_{ij} = 1|X_{ij} = x) \ge \lambda > 0$. Then for large enough m, equation A.2.4 holds for the missing edges example as well.

Lastly, we discuss respondent driven sampling. In this setting, community membership can be defined based on a partition of the covariates, thus allowing for an observable trait in the graph, a similar strategy is adopted by Roch and Rohe (2018).

Example A.2.3 (Respondent driven sampling). Let $i \in \{1, 2, ..., m\}$ denote the indices of a sample of individuals obtained through respondent driven sampling. An initial number of individuals are recruited as seeds, and subsequent individuals are recruited via referrals from the others in a population. Tran and Vo (2021) develop a consistent estimator for the model parameters of the stochastic blockmodel.

Let \widetilde{G}_m be the subgraph of G_n sampled from a set of nodes $\{1, 2, ..., m\}$. Let M_k denote the number of individuals in the subsample of type k and let $M_{kk'}^{\leftrightarrow}$ denote the number of connected individuals in the subgraph \widetilde{G}_m .

The cross-type probabilities can be estimated as follows:

$$\widehat{\mathbf{P}}_{kk'} = \begin{cases} \frac{M_{kk'}^{\leftrightarrow}}{M_k M_{k'}} & When \ k \neq k' \\ \\ \frac{M_{kk}^{\leftrightarrow}}{M_k (M_k - 1)} & otherwise \end{cases}$$

Tran and Vo (2021) illustrate the consistency of these parameters (Theorem 4.2 in their paper), in particular $|\widehat{\mathbf{P}}_{kk'} - \mathbf{P}_{kk'}| = O_P(m^{-1})$

A.2.3.2 Estimation of Other Network Models

Though we emphasise the estimation of the stochastic blockmodel, there are several other methods available for estimation of the network formation model. These include the beta model of Chatterjee and Diaconis (2011), in which the graph generation model consists of two model parameters ν_i , ν_i possibly altered through some additional dyadic covariates X_{ij}^*

$$P(G_{ij} = 1|\theta_0) = \widetilde{f}(\nu_i + \nu_j + \beta^T X_{ij}^*)$$

where \tilde{f} is a link function. Alternatively one can consider the latent space model of Hoff et al. (2002) which include latent positions on some unobserved manifold \mathcal{M}^p .

$$P(G_{ij} = 1|\theta_0) = f(\nu_i + \nu_j + d_{\mathcal{M}^p}(Z_i, Z_j))$$

In each of these cases Breza et al. (2023) illustrate consistent estimation rates in the $||\hat{\theta} - \theta_0||_{\infty} = \mathcal{O}_P\left(\sqrt{\frac{\log(n)}{n}}\right)$ with the use of aggregated relational data. Since this represents the coarsest datatype we expect similar rates to hold for subgraph sampling and respondent driven sampling. Though this rate is too slow for the to ignore the effect of the estimation of the graph model, in examples where one expect a high level of correlation among the outcomes it can be practical to use these methods.

A.2.4 An EM algorithm for Logistic Regression

Here we elaborate on the computation of a Z estimator. In general, an estimator may require specific implementation, we provide an illustrative example with logistic regression. Recall the characterization of the average estimating function $m_i(Y_i, \mathbf{a}, \mathbf{X}; \beta, \theta) = \mathbb{E}[\widetilde{m}(Y_i, S_i(\mathbf{X}, G), V_i(\mathbf{a}, G); \beta) | \mathbf{Y}, \mathbf{a}, \mathbf{X}; \theta]$. Under this model, $P(Y_i = 1 | S_i(\mathbf{X}, G), V_i(\mathbf{a}, G)) = \Lambda(\widetilde{h}(S_i, V_i)^T \beta)$. In order to compute the new estimating function, we need to be able to consider the distribution of the graph, conditional on the observed outcome Y_i . Specifically.

$$P(G|Y_i, \mathbf{a}, \mathbf{X}, \beta, \theta) = \frac{P(Y_i|G, \mathbf{a}, \mathbf{X}; \beta)P(G|\mathbf{a}, \mathbf{X}, \theta)}{P(Y_i|\mathbf{a}, \mathbf{X}, \beta, \theta)}$$
$$= \frac{P(Y_i|S_i(\mathbf{X}, G), V_i(\mathbf{a}, G); \beta)P(G|\theta)}{P(Y_i|\mathbf{a}, \mathbf{X}, \beta, \theta)}$$

In a standard missing data problem, one would impute the missing covariates directly, however, due to the dependence through the graph, this can be very difficult to achieve in practice. However, it will be straightforward to sample from the graph model $P(G|\theta)$. Using a simple approach, we can compute the maximizer exploiting standard software methods using an EM algorithm (Dempster et al., 1977; Wu, 1983). Suppose that we draw a sample of graphs from the generative model $\{G^{(l)}\}_{l=1}^{L} \sim_{iid} P(G|\theta)$.

Let $w_i(Y_i, G; \beta)$ define the weight of an observation.

$$w(Y_i, G; \beta) = \frac{P(Y_i | S_i(\mathbf{X}, G), V_i(\mathbf{a}, G); \beta)}{P(Y_i | \mathbf{a}, \mathbf{X}, \beta, \theta)}$$
$$\approx \frac{P(Y_i | S_i(\mathbf{X}, G), V_i(\mathbf{a}, G); \beta)}{\frac{1}{L} \sum_{l=1}^{L} P(Y_i | S_i(\mathbf{X}, G^{(l)}), V_i(\mathbf{a}, G^{(l)}); \beta)}$$

We next construct the EM algorithm as follows.

In practice, this allows for one to use standard solvers for the (M-step), after sampling a single time with the (E-step).

Additionally, one can include correlations across the observations Y_i through the use of a generalized estimating equation approach. In other generalized linear models, additional assumptions may be required in order to model the full conditional distribution $P(Y_i|S_i(\mathbf{X}, G), V_i(\mathbf{a}, G); \beta)$ such as a dispersion component.

- 1: Sample $\{G^{(l)}\}_{l=1}^{L} \sim_{\text{iid}} P(G|\widehat{\theta})$ denote a sample from the graph model and initialize parameters $\widehat{\beta}^{(0)}$
- 2: for $t \in \{1, 2, ..., T\}$ do
- 3: (E-step) Compute the weighted empirical estimating function

$$m_n^{(t)}(\mathbf{Y}|\mathbf{a}, \mathbf{X}, \beta, \widehat{\theta}) = \frac{1}{L} \frac{1}{n} \sum_{l=1}^{L} \sum_{i=1}^{n} \widetilde{m}(Y_i, S_i(\mathbf{X}, G^{(l)}), V_i(\mathbf{a}, G^{(l)}); \beta) w(Y_i, G^{(l)}; \widehat{\beta}^{(t-1)})$$

4: (M-step) Solve the new estimating function by solving:

$$m_n^{(t)}(\mathbf{Y}|\mathbf{a}, \mathbf{X}, \widehat{\beta}^{(t)}, \widehat{\theta}) = 0$$

5: end for

A.2.5 Plug-in estimates of the Causal parameter

For many problems, the parameter of interest is a causal query conditional on the complete graph G as described in section 2.3. For example, one may care about the expected number of adoptions after seeding an individual in block k v.s. block k'. In this section, we illustrate how to construct an estimate of the causal parameter $\Psi(\mathbf{a}|G)$ using our conditional model estimation procedure.

Let $\Psi(\mathbf{a}|\theta_0) = \mathbb{E}[\Psi(\mathbf{a}|G)|\mathbf{a}, \mathbf{X}, \theta_0]$ be the average causal effect of policy \mathbf{a} over all draws of the graph model θ_0 . We will establish conditions under which these two quantities are close to one another.

Recall the true conditional mean function $\mathbb{E}[Y|S_i = s, V_i = v] = h_0(s, v)$. Under a correctly specified conditional model, $h_0(s, v) = h(s, v; \beta_0)$, and $\Psi(\mathbf{a}|\theta_0) = \Psi(\mathbf{a}|\beta_0, \theta_0)$ where

$$\Psi(\mathbf{a}|\beta,\theta) = \frac{1}{n} \sum_{i=1}^{n} \mathbb{E}[f(V_i, S_i; \beta) | \mathbf{a}, \mathbf{X}, \theta].$$
(A.2.5)

In order to estimate $\Psi(\mathbf{a}|G)$ we plug-in the estimates for the mean model and network

model $\Psi(\mathbf{a}|\hat{\beta},\hat{\theta})$. We next discuss the asymptotics of the plug-in estimate.

Lemma A.2.5 (Inference for a plug-in causal parameter). Assume the conditions of 3.1. Further, assume:

$$\sup_{\beta} |\mathbb{E}[h(S_i(\mathbf{X};G), V_i(\mathbf{a}|G); \beta)|\mathbf{a}, \mathbf{X}, \theta] - \mathbb{E}[h(S_i(\mathbf{X};G), V_i(\mathbf{a}|G); \beta)|\mathbf{a}, \mathbf{X}, \theta']| \le b_i ||\theta - \theta'||$$
(A.2.6)

where $b_i \leq M < \infty$. Denote

$$Q_n(\beta) := \frac{1}{n} \sum_{i=1}^n \frac{\partial}{\partial \beta'} \mathbb{E}[h(S_i(\mathbf{X}; G), V_i(\mathbf{a}|G); \beta') | \mathbf{a}, \mathbf{X}, \theta_0] \Big|_{\beta' = \beta} \in \mathbb{R}^{1 \times p}$$

and

$$\widetilde{\omega}_n := Q_n(\beta_0) D_n(\beta_0) \Gamma_n D_n(\beta_0)^T Q_n(\beta_0)^T.$$

If $s(n) = o(\sqrt{\widetilde{\omega}_n})$. Then

$$\widetilde{\omega}_n^{-1/2}(\Psi(\widehat{\beta},\widehat{\theta}) - \Psi(\beta_0,\theta)) \to_d N(0,1)$$
(A.2.7)

This lemma is essentially an application of the delta method, with the additional caveat that we estimate θ before the plug-in estimate. As before, this requires a fast estimate of the graph generative model parameter, but we add the slightly different assumption (eq. (A.2.6)) that the smoothness in the model class is over the conditional response models $\mathbb{E}[h(S_i, V_i; \beta)|\theta]$, rather than the estimating function $\widetilde{m}(Y, S, V|\beta, \theta)$.

Convergence of the causal parameter to the average over graphs

As we have previously discussed, we can only hope to estimate $\Psi(\mathbf{a}|\theta_0)$ as we do not have access to the full graph G. We next introduce a simple conditions under which the parameter $\Psi(\mathbf{a}|G)$ is close to its average over draws of the graph $G \sim \theta_0$, $\Psi(\mathbf{a}|\theta_0)$.

Assumption A.2.6 (v_n -response dependence). For any graph draw G let $G'^{(ij)}$ denote the graph G with the ij entry swapped from 0 to 1 or vice versa. Let $c_{ij,n}$ denote the bounds of

the differences such that

$$\left|\frac{1}{n}\sum_{i=1}^{n}h_{0}(S_{i}(\mathbf{X},G),V_{i}(\mathbf{a},G)) - h_{0}(S_{i}(\mathbf{X},G^{\prime(ij)}),V_{i}(\mathbf{a},G^{\prime(ij)}))\right| \le c_{ij,n}$$
(A.2.8)
And let $v_{n}^{2} = \sum_{ij:i \ne j} c_{ij,n}^{2}$

Lemma A.2.7. Under Assumption A.2.6

$$\Psi(\mathbf{a}|G) - \Psi(\mathbf{a}|\theta_0) = O_P(v_n)$$

The proof is a one-line application of McDiarmid's inequality. Previous related work such as Breza et al. (2023) typically assume that such a quantity is consistent, however here we quantify the rate here. We next highlight an example;

Example A.2.4 (Conditional Mean Function Example). We abbreviate G = G and $G' = G'^{(kl)}$. Let $h_0(S_i(\mathbf{X}, G), V_i(\mathbf{a}, G)) = \beta_0 + \beta_1 a_i + \beta_2 X_i + \beta_3 \sum_{l \neq i} \frac{X_l G_{kl}}{n} + \beta_4 \sum_{l \neq k} \frac{a_l G_{il}}{n}$ denote a linear response function dependent on the density of connected neighbors. Suppose that the covariate values are bounded $|X_i| \leq M < \infty$. Then:

$$\left| \frac{1}{n} \sum_{k=1}^{n} h_0(S_i(\mathbf{X}, G), V_i(\mathbf{a}, G)) - h_0(S_i(\mathbf{X}, G'), V_i(\mathbf{a}, G')) \right|$$

= $\left| \frac{1}{n} \sum_{k=1}^{n} \sum_{l \neq k} \beta_3 \frac{X_l(G_{kl} - G'_{kl})}{n} + \beta_4 \frac{a_l(G_{kl} - G'_{kl})}{n} \right|$
 $\leq \left| \frac{1}{n} \frac{X_j + X_i}{n} + \frac{|a_j| + |a_i|}{n} \right|$
 $\leq (2M+2)/n^2$

Applying A.2.7 illustrates that: $\Psi(\mathbf{a}|G) - \Psi(\mathbf{a}|\theta_0) = O_p(n^{-1})$. Hence in order to estimate the expected average outcome, all we need is a consistent estimate of the model parameters β_0 .

A.2.6 Optimal design for a Z-estimator

Here we illustrate the optimal design approach for Z-estimators. In this example, the variance itself may depend on the a parameter β , and thus one can include a working candidate for the parameter β' . In general, one could also propose a feasible range of the working parameters β' and consider the worst case variance in that range.

e.		
----	--	--

- 1: **Inputs:** Working covariance Γ_n , model estimate $\hat{\theta}$, working parameter β'
- 2: Sample L draws from the graph model $\{\widehat{G}^{(l)}\}_{l=1}^L \sim \widehat{\theta}$
- 3: Sample R treatments $\{\mathbf{a}_r\}_{r=1}^R$ according to the block-saturation levels τ .
- 4: for $r \leftarrow 1$ to R do

5: Compute
$$\widehat{D}_r(\mathbf{a}) = \frac{1}{nL} \sum_{l=1}^L \sum_{i=1}^n \nabla_\beta m_i(Y_i, S_i V_i; \widehat{G}^{(l)}, \beta')$$

6: Compute the variance for a single draw \mathbf{a}_r :

$$v^{\phi}(\mathbf{a}_r;\widehat{\theta}) = \phi^T \widehat{D}_r(\mathbf{a})^{-1} \Gamma_n \widehat{D}_r(\mathbf{a})^{-1T} \phi$$

7: end for

8: Average over each of the draws $\overline{\upsilon}(\tau; \widehat{\theta}) = \sum_{r=1}^{R} \upsilon^{\phi}(\mathbf{a}_r; \widehat{\theta})$

A.2.7 Experimental Design Variance Minimization With Model Uncertainty

As an extension of our variance minimizing procedure, we can incorporate the uncertainty in our estimates of the model parameters. For instance, consider the following parametric bootstrap approach for estimating the model parameters of the stochastic blockmodel when using ARD.

For example, consider a scenario where we utilize the stochastic blockmodel and we

collect ARD. Denote $\hat{\theta} = (\{\hat{Z}_i\}_{i=1}^n, \hat{\mathbf{P}})$ the initial estimate of the model as computed from Lemma 3.4. We can construct a sampling distribution of $\hat{\theta}^{(b)}$ using the following procedure. Let X_{it}^* denote the ARD responses of the number of connections individual *i* has to someone of trait *t* and let $T_i \in \{0, 1\}^T$ denote the trait memberships of the corresponding individuals.

Algorithm 7 Bootstrap ARD algorithm.

- 1: Estimate $\widehat{\theta}$ from \mathbf{X}^*
- 2: for $b \in \{1, 2, \dots, B\}$ do
- 3: Sample $G^{(b)} \sim \hat{\theta}$
- 4: Construct the ARD vector based on the resampled responses $X_{it}^{*(b)}$ using counts according to connections of $G^{(b)}$ to the nodes with corresponding traits $\{T_i\}_{i=1}^n$
- 5: Estimate $\hat{\theta}$ from $\mathbf{X}^{*(b)}$

6: end for

This approach can work for any procedures which can allow for a sampling distribution of the model parameters $\{\widehat{\theta}^{(b)}\}_{b=1}^{B}$. For example Baraff et al. (2016) considers a nonparametric bootstrap for respondent driven sampling.

In all such cases we would like to include the uncertainty in $\hat{\theta}$ to the saturation assignment, we apply Algorithm 3 (or Algorithm 6) to each of the *b* draws. Using the distribution of variances obtained over the *b* draws, one can compute average or upper confidence bounds on the variance. For example in the simulation illustrated in section A.4.2 we select saturations based on the 2-standard deviation upper confidence bound of the average variance across $b \in \{1, 2, ..., B\}$.

A.3 Proofs of Theorems

In this section, we introduce the proofs for the results in the main paper as well as the additional theoretical results presented in section A.2.

A.3.1 Proof of Lemma 3.4

Proof. Under the stochastic blockmodel assumption, the true latent traits are some discrete type $k_i \in \{1, 2, ..., K\}$. Then the mean connection probability Z_{ck} is simply a mixture over the connection probabilities, weighted by $P(k_j = k'|t_j = t)$. Let N_k denote the set of individuals with group k membership. Furthermore, let $n = |N_k|$. Denote analogous quantities for the trait memberships N_t as well as the intersection of k and t by N_{kt} . When we have a correct clustering. Denote $\widehat{P}_{kt} = \frac{1}{n_k} \sum_{i \in N_k} \frac{1}{n} \widetilde{Y}_{it}$. Assuming independent samples conditional on the graph clusters, let $P_{kt} = \frac{1}{n_t} \sum_{i \in N_{kt}} P_{kk'}$ denote the mean probability of connection averaged over the clusters conditional on their latent traits. Let $\omega_{kt} = \frac{n_{kt}}{n_t}$.

We can express $\tilde{P}_{kt} = P(G_{ij} = 1 | k_i = k, t_j = t)$ as a weighted sum of the connection probabilities from the constituent distributions. If the true clusters are known, then these proportions ω_{kt} are known exactly from the data. Then

$$\begin{split} \widetilde{P}_{kt} &= P(G_{ij} = 1 | k_i = k, t_j = t) \\ &= \sum_{k'=1}^{K} P(G_{ij} = 1 | k_i = k, k_j = k', t_j = t) P(k_j = k' | t_j = t) \\ &= \sum_{k'=1}^{K} P(G_{ij} = 1 | k_i = k, k_j = k', t_j = t) \omega_{k't} \\ &= \sum_{k'=1}^{K} P(G_{ij} = 1 | k_i = k, k_j = k') \omega_{k't} \\ &= \sum_{k'=1}^{K} P_{kk'} \omega_{k't} \end{split}$$

Expressing this relationship over the whole set of matrices, we have:

$$\widetilde{P} = \Omega P$$

Where $\Omega_{tk} = \frac{n_{tk}}{n_k}$. We can solve this system as long as the columns of Ω are linearly independent. Therefore:

$$P = (\Omega^T \Omega)^{-1} \Omega^T \widetilde{P}$$

We next bound the estimation error in Frobenius norm of the true cross-cluster probabilities

$$\begin{split} ||\widehat{P} - P||_{F} &= ||(\Omega^{T}\Omega)^{-1}\Omega^{T}(\widehat{\widetilde{P}} - \widetilde{P})||_{F} \\ &\leq ||(\Omega^{T}\Omega)^{-1}\Omega^{T}||_{F}||(\widehat{\widetilde{P}} - \widetilde{P})||_{F} \\ &\leq \sqrt{||(\Omega^{T}\Omega)^{-1}\Omega^{T}||_{F}^{2}}||(\widehat{\widetilde{P}} - \widetilde{P})||_{F} \\ &\leq \sqrt{\mathrm{Tr}\left((\Omega^{T}\Omega)^{-1}\Omega^{T}\Omega(\Omega^{T}\Omega)^{-1}\right)}||(\widehat{\widetilde{P}} - \widetilde{P})||_{F} \\ &= \sqrt{\mathrm{Tr}\left((\Omega^{T}\Omega)^{-1}\right)}||(\widehat{\widetilde{P}} - \widetilde{P})||_{F} \end{split}$$

Since we assume that Ω 's column's are linearly independent, then $\Omega^T \Omega$ is invertible. Therefore, what remains is bounding the Frobenius norm of $||(\widehat{\widetilde{P}} - \widetilde{P})||_F$.

For each element, let

$$\widehat{\widetilde{P}}_{tk} = \frac{1}{n_k n_t} \sum_{i \in N_k} \widetilde{Y}_{ik}$$
$$= \frac{1}{n_k n_t} \sum_{i \in N_k} \sum_{j \in N_t} G_{ij}$$

Therefore, applying Hoeffding's inequality

$$P(|\widetilde{\widetilde{P}}_{tk} - \widetilde{P}_{tk}| \ge \epsilon) \le 2 \exp\left(-2\epsilon^2 n_k n_t\right)$$

Letting $\rho_k = \frac{n_k}{n}, \rho_t = \frac{n_t}{n}$, then

$$P(|\widehat{\widetilde{P}}_{tk} - \widetilde{P}_{tk}| \ge \epsilon) \le 2 \exp\left(-2\epsilon^2 \rho_k \rho_t n^2\right)$$

Therefore, by a union bound,

$$P(\max_{k,t} |\widehat{\widetilde{P}}_{tk} - \widetilde{P}_{tk}| \ge \epsilon) \le 2KT \exp\left(-2\epsilon^2 \rho_k \rho_t n^2\right)$$
$$\implies P(\sum_{k,t} |\widehat{\widetilde{P}}_{tk} - \widetilde{P}_{tk}| \ge KT\epsilon) \le 2KT \exp\left(-2(KT)^2 \epsilon^2 \rho_k \rho_t n^2\right)$$

Therefore,

$$||\widehat{\widetilde{P}}_{tk} - \widetilde{P}_{tk}||_1 = \mathcal{O}_P(\frac{KT\sqrt{\log(KT)}}{n})$$

Hence

$$||\widehat{P} - P||_2 = \mathcal{O}_P(\frac{KT\sqrt{\log(KT)}}{n})$$

Lastly, we show that as n grows, the probability of achieving a correct clustering of the true block memberships approaches 1. Recall that $n_t = \rho_t n$, and let $\underline{\rho_T} = \min_t \rho_t$. By Hoeffding's inequality: $P(||X_i^{\dagger} - Z_{k_i}|| > \epsilon_n) \leq 2 \exp(-2\epsilon_n^2 n/\underline{\rho_T})$. Taking a union bound over all response vectors, $P(\max_i ||X_i^{\dagger} - Z_{k_i}|| > \epsilon_n) \le 2n2 \exp(-2\epsilon_n^2 n/) \to 0$ for all $\epsilon_n = o(\sqrt{\log(n)/n}).$

Therefore, as n grows, the normalized response vectors in each cluster become well separated, and once $\epsilon_n < \min ||Z_k - Z_{k'}||/2$, then all clusters will be well separated and naively hierarchical agglomerative clustering will consistently group the blocks together for K clusters. Therefore for example, if we let $\epsilon_n = \log(n)n^{-1/2}$, then $P(\max_i\{\hat{k} \neq k\} = O(\frac{1}{n}))$. Of course the labels learned are only consistent up to permutation. We exploit the fact that as referred to in Breza et al. (2023), the clustering problem gets easier as the sample size grows. Let \mathcal{E} be the event that $\hat{k}_i = k_i$ up to permutation for all $i \in \{1, 2, \ldots, n\}$, i.e. $P(\max_i |\hat{k}_i = k_i| > 0) = 1 - P(\mathcal{E}) \leq \frac{1}{n}$. Since the estimators are not necessarily independent of the event of perfect classification.

$$\begin{split} P(||\widehat{P}_{\widehat{\mathbf{k}}} - P_{\mathbf{k}}|| > \epsilon) &= P(||\widehat{P}_{\widehat{\mathbf{k}}} - P_{\mathbf{k}}|| > \epsilon |\mathcal{E})P(\mathcal{E}) + P(||\widehat{P}_{\widehat{\mathbf{k}}} - P_{\mathbf{k}}|| > \epsilon |\mathcal{E}^{c})P(\mathcal{E}^{c}) \\ &\leq P(||\widehat{P}_{\widehat{\mathbf{k}}} - P_{\mathbf{k}}|| > \epsilon, \mathcal{E}) + P(\mathcal{E}^{c}) \\ &\leq P(||\widehat{P}_{\widehat{\mathbf{k}}} - P_{\mathbf{k}}|| > \epsilon, \mathcal{E}) + \frac{1}{n} \\ &= P(||\widehat{P}_{\mathbf{k}} - P_{\mathbf{k}}|| > \epsilon, \mathcal{E}) + \frac{1}{n} \text{ Since } \mathcal{E} \text{ indicates the correct classification} \\ &\leq P(||\widehat{P}_{\mathbf{k}} - P_{\mathbf{k}}|| > \epsilon) + \frac{1}{n} \\ &\leq \sqrt{\operatorname{Tr}\left((\Omega^{T}\Omega)^{-1}\right)} 2KT \exp\left(-2(KT)^{2}\epsilon^{2}\rho_{k}\rho_{t}n^{2}\right) + \frac{1}{n} \end{split}$$

Therefore

$$||\widehat{P}_{\widehat{\mathbf{k}}} - P_{\mathbf{k}}|| = \mathcal{O}_P(\frac{KT\sqrt{\log(KT)}}{n})$$

A.3.2 Proof of theorem 3.2

Proof. We emphasise that in general, the outcomes \mathbf{Y} may be dependent, and this is reflected through correlation in the estimating functions (or the residuals in the case of OLS). We will partition the proof into two sections. First, we will prove the consistency of the estimator $\hat{\beta}$ and secondly, we will prove the central limit theorem.

Consistency: The following result hinges on a typical consistency proof for the M or Z estimators using a structure similar to those found in Chapter 5 of Vaart (1998). First, we denote that:

$$m_n(\mathbf{Z}; \widehat{\beta}, \widehat{\theta}) - g_n(\mathbf{Z}; \widehat{\beta}) = m_n(\mathbf{Z}; \widehat{\beta}, \widehat{\theta}) - m_n(\mathbf{Z}; \widehat{\beta}, \theta_0)$$
$$\leq b_n(\mathbf{Z}) ||\widehat{\theta} - \theta_0||$$
$$= O_P(1)o_P(s(n))$$
$$= o_P(s(n))$$

Next, we can see that, based on this expansion,

$$m_n(\mathbf{Z}; \widehat{\beta}, \widehat{\theta}) = 0$$

$$\implies 0 = (m_n(\mathbf{Z}, \widehat{\beta}, \widehat{\theta}) - g_n(\mathbf{Z}, \widehat{\beta})) + g_n(\mathbf{Z}; \widehat{\beta})$$

$$= o_P(s(n)) + g_n(\mathbf{Z}; \widehat{\beta}) \text{ By A2}$$

At this point, we can now treat this as a standard Z-estimation problem. Therefore, by A2 and A1, then $\hat{\beta}$ is a solution to the estimating function g and is therefore consistent by an application of Theorem 5.9 of Vaart (1998).

Asymptotic Normality: We illustrate asymptotic normality through a Taylor series

expansion argument. As we saw in the consistency part of the proof

$$g_n(\mathbf{Z};\widehat{\beta}) = o_P(s(n))$$

For brevity in notation, we suppress the dependence on \mathbf{Z} , which is implicit for functions, with the subscript *n*. Using a Taylor expansion around β_0 , and let $\tilde{\beta}_j \in [\beta_{0,j}, \hat{\beta}_j]$ for $\beta_{0,j} \leq \hat{\beta}_j$ and $\tilde{\beta}_j \in [\hat{\beta}_j, \beta_{0,j}]$ otherwise.

$$g_n(\widehat{\beta}) = g_n(\beta_0) + D_n(\mathbf{Z};\beta_0)(\widehat{\beta} - \beta_0) + \sum_{jk} \frac{\partial^2}{\partial \beta_j \beta_k} g_n(\mathbf{Z};\widetilde{\beta})(\widehat{\beta}_j - \beta_{0,j})(\widehat{\beta}_k - \beta_{0,k})$$
$$= g_n(\beta_0) + D_n(\mathbf{Z};\beta_0)(\widehat{\beta} - \beta_0) + o_P(s(n) + ||\widehat{\beta} - \beta_0||)$$

by the application of the consistency and A2. Therefore, we focus on main terms. By Assumption C1.

Therefore:

$$\Gamma_n^{-1/2} D_n(\mathbf{Z};\beta_0)(\widehat{\beta}-\beta_0) = \Gamma_n^{-1/2} g_n(\beta_0) + o_p(\frac{s(n)}{r(n)})$$

Noting that $D_n(\beta_0) - D(\beta_0) = o_P(1)$, by an application of Slutsky's lemma:

$$\Gamma_n^{-1/2} D(\beta_0) (\widehat{\beta} - \beta_0) \to_d N(0, I_p)$$

and therefore, the proof is complete.

A.3.3 Proof of Lemma 3.5

We first include a useful lemma for bounding the approximation of the error of the graphon model.

Lemma A.3.1 (Lemma 2.1 of Gao et al. (2015)). Denote $k_i \in \{1, 2, ..., K\}$ are the block memberships of a stochastic-blockmodel with average connection probabilities across blocks

 $\overline{\eta}_{ij} = P_{k,k'} = \frac{1}{n_k n_{k'}} \sum_{i,j:k_i=k,k_j=Z_i} \sum_{l:Z_l=Z_j} \eta_{kl}$. If the true graphon $g \in \mathcal{H}_{\alpha}(M)$, then, there exists some membership vector \mathbf{k} and corresponding average across block probabilities P_0 such that:

$$\frac{1}{n^2} \sum_{ij} (\eta_{ij} - \overline{\eta}_{ij})^2 \le M^2 \left(\frac{1}{K^2}\right)^{\alpha \wedge 1}$$

We now proceed with a the proof of the lemma.

Proof. We firstly use a Taylor expansion of $L_n(\beta_0, \eta_*)$ where $\tilde{\beta}$ is an element-wise intermediate value of β and $\tilde{\beta}$

$$L_{n}(\beta_{0},\eta_{*}) = L_{n}(\beta_{*},\eta_{*}) + \frac{\partial}{\partial\beta}L_{n}(\beta,\eta_{*})\Big|_{\beta=\beta_{*}}(\beta_{0}-\beta_{*})$$
$$+ \sum_{jk}\frac{\partial^{2}}{\partial\beta_{j}\partial\beta_{k}}L_{n}(\widetilde{\beta},\eta_{*})(\beta_{0j}-\beta_{*j})(\beta_{0k}-\beta_{*k})$$
$$\frac{\partial}{\partial\beta}L_{n}(\beta,\eta_{*})\Big|_{\beta=\beta_{*}}(\beta_{0}-\beta_{*}) = -L_{n}(\beta_{0},\eta_{*}) + \sum_{jk}\frac{\partial^{2}}{\partial\beta_{j}\partial\beta_{k}}L_{n}(\widetilde{\beta},\eta_{*})(\beta_{0j}-\beta_{*j})(\beta_{0k}-\beta_{*k})$$

Since we assume $L_n(\beta, \eta_*)$ is twice continuously differentiable in β , and \mathcal{B} is compact, then $\frac{\partial^2}{\partial \beta_j \partial \beta_k} L_n(\widetilde{\beta}, \eta_*)$ is bounded. Therefore,

$$||\beta_0 - \beta_*||_2 \le \frac{|L_n(\beta_0, \eta_*)|}{\lambda \sqrt{p}} + O(||\beta_0 - \beta_*||_2^2)$$

Lastly, by our continuity assumptions, $|L_n(\beta_0, \eta_*)| \leq L||\eta_0 - \eta_*||_2/n \leq LMK^{-(\alpha \wedge 1)}$. After applying this, our proof is complete.

A.3.4 Proof of Lemma A.2.4

Proof. The proof is straightforward application of Hoeffding's inequality. Given an m node subsample of the full graph, and given their known types. Since $\widehat{\mathbf{P}}_{kk'} = \frac{1}{\rho_{kk'}m} \sum_{i,j} G_{ij}I(k_i = k, k_j = k')$, then the final result is a direct application of Hoeffding's inequality.

For the missing data case, we can plug-in the estimate of the edge sampling $P(G_{ij} = 1|X_{ij} = x)$ in order to correct for the missingness of the edges. If $\sup_x |\hat{P}(G_{ij} = 1|X_{ij} = x) - P(G_{ij} = 1|X_{ij} = x)| = o_P(m^{-1})$ then the estimation of the propensity is negligible and we can correct for the missingness of edges.

A.3.5 Proof of theorem A.2.3

Proof. We first we expand the form of the OLS estimator.

$$\begin{split} \widehat{\beta}_{lm} &= \mathsf{H}_{n}^{-1}(\widehat{\theta}) \frac{1}{n} \sum_{i=1}^{n} \widetilde{H}_{i}(\widehat{\theta}) Y_{i} \\ &= \mathsf{H}_{n}^{-1}(\widehat{\theta}) \frac{1}{n} \sum_{i=1}^{n} \widetilde{H}_{i}(\widehat{\theta}) (\widetilde{H}_{i}^{T}(\theta_{0})\beta_{0} + u_{i}) \\ &= \mathsf{H}_{n}^{-1}(\widehat{\theta}) \frac{1}{n} \sum_{i=1}^{n} \widetilde{H}_{i}(\widehat{\theta}) (\widetilde{H}_{i}^{T}(\widehat{\theta})\beta_{0} + (\widetilde{H}_{i}^{T}(\theta_{0}) - \widetilde{H}_{i}^{T}(\widehat{\theta}))\beta_{0} + u_{i}) \\ &= \beta_{0} + \underbrace{\mathsf{H}_{n}^{-1}(\widehat{\theta}) \frac{1}{n} \sum_{i=1}^{n} \widetilde{H}_{i}(\widehat{\theta}) (\widetilde{H}_{i}^{T}(\theta_{0}) - \widetilde{H}_{i}^{T}(\widehat{\theta}))\beta_{0}}_{(A)} \\ &+ \underbrace{\mathsf{H}_{n}^{-1}(\widehat{\theta}) \frac{1}{n} \sum_{i=1}^{n} (\widetilde{H}_{i}(\widehat{\theta}) - \widetilde{H}_{i}(\theta_{0}))u_{i}}_{(B)} + \underbrace{\mathsf{H}_{n}^{-1}(\widehat{\theta}) \frac{1}{n} \sum_{i=1}^{n} \widetilde{H}_{i}(\theta_{0})u_{i}}_{(C)} \end{split}$$

We next bound terms (A) and (B) after which, the asymptotic distribution of (C) will be apparent.

We note that the Hessian evaluated at the true model parameters can be evaluated $H_n(\widehat{\theta}) = H_n(\theta_0) + o_P(s(n))$ by assumptions E2 and E1. By the continuous mapping theorem $H_n(\widehat{\theta}) = H_n(\theta_0) + o_P(s(n))$. We see that $(A) = o_P(s(n))$ by assumptions E4, E2 and E1. Next, by the stochastic boundedness of the error E5 and applying Hölder's inequality.

$$\frac{1}{n}\sum_{i=1}^{n} (\widetilde{H}_{i}(\widehat{\theta}) - \widetilde{H}_{i}(\theta_{0})) \leq (\frac{1}{n}\sum_{i=1}^{n}|u_{i}|)\max_{i}||\widetilde{H}_{i}(\widehat{\theta}) - \widetilde{H}_{i}(\theta_{0})||$$
$$= o_{P}(s(n))$$

Therefore:

$$\Gamma_n^{-1/2} \mathsf{H}_n(\widehat{\theta})(\widehat{\beta}_{lm} - \beta_0) = \Gamma_n^{-1/2} \sum_{i=1}^n \widetilde{H}_i(\theta_0) u_i + o_P\left(\frac{s(n)}{r(n)}\right) \to_d N(0, I_p)$$

by F1 and Slutsky's Lemma, completing the proof.

A.3.6 Proof of Lemma A.2.5

Proof. The proof follows from an application of the delta method, with the additional caveat that we must account for the estimation of the model parameters θ_0 In this case:

$$\begin{aligned} |\Psi(\widehat{\beta},\widehat{\theta}) - \Psi(\widehat{\beta},\theta_0)| &\leq \frac{1}{n} \sum_{i=1}^n b_i ||\widehat{\theta} - \theta_0|| \\ &= o_P(s(n)) \end{aligned}$$

The remainder of the proof follows from a simple application of the delta method using the plug-in estimator $\Psi(\hat{\beta}, \theta_0)$. See Theorem 3.1 of Vaart (1998).

A.4 Additional Simulations

Here we provide additional details with respect to several aspects of our methodology. We also include further details on several of the details for the implementation of competing methods in section A.4.5.

A.4.1 Coverage of the GATE

In our simulation setup in section 5.1 we can also compute confidence intervals based on the regression $Y_i = \beta^T \mathbb{E}[\tilde{h}(S_i, V_i)] + \epsilon_i$ where we apply the Eicker-Huber-White sandwich estimator of the variance. We then compute the corresponding plug-in estimator of the variance using the covariates observed and Lemma A.2.5. Since the covariates in the true

regression model behave like averages over the graph, we expect Lemma A.2.7 to hold and therefore the difference between the GATE for any one draw of the graph, and the true GATE is very small. We see in Figure 6 that the coverage tends to be larger than the nominal 95%, though in general, due to model misspecification of the true-graph, there can be additional uncertainty due to the misspecification of the graph model. However, we see in this simple example that the coverage performs well with an off-the-shelf implementation.



Figure 6: Coverage of the GATE using Eicker-Huber-White estimates of the variance.

A.4.2 Experimental Design: Local Diffusion

We next consider an example using a local diffusion process. We suppose that seed nodes are placed at time 0 and that outcomes are measured at time T = 1, allowing for diffusion to only take place to the immediate neighbors with a fixed probability q. In this case, for nonseed nodes the probability of infection is related to the total number of treated neighbors
through the following link function. Under this model let $V_i \in \{0, 1\}$ denote the exposure as to whether one of their neighbors have received the treatment, i.e. $V_i = I(\sum_j G_{ij}a_j > 0)$. Then

$$\mathbb{E}[Y_i|V_i, S_i] = qV_i$$

In this experiment, a single individual is seeded in each network. Our goal is to identify the best individuals in each of the network to seed and rank them by the expected variance of the estimator. We compare this to random seeding of individuals in the network as well as seeding by only the highest degree nodes. We use the networks constructed by the union of all connections of Banerjee et al. (2019). We construct estimates of the stochastic blockmodel as the partial data example using K = 3 in each case. We construct the traits using ARD responses based on number of connections with the following traits outlined in the Appendix in section A.4.5.2. We also include an alternative where a beta-model (Chatterjee and Diaconis, 2011) is used in place of the SBM for the degree seeding where further details on estimation are included in section A.4.5.1. We then draw samples of the graph using the parametric bootstrap to obtain a resampled distribution of ARD $\{\mathbf{X}^{*(b)}\}_{b=1}^{B}$ for B = 1000. We identify the optimal treatment block for each parameter according to section A.2.7. We simulate 1000 draws of the draws in the diffusion process for each true, and plot the associated bias and RMSE of the seeding strategies in Figure 7 with a true diffusion parameter q = 0.2.

In the full data case, the optimal strategy would be to seed the highest degree node in each of the networks and measure whether each of their neighbors are infected at time T = 1. However, this poses a problem for the stochastic blockmodel as we are essentially picking an outlier to seed, which is different than a typical member of the block over draws of the process. This can be corrected for using a model which accounts for degree heterogeneity, in our case, the beta model. In our optimal seeding strategy, we find that the RMSE is lower in both the degree optimized strategy with the beta model, as well as the block optimized strategy with the SBM, than even the full data version with a completely randomized allocation, hence highlighting the role of the interplay of the model of the graph and the experimental design. This behavior is observed in Figure 7.



(a) Bias of Full and Partial Data Diffusion Param-(b) RMSE of Full and Partial Data Diffusion Pa-eter Estimatesrameter Estimates

Figure 7: RMSE and bias of estimating parameter q using random seeding, and the optimal seed for each village.

A.4.3 Estimated outcome model

In this example, we consider a problem of optimal treatment assignment after the outcome model is estimated. We consider an example where an outcome model is estimated and transported to a new population. In this example we suppose that there is some benefit $\beta_1 > 0$ to receiving a treatment, and some smaller benefit based on the fraction of the neighbors treated $0 < \beta_2 < \beta_1$. We wish to assign treatments in a way that will maximize the expected outcome $\Psi(\mathbf{a}|G)$ for each network.

$$Y_i = \beta_0 + \beta_1 a_i + \beta_2 q_1 + \epsilon_i$$

Where $q_i := \frac{1}{d_i} \sum_{j=1}^n G_{ij} a_j$ denotes the normalized number of treated neighbors. We simulate the data with $\beta_0 = 1$, $\beta_1 = 1$ and $\beta_2 = 1/2$ with $\sigma_i \sim N(0, 1)$. We choose this form of a response function since it will be simple to solve with an off the shelf mixed-integer programming approach using CVXR (Fu et al., 2020).

We suppose that in each example there is only a budget for $B \in \{10, 20, 40, 80\}$ treatments for each of the villages. The goal is to maximize the overall expected outcome. We consider the following competing procedures. In this case, we suppose that we have a single pilot network where we can learn the model and the goal is to maximize the benefit on the remaining networks. We use the same gossip diffusion networks as in sections A.4.2 and 5.2.

We compare the following seeding strategies.

- 1. Random assignment to all individuals in the network
- 2. Equal assignment amongst clusters.
- 3. Assign treatments ordered by the highest degree of the nodes.
- 4. Maximize the total expected outcome by maximizing $\max_{\mathbf{a};||a||_1 \leq B} \Psi(\mathbf{a}; \hat{\beta}, \hat{\theta})$

Let $\mathbb{E}[Y_i|\mathbf{a}] = \beta_0 + \beta_1 a_i + \beta_2 (1-a_i) \sum_{k'=1}^K \widehat{P}_{\hat{k}_i k'} n_{t,k}$ and let $n_{t,k} = \sum_{j:k_j=k} a_j$. Therefore, the objective function.

$$\Psi(\mathbf{a}|\beta,\theta) = \beta_0 + \frac{1}{n}\beta_1 \mathbf{1}^T \mathbf{n}_t + \frac{1}{n}\beta_2 \zeta^T \mathbf{n}_t$$



Figure 8: Our method, model based optimal treatment allocation (Model Opt) compared to random assignment and assignment to largest and smallest clusters respectively. The larger the values represent larger average outcomes in each of the networks. Curves are fit using cubic splines. The model based optimal design tends to give a higher value at each of the sample sizes at each treatment budget. For example, at a sample size of 150 and a treatment budget of 10, our methods leads to a 30% increase in the average outcome.

where $\zeta = \frac{1}{d_i} \sum_{i=1}^{n} \mathbf{P}_{k_i, \cdot}$ and $\mathbf{n}_t = (n_{t,1}, n_{t,2}, \dots, n_{t,K})$. In general, given a conditional model, one may fine tune the optimization approach to the particular challenges of evaluating the optimal treatment allocation. We partition each network into 6 blocks.

We plot the expected average outcome under each of the treatment allocations for the remaining 68 networks after learning a model from the first pilot network. We repeat this for the total number of treatments $B \in \{10, 20, 40, 80\}$.

In Figure 8 we find that based on our method, we can achieve higher average outcomes than simple models based on the block positioning alone, emphasizing the importance of considering the potential outcome model when optimal targeting.

A.4.4 Inference for evidence of complex contagion with partial network data

We can also replicate the results of Beaman et al. (2021)'s study on the evidence of pitplanting. They consider 3 measures of information diffusion. Firstly, if an individual has heard of pitplanting, second, if they know how to pitplant, and thirdly whether they adopt pitplanting in their practice. In order to control for one's position in the network, the authors consider the distance between the optimal seeds using two other targeting methods, simple diffusion, and geo-targeting as well as complex contagion. They then compare the increased odds of con

$$Y_{iv} = \alpha + \beta_1 I (1TSeeds) + \beta_2 I (2TSeeds) + \beta_3 I (1Simple)_{iv} + \beta_4 I (2Simple)_{iv} + \beta_5 I (1Complex)_{iv} + \beta_6 I (2Complex)_{iv} + \beta_7 I (1Geo)_{iv} + \beta_8 I (2Geo)_{iv} + \delta_v + \epsilon_{iv}$$

Again, we generate synthetic covariates and apply a stochastic blockmodel in order to estimate K = 8 blocks within each of the networks. We plot the coefficients for the connection to exactly 1 seed, 2 seeds and within radius 2 of at least 1 seed in Figure 9. We note that we run the same regression as in Beaman et al. (2021), however, some since the full network data includes some additional noise top preserve anonymity, we do not have the exact same estimates of the coefficients as in their paper, however, the conclusions are substantively the same.



Figure 9: Replication of regression coefficients using aggregated relational data and associated 95% confidence intervals.

A.4.5 Additional Experimental Details

To aid in reputability, we include additional details regarding the implementation of our methods as well as competing methods.

A.4.5.1 Beta Model Estimation

Another common model utilized for random graph formation is the beta model coined by Chatterjee et al. (2011). Namely these are a class of models that can be learned based on their degree sequence. We consider a version where each node has an affinity parameter ν_i and the probability of connection between each pair of nodes is $P(G_{ij} = 1) = \nu_i \nu_j$. Let $\nu_n = \sum_{i=1}^n \nu_i$ Therefore, $\mathbb{E}[d_i = d] = \sum_{j \neq i} P(G_{ij} = 1) = \nu_i (\nu_n - \nu_i)$. The set of parameters $\{\nu_i\}_{i=1}^n$ can be estimated using an iterative solution to the fixed point equation:

$$\nu_i^{(t+1)} = d_i / (\nu_n^{(t)} - \nu_i^{(t)})$$

A.4.5.2 ARD Questions

We utilize the measured traits to construct responses for ARD questions for each individual for the networks in Banerjee et al. (2019). The constructed ARD include traits which ask "How many people do you know ..."

- that are in each sub-caste?
- that are Farmers, Shop owners, Domestic workers etc. ?
- that own their house?
- that have a house with at least 3 rooms?
- that have access to electricity?

For the estimation of the GATE using Banerjee et al. (2013), we use Leiden clustering and denote the clusters traits. When replicating the results of Beaman et al. (2021), only a subset of nodes have available covariate. As was done in our examples with Banerjee et al. (2013), we construct synthetic traits using the clusters observed from Leiden clustering for K = 10. ARD is then constructed based on the connections to nodes of each trait.

A.4.5.3 GATE Estimators

The two estimators we compare for estimation of the global average treatment effect are the difference in means estimator $\hat{\tau}_{DM}$ and the Horvitz-Thompson estimator $\hat{\tau}_{HT}$. Let E_{i0} and E_{i1} denote the events that all neighbours of *i* are untreated (including *i* themselves) and treated respectively.

$$\widehat{\tau}_{DM} = \frac{1}{n_1} \sum_{i=1}^n Y_i a_i - \frac{1}{n_0} \sum_{i=1}^n Y_i (1 - a_i)$$
$$\widehat{\tau}_{HT} = \frac{1}{n} \sum_{i=1}^n \frac{Y_i I(E_{i1})}{P(E_{i1})} - \frac{Y_i I(E_{i0})}{P(E_{i0})}$$

In general, the Horvitz-Thompson estimator will be unbiased, however, it can often suffer from high variance for two reasons. Firstly, the probabilities of the events that all nodes are treated may be exceedingly low, inflating this variance, and also, relatively few nodes receive the exposures under which all of their neighbours are treated or none of them are.

In the case where the spillover effects are relatively mild, often a difference in means approach to the estimator is preferred. The effect of cluster randomization on the MSE of this estimator has been further studied in the complete network Brennan et al. (2022); Viviano (2020).