

Dualities and Categorical Structures from 2D Up

Veronica Pasquarella

Department of Applied Mathematics and Theoretical Physics (DAMTP)

University of Cambridge,

Wilberforce Road, CB3 0WA, Cambridge, UK

E-mail: vp360@damtp.cam.ac.uk

ABSTRACT: A fundamental step towards studying string theory vacua, and, ultimately, their stability, is that of understanding the underlying mathematical structure of the QFT resulting from its dimensional reduction on Calabi-Yau (CY) manifolds, the latter being complex manifolds admitting a category theory description. The present work aims at addressing this question in the case of lower-dimensional gravitating systems, and supersymmetric quiver gauge theories, highlighting the role played by dualities and categorical structures towards studying analytical structures of amplitudes. (Based on the author's PhD thesis).

KEYWORDS: factorisation homology, higher-categorical structures, geometric invariant theory, supersymmetric quiver gauge theories, lower-dimensional holographic duality and vacuum transitions

Contents

I	Introduction	1
1	What this work is about	1
1.1	Motivations and Future Perspectives	1
1.2	Outline	4
2	Roadmap	11
2.1	Roadmap	11
2.1.1	Fields and Interactions	12
2.1.2	Amplitudes and Hilbert series	12
2.2	(Some) Open Questions in Quantum Field Theory	14
2.3	Where to next?	16
3	Relative QFTs	17
3.1	Rings, homologies, cohomologies, and modules	17
3.2	Abstract topological symmetry data in field theory	17
3.2.1	Fiberings over BG	18
3.3	Preliminary examples: the σ -model and gauge theories	19
3.4	Theory X	23
3.5	Main Example: the Coulomb branch chiral algebra	25
3.5.1	Star-shaped quivers	26
3.6	Lagrangian submanifolds of a symplectic manifold	28
3.6.1	Hyperkähler quotients	28
3.7	Abelianisation	29
3.7.1	Towards Category Theory	30
II	Higher-Categories and Dualities	30
4	Higher-categorical structures	31
4.1	Higher-categorical structures	32
4.1.1	Defect fusion categories	33
4.1.2	Example of higher-categorical structures	36
4.2	String Theory and Quantum Mechanics (QM) as Functorial Field Theories (FFT)	38
4.2.1	Calabi-Yau (CY) categories	40

5	Main Example: Cochain level theories	41
5.1	Cochain complexes	41
5.2	Moore-Segal bordism operators	43
5.2.1	Cochain level theories	43
5.3	The Moore-Segal setup	43
5.4	Moore-Tachikawa varieties	45
5.4.1	Categorical structure assuming duality	45
5.4.2	Duality from an algebraic perspective	48
5.5	Examples	51
6	Dualities among Algebraic Varieties	51
6.1	Equivariant cohomology, fixed point localisation and convolution product	52
6.2	Mirror symmetry in Moore-Tachikawa varieties	54
6.2.1	The Kostant-Whittaker symplectic reduction	55
6.3	Homological mirror symmetry	56
6.3.1	Relation to our main example	57
6.3.2	Lagrangian submanifolds	57
6.4	3D mirror symmetry	58
6.4.1	Categorical formulation	58
6.5	From Homological mirror symmetry to 3D mirror symmetry	59
6.5.1	Relation to our main example	60
7	Holographic Dualities and Beyond	62
7.1	Holographic duality	62
7.1.1	Liouville Theory	64
7.2	The Chern-Simons/Wess-Zumino-Witten (CS/WZW) correspondence	66
7.3	Jackiw-Teitelboim (JT)-gravity from CS-theory	67
7.3.1	(P)SU(N) in 4D $\mathcal{N} = 1$ Super Yang Mills	70
7.4	From brane webs to quiver gauge theories	71
7.5	Beyond holographic duality: adding boundaries	72
7.5.1	Islands in JT-gravity	73
7.5.2	Conformal interfaces as boundary states	73
III	From knot invariants to QFTs	76
8	Summary and Motivations of Part III	77
8.1	Motivations	77
8.2	Outline and main results	78
9	Quantum Field Theory from Category Theory	79
9.1	The Freed-Moore-Teleman construction	80
9.1.1	Gauging in relative QFTs	80
9.1.2	Absolute 4D theories from 6D $\mathcal{N} = (2, 0)$ SCFTs	82

9.1.3	From relative to absolute	83
9.2	Fiber functors from non-invertible symmetries	87
9.2.1	Intrinsic non-invertible symmetries and fiber functors	88
9.2.2	Quantum dimension from relative gaugeable algebra	93
9.3	Abelianisation (once more)	101
10	Instantons and Scattering Amplitudes	104
10.1	Instantons	104
10.1.1	Instantons in QM	104
10.1.2	Instantons in QFT	106
10.1.3	Instantons in 4D supersymmetric gauge theories	106
10.1.4	Instantons in oriented string theory: the ADHM construction	107
10.2	The WKB approximation	107
10.2.1	WKB for Hitchin systems	109
10.2.2	WKB for vacuum transitions	109
11	Conclusions and Outlook of Part III	110
IV	Holographic interpretation of 2D vacuum transitions	111
12	Summary and Motivations of Part IV	111
12.1	Motivations	111
12.2	Outline and main results	116
13	Euclidean transitions in 2D	117
13.1	Vacuum transitions in 2D without gravity	117
13.2	Coleman-De Luccia from the Almheiri-Polchinski setup	119
13.3	Brown-Teitelboim in 2D from Almheiri-Polchinski	127
14	Lorentzian transitions in 2D	131
14.1	Actions with vanishing constant energies	132
14.2	Transitions among de Sitter, anti-de Sitter, and Minkowski vacua	138
15	Lorentzian transitions with black holes	140
15.1	Actions with non-vanishing constant energies	140
15.2	Transitions with conical defects	142
15.3	Flat limit	144
16	Holographic interpretation	145
16.1	Islands and horizons in holographic black hole evaporation	146
16.2	Holographic embedding of vacuum transitions	149
16.2.1	Local transitions, entanglement and islands in wedge holography	152
16.2.2	A BCFT ₂ perspective: transitions between end-of-the-world branes	160
16.2.3	An internal CFT ₂	161

17 Conclusions and Outlook of Part IV	163
V Beyond Categorical Dualisability	165
18 Summary and Motivations of Part V	165
18.1 Motivations	165
18.2 Outline and main results	168
19 Categorical Dualisability and full extendibility	169
19.1 Factorisation Homology in context	169
19.1.1 Hochschild cohomology and cochain complexes	170
19.1.2 Factorisation homology of stratified spaces	171
19.1.3 Non-contractible loops	174
19.2 Dualisability and full-extendibility	174
19.2.1 Extending TQFTs	174
19.3 Comparison with our main example	175
19.3.1 Dualisability in categorical structures	175
19.3.2 The case of the Hilbert series	176
20 Drinfeld Centers from Magnetic Quivers	177
20.1 Gauging and Hasse diagrams	178
20.2 Drinfeld centers and mirror symmetry	180
20.3 Drinfeld centers from magnetic quivers of 3D $\mathcal{N} = 4$ gauge theories	180
21 Moore-Tachikawa varieties: beyond duality	184
21.1 Categorical structure without duality	185
21.2 Interesting shortcomings	186
21.3 Back to the identity from moduli spaces	188
21.3.1 Hilbert series of Hochschild cohomology of block algebras	189
21.3.2 Hitchin systems and mirror quivers	189
21.3.3 Looking for the identity	190
21.4 Where to next?	191
21.5 Conclusions and Outlook	192
22 Conclusions and Outlook of Part V	192
VI Conclusions and Outlook	193
23 Conclusions and Outlook	193
A Loop groups	197

B	Supersymmetric Field Theories	198
B.1	Physical Lagrangians	198
B.2	UV/IR flows and their parametrisation	199
B.3	The Witten index	199
B.3.1	R-symmetry, supergroups, superalgebras and twisting	200
B.3.2	The importance of flavour to extremise \mathcal{Z}	202
B.4	SUSY enhancement from branes at orbifolds	202
B.4.1	ABJM and mirror symmetry	203
C	Higher-categorical symmetries from topological defects	203
C.1	Generalised gauging and localised symmetries	205
C.2	Fusion rules	206
D	Topological Wick rotation and holographic dualities	207
E	Quiver gauge theories	208
E.1	Graphs	208
E.2	Dynkin diagrams	209
E.3	Framed Nakajima quiver varieties	210
F	Derivation of generalised black hole (BH) entropy	214
F.1	”Replica trick” derivation	214
F.2	Rényi and von Neumann entropies	214
G	Topological Quantum Field Theories (TQFTs)	218
G.1	Attempts and advancements	218
G.2	The Atiyah-Segal axioms	219
G.2.1	Segal’s Conformal Field Theory (CFT) axioms	219
G.2.2	The Atiyah-Segal axioms	220
G.3	Schwarz-type TQFTs	220
G.3.1	Chern-Simons (CS) Theory	220
G.4	Topological String Theory	221
G.4.1	The A- and B-model	221
H	QFT and the Jones polynomial	222
H.1	QFT and the Jones polynomials	222
H.2	Schemes or algebraic stacks	222
H.3	Morse Theory and Classifying Spaces	222
H.4	Donaldson invariants	223
H.5	Seiberg-Witten theory	224
H.6	From CS to Morse theory	224
H.6.1	Morse Theory	224
H.6.2	Relation to Moduli Spaces of SQFTs	225
H.6.3	The Cobordism Hypothesis	227

I	4D descendants of 6D $\mathcal{N} = (2, 0)$ SCFTs	227
I.1	Class \mathcal{S}	227
I.2	Vacuum structure: Coulomb, Higgs and mixed branches	229
I.3	Metrics	231
I.4	The Alday-Gaiotto-Tachikawa (AGT) correspondence: a brief overview	233
I.5	3D SCFTs on punctures	236
I.6	The Geometric Langlands Program (GLP)	238
J	Categories, knots, and entanglement	242
J.1	Higher-categorical structure of (linked minimal models) knots	242
J.2	Spiral maps, fusion multiplicities and holomorphic blocks	246
J.3	The total quantum dimension	248
K	Sigma-models	249
K.1	The sigma-model	249
K.2	Riemann surfaces	250
K.3	Moduli spaces	250
K.4	The nonlinear sigma-model	250
K.5	Principle bundles and connections	252
K.6	Mathematical Gauge theory	252
K.7	The gauged sigma-model	253
L	(A)dS₃/$T\bar{T}$-deformed CFTs	254
L.1	$\mathcal{Z}_{S^2}{}^{T\bar{T}}$ and $S_{T\bar{T}}$ for AdS	256
L.2	$\mathcal{Z}_{S^2}{}^{T\bar{T}}$ and $S_{T\bar{T}}$ for dS	257
L.3	Comparing $S_{T\bar{T}}$ for AdS and dS	258
L.4	Cutoff dependence of S_{EE}	259
L.5	Key points of $T\bar{T}$ -deformations	259
M	Von Neumann algebras (VNAs)	260
M.1	Types of VNAs	260
M.2	Standard form of a II_1 factor	260
M.3	Representations and von Neumann dimensions	261
M.4	The Jones index	261
N	Other appendices to Part IV	262
N.1	Twist operators and Rényi entropies	262
N.2	$S_{tot}^{\text{AdS} \rightarrow \text{AdSBH}}$	264
N.3	No obstructions from the Generalised Second Law (GSL)	265
N.4	Roots of a quartic	266

O Geometric Invariant Theory (GIT)	267
O.1 Equivariant cohomology	267
O.2 The GIT construction	268
O.3 Moduli spaces: Coulomb and Higgs branches	270
O.4 The monopole formula	270
P Magnetic quivers	271
P.1 Magnetic Quivers: a unifying framework	271
P.2 Intersecting cones	272
P.2.1 Magnetic quivers and Hasse diagrams	274
P.2.2 Rank-1 and rank-2 examples	274

Part I

Introduction

1 What this work is about

The present Thesis is, at the same time, a continuation of an original work I did with my supervisor, Professor Fernando Quevedo, [7], and the birth of my research independence in the world of Physical Mathematics.

Our original aim was that of identifying a way for describing transitions in between different vacua within the String Theory Landscape, and ultimately using this as a criterion for establishing which String Theory models could more accurately describe the world in which we live.

Identifying a rigorous formalism for describing vacuum decay processes has been a great matter of debate for almost half a century, and part of our joint collaboration was attempting to provide further insight into which prescription could better suit our purposes.

As a step forward towards achieving such aim, we carried out an analytical study of the underlying mathematical structure of field theories. In doing so, the main tools adopted in our treatment are dualities and higher-categorical structures.

This first section lays out the motivations for using these tools in addressing this question, as well as indicating future research directions that will be pursued by the author. We conclude the section outlining the content of the present work, highlighting some key findings, referring to their respective sections for detailed explanation.

1.1 Motivations and Future Perspectives

String theory¹ is a promising candidate unifying theory of fundamental interactions. In its fermionic formulation, it involves ten dimensions, of which nine are spatial and one is

¹See for instance [8–17].

temporal. Crucially, such formulation strongly relies upon Supersymmetry (SUSY), [18], which is a correspondence in between bosonic and fermionic particles.

According to present-day evidence, SUSY is certainly broken in our world, and plenty of effort has been made in identifying traces of such symmetry breaking from the observational side², eventually leading to the extension of the Standard Model (SM), [28], by encoding additional particles and interactions. Understanding how the Standard Model can be embedded in String Theory is certainly a step forward towards identifying patterns of SUSY breaking in our world. Albeit being broken, SUSY certainly plays a crucial role in understanding which String Theory model is most appropriate for the SM to arise as an effective field theory (EFT), precisely by studying its breaking.

String theory vacua³, where the SM would eventually be embedded, are parametrised by the so-called moduli space, and the latter is known to admit a categorical algebro-geometric description. Crucially, great advancements in the latter took place simultaneously to the birth and development of SUSY and String Theory. In recent years, joint collaboration in between pure mathematicians and mathematical physicists, primarily Freed, Hopkins, Lurie, Moore, Segal, and Teleman⁴, shed light on new interesting perspectives which happen to be particularly relevant for the objectives outlined above. The main feature of their approach is that of applying the powerful tool of higher-categorical structures within the context of gauge theories and their supersymmetric counterparts, with the ultimate aim being that of explaining what is the underlying mathematical structure of a quantum field theory (QFT), [105–107]. Their work is based upon identifying underlying topological structures from which partition functions, and correlation functions can be derived. Furthermore, their formalism happens to be particularly efficient for keeping track of the spectrum (i.e. the particle-field content) of a given QFT, and how the latter changes under gauging of the global symmetries of the theory one started from.

Up to now, most of the mathematical control in describing higher-dimensional field theories has been for descendants of maximally-supersymmetric 6D $\mathcal{N} = (2, 0)$ superconformal field theories (SCFTs), [108–111], from which a rich web of dualities arises upon dimensional reduction to, either, 4 or 3 spacetime dimensions⁵. Such descendants feature fewer supersymmetries with respect to their non-Lagrangian parent theory. In particular, the most studied are 4D $\mathcal{N} = 2$ (also known as class \mathcal{S} theories), and 3D $\mathcal{N} = 4$ SCFTs, exhibiting exactly half of the amount of supersymmetry of their parent 6D theory.

From the work of Alday, Gaiotto and Tachikawa (AGT), [135], it is known that class \mathcal{S} theories can be equally described in terms of a 2D topological field theory (TFT) living on the Riemann surface on which dimensional reduction has been performed. Such a 2D TFT is defined to be a functor in between 2-categories. Such 2-categories were first studied by Moore and Tachikawa, [136], which, in turn, relied upon the pioneering work of Moore

²Among the most recent advancements in the field, we refer the interested reader to the following works [21–27, 595, 596].

³This has been a very active area of research on the past three decades. Some of the most relevant works in this regard are [29–66].

⁴For the purpose of our treatment, the most important works by these authors are the following: [68–80, 82–89, 91–103].

⁵See for instance [112, 122–124, 126–128, 130, 132, 133, 133, 135, 136, 310, 389].

and Segal, [137]. The main point of their work is that, for a certain class \mathcal{S} theory to be uniquely defined (i.e. for its spectrum of operators to be uniquely determined), one needs to fully specify the source and target 2-categories of the 2D TFT corresponding to the given 4D $\mathcal{N} = 2$ SCFT in question. In technical terms, the objects of the source 2-category are circles, and the 1-morphisms between them are bordisms. The target 2-category is obtained by acting with the functor on these objects and 1-morphisms, resulting in the gauge group of the 4D dual theory, and the bordism operators (also known as the Moore-Segal bordism operators), with the latter being related to algebraic varieties defining moduli spaces in algebraic geometry, i.e. the Higgs and Coulomb branches.

Higgs and Coulomb branches of supersymmetric gauge theories are fascinating objects to explore, since they shed light on dualities characterising the theory they are associated with. From a pure mathematical point of view, they can be realised in terms of the so-called geometric invariant theory (GIT) construction, and quotienting⁶. For the case of 3D $\mathcal{N} = 4$ SCFTs, exchange in between Higgs and Coulomb branches corresponds to a statement of 3D mirror symmetry between two theories (first studied by Intriligator and Seiberg from a pure theoretical-physics point of view, [146]). This 1-to-1 correspondence extends to the calculation of invariants, most importantly the Hilbert Series, accounting for all the gauge-invariant operators of the theory.

However, as also pointed out by Moore, Segal and Tachikawa, their analysis lacked a rigorous mathematical treatment, most importantly for the case in which the target category of the functor is a hyperkähler quotient⁷, which is indeed the case when describing algebraic varieties such as the Higgs branch of a given theory. The work I have conducted so far, provides a step forward towards achieving a rigorous mathematical formulation of such 2-categorical structures for the reason that I will now outline.

As previously explained, TFTs play a crucial role in bridging the gap between QFT and Mathematics⁸. Because of this, the formalism outlined above suggests to always recast a given QFT to a TFT description. For example, 3D $\mathcal{N} = 4$ SCFTs can be turned into 3D TFTs by performing a topological twist. Unlike ordinary 3D TFTs, which are usually trivial in nature in the sense that there are no local degrees of freedom, topologically-twisted TFTs come with nontrivial boundaries, called Lagrangian submanifolds. Importantly, the latter geometrize the boundary conditions of the field content of the theory in question, and, upon studying the intersection (i.e. the 1-morphisms) in between these Lagrangian submanifolds, one can reformulate the statement of 3D mirror symmetry of the parent 3D $\mathcal{N} = 4$ SCFT as that of homological mirror symmetry, which is essentially a duality in between 2-categories. The steps described so far clearly show the advantage of bringing the description down to that the 2-category of Lagrangian submanifolds, which is exactly what we need to keep track of the field content of the theory we started with.

The 3D mirror symmetry/homological mirror symmetry correspondence has been ex-

⁶There is a vast literature in this regard, but we will mostly be focussing on the following works in due course [138, 140, 142–144, 144, 598, 599, 614].

⁷Clarification on the terminology adopted will be provided in Part V of the present work.

⁸For some of the pioneering, as well as most recent advancements in the field, we refer the interested reader to the following works [170–179, 181–186, 188–190, 609].

tensively studied for the specific case in which the correspondence in between the Coulomb and Higgs branch in 3D is 1-to-1, [147, 192, 193]. However, recent developments within the context of supersymmetric quiver gauge theories, mostly led by Hanany and collaborators,⁹ have shown that there are cases in which Higgs branches of fermionic theories can be broken down into several constituents, each one associated with the Coulomb branch of a certain 3D $\mathcal{N} = 4$ SCFT (a.k.a. the cone structure), with the latter being described by the so-called magnetic quivers. Such correspondence was primarily identified by studying the Hilbert series as a geometric invariant. In one of my recent works, [3], I explain how the emergence of this multi-cone structure in the Higgs branch is equivalent to a generalised statement of 3D mirror symmetry, in turn descending from relaxing constraining assumptions on which homological mirror symmetry is based. As a follow up, in [4] and [3], I propose a generalisation of the Moore-Tachikawa varieties for the case in which the target category of the 2D TFT is a hyperkähler quotient. As proved in the former of these two papers, achieving this requires generalising the bordism operators of Moore and Segal to the case involving lack of reparametrisation-invariance on the Riemann surface, ultimately enabling to relate this to the issue of defining a Drinfeld center for composite class \mathcal{S} theories.

So far, my main focus has been on class \mathcal{S} theories, and 3D $\mathcal{N} = 4$ SCFTs, both featuring 8 supercharges. To what extent the generalised 3D mirror symmetry/homological mirror symmetry correspondence can be extended to algebraic varieties associated with theories featuring different amounts of supersymmetries is part of my currently ongoing research. The ultimate aim would be to extend this formalism to quiver gauge theories describing the SM and its possible extensions, eventually shedding light on the mathematical description of SUSY breaking mechanisms leading to observable EFTs.

In summary, the key take-home message of my work is that I have identified an essential question to be addressed, which can simply be stated as follows: generalising the mutual relation between homological mirror symmetry and 3D mirror symmetry, and its impact towards understanding the structure of quantum field theory, as well as the embedding of the SM in String Theory models, by means of categorical algebraic geometry.

1.2 Outline

The concluding part of this first section outlines the roadmap that we will be taking in exploring the topics listed in the preliminary part of the Introduction. Given the richness and breadth of tools encountered in our journey, it would be rather challenging to sketch an accurate sequence of items in a unique diagram. We attempted to provide an oversimplified version of this in figures 1 and 2, but we wish to emphasise that this is by no means exhaustive. The reason for adding these diagrams at this stage is that we believe they might turn out useful to the reader as a guide to refer to when delving in the deeper analysis treated in the forthcoming sections.

As previously explained, our work is motivated by identifying a String Theory model that best suits the description of our world as an EFT. Albeit being a very promising

⁹See for instance [195–200].




Figure 1. Summary diagram of the main components and motivations of this work. With it, we wish to highlight the top-down approach employed in our treatment, baring in mind the diagram on the left hand side of figure 2.

candidate as a unifying theory, String Theory still features facets to be explored. Among the most pressing is its non-perturbative formulation. Notoriously, the latter was among the first motivations of the Holographic Principle, [201], which relates strong and weak coupling limits of two different theories, one living at the boundary of the other. First inspired by black hole mechanics, the Holographic Principle relates the entropy of a certain volume,




Figure 2. The diagram on the left hand side reproduces the top-down route connecting the main ideas underlying our treatment. The diagram on the right hand side, instead, outlines some more specific details of what we will be encountering in the following sections, and is mainly presented here as a guide the reader can refer to when delving in the more detailed analysis to come.

with the number of degrees of freedom living on its boundary. From statistical mechanics, we know that the entropy is nothing but a functional descending from extremisation of the partition function associated to a given theory.

The AdS/CFT correspondence is one of the most notorious realisations of the Holographic Principle, has often proved to be crucial in circumventing issues encountered in analytical calculations. According to the Anti-de Sitter/Conformal Field Theory (AdS/CFT) correspondence, one of the two theories in question is a theory of quantum gravity theory¹⁰. Hence, studying the correspondence is expected to offer a promising way for describing a quantum gravity theory that most faithfully resembles our world. Clearly, the main shortcoming of the $\text{AdS}_5 \times S^5/4\text{D } \mathcal{N} = 4$ SYM correspondence, is that it features five extended spacetime dimensions. This prompted many subsequent formulations, attempting to explain what the correspondence should be when dealing with and AdS_4 spacetime factor, instead. The pioneering works of Aharony, Bergman, Jafferis, and Maldacena (ABJM), showed that such a theory is dual to Chern-Simons (CS) matter theory.

Regardless of the particular string theory model adopted, the vacuum descending from dimensional reduction of the 10-dimensional fermionic string, is an AdS spacetime, with uplifting techniques leading to, either, Minkowski or de Sitter (dS) solutions. From experimental observation, we know our universe is of the dS-type, and we therefore expect to be able to achieve this in a way which is compatible with the Holographic Principle. The only advancements that have been made so far in extending the AdS/CFT correspondence to positive signs of the cosmological constant, have been in lower-dimensional

¹⁰Indeed, in Maldacena's first proposal, this corresponds to type-IIB string theory on an $\text{AdS}_5 \times S^5$ background.




Figure 3. This list succinctly outlines the main achievements of Parts II, III, V of the present work. We kindly suggest the interested reader to refer to it while reading later sections. The important thing we wish to highlight is how all our major results are related to the main example described in the second half of section 3. Please note that Part IV involves a separate treatment, analysing other aspects of lower-dimensional QFTs, mostly related to holographic duality and its generalisation, which are not listed in this table.

setups. Interestingly, it so too happens that proceedings (albeit not yet a full resolution) towards solving the information loss paradox, have also taken place in the setup of lower-dimensional gravity, specifically Jackiw-Teitelboim (JT), and its interplay with QM. The topological nature of JT-gravity makes it a very particular theory, and understanding to what extent its features can be extended to its 4D counterpart is still being investigated.

Nevertheless, JT-gravity is also known to arise as the semiclassical limit of Liouville Theory, which is, in turn, a prototypical example of the bosonic string theory action, as well as being itself the dual of gravity in AdS_3 . One would therefore expect that results achieved in understanding this model from a holographic point of view should, in turn, be able to shed light on higher dimensional setups as well.

As already mentioned, the power of the Holographic Principle is in it being a duality mapping calculations in between different theories. Vacuum decay in between string theory vacua is a non-perturbative effect. Nonperturbative effects in string theory vacua, also known as *instantons*, are defects interpolating between different vacua of the theory in question, and have been amply studied within the context of quantum mechanics (QM), [202], QFT, and their 4D supersymmetric counterparts. Given the lack of a complete

non-perturbative formulation of String Theory, it is reasonable to propose that one might address the same question from the perspective of a non-gravitating theory living on the boundary of the given quantum gravity (QG) model in question.

Furthermore, the AdS/CFT correspondence strongly relies upon the boundary theories being supersymmetric. On the other hand, present-day observations clearly support the statement of SUSY being broken, and we would therefore expect that, if it were possible to assign a holographic dual to our 4-dimensional dS universe, then it must feature a non-supersymmetric QFT on its boundary.

We are therefore drawn to the final standing question: what is a QFT?

Our work provides some further advancements in answering this question. In exploring the importance of 2D structures towards describing higher-dimensional QFTs, we rely upon the joint role played by dualities and categories. We will now briefly explain what we mean by this in this introduction, while referring to the respective sections of this work for a more detailed outline. We believe that any advancement in explaining the decay of String Theory vacua, as well as identifying the most promising String Theory model completion of the SM requires extending a categorical algebraic geometry techniques to QFTs with reducing amounts of supersymmetries.

- The dualities we will be dealing with are holographic dualities (and their generalisations), [203], 3D mirror symmetry, [146], and homological mirror symmetry, [147]¹¹.
- The second crucial ingredient in our treatment are categories. In particular, we will be focussing on the importance of the formalism they provide for describing dualities.

The present work was structured in six parts:

1. At first, we introduce the notion of relative QFTs, with detailed explanation for the case of sigma models, gauge theories, and 6D SCFTs. We then turn to the main case of interest to us, namely Coulomb branches of star-shaped quiver gauge theories, introducing the Kirwan map, and the notion of abelianisation, naturally paving the way to the introduction of category theory, which plays a crucial role in the remainder of our work.
2. Part II introduces the main ingredients of our treatment, namely categories and dualities. After having introduced higher-categorical structures, we specify how they are realised in the main example outlined in Part II thanks to the work of Moore and Tachikawa, [136]. Moreover, category theory is particularly useful for describing dualities, which is another crucial ingredient characterising this thesis. The dualities we encountered are:

¹¹Among the many relevant works on the topic, we mention the following [148–168, 450].

- Homological mirror symmetry.
- 3D mirror symmetry.
- The Alday-Gaiotto-Tachikawa (AGT) correspondence.
- Holographic duality.

Apart from revising well-known results in the literature, we highlight the importance of going beyond full categorical dualisability in the context of Moore-Tachikawa varieties, to which Part V was devoted. In doing so, we relate their formalism to that of SymTFT constructions of Freed, Moore and Teleman, [102], briefly overviewed in Part III.

3. Part III explains how QFTs arise from knot invariants. Specifically, we overview the Symmetry Topological Field Theory (SymTFT) construction of Freed, Moore, and Teleman focussing on the dimensional reduction of non-Lagrangian theories, specifically 6D $\mathcal{N} = (2, 0)$ SCFTs, [108, 109], and the categorical structure of the resulting class \mathcal{S} theories, [1]. Making use of the AGT correspondence, [135], we turn to a description in terms of 2D topological theories defined as functors between 2-categories.

This Part mostly centers around:

- The algebro-geometric formulation of moduli spaces, and their construction by means of the Geometric Invariant Theory (GIT) construction. Importantly, we relate this construction to the abelianisation procedure outlined in our main example in section 3.5, paving the way to further analysis in Part V.
- Turning from QM to Symplectic Geometry by means of the WKB approximation.

Both topics have thoroughly been addressed in the literature, and we therefore devote some sections in the appendix to explaining them in greater depth.

4. Part IV builds from open questions pointed out in the first paper by the author and her supervisor, [7]. In addressing this topic, we restrict to lower-dimensional setups in order to deal with open issues usually arising in higher-dimensional settings, extensively addressed in the literature, [208–210]. In doing so, we built connection with setups involving the emergence of islands, as well as recent developments in the context of von Neumann algebras for calculating generalised entropies in gravitating systems and QFT, [211–217]. The interesting connection with the remainder of our work is twofold: firstly, the need to resort to a certain dual picture in order to interpret the analytic results of the calculations performed; secondly, the richer structure resulting from the lower-dimensional analysis, emphasises the role played by internal structures in higher-dimensional theories, therefore suggesting alternative perspectives might be needed in order to address other open issues in QFT.

5. Part V results from the merge of three single-authored papers, [3–5]. The key findings in this Part are:

- Moore-Tachikawa varieties without dualities are naturally realised in supersymmetric quiver gauge theories, and we therefore put forward a correspondence in between such theories, magnetic quivers, and Drinfeld centers, [3]. As a result of the formalism adopted (namely the definition of Moore-Segal bordism operators, [4, 137]) applied to Moore-Tachikawa varieties without categorical duality, [136], we provide further evidence of our conjectured relation in between homological mirror symmetry and 3D mirror symmetry, first presented in [3].
- The last paper written by the author prior to submitting this Thesis, [?], nicely ties together the topics covered in [3, 4], while also preparing the stage for the future research the author will be conducting as a postdoc¹². In particular, this part takes a more algebro-geometric approach in explaining the mathematical structure underlying supersymmetric quiver gauge theories, with a particular focus on dualities and their mutual relations in terms of higher-categories. As will be explained in depth, the crucial role is played by factorisation homology, [220]. We will not attempt to provide an exhaustive definition of it in this introduction, and, instead, refer the reader to the related section in the core of this work. What we can anticipate, though, is that factorisation homology is related with the counting of ground state degeneracy, and, when applied to the AGT correspondence, provides a tool for evaluating the 4-sphere partition function of class \mathcal{S} theories. Specifically, as we shall see, factorisation homology allows us to relate categorical dualisability and full-extendibility¹³. Within the context of non-Lagrangian theories, such as 6D $\mathcal{N} = (2, 0)$ SCFTs this is particularly relevant, since it introduces the notion of ring objects, of which Hilbert series are one of the examples of most interest to us. As an algebraic variety, we focus on the calculation of the Hilbert series on Coulomb and Higgs branches, with the latter being related by the geometric invariant theory (GIT) quotient construction, [613]. As such, the GIT quotient provides a recipe for realising 3D mirror symmetry; we therefore show how it can be suitably extended for cases where mirror duals are not necessarily in a 1-to-1 correspondence. In relation to [222], this nicely fits with the magnetic quiver prescription, for describing the Higgs branch of certain supersymmetric quiver gauge theories with non-primitive ideals¹⁴. We then show how the 2D description of disk algebras can be suitably adopted for describing Hitchin systems of class \mathcal{S} theories. We concluded outlining future directions that are currently under investigation by the same author.

¹²Partly outlined in [6].

¹³And is therefore in line with the connection between dualities and categorical structures.

¹⁴Namely the 2-cone structure, [223].

6. Part VI closes outlining the main achievements of the present work, with an outlook on some key questions the author wishes to address in her future research.

We believe our treatment opens the way to further investigating the mutual relation in between homological mirror symmetry and 3D mirror symmetry with the ultimate aim of aiding at understanding the underlying mathematical structure of QFT. These are currently under investigation by the same author, and we plan to report about any advancement in the field in the near future.

2 Roadmap

2.1 Roadmap

This first section is devoted to outlining the path we will be adopting for addressing some of the issues mentioned in the Introduction. The diagram of figure 2.1 simplifies this, and we will start addressing the topics outlined there starting from section 3, continuing in an upcoming work [597]. For those who are already familiar with the terminology adopted in figure 2.1, the first few sections of this work are purely revision. However, they might still find some interesting connections, which, to the best of our knowledge, haven't been spelled out in the literature with the same emphasis and perspective.




Figure 4. Roadmap for the present and upcoming work.

Essentially, Part I can be thought of as an extremely concise summary of the basic knowledge gathered by a mathematical physicist in the attempt to explain a possible

roadmap towards deepening the understanding of some major open questions in Theoretical Particle Physics from a Mathematician’s point of view.

As an advanced notice, we advise the reader not to be misguided by the oversimplified explanation of the first few pages of this work: technicalities will soon arise.

In this preliminary section, we introduce the unfamiliar reader to:

1. The basic building blocks of QFT¹⁵.
2. Some key theoretical tools adopted by Phenomenologists for analysing them.

2.1.1 Fields and Interactions

Quantum fields are mutually-interacting entities exhibit wave-particle duality. QFT is arguably the most successful scientific theory so far, with the Standard Model (SM) of Particle Physics being a prototypical example. In some cases, QFT predicts to great accuracy observational outcomes, and has also been used in the context of pure mathematics.

With the advent of the quantum revolution, fields were promoted to being fundamental constituents, whereas all fundamental particles were understood to arise as a quantum mechanical (QM) effect. This extends to all particles known so far, namely quarks, leptons, gluons, W- and Z-bosons, and the Higgs boson.

The SM is an example of a QFT containing all the particles known up to now, with their associated three force fields¹⁶ Extensions of the SM, referred to as Beyond the Standard Model (BSM) setups, are currently an active area of research, providing extensions of the SM by adding interactions and/or particles.

A given QFT is defined in terms of a Lagrangian density, which, upon integrated over spacetime, defines the action, S , which is, essentially, a functional of the fields. The Variational Principle constraints the action in such a way that its extremisation leads to the equations of motion of the respective fields. Importantly, the action is needed for defining the partition function, i.e. the expectation value of the identity, as well as any other expectation value of operators¹⁷ accounted for by a given effective field theory (EFT).

2.1.2 Amplitudes and Hilbert series

Amplitudes

Scattering amplitudes are among the most important tools in Particle Physics. They are the building blocks for defining transition amplitudes of a certain set of states to another. In the context of collider physics, they are an essential theoretical background tool for predicting and analysing particle collisions and decays. Importantly, if new physics were to be discovered, it must be signalled by a mismatch in between experimental outcomes and scattering calculations, that could either feature as the detection of new particles, and/or the identification of a new interaction (and therefore a new fundamental force).

¹⁵We strongly suggest the interested reader to refer to the work of pioneers in the field for a complete treatment of QFT, specifically [105–107].

¹⁶The matter fields comprised in it come in 3 families, plus the Higgs boson.

¹⁷The latter are also known as correlation functions.




Figure 5. A simplified depiction of a scattering process. The S -matrix defines the amplitude from the incoming states (black lines on the left) to the outgoing states (black lines on the right). The shaded circle in the middle encodes the information regarding the interactions and operator content characterising a certain EFT.

In a nutshell, if our theoretical knowledge of particle physics were complete, we should be able to explain all experimental outcomes, recasting them to the Lagrangian theories at hand. However, it so happens that there are some crucial unanswered questions, that clearly signal the need for a deeper mathematical understanding of the models developed so far. Those that motivated our work in the first place are the following:

1. The mass hierarchy problem.
2. The lightness of the Higgs mass.

To help the unfamiliar reader, we will come back to explaining these issues in section 2.2.

Hilbert series and effective operators

The Hilbert series, HS, is defined as follows

$$\text{HS}[\phi] = \prod_{i=1}^n \int_{\mathcal{G}_i} d\mu_i \text{PE}[\varphi, R] \quad (2.1)$$

where \mathcal{G}_i denotes a given gauge group, φ is a *spurion* variable¹⁸, $d\mu_i$ denotes the Haar measure, and $\text{PE}[\varphi, R]$ the *plethystic exponential*, which, in turn, is defined as follows¹⁹,

$$\text{PE}[\phi, R] = \exp \left[\sum_{r=1}^{\infty} \frac{\phi^r \chi_R(z_j^r)}{r} \right], \quad \text{PE}[\psi, R] = \exp \left[\sum_{r=1}^{\infty} (-1)^{r+1} \frac{\psi^r \chi_R(z_j^r)}{r} \right] \quad (2.2)$$

for bosons and fermions, respectively²⁰. We will be entering the details of the calculation of (2.1) in an upcoming work, [597]. For the purpose of this introductory work, it is enough to know that HS is constructed in terms of the degrees of freedom (namely the particle content) and their transformation properties under the symmetries of a given theory. This is enough to say that, in order to build the Hilbert series, we need to know what the complete

¹⁸A spurion could either be a scalar, a fermion, or a gauge field.

¹⁹See for instance [606].

²⁰ χ here denotes the character, and depends in the representation, R .

set of operators of a given theory is. Completeness of the spectrum of operators is a major sought-after objective in particle physics, since it paves the way towards identifying possible extensions of the fundamental laws of nature. Given the relation in between scattering amplitudes and HS, a mismatch with predictions for a given theory signal the need to extend the effective field theory of reference. If the theory in question is taken to be the Standard Model (SM) of Particle Physics, its resulting extensions are referred to as Beyond the Standard Model (BSM) effective field theories, [604–606].

2.2 (Some) Open Questions in Quantum Field Theory

Albeit having been successfully verified experimentally, the SM of Particle Physics is clearly incomplete. The Higgs boson, [611, 612], responsible for the mass of fundamental particles, was the last piece of a jigsaw to be found, yet the first of a more complicated one. The Higgs boson is a scalar responsible for the mass gain of fundamental particles, via the so-called Higgs mechanism²¹. Undoubtedly one of the major theoretical physics achievements of the past century, the model is clearly incomplete. The two major issues we indicate are the mass hierarchy problem and the lightness of the Higgs boson mass. Each one of them is the object of active investigation by world-leading experts in the field, and by no means we are attempting to provide a full explanation of either of them in our treatment. Nevertheless, we consider them as part of the motivations leading to the need for furthering the understanding of the underlying mathematical structure of QFT.

The Mass Hierarchy Problem

Essentially, this is a statement regarding the unexplainable discrepancy between the energy scales at which the fundamental interactions decouple. A successful unifying theory should be able to explain why this symmetry breaking pattern.

Lightness of the Higgs Boson and Supersymmetry

Another open question highlighting the fact that the SM is still incomplete is that the Higgs mass is too light. Given that it is the only spinless particle in the SM, theorems imply it should be extremely massive. One of the possible reasons explaining the very light mass of the Higgs was thought to be Supersymmetry (SUSY), proposing that, in order to compensate the very light mass observed, new particles (referred to as superpartners) should be detected at the Large Hadron Collider (LHC). Up to now, though, these superpartners haven't been observed by collider experiments at the energies predicted by the theory. This inevitably leads to the question, to what extent is SUSY able to provide a consistent theory of everything, given that String Theory so strongly relies upon it?

We believe that SUSY and String Theory are very promising candidates for a unifying theory of everything, but, in order to overcome the apparent shortcomings listed above, we need to address the deep question highlighted in the introduction, namely: "What is (the mathematical structure of) a QFT?". The present work is meant to provide further

²¹We refer the interested reader to the original work by Professor Peter Higgs for a detailed explanation of the process, [611, 612].

insights in addressing this question, by making use of mathematical advancements in the field of categorical algebraic geometry. Our findings shed light on a new perspective that is proposed to be a promising alternative top-down construction of QFTs. The ultimate aim would be that of identifying the mathematical structure underlying the SM, and, with it, its embedding in the most suitable string theory vacuum possible, to which we plan to report in upcoming work [597].

Albeit being among the most successful accomplishments in theoretical physics, QFT, in its present-day formulation, is mathematically incomplete. To see why this is the case, let us take a step back, explaining a few preliminary features of the theory, and where the issues arise, [204, 205].

In a QFT, at every point in spacetime one can think of there being an operator, i.e. an infinite-dimensional matrix, acting on a Hilbert space. The mathematical issue arises when attempting to reconcile this picture with the continuum of spacetime to describe our universe. In the quantum picture, the continuum limit is expected to arise upon bringing any two points infinitesimally-close to each other. From a computational perspective, though, this gives rise to divergences, that can only be overcome by supposing spacetime is discretised. In mathematical language, this discretisation turns spacetime into a *lattice*, thereby enabling to keep a finite distance in between any two points in spacetime. However, according to the Nielsen-Ninomiya Theorem, this does not hold for any QFT, and, in particular, it doesn't hold for the SM.

The great advantage of lattice field theories, is that they enable to have very good mathematical and analytic control, meaning that it is practically impossible for them to describe inconsistent theories. It is therefore tantalising to believe that, if we were able to place the SM on a lattice, we might be able to identify what it is that we are currently missing of QFT as it stands for it to be regarded to as a mathematically rigorous theory.

The reason why the SM cannot (yet) be reconciled with the lattice description is the fact that it is practically impossible to express parity violation in a mathematically rigorous way. In QFT, this is overcome by imposing the so-called *anomaly cancellation*. Importantly, the latter are closely related to the topology of spacetime, and therefore can only arise in the continuum²².

Another open question highlighting the fact that the SM is still incomplete is that the Higgs mass is too light. Given that it is the only spinless particle in the SM, theorems imply it should be extremely massive. One of the possible reasons explaining the very light mass of the Higgs was thought to be Supersymmetry, proposing that, in order to compensate the very light mass observed, new particles should be detected at the LHC. Up to now, the predicted particles still haven't been observed by collider experiments at the energies predicted by the theory. This inevitably leads to the question, to what extent is SUSY able to provide a consistent theory of everything, given that String Theory so strongly relies upon it?

We believe that SUSY and String Theory are very promising candidates for a unifying theory of everything, but, in order to overcome the apparent shortcomings listed above,

²²For an alternative prescription, see [622].

we need to address the deep question highlighted in the introduction, namely: "What is (the mathematical structure of) a QFT?". The present work is meant to provide further insights in addressing this question, by making use of mathematical advancements in the field of higher-categorical structures. Our findings shed light on a new perspective that is proposed to be a promising alternative top-down construction of QFTs. The ultimate aim would be that of identifying the mathematical structure underlying the SM, and, with it, its embedding in the most suitable string theory vacuum possible.

2.3 Where to next?

Key advancements towards achieving a rigorous mathematical formulation of QFTs strongly relies upon a higher-categorical prescription. In particular, the symmetries of a QFT can be thought of as defects living in different categories connected by functors.

Overall, the present work clearly shows the richness and in-depth understanding that underlying mathematical structures and dualities are able to provide in, both, gravitating systems and QFTs alike.

Part of this work explores the role played by black holes in transitions involving 2D theories of gravity. In particular, we build connection with the von Neumann algebras, recently put forward by Witten for calculating the entropy of states in QFT and gravity. We therefore show how generalisations of the holographic correspondence, and quantum information tools enable to overcome issues encountered in higher-dimensional setups.

The complementary part of our treatment deals with interesting cases where lower-dimensional descriptions of higher-dimensional theories are needed. Focussing on dimensional reduction of non-Lagrangian theories, specifically 6D $\mathcal{N} = (2, 0)$ SCFTs on the Riemann surface, and exploiting the Alday-Gaiotto-Tachikawa correspondence, we map the 4D description of class \mathcal{S} theories to their 2D TFT duals. Viewing the 2D theory as a functor in between 2-categories, we propose a generalisation of Moore-Segal bordism operators as needed for describing Moore-Tachikawa varieties for the case not involving categorical duality, ultimately relating this to joined systems of class \mathcal{S} theories joined by non-invertible (categorical) defects. By means of algebro-geometric tools, we show how this formalism nicely reconciles with the calculation of Hilbert series for theories admitting a magnetic quiver description. As a byproduct of the tools adopted, our findings provide further evidence of a generalised correspondence in between homological mirror symmetry and 3D mirror symmetry.

In our analysis, we encountered two major setups built upon the notion of higher-categories, namely symmetry TFTs (SymTFT) and topological orders (TO). Plenty of effort has been made towards building a correspondence in between the two, most recently in [376]. Our findings provide further support towards strengthening the connection between the two description.

En-passing, we commented on a proposal regarding the realisation of such composite absolute theories and the definition of a fiber functor intrinsically related to the notion of a partition function for a 3D theory rather than a 4D theory.

We conclude by stressing that our analysis is mostly motivated by furthering the understanding of 6D $\mathcal{N} = (2, 0)$ SCFTs, from, both, a mathematical and physical point of

view. To what extent these findings might be mapped to other setups is currently under investigation.

3 Relative QFTs

The present section consists of two parts:

1. At first, we introduce the notion of relative QFT, and how this applies to sigma models, gauge theories, and 6D $\mathcal{N} = (2, 0)$ SCFTs. This is essentially a brief overview of [103].
2. Then, we turn to our main example in this context, namely Coulomb branches of star-shaped quivers²³, which will be useful to keep as a reference for later sections and Parts of this work. Most importantly, we introduce the notion of *abelianisation*, which will be of great importance throughout our treatment. As we shall see, this naturally paves the way towards introducing category theory tools.

3.1 Rings, homologies, cohomologies, and modules

In this section we start by introducing some preliminary notation that will be needed for the remainder of our treatment. Most of this is well-known to experts in the field, but we report the main definition just for completeness.

Rings are algebraic structures generalising fields, where multiplication need not be commutative, and inverses might not exist. It is a set endowed with two binary operations called addition and multiplication, such that the ring is an abelian group with respect to the addition operation, and has a multiplicative identity element.

Homologies were originally defined in Algebraic Topology, as a general way of associating a sequence of algebraic objects, such as abelian groups or modules, with other mathematical objects, such as topological spaces.

Modules generalise the notion of vector space, in which the field of scalars is replaced by a ring. A module also generalises the notion of abelian groups, since the latter are modules over the ring of integers. Modules are closely related to representation theory of groups, and are also widely used in homological algebra, Algebraic Geometry, and Algebraic Topology.

3.2 Abstract topological symmetry data in field theory

To a given QFT, F , fixing $N \in \mathbb{Z}^{\geq 0}$, one can assign topological data²⁴, called quiche, namely a pair (σ, ρ) in which

$$\sigma : \text{Bord}_{N+1}(F) \rightarrow \mathcal{C} \tag{3.1}$$

²³This terminology will be explained in due course.

²⁴For a brief overview of topological field theories (TFTs), we suggest the interested reader to appendix G, which is mostly overview of well-known background material.

is an $N + 1$ -dimensional TFT and ρ is a right topological σ -module. The quiche is N -dimensional, hence it shares the same dimensionality as the theory on which it acts. Let F be an N -dimensional field theory. A (σ, ρ) -*module structure* on F is a pair (\tilde{F}, θ) , in which \tilde{F} is a left σ -module and θ is an isomorphism

$$\theta : \rho \otimes_{\sigma} \tilde{F} \xrightarrow{\cong} F \quad (3.2)$$

of absolute N -dimensional theories. The LHS of (9.1) defines the dimensional reduction, i.e. the sandwich we were referring to earlier. σ needs only be a *once-categorified* N -dimensional theory, whereas ρ and \tilde{F} are relative theories.

Examples of relative theories are, [103]:

1. 2D chiral WZW CFTs based on a compact Lie group G and level k , w.r.t. 3D topological CS theory associated to the (G, k) -data of its 2D boundary.
2. Gauge theories.
3. 6D $\mathcal{N}=(2,0)$ SCFTs w.r.t. 7D TQFT in the bulk.

More generally, *relative theories* are models whose fields form a fibration

$$\mathcal{F} \longrightarrow \mathcal{M} \xrightarrow{\pi} \mathcal{B} \quad (3.3)$$

where \mathcal{B} and \mathcal{F} denote the base and the fiber, respectively. Relative fields are the fibers of π . The path integral over the base reduces a finite sum. For the theory to be absolute, the same should hold for \mathcal{F} as well.

As originally outlined in [103], QFT F relative to α is a homomorphism

$$\tilde{F} : \tau_{\leq n} \alpha \rightarrow \mathbf{1} \quad (3.4)$$

where *alpha* is the bulk theory w.r.t. F .

3.2.1 Fiberings over BG

Symmetries in field theory are expressed in terms of topological defects acting as operators on state spaces of the theory in question. The idea at the heart of this are the notions of abstract groups and abstract algebras. In their work, [102], Freed, Moore and Teleman propose an abstract symmetry structure in the context of field theory and its concrete realisation.

A field theory is analogous to a linear representation of a Lie group or a module over an algebra. Hence their definition of the boundary theory as a module. The essential content of their definition is that of expressing a QFT, F , as a sandwich $\rho \otimes_{\sigma} \tilde{F}$, with ρ a regular Dirichlet BC (right boundary theory for σ) and \tilde{F} the left boundary theory for σ (typically not topological).

Altogether, (σ, ρ) constitutes the n -dimensional *quiche*, defining the abstract symmetry structure of a given QFT. Defects away from \tilde{F} are topological and belong to (σ, ρ) . (σ, ρ) -defects are analogous to elements of an abstract algebra (connection with GKS). The

novelty of the treatment by Freed, Moore and Teleman is the fact that the sandwich representation develops a calculus of topological defects acting on a QFT (based on fully local TFT). The TFT imposes strong finiteness constraints on the QFT.

Let G be a finite group. A classifying space BG is derived from a contractible topological space EG equipped with a free G -action by taking the quotient. The homotopy type of BG is independent of choices. If X is a topological space equipped with a G -action, then the Borel construction, [623] is the total space of the fiber bundle.

$$\begin{array}{ccc}
X_G & \equiv & EG \times_G X \\
\downarrow \pi & & \\
& & BG
\end{array} \tag{3.5}$$

with fiber X . If $* \in BG$ is a point, and we choose a basepoint in the G -orbit in EG labelled by $*$, then the fiber $\pi^{-1}(*)$ is canonically identified with X . The abstract group symmetry data is the pair $(BG, *)$, and a realisation of the symmetry $(BG, *)$ on X is a fiber bundle over BG together with an identification of the fiber over $* \in BG$ with X .

3.3 Preliminary examples: the σ -model and gauge theories

In light of the techniques introduced in the following Parts of this work, we now introduce the notion of moduli space, as denoting the minima of the theory in question, namely the least energy field configurations²⁵. Albeit the sample theories presented in this section are quite trivial, they are sufficient for building intuition for what lies ahead in our treatment.

1) Sigma models

Data: G a finite group and M a smooth manifold with a free left-acting G and quotient \bar{M} (equivalently the gauged σ model on M).

Denoting with $BG(X)$ the collection of principal G -bundles over X ,

A principal G -bundle is a covering space $P \rightarrow X$ and a free- G action on P such that $P \rightarrow X$ is a quotient map for the G -action. We denote the collection of principal G -bundles over X with $BG(X)$, with the latter being a groupoid, and, as such, is a category in which every morphism is invertible.

A *symmetry*

$$\varphi : (P \rightarrow X) \rightarrow (P' \rightarrow X) \tag{3.6}$$

is a diffeomorphism

$$\varphi : P \rightarrow P' \tag{3.7}$$

²⁵For a brief overview of moduli spaces in some basic examples, we refer the interested reader to appendix [K](#).

commuting with G and covering id_X . The automorphism group of $P \rightarrow X$ is the group of gauge transformations.

The path-integral over $BG(Y)$ is an integral over the equivalence classes of G -bundles.

Fixing $(P \rightarrow X) \in BG(X)$, a relative field over $(P \rightarrow X)$ is a pair (f, θ) consisting of a smooth map $f : X \rightarrow \bar{M}$ and an isomorphism $\theta : (P \rightarrow X) \rightarrow f^*(M \rightarrow \bar{M})$ of G -bundles over X . Given that relative fields feature no automorphisms, they are *rigid*, and therefore constitute a space rather than a category.

Given a pair of theories, (α, F) , with α topological and defined on all manifolds, and F a Riemannian theory, using the knowledge of the σ -model F to predict the structure of α .

The relative path-integral $F(X)$ on a closed D -dimensional manifold is an integral over relative fields, hence a function

$$F(X) : BG(X) \rightarrow \mathbb{C} \quad (3.8)$$

As such, $F(X)$ is invariant under symmetries of $BG(X)$, and is therefore expressed as a function on equivalence classes, $H^1(X; G)$, with the latter denoting the isomorphic classes of principal G -bundles over X . From this, (3.8) can be reexpressed as

$$F(X) : H^1(X; G) \rightarrow \mathbb{C} \quad (3.9)$$

If F is to be a QFT relative to a $D + 1$ -dimensional bulk α , in the sense that

$$F(X) : \mathbb{C} \rightarrow \alpha(X) \quad (3.10)$$

then, $F(X)$ can be identified as an element of the vector space $\alpha(X)$, leading to the conclusion that $\alpha(X)$ is the free vector space on the finite set $H^1(X; G)$.

Next, considering a closed $D - 1$ -dimensional manifold, Y , the relative canonical quantisation of $F(Y)$ is obtained by performing the quantisation on the fibers of the map

$$\text{Map}(Y, \bar{M}) \rightarrow BG(Y) \quad (3.11)$$

such that the vector bundle is

$$F(Y) \rightarrow BG(Y) \quad (3.12)$$

The bundles are grouped into isomorphism classes. Points featuring in each isomorphism class denote the G -bundles $Q \rightarrow Y$, and the arrows the isomorphisms between the G -bundles. Selecting a representative bundle, $Q \rightarrow Y$, in each isomorphism class, the groupoid $BG(Y)$ corresponds to a simpler groupoid with a finite set of $H^1(Y; G)$ and no morphisms between them.

If $m \in H^1(Y; G)$ is the class of a G -bundle $Q \rightarrow Y$, then the automorphism group of m is the group of gauge transformations of $Q \rightarrow Y$, and the equivariant bundle, (3.12), decomposes into topological vector spaces indexed by pairs (m, e) in which $m \in H^1(Y; G)$ and e is a complex irreducible representation of the automorphism group of m . The partition function on Y




Figure 6. A depiction of the vector bundle.

$$F(Y) : \mathbf{1}(Y) \rightarrow \alpha(Y) \quad (3.13)$$

may be identified with an object belonging to the linear category $\alpha(Y)$, leading to the prediction that $\alpha(Y)$ is the free Vect_{top} -module with basis pairs (m, e) .

For the case in which X is a 1D theory, namely $X \equiv S^1$, $H^1(S^1; G)$ is the set of *conjugacy classes* in G . If $Y \equiv \text{pt}$, then $H^1(\text{pt}; G)$ has a single element, representing the trivial bundle with automorphism group G . Consequently, $\alpha(\text{pt})$ defines the category of representations of G .

In this case, α is the finite 2D gauge theory with gauge group G , and can be defined on any manifold of $D \leq 2$ by a finite path-integral. Given that the theory has a vanishing Lagrangian, the partition function on a closed 2D manifold, M_2 , reduces to the weighted counting of G -bundles

$$\alpha(M_2) = \sum_{[R \rightarrow M_2]} \frac{1}{|\text{Aut}(R \rightarrow M_2)|} \quad (3.14)$$

The finite path integral over $X \equiv S^1$ is the space of invariant sections of the trivial equivariant line bundle $BG(S^1)$. On $Y \equiv \text{pt}$, instead, the exponential of the action has constant value in the linear category Vect_{top} , hence the path integral defines the subcategory of invariants under G , namely the category of representations of G .

From relative to absolute

An *absolute* theory can be recovered from a relative one under gauging by a finite subgroup $G' \subset G$. In such case, the σ -model on a closed D -dimensional manifold can be expressed in terms of the relative theory as follows

$$f_{G'}(X) = \sum_{m' \in H^1(X; G')} \frac{1}{|Z_{G'}(m')|} F(X; m') \quad (3.15)$$

where the sum runs over the equivalence classes of G' -bundles $P \rightarrow X$, and $Z_{G'}(m')$ denotes the automorphism group of a representative of the equivalence class m' .

Following similar arguments, the quantum topological vector space on a closed $D - 1$ -dimensional manifold Y can be expressed as

$$f_{G'}(Y) = \bigoplus_{m' \equiv [Q' \rightarrow Y] \in H^1(X; G')} F(Y; Q \rightarrow Y)^{\text{Aut}(Q' \rightarrow Y)} \quad (3.16)$$

2) Gauge theories

A very similar analysis holds for the case of gauge theories, with all the fields' categorical levels enhanced by 1 unit, due to the presence of an additional symmetry.

In this case, the theory is specified by the following data: a Lie algebra, \mathfrak{g} , a covering homomorphism $G \rightarrow \tilde{G}$ of compact connected Lie groups with kernel G_o a finite central abelian subgroup of G . For example, if $G_o \subset G$ and $\tilde{G} \equiv G \setminus G_o$ is the adjoint group, this gives a canonical choice of data associated to a compact semisimple Lie algebra.

Assuming $\tilde{P} \rightarrow X$ denotes a principal \tilde{G} bundle, the obstruction to lifting to a principal G -bundle is measured by a G_o -gerbe, $\mathcal{G}(\tilde{P}) \rightarrow X$, and with $B^2 G_o(X)$ the collection of G_o -gerbes over X . $B^2 G_o(X)$ is a 1-category. Its equivalence classes constitute the cohomology group $H^2(X; G_o)$. The group of equivalence classes of automorphisms between objects is $H^1(X; G_o)$, and the group of automorphisms of automorphisms is $H^0(X; G_o)$.

For gauge theories, a relative field can be defined as $(\mathcal{G} \rightarrow X) \in B^2 G_o(X)$. A relative field over $(\mathcal{G} \rightarrow X)$ is a pair $(\tilde{\Theta}, \theta)$ consisting of a \tilde{G} -connection $\tilde{\Theta}$ and an isomorphism $\theta : \mathcal{G} \rightarrow \mathcal{G}(\tilde{\Theta})$ of G_o -gerbes. $(\tilde{\Theta}, \theta)$ form an ordinary grupoid, with no automorphisms between automorphisms. The fields are \tilde{G} -connections with G -gauge transformations.

The quantum theories (α, F) that can be built from these fields are a $D+1$ dimensional topological theory and a D -dimensional relative theory F . α is now defined as a path-integral over the G_o -gerbes, specifically

$$\alpha(M_2) \equiv \frac{|H^2(X; G_o)| |H^0(X; G_o)|}{|H^1(X; G_o)|} \quad (3.17)$$

Consequently, $\alpha(X)$ is the vector space of complex-valued functions on $B^2(X)$. The relative theory F is an element of $\alpha(X)$ since

$$F(X) \equiv H^2(X; G_o) \rightarrow \mathbb{C} \quad (3.18)$$

$\alpha(Y)$, instead, defines the linear category of vector bundles over $B^2 G_o(Y)$. The relative theory F determines a particular vector bundle $F(Y) \rightarrow B^2 G_o(Y)$.

For any finite abelian group AA , its Pontryagin dual is defined as follows

$$A^\vee \stackrel{\text{def.}}{=} \text{Hom}(A, G_o) \quad (3.19)$$

Once having chosen the basepoints, $F(Y) \rightarrow B^2 G_o(Y)$ determines the topological vector spaces $F(Y; m, e)$ for $m \in H^2(Y; G_o), e \in H^1(Y; G_o)^\vee$, where (m, e) denote the classes of discrete magnetic and electric fluxes.

From relative to absolute

Once more, an absolute gauge theory can be obtained by gauging by a subgroup of G , leading to a relic gauge group $G \setminus G'_o$, $G'_o \subset G_o$, and its expression reads

$$f_{G'_o}(X) = \sum_{m' \in H^2(X; G_o)} \frac{|H^0(X; G'_o)|}{|H^1(X; G'_o)|} F(X; m'). \quad (3.20)$$




Figure 7. Specifying the topological boundary conditions turns a relative QFT into an absolute one.

3.4 Theory X

We now turn to the case of 6D $\mathcal{N} = (2, 0)$ SCFTs. The data defining the theory in question are:

1. A reductive Lie algebra, \mathfrak{g} , with an invariant inner product, \langle, \rangle , such that all coroots have square of length 2. A reductive real Lie algebra is defined as the direct sum

$$\mathfrak{g} = \mathfrak{z} \oplus \mathfrak{g}', \quad (3.21)$$

where \mathfrak{z} denotes its center, and $\mathfrak{g}' \equiv [\mathfrak{g}, \mathfrak{g}]$ its semisimple subalgebra. A Cartan subalgebra $\mathfrak{h}' \subset \mathfrak{g}'$ determines the coroot lattice, which is a full sublattice $\Gamma' \subset \mathfrak{h}'$.

2. A full lattice $\Gamma \supset \Gamma'$ in $\mathfrak{h} = \mathfrak{z} \oplus \mathfrak{h}'$, such that the inner product is integral and even on Γ .

Given this data, one can define $\Lambda \supset \Gamma$ as the dual lattice to Γ in \mathfrak{h} , with Λ being the subset of vectors $\eta \in \mathfrak{h}$ such that

$$\langle \eta, \Gamma \rangle \subset \mathbb{Z}. \quad (3.22)$$

Then, the quotient

$$\pi \stackrel{def.}{=} \Lambda/\Gamma \tag{3.23}$$

is a finite abelian group equipped with a perfect pairing

$$\pi \times \pi \rightarrow \mathbb{Q}/\mathbb{Z} \subset \mathbb{T} \tag{3.24}$$

induced from the inner product. This pairing induces an isomorphism

$$\pi \simeq \pi^\vee, \tag{3.25}$$

implying π is Pontrjagin self-dual.

The expectation outlined by Freed and Teleman on the basis of Witten's pioneering work, is therefore that, given

$$(\mathfrak{g}, \langle, \rangle, \Gamma), \tag{3.26}$$


there exists a finite 7D TQFT

$$\alpha_{\mathfrak{g}} \stackrel{def.}{=} \alpha(\mathfrak{g}, \langle, \rangle, \Gamma), \tag{3.27}$$

and a 6D QFT

$$X_{\mathfrak{g}} \stackrel{def.}{=} X_{(\mathfrak{g}, \langle, \rangle, \Gamma)}, \tag{3.28}$$

relative to $\alpha_{\mathfrak{g}}$, with $\alpha_{\mathfrak{g}}$ a bordism mutli-category of manifolds with unspecified tangential structure. This theory is constitutes a 7D analogue of CS theory for torus groups. On the other hand, the 6D theory, $X_{\mathfrak{g}}$, is meant to be a geometric bordism category with a conformal structure.



$X_{\mathfrak{g}}$

Figure 8. Theory X , here denoted $X_{\mathfrak{g}}$, lives on the boundary of a 7D bulk TQFT, $\alpha_{\mathfrak{g}}$. No topological boundary theory can be defined to make the theory absolute.

For X a closed oriented 6-manifold, the cup product and pairing (3.24) combine to give a non-degenerate skew-symmetric pairing

$$H^3(X; \pi) \times H^3(X; \pi) \rightarrow \mathbb{T} \quad (3.29)$$

on the finite group $H^3(X; \pi)$. Upon taking Y to be an oriented a 5-dimensional manifold, $H^3(Y; \pi)$ defines the group of self-dual fluxes, and, from the pairing and Poincaré-Pontrjagin duality, the following isomorphism follows

$$H^3(Y; \pi) \simeq H^3(Y; \pi^\vee) \simeq H^2(Y; \pi)^\vee, \quad (3.30)$$

where self-duality identifies the electric and magnetic groups. Then, the linear category $\alpha_{\mathfrak{g}}$ is the free Vect_{top} -module with basis $H^3(Y; \pi)$, and, to each self-dual flux, $\sigma \in H^3(Y; \pi)$, should correspond a quantum topological vector space, $X_{\mathfrak{g}}(Y; \sigma)$.

3.5 Main Example: the Coulomb branch chiral algebra

An example of (3.5) is

$$\begin{array}{ccc} \mathcal{M}_C & \longleftarrow & T_C^\vee \\ & & \downarrow \pi \\ & & \mathfrak{t}_C / W \end{array} \quad (3.31)$$

where \mathfrak{t}_C denotes the complexified Cartan subalgebra of G , $\mathfrak{t}_C \subset \mathfrak{g}_C$, W is the Weyl group, and T_C^\vee is the complexified dual of the maximal torus. $T_C^\vee \simeq \mathbb{C}^{\text{rank}(G)}$ is a holomorphic Lagrangian torus with respect to the holomorphic symplectic structure. The base \mathfrak{t}_C / W is parametrised by the diagonal expectation value of a complex scalar vectormultiplet, [104],

$$\varphi \in \mathfrak{t}_C \subset \mathfrak{g}_C. \quad (3.32)$$

Global coordinates on the base are given by polynomials in φ which are gauge-invariant operators of a non-abelian theory. φ is referred to as a *generic* point of the Coulomb branch if:

1. It fully breaks gauge symmetry to the torus (practically turning the W-boson massive),

$$M_\alpha \equiv \langle \alpha, \varphi \rangle \neq 0, \quad (3.33)$$

for all α , roots of G .

2. Giving a non-zero effective mass to every hypermultiplet,

$$M_\lambda \equiv \langle \lambda, \varphi \rangle \neq 0, \quad (3.34)$$

for all λ , weights of R .

Hence, a generic point of \mathfrak{t}_C lies in the complement of all weight and root hyperplanes, [104].

\mathcal{M}_C is a complex symplectic manifold, on which one can calculate the chiral ring $\mathbb{C}[\mathcal{M}_C]$ and its Poisson structure. The chiral-ring operators lie in the cohomology of a topological supercharge, and therefore lie within the topological subsector of the 3D gauge theory in question. Consequently, the product of chiral operators is topologically protected and can be calculated by means of TQFT techniques.

The chiral algebra \mathcal{A} can be defined as the local chiral ring in the neighbourhood of a generic point φ on the base of the Coulomb branch when W-bosons and hypermultiplets become massless,

$$\mathcal{M}_C^{\text{abel.}} \stackrel{\text{def.}}{=} \pi^{-1}((\mathfrak{t}_C \setminus \Delta \cup \Delta_R)/W) \subset \mathcal{M}_C \quad (3.35)$$

where

$$\Delta \stackrel{\text{def.}}{=} \bigcup_{\text{roots } \alpha} \{M_\alpha(\varphi) = 0\} \subset \mathfrak{t}_C \quad , \quad \Delta_R \stackrel{\text{def.}}{=} \bigcup_{\text{weights } \lambda} \{M_\lambda(\varphi) = 0\} \subset \mathfrak{t}_C. \quad (3.36)$$

From this, it follows that the algebra \mathcal{A} is defined from the covering of (3.35) as

$$\mathcal{A} \stackrel{\text{def.}}{=} \mathbb{C}[\tilde{\mathcal{M}}_C^{\text{abel}}] \quad (3.37)$$

from which it follows that

$$\mathbb{C}[\mathcal{M}_C] \hookrightarrow \mathcal{A}. \quad (3.38)$$

The generators of \mathcal{A} come in two main types, [104]:

1. Polynomials in $\varphi, M_\alpha^{-1}, M_\lambda^{-1}$.
2. Abelian monopole operators for any cocharacter $A \in \text{Hom}(U(1), T) \simeq \mathbb{Z}^{\text{rank}(G)}$.

Gathering these properties together, the abelianised embedding of the chiral ring operators can succinctly be denoted as follows

$$\mathcal{A} = \mathbb{C} \left[\varphi, \{M_\alpha^{-1}\}, \{M_\lambda^{-1}\}, \{v_A\} \right] / \text{rel.} \quad (3.39)$$

3.5.1 Star-shaped quivers

The Coulomb branch

For the case of interest to us, we focus on star-shaped quivers, conventionally denoted by $\mathcal{T}_{N,\mathbf{k}}$, where N dictates the number of gauge nodes in the quiver, and \mathbf{k} the number of legs. To see how we have outlined above practically leads to the chiral ring defining the Coulomb branch, lets assume that $N = 2$ and $\mathbf{k} = 3$. In this case the Coulomb branch is flat

$$\mathcal{M}_C \simeq \mathbb{C}^8. \quad (3.40)$$

This is dual to a 4D class \mathcal{S} theory with A_1 structure, namely the Grassmannian. The gauge group of the theory in question, $\mathcal{T}_{2,3}$, reads

$$G = \left[U(2) \times U(1)^3 \right] / U(1)_{\text{diag}}. \quad (3.41)$$

Monopole charges are labelled by, [104],

$$\text{cochar}(G) = \text{Hom}(U(1), T) = \mathbb{Z}^5 / \mathbb{Z}_{\text{diag}} \quad (3.42)$$

assigning 5 integers as follows

$$(A_1, A_2; B_1, B_2, B_3) \in \text{cochar}(U(2)) \times \text{cochar}(U(1)^3) / A_{\text{diag}}. \quad (3.43)$$

By definition, two cocharacters are equivalent if they differ by an integer multiple of A_{diag} . On the other hand, the weights are defined as follows

$$\lambda = \text{weights}(G) \stackrel{\text{def.}}{=} \text{Hom}(T, U(1)) \quad (3.44)$$

such that

$$\text{weights}(G) \times \text{cochar}(G) \longrightarrow \mathbb{Z} \quad \text{and} \quad \langle \lambda, A_{\text{diag}} \rangle = 0 \quad \forall \lambda. \quad (3.45)$$

The matter representation is

$$\mathcal{R} = R \oplus R^*. \quad (3.46)$$

$\mathcal{T}_{2,3}$ has an $SU(2)^3$ flavour symmetry acting on its Coulomb branch, and a corresponding $SL(2, \mathbb{C})^3$ symmetry in the chiral ring. This should be generated by three $\mathfrak{sl}(2, \mathbb{C})^*$ -valued complex moment maps μ_a , $a = 1, 2, 3$.

This is a particular example, where the Coulomb branch of the star-shaped quiver coincides with the Higgs branch of the 4D class \mathcal{S} theory $\mathcal{T}_2[\Sigma_{0,3}]$, hence it simply reads

$$\mathcal{M}_H \simeq \mathbb{C}^8. \quad (3.47)$$

The Higgs branch

\mathbf{k} denotes the number of punctures on sphere. The Higgs branch is defined once having identified all the pair of pants decompositions of $\Sigma_{0,\mathbf{k}}$, namely

$$\Sigma_{0,\mathbf{k}} \simeq \bigcup_{\alpha=1}^{k-2} \Sigma_{0,3}^{(\alpha)}, \quad (3.48)$$

from which the Higgs branch is extracted as a holomorphic symplectic quotient

$$\mathcal{M}_H^{4D} \simeq \left[\prod_{\alpha=1}^{k-2} \mathbb{C}^8 \right] // SL(2, \mathbb{C})^{k-3}. \quad (3.49)$$

However, the disadvantage of this case is that the decomposition is not unique, and the operators and relations are not manifest.

1- and 2-legged star-shaped quivers

For $\mathbf{k} = 1, 2$, the Coulomb branches simply read

$$\mathcal{M}_C \simeq T^*SL(2, \mathbb{C}) \quad , \quad \mathcal{M}_C \simeq T^*SL(2, \mathbb{C}) //_{\psi} \mathfrak{N} \quad (3.50)$$

where $T^*SL(2, \mathbb{C})$ is the cotangent bundle of the complex group $SL(2, \mathbb{C})$, whose points are parametrised by a triple (μ_L, g, μ_R) , such that

$$T^*SL(2, \mathbb{C}) \equiv \left\{ (\mu_L, g, \mu_R) \mid \mu_{L,R} \in \mathfrak{sl}(N, \mathbb{C})^* , g \in SL(N, \mathbb{C}) , \mu_L g = g \mu_R \right\}, \quad (3.51)$$

with $\mu_{L,R}$ complex moment maps for the cation of the left and right multiplication of $SL(N, \mathbb{C})$ on itself, extending to $T^*SL(2, \mathbb{C})$ as complex Hamiltonian actions²⁶.

3.6 Lagrangian submanifolds of a symplectic manifold

We now outline some key terminology needed for later purposes in our treatment. In particular we introduce Lagrangian submanifolds of a symplectic manifold in the context of quiver gauge theories. The remainder of this section is mostly a review of [104].

3.6.1 Hyperkähler quotients

Let X be a Kähler manifold presented as a Kähler quotient of \mathbb{C}^n by the linear action of a compact group G . The hyperkähler analogue, \mathfrak{M} , of X is a hyperkähler quotient of the cotangent bundle $T^*\mathbb{C}^n$ by the induced G -action. Special instances of this construction include hypertoric and quiver varieties, [136].

Let M be a hyperkähler manifold equipped with a hyperhamiltonian action of a compact Lie group G , and moment maps μ_a , $a = 1, 2, 3$. M is a symplectic manifold, and X is a Lagrangian submanifold with respect to M ,²⁷

Suppose that

$$\xi = \xi_R \oplus \xi_C \quad (3.52)$$

is a central regular value of $\mu_{\text{HK}}^{-1}(\xi)$. Then there is a unique hyperkähler structure on the hyperkähler quotient

$$\mathfrak{M} = M //_{\xi} G \stackrel{\text{def.}}{=} \mu_{\text{HK}}^{-1}(\xi) / G, \quad (3.53)$$

which is the hyperkähler analogue of a kähler quotient

$$X = \mathbb{C}^n // G, \quad (3.54)$$

where

$$\mu_{\text{HK}} \equiv \mu_R \oplus \mu_C : M \rightarrow \mathfrak{g}^* \oplus \mathfrak{g}_C^* \quad (3.55)$$

²⁶Essentially, $\mu_{L,R}$ respectively denote the left- and right-invariant trivialisation of the cotangent bundle.

²⁷More of this will be explained in the context of 6.5.

with associated symplectic and holomorphic symplectic forms, $\omega_{\mathbb{R}}^{\xi}$ and $\omega_{\mathbb{C}}^{\xi}$, such that they pull back to the restrictions of $\omega_{\mathbb{R}}$ and $\omega_{\mathbb{C}}$ to $\mu_{\text{HK}}^{-1}(\xi)$.

Restricting to the case in which $M = T^*\mathbb{C}^n$, (3.55) reduces to

$$\mu_{\mathbb{R}} \oplus \mu_{\mathbb{C}} : T^*\mathbb{C}^n \rightarrow \left(\mathfrak{su}(2)^* \oplus (\mathfrak{t}^n)^* \right) \oplus \left(\mathfrak{sl}(2, \mathbb{C})^* \oplus (\mathfrak{u}(1)^n)_{\mathbb{C}}^* \right), \quad (3.56)$$

in which case the hyperpolygon space is the hyperkähler quotient

$$\mathfrak{M}(\alpha) \stackrel{\text{def.}}{=} T^*\mathbb{C}^n //_{(\alpha, 0)} G = \left(\mu_{\mathbb{R}}^{-1}(\alpha) \cap \mu_{\mathbb{C}}^{-1}(0) \right) / G, \quad (3.57)$$

which is a smooth, noncompact hyperkähler manifold of complex dimension $2(n-3)$.

3.7 Abelianisation

To calculate the circle-equivariant cohomology of a hyperkähler space \mathfrak{M} , we need the Kirwan map

$$\kappa_G : H_{S^1 \times G}^*(M) \longrightarrow H_{S^1}^*(M//G) \quad (3.58)$$

$$\hat{H}_{S^1}^*(M//G) \simeq \frac{\hat{H}_{S^1}^*(M//T)^W}{\text{ann}(e)}, \quad (3.59)$$

where T is the maximal torus, W is the Weyl group of G , and

$$e = \prod_{\alpha \in A} \alpha(x - \alpha) \in (\text{Sym } \mathfrak{t}^*)^W \otimes \mathbb{Q}[x] \simeq H_{S^1 \times T}(\text{pt})^W \subseteq \hat{H}_{S^1 \times T}(\text{pt})^W. \quad (3.60)$$

The main goal of his work is that of stating and proving an abelianisation theorem for hyperkähler quotients. This features two main obstacles:

1. Hyperkähler quotients are rarely compact.
2. Surjectivity of the Kirwan map from $H_G^\bullet(X)$ to $H^\bullet(X//G)$.

The analogous map for circle compact hyperkähler quotients is conjecturally surjective, but only a few cases are known to be so. They adopt a different method, instead, weaker than surjectivity. But this is true for the case of Nakajima quiver varieties:

$$\kappa_G : H_{S^1 \times G}^\bullet(M) \rightarrow H_{S^1}^\bullet(M//G). \quad (3.61)$$

Suppose that $M//G$ and $M//T$ are both circle compact. If $\gamma \in \hat{H}_{S^1 \times G}^\bullet(X)$, then integration over the Kirwan map leads to

$$\int_{X//G} \hat{\kappa}_G(\gamma) = \frac{1}{|W|} \int_{X//T} \hat{\kappa}_G(\gamma) \circ r_T^G(\gamma) \cdot e, \quad (3.62)$$

where

$$r_T^G : \hat{H}_{S^1 \times G}^\bullet(M) \rightarrow \hat{H}_{S^1 \times T}(M)^W. \quad (3.63)$$

Importantly in this case, the rational Kirwan map is surjective, implying the following chain of isomorphisms

$$\hat{H}_{S^1}^\bullet(M//G) \simeq \frac{\hat{H}_{S^1}^\bullet(M//T)^W}{\text{ann}(e)} \simeq \left(\frac{\hat{H}_{S^1}^\bullet(M//T)}{\text{ann}(e')} \right)^W \quad (3.64)$$

where

$$e = \prod_{\alpha \in A} \alpha(x - \alpha) \in (\text{Sym } \mathfrak{t}^*)^W \otimes \mathbb{Q}[x] \simeq H_{S^1 \times T}(\text{pt})^W \subseteq \hat{H}_{S^1 \times T}(\text{pt})^W. \quad (3.65)$$

$$e' = \prod_{\alpha \in \Delta^-} \alpha \prod_{\alpha \in \Delta} (x - \alpha) \in (\text{Sym } \mathfrak{t}^*)^W \otimes \mathbb{Q}[x] \simeq H_{S^1 \times T}(\text{pt})^W \subseteq \hat{H}_{S^1 \times T}(\text{pt})^W. \quad (3.66)$$

$$M//G = \mu_G^{-1}(\xi, 0)/G \quad , \quad M//T = \mu_T^{-1}(\text{pr}(\xi), 0)/T \quad (3.67)$$

and

$$\text{pr} : \mathfrak{g}^* \rightarrow \mathfrak{t}^* \quad (3.68)$$

is the natural projection. Then T acts on M with hyperkähler moment map

$$\mu_T = \text{pr} \circ \mu_{\mathbb{R}} \oplus \text{pr}_{\mathbb{C}} \circ \mu_{\mathbb{C}} : M \rightarrow \mathfrak{t}^* \oplus \mathfrak{t}_{\mathbb{C}}^*. \quad (3.69)$$

Let $\xi \in \mathfrak{g}^*$ be central elements such that $(\xi, 0)$ is a regular value of μ_G and $(\text{pr}(\xi), 0)$ is a regular value of μ_T . By further assuming that G acts freely on $\mu_G^{-1}(\xi, 0)$, and T acts freely on $\mu_T^{-1}(\text{pr}(\xi), 0)$. Upon taking (3.67) to be hyperkähler quotients of M by G and T , respectively, since μ_G and μ_T are circle equivariant, then the action of S^1 on M descends to the action of S^1 on M descends to the action on the hyperkähler quotients. $M//T$ also inherits an action of the Weyl group W of G .

3.7.1 Towards Category Theory

Understanding the impact of what we have outlined up to now in the context of this project, we need a crucial ingredient, namely that of higher-categorical structures, to which Part II of this work is devoted. As we shall see there, they provide a natural language for describing interesting dualities arising in supersymmetric quantum field theories. In particular, we will be focussing on the category of circle compact manifolds.

Part II

Higher-Categories and Dualities

4 Higher-categorical structures

Part II of our treatment introduces two main tools needed in the remainder of our work, namely higher-categorical structures and dualities. Given the breadth of topics either terms encompass, we now pause momentarily to briefly outline what are the types of most interest to us, with the purpose of guiding the reader through the remaining sections of this Part. Importantly, this will build understanding of the main tools needed for addressing questions raised in later sections.




Figure 9. Main cases of higher-categorical structures, and tools adopted for describing them. Each one of them will arise in due course in the remainder of our treatment.

Figure 19 summarises the key examples we are interested in:

1. Higher-categorical structures describing String Theory, QFT, and QM, mostly relying upon functorial field theory (FFT), and algebro-geometric tools.
2. Holographic dualities (such as the AdS/CFT correspondence, CS/WZW-models, wedge holography, and generalisations thereof, in turn accounting for the emergence of islands), and categorical dualities (3D mirror symmetry, and homological mirror symmetry).

As explained in the introductory part of this work, understanding the underlining mathematical structure of QFT is one of the core open questions in the realm of physical mathematics. Higher-categorical structures have proved being extremely efficient in achieving this for gauge theories and their supersymmetric counterparts. In particular, this formalism happens to be particularly efficient for keeping track of the spectrum (i.e. the particle-field content) of a given QFT, and how the latter changes under gauging of the global symmetries of the theory one started from.

Up to now, most of the mathematical control in describing higher-dimensional field theories has been for descendants of maximally-supersymmetric 6D $\mathcal{N} = (2, 0)$ superconformal field theories (SCFTs), from which a rich web of dualities arises upon dimensional reduction to, either, 4 or 3 spacetime dimensions. Such descendants feature fewer supersymmetries with respect to their non-Lagrangian parent theory. In particular, the most studied are 4D $\mathcal{N} = 2$ (also known as class \mathcal{S} theories), and 3D $\mathcal{N} = 4$ SCFTs, exhibiting exactly half of the amount of supersymmetry of their parent 6D theory. In this concluding part of our work, we will be restricting to these, [108–112].

In a later section of this Part, we will be focusing on the applicability of category theory to the specific case of theories descending from dimensional reduction of 6D $\mathcal{N} = (2, 0)$ SCFTs, specifically class \mathcal{S} theories, which, according to the Alday-Gaiotto-Tachikawa (AGT)-correspondence, can be equivalently described in terms of a 2D TFT living on the Riemann surface with respect to which dimensional reduction has been performed. Because of this, class \mathcal{S} theories constitute a prototypical example of theories bridging between the two upper nodes of the diagram in figure 19. This part of our work also introduces TQFTs and varieties arising in algebraic geometry as two main examples of higher-categorical structures.

In section 3, we have gathered sufficient evidence supporting the utility of applying category theory for describing simple examples of QFT models. We now switch gears, and turn to a more algebro-geometric analysis of the setup outlined so far. In doing so, we will mostly be relying on [598–601], shedding new light and emphasising techniques that should be further developed and applied to Beyond the Standard Model setups²⁸.

4.1 Higher-categorical structures

For a QFT to be well defined, its spectrum of allowed operators must be compatible with the symmetries of the given theory, [372]. This statement has been a very active area of research, and mostly motivated furthering a full mathematical formulation of QFTs. Category theory has been pointed out as a promising candidate in pursuing such aim.

Following suite, the present work aims at exploring the higher-categorical theory of and descending from 6D $\mathcal{N} = (2, 0)$ SCFTs with the aim of identifying a categorical theory quantity enabling to distinguish between intrinsic and non-intrinsic non-invertible symmetries in class \mathcal{S} theories obtained by dimensional reduction of their parent theory.

Before delving into the specifics of the 6D theory, to which section 9.2 is devoted, the current section aims at providing a brief overview of some essential tools we will need in the remainder of the paper, as well as further motivating the use of such mathematical tools within the following theoretical physics treatment. This section is structured as follows:

- Section 4.1 provides a brief overview of some key features of higher-categorical structures that are relevant for our analysis. In the first part of this section, we briefly recall the relation between the total quantum dimension, fusion and braiding structures.

²⁸More details in this regard will appear in an upcoming work by the same author, [597].

- As an example of a higher-categorical structure, we first describe the Chern-Simons theory for a flux-charge-quasiparticle setup, and how gauging a subgroup of the anyonic symmetry affects the total quantum dimension. The section concludes with an explanation for how to gauge a higher-categorical structure by means of algebraic gauging, [607]. This is really meant to be a warm-up example to the follow-up example of interest to us, namely string theory. For the purpose of our work, its functorial field theory formulation turns out being particularly significant. As we shall see, our analysis extends that of Moore and Segal, [137], and Moore and Tachikawa, [136].

4.1.1 Defect fusion categories

Categories are defined as sets of objects related to each other by morphisms. A 2-category is a collection of categories related to each other by *functors*. Groups constitute a special type of categories, where only one element is present, and every morphism is an isomorphism.

(n, r) -categories are generalisations of, both, n -categories and n -grupoids. As n increases, there are many possibilities, until there are infinitely many of (∞, r) -categories. An $(\infty, 1)$ -category is an ∞ -category in which all n 0morphisms for $n \geq 2$ are equivalences. This is the joint generalisation of the notion of category and ∞ -grupoid.

Among all (n, r) -categories, $(\infty, 1)$ -categories are special in that they are the simplest structures that at the same time admit a higher version of category theory, and encode information about higher-equivalences. In general, to understand (n, r) -categories, one requires the notion of the underlying $(n, 1)$ -, hence, the $(\infty, 1)$ -category, which already captures most of the information of interest.

A *higher*-category, or n -category, is an $(\infty, 1)$ -category enriched with $(n-1)$ -categories, with n denoting the dimensionality.

A category, \mathcal{C} , is *monoidal* (MC) if it comes equipped with a bi-functor

$$\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}. \quad (4.1)$$

For \mathcal{C} semisimple, the tensor product of any two simple objects $x, y \in \mathcal{C}$ is

$$x \otimes y \simeq \bigoplus_{z \in \text{Irr}(\mathcal{C})} N_{xy}^z \cdot z \quad , \quad N_{xy}^z \in \mathbb{N}, \quad (4.2)$$

where $\text{Irr}(\mathcal{C})$ denotes the irreducible representations, $\left\{ N_{xy}^z \right\}_{x, y, z \in \text{Irr}(\mathcal{C})}$ denote the *fusion rules* of \mathcal{C} . Equivalently, the isomorphism classes of simple objects \mathcal{C} generate the *fusion ring* or *Grothendieck ring* of \mathcal{C} , where multiplication is given by (4.1), and N_{xy}^z as structure constants.

Given $x, y \in \mathcal{C}$

$$\text{Hom} : x, y \rightarrow \text{Hom}(x, y) \quad (4.3)$$

is an $(n-1)$ -category, and for $\mathbf{1} \in \mathcal{C}$

$$\text{Hom} : \mathbf{1}, \mathbf{1} \rightarrow \text{Hom}(\mathbf{1}, \mathbf{1}) \stackrel{\text{def.}}{=} \Omega \mathcal{C}, \quad (4.4)$$

$\Omega \mathcal{C}$ is a monoidal $(n-1)$ -category. More generally, an (n, r) -category is a higher-category such that:

1. All k -morphisms for $k > n$ are trivial.
2. All k -morphisms for $k > r$ are invertible.

Braiding and ribbon structure

A modular category (MC) is *braided* if, in addition to (4.1) it is also equipped with a natural isomorphism

$$c_{x,y} : x \otimes y \rightarrow y \otimes x, \quad (4.5)$$

called *braiding*. Given two simple objects $x, y \in \mathcal{C}$, the *S-matrix* is defined as

$$S \stackrel{\text{def.}}{=} (S_{xy})_{z \in \text{Irr}(\mathcal{C})}, \quad (4.6)$$

with components defined from (4.5)

$$S_{xy} \stackrel{\text{def.}}{=} \text{Tr} (c_{y,x^*} \circ c_{x^*,y}) \equiv \text{Tr} (c_{y^*,x} \circ c_{x,y^*}). \quad (4.7)$$




Figure 10. The S-matrix corresponds to the Hopf link, and encodes the information of mutual-statistics. As shown in these pictures, the linking between particle worldlines encodes information of mutual-statistics.

For \mathcal{C} a braided fusion category and $x \in \mathcal{C}$, the 2 morphisms

$$x \xrightarrow{\text{id}_x \otimes b_x} x \otimes x \otimes x^* \xrightarrow{c_{x,x} \otimes \text{id}_{x^*}} x \otimes x \otimes x^* \xrightarrow{\text{id}_x \otimes b_x^\dagger} x \quad (4.8)$$

$$x \xrightarrow{d_x^\dagger \otimes \text{id}_x} x^* \otimes x \otimes x \xrightarrow{\text{id}_{x^*} \otimes c_{x,x}} x^* \otimes x \otimes x \xrightarrow{b_x \otimes \text{id}_x} x \quad (4.9)$$

are both equivalent to

$$\theta_x : x \rightarrow x, \quad (4.10)$$

defining the *twist* or *topological spin* of x . Importantly, $\theta_x \neq \text{id}_x$, since the worldline should be viewed as a ribbon, hence a line equipped with a framing. Both processes described in figure 11 are therefore equivalent to twisting the ribbon counterclockwise. The twisting rotates the topological defect by 360° , reason why θ_x is called the topological spin of x .



Figure 11. The ribbon structure describing self-statistics is here drawn as the worldline of a particle excitation x winding around itself. Such information is encoded in the T -matrix.

$\forall x \in \mathcal{C}$, since $\text{Hom}_{\mathcal{C}}(x, x) \simeq \mathbb{C}$,

$$\theta_x \equiv T_x \cdot \text{id}_x \quad , \quad T_x \in \mathbb{C}, \quad (4.11)$$

where the T -matrix of the unitary braided fusion category is defined as

$$T \stackrel{\text{def.}}{=} (T_x \delta_{xy})_{x, y \in \text{Irr}(\mathcal{C})}, \quad (4.12)$$

encoding the information of self-statistics.

Quantum double category of a finite group

\forall finite group G , there is a unitary MTC \mathcal{D}_G called *double quantum category of G* , [353], such that:

1. An object in \mathcal{D}_G is a vector space V equipped with a G -action

$$\rho : G \longrightarrow GL(V), \quad (4.13)$$

and a G -grading $V \equiv \bigoplus_{g \in G} V_g$ such that

$$\rho(g) : V_h \longrightarrow V_{ghg^{-1}} \quad , \quad \forall g, h \in G. \quad (4.14)$$

2. A morphism in \mathcal{D}_G is a \mathbb{C} -linear map that is, both, a morphism in $\text{Rep}(G)$ and a morphism in $\text{Vec}(G)$.
3. Given $g \in G$, we denote its conjugacy class as $[g]$, and its centraliser as $Z(g)$. If π is an irreducible $Z(g)$ -representation, the induced representation $\text{Ind}_{Z(g)}^G(\pi) \stackrel{\text{def.}}{=} X_{(g, \pi)}$ is a G -representation admitting a canonical $G/Z(g) \simeq [g]$ -grading. $X_{(g, \pi)}$ is a simple object in \mathcal{D}_G .

4. The isomorphism classes of objects in \mathcal{D}_G are labelled by pairs $([g], [\rho])$ where $[g]$ is the conjugacy class in G and $[\rho]$ is an isomorphism class of irreducible representations of $Z(g)$, where $Z[g]$ only depends on the conjugacy class of g .

The properties of the S and T matrices defined in (4.6) and (4.12), and their generalisations to the case of quantum double categories have been extensively described in previous works (see, for example, [353]). For the purpose of the present work, it is important to stress that the S and T matrices are related to the generators of the $SL(2, \mathbb{Z})$ group

$$s \stackrel{\text{def.}}{=} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad t \stackrel{\text{def.}}{=} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad (4.15)$$

such that

$$(st)^3 \equiv s^2, \quad s^4 \equiv 1. \quad (4.16)$$

under a projective $SL(2, \mathbb{Z})$ -representation

$$s \mapsto \frac{S}{\sqrt{\dim(\mathcal{C})}}, \quad t \mapsto T. \quad (4.17)$$

The quantity $\mathcal{D} \stackrel{\text{def.}}{=} \dim(\mathcal{C})$ is the *total quantum dimension* of the category \mathcal{C} . For the case in which \mathcal{C} is made up of purely simple objects, the total quantum dimension of a given superselection sector simply reads

$$\mathcal{D}_x \stackrel{\text{def.}}{=} \sqrt{\sum_i \dim^2(x_i)} = \sqrt{\sum_i S_{1x_i}^2}, \quad (4.18)$$

where x_i are the elements of the category and $S_{1x_i}^2$ the square of the S -matrix evaluated between the identity element, $\mathbf{1}$, and any other non-trivial element, x_i .

4.1.2 Example of higher-categorical structures

CS theory

We now turn to describing the main example of MTC of interest to us. We will assume that the total category associated to a theory \mathfrak{T} with gauge group G , $\mathcal{C}_{\mathfrak{T}}$ is given by

$$\mathcal{C}_{\mathfrak{T}} \stackrel{\text{def.}}{=} \{ \mathcal{C}_{point}, \mathcal{C}_{flux}, \mathcal{C}_{Hopf} \}, \quad (4.19)$$

where each subcategory is a different superselection sector of the theory, defined as follows:

1. $\mathcal{C}_{point} \equiv \{1, a, \dots\}$, $a \equiv R \in (G)_{IRREP}$, are point excitations, corresponding to the irreducible representations of the group G . The quantum dimension of each element of the category is $d_a \stackrel{\text{def.}}{=} \dim R$.

2. $\mathcal{C}_{flux} \equiv \{1, \mu, \dots\}$ is the superselection sector of pure fluxes, and therefore corresponds to the conjugacy classes of the finite group G , $C \in (G)_{cj}$. Their quantum dimensions are $d_\mu \stackrel{def.}{=} \sqrt{|C|}$, with C denoting the number of group elements in a given conjugacy class.
3. $\mathcal{C}_{Hopf} \equiv \{1, \eta, \dots\}$ is the collection of Hopf link excitations, with $\eta \stackrel{def.}{=} (C_{(g,h)}, R)$, $gh \equiv hg$, $C_{(g,h)} \equiv \{(tgt^{-1}, tht^{-1} | t \in G)\}$, $R \in E_{(g,h)} \equiv \{t \in G | (g, h) \equiv (tgt^{-1}, tht^{-1})\}$, and quantum dimensions

$$d_\eta \stackrel{def.}{=} \frac{|G|}{|E_{(g,h)}|} \dim R \quad (4.20)$$

The 1-morphisms between these categories are

$$\varphi : \mathcal{C}_{flux} \hookrightarrow \mathcal{C}_{loop} \subset \mathcal{C}_{Hopf} \quad (4.21)$$

$$\phi : \mathcal{C}_{point} \hookrightarrow \mathcal{C}_{loop} \subset \mathcal{C}_{Hopf} \quad (4.22)$$

$$v : \mathcal{C}_{Hopf} \longrightarrow \mathcal{C}_{Hopf} \quad , \quad v(\eta) \equiv \eta^V \quad (4.23)$$

The aforementioned properties suggest the physical interpretation of the superselection sectors in (4.19) as the electric, magnetic and dyonic charges characterising the spectrum of the theory \mathfrak{T} . As a result, the objects living in $\mathcal{C}_{\mathfrak{T}}$ are defined by a 3-tuple $\chi \equiv ([M], \lambda, \rho)$, corresponding to a flux-quasiparticle-charge composite.

As mentioned earlier on in the introduction, application of categories in physics began in the study of 2D CFTs and $D+1$ TQFTs. Their mathematical formulation was originally motivated by the idea of formalising the factorisation property of path-integrals as a certain monoidal functor defined on a cobordism category. Thanks to the pioneering work of Moore and Seiberg on *modular tensor categories* in 2D RCFTs, Rashetikhin and Turaev reformulated their discovery in present-day categorical formulation, and used it to give the 2+1 D RT TQFT formulation, eventually leading to the study of topological excitations (or anyons) in a 2D topological order. Anyons are quasiparticles, and as such can be thought of as being the physical counterpart of (4.19). Systems exhibiting anyon symmetries with gauge group G in a given topological phase efficiently be related to Chern-Simons (CS) theory.

For such correspondence to hold, the CS theory needs to account for the topological information defined by the braiding statistics and quasiparticle fusion rules (cf. section 4.1.1). For example, an abelian phase in (2+1)D can be characterised by a QFT with partition function defined as follows

$$\mathcal{Z}[\mathcal{J}] \equiv \int \left[\mathcal{D}\alpha(\vec{r}) \right] \exp(iS[\mathcal{J}]), \quad (4.24)$$

involving an N -component set of $U(1)$ gauge fields $\alpha \equiv (\alpha_1, \dots, \alpha_N)$ with action

$$S[\mathcal{J}] \equiv \frac{1}{4\pi} \int \left(K_{IJ} \alpha^I \wedge d\alpha^J + \alpha^I \mathcal{J}_I \right). \quad (4.25)$$

Quasiparticles, $\psi^{\vec{a}}$, are sources for $\mathcal{J}_1^{a_1}, \dots, \mathcal{J}_N^{a_N}$, labelled by $\vec{a} \equiv (a_1, \dots, a_N)$ on a lattice $\Gamma^* \equiv \mathbb{Z}^N$. At long distance, nearby quasiparticles can form a single entity, leading to the definition of a fusion structure. The K -matrix in (4.25) dictates the braiding statistics of quasi-particles

$$\mathcal{D} S_{ab} \equiv e^{2\pi i a^T K^{-1} b}, \quad \mathcal{D} \equiv \sqrt{|\det(K)|} \equiv \sqrt{\mathcal{A}} \quad (4.26)$$

with \mathcal{D} the total quantum dimension. The exchange statistics, instead, is defined by

$$\theta_a \equiv e^{2\pi i a^T K^{-1} b}, \quad (4.27)$$

corresponding to the spin of the quasi particle. The topological phase generated by gauging the anyonic symmetry, is referred to as *twisted liquid*. The latter are generalisations of (2+1)D discrete gauge theories. Their quasiparticles are compositions of fluxes and charges associated to the gauged anyonic symmetry, as well as a superselection sector of the original topological state. Their total quantum dimensions are related as follows

$$\mathcal{D}_{TL} \equiv \mathcal{D}_o |G|, \quad (4.28)$$

with $|G|$ the order of the anyonic symmetry group G being gauged, and \mathcal{D}_o the quantum dimension of the pre-gauged theory.

An important remark is of order, though. A CS description of a topological phase (cf. (4.25)) is not unique. This follows from the fact that the K matrix encodes the same fusion and braiding structure even after undergoing a basis transformation. The set of automorphisms, $\text{Aut}(K)$, preserving K correspond an anyonic symmetry operation permuting the anyons with the same fusion properties and spin-statistics. As such, it classifies the global symmetries of the TQFT associated to the CS action, S_{CS} . Once having modded-out the trivial relabellings, $\text{Inn}(K)$, we are left with the outer automorphisms

$$\text{Outer}(K) \stackrel{\text{def.}}{=} \frac{\text{Aut}(K)}{\text{Inner}(K)}, \quad (4.29)$$

which will play a key role in the remainder of the present work.

These results about the total quantum dimension in gauged topological orders have been thoroughly investigated in recent years. In the next section, we shall see how the presence of intrinsic non-invertibles arising in the gauge theory lead to a different expression for the total quantum dimension, thereby signalling that the multiplicity of some superselection sectors can be greater than unity.

4.2 String Theory and Quantum Mechanics (QM) as Functorial Field Theories (FFTs)

As previously mentioned, the previous section is really meant to be a warm-up example to the follow-up example of interest to us, namely string theory. For the purpose of our work,

its functorial field theory formulation turns out being particularly significant. As we shall see, our analysis extends that of Moore and Segal, [137], and Moore and Tachikawa, [136].

String Theory should be interpreted as a functor

$$\mathcal{F} : \text{Cob}_2 \rightarrow \text{Vect}, \quad (4.30)$$

where vector spaces describe the quantum states of strings and the operators describe how the possible interaction processes that can happen between strings affect the quantum state. Cob_2 denotes the geometry of spacetime, whereas Vect denotes the linear algebra of quantum physics.




Figure 12. String Theory as cobordism functor (on the LHS). On the RHS, instead, the different string theory sectors emerging as, either, low-energy limits (for 11D SUGRA) or from different compactifications of M-theory (R.H.S.). The different sectors differ by means of the p-forms that can be defined in the 10 or 11D spacetimes, thereby constraining the types of objects to which they couple, i.e. Dp-branes, which can really be thought of as the string theory analogues of the Wilson and 't Hooft operators, Some sectors are related by S and T -duality, denoted by the red and orange continuous lines, respectively.

The functorial approach can similarly be extended to QFT (with QM as a particular case). The most subtle part being that of completely identifying the exact structure of, both, the source and target category.

Usually, QM is described by:

1. A set, V , of possible quantum states of a particle (described by vectors in a complex inner product space).
2. A set, U_t , of operators on V , one for each $t \in \mathbf{R}_{\geq 0}$, equipped with an identity element, $U_o \equiv I$, and group composition law

$$U_{t+s} = U_t \circ U_s, \quad (4.31)$$

with U_t denoting the evaluation of a state from time 0 to time t .

4.2.1 Calabi-Yau (CY) categories

Calabi Yau categories were originally studied as an abstract version of the derived category of coherent sheaves on a CY manifold, [625–627]. We now provide a short digression explaining this terminology.

An algebraic scheme is *locally noetherian* if it can be covered by a family U_α of open subsets of the form

$$U_\alpha = \text{Spec } R_\alpha, \quad (4.32)$$

where R_α are noetherian rings.

When X is a noetherian scheme, one has the full *abelian subcategory of coherent sheaves*, corresponding to locally finitely-generated modules, [304],

$$\text{Coh}(O_X) \subset \text{QCoh}(O_X). \quad (4.33)$$

One can define from these the triangulated subcategory

$$D(\text{Coh}(O_X)) \subset D(\text{QCoh}(O_X)). \quad (4.34)$$

Denoting with $D^b(\text{Coh}(O_X))$ the bounded derived category of $\text{Coh}(O_X)$, the fully-faithful functor

$$D^b(\text{Coh}(O_X)) \hookrightarrow D(\text{Mod}(O_X)), \quad (4.35)$$

identifies $D^b(\text{Coh}(O_X))$ with the full subcategory $D(\text{Mod}(O_X))$ of bordered complexes whose cohomology objects are coherent sheaves.

A CY category is defined as a Vect-enriched²⁹ category \mathcal{C} equipped for each object $c \in \mathcal{C}$ with a trace-like map

$$\text{Tr}_c : \mathcal{C}(c, c) \rightarrow k \quad (4.37)$$

to the ground field, k , such that, for any $d \in \mathcal{C}$, the induced pairing

$$\langle -, - \rangle_{c,d} : \mathcal{C}(c, d) \otimes \mathcal{C}(d, c) \rightarrow k \quad (4.38)$$

given by

$$\langle f, g \rangle = \text{Tr}(g \circ f) \quad (4.39)$$

is symmetric and non-degenerate.

Examples of CY-categories are:

²⁹Given a monoidal category, K , i.e. one such that

$$\circ : \text{Hom}(y, z) \otimes \text{Hom}(x, y) \rightarrow \text{Hom}(x, z) \quad (4.36)$$

an *enriched category*, \mathcal{C} over K features a collection of objects $\text{ob}(\mathcal{C})$, such that, for any pair $x, y \in \text{ob}(\mathcal{C})$, $\text{Hom}(x, y) \in K$.

1. From CY varieties, [625–627].
2. Fukaya categories associated with a symplectic manifold, X , [153].

We will be coming back to these in section 6.3. For the moment, we will simply say that the latter have been used to classify topological conformal field theories (TCFTs), originally arising as an abstract version of CFTs constructed from sigma-models whose targets are CY-manifolds.

5 Main Example: Cochain level theories

We now start putting to practice the formalism outlined in section 4 to our main example, namely the algebraic varieties introduced in section 3. The reason for doing so is to explain the relation in between them and Moore-Tachikawa varieties. As noted by the authors themselves, [136], though, their treatment calls for a categorical description of the hyperkähler generalisation of the target category. We anticipate that the answer to such generalisation can be achieved by making use of equivariant cohomology as in the approach of Teleman, [192]. In Part III we will be connecting this setup to the emergence of peculiar higher-categorical structures in class \mathcal{S} theories and their 2D TFT duals.

The present section is structured as follows:

1. At first, we overview some key features of equivariant cohomology and the convolution product, both within the context of quiver gauge theories, [598, 599]
2. Then, we introduce Moore-Tachikawa varieties and their 2D TFT counterpart³⁰, connecting categorical structures and quiver varieties, [136]. Making use of [600], we emphasise how and when 3D mirror symmetry manifests from a categorical perspective, specifically in terms of bordisms, making use of the Kostant-Whittaker symplectic reduction, [598].
3. Last but not least, we close the section highlighting the importance of abelianisation for embedding the ring homology of the theory in question. In Part III, we will see how this is in turn related to (un)gauging categorical SymTFTs.

5.1 Cochain complexes

A *cochain complex*, (A^\bullet, d^\bullet) , is an algebraic structure that consists of a sequence of abelian groups (or modules), A^\bullet , and a sequence of homomorphisms between consecutive groups, d^\bullet , such that the image of each homomorphism is included in the kernel of the next. To a chain complex, (A_\bullet, d_\bullet) , there is an associated homology, which describes how the images are included in the kernels. A cochain complex is similar to a chain complex, except that its homomorphisms are in the opposite direction. The homology of a cochain complex is called its *cohomology*

³⁰For completeness, we provide some further details in appendix 5.4.




Figure 13. This diagram shows how the main example introduced in section 3.5 fits within the 2-categorical description of Moore-Tachikawa varieties, together with their generalisation, and the gauging prescription within the SymTFT setup.

$$H(\mathcal{C}) \stackrel{def.}{=} \text{Ker}(\mathcal{Q}) / \text{Im}(\mathcal{Q}). \quad (5.1)$$

The n^{th} homology group, $H_n(H^0)$ is

$$H_n \stackrel{def.}{=} \text{Ker } d_n / \text{Im } d_{n+1}. \quad (5.2)$$

The central object in closed string theory is the vector space $\mathcal{C} \equiv \mathcal{C}_{S^1}$ of states of a single parametrised string. \mathcal{C} denotes the cochain complex in this case, [137]. The latter comes equipped with a grading given by the ghost number, and an operator $\mathcal{Q} : \mathcal{C} \rightarrow \mathcal{C}$ called the BRST operator, raising the ghost number by 1, and such that $\mathcal{Q}^2 \equiv 0$.

5.2 Moore-Segal bordism operators

5.2.1 Cochain level theories

Cochain level theories constitute the most important generalisation of the open and closed TFT construction, reason why we will be devoting the present section to briefly overviewing some of its key features that are mostly needed for our treatment. The reason for doing so is that these mathematical structures enable to determine the set of possible D-branes given a closed string background, [137]. In their work, [137] address this problem from the point of view of a 2D TFT, (18.2), where the whole content of the theory is encoded in a finite-dimensional commutative Frobenius algebra.

The present section is therefore structured as follows:

1. At first, we briefly overview cochain complexes as the essential mathematical tools needed for translating the setup of our previous work, [1], in the formalism of Moore and Segal.
2. We then turn to highlighting the construction of cobordism operators, [137], emphasising its dependence on the conformal structure of the Riemann surface.

5.3 The Moore-Segal setup

The most general finite-dimensional commutative algebra over the complex numbers is of the form

$$\mathcal{C} \stackrel{\text{def.}}{=} \bigoplus_x \mathcal{C}_x \quad , \quad x \in \text{Spec}(\mathcal{C}), \quad (5.3)$$

with

$$\mathcal{C}_x \stackrel{\text{def.}}{=} \mathbb{C} \mathcal{E}_x \oplus m_x, \quad (5.4)$$

where \mathcal{E}_x is an idempotent, and m_x a nilpotent ideal. If \mathcal{C} is a Frobenius algebra, then so too is each \mathcal{C}_x .

In their treatment, [137] restrict to the semisimple³¹ case. Semisimplicity admits many equivalent definitions:

1. The presence of simultaneously-diagonalisable fusion rules.
2. There exists a set of basic idempotents \mathcal{E}_x such that

$$\mathcal{C} \stackrel{\text{def.}}{=} \bigoplus_x \mathbb{C} \mathcal{E}_x \quad , \quad x \in \text{Spec}(\mathcal{C}), \quad \text{with} \quad \mathcal{E}_x \mathcal{E}_y \equiv \delta_{xy} \mathcal{E}_y. \quad (5.5)$$

3. \mathcal{C} is the algebra of complex-valued functions on the finite set of characters of \mathcal{C} , $X \in \text{Spec}(\mathcal{C})$.

³¹Despite appearing quite restrictive, committing to semisimplicity is enough to shed light on the essential structure of the theory. According to [137], to go beyond it, the appropriate objects of study, are cochain-complex valued TFTs rather than non-semisimple TFTs in the usual sense.

For any pair of boundary conditions, a, b , the corresponding cochain complex for a semisimple category is defined as follows, [137],

$$\mathcal{O}_{aa} \simeq \bigoplus_x \text{End}(W_{x,a}), \quad (5.6)$$

$$\mathcal{O}_{ab} \simeq \bigoplus_x \text{Hom}(W_{x,a}; W_{x,b}), \quad (5.7)$$

where $W_{x,a}$ is a vector space associated to every idempotent \mathcal{E}_x .

Cobordism operators

We now turn to the key elements for our analysis, namely cobordism operators. In doing so, we will be using the definition provided by [137], highlighting the crucial property that will be mostly needed in sections 20 and 21. A cobordism Σ from p circles to q circles gives an operator

$$U_{\Sigma, \alpha} : \mathcal{C}^{\otimes p} \rightarrow \mathcal{C}^{\otimes q}, \quad (5.8)$$

which depends on the conformal structure α on Σ . This operator, (5.8), is a cochain map, but its crucial feature is that, changing the conformal structure α on Σ , changes $U_{\Sigma, \alpha}$ only by a cochain homotopy.

To describe how $U_{\Sigma, \alpha}$ varies with α , if \mathcal{M}_Σ is the moduli space of conformal structures on the cobordism Σ which are the identity on the boundary circles, there is a resulting cochain map

$$U_\Sigma : \mathcal{C}^{\otimes p} \rightarrow \Omega(\mathcal{M}_\Sigma; \mathcal{C}^{\otimes q}), \quad (5.9)$$

with the target denoting the de Rham complex of forms on \mathcal{M}_Σ with values in $\mathcal{C}^{\otimes q}$.

An alternative, totally equivalent, definition is that of the following cochain map

$$U_\Sigma : C_\bullet(\mathcal{M}_\Sigma) \rightarrow (\mathcal{C}^{\otimes p})^* \otimes \mathcal{C}^{\otimes q}. \quad (5.10)$$

In the next sections, we will show that lack of reparametrisation-invariance on the Riemann surface implies interesting mathematical and physical features of the resulting theory of interest.

Key points

The main points to keep in mind throughout the reminder of our treatment are the following:

- Cochain level theories provide the natural mathematical formalism for describing absolute theories obtained by partial gaugings of the SymTFT in the Freed-Moore-Teleman setup.
- The definition of cobordism operators associated to such complex cochain structure follows from the assumption that the Riemann surface is reparametrisation-invariant.

Quivers

Equipped with the tools revised in the previous sections, we are now ready to begin the analysis outlined in the roadmap illustrated in figure 2.1. The top entry of our diagram reads an example of a 2-category, namely that of Moore-Tachikawa varieties³². The reason why we are referring to them as our main example is their relation to Coulomb branches of star-shaped quivers, [600], to which we can apply the analysis of section 3. The setups we are dealing with are highly supersymmetric, thereby describing theories that are very far from the SM of particle physics, even in its minimally supersymmetric realisation (MSSM). However, the purpose of this article is that of highlighting features in categorical structures of supersymmetric gauge theories that could come to aid in furthering the understanding of extensions of the SM to possible completions in unifying theories, in turn descending from a supersymmetric parent setting, for the reasons outlined in section 3.4. Furthermore, it is important to remark that, the quiver description of the SM and its extensions is an essential step towards embedding them in any string theory setup, and we refer the reader to some key works on this regard, such as [32, 608].

5.4 Moore-Tachikawa varieties

Having outlined the importance of reparametrisation-invariance in the definition of bordism operators, [137], we now turn to the particular application in describing maximal dimensional Higgs branches of class \mathcal{S} theories, as first proposed by [136]. Our major contribution in the present section will be highlighting where upgrades to the categories defined in [137] are needed for dealing with setups as the ones associated to the correspondence depicted in figure 64, namely those leading to the emergence of composite class \mathcal{S} theories separated by a non-invertible defect.

This section is structured as follows:

1. At first, we briefly overview the source and target categorical structure proposed in [136] assuming duality.
2. We then explain what categorical duality means from an algebraic perspective.
3. We conclude the section indicating the relation between Moore-Tachikawa varieties and Coulomb branches of quiver gauge theories as an interesting realisation of 3D mirror symmetry, and how the categorical generalisation proposed in this work suggests interesting applications to quiver varieties that will be addressed in more detail in Part III.

5.4.1 Categorical structure assuming duality

According to the AGT correspondence, to a given class \mathcal{S} theory, one can assign a 2D TFT valued in a symmetric monoidal category, [136],

³²To facilitate the reader, we have added an appendix with further details to be read, albeit we highly recommend to go back to the original paper [136] for a more thorough overview.




Figure 14. Partial reproduction of a diagram displayed in [112]. The first part of our treatment focuses on the functorial field theory description of class \mathcal{S} theories and their Higgs branches in terms of 2D TFT cobordism constructions.

$$\eta_{G_{\mathbb{C}}} : \text{Bo}_2 \rightarrow \text{HS} \quad (5.11)$$

The existence of this 2D TFT relies on the the source and target categories satisfying a certain list of properties, [136]. We will not reproduce all of them in our treatment, and refer the interested reader to the original work of Moore and Tachikawa for a detailed explanation. In this first part of the section, we will only point out some of the crucial assumptions made in their work for reasons that will become clear in the following pages.

Duality

For the purpose of our work, the crucial assumption made in [136] is the duality structure of the source category Bo_2 . As explained in [136], duality implies that the 2-category Bo_2 is fully specified by its objects, S^1 , and 1-morphisms, namely the bordisms depicted in figure 15. The middle bordism, i.e. the one labelled V , is the identity bordism. One can easily see this by noticing that V is topologically equivalent to a cylinder whose edges are the red circles, i.e. the object of 2-category Bo_2 (the closed string we were referring to in section 5.2.1).

For $\eta_{G_{\mathbb{C}}}$ to be well defined, the source and target categories are required to satisfy certain sewing relations, [136, 137]. This practically means that, compositions between morphisms should close. In particular, the identity itself can be defined in terms of composite homomorphisms as follows,

$$U_{G_{\mathbb{C}}} \circ W_{G_{\mathbb{C}}} \equiv T^* G_{\mathbb{C}}, \quad (5.12)$$

where $T^*G_{\mathbb{C}} \equiv V_{G_{\mathbb{C}}}$. Indeed, one can easily see that combining the first and third bordisms in figure 15, is topologically equivalent to V .³³




Figure 15. Basic bordisms assuming duality of, both, the source and target categories leading to the definition of the identity element, $V_{G_{\mathbb{C}}}$, and the maximal dimensional Higgs branch, $W_{G_{\mathbb{C}}}$.

We therefore wish to highlight the following

Main point: Duality ensures the presence of an identity associated to a certain gauge group, $G_{\mathbb{C}} \equiv \eta_{G_{\mathbb{C}}}(S^1)$, (18.3).

Key axiom

(5.12) is essential for us in relating the formalism of [136, 137] to the setup of figure 64. In particular, it is what leads to the definition of the triple featuring on the RHS of (18.1). To see this explicitly, let us recall a crucial axiom required to be satisfied by (5.11), and, therefore, in turn by (5.12), [136].

For $X \in \text{Hom}(G'_{\mathbb{C}}, G_{\mathbb{C}})$ and $Y \in \text{Hom}(G_{\mathbb{C}}, G''_{\mathbb{C}})$, their composition

$$Y \circ X \in \text{Hom}(G'_{\mathbb{C}}, G''_{\mathbb{C}}) \quad (5.13)$$

is identified with the holomorphic symplectic quotient

$$\begin{aligned} Y \circ X &\stackrel{\text{def.}}{=} X \times Y // G_{\mathbb{C}} \\ &= \{(x, y) \in X \times Y \mid \mu_X(x) + \mu_Y(y) = 0\} / G_{\mathbb{C}}, \end{aligned} \quad (5.14)$$

where

$$\mu_X : X \longrightarrow \mathfrak{g}_{\mathbb{C}}^* \quad , \quad \mu_Y : Y \longrightarrow \mathfrak{g}_{\mathbb{C}}^* \quad (5.15)$$

are the moment maps of the action of $G_{\mathbb{C}}$ on X and Y , with $\mathfrak{g}_{\mathbb{C}}$ the Lie algebra associated to $G_{\mathbb{C}}$. The identity element

$$T^*G_{\mathbb{C}} \stackrel{\text{def.}}{=} \text{id}_{G_{\mathbb{C}}} \in \text{Hom}(G_{\mathbb{C}}, G_{\mathbb{C}}) \quad (5.16)$$

comes with a Hamiltonian $G_{\mathbb{C}} \times G_{\mathbb{C}}$ action. As also explained in [136], to see that $T^*G_{\mathbb{C}}$ acts as the identity, it is enough to consider a composition of homomorphisms, $T^*G_{\mathbb{C}} \circ X$.

³³Indeed, V is topologically equivalent to the cylinder, i.e. the cobordism between S^1 and itself.

Identifying $T^*G_{\mathfrak{c}} \simeq G_{\mathfrak{c}} \times \mathfrak{g}_{\mathfrak{c}}$, and an element of $T^*G_{\mathfrak{c}}$ as (g, a) . The moment map condition, (5.14), reduces to

$$a + \mu(x) = 0, \quad (5.17)$$

from which a can be removed. Consequently, the induced 2-form on the solution space is $G_{\mathfrak{c}}$ -invariant and basic. Upon taking the quotient with respect to $G_{\mathfrak{c}}$, we can gauge g to 1, leading to a holomorphic isomorphism with the original X space with its symplectic form. Note that this is exactly like the constraint found in our main example (3.51) within the context of Coulomb branches of star-shaped quivers.

The categorical quotient taken in defining the composition (5.14) is equivalent to the one that a given absolute theory should be equipped with to potentially gauge away its entire operator content, while leaving only the identity in the spectrum³⁴. Indeed, this is true as long as the embeddings of the subalgebras associated to $G'_{\mathfrak{c}}$ and $G''_{\mathfrak{c}}$ are subsets of each other within the mother algebra $\mathfrak{g}_{\mathfrak{c}}$. However, we are interested in describing more general setups, where the embeddings of the algebras are intersecting albeit not one included within the other. In the remainder of our treatment, we will explain that, for this to be described in the formalism of [136, 137], the standard identity element associated to the gauge group $G_{\mathfrak{c}}$ and embedding Lie algebra $\mathfrak{g}_{\mathfrak{c}}$ needs to be removed from Bo_2 , while being replaced by a new composite bordism, and propose the definition of a new functor.

5.4.2 Duality from an algebraic perspective

Before turning to explaining what are the changes that the source and target categories should undergo, we will briefly pause for a digression explaining how reparametrisation-invariance on the Riemann surface involved in the definition of the bordism operators, outlined in section 5.2.1, is strongly related to the aforementioned duality assumption.

As explained in [136], the crucial point is that, thanks to the duality propriety of the source 2-category Bo_2 , the identity element in the target category, $T^*G_{\mathfrak{c}}$, is reparametrisation-invariant. In particular, one could compose the identity morphisms as follows³⁵

$$T^*G_{\mathfrak{c}}^{a'} \circ T^*G_{\mathfrak{c}}^a \equiv \left(T^*G_{\mathfrak{c}}^a \times T^*G_{\mathfrak{c}}^{a'} \right) // G_{\mathfrak{c}}, \quad (5.18)$$

where a, a' denote the choice of the conformal structure. In particular, one can think of them as denoting the area of the bordism.

From the considerations made above, it therefore follows that one could rephrase (5.18) as the definition of the Drinfeld center for the composite system made up of two class \mathcal{S} theories (associated to the two gauge groups involved) separated by an invertible defect, with the latter ensuring reparametrisation invariance of the Riemann surface [137].

Concretely, under the duality assumption, one could gauge away either of the two groups, while being left with the following

³⁴cf. explanation in the Introduction. This is basically what leads to the definition of the fiber functor, moment map and Drinfeld center.

³⁵Making use of the axiom (5.14).

$$\mathbf{1}_{G_C^{a'}} \circ T^* G_C^a \equiv \left(T^* G_C^a \times \mathbf{1}_{G_C^{a'}} \right) // G_C. \quad (5.19)$$

If the conformal structure were the same on the two sides, then it would be the same, either with or without the composition rule. If

$$G_C^a \times G_C^{a'} \equiv G_C^{a+a'} \equiv G_C, \quad \forall a, a', \quad (5.20)$$

then (5.18) can be recast to the following

$$\boxed{T^* G_C^a \equiv T^* G_C^a // G_C}, \quad (5.21)$$

which is equivalent to a statement of S-duality³⁶. Once more, we highlight that this is possible because the source category for the 2D TFT associated to the group G_C contains the identity element. But, in case this is not true³⁷, (5.21) needs to be changed accordingly, which one could think of as a generalisation of an S-duality statement. Indeed, if the group composition rules do not hold,

$$G_C^{a+a'} \neq G_C, \quad \forall a, a', \quad (5.22)$$

we get something that is not simply the ordinary S-dual theory, (5.21).

The main purpose of our work is basically to go backwards, starting from the LHS of (5.18) and determining what the RHS should be. Most importantly, we need to:

1. Identify G_C in the new theory obtained by composing the two theories on the LHS, each one characterised by a different choice of conformal structure on the Riemann surface.
2. Equivalently to 1., reconstruct $T^* G_C$, namely the identity of the composite theory.
3. We highlight that the most important generalisation of the 2D TFT (5.14) that one should really be using for the case of interest to us is instead the following

$$\boxed{\tilde{\eta}_{G_C} : \text{Bo}_2 \setminus V \rightarrow \text{HK}} \quad (5.23)$$

which, as already pointed out in [136], requires removing the identity element from the source category. Its effect on the target is to turn it into a hyperkähler quotient. Its connection with theoretical physics³⁸ is the main focus of section 21.

From step 1., an important observation is in order. G_C acts on the two factors on the RHS of (5.18) in separate ways. This is part of the meaning of the generalisation of

³⁶Note that the conformal structure a is the same on both sides.

³⁷Such as the case in which the Riemann surface is no longer reparametrisation-invariant.

³⁸Already presented in [3].

S-duality that we were previously referring to. Indeed, the categorical quotient $//G_{\mathbb{C}}$ tells us what the identity is as a result of gauging a certain subalgebra. This is obtained by taking the 1-morphisms on either side of the correspondence and taking their nontrivial composition w.r.t. $T^*G_{\mathbb{C}}$, with the latter being the identity in the target category. But the latter was assumed to be removed. The immediate suggestion to circumvent this shortcoming is that the functors defining the identity element of each individual theory on the LHS of (5.18) is different w.r.t. the one on the RHS. In section 21 we therefore propose the generalisation of 3D mirror symmetry as the need to define two different 2D TFT functors associated to the left and right hand sides of (5.18).

Algebraic Varieties

In the concluding part of this section, we highlight an interesting application of (5.23) in the context of quiver gauge theories, which will be explained in more detail in the concluding sections of this work, in particular leading towards a generalisation 3D mirror symmetry.

The Higgs branches described by Moore and Tachikawa are known to have been reproduced by [598] as the Coulomb branches of 3D $\mathcal{N} = 4$ supersymmetric quiver gauge theories. Such correspondence is therefore equivalent to a statement of 3D mirror symmetry. The purpose of [4] is to explain what the 3D dual of a theory described by (5.23) is in terms of Coulomb branches of 3D $\mathcal{N} = 4$ quiver gauge theories. In this way, we expect to be able to prove the statements made in [3].

If V has been removed from the source, one should expect there to be more than one 2D TFT of the kind (5.11) associated to two different gauge groups whose embedding in the gauge group associated to the original TFT with identity element V is not simply a cochain complex. Correspondingly, this also means that there is more than one 1-morphism $U_{G_{\mathbb{C}}}$.

Given that the identity of the embedding theory is defined as follows

$$\boxed{U_{G_{\mathbb{C}}} \circ W_{G_{\mathbb{C}}} \equiv T^*G_{\mathbb{C}}} \quad , \quad \boxed{\eta_{G_{\mathbb{C}}}(V) \equiv T^*G_{\mathbb{C}}} . \quad (5.24)$$

and that

$$U_{G_{\mathbb{C}}} \stackrel{def.}{=} G_{\mathbb{C}} \times S_n \subset G_{\mathbb{C}} \times \mathfrak{g}_{\mathbb{C}} \simeq T^*G_{\mathbb{C}}, \quad (5.25)$$

with S_n is the Slodowy slice at a principal nilpotent element n . The physical theories of class \mathcal{S} predict the existence of a variety $W_{G_{\mathbb{C}}}$ satisfying the properties needed to define a TFT, $\eta_{G_{\mathbb{C}}}$. From the duality assumption, it follows that the dimensionalities of the two varieties are related as follows

$$\dim_{\mathbb{C}} U_{G_{\mathbb{C}}} \stackrel{def.}{=} \dim_{\mathbb{C}} G_{\mathbb{C}} + \text{rank } G_{\mathbb{C}}. \quad (5.26)$$

$$\dim_{\mathbb{C}} W_{G_{\mathbb{C}}} \stackrel{def.}{=} 3 \dim_{\mathbb{C}} G_{\mathbb{C}} - \text{rank } G_{\mathbb{C}}. \quad (5.27)$$

However, if the identity needs to be removed from Bo_2 , $T^*G_{\mathbf{c}}$ is not the identity and, in particular (5.27) needs to be redefined precisely because the source is no longer a dual category.

As a concluding remark to what we have just said, in [137] they conjecture the following property for the moment maps associated to the G^3 action on the Higgs branch $W_{G_{\mathbf{c}}}$

$$\boxed{\mu_i : W_{G_{\mathbf{c}}} \rightarrow \mathfrak{g}_{\mathbf{c}}}^* \quad , \quad i = 1, 2, 3. \quad (5.28)$$

This is crucial to our analysis since (5.28) can be inverted to obtain the Higgs branch as a hyperkähler quotient

$$\boxed{W_{G_{\mathbf{c}}} \equiv \mu^{-1}/G^3} . \quad (5.29)$$

However, for the case in which the identity is removed from the source category, (5.29) does not hold anymore precisely because of the lack of permutational symmetry arising in the quotient. In section 21 we will be explaining how to define $W_{G_{\mathbf{c}}}$ and its dimensionality for the case involving categories without a duality structure.

5.5 Examples

These are exactly the examples we encountered in section 3.5, when dealing with Coulomb branches of star-shaped quivers.

Key points

The main points to bare in mind from this section are the following:

- Categorical duality ensures the presence of an identity object.
- 3D mirror symmetry requires reparametrisation-invariance.

6 Dualities among Algebraic Varieties

In this section we describe how abelianisation fits in the algebro-geometric description of 3D mirror symmetry. In doing so, we explicitly explain the analogy in between different treatments, [136]. The main aim for doing so at this stage is that of introducing the reader to some crucial notions that will later be developed in more exotic setups, to which Part III and V are devoted.

The present section is structured as follows:

1. At first, we overview some key features of equivariant cohomology and the convolution product, both within the context of quiver gauge theories, [598, 599].
2. Then, making use of [600], we emphasise how and when 3D mirror symmetry manifests from a categorical perspective, specifically in terms of bordisms, making use of the Kostant-Whittaker symplectic reduction, [598].

3. Last but not least, we close the section highlighting the importance of abelianisation for embedding the ring homology of the theory in question, explaining how this is in turn related to (un)gauging the categorical SymTFT introduced in section 13. In doing so, we also overview the work of Teleman, [192], thereby connecting it with Moore-Tachikawa varieties, and Coulomb branches of star-shaped quivers.

6.1 Equivariant cohomology, fixed point localisation and convolution product

Equivariant cohomology, [386], also known as Borel-Moore cohomology, [623], is a cohomology theory from algebraic topology which applies to topological spaces with a group action. It can be viewed as a generalisation of group cohomology. More explicitly, the equivariant cohomology ring of a space X with action of a topological group G is defined as the ordinary cohomology ring with coefficient ring Λ of the homotopy quotient $EG \times_G X$

$$H_G^\bullet(X; \Lambda) = H^\bullet(EG \times_G X; \Lambda). \quad (6.1)$$

The Kirwan map

In section 3 we saw that the Kirwan map, first introduced in [138], states

$$H_G^\bullet(M) \rightarrow H^\bullet(M//_p G), \quad (6.2)$$

where M is a Hamiltonian G -space, i.e. a symplectic manifold acted on by a Lie group G with a moment map

$$\mu : M \rightarrow \mathfrak{g}^*. \quad (6.3)$$

$H^\bullet(M)$ is an equivariant cohomology ring of M , i.e. the cohomology ring of the homotopy quotient $EG \times_G M$ of M by G .

$$M//_p G = \mu^{-1}(p)/G \quad (6.4)$$

is the symplectic quotient of M by G at a regular central value $p \in Z(\mathfrak{g}^*)$ of μ . It is defined as the map of equivariant cohomology induced by the inclusion

$$\mu^{-1}(p) \hookrightarrow M \quad (6.5)$$

followed by the canonical isomorphism

$$H_G^\bullet(\mu^{-1}(p)) \simeq H^\bullet(M//_p G). \quad (6.6)$$

Fixed point localisation and convolution product

In BFN, the embedding of algebras is performed under the so called *convolution product*. To see how this is defined, we start off with the convolution diagram for the affine Grassmannian, [598, 599]

$$\mathrm{Gr}_G \times \mathrm{Gr}_G \xleftarrow{p} G_\kappa \times \mathrm{Gr}_G \xrightarrow{q} \mathrm{Gr}_G \tilde{\times} \mathrm{Gr}_G \xrightarrow{m} \mathrm{Gr}_G, \quad (6.7)$$

where p, q are projections, m is a multiplication, and

$$\mathrm{Gr}_G \tilde{\times} \mathrm{Gr}_G = G_\kappa \times_{G_\mathcal{O}} \mathrm{Gr}_G. \quad (6.8)$$

Hence, given any two $G_\mathcal{O}$ -equivariant perverse sheaves $\mathcal{A}_1, \mathcal{A}_2$, the pullback

$$p^* (\mathcal{A}_1 \boxtimes \mathcal{A}_2) \equiv \mathcal{A}_1 \tilde{\boxtimes} \mathcal{A}_2 \quad (6.9)$$

descends to $\mathrm{Gr}_G \tilde{\times} \mathrm{Gr}_G$ by equivariance, and we can define the convolution product as follows

$$\mathcal{A}_1 \star \mathcal{A}_2 = m_* (\mathcal{A}_1 \tilde{\boxtimes} \mathcal{A}_2). \quad (6.10)$$

The latter defines a symmetric monoidal structure on the category of $G_\mathcal{O}$ -equivariant perverse sheaves on Gr_G , and is equivariant to the monoidal category of finite-dimensional representations of the Langlands dual of G .

(6.8) is the convolution diagram for the case in which G is a complex reductive group and representation $\mathbf{N} = 0$. For arbitrary \mathbf{N} , we can generalise the convolution diagram, achieving a noncommutative algebraic structure. In order to do so, we define a triple $\mathcal{R} = (\mathcal{P}, \varphi, s)$, where \mathcal{P} is a G -bundle on a formal disk $D = \mathrm{Spec} \mathbf{C}[[z]]$, φ is its trivialisation over the punctured disk $D^* = \mathrm{Spec} \mathbf{C}((z))$, and s is a section of the associated vector bundle $\mathcal{P} \times_G \mathbf{N}$ such that it is sent to a regular section of the trivial bundle under φ .

Coulomb branches

Given a complex vector space, V , on a connected reductive algebraic group, G , with a fixed faithful linear action on V , the *Higgs branch* is defined as follows

$$\mathcal{M}_H = \mu^{-1}(0) // G = \mathrm{Spec} \left(\mathbf{C} \left[\mu^{-1}(0) \right]^G \right) \quad (6.11)$$

with

$$\mu : T^*V \rightarrow \mathfrak{g} \quad (6.12)$$

denoting the moment map, whereas the *Coulomb branch*, \mathcal{M}_C , is the spectrum of a ring constructed as a convolution algebra in the homology of the affine Grassmannian, where the choice of the representation V is incorporated as certain quantum corrections to convolution in homology, which are kept track of by an auxiliary vector bundle, [598, 599]. \mathcal{M}_H and \mathcal{M}_C are conjectured to be symplectic duals to each other, with the most important relation in between them being Koszul duality between the categories of the quantised varieties, [603].

Let G be a simply-connected complex semisimple group, and let G_{ad} be its adjoint form. The group G_{ad} acts on G by conjugation, and G contains a transverse slice Σ for

this action, which was first introduced by Steinberg, [624]. The resulting multiplicative universal centraliser is the smooth affine variety

$$Z_G = \left\{ (a, h) \in G_{ad} \times \Sigma \mid a \in G_{ad}^h \right\}. \quad (6.13)$$

When G is simply-laced, Bezrukavnikov, Finkelberg, and Mirković, [386], showed that its coordinate ring is isomorphic to the equivariant K-theory of the affine Grassmannian of the Langlands dual group G^\vee , therefore, in this case, Z_G is an example of a Coulomb branch as defined by BFN [598, 599].

6.2 Mirror symmetry in Moore-Tachikawa varieties

We now turn to the specific case of interest to us, namely Moore-Tachikawa varieties, and their relation with the work of [102, 192, 600, 601]. First and foremost, we will briefly overview how the Higgs and Coulomb branches arise in this setup.

Given a reductive connected group, G , there is a functor defined as follows

$$\eta_{G_{\mathcal{C}}} : \text{Bo}_2 \rightarrow \mathcal{C} \quad (6.14)$$

where \mathcal{C} is a hyperbolic symplectic manifold, whose objects are affine algebraic groups, and whose morphisms

$$\text{Hom}_{\mathcal{C}}(G_1, G_2) \quad (6.15)$$

are isomorphism classes of symplectic³⁹ varieties with Hamiltonian $G_1 \times G_2$ -action. $\eta_{G_{\mathcal{C}}}$ is fully specified by its action on objects and morphisms, namely

$$\eta_{G_{\mathcal{C}}}(S^1) \mapsto G \quad , \quad \eta_{G_{\mathcal{C}}}(S^2) \mapsto Z_G \quad (6.16)$$

$$\eta_{G_{\mathcal{C}}}(S^1 \times S^1) \mapsto T^*G$$

where Z_G is the *universal centraliser*, consisting of all the pairs (x, g) with $x \in \mathfrak{g}$, \mathfrak{g} Lie, regular, $g \in G$ and $ad(x)_{\mathfrak{g}} = x$. Given $K \subset \mathfrak{g}$ a Kostant slice, and (e, h, f) a principal \mathfrak{sl}_2 -triple in \mathfrak{g} , the Kostant slice can be rewritten as follows

$$K = e + Z(f) \quad (6.17)$$

with $Z(f)$ the centraliser of f in \mathfrak{g} , $\mathfrak{g} \simeq \mathfrak{g}^*$. For a given Kostant slice, the universal centraliser can therefore be redefined as follows

$$Z_G = \{(g, x) \mid g \in G, x \in K, ad_{\mathfrak{g}}(x) = x\}. \quad (6.18)$$

What we have outlined so far can be suitably adapted to the case of interest to us, namely when Σ is a Riemann surface whose boundary has n components, from which one gets a symplectic variety X with a G^n -action. This symplectic variety, X , is often times referred to as the *Higgs branch*.

³⁹By this we mean affine, Poisson, operationally symplectic, with a Hamiltonian chosen moment map.

6.2.1 The Kostant-Whittaker symplectic reduction

The action of the functor explicited in (6.16), however, is not exhaustive. Indeed, there is one more piece of information that is needed in order for the 2D TFT to be fully determined, and this is its action on the bordism corresponding to the three-punctured sphere.

To see how this relates to quivers, let us consider the case in which X is a symplectic variety with G -action. Let $N \subset G$ be a maximal map, and

$$\chi: N \longrightarrow \mathbf{C} \quad (6.19)$$

a non-degenerate homomorphism such that

$$N \Big/ [N, N] \simeq \bigoplus_{\alpha_i} \mathbf{C}_{\alpha_i}, \quad (6.20)$$

where α_i denote the simple roots. Then, the Kostant-Whittaker reduction on the symplectic variety corresponds to the quotient

$$KW(X) = \mu_N^{-1}(\chi) \Big/ N, \quad (6.21)$$

and the universal centraliser becomes

$$Z_G \simeq KW_{G \times G}(T^*G). \quad (6.22)$$

Furthermore, we can take

$$G \times K = KW_{G^V}(T^*G) \quad , \quad KW_G(G \times K) = Z_G, \quad (6.23)$$

meaning that filling in a hole on the sphere corresponds to performing a KW_G reduction. Note that this is a statement of 3D mirror symmetry.

According to the Moore-Tachikawa conjecture, overviewed in slightly more detail in appendix 5.4, a 2D TFT is an example of a 2-categorical structure, that is the Alday-Gaiotto-Tachikawa dual of a supersymmetric 4D theory arising from dimensional reduction of 6D $\mathcal{N} = (2, 0)$ SCFTs of type A_{N-1} on a sphere with N maximal punctures. These theories, oftentimes denoted $\mathcal{T}_N[\Sigma_{0,k}]$, upon being further dimensionally reduced on a circle, $\mathcal{T}_N[\Sigma_{0,k} \times S^1]$, are known to be related to Sicilian 3D $\mathcal{N} = 4$ theories, the so-called star-shaped quivers, upon taking the $\lim_{S^1 \rightarrow 0}$, [600].

In [616], the authors calculated the Hilbert series of the chiral ring

$$\mathbf{C}[\mathcal{M}_C] \simeq \mathbf{C}[\mathcal{M}_H^{4D}] \quad (6.24)$$

exactly with the intention of obtaining new information about the structure of the Higgs branches of the Higgs branch of the 4D theory, \mathcal{M}_H^{4D} , via a direct analysis of the corresponding Coulomb branches \mathcal{M}_C of $\mathcal{T}_{N,k}$, for general N and k . In addition to this, our work wishes to emphasise the importance of the calculation of the Hilbert series together with the correct identification of the ring elements and relations in between them, which

plays a crucial role in developing connection with Particle Physics and calculations therein. In achieving the second objective, a crucial role is played by the so-called *abelianisation* procedure, first introduced by [601], and subsequently emphasised also in [600].

6.3 Homological mirror symmetry

We now turn to a more general setup, first outlined by [192]. In our brief overview, we highlight the importance of its relation with our main example, 3, and how it fits within the context of abelianisation. As we shall see, the key expression that we need to consider is (3.51).

At first, let us briefly overview the notion of homological mirror symmetry in its first formulation, [147].

Homological mirror symmetry, consists in a proposed agreement between two categories, namely:

- Fukaya’s A_∞ -category, $\mathcal{F}(X)$, on the symplectic side,
- The derived category with Yoneda structure, $D^b \mathfrak{Coh}(X^\vee)$, on the complex side.

For completeness, we now turn to explain this terminology.

Fukaya categories

A symplectic manifold, X , is a smooth manifold of even dimension equipped with a non-degenerate symplectic form $\omega \in \Omega_{cl}^2(X)$. A Fukaya category, [153], of a symplectic manifold, X , is an A_∞ -category with Lagrangian submanifolds $L \subset X$ as objects, whose intersections define the Hom-space. An A_∞ -category is a category with associativity condition (A) relaxed without bound on degrees of homotopies (∞). They are linear categories, i.e. their Hom-objects are chain complexes.

Yoneda structure

Yoneda structure, [381], provides in a 2-categorical setting the axiomatic description of the formal properties of the usual presheaf construction and Yoneda embedding of locally smaller categories.

The axioms of Yoneda structure capture the properties of the presheaf construction with CAT replaced by a general 2-category, \mathcal{K} . In order to handle size issues, a class of admissible or legitimate 0-cells is singled-out in $|\mathcal{K}|$. In order to handle size issues, as well as a class of 1-cells that behave well w.r.t. this class and the presheaf construction. In fact, it suffices to describe the admissible 1-cells, since one can identify the admissible 0-cells with the admissible identity 1-cells.

Let \mathcal{K} be a 2-category, and \mathbb{A} a class of 1-cells. The 1-cells of $f \in \mathbb{A}$ are called *admissible* if for all $f \in \mathbb{A}$ and composable 1-cells $g \in \mathcal{K}$, $f \circ g \in \mathbb{A}$. In this setup, a 0-cell $C \in |\mathcal{K}|$ is admissible if id_C is.

A *Yoneda structure* is therefore defined as a pair $(\mathcal{P}, \mathbb{A})$, where \mathcal{P} is a pre-sheaf construction for \mathbb{A} assigns to every admissible object $A \in |\mathbb{A}|$ and object $\mathcal{P}A \in |\mathcal{K}|$ called its object of presheaves, and an admissible 1-cell

$$y_A : A \rightarrow \mathcal{P}A \tag{6.25}$$

called its Yoneda morphism subject to three conditions:

Relation to Factorisation Homology

As we shall see later on in section 19, this formulation is compatible with that of factorisation homology for class \mathcal{S} theories, first outlined in our work, [5].

6.3.1 Relation to our main example

The main point, to which we will be coming back later on in Part V is that the morphisms in the category of compact contractible circles encountered in Moore-Tachikawa varieties are the 1-cells featuring in the Yoneda structure description.

6.3.2 Lagrangian submanifolds

A Lagrangian submanifold of a symplectic manifold⁴⁰, $L \subset X$, is a submanifold which is a maximal isotropic submanifold on which $\omega = 0$. They constitute the leaves of real polarisations, and are, therefore, crucial elements of symplectic geometry.

There is an equivalence of categories between the category of \mathbb{C} -linear representations of a quiver \mathcal{Q} and the category of left $\mathbb{C}\mathcal{Q}$ -modules.




Figure 16. The upper and lower blue arrows correspond to geometric and algebraic resolutions of \mathfrak{M}^0 , respectively.

Moment map and higher homologies

In general, homologies of derived representation schemes can be highly nontrivial. However, in this particular case, one can identify a necessary and sufficient condition for the vanishing of the higher homologies based on the flatness of μ , [344]. In particular, in such reference, it was shown that the derived representation scheme $\text{DRep}_{v,w}(\mathcal{A})$ has vanishing higher homologies if and only if $\mu^{-1}(0) \subset M(\mathcal{Q}, v, w)$ is a complete intersection, which happens only if the moment map is flat, [365]. As we shall see, the requirement of the algebraic variety to be a complete intersection is crucial for the purpose of our treatment.

⁴⁰Note that here we are referring to Calabi-Yau 3-folds, hence X is of even dimensions.

In particular, it ensures the emergence of a 2-categorical structure, whose importance will be the core topic of section 15.

Prior to that, we devote the following subsection to explain the role played by magnetic quivers for describing generalisations of homological mirror symmetry in quiver gauge theories with 8 supercharges.

6.4 3D mirror symmetry

We now briefly overview one of the main dualities of interest to us, namely 3D mirror symmetry. The literature exploring this topic is extremely vast, and we are aware of the fact that our reference list is by no means complete. To the extent possible, we will be referring to the works that are more closely related to the purpose and aims of our work. From the physical point of view, the pioneering work dates back to the joint work of Intriligator and Seiberg, [146], which in turn is related to that of CY-manifolds, developed in the mathematical community.

As initially quoted in section 4.2.1, CY-manifolds are the target of functors defining CY-categories.

The aim of this and the next section is that of geometrising the 3D mirror symmetry outlined in our main example in section 3.5 as a duality exchanging Coulomb and Higgs branches of 3D $\mathcal{N} = 4$ SCFTs. This aim was already achieved by Kapustin, Rozansky, and Saulina, [383], who were the first to address this in the context of Rozansky-Witten (RW) theories, [343].

The simplest topological BCs in the RW model with target X correspond to complex Lagrangian submanifolds of X corresponding to the fact that one associates a boundary condition to a fibration over a complex Lagrangian submanifold whose fibration is a CY-manifold. The set of BCs for the RW model with target X should be thought of as a set of objects of a 2-category.

6.4.1 Categorical formulation

Gromov-Witten (GW) theory is among the most studied 2D TQFTs, [155]. It assigns to a compact symplectic manifold X a space of states $H^\bullet(X)$. Surfaces with points labelled by states give correlators counting pseudo-holomorphic maps to X with coincidence conditions.

Mirror symmetry reduces GW theory to calculations in the complex geometry of a mirror manifold, X^V , introduced by Kontsevich, [147].

The key idea is that, including all surfaces with corners, and BCs (branes) forming a linear category with structure, this category should be able to determine all invariants.

Mathematically, the description starts with a group action acting on the generating category (the Fukaya category), which is factored through the topology of the group. Their mirror description involves holomorphic symplectic manifolds and Lagrangians related to the Langlands dual group. As an application: the complex mirrors of flag varieties proposed by Marsh and Rietsch, [618], ⁴¹.

⁴¹We refer to section 3.5 for preliminary definitions.

The aim is that of studying equivariance in the higher-algebra surrounding TQFTs. The homological information lies in the neighborhood of a Lagrangian within X (a symplectic manifold), whereas most of the interesting physics happens elsewhere. The emerging geometric picture is the following: representations admit a character theory, but characters are now coherent sheaves on a manifold related to the conjugacy classes, instead of functions. The manifold in question is the Bezrukavnikov-Finkelberg-Mirkovic (BFM) space of the Langlands dual Lie group, G^V , and is closely related to the space of conjugacy classes in the complex group $G_{\mathbb{C}}^V$. Multiplicity spaces of G -invariant maps between linear representations are now replaced by multiplicity categories, whose dimensions are the Hom-spaces in the category of coherent sheaves.

There is a preferred family of simple representations that foliate the BFM space. Every such representation is symplectically-induced from a 1D representation of a certain Levi subgroup of G : it is the Fukaya category of a flag variety of G (similarly to holomorphic induction, the Borel-Weil construction of irreducible representations of G). This is the same governing structure emerging in KRS, where BCs of the 3D TQFT are associated to a holomorphic Lagrangian submanifold.

Meta-statement, [192]: Pure topological gauge theory in 3D for a compact Lie group G is equivalent to the Rozansky-Witten theory for the BFM space of the Langlands dual Lie group G^V .

2D TQFT can be thought of as a higher analogue of cohomology (Fukaya-Floer theory of a symplectic manifold is like a refinement of ordinary cohomology), according to which gauged TQFTs are described in terms of equivariant cohomology. This describes categorification of representation theory of a compact Lie group G , with representations being topological.

In the RW model, one must describe geometrically two functors from the 2-category of linear categories with G -action to linear categories:

1. The forgetful functor, remembering the underlying category (describing the pre-gauged TQFT). This is represented by a regular representation of G , i.e. categories of coherent sheaves over the Lagrangians $\text{Spec}H_*(\Omega G)$.
2. The invariant category, generating the gauged TQFT, represented by a trivial representation of G , $\mathfrak{t}_{\mathbb{C}}/W \equiv T_1^*T_{\mathbb{C}}^V/W$.

6.5 From Homological mirror symmetry to 3D mirror symmetry

The concluding part of this section is a very succinct overview of some key features emerging in the work of [192], which is of particular importance to us. As we shall see, his work is deeply related to that of Moore and Tachikawa, thanks to their relation to Coulomb branches of star-shaped quivers, briefly overviewed in section 3.5.

To the best of our knowledge, the correspondence in between these three setups has only been spelled out in our work, [3]. The crucial step in understanding these different setups is looking at how mirror symmetry is realised in each case. As we shall see, this

builds a dictionary that can be used for furthering the study of more exotic underlying mathematical structures.

The 2-category of KRS and the Bezrukavnikov-Finkelberg-Mirkovic (BFM) space

The image of a point $Z(\text{pt})$ is an object in the 3D bordism 3-category. Lurie’s generator for pure 3-dimensional gauge theory should have categorical depth 2. His proposal for such generator is a 2-category associated to a certain holomorphic symplectic manifold, whose existence had already been conjectured by KRS and KR. When X is compact, this 2-category should generate the RW-theory of X . In particular, its Hochschild cohomology (a 1-category with braided tensor structure), should be a dg-refinement of the derived category of coherent sheaves on X , $\sqrt{\mathfrak{Coh}(X)} \cong \text{KRS}(X)$. Among objects in $\text{KRS}(X)$ are smooth holomorphic Lagrangians $L \subset X$. More general objects are coherent sheaves of \mathcal{O}_L -linear categories on such L . Given two Lagrangians, $L, L' \in X$, $\text{Hom}(L, L')$ is a sheaf of categories supported on $L \cap L'$, and a $(\mathfrak{Coh}(L), \otimes) - (\mathfrak{Coh}(L'), \otimes)$ bimodule.

Localising at L , we choose a formal neighbourhood identified symplectically with T^*L such that we can regard (locally) as the graph of a differential $d\Psi$ for a potential function $\Psi : L \rightarrow \mathbb{C}$. Locally, when this identification is valid, $\text{Hom}(L, L') \cong \text{MF}(L, \Psi)$. The corresponding figure is 17.




Figure 17. BFM space for an abelian gauge group, [192].

For the non-abelian case, instead, this description in terms of complex manifolds and potentials is inadequate, since the BFM space is not quite a cotangent bundle. The corresponding figure is 18.

6.5.1 Relation to our main example

The Hom category $\text{Hom}(L, L')$ for two Lagrangians $(L, L') \subset X$ with finite intersection supplies a 2D TFT for framed surfaces (following from matrix factorisations). X itself aims at defining RW-theory, and each L, L' is a BC for it.

$$L, L' : \text{Id} \rightarrow \text{RW}_X \tag{6.26}$$




Figure 18. BFM space for a non-abelian gauge group, [192].

are morphisms from the trivial 3D theory Id to RW_X viewed as functors from Bord_2 to the 3-category of linear 2-categories and $\text{Hom}(L, L')$ of natural transformations between these morphisms is the generator of the theory in [192]. Note that this is equivalent to the definition of the triple (3.51) in our main example in section 3, since the RW_X -theory lives on a Calabi-Yau manifold, and the Lagrangians L, L' act as the left- and right- moment maps projecting the identity element Id . This basically shows that the method put forward by [192] is an abelianisation procedure enabling to foliate the BFM space for any given pair of Lagrangian submanifolds.

The Hochschild homology

$$\text{HH}_{\dim L} \text{Hom}(L, L') \simeq \Gamma \left(L \cap L'; (\omega \otimes \omega')^{1/2} \right) \quad (6.27)$$

A domain wall between different TQFTs generalises the notion of BCs. This is an adjoint pair of functors between the TQFTs meeting certain dualisability conditions. A BC is a domain wall with a trivial TQFT. Just as a holomorphic Lagrangian is expected to define a BC for RW_X , a holomorphic Lagrangian correspondence

$$X \leftarrow C \rightarrow Y \quad (6.28)$$

should define a domain wall between RW_X and RW_Y . This can be used for comparing gauge theories of different groups.

$$\text{BFM}(G^V) \leftarrow \text{BFM}(G^V) \times_{\mathfrak{t}_C/W} \mathfrak{t}_C \rightarrow T^*T_C^V \quad (6.29)$$

The correspondence matches an adjoint pair of restriction-induction functors between categorical T - and G -representations. Induction from a category \mathcal{C} with topological T -action is affected by string topology with coefficients in the flag variety G/T :

$$\text{Ind}(\mathcal{C}) \equiv C_* \Omega G \otimes_{C_* \Omega T} \mathcal{C} \quad (6.30)$$

Restricting the G -action to T , the flag manifold G/L is a transformation from L -gauge theory to a T -gauge theory, given by composition of the symplectic induction and string topology domain walls

$$T(L^V) \xrightarrow{\text{SInd}} T(G^V) \xrightarrow[\text{Today}]{} \text{BFM}(G) \xrightarrow{\text{ST}} \text{BFM}(T^V) \equiv T^* T_C^V \quad (6.31)$$

The flag manifold G/L is therefore a transformation from L -gauge theory to T -gauge theory given by the composition of symplectic induction and string topology domain walls.

One of the main aims of the work by KRS was that of furthering the understanding of 3D mirror symmetry in terms of homological mirror symmetry between categories of Lagrangian submanifolds. In the concluding part of this work, we show that this correspondence can equivalently be realised in terms of the magnetic quiver prescription, which can be further extended to the case involving non-invertible defects in presence of two cones⁴².

7 Holographic Dualities and Beyond




Figure 19. Main types of dualities arising in due course in the remainder of our treatment.

7.1 Holographic duality

Inspired by black hole mechanics, the Holographic Principle was first proposed by Gerard 't Hooft, and subsequently given a more precise string-theoretic interpretation by Susskind. It is a property of String Theory as well as a supposed property of QG, stating that, the description of a volume of space can be encoded in its boundary. The AdS/CFT correspondence provides a non-perturbative formulation of String Theory with certain boundary

⁴²We will come back to this in section 20.

conditions, and constitutes one of the most successful realisations of the Holographic Principle.

One of the original motivations for studying the AdS/CFT correspondence was developing a non-perturbative formulation of string theory.

The 4D $\mathcal{N} = 4$ SYM/ string theory on $\text{AdS}_5 \times S^5$ case

For the particular case of 4D $\mathcal{N} = 4$ SYM/ string theory on $\text{AdS}_5 \times S^5$, symmetries play a crucial role in establishing the holographic correspondence. Specifically, the isometry invariance group of AdS_5 , namely $SO(4, 2)$, corresponds to the symmetry group of CFT_4 s, whereas the isometry group of S^5 , $SO(6) \simeq SU(4)$, corresponds to the R-symmetry group of $\mathcal{N} = 4$ SYM.

This statement can be generalised to arbitrary dimensions, in the sense that the isometry group of AdS_{D+1} is $SO(D, 2)$, corresponding to the conformal group in $(D - 1, 1)$ dimensions. The boundary of AdS_{D+1} is \mathbb{R}^D in Poincare' coordinates, and $\mathbb{R}_t \times S^{D-1}$ in global coordinates, which is conformal to \mathbb{R}^D . Figure 20 shows this.




Figure 20.

A crucial feature of the correspondence is the fact that AdS acts as a quantum mechanical box, meaning light can travel from the bulk to the boundary in finite time, implying the two are in causal contact with each other. Consequently to this, unlike the case of Minkowski spacetime, the notion of S-matrix falls short from being well-defined in AdS. On the other hand, the well-defined observables at hand are correlation functions of fields, with sources placed on the conformal boundary.

For the purpose of our treatment, we only need to state what the key points of the holographic correspondence imply from the point view of the bulk-boundary interaction, which is what we will now turn to.

In its original formulation, one expects there to be a correspondence in between the theory of open strings living on N D -branes, i.e. $\mathcal{N} = 4$ SYM with $SU(N)$ gauge group, and the gravitating theory corresponding to fields of the Hawking radiation existing in the background curved by the D -branes. Indeed, the Hawking radiation process plays a crucial role in the AdS/CFT correspondence. Considering the D -branes as endpoints of open strings, then string theory with $D3$ -branes has three ingredients:

1. The open strings existing on the $D3$ -branes, giving a theory that reduces to 4D $\mathcal{N} = 4$ SYM in the low-energy limit.

2. The closed strings existing in the bulk spacetime, giving a theory that is SUGRA coupled to the massive modes of the string. In the low-energy limit, only SUGRA will remain.
3. The interactions in between the two theories, of which Hawking radiation is a manifestation.




Figure 21.

Generalisations

To better understand the quantum aspects of our 4D world, some have considered lower-dimensional mathematical models, where spacetime is assumed to be 3-dimensional. The advantage of such a lower-dimensional setup is the fact that the description of the gravitational field is greatly simplified, and one can therefore use standard QFT techniques, circumventing the need to resort to string theory in 4D.

Two of the most important proposals for such lower-dimensional correspondence are the following:

1. In 1986, Brown and Henneaux, [207], realised that quantum gravity in 3D is closely related to a particular CFT_2 , namely Liouville Field Theory.
2. Edward Witten related 3D gravity in AdS to a CFT with monster group theory.

Both cases provide examples of the AdS/CFT correspondence that do not require the full apparatus of string or M-theory.

7.1.1 Liouville Theory

Liouville Theory describes the dynamics of a field, φ on a 2D space, subject to the potential

$$V(\varphi) = e^{2b\varphi}, \quad (7.1)$$

where b plays the role of the coupling constant. Its action reads

$$S[\varphi] = \frac{1}{4\pi} \int d^2x \sqrt{g} \left(g^{\mu\nu} \partial_\mu \varphi \partial_\nu \varphi + Q\mathcal{R}\varphi + \lambda' e^{2b\varphi} \right), \quad (7.2)$$

where $g^{\mu\nu}$ is the 2D metric, \mathcal{R} the Ricci scalar, and λ' is often referred to as the cosmological constant.

The equations of motion from (7.3) read

$$\frac{1}{\sqrt{g}} \partial_\mu (\sqrt{g} g^{\mu\nu} \partial_\nu \varphi) = \frac{1}{2} Q\mathcal{R} + \lambda' b e^{2b\varphi}. \quad (7.3)$$

In Euclidean signature, (7.3) reduces to

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) \varphi = \frac{1}{2} Q\mathcal{R} + \lambda' b e^{2b\varphi}. \quad (7.4)$$

Assuming the Euclidean metric takes the following form

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = dzd\bar{z}, \quad (7.5)$$

the components of the energy-momentum tensor read

$$T_{z\bar{z}} = T_{\bar{z}z} = 0 \quad (7.6)$$

and

$$T_{zz} = (\partial_z \varphi)^2 + Q\partial_z^2 \varphi, \quad T_{\bar{z}\bar{z}} = (\partial_{\bar{z}} \varphi)^2 + Q\partial_{\bar{z}}^2 \varphi, \quad (7.7)$$

with

$$\partial_{\bar{z}} T_{zz} = 0, \quad \partial_z T_{\bar{z}\bar{z}} = 0. \quad (7.8)$$

Each one of the non-vanishing components (7.7) generates a Virasoro algebra with central charge

$$c = 1 + 6Q^2. \quad (7.9)$$

For such Virasoro algebras, $e^{2\alpha\varphi}$ is a primary field with conformal dimension

$$\Delta = \alpha(Q - \alpha). \quad (7.10)$$

For the theory (7.3) to be conformally-invariant, the field $e^{2b\varphi}$ must be exactly marginal, i.e. it must have conformal dimension such that

$$\Delta(b) = 1 \quad \Rightarrow \quad Q = b + \frac{1}{b}. \quad (7.11)$$

The last condition in (7.11) ensures (c, Δ) is invariant under the duality

$$b \longrightarrow \frac{1}{b}. \quad (7.12)$$

Relation to other theories

Liouville theory admits two different supersymmetric extensions, namely with $\mathcal{N} = 1$ and $\mathcal{N} = 2$. Furthermore, it can be obtained from the WZW model.

Liouville theory also appears in the context of bosonic string theory. In its semiclassical limit it coincides with Jackiw-Teitelboim (JT)-gravity, which is another formulation of 2D gravity. To see how JT-gravity can be recovered from Liouville theory, we start from a double Liouville theory built combining a spacelike and a timelike Liouville theory.

To begin with, we start by defining the following CFT action

$$S_L = \int d^2x \sqrt{g} \left[(\partial_\mu \phi)^2 - (\partial_\mu \chi)^2 + \mu e^{2b\phi} - \mu e^{-2b\chi} \right] + 2 \int d\theta (Q\phi + q\chi) \mathcal{K}^{(0)}. \quad (7.13)$$

Under the field redefinition

$$\Phi \stackrel{def.}{=} -\frac{1}{2b}(\phi + \chi) \quad , \quad \rho \stackrel{def.}{=} \frac{b}{2}(\phi - \chi), \quad (7.14)$$

(7.13) reduces to

$$S = \int d^2x \sqrt{g} \left[-4(\partial\rho)(\partial\Phi) - 2\mu e^{2\rho} - \sinh(2b^2\Phi) \right] - 4 \int d\theta \Phi \mathcal{K}^{(0)}. \quad (7.15)$$

Taking ρ to be the conformal factor of the JT-gravity metric

$$ds_{JT}^2 = e^{2\rho} (dr^2 + r^2 d\theta^2), \quad (7.16)$$

(7.15) ultimately becomes

$$S = -2 \int d^2x \sqrt{g_{JT}} \left[\Phi \mathcal{R} + \mu e^{2\rho} - \sinh(2b^2\Phi) \right] - 4 \int d\theta \sqrt{\gamma_{JT}} \Phi \mathcal{K}, \quad (7.17)$$

which in the semiclassical limit (namely $\lim_{b \rightarrow 0}$ with $\mu b^2 = 1$) is exactly equivalent to JT-gravity.

7.2 The Chern-Simons/Wess-Zumino-Witten (CS/WZW) correspondence

One realisation of the holographic principle in QFT between 3D G -CS theory as the bulk field theory, and the 2D WZW model on a suitable Lie group G as the boundary theory. This case was known and understood well before the holographic principle was first formulated.

Unlike the AdS/CFT correspondence, this is an actual theorem, stating the normal equivalence in between the space of quantum states of CS theory on a surface Σ , and the space of conformal blocks on the WZW model.

The FRS construction/classification provides a rigorous classification of all rational full CFT_2 s, where full means that these theories are defined at all genera, namely on all 2D cobordisms.

The construction of CFT_2 s relies on the following:

1. A local geometric aspect.
2. A global topological aspect. Taking in the spaces of conformal blocks one element such that this choice globally fits together under gluing of 2D cobordisms defined the sewing constraint.

Because of this, the FRS formalism provides a mathematical formulation of the holographic correspondence.

Given a vertex operator algebra (VOA) that controls the CFT locally, then, for RCFTs, their representation category is a MTC (in turn obtained from, either, a Hopf algebra, or loop groups representations), and the Reshetikhin-Turaev or Turaev-Viro construction defines the 3D TQFTs, namely the 3D CS theory.

Full CFT_2 s are relevant in String Theory, since these correspond to the vacua that string perturbation theory series are perturbing about. The FRS classification is part of the mathematical identification of the landscape of string theory vacua,

RCFT₂s are a small part of all possible CFT_2 s (indeed, usually, any deformed CFT_2 is non-rational). However, RCFT₂s (namely the WZW-model) plays an important role as the internal parts of string theory KK-compactifications.

7.3 Jackiw-Teitelboim (JT)-gravity from CS-theory

In its first-order formulation, the Einstein-Hilbert action for 3D $SL(2, \mathbf{R})$ CS gravity is defined as the difference of two CS actions

$$S_{EH} = S_{CS}[A] - S_{CS}[\bar{A}]. \quad (7.18)$$

For AdS_3 in Lorentzian signature, the connections A and \bar{A} are $\mathfrak{sl}(2, \mathbf{R})$ -valued 1-forms. In the fundamental representation, the generators of the Lie algebra $\mathfrak{sl}(2, \mathbf{R})$ read

$$L_o = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{pmatrix}, \quad L_1 = \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix}, \quad L_{-1} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad (7.19)$$

obeying the commutation relation

$$[L_m, L_n] = (m - n) L_{m+n}. \quad (7.20)$$

The e.o.m. for the connections in (7.3) read

$$F = dA + A \wedge A = 0, \quad \bar{F} = d\bar{A} + \bar{A} \wedge \bar{A} = 0 \quad (7.21)$$

whose solution is pure gauge

$$A = g^{-1} dg, \quad \bar{A} = \bar{g}^{-1} d\bar{g} \quad g, \bar{g} \in \mathfrak{sl}(2, \mathbf{R}). \quad (7.22)$$

In terms of (t, z, r) -coordinates, where r denotes the holographic direction, upon choosing suitable boundary conditions, (7.22) reads as follows

$$A = b(r)(d + a(z))b(r) \quad , \quad b(r) = r^{L_o} \quad (7.23)$$

$$\bar{A} = \bar{b}(r)(d + \bar{a}(z))\bar{b}(r)^{-1} \quad , \quad \bar{b}(r) = r^{\bar{L}_o}. \quad (7.24)$$

In presence of a boundary, the CS action needs to be equipped with a suitable boundary term, ensuring admits a well-defined variational principle

$$\delta S_{EH} = \delta S_{CS}[A] - \delta S_{CS}[\bar{A}] \quad (7.25)$$

$$\Rightarrow \delta S_{EH} + \delta S_{bdy} = \frac{1}{4\pi G} \int_{\partial M_3} \epsilon_{ab} f^a \wedge \delta e^b, \quad (7.26)$$

with e^a the source and f^a the expectation value of the dual operator. In particular, the operator dual to the boundary vielbein is the stress tensor

$$\delta S = 4 \int_{\partial M_3} d^2 x (\det e) T_a^i \delta e^b. \quad (7.27)$$

Interestingly, upon choosing a particular boundary term and subsequently taking the dimensional reduction of either CS action, we end up with a BF-theory

$$\begin{aligned} 8\pi G S_{CS} &= \frac{1}{2} \int_{M_3} d^3 x \epsilon^{\mu\nu\rho} \text{Tr} \left(A_\mu \partial_\nu A_\rho + \frac{2}{3} A_\mu A_\nu A_\rho \right) \\ &= \int_{M_3} d^3 x \text{Tr} (A_\varphi F_{tr} + A_r \partial_\varphi A_t) + \frac{1}{2} \oint_{\partial M_3} \text{Tr} A_t^2. \end{aligned} \quad (7.28)$$

The corresponding 2D theory living on the conformal boundary is obtained upon taking the $\lim_{r \rightarrow 0}$ of (7.28). Imposing the following boundary conditions

$$A_t = A_\varphi \Big|_{\partial M_3} \quad \text{such that} \quad \phi \equiv A_\varphi \quad \text{and} \quad \partial_\varphi = 0, \quad (7.29)$$

(7.28) turns into

$$\boxed{S_{BF} = \int_{M_2} \text{Tr} (\phi F) + \frac{1}{2} \oint_{\partial M_2} \text{Tr} (\phi^2)}, \quad (7.30)$$

where the first term is the usual BF-term for the gauge group $G = SL(2, \mathbf{R})$, whereas the second term is the Schwarzian-type boundary term.

The degrees of freedom in this resulting action, (7.30), are: a gauge field A_μ , with associated field strength $F_{\mu\nu}$, along with an $SL(2, \mathbf{R})$ -valued scalar field ϕ . The latter corresponds to a boundary degree of freedom (also known as the particle-on-a-group).

subsectionBranes in QFTs and String Theory

D-branes

In the cobordism formulation of a higher-dimensional QFT, a D-brane is a type of data assigned to the QFTs boundaries of cobordisms.

Key examples include the following:

1. RCFT₂s (the full rational CFT₂, are classified using the FRS-formalism⁴³, by means of:
 - A MTC, \mathcal{C} , the category of representations of the VOA of the CFT₂.
 - A special symmetric Frobenius algebra object, A , internal to \mathcal{C} .

Hence, a D-brane of the theory is an A -module in \mathcal{C} .

2. The $(\infty, 1)$ -categorical formulation of TFT₂s, leads to the A- and B-models. The A-model's branes form the Fukaya category of the target space. The B-model's branes form a stable $(\infty, 1)$ -category of complexes. The WZW-model is a particular example of this, where the target space is a simple Lie group $X \equiv G$, and the background field is a circle 2-bundle with connection on G , representing the background field, known as the Kapustin-Rozansky (KR) field. The study of Kontsevich's homological algebra reformulation of mirror symmetry, and the study of derived D-brane categories constituted a field of pure maths on its own.
3. In type-IIA SUGRA, they are the D0-, D2-, D4-, D6-, D8-, F1-, NS5-branes.
4. In type-IIB SUGRA, they are D1-, D3-, D5-, D7-, F1-, NS5-, (p, q) -5, $[p, q]$ -7 branes.
5. In M-theory, M2-, M5-, M9-, M-wave.

The topological part of D-brane actions behave as generalised symmetry operators. In particular, they exhibit:

1. Confinement/deconfinement transition, via the Hanany-Witten effect via D3-brane nucleation.

The Hanany-Witten mechanism, [610], is realised in brane web setups. As shown in figure 1, it consists in the extension of F1-branes between D3- and D5-branes as they pass across each other. This behaviour is naturally arising within the context of 4D QFTs with mixed t'Hooft anomaly. We will briefly outline this example in Part III of the present treatment.

2. Fusion rules via the Myers effect. This is an additional feature connecting braneweb constructions with symmetry generators in 4D QFTs. In particular, the Myers effect, well known in the context of D0-D2 branes, is analogous to the fusion rules arising from gauging higher-form symmetries in QFT.

⁴³FRS stands for Fuchs, Runke, and Schweigert, who provided a rigorous construction and classification of all rational full 2D CFTs, where full means that these theories are defined at all genera, namely on all 2D cobordisms.




Figure 22.

7.3.1 (P)SU(N) in 4D $\mathcal{N} = 1$ Super Yang Mills

To put in practice what we have said up to now, we briefly reproduce an example outlined in [617], in the context of symmetries of 4D QFTs in relation to D-branes structures. The 4D theory is a gauge theory with a 0-form symmetry, \mathbb{Z}_M , with background field A_1 , and a 1-form symmetry, Z_M , with background field B_2 , related by a mixed 't Hooft anomaly

$$\mathcal{A} = -\frac{2\pi}{M} \int A \cup \frac{\mathcal{P}(B_2)}{2}, \quad (7.31)$$

where $\mathcal{P}(B_2)$ is the Pontrjagin square of B_2 .

The generators of the 0- and 1-form symmetries are generated by 3D and 2D topological defects $D_3^g(M_3), D_2^h(M_2)$, respectively, such that their fusion rules obey group-like structure relations

$$D_p^{g_1}(M_p) \otimes D_p^{g_2}(M_p) = D_p^{g_1 g_2}(M_p). \quad (7.32)$$

Due to the anomaly (7.31), the generators of the 0-form symmetry transform nontrivially, namely

$$D_3^g(M_3) \rightarrow D_3^g(M_3) \exp\left(\int_{M_4} -\frac{2\pi i}{M} \frac{\mathcal{P}(B_2)}{2}\right), \quad (7.33)$$

with $\partial M_4 = M_3$. In particular, when gauging the 1-form symmetry, (7.33) is inconsistent. To solve this problem, we need to dress $D_3(M_3)$ with a minimal TQFT $\mathcal{A}^{M,p}$ featuring a 1-form symmetry Z_M that cancels the anomaly. For this example, this is a $U(1)_M$ theory, and the dressed defect is redefined as follows

$$\mathcal{N}_3^{(1)} = D_3^{(1)} \otimes \mathcal{A}^{M,1}, \quad (7.34)$$

with non-invertible fusion rules

$$\mathcal{N}_3^{(1)} \otimes \mathcal{N}^{(1)} = \mathcal{A}^{M,2} \mathcal{N}_3^{(2)}, \quad \mathcal{N}_3^{(1)} \otimes \mathcal{N}^{(1)\dagger} = \sum_{M_2 \in H_2(M_3, \mathbb{Z}_M)} \frac{(-1)^{Q(M_2)} D_2(M_2)}{|H^0(M_3, \mathbb{Z}_M)|}, \quad (7.35)$$

where

$$D_2(M_2) = e^{i2\pi \int_{M_2} \frac{b_2}{M}}, \quad (7.36)$$

with b_2 the gauge field for the 1-form symmetry. In relation to type-IIB string theory, these symmetry operators $\mathcal{N}_3^{(1)}$ are associated to D5-branes, whereas 't Hooft operators are associated to D3-branes, [617]. We will be referring to this example for later convenience in Part III when introducing Symmetry Topological Field Theories (SymTFTs).

7.4 From brane webs to quiver gauge theories



Figure 23.

Quivers encode the Lagrangian of a supersymmetric gauge theory. They take different forms according to the number of supercharges. For the case in which there are 8 of them, quiver diagrams encode the gauge symmetries, flavour symmetries, and matter field content. The fields featuring in a certain theory are contained in vector multiplets and hypermultiplets. The field content changes according to the dimensionality of the theory in question.

The quiver diagram encodes the representations of the gauge and matter fields under which they transform. Specifically, the vector multiplets transform under the adjoint representation of the gauge group representation, whereas the hypermultiplets transform under the bifundamental representation of the gauge group and the flavour group.




Figure 24.

Importantly, from the quiver, we can read the superpotential. In figure 7.4 we show the example of a framed Nakajima quiver variety. In this case, the corresponding superpotential reads as follows

$$\mathcal{W} = \text{Tr} \left(A_i^\alpha \Phi_j^i B^{j\beta} \right), \quad (7.37)$$

where A_j^α, Φ_j^i and $B^{j\beta}$ denote the chiral multiplets, the anti-chiral multiplets, and the adjoint chirals, respectively. The indices i, j are related to the gauge group, whereas the indices α, β are related to the flavour group.

7.5 Beyond holographic duality: adding boundaries

Boundary Conditions (BCs) of $\mathcal{N} = 4$ SCFTs

$AdS_4 \times S_2^{(A)} \times S_2^{(B)}$ fibered over a Riemann surface, Σ , the corresponding metric solution reads

$$ds^2 = f_4^2 ds_{AdS_4}^2 + f_1^2 ds_{S_2^{(A)}}^2 + f_2^2 ds_{S_2^{(B)}}^2 + 4\rho^2 (dr^2 + r^2 d\theta^2), \quad (7.38)$$

where the metric functions f_i, ρ are functions of h_1, h_2 defined on Σ , via (r, θ) . As explained in [290], the field theory in question is uniquely defined by the following set of parameters

$$N_{D5}^{(A)} \stackrel{def}{=} \frac{1}{\sqrt{g}} c_A, \quad , \quad N_{BS5}^{(B)} \stackrel{def}{=} \sqrt{g} d_B \quad (7.39)$$

$$L_A \stackrel{def}{=} \sqrt{g} l_A + \frac{2}{\pi} \sum_B N_{D5}^{(B)} \tan^{-1} \left(\frac{l_A}{k_B} \right), \quad K_B \stackrel{def}{=} \frac{1}{\sqrt{g}} k_B + \frac{2}{\pi} \sum_A N_{NS5}^{(B)} \tan^{-1} \left(\frac{k_B}{l_A} \right). \quad (7.40)$$

At this point, one could further redefine \hat{L}_i as the number of D3-branes ending from the right on the i^{th} D5-brane plus the number of NS5-branes to the left of this D5-brane, and with \hat{K}_i the number of D3-branes ending on the i^{th} NS5-brane plus the number of D5-branes to the left of this NS5-brane, leading to the brane-web intersection reported in figure 53.




Figure 25.

In a SCFT with $\mathcal{N} = 4$ supersymmetry, the Higgs and Coulomb branch are both conical and meet at their common origin. In general, there might also be mixed branches.

$\frac{1}{2}$ -BPS BCs in $\mathcal{N} = 4$ SYM

$\frac{1}{2}$ -BPS BCs in $\mathcal{N} = 4$ SYM with gauge group G admit a general classification in terms of a triple, (ρ, H, \mathcal{B}) , [126], where

$$\rho : \mathfrak{su}(2) \rightarrow \mathfrak{g} \tag{7.41}$$

denotes the embedding of the Lie algebra of $SU(2)$ in the Lie algebra of G , H is a subgroup of G resulting from the symmetry breaking induced by (7.41), whereas \mathcal{B} denotes the 3D boundary field theory with symmetry H .

7.5.1 Islands in JT-gravity

The holographic description of the entanglement entropy for Hawking radiation in arbitrary spacetime dimensions can be described in 3 different ways. This follows from the extension of the 2d JT-gravity model to arbitrary dimensions relying upon the Randall-Sundrum model with a brane embedded in AdS_{d+1} . Maldacena et al.'s treatment [211] is equivalent to the brane (EFT) description. The bulk perspective shows that it is indeed reasonable to extend JT-gravity results to higher-dimensions relying upon dualities. In particular, wedge holography [236] looks very promising for addressing vacuum transitions in higher dimensions.

1. Boundary description




Figure 26. Both of the QMs and field theory systems are prepared in independent thermofield double states.

2. Brane description




Figure 27. Introducing a 2D holographic geometry (JT-gravity), dual to the entangled state of QM_L and QM_R . The CFT_2 coupling to JT-gravity is the same as the one featuring in the bath.

3. Bulk description

7.5.2 Conformal interfaces as boundary states

The idea of the replica trick, at first applied within the context of statistical field theory and condensed matter physics, basically consists in taking n -copies of a certain system and




Figure 28. The doubly-holographic description, where the holographic CFT_2 is replaced by a bulk region with geometry of an AdS_3 black hole.

relating its behaviour to that of the original system, i.e for $n = 1$. The reason for doing so is that, in principle, there might be noncommutative behaviour emerging from permutations of different subregions of the n -copied system, which in turn would highlight the presence of interesting entangling features that are in turn scale-dependent, i.e. their behaviour is determined by the size of one copy.

Translating this idea in a suitable language for our purposes, i.e. describing quantum transition in between different spacetimes,⁴⁴ there is one additional key point to bare in mind. Quantum transitions are really meant to describe *local* spacetime transformations. As an example, describing uptunneling from AdS would really mean considering a portion of the domain at a given instant of time (in global coordinates) and then defining its corresponding density matrix by post-selection according to Takayanagi et al.'s prescription discussed in the previous section. The final state $|\varphi\rangle$ will be defined such that the covariant holographic entanglement entropy satisfies the 2-step minimisation-extremisation (min-extr) of the expectation value of $Area(\Gamma_A)$. ρ_A is therefore the density matrix associated to the states defined on such subregion at the boundary of AdS. Being this a portion of the entire domain at r_∞ , we can think of A as arising by quotienting w.r.t. Z_n -symmetry, with n determined by the particular choice of the domain A . However, as also explained in the appendix, even for the simplest $U(1)$ -case, an anomaly emerges, since the symmetry is broken in the quantum theory by the state and the partition function. Restoring it at the level of the partition function requires adding Wilson loops, i.e. additional operators to the quantum theory. In turn, these operators can be thought of as the lower-dimensional counterparts of background fields living in one spatial dimension higher. Their coupling will ultimately lead to the definition of Chern-Simons terms, i.e. defects added to the bulk theory in order to compensate for the anomaly featuring in the original QM theory. This procedure is known as the *anomaly-inflow mechanism*.

Our main interest is addressing the issue from the quantum gravity perspective. In order to do so, we rely upon the holographic dictionary of the AdS_{d+1}/CFT_d correspondence, at first starting with $d = 2$, from which some major considerations can be extracted and later on extended to higher-dimensional setups.

⁴⁴and as it will become clear later on in the treatment.

We start by considering 2 CFTs separated by a conformal interface, which basically provide a special example of extended operators, such as Wilson lines, D-branes, or other extended sources. Upon exchanging the role of space and time, these conformal boundaries can be mapped to boundary states, following the so-called *folding trick*. As a result of this procedure, we end up with a BCFT. The boundary conditions defined at the interface can be suitably chosen accordingly to the setup we wish to describe. 2 particular examples are provided by the totally-reflecting and the topological BCs.

The boundary arising from the folding trick can really be thought of as a special case of a *defect*, for which an entropy can be derived from the entanglement entropy of a (1 + 1) CFT. Specifically, the entanglement entropies with and without the boundary defect respectively read

$$S_1 = \frac{c}{6} \ln \frac{L}{\epsilon} + \ln(g_B) + \alpha_0 \quad , \quad S_2 = \frac{c}{6} \ln \frac{L}{\epsilon} + \alpha_0 \quad (7.42)$$

where c is the central charge of the CFT₂, ϵ is the UV cutoff introduced for regularisation purposes, L is the length of the spatial region on which the entanglement entropy is being evaluated, α_0 is an overall constant, and g_B is the ground state degeneracy. Upon taking the difference in between the two quantities in (7.42), we are left with the entropy of the defect

$$S_{\text{bdry}} = S_1 - S_2 = \ln(g_B) \quad . \quad (7.43)$$

The defect, therefore, contributes with an additional term to the partition function, which asymptotically reads

$$S = \ln g_B \quad , \quad g_B = \langle 1 \rangle = \langle 0 | D | 0 \rangle \quad (7.44)$$

with D defining the *defect operator*

$$D = g|0 \rangle \langle 0| + \dots \quad . \quad (7.45)$$

In string theory, e.g., the role of the g -factor is played by the tension of a D-brane. It is interesting to note that this is exactly the term entering in the Hamiltonian formulation of quantum transitions by implementing the junction conditions, with the brane tension separating different spacetimes.

The *degeneracy* of the ground state is therefore defined by the overlap between the boundary state and the vacuum

$$g_B = \langle 0 | B \rangle \quad . \quad (7.46)$$

The *g-theorem* states that the value of g_B must decrease under the boundary RG-flow [206].

Given g_B , the *distance* in between the two original CFTs is defined as follows

$$d(1, 2) = \min \sqrt{\ln g_B} \quad . \quad (7.47)$$

From this, we therefore understand that BCFTs, such as the case of interest for us being $\rho_A = \text{Tr}_B \rho$, are incredibly rich in structure and carry many interesting and hidden features that will turn out essential in the forthcoming sections. In particular, the *boundaries* of the BCFT carry d.o.f. associated to the degeneracy of the ground state and therefore signal whether a defect needs to be added to the path-integral evaluation when taking n -copies of the given subregion where the BCFT lives. Such defect is, in turn associated to an anomaly induced by the anomaly-inflow mechanism of a bulk background field which needs to be added to the quotiented geometry.

In terms of the von Neumann formulation, the entanglement entropy reads

$$S_A = -\text{Tr}_A (\rho_A \ln(\rho_A)) = - \lim_{n \rightarrow 1} \frac{d}{dn} \text{Tr}_A(\rho_A^n) \quad , \quad (7.48)$$

where in the last step of (7.48) we implemented the *replica trick*.

For integer values of n -replicas, $\text{Tr}(\rho_A^n)$ is the partition function on a Riemann surface, i.e.

$$\text{Tr}_A(\rho_A^n) \sim \mathcal{Z}_n = \int [d\varphi_1 \dots d\varphi_n]_{\mathcal{M}_n} \exp \left[- \int_{\mathcal{M}_n} d^2x \mathcal{L}^{(n)}[\varphi_1, \dots, \varphi_n](x) \right] \quad . \quad (7.49)$$

Invariance of the boundary theory under Z_n -symmetry corresponds to the Lagrangian being preserved under permutations of the n -copies of the QFT, i.e.

$$\mathcal{L}^{(n)}[\varphi_1, \dots, \varphi_n](x) \dots, \sigma \varphi_n](x) = \mathcal{L}[\varphi_1](x) + \dots + \mathcal{L}[\varphi_n](x) \quad , \quad (7.50)$$

with $\sigma \varphi_i = \varphi_{i+1 \bmod n}$. The action of σ belonging the symmetry group Z_n and its inverse σ^{-1} are associated to a *twist fields*, whose correlation function is proportional to (7.49). The partition function of the n -copied system can therefore be used for redefining the entanglement entropy in the $\lim_{n \rightarrow 1}$, and (7.48) therefore reduces to

$$S_A = - \lim_{n \rightarrow 1} \frac{d}{dn} \mathcal{Z}_n \quad . \quad (7.51)$$

With this setup, we are now ready to derive the generalised expression for the entanglement entropy, where the contribution from quantum extremal surfaces adds to the von Neumann entropy.

Part III

From knot invariants to QFTs

8 Summary and Motivations of Part III

8.1 Motivations

Since first encountered, [108], 6D $\mathcal{N} = (2, 0)$ SCFTs, also referred to as *theory X*,⁴⁵, have attracted great interest in, both, the mathematics and high-energy physics community alike. From the high-energy physics point of view, they are essential for deriving lower-dimensional SCFTs, [109, 112, 310, 378], whose features are much better understood⁴⁶. On the other hand, from a purely mathematical perspective, they provide an extremely interesting and rich setup where to exploit the present-day knowledge of representation theory, [110, 111].

In recent years, the two communities have been brought together by fascinating developments on either side, [312], highlighting the importance of bridging the gap in between different formalisms, with category theory playing a key role in pursuing this task. Indeed, the pioneering works of [313–315] led to the discovery of modular tensor categories in anyon gauging theory. Since then, application of such deep mathematical language inspired applicability to higher-dimensional QFTs, by identifying their symmetries with topological defects, [316]. Of particular importance to us is the richness of higher-categorical theory applied to topological orders (TOs), [352–360], and symmetry topological field theories (SymTFTs), [102, 103].

Embracing the thrust of recent developments in the field, the present work is meant to be the first of a series of papers where the author applies the techniques of category theory to furthering the understanding of 6D $\mathcal{N} = (2, 0)$ SCFTs and theories derived from them. In doing so, we will attempt, wherever possible, to provide an understanding of the procedure as well as of the outcome of our findings, from, both, a mathematician’s and a theoretical physicist’s point of view. Hopefully, this will further motivate both communities to benefit from each other’s influence.

This part of our work aims at furthering the understanding of the categorical structure arising from dimensional reduction of 6D $\mathcal{N} = (2, 0)$ SCFTs. In particular, our aim is that of identifying a criterion to distinguish between intrinsic and non-intrinsic non-invertible symmetries⁴⁷ in terms of the quantum dimension⁴⁸. In doing so, we rely upon their description in terms of relative field theories, as described in [367] in terms of the Freed-Moore-Teleman setup, [102, 103].

⁴⁵Please see appendix 3 for some brief overview of its importance in pure mathematics and the Geometric Langlands Program.

⁴⁶We review some key features of its 4D descendants in appendix I.

⁴⁷See, for example, [329, 332, 333, 366–371, 539] for explanation on the current state of the art.

⁴⁸The total quantum dimension is defined from the braiding and fusion rules of the underlying categorical structure of the topological symmetries of the theory in question. Further explanation will be provided in due course.

As previously argued, class \mathcal{S} theories are special in the sense that they are absolute, and admit a quiver gauge theory description. Their TFT counterparts are the so-called minimal models, which, in turn are characterised by a higher-categorical structure, as reviewed in section 14. The multiplicities of the different superselection sectors might exceed unity, thereby signalling the possibility of information storage, correspondingly leading to a decrease in \mathcal{D} . Given that the definition of the entanglement entropy involves a partial tracing of the total boundary state, it might be that the Rényi entropy shows a characteristic periodicity \mathfrak{p} signalling that the state defining the reduced density matrix can be factorised under suitable redefinition of the modular operators involved in the definition of the link state itself.

Our main proposal is that the total quantum dimension of the gaugeable algebra implementing the gauging in the bulk Sym TFT is a key quantity to distinguish in between non-invertibility being intrinsic or non-intrinsic. Specifically, for the intrinsic case, multiplicity is greater w.r.t. the non-intrinsic case, thereby signalling the possibility for additional d.o.f. to be stored in certain superselection sectors of the resulting absolute theory.

Our results extend arguments proposed in [333], where the authors also proposed a way of distinguishing intrinsic from non-intrinsic non-invertibility in 2D by means of the quantum dimension of the non-invertible defect.

8.2 Outline and main results

The present Part contains the main findings first presented in an original work by the same author, and serves as a bridge between the preliminary sections of the present work and the remainder of this thesis. The content of the following two sections is as follows:

1. In section 9 we introduce the symmetry topological field theory (SymTFT) construction proposed by Freed, Moore, and Teleman for describing supersymmetric gauge theories descending from dimensional reduction of 6d $\mathcal{N} = (2, 0)$ SCFTs. In particular, we focus on the non-invertible defects arising in between class \mathcal{S} theories. Our main focus is on the following:
 - We highlight the importance of defining the Drinfeld center for the case in which non-invertibility is non-intrinsic, and relate this to the need of generalising the 3D mirror symmetry/homological mirror symmetry correspondence explained in section 6. This prepares the stage for the more detailed analysis to which sections 19, 5, 20, and 21 are devoted.
 - Exploiting the relation in between the fusion rules of such defects and the partition function on the absolute 4D theory, we show that the quantum dimension of the gaugeable algebra in the intrinsic case always exceeds that featuring in the non-intrinsic case. We thereby propose this as a general criterion to be applied to class \mathcal{S} theories descending from 6D $\mathcal{N} = (2, 0)$ SCFTs.
2. In section 10, we then turn to one of the key topics bridging with the remainder of our treatment, namely instantons. After having overviewed how they arise in QQM,

QFT, 4D supersymmetric gauge theories, and oriented string theory, we explain their relation to the ADHM construction, and the Higgs branch moduli space. In the concluding part of the same section, we then introduce the WKB approximation as a method used for determining approximate solutions to wall-crossing setups, with particular focus on two settings of interest for us, namely Hitchin systems and vacuum transitions.

9 Quantum Field Theory from Category Theory

We now turn to explaining how QFT can be mathematically described in terms of Category Theory.

To begin with, we start off by listing some of the ingredients that are most relevant to our analysis⁴⁹, briefly outlined in the table below.

<u>Particle Physics</u>	<u>Category Theory</u>
Gauge Theory	Algebras
Operators	Homologies
Quivers	Quivers/Algebraic varieties
Hilbert Series	Hilbert Series
SUSY	SUSY

This shows some quantities/entities encountered in the realm of Particle Physics and their Category Theory counterpart. We have highlighted two of them in red. The reason for this will be more clear in Part V, but the familiar reader will certainly notice that some motivation was already hinted at in section 3. For the moment, we simply recall what was mentioned in the Introduction, namely that the Hilbert series and ring homologies are key complementary ingredients towards understanding a given QFT. Among the two, though, the latter is the one that is more directly related to understanding the underlying categorical structure of the theory in question, and therefore constitutes the ultimate objective one wishes to obtain. On the other hand, the Hilbert series, is obtained from ring homologies, but hides some key categorical features that are only accessible via the homological perspective.⁵⁰

This intermediate section is mostly an overview of [102, 102, 103], and is needed in order to understand the analysis carried out in the concluding part of our work, J. This section is structured as follows:

1. At first, we review mathematical gauge theory and the definition of moduli spaces therein, [102].

⁴⁹As we shall see, more richness will be added in due course in our treatment. The aim of the table is simply that of helping the reader in gaining intuition for the setup and the motivations behind the procedure adopted.

⁵⁰Indeed, it is known that different ring homologies could lead to the same Hilbert series, [616].

2. Then, we recall the notion of relative field theories, [103], which can also be adapted to gauge theories.
3. In conclusion, we explain how this fits with the higher-categorical prescription of Freed, Moore and Teleman, [102], and how gauging can be implemented in terms of gaugeable homomorphisms in the Symmetry Topological Field Theory categorising the symmetries of a given QFT, [607].

For reasons that will become clear by the end of this section, the crucial part of the higher-categorical perspective can be summarised as follows:

- Keeping track of the identity.
- Embedding the gauged algebra.

Most importantly, they both lie within the notion of *abelianisation*. In section J we will see explicitly how they arise within the context of supersymmetric quiver varieties, [600, 601].

9.1 The Freed-Moore-Teleman construction

9.1.1 Gauging in relative QFTs

In the formulation of [103], a *relative* field theory, \tilde{F} , requires additional topological data in order to be fully specified. Such data is encoded in a pair (σ, ρ) , referred to as *quiche*. σ is the *symmetry topological field theory* (SymTFT), whereas ρ geometrises the choice of boundary conditions for the fields defining the relative theory \tilde{F} . The overall system, depicted in figure 29, gives rise to an *absolute* QFT, F_ρ .




Figure 29. The Freed-Moore-Teleman setup, with \tilde{F} denoting a relative QFT. Specifying the topological data (σ, ρ) , the resulting theory, F_ρ is absolute.

In mathematical terms, the description outlined above can be formulated in terms of bordism in the following way. Fixing $N \in \mathbb{Z}^{\geq 0}$, then a quiche is a pair (σ, ρ) in which

$\sigma : \text{Bord}_{N+1}(F) \rightarrow \mathcal{C}$ is an $N + 1$ -dimensional TFT and ρ is a right topological σ -module. The quiche is N -dimensional, hence it shares the same dimensionality as the theory on which it acts. Let F be an N -dimensional field theory. A (σ, ρ) -module structure on F is a pair (\tilde{F}, θ) , in which \tilde{F} is a left σ -module and θ is an isomorphism

$$\theta : \rho \otimes_{\sigma} \tilde{F} \xrightarrow{\cong} F_{\rho}, \quad (9.1)$$

of absolute N -dimensional theories, with (9.1) defining the dimensional reduction leading to the absolute theory. σ needs only be a *once-categorified* N -dimensional theory, whereas ρ and \tilde{F} are relative theories.

In theories admitting a higher-categorical structure as (4.19), topological defects in different superselection sectors have different dimensionality. Consequently, the fusion rules are categorical, in the sense that they usually don't follow group-like composition laws. Because of this these defects are also called categorical or non-invertible.

Prior to delving into the specific case of interest to us, namely relative QFTs, we first wish to provide some further explanation for what gauging a categorical structure actually means. In doing so, we refer to the work of many experts in the field, and, in particular [607]. As explained in such reference, for any fusion n-category \mathfrak{G} , any fiber functor

$$\mathcal{F} : \mathfrak{G} \rightarrow \text{nVec}, \quad (9.2)$$

selects nVec as the image of a gaugeable algebra living in \mathfrak{G} , corresponding to a projection on the identity. The gauging process, can therefore be defined as a map

$$\mu : \mathfrak{G} \rightarrow \mathcal{A}_C, \quad (9.3)$$

with \mathcal{A}_C the algebra of invertible topological operators in \mathfrak{G} . In terms of the example reproduced in section 7.3.1, \mathcal{A}_C refers to the algebra that dresses the symmetry operators once having implemented the gauging to ensure gauge invariance. Hence, μ really corresponds to the projection of the identity to a subalgebra of the total algebra, \mathcal{A} , featuring in the pre-gauged theory, with $\mathcal{A}_C \subset \mathcal{A}$. For later purposes, we highlight the relation in between this description with that introduced in section 3 in relation to Moore-Tachikawa varieties. As also defined there, the crucial role from the category theory point of view really consists in defining the cotangent bundle as a triple of elements, two of which are left- and right- acting moment maps, mutually constrained by the main axiom outlined by Moore and Tachikawa. As explained in section 5.4.1, this follows from full-dualisability of the 2-categories involved in defining the 2D TFT associated to a given class \mathcal{S} theory⁵¹. Given (9.3), the norm element

$$N \stackrel{def}{=} \bigoplus_{g \in \mathfrak{G}} \mu(g). \quad (9.4)$$

carries the structure of an n-categorical idempotent, also known as *gaugeable algebra*, depicted in black in figure 30. The requirement for (9.4) to be a higher-idempotent is needed

⁵¹As explained in Part V, this follows from the fact that the category of contractible cycles associated to a given algebraic variety admits a Kirwan map ensuring abelianisation in terms of equivariant cohomology.

to ensure the flooding doesn't depend on the specific features of the network being adopted to perform the gauging. The algebra of topological operators that are left are denoted by $\mathcal{A} // \mathfrak{G}$. The equivalence of the second and third picture from the left in figure 30 follows from N being a higher-gaugeable algebra. As we shall see, this pattern emerges when gauging the 5D SymTFT of class \mathcal{S} theories, where the objects of the 2-category in question will be Wilson surfaces charged under the 1-form symmetry being gauged.




Figure 30. Gauging corresponds to condensing an algebra in a TFT. Idempotency ensures the resulting theory can be effectively thought of as featuring a unique defect, as shown on the RHS.




Figure 31. Condensing two different subalgebras, $\mathcal{A}_1, \mathcal{A}_2 \subset \mathcal{A}$, the resulting theory corresponds to one with a changed phase with a gauged defect resulting from a relative gaugeable algebra, \mathcal{A}_{12} ending in the bulk. The defect at the endpoint is nontrivial, and can therefore be thought of as a $\text{Hom}(\mathbf{1}_c, \mathcal{A}_{12})$.

Unlike 30, 31 does not admit a straightforward expression as (9.2).

9.1.2 Absolute 4D theories from 6D $\mathcal{N} = (2, 0)$ SCFTs

The main features of 6D $\mathcal{N}=(2,0)$ SCFTs are that:

- They admit no Lagrangian formulation, [103, 110].
- The theory depends on the Lie algebra rather than the Lie group.
- Ordinary⁵² 5D and 4D QFTs can be obtained by dimensional reduction on a Riemann surface, $\Sigma_{g,n}$.

⁵²By this we mean that the lower-dimensional theories depend on the specific choice of the Lie group.




Figure 32. Adaptation of Freed-Moore-Teleman to the case involving twisted gauged defects. The figure on the far right corresponds to the case of interest to us, namely a configuration involving two different absolute 4D gauge theories separated by a defect. As we shall see, such defect is intrinsically non-invertible, corresponding to the presence of a relative uncondensed subalgebra, dressing \mathcal{N}_3 in a nontrivial way. In higher-categorical terms, it corresponds to a fusion tensor category implementing the morphisms between the operators charged under the gauged symmetry.

- It is a *relative* w.r.t. an *extended* QFT in 1-dimension higher.

Examples of relative theories are models whose fields form a fibration, [103],

$$\mathcal{F} \longrightarrow \mathcal{M} \xrightarrow{\pi} \mathcal{B} \quad (9.5)$$

where \mathcal{B} and \mathcal{F} denote the base and the fiber, respectively. Relative fields are the fibers of π . The path integral over the base reduces a finite sum. For the theory to be absolute, the same should hold for \mathcal{F} as well.

When compactifying the 6D theory on $\Sigma_{g,n}$ without punctures:

1. If $g \equiv 1$ $\Sigma_g \equiv T^2$, the resulting class \mathcal{S} theories are $\mathcal{N}=4$ SYM with gauge algebra $\mathfrak{su}(N)$.
2. For $g > 1$, the resulting theories are models with an S -duality frame, interpreted as conformal gaugings of $2g - 2$ trinion T_N theories coupled to $3g - 3$ gauge groups with $\mathfrak{su}(N)$ gauge algebras.

9.1.3 From relative to absolute

The procedure described in section 9.1.1 cannot be applied to the case of maximally supersymmetric 6D theories. The reason being that, while a bulk SymTFT can be defined, thanks to the Milnor-Moore theorem, [311], specifying any topological boundary conditions would inevitably imply supersymmetry breaking⁵³, [103]. Nevertheless, such 6D theories provide the natural realm from which one can realise lower-dimensional absolute theories,

⁵³To the best of our knowledge, the only exception to this is dealt with in [620].

as well as constituting an extremely rich setting where to explore and put to practice mathematical techniques provided by representation theory, [111].

In this subsection we briefly overview the work of [367], and propose a criterion for distinguishing intrinsic from non-intrinsic non-invertible defects.

Assuming the 6D $\mathcal{N} = (2, 0)$ SCFT is defined on $Y = X \times \Sigma_g$, with X a compact 4D spacetime, torsion-free, and with trivial 1st cohomology, $H^1(X, \mathbb{Z}_N)$, the 6D self-dual fluxes belong to⁵⁴

$$H^3(Y, \mathbb{Z}_N) \simeq H^1(\Sigma_g, \mathbb{Z}_N) \otimes H^2(X, \mathbb{Z}_N). \quad (9.6)$$

As far as the 6D theory is concerned, this is the only information at our disposal, given that the theory is non-Lagrangian, [103, 110]. However, for the 4D theory to be fully defined, additional data is needed. In particular, one needs to identify a *maximal isotropic sublattice* L of $H^1(\Sigma_g, \mathbb{Z}_N)$ w.r.t. the canonical pairing induced by the intersection pairing on Σ_g , thereby leading to

$$\mathcal{L} \simeq L \otimes H^2(X, \mathbb{Z}_N) \quad (9.7)$$

as the maximal isotropic lattice of $H^3(Y, \mathbb{Z}_N)$. (9.7) enables to distinguish between different global forms of class \mathcal{S} theories. Having fixed \mathcal{L} , the fluxes in its complement

$$\mathcal{L}^\perp \stackrel{\text{def.}}{=} H_3(X_6, \mathbb{Z}_N)/\mathcal{L} \quad (9.8)$$

parametrise the possible partition functions for the 4D theory with inequivalent 1-form symmetry backgrounds along X .

The 6D $\mathcal{N}=(2,0)$ theory, assigns to the 6D manifold Y a partition vector rather than a partition function, with the former belonging to the Hilbert space constituting a representation of a Heisenberg algebra of non-commuting discrete 3-form fluxes in \mathbb{Z}_N .

For fixed \mathcal{L} , the partition vector takes the form

$$|\mathcal{Z}(Y)\rangle \equiv \sum_{v \in \mathcal{L}^\perp} \mathcal{Z}_v(Y) |\mathcal{L}; v\rangle, \quad (9.9)$$

with $\mathcal{Z}_v(Y)$ denoting the 6D conformal blocks, which also encode the *global structures* of the 4D theory. Under a change of basis, (9.9) turns into

$$|\mathcal{Z}(Y)\rangle \equiv \sum_{v' \in \mathcal{L}'^\perp} \mathcal{Z}_v(Y) \sum_{v \in \mathcal{L}^\perp} R_v^{v'} |\mathcal{L}; v\rangle, \quad (9.10)$$

leading to the following relation between conformal blocks

$$\mathcal{Z}_v(Y) \equiv \sum_{v' \in \mathcal{L}'^\perp} R_v^{v'} \mathcal{Z}_{v'}(Y). \quad (9.11)$$

⁵⁴ \mathbb{Z}_N denotes the *defect group* of the 6D theory.

The 7D and 5D SymTFTs

The Milnor-Moore Theorem states that, the universal enveloping algebra of a Lie algebra is a Hopf algebra whose primitive elements are the elements of the original Lie algebra. A Hopf algebra on an associative algebra turns the category of modules into a monoidal category equipped with a fiber functor. This is a statement of Tannaka duality.

Putting this together for the case of 6D $\mathcal{N} = (2,0)$ SCFTs, the bulk SymTFT is a pointed braided tensor category where the braiding is in bijection with 3-cocycles.

Following the arguments outlined in section 4.1.2, one can take the 7D TQFT in the bulk of the relative 6D $\mathcal{N} = (2,0)$ theory, to be a CS theory with action

$$S_{7D} = \frac{N}{4\pi} \int_{W_7} dc \wedge c \quad , \quad c \in H^3(W_7, U(1)), \quad (9.12)$$

and Wilson surfaces

$$\Phi_q(\mathcal{M}_3) \stackrel{def.}{=} e^{iq \int_{\mathcal{M}_3} c} \quad , \quad q \in \mathbb{Z}_N \quad , \quad \mathcal{M}_3 \in H^3(W_7, U(1)). \quad (9.13)$$

Taking two such Wilson surfaces $\Phi_q(\mathcal{M}_3), \Phi_{q'}(\mathcal{M}'_3)$, with $\mathcal{M}_3, \mathcal{M}'_3$ forming a Hopf link in W_7 , and inserting them in the path integral, amounts to changing the action (9.12) by adding the holonomy terms associated to the operator insertions

$$\begin{aligned} S_{7D} &= \frac{N}{4\pi} \int_{W_7} dc \wedge c + q \int_{\mathcal{M}_3} c + q' \int_{\mathcal{M}'_3} c \\ &= \frac{N}{4\pi} \int_{W_7} dc \wedge c + \int_{W_7} \left(q\omega_{\mathcal{M}_3} + q'\omega_{\mathcal{M}'_3} \right) \wedge c, \end{aligned} \quad (9.14)$$

with $\omega_{\mathcal{M}_3}, \omega_{\mathcal{M}'_3}$ denoting the Poincaré duals of $\mathcal{M}_3, \mathcal{M}'_3$, respectively. Integrating-out c , and imposing

$$dc \equiv -\frac{2\pi}{N} \left(q\omega_{\mathcal{M}_3} + q'\omega_{\mathcal{M}'_3} \right), \quad (9.15)$$

defining V_4 such that

$$\partial V_4 \equiv \left(q\omega_{\mathcal{M}_3} + q'\omega_{\mathcal{M}'_3} \right), \quad (9.16)$$

implies

$$c \equiv -\frac{2\pi}{N} \text{PD}(V_4). \quad (9.17)$$

From this, (9.14) reduces to

$$S_{7D} = \frac{2\pi}{N} qq' \text{link}(\mathcal{M}_3, \mathcal{M}'_3), \quad (9.18)$$

where the linking is the Hopf link occurring in W_7 . It features in (9.18) due to the fact that V_4 is a Seifert surface for the combination $q\mathcal{M}_3 + q'\mathcal{M}'_3$, meaning that, every time \mathcal{M}_3 pierces V_4 , it links \mathcal{M}'_3 once. It thereby follows that, the Hopf link of Wilson surfaces

supported on \mathcal{M}_3 and \mathcal{M}'_3 in W_7 can be thought of as an operator insertion in the 7D path integral,

$$\langle \Phi_q(\mathcal{M}_3)\Phi_{q'}(\mathcal{M}'_3)\dots \rangle \equiv e^{\frac{2\pi i}{N}qq'\text{link}(\mathcal{M}_3,\mathcal{M}'_3)} \langle \dots \rangle, \quad (9.19)$$

where the phase on the RHS being the analog of the S -matrix element S_{ab} between anyons of charges a, b in $U(1)_k$ CS theory in 3D, namely $S_{ab} \equiv e^{2\pi iab/k}$.

For a given $X_6 \equiv \Sigma_{g,n} \times X_4$, there is a particular class of 7-manifolds W_7 obtained by taking $W_7 \equiv V_{g,n} \times X_4$, with $V_{g,n}$ a 3-manifold with $\partial V_{g,n} \equiv \Sigma_{g,n}$.

For any Riemann surface $\Sigma_{g,0}$, there are many inequivalent 3-manifolds with $\Sigma_{g,0}$ as its boundary. One such example are handlebodies. To construct a handlebody, choose a set g of *meridians*, $\{\mu_i\}, i \equiv 1, \dots, g$, i.e. a set of generators of $H_1(\Sigma_{g,0}, \mathbb{Z})$ which become trivial as elements of $H_1(V_{g,0}, \mathbb{Z})$. The remaining g generators of $H_1(\Sigma_{g,0}, \mathbb{Z})$ lift to generators of $H_1(V_{g,0}, \mathbb{Z})$, and are referred to as *longitudes*, $\{\lambda_i\}, i \equiv 1, \dots, g$. Not every choice of g generators gives rise to a legitimate set of meridians (i.e. vanishing cycles). For a specific choice of meridians to be legitimate, it must correspond to a maximal isotropic sublattice

$$L \in H_1(\Sigma_{g,0}, \mathbb{Z}). \quad (9.20)$$

The handlebody specified by the choice of meridians L will be denoted by $V_{g,0}^L$. Given $V_{g,0}^L$, there are multiple choices of longitudes, differing by shifts in meridians

$$\lambda'_i \equiv \lambda_i + \sum_{j=1}^3 k_{ij} \mu_j \quad , \quad k_{ij} \in \mathbb{Z}, \quad (9.21)$$

satisfying the following constraint

$$\langle \mu_i, \lambda_j \rangle \equiv - \langle \lambda_j, \mu_i \rangle \equiv \delta_{ij}. \quad (9.22)$$

7D CS theory on a handlebody can be formulated by starting with a definition of the fields which reads

$$b_i \stackrel{def.}{=} \oint_{\mu_i} c \quad , \quad \hat{b}_i \stackrel{def.}{=} \oint_{\lambda_i} c \quad , \quad i = 1, \dots, g. \quad (9.23)$$

The meridians are naively contractible by definition. However, this does not necessarily imply the vanishing of b_i , since there might be Wilson surfaces supported on λ_i and μ_i that are nontrivially linking, effectively making μ_i non-contractible.

The Wilson lines wrapping meridians and longitudes can be expressed as follows

$$\Phi_i(\mathcal{M}_2) \stackrel{def.}{=} \Phi_i(\mathcal{M}_2 \times \mu_i) \equiv e^{\frac{2\pi i}{N} \oint_{\mathcal{M}_2} b_i} \quad , \quad \hat{\Phi}_i(\mathcal{M}_2) \stackrel{def.}{=} \hat{\Phi}_i(\mathcal{M}_2 \times \lambda_i) \equiv e^{\frac{2\pi i}{N} \oint_{\mathcal{M}_2} \hat{b}_i} \quad , \quad (9.24)$$

which might be nontrivially linking in the 5D theory resulting from dimensional reduction of the 7D TFT on $\Sigma_{g,0}$. Compactifying the theory on $\Sigma_{g,0}$ in presence of a series of Wilson 3-surfaces on $\lambda_j \times \mathcal{M}'_{2,j}$, the linking equation becomes an operator equation in 4D

$$\Phi_i(\mathcal{M}_2) \equiv e^{\frac{2\pi i}{N} \sum_{j=1}^g \langle \mu_i, \lambda_i \rangle \int_{\mathcal{M}_2} B_j}, \quad (9.25)$$

where $B_j \in H^2(X_4, \mathbb{Z}_N)$ denotes the Poincaré dual of $\mathcal{M}'_{2,j}$. Hence, in any 4D calculation involving Φ_i , the latter can be moved to the right of $\hat{\Phi}_j$, and subsequently be shrunk to a point by introducing the phase (9.25). Hence, the insertion of longitudinal Wilson lines allows us to make $b_i \neq 0$, while still keeping it constant, thereby defining a background field.

At this point, the Sym TFT can be obtained by splitting the handlebody in the following way:

$$V_{g,0} \stackrel{def.}{=} \begin{cases} V_{g,0}^{L,in} & \text{for } y \geq y_* \\ V_{g,0}^{ext} \stackrel{def.}{=} \Sigma_{g,0} \times [0, y_*] & \text{for } 0 \leq y \leq y_* \end{cases}. \quad (9.26)$$

Importantly, it features two boundaries (cf. figure 33):

- One, at $y = 0$, is where the 6D $\mathcal{N}=(2,0)$ theory lives.
- The other, at $y = y_*$, is where the longitudinal Wilson surfaces fixing the background fields are placed.

9.2 Fiber functors from non-invertible symmetries

The main motivation of the present section is that of studying the defects interpolating different 4D class \mathcal{S} theories descending from dimensional reduction of 6D $\mathcal{N}=(2,0)$ SCFTs. In particular, explaining how such defects emerge under gauging of multiple subalgebras, we propose the quantum dimension of the relative gaugeable subalgebra as the quantity needed to distinguish whether such interpolating defect is intrinsically or non-intrinsically non-invertible, thereby proposing the 4D counterpart of arguments proposed in [333] for the 2D case.

The present section is therefore structured into 4 main parts:

- At first, we describe the procedure, [374, 375, 607], and its adaptation to the Freed-Teleman, [103], and Freed-Moore-Teleman construction, [102], of relative field theories and topological symmetries of QFTs, respectively, with particular emphasis on the case of multiple simultaneous gauging of different subalgebras.
- We then briefly overview the procedure adopted by [367] for realising absolute 4D SCFTs from $\mathcal{N} = (2,0)$ SCFTs, explicitly rephrasing it in terms of the multiple gauging procedure described in section 9.1.1.
- When describing such interpolating defects in terms of gauging of homomorphisms, they naturally admit an interpretation in terms of fusion tensor categories interpolating between different 2-categories, to a total quantum dimension can be assigned.




Figure 33. In presence of the splitting (9.26), the state $\langle L |$ separates the anomaly TFT from the Sym TFT. The picture above is valid for $\forall g$, but has been drawn with $g = 1$ for simplicity.

Section 14 closes with the relation between the issue in defining a fiber functor in presence of non-invertible defects interpolating between different absolute theories.

- Making use of the gauging procedure explained in section 9.1.1, we propose a criterion for distinguishing between the two types of non-invertible defects in terms of the total quantum dimension of the relative gauged algebra implementing the gauging.

9.2.1 Intrinsic non-invertible symmetries and fiber functors

On the handlebody, one can perform two different operations:

1. A *modular transformation* of $\Sigma_{g,0}$, acting on the entire handlebody and generically changing the period matrix of $\Sigma_{g,0}$ at each cross section. The theory in the bulk of the handlebody is topological, implying only $\Sigma_{g,0}$ is affected by such transformations, which is where the 6D $\mathcal{N} = (2,0)$ theory lives.
2. Excising an *inner* handlebody, in the setup where the splitting (9.26) takes place. The operation of interest consists in gluing together the 2 parts $V_{g,0}^{L,in}$ and $V_{g,0}^{ext}$ with a nontrivial element of the modular group $Sp(2g, \mathbb{Z}_N)$, corresponding to a surgery operation on the 3-manifolds.

Given that the latter takes place in the interior of the handlebody, it doesn't affect $\Sigma_{g,0}$ where the 6D theory lives, hence, the period matrix remains unchanged in the region the theory is sensitive to. However, this surgery operation changes the global form of L , since it changes which are the contractible cycles in the handlebody. Hence, surgery is implemented by an element F of the modular group which only affects the inner boundary of the handlebody, thereby changing the global form. This operation may in turn be combined with a modular transformation acting on the entire handlebody, reverting the interior of the handlebody back to its original form, but now changing the period matrix of the Riemann surface.

Choosing a period matrix that remains invariant under the action of F , the geometry with the combined surgery plus modular transformation has the same boundaries as the original geometry. Importantly, though, the full geometry is no longer the same, due to the internal twist.

Under dimensional reduction on $\Sigma_{g,0}$, (9.14) becomes

$$S_{5D} \equiv \frac{2\pi}{N} \sum_{i=1}^g \int_{X_4 \times [0, y_*]} b_i \cup \delta \hat{b}_i. \quad (9.27)$$

The fact that b_i, \hat{b}_i are dynamical, implies λ_i, μ_i are both nontrivial in $V_{g,0}^{ext}$. This follows from the fact that L specifies which directions of the Riemann surface become trivial in the handlebody, namely the meridians, and which of the dynamical fields of the BF -theory become background. To each L corresponds a different choice of Dirichlet boundary conditions for different sets of fields in the BF -theory. As explained in [367], the 7D CS theory on a Riemann surface can only capture the full Sym TFT iff all the non-invertible defects are non-intrinsic.

The 5D Wilson lines are defined from the 7D Wilson surfaces as follows

$$\Phi_{\vec{n}}(\mathcal{M}_2) \stackrel{def.}{=} \Phi(\mathcal{M}_2 \times \gamma_{\vec{n}}), \quad (9.28)$$

with 1-cycles

$$\gamma_{\vec{n}} \stackrel{def.}{=} \sum_{j=1}^g e_j \lambda_j + \sum_{j=1}^g m_j \mu_j, \quad \vec{n} \stackrel{def.}{=} (e_1, \dots, e_g; m_1, \dots, m_g). \quad (9.29)$$

Explicitly, (9.28) can be expressed as follows

$$\Phi_{\vec{n}}(\mathcal{M}_2) \equiv e^{\frac{2\pi i}{N} \frac{1}{2} e \cdot m(\mathcal{M}_2, \mathcal{M}_2)} \prod_{i=1}^g \Phi^{e_i}(\mathcal{M}_2 \times \gamma_{\vec{n}}) \prod_{i=1}^g \Phi^{m_i}(\mathcal{M}_2 \times \gamma_{\vec{n}}), \quad (9.30)$$

with each electric and magnetic component, in turn descending from a 3D Wilson surface in 7D associated to a cycle $\gamma_{\vec{n}}$.

Surgery defects in 7D correspond to loci across which F acts on $\gamma_{\vec{n}}$ as follows

$$F(\gamma_{\vec{n}}) \equiv \gamma_{F\vec{n}}. \quad (9.31)$$

In 5D, they reduce to gaugeable defects implementing morphisms in a higher-categorical structure

$$\mathcal{C}_F(X_4) : \Phi_{\vec{n}}(\mathcal{M}_2) \mapsto \Phi_{F\vec{n}}(\mathcal{M}_2) \quad , \quad \mathcal{M}_2 \subset X_4. \quad (9.32)$$




Figure 34. The picture on the LHS shows the identification of a morphism between objects as a defect in a higher-categorical structure. On the RHS, the interface turns into a boundary defect once having performed the folding trick.

As suggested by (9.32), F acts on the labellings of the objects living in the theory. These objects are dyonic, with nontrivial linking. An explicit expression for the morphism (9.32) as a defect follows by performing the folding trick on the surgery point in the 5D configuration depicted in figure 34

$$\mathcal{C}(X_4) = \frac{|H^0(X_4, \mathbb{Z}_p)|}{|H^1(X_4, \mathbb{Z}_p)|} \sum_{\mathcal{M}_2 \in H_2(X_4, \mathbb{Z}_p^{2g})} \exp\left(\frac{2\pi i}{p} \langle F\mathcal{M}_2, \mathcal{M}_2 \rangle\right) \Phi((\mathbb{1} - F)\mathcal{M}_2), \quad (9.33)$$

implying that, $\forall F \in Sp(2g, \mathbb{Z}_p)$, there is a codimension-1 defect implementing the symmetry associated to F in the bulk 5D TQFT. Such defect is invertible, and therefore its fusion rules simply read

$$\boxed{\mathcal{C}(X_4) \times \bar{\mathcal{C}}(X_4) \equiv \mathcal{A} \equiv \mathbf{1}}. \quad (9.34)$$

where \mathcal{A} corresponds to the condensed algebra implementing the gauging. Notice that the defect (9.33) is equivalent to the one arising from gauging by algebraic gauging in a topological order, as explained in section 9.1.1 and depicted in figure 30. The crucial meaning of equation (9.34) is the fact that, gauging is effectively acting as a redefinition of the identity from the mother theory to the gauged one. In particular, the last equality claims that the condensed algebra in the mother theory is projected to the identity of the gauged theory. Pictorially, this corresponds to the configuration on the LHS of 35.




Figure 35. Gauging the defects living in the SymTFT upon implementing an outer automorphism, changes the boundary conditions of the resulting absolute theory. In presence of a unique gauging, the resulting configuration is depicted on the LHS. On the other hand, upon condensing two different subalgebras of the original bulk theory, leads to a bulk gauging defect effectively ending in the bulk, as shown in the figure in the middle. This is equivalent to realising a configuration comprising two different boundary conditions separated by a topological defect (as shown on the RHS).

Gauged defects associated to F , can in turn give rise to $|F|$ -ality defects in the boundary 4D gauge theory. Such defects can be constructed from (9.33) by admitting Dirichlet boundary conditions for the gauged defects, allowing them to terminate in the bulk, [333, 341, 342]. Practically, this corresponds to a 3D-manifold attached to a 4D manifold, which, combined together, give rise to a *twist defect*, $T_F(\mathcal{M}_3, \mathcal{M}_4)$, whose expression is obtained from (9.33) once rewritten in terms of relative cohomology. From the perspective of the gauging procedure, this amounts to considering simultaneous gauging of two different subalgebras within the gaugeable subalgebra of the mother theory. In terms of its SymTFT characterisation, this would correspond to the gaugeable defect effectively ending in the bulk SymTFT as shown in the middle of figure 35. Equivalently, this is equivalent to describing a relative field theory with two boundary conditions interpolated by a topological defect.

The gauged defect can be reexpressed in terms of the quantum dimensions of the underlying anyon description by noticing that the action of F is equivalent to that of the outer automorphism keeping track of the anyonic symmetry being gauged⁵⁵. This follows from the fact that the elements of the sum appearing in the gauged defect, (9.33), can be broken down into two main parts, namely

$$\exp\left(\frac{2\pi i}{p} \langle F\mathcal{M}_2, \mathcal{M}_2 \rangle\right) \Phi((1-F)\mathcal{M}_2) \equiv S_{\vec{a} \ F\vec{a}} \mathcal{G}_{F\vec{a}}, \quad (9.35)$$

with $S_{\vec{a}, F\vec{a}}$ and $\mathcal{G}_{F\vec{a}}$ denoting the S -matrix entries for the associated anyonic charges, and the exceptional fibration operator, respectively. Extending these arguments, the sum over the 2-manifolds \mathcal{M}_2 can be replaced by a sum over the anyonic charges compatible with the symmetries of the given theory. This way, (9.33) can be rewritten as follows

⁵⁵As explained in section 4.1.2 and 9.1.1.




Figure 36. The gauged defect $\mathcal{C}_F(X_4)$ implements the outer automorphism in the 5D TQFT. Applying twisted BCs, the gauged defect can effectively terminate in the 5D bulk, and is therefore topological. At this point, the width of the 5D theory can be taken to vanish, resulting in the configuration on the RHS, with the two phases of the 4D theory separated by a topological non-invertible defect, $\mathcal{N}_F(\mathcal{M}_3)$.

$$T_F(\mathcal{M}_3, \mathcal{M}_4) \stackrel{def.}{=} \frac{|H^0(X_4, \mathbb{Z}_p)|^{2g}}{|H^1(X_4, \mathbb{Z}_p)|^{2g}} \sum_{\vec{a}} S_{\vec{a}, F\vec{a}} \mathcal{G}_{F\vec{a}}, \quad (9.36)$$

where $\vec{a} \stackrel{def.}{=}} ([a], \rho(a))$, with $[a], \rho(a)$ denoting the conjugacy class and the representation of the corresponding anyon charges, respectively. Given that the defect is now topological, it can be moved to the 4D boundary, defining an interface for the 4D gauge theory

$$\mathcal{N}(\mathcal{M}_3) \stackrel{def.}{=} \lim_{y \rightarrow 0} T_F(\mathcal{M}_3, \mathcal{M}_4), \quad (9.37)$$

obeying non-invertible fusion rules. From the 4D perspective of the absolute theory on the RHS of figure 36, $\mathcal{N}(\mathcal{M}_3)$ is *intrinsically*-non-invertible, namely, it cannot be rendered invertible by acting with $SL(2, \mathbb{Z})$ -transformations. This is equivalent to stating that the action of F cannot be reversed simply by acting with a modular transformation. Non-invertibility implies its fusion rules now read

$$\boxed{\mathcal{N}(\mathcal{M}_3) \times \bar{\mathcal{N}}(\mathcal{M}_3) = \mathcal{A}_{X_4}}, \quad (9.38)$$

with \mathcal{A}_{X_4} denoting the relative gaugeable algebra in the absolute theory.

Crucially, the RHS of equation (9.38) is not simply the identity of a 4D absolute theory, corresponding to the fact that there is no direct counterpart of equation (9.2) upon performing a double gauging involving two distinct subalgebras of the gaugeable algebra characterising the mother SymTFT. Given that the latter is basically equivalent to the existence of a well-defined partition function for the resulting absolute theory, we claim that, if might be able to define a counterpart of the functor (9.2) for the case depicted at the centre and RHS of figure 35, by composing the following chain of functors




Figure 37. A qualitative diagram depicting the counterpart of the fiber functor’s domain for the case of figure 35. The black arrows indicate the choice of different gaugeable subalgebras, leading to different 4D absolute theories, labelled by 2-categories \mathcal{C}, \mathcal{D} .

where \mathcal{A}_{4D} is the gaugeable algebra leading to a single 4D absolute theory, whereas \mathcal{A}_1 and \mathcal{A}_2 correspond to different subalgebras of \mathcal{A}_{4D} leading to different 2-categories of underlying charges associated to the gauged symmetry characterising the two distinct 4D absolute theories. This is current work in progress by the same author, and we plan to report about it in a forthcoming work. The main message we want to convey with this is that the fiber functor that can be defined for the configuration at the centre and RHS of figure 35 leads to the partition function of a 3D theory.

9.2.2 Quantum dimension from relative gaugeable algebra

In the concluding part of this section, we make use of the tools outlined in the previous parts of the present treatment, leading to the proposal of a probe quantity enabling to distinguish whether the defect interpolating in between class \mathcal{S} theories is intrinsically or non-intrinsically non-invertible. The starting point will be the setting of section 9.1.1, where we revised the gauging procedure, implemented by flooding the topological order corresponding to the 2-category of Wilson surfaces living in the bulk SymTFT with a tensor network of an idempotent element, namely the norm element (9.4).

Practically, the two phases (prior and after gauging) can be thought of as arising from a phase transition in between two MTCs, \mathcal{B}_{ung} and \mathcal{B}_g , corresponding to the ungauged and the gauged phases, respectively, [374]. Imposing natural physical requirements, it is possible to derive a relation between the anyons in the \mathcal{B}_g phase and those in the \mathcal{B}_{ung} phase. The vacuum, or tensor unit \mathcal{A} in \mathcal{B}_g is necessarily a connected commutative separable algebra in \mathcal{B}_{ung} , and the category \mathcal{B}_g is equivalent to the category of local \mathcal{A} -modules as MTCs. This gauging produces a gapped domain wall (DW) with wall excitations given by the category of \mathcal{A} -modules in \mathcal{B}_{ung} . The domain wall separating them, is, in turn a category, $\mathcal{A} \equiv \mathcal{C}(X_4)$, as shown in figure 38.

For any pair of elements, M, N ,

$$\text{Hom}_{\mathcal{B}_g}(M, N) \hookrightarrow \text{Hom}_{\mathcal{B}_{ung}}(M, N). \quad (9.39)$$

Assuming vacuum degeneracy being trivial,

$$\text{Hom}_{\mathcal{A}}(\mathbf{1}_{\mathcal{A}}, \mathbf{1}_{\mathcal{A}}) \simeq \mathbb{C}, \quad (9.40)$$

\mathcal{A} must be a UFC with a unique spherical structure. All objects in \mathcal{A} should come from objects in \mathcal{B}_{ung} , and should satisfy the following properties:




Figure 38. This picture shows the categories of anyons associated to the ungauged and gauged theories, denoted by \mathcal{B}_{ung} and \mathcal{B}_g , respectively. Gauging can be visualised as the domain wall effectively separating two different topological phases.

1. \mathcal{A} is a subcategory of \mathcal{B}_{ung}

2.

$$\mathrm{Hom}_{\mathcal{A}}(M, N) \hookrightarrow \mathrm{Hom}_{\mathcal{B}_{ung}}(M, N) \quad (9.41)$$

3. $\mathbf{1}_{ung}$ should condense into B , such that

$$\iota_B : \mathbf{1}_{\mathcal{B}_{ung}} \rightarrow B \in \mathcal{B}_{ung} \quad (9.42)$$

$\mathbf{1}_{\mathcal{B}_g} \equiv \mathcal{A}$ should fuse into the vacuum on the wall when moving \mathcal{A} close to the wall, such that, in \mathcal{B}_{ung} , the corresponding map reads

$$\iota_A^B : A \rightarrow B \quad (9.43)$$

What we have just said can be succinctly re-expressed as follows

$$\boxed{\mathbf{1}_{\mathcal{B}_g} \equiv \mathcal{A} \subset \mathcal{B}_{ung}}. \quad (9.44)$$

After what we have just said, the key point to focus on is the fact that the gauging procedure, the identity operator changes, in the sense that, from being a simple element in the category \mathcal{B}_{ung} , it is nontrivial in the category of the gauged theory, \mathcal{B}_g , since $\mathbf{1}_{\mathcal{B}_g} \equiv \mathcal{A}$.

The analogy we have drawn so far with the example shown in figure (9.44) taken from [374], corresponds to the case of a single gauging of the 5D SymTFT, leading to the LHS of our figure 35. For the case involving a double gauging, instead, such prescription needs to be suitably adapted. The motivation still follows the arguments presented in [374]. We now argue why this is indeed the case specifying to the example described in section 9.1.2.

Recall that $\mathcal{C}(X_4)$ is the operator implementing the outer automorphism on the category of defects defining the 5D SymTFT, and corresponds to the idempotent object that is constituted of the algebra elements that are flooding the SymTFT in order to implement the gauging. Denoting by $\mathcal{C}_g, \mathcal{C}_{ung}$ the 2-category of defects characterising the 4D absolute theories, if the two phases are related by a single bulk gauging, it follows that

$$\dim_{\mathcal{C}_g} \mathbf{1}_{\mathcal{C}_g} \equiv \frac{\dim_{\mathcal{C}_{ung}} \mathcal{A}_{4D}}{\dim_{\mathcal{C}_{ung}} \mathcal{A}_{4D}} \equiv 1. \quad (9.45)$$

But this is only from the point of view of the gauged theory, \mathcal{C}_g . On the other hand, \mathcal{A}_{4D} is a graded object in \mathcal{C}_{ung} , and therefore

$$\dim_{\mathcal{C}_{ung}} \mathcal{A}_{4D} \neq 1. \quad (9.46)$$

However, our aim is that of describing a composite configuration for the absolute theory of the kind depicted on the LHS of figure 41, where, as explained in sections 9.1.2 and 14, both phases really descend from the gauging of different subalgebras within the gaugeable algebra of the original ungauged SymTFT⁵⁶. The different 4D absolute gauge theories are interpolated by a non-invertible defect defined as

$$\mathcal{N}(\mathcal{M}_3) \stackrel{def.}{\equiv} \mathcal{D}_3 \mathcal{A}_{\epsilon\rho} \quad (9.47)$$

where $\mathcal{A}_{\epsilon\rho}$ plays the role of a relative gaugeable algebra that cannot be identified with the vacuum on either side.




Figure 39. Composite 4D absolute theories separated by an intrinsic non-invertible defect on the LHS. Rephrased in terms of the higher-categorical structure underlying the resulting theory, the algebra dressing the interpolating defects \mathcal{N}_3 is a homomorphism between the objects of a 2-category.

Consistency of the configuration associated to the LHS of figure 41, is mapped to a statement of Witt equivalence between the underlying 2-categories depicted on the RHS of the same figure, [374],

$$\mathcal{C} \boxtimes \bar{\mathcal{D}} \simeq \mathfrak{Z}(\mathcal{A}_{\epsilon\rho}), \quad (9.48)$$

where $\mathcal{A}_{\epsilon\rho}$ denotes the relative gaugeable algebra featuring in (9.47).

⁵⁶This is our proposal on the basis of [374, 607].

The relative gaugeable algebra, $\mathcal{A}_{\epsilon\rho}$, constitutes a fusion tensor category⁵⁷ defining morphisms between any pair of elements $L, L' \in \mathcal{C}, \mathcal{D}$.

$$\mathcal{A}_{\epsilon\rho}^{L,L'} \stackrel{def.}{=} \left\{ \mathcal{A}_1^{L,L'}, \mathcal{A}_0^{L,L'} \right\}, \quad (9.49)$$

whose total quantum dimension can be compared to that of the identity, i.e. of the algebra that has been condensed to achieve the reference gauge theory, with 2-category \mathcal{C} .




Figure 40. The algebra for the underlying category, denoted by \mathcal{A} can be partially condensed, under suitable choice of subalgebras $\mathcal{A}_1, \mathcal{A}_2 \subset \mathcal{A}$, such that they lead to different gauge theories, whose 2-categories are labelled by \mathcal{C}, \mathcal{D} .




Figure 41. The composite system (40) can be thought of as describing a thickened domain wall, ${}_{\mathcal{A}_1} \mathcal{A}_{\mathcal{A}_2} \stackrel{def.}{=} \mathcal{A}_1 \otimes_{\mathcal{A}} \mathcal{A}_2$.

From the point of view of the absolute 4D theory, the non-invertible defect can be thought of as a thickened domain wall obtained by condensing two different subalgebras

⁵⁷With \mathcal{A}_1 and \mathcal{A}_0 denoting the object and the morphisms, respectively.

within the original condensing algebra \mathcal{A} , as shown in figure 41. Given that the target absolute theory we want to achieve is that of figure 40, then the corresponding 2-categories of defects can be obtained starting from

$$\mathcal{A} \equiv \mathcal{D} \boxtimes \mathfrak{Z}(\mathcal{E}), \quad (9.50)$$

$$G : \mathcal{C} \boxtimes \mathfrak{Z}(\mathcal{D}) \simeq \mathcal{D} \boxtimes \mathfrak{Z}(\mathcal{E}), \quad (9.51)$$

and the fiber functors

$$\mathcal{F}_{\mathcal{E}} : \mathfrak{Z}(\mathcal{E}) \rightarrow \mathcal{E} \quad , \quad \mathcal{F}_{\mathcal{D}} : \mathfrak{Z}(\mathcal{D}) \rightarrow \mathcal{D} \quad (9.52)$$

with corresponding duals $\mathcal{F}_{\mathcal{E}}^V, \mathcal{F}_{\mathcal{D}}^V$.

The fact that $\mathcal{N}(\mathcal{M}_3)$ is intrinsically non-invertible corresponds to saying that the outer automorphism implemented by the action of F cannot be undone by a combination of modular transformations, as described in section 14. In particular, $\mathcal{A}_{\epsilon\rho}$ is different from the identity on either side of the defect, and, therefore, $\mathcal{D}_{\mathcal{A}_{\epsilon\rho}} \neq \mathcal{D}_{\mathbf{1}}$.

Proof

Take \mathcal{B} to be a semisimple abelian category over \mathbb{C} ; assuming it is a rigid braided tensor category. I is the set of isomorphism classes of irreducible objects in \mathcal{B} , with representatives $V_i, \forall i \in I$. Assuming the spaces of morphisms are finite-dimensional. $\mathbf{1} \in \mathcal{B}$ is assumed to be a simple object corresponding to the labelling $i \equiv 0 \in I \Rightarrow V_0 \stackrel{def.}{\equiv} \mathbf{1}$. A \mathcal{B} -algebra is an object $\mathcal{A} \in \mathcal{C}$ with morphisms $\mu : \mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$ and $\iota_{\mathcal{A}} : \mathbf{1} \hookrightarrow \mathcal{A}$, with

$$\dim \text{Hom}_{\mathcal{B}}(\mathbf{1}, \mathcal{A}) \equiv \mathbf{1}. \quad (9.53)$$

Given \mathcal{B} and a \mathcal{B} -algebra \mathcal{A} , $\text{Rep}\mathcal{A}$ is defined as pairs (V, μ_V) , where $V \in \mathcal{B}$ and $\mu_V : \mathcal{A} \otimes V \rightarrow V$ morphisms in \mathcal{B} .

For example, if G is a finite group and \mathcal{B} is the category of finite-dimensional complex representations of G , semisimple \mathcal{B} -algebras correspond to different semigroups in G (this is what we need to implement gauging in the Freed-Moore-Teleman setup).

Here, in $\text{Rep}\mathcal{A}$, \mathcal{A} is the unit object. For the case where there are two different absolute theories separated by an intrinsically non-invertible defect, the setup of joint absolute theories results from a double gauging procedure [Liang Kong] and it makes sense to consider the composite object $X \otimes_{\mathcal{A}} Y$ with \mathcal{A} the gaugeable algebra in \mathcal{C} ; so, we have that $X, Y \in \text{Rep}\mathcal{A}$. But, at the same time, X and Y are also elements of two different subalgebras within \mathcal{A} corresponding to two different gaugings. E.g. $X \in \text{Rep}\mathcal{A}_1$ and $Y \in \text{Rep}\mathcal{A}_2$, with $\mathcal{A}_1, \mathcal{A}_2 \subset \mathcal{A}$. In turn, \mathcal{A}_1 is the unit element in $\text{Rep}\mathcal{A}_1$ and \mathcal{A}_2 is the unit element in $\text{Rep}\mathcal{A}_2$.

Hence, if we consider the original mother algebra \mathcal{A} with its characteristic elementary objects V_i , and assuming every one of them has multiplicity 1 within \mathcal{A} , we get

$$\dim_{\mathcal{A}}(X \otimes_{\mathcal{A}} Y) \equiv \dim_{\mathcal{A}}(X) \dim_{\mathcal{A}}(Y) \quad (9.54)$$

and

$$\dim_{\mathcal{A}}(X) \equiv \frac{\dim_{\mathcal{B}}(X)}{\dim_{\mathcal{B}}(\mathcal{A})}. \quad (9.55)$$

Furthermore, we have that

$$\dim_{\mathcal{A}}(X \otimes_{\mathcal{A}} Y) \equiv \frac{\dim_{\mathcal{B}}(X) \dim_{\mathcal{B}}(Y)}{\dim_{\mathcal{B}}(\mathcal{A})}. \quad (9.56)$$

Extending equation (9.56) to the case in which X, Y are replaced by the $\mathcal{A}_1, \mathcal{A}_2$ subalgebras themselves, we can look at the following ratio

$$\frac{\dim_{\mathcal{B}}(\mathcal{A}_1 \otimes_{\mathcal{A}} \mathcal{A}_2)}{\dim_{\mathcal{B}}(\mathcal{A}_1)} \equiv \frac{\mathcal{D}_{\mathcal{A}}^{intr.}}{\mathcal{D}_{\mathcal{A}}^{non-intr.}}, \quad (9.57)$$

where on the RHS we identified the numerator and denominator with the quantum dimension of algebras associated to the intrinsic or non-intrinsic non-invertible defects separating absolute theories resulting from the Freed-Moore-Teleman construction discussed in the first part of section 9.1.1. Making use of (E.4) and (E.5), (9.58), we get

$$\frac{\dim_{\mathcal{A}}(\mathcal{A}_1 \otimes_{\mathcal{A}} \mathcal{A}_2) \dim_{\mathcal{B}}(\mathcal{A})}{\dim_{\mathcal{A}}(\mathcal{A}_1) \dim_{\mathcal{B}}(\mathcal{A})} \equiv \frac{\dim_{\mathcal{A}}(\mathcal{A}_1) \dim_{\mathcal{A}}(\mathcal{A}_2)}{\dim_{\mathcal{A}}(\mathcal{A}_1)} \equiv \dim_{\mathcal{A}}(\mathcal{A}_2). \quad (9.58)$$

Given that

$$\dim_{\mathcal{A}}(\mathcal{A}_2) \stackrel{def.}{=} \sum_i d_i V_i \quad (9.59)$$

with $V_i \in \mathcal{A}_2$ being simple objects in \mathcal{A}_2 (similar arguments hold for \mathcal{A}_1 , since both are assumed to be subalgebras in \mathcal{A} , whereas

$$d_i \stackrel{def.}{=} \dim_{\mathcal{A}} V_i, \quad (9.60)$$

it therefore follows that

$$\boxed{\frac{\dim_{\mathcal{B}}(\mathcal{A}_1 \otimes_{\mathcal{A}} \mathcal{A}_2)}{\dim_{\mathcal{B}}(\mathcal{A}_1)} \equiv \frac{\mathcal{D}_{\mathcal{A}}^{intr.}}{\mathcal{D}_{\mathcal{A}}^{non-intr.}} > 1}. \quad (9.61)$$

For the configuration of interest, namely two different absolute theories obtained by gauging of two different subalgebras $\mathcal{A}_1, \mathcal{A}_2 \subset \mathcal{A}$, separated by an intrinsic non-invertible defect, the numerator in (9.61) corresponds to the nontrivial fusion.

We are therefore lead to conclude that the total quantum dimension of the relative gaugeable algebra $\mathcal{A}_{\epsilon\rho}$ is greater w.r.t. that of the condensed algebra leading to the single gauged phase, \mathcal{C} , and we propose the following

$$\mathcal{N}(\mathcal{M}_3) \times \bar{\mathcal{N}}(\mathcal{M}_3) \equiv \mathcal{A}_{4D} \quad (9.62)$$

for class \mathcal{S} theories, whereas the denominator in (9.61) corresponds to the quantum dimension of the algebra that has been condensed in the SymTFT to obtain a specific absolute theory, namely

$$\mathcal{D}_3 \times \bar{\mathcal{D}}_3 \equiv \mathbf{1} \equiv \mathcal{A}_1 \quad (9.63)$$

Gathering these results together, we can rewrite equation (9.61) in terms of the notation adopted in the first part of this section, leading to the following result

$$\boxed{\frac{\langle \mathcal{N}(\mathcal{M}_3) \times \bar{\mathcal{N}}(\mathcal{M}_3) \rangle_{\mathcal{B}}}{\langle \mathcal{D}_3 \times \bar{\mathcal{D}}_3 \rangle_{\mathcal{B}}} \equiv \frac{\langle \mathcal{A}_{\epsilon\rho} \rangle_{\mathcal{B}}}{\langle \mathbf{1}_{\mathcal{B}} \rangle_{\mathcal{B}}} \equiv \left(\frac{\mathcal{D}_{\mathcal{A}}^{intr.}}{\mathcal{D}_{\mathcal{A}}^{non-intr.}} \right)_{X_4} > 1}, \quad (9.64)$$

with the last relation indicating that such ratio can be used as a parameter probing whether the two gauge theories are separated by an intrinsic non-invertible symmetry. Consequently, the following relation

$$\boxed{\mathcal{D}_{\mathcal{A}_{\epsilon\rho}(X_4)}^{intr.} > \mathcal{D}_{\mathcal{A}_{\epsilon\rho}(X_4)}^{non-intr.}} \quad (9.65)$$

implies that the fusion category associated to intrinsic non-invertible defects is characterised by a bigger quantum dimension, implying such configuration (arising from double gauging) is able to store a higher number of degrees of freedom w.r.t. the case in which there is a unique gauging taking place⁵⁸.

Key points

1. For class \mathcal{S} theories, $\mathcal{D}_{\mathcal{A}_{\epsilon\rho}}$ enables to probe whether a defect is intrinsically or non-intrinsically non-invertible.
2. $\mathcal{D}_{\mathcal{A}_{\epsilon\rho}(X_4)}^{intr.} > \mathcal{D}_{\mathcal{A}_{\epsilon\rho}(X_4)}^{non-intr.}$, and therefore the former is a theory equipped with a higher-categorical structure enabling to store more d.o.f. in certain superselection sectors.
3. For theories descending from 6D $\mathcal{N} = (2,0)$ SCFTs, if the gauging procedure leads to the presence of intrinsic non-invertible defects, the total quantum dimension of the algebra implementing the non-invertibility of such defects exhibits an increase in its total quantum dimension, signalling the relative gaugeable algebra $\mathcal{A}_{\epsilon\rho}$ is characterised by superselection sectors admitting a richer structure w.r.t. the condensed algebra leading to a single absolute 4D gauged theory.

⁵⁸Note that this criterion is perfectly in agreement with Remark 2.5 in [621].

At first sight, the title chosen for this article might appear provocative, and the length of the paper is certainly not fair towards the rich and deep advancements made by the phenomenological community so far⁵⁹. However, we stress that our aim is that of introducing important open questions in the realm of Particle Physics in a mathematical language approachable by categorical algebraic geometers.

Actually, this work is really meant to be the first of a series of papers by the same author continuing from past developments, bridging the gap between Pure Mathematics and Theoretical Particle Physics achievements. The author’s engagement with experts from both lines of research is what motivated this project in the first instance. In upcoming work, [597], we will be developing this description in terms of algebraic and Hilbert series calculation in greater detail. For the moment, we anticipate that the purpose of this introductory article is mainly fourfold:

- To illustrate (some) key fundamental problems in Theoretical Particle Physics necessitating further mathematical understanding.
- Opening the scene to well-established techniques in representation theory and categorical algebraic geometry to help in this regard.
- For those who are not experts in the field, serve as an oversimplified explanation of the powerful formalism put forward by Braverman, Finkelberg, and Nakajima (BFN), [598, 599]. In doing so we will explicitly show the relation in between their work, that of Freed, Moore, and Teleman (FMT), [102], and Teleman’s, [192], which, to the best of our knowledge, has only previously been referred to in [3].
- Motivating Phenomenologists to engage more with Pure Mathematicians, fostering mutual inspiration for common aims.

The main objective of this paper is that of explaining the higher categorical structures that are needed for describing the invariants associated to specific supersymmetric quiver gauge theories, with a particular focus on dualities and their mutual relations in terms of higher-categories. We believe that deepening its understanding and generalisations thereof, could lead to uncharted corners of mathematics that could be used for embedding of the Standard Model (SM) of Particle Physics, [28] in possible supersymmetric unifying theories, [18].

The present work is structured as follows:

1. Section 14 explains the motivations and outline of the present and forthcoming papers [597]. In doing so, we briefly introduce scattering amplitudes as crucial tools within the context of particle physics, explaining their connection with Hilbert series. We then briefly explain some of the key open questions in Theoretical Particle Physics

⁵⁹For a detailed overview of the state of the art and major developments in the field, we refer the interested reader to the rich literature at our disposal, including [21–27, 592–595], and [32, 48, 50, 58, 60, 61, 596] for slightly more theoretically-oriented perspectives in this regard, especially in relation with string theory, [8–15, 17].

necessitating further mathematical understanding, specifically the hierarchy problem, and the lightness of the Higgs mass (and its relation to Supersymmetry (SUSY), [18]).

2. Section 13 is mostly a revision of [102, 103], and we use it at this stage for multiple reasons. First of all because it allows to introduce gauge theories and Lagrangians in a mathematical language, which in turn is suitable to make connection with the higher-categorical Symmetry Topological Field Theory (SymTFT) prescription put forward by Freed, Moore, and Teleman, (FMT), [102], in the context of relative quantum field theories (QFTs).
3. The natural way of describing gauge theories is by means of quivers. The Standard Model itself admits such a representation. It is also the case, though, that quivers are crucial for describing certain algebraic varieties associated with highly supersymmetric setups. Section J is therefore devoted to explaining in a simplified way the well-established BFN construction leading to ring homologies. The theories in question are 3D $\mathcal{N} = 4$ supersymmetric quiver gauge theories. This section highlights the importance that categorical algebraic geometry plays in identifying ring homologies accounting for the complete spectrum of a given theory, therefore giving a complementary insight with respect to the Hilbert series alone. In so doing, we will highlight the importance of *abelianisation*, [600, 601]. In doing so we will explicitly show the relation in between their work, that of Freed, Moore, and Teleman (FMT), [102], and Teleman's, [192], which, to the best of our knowledge, has only previously been mentioned in [3].
4. From the discussion of the previous section, the concluding section, 21.4, paves the way to forthcoming work [597], where a more detailed analysis with regard to the algebraic structure and Hilbert series of (Beyond) the Standard Model scenarios will be carried out. Additional complementary directions addressed by the same author will be developing ring homologies for Moore-Tachikawa varieties, [136], beyond categorical duality⁶⁰, and understanding their implications for Koszul duality, [603]. We plan to report of any advancements in this regard in the near future, [597].

9.3 Abelianisation (once more)

The basic idea of abelianisation is to embed the Coulomb branch chiral ring $\mathbf{C}[\mathcal{M}_C]$ of a non-abelian theory into a larger, semisimple abelian algebra, \mathcal{A} ,

$$\boxed{\mathbf{C}[\mathcal{M}_C] \hookrightarrow \mathcal{A}}. \tag{9.66}$$

Essentially, abelianisation corresponds to fixed-point localisation in the equivariant homology, which we will briefly explain momentarily. The embedding (J.5) allows to:

1. Verifying relations among elements of $\mathbf{C}[\mathcal{M}_C]$.

⁶⁰Thereby following up from the analysis carried out in [4].

2. Identifying the Poisson structure on $\mathbf{C}[\mathcal{M}_C]$ and its deformation quantisation.
3. Extend the algebra $\mathbf{C}[\mathcal{M}_C]$ over twistor space, ultimately enabling to access the hyperkähler structure on \mathcal{M}_C .

In the context of BFN, this arises when considering a 3D $\mathcal{N} = 4$ gauge theory on $\mathbb{C} \times \mathbb{R}$ spacetime with $\frac{1}{2}$ -BPS boundary conditions \mathcal{B} near spatial infinity on \mathbb{C} .




Figure 42. Picture reproduced from the work of [600], encoding the key idea outlined in their treatment, and from which our work was in turn inspired.

Working in the cohomology of the Rozansky-Witten theory, \mathcal{Q}_{RW} , the fields may be localised in the QM to sections of bundles that solve the \mathcal{Q}_{RW} BPS equations. This yields the gauged $\mathcal{N} = 4$ QM on

$$\mathcal{M}_{[\mathbf{C}]} = \{ (E, X) \}, \quad (9.67)$$

where E is a holomorphic $G_{\mathbf{C}}$ -bundle on \mathbf{C} , and X is a holomorphic section of an associated R -bundle with gauge group, $calG$ is the holomorphic $G_{\mathbf{C}}$ gauge transformation on \mathbf{C} .

The Hilbert space of the space of the QMs in the \mathcal{Q}_{RW} twist should be the \mathcal{G} -equivariant de Rham cohomology of $\mathcal{M}_{[\mathbf{C}]}$. Turning on the Omega background further, corresponds to working equivariantly w.r.t. the $U(1)_{\epsilon}$ spatial rotation group of \mathbf{C} , which is an ordinary symmetry of the moduli space, the Hilbert space is

$$\mathcal{H} \stackrel{def.}{=} H_{G \times U(1)_{\epsilon}}^{\bullet} \left(\mathcal{M}_{[\mathbf{C}]} \right). \quad (9.68)$$

$\mathcal{M}_{[\mathbf{C}]}$ is contractible to a point where E is the trivial bundle and X is the zero-section. \mathcal{G} is contractible to G , therefore the Hilbert space reduces to

$$\mathcal{H} \simeq H_{G \times U(1)_{\epsilon}}^{\bullet} (\text{pt.}) = \mathbf{C}[\varphi, \epsilon]^W, \quad (9.69)$$

namely Weyl-invariant polynomials in the equivariant weights $\varphi \in \mathfrak{t}_{\mathbf{C}}$ and ϵ .

We can therefore write the BFN construction, which essentially reads as follows

$$\mathbf{C}_{\epsilon} [\mathcal{M}_C] \equiv H_{g \times g' \times U(1)_{\epsilon}}^{\bullet} \left(\mathcal{M}_{[\mathbf{C} \cup \mathbf{C}]} \right) \simeq H_{G \times U(1)_{\epsilon}}^{\bullet} \left(\mathcal{M}_{[\mathbf{C} \cup \mathbf{C}]} / \mathcal{G}' \right), \quad (9.70)$$

where $\mathcal{G}, \mathcal{G}'$ are the groups of regular holomorphic gauge transformations on the top and bottom copies of \mathbf{C} , and $U(1)_{\epsilon}$ is the usual spatial rotation group. Importantly, this space,

(9.70), is non-contractible, and has highly nontrivial topology since it contains monopole operators. OPEs of local operators in the algebra $\mathbf{C}_\epsilon[\mathcal{M}_\epsilon]$ naturally correspond to the convolution product rather than the cup product.

We are now ready to explain a particularly useful tool in the context of equivariant cohomology, namely fixed-point localisation. Letting $T \subset G$ denote the maximal torus, one finds that $T \times U(1)_\epsilon$ fixed points of $\mathcal{M}_{[\mathbf{C} \cup \mathbf{C}]} / \mathcal{G}'$ are isolated points described as pairs $(E, X), (E', X')$ where $X = X' = 0$ are zero sections, E is trivial, and E' is obtained from E by a gauge transformation

$$g(z) = z^A \quad , \quad A \in \text{cochar}(G), \quad (9.71)$$

meaning fixed points are labelled by cocharacters corresponding to abelian monopole operators.

If \mathcal{F} is the fixed point set of the $T \times U(1)_\epsilon$ action on $\mathcal{M}_{[\mathbf{C} \cup \mathbf{C}]} / \mathcal{G}'$, $\mathcal{F} \simeq \text{cochar}(G)$ is isomorphic to the cocharacter lattice.

The equivariant cohomology of the fixed point set contains a copy of

$$H_{T \times U(1)_\epsilon}^\bullet(\text{pt}) = \mathbf{C}[\varphi, \epsilon] \quad (9.72)$$

for every point in \mathcal{F} , i.e.

$$H_{T \times U(1)_\epsilon}^\bullet(\mathcal{F}) \simeq \mathbf{C} \left[\varphi, \epsilon, \{v_A\}_{A \in \text{cochar}(G)} \right]. \quad (9.73)$$

In its localised version, one gets that

$$\mathcal{A}_\epsilon \subset H_{T \times U(1)_\epsilon}^\bullet(\mathcal{F})^{\text{loc}}. \quad (9.74)$$

The crucial difference in between the two sides in (9.74) is that the localised cohomology indiscriminately inverts all weights, whereas in \mathcal{A}_ϵ we only invert part of them. The localisation theorem provides the following algebraic embedding

$$\mathbf{C}_\epsilon \left[\mathcal{M}_{[\mathbf{C}]} \right] = H_{G \times U(1)_\epsilon}^\bullet \left(\mathcal{M}_{[\mathbf{C} \cup \mathbf{C}]} / \mathcal{G}' \right) \hookrightarrow H_{T \times U(1)_\epsilon}^\bullet(\mathcal{F})^{\text{loc}}, \quad (9.75)$$

which is the abelianisation we introduced in (J.5). Importantly, the RHS of (9.75) is the mother algebra we were looking for, the un-gauged algebra referred to in section 13, and the key entity encoding essential information about the QFT in question. In conclusion to this section, we wish to highlight that, for any successful QFT to be fully specified, one should really be able to determine such an embedding. Any shortcomings in achieving it, should be taken as signalling a yet incomplete understanding of the underlying mathematical structure of the theory in question.

10 Instantons and Scattering Amplitudes

10.1 Instantons

Instantons are solutions to the classical equations of motion in Euclidean spacetime with a finite non-zero action. They arise in, both, QFT and QM, and can be thought of as critical points of the action.

In QFT, instantons:

1. Appear in the path-integral as leading quantum corrections to the classical behaviour of the system, and
2. Can be used to study tunnelling in various systems, such as Yang-Mills (YM) theory.

In mathematical terms, a YM instanton is a self-dual or anti-self-dual connection in a principal bundle over a 4D Riemann manifold, and are topologically nontrivial solutions to the YM equations minimising the energy functional.

10.1.1 Instantons in QM

An example of a system with an instanton effect is the double-well potential, for example

$$V(z) = (z^2 - 1)^2, \quad (10.1)$$

in which case the classical minima lie at $z = \pm 1$, corresponding to the two lowest energy states. In QM, instead, the energy eigenstates are found by solving the Schrödinger equation

$$-\frac{\hbar^2}{2m} \frac{\partial}{\partial z^2} \psi + V\psi = E\psi. \quad (10.2)$$

In this case, though, the lowest-energy eigenvalue is unique, since the ground state wavefunction localises at both classical minima, $z = \pm 1$, due to quantum interference, which justifies introducing the notion of tunnelling in the first place. In particular, instantons enable to explain why this is so in the semi-classical approximation of the path-integral formulation in Euclidean time.

In the path-integral formulation of the tunnelling process, the transition amplitude reads as follows

$$\begin{aligned} K(z_1, z_2; t) &= \left\langle z = z_1 \left| e^{-\frac{iHt}{\hbar}} \right| z = z_2 \right\rangle \\ &= \int d[z(t)] e^{\frac{iS}{\hbar}}. \end{aligned} \quad (10.3)$$

Under Wick-rotation,

$$\begin{aligned} \tilde{K}(z_1, z_2; t) &= \left\langle z = z_1 \left| e^{-\frac{Ht}{\hbar}} \right| z = z_2 \right\rangle \\ &= \int d[z(t)] e^{-\frac{S_E}{\hbar}}, \end{aligned} \quad (10.4)$$




Figure 43.

with Euclidean action

$$S_E = \int_{\tau_1}^{\tau_2} \left(\frac{1}{2} m \left(\frac{dz}{d\tau} \right)^2 + V \right) d\tau. \quad (10.5)$$

Specifying to the double-well potential (10.1.1), setting $z_2 = -z_1 = 1$, (10.5) becomes

$$\begin{aligned} S_E &= \int_{\tau_1}^{\tau_2} \left[\frac{1}{2} \left(\left(\frac{dz}{d\tau} \right)^2 - \sqrt{2V} \right)^2 + \sqrt{2} \frac{dz}{d\tau} \sqrt{V} \right] d\tau \\ &= \int_{\tau_1}^{\tau_2} \frac{1}{2} \left(\left(\frac{dz}{d\tau} \right)^2 - \sqrt{2V} \right)^2 d\tau + \sqrt{2} \int_{-1}^1 (1 - z^2) dz \\ &\geq \frac{4\sqrt{2}}{3}. \end{aligned} \quad (10.6)$$

The inequality in the last line of (10.6) is saturated by the solution of the following differential equation (arising by setting the first integral in (10.6) to zero)

$$\frac{dz}{d\tau} = \sqrt{2V} \quad (10.7)$$

subject to the boundary conditions

$$z(\tau_1) = -1, \quad z(\tau_2) = 1. \quad (10.8)$$

The instanton solution interpolating between the two classical minima therefore reads

$$z(\tau) = \tanh(\tau - \tau_o), \quad (10.9)$$

where $\tau_o = \text{const.}$, and interpolates between the two asymptotic values set by the boundary conditions (10.8).




Figure 44.

10.1.2 Instantons in QFT

Instantons also arise when studying the vacuum structure of QFTs, where the true vacuum can be defined as an overlap of several topologically-inequivalent sectors (also known as topological vacua). Among the most known examples are QFTs with non-abelian gauge group, i.e. Yang-Mills theories. In such case, the inequivalent topological sectors are classified by means of the third homotopy group of $SU(2)$,

$$\pi_3(S^3) = \mathbb{Z}, \quad (10.10)$$

whose group manifold is the 3-sphere, S^3 . From (10.10), it therefore follows that there are infinitely-many topologically-inequivalent vacua, labelled by the Pontrjagin index, N .

In this case, an instanton is interpreted as a tunnelling effect between these inequivalent topological vacua. More concretely, if an instanton is assigned a Pontrjagin index Q , then it mediates between the topological vacua $|N\rangle$ and $|N+Q\rangle$, whereas the true vacuum of the theory would be expressed as the following linear supersposition

$$|\theta\rangle \stackrel{def.}{=} \sum_{N=-\infty}^{+\infty} e^{i\theta N} |N\rangle. \quad (10.11)$$

10.1.3 Instantons in 4D supersymmetric gauge theories

Supersymmetric gauge theories often obey nonrenormalisation theorems, with the latter restricting the allowed kinds of quantum corrections. Many such theorems only apply to perturbative corrections, reason why instantons, that are non-perturbative in nature, turn out being the only corrections possible.

Many instanton calculations were performed in supersymmetric theories during the 1970s and 1980s⁶¹.

⁶¹See for instance [67].

According to the amount of supersymmetry that is present, different corrections can be considered:

1. $\mathcal{N} = 1$ SUSY gauge theories: instantons can modify the superpotential, potentially lifting the vacua.
2. $\mathcal{N} = 2$: the superpotential receives no quantum correction, but the corrections to the metric of the space of vacua from instantons can be nontrivial.
3. $\mathcal{N} = 4$: instantons lead to no quantum corrections, neither in the superpotential, nor the Kähler metric.

10.1.4 Instantons in oriented string theory: the ADHM construction

A Dp -brane is a gauge theory instanton in the world volume $(p + 5)$ -dimensional $U(N)$ gauge theory on a stack of N $D(p + 4)$ -branes.

YM-instantons are used for studying the vacuum structure of YM theory and classifying 4-manifolds. A method for studying self-dual YM-instantons is due to ADHM, who first performed this calculation for classical gauge groups, $SU(N)$, $SO(N)$, and $Sp(N)$. Douglas, [484], and Witten, [485], found that the ADHM construction can be realised in string theory. In particular, the moduli space of instantons on \mathbb{R}^4 is equivalent to the Higgs branch of supersymmetric gauge theories on a system of Dp - $D(p + 4)$ branes. These are quiver gauge theories with 8 supercharges (e.g. 4D $\mathcal{N} = 2$ for $p = 3$).

The moduli space of instantons is the Higgs branch of SUSY gauge theories. Gauge theories for classical gauge groups can be embedded as worldvolume theories of Dp -branes in backgrounds of $D(p + 4)$ -branes.

Self-dual instantons on $\mathbb{C}P^2$ or \mathbb{C}^2 are described an ADHM-like construction allowing to calculate the Hilbert series of the moduli space⁶².

10.2 The WKB approximation

The aim is that of understanding the geometric structures naturally arising when trying to solve the Schrödinger equations. It involves symplectic geometry, spectral theory, and microlocal analysis.

The WKB method is a technique adopted for finding approximate solutions to the Schwarzschild equation. More sophisticated developments of this have led to other techniques such as microlocal analysis, [194].

For a given Hamiltonian

$$H(q, p) = \frac{p^2}{2m} + V(q) \tag{10.12}$$

the Hamiltonian flow equations read

$$\dot{q} = \frac{p}{m} \quad , \quad \dot{p} = -V'(q). \tag{10.13}$$

⁶²For a more detailed explanation and overview of the Hilbert series in this context, please see appendix P.

Fixing $\hbar \in \mathbf{R}_+$, the Schrödinger equation reads as follows

$$i\hbar \frac{\partial \psi}{\partial t} = \hat{H} \psi \quad \text{with } \hat{H} = -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + m. \quad (10.14)$$

Looking for stationary solutions of the form

$$\psi(x, t) = \varphi(x) e^{-i\omega t}, \quad (10.15)$$

and assuming

$$\varphi(x) = e^{ix\xi}, \quad (10.16)$$

namely, as V varies with x , so too does ξ , the WKB approximation leads to a solution of the form

$$\varphi(x) = e^{iS(x)/\hbar}, \quad (10.17)$$

with $S(x)$ the phase function solving the Hamilton-Jacobi (HJ) equations obtained from

$$(\hat{H} - E)\varphi = \left[\frac{(S'(x))^2}{2m} + (V - E) - \frac{i\hbar}{2m} S''(x) \right] e^{iS(x)/\hbar} \quad (10.18)$$

assuming $\hbar \ll 1$,

$$H(x, S'(x)) = \frac{S'(x)^2}{2m} + V(x) = E \quad (10.19)$$

$$\Rightarrow S'(x) = \pm \sqrt{2m(E - V(x))}. \quad (10.20)$$

Given the classical phase plane $\mathbb{R}^2 \simeq T^*\mathbb{R}$ with coordinates (q, p) , the differential $dS = S' dX$ can be viewed as a mapping

$$dS : \mathbf{R} \rightarrow T^*\mathbf{R}, \quad (10.21)$$

where $p \equiv S'$. Then, S satisfies the HJ equation if and only if the images of dS lie in the level manifold $H^{-1}(E)$.

These arguments can be extended to \mathbf{R}^n , at which point we get

$$S : \mathbf{R}^n \rightarrow \mathbf{R} \quad (10.22)$$

as a solution of the HJ equation.

In the remainder of this section, we briefly overview the applicability of the WKB approximation to two setups that are most important for our work, namely Hitchin systems and vacuum transitions.

10.2.1 WKB for Hitchin systems

Given the construction of the coordinates⁶³ of appendix I.3 associated to the metric on the Riemann surface for class \mathcal{S} theories, one can look for particular asymptotic solutions of the connection \mathcal{A} such that

$$\lim_{\zeta \rightarrow 0} \mathcal{A} = \frac{R\varphi}{\zeta}. \quad (10.23)$$

At every point of the Riemann surface, φ takes two eigenvalues, $\pm\lambda$, with λ being a multi-valued 1-form on the Riemann surface. The WKB approximation then states that, if one chooses a gauge such that

$$\varphi = \begin{pmatrix} \lambda & 0 \\ 0 & -\lambda \end{pmatrix}, \quad (10.24)$$

then there are two independent approximate \mathcal{A} -flat sections of the type

$$\psi^{(1)} \sim \begin{pmatrix} e^{-\frac{R}{\zeta} \int \lambda} \\ 0 \end{pmatrix}, \quad \psi^{(2)} \sim \begin{pmatrix} 0 \\ e^{-\frac{R}{\zeta} \int \lambda} \end{pmatrix}, \quad (10.25)$$

and one would therefore expect that, in the $\lim_{\zeta \rightarrow 0}$, the calculation reduces to that of periods of the 1-form λ on the Riemann surface⁶⁴.

10.2.2 WKB for vacuum transitions

The treatment of the first part of this subsection extends to the case of QM tunnelling between different minima of a given theory. To see this explicitly, it is sufficient to consider the simplest setting possible, where the field theory in question is defined by the action

$$S = \int d^4x \left(-\frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi - V(\varphi) \right). \quad (10.26)$$

The corresponding energy of the system therefore reads

$$E[\varphi(x)] = \int d^3x \left(\frac{1}{2} (\nabla \varphi)^2 + V(\varphi) \right). \quad (10.27)$$

The specific profile of the potential, $V(\varphi)$, is obviously going to dictate the types (and number) of vacua available in a give theory.

Albeit (10.26) explicitly being higher-dimensional, under the assumption of spherical symmetry, the nucleation process can be recast to that of QM tunnelling, whose unique degree of freedom is the radius of the new vacuum bubble emerging in the background vacuum. Figure 45 is useful in understanding what we have just said. In the under-the-barrier region, the solution to the HJ equations, (10.28), becomes imaginary, leading to the definition of the *bounce*

⁶³We refer the reader to the respective section in the appendix for a more detailed analysis of the specific setup, which is a review of the original work by Gaiotto, Moore, and Neitzke.

⁶⁴More detailed explanation on this terminology is provided in section I.3.




Figure 45.

$$B \stackrel{def.}{=} iS = 2i \int_{z_1}^{z_2} \sqrt{2m(E - V(x))} \quad (10.28)$$

In the following Part, we will be analysing in great depth the case of vacuum transitions for 2D field theories with gravity, where the procedure just outlined is also known as the Coleman-De Luccia vacuum decay description. For reasons that we will be explaining in due course, though, we will also be outlining the importance of identifying alternative methods for calculating decay rates. The need to do so is multifold. If we were to pick one of the most crucial points to be addressed, is the absence of a closed-form expression for the String Theory Landscape potential. More of this will be motivated in section 12, as introductory to Part IV.

11 Conclusions and Outlook of Part III

Key advancements towards achieving a rigorous mathematical formulation of QFTs strongly relies upon a higher-categorical prescription. In particular, the symmetries of a QFT can be thought of as defects living in different categories connected by functors.

In this Part of our work, we identified a criterion to distinguish between intrinsic and non-intrinsic non-invertible symmetries arising from the categorical structure of 6D $\mathcal{N} = (2, 0)$ SCFTs. In particular we were able to derive a relation in terms of the quantum dimension of the condensing algebra implementing the gauging in the bulk SymTFT. In doing so, we relied upon the description in terms of relative field theories, as described in [367] in terms of the Freed-Moore-Teleman setup, [102, 103], together with the gauging prescription, [607]. For the intrinsic non-invertible case, multiplicity is greater w.r.t. the non-intrinsic case, thereby signalling the possibility for additional d.o.f. to be stored in certain superselection sectors of the resulting absolute theory.

Our results extend arguments proposed in [333], where the authors also proposed a way of distinguishing intrinsic from non-intrinsic non-invertibility in 2D by means of the quantum dimension of the non-invertible defect.

In our analysis, we encountered two major setups built upon the notion of higher-categories, namely symmetry TFTs (SymTFT) and topological orders (TO). Plenty of effort has been made towards building a correspondence in between the two, most recently in [376]. Our findings provide further support towards strengthening the connection between the two description.

We conclude this Part by stressing that our analysis is mostly motivated by furthering the understanding of 6D $\mathcal{N} = (2, 0)$ SCFTs, from, both, a mathematical and physical point of view. To what extent these findings might be mapped to other setups is currently under investigation. In an upcoming work [597], we will report on further advancements building on these findings.

Part IV

Holographic interpretation of 2D vacuum transitions

12 Summary and Motivations of Part IV

12.1 Motivations




Figure 46. The string theory landscape.

As already mentioned in the introductory part of the present work, the original motivation to our work was that of studying the stability of vacua arising in the string theory landscape. The main reason for this being that of identifying a string theory model that

would better suit the description of our universe in its present-day status, as well as describing its evolution according to well-established cosmological models.

A faithful description of the stability of such vacua, requires identifying the fundamental degrees of freedom characterising the minima of the string theory landscape, and plenty of effort has been made so far in doing so, by means of, both, analytic and numerical techniques⁶⁵. However, to the best of our knowledge, no systematic method has yet been identified as effectively describing the decay of such vacua into others, and the work in collaboration with my supervisor constitutes a step forward towards this aim.

Figure 46 was purposely meant to show the randomness with which different vacua build the string theory landscape, highlighting the lack of a unique potential accounting for all of them at once.

Identifying a rigorous formalism towards describing vacuum decay processes has been a great matter of debate for almost half a century, [208–210], and part of our joint collaboration was attempting to provide further insights into which description could better suit our purposes.

Together with the quantum study of black hole physics, the study of vacuum transitions provides a rich arena to explore quantum aspects of gravity, and may shed some light towards a proper understanding of quantum gravity. Recent progress in the study of black hole information (for a review see [211]) has been achieved, in great part, studying concrete examples in 2D, and it is natural to explore if the concepts used there, such as quantum extremal surfaces, generalised entropies and islands may also play a role in addressing questions regarding other physical systems involving quantum aspects of gravity, namely, early universe cosmology and vacuum transitions.

The following four sections of this work aim at providing a general perspective towards understanding vacuum transitions in 2D. The reason for doing so is multi-fold. Among the most important, we highlight the following:

1. To understand, from the simple 2D set-up, the stability of vacua with different values and signs of the vacuum energy, in particular having in mind the string theory landscape.
2. To study the possible emergence of unitarity-violating behaviour, such as encountered in the black hole information paradox [212–214], in a different physical system.
3. To identify potential holographic interpretations of vacuum transitions, [203, 224], and their generalisations to higher-dimensional holographic embeddings and defect field theories.
4. To explore the possibility of assigning an effective entropy to transition amplitudes for spacetimes with arbitrary sign of the cosmological constant. The main motivation for this comes from recent developments towards studying the emergence of spacetime from entanglement [225–228].

⁶⁵Among the most important works in this regard, we refer the interested reader to the following [29–66].

Different approaches, same aim

Throughout the last four decades, the study of quantum transitions in 4D has been addressed by means of three main approaches, differing in terms of the way the cosmological constants are being defined, and the use of Euclidean or Lorentzian techniques. In order of appearance in the literature, they are:

- Coleman-de Luccia (CDL) [208], describing transitions between different local minima in a scalar field potential, following a Euclidean approach.
- Brown-Teitelboim (BT) [209], Euclidean vacuum transitions mediated by brane nucleation.
- Fischler-Morgan-Polchinski (FMP) [210], transitions between two spacetimes with different cosmological constants, by means of the Hamiltonian formalism in Lorentzian signature.

In each case, the quantity of interest is the transition amplitude, Γ . In the Euclidean methods, this is obtained from the *bounce*, B , defined as the difference between the Euclidean action evaluated on the instanton $S_E|_{inst}$ and on the background $S_E|_{bckgr}$:

$$\Gamma \sim \exp(-B) \quad , \quad B \stackrel{def.}{=} S_E|_{inst} - S_E|_{bckgr}. \quad (12.1)$$

The main motivation for this approach follows from the relation between the WKB approximation and vacuum transitions, with the latter being viewed as the quantum mechanical process of a particle crossing a potential barrier.

The Lorentzian method of FMP, instead, proposes an alternative definition for Γ ; the transition amplitude from a spacetime \mathcal{M}_1 to another \mathcal{M}_2 , is defined as the relative ratio of two probabilities,

$$\Gamma_{1 \rightarrow 2} \stackrel{def.}{=} \frac{P_{\text{nothing} \rightarrow \mathcal{M}_1 / \text{Wall} / \mathcal{M}_2}}{P_{\text{nothing} \rightarrow \mathcal{M}_1}} = \frac{|\Psi_{\text{nothing} \rightarrow \mathcal{M}_1 / \text{Wall} / \mathcal{M}_2}|^2}{|\Psi_{\text{nothing} \rightarrow \mathcal{M}_1}|^2} \quad , \quad (12.2)$$

with both nucleations out of nothing being identified with Hartle-Hawking (HH)-like states [229]. This is a conditional probability, with the numerator and denominator corresponding to the end point of the transition and the original background spacetime, respectively.

In an earlier paper, [7] (see also [230, 231]), we recovered the original BT result for $dS_4 \rightarrow dS_4$ ⁶⁶, by means of the FMP method, with total action given by

$$\log \Gamma_{dS \rightarrow dS} = \frac{\eta\pi}{2G} \left[\frac{[(H_o^2 - H_I^2)^2 + \kappa^2(H_o^2 + H_I^2)]R_o}{4\kappa H_o^2 H_I^2} + \frac{1}{2H_o^2} - \frac{1}{2H_I^2} \right] \quad , \quad (12.3)$$

Here H_o, H_I refer to the Hubble parameter for the spacetimes outside and inside the bubble, κ is the tension of the bubble, R_o its radius at nucleation and $\eta = \pm 1$. Under mutual exchange of the 2 vacua, the 1st term in (12.3), coming from the brane, is symmetric,

⁶⁶The outer and inner vacua are denoted by o, I , respectively.

whereas the other 2 terms just flip their sign. From this follows that, the ratio between the direct and reverse transition reads

$$\frac{\Gamma_{o \rightarrow I}}{\Gamma_{I \rightarrow o}} = e^{\frac{\eta\pi}{2G} \left(\frac{1}{H_o^2} - \frac{1}{H_I^2} \right)} = e^{\eta(S_o - S_I)} \quad (12.4)$$

For $\eta = 1$, since S_o and S_I are the entropies of the corresponding de Sitter spacetimes, this is a statement of *detailed balance*. According to [233], such entropic argument cannot be extended to spacetimes other than dS⁶⁷, due to the topological change the spacetime would undergo as a consequence of the change in sign of its curvature. Furthermore, upon taking a vanishing background cosmological constant, (12.3) diverges, thereby forbidding up-tunnelling from a flat spacetime. In such case, (12.4) would thereby suggest that the background entropy is divergent. However, this is clearly in contrast with the common lore that the Minkowski vacuum should have vanishing entropy instead.

Following the steps of [210], whose work was in turn motivated by [234]⁶⁸, we showed in [7] that this issue can be addressed by introducing a Schwarzschild black hole in the background, and realising Minkowski as the limit of the vanishing mass of the black hole itself. Indeed, the total action, in such case, reads

$$S_{\text{Sch} \rightarrow \text{dS}} = \frac{\eta\pi}{2G} [S_{\text{bounce}} - S_{\text{bckg}}] = \frac{\eta\pi}{2G} [S_{\text{bounce}} - 4G^2 M^2] \quad (12.5)$$

and clearly remains finite in the formal limit ⁶⁹ $\lim_{M \rightarrow 0}$.

The question of whether, and how, we can possibly reconcile these apparently contrasting behaviours partly motivated the present work, with the possibility to correctly assign an entropy to the spacetimes involved. Before embarking in the analysis outlined in the present work, our original interpretation was that transitions were taking place in *local* regions of spacetime; as such, topologically-inspired arguments for the inapplicability of detailed balance could be dropped. But still, a probably more interesting question emerged: can we assign an entropic interpretation to the direct amplitude itself and not only to the ratio of the amplitudes, independently of detailed balance? As we shall see, the answer will turn out to be affirmative in 2D.

In achieving this result, holography will turn out to play a key role. Since its 1st formulation, [203], there have been many generalisations: AdS/BCFT, [236], wedge holography (codim-2), [237], $T\bar{T}$, [238–242], the dS/CFT, [243], and dS_{D+1}/dS_D correspondence, [244], etc. Last but not least, the *island*, [245–247], pushing the holographic duality to

⁶⁷The definition of the entropy for the static dS patch is a longstanding matter of debate. Motivated by our findings, we will be arguing that the entropy definition arising from the study of vacuum transitions is only meaningful when considering a finite portion of the dS patch. Our findings are therefore in agreement with those of others, such as, e.g. [232], where the authors outline the need to introduce a boundary to dS for assigning a notion of entropy to it.

⁶⁸The configuration described by Farhi, Guth and Guven, [234], has recently been addressed in the literature, [235], and claimed to be unsuitable for the application of detailed balance. However, we will be able to prove that the claims made in [235] are not applicable in our setup given the locality of the processes being described.

⁶⁹Note that this limit is only an approximation since the lower bound on a black hole mass is the Planck scale.

its currently most extreme formulation, since it involves the entanglement of spatially-disconnected regions. Throughout our treatment we will encounter applications⁷⁰ of all such cases within the context of 2D vacuum transitions, with the different methods showing complementary holographic interpretations, thereby consistently reconciling all predictable disagreements.⁷¹

Our main result in this regard is having found expressions of the form (12.3) for vacuum transition rates in the 2-dimensional case. We will perform the calculation in all three formalisms outlined above (CDL, BT and FMP), providing their corresponding holographic interpretation. Given that the BT case has already been addressed by the authors in their original work, [209], we will be using it as a “standard ruler” to check the results obtained by other means. In doing so, however, we will not simply take it for granted, but, rather, derive it from the suitably adjusted setup of CDL in 2D, which, to the best of our knowledge, has not been dealt with explicitly in the literature. As outlined in section 13, the common starting point for both Euclidean methods will be the Almheiri-Polchinski action, [268],

$$S_{2D} = \int d^2x \sqrt{-g} [\phi^2 \mathcal{R} + \lambda \phi'^2 - U(\phi)] \quad (12.6)$$

which resembles the starting point of [208] for decays in the presence of gravity.

Due to the specific features of the 2D setup, though, the Ricci scalar or the scalar kinetic term can be eliminated by a suitable rescaling of the metric and of the potential. As we shall see, removing the ϕ'^2 -term we can calculate the transition amplitude either with or without the *thin-wall* approximation, proper to CDL. On the other hand, upon removing the Ricci scalar term, and adding suitable boundary terms ensuring the action satisfies a variational principle, we recover the gravitational action of BT. The importance of the addition of the boundary terms will be highlighted in due course, but it is worth stressing that, for the case in which the kinetic term is suppressed, they trivially vanish in either vacuum. Correspondingly, this is mapped to the fact that the calculation by means of the CDL method in 2D describes transitions in *absence* of gravity. From the holographic interpretation, we will see that, indeed, the total bounce is reminiscent of the entropy of an *internal* CFT₂, which geometrises the RG-flow interpolating between different values of the dilaton.

For the case of BT, instead, the addition of the boundary terms, accounting for the presence of gravity, is encoded in boundary entropies of defect CFT₁ s placed at the end-points of the *bulk* CFT₂, which in turn are dual to the end-of-the-world (ETW) branes, [269, 270], where the spacetimes undergoing the transition live (cf. figure 89). A more detailed explanation of figure 89 will be provided in section 13.

⁷⁰For other recent applications of entanglement and islands within a cosmological setup, please see, e.g. [248–264].

⁷¹This shows that the holographic obstructions towards realising specific cosmological and gravitational processes (as in [265–267]), can be overcome.




Figure 47. The CFT_2 geometrises the RG-flow defining B_{wall} , and in this sense is *internal*. The CFT_1 s are the duals of the AdS_2 spacetimes involved in the transition living on the 2 ETW branes. Their opening angles, $\Theta_{1,2}$, and corresponding tensions, result in the value of Λ_{\pm} on the 2 sides of the wall. A more detailed explanation is provided in section 13

12.2 Outline and main results

In summary, Part IV of this thesis is structured as follows: in section 13 we extend the CDL formalism to 2D, which, to the best of our knowledge, has not been explicitly done in the literature. In doing so, we use the JT-gravity theory analysed by Almheiri and Polchinski [268]. Thanks to the nature of the 2D setup, under suitable Weyl transformation, we can trade the kinetic term for the dilaton with the Ricci scalar. We thereby perform the calculation in both ways, showing that the case without the kinetic term describes transitions in absence of gravity. Upon removing the Ricci term, instead, the action can be brought back to the same form as in BT, up to a boundary term whose role is essential to account for the gravitational interaction. The second part of the section provides a brief overview of the 2D treatment of BT, as already addressed by the authors in their original work, [209], also highlighting some key features that will turn out useful in following sections.

In section 14, we turn to the Hamiltonian formalism of [210] applied to 2D JT-gravity [271–273], and determine the transition rates in absence of black holes, finding that uptunnelling from and downtunnelling to Minkowski spacetimes are not allowed.

In section 15, we generalise the results of section 14 by adding a constant term in the action that allows for non-extremal black holes to emerge, as long as their mass lies within a certain range to be specified in due course. The range emerges from requiring the existence of 2 distinct physical turning points. We conclude that, upon adding a black hole of suitable mass on either side, the flat spacetime limit does not violate unitarity, and we are still left with a well defined transition amplitude.

In section 16 we provide a possible holographic interpretation of the total bounces and actions calculated throughout our work, showing mutual compatibility and complementarity. Altogether, this leads to an exhaustive explanation of figure 89. In particular, we find that the corresponding expression for the transition rate in presence of gravity, and in absence of black holes, is given by the difference of entropies of $T\bar{T}$ -deformed CFTs, hence proving the locality of the nucleation process. Upon adding black holes, within a

suitable mass range, instead, the total action can be expressed as the difference of generalised entropies, with an island emerging beyond a critical value of the black hole mass. In particular, we show that, whenever an island is present, uptunnelling is always possible. Furthermore, the results obtained by means of the FMP method are found to agree with the expression provided by [274] for describing mutual approximation of boundary states belonging to different CFTs under suitable parametric redefinition. The BT results, which in section 15 were proved to be equivalent to the FMP results in absence of black holes, can be expressed in terms of entropies of BCFT₂s with 2 nontrivial boundary conditions dual to ETW branes. Last but not least, the CDL result can be recast in the form of an entropy product of a CFT₂, thereby showing agreement with the expectations following from the analytic behaviour encountered in section 13. These findings further support the interpretation of the dilatonic field as playing the role of an *internal* CFT₂ on which the entanglement entropy is being evaluated.

One of the key results of our work can therefore be synthesised by saying that the total action (or bounce) associated to the direct vacuum transition process carries an *internal* entropic interpretation. In particular, for the BT and FMP cases, they can always be expressed as the difference of generalised entropies. However, only the formalism of [210] provides the right setup for an island to emerge. Following a brief summary of our findings, at the end of section 16, we also point out:

1. The emergence of an entropic hierarchy, similar to the one featuring among the von Neumann algebras recently analysed by Witten et al. within the context of the algebra of observables, [215, 216],
2. The importance of scales throughout our analysis, by making a comparison with recent developments in the literature, specifically Susskind [276], Schlenker and Witten, [277].

13 Euclidean transitions in 2D

In the first section of this Part, we analyse 2D vacuum transitions by means of the Euclidean methods of CDL and BT, with the latter being essentially already known from the original work of [209]. Due to the specifics of the 2D setup, both results can be obtained from the same formalism, namely that of Almheiri and Polchinski, [268]. However, along the way, we encounter significant differences, resulting in different final expressions for the bounces. A physical explanation for this will be outlined throughout the treatment, leaving a more detailed justification to section 16, where, by means of holographic arguments, we will be showing the complementarity of the methods applied.

13.1 Vacuum transitions in 2D without gravity

Let us start considering the simplest case of vacuum transitions in 2D without including gravity. As already argued by the authors of [208], the final amplitude does not depend on the particular profile of the potential. However, their calculation still relies upon the potential exhibiting specific features. In particular, for the purpose of describing the process,

it must have at least two minima, corresponding to the vacua involved in the transition. The simplest starting point is therefore to assume a double well potential, such as the one depicted in figure 48, whose analytic expression can be split into two parts as follows,

$$V(\phi) \stackrel{def.}{=} V_o + V_\varepsilon, \quad (13.1)$$

where V_o is the ordinary degenerate double well potential satisfying the following properties

$$V_o(\phi_+) \equiv V_o(\phi_-) \quad , \quad \left. \frac{dV_o}{d\phi} \right|_{\phi_\pm} = 0, \quad (13.2)$$

with an additional symmetry breaking term proportional to the energy difference between the minima,

$$\varepsilon \stackrel{def.}{=} V(\phi_+) - V(\phi_-) \equiv \Lambda_o - \Lambda_I. \quad (13.3)$$

The interpolation between the two minima is mediated by the scalar field, as depicted in figure 48, since the latter is characterised by a nontrivial spatial profile




Figure 48. The transition between the minima of the potential is independent of the specific profile of V , and is only a function of $\Lambda_{I,o}$ and the tension of the wall.

According to the Euclidean procedure, the decay rate, Γ , is defined from the bounce, B , as follows

$$\Gamma \sim e^{-B} \quad , \quad B \stackrel{def.}{=} S_E|_{inst} - S_E|_{bckgr}, \quad (13.4)$$

where S_E denotes the Euclidean action associated to a given theory. Specifically, there are 3 main contributions,

$$B_{tot}^{CDL} = B_{out} + B_{wall} + B_{inside}. \quad (13.5)$$

For a scalar field admitting an $O(2)$ -symmetric instanton solution $ds^2 = d\rho^2 + \rho^2 d\theta^2$, the corresponding Euclidean action reads:

$$S_E = \int dx^2 \left[\frac{1}{2} (\partial_\mu \phi)^2 + V(\phi) \right] = \pi \int d\rho \rho \left[\frac{1}{2} (\phi')^2 + V(\phi) \right]. \quad (13.6)$$

where $' \equiv \frac{d}{d\rho}$. The equation of motion for ϕ , in the thin-wall approximation, reduces to:

$$\phi' \phi'' - V'_o = \left(\frac{1}{2} (\phi')^2 - V_o \right)' = 0, \quad (13.7)$$

which is exactly integrable once having chosen suitable boundary conditions, $\phi(\rho = \infty) = \phi_+$.

$$\phi' = \sqrt{2(V_o(\phi) - V_o(\phi_{\pm}))} \quad \Rightarrow \quad \rho - \bar{\rho} = \int_{\bar{\phi}}^{\phi} \frac{d\phi}{\sqrt{2(V_o(\phi) - V_o(\phi_{\pm}))}} \quad (13.8)$$

with $\bar{\phi} \stackrel{def.}{=} \frac{\phi_+ + \phi_-}{2}$. Equation (13.8) defines the turning point $(\bar{\rho}, \bar{\phi})$. The total bounce is simply given by the interior and the wall

$$B_{tot} = B_{in} + B_{wall} = -\pi \bar{\rho}^2 \varepsilon + 2\pi \bar{\rho} S_1, \quad (13.9)$$

where, making use of (13.8), the tension of the wall is defined as

$$S_1 = \int d\rho [2(V_o(\phi) - V_o(\phi_+))]. \quad (13.10)$$

Equation (13.9) is extremised at $\bar{\rho} = \frac{S_1}{\varepsilon}$, at which the total extremised bounce reads:

$$\boxed{B_{tot}^{extr} = \frac{\pi S_1^2}{\varepsilon}} \quad (13.11)$$

The presence of a single spatial dimension allows for a direct identification of the nucleation process described by (13.11) with the Schwinger pair production, [278], whose bounce reads


$$B_{Schw} = \frac{\pi m^2}{|eE_{on}|} = \pi m \bar{\rho}_{BT}, \quad (13.12)$$

with $\bar{\rho}_{BT} \stackrel{def.}{=} \frac{m}{|eE_{on}|}$ being the value of the turning point extremising the bounce of the corresponding process. Equation (13.12) suggests the following parametric identification:

$$m \leftrightarrow S_1 \quad , \quad \varepsilon \leftrightarrow |eE_{on}|, \quad (13.13)$$

where m is the mass of the particles, whereas $|eE_{on}|$ is the difference in energy provided by the electric field prior and after pair creation, and can therefore be associated to the energy difference between the minima in V .

Figure 49. Brane nucleation in one spatial dimension, with $\Lambda_{\pm} \stackrel{def.}{=} V(\phi_{\pm})$ is analogous to the Schwinger process upon interpreting the brane with tension σ as the particle/antiparticle pair (denoted by the two red dots).



13.2 Coleman-De Luccia from the Almheiri-Polchinski setup

We will now turn to the case with gravity. The Euclidean action for gravity coupled to a scalar field should contain a potential and a kinetic term. However, in 2D gravity, the Ricci scalar needs to be coupled to the dilaton to ensure that the theory is nontrivial, as first formulated by Jackiw and Teitelboim in [271]. This is a first major difference with respect to the 4D treatment, from which a dynamical coupling between ϕ and ρ is expected to arise at the level of the equations of motion.

Light-cone and cartesian coordinates

Given the conformal metric

$$ds^2 = -e^{2\omega} dx^+ dx^- = 4 e^{2\omega} (-dt^2 + dz^2) \quad (13.14)$$

with $x^\pm \stackrel{def.}{=} t \pm z$ and $2 \partial_\pm = \partial_t \pm \partial_z$, the Ricci scalar in 2D is given by the following expression:

$$\begin{aligned} \mathcal{R} &= -4 e^{-2\omega} (\partial_+ \partial_- \omega - 2 \partial_+ \omega \partial_- \omega) \\ &= -2 [\partial_t (e^{-2\omega} \partial_t \omega) - \partial_z (e^{-2\omega} \partial_z \omega)]. \end{aligned}$$

In Lorentzian signature, the 2D action in terms of x^\pm reads

$$S_{2D} = \int d^2x \sqrt{-g} [\phi^2 \mathcal{R} + \lambda \partial_+ \phi \partial_- \phi - U(\phi)]. \quad (13.15)$$

Given the particular nature of the 2D setup, we can proceed in two ways: either removing the kinetic term or performing a Weyl rescaling of the metric and setting $\mathcal{R} = 0$. Both procedures will be explored in turn in the present section.

Removing the kinetic term

The kinetic term can be removed under suitable rescalings

$$\rho \rightarrow \phi^{-\lambda/4} \rho \quad , \quad U \rightarrow \phi^{-\lambda/2} U \stackrel{def.}{=} U_o, \quad (13.16)$$

such that the action resembles that of JT-gravity. Once integrated by parts, equation (13.15) is simply

$$S_{2D} = -8 \left. (\phi^2 \partial_- \omega) \right|_{x^+ = \text{const.}} - \int d^2x [8 (\partial_+ \phi^2) (\partial_- \omega) + U_o e^{2\omega}], \quad (13.17)$$

from which the e.o.m. for ϕ^2 and ω can be extracted

$$4\partial_+ \partial_- \omega = \frac{e^{2\omega}}{2} \partial_{\phi^2} U_o \equiv T_{+-}, \quad 4\partial_- \partial_+ \phi^2 = U_o e^{2\omega} \equiv T_{-+}, \quad (13.18)$$

corresponding to the off-diagonal components of the energy momentum tensor $T_{\mu\nu}$. The remaining components, instead, read:

$$T_{\pm\pm} = 0 \quad \Rightarrow \quad \partial_\pm (e^{-2\omega} \partial_\pm \phi^2) = 0. \quad (13.19)$$

In cartesian coordinates, the Lorentzian action reads:

$$S_{2D} = S_{bdy} - \int dt dz [2\phi^2 \partial_z^2 \omega + U_o e^{2\omega}], \quad (13.20)$$

where $\omega = \omega(z)$, $\phi^2 = \phi^2(z)$. (13.20) corresponds to the starting point for the extension of CDL to 2D, and the constraints (13.19) reduce to [268],

$$\partial_z (e^{-2\omega} \partial_z (\phi^2)) = 0. \quad (13.21)$$

Wick rotation and polar coordinates

Following [208], we turn to Wick-rotated and polar coordinates:

$$ds^2 = e^{2\omega} (dt^2 + dz^2) = \rho^2 (dr^2 + r^2 d\theta^2), \quad (13.22)$$

where,

$$\begin{cases} t = r \cos \theta \\ z = r \sin \theta \end{cases}, \quad r \stackrel{\text{def.}}{=} \sqrt{t^2 + z^2}, \quad \theta \stackrel{\text{def.}}{=} \tan^{-1} \frac{z}{t}. \quad (13.23)$$

The Euclidean action reads:

$$S_E = 4\pi \phi^2 r \omega' \Big|_{r_i}^{r_f} - 2\pi \int dr \left[2 (\phi^2)' r \omega' - r U_o e^{2\omega} \right], \quad (13.24)$$

where the 2π factor comes from the integration over θ and $' \stackrel{\text{def.}}{=} \partial_r$, with the equations of motion (e.o.m.) for ϕ^2 and ρ now being:

$$2 (r \omega')' = -r e^{2\omega} \partial_{\phi^2} U_o, \quad (r(\phi^2)')' = -r e^{2\omega} U_o. \quad (13.25)$$

Notice that the latter is exactly the e.o.m. for the scalar field obtained by CDL in the thin-wall approximation. Indeed, the only term that is missing w.r.t. the full equations is the $\frac{\rho' \phi'}{\rho}$ -term that can be tuned to zero in the thin-wall approximation. Assuming θ -independence of ϕ and ω , (13.21) reads

$$\partial_r \left(\frac{e^{-2\omega}}{r} \partial_r \phi^2 \right) = 0. \quad (13.26)$$

To obtain an explicit expression for (13.5), we need to specify what we mean by the minima of the potential and the wall separating them within the Almheiri-Polchinski setup. We now turn to describing both, one at a time.

$(\phi^2)' = 0$: defining the vacua

For $\phi^2 = \text{const.}$, (13.25) implies $U_o = 0$ and (13.26) is trivially satisfied. Redefining $u \stackrel{\text{def.}}{=} \ln r$, (13.25) becomes

$$2 \ddot{\omega} = -r^2 e^{2\omega} \partial_{\phi^2} U_o = -\partial_{\phi^2} U_o e^{2(\omega+u)}, \quad (13.27)$$

where $\cdot \stackrel{\text{def.}}{=} \partial_u$. Redefining $\omega + u \stackrel{\text{def.}}{=} f$, equation (13.27) becomes

$$2 \ddot{f} = -\partial_{\phi^2} U_o e^{2f}. \quad (13.28)$$

The first equation in (13.25) shows that $\partial_{\phi^2} U_o \stackrel{\text{def.}}{=} 2\Lambda = \text{constant}$. It defines the cosmological constant of the 2D spacetime. We can therefore determine the solutions corresponding to dS and AdS by respectively taking positive or negative values of such constant:

1) $\Lambda > 0$ equation (13.28) is solved by

$$\rho = e^\omega = e^{f-u} = \frac{2a}{\sqrt{2\Lambda} \cosh(au + b) r}. \quad (13.29)$$

2) $\Lambda < 0$ equation (13.28) is solved by

$$\rho = e^\omega = e^{f-u} = \begin{cases} \frac{a}{\sqrt{2\Lambda} (au+b) r} \\ \frac{a}{\sqrt{2\Lambda} \sin(\text{h})(au+b) r} \\ \frac{a}{\sqrt{2\Lambda} \sin(au+b) r}. \end{cases} \quad (13.30)$$

The three cases outlined in 13.30 correspond to the Poincaré patch of AdS₂, global AdS₂ and an AdS₂ black hole, respectively. Figure 50 helps visualising the regions covered by the different coordinate systems. For either sign of the cosmological constant, the radial coordinate can be redefined, such that the line element takes the following form:

$$ds^2 = d\hat{r}^2 + f(\hat{r}) d\theta^2. \quad (13.31)$$

The corresponding redefinitions are

$$r = \left(\frac{1}{K} \tan \left(\frac{\sqrt{2\Lambda} \hat{r}}{2} \right) \right)^{1/a} \quad \text{for } \Lambda > 0, \quad (13.32)$$

and

$$\frac{\ln |a \ln |r| + b|}{\sqrt{2\Lambda}} = \hat{r} \quad , \quad r = \left(\frac{1}{K} \tanh \left(\frac{\sqrt{2\Lambda} \hat{r}}{2} \right) \right)^{1/a} \quad \text{for } \Lambda < 0, \quad (13.33)$$

for the $1/r \ln r$ and $\frac{1}{\sinh}$ cases, respectively.




Figure 50. The picture on the LHS shows the different regions of AdS₂ covered by the global (the whole rectangular strip), Poincaré (on the RHS of the red lines) and BTZ black hole (the shaded blue region). For the purpose of interest to us, we will only be using the latter, given that the first two are associated to the infinite regions of spacetime. The figures in the middle and on the right, show that the BTZ metric effectively covers only a portion of the Poincaré disk.

For $a = 1, b = 0$, (13.29) and (13.30) reduce to:

$$\rho_{\text{dS}_2} = \frac{2}{\sqrt{2\Lambda} (r^2 + 1)} \quad , \quad \rho_{\text{AdS}_2} = \begin{cases} \frac{a}{\sqrt{2\Lambda} \ln |r| r} \\ \frac{2i}{\sqrt{2\Lambda} (r^2 - 1)} \\ \frac{2i}{\sqrt{2\Lambda} (r^{2i} - 1)}. \end{cases} \quad (13.34)$$

Under a suitable coordinate transformation, these results are equivalent to the ones obtained by Almheiri and Polchinski. The first and third solutions in (13.30) are not relevant for our purposes, given that they correspond to the Poincarè patch, and global AdS. In figure 50, they correspond to the region inside the red triangle and the green shaded strip, respectively. Instead, we will restrict to the $\rho \sim \frac{1}{\sinh}$ case, corresponding to the BTZ black hole, [279], (the region outside the black hole is shaded in blue), which is a quotient of the Poincarè patch.

$(\phi^2)' \neq 0$: defining the wall

We now turn to the case in which $(\phi^2)' \neq 0$. Integrating over (13.26) twice, we find, [268]

$$(\phi^2)' = r c_1 e^{2\omega} \quad \Rightarrow \quad \phi_+^2 - \bar{\phi}^2 = \frac{c_1}{2} \int_0^{\bar{r}} dr r e^{2\omega}, \quad (13.35)$$

where $\bar{\phi}^2 \stackrel{def.}{=} \frac{\phi_+^2 + \phi_-^2}{2}$ is the mean value of the field between the two minima. This defines the turning point $(\bar{\phi}^2, \bar{r})$ in analogy with the Euclidean formalism of [208]. Up to now, the constant c_1 is unspecified. However, its importance is central in our treatment, as will be explained later.

The radius of the nucleated bubble is parameterised by the \hat{r} -variable, defined in (13.32) and (13.33) for dS₂ and AdS₂, respectively. Setting $a = 1, b = 0$, it is a monotonic function and $\text{sgn } \hat{r} \equiv \text{sgn } r$. It therefore follows that we can trade the integration over \hat{r} with that over r without loss of generality. From the equations of motion, we know that $U_o = 0$ at any minimum, but different values of ϕ^2 come with a corresponding value of c_1 triggering the flow described by (13.35). Equation (13.35) describes the flow from one value of the dilaton to that at the turning point




Figure 51. The brane in 2D is mapped to a particle/antiparticle pair (blue dots). The angle $\alpha \stackrel{def.}{=} \sqrt{\frac{\Lambda}{2}} \hat{r}$, denotes the location of the brane separating the background (green) from the interior of the bubble (purple). The corresponding value of r is obtained by the intersection shown in the figure, namely from $\tan(\text{h})\alpha$. Note that $\bar{r} = 1$ corresponds to $\alpha = \frac{\pi}{4}$.

$$\phi_+^2 - \phi_-^2 = c_1^+ \int_0^{\bar{r}} dr r e^{2\omega} = \frac{c_1^+}{\Lambda_+} \int_0^{\bar{r}} dr \frac{2r}{(r^2 - 1)^2} = -\frac{c_1^+}{\Lambda_+} \left[\frac{1}{\bar{r}^2 - 1} + 1 \right]. \quad (13.36)$$

Equation (13.36) can also be interpreted as describing the running of the cosmological constant along the RG-flow originating from ϕ_+^2 and parameterised by c_1^+ . Upon integrating over the reverse flow, defined by $c_1^- \equiv -c_1^+$, we get

$$\phi_-^2 - \phi_+^2 = -\frac{c_1^-}{\Lambda_-} \left[\frac{1}{\bar{r}^2 - 1} + 1 \right]. \quad (13.37)$$

Consistency between (13.36) and (13.37) leads to a constraint for the turning point \bar{r} , which in turn places constraints on the parameters of the theory of either vacua. A similar analysis can be carried out for dS and mixed AdS/dS transitions, and the respective constraints arising from this procedure can be summarised as follows

$$\boxed{\begin{aligned} \bar{r}^2 \Big|_{\text{AdS}_2 \rightarrow \text{AdS}_2} &= 1 + \frac{\Lambda_-}{\Lambda_+} \quad , \quad \bar{r}^2 \Big|_{\text{dS}_2 \rightarrow \text{dS}_2} = \frac{\Lambda_+}{\Lambda_-} - 1 \\ \bar{r}^2 \Big|_{\text{AdS}_2 \leftrightarrow \text{dS}_2} &= \frac{1}{2} \left(-\frac{\Lambda_+ + \Lambda_-}{\Lambda_+ - \Lambda_-} \pm \sqrt{\left(\frac{\Lambda_+ + \Lambda_-}{\Lambda_+ - \Lambda_-}\right)^2 + 8} \right). \end{aligned}} \quad (13.38)$$

For $\text{dS}_2 \rightarrow \text{dS}_2$ processes, the constraint $\bar{r} > 0$ requires $\Lambda_+ > \Lambda_-$, implying only duntunelling is allowed, whereas for transitions of the $\text{AdS}_2 \leftrightarrow \text{dS}_2$ kind, only the the + solution ensures \bar{r} is physical.

Minkowski vacua

Before turning to the evaluation of the bounces, we wish to highlight that the system of equations (13.25) and (13.26), together with the following expression valid at local minima of the potential

$$U_o \Big|_{min} = \phi^2 \partial_{\phi^2} U_o + 2c_1 = 0 \quad (13.39)$$

implies that the Minkowski vacuum cannot provide a stable solution originating a nontrivial flow. Indeed, from (13.39), for $\Lambda = 0$, it follows that $c_1 = 0$, implying the flow generated by such vacuum defines a flat direction in the space of metrics and does not end in a vacuum other than Minkowski itself. Because of this, at least for the moment, we will not be analysing any transitions involving such spacetimes, and we will only be looking at the flat limit of the bounces by taking $\lim_{\Lambda_{\pm} \rightarrow 0}$.

The total bounce

The total bounce is the result from the sum of 2 contributions,

$$B_{tot} = B_{wall} + B_{in}, \quad (13.40)$$

given that $B_{out} = 0$. The scalar field is constant in a local minimum of the potential, and therefore $U_o = 0$ due to (13.25). From (13.25), the Euclidean actions for (A)dS₂, read

$$S_E \Big|_{\text{dS}_2} = -2\pi \phi_-^2 \int_0^{\bar{r}} dr r e^{2\omega} \partial_{\phi^2} U_o = 4\pi \phi_-^2 \left[\frac{1}{\bar{r}^2 + 1} - 1 \right], \quad (13.41)$$

$$S_E \Big|_{\text{AdS}_2} = -2\pi \phi_-^2 \int_0^{\bar{r}} dr r e^{2\omega} \partial_{\phi^2} U_o = 4\pi \phi_-^2 \left[\frac{1}{\bar{r}^2 - 1} + 1 \right], \quad (13.42)$$

from which the inner bounces associated to any possible transition between (A)dS₂ spacetimes, can be evaluated. Inside the wall, the bounce reads

$$\begin{aligned} B_{wall} &= 2\pi \int_0^{\bar{r}} dr r e^{2\omega} [-\phi_-^2 \partial_{\phi^2} U_o + U_o(\phi) + \phi_+^2 \partial_{\phi^2} U_o(\phi_+) - U_o(\phi_+)] \\ &= -2\pi \phi_-^2 U_o \Big|_{\phi_+^2}^{\phi_-^2} + \frac{4\pi}{c_1} \int_{\phi_+^2}^{\phi_-^2} d(\phi^2) [U_o(\phi) + \phi_+^2 \Lambda_+] \\ &= \frac{4\pi}{c_1} S_1 + 4\pi (\phi_-^2 - \phi_+^2), \end{aligned}$$

where

$$S_1 \stackrel{def.}{=} \int_{\phi_+^2}^{\phi_-^2} d(\phi^2) U_o(\phi) \quad \text{with} \quad U_o = -\frac{e^{-2\omega}}{r} (r(\phi^2)')', \quad (13.43)$$

where the last expression follows from the equation of motion for ϕ^2 . According to the specific configuration being studied, there are 4 possible cases:

$$B_{wall} = 4\pi \Delta\phi^2 \cdot \begin{cases} \left(2\frac{\phi_-^2}{\phi_+^2} + 1\right) \\ \left(2\frac{\phi_-^2}{\phi_+^2} - 1\right) \\ \left(2\frac{\phi_-^2}{\phi_+^2} + 1\right) \\ \left(2\frac{\phi_-^2}{\phi_+^2} - 1\right) \end{cases} \Rightarrow B_{tot} = 4\pi \Delta\phi^2 \cdot \begin{cases} \left(\frac{\phi_-^2}{\phi_+^2} + 2\right) & [\text{AdS}_2 \rightarrow \text{AdS}_2] \\ \left(3\frac{\phi_-^2}{\phi_+^2} - 2\right) & [\text{dS}_2 \rightarrow \text{dS}_2] \\ \left(3\frac{\phi_-^2}{\phi_+^2} + 2\right) & [\text{AdS}_2 \rightarrow \text{dS}_2] \\ \left(\frac{\phi_-^2}{\phi_+^2} - 2\right) & [\text{dS}_2 \rightarrow \text{AdS}_2] \end{cases} \quad (13.44)$$

where $\Delta\phi^2 \stackrel{def.}{=} \phi_-^2 - \phi_+^2$.

Note that, upon taking $\lim_{\Lambda_{\pm} \rightarrow 0}$, all transition amplitudes vanish. In particular, there is no 2D counterpart for, either, $\text{dS}_4 \rightarrow \text{Mink}_4$ nor $\text{Mink}_4 \rightarrow \text{AdS}_4$ processes, which were originally addressed in [208].

CDL in the thin-wall approximation

In the above derivation of the total bounces, a closed form expression could be found without needing to resort to the thin-wall approximation. It is therefore interesting to see what the result might be for the case in which we were to apply the standard procedure as in [208]. The starting point is still an O(2)-symmetric instanton solution

$$ds^2 = d\xi^2 + \rho^2(\xi)d\theta^2. \quad (13.45)$$

The radial coordinate used in our derivations, r , is related to ξ in (13.45) as follows⁷²

$$\hat{r} = \sqrt{\frac{2}{\Lambda}} \tan^{-1} r \quad \Rightarrow \quad d\hat{r} = \sqrt{\frac{2}{\Lambda}} \frac{dr}{1+r^2} = \sqrt{\frac{2}{\Lambda}} \cos^2 \left(\sqrt{\frac{\Lambda}{2}} \xi \right) dr = d\xi, \quad (13.46)$$

and to the function ρ_{CDL} in (13.45) as

$$\rho_{CDL} = \sqrt{\frac{2}{\Lambda}} \frac{r}{1+r^2}. \quad (13.47)$$

In particular, we will be looking at the case of $dS_2 \rightarrow \text{Mink}_2$ transitions. In order to do so, we need to re-express the Euclidean action

$$S_E = 2\pi \int_0^{\bar{r}} dr [2\phi^2(r\omega')' + rU_o e^{2\omega}], \quad (13.48)$$

in terms of the new variables (\hat{r}, ρ) or (ξ, ρ) . For Minkowski, $\hat{r} \equiv r \equiv \xi \equiv \rho$. The Euclidean action therefore simply vanishes on the inside. For dS , we know that $e^{2\omega} = \frac{2}{\Lambda(r^2+1)^2}$, as previously derived. On the inside, the dilaton is constant, and the Euclidean action reduces to

$$S_{in} = 2\pi\phi^2 r\omega' \quad \text{with} \quad \omega' = \frac{(e^\omega)'}{e^\omega} = -2\rho_{CDL} \sqrt{\frac{\Lambda}{2}}. \quad (13.49)$$

For ease of notation, we will simply write the radial coordinate as ρ . Inverting the last equality in (13.49), we get

$$r = \frac{1 - \sqrt{1 - 2\Lambda\rho^2}}{\sqrt{2\Lambda\rho}} \quad \Rightarrow \quad S_{in} = -2\pi\phi^2 \left(1 - \sqrt{1 - 2\Lambda\bar{\rho}^2}\right). \quad (13.50)$$

The wall's bounce, instead, reads

$$B_{wall} = \pi\bar{\rho} \bar{\phi}^2 S_1 + 4\pi(\phi_-^2 - \phi_+^2), \quad (13.51)$$

where we made use of the thin-wall approximation for extracting $\bar{\rho}$ from the integral and have also defined the tension of the wall in terms of the rescaled potential in the Einstein frame, U_o^{new}

$$S_1 = 2 \int d\xi U_o^{new} = 2 \int d\xi [\phi^{-2}U_o - \partial_{\phi^2}U_o]. \quad (13.52)$$

For the case of a $dS_2 \rightarrow \text{Mink}_2$ transition, the total bounce therefore reads

$$B_{tot} = -2\pi\phi_+^2 \left(1 + \sqrt{1 - 2\Lambda\bar{\rho}^2}\right) + \pi\bar{\rho} \bar{\phi}^2 S_1, \quad (13.53)$$

⁷²Notice that for AdS the same arguments follow, the only difference being that trigonometric functions turn into hyperbolic functions due to the negative sign of Λ .

and it is extremised at

$$\bar{\rho} = \frac{S_1 \bar{\phi}^2}{4\Lambda \phi_+^2 \sqrt{1 + \frac{S_1^2 \bar{\phi}^4}{8\Lambda \phi_+^4}}}. \quad (13.54)$$

$$\Rightarrow \boxed{B_{tot}^{\text{dS}_2 \rightarrow \text{Mink}_2} \Big|_{extr} = 2\pi \phi_+^2 \sqrt{1 + \frac{S_1^2 \bar{\phi}^4}{8\Lambda \phi_+^4}} \left(1 - \frac{3}{1 + \frac{S_1^2 \bar{\phi}^4}{8\Lambda \phi_+^4}} \right)}. \quad (13.55)$$

In a similar fashion, for the generic (A)dS₂ → (A)dS₂ case, the total bounce reads

$$\boxed{B_{tot}^{(\text{A})\text{dS}_2 \rightarrow (\text{A})\text{dS}_2} = 2\pi \phi_-^2 \left(1 + \sqrt{1 - 2\Lambda_- \bar{\rho}^2} \right) - 2\pi \phi_+^2 \left(1 + \sqrt{1 - 2\Lambda_+ \bar{\rho}^2} \right) + \pi \bar{\rho} \bar{\phi}^2 S_1}. \quad (13.56)$$

13.3 Brown-Teitelboim in 2D from Almheiri-Polchinski

We now turn to the alternative path for determining the bounce describing 2D transitions, namely by removing the Ricci scalar and restoring the kinetic term in the AP action:

$$S_{AP} = \int dx^2 [\phi^2 \mathcal{R} + U_o] e^{2\omega}. \quad (13.57)$$

The ϕ'^2 -term can be restored by reabsorbing ϕ^2 in the metric. In particular, under the rescalings $\rho \rightarrow \phi \rho$ and $U_o \rightarrow \phi^{-2} U_o \stackrel{def.}{=} 16\Lambda_{BT}$, (13.57) turns into

$$S_{AP} = 8\pi \int dz \left[-u'^2 + 4\Lambda_{BT} e^{2f} \right], \quad (13.58)$$

with $f \stackrel{def.}{=} \ln \phi + \omega \stackrel{def.}{=} u + \omega$. Before delving into further calculations, some important remarks are in order. The equations of motion undoubtably play a key role within the CDL formalism, as can be appreciated by direct inspection of their original work in 4D, [208]. When performing the calculation for transition amplitudes within the AP setup (once having removed the kinetic term for the scalar field), the crucial relation that allowed us to circumvent the need to resort to the thin-wall approximation (upon which [208] rely) was the change of variables resulting from requiring vanishing diagonal components for the energy momentum tensor, (13.19). As previously argued, in polar coordinates, they reduce to

$$\partial_r \left(\frac{e^{-2\omega}}{r} \partial_r \phi^2 \right) = 0 \quad \Rightarrow \quad \partial_z (u \partial_z u) = (uu')' = 0, \quad (13.59)$$

where $z \stackrel{def.}{=} \ln r$. At the same time, the term on the RHS in (13.59) also emerges upon integrating by parts the kinetic term in (13.58). Correspondingly, we are led to consider the following quantity, $M_{ADM} \stackrel{def.}{=} uu' \Big|_{bdy}$, which can be identified with the ADM mass of

the spacetime. When removing the kinetic term in \mathcal{L}_{AP} , the dilaton is constant in each vacuum solution, therefore M_{ADM} is trivially vanishing to start with, as follows from the equation of motion⁷³ On the other hand, when performing the Weyl transformation of the metric leading to (13.58), this is no longer true. Indeed, when using (13.58) to describe transitions, $\Delta M_{ADM} \stackrel{def.}{=} uu' \Big|_{bck}^{inst} \neq 0$. We thereby need to add such boundary term by hand to ensure energy conservation throughout the nucleation process as a whole.

From the considerations we have just made, (13.58) should therefore be upgraded in the following way

$$\begin{aligned} S_{BT}^{grav} &= S_{AP} + 16\pi uu' \Big|_{bdy} \\ &= 8\pi \int dz \left[-u'^2 + 4\Lambda_{BT} e^{2f} \right] + 16\pi uu' \Big|_{bdy}, \end{aligned} \quad (13.60)$$

which is exactly equivalent to the BT action describing brane nucleation in 2D in presence of gravity, in which case the total bounce in conformal gauge (with metric function parameterised by e^φ) reads, [209],

$$B_{TOT} = 2\pi m\bar{\rho} + \frac{\pi}{2k} \int_{\bar{r}_o}^{\infty} dr r \left[-(\varphi_{,r})^2 \Big|_{bck}^{inst} + 4\Lambda_I e^{\varphi_{inst}} - 4\Lambda_o e^{\varphi_{bck}} \right] + \frac{\pi}{k} r \varphi_{,r} \varphi \Big|_{bck}^{inst}, \quad (13.61)$$

where the *effective* cosmological constants are defined as follows

$$\Lambda_{o,I} \stackrel{def.}{=} \lambda + \frac{1}{2}k E_{o,I}^2. \quad (13.62)$$

In BT's notation, k is proportional to G_N and is kept fixed throughout their treatment. The dynamics is thereby carried by the change in the electric field, as follows from (13.77). On the other hand, on the case of AP, as discussed earlier on in the section, we can think of k as being the running coupling ensuring the interpolation in between the two vacua parameterised by different values of the dilatonic field. Indeed, in this sense we argued earlier on that transitions in AP are equivalent to RG-flows.

The equations of motion for ϕ is:

$$-u'' = 4\Lambda_{BT} e^{2\omega+2u}. \quad (13.63)$$

It admits a solution of the kind

$$u(z) = -\ln \left| \cos(h) \left(-2\sqrt{\Lambda_{BT}} e^{2\omega} z + \text{const.} \right) \right|. \quad (13.64)$$

Fixing $\omega \equiv 0$ and $\text{const.} \equiv 0$, the profile for the scalar field reads:

$$\phi(z) = \frac{1}{\cos(h) \left(-2\sqrt{\Lambda_{BT}} z \right)}, \quad (13.65)$$

⁷³As we shall see in greater detail in section 16, this is consistent with the fact that such types of transitions take place in absence of gravity.

which is either a trigonometric or hyperbolic function, according to the sign of Λ_{BT} . Upon integrating the first term in (13.60) by parts, and using (13.63), we get

$$S = 8\pi \int_{\bar{z}}^{z_h} dz u'' (2u - 1), \quad (13.66)$$

where \bar{z} and z_h denote the location of the turning point and the horizon, respectively. Specifically, for AdS, the latter is simply $z_h \equiv 0$, whereas for dS it is finite. For the instanton to exist, $\bar{z} > z_h$ is needed, and (13.66) therefore reads:

$$S_{tot} = -16\pi \left[\sqrt{\Lambda_{BT}} \tan(h) \left(-2\sqrt{\Lambda_{BT}} z \right) \ln \left| \cos(h)^2 \left(-2\sqrt{\Lambda_{BT}} z \right) \right| + 4\Lambda_{BT} z + 3\sqrt{\Lambda_{BT}} \tan(h) \left(-2\sqrt{\Lambda_{BT}} z \right) \right] \Big|_{\bar{z}}^{z_h}. \quad (13.67)$$

Equation (13.67) holds for, both, the instanton and background solutions, the only difference being the value of Λ_{BT} . From now on it is convenient to analyse the AdS and dS cases separately. 1) AdS₂ → AdS₂

In this case $z_h \equiv 0$. Redefining

$$\sqrt{\Lambda_{\pm}} \sqrt{1 \pm \phi_{\pm}^2} \stackrel{def.}{=} \phi_{\pm}^{new} \Rightarrow \phi_{\pm}^2 = \pm \left(1 - \frac{\phi_{\pm}^{new 2}}{\Lambda_{\pm}} \right), \quad (13.68)$$

ensures that now ϕ_{tp} is the same on either side at \bar{z} , hence

$$B_{in}^{AdS_2 \rightarrow AdS_2} = 16\pi \left[\phi_{tp}^{new} \ln \left| \frac{1 - \frac{\phi_{tp,-}^{new 2}}{\Lambda_-}}{1 - \frac{\phi_{tp,+}^{new 2}}{\Lambda_+}} \right| + 3(\Lambda_- - \Lambda_+) \bar{z} \right]. \quad (13.69)$$

Which follows from $\phi_{tp,-}^{new} \equiv \phi_{tp,+}^{new}$, by means of (13.68), whereas the wall's bounce reads:

$$B_{wall} = 8\pi \int dz \left[-u'^2 + 4\Lambda_{BT} e^{u2} \right] \stackrel{def.}{=} 32\pi \sigma = 32\pi m \bar{z}, \quad (13.70)$$

where σ denotes the tension of the brane and $m \stackrel{def.}{=} \Lambda_+ - \Lambda_-$. The total bounce therefore reads:

$$B_{tot}^{AdS_2 \rightarrow AdS_2} = 16\pi \left[\phi_{tp}^{new} \ln \left| \frac{1 - \frac{\phi_{tp,-}^{new 2}}{\Lambda_-}}{1 - \frac{\phi_{tp,+}^{new 2}}{\Lambda_+}} \right| - \frac{1}{3} m \bar{z} \right]. \quad (13.71)$$

2) dS₂ → dS₂ Similar considerations can be performed for the case of dS. The main difference is that now the horizons contribute with nontrivial terms to the actions. In particular, being $z_h \neq 0$, we get

$$\boxed{
\begin{aligned}
B_{tot}^{\text{dS}_2 \rightarrow \text{dS}_2} = 16\pi & \left[\phi_{tp}^{new} \ln \left| \frac{1 - \frac{\phi_{tp,-}^{new 2}}{\Lambda_-}}{1 - \frac{\phi_{tp,+}^{new 2}}{\Lambda_+}} \right| + 3 \left(\Lambda_- - \Lambda_+ + \frac{m}{6} \right) \bar{z} + \right. \\
& \left. + \phi_{h,-}^{new} \left(2 - 3\sqrt{1 - \frac{\phi_{h,-}^{new 2}}{\Lambda_-}} \right) - \phi_{h,+}^{new} \left(2 - 3\sqrt{1 - \frac{\phi_{h,+}^{new 2}}{\Lambda_-}} \right) \right].
\end{aligned}
} \tag{13.72}$$

Once more, the terms linear in \bar{z} provide an energy conservation relation and are therefore vanishing. The terms featuring in the last line, instead, provide the contributions from the horizons. The results (13.71), (13.72) agree with the expression of the total bounce for type-1 instantons in [209]

$$\boxed{
B = 2\pi m \bar{\rho} - \frac{4\pi}{k} \ln \frac{r_i}{r_o} + \frac{2\pi \bar{\rho}}{k} (\Lambda_i r_i - \Lambda_o r_o) .
} \tag{13.73}$$

prior to extremising with respect to $\bar{\rho}$. In (13.73), we made use of the same notation as can be found in the literature, with

$$\bar{r}_{I,o} \stackrel{\text{def.}}{=} \frac{2}{\Lambda_{I,o} \bar{\rho}} \left[1 - \sqrt{1 - \Lambda_{I,o} \bar{\rho}} \right]. \tag{13.74}$$

Importantly, the last 2 terms in (13.73) arise only for the $\text{dS}_2 \rightarrow \text{dS}_2$ case, and are absent for $\text{AdS}_2 \rightarrow \text{AdS}_2$, consistently with the results obtained in (13.72) and (13.71), respectively. Compatibility with our results follows from the following identifications

$$\bar{z} \longleftrightarrow \bar{\rho} \quad , \quad \phi_{tp}^{new} \stackrel{\text{def.}}{=} \frac{1}{k} = \frac{E_{o,I}^2}{2(\Lambda_{o,I} - \lambda)} \quad , \quad \frac{\phi_{\pm}^{new 2}}{\Lambda_{\pm}} \stackrel{\text{def.}}{=} 1 + \frac{\bar{r}_{i,o}^2 \Lambda_{\pm}}{4}. \tag{13.75}$$

Key points and remarks

- Unlike the ordinary Schwinger process, in presence of gravity, the Euclidean action needs to be dressed with a boundary term corresponding to the ADM mass

$$S_{BT}^{grav} = S_{AP} + 16\pi u u' \Big|_{bdy} . \tag{13.76}$$

- In BT, the dynamical process of brane nucleation is encoded in the change in the electric field, while keeping k (i.e. G_N) fixed, as follows from (13.77). On the other hand, on the case of AP, k is related to the running coupling, ϕ , ensuring the interpolation in between the two vacua parameterised by different values of the dilatonic field.
- Up-tunneling falls into the forbidden region of parameter space of the type-1 instanton, since, this configuration fails to satisfy the energy conservation relation associated to the brane nucleation process.

- Interestingly, for either sign of the cosmological constant, the flat limit cannot be taken. This follows from the fact that such processes correspond to topology change, and would thereby fall within a different instanton type. The reason for this can be traced back to the extremality of the spacetimes involved, due to the definition of effective cosmological constants,

$$\Lambda_{o,I} \stackrel{def.}{=} \lambda + \frac{1}{2}k E_{o,I}^2. \quad (13.77)$$

- As already highlighted in [209], there are two different ways of calculating the bounce action. Either substituting the singular solutions to the field equations in the Euclidean action, or, alternatively, integrating over the 2 bulk spacetimes separately, in which case the δ -function in the fields would be replaced by the Gibbons-Hawking boundary terms, with the membrane providing the natural boundary for both spacetimes. The 2^{nd} procedure agrees with the first upon extremising the bounce with respect to the membrane size, $\bar{\rho}$. According to [209], this is preferable, since it ensures that the extremisation of the bounce on the classical instanton solution is not an artefact of the formalism⁷⁴.

14 Lorentzian transitions in 2D

As proved in an earlier work, [7], vacuum transitions in 4D can be described in three equivalent ways, namely by means of the CDL, BT and FMP formalisms. The importance of their agreement in the well known case of dS_4 spacetime, is what led us to explore to what extent the obstruction to other types of transitions may be overcome. In particular, the role played by black holes in [7], inspired by the original work of FMP, which had no counterpart in the BT and CDL methods, turned out to be crucial for enabling to define transitions involving Minkowski spacetimes.

Given these motivations, we now turn to extending the Hamiltonian method of Fischler, Morgan and Polchinski (FMP) [210] to its 2D counterpart, namely Jackiw-Teitelboim (JT)-gravity [271, 272], whose action, up to boundary terms, reads

$$S_{JT} = \frac{\phi_o}{2\kappa^2} \int d^2x \sqrt{-g} \mathcal{R} + \frac{1}{2\kappa^2} \int d^2x \sqrt{-g} \phi (\mathcal{R} - 2\Lambda), \quad (14.1)$$

with ϕ_o being a constant. The coupling of the dilaton ϕ to the Ricci scalar ensures gravity is nontrivial. We can think of (14.1) as being the 2D version of the Einstein-Hilbert action with a cosmological constant describing gravity in arbitrary dimension. ϕ plays the role of a Lagrange multiplier and ensures curvature is fixed. From a Field Theory point of view, the expression (14.1) is that of a renormalised theory, with the dilaton accounting for the renormalisation of Newton's constant.

The setup consists of pure gravity with spherical symmetry and a wall (the bubble's surface) at $z = \hat{z}$, separating two spacetimes with two different cosmological constants

⁷⁴In section 14 we will argue that the starting point in the Hamiltonian formalism of FMP is precisely the first of the two methods outlined in [209].

Λ_+, Λ_- , similarly to Brown and Teitelboim [209]. Given the above considerations, our proposed initial Lagrangian density, thereby takes the general form:

$$\begin{aligned} \mathcal{L} = & \sqrt{-g}\phi [\mathcal{R} - 2\Lambda_+\Theta(z - \hat{z}) - 2\Lambda_-\Theta(\hat{z} - z)] - \delta(z - \hat{z})\sigma\sqrt{(N^t)^2 + L^2(N^z + \dot{\hat{z}})^2} + \\ & + \sqrt{-g}[\phi_o\chi_{\mathcal{M}} - \mathcal{B}_+\Theta(z - \hat{z}) - \mathcal{B}_-\Theta(\hat{z} - z)], \end{aligned}$$

i.e. the action for JT-gravity coupled to matter, with the latter being provided by the brane term, the two cosmological constants Λ_+, Λ_- and two additional constants⁷⁵, \mathcal{B}_\pm . Following FMP we start with the metric:

$$g_{\mu\nu} = \begin{pmatrix} -(N^t)^2 + (N^z)^2 & LN^z \\ LN^z & L^2 \end{pmatrix}. \quad (14.2)$$

14.1 Actions with vanishing constant energies

First, we will be considering the $\mathcal{B}_\pm = 0$ case. The $\mathcal{B} \neq 0$ case will be dealt with in section 15. Substituting (14.2) in (14.2) with $\mathcal{B}_\pm = 0$ and integrating by parts, we get⁷⁶

$$\begin{aligned} \mathcal{L} = & \mathcal{L}_{BT_s} + 2\phi \left(\frac{N^{tt}}{L} \right)' - \frac{2\dot{\phi}\dot{L}}{N^t} - \frac{2\phi'N^z(N^{zt} - \dot{L})}{LN^t} + \\ & - 2\Lambda_+\Theta(z - \hat{z}) - 2\Lambda_-\Theta(\hat{z} - z) - \delta(z - \hat{z})\sigma\sqrt{(N^t)^2 + L^2(N^z + \dot{\hat{z}})^2} \\ = & \mathcal{L}_{BT_s} + \pi_L\dot{L} + \pi_\phi\dot{\phi} - N^t\mathcal{H}_t - N^z\mathcal{H}_z, \end{aligned}$$

where the total derivative terms read

$$\mathcal{L}_{BT_s} = -2 \left(\frac{\phi N^{tt}}{L} \right)' + \left(\frac{2\phi\dot{L}}{N^t} \right)' + \left(\frac{2N^z(N^{zt} - \dot{L})}{LN^t} \right)'. \quad (14.3)$$

The conjugate momenta are:

$$\pi_L = 2 \left(\frac{\phi'N^z}{LN^t} - \frac{\dot{\phi}}{N^t} \right), \quad \pi_\phi = \frac{2(N^{zt} - \dot{L})}{N^t}, \quad \pi_{\hat{z}} = -\frac{\sigma L^2(N^z + \dot{\hat{z}})\delta(z - \hat{z})}{\sqrt{(N^t)^2 + L^2(N^z + \dot{\hat{z}})^2}}, \quad (14.4)$$

The Hamiltonian and total momentum are:

⁷⁵These act as deformation parameters in the 2D theory, contributing to the Hamiltonian of the system in the form of constant energy terms. As we shall see in section 15, they play the role of black hole masses, and will be subject to constraints specified in due course.

⁷⁶Notice that in the calculations we are omitting the topological term but we will add it back to the final expression for the action.

$$\begin{aligned}
-\mathcal{H}_t &= \frac{\delta \mathcal{L}}{\delta N^t} = -2 \left(\frac{\phi'}{L} \right)' - \frac{2\dot{\phi}}{N^t} (N^{z'} - \dot{L}) + \frac{2\phi' N^z (N^{z'} - \dot{L})}{N^{t2} L} + \\
&\quad - 2\phi L [\Lambda_+ \Theta(z - \hat{z}) + \Lambda_- \Theta(\hat{z} - z)] - \frac{\sigma N^t \delta(z - \hat{z})}{\sqrt{(N^t)^2 + L^2 (N^z + \dot{\hat{z}})^2}}, \\
-\mathcal{H}_z &= \pi'_L - \phi' \frac{\pi_\phi}{L} - \delta(z - \hat{z}) \pi_{\hat{z}}.
\end{aligned} \tag{14.5}$$

Away from the wall, the momentum constraint reads

$$\pi'_L = \phi' \frac{\pi_\phi}{L}, \tag{14.6}$$

in terms of which \mathcal{H}_t can be re-expressed⁷⁷, thereby leading to the following expression

$$\boxed{(\pi_L^2) \Big|_{\hat{z} \pm \epsilon} = 4 \left[\left(\frac{\phi'}{L} \right)^2 \Big|_{\hat{z} \pm \epsilon} + \mathcal{C}_\pm + \Lambda_\pm \phi^2 \right]}, \tag{14.7}$$

with \mathcal{C}_\pm denoting an integration constants. A key feature emerging from this analysis is that (14.7) is a dimensionful relation. This follows from the fact that $[\phi] = 0$ in 2D, hence $[\mathcal{C}_\pm] = 2$. The junction conditions can be extracted upon integrating (14.5) across the brane. In the rest frame of the latter, we get

$$\begin{cases} -\int_{\hat{z}-\epsilon}^{\hat{z}+\epsilon} dz \mathcal{H}_z = \pi_L \Big|_{\hat{z}-\epsilon}^{\hat{z}+\epsilon} - \hat{\pi}_{\hat{z}} = 0 \\ -\int_{\hat{z}-\epsilon}^{\hat{z}+\epsilon} dz \mathcal{H}_t = -2 \frac{\phi'}{L} \Big|_{\hat{z}-\epsilon}^{\hat{z}+\epsilon} - \frac{\sigma N^t}{\sqrt{(N^t)^2 + L^2 (N^z + \dot{\hat{z}})^2}} = 0 \end{cases} \xrightarrow{N^t=1, N^z=0} \begin{cases} \frac{\phi'}{L} \Big|_{\hat{z}_-}^{\hat{z}_+} = -\frac{\sigma}{2} \\ \pi_L \Big|_{\hat{z}_-}^{\hat{z}_+} = 0, \end{cases}$$

where the extrinsic curvatures on either side read

$$\boxed{\frac{\phi'}{L} \Big|_{\hat{z}_\pm} = \frac{(\Lambda_+ - \Lambda_-) \phi^2}{\sigma} \mp \frac{\sigma}{4} + \frac{\mathcal{C}_+ - \mathcal{C}_-}{\sigma}}. \tag{14.8}$$

Equating the new expression for π_L with the one obtained from the variation of \mathcal{L} , we get

$$\pi_L \Big|_{\hat{z}_+} = -2\dot{\phi}. \tag{14.9}$$

The constants $\mathcal{C}_+, \mathcal{C}_-$ can be absorbed by a rescaling of ϕ and π_L . We will partially use this freedom to choose $\mathcal{C}_+ = \mathcal{C}_-$. Squaring both sides and redefining, where the b -subindex

⁷⁷Imposing the Hamiltonian constraint is equivalent to selecting a specific energy for the configuration described by the Lagrangian of the theory. Because of this, the Lorentzian approach of FMP is naturally providing a microcanonical description of vacuum decays. The improvement of their method with respect to those outlined in section 13 is comparable to the one in [280], where the authors provide a microcanonical generalisation of the canonical treatment analysed in [281].

denotes the value of the dilaton at the boundary. $\Phi_b = \phi_b/\sqrt{|\mathcal{C}|}$ and solving this quadratic equation with respect to $\dot{\Phi}_b^2$ leads to the energy conservation relation

$$\boxed{\dot{\Phi}_b^2 + V_{eff} = -1}, \quad (14.10)$$

with effective quadratic potential defined as

$$\boxed{V_{eff} = -\frac{1}{4} \left[\left(\frac{\Lambda_+ - \Lambda_- - \frac{\kappa^2}{4}}{\kappa\sqrt{|\mathcal{C}|}} \right)^2 + \Lambda_+ \right] \Phi_b^2 - \frac{3}{4}}, \quad (14.11)$$

where we have used that $\sigma = \kappa\phi_b$. The value of the dilaton at the turning point is found by setting $\dot{\Phi}_b = 0$, i.e. when the classical motion is reversed, and solving for the unique turning point

$$V_{eff} = -1 \quad \Rightarrow \quad \Phi_o = \left[\left(\frac{\Lambda_+ - \Lambda_- - \frac{\kappa^2}{4}}{\kappa\sqrt{|\mathcal{C}|}} \right)^2 + \Lambda_+ \right]^{-\frac{1}{2}}. \quad (14.12)$$

Notice the similarity with the 4D case from FMP where the dilaton is playing the role of the compactification radius.

Metric and dilaton profiles

We now determine the solutions to the equations of motion arising from the variational principle. These are needed for evaluating the bulk and boundary actions, addressed next. In conformal gauge, the constraint $\mathcal{R} = 2\Lambda$ can be solved in order to recover an expression for L in terms of the z -coordinate. In the Lagrangian density we started from, $\phi(\mathcal{R} - 2\Lambda_{\pm})$ plays the role of a perturbation around the background action which, for the 2D case is purely topological. For the given gauge choice, the metric and dilatonic profile can be expressed as

$$ds^2 = \frac{1}{z^2} (-dt^2 + dz^2) \quad , \quad \text{with} \quad L = \frac{1}{z} \quad , \quad \phi = \text{const}. \quad (14.13)$$

The e.o.m. arising from \mathcal{L}_{JT} , i.e.

$$\phi \left(-\frac{L''}{L} + \frac{2L'^2}{L^2} \right) = 2\phi\Lambda_{\pm}, \quad \phi'' = 2\phi\Lambda_{\pm}L^2, \quad (14.14)$$

are solved by

$$\boxed{L = \frac{c}{\sinh(bz)} \quad , \quad \phi = 2a \coth(bz)}, \quad \text{with} \quad \Lambda_{\pm} \stackrel{\text{def.}}{=} \frac{b^2}{c^2}. \quad (14.15)$$

For the case of pure AdS₂ in Poincarè coordinates, (14.13), the metric diverges at the location of the conformal boundary, placed at $z = 0$. The latter can be mapped to the horizon at $\rho = \gamma$ via the following coordinate transformation

$$\boxed{z \stackrel{\text{def.}}{=} \frac{1}{b} \coth^{-1} \left(\frac{\rho}{\gamma} \right)} \Rightarrow dz = \frac{\gamma}{b} \frac{d\rho}{\gamma^2 - \rho^2}, \quad (14.16)$$

whereas the dilatonic profile becomes linearly-dependent on the redefined spatial coordinate

$$\boxed{\phi = 2a \frac{\rho}{\gamma}}. \quad (14.17)$$

Notice that, going from ρ to z in (14.16), is analogous to the analytic extension of the metric beyond the horizon. In particular, such transformation takes place by introducing an additional scale, μ . Upon substituting (14.16) in the original line element (14.13), we get

$$ds^2 = \frac{\rho^2 - \gamma^2}{\gamma^2} \left(-dt^2 + \frac{\gamma^2}{b^2} \frac{d\rho^2}{(\rho^2 - \gamma^2)^2} \right). \quad (14.18)$$

The main feature of this result is the explicit emergence of a horizon at $\rho = \gamma$ if $\gamma^2 > 0$. For the $\gamma^2 < 0$ case, instead, there is no horizon. Rewriting (14.18) in terms of ϕ , with $\gamma = 1/b$, we get

$$\boxed{ds^2 = \left(\frac{\phi^2}{4a^2} - 1 \right) \left(-dt^2 + \frac{1}{4a^2 b^2} \frac{d\phi^2}{\left(\frac{\phi^2}{4a^2} - 1 \right)^2} \right)}. \quad (14.19)$$

The vanishing of the conjugate momentum with respect to L imply:

$$\boxed{\phi^2 = -\frac{\mathcal{C}}{\Lambda_{\pm}} \stackrel{\text{def.}}{=} \mu_{\pm}^2}, \quad (14.20)$$

where the system is understood to be evaluated at the turning point in the $N^z = 0$ gauge. Equation (14.20) enables to identify the parameter in (14.19) $2a = \sqrt{-\mathcal{C}/\Lambda_{\pm}} = \mu_{\pm}$ as the cosmological horizon in a dS₂ metric. Furthermore, from (14.15), we get:

$$\phi' = 2ab(1 - \coth^2(bz)) = 2ab - \frac{b}{2a}\phi^2 = \frac{b}{2a}(4a^2 - \phi^2), \quad (14.21)$$

from which it can be appreciated that the change in sign of the extrinsic curvature (ϕ') takes place when crossing the cosmological horizon at $\phi = 2a$.

Bulk and boundary actions

In the Hamiltonian formalism, the transition rate is defined in terms of the extremised total action including, both, bulk and wall terms

$$\Gamma(A \rightarrow A \oplus W \oplus B) = \exp[-S_{tot} + S_{bckgr}], \quad (14.22)$$

where A, B denote the spacetimes involved separated by the brane W , and

$$S_{tot}^{A/B} = S_{top} + S_{bulk} + S_{brane} \quad , \quad S_{bckgr} \stackrel{def.}{=} \frac{\phi_{bckgr}}{16\pi G} \chi_{\mathcal{M}}. \quad (14.23)$$

The topological term S_{top} cancels with the background contribution when evaluating the transition rate. The extremised action is obtained by integrating over the Hamilton-Jacobi equations. By imposing the secondary constraints derived above, the variation of the bulk action therefore reads

$$\delta S_{bulk} = \int dz [\delta L(z)\pi_L + \delta\phi(z)\pi_\phi] = 2 \int dz \delta L(z)\pi_L \quad (14.24)$$

$$\begin{aligned} \Rightarrow S_{bulk}^{(+)} &= \frac{2\eta}{G} \int dz \int dL \sqrt{\left(\frac{\phi'}{L}\right)^2 + \mathcal{C} + \Lambda_+ \phi^2} \\ &= \frac{2\eta}{G} \int_0^{\hat{z}} dz \left[L \sqrt{\left(\frac{\phi'}{L}\right)^2 + \mathcal{C} + \Lambda_+ \phi^2} - \phi' \cosh^{-1} \left(\frac{\phi'}{L\sqrt{-\mathcal{C} - \Lambda_+ \phi^2}} \right) \right]. \end{aligned} \quad (14.25)$$

Similarly, for the other vacuum,

$$S_{bulk}^{(-)} = \frac{2\eta}{G} \int_{\hat{z}}^{\infty} dz \left[L \sqrt{\left(\frac{\phi'}{L}\right)^2 + \mathcal{C} + \Lambda_- \phi^2} - \phi' \cosh^{-1} \left(\frac{\phi'}{L\sqrt{-\mathcal{C} - \Lambda_- \phi^2}} \right) \right]. \quad (14.26)$$

The bulk action is extremised when the integrand featuring in the first term of (14.25) and (14.26) vanishes, therefore leaving with the corresponding contribution from the value of the dilaton at the turning point and at the horizons characterising spacetimes being involved. For such values, the argument of \cosh^{-1} is imaginary, therefore, we can rewrite it as \cos^{-1} , which provides an overall π factor. The extremes of integration for, both, (14.25) and (14.26) are dictated by the relative position of the innermost horizon in a given spacetime with respect to the turning point ϕ_o , whose expression was explicitly derived in the previous subsection, cf. eq. (14.12). From equations (14.21) we therefore get the following contributions

$$S_{bulk}^{(-)} = \frac{2\pi\eta}{G} \int_{z_{tp}}^{z_{h,-}} dz \phi' = \frac{2\pi\eta}{G} \int_{\mu_-}^{\phi_o} \epsilon_- d\phi = \frac{2\pi\eta}{G} \left[\phi_o \Theta \left(-\frac{\phi'}{L} \Big|_{-} \right) - \mu_- \right], \quad (14.27)$$

with $\Theta \left(-\frac{\phi'}{L} \Big|_{-} \right)$ being the Θ -function that is non-vanishing only if the extrinsic curvature on the inner side of the brane is negative, thereby indicating that the turning point is placed beyond the horizon. The same argument follows for the bulk integral over the outer vacuum. For the case involving pure AdS, instead, due to the absence of horizons, only the ϕ_o -terms will be contributing as relevant extremes of integration. Furthermore, precisely because of this, the extrinsic curvature on either side will also preserve the sign over the entire spacetime, thereby leading to the mutual cancellation of the Θ -terms. Combining

the claims just made, the corresponding result for mixed AdS/dS configurations follows. For the 3 cases of interest, the total bulk action therefore reads

$$\begin{aligned}
S_{bulk}^{dS \rightarrow dS} &= S_{bulk}^{(+)} + S_{bulk}^{(-)} = \frac{2\pi\eta}{G} \left[\phi_o \Theta \left(-\frac{\phi'}{L} \Big|_{-} \right) - \mu_- - \phi_o \Theta \left(-\frac{\phi'}{L} \Big|_{+} \right) + \mu_+ \right] \\
S_{bulk}^{AdS \rightarrow AdS} &= S_{bulk}^{(+)} + S_{bulk}^{(-)} = 0 \\
S_{bulk}^{dS \rightarrow AdS} &= S_{bulk}^{(+)} + S_{bulk}^{(-)} = \frac{2\pi\eta}{G} \left[\phi_o \Theta \left(-\frac{\phi'}{L} \Big|_{-} \right) - \phi_o \Theta \left(-\frac{\phi'}{L} \Big|_{+} \right) + \mu_+ \right].
\end{aligned}
\tag{14.28}$$

The brane action is obtained by considering the variation of (14.24) at the location of the wall $z = \hat{z}$. From the junction conditions, $\hat{\pi}_L \Big|_{+} = \hat{\pi}_L \Big|_{-}$, it follows that only the second term in, both, (14.25) and (14.26) contributes nontrivially

$$\begin{aligned}
S_{brane} &= \frac{2}{G} \int_0^{\phi_o} d\phi_b \left[\cosh^{-1} \left(\frac{\phi'_b(\hat{z} - \epsilon)}{L\sqrt{-\mathcal{C} - \Lambda_- \phi_b^2}} \right) - \cosh^{-1} \left(\frac{\phi'_b(\hat{z} + \epsilon)}{L\sqrt{-\mathcal{C} - \Lambda_+ \phi_b^2}} \right) \right] \\
&= \frac{2}{G} \int_0^{\phi_o} d\phi_b \left[\cosh^{-1} \left(\frac{A_1 \phi_b}{\sqrt{1 - \frac{\Lambda_-}{|\mathcal{C}|} \phi_b^2}} \right) - \cosh^{-1} \left(\frac{A_2 \phi_b}{\sqrt{1 - \frac{\Lambda_+}{|\mathcal{C}|} \phi_b^2}} \right) \right],
\end{aligned}$$

where

$$A_1 \stackrel{\text{def.}}{=} \frac{\Lambda_+ - \Lambda_- + \frac{\kappa^2}{4}}{\kappa\sqrt{-\mathcal{C}}}, \quad A_2 \stackrel{\text{def.}}{=} \frac{\Lambda_+ - \Lambda_- - \frac{\kappa^2}{4}}{\kappa\sqrt{-\mathcal{C}}}.
\tag{14.29}$$

The brane action associated to the cases listed in (14.28) reads:

$$\begin{aligned}
S_{brane}^{dS/dS} &= \frac{2\pi\eta}{G} \left[\phi_o \Theta \left(-\frac{\phi'}{L} \Big|_{-} \right) - \phi_o \Theta \left(-\frac{\phi'}{L} \Big|_{+} \right) + \right. \\
&\quad \left. + \frac{\mu_-}{2\pi} \ln \left| \frac{\mu_- A_1 + 1}{\mu_- A_1 - 1} \right| - \frac{\mu_+}{2\pi} \ln \left| \frac{\mu_+ A_2 + 1}{\mu_+ A_2 - 1} \right| \right], \\
S_{brane}^{AdS/AdS} &= \frac{\eta}{G} \left[\mu_- \ln \left| \frac{\mu_- A_1 + 1}{\mu_- A_1 - 1} \right| - \mu_+ \ln \left| \frac{\mu_+ A_2 + 1}{\mu_+ A_2 - 1} \right| \right].
\end{aligned}
\tag{14.30}$$

The Θ -terms in (14.30) cancel each other out for the same argument outlined when dealing with the bulk action. For the case of mixed transitions of the dS \rightarrow AdS-kind, the brane action would be structurally equivalent to that of the dS \rightarrow dS case, once having suitably accounted for the difference in sign of the cosmological constants involved.

14.2 Transitions among de Sitter, anti-de Sitter, and Minkowski vacua

In summary we can write the transitions so far as:

$$\begin{aligned}
 \ln \Gamma_{\text{AdS} \rightarrow \text{AdS}} &= \frac{\eta}{G} \left[\mu_- \ln \left| \frac{\mu_- A_1 + 1}{\mu_- A_1 - 1} \right| - \mu_+ \ln \left| \frac{\mu_+ A_2 + 1}{\mu_+ A_2 - 1} \right| \right] \\
 \ln \Gamma_{\text{dS} \rightarrow \text{dS}} &= \frac{2\pi\eta}{G} \left[\mu_+ - \mu_- + \frac{\mu_-}{2\pi} \ln \left| \frac{\mu_- A_1 + 1}{\mu_- A_1 - 1} \right| - \frac{\mu_+}{2\pi} \ln \left| \frac{\mu_+ A_2 + 1}{\mu_+ A_2 - 1} \right| \right] \\
 \ln \Gamma_{\text{dS} \rightarrow \text{AdS}} &= \frac{2\pi\eta}{G} \left[\mu_+ + \frac{\mu_-}{2\pi} \ln \left| \frac{\mu_- A_1 + 1}{\mu_- A_1 - 1} \right| - \frac{\mu_+}{2\pi} \ln \left| \frac{\mu_+ A_2 + 1}{\mu_+ A_2 - 1} \right| \right].
 \end{aligned} \tag{14.31}$$

Their equivalence with the corresponding expressions in BT becomes manifest under the following parametric redefinitions

$$\bar{z} = -\frac{1}{4\sqrt{\Lambda_+}} \ln \left| 1 - \frac{\phi_{tp,+}^{new 2}}{\Lambda_+} \right|, \quad \phi_{tp}^{new 2} \stackrel{def.}{=} \Lambda_{\pm} A_{1,2} \mu_{\pm} = A_{1,2} \sqrt{\frac{-\mathcal{C}}{\Lambda_{\pm}}} \Lambda_{BT}. \tag{14.32}$$

Comparing dS, AdS and Minkowski transitions

Some important observations can be drawn when comparing (14.31).

- Horizons' contribution

The first main difference between the AdS and dS transitions is the horizon contribution to the amplitude, which is crucial when considering the flat limit. A detailed explanation and interpretation of these findings will be outlined in section J.

- Bounds on wall tension

For the $\text{AdS}_2 \rightarrow \text{AdS}_2$ case, the classical turning point ϕ_o is real if and only if the tension of the brane is constrained within the following range

$$\kappa < 2 \left| \sqrt{-\Lambda_+} - \sqrt{-\Lambda_-} \right|. \tag{14.33}$$

- The flat limit from AdS

Interestingly, upon taking, either, $\Lambda_{\pm} = 0$ in $\text{AdS}_2 \rightarrow \text{AdS}_2$ processes, the amplitude is still finite

$$\boxed{
\begin{aligned}
S_{tot}^{\text{Mink.} \rightarrow \text{AdS}} &= \frac{\eta}{G} \mu_- \ln \left| \frac{\mu_- A_1 + 1}{\mu_- A_1 - 1} \right| \\
S_{tot}^{\text{AdS} \rightarrow \text{Mink}} &= -\frac{\eta}{G} \mu_+ \ln \left| \frac{\mu_+ A_2 + 1}{\mu_+ A_2 - 1} \right|
\end{aligned}
}
\quad \text{with } A_1 \stackrel{\text{def.}}{=} \frac{-\Lambda_- + \frac{\kappa^2}{4}}{\kappa\sqrt{-\mathcal{C}}}, \quad A_2 \stackrel{\text{def.}}{=} \frac{\Lambda_+ + \frac{\kappa^2}{4}}{\kappa\sqrt{-\mathcal{C}}}.$$
(14.34)

Therefore up and down-tunneling are allowed in this case. Finiteness follow from the fact that the brane action vanishes on the side where the cosmological constant is turned off.

- Detailed balance

Under the exchange $\mu_+ \leftrightarrow \mu_-$, with $\mathcal{C}_+ = \mathcal{C}_-$, we get $A_1 \leftrightarrow A_2$ therefore, from (14.30), we obtain:

$$\boxed{
\frac{\Gamma_{1 \rightarrow 2}}{\Gamma_{2 \rightarrow 1}} = e^{\frac{8\eta\pi}{G}(\phi_+^h - \phi_-^h)} = e^{\eta(S_1 - S_2)},
}$$
(14.35)

proving that the principle of detailed balance is satisfied, once we identify the value of the dilaton at the horizon with the entropy. It is interesting to notice that detailed balance is still working even in this setup, thereby consistently proving that the value of the dilaton at the horizon is playing the role of the black hole entropy. However, the particular nature of the cosmological horizon leads to unitarity issues upon taking the flat spacetime limit from a pure dS setup.

- The flat limit from dS

The main difference with respect to the AdS case is that, this time, the definition of the turning point, ϕ_o , is not subject to any parametric constraint, since $\Lambda_{\pm} > 0$. However, the presence of an additional term proportional to the cosmological horizon in S_{TOT} , inevitably leads to a divergent action upon taking the flat limit on either side of the brane

$$\boxed{
\begin{aligned}
S_{tot}^{\text{Mink.} \rightarrow \text{dS}} &= \frac{2\eta\pi}{G} \left[-\mu_- + \frac{\mu_-}{2\pi} \ln \left| \frac{\mu_- A_1 + 1}{\mu_- A_1 - 1} \right| + \mu_+ \right] \rightarrow \infty \\
S_{tot}^{\text{dS} \rightarrow \text{Mink.}} &= \frac{2\eta\pi}{G} \left[\mu_+ - \frac{\mu_+}{2\pi} \ln \left| \frac{\mu_+ A_2 + 1}{\mu_+ A_2 - 1} \right| - \mu_- \right] \rightarrow -\infty,
\end{aligned}
}$$
(14.36)

for up and down-tunneling, respectively. In particular, this proves that there is no 2D counterpart of the CDL transition for dS \rightarrow Mink. It is an important feature of the 2D theory that the divergence does not cancel, thereby signalling the need for a deeper understanding. Indeed, when taking the $\lim_{\mu_+ \rightarrow \infty}$, we are really moving very far

away with respect to the perturbative regime where the near-horizon approximation of JT-gravity is valid. As the horizon moves deeper inside the IR bulk, so too does the turning point. However, ϕ_o will at most asymptote to a finite value, ultimately reemerging from the horizon in the form of a naked singularity. At this point, the causal structure of the spacetime prevents the nucleation of a new bulk phase.

15 Lorentzian transitions with black holes

15.1 Actions with non-vanishing constant energies

We will now turn to the more general case in which a constant term (with respect to ϕ) is being added to the Lagrangian density, which now reads

$$\mathcal{L} = \sqrt{-g}\phi [\mathcal{R} - 2\Lambda_+\Theta(z - \hat{z}) - 2\Lambda_-\Theta(\hat{z} - z)] - \delta(z - \hat{z})\sigma\sqrt{(N^t)^2 + L^2(N^z + \dot{\hat{z}})^2} + \sqrt{-g}[\phi_o\chi_{\mathcal{M}} - \mathcal{B}_+\Theta(z - \hat{z}) - \mathcal{B}_-\Theta(\hat{z} - z)].$$

The additional \mathcal{B} -terms play the same role as the black hole mass in the 4D analysis carried out in [210]. Indeed, such term contributes to the 2D metric function by means of a linear term in ϕ , therefore featuring the appropriate scaling for a 2D black hole. However, unlike the 4D case, where M emerges as an integration constant, the lower dimensionality of the current analysis requires it to be defined as a parameter of the theory.

The procedure for determining the total action is very similar to the one carried out in section 14, hence we will just highlight the main differences. The Hamiltonian constraint picks up additional \mathcal{B}_\pm terms, whereas the momentum constraint remain unaltered. This follows from the fact that the determinant of the metric is N^z -independent. In particular, the latter ensures that the relation in between the conjugate momenta π_L and π_ϕ is preserved. Upon integrating the Hamiltonian constraint across the wall, the conjugate momentum with respect to the metric reads

$$\boxed{(\pi_L^2)\Big|_{\hat{z}_\pm} = 4 \left[\left(\frac{\phi'}{L}\right)^2\Big|_{\hat{z}_\pm} + \mathcal{C}_\pm + \mathcal{B}_\pm\phi + \Lambda_\pm\phi^2 \right]}, \quad (15.1)$$

and the extrinsic curvatures

$$\frac{\phi'}{L}\Big|_{\hat{z}_\pm} = \frac{(\Lambda_+ - \Lambda_-)\phi^2}{\sigma} \mp \frac{\sigma}{4} + \frac{\mathcal{C}_+ - \mathcal{C}_-}{\sigma} + \frac{(\mathcal{B}_+ - \mathcal{B}_-)\phi}{\sigma}, \quad (15.2)$$

from which the junction conditions follow similarly to the $\mathcal{B}_\pm = 0$ case. Upon performing the same procedure outlined in section 14, and still requiring $\mathcal{C}_+ = \mathcal{C}_-$, the effective potential reads

$$V_{eff} = -\frac{1}{4} \left[\left(\left(\frac{\Lambda_+ - \Lambda_- - \frac{\sigma^2}{4}}{\sigma} \right) \Phi_b^2 + \frac{(\mathcal{B}_+ - \mathcal{B}_-)\Phi_b}{\sqrt{|\mathcal{C}|\sigma}} \right)^2 - 1 + \frac{\mathcal{B}_+}{\sqrt{|\mathcal{C}|}}\Phi_b + \Lambda_+\Phi_b^2 \right] - 1. \quad (15.3)$$

We will restrict to the case $\mathcal{B}_+ = \mathcal{B}_- = \mathcal{B}$. The general case $\mathcal{B}_+ \neq \mathcal{B}_-$ can be considered without changing the conclusions. For $\mathcal{B}_+ = \mathcal{B}_-$, the turning points are

$$\boxed{\Phi_{1,2} \stackrel{\text{def.}}{=} -\frac{\mathcal{B}\Phi_o^2}{2\sqrt{|\mathcal{C}|}} \left[1 \pm \sqrt{1 + \frac{4\mathcal{C}}{\mathcal{B}^2\Phi_o^2}} \right]}, \quad (15.4)$$

which, in the $\mathcal{B} \rightarrow 0$ limit, consistently reduces to $|\phi_1| = |\phi_2| = \phi_o$, thereby correctly recovering the result obtained in section 14. From (15.4), the existence of 2 physical turning points is ensured upon requiring the following parametric constraints

$$\boxed{-\frac{1}{4} < \frac{\mathcal{C}}{\mathcal{B}^2\Phi_o^2} < 0, \quad \text{with } \mathcal{B}, \mathcal{C} < 0,} \quad (15.5)$$

which can also be rewritten as follows

$$\boxed{-2\sqrt{|\Lambda\mathcal{C}|} < \mathcal{B} < 0.} \quad (15.6)$$

In particular, (15.6) defines a bound for \mathcal{B} , which, as will be argued in section 15.3, plays an essential role in our analysis. Indeed, as shown next, this parameter is related to the black hole mass.

Metric and dilaton profiles

Let us generalise the results of section 14.1 to the case $\mathcal{B} \neq 0$. Here, the dilatonic e.o.m. pick up an additional constant term, and, consequently, so too does the field,

$$\phi'' = 2\phi L^2 \Lambda_{\pm} \pm \mathcal{B} L^2 \quad \Rightarrow \quad \phi = 2a \coth(bz) \pm \frac{\mathcal{B}}{2\Lambda_{\pm}}. \quad (15.7)$$

Under the coordinate transformation (14.16), the line element turns into

$$\boxed{ds^2 = \frac{\left(\phi \pm \frac{\mathcal{B}}{2\Lambda_{\pm}}\right)^2 - 4a^2}{4a^2} \left(-dt^2 + \frac{4a^2}{b^2} \frac{d\phi^2}{\left(\left(\phi \pm \frac{\mathcal{B}}{2\Lambda_{\pm}}\right)^2 - 4a^2\right)^2} \right)}, \quad (15.8)$$

which is exactly of the S(A)dS-kind. The parameters featuring in (15.8) can be related to the ones we have and will be using in the expressions for the total actions via the following identification

$$\frac{\mathcal{B}^2}{4\Lambda_{\pm}^2} - 4a^2 = \frac{\mathcal{C}}{\Lambda_{\pm}} \quad \Rightarrow \quad \mathcal{B} = \pm 2\Lambda_{\pm} \sqrt{4a^2 + \frac{\mathcal{C}}{\Lambda_{\pm}}}. \quad (15.9)$$

Note that, in this case, the metric has 2 horizons, associated to the roots of the metric function in (15.8) that read

$$\phi_{h,1,2}^{\pm} = -\frac{|\mathcal{B}|}{2\Lambda_{\pm}} \left[\text{sign}_{\pm}(\mathcal{B}) \pm \frac{4a\sqrt{\Lambda_{\pm}}}{\mathcal{B}} \right] \stackrel{(15.9)}{=} -\frac{|\mathcal{B}|}{2\Lambda_{\pm}} \left[\text{sign}_{\pm}(\mathcal{B}) \pm \sqrt{1 - \frac{4C\Lambda_{\pm}}{\mathcal{B}^2}} \right]. \quad (15.10)$$

This shows the importance of introducing \mathcal{B} . Which sign is being identified with the innermost or outermost horizon of the given spacetimes entirely depends on the signs of the parameters of the theory $(\mathcal{B}, \Lambda_{\pm})$. In particular, \mathcal{B} will have opposite sign on the 2 sides to ensure the presence of 2 horizons for the $\Lambda_{\pm} > 0$ case. Setting the outer horizon as a reference, the innermost horizon for $\mathcal{B} < 0$ ⁷⁸ is given by the choice of the + root. As a consequence of this, the outermost horizon on the inner vacuum is given by the - sign, with $\mathcal{B} > 0$. The change in sign for \mathcal{B} in the 2 vacua ensures that we can systematically use the same coordinate ϕ on both sides of the brane. Indeed, this is not something surprising, given that the value of the dilaton changes sign in different causal patches of a maximally-extended metric, and therefore it is in agreement with the description outlined above given that the nucleation process requires crossing the horizon of a given spacetime in static coordinates.

In the 4D case, FMP showed that the black hole mass arises as an integration variable, while in the 2D case we show that it is part of the theory, since it features at the level of the Lagrangian density. However, one should not be misled by this, in the sense that, unlike the Euclidean formalisms referred to in section 2, the main feature of the Hamiltonian method is precisely that of implementing constraints at the level of the Lagrangian, such that the total action describing the transition is effectively accounting for a joined system of two vacua separated by a wall. The black hole mass will therefore enter in the joined system's Lagrangian inside the Hamiltonian constraint, and the matching conditions will therefore act as a selection rule assigning to each spacetime a notion of state labelled by their respective value of the black hole mass, \mathcal{B}_{\pm} .

15.2 Transitions with conical defects

The evaluation of the extremised total action can be carried out in complete analogy with the procedure applied in 14. The bulk and brane actions read

$$S_{bulk}^{(\pm)} = \frac{2\eta}{G} \int dz \left[L \sqrt{\left(\frac{\phi'}{L}\right)^2 + C + \mathcal{B}\phi + \Lambda_{\pm}\phi^2} - \phi' \cosh^{-1} \left(\frac{\phi'}{L\sqrt{-C - \mathcal{B}\phi - \Lambda_{\pm}\phi^2}} \right) \right],$$

$$S_{brane} = \frac{2\eta}{G} \int_{\phi_1}^{\phi_2} d\phi_b \left[\cosh^{-1} \left(\frac{\phi'_b(\hat{z} - \epsilon)}{L\sqrt{-C - \mathcal{B}\phi_b - \Lambda_- \phi_b^2}} \right) - \cosh^{-1} \left(\frac{\phi'_b(\hat{z} + \epsilon)}{L\sqrt{-C - \mathcal{B}\phi_b - \Lambda_+ \phi_b^2}} \right) \right].$$

⁷⁸This condition follows from requiring both turning points to be physical, see eq. (15.4).

Once more, extremisation of the bulk actions occurs at the horizons according to the range of the mass parameter. This time, we have two horizons on either side, defined by

$$\phi_{h,1,2}^+ \stackrel{\text{def.}}{=} -\frac{|\mathcal{B}|}{2\Lambda_+} \left[\text{sgn}(\mathcal{B}) \mp \sqrt{1 - \frac{4\Lambda_+ \mathcal{C}}{\mathcal{B}^2}} \right], \quad \phi_{h,1,2}^- \stackrel{\text{def.}}{=} -\frac{|\mathcal{B}|}{2\Lambda_+} \left[\text{sgn}(\mathcal{B}) \mp \sqrt{1 - \frac{4\Lambda_- \mathcal{C}}{\mathcal{B}^2}} \right], \quad (15.11)$$

with $\mathcal{C}_\pm < 0$. Given that $\Lambda_\pm \geq 0$, for both horizons to be physical, it must be that $\text{sgn}(\mathcal{B}) = -1$. The bulk action is extremised at the outermost and innermost horizons on the 2 sides of the brane, respectively, (15.11), and at the turning point ϕ_2 , provided the argument of \cosh^{-1} changes sign in S_{bulk} .

The total bulk action reads

$$S_{\text{bulk}} = \frac{2\eta}{G} \left[-\phi_{h,2}^- + \phi_{h,1}^+ + \phi_2 \left[\Theta \left(-\frac{\phi'}{L} \Big|_- \right) - \Theta \left(-\frac{\phi'}{L} \Big|_+ \right) \right] \right], \quad (15.12)$$

and therefore we can write the transitions as:

$$\boxed{S_{\text{tot}}^{(\text{A})\text{dS} \rightarrow (\text{A})\text{dS}} = \frac{2\eta}{G} \left[-\frac{\mathcal{B}}{2\Lambda_+} \left[1 - \sqrt{1 - \frac{4\mathcal{C} \Lambda_+}{\mathcal{B}^2}} - \ln \left| \frac{y_4}{y_3} \sqrt{\frac{1 - y_3^2}{1 - y_4^2}} \right| \right] + \frac{\mathcal{B}}{2\Lambda_-} \left[1 - \sqrt{1 - \frac{4\mathcal{C} \Lambda_-}{\mathcal{B}^2}} - \ln \left| \frac{y_2}{y_1} \sqrt{\frac{1 - y_1^2}{1 - y_2^2}} \right| \right] + -\phi_2 \ln \left| \frac{A_2 y_2}{A_4 y_4} \sqrt{\frac{1 - y_4^2}{1 - y_2^2}} \right| + \phi_1 \ln \left| \frac{A_2 y_1}{A_4 y_3} \sqrt{\frac{1 - y_3^2}{1 - y_1^2}} \right| + -\frac{1}{2} \left[\sqrt{\frac{\mathcal{B}^2}{4\Lambda_-} - \mathcal{C}} \ln \left| \frac{1 - y_2}{1 + y_2} \frac{1 + y_1}{1 - y_1} \right| - \sqrt{\frac{\mathcal{B}^2}{4\Lambda_+} - \mathcal{C}} \ln \left| \frac{1 - y_4}{1 + y_4} \frac{1 + y_3}{1 - y_3} \right| \right] \right],$$

where $y_{1,2} \equiv y_{3,4} = y(\phi_{1,2})$,

$$y \stackrel{\text{def.}}{=} \sqrt{\frac{\Lambda}{\frac{\mathcal{B}^2}{4\Lambda} - \mathcal{C}}} \left(\frac{\mathcal{B}}{2\Lambda} + \phi_b \right), \quad \phi_{1,2} = -\frac{\mathcal{B}\phi_o^2}{2\sqrt{|\mathcal{C}|}} \left[1 \pm \sqrt{1 + \frac{4|\mathcal{C}|}{\mathcal{B}^2\phi_o^2}} \right] \quad (15.13)$$

and, for pure notational convenience, the following parametric redefinitions were performed

$$A_1 = \frac{\Lambda_+ - \Lambda_- + \frac{\kappa^2}{4}}{\kappa\sqrt{\mathcal{C} - \frac{\mathcal{B}_-^2}{4\Lambda_-}}}, \quad A_3 = \frac{\Lambda_+ - \Lambda_- - \frac{\kappa^2}{4}}{\kappa\sqrt{\mathcal{C} - \frac{\mathcal{B}_+^2}{4\Lambda_+}}} \quad (15.14)$$

$$A_2 \stackrel{\text{def.}}{=} \frac{A_1}{\sqrt{\Lambda_-}} \sqrt{\frac{\mathcal{B}^2}{4\Lambda_-} - \mathcal{C}}, \quad A_4 \stackrel{\text{def.}}{=} \frac{A_3}{\sqrt{\Lambda_+}} \sqrt{\frac{\mathcal{B}^2}{4\Lambda_+} - \mathcal{C}}.$$

Setting $\mathcal{B} = 0$, (15.13) reduces to the results obtained in section 14. Upon exchanging $\Lambda_- \longleftrightarrow \Lambda_+$, the turning points on the 2 sides are correspondingly mapped to $y_2 \rightarrow -y_4$

and $y_1 \rightarrow -y_3$. From this follows that the arguments of all the \ln -terms are inverted. The ratio of direct and reverse processes, lead to an expression for detailed balance which reads

$$\boxed{\ln \frac{\Gamma_{\Lambda_+ \rightarrow \Lambda_-}}{\Gamma_{\Lambda_- \rightarrow \Lambda_+}} = \frac{4\eta}{G} [\phi_h^+ - \phi_h^-]}, \quad (15.15)$$

with ϕ_h^\pm denoting the value of the dilaton at the BH horizons on either side. Notice that this holds for, either, (A)dS \rightarrow (A)dS.

15.3 Flat limit

In the last part of this section, we will show that, upon taking the flat limit on either side of the brane, the 2D transition amplitude is still well defined. Let us consider the $\Lambda_- \equiv 0$ case. In the flat limit, the value of the dilaton at the outermost horizon on the inner vacuum's side becomes constant

$$\lim_{\mu_- \rightarrow \infty} \frac{\mathcal{B}}{2\Lambda_-} \left[1 - \sqrt{1 - \frac{4\mathcal{C}\Lambda_-}{\mathcal{B}^2}} \right] \equiv \boxed{\frac{\mathcal{C}}{\mathcal{B}} \stackrel{\text{def.}}{=} \phi_f}. \quad (15.16)$$

The constraint on the black hole mass ensuring the existence of 2 physical turning points, (15.5), (derived in section 15.1) can be re-expressed in terms of (15.16) as follows

$$\boxed{0 > \mathcal{B} > -\frac{\phi_f}{4\Phi_o^2}}. \quad (15.17)$$

In the $\lim_{\mathcal{B} \rightarrow 0}$, (15.16) diverges and therefore we correctly recover the behaviour of the divergent dilaton at the horizon encountered in section 14. The rate (including background subtraction) becomes:

$$\boxed{B_{tot}^{\text{Mink.} \rightarrow \text{dS}} = \frac{2\eta}{G} \left[-\phi_{h,2}^- + \phi_f - \sqrt{\frac{\mathcal{B}^2}{4\Lambda_-} - \mathcal{C}} \left[y_2 \ln \left| \frac{A_2 y_2}{\sqrt{1-y_2^2}} \right| - y_1 \ln \left| \frac{A_2 y_1}{\sqrt{1-y_1^2}} \right| \right] + \right. \\ \left. - \sqrt{\frac{\mathcal{B}^2}{4\Lambda_-} - \mathcal{C}} \left[\tanh^{-1} y_2 - \tanh^{-1} y_1 \right] \right],}$$

and is symmetric under the exchange of inner and outer vacua. Detailed balance is still satisfied:

$$\boxed{\ln \frac{\Gamma_{\text{Mink.} \rightarrow (\text{A})\text{dS}}}{\Gamma_{(\text{A})\text{dS} \rightarrow \text{Mink.}}} = \frac{4\eta}{G} [\phi_f - \phi_h]} \quad (15.18)$$

Summary of results and comparison with the 4D case

The above calculations show that:

- The constant \mathcal{B} -term added to \mathcal{L} is playing the role of the black hole mass. This introduces a second turning point from the 2D dS point of view, as long as (15.5) is satisfied.
- The flat spacetime limit for (A)dS \rightarrow (A)dS transitions, in the presence of a black hole, is finite. The dilaton at the horizon reaches a constant value, consistent with unitarity.
- In particular, the Mink \rightarrow AdSBH transition we obtain is in agreement with arguments provided in [224], supporting compatibility of the process with the holographic principle.

The key differences with respect to the 4D case analysed in [7] are:

- The divergence of $dS_2 \rightarrow$ Mink₂ actions in absence of black holes
- The black hole mass is a parameter of the theory, explicitly featuring in the Lagrangian density, and is not an integration constant, albeit subject to the constraint (15.5).

16 Holographic interpretation

The importance played by black holes in ensuring the finiteness of transition amplitudes upon taking the flat limit, motivates seeking for a deeper physical interpretation of our findings. In doing so we rely upon holography, with the main reason for this being that holographic techniques in 2D have played a key role for addressing the information loss paradox within the context of black hole physics, [245]. The importance of understanding decay processes involving Minkowski and AdS spacetimes from a holographic perspective was already addressed in [224], and one of our findings is proving agreement between the arguments outlined in [224] and ours in the lower-dimensional setup.

In the present section, we argue that 2D vacuum transitions can be embedded within a similar holographic setup as the one describing black hole evaporation in braneworld models. The main reason being the role played by JT-gravity in both settings. In pursuing such task, we argue that:

- An entropy interpretation may be assigned to the total action or bounce and not only to the ratio of decay rates as in detailed balance.
- Unitarity issues emerging in the flat limit can be overcome in the Hamiltonian formalism in presence of non-extremal spacetimes in a way which is analogous to the

island proposal. This follows from the fact that the total action can be rewritten as a difference of generalised entropies, each one defined as [211, 245]⁷⁹

$$S_{\text{gen}}(R) = \min_{\partial I} \text{extr} \left[\frac{\text{Area}(\partial I)}{4G_N} + S_{EE}(I \cup R) \right]. \quad (16.1)$$

- Transitions described by means of the BT and FMP methods are actually *local*.
- For the case of vacuum transitions, the entanglement in 16.1 is *internal*.
- From the boundary perspective, a vacuum transition in the bulk may correspond to a phase transition in the boundary such as a deconfinement to confinement transition.

16.1 Islands and horizons in holographic black hole evaporation

We will start by briefly reviewing the recent work on black holes and the information paradox that will be useful for addressing the vacuum transitions in a similar way. Our first motivation in searching for a holographic embedding of vacuum transitions, is the importance of the role played by horizons, with the Holographic Principle being its prototypical example. Before delving into a detailed analysis of our findings, we first provide some further arguments in support of the similarity between the information loss paradox, and the divergence of the entropy associated to the cosmological horizon in the static dS patch. To the best of our knowledge, the arguments we propose provide an original motivation towards applying new holographic techniques, as outlined from section 16.2 onwards.

Holographic black hole evaporation and gravitating baths

In the first formulation of the information paradox, namely within the context of the evaporating (1-sided) black hole, spacetime is asymptotically flat. In its corresponding holographic embedding, this assumption is mapped to having a *non-gravitating bath*. In the KR/HM-construction realising it, [211, 245, 269] (as depicted on the RHS of figure 52), the bath (denoted in blue) lives on the conformal boundary at $z = 0$, and, in this sense, is non-gravitating.




Figure 52. The dashed vertical line (on the LHS) denotes the *Cardy branes* within the KR/HM-construction, [211, 245, 269], separating the causal wedge of the *bulk* CFT_D from that of the black hole QMs living on the defect BCFT_{D-1}. The purple shaded regions on the RHS denote the entanglement wedges for the radiation region $A \subset R$ as time evolves. The orange curves on the RHS correspond to the RT-surfaces associated to the choice of the boundary subregion A .

⁷⁹This expression turns out to be applicable to dS₂ spacetimes as well, and our results are therefore in agreement with those of [252].

The KR/HM-construction is able to simultaneously realise three different, albeit physically equivalent, descriptions of the same system, [245], namely:

1. From the *bulk*, the system is described by an AdS_{D+1} with an ETW brane.
2. From the *brane*, it is given by a CFT_D with a UV-cutoff + gravity on the ETW brane (asymptotically AdS_D) which in turn is coupled with transparent boundary conditions to a BCFT_D .
3. From the *boundary*, it is a BCFT_D with nontrivial BCs.

The key quantity to keep track of is the ratio between the central charges of the bulk and defect CFTs defining the BCFT_D , [291],

$$F \stackrel{\text{def.}}{=} \frac{c_{bdy}}{c_{bulk}} = \frac{6 \ln g}{c_{bulk}}, \quad (16.2)$$

from which the tension of the brane, T , and brane angle, Θ , can be determined as follows, [291],

$$e^F = \frac{1+T}{\sqrt{1-T^2}} = \frac{1+\sin\Theta}{\cos\Theta}, \quad 1-e^{-2F} = \frac{2\sin\Theta}{1+\sin\Theta}, \quad (16.3)$$

with BH evaporation being described by the mutual exchange of degrees of freedom between the bath and the black hole. Constraints on the CFT parameters are holographically dual to the allowed ranges for T and Θ by virtue of (16.3). From this follows that t_{Page} is mapped to a critical value of F , beyond which an island is expected to arise. The second orange line in figure 53, which joins R with I , becomes the dominant channel⁸⁰ for $t > t_{Page}$. The endpoint of I is the quantum extremal surface (QES) contributing with the area law term to S_{gen}

$$S_{gen}(R) = \min_{\partial I} \text{extr} \left[\frac{\text{Area}(\partial I)}{4G_N} + S_{EE}(I \cup R) \right]. \quad (16.4)$$

The main caveat featuring in the non-gravitating bath setup, namely the presence of a massive graviton [283], $m_{grav}^2 \sim c_{bulk}/c_{bdy}$, can be overcome by introducing a gravitating bath, as shown on the RHS of figure 53. The 2 configurations can be smoothly interpolated under the action of an RG-flow driving the conformal boundary at a finite cutoff in the AdS bulk, thereby rendering it gravitating. In terms of the parameter F , (21.10), the RG-flow corresponds to the limit in which $c_{bdy} \gg c_{bulk}$ such that, to sufficiently good approximation, the BCs of the BCFT_D encode all the degrees of freedom of the boundary theory. Correspondingly, the only part of the conformal boundary that is left is a codimension-2 theory, CFT_{D-1} , located at $z = \epsilon$, as shown on the RHS of figure 53, where the issue of the *flat entanglement spectrum* can be circumvented if calculating the entanglement on a subsystem of the relic codimension-2 boundary theory, [283].

Cosmological horizon

Equipped with this insight, we now turn to the case of the cosmological horizon, the main point of this digression being that of outlining the similarity between the unitarity issues in global dS and black hole evaporation in asymptotically flat spacetimes.

The cosmological horizon is placed at a finite distance⁸¹, $r_h = 1/H$, with respect to an observer located at the origin of the static patch, thereby implying that there is only a finite amount of

⁸⁰The study of phase transitions in holography initiated from the first work of [292].

⁸¹Each point along the red line in figure 54 is a codimension-2 surface of radius r_h .




Figure 53. Non-gravitating (left) and a gravitating bath (right) KR/HM-constructions.




Figure 54. The picture on the left is at a fixed time slice. The plane is where the CFT lives and the bulk is AdS, with z realised holographically. The subregion of bulk spacetime lying between A and Γ_A defines the *entanglement wedge* of A . Global dS_D for $D > 2$ (on the RHS). An observer at $r = 0$ can only access degrees of freedom within the cosmological horizon $r = 1/H$.

spacetime that she/he can be in causal contact with. Because of this, one might be led to conclude that the area of the cosmological horizon should somehow be related to the entropy of the static patch itself, as the Holographic Principle would suggest. However, it is also known that the dS Hilbert space is infinite-dimensional, [285]. The reason for this follows from the fact that the dS isometry group is noncompact, and therefore has no finite dimensional unitary representations. This provides one of the main obstacles towards extending the holographic dictionary of [203] to the case of a positive cosmological constant.

The configuration just described seems to share similar unitarity issues as encountered for the case of the evaporating black hole. Assuming the horizon surrounding the static patch arises from partial tracing over the global spacetime, c.f. figure 54, when taking $\lim_{\Lambda \rightarrow 0}$, the cosmological horizon will correspondingly be pushed towards \mathcal{I}^+ , at which the entropy should vanish. Intuitively, this is in agreement with the partial-tracing argument outlined above, since, as the cosmological horizon is pushed further away from the observer in the static patch, the degrees of freedom which were previously “hidden” (i.e. traced-over), are expected to re-enter the horizon, thereby becoming accessible to the observer. However, this falls short from being true given the divergence of the entropy defined by the area-law, which, e.g., in 4D scales as $\sim \mathcal{O}(\frac{1}{H^2})$. The area of the codimension-2 surface located at such distance can be interpreted by holographic arguments as the entropy of the static patch, which, in the von Neumann formulation, follows from having traced over the spacetime beyond the cosmological horizon, in analogy with the black hole picture. As such, it is not a QES.

Given the monotonic behaviour of, both, the background action for $dS \rightarrow dS$ transitions (as found in [7]) and the von Neumann entropy, we suggest that the divergence issue arising from the flat spacetime limit of de Sitter might be solved by introducing a more suitable definition of the de Sitter entropy, in a similar fashion as S_{gen} enables to recover unitarity within the context of black hole evaporation. As anticipated at the beginning of this section, this identification turns out to be possible⁸².

16.2 Holographic embedding of vacuum transitions

Here we present our proposal for the holographic description of the vacuum transitions. As highlighted throughout the previous sections, the three methods used for deriving transition amplitudes exhibit unexpected behaviours. In the present section, we wish to provide a unifying holographic framework describing all processes, mostly motivated by the fact that:

1. The action for the system of 2D spacetimes joined by a wall is analogous to the JT-gravity coupled to a CFT_2 setup adopted to formulate the island proposal within the context of the information paradox.
2. The *bulk-brane-boundary* description of the same phenomenon in the KR/HM setup accounts for complementary features of the same underlying process.

It is indeed in the spirit of the BCFT formulation of the information paradox, that we propose a holographic embedding of the transitions described analytically in the first three sections of the present work, after having implemented suitable adaptations due to the specifics of the configurations being analysed.




Figure 55. The LHS shows a generalisation of figure 53 for vacuum transitions. A deformation drives the RG-flow on the boundary, assuming $c_{bulk} \ll c_{bdy}$ at both endpoints. In the IR of the RG-flow (as shown on the RHS), the vacuum transition takes place in between subregions of different AdS_D 's, each one lying on a different ETW. More details will be outlined in the concluding part of this section.

Figure 55 shows our proposal for the holographic embedding of 2D vacuum transitions:

- Given that the process involves two different spacetimes, each one being associated to a different JT-action, the holographic embedding of the transition can be described by two ETW branes (each one corresponding to one of the two vacua, $ETW_{1,2}$) with the brane separating them (mediating the decay process) being geometrically realised by a composite CFT (denoted by $\overline{CFT}_A \otimes CFT_B$ in figure 55), which can be achieved by performing the folding trick.

⁸²Interesting new developments towards realising a dS_4/CFT_3 correspondence have recently been presented in [584], where the interpretation of the maximal entanglement between the north and south pole in global dS shares the same motivation as the argument presented in our work.

- The radiation region lies on the conformal boundary (denoted by the tensor product of CFTs arising from the folding trick), which in turn is brought at a finite cutoff, ϵ_{IR} , with nontrivial BCs, the latter denoted by $BCFT_{D-1}, BCFT'_{D-1}$. The theory living on the gravitating bath interpolates between the values of the D -dimensional cosmological constants. As such, the wall plays the role of the entangling surface in between the spacetimes involved, with the entanglement being *internal*.
- Attached to the radiation region are the two ETW branes. Each one of them accommodates one of the two spacetimes, (A)dS $_D$.
- The composite CFT on the gravitating bath (depicted on the LHS of figure 55) is ultimately driven by an irrelevant operator to the configuration on the RHS of figure 55, which is reminiscent of the wedge holographic construction, [237].
- As explained at multiple stages throughout our treatment, the total bounce and action for a given vacuum transition, result once having suitably accounted for background subtraction. As we shall see, it is particularly instructive to assign a corresponding holographic realisation of the background configuration as well. Our proposal is represented in figure 56 on the RHS. We now turn to explaining it in more detail, albeit most arguments follow through from the ones outlined when explaining the RHS of figure 55. The background action is a pure CFT. As such, it can be equally described independently of the choice of the UV cutoff. Hence, starting from a pure AdS $_{D+1}$ /CFT $_D$ setup, with the conformal boundary defined on the whole real line at $z = 0$, we can perform a suitable conformal transformation enabling to restrict the boundary domain to an interval with BCs dual to two ETW branes characterised by the same D -dimensional cosmological constant. Given that the theory is still conformal, the width of the segment can be arbitrary; for the purpose of interest to us, we need to bring the conformal boundary at the same IR cutoff as the two-spacetime joined system on the RHS of figure 55, such that the background subtraction exactly overlaps with the instanton.




Figure 56. The background configuration, with respect to which the decay rate is normalised, is drawn on the right. From the boundary theory, CFT $_{D-1}$, this corresponds to the deconfined phase of the theory.

Concrete proposal for the vacuum transition from the CFT side

If there is a holographic dual formulation of vacuum transitions, a natural question to ask is: what is the corresponding physical effect that happens on the CFT side that describes the vacuum transition in the bulk. Here we proposed that on the CFT side, the vacuum transition corresponds to a field theoretical phase transition such as a confinement/de-confinement phase transition. For this we need to identify the dual of the background spacetime and the dual of the final composite

spacetime. For concreteness let us concentrate on a bulk AdS. As we described before the vacuum transition corresponds not to the decay of a full AdS but a portion of an AdS. For the dual we may represent this in terms of a double-well scalar potential that originally has two vacua related by $\phi \rightarrow -\phi$ separated by a domain wall. Modding out by the $\phi \rightarrow -\phi$ symmetry this reduces the bulk spacetime to a portion with a boundary. This can provide a description of the dual of the original AdS. The description of the final spacetime can be seen in this way as a similar scalar potential but with the two minima being non-degenerate corresponding to two different spacetimes joined by the boundary. The question is how are the theories in the two boundaries related.

In figure 56 the background can be thought of as being given by a degenerate double well potential, with the two vacua identified. In their holographic realisation in terms of ETW branes (as shown on the RHS of figure 56), only one of them will be undergoing the transition, leading to the configuration depicted on the RHS of figure 55.

The interpolation between the deconfined and the confined phase of the theory living on the boundary can be realised by adding to the deconfined phase corresponding to figure 56, a deformation, leading to degeneracy breaking between the two vacua, as shown on the RHS of figure 57. Given that our analysis focuses on 2D, $T\bar{T}$ -deformations will turn out being the relevant ones.

The reason why we can effectively interpret vacuum transitions in terms of a deconfinement/confinement transition can be understood making use of the analysis carried out in [585] and [586]. In the former, it is shown that the difference between the two phases is dictated by the different dependence of the entanglement entropy on N , which is the key parameter in the holographic dictionary⁸³. In particular, following the RT prescription, this amounts to a difference in the behaviour of the expectation value of the Wilson line connecting two arbitrary points on the conformal boundary: for the deconfined phase, $S_{EE} \sim \mathcal{O}(N^2)$, whereas for the confined phase, $S_{EE} \sim \text{const}$. What this practically means is that, the former is actually associated to a conformal theory at $z = 0$, for which the Wilson line diverges unless a UV-cutoff is introduced. Bulk reconstruction, in such setup, allows to recover the entire (infinite) AdS_{D+1} bulk volume upon removing the cutoff. On the other hand, the confined phase is characterised by a constant value of the entanglement entropy with respect to N , meaning that bulk reconstruction can effectively account for a fixed AdS_{D+1} bulk volume. In figure 56, this corresponds to having brought the conformal boundary at ϵ_{IR} , resulting only in a finite (A)dS_D volume being involved in the process.

In a similar, and somewhat complementary fashion, the CFT description of the processes dealt with in our treatment find realisation in a recent work [586], where scalar potentials of the kind depicted in figure 57 have been identified with degenerate and non-degenerate vacua in studies of 2d QCD. In [586] the degenerate case can be identified with a deconfinement phase, whereas the non-degenerate case with a confined phase⁸⁴. In their treatment, the expectation value of the Wilson loop separating different vacua exhibits, either, an area- or perimeter-law like behaviour according to, whether, the system is in the confined or deconfined phase, respectively. More efficiently, such classification can be further performed by analysing the ratio of partition functions of the gauged and ungauged theory in the infinite volume⁸⁵ limit. If such ratio is order unity, it means the system is deconfined. On the other hand, if the ratio vanishes it is confined.

In drawing a comparison between the processes dealt with in the present work and those of Komargodski et al., it is important to notice that, in such reference, their analysis embraces, both, vacua and universes. While often used interchangeably, the main difference between vacua and universes is that the latter are separated by infinite barriers, whereas the former might allow finite tension domain walls interpolating between different superselection sectors of the same universe,

⁸³Indeed, N is related to the central charge and the AdS radius.

⁸⁴See also [251] for an interesting discussion of confinement and cosmology.

⁸⁵Namely the large- N limit.




Figure 57. Under the action of an irrelevant operator, the boundary theory can be smoothly interpolated from the deconfined (left) to the confined phase (right). The latter is the field theory realisation of the RHS of figure 55. In such description, the interpolating scalar, ϕ , plays the role of a monopole field in a deconfinement/confinement transition, in a similar fashion as in a previous work by Komargodski et al.

and are therefore the ones relevant for our treatment. The systems analysed in Komargodski et al.'s work exhibit multiple vacua, distributed between different universes, and study confinement and deconfinement of Wilson lines interpolating between them. Their counterpart for transitions of the kind associated to the RHS of figure 57, are therefore those involving confined Wilson lines interpolating between vacua belonging to the same universe.

Clearly more work needs to be done in order to fully describe the vacuum transitions from the CFT side.

16.2.1 Local transitions, entanglement and islands in wedge holography

As a first concrete application of the tools outlined in the first part of this section, we now turn to the holographic interpretation of the FMP results obtained in sections 14 and 15. In particular, we prove that:

- Under suitable parametric redefinition, the results obtained by means of the FMP method are found to agree with the expression provided by [274] for describing mutual approximation of boundary states belonging to different CFTs.
- The corresponding expression for the transition rate in presence of gravity, and in absence of black holes, is given by the difference of entropies of $T\bar{T}$ -deformed CFTs, hence proving the locality of the nucleation process. Given its interpretation as being an internal entanglement, such transitions provide an example of an $\text{AdS}_2/\text{CFT}_1 \subset \text{AdS}_3/\text{CFT}_2$.⁸⁶
- Upon adding black holes, instead, the total action can be expressed as the difference of generalised entropies, with an island emerging beyond a critical value of the black hole mass. As such, this is an example of an $\text{AdS}_2/\text{CFT}_1 \not\subset \text{AdS}_3/\text{CFT}_2$.

ICFTs and wedge holography

In this subsection, we prove that quantum transitions involving different subregions of spacetime can be described by means of dual CFTs interacting via an interface, [274]. The authors of the

⁸⁶The lower-dimensional holographic setup involved in these processes shares similar features as to the one involving spacetimes emerging from matrix QMs, as analysed, e.g. in [297], and their higher dimensional counterparts in [298].

latter describe mutual approximation of boundary states, $\Psi_{A,B}$, belonging to the Hilbert spaces of different CFTs, (as shown in figure 58).

The correspondence adopted in [274], specifying to the case of $\text{AdS}_3/\text{CFT}_2$, relates the following set of parameters on either side of the interface

$$\boxed{L_1, L_2, \kappa \longleftrightarrow c_A, c_B, c_{bdy} \sim \ln g,} \quad (16.5)$$

with $L_{1,2}$ being the AdS radii, κ the tension of the wall separating the bulk spacetimes, $c_{A,B}$ the central charges, and $\ln g$ denoting the entropy of the ICFT, S_{ICFT} , with g defining the degeneracy of the ground state.




Figure 58. Replacing a small portion of the background CFT_A with a different theory, CFT_B implies a change in the energy spectrum. Because of this, for a given boundary state, the corresponding S_{EE} leads to a different effective bulk-reconstruction of the AdS dual.

Our first finding is that $\text{AdS}_2 \rightarrow \text{AdS}_2$ transitions are equivalent to the case where the mutual approximation described in [274] only involves the ground states. Indeed, under suitable parametric redefinition, the action for the transition coincides with the one obtained by [274], as explicitly shown below

$$\begin{aligned} S_{tot}^{\text{AdS} \rightarrow \text{AdS}} &= \frac{\eta}{G} \left[\mu_- \ln \left| \frac{\mu_- A_1 + 1}{\mu_- A_1 - 1} \right| - \mu_+ \ln \left| \frac{\mu_+ A_2 + 1}{\mu_+ A_2 - 1} \right| \right] \\ &= \frac{2\eta}{G} \left[(\mu_+ - \mu_-) \tanh^{-1} \left(\frac{\Delta_-}{\kappa} \right) - (\mu_+ + \mu_-) \tanh^{-1} \left(\frac{\kappa}{\Delta_+} \right) \right] \\ &= 2\eta S_{ICFT} = 2\eta \ln g(\kappa) = -2\eta F_\partial, \end{aligned}$$

where we defined

$$\Delta_\pm \stackrel{\text{def.}}{=} 2 \left(\sqrt{-\Lambda_+} \pm \sqrt{-\Lambda_-} \right), \quad (16.6)$$

and $g(\kappa)$ denotes the ground state degeneracy and F_∂ the *boundary free energy*, which is a monotonic function of κ . The entire transition is hence described by means of a codimension-2 theory as depicted in figure 59, with the bulk emerging via wedge holography.

Making use of the following parametric redefinitions

$$\mu \stackrel{\text{def.}}{=} \sqrt{\frac{\lambda c}{12}}, \quad \phi_o \stackrel{\text{def.}}{=} r, \quad \lambda \stackrel{\text{def.}}{=} 2\sqrt{-c\Lambda}, \quad (16.7)$$

the total action can be rewritten as

$$\boxed{S_{tot}^{(\text{A})\text{dS} \rightarrow (\text{A})\text{dS}} = 2\pi\eta \left[S_{TT}^- - S_{TT}^+ \right]_{\text{univ.}} = -2\pi\eta F_\partial = 2\pi\eta S_{ICFT},} \quad (16.8)$$




Figure 59. As shown in [274], the trajectory of the interface separating pure AdS₃ phases has no turning point (LHS), and, correspondingly, the degeneracy of the ground state is a monotonic function of the tension κ , as found in section 13. No island emerges in absence of conical defects (cf. picture on the RHS). The bulk spacetime is completely determined by means of wedge holography from the CFT_{D-1} . S_{EE} is evaluated on the codimension-2 surface at the conformal boundary.

where

$$\begin{aligned}
S_{T\bar{T}} &= \left(1 - \frac{r}{2} \frac{\partial}{\partial r}\right) \ln \mathcal{Z}_{S^2}^{AdS, T\bar{T}} = \pi \frac{c}{3} \sinh^{-1} \left(\sqrt{\frac{12}{c\lambda}} r \right) \\
S_{r\bar{r}} &= \left(1 - \frac{r}{2} \frac{\partial}{\partial r}\right) \ln \mathcal{Z}_{S^2}^{dS, T\bar{T}} = \frac{c}{6} + \frac{c}{3} \ln \left| \sqrt{\frac{\lambda c}{12}} \frac{1}{r} \right| + \frac{c}{3} \tan^{-1} \left(\frac{1}{\sqrt{\frac{\lambda c}{12r^2} - 1}} \right) + \text{const.} \\
&= S_\epsilon + S_{\text{univ.}},
\end{aligned}$$

for the duals of AdS₃ and dS₃, respectively, [240, 241]. $S_{\text{univ.}}$ and S_ϵ denote the *universal* and *cutoff-dependent* parts of (16.9), respectively, with

$$S_\epsilon \stackrel{\text{def.}}{=} \frac{c}{3} \ln \left| \sqrt{\frac{\lambda c}{12}} \frac{1}{r} \right|, \quad \text{and} \quad \epsilon \stackrel{\text{def.}}{=} \sqrt{\frac{\lambda c}{12}}. \quad (16.9)$$

According to the Casini-Huerta-Myers prescription, the universal part is the one contributing to the definition of the boundary free energy for a BCFT, and therefore this is the only term we need to retain for our purposes. Indeed, S_ϵ can be removed by simply setting $r = \epsilon$, namely choosing the cutoff to be equal to the localisation radius r . From (16.8), we deduce that:

1. This chain of equivalence relations, (16.8), suggests an interesting direct correspondence between the mutual approximation of CFT states, [274], and $T\bar{T}$ -deformed CFTs. By virtue of the identification with S_{gen} , equation (16.6) implies the absence of a QES, and therefore of an island, consistently with the fact that the spacetimes involved in the transitions have no event horizons.
2. The direct relation between the extremised action and $S_{T\bar{T}}$ in 2D, ensures the locality of the process being described. However, as also suggested from (16.7), upon taking the flat limit on either side, the cutoff radius diverges ($r \rightarrow \infty$), and so too does the turning point, meaning the process is forbidden. We therefore conclude that, as long as the parameters of the theory are kept constant, no issue arises, and the transition remains local.
3. The cutoff nature of the cosmological horizon suggested by (16.8), proves that its contribution to S_{gen} should be understood as being part of $S_{T\bar{T}}$. The motivation that led us to connect this issue with the one encountered in the black hole information paradox, precisely resided

in the need to correctly define the dS entropy. Indeed, its divergence resembles that of the monotonic von Neumann entropy in the BH evaporation process, and, holographically, this corresponds to the standard divergence of S_{EE} upon removing the UV-cutoff. Just as in the black hole evaporation process the event horizon is not the QES extremising S_{gen} , so too can be claimed for the cosmological horizon, therefore proving the absence of islands in pure de Sitter spacetimes.

4. Localisation techniques play a key role in evaluating the partition function for a given CFT. Given its key role in determining S_{EE} , the extremisation procedure rooted in the formalism is compatible with the method defining S_{TOT} by means of the FMP method. In this section, we prove that these quantities are indeed proportional to each other.

The emergence of the island

In presence of a non-extremal black hole, the total action for $\text{AdS}_2 \rightarrow \text{AdSBH}_2$ reads:

$$S_{tot}^{\text{AdS} \rightarrow \text{AdSBH}} = \frac{2\pi\eta}{G} \left[\frac{\mathcal{B}}{2\Lambda_-} \left[1 - \sqrt{1 - \frac{4\mathcal{C}\Lambda_-}{\mathcal{B}^2}} - \ln \left| \frac{y_2}{y_1} \sqrt{\frac{1-y_1^2}{1-y_2^2}} \right| \right] + \right. \\ \left. - \phi_2 \ln \left| \frac{A_2 y_2 \sqrt{1-y_4^2}}{A_4 y_4 \sqrt{1-y_2^2}} \right| + \phi_1 \ln \left| \frac{A_2 y_1 \sqrt{1-y_3^2}}{A_4 y_3 \sqrt{1-y_1^2}} \right| + \right. \\ \left. - \frac{1}{2} \sqrt{\frac{\mathcal{B}^2}{4\Lambda_-} - \mathcal{C}} \ln \left| \frac{1-y_2}{1+y_2} \frac{1+y_1}{1-y_1} \right| + \mu_+ \ln \left| \frac{1-y_4}{1+y_4} \frac{1+y_3}{1-y_3} \right| \right],$$

where the notation is the same as in section 15. (16.10) coincides with the result by [274] for the case of an excited state of the new CFT approximating the ground state of the background theory under the following identifications

$$S_{brane} = (1-\mu)\Delta t_2 - \Delta t_1 \\ = -\frac{1}{2} \sqrt{\frac{\mathcal{B}^2}{4\Lambda_-} - \mathcal{C}} \ln \left| \frac{1-y_2}{1+y_2} \frac{1+y_1}{1-y_1} \right| + \mu_+ \ln \left| \frac{1-y_4}{1+y_4} \frac{1+y_3}{1-y_3} \right|, \\ S_{hor}^{(-)} = (1-\mu)\frac{\beta}{2} = -\frac{1}{2} \sqrt{\frac{\mathcal{B}^2}{4\Lambda_-} - \mathcal{C}},$$

with the latter, (16.10), being the 2^{nd} term featuring in the total action (16.10). The terms proportional to $\phi_{1,2}$ are $\sim \mathcal{O}(\mathcal{B})$, and so too are the first and third terms featuring in the first line of the expression for the total action. The results in [274] are in agreement with ours upon expanding the total action with respect to the deformation parameter, \mathcal{B} . As also mentioned by the authors of [274], their expression does not contain terms that vanish upon removing the cutoff, thereby explaining the missing terms in performing the comparison. We also made a consistency check with the results obtained in section 15, and found that subleading terms in the l.o. \mathcal{B} -expansion vanish in the $\lim_{\mathcal{B} \rightarrow 0}$.

The integration variables in the descriptions outlined in section 15 and [274] are related as follows

$$y \stackrel{\text{def.}}{=} \sqrt{\frac{1-\mu}{\lambda}} r. \quad (16.10)$$

In terms of which, the 3D BTZ geometry, defining an orbifold of the universal cover of the Poincaré metric, can be associated with the line element




Figure 60. By adding a black hole in the 3D bulk (LHS), instead, a turning point emerges, resulting in the ETW brane closing up and rejoining the conformal boundary. Correspondingly, an island emerges in the wedge holographic picture (RHS). This is the key difference with respect to figure 59 on the LHS, where there is no bulk turning point, and therefore, the RT surface doesn't intersect the ETW brane.

$$ds^2 = -\frac{r^2 - r_h^2}{L^2} dt^2 + \frac{L^2}{r^2 - r_h^2} dr^2 + \frac{r^2}{L^2} dx^2, \quad (16.11)$$

and becomes 2D conformally flat at the cutoff $r = r_c$, where the $T\bar{T}$ -deformed CFT of inverse temperature $\beta \stackrel{\text{def.}}{=} 2\pi L^2/r_h$ is located. Notice that $\lim_{r_h \rightarrow 0} \beta = \infty$, as expected for the $T=0$ theory. Recall that the variable y , featuring in (16.10), introduced in section 15 parametrises the value of the dilaton, ϕ_b , on the brane interpolating between the 2D vacua involved in the decay process. As previously argued, in the holographic interpretation of the process, the 1D brane is mapped to a bulk CFT_2 on which $S_{T\bar{T}}$ is evaluated.

In the same fashion, one might be led to believe that the role played by the cutoff r_c is analogous to ϕ_b , since all other parameters in the definition of y are held fixed by the formalism. However, we will now be arguing that extra care is needed in building the correspondence between the 2 setups. For $\mathcal{B} = 0$, $y = \frac{\phi_b}{\mu}$, and, due to the absence of horizons, $y_{tp} = 1$ is unique. On the other hand, in presence of a black hole, the following identification needs to be made

$$\tanh x \stackrel{\text{def.}}{=} y \Rightarrow \sinh x = \frac{r_h}{r_c} = \frac{\sqrt{1-y^2}}{y} \Rightarrow r_c \stackrel{\text{def.}}{=} r_h \frac{y}{\sqrt{1-y^2}} \xrightarrow{y \rightarrow 1} \infty, \quad (16.12)$$

from which the following information can be extracted:

1. If $\mathcal{B} = 0$, then $y_{tp} = 1$ is unique, and $r_c = \infty$. Correspondingly, (16.11) reduces to a pure AdS_3 metric, and the bulk configuration is the same as the one drawn on the LHS of figure 59.
2. As proved in section 15.3, $y_{tp} = 1$ is also the limiting value of the turning points upon taking the flat limit on either side of the 2D transition. The last passage in (16.12) thereby shows that the effect of removing the cutoff in [274] is equivalent to taking $\lim_{\mathcal{B} \rightarrow 0}$ given the identification of the turning point y_{tp} with r_c .
3. These observations imply the need to add a black hole in the 3D bulk in order to keep r_c finite, thereby ensuring the ETW brane separating the two phases can actually close up, as shown on the RHS of figure 60.

In order to provide further justification for the claims made so far, we will now show that:

- Transition in between pure (A)dS spacetimes described by (16.8) can also be obtained in presence of a suitable value of $\mathcal{B} \neq 0$, thereby justifying the identification of the black hole mass with the $T\bar{T}$ -deformation ensuring the locality of the transition taking place⁸⁷.
- For arbitrary values of \mathcal{B} , the brane action undergoes a phase transition when the black hole mass decreases below a certain critical value.

How the number of turning points depends on \mathcal{B}

For particular values of \mathcal{B} , the turning point reduces to

$$\phi_2 = \begin{cases} 0 & \text{for } \mathcal{B} = 0 & \equiv \text{ pure dS, 1 horizon, } y_2 = 0 \\ \mu_{\pm} & \text{for } \mathcal{B} = -2\sqrt{|\mathcal{C}|\Lambda} & \equiv \text{ extremal dSBH, 2 degenerate horizons, } y_2 = 1 \\ \in]0, \mu_{\pm}[& \text{for } -2\sqrt{|\mathcal{C}|\Lambda} < \mathcal{B} < 0 & \equiv \text{ non-extremal dSBH, 2 horizons, } y_2 \neq \{0, 1\}. \end{cases}$$

This shows that, in terms of the coordinate y , there is no turning point in absence of a conformal symmetry breaking term in \mathcal{L} , thereby forbidding the transition to take place. On the other hand, for $\mathcal{B}_{\text{dS}} = -2\sqrt{|\mathcal{C}|\Lambda}$, the turning point is unique, and therefore coincides with the results obtained in section 14. For any other value ranging in between, i.e. $\mathcal{B} \in]-2\sqrt{|\mathcal{C}|\Lambda}, 0[$, there are 2 physical turning points in terms of ϕ_b , consistently with the constraint derived in section 15, (15.6).

Having said this, we therefore conclude that:

- The role of \mathcal{B} is effectively equivalent to that of an irrelevant operator on a CFT₂.
- The results obtained in section 14 should be understood of as arising from the IR limit of the RG-flow of an irrelevant operator, whose imprint in the transition amplitude is given by the finiteness of the turning point at which nucleation takes place.
- The cosmological horizons and the cutoff radius at which localisation takes place are proportional to each other, implying the flat limit is equivalent to removing the cutoff altogether; this explains the divergences encountered in section 14.
- According to the value of \mathcal{B} , the nucleation process features a different number of physical turning points. Only in presence of two distinct physical values of the turning points, the flat limit can be taken (as explained in section 15.3).
- In light of further considerations made in the remainder of our work, it is important to notice that, due to the specifics of the formalism, the parameter \mathcal{B} is not simply part of the theory, but, rather specifies the state involved in the transition. This follows from the Hamiltonian constraint and the definition of the transition amplitude.

Relation to S_{gen} and phase transition of S_{brane}

The systems described in section 15 experience a phase transition as the black hole mass decreases, thereby providing further evidence of the fact that, in some cases, 2D vacuum transition can only

⁸⁷Our finding is hence in agreement with [301], where the authors show that dS black holes behave as localized, constrained, states, and therefore feature a lower entropy with respect to pure dS.

take place in presence of islands. The actions obtained in sections 14 and 15 can be recast to the difference of two generalised entropies, (16.4),

$$S_{tot}^{(A)dS \rightarrow (A)dS} = 2\pi\eta [S_{gen} - S_{gen}] = 2\pi\eta \left[\frac{\phi_{h,+}}{G} - \frac{\phi_{h,-}}{G} + S_{EE,-}^{(A)dS,TT} - S_{EE,+}^{(A)dS,TT} \right]. \quad (16.13)$$

In agreement with arguments outlined in [277], the island only emerges beyond a certain threshold value of the black hole mass, which, as previously shown, is accounted for by the parameter \mathcal{B} . Each part of the brane action is an entanglement entropy of two disjoint intervals on a CFT_2 with deformation parameter \mathcal{B} . Expanding to leading order in \mathcal{B} , both terms can be brought to the following form

$$S_{EE} = S_{EE}^{(2)} \Big|_{\mathcal{B}=0} + \delta S_{EE}^s \\ \sim -\frac{8c_-}{3} \ln \left| \frac{1}{\epsilon} \sinh \left(\frac{2\pi r_-}{\beta_-} \right) \right| - \left(1 - \frac{\phi_o^2}{\mu_-^2} \right) \frac{\mathcal{B}c_-^2}{9\Lambda_-} \coth^2 \left(\frac{2\pi r_-}{\beta_-} \right) + \frac{\frac{c_-}{3}}{\sinh^2 \left(\frac{2\pi r_-}{\beta_-} \right)}.$$

In the second line of (16.14), we redefined the parameters featuring in our original expressions in order to make the comparison with the standard CFT calculation more explicit. Notice that this interpretation is fully legitimate, since the parameter \mathcal{B} specifies the states on either vacuum being involved in the transition process.




Figure 61. A 3D plot of $S_{EE}(\mathcal{B}, x)$. x denotes the portion of the CFT_2 on which S_{EE} is evaluated. As \mathcal{B} changes, so too does the slope of (16.14), as shown on the LHS. The figure on the RHS shows the smooth interpolation between the monotonic behaviour of S_{EE} as one would expect if evaluated on the disconnected channel over 2 intervals, to a profile admitting a minimum. The phase transition occurs at $\mathcal{B} < 0$, corresponding to an upper bound to the black hole mass below which an island needs to be taken into account.

In section 15, we showed that $\mathcal{B} < 0$ and, from section 14, we know that $\phi_o^2 > \mu_-^2$, therefore $y > 0$ is the range of interest for the decay process described by (16.13). As shown on the RHS of

figure 61, the entanglement entropy, S_{EE} , experiences a phase transition, for $y > 0$. In particular, the latter takes place at $y = 1$. It is possible to argue that this is indeed consistent with our finding in section 15 that the black hole mass features in the Lagrangian density as a $T\bar{T}$ -deformation, which needs to be constrained between a certain range, to ensure the existence of two turning points, (15.5).

Indeed, figure 61 further supports our argument that, the phase transition undergone by the brane action corresponds to a Page transition beyond a suitable value of \mathcal{B} , with the latter playing a similar role as t_{Page} within the context of the information loss paradox. Consequently, this justifies the emergence of the island in (16.13) upon interpreting the total action as being the difference of generalised entropies.

An emergent holographic embedding of 2D vacuum transitions with gravity

In summary, from the holographic interpretation of the FMP results we proved that:

- The flexibility of the FMP method relies upon the fact that such kinds of *exactly integrable deformations* ($T\bar{T}$ s) can be accounted for at the level of the Lagrangian density, as explicitly outlined in section 15. Because of this, this setup can potentially accommodate an island, as long as the black hole mass lies within a certain range.
- In absence of black holes, the total action for the transition is proportional to the defect entropy, S_{ICFT} , which is codimension-2 with respect to the auxiliary AdS_3 bulk. This can also be re-expressed as a difference of S_{EE}^{TT} s, thereby ensuring the locality of the transition as well as the fact that S_{TOT} is given in terms of *internal* entanglement entropies evaluated on a gravitating bath.
- In presence of black holes, S_{brane} behaves as a quantity experiencing a phase transition beyond a certain value of \mathcal{B} , which can be thought of as playing the role of t_{Page} within the island formulation. Furthermore, given the fact that the S_{EE} s defining S_{brane} are basically describing an *internal* entanglement, due to the role of the dilaton in the given setup, our findings are compatible with the claims of [282].

Having said this, we are now able to further motivate our proposal, i.e. figure 55, as being an appropriate adaptation of figure 53 for describing vacuum transitions in presence of gravity.

- The total action for $\text{AdS}_2 \rightarrow \text{AdS}_2$ transitions calculated by means of the FMP method, is a codimension-2 quantity with respect to its AdS_3 embedding, and therefore is to be associated to the IR-limit of an RG-flow, in agreement with wedge-holography. The same picture can be drawn for transitions involving dS spacetimes, given their realisation via $T\bar{T}$ -deformations in the IR of an AdS.
- The portions of spacetime involved in the transition lie on different ETW branes, which, in turn, are holographically dual of the defects BCFT_{D-1} s. The interval in between the two defect theories now accommodates a composite CFT_D with nontrivial BCs provided by the defect theories themselves.
- Under the assumption that $c_{bdy} \gg c_{bulk}$, the RG-flow will drive the pair of defect CFT_{D-1} s in the UV to a single CFT_{D-1} in the IR. An island emerges if the resulting CFT_{D-1} alone is unable to realise the bulk AdS_3 by wedge holography.

Implications for up-tunnelling

Given the chain of relations (16.8), which in turn relies upon the parametric identification (16.7), we can now appreciate the fact that the obstructions to up-tunneling outlined in section 14 share a common origin. In particular, we notice that:

- The vanishing brane action in the flat limit, is compatible with the fact that $S_{EE}^{TT} \rightarrow 0$ as $\epsilon \equiv \frac{r}{\sqrt{c\lambda}} \rightarrow 0$, thereby further justifying (16.8).
- The divergence of the bulk action in the flat limit of dS can be recast to the need for redefining the universal part of S_{EE}^{TT} for dS including the $\frac{c}{6}$ term, which, for the decay process analysed in section 14 corresponds to μ , namely the value of the dilaton at the horizon. Given the cutoff nature of the cosmological horizon for pure dS spacetimes (due to (16.8)), it therefore follows that, upon taking the flat limit, the transition is no longer local.
- AdS₂ \rightarrow AdS₂ up-tunneling, AdS₂ \rightarrow Mink₂ and Mink₂ \rightarrow dS₂ can only be achieved as long as we are able to ensure the locality of the decay processes, or, equivalently, the finiteness of the turning point associated to them. For this to happen, the localisation radius, r , and the cosmological constant of a given vacuum need to disentangle. This can be achieved by introducing an additional scale, ultimately ensuring the finiteness of the transition under arbitrary change of the cosmological constant.
- This new scale, namely the black hole mass, is responsible for an additional contribution to the total action which cannot be reabsorbed within the definition of S_{EE}^{TT} , thereby signalling the emergence of a QES in (16.13), and proving the need for an island to be present for up-tunneling to take place.

16.2.2 A BCFT₂ perspective: transitions between end-of-the-world branes

In section 15 we proved agreement between the BT and FMP results in absence of black holes. This subsection is meant to provide further supportive evidence of their matching from a holographic point of view, showing that transitions described via the BT formalisms provide an example of an AdS₂/CFT₁ \subset AdS₃/CFT₂. In doing so, we will be focussing on AdS₂ \rightarrow AdS₂ processes, showing that the total bounce can be mapped to the entanglement entropy on a BCFT₂.

As already explained in section 13, in the BT setup, the starting point is analogous to that of a type-II Einstein-Maxwell-dilaton gravity theory in 2D, which was proved to arise from dimensional reduction of a BTZ black hole in [299].

In performing this comparison, the key quantity of interest is the boundary free energy, F_{∂} , [289], which, for a BCFT₂ reads

$$-F_{\partial} = S_{CFT_2} + S_{bdy} = \frac{T}{\sqrt{1-T^2}} - \ln \left| \frac{1-T}{1+T} \right|, \quad (16.14)$$

where T is the tension of the ETW brane and is related to its opening angle with respect to the Cardy brane as $\Theta \stackrel{def.}{=} \tan^{-1} T$, as depicted on the LHS of figure 63.

For type-1 instantons, (16.16) is nontrivial for both vacua, i.e. $T, \Theta \neq 0$, and the total bounce can be re-expressed as follows:

$$B_{type-1} = 2\pi S_{BCFT_2} = 2\pi \left[S_{CFT_2} + S_{bdy}^- - S_{bdy}^+ \right], \quad (16.15)$$

where




Figure 62. The CFT_1 s are dual to the AdS_2 spacetimes involved in the transition living on the two ETW branes and the wedges of opening angles $\Theta_{1,2}$. Their corresponding tensions, result in the value of Λ_{\pm} on the two sides of the wall.

$$r_o^{-1} \stackrel{def.}{=} \bar{\rho} \quad , \quad r_H \stackrel{def.}{=} \sqrt{\Lambda_{o,I}} \quad , \quad T \stackrel{def.}{=} \sqrt{1 - \frac{r_H^2}{r_o^2}} \quad , \quad S_{bdy} = -2 \ln \left| \frac{1 - \sqrt{1 - \Lambda \bar{\rho}^2}}{\sqrt{\Lambda} \bar{\rho}} \right| .$$

Equation (16.15) indicates that either vacuum can be associated to a defect CFT_1 living at the endpoints of a bulk CFT_2 . The overall system therefore defines a $BCFT_2$ as depicted on the RHS of figure 62, with S_{bdy} being evaluated on the lower-dimensional defect CFT s. In particular, there are two different values of T , one for each vacuum. The mismatch in between the two gives rise to the holographic central charge, which counts the degrees of freedom of the ground state of the codimension-2 theory living on the boundary.

16.2.3 An internal CFT_2

We now turn to the holographic interpretation of the results obtained in section 13 by means of the CDL method, further supporting the emergent analytic complementarity between the CDL and BT formalisms in 2D. In doing so, two main features will become manifest:

1. The entanglement along the dilatonic direction is effectively *internal*.
2. The CDL method (in 2D) describes vacuum decays in *absence* of gravity.

A CFT_2 is uniquely defined by its central charge and the conformal dimensions of the operators living in the theory, (c, Δ_i) . Combined together, they define the energy spectrum of the states of a given CFT .

From the fundamental parameters of the CFT_2 , the entropy of an excited state is defined in terms of the *Cardy formula*,

$$S_{Cardy} = 2\pi \sqrt{\frac{c}{6} \left(\Delta - \frac{c}{24} \right)} , \quad (16.16)$$

where Δ and $\Delta_o = \frac{c}{24}$ denote the black hole and ground state energies, respectively.

For the case of a CFT_2 where the degeneracy of the ground state is 1, the following definitions hold

$$S_{CFT_2} = \frac{c}{3} \sqrt{\mu - 1} \quad , \quad E_o = \frac{c}{12} \mu \quad , \quad c = 6 \frac{\partial S_{CFT}^2}{\partial E_o} , \quad (16.17)$$

with $\mu \stackrel{def.}{=} \frac{\Delta}{c}$ being related to the black hole mass in the excited state. Focussing on the $AdS_2 \rightarrow AdS_2$ process, with associated extremised bounce that can be rewritten as follows

$$B_{tot}^{\text{AdS}_2 \rightarrow \text{AdS}_2} = 4\pi S_{\pm}^2 = 4\pi c_{hol}^{CFT_2} M_{BH}, \quad (16.18)$$

$$c_{hol}^{CFT_2} \stackrel{\text{def.}}{=} \phi_-^2 + 2\phi_+^2, \quad M_{BH} \stackrel{\text{def.}}{=} \frac{\phi_-^2}{\phi_+^2} - 1, \quad S_{\pm} \stackrel{\text{def.}}{=} \sqrt{\phi_-^2 + 2\phi_+^2} \sqrt{\frac{\phi_-^2}{\phi_+^2} - 1}, \quad (16.19)$$

with M_{BH} being a conserved charge along the flow and $c_{hol}^{CFT_2}$ its conjugate chemical potential. The last passage in (16.18) follows from the relation between the entropy product $S_+ S_-$ and the central charge. Our result (16.19) goes beyond the original formulation of c by Cardy because it is associated to the defect lying in between the two vacua.

A key feature of (16.18) is the fact that it defines an entropy product, and (apparently) not a difference in between entropies, as one might have expected to start with. The main reason for this is that the background configuration in CDL is providing a reference state, the *ground state* of the CFT_2 , with respect to which the newly-nucleated spacetime behaves as an excited state, namely an extremal black hole. Comparison with (16.17) enables us to deduce that

$$\boxed{\frac{\phi_-^2}{\phi_+^2} \stackrel{\text{def.}}{=} \mu}, \quad (16.20)$$

indicating that the newly nucleated vacuum is to be identified with an extremal black hole spacetime embedded in the background vacuum. Extremality follows from the fact that the mass of the BH is related to the cosmological constant itself. Indeed, from (16.19), M_{BH} is a function of Λ_{\pm} .

Relying upon arguments inspired by black hole microstate cosmology, [302], a further remark is in order. We have shown that transitions of the kind $\text{AdS}_2 \rightarrow \text{Mink}_2$ are forbidden. This is perfectly in agreement with the identification we have just made, namely (16.20), since, from the CFT point of view, we can think of Mink_2 as being obtained from $\lim_{c \rightarrow \infty}$ of a CFT_2 . The nucleation process would therefore correspond to the ground state of a CFT_2 insertion attempting to mimic an excited state of the background CFT_2 . However, by definition, there is no horizon in the new vacuum that could attempt to screen the infinite number of degrees of freedom associated to a divergent central charge, and therefore the relative entropy (16.17) cannot be defined. In conclusion, due to the extremality nature of the spacetime suggested by (16.20), such transition is forbidden, since it corresponds to an overcounting of the microstates defining the entropy.

Final remarks

To finish this section we briefly collect some relevant points and mention the potential relation of our results with other recent developments.

- In summary, in absence of black holes, the three methods lead to the following results in 2D

$$\begin{cases} B_{CDL/AP}^{wo.kt} = 2\pi S_{CFT_2}^2 = 2\pi c_{hol}^{CFT_2} M_{BH} \\ S_{FMP}^{\mathcal{B}=0} = B_{BT} = B_{CDL/AP}^{w.kt} + M_{+}^{-} = 2\pi S_{BCFT_2} = 2\pi [S_{CFT_2} + S_{bdy}^{-} - S_{bdy}^{+}] \end{cases}, \quad (16.21)$$

where $B_{CDL/AP}^{wo.kt}$ and $B_{CDL/AP}^{w.kt}$ denote the bounces calculated in absence and in presence of the kinetic term for ϕ , respectively, whereas

$$S_{bdy} = \ln \langle 0|\mathcal{B} \rangle \stackrel{def.}{=} \ln g = \ln \left| \frac{1+T}{1-T} \right|, \quad (16.22)$$

in turn highlighting the consistency and mutual complementarity of their holographic interpretations.

- The identifications featuring in (16.21), signal the emergence of an *entropic hierarchy*, resulting in an interesting correspondence between our findings and those of [215, 216] within the realm of von Neumann algebras (VNAs)⁸⁸. In particular, the fact that the CDL calculation describes transitions in absence of gravity and the total bounce can be recast in the form of a relative entropy, is compatible with the fact that a type-III VNA only admits a relative entropy definition, rather than a von Neumann entropy. Turning on gravity, namely going beyond large- N , we have the BT and FMP results (in absence of black holes), therefore corresponding to type-II _{∞} VNAs. Upon adding a black hole, instead, which at the level of the Lagrangian density corresponds to turning on an irrelevant deformation, the UV theory is different with respect to that of the case with $\mathcal{B} = 0$, leading us to conclude that the projection operator ensuring the reduction from type-II _{∞} to type-II₁ is mapped to the choice of the deformation in the FMP setup, hence to the choice of the black hole mass.
- Vacuum transitions are defined in terms of amplitudes, and we have gathered supportive evidence throughout this work that such processes actually resemble a statement of mutual approximation of CFT states, [274]. We emphasised that the emergence of the island can only be achieved in presence of non-extremal black holes, and that, because of this, upon taking the flat limit, the corresponding action remains finite and experiences a phase transition beyond a critical value of the black hole mass. Rephrasing this in the language [276, 277], the flat limit corresponds to the large- N limit of the dual gravitational theory at hand. Inspired by the arguments of [277], regarding the importance of scale separation for reconciling connected amplitudes from disconnected boundaries (CADBs) with holography, we therefore claim that, below a critical value of the black hole mass, the flat limit leads to a divergent action due to the lack of scale separation between the cosmological constant and the black hole mass.

17 Conclusions and Outlook of Part IV

Part IV of the present work was devoted to understanding the subject of vacuum transitions in the simple setting of 2-dimensional gravity. Due to the peculiarities of the latter, the general analysis is not a straightforward extension of the 4D cases. Our main results are the calculation of transition amplitudes obtained following three known approaches, namely the two Euclidean methods of CDL and BT, and the Lorentzian one of FMP. We presented explicit expressions for the transition rates in each method, and for the different signs of the vacuum energies, and took the first steps towards understanding the results. Interestingly, in comparing the Euclidean and Lorentzian prescriptions, we find many similarities, as well as differences.

Specifying to the case of (A)dS₂ \rightarrow (A)dS₂ transitions, we derived the corresponding expressions for the total bounce or action in the AP setup following the CDL method in thick, (13.44), and thin-wall approximation, (13.56), the BT, (13.71), and FMP method without black holes, (14.31). As emphasised in section 14, the BT and FMP results agree under suitable parametric redefinition, thereby proving expected agreement in between the 2 methods. Such redefinition, though, is highly

⁸⁸See also [591] for an alternative perspective in the study of the Hilbert space of dS in arbitrary dimensions, mostly motivated by the bootstrap program. We thank the anonymous referee for bringing this article to our attention.

nontrivial, and can be recast to the different way in which the cosmological constants are being defined in the 2 setups, which in turn is due to the difference in between the Lagrangian densities defining the 2 theories.

On the other hand, the result obtained by means of the CDL setup in the thin-wall approximation, (13.56), is equivalent to part of the BT result. In particular, all terms can be identified (under suitable redefinition), apart from the ones arising from the boundary terms associated to the ADM mass. As also argued in section 13, these provide a nontrivial contribution to the total bounce upon turning on gravity. However, as a consequence of the particular nature of the 2D theory, the equation of motion for the AP setup (when removing the kinetic term for the dilaton from the Lagrangian density), imply $M_{ADM} = 0$. Therefore, 2D transitions described by means of the thin-wall approximated CDL method, should really be interpreted as describing nucleation processes in absence of gravity.




Figure 63. The 3 holographic descriptions are complementary to each other.

Furthermore, the thick-wall analysis leads to an expression for B_{TOT} with unique features, in the sense that there seems to be no suitable reparametrisation enabling to recast it in any of the expressions obtained by means of other procedures. However, as argued in section 14, under suitable parametric redefinitions, we can still provide a holographic interpretation of our result, which is complementarily in agreement with the others.

We may summarise the holographic interpretation of our results as follows:

- In section 16 we provide a possible holographic interpretation of the total bounces and actions calculated throughout our work, showing mutual compatibility and complementarity. Figure 63 shows how these processes can be understood as taking place in a KR/HM setup with two ETW branes and one *internal* gravitating bath, geometrising the brane mediating the nucleation process. In particular, we find that the corresponding expression for the transition rate in presence of gravity, and in absence of black holes, is given by the difference of entropies of $T\bar{T}$ -deformed CFTs, hence illustrating the *locality* of the nucleation process.
- Upon adding black holes of suitable mass, (15.5), instead, the total action can be expressed as the difference of generalised entropies, with an island emerging beyond a critical value of the black hole mass. Furthermore, under suitable parametric redefinition, the results obtained by means of the FMP method are found to agree with the expressions derived in [274] for describing mutual approximation of boundary states belonging to different CFTs.
- We propose that the vacuum transitions in the bulk correspond to a deconfinement/confinement transition on the CFT side although further study needs to be made towards this correspondence.
- The BT results, which were shown to be equivalent to those obtained through the FMP method in absence of black holes, can be expressed in terms of entropies of BCFT₂'s with

2 nontrivial boundary conditions dual to ETW branes. Furthermore, the CDL result can be recast in the form of an entropy product of a CFT_2 thereby showing agreement with the expectations following from the analytic behaviour encountered in section 13.

- One of our results is that the total action (or bounce) associated to the decay process carries an *internal* entropic interpretation. In particular, for the BT and FMP cases, they can always be expressed as the difference of generalised entropies. Only the latter, however, provides the right setup for an island to emerge. In particular, the black hole mass, which in 2D is an explicit parameter of the theory, is, in a way, playing the role of time in the Hawking evaporation process, and is responsible for S_{brane} undergoing a phase transition, as explained in section J.

Understanding the behaviour of the same physical process by means of different formalisms is equally important in all fields of research, and appears to be particularly promising within the context of quantum gravity. Recent progress towards understanding the behaviour of the partition function for QG within, either, the canonical or microcanonical ensemble, [280, 281], as well as the study of the phases of gauge theories and their holographic duals, [306], are certainly among the most promising directions deserving more investigation. Clearly our work leaves many open questions. We hope that our results will be useful for addressing further questions related to vacuum transitions, early universe cosmology, and holography.

Understanding the behaviour of the same physical process by means of different formalisms is equally important in all fields of research, and appears to be particularly promising within the context of quantum gravity. Recent progress towards understanding the behaviour of the partition function for QG within, either, the canonical or microcanonical ensemble, [280, 281], as well as the study of the phases of gauge theories and their holographic duals, [306], are certainly among the most promising directions deserving more investigation. We hope the results found through this investigation will be useful for addressing further questions related to vacuum transitions, early universe cosmology, and holography.

Part V

Beyond Categorical Dualisability

18 Summary and Motivations of Part V

18.1 Motivations

For a theory, \mathcal{T} , to be absolute, the following triple needs to be defined

$$\boxed{\mathcal{T} \longleftrightarrow (\mathcal{F}, \mu, \mathfrak{Z})}, \quad (18.1)$$

where \mathcal{F} is the fiber functor, μ the moment map, and \mathfrak{Z} the Drinfeld center⁸⁹. Consistency of the underlying mathematical structure requires these three quantities to be mutually related. Indeed, upon defining any one of them, the other two should automatically follow. The purpose of [1, 3] and the present work is to show that an apparent shortcoming in defining such triple corresponds to the emergence of interesting physics, rather than being a fault of the sought after absolute theory. In

⁸⁹We refer to section 9 for a detailed explanation of this terminology.

particular, we will show that this can be used to explain the emergence of non-invertible symmetries separating different class \mathcal{S} theories, [1, 366–371].

The crucial references we rely upon are the works of Moore and Segal, [137], and Moore and Tachikawa, [136]. As briefly reviewed in the following sections, [136] proposes a redefinition of class \mathcal{S} theories (cf. figure 14) in terms of a 2D TFT, namely the functor

$$\eta_{G_c} : \text{Bo}_2 \rightarrow \text{HS} \quad (18.2)$$

with Bo_2 and HS denoting the bordism 2-category and the holomorphic symplectic 2-category, respectively⁹⁰, associated to a given 4D $\mathcal{N} = 2$ SCFT. The definition of (18.2) strongly relies upon assuming, both, the source and target categories, enjoy a duality structure which, in turn follows from the presence of an identity element in both categories. In [136], the authors show that, under the duality assumption, for the categories in (18.2) to be well-defined, it is enough to specify their objects and 1-morphisms. Essentially, the objects of Bo_2 are circles, S^1 , and the 1-morphisms are cobordisms between different disjoint unions of circles and the empty set. Their respective counterpart on the holomorphic symplectic side correspond to the gauge group, [136],

$$\eta_{G_c} (S^1) \stackrel{\text{def.}}{=} G_c, \quad (18.3)$$

and the cobordism operators, [136],

$$\eta_{G_c} (\text{Hom} (S^1, \emptyset)) \stackrel{\text{def.}}{=} U_{G_c} \quad (18.4)$$

$$\eta_{G_c} (\text{Hom} (S^1 \sqcup S^1, \emptyset)) \stackrel{\text{def.}}{=} V_{G_c} \quad (18.5)$$

$$\eta_{G_c} (\text{Hom} (S^1 \sqcup S^1 \sqcup S^1, \emptyset)) \stackrel{\text{def.}}{=} W_{G_c}. \quad (18.6)$$

respectively.

Importantly for us, the Moore-Tachikawa varieties described by (18.3), (18.4), (18.5), and (18.6) constitute the quiver gauge theory⁹¹ realisation of [137], where Moore and Segal propose the mathematical formalism needed for addressing the following question: given a certain closed string theory background, what is its corresponding D-brane content⁹²?

Our aim is that of explaining how and why one needs to generalise the construction of [136] from a higher-categorical point of view, and what its implications are on the theoretical physics side. In doing so, we highlight the crucial properties and axioms satisfied by the cobordism operators outlined in [136, 137], and how they can be generalised to account for more general setups from, both, the mathematical and theoretical physics perspectives.

In particular, we emphasise the dependence of (18.2) on the conformal structure of the Riemann surface on which the compactification of the 6D $\mathcal{N} = (2, 0)$ SCFT has been performed to achieve a certain class \mathcal{S} theory and how lack of reparametrisation invariance, corresponding to the absence of the identity element in its source and target categories, [136], signals the presence of (non-invertible) categorical symmetries separating different absolute theories.

At the heart of this is the correspondence sketched in figure 64.

In [1] we explained how gauging a Symmetry Topological Field Theory (SymTFT) enables to change the boundary conditions of the fields living in the absolute field theory resulting from the

⁹⁰For more details, we refer the reader to section 4.1.

⁹¹Among the cobordism operators is the Higgs branch of class \mathcal{S} theories, (18.6).

⁹²Note that this is essentially the same question addressed in [372].




Figure 64. Adaptation of a correspondence first proposed in [1] playing a key role towards generalising [136] to the hyperkähler target category case. As explained in the following sections, this also requires the generalisation of cobordism operators, [137], due to the lack of reparametrisation-invariance on the Riemann surface on which the compactification of the 6D $\mathcal{N} = (2, 0)$ SCFT is performed.

Freed-Moore-Teleman construction, [102, 103, 374, 607]. To each gauging corresponds a choice of triples, (18.2), and, for any absolute theory, it is enough to define one of the three entries on the RHS of (18.2) to determine the other two. For the purpose of this article, we will mostly focus on the second, namely the moment map, defined as follows

$$\mu : \mathfrak{G} \rightarrow \mathcal{A}, \quad (18.7)$$

where \mathfrak{G} is an n -categorical structure, and \mathcal{A} is the algebra of invertible topological defects associated to the action $\mu(\mathfrak{G})$. Gauging means taking the categorical quotient with respect to \mathfrak{G} (in notation $//_{\mu} \mathfrak{G}$), and projecting its image under μ to the identity⁹³. Practically, one could perform a total gauging of the theory by choosing \mathcal{A} such that the overall spectrum of the theory of the gauged theory is only the (new) identity element. For the purpose of our work, instead, we are interested in understanding mathematical structures arising by gauging with respect to different subalgebras within \mathcal{A} , that are mutually intersecting, albeit not contained within each other. The ultimate aim is that of explaining the emergence of (non-invertible) categorical symmetries in certain supersymmetric quiver gauge theories once described in terms of Coulomb branches of magnetic quivers of 3D $\mathcal{N} = 4$ quiver gauge theories, which is the main focus of an upcoming paper by the same author, [597]. In such analysis we will be applying some of the findings of [377, 598].

In [3], we argued in favour of the following statements:

- Lack of reparametrisation-invariance of bordism operators signals the presence of intrinsic non-invertible defects separating different class \mathcal{S} theories.
- Defining the Drinfeld center for a system of composite class \mathcal{S} theories separated by non-invertible defects constitutes a nontrivial generalisation of an S-duality statement.

Mostly relying upon [137], we highlighted the importance of the construction of cobordism operators, emphasising their dependence on the conformal structure of the Riemann surface. We then turned to the discussion of a particular 2D TFT valued in a symmetric monoidal category, namely the maximal dimension Higgs branch of class \mathcal{S} theories. After briefly reviewing the properties outlined in [136], we proposed their generalisation for the case in which the target category of the η_{G_c} functor is a hyperkähler quotient. We concluded outlining the possible extension of this

⁹³We thank Nathan Seiberg for instructive discussion regarding the appropriateness of the terminology to be adopted in describing this formalism.

treatment towards a mathematical formulation of magnetic quivers within the context of Coulomb branches of 3D $\mathcal{N} = 4$ quiver gauge theories.

In this new work, we take a more algebro-geometric approach in explaining the mathematical structure underlying supersymmetric quiver gauge theories, with a particular focus on dualities and their mutual relations in terms of higher-categories.

As will be explained in depth, the crucial role is played by factorisation homology. We will not attempt to provide an exhaustive definition here, and refer the reader to the related section in the core of this work, instead. What we can anticipate, though, is that factorisation homology is related with the counting of ground state degeneracy, and, when applied to the AGT correspondence, provides a tool for evaluating the 4-sphere partition function of class \mathcal{S} theories. Specifically, as we shall see, factorisation homology allows us to relate categorical dualisability and full-extendibility⁹⁴. Within the context of non-Lagrangian theories, such as 6D $\mathcal{N} = (2, 0)$ SCFTs, this is particularly relevant since it introduces the notion of ring objects, of which Hilbert series are one of the examples of most interest to us.

As an algebraic variety, we focus on the calculation of the Hilbert series on Coulomb and Higgs branches, with the latter being related by the geometric invariant theory (GIT) construction, [613]. GIT being a recipe for realising 3D mirror symmetry, [146], we show how it can be suitably extended for cases where mirror duals are not necessarily in a 1-to-1 correspondence. In relation to Part II, this nicely fits with the magnetic quiver prescription for describing the Higgs branch of certain supersymmetric quiver gauge theories with non-primitive ideals⁹⁵.

We then show how the 2D description of disk algebras can be suitably adopted for describing Hitchin systems associated to joint class \mathcal{S} theories. We conclude outlining future directions that are currently under investigation by the same author.

Following our treatment in [3] we explain the higher categorical structures that are needed for describing the invariants associated to specific supersymmetric quiver gauge theories admitting a magnetic quiver description.

The key tool we will be using in achieving such aim is factorisation homology, of which a more detailed definition will be provided in the core of the paper. For the purpose of the introduction, we might just say that its role is that of assigning invariants to topological manifolds.

18.2 Outline and main results

Part V of this work is structured as follows:

1. section 19 introduces factorisation homology and its applicability within the context of class \mathcal{S} theories by means of the AGT correspondence. The section ends recapitulating the interplay between dualisability and full-extendibility of TQFTs, with a particular focus of Moore-Tachikawa varieties⁹⁶.
2. Section 20 explains the relation in between the Drinfeld center and magnetic quivers, combining sections 5 and 9.
3. In section 21 we propose the generalisation of Moore-Tachikawa varieties for the case in which the target category of the η_{G_c} functor is a hyperkähler quotient. We conclude outlining the

⁹⁴And is therefore in line with the connection between dualities and categorical structures.

⁹⁵Namely the 2-cone structure, [223].

⁹⁶In section 5, we briefly overviewed cochain level theories as the most important generalisation of the open and closed TFT construction. Mostly relying upon [137], we highlighted the importance of the construction of the cobordism operator, highlighting their dependence on the conformal structure of the Riemann surface. We then turn to the discussion of a particular 2D TFT valued in a symmetric monoidal category, namely the maximal dimension Higgs branch of class \mathcal{S} theories.

possible extension of this treatment towards a mathematical formulation of magnetic quivers within the context of Coulomb branches of 3D $\mathcal{N} = 4$ quiver gauge theories which will be addressed in an upcoming work by the same author, [597]. The concluding part of this section recapitulates the essential tools required for performing GIT quotient constructions, highlighting its importance for realising 3D mirror symmetry. In particular, we focus on the calculation of the Hilbert series as specific examples of algebraic varieties of crucial interest in the study of categorical dualities for supersymmetric quiver gauge theories. We conclude showing how Coulomb and Higgs branches from class \mathcal{S} Hitchin systems carry the same information as the gaugeable algebras leading to SymTFT constructions, discussed in Part III. We conclude outlining interesting open questions and future directions of investigation by the author.

19 Categorical Dualisability and full extendibility

The present section is structured as follows:

1. Section 19.1 is devoted to explaining the role played by factorisation homology in describing the invariants of a given TFT. In particular, this prepares the stage for the following sections, where we will be dealing with the applicability of this formalism to the specific case of class \mathcal{S} theories⁹⁷.
2. In section 19.2, we then turn to explaining the relation between categorical dualisability and full-extendibility. In particular, this nicely relates to the notion of Drinfeld centers and gauging categorical structures.

19.1 Factorisation Homology in context

The present section is mostly revision of well-known literature⁹⁸. It provides a brief mathematical digression, needed for introducing factorisation homology, and its utility for evaluating the index of a 2D TQFT. In doing so, we explain the notions of Hochschild cohomology, stratified spaces, and disk algebras. In the following sections we will see that, combining this formalism to algebro-geometric techniques, leads to interesting new findings.

Cohomology and Homotopy Equivalence

In homology theory and algebraic topology, cohomology is a general term for a sequence of abelian groups, usually one associated with a topological space, often defined from a cochain⁹⁹ complex. Cohomology can be viewed as a way of assigning richer algebraic invariants to a space with respect to homology.

Some versions of cohomology arise by dualising the homological construction. Some of their properties are formulated in a similar fashion with respect to their homological counterpart. However, cohomology enjoys a richer structure, which is lacking in homology, which we now turn to explain.

For a given topological space X , and a commutative ring, R , the cup product is a bilinear map

$$H^i(X, R) \times H^j(X, R) \rightarrow H^{i+j}(X, R). \quad (19.1)$$

⁹⁷Note the importance of having introduced the notion of contractible cells in section 3 in relation to abelianisation and equivariant cohomology.

⁹⁸Specifically, we suggest the interested reader to consult the following reference [137].

⁹⁹Cochains are functions on the group of chains in the homology theory.

This product defines the cohomology ring of X as follows

$$H^\bullet(X, R) \equiv \bigoplus_i H^i(X, R). \quad (19.2)$$

It is graded commutative in the sense that, given $u \in H^i(X, R)$, $v \in H^j(X, R)$,

$$uv = (-1)^{ij} vu. \quad (19.3)$$

It follows that, if two spaces are homotopy equivalent, their cohomology rings are isomorphic.

Homotopy equivalence can be expressed in terms of looping by representation spheres. In particular, a homotopy equivalence between topological spaces X and Y is a morphism $f : X \rightarrow Y$ which has a homotopy inverse, hence such that there exists a morphism $g : Y \rightarrow X$ and homotopies

$$g \circ f \sim \mathbf{1}_X \quad \text{and} \quad f \circ g \sim \mathbf{1}_Y, \quad (19.4)$$

thereby making homotopy equivalence very similar to the notion of categorical equivalence.

For a given topological space, X , elements of $H^i(X)$ can be thought of as represented by codimension- i subspaces of X that can move freely on X . For the case at hand, an element of $H^i(X)$ is defined by a continuous map

$$f : X \rightarrow M, \quad (19.5)$$

with pullback

$$f^* : H^\bullet(M, R) \rightarrow H^\bullet(X, R), \quad (19.6)$$

and a closed codimension- i submanifold N of M with an orientation in the normal bundle. As such, the class $f^*([N]) \in H^i(X)$ can be thought of to lie in the subspace $f^{-1}(N)$ of X , since $f^*([N])$ restricts to 0 in the cohomology of the open subset $X - f^{-1}(N)$. The cohomology class $f^*([N])$ is free to move in X in the sense that N could be replaced by any continuous deformation of N inside M .

19.1.1 Hochschild cohomology and cochain complexes

Given a finite group action on a suitably enhanced triangulated category linear over a field, [385], Hochschild cohomology and group actions establish a formula for the Hochschild cohomology of the category of invariants, assuming the order of the group is coprime to the characteristic of the base field.

For a k -linear category with G -action,

$$\bigoplus_{g \in G} \phi_g : \mathcal{C} \rightarrow \mathcal{C} \quad (19.7)$$

induces the norm functor

$$N_m : \mathcal{C}_G \rightarrow \mathcal{C}^G \quad (19.8)$$

$$x \mapsto \sum_{g \in G} g(x)$$

characterised by the existence of a factorisation property

$$N_m \circ q : \mathcal{C} \rightarrow \mathcal{C}^G. \quad (19.9)$$

Importantly, N_m is an equivalence.

If \mathcal{C}, \mathcal{D} are k -linear categories, then the k -linear exact functors from \mathcal{C} to \mathcal{D} form the objects of the k -linear category $\text{Fun}_k(\mathcal{C}, \mathcal{D})$.

Let \mathcal{C} be a k -linear category, and let $\phi : \mathcal{C} \rightarrow \mathcal{C}$ be an endofunctor. The Hochschild cochain complex of \mathcal{C} with coefficients in ϕ is defined as

$$HC^\bullet(\mathcal{C}, \phi) \stackrel{\text{def.}}{=} \text{Map}_{\text{Fun}_k(\mathcal{C}, \mathcal{C})}(\mathbf{1}_{\mathcal{C}}, \phi) \in \text{Vect}_k. \quad (19.10)$$

The Hochschild cohomology $HH^\bullet(\mathcal{C}, \phi)$ of \mathcal{C} with coefficients in ϕ is the cohomology of this complex. For the case in which $\phi \equiv \mathbf{1}_{\mathcal{C}}$,

$$HC^\bullet(\mathcal{C}) \stackrel{\text{def.}}{=} HC^\bullet(\mathcal{C}, \mathbf{1}_{\mathcal{C}}) \quad , \quad HH^\bullet(\mathcal{C}) \stackrel{\text{def.}}{=} HH^\bullet(\mathcal{C}, \mathbf{1}_{\mathcal{C}}) \quad (19.11)$$

define the Hochschild cochain complex and the Hochschild cohomology of \mathcal{C} , respectively. The latter comes equipped with a natural algebra structure, induced by composition in $\text{Map}_{\text{Fun}_k(\mathcal{C}, \mathcal{C})}(\mathbf{1}_{\mathcal{C}}, \mathbf{1}_{\mathcal{C}})$.

Note that, given a k -linear category \mathcal{C} with an action by a finite group G , there is an induced action of $G \times G$ on $\text{Fun}_k(\mathcal{C}, \mathcal{C})$ consisting in the following. An element $(g_1, g_2) \in G \times G$ acts on $\text{Fun}_k(\mathcal{C}, \mathcal{C})$ by sending $F : \mathcal{C} \rightarrow \mathcal{C} \in \text{Fun}_k(\mathcal{C}, \mathcal{C})$ to $\phi_{g_2} \circ F \circ \phi_{g_1}^{-1}$. Via the diagonal embedding $G \subset G \times G$, this restricts to the conjugation action of G on $\text{Fun}_k(\mathcal{C}, \mathcal{C})$.

A main theorem in [385] states that, given \mathcal{C} a k -linear category with an action by a finite group G , and assuming the order of G is coprime to the characteristic of K , then there is an isomorphism

$$HH^\bullet(\mathcal{C}^G) \simeq \left(\bigoplus_{g \in G} HH^\bullet(\mathcal{C}, \phi_g) \right)^G, \quad (19.12)$$

with $\phi_g : \mathcal{C} \rightarrow \mathcal{C}$ an autoequivalence corresponding to $g \in G$, and the G -action on the right side induced by the conjugation action of G on $\text{Fun}_k(\mathcal{C}, \mathcal{C})$. In particular,

$$HH^\bullet(\mathcal{C})^G \hookrightarrow HH^\bullet(\mathcal{C}^G). \quad (19.13)$$

We will come back to this in section 21 when discussing the GIT quotient and the Kirwan map.

19.1.2 Factorisation homology of stratified spaces

Factorisation homology satisfies a defining property called \otimes -excision, which determines $\int_M \mathcal{A}$ up to equivalence,

$$\boxed{\int_M \mathcal{A} \simeq (H, u_\Sigma)}, \quad (19.14)$$

with such property being a special case of the pushforward property. The RHS of (19.14) consists of the Hilbert space of the TO, H , and the ground state degeneracy, u_Σ . (19.14) allows to reduce the computation of the factorisation homology of a surface to that of a lower-dimensional manifold, and, eventually, to that of a 0-dimensional manifold.

The only known global observable on Σ for an anomaly-free TO is the ground state degeneracy. The theory of factorisation homology gives a powerful tool for calculating it. When a closed stratified surface is decorated by anomaly-free topological defects of codimension 0,1,2, the theory of factorisation homology gives us a powerful tool for calculating the degeneracy of the ground state. As an important bi-product, FH provides a powerful tool for calculating global observables on surfaces for anomalous TOs. Indeed, relation between FH and TOs enables to make direct contact with the theoretical physics setup of interest to us.

There is a well-defined fully faithful functor

$$\mathfrak{Z} : \text{UMTC}^{\text{ind}} \rightarrow \text{UMTC} \quad (19.15)$$

acting on objects and morphisms as follows

$$\mathcal{C} \rightarrow \mathfrak{Z}(\mathcal{C}) \quad , \quad {}_c\mathcal{M}_{\mathcal{D}} \mapsto \mathfrak{Z}(\mathcal{M}) \stackrel{\text{def.}}{=} \text{Fun}_{c|\mathcal{D}}(\mathcal{M}, \mathcal{M}). \quad (19.16)$$

For \mathcal{C} a UMTC, (19.15) defines the Drinfeld center of \mathcal{C} , $\mathfrak{Z}(\mathcal{C})$, a canonical braided monoidal category such that

$$\mathfrak{Z}(\mathcal{C}) \simeq \mathcal{C} \boxtimes \bar{\mathcal{C}}. \quad (19.17)$$

More explicitly, Th.2.5 in [374] implies that, given $\mathcal{C}, \mathcal{D}, \mathcal{E}$ UMFCs, \mathcal{M} a multi-fusion $\mathcal{C} - \mathcal{D}$ -bimodule, and \mathcal{N} a multi-fusion $\mathcal{D} - \mathcal{E}$ -bimodule, the assignment

$$f \boxtimes_{\mathfrak{Z}(\mathcal{D})} g \mapsto f \boxtimes_{\mathcal{D}} g \quad (19.18)$$

defines an equivalence between two multi-fusion $\mathfrak{Z}(\mathcal{C}) - \mathfrak{Z}(\mathcal{E})$ -bimodules:

$$\text{Fun}_{c|\mathcal{D}}(\mathcal{M}, \mathcal{M}) \boxtimes_{\mathfrak{Z}(\mathcal{D})} \text{Fun}_{\mathcal{D}|\mathcal{E}}(\mathcal{N}, \mathcal{N}) \simeq \text{Fun}_{c|\mathcal{E}}(\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N}, \mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N}) \quad (19.19)$$

from which the following two corollaries follow:

$$\mathcal{C} \boxtimes_{\mathfrak{Z}(\mathcal{C})} \mathcal{D}^{\text{rev}} \simeq \text{Fun}_H(\mathcal{M}, \mathcal{M}) \quad (19.20)$$

$$\boxtimes_{(c_0, \dots, c_{n-1})}^{\circ}(\mathcal{M}_1, \dots, \mathcal{M}_n) \simeq \text{Fun}_H(\mathcal{P}, \mathcal{P}), \quad (19.21)$$

with \mathcal{P} a unique unitary category such that this correspondence holds.

Next, consider $\text{Mfld}_n^{\text{or}}$ to be the topological category whose objects are oriented n -manifolds whose objects are oriented n -manifolds without boundary. For any two oriented manifolds M, N , the morphism $\text{Hom}_{\text{Mfld}_n^{\text{or}}}(M, N)$ is the space of all orientation-preserving embeddings $e : M \rightarrow N$ endowed with the compact-open topology.

Denoting with $\mathcal{M}\text{fld}_n^{\text{or}}$ the symmetric monoidal ∞ -category associated to the topological category $\text{Mfld}_n^{\text{or}}$, the symmetric monoidal ∞ -category $\text{Disk}_n^{\text{or}}$, is the full subcategory of $\mathcal{M}\text{fld}_n^{\text{or}}$ whose objects are disjoint union of finitely-many n -dimensional Euclidean spaces equipped with standard orientation.

If \mathcal{V} is a symmetric monoidal ∞ -category, M an oriented manifold, and

$$\boxed{\mathcal{A} : \text{Disk}_n^{\text{or}} \rightarrow \mathcal{V}} \quad (19.22)$$

an n -disk algebra in \mathcal{V} , the factorisation homology of M with coefficients in \mathcal{A} is an object of \mathcal{V} given by

$$\int_M \mathcal{A} \stackrel{\text{def.}}{=} \text{Colim.} \left(\left(\text{Disk}_n^{\text{or}} \right)_{/M} \rightarrow \text{Disk}_n^{\text{or}} \xrightarrow{\mathcal{A}} \mathcal{A} \right), \quad (19.23)$$

with $\left(\text{Disk}_n^{\text{or}} \right)_{/M}$ denoting the category of n -disks embedded in M . If $\mathbf{1}_n$ is the trivial n -disk algebra assigning to each \mathbf{R}^n $\mathbf{1}_{\mathcal{V}} \in \mathcal{V}$, it follows that

$$\int_M \mathbf{1}_n \simeq \mathbf{1}_{\mathcal{V}}. \quad (19.24)$$

The canonical morphism $\mathbf{1}_n \rightarrow \mathcal{A}$ then induces a morphism $\mathbf{1}_v \rightarrow \int_M \mathcal{A}$, implying the factorisation homology $\int_M \mathcal{A}$ is not just an object of \mathcal{V} , but also comes equipped with a 0-disk algebra structure, the latter playing a crucial role for ground state degeneracy calculations.

Explicit calculation of factorisation homology as a 0-disk algebra is very useful. Indeed, factorisation homology on a closed stratified surface Σ with anomaly-free coefficient \mathcal{A} , (H, u_Σ) , exactly gives the ground state degeneracy of the same surface decorated by anomaly-free topological defects of codimension 0, 1, and 2 that are associated to \mathcal{A} , namely u_Σ .

Examples of stratified factorisation homology

Given a stratified 2-disk, M_1 , with \mathcal{A} a 2-disk algebra assigned to 2-cells, 1-disk algebras associated to 1-cells, and \mathcal{P} as the unique 0-algebra assigned to the unique 0-cell, examples of its factorisation homology read as follows

$$\int_{M_1} \mathcal{A}_1 \simeq \mathcal{A} \quad , \quad \int_{M_1 \setminus \{0\}} \mathcal{A}_1 \simeq HH_\bullet(\mathcal{A}) \stackrel{def.}{\equiv} \mathcal{A} \otimes_{\mathcal{A} \otimes \mathcal{A}^{op}} \mathcal{A}, \quad (19.25)$$

with $HH_\bullet(\mathcal{A})$ denoting the Hochschild cohomology.

For anomaly-free coefficient systems,




Figure 65. A stratified 2-disk with 2-cells labelled by UMTCs \mathcal{C} and \mathcal{D} . The unique 1-cell is labelled by a closed multi-fusion $\mathcal{C} - \mathcal{D}$ -bimodule, \mathcal{M} . The two 2-cells are labelled by (\mathcal{D}, d) , (\mathcal{M}, m) , for $d \in \mathcal{D}$ and $m \in \mathcal{M}$, respectively.

$$\boxtimes_{(c_0, \dots, c_{n-1})}^{\circ} (\mathcal{M}_1, \dots, \mathcal{M}_n) \simeq \text{Fun}_H(\mathcal{P}, \mathcal{P}). \quad (19.26)$$

The unique 1-cell labelled by a closed multi-fusion $\mathcal{D} - \mathcal{C}$ -bimodule \mathcal{M} . The two 0-cells are labelled by (\mathcal{D}, d) and (\mathcal{M}, m) , for $d \in \mathcal{D}$, and $m \in \mathcal{M}$, respectively. The closed multi-fusion $\mathcal{C} - \mathcal{D}$ -bimodule structure on \mathcal{M} induces two monoidal functors

$$\mathcal{C} \xrightarrow{L} \mathcal{M} \xleftarrow{R} \mathcal{D}. \quad (19.27)$$

Given $(\mathbb{R}^2; \Gamma_0)$ a stratified 2-disk, the FH reads as follows

$$\int_{(\mathbb{R}^2; \Gamma_0)} (\mathcal{C}, \mathcal{D}; \mathcal{M}; (\mathcal{D}, d), (\mathcal{M}, m)) \simeq (\mathcal{C}, L^\vee(m \otimes R(d))), \quad (19.28)$$

with L the right-adjoint functor of L . If $\mathcal{D} \equiv \mathcal{C}_{\mathcal{A}}^0$, $\mathcal{M} \equiv \mathcal{C}_{\mathcal{A}}$ for a connected commutative separable algebra \mathcal{A} in \mathcal{C} , $d \equiv \mathbf{1}_{\mathcal{D}}$, and $m \equiv \mathbf{1}_{\mathcal{M}}$, then (19.28) reduces to

$$\int_{(\mathbb{R}^2; \Gamma_0)} \left(\mathcal{C}, \mathcal{C}_{\mathcal{A}}^0; \mathcal{C}_{\mathcal{A}}; (\mathcal{C}_{\mathcal{A}}^0, d), (\mathcal{C}_{\mathcal{A}}, \mathbf{1}_{\mathcal{C}_{\mathcal{A}}^0}) \right) \simeq (\mathcal{C}, \mathcal{A}). \quad (19.29)$$

19.1.3 Non-contractible loops

In presence of non-contractible loops,

$$\int_{(S^2; S^1)} (\mathcal{C}, \mathcal{D}; \mathcal{K}; \emptyset) \simeq (H, \text{Hom}_{\mathcal{C} \boxtimes \mathcal{D}}(\mathbf{1}_{\mathcal{C} \boxtimes \mathcal{D}}, \mathcal{A})) \quad (19.30)$$

with

$$\mathcal{A} \stackrel{\text{def.}}{=} (L \boxtimes R)^V(\mathbf{1}_{\mathcal{K}}) \quad (19.31)$$

a commutative separable algebra in $\bar{\mathcal{C}} \boxtimes \mathcal{D}$. If \mathcal{K} is not fusion, then

$$\dim \text{Hom}_{\mathcal{C} \boxtimes \mathcal{D}}(\mathbf{1}_{\mathcal{C}} \boxtimes \mathbf{1}_{\mathcal{D}}, \mathcal{A}) = \dim \text{Hom}_k(\mathbf{1}_{\mathcal{K}}, \mathbf{1}_{\mathcal{K}}) > 1. \quad (19.32)$$

When this happens, the corresponding topological order is not stable, and triggers an RG-flow. Via factorisation homology, one defines the W -matrix, $W^{\mathcal{M}}$, for \mathcal{M} as follows

$$(L_{\mathcal{M}} \boxtimes R_{\mathcal{M}})^V(\mathbf{1}_{\mathcal{M}}) \simeq \bigoplus_{i \in O(\mathcal{C}), j \in O(\mathcal{D})} W_{ij}^{\mathcal{M}} i \boxtimes j^* \quad (19.33)$$

$$(L_{\mathcal{M}} \boxtimes R_{\mathcal{M}})^V(\mathbf{1}_{\mathcal{M} \boxtimes \mathcal{D} \boxtimes \mathcal{N}}) \simeq \bigoplus_{i \in O(\mathcal{C}), j \in O(\mathcal{E})} (W^{\mathcal{M}} W^{\mathcal{N}})_{ij} i \boxtimes j^*, \quad (19.34)$$

given $(\mathbf{R} \times S^1; S^1 \cup \dots \cup S^1)$ a stratified open cylinder with 2-cells labelled by UMTCs $\mathcal{C}_0, \dots, \mathcal{C}_n$, and 1-cells (namely non-contractible loops) labelled by closed multi-fusion $\mathcal{C}_{i-1} - \mathcal{C}_i$ -bimodules \mathcal{M}_i , $i = 1, \dots, n$, and no 0-cells.

Crucially, the above leads the definition of the ground state degeneracy, namely the index we are looking for

$$\boxed{\dim u \stackrel{\text{def.}}{=} \text{Tr} \left(W_1^{\mathcal{M}} W_2^{\mathcal{M}} \dots W_n^{\mathcal{M}} \right)}. \quad (19.35)$$

From the AGT correspondence, we know that this is equal to the partition function of a class \mathcal{S} theory.

As we will see in the upcoming sections, we can map the issue of gluing different class \mathcal{S} theories separated by a non-invertible defect, [1, 367], to that of defining the ground state degeneracy of a 2D TO in presence of non-contractible loops.

19.2 Dualisability and full-extendibility

We now turn to overviewing the relation between categorical dualisability and full-extendibility of a 2D TQFT. As we shall see, when applying this to a specific algebraic variety, namely the Hilbert series, homological mirror symmetry, in its categorical formulation, results as a biproduct of full-extendibility.

19.2.1 Extending TQFTs

Lurie proved that a TQFT is fully determined by what it assigns to a point. Furthermore, fully extended TQFTs with values in a symmetric monoidal (∞, n) -category, \mathcal{C} , are equivalent to fully dualisable objects in \mathcal{C} .

For a 2D TQFT, there are two main cases, [391]:

1. The TQFT is fully-extended, in which case the 2-functor reads as follows:

$$\mathcal{Z} : \text{Bord}_{2,1,0}^\sigma \rightarrow \mathcal{B}, \quad (19.36)$$

where $\text{Bord}_{2,1,0}^\sigma$ denotes the bicategories of points, 1-manifolds with boundaries, and 2-manifolds with corners. The target, \mathcal{B} , is most commonly associated with the bicategory of finite-dimensional \mathbf{k} -algebras, bimodules, and bimodule maps, $\text{Alg}_{\mathbf{k}}$. From the cobordism hypothesis, it follows that the extended framed TQFTs with values in $\text{Alg}_{\mathbf{k}}$ are classified by finite-dimensional separable \mathbf{k} -algebras.

2. The TQFT is not fully-extended, instead, if

$$\mathcal{Z} : \text{Bord}_{2,1}^{\text{or}} \rightarrow \mathcal{V}, \quad (19.37)$$

with \mathcal{V} a symmetric monoidal 1-category.

A non-extended 2D TQFT can be extended to the point if there is a symmetric monoidal bicategory, \mathcal{B} , and an extended TQFT such that

$$\mathcal{V} \simeq \text{End}_{\mathcal{B}}(\mathbf{1}_{\mathcal{B}}) \quad \text{and} \quad \mathcal{Z}_{ne} \simeq \mathcal{Z} \Big|_{\text{End}_{\text{Bord}_{2,1,0}^\sigma}(\emptyset)}, \quad (19.38)$$

where $\mathbf{1}_{\mathcal{B}} \in \mathcal{B}$ and $\emptyset \equiv \mathbf{1}_{\text{Bord}_{2,1,0}^\sigma(\emptyset)}$.

The prescription (19.38) is clearly not unique. In particular, it depends on the target \mathcal{B} .

If a non-extended 2D TQFT is a restriction of an appropriate defect TQFT $\mathcal{Z}_{ne}^{\text{def}}$, then the non-extended TQFT can be extended to the point as long as we associate it with the bicategory $\mathcal{B}_{\mathcal{Z}_{ne}^{\text{def}}}$.

19.3 Comparison with our main example

We now turn to explaining why this closely resembles the gauging prescription of SymTFTs, [102]. In order to do so, we build connection with our treatment and main example outlined in Part I and Part II of this work, namely algebraic varieties and ring homologies.

In building the understanding for how the setup of subsection 19.1.3 is of use for relating 3D mirror symmetry and homological mirror symmetry, we first need a brief mathematical digression, presented in the remainder of this section. Combining these tools with those of section 5, will be the main focus of sections 20 and 21.

19.3.1 Dualisability in categorical structures

A dualisable object in a symmetric monoidal (∞, n) -category, \mathcal{C} , is fully dualisable if the structure maps of the duality unit and counit each admit adjoints, as well as higher-morphisms between them, up to level $(n - 1)$.

By the Cobordism Hypothesis Theorem, Lurie proved that symmetric monoidal (∞, n) -functors out of the (∞, n) -category of cobordisms are characterised by their value at the point, which is a fully-dualisable object.

The notion of dualisability and full dualisability can be extended to the category of chain complexes, where an object is said to be dualisable if and only if it is a bounded chain complex of dualisable modules.

19.3.2 The case of the Hilbert series

We now turn to the higher-categorical structure most relevant for our treatment, whose dualisability is the core topic of the present and following sections.

In [385], the author explores many applications of representation theory of categories to sequences of groups arising in topology, algebra, and combinatorics.

Given $(V_n)_{n \in \mathbb{N}}$ a sequence of representations of groups G_n over a field k , it can be lifted to a category \mathcal{C} whose objects (or isomorphism classes) are indexed by natural numbers, such that the automorphism group of the n^{th} object of \mathcal{C} is G_n . A \mathcal{C} -representation, or \mathcal{C} -module, is a functor $\mathcal{C} \rightarrow \text{Mod}k$. One attempts to use the representation theory of \mathcal{C} to show that the whole sequence of representations is determined by a finite amount of data, and to discover universal patterns which the sequence must satisfy.

These patterns are often expressed in terms of generating functions. For example, one invariant of a sequence of representations is its sequence of dimensions, $(\dim V_n)_{n \in \mathbb{N}}$. Such data can be recorded as a generating function, called the Hilbert series of $(V_n)_{n \in \mathbb{N}}$

$$\text{HS}_V(t) \stackrel{\text{def.}}{=} \sum_n \dim_k V_n t^n \in \mathbb{Z}[[t]]. \quad (19.39)$$

In most cases, theorems from the representation theory of \mathcal{C} imply that $\text{HS}_V(t)$ is rational, with denominator of a certain type. For example, considering the category of finite dimensional vector spaces over \mathbb{F}_q and linear injections between them, if V is a finitely generated VI_q module, and $k \neq \text{char } \mathbb{F}_q$, then $\text{HS}(t)$ is rational, with denominator

$$w_d(t) \stackrel{\text{def.}}{=} \prod_{j=0}^{d-1} (1 - q^j t). \quad (19.40)$$

Each factor in (19.40) is a Whitney polynomial of the poset of subsets of a d -element set.

Building connection with categorical structures, we use the notion of *posets*. In order theory, a poset is a partially-ordered set, namely an arrangement such that, for certain pairs of elements, one precedes the other. The adjective partial denotes the fact that not every pair of elements needs to be comparable; indeed, there may be pairs for which neither element precedes the other. Partial order, thus generalises total orders, in which every pair is comparable. Given this, one can further define the *order complex* associated to a poset (S, \leq) which has the set S as vertices, and the finite chains of (S, \leq) as faces. To a poset, one can associate a poset topology, namely a topological structure on the poset of finite chains of (S, \leq) , ordered by inclusion. Using methods from poset topology, it is possible to construct a chain complex of \mathcal{C} -modules $k_d(V)$, $\forall d \in \mathbb{N}$, such that exactness of $k_d(V)$ categorifies the equation

$$w_d(t) \text{HS}_V(t) \equiv p(t), \quad (19.41)$$

for $p(t)$ a polynomial. If V is finitely generated, either k_d or a power of it applied to V is exact. In particular, categorification explains why the denominator (19.40) appears. This goes back to the construction outlined in our main example when integrating over the Kirwan map, (3.62).

Poset homology

Poset homology are chain complexes classifying rationality. Extended literature on the subject explains there is a method for assigning a chain complex to a module over a combinatorial category, such that, if the complex is exact, the module has a rational Hilbert series. Furthermore, these complexes are known to satisfy Homology-vanishing theorems. Hasse diagrams, discussed in [3], are an example of chain complexes associated to poset representations. Homological mirror symmetry




Figure 66. The upper and lower blue arrows correspond to geometric and algebraic resolutions of \mathfrak{M}^0 , respectively.

as an equivalence in between Hilber series on two categories, and rationality, are a direct biproduct of the defining features of poset homology, and constitutes a particular example of dualisable categorical structures, [147].

There is an equivalence of categories between the category of \mathbb{C} -linear representations of a quiver \mathcal{Q} and the category of left $\mathbb{C}\mathcal{Q}$ -modules.

Moment map and higher homologies

In general, homologies of derived representation schemes can be highly nontrivial. However, in this particular case, one can identify a necessary and sufficient condition for the vanishing of the higher homologies based on the flatness of μ , [344]. In particular, in such reference, it was shown that the derived representation scheme $\mathrm{DRep}_{v,w}(\mathcal{A})$ has vanishing higher homologies if and only if $\mu^{-1}(0) \subset M(\mathcal{Q}, v, w)$ is a complete intersection, which happens only if the moment map is flat, [365]. As explained in [3, 4], the requirement of the algebraic variety to be a complete intersection is crucial for the purpose of our treatment. In particular, it ensures the emergence of a 2-categorical structure, whose importance will be the core topic of sections 20 and 21.

20 Drinfeld Centers from Magnetic Quivers

We now turn to address a key issue highlighted in section 9.1.1, namely how to define a fiber functor associated to multiple gaugings not isomorphic to each other. For completeness, prior to explaining how this can be solved combining magnetic quivers and Drinfeld centers¹⁰⁰, we will first review the features of the setup outlined in section 9 needed in the present treatment. This section is structured into three parts:

- Relying upon the description of the HB and CBs as algebraic varieties (cf. section P.2), we explain how gauging can be related to the poset ordering leading to the construction of Hasse diagrams, thanks to the unifying role of the moment map.
- We then turn to explaining how the identification of such moment map ensures the quiver gauge theory enjoys a generalisation of homological mirror symmetry, with the latter corresponding to the presence of a Drinfeld center and associated fiber functor for a given 2-categorical structure, related to Rozansky-Witten theory, [343].

¹⁰⁰The definition of the latter will be provided in due course.

- We conclude opening a connection between the topics outlined in the present work and those of [192, 193].

The relation between posets, complex cohomology, and Drinfeld centers, will be explored in greater mathematical detail in section 21. This relation is intrinsic in this section, thanks to the connections outlined in section P.1; however, we are confident that a more mathematical treatment could also lead to further interesting insights that are worth exploring.




Figure 67. Condensing two different subalgebras, $\mathcal{A}_1, \mathcal{A}_2 \subset \mathcal{A}$, the resulting theory corresponds to one with a changed phase with a gauged defect resulting from a relative gaugeable algebra, \mathcal{A}_{12} ending in the bulk. The defect at the endpoint is nontrivial, and can therefore be thought of as a $\text{Hom}(\mathbb{1}_C, \mathcal{A}_{12})$.

20.1 Gauging and Hasse diagrams

We first start by recalling the configuration of interest outlined in section 9, motivating the importance of the Drinfeld center in geometric representation theory, [193].

Fiber functors




Figure 68. Adaptation of Freed-Moore-Teleman to the case involving twisted gauged defects. The figure on the far right corresponds to the case of interest to us, namely a configuration involving two different absolute 4D gauge theories separated by a defect. As we shall see, such defect is intrinsically non-invertible, corresponding to the presence of a relative uncondensed subalgebra, dressing \mathcal{N}_3 in a nontrivial way. In higher-categorical terms, it corresponds to a fusion tensor category implementing the morphisms between the operators charged under the gauged symmetry.

In our previous work, [1], we already argued that, unlike figure 30, the third picture from the left in figure 68 does not admit a straightforward expression for the fiber functor as (9.2). Nevertheless,




Figure 69. This picture reproduces an example of a correspondence, explained in section 14.

we claimed that a composite fiber functor could still be assigned thanks to the identification of the underlying algebraic structure of the composite theories

$$\boxed{{}_{\mathcal{A}_1} \mathcal{A}_{\mathcal{A}_2} \stackrel{def.}{\equiv} \mathcal{A}_1 \otimes_{\mathcal{A}} \mathcal{A}_2} \quad (20.1)$$

Our proposal in [1] was that the corresponding fiber functor should really be defining the partition function of a 3D theory, whose identification we anticipated being the core topic of the present work. The remainder of this section is devoted to prove that the resulting 3D theory is a 3D $\mathcal{N} = 4$ quiver gauge theory whose HB, when viewed as a complete intersection algebraic variety, enables to define the Drinfeld center of its underlying 2-categorical structure. In doing so, we need to completely specify its 1- and 2-morphisms, namely the generators and the relations in between them, both needed in the evaluation of the HB Hilbert series. As explained in section 19.1, the HS is nothing but the sum of gauge-invariant algebraic operators living in the chiral ring of the theory.

The Drinfeld center is defined as the 0^{th} -Hochschild cohomology, namely the endomorphisms of the identity element of a given category, \mathcal{C} :

$$\boxed{\mathfrak{Z}(\mathcal{C}) \stackrel{def.}{\equiv} \text{End}(\mathbf{1}_{\mathcal{C}}) \equiv \text{Hom}(\mathbf{1}_{\mathcal{C}}, \mathbf{1}_{\mathcal{C}}) \equiv HH^{\bullet 0}(\mathcal{C})} \quad (20.2)$$

The existence of the Drinfeld center and that of the fiber functor are mutually related by definition, since

$$\boxed{\mathcal{F} : \mathfrak{Z}(\mathcal{C}) \longrightarrow \mathcal{C}} \quad (20.3)$$

From what we have just outlined, it clearly follows that, the issue of identifying the Drinfeld center is equivalent to that of assigning a fiber functor projecting to the category \mathcal{C} . For the purpose of our treatment, namely building connection with geometric representation theory, \mathcal{C} is really meant to be the category of representations, with the latter being described in terms of 2D TFTs, namely Lagrangian submanifolds living in a symplectic manifold X , playing the role of boundary conditions for 3D Rozansky-Witten theory, RW_X . the meaning of this will be explained in section 20.2.

Relation to Hasse diagrams

The relation in between gauging and the construction of the Hasse diagram basically follows from the underlying role played by the moment map. In the former, it comes hand-in-hand with the definition of the fiber functor, meaning a partition function can be assigned to the resulting absolute theory, whereas, in the context of quiver gauge theories, the flatness of the moment map ensures a Hilbert series can be given, thereby fully specifying the generators and relations leading to the Hasse diagram construction itself. Importantly, the poset structure of the Hasse diagram admits a complex cochain description, thereby opening the possibility of studying this from a more mathematically rigorous point of view.

20.2 Drinfeld centers and mirror symmetry

As a key result of our work, we therefore wish to highlight the following statement as being applicable to the theories specified in the previous section P.1, namely quiver gauge theories with 8 supercharges,

Key point: identifying the Drinfeld center is equivalent to assigning a Hilbert series with irreducible representations, i.e. specifying the cones in the Hasse diagram, and their intersection.


The remainder of the present section is devoted to proving such assertion. In doing so, we mostly refer to [192, 193], which was one of the first motivations of this work. A more detailed mathematical description of the correspondence with such references, as well as with [346], is the core topic of the remaining sections. For the moment, we wish to highlight that such a correspondence exists, and can be understood from a mathematical physicist’s perspective. As a bi-product, we show how magnetic quivers can be combined with higher categories to extend the setup of [192, 193] for describing higher dimensional QFTs admitting a quiver gauge theory description.

Prior to doing so, we will first briefly recapitulate the notion of homological mirror symmetry, introduced in section E.3

Homological mirror symmetry

For our purposes, we are mostly interested in a geometrical representation theory realisation of homological mirror symmetry, for the same motivations outlined in [192]. As previously explained in [364], 3D mirror symmetry can be rephrased in terms of topological representation theory thanks to the Drinfeld center, [193]. We now briefly overview such argument, in order to make contact with the setting we are dealing with. For the case in which the two Lagrangians are related by mirror symmetry, such as is the case between Neumann and Dirichlet boundary conditions, the LHS of figure 70 is such that the intersection is unable to account for the Drinfeld in a straightforward manner, signalling the presence of a non-invertible defect separating the resulting absolute theories.

$$\text{Vect} \langle G \rangle \otimes_3 \text{Rep}(G) \equiv \text{Vect} \tag{20.4}$$



20.3 Drinfeld centers from magnetic quivers of 3D $\mathcal{N} = 4$ gauge theories

In this concluding section, we combine the tools encountered throughout our treatment, drawing additional major conclusions, as well as setting the stage for the core topic of the next section, first presented by the author in [3].

Figure 70. The black arrow in the diagram on the RHS denotes the fiber functor mapping from the Drinfeld center to the category of representation. Correspondingly, on the LHS, the Drinfeld center is the green shaded region whose boundaries are the regular and trivial representations, denoted in blue and red, respectively. For the case in which the two are related by ordinary mirror symmetry, their intersection contains the same information as \mathfrak{Z} , and therefore a fiber functor which is fully forgetful can be assigned.

2-fiber products

Some preliminary mathematical tools. If \mathcal{C} is a 2-category of categories, then there is a notion of a 2-fiber product, $\mathcal{C}_1 \times_{\mathcal{D}} \mathcal{C}_2$, also denoted by a diagram

$$\mathcal{C}_1 \xrightarrow{f} \mathcal{D} \xleftarrow{g} \mathcal{C}_2 \quad (20.5)$$

is the category of triples (c_1, c_2, ϕ) , where $c_1 \in \mathcal{C}_1$, $c_2 \in \mathcal{C}_2$, and $\phi : f(c_1) \xrightarrow{\sim} g(c_2)$ is an isomorphism of their images \mathcal{D} . For an object $P \in \mathcal{C}$, a 2-fiber product $X \times_S Y$ associated to the diagram

$$X \xrightarrow{f} S \xleftarrow{g} Y \quad (20.6)$$

is a quadruple (P, p, q, ϕ) , with 1-morphisms $p : P \rightarrow X, q : P \rightarrow Y$, and a 2-isomorphism

$$\phi : f \circ p \simeq g \circ q \quad (20.7)$$

such that $\forall Z \in \mathcal{C}$, the natural functor

$$\mathrm{Hom}_{\mathcal{C}}(Z, P) \rightarrow \mathrm{Hom}_{\mathcal{C}}(Z, X) \times_{\mathrm{Hom}_{\mathcal{C}}(Z, S)} \mathrm{Hom}_{\mathcal{C}}(Z, Y) \quad (20.8)$$

is an equivalence of categories.

Rozansky-Witten Theory and the Drinfeld center

One of the examples encountered in the previous section, namely $SU(2)$ with $N_f = 2$, is the simplest case involving two cones with nontrivial intersection in its Hasse diagram. As such, it is an example where ordinary mirror symmetry in the sense of (P.16) does not apply for the 3D $\mathcal{N} = 4$ theory arising under dimensional reduction. Nevertheless, a suitable mirror dual can still be assigned once the generators of the two cones featuring in the Hasse diagram, together with their intersection, are correctly accounted for in the calculation of the Hilbert series (HS), i.e. the partition function counting the gauge-invariant operators living in the chiral ring. In the previous section we explained how the magnetic quiver prescription enables to identify the HS correctly for cases where ordinary mirror symmetry does not apply.

To make contact with the calculation of Drinfeld centers, let us remind ourselves of the case of 4D $SU(2)$ $\mathcal{N} = 2$, first studied by Seiberg and Witten within the context of its 3D reduction. Its low-energy limit is described as a sigma-model in the space of vacua, corresponding to the Atiyah-Hitchin moduli space of charge 2 monopoles. The resulting 3D $SU(2)$ theory was then identified with the Rozansky-Witten theory (RW) of the hyper-kähler Atiyah-Hitchin manifold. Later developments of this by Kapustin, Rozansky, [382], and later with Saulina, [383], addressed their 2-category of branes, with the latter containing smooth holomorphic Lagrangians, L .

In the RW model, one must geometrically describe two functors from the 2-category of linear categories with G -action to linear categories:

1. the forgetful functor, keeping track of the underlying category (describing the pre-gauged TQFT), represented by the regular representation of G ,




Figure 71. Adaptation of the RW_X theory (on the LHS) to the quiche of a relative field theory (RHS).

2. the invariant category, generating the gauged TQFT, associated to the trivial representation of G .

This already encodes the idea of mirror symmetry, whose generalisation is realised thanks to the composite fiber functor described in [192], and overviewed in the concluding part of this section in relation to our analysis.

The objects in the 2-category of KRS are branes in the BFM space whose boundaries are Lagrangians $L, L' \subset X$, X a symplectic manifold, specifying the RW_X according to the choice of X . L, L' are 1-morphisms generating the theory upon acting on the trivial theory

$$\boxed{L, L' : \text{Id} \longrightarrow RW_X} \quad (20.9)$$




Figure 72. The BFM space for an abelian (LHS) and non-abelian (RHS) gauge theory, [192]. The KRS objects living in them are the objects in the 2-category referred to in the text and the trivial and regular representations are the Lagrangian submanifolds L, L' . Their intersection are Hom-spaces.

The 2-morphisms are denoted by $\text{Hom}(L, L')$. For the case in which L, L' are related by ordinary mirror symmetry, the 2-morphisms encode all the information, and the 2-category can be completely reconstructed from them. For the case of non-abelian gauge groups, a generalisation of the notion of mirror symmetry is needed. Practically, this means we need to specify, both, the 1-morphisms and the 2-morphisms to reconstruct the bulk of the BFM space, and completely describe the KRS branes living in them.

In connection with our previous work, [1], it is instructive to visualise what we have just said as shown in figure 71. Fully specifying the topological boundary conditions leads to an absolute theory. In case $\text{Hom}(L, L')$ leads to a non-invertible defect, the fiber functor, hence the Drinfeld center in figure 70 needs a 2-fiber product structure to be fully specified, as also explained in [1]. In particular, this highlights the importance of specifying the entire underlying 2-categorical structure, namely the Lagrangian submanifolds as well as the homomorphisms between them. This ensures the objects in the 2-category of KRS, can be fully specified. From the arguments outlined in the preliminary part of this work, we are therefore led to conclude that the configuration on the RHS of figure 71 is dual to another theory with the same Drinfeld center and only one Lagrangian submanifold on the topological boundary condition, thereby providing a sample realisation of generalised homological mirror symmetry.

Symplectically induced representations

In conclusion, we outline more explicitly the connection with the work of [192], setting the stage for a more detailed mathematical treatment, which is the main focus of [3].




Figure 73. The composition of fiber functors associated to double symplectic inductions leading to the regular and trivial representations.

One of the main aims of [192] is that of identifying a preferred family of simple representations, foliating the BFM space, ultimately describing the space of possible representations. The prescription outlined in such reference is motivated by the realisation of homological mirror symmetry. The construction of the preferred family of foliations also requires making use of a 2-fiber product, defined as follows

$$\boxed{\mathcal{T} \times_{T_{reg}^* G_c} Z_{reg}} \quad (20.10)$$

with \mathcal{T} and Z_{reg} denoting the trivial and regular fibers, respectively. (20.10) results from the composition outlined in figure 73, where

$$BFM(G) \stackrel{def.}{=} Z_{reg}/G_c \quad , \quad T(G) \stackrel{def.}{=} (N, \chi) \backslash T^* G_c // (N, \chi) \equiv N \setminus \mathcal{T} / N, \quad (20.11)$$

with

$$\chi : n \longrightarrow \mathbb{C}^\times \quad (20.12)$$

the regular character. Every such representation is symplectically induced from a 1D representation of a certain Levi subgroup G . As such, the preferred family of foliations resulting from (20.10) is the Fukaya category of a flag variety of G .

Given the considerations made when describing figure 71 in the context of RW-theory, we therefore notice that the aim of describing objects in the bulk of figure 72 is equivalent to that of defining the Drinfeld center for two absolute theories separated by a non-invertible defect, as encountered in [1]. Our main assertion is that, thanks to the prescription outlined in section P.2, in presence of multiple symplectically-induced representations, the resulting composite fiber functor defines the Drinfeld center of a mirror 3D $\mathcal{N} = 4$ gauge theory, resulting from the union of multiple Coulomb branches of magnetic quivers associated to 3D $\mathcal{N} = 4$ theories, thereby constituting a generalised realisation of homological mirror symmetry.

A more mathematical reformulation of this is the core topic of upcoming sections.

Key points

We can summarise the main results of sections 19 and 20 as follows:

- The 2-categorical structure description of the HB Hasse diagram of quiver gauge theories with 8 supercharges is equivalent to that of KRW. Importantly, this follows from the dimensional reduction of 4D $\mathcal{N} = 2$ $SU(2)$ gauge theory with $N_f = 2$, as the most basic case with HB Hasse diagram consisting of two intersecting cones.
- Importantly, the flatness of the moment map, ensuring the vanishing of higher homology, [344], corresponds to the possibility of fully specifying the HS associated to the HB of a given theory. We emphasise that this is equivalent to stating the existence of a fiber functor in a 2-category, ultimately leading to the definition of a partition function of an absolute theory, according to the prescription of [102].
- The identification of the MQs whose CBs equal the HB of the original theory, corresponds to the identification of the Drinfeld center of a 3D theory, with the latter being a RW theory, RW_x , whose BCs are Lagrangian submanifolds within a certain symplectic manifold, X . The complete reconstruction of the BFM space, where the objects of the 2-category live, corresponds to defining the Drinfeld center w.r.t. the given Lagrangian submanifolds and their intersection, i.e. the 1- and 2-morphisms characterising the 2-category in question.
- The fiber functor defining the partition function of interest is really obtained as a 2-fiber product, and is therefore compatible with the result of [192], motivated by generalising homological mirror symmetry.

21 Moore-Tachikawa varieties: beyond duality

We now have all the tools needed for addressing an open question raised in section 5, namely extending Moore-Tachikawa varieties beyond categorical duality. In doing so, we will be exploiting tools from algebraic geometry and higher-category theory.

In particular, we wish to stress the importance of dealing with the hyperkähler structure of the target 2-category of 2D TFTs describing Moore-Tachikawa varieties, and how this is related to the emergence of new higher-categorical structures requiring generalisation of the ordinary homological mirror symmetry/3D mirror symmetry correspondence.

The starting point is the following relation in between class \mathcal{S} theories and the gluing of their AGT dual 2D TFTs, [136],

$$\boxed{S_L \times_{G,q} S_R = S(C_L \times C_R, D_{LR})}, \quad (21.1)$$

where $(C_L, D_L), (C_R, D_R)$ are decorated 2-surfaces. The AGT correspondence states the equivalence holds in general. However, Moore and Tachikawa pointed out some interesting shortcomings for the case in which the 2-surface C is a sphere with two punctures. In such case, they claim that one does not achieve a genuine 4D class \mathcal{S} theory. The main aim of the present section is that of proving that this issue could be overcome by making use of a generalised notion of homological mirror symmetry/3D mirror symmetry correspondence, which, from the perspective of Moore-Tachikawa varieties, amounts to accounting for the hyperkähler structure of the target 2-category of the 2D TFT AGT-dual to the composite class \mathcal{S} theory on the LHS of (21.1).

Prior to delving into explaining a possible way of constructing a 2D TFT whose target category is a hyperkähler quotient, we briefly overview the work of [136] who first pointed out the hyperkähler structure needed generalisation of the target category of the 2D TFT η_{C_c} . We briefly overview some specific examples they outlined, explaining the relation with our treatment in Part I when dealing with star-shaped quivers.

Upon removing the identity from the source category, a morphism is now defined as a pair (C, a) , when C is an orientable 2-manifold with boundaries, and a is a positive real number which can be thought of as the area of C .

The composition of the two morphisms adds the area. For example, we see that

$$\lim_{a \rightarrow 0} (W, a) = \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2, \quad (21.2)$$

which is equivalent to the Coulomb branch encountered in Part I. Similarly, for $\mathfrak{g} = A_2$, $\eta_{A_2}(W, a)$ is equivalent to the minimal nilpotent orbit of $E_6(\mathbb{C})$ as the hyperkähler manifold in the $\lim_{a \rightarrow 0}$. At finite a , the hyperkähler metric is only invariant under $SU(3)^3$ and not the whole E_6 .

21.1 Categorical structure without duality

As explained in section 5, removing the identity element in Bo_2 requires having to introduce at least two different functors $\eta_{G'_c}, \eta_{G''_c}$, whose action on the circle and its bordism reads as follows

$$\eta_{G'_c}(S^1) \equiv G'_c, \quad \eta_{G''_c}(S^1) \equiv G''_c \quad (21.3)$$

$$\eta_{G'_c}(U) \equiv U_{G'_c} \equiv G'_c \times S_n \subset G'_c \times \mathfrak{g}'_c \simeq T^*G'_c, \quad (21.4)$$

$$\eta_{G''_c}(U) \equiv U_{G''_c} \equiv G''_c \times S_n \subset G''_c \times \mathfrak{g}''_c \simeq T^*G''_c, \quad (21.5)$$

$$\eta_{G'_c}(V) \equiv V_{G'_c} \stackrel{\text{def.}}{\equiv} T^*G'_c, \quad (21.6)$$

$$\eta_{G''_c}(V) \equiv V_{G''_c} \stackrel{\text{def.}}{\equiv} T^*G''_c, \quad (21.7)$$

where we are assuming

$$\mathfrak{g}'_c \cap \mathfrak{g}''_c \neq \{\emptyset\}, \quad \text{and} \quad \mathfrak{g}'_c \cup \mathfrak{g}''_c \equiv \mathfrak{g}_c, \quad (21.8)$$

$$\mathfrak{g}'_c \not\subset \mathfrak{g}''_c, \quad \text{and} \quad \mathfrak{g}''_c \not\subset \mathfrak{g}'_c. \quad (21.9)$$

(21.8) and (21.9) imply that the two subalgebras involved, $\mathfrak{g}'_{\mathcal{C}}, \mathfrak{g}''_{\mathcal{C}}$ are associated to different subgroups, $G'_{\mathcal{C}}, G''_{\mathcal{C}}$, and that the identity elements differ, $T^*G'_{\mathcal{C}} \neq T^*G''_{\mathcal{C}}$, even under reparametrisation of the Riemann surface. For each one of the 2D TFTs, $\eta_{G'_{\mathcal{C}}}, \eta_{G''_{\mathcal{C}}}$ one could use the formalism of [136, 137], describing two different class \mathcal{S} theories, both descending from 6D $\mathcal{N} = (2, 0)$ by dimensionally reducing on a Riemann surface without reparametrisation invariance. However, given the assumption that $(G'_{\mathcal{C}}, \mathfrak{g}'_{\mathcal{C}}), (G''_{\mathcal{C}}, \mathfrak{g}''_{\mathcal{C}})$ can be embedded in a unique $(G_{\mathcal{C}}, \mathfrak{g}_{\mathcal{C}})$, it is natural to ask what should the triple on the RHS of (18.1) be for the resulting theory to be absolute?

We know that the fiber functor and Drinfeld centers for a given absolute theory are defined as follows

$$\mathcal{F} : \mathfrak{Z}(\text{Bo}_2) \rightarrow \text{Bo}_2, \quad (21.10)$$

$$\mathfrak{Z}(\text{Bo}_2) \equiv \text{End}_{\text{Bo}_2}(T^*G_{\mathcal{C}}) \equiv \text{Hom}_{\text{Bo}_2}(T^*G_{\mathcal{C}}, T^*G_{\mathcal{C}}), \quad (21.11)$$

both of which crucially rely upon the presence of an identity in the source and target of $\eta_{G_{\mathcal{C}}}$. Apparently, we run into a contradiction, since the identity element $T^*G_{\mathcal{C}}$ has been removed by assumption, thereby implying $\mathfrak{Z}(\text{Bo}_2)$ cannot be defined in the ordinary way. On the other hand, one could define the new identity as being a composite object defined in the following way

$$\boxed{T^*\tilde{G}_{\mathcal{C}} \stackrel{\text{def.}}{=} T^*G'_{\mathcal{C}} \otimes_{T^*G_{\mathcal{C}}} T^*G''_{\mathcal{C}}}, \quad (21.12)$$

and therefore our proposal for (21.10) and (21.11) reads as follows

$$\boxed{\mathcal{F} : \mathfrak{Z}(\tilde{\text{Bo}}_2) \rightarrow \tilde{\text{Bo}}_2}, \quad (21.13)$$

$$\boxed{\mathfrak{Z}(\tilde{\text{Bo}}_2) \equiv \text{End}_{\tilde{\text{Bo}}_2}(T^*G'_{\mathcal{C}} \otimes_{T^*G_{\mathcal{C}}} T^*G''_{\mathcal{C}})}, \quad (21.14)$$

with

$$\tilde{\text{Bo}}_2 \stackrel{\text{def.}}{=} \text{Bo}_2 / V. \quad (21.15)$$

In the following subsection, we will explain why (21.13) and (21.14) are reasonable proposals for defining an absolute theory in absence of a categorical duality structure.

21.2 Interesting shortcomings

As explained in [1], the loss of reparametrisation invariance of the Riemann surface signals the presence of an intrinsically-non-invertible defect between different class \mathcal{S} theories. We will now show that the setup described in section 21.1 is equivalent to that of [1].

The starting point in our argument is the conjecture (5.28) and (5.29). In absence of categorical duality of, both, source and target, the 2D TFT associated to the maximal dimensional Higgs branch of [136] is no longer associated to a moment map that is equivalent for all the constituent S^1 's, thereby violating the conjecture made by [136]. Explicitly,

$$\boxed{W_{G_{\mathcal{C}}} \neq \mu^{-1}/G^3}. \quad (21.16)$$

This is because the algebraic variety associated to the Higgs branch W_{G_c} is not a hyperkähler quotient. In particular, it is associated to a non-primitive ideal.

Expanding further on this topic, the crucial point is that, when giving up the duality propriety, there is no longer the identity element in the source category, T^*G_c , but, rather, there is one identity for each underlying constituent, (21.4) and (21.5).

In order to determine how the resulting composite identity element should look like, we need to briefly recall what was outlined in section 5.4.2. Given two different morphisms, and taking their composition

$$T^*G_c^{a'} \circ T^*G_c^a \equiv (T^*G_c^a \times T^*G_c^{a'}) // G_c, \quad (21.17)$$

we know that, under the duality assumption, the axiom (5.18) comes with two moment maps, (5.14), each one describing the embedding of the individual morphisms within the algebra of the mother theory. As explained in section 5.4.2, one actually uses the mutual relation in between such moment maps to prove that T^*G_c behaves as the identity. As also claimed in the previous section, (21.17) expresses the need to define a Drinfeld center for the composite system made up of two class \mathcal{S} theories. In presence of reparametrisation invariance, such theories can be thought of as being separated by an invertible defect, from which (21.17) can be reduced to a statement of S-duality.

On the other hand, in absence of reparametrisation-invariance, the resulting class \mathcal{S} theories would, by definition, be separated by an intrinsically non-invertible defect, with the latter being responsible for the lack of reparametrisation invariance of the Riemann surface [137].

Our main question is to find the Drinfeld center for a given Bo_2

$$\mathcal{F}: \mathfrak{Z}(\text{Bo}_2) \rightarrow \text{Bo}_2, \quad (21.18)$$

and we know that, to a given fiber functor, \mathcal{F} , there is an associated moment map

$$\mu: \mathfrak{G} \rightarrow \mathcal{A}_c. \quad (21.19)$$

with \mathfrak{G} being Bo_2 in this case, and \mathcal{A} the algebra of invertible topological defects that projects to the identity T^*G_c under complete gauging of the theory. This is equivalent to stating that T^*G_c is the identity element once having projected over \mathcal{A}_c

$$\mathcal{A} //_{\mu} \mathfrak{G} \quad \text{with} \quad \mu: \mathfrak{G} \rightarrow \mathcal{A}_c \quad (21.20)$$

choosing the definition of the identity in the following way

$$T^*G_c \simeq \mathbf{1}_{G_c} \equiv \mathcal{A} //_{\mu} \mathfrak{G}. \quad (21.21)$$

For the purpose of our work, $\mathfrak{G} \stackrel{\text{def.}}{=} \text{Bo}_2$, therefore

$$T^*G_c \simeq \mathbf{1}_{G_c} \equiv \mathcal{A} //_{\mu} \text{Bo}_2 \quad (21.22)$$

However, in the case of section 21.1, there are two different moment maps involved, one for each choice of conformal structure on the Riemann surface, that are not mutually related by the moment map constraint following from the axiom (5.18). We therefore need to find the moment map (and corresponding gauge group \tilde{G}_c)

$$\tilde{\mu}: \tilde{\mathfrak{G}} \rightarrow \tilde{\mathcal{A}}_c, \quad (21.23)$$

whose identity element

$$\tilde{\mathcal{A}}_c \stackrel{def.}{=} \mathcal{A}_1 \otimes_{T^*G_c} \mathcal{A}_2 \quad (21.24)$$

constitutes the identity of the composite theory. If T^*G_c is the identity that has been removed from a particular source category, it still exists, but is no longer the identity present in $\tilde{\mathcal{B}}\mathcal{O}_2$ of a given $\tilde{\eta}_{G_c}$. From the RHS of (21.17), the identity of $\tilde{\mathcal{B}}\mathcal{O}_2$ therefore reads

$$T^*\tilde{G}_c \stackrel{def.}{=} T^*G'_c \otimes_{T^*G_c} T^*G''_c, \quad (21.25)$$

such that its Drinfeld center can be determined.

Defining $\tilde{\mu} : \tilde{\mathcal{B}}\mathcal{O}_2 \rightarrow \tilde{\mathcal{A}}$ as the moment map associated to the composite theory, the corresponding fiber functor can be explicitly rewritten as follows

$$\mathcal{F} : \mathfrak{Z}\left(\tilde{\eta}_{G_c}^{-1}\left(T^*\tilde{G}_c\right)\right) \rightarrow \tilde{\eta}_{G_c}^{-1}\left(T^*\tilde{G}_c\right). \quad (21.26)$$

ultimately enabling us to reformulate the problem of finding the Drinfeld center to that of identifying \mathcal{F} for a given μ .

Key points

The main points are the following:

- Lack of reparametrisation-invariance of bordism operators signals the presence of intrinsic non-invertible defects separating different class \mathcal{S} theories.
- Defining the Drinfeld center for a system of composite class \mathcal{S} theories separated by non-invertible defects constitutes a nontrivial generalisation of an S-duality statement.

21.3 Back to the identity from moduli spaces

In this concluding section, we gather the key tools outlined in this work, highlighting some main features leading towards the relation between homological mirror symmetry and 3D mirror symmetry.

3D mirror symmetry is defined as a duality exchanging Coulomb and Higgs branches, [146]. Making use of their algebro-geometric definition, briefly overviewed in section P.1, together with our previous work, [4], we will be arguing that *ordinary* 3D mirror symmetry, formulated as a 1-to-1 correspondence between a Coulomb and a Higgs branch, can be generalised to cases involving more theories (namely more Coulomb branches for a given Higgs branch). Specifically, we will be referring to the case of supersymmetric quiver gauge theories admitting a magnetic quiver description, and whose Higgs branch splits into two cones. This will ultimately provide further evidence to the claims made in [3].

As we shall see, the quantity that will enable us to probe such duality structure is the categorical interpretation of Hochschild cohomologies of block algebras, 19.1.2.

The present section is structured into three parts:

1. At first, we briefly overview the definition of block algebras, their associated Hochschild cohomology, [619], and the relation with the Drinfeld center, the latter following from our preliminary discussion in section 19.2.

2. We then turn to Coulomb and Higgs branches arising from Hitchin system for class \mathcal{S} theories, [389]. In particular, we highlight the fact that determining the Hitchin base, namely the Coulomb branch of the moduli space of vacua of class \mathcal{S} theories, from the Higgs branch, is equivalent to specifying the identity in the embedding category, in complete analogy with the criterion outline in section 19.1 and 19.2.
3. Ultimately, we combine the tools outlined in the remainder of the present treatment, together with our previous work, [4], explaining how in presence of algebraic varieties related to Moore-Tachikawa varieties without categorical duality, admit a generalised notion of 3D mirror symmetry thanks to the analogy with algebraic gauging in SymTFT constructions leading to intrinsically-non-invertible defects.

21.3.1 Hilbert series of Hochschild cohomology of block algebras

In this first subsection, we will briefly overview a key result of [619], combining it with material outlined in section 19.1.

Let p be a prime, k an algebraically closed field of characteristic p , G a finite group, and B a block algebra of kG . A defect group of B is a minimal subgroup P of G such that B is isomorphic to a direct summand of $B \otimes_{kP} B$ as a $B - B$ -bimodule. The defect groups of B form a G -conjugacy class of p -subgroups of G , and the defect of B is the integer $d(B)$ such that $p^{d(B)}$ is the order of the defect group of B .

Donovan's conjecture predicts that there should only be finitely-many Morita equivalence classes of block algebras of Hochschild cohomology algebras of block algebras with a fixed defect d . For any integer $n \in \mathbb{N}$, $\dim HH^n(B)$ is bounded in terms of the defect groups of the block algebra B .

The first result in [619] adds to this the fact that there is a bound on the degrees of the generators and relations of the Hochschild cohomology $HH^\bullet(B)$, which only depends on the defect of B . In [619], this comes in three theorems, ultimately claiming that, the Hilbert series of the Hochschild cohomology of a block and its defect determine each other up to finitely many possibilities. In particular,

$$HH^{\bullet 0}(B) \equiv \mathfrak{Z}(B), \quad (21.27)$$

where $\mathfrak{Z}(B)$ is the Drinfeld center.

21.3.2 Hitchin systems and mirror quivers

We now turn to the more specific setup of interest to us, namely that of class \mathcal{S} theories. In particular, we are interested in their moduli space of vacua, as well as their description in terms of 2D TQFTs, thanks to the AGT correspondence.

Hitchin systems

A Hitchin systems is an integrable system depending on the choice of a complex reductive group and a compact Riemann surface. It lies at the interface of algebraic geometry, Lie algebra theory, and integrable systems. It also plays a crucial role in the geometric Langlands program.

The Hitchin equations consist in a system of PDEs on the Riemann surface, C , concerning a triple (E, D, φ) , [389], where:

1. E is a G -bundle on C .
2. D is a G -connection in E .
3. $\varphi \in \Omega^1(\text{End } E)$.

Spectral curves and Hitchin fibrations

The Higgs branch moduli space, \mathcal{M}_H , is an example of one such complex integrable system. In particular, it is a fibration over a complex base space, \mathcal{B} , where the generic fiber is a compact complex torus.

For gauge groups $G = SU(N)$ or $G = PSU(N)$, given a Higgs bundle (E_h, ϕ) , then the eigenvalues of ϕ in the standard representation of G give an N -sheeted branch cover of C , [389],

$$\Sigma \stackrel{def.}{=} \left\{ \left(z \in C, \lambda \in T_z^* C \right) : \det(\phi(z) - \lambda) = 0 \right\} \subset T^* C, \quad (21.28)$$

which is the spectral curve corresponding to the Higgs bundle, (E_h, ϕ) .

The moduli space of harmonic bundles carries a natural family of complex structures J_ζ , parameterised by $\zeta \in \mathbf{CP}^1$. This follows from the fact that \mathcal{M} carries a natural hyperkähler metric, [389]. To see this, fix a G -bundle E . Then let \mathcal{C} denote the space of pairs (D, φ) without imposing the Hitchin equations. By definition, \mathcal{C} is an infinite-dimensional affine space, with a natural hyperkähler structure. Moreover, \mathcal{C} is naturally acted on by the gauge group \mathcal{G} . This action preserves the hyperkähler structure and has a moment map $\vec{\mu}$. The Hitchin equations imply that all 3 components of $\vec{\mu}$ must vanish. Denoting by \mathcal{G} the smooth sections of $\text{Aut } E$, we therefore have

$$\mathcal{M} \equiv \mu^{-1}(0)/\mathcal{G}, \quad (21.29)$$

namely the hyperkähler quotient

$$\mathcal{M} \equiv \mathcal{C}/\mathcal{G}, \quad (21.30)$$

implying \mathcal{M} is in turn hyperkähler. Every hyperkähler manifold carries a canonical family of complex structures parametrised by $\zeta \in \mathbf{CP}^1$, in this case denoted by J_ζ .

Let \mathcal{B} denote the space of all N -sheeted branched covers $\Sigma \subset T^*C$ of C . In this case, \mathcal{B} is a finite-dimensional complex vector space. Passing from the Higgs bundle to Σ gives a projection known as the Hitchin fibration

$$\mathcal{M}_H \rightarrow \mathcal{B}, \quad (21.31)$$

where \mathcal{B} is the Hitchin base.

As thoroughly explained in [389], \mathcal{B} is the Coulomb branch of the moduli space of vacua of the class \mathcal{S} theory in question, conventionally denoted $X_{\mathfrak{g}}(C)$. At any nonsingular point in the Coulomb branch, there is an electromagnetic charge lattice

$$\Gamma \rightarrow \mathcal{B}_{reg} = \mathcal{B}/\mathcal{B}_{sing}. \quad (21.32)$$

At every point of base, $u \in \mathcal{B}$, there is a spectral curve, Σ_u , thereby determining the local lattice system as the first cohomology, $H_1(\Sigma_u, \mathbf{Z})$.

21.3.3 Looking for the identity

In this concluding part of the section, we are now ready to gather all the ingredients outlined in the previous part of this work, to show how it relates to the formalism outlined in [4].

In doing so, we mostly need to recall the notion of the identity in Moore-Tachikawa varieties, and their generalisation for the case in which the target 2-category is a hyperkähler quotient

$$T^* \tilde{G}_C \stackrel{def.}{=} T^* G'_C \otimes_{T^* G_C} T^* G''_C. \quad (21.33)$$

As explained in [4], in such case, the identity is naturally removed by construction, and this clearly fits with taking the categorical quotient according to the GIT prescription, (21.30). From the work of Moore and Tachikawa, we also know that such categories are non-dualisable, and, consequently, not associated to a fully-extended TQFT.

As explained in section 19.2, full-extendibility can be recovered only upon specifying the identity element in the fully extended mother category from which the GIT quotient descends, (19.38). In particular, as suggested from (21.33), the identity itself might be a composite object, as the one arising when performing double gauging of a SymTFT leading to non-invertible defects, [1].

This partial gauging and gluing of different theories is precisely of the type occurring in the context of supersymmetric quiver gauge theories whose Higgs branch admits a magnetic quiver description, [222], as well as 3D mirrors of Sicilian theories, [377]. We plan to report a more detailed treatment for these specific examples in due course, [597].

At first, we briefly overviewed cochain level theories as the most important generalisation of the open and closed TFT construction, emphasising its relation to the SymTFT construction leading to absolute theories. Mostly relying upon [137], we highlighted the importance of the construction of cobordism operators, emphasising their dependence on the conformal structure of the Riemann surface. In section 4.1 we then turned to the discussion of a particular 2D TFT valued in a symmetric monoidal category, namely the maximal dimension Higgs branch of class \mathcal{S} theories. After briefly reviewing the properties outlined in [136], we propose their generalisation for the case in which the target category of the η_{G_c} functor is a hyperkähler quotient. We concluded outlining the possible extension of this treatment towards a mathematical formulation of magnetic quivers within the context of Coulomb branches of 3D $\mathcal{N} = 4$ quiver gauge theories which will be addressed in an upcoming work by the same author.

21.4 Where to next?

With the discussion outlined in Part II, III and V, we have shown the importance of abelianisation in geometric representation theory. Specifically, we highlighted the common ground between [102, 192, 600].

As explained at the very beginning of our work, one of our key motivations is the need to deepen the mathematical toolbox adopted by Phenomenologists in answering open questions in Particle Physics. Recent developments in the Pheno community have started relying upon the calculation of the Hilbert series, most importantly [604–606], successfully encoding key fundamental processes, such as the Higgs mechanism. The crucial point of our analysis, though, is that, albeit Hilbert series calculations are indeed very useful for keeping track of the degrees of freedom and symmetries of a given theory, in principle they are not enough to determine the exact operator content of the theory in question.

On the other hand, ring homologies, on which Hilbert series are calculated, encode crucial properties of the QFT at hand, enabling a much deeper understanding of its underlying categorical structure. The key take-home message of this work is the crucial role played by abelianisation for identifying the underlying mathematical structure of a given QFT. This concept will be investigated further for the specific case of the SM and its extensions in future work, [597].

Additional complementary directions addressed by the same author will be developing ring homologies for Moore-Tachikawa varieties beyond categorical duality¹⁰¹, and understanding their implications for Koszul duality. We plan to report of any advancements in this regard in the near future.

¹⁰¹Thereby following up from the analysis carried out in [4].

21.5 Conclusions and Outlook

With this work, the aim of the author was that of continuing from past developments, bridging the gap in between the Pure Mathematics and the Theoretical Particle Physics. The purpose of this thesis was mainly:

- To illustrate (some) key fundamental problems in Theoretical Particle Physics necessitating further mathematical understanding.
- Opening the scene to well-established techniques in representation theory and categorical algebraic geometry to help in this regard.
- For those who are not experts in the field, serve as an oversimplified explanation of the powerful formalism put forward by BFN, and connecting it with Moore-Tachikawa varieties.
- Motivating Phenomenologists to engage more with Pure Mathematicians.

The main objective of this Part was that of explaining the higher categorical structures that are needed for describing the invariants associated to specific supersymmetric quiver gauge theories, with a particular focus on dualities and their mutual relations in terms of higher-categories. We plan of reporting on any future developments in this regard in the near future, [597].

22 Conclusions and Outlook of Part V

Part V of this work is structured as follows:

1. After a preliminary list of definitions, section 19 introduces factorisation homology and its applicability within the context of class \mathcal{S} theories by means of the AGT correspondence. The section ends recapitulating the interplay between dualisability and full-extendibility of TQFTs, with a particular focus of algebraic varieties. This follows the treatment of section 5, which briefly overviewed cochain level theories as the most important generalisation of the open and closed TFT construction. Mostly relying upon [137], we highlight the importance of the construction of the cobordism operator, highlighting its dependence on the conformal structure of the Riemann surface. We then turn to the discussion of a particular 2D TFT valued in a symmetric monoidal category, namely the maximal dimension Higgs branch of class \mathcal{S} theories.
2. Section 20 explains the relation in between the Drinfeld center and magnetic quivers, combining sections 5 and 9.
3. In section 21 we propose the generalisation of Moore-Tachikawa varieties for the case in which the target category of the η_{G_c} functor is a hyperkähler quotient. We conclude outlining the possible extension of this treatment towards a mathematical formulation of magnetic quivers within the context of Coulomb branches of 3D $\mathcal{N} = 4$ quiver gauge theories which will be addressed in an upcoming work by the same author, [597]. The concluding part of this section recapitulates the essential tools required for performing GIT quotient constructions, highlighting its importance for realising 3D mirror symmetry. In particular, we focus on the calculation of the Hilbert series as specific examples of algebraic varieties of crucial interest in the study of categorical dualities for supersymmetric quiver gauge theories. We conclude showing how Coulomb and Higgs branches from class \mathcal{S} Hitchin systems carry the same information as the gaugeable algebras leading to SymTFT constructions, discussed in Part III. We conclude outlining interesting open questions and future directions of investigation by the author.

Part VI

Conclusions and Outlook

23 Conclusions and Outlook

String theory is a promising candidate unifying theory of fundamental interactions. In its fermionic formulation, it involves ten dimensions, of which nine are spatial and one is temporal. Crucially, such formulation strongly relies upon Supersymmetry (SUSY), which is a correspondence in between bosonic and fermionic particles.

According to present-day evidence, SUSY is certainly broken in our world, and plenty of effort has been made in identifying traces of such symmetry breaking from the observational side, eventually leading to the extension of the Standard Model (SM) by encoding additional particles and interactions. Understanding how the Standard Model can be embedded in String Theory is certainly a step forward towards identifying patterns of SUSY breaking in our world. Albeit being broken, SUSY certainly plays a crucial role in understanding which String Theory model is most appropriate for the SM to arise as an effective field theory (EFT), precisely by studying its breaking.

String theory vacua, where the SM would eventually be embedded, are parametrised by the so-called moduli space, and the latter is known to admit a categorical algebro-geometric description. Crucially, great advancements in the latter took place simultaneously to the birth and development of SUSY and String Theory. In recent years, joint collaboration in between pure mathematicians and mathematical physicists, primarily Freed, Hopkins, Lurie, Moore, Segal, and Teleman, shed light to new interesting perspectives which happen to be particularly relevant for the objectives outlined above. The main feature of their approach is that of applying the powerful tool of higher-categorical structures within the context of gauge theories and their supersymmetric counterparts, with the ultimate aim being that of explaining what is the underlying mathematical structure of a quantum field theory (QFT). Their work is based upon identifying underlying topological structures from which partition functions, and correlation functions can be derived. Furthermore, their formalism happens to be particularly efficient for keeping track of the spectrum (i.e. the particle-field content) of a given QFT, and how the latter changes under gauging of the global symmetries of the theory one started from.

Up to now, most of the mathematical control in describing higher-dimensional field theories has been for descendants of maximally-supersymmetric 6D $\mathcal{N} = (2, 0)$ superconformal field theories (SCFTs), from which a rich web of dualities arises upon dimensional reduction to, either, 4 or 3 spacetime dimensions. Such descendants feature fewer supersymmetries with respect to their non-Lagrangian parent theory. In particular, the most studied are 4D $\mathcal{N} = 2$ (also known as class \mathcal{S} theories), and 3D $\mathcal{N} = 4$ SCFTs, exhibiting exactly half of the amount of supersymmetry of their parent 6D theory.

From the work of Alday, Gaiotto and Tachikawa (AGT), it is known that class \mathcal{S} theories can be equally described in terms of a 2D topological field theory (TFT) living on the Riemann surface on which dimensional reduction has been performed. Such a 2D TFT is defined to be a functor in between 2-categories. Such 2-categories were first studied by Moore and Tachikawa, which, in turn, relied upon the pioneering work of Moore and Segal. The main point of their work is that, for a certain class \mathcal{S} theory to be uniquely defined (i.e. for its spectrum of operators to be uniquely determined), one needs to fully specify the source and target 2-categories of the 2D TFT corresponding to the given 4D $\mathcal{N} = 2$ SCFT in question. In technical terms, the objects of the source 2-category are circles, and the 1-morphisms between them are bordisms. The target 2-category is obtained by acting with the functor on these objects and 1-morphisms, resulting in

the gauge group of the 4D dual theory, and the bordism operators (also known as the Moore-Segal bordism operators), with the latter being related to algebraic varieties defining moduli spaces in algebraic geometry, i.e. the Higgs and Coulomb branches.

Higgs and Coulomb branches of supersymmetric gauge theories are fascinating objects to explore, since they shed light on dualities characterising the theory they are associated with. From a pure mathematical point of view, they can be realised in terms of the so-called geometric invariant theory (GIT) construction, and quotienting. For the case of 3D $\mathcal{N} = 4$ SCFTs, exchange in between Higgs and Coulomb branches corresponds to a statement of 3D mirror symmetry between two theories (first studied by Intriligator and Seiberg from a pure theoretical-physics point of view). This 1-to-1 correspondence extends to the calculation of invariants, most importantly the Hilbert Series, accounting for all the gauge-invariant operators of the theory.

However, as also pointed out by Moore, Segal and Tachikawa, their analysis lacked a rigorous mathematical treatment, most importantly for the case in which the target category of the functor is a hyperkähler quotient, which is indeed the case when describing algebraic varieties such as the Higgs branch of a given theory. The work we have conducted so far provides a step forward towards achieving a rigorous mathematical formulation of such 2-categorical structures for the reason that we will now outline.

As previously explained, TFTs play a crucial role in bridging the gap between QFT and Mathematics. Because of this, the formalism outlined above suggests to always recast a given QFT to a TFT description. For example, 3D $\mathcal{N} = 4$ SCFTs can be turned into 3D TFTs by performing a topological twist. Unlike ordinary 3D TFTs, which are usually trivial in nature in the sense that there are no local degrees of freedom, topologically-twisted TFTs come with nontrivial boundaries, called Lagrangian submanifolds. Importantly, the latter geometrize the boundary conditions of the field content of the theory in question, and, upon studying the intersection (i.e. the 1-morphisms) in between these Lagrangian submanifolds, one can reformulate the statement of 3D mirror symmetry of the parent 3D $\mathcal{N} = 4$ SCFT as that of homological mirror symmetry, which is essentially a duality in between 2-categories. The steps described so far clearly show the advantage of bringing the description down to that the 2-category of Lagrangian submanifolds, which is exactly what we need to keep track of the field content of the theory we started with.

The 3D mirror symmetry/homological mirror symmetry correspondence has been extensively studied for the specific case in which the correspondence in between the Coulomb and Higgs branch in 3D is 1-to-1. However, recent developments within the context of supersymmetric quiver gauge theories, mostly led by Hanany and collaborators, have shown that there are cases in which Higgs branches of fermionic theories can be broken down into several constituents, each one associated with the Coulomb branch of a certain 3D $\mathcal{N} = 4$ SCFT (a.k.a. the cone structure), with the latter being described by the so-called magnetic quivers. Such correspondence was primarily identified by studying the Hilbert series as a geometric invariant. In one of our recent works [3], we explained how the emergence of this multi-cone structure in the Higgs branch is equivalent to a generalised statement of 3D mirror symmetry, in turn descending from relaxing constraining assumptions on which homological mirror symmetry is based. As a follow up, in [4] we proposed a generalisation of the Moore-Tachikawa varieties for the case in which the target category of the 2D TFT is a hyperkähler quotient. As proved in the former of these two papers, achieving this requires generalising the bordism operators of Moore and Segal to the case involving lack of reparametrisation-invariance on the Riemann surface, ultimately enabling to relate this to the issue of defining a Drinfeld center for composite class \mathcal{S} theories.

So far, our main focus has been on class \mathcal{S} theories, and 3D $\mathcal{N} = 4$ SCFTs, both featuring 8 supercharges. To what extent the generalised 3D mirror symmetry/homological mirror symmetry correspondence can be extended to algebraic varieties associated with theories featuring different

amounts of supersymmetries is part of my currently ongoing research. The ultimate aim would be to extend this formalism to quiver gauge theories describing the SM and its possible extensions, eventually shedding light on the mathematical description of SUSY breaking mechanisms leading to observable EFTs.

This work explored the importance of 2D structures towards describing higher-dimensional quantum field theories (QFTs). In doing so, we relied upon the joint role played by dualities and categories.

The present work was structured in six parts:

1. At first, we introduced the notion of relative QFTs, with detailed explanation for the case of sigma models, gauge theories, and 6D SCFTs. We then turned to the main case of interest to us, namely Coulomb branches of star-shaped quiver gauge theories, introducing the Kirwan map, and the notion of abelianisation, naturally paving the way to the introduction of category theory, which played a crucial role in the remainder of our work.
2. Part II introduced the main ingredients of our treatment, namely categories and dualities. After having introduced higher-categorical structures, we provided two specific examples that were recursively encountered throughout our treatment, specifying how they are realised in the main example outlined in Part II thanks to the work of Moore and Tachikawa, [136]. Category theory is particularly useful for describing dualities, which is another crucial ingredient characterising this thesis. The dualities we encountered are:
 - Homological mirror symmetry.
 - 3D mirror symmetry.
 - The Alday-Gaiotto-Tachikawa (AGT) correspondence.
 - Holographic duality.

Apart from revising well-known results in the literature, we highlight the importance of going beyond full categorical dualisability in the context of Moore-Tachikawa varieties, to which Part V was devoted. In doing so, we related their formalism to that of SymTFT constructions of Freed, Moore and Teleman, [102], briefly overviewed in Part III.

3. Part III explained how QFTs arise from knot invariants. Specifically, we explained the Symmetry Topological Field Theory (SymTFT) construction of Freed, Moore, and Teleman. In particular, we focussed on the dimensional reduction of non-Lagrangian theories, specifically 6D $\mathcal{N} = (2, 0)$ SCFTs, [108, 109], and the categorical structure of the resulting class \mathcal{S} theories, [1]. Making use of the AGT correspondence, [135], we turned to a description in terms of 2D topological theories defined as functors between 2-categories.

This Part mostly centered around:

- The algebro-geometric formulation of moduli spaces, and their construction by means of the Geometric Invariant Theory (GIT) construction. Importantly, we relate this construction to the abelianisation procedure outlined in our main example in section 3.5, paving the way to further analysis in Part V.
- Turning from QM to Symplectic Geometry by means of the WKB approximation.

Both topics have thoroughly been addressed in the literature, and we therefore devoted some sections in the appendix to explaining them in greater depth.

4. Part IV built from open questions pointed out in the first paper by the author and her supervisor, [7]. In addressing this topic, we restrict to lower-dimensional setups in order to deal with open issues usually arising in higher-dimensional settings, extensively addressed in the literature, [208–210]. In doing so, we built connection with setups involving the emergence of islands, as well as recent developments in the context of von Neumann algebras for calculating generalised entropies in gravitating systems and QFT, [211–217]. The interesting connection with the remainder of our work is twofold: firstly, the need to resort to a certain dual picture in order to interpret the analytic results of the calculations performed; secondly, the richer structure resulting from the lower-dimensional analysis, emphasises the role played by internal structures in higher-dimensional theories, therefore suggesting alternative perspectives might be needed in order to address other open issues in QFT.
5. Part V results from the merge of three single-authored papers, [3–5]. The key findings in this Part are:
 - Moore-Tachikawa varieties without dualities are naturally realised in supersymmetric quiver gauge theories, and we therefore put forward a correspondence in between such theories, magnetic quivers, and Drinfeld centers, [3]. As a result of the formalism adopted (namely the definition of Moore-Segal bordism operators, [4, 137]) applied to Moore-Tachikawa varieties without categorical duality, [136], we provide further evidence of our conjectured relation in between homological mirror symmetry and 3D mirror symmetry, first presented in [3].
 - The last paper written by the author prior to submitting this Thesis, [5], nicely ties together the topics covered in [3, 4], while also preparing the stage for the future research the author will be conducting as a postdoc¹⁰². In particular, this part takes a more algebro-geometric approach in explaining the mathematical structure underlying supersymmetric quiver gauge theories, with a particular focus on dualities and their mutual relations in terms of higher-categories. As will be explained in depth, the crucial role is played by factorisation homology, [220]. We will not attempt to provide an exhaustive definition of it in this introduction, and, instead, refer the reader to the related section in the core of this work. What we can anticipate, though, is that factorisation homology is related with the counting of ground state degeneracy, and, when applied to the AGT correspondence, provides a tool for evaluating the 4-sphere partition function of class \mathcal{S} theories. Specifically, as we shall see, factorisation homology allows us to relate categorical dualisability and full-extendibility¹⁰³. Within the context of non-Lagrangian theories, such as 6D $\mathcal{N} = (2, 0)$ SCFTs this is particularly relevant, since it introduces the notion of ring objects, of which Hilbert series are one of the examples of most interest to us. As an algebraic variety, we focus on the calculation of the Hilbert series on Coulomb and Higgs branches, with the latter being related by the geometric invariant theory (GIT) quotient construction, [613]. As such, the GIT quotient provides a recipe for realising 3D mirror symmetry; we therefore show how it can be suitably extended for cases where mirror duals are not necessarily in a 1-to-1 correspondence. In relation to [222], this nicely fits with the magnetic quiver prescription, for describing the Higgs branch of certain supersymmetric quiver gauge theories with non-primitive ideals¹⁰⁴. We then show how the 2D description of disk algebras can be suitably adopted for de-

¹⁰²Partly outlined in [6].

¹⁰³And is therefore in line with the connection between dualities and categorical structures.

¹⁰⁴Namely the 2-cone structure, [223].

scribing Hitchin systems of class \mathcal{S} theories. We concluded outlining future directions that are currently under investigation by the same author.

6. Part V closes outlining the main achievements of the present work, with an outlook on some key questions the author wishes to address in her future research.

We believe our treatment opens the way to further investigating the mutual relation in between homological mirror symmetry and 3D mirror symmetry with the ultimate aim of aiding at understanding the underlying mathematical structure of QFT. These are currently under investigation by the same author, and we plan to report about any advancement in the field in the near future.

Acknowledgements

These acknowledgements are way far from reflecting my sincere gratitude towards the many people who trusted my passion, instinct, and dedication towards research. The names listed below are only part of the ones I wish to thank, and to whom I will be forever grateful.

First and foremost, my supervisor, Professor Fernando Quevedo. To summarise my working experience with him, I would say our collaboration has been a unique blend of creativity and perseverance, from my side, and patience and wisdom, from his. Thank you for following my progress towards becoming an independent researcher: it has been an honour to be working with you.

I also wish to thank Ahmed Almheiri, Shanta de Alwis, Tudor Dimofte, Iñaki García Etxebarria, Victor Gorbenko, David Jordan, Yolanda Lozano, Juan Maldacena, Gregory Moore, Hiraku Nakajima, Carlos Nuñez, Fernando Quevedo, Jorge Santos, Nathan Seiberg, Marcus Sperling, Constantin Teleman, Mark Van Raamsdonk, and Edward Witten for insightful discussions and questions raised at different stages of this work.

A special acknowledgement goes to, both, the Hodge Institute in Edinburgh, and the Yau Center for Mathematical Sciences, Tsinghua University in Beijing, for their hospitality, where parts of this work were completed and first presented. In particular, many thanks to my respective hosts, Tudor Dimofte, David Jordan, Nicolai Reshetikhin and Shing-Tung Yau.

I also wish to thank the organisers of all the events that I attended throughout my PhD, allowing this work to be presented at different stages, and the partial support by the STFC Consolidated HEP theory grant ST/T000694/1 through DAMTP towards covering tuition fees.

Being part of the DAMTP HEP Group made me feel at home while abroad, and I will be forever grateful for their warm welcome, which has certainly stimulated my research advancements in many ways. Many thanks to all other members of the HEP-Gang.

Last but not least, I will be forever grateful to the place that raised me during the first six years at university, the beautiful Trieste, and to the sailor whose dream made mine come true.

A Loop groups

A *loop group* is a group of loops in a topological group G with multiplication defined pairwise. It is a group of continuous mappings from a manifold M to a topological group G . Assuming $M = S^1$, the loop group is therefore formally defined as follows

$$LG \stackrel{def.}{=} \{\gamma : S^1 \rightarrow G \mid \gamma \in C(S^1, G)\}. \quad (\text{A.1})$$

Parametrising S^1 with θ , we get that

$$\gamma: \theta \in S^1 \mapsto \gamma(\theta) \in G, \quad (\text{A.2})$$

from which multiplication in LG is defined as follows

$$(\gamma_1 \gamma_2)(\theta) = \gamma_1(\theta) \gamma_2(\theta). \quad (\text{A.3})$$

Associativity follows from the associativity property in G . The inverse is given by

$$\gamma^{-1}: \gamma^{-1}(\theta) \equiv \gamma(\theta)^{-1}, \quad (\text{A.4})$$

and the identity is defined as follows

$$e: \theta \mapsto e \in G. \quad (\text{A.5})$$

An important example of loop groups is the group of based loops on G , ΩG . It is the kernel of the evaluation map

$$\begin{aligned} e_1: LG &\rightarrow G \\ \gamma &\mapsto \gamma(1), \end{aligned} \quad (\text{A.6})$$

and as such it is a normal subgroup of LG . G can also be embedded in LG , as the subspace of constant loops, leading to the split-exact sequence

$$1 \rightarrow \Omega G \rightarrow LG \rightarrow G \rightarrow 1. \quad (\text{A.7})$$

The space LG , therefore naturally splits as a semi-direct product

$$LG = \Omega G \rtimes G. \quad (\text{A.8})$$

B Supersymmetric Field Theories

B.1 Physical Lagrangians

A theory admitting a Lagrangian formulation, requires the Lagrangian to be physical, namely it must be:

1. real
2. local
3. Poincare'-invariant.

The Poincare' group, P^n is a subgroup of the global symmetry group of the theory, and, according to the Coleman-Mandula Theorem, the whole symmetry group of the theory takes the following form

$$P^n \times G, \quad (\text{B.1})$$

with G denoting a compact Lie group. By definition, symmetries give rise to conserved quantities. For the case at hand, (B.2) leads to conservation of energy and momentum, for the Poincare' group, and electric charges and quantum numbers for the compact Lie group.

The constraints of the Coleman-Mandula Theorem pertinent to bosonic theories can be relaxed in presence of fermions by extending the Poincare' group to the super-Poincare' group, $P^{n|s}$, which

is a super Lie group¹⁰⁵. The supersymmetric extension of the Coleman-Mandula Theorem, results in stating that the only allowed super Lie groups are of the kind

$$P^{n|s} \times G. \tag{B.2}$$

For the sake of completeness, we will briefly outline the main definitions of the terminology adopted in this section, [86].

amount of SUSY	max. dim.	σ -model	gauge theory
1	2	sp.f.	sp.f.
2	3	sp.f.	sp.f.
4	4	sp.f.(K)	sp.f.
8	6	sp.f.(hK)	sp.f.
16	10	-	existing

Figure 74.

A *super Lie group* is a group object in the category of supermanifolds. A *super Lie algebra* is a $\mathbb{Z}/2\mathbb{Z}$ -graded vector space

$$\mathfrak{g} = \mathfrak{g}^0 \oplus \mathfrak{g}^1, \tag{B.3}$$

equipped with a bracket operation

$$[\cdot, \cdot] : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}, \tag{B.4}$$

which is skew-symmetric and satisfies the Jacobi identity, with

$$[\mathfrak{g}^1, \mathfrak{g}^1] \in \mathfrak{g}^0. \tag{B.5}$$

The role of the super Poincare' group is that of restricting the possible vector spaces (i.e. particle content) and masses that can occur in a given theory. Mathematically speaking, an irreducible representation of $P^{n|s}$ may be realised as the space of sections of a homogeneous $\mathbb{Z}/2\mathbb{Z}$ -graded vector bundle over a mass shell. The extension to the case of interacting theories follows suite by noticing that one may use the free-field approximation at any vacuum, giving rise to a certain $P^{n|s}$ representation, which, in turn, follows from the free theory constraints.

B.2 UV/IR flows and their parametrisation

B.3 The Witten index

Supersymmetric theories are characterised by the presence of supersymmetric charge operators, \mathcal{Q} , such that bosons, $|B\rangle$, and fermions, $|F\rangle$, of energy E , are transformed into each other

¹⁰⁵Here, $P^{n|s}$ denotes a supersymmetric extensions of the Poincare' algebra $P^n \equiv \text{Lie}(P^n)$, whose odd part has dimension s .

$$\mathcal{Q}|B\rangle = \sqrt{E}|F\rangle \quad , \quad \mathcal{Q}|F\rangle = \sqrt{E}|B\rangle . \quad (\text{B.6})$$

(B.6) is equivalent to the statement that non-zero energy states come in Bose-Fermi pairs. This simple statement suggests how to establish whether SUSY is broken or not, namely by counting the zero-energy states. The key element for doing so was first proposed by Witten in [587], and reads

$$\mathcal{I} \stackrel{\text{def.}}{=} \text{Tr}(-1)^F e^{-\beta H} \quad (\text{B.7})$$

with F, β and H denoting the fermion number, the inverse temperature and the energy of the state, respectively. Due to (B.6), (B.7) is topological in nature and, by definition, only depends on the number of ground states. In particular, if $\mathcal{I} \neq 0$, the theory is certainly supersymmetric. Furthermore, it is independent w.r.t. β or any other parameter of the theory. Interestingly, though, there is a way in which \mathcal{I} can be effectively influenced by the features of a specific theory, namely if the number of ground states changes. As we shall see, one way of achieving this is via *spectral flow*, [588]. The latter will be the main focus of the next sections. However, for consistency, we already provide some qualitative arguments for its importance already at this stage.

Essentially, the spectral flow is implemented by an element of the superconformal algebra, enabling to interpolate between states defined as path-integrals with antiperiodic (fermionic) and periodic (bosonic) boundary conditions, i.e. between the NS- and R-sectors of a certain SUSY theory. Because of this, the action of the spectral flow is equivalent to changing the ground state degeneracy of the original SUSY theory.

B.3.1 R-symmetry, supergroups, superalgebras and twisting

Throughout this work, we will often be referring to many types of dualities. Even though they might seem every different in nature, one of the aims of our analysis will be furthering the mutual relation in between them, highlighting how and what one can tell about the others.

Among the first examples of dualities relevant for our purposes are holography, [201], and modular transformations. The former was originally formulated as a correspondence in between type-IIB string theory on an $\text{AdS}_5 \times S^5$, dual to 4D $\mathcal{N} = 4$ SYM. The holographic dictionary comprises the equivalence of the partition function on the field theory and gravity sides as well as the correspondence in between their symmetries, given by the superalgebra $PSU(2, 2|4)$. The bosonic subgroup is given by

$$SO(4, 2) \times SO(6) \subset SU(2, 2) \times SU(4) \subset PSU(2, 2|4) \quad (\text{B.8})$$

consistently with the isometries of AdS_5 and S^5 , as well as the conformal symmetry and the R-symmetry of the CFT_4 . By definition, an *R-symmetry* is one that does not commute with SUSY. As we shall see, its role will be essential throughout our treatment.

Modular transformations, instead, come in 2 kinds, denoted S and T . In string theory setups, they amount to changing the Yang-Mills coupling, τ_{YM} , as follows

$$\tau_{YM} \xrightarrow{S} -\frac{1}{\tau_{YM}} \quad \tau_{YM} \xrightarrow{T} \tau_{YM} + 1 \quad (\text{B.9})$$

At the level of the objects living in a certain string theory setup, modular transformations enable to map one theory into the other by acting on the objects with S, T . In particular, S relates strong and weak coupling, and therefore leads to a similar setup as the spectral flow mentioned earlier on B.3.

One way of testing these dualities, is by looking for quantities that are protected by supersymmetry. As also mentioned in B.3, one example is the matching of the spectrum of states characterising the Dp-branes. For example, the extended (i.e. $\mathcal{N} > 1$) superalgebra of 4D massive particles at rest reads

$$\{\mathcal{Q}_\alpha^I, \mathcal{Q}_\beta^{\dagger J}\} = 2M\delta^{IJ}\delta_{\alpha\beta} + 2iZ\Gamma_{\alpha,\beta}^0 \quad (\text{B.10})$$

with $I, J = 1, \dots, \mathcal{N}$, labelling the number of supersymmetries, $\alpha, \beta = 1, 2, 3, 4$, the components of the Majorana spinor supercharge, and $Z^{IJ} = -Z^{JI}$ denoting the *central charge* matrix. The components of the latter constitute a set of conserved quantities that commute with the other generators of the algebra. They are electric and magnetic charges coupling to the gauge fields belonging to the supergravity multiplet. Under the action of a unitary operator, U , it can be brought down to a block-diagonal form. The structure of the superconformal algebra therefore implies that the $2\mathcal{N} \times 2\mathcal{N}$ matrix on the RHS of B.13

$$\begin{pmatrix} M & Z \\ Z^\dagger & M \end{pmatrix} \quad (\text{B.11})$$

should be positive-definite with nonnegative eigenvalues $M \pm |Z_i|$, where $i = 1, \dots$, and $|Z_1| \geq |Z_2| \geq \dots$. This results in the so-called *BPS – bound*

$$\boxed{|Z_1| \leq M} \quad (\text{B.12})$$

Saturation of the BPS bound leads to a supermultiplet shortening, due to the emergence of zeros in the superalgebra when $M \equiv |Z_1|$. The bigger the number of equal central charges, the smaller the amount of SUSY left. Interestingly, BPS states, i.e. those with $M \equiv |Z_1|$, are stable and independent from the parameters of the theory as long as supersymmetry is preserved, and therefore shows exactly the same property as the Witten index, (B.7). As already argued in B.3, this quantity, namely the BPS states, can be used for bridging the study of the theory from weak to strong coupling.

However, the independence of (B.12) from the specifics of the theory is lost once another representation becomes degenerate with the BPS multiplet, such as occurring due to the Higgs mechanism.

For the purpose of our analysis, it is convenient to rewrite the superconformal algebra (B.13) in terms of the R -charges

$$\{\mathcal{Q}_\alpha^I, \mathcal{Q}_\beta^{\dagger J}\} = 2M_{\alpha\beta}\delta^{IJ} + 2i\omega_{\alpha\beta}\epsilon^{IJ}R \quad (\text{B.13})$$

with $\omega_{\alpha\beta}$ defining the symplectic form and ϵ^{IJ} the Levi-Civita symbol. For $\mathcal{N} = 2$, $M_{\alpha\beta} \in USp(2|2)$ and R being the $U(1)$ superconformal R -symmetry. Given that R defines the central extension of the superalgebra, it follows that

$$[R, \mathcal{Q}_\alpha^I] = \epsilon^{IJ}\mathcal{Q}_\alpha^J \quad (\text{B.14})$$

The lack of commutability in between the 2 sets of operators belonging to the extended superalgebra plays an essential role in the following analysis. In particular, it ensures that, as long as a relic R -symmetry is preserved, so too is SUSY (either totally or partially). These arguments therefore highlight the role played by the R -charges for protecting the index. As we shall see, (B.14) is at the heart of the noncommutativity in between 2 operations: the spectral flow and twisting.

B.3.2 The importance of flavour to extremise \mathcal{Z}

In [589, 590], for 2 and 4D respectively, it has been shown that the partition function of a CFT can be determined by means of localisation techniques starting from the knowledge of a UV-Lagrangian of the theory as long as the coupling to the background curvature of the sphere on which localisation is performed preserves the superconformal algebra. Equivalently, this statement amounts to ensuring the superconformal algebra contains the exact R -charges of the IR-theory, thereby ensuring that, along the spectral flow, the index accounts for the correct number of ground states of the theory.

The present work aims towards analysing the effect of *partial gauging* on, both, modular and holographic dualities in the attempt to provide a unique framework towards understanding the information loss paradox from a higher-dimensional realisation encompassing string theory and defect field theories. In pursuing such task, we will recast the issue of evaporating black hole microstate counting in the language of fusion of non-invertible defects.

B.4 SUSY enhancement from branes at orbifolds

Starting from a minimal model exhibiting the same SUSY enhancement w.r.t. the ABJM-theory, namely $\mathcal{N} = 2$ Chern-Simons theory with gauge group $U(N) \times U(N)$, and $SU(2) \times U(1)$ global symmetry. The field content consists of:

- 2 gauge fields: A_μ, \hat{A}_μ
- bi-fundamental bosons and fermions, Z^A, ψ^A , with $A = 1, 2$, and their hermitian conjugates.

Upper and lower indices denote the $\mathbf{2}$ and $\bar{\mathbf{2}}$ -representations of the global $SU(2)$, respectively. Furthermore, the gauge and matter fields are equipped with gauge group indices, a, b, \hat{a}, \hat{b} , for $U(N) \times U(N)$. Explicitly, they would read $A^a_{\mu b}$, $\hat{A}^{\hat{a}}_{\mu \hat{b}}$, $Z^{A, a}_{\hat{b}}$, $\psi^{A, a}_{\hat{b}}$, etc.

For the given field content, the corresponding $\mathcal{N} = 2$ action reads

$$S = \int d^3x (\mathcal{L}_o + \mathcal{L}_{CS} - V_{ferm} - V_{bos}) \quad (\text{B.15})$$

The kinetic terms are

$$\mathcal{L}_o = \text{Tr} \left(-D_\mu Z_A^\dagger D^\mu Z^A + i\psi^{\dagger A} \gamma^\mu D_\mu \psi_A \right) \quad (\text{B.16})$$

with covariant derivative

$$D_\mu = \partial_\mu + i[A_\mu, \] \quad (\text{B.17})$$

whereas the Chern-Simons terms read

$$\mathcal{L}_{CS} = \epsilon^{\mu\nu\rho} \text{Tr} \left(A_\mu \partial_\nu A_\rho + \frac{2i}{3} A_\mu A_\nu A_\rho - \left(A_\mu \longleftrightarrow \hat{A}_\mu \right) \right) \quad (\text{B.18})$$

The total action (B.15) is invariant under SUSY transformations. In particular, we have that

$$\delta Z^A = i\epsilon^\dagger \epsilon^{AB} \psi_B \quad (\text{B.19})$$

meaning that supersymmetry interpolates between different representations of the fields. Furthermore, the action has an additional $\mathcal{N} = 2$ SUSY dependent on the gauge group. In order to see this explicitly, we need to introduce a new local operator which enables for the representation

to remain the same upon turning bosons to fermions and viceversa. As already shown in the literature, such local operator, T_{ab} plays the role of a monopole operator ensuring gauge invariance of the fundamental fields of the theory. Correspondingly, the SUSY transformation rules turn into

$$\delta Z_b^{Aa} = iT_{ac}\psi_b^{\dagger Ac}\epsilon \quad (\text{B.20})$$

Having introduced a new operator in the spectrum, gauge invariance is restored ensuring the supersymmetric transformations preserve the representation of the fields.

B.4.1 ABJM and mirror symmetry

ABJM is a specific example of generalised Gaiotto-Witten (GW) setups. Indeed, it can be thought of as arising from the RG-flow of a 4D $\mathcal{N} = 4$ SYM on an interval with S-dual BCs.

However, its 1st formulation, was mainly motivated by the search for an AdS₄/CFT₃ correspondence, that could potentially be applied to string theory compactification setups. It was therefore As such, it constitutes a concrete setup where non-invertibles can be put to test.

C Higher-categorical symmetries from topological defects

To each D -dimensional QFT, \mathcal{T} , it is possible to assign a $(D - 1)$ -dimensional category, $\mathcal{C}_{\bar{x}}$, defined as *higher-category* if $D > 2$. $\mathcal{C}_{\bar{x}}$ features D -levels:

1. 1st level: 0-morphisms, namely the objects living in $\mathcal{C}_{\bar{x}}$
2. 2nd level: 1-morphisms, defining the relations between the objects
3. 3rd level: 2-morphisms between the 1-morphisms.
4. iteratively up to $D - 1$.




Figure 75. Fusion of objects in $\mathcal{C}_{\bar{x}}$.

The objects are codimension-1 topological defects

$$D_{D-1} \stackrel{def.}{=} \bigoplus_i n_i D_{D-1}^{(i)} \quad , \quad n_i \geq 0 \quad (\text{C.1})$$




Figure 76. Generalised fusion between 1-morphisms.

where $D_{D-1}^{(i)}$ are codimension-1 topological defects, and $2 \leq i \leq D$ denotes the $(i-1)$ -morphisms between $(i-2)$ -morphisms. $\sum_i n_i$ defining the total number of vacua such that in n_i of them, D_{D-1} behaves as $D_{D-1}^{(i)}$. For the case of a simple object, there is a unique vacuum, and the defect carries a single topological local operator on its worldvolume.

Under fusion, the objects give rise to a *monoidal* structure

$$D_{D-1}^{(1)} \otimes D_{D-1}^{(2)} \equiv D_{D-1}^{(12)} \quad (\text{C.2})$$

as depicted on the LHS of figure 75.



Figure 77. If $D_p^{(1)}$ and $D_p^{(2)}$ are related by gauging, wrapping the $D_{p-1}^{(1,2)}$ defect around an S^{p-1} , bridges the LHS and the RHS of this figure. The algebra can therefore be placed on the defect for defining interfaces.

The fusion structure (C.2) descends to a fusion structure of objects of a higher-category of symmetries localised along D_{D-1}

$$D_{D-2}^{(1,2)} \otimes_{D_{D-1}^{(2)}} D_{D-2}^{(2,3)} \equiv D_{D-2}^{(2,3)} \circ D_{D-2}^{(1,2)} \quad (\text{C.3})$$

For the fusion (C.3) to define a 1-morphism, $\mathcal{C}_{\bar{x}}$ is required to admit the existence of 2-morphisms, namely the theory has to be, at least, 3D. However, this does not prevent the possibility of defining other types of fusion rules, independently of the existence of higher-morphisms, such as the one depicted in figure 76.

C.1 Generalised gauging and localised symmetries

Let us consider the defects $D_p^{(1)}, D_p^{(2)}, D_{p-1}^{(1,2)}$, with the latter being defined such that, when wrapping an S^{D-1} , $D_p^{(2)}$ vanishes. This corresponds to saying that $D_p^{(1)}$ and $D_p^{(2)}$ are related by gauging. Practically, this relation implies that, for any given representative of $D_p^{(1)}$, we obtain a representative of $D_p^{(2)}$ upon gauging the worldvolume theory of $D_p^{(1)}$. Furthermore, all topological subdefects of $D_p^{(2)}$ can be obtained from those of $D_p^{(1)}$.

This observation enables one to define a generalised gauging construction for any subcategory $\mathcal{C}_{\bar{x}, D_p} \subset \mathcal{C}_{\bar{x}}$. W.r.t. the original categorical theory $\mathcal{C}_{\bar{x}}$, the D_p -defect is a $(D-p-1)$ -morphism. The objects of the subcategory $\mathcal{C}_{\bar{x}, D_p}$ are $(D-p)$ -morphisms of $\mathcal{C}_{\bar{x}}$ from D_p to itself. Its 1-morphisms are $(D-p+1)$ -morphisms of $\mathcal{C}_{\bar{x}}$. The fusion structure \otimes of $\mathcal{C}_{\bar{x}, D_p}$ descends from \otimes_{D_p} on $\mathcal{C}_{\bar{x}}$, and, consequently, its fusion rules as well.

For $p=2$, the gauging is performed by acting with an algebra, $A^{(1,2)}$, living in the 1-category $\mathcal{C}_{\bar{x}, D_2^{(1)}}$. The algebra itself consists of the following elements

$$A^{(1,2)} \stackrel{def.}{=} \left\{ A_1^{(1,2)}, A_0^i \right\} \quad , \quad i \equiv \alpha, \beta, \gamma, \delta \quad (\text{C.4})$$

namely an object¹⁰⁶

$$A_1^{(1,2)} \stackrel{def.}{=} D_1^{(1,2)} \otimes D_1^{(2,1)} \quad (\text{C.5})$$

and canonical morphisms

$$\begin{aligned} A_0^\alpha : A_1^{(1,2)} \otimes A_1^{(1,2)} &\rightarrow A_1^{(1,2)} \quad , \quad A_0^\beta : A_1^{(1,2)} \rightarrow A_1^{(1,2)} \otimes A^{(1,2)} \\ A_0^\gamma : A_1^{(1,2)} &\rightarrow 1_{D_2^{(1)}} \quad , \quad A_0^\delta : 1_{D_2^{(1)}} \rightarrow A_1^{(1,2)}. \end{aligned} \quad (\text{C.6})$$

Category of lines after gauging

After gauging, the resulting defect can be expressed in terms of (C.4)

$$D_2^{(2)} \equiv \frac{D_2^{(1)}}{A^{(1,2)}}. \quad (\text{C.7})$$

The gauging of $\mathcal{C}_{\bar{x}, D_2^{(1)}}$ is performed upon inserting a mixture of topological defects, constructed from $A^{(1,2)}$, along the entire domain of the line defect $D_1^{(1,2)}$.

For the case of arbitrary p , the gauging is described by a $(p-1)$ -algebra in the $(p-1)$ -category $\mathcal{C}_{\bar{x}, D_p^{(1)}}$

$$A^{(1,2)} \stackrel{def.}{=} \left\{ A_{p-1}^{(1,2)}, A_{p-2}^i \right\} \quad , \quad 1 \leq i \leq p-1 \quad (\text{C.8})$$

where the morphisms (denoted by i) describe all possible creation and annihilation channels. $D_p^{(2)}$ is obtained by placing a mixture of elements of $A_p^{(1,2)}$ along $D_p^{(1)}$.

¹⁰⁶Constituting a gaugeable 1-form symmetry. This is because the algebra described here is an example of an algebra that can be projected to the identity to achieve a gauged theory, namely quotienting the higher-category of topological defects in the SymTFT of the QFT in question, [607].

C.2 Fusion rules

The importance of having introduced the algebra to implement the gauging, manifests when looking at the fusion rules between the defects characterising a given theory. For ease of notation, we will now argue on the basis of a category $\mathcal{C}_{\text{id}, \mathfrak{T}}$, and the fusion of 2D objects

$$D_2^{(O)} \otimes D_2^{(O')} = \bigoplus_i D_2^{(i)}, \quad (\text{C.9})$$

with O, O' denoting different orbits. The line operators describing the 1-morphisms

$$D_2^{(O)} \otimes D_2^{(O')} \rightarrow D_2^{(i)} \quad (\text{C.10})$$

arrange into orbits, O'' , under the action of the gauge group G , such that the fusion can be re-expressed in terms of them

$$D_2^{(O)} \otimes D_2^{(O')} = \bigoplus_{O''} D_2^{(O'')} \quad (\text{C.11})$$

where the RHS is a representative of the equivalence class of the fusion on the LHS in the quotient theory \mathfrak{T}/G . The generalised gauging on top of each $D_2^{(O'')}$ term is encoded by an algebra $A^{(O''), (OO')}$ in the 1-category $\mathcal{C}_{\text{id}, \mathfrak{T}/G, D_2^{(O'')}}$, with object

$$A_1^{(O''), (OO')} \equiv \bigoplus_i n_i D_1^{(i)} \quad (\text{C.12})$$

where n_i denotes the dimension of the vector space constituted by local operators living at the end of $D_1^{(i)}$ along $D_1^{(OO'), (O'')}$. Consequently, after gauging, the fusion of surface defects, (C.9), turns into

$$D_2^{(O)} \otimes D_2^{(O')} = \bigoplus_{O''} \frac{D_2^{(O'')}}{A_1^{(O''), (OO')}} \quad (\text{C.13})$$

with the dependence on the algebra describing the gauging on top of each elementary surface term. This is referred to as the *Dressing* of the topological operators. Similar considerations can be made for the case of line defects.

For the purpose of this work, we will be mostly focussing on duality, triality and N -ality defects. By definition, the first 2 are associated to theories with $\mathbb{Z}_N^{(1)}$ 1-form symmetries, whereas the latter emerge in presence of a $\mathbb{Z}_N^{(1)} \times \mathbb{Z}_N^{(1)}$ symmetry. It is worth mentioning already at this stage that their fusion rules are basically providing a generalisation of (C.4). Specifying to the case of (3+1) TQFTs, the fusion of 3D duality/triality defects, $\mathcal{N}(\mathcal{M}_3) \stackrel{\text{def.}}{=} D_3(\mathcal{M}_3)A_3$ reads

$$\mathcal{N}(\mathcal{M}_3) \times \bar{\mathcal{N}}(\mathcal{M}_3) \equiv \mathcal{C}(\mathcal{M}_3) \quad (\text{C.14})$$

with $\mathcal{C}(\mathcal{M}_3)$ denoting the *gauging defect* defined as

$$\mathcal{C}(\mathcal{M}_3) \stackrel{\text{def.}}{=} \frac{1}{|H^0(\mathcal{M}_3, G)|} \sum_{\Sigma_2 \in H_2(\mathcal{M}_3, G)} L(\Sigma_2) \quad (\text{C.15})$$

From (C.14), $\mathcal{N}(\mathcal{M}_3)$ is often referred to as the square root of the gauging. As a further major consideration worth mentioning is that the quantum dimensions of the duality and triality defects on S^3 coincide.



Figure 78. TWR as an example of holographic duality.

D Topological Wick rotation and holographic dualities

The AdS/CFT correspondence is an example of holographic duality. Two main features characterising such correspondence are that:

1. Gauge symmetries in AdS \leftrightarrow global symmetries of the CFT.
2. Correlation functions are only defined when boundary conditions of the AdS theory are specified.

These same properties are shared by other holographic dualities. A new type of holographic duality is the one given by topological Wick rotation.

The holographic dictionary encoded in figure 78 reads as follows:

1. an $N + 1$ dimensional topological order with a gapped boundary is dual to an N dimensional liquid with an internal symmetry of finite type. \mathcal{S} is the category of topological defects in the bulk. On the RHS it is the superselection sector of states. $\mathfrak{Z}(\mathcal{S})$ is the category of topological defects in the bulk and naturally acts on \mathcal{S} . $\mathfrak{Z}(\mathcal{S})$ is the category of topological operators (non-local operators invariant under LOA).
2. ${}^{\mathfrak{Z}(\mathcal{S})}\mathcal{S}$ is an *enriched category*.

Consider gapped/gapless edges of a 2+1 dimensional topological order (\mathcal{C}, c) , with $a, b, c \in \mathcal{S}$ labelling the topological line defects. The macroscopic observables of the 1+1 dimensional edge form an enriched fusion category ${}^{\mathcal{B}}\mathcal{S}$. The spaces of fields living on the 0 dimensional defect junction $\{M_{a,b}\}$ lead to the OPE

$$M_{b,c} \otimes_{\mathcal{C}} M_{a,b} \xrightarrow{\circ} M_{a,c} \quad (\text{D.1})$$

$\{M_{a,b}\}_{a,b \in \mathcal{S}}$ is similar to a category $\{\text{hom}(a, b)\}_{a,b \in \mathcal{C}}$. $M_{a,b} \circ \in \mathcal{B}$ defines an enriched fusion category. Introducing an abstract MTC \mathcal{B} and a braided equivalence $\phi: \text{Mod}_V \xrightarrow{\cong} \mathcal{B}, (\phi, {}^{\mathcal{B}}\mathcal{S})$ can encode different enrichments ${}_{\phi}^{\text{Mod}_V}\mathcal{S}$. Correlation functions and OPEs among defect fields can be recovered from the triple $(V, \phi, {}^{\mathcal{B}}\mathcal{S})$. The gapped/gapless boundaries of a 2+1 dimensional topological order (\mathcal{C}, c) can be completely characterised or classified by the triple $(V, \phi, {}^{\mathcal{B}}\mathcal{S})$, where:

1. V is the chiral/non-chiral symmetry which
 - for a chiral gapless boundary, corresponds to a rational VOA of central charge c
 - for a non-chiral gapless boundary, it is a rational full field algebra
 - when $V \equiv \mathbb{C}$, it is described by a gapped boundary.
2. $\phi: \text{Mod}_V \xrightarrow{\cong} \mathcal{B}$ is a braided equivalence.

3. \mathcal{S} is a fusion category equipped with a braided equivalence $\mathcal{C} \boxtimes \bar{\mathcal{B}} \simeq \mathfrak{Z}(\mathcal{S})$, which determines ${}^{\mathcal{B}}\mathcal{S}$ via the so-called *canonical construction*.

All gapped/gapless edges of 2+1 dimensional topological orders can be determined by performing a TWR and adding the chiral/non-chiral symmetry data by hand $(V, \text{Mod}_V \xrightarrow{\phi} \mathcal{B}, {}^{\mathcal{B}}\mathcal{S})$. ${}^{\mathcal{B}}\mathcal{S}$ is the *topological skeleton* (encoding the topological or categorical data), whereas (V, ϕ) constitutes the local quantum symmetry of the edge (e.g. the VOA or the full field algebra or the conformal net of local operator algebras (LOA)). ${}^{\mathcal{B}}\mathcal{S}$ is an *abstract enriched FC*, acquiring precise meaning only once \mathcal{B} is identified with Mod_V via the braided equivalence $\phi : \text{Mod}_V \xrightarrow{\simeq} \mathcal{B}$, suggesting that a gapped/gapless quantum liquid χ can be described by a 2-step data $\chi \equiv (\chi_{lqs}, \chi_{sk})$, since the same ${}^{\mathcal{B}}\mathcal{S}$ can be realised by different gapped phases. Examples of these constructions are 2D WZW CFTs from 3D Chern-Simons theories. Constructing correlators in 2D CFTs from 3D TFTs. There is a correspondence between Lagrangian algebras in 3D anyon gauging and modular-invariant partition functions in 2D CFTs. Dualities are explicitly constructed from classical statistical models. Correlators mapping 3D TQFTs to 2D RCFTs. String-net construction of 2D correlators.




Figure 79.

Theorem (Kong, Huang)

There is a holographic duality between $(\mathfrak{Z}(\mathcal{S}), \mathcal{S})$ and ${}^{\mathcal{B}}\mathcal{S}$. An $N+1$ dimensional topological order with higher-dimensional symmetry \mathcal{R} can be completely characterised:

- a fusion n -category \mathcal{A} equipped with a braided embedding $\mathcal{R} \hookrightarrow \mathfrak{Z}_1(\mathcal{A})$
- a braided equivalence $\phi : \mathfrak{Z}_1(\mathcal{R}) \xrightarrow{\simeq} \mathfrak{Z}_1(\mathcal{A})$.

Generalising \mathcal{R} to any fusion $(N-1)$ -category, one obtains a characterisation of N dimensional gapped quantum liquids by an $N+1$ dimensional topological order with gapped boundary. TWR suggests that this holographic duality is actually functorial. TWR gives an equivalence between the category of $N+1$ dimensional non-chiral topological orders with gapped boundary and the category of N dimensional topological liquids

$$(\mathfrak{Z}_1(\mathcal{A}), \mathcal{A}) \longrightarrow {}^{\mathfrak{Z}_1(\mathcal{A})}\mathcal{A} \quad (\text{D.2})$$

This holographic dictionary automatically includes a dictionary map of dualities because many dualities can be realised by invertible DWs, e.g. e-m, and K-W duality.

One of the main goals behind all these studies is the mathematical theory of higher-dimensional CFTs.

E Quiver gauge theories

E.1 Graphs

A graph, Γ , gives a category, \mathcal{C}_Γ , with 3 objects: A, B , and C . $\mathcal{C}_\Gamma(A, B)$ is just the single arrow from A to B .




Figure 80. Examples of type-A and type-D quivers.




Figure 81.




Figure 82. Examples of type-E quivers.

E.2 Dynkin diagrams

The main focus of our analysis will be the understanding of the behaviour of 4D $\mathcal{N} = 2$ theories. In doing so, we wish to bring the attention of the reader towards the construction of the 4D theory itself, which already accounts for very useful insight towards understanding major issues we wish to address.

The 4D $\mathcal{N} = 2$ theories that are usually being dealt with, are assumed to possess S -duality as inherited from their mother theory, namely 6D $\mathcal{N} = (2, 0)$ SCFT theories, upon dimensional reduction and topological twist on a Riemann surface Σ_2 . Albeit not being this the unique way of deriving such 4D theories, yet they provide the most generic way, which, in turn, allows for a plethora of different descending theories. These 6D SCFTs are very particular UV theories. They were discovered as the decoupling limit of type-IIB on an orbifold singularity. The resolution of the singularity leads to a 6D SCFT theory with characterised according to the ADE classification, with A, D, E denoting the corresponding types of Dynkin diagrams.

E.3 Framed Nakajima quiver varieties

In this first subsection, we briefly overview the features of framed Nakajima quiver varieties, [344], and their geometric and algebraic resolutions.




Figure 83. An example of a framed Nakajima quiver variety. This will be a prototypical example of the theories we will be addressing in the present work, namely for quiver gauge theories with 8 supersymmetries. The unframed quiver would be the gauge node without matter. Here we have set $G_v \equiv SU(N)$. However, in principle, other choices could have been taken.

Geometric (Nakajima) resolution

Quiver varieties are varieties of quiver representations of a quiver: one fixes a vector space at each vertex, then considers the linear space of representations associated to each arrow of the quiver linear map. A framed version of this was initially introduced by Kronheimer and Nakajima, amounting to doubling the set of vertices, and drawing a new arrow from each new vertex to its corresponding old one. An example of the resulting quiver (which is of the prototypical type of interest for us) is depicted¹⁰⁷ in figure 83.

Framed representations appear naturally in ADHM quiver construction of self-dual or anti-self-dual YM on S^4 . This is also interesting from the point of view of representation theory of Lie algebras because dimension vectors of the framed vertices appear as highest weights of the representations. There are different notions of framing. Using Nakajima’s version for quiver varieties, the framed quiver is doubled, meaning each arrow gets doubled by an arrow that goes in the opposite direction. The linear space of representations becomes a linear cotangent bundle

$$M(\mathcal{Q}, v, w) \stackrel{def.}{=} T^* L(\mathcal{Q}^{fr}, v, w), \tag{E.1}$$

where $\mathcal{Q}, \mathcal{Q}^{fr}, v, w$ respectively denote the original quiver, the framed quiver, the number of original vertices, and the number of framed vertices. The gauge group is a general linear group on the original vertices $G \equiv G_v$, and there is a moment map

$$\boxed{\mu : M(\mathcal{Q}, v, w) \rightarrow \mathfrak{g}^*}. \tag{E.2}$$

Nakajima quiver varieties are Hamiltonian reductions of the following action

$$G \curvearrowright M(\mathcal{Q}, v, w), \tag{E.3}$$

and come into two types:

¹⁰⁷The nomenclature featuring in the framed quiver will be explained in section P.2, but we emphasise that this is standard notation in the quiver literature.

1. the *affine* Nakajima variety, defined as the partial character variety

$$\mathfrak{M}^0(\mathcal{Q}, v, w) \equiv \mu^{-1}(0)//G, \quad (\text{E.4})$$

2. *quasi-projective*

$$\mathfrak{M}^\chi(\mathcal{Q}, v, w) \equiv \mu^{-1}(0)//_\chi G. \quad (\text{E.5})$$

The gauge group by which we take the quotient is

$$G \equiv G_v \stackrel{\text{def.}}{=} \prod_a GL_{v_a}(\mathbb{C}) \subset G_v \times G_w \quad (\text{E.6})$$

For any choice of the nontrivial character

$$\chi: G \rightarrow \mathbb{C}^\times \quad (\text{E.7})$$

there is a proper Poisson morphism

$$p: \mathfrak{M}^\chi(\mathcal{Q}, v, w) \rightarrow \mathfrak{M}^0(\mathcal{Q}, v, w), \quad (\text{E.8})$$

which is as a symplectic resolution of the singularities of \mathfrak{M}^0 .

In both cases, (E.4) and (E.5), $\mu^{-1}(0)$ denotes the fiber of zero through the moment map. The latter can be used as a representation scheme for the path algebra, \mathcal{A} , modulo the ideal, \mathcal{I}_μ ,

$$\mathcal{A} \stackrel{\text{def.}}{=} \overline{\mathbb{C}\mathcal{Q}^{\text{fr}}}/\mathcal{I}_\mu, \quad (\text{E.9})$$

of the framed doubled quiver

$$\mu^{-1}(0) \equiv \text{Rep}_{\mathbb{C}^v \otimes \mathbb{C}^w}(\mathcal{A}) \stackrel{\text{def.}}{=} \text{Rep}_{v,w}(\mathcal{A}), \quad (\text{E.10})$$

where we made use of the following shorthand notation

$$\mathbb{C}^v \stackrel{\text{def.}}{=} \bigoplus_a \mathbb{C}^{v_a}, \quad \mathbb{C}^w \stackrel{\text{def.}}{=} \bigoplus_a \mathbb{C}^{w_a}, \quad (\text{E.11})$$

For the purpose of our work, the main result of [344] is being able to relate such varieties to derived representation schemes, with the latter being obtained by means of an alternative resolution of the affine Nakajima quiver variety. The underlying reason for the importance of this resides with it being a sample realisation of homological mirror symmetry¹⁰⁸.

Prior to explaining the alternative resolution leading to derived representation schemes, we wish to emphasise that the correspondence that we are looking for, summarised in figure 66, [344], emerges from comparing invariants, and the required conditions for them to match at the two endpoints of the red arrow. From the symplectic resolution point of view, this requires flatness of the moment map, as explained in [344], following [365]. We refer the interested reader to such references for more detailed explanation regarding the definition and properties of flat moment maps. For our purposes, the crucial point is that when the invariants in the symplectic and derived scheme description match, we are dealing with a complete intersection¹⁰⁹.

$$\begin{array}{ccc}
& & L\Gamma(-) \\
\text{Ho}(\text{DGAff}_k^{op}) & \xrightleftharpoons{\quad} & \text{Ho}(\text{CDGA}_k^+) \\
& & \text{RSpec}
\end{array}$$

Figure 84.

Algebraic resolution

The technique of algebraic resolution was first put forward by [615], and consists in resolving the singularities of the representation schemes by introducing homological algebra.

Prior to explaining derived representation schemes, let us first briefly overview the notion of a differential graded (dg)-schemes. A dg-scheme is a pair $X \stackrel{def.}{=} (X_o, \mathcal{O}_{X,\bullet})$, with $X_o \equiv M(\mathcal{Q}, v, w)$ denoting the vector space of linear representations, and $\mathcal{O}_{X,\bullet}$ a sheaf of dg-algebras such that their zeroth homology reads as follows

$$\pi_0(X) \stackrel{def.}{=} \text{Spec}(H_0(\mathcal{O}_{X,\bullet})) \equiv \mu^{-1}(0). \quad (\text{E.12})$$

The derived representation scheme of the relative algebra¹¹⁰, \mathcal{A} , in a vector space V , is the object, [344],

$$\text{DRep}_V(\mathcal{A}) \in \text{Ho}(\text{DGAff}_k), \quad (\text{E.13})$$

obtained applying the following composition of functors

$$\text{DRep}_V(-) : \text{Alg}_S \longrightarrow \text{Ho}(\text{DGA}_S^+) \xrightarrow{L(-)_V} \text{Ho}(\text{CDA}_k^+) \xrightarrow{\text{RSpec}} \text{Ho}(\text{DGAff}_k), \quad (\text{E.14})$$

where

$$L(-) : \text{Ho}(\text{DGA}_S) \rightarrow \text{Ho}(\text{CDGA}_S) \quad (\text{E.15})$$

is the derived representation functor acting on the homotopy category, Ho , whose homology is the representation homology of a differential graded algebra, \mathcal{A} ,

$$H_\bullet(\mathcal{A}, V) \stackrel{def.}{=} H_\bullet(L(\mathcal{A})_V). \quad (\text{E.16})$$

On the other hand, RSpec in (E.14) denotes the derived spectrum, defining an equivalence on the homotopy categories

Altogether, (E.14) leads to

$$\text{DRep}_V(\mathcal{A}) = \text{RSpec}(L(\mathcal{A}_V)) = \text{Rep}_V(\mathcal{A}_{cof}) \quad (\text{E.17})$$

with

$$\mathcal{A}_{cof} \xrightarrow{\sim} \mathcal{A} \quad (\text{E.18})$$

¹⁰⁸We will be explaining this in due course.

¹⁰⁹The meaning of the latter will be explained in section P.2.

¹¹⁰Associated to the choice of a path in the framed quiver.

a cofibrant replacement. Different choices of cofibrant replacements lead to different models of (E.17). For the specific case of framed Nakajima quiver varieties, the derived representation scheme reads, [344],

$$\mathrm{DRep}_{v,w}(\mathcal{A}) = \mathrm{Spec}(L(\mathcal{A}_{v,w})) \in H_0(\mathrm{DGAff}_{\mathbb{C}}), \quad (\text{E.19})$$

with representation homology

$$H_{\bullet}(\mathcal{A}, v, w) \equiv H_{\bullet}(L(\mathcal{A})_{v,w}) \in \mathrm{CDGA}_{\mathbb{C}}^+, \quad (\text{E.20})$$

a graded commutative algebra. Denoting by $\{U_{\lambda}\}$ the irreducible representations of G , (E.20) explicitly reads as follows

$$H_{\bullet}(\mathcal{A}, v, w) \stackrel{\text{def.}}{=} \bigoplus_{\lambda} \mathrm{Hom}_G(U_{\lambda}, H_{\bullet}(\mathcal{A}, v, w)) \otimes U_{\lambda}, \quad (\text{E.21})$$

with $\mathrm{Hom}_G(U_{\lambda}, H_{\bullet}(\mathcal{A}, v, w))$ denoting its isotypical components, namely modules over

$$H_0(\mathcal{A}, v, w)^G \equiv \mathcal{O}(\mu^{-1}(0))^G. \quad (\text{E.22})$$

For each irreducible representation of G_v , the Euler character is defined as follows

$$\begin{aligned} \chi_T^{\lambda}(\mathcal{A}, v, w) &\stackrel{\text{def.}}{=} \sum_{i=0}^{\infty} (-1)^i [\mathrm{Hom}_G(U_{\lambda}, H_i(\mathcal{A}, v, w))] \\ &= \sum_{i=0}^{\infty} (-1)^i [H_i(L(\mathcal{A}))_{\lambda, v, w}^{G_v}] \in K_T(\mathfrak{M}^0), \end{aligned} \quad (\text{E.23})$$

where $T \stackrel{\text{def.}}{=} T_w \times T_l$ accounts for the framing, and

$$H_i(\pi) : H_i(\mathcal{A}_{\text{cof}}) \rightarrow H_i(\mathcal{A}) \quad (\text{E.24})$$

is an isomorphism such that \mathcal{A}_{cof} has no higher homologies, namely,

$$\begin{cases} H_0(\pi) : H_0(\mathcal{A}_{\text{cof}}) \xrightarrow{\sim} \mathcal{A} \\ H_i(\mathcal{A}_{\text{cof}}) \equiv 0, \forall i \geq 1. \end{cases} \quad (\text{E.25})$$

and

$$\pi : \mathcal{A}_{\text{cof}} \longrightarrow \mathcal{A} \quad (\text{E.26})$$

defines an acyclic fibration in $\mathrm{DGA}_{\mathbb{C}}^+$.

Hilbert series

One of the key results of [344] is that agreement in between invariants calculated from the Nakajima and derived representation schemes requires flatness of the moment map on the symplectic side, and vanishing higher homologies in the derived case. Importantly for us, such agreement involves integrated characters, also known as the Hilbert series (HS). The latter is a map which reads

$$\mathrm{HS}_T : \begin{cases} K_T(\mathfrak{M}^0) \longrightarrow K_T(\mathrm{pt}), \text{ if } \mathfrak{M}^0 \text{ is compact} \\ K_T(\mathfrak{M}^0) \longrightarrow \mathrm{Frac}(T), \text{ otherwise.} \end{cases} \quad (\text{E.27})$$

From this, the resulting Weyl integral formula reads

$$\text{HS}_T \left(\chi_T^G(\mathcal{A}, v, w) \right) \equiv \frac{1}{|G|} \int_G dg \text{HS}_{G \times T} \left(\chi_T^G(\mathcal{A}, v, w) \right). \quad (\text{E.28})$$

Upon choosing the variables in the maximal torus of the gauge group, $x \in T_v \subset G$, and equivariant variables $t \equiv (a, l) \in T \equiv T_w \times T_l$, (E.28) can be re-expressed as follows

$$\text{HS}_T \left(\chi_T^G(\mathcal{A}, v, w) \right) \equiv \frac{1}{|W|} \int_{T_v} dx \text{WF}(x) \frac{\prod_i (1 - l_1 l_2 b_i)}{\prod_j (1 - a_j)}, \quad (\text{E.29})$$

with b_i, a_i respectively denoting the weights of $M(\mathcal{Q}, v, w)^*$ and \mathfrak{g} .

F Derivation of generalised black hole (BH) entropy

F.1 "Replica trick" derivation




Figure 85.




Figure 86.

F.2 Rènyi and von Neumann entropies

Given 2 systems A and B both equipped with a field theory which, for the latter it is assumed to be, in turn, coupled to JT-gravity,

$$S_B = \ln Z_B = \frac{\phi_o}{4\pi} \left[\int_B R + 2 \int_{\partial B} K \right] + \frac{\Phi}{4\pi} \int_B (R - \Lambda) + \frac{\Phi_b}{2\pi} \int_{\partial B} K + \ln Z_{CFT}[g] \quad (\text{F.1})$$

the Rènyi entropies read

$$S = \text{tr} \rho_A^n = \sum_{i_1 \dots i_n} p_{i_1} \dots p_{i_n} \langle \psi_{i_1} | \psi_{i_2} \rangle \langle \psi_{i_2} | \psi_{i_3} \rangle \dots \langle \psi_{i_n} | \psi_{i_1} \rangle \quad (\text{F.2})$$




Figure 87. Taking n -copies of the gravitating universe over which the path integral is performed. The pictures at the centre and on the RHS, instead, show the completely disconnected and completely connected saddles, the latter being associated to Euclidean wormholes.

The path integral is assumed to be performed over all the possible topologies consistent with the same boundary conditions.

$$\text{tr} \rho_A^n = \frac{1}{Z_1^n} \left[e^{-I_{grav}[M_{disc}]} Z_{CFT}[M_{disc}] + e^{-I_{grav}[M_{conn}]} Z_{CFT}[M_{conn}] \right] \quad (\text{F.3})$$

with

$$M_1 = e^{-I_{grav}[M_{conn}]} Z(\beta) \quad (\text{F.4})$$

being the saddle point geometry for the gravitating system B .

$$\begin{aligned} \Rightarrow \text{tr} \rho_A^n &= \frac{1}{Z_1^n} \left[e^{-n I_{grav}[M_1]} \sum_i e^{-n \beta E_i} + e^{-I_{grav}[M_{conn}]} Z_{CFT}[M_{conn}] \right] = \\ &= \frac{1}{Z_1^n} \left[Z(n\beta) + e^{-I_{grav}[M_{conn}]} Z_{CFT}[M_{conn}] \right] \quad . \end{aligned}$$

To build M_{conn} , we define C_x as a portion of a Cauchy slice of B with linear size $2\pi x$. Then, by gluing cyclically n -disconnected components of M_{disc} along C_x forming a \mathbf{Z}_n -symmetric Euclidean wormhole topology obeying the gravitational e.o.m., the Rènyi entropy can be rewritten as follows

$$\text{tr} \rho_A^n = \frac{Z(n\beta)}{Z_1^n(\beta)} \left[Z(n\beta) + e^{-I_{grav}[M_{conn}]} Z_{CFT}[M_{conn}] \right] \quad (\text{F.5})$$

In this subsection, we apply the machinery outlined in the previous pages for deriving the explicit expression of the generalised entanglement entropy, showing the emergence of the contribution of the quantum extremal surface resulting from the quotiented boundary geometry.

As already mentioned, AdS/CFT and, more generally, the Holographic Principle, claims that




Figure 88. These pictures shows the emergence of fixed points under Z_n -symmetry of the boundary. On the right, the fixed point is depicted in red. Under the action of the generators of Z_n -symmetry, the green subregion is mapped to equivalent subregions. The nontrivial fixed point indicates the emergence of a nontrivial boundary for the CFT, which in turn carries d.o.f. associated to the degeneracy of the ground state, as already outlined in previous subsection.

$$Z_{bdry}(M, J) = Z_{bulk}(M, J) \quad , \quad (\text{F.6})$$

i.e., the bulk string partition function is equivalent to the generating functional of its boundary CFT dual.

Expecting a UV completion of the bulk theory, i.e. not pure gravity. However, when G_N is small, it is possible to approximate it as

$$Z_{bulk} = \sum_g e^{-I(g)} Z_{matter}(g) \quad (\text{F.7})$$

where g is a classical background with geometry g and

$$\frac{\delta}{\delta g} \left[e^{-I(g)} Z_{matter}(g) \right] = 0 \iff R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = 8\pi G \langle T_{\mu\nu} \rangle \quad . \quad (\text{F.8})$$

The partition function is determined by the saddle point extremising (F.8). In general, these solutions are hard to find even for AdS_3 , and even more for higher-dimensions. Finding g for arbitrary n turns out to be possible through the aid of the replica trick, which in turn relies upon the assumption that the boundary has a Z_n -symmetry, which in turn corresponds to a cyclic permutation of the $\hat{\rho}$ featuring in the boundary path-integral.

Assumptions:

1. in the dominant saddle, this Z_n -symmetry is also a symmetry of the bulk, i.e. not spontaneously-broken
2. the dominant saddles are all we need, i.e. it provides a good analytic continuation, but there might be jumps in n leading to discontinuities.

As a matter of fact, we can therefore restrict our treatment to *quotiented geometries*. In addition, we need to take into account the *fixed points*¹¹¹ of the quotiented geometry (i.e. the branch cuts). In the bulk a surface with a similar behaviour must also be defined, such that the bulk geometry is smooth everywhere except at the fixed points.

¹¹¹uniquely identified

Quotienting by Z_n , we end up with a conical singularity. Writing up the semi-classical action in terms of the data of the quotiented geometry, equation (F.7) becomes

$$Z_n = e^{-I_n(g^{(n)}, X^{(n)})} Z_{matter,n}(g^{(n)}, X^{(n)}) \quad , \quad Z_{matter,n} = \text{Tr}_{\bar{b}}(\rho_b^n) \quad . \quad (\text{F.9})$$

The full unquotiented geometry is a saddle-point of the gravitational action, therefore satisfies the extremisation

$$\frac{\delta}{\delta g} \left[e^{-I(g)} Z_{matter}(g) \right] = 0 \quad (\text{F.10})$$

$$\iff \frac{\delta}{\delta g^{(n)}} \left[e^{-I(g^{(n)})} Z_{matter,n}(g^{(n)}) \right] = 0 \quad , \quad \frac{\delta}{\delta X^{(n)}} \left[e^{-I_n(g^{(n)})} Z_{matter,n}(g^{(n)}) \right] = 0 \quad . \quad (\text{F.11})$$

A possible solution to (F.11) is of the form

$$Z_n = \exp \left[-n I_1(g^{(n)}) - (n-1) \frac{A(X^{(n)})}{4G_N} \right] \text{Tr}(\rho_b^n) \quad . \quad (\text{F.12})$$

At this point, we can ease the assumption that n is an integer, in particular $n = 1 + \epsilon$, with $\epsilon \ll 1$. Working to leading-order in ϵ , the partition function becomes

$$\begin{aligned} Z_{1+\epsilon} &= e^{-I(g^{(1)})} Z_{matter} \left[1 - \epsilon I_1(g^{(1)}) - \epsilon \frac{\partial g^{(n)}}{\partial N} \frac{\delta I_1(g^{(1)})}{\delta g^{(n)}} - \epsilon \frac{A(X^{(1)})}{4G_N} + \right. \\ &\quad \left. + \frac{\epsilon}{Z_{matter}} \frac{\partial g^{(n)}}{\partial N} \frac{\delta Z_{matter}(g^{(1)})}{\delta g^{(n)}} + \frac{\epsilon}{Z_{matter}} \text{Tr}(\rho_b \ln \rho_b) \right] \quad . \quad (\text{F.13}) \end{aligned}$$

At $n = 1$, i.e. with one copy of the geometry, the action does not depend on the fixed point and $\text{Tr} \rho_b$ doesn't depend on b . However, the location of $X^{(1)}$ is not determined at $n = 1$, but at leading-order- ϵ corrections. (F.13) can be further simplified by applying the semi-classical Einstein equations that read

$$\frac{\delta I^{(1)}}{\delta g} + \frac{1}{Z_{matter}} \frac{\delta Z_{matter}}{\delta g} = 0 \quad , \quad (\text{F.14})$$

therefore, we are left with

$$Z_1^{1+\epsilon} = e^{-I_1(g^{(1)})} Z_{matter} \left[1 - \epsilon I_1(g^{(1)}) + \epsilon \ln(Z_{matter}) \right] \quad . \quad (\text{F.15})$$

For the von Neumann entropy

$$\frac{Z_{1+\epsilon}}{Z_1^{1+\epsilon}} = 1 - \epsilon \frac{A(X)}{4G_N} + \epsilon \text{Tr}(\hat{\rho}_b \ln \rho_b) - \epsilon \text{Tr}(\hat{\rho}_b \ln Z_{matter}) \quad (\text{F.16})$$

$$= 1 - \epsilon \frac{A(X)}{4G_N} + \epsilon \text{Tr}(\hat{\rho}_b \ln \hat{\rho}_b) \quad (\text{F.17})$$

$$\Rightarrow S = \lim_{\epsilon \rightarrow 0} \left[-\frac{1}{\epsilon} \ln \frac{Z_{1+\epsilon}}{Z_1^{1+\epsilon}} \right] = \frac{A(X)}{4G_N} + S_{bulk}(X) \quad . \quad (\text{F.18})$$

For $n = 1$, X is nothing but the quantum extremal surface satisfying (F.10). Furthermore, among all possible X s choose the smallest. This two-step (extr-min) procedure is what defines the quantum extremal surface prescription. Notice that the deep point of the replica trick consists in treating n as an arbitrary parameter and expanding it around $n = 1$, such that $g^{(1)}$ parametrises the

background metric without backreactions dominates the path-integral. On the other hand, Z_{matter} across the whole derivation, is really meant to be the full path-integral for the matter content. It is therefore takes into account of all fluctuations in the matter sector, which is what enables to achieve the 1-loop correction to the pure gravity result in (F.18).

G Topological Quantum Field Theories (TQFTs)

The present section introduces topological field theories and their String Theory counterparts, namely Topological String Theory. The reason for introducing this is multifold. As already mentioned in the introduction, they constitute an essential tool for probing the mathematical structure of QFTs. The most important advancements in the field took place in 2 and 3-dimensions. Most importantly, the pioneering works of Atiyah, Schwarz, Segal and Witten¹¹². Sticking to such lower-dimensional setups might seem reductive, especially when having in mind higher-dimensional field theories. However, the great advantage of the 2 and 3D settings have proved to give rise to an extremely rich variety of studies, whose findings can be interestingly applied to higher dimensions as well. As mentioned in the Introduction, the AGT correspondence is a clear example of the utility of 2D TFTs for describing 4D $\mathcal{N} = 2$ SCFTs¹¹³.

G.1 Attempts and advancements

Topological field theories (TFTs) are a bridge between QFT and maths. In gauge theory and mathematical physics, a TQFT is a QFT calculating topological invariants. They are related to knot theory, the theory of 4-manifolds, and the theory of moduli spaces in algebraic geometry [147, 171, 445, 568].




Figure 89.

The adjective topological refers to the fact that correlation functions are independent from the spacetime metric. On Minkowski spacetime, topological theories are trivial when contracted to a point. Because of this, they need to be placed on a Riemann surface instead. They come in two main types:

1. Schwarz-type TQFTs, e.g. the BF-model

$$S = \int_M BF, \tag{G.1}$$

where B is an auxiliary field, whereas F is the derivative of a connection, A . This action is explicitly topological, given that the metric does not feature at all. The alternative formulation by Schwarz reads as follows

¹¹²The pioneering works on the subject are the following [170–177].

¹¹³We will come back to this in greater detail in Part V.

$$S = \int_M A \wedge dA. \tag{G.2}$$

An example of this is Chern-Simons theory, applied to knot invariants, to which we shall turn in the following section.

2. Witten-type TQFTs [topological 4D Yang Mills theory in 4D]. For a given 2-form, B , the corresponding action reads as follows

$$S = \int_M B \wedge \delta B, \tag{G.3}$$

- S has a symmetry, i.e. $\delta S = 0$ under it.
- $\delta^2 = 0$, i.e. the symmetry is exact.
- There are observables, $\{\mathcal{O}_i\}$, such that $\delta \mathcal{O}_i = 0$.
- $T^{\alpha\beta} = \delta G^{\alpha\beta}$ for any arbitrary tensor $G^{\alpha\beta}$.

2D TFTs are mostly related to mirror symmetry. On the other hand, 3D TFTs are much less understood, apart from Chern-Simons theory, which has no nontrivial local observables given that there are no boundary conditions. The only topological observables are Wilson loops (localised on 1D closed submanifolds). Kapustin, Rozansky and Saulina (KRS) were the first to have introduced nontrivial boundary conditions for a 3D TFT. This theory as a 3D sigma-model, first introduced by Rosanzky and Witten (a.k.a. the RW-model). The target of the TFT is required to be hyperkähler, X , or, more generically, a holomorphic symplectic manifold. Upon dimensional reduction on an S^1 , it yields the B-model.

A physical motivation for studying the RW-model is its relation with 3D mirror symmetry. In particular, understanding the RW-model, is a step forward towards understanding 3D mirror symmetry.

As we will see in greater detail, the boundaries of KRS are Langrangian submanifolds of X , with the latter building relation with derived algebraic geometry.

G.2 The Atiyah-Segal axioms

The Atiyah-Segal axioms are particularly useful for understanding the mathematical structure of Schwarz-type QFTs. The key point of such axioms is that a TQFT is a functor from a bordism category to the category of vector spaces.

G.2.1 Segal's Conformal Field Theory (CFT) axioms

The object of Segal's work, is that of presenting a mathematical formulation of CFT_{2S} , and describing basic examples already known from previous literature. Furthermore, he also presents five examples of areas of pure mathematics where CFTs play a crucial role:

1. The monster group.
2. The representation theory of loop groups, and the group $\text{Diff}(S^1)$ of diffeomorphisms on the circle.
3. How representations of $\text{Diff}(S^1)$ are related to the geometry of moduli spaces of the Riemann surfaces. (Mumford's classification of holomorphic line bundles descends from this.)
4. Some of V. Jones' new representations of braid groups, and his classification of subfactors in von Neumann algebras.
5. Elliptic cohomology theory of Lendweber-Stong, [191].

G.2.2 The Atiyah-Segal axioms

The key point of such axioms is that a TQFT is a functor from a bordism category to the category of vector spaces.

Take a commutative ring, Λ , with the identity, $\mathbf{1}$. Then, a D -dimensional TQFT¹¹⁴ is defined from Λ according to the following axioms:

- The Homotopy axiom A finitely-generated Λ -module, $\mathcal{Z}(\Sigma)$ associated to each oriented closed smooth d -dimensional manifold, Σ .
- The Additive axiom An element $\mathcal{Z}(M) \in \mathcal{Z}(\partial M)$ associated to each oriented smooth $(d+1)$ -dimensional manifold M .
- \mathcal{Z} is functorial with respect to orientation-preserving diffeomorphisms of Σ and M .
- \mathcal{Z} is involutory, i.e. $\mathcal{Z}(\Sigma^*) = \mathcal{Z}(\Sigma)^*$.
- \mathcal{Z} is multiplicative.
- $\mathcal{Z}(\emptyset) = \Lambda$ for the d -dimensional manifold, and $\mathcal{Z}(\emptyset) = \mathbf{1}$ for the $(d+1)$ -dimensional empty manifold.
- The Hermitian axiom $\mathcal{Z}(M^*) = \overline{\mathcal{Z}(M)}$.

In practice, Σ and $\mathcal{Z}(\Sigma)$ respectively correspond to the physical space and the Hilbert space of the QFT, with associated vanishing Hamiltonian, $H = 0$. The latter property follows from the topological nature of the theory in question, implying there is no real dynamics in the $\Sigma \times I$ -cylinder. However, there can still be nontrivial propagation from Σ_o to Σ_l through an intermediate, topologically nontrivial, manifold M such that

$$\partial M = \Sigma_o^* \cup \Sigma_l. \quad (\text{G.4})$$

If $\partial M \equiv \Sigma$, then the distinguished vector in the Hilbert space $\mathcal{Z}(M)$ is the vacuum state defined by M . For a closed manifold M , $\mathcal{Z}(M)$ is nothing but the partition function, namely the expectation value of the identity operator.

The additive axiom can be understood by taking into account the fact that finite dimensional vector spaces for a compact closed category. For genuine TFTs, all spaces of quantum states are finite-dimensional, hence one could equivalently consider the linear dual spaces, implying the propagator (or correlator) map

$$Z(M) : Z(\partial_{in} M) \rightarrow Z(\partial_{out} M) \quad (\text{G.5})$$

becomes a linear map

$$\mathbb{C} \rightarrow Z(\partial_{out} M) \otimes Z(\partial_{in} M)^* = Z(\partial M). \quad (\text{G.6})$$

G.3 Schwarz-type TQFTs

G.3.1 Chern-Simons (CS) Theory

A 3D TQFT of Schwarz-type, developed by Edward Witten, and named after Chern and Simons who introduced the Chern-Simons 3-form. The latter is defined as follows: given a manifold and a Lie algebra valued 1-form, A , over it, one can define a family of p -forms as

¹¹⁴Where $D = d + 1$, d denoting the number of spatial dimensions.

$$d\omega_{2k-1} = \text{Tr} \left(F^k \right), \quad (\text{G.7})$$

where the right-hand-side is proportional to the k^{th} Chern-character of the connection A . For the specific case where $p = 3$, (G.7) reads

$$\text{Tr} \left[F \wedge A - \frac{1}{3} A \wedge A \wedge A \right] = \text{Tr} \left[dA \wedge A + \frac{2}{3} A \wedge A \wedge A \right]. \quad (\text{G.8})$$

In gauge theory, the integral of the Chern-Simons form is a global geometric invariant, modulo addition of an integer.

G.4 Topological String Theory

Topological String Theory (TST) is a version of String Theory that follows the idea of TQFTs. There are 2 main versions of it, conventionally referred to as the topological A- and B-model.

Calculations in TST encode all holomorphic quantities that are protected by spacetime SUSY. Various such calculations are related to CS-theory, Gromov-Witten (GW)-invariants, mirror symmetry, and the Geometric Langlands Program, just to name a few.

The operators in a TST represent the algebra of operators in the full string theory preserving a certain amount of SUSY.

Abelian gauge theories usually appear in String and M-theory, where abelian means that, in these theories, the space of gauge inequivalent fields forms an abelian group.

A central extension of a group G by an abelian group A is a group \tilde{G} such that

$$0 \rightarrow A \rightarrow \tilde{G} \rightarrow G \rightarrow 0, \quad (\text{G.9})$$

with A being in the center of \tilde{G} . \tilde{G} can be thought of as a set of pairs (g, a) with twisted multiplication

$$(g_1, a_1) (g_2, a_2) = (g_1 g_2, c(g_1, g_2) a_1 a_2). \quad (\text{G.10})$$

From a purely mathematical perspective, it is natural to question about the implications of this for the formulation of topological D-branes in terms of derived categories.

G.4.1 The A- and B-model

- The A-model

The topological A-model comes with a 6-dimensional Kähler target space. Such theory is equipped with fundamental strings wrapping around 2D holomorphic curves, as well as D2-branes wrapping Lagrangian submanifolds of spacetime. The worldvolume theory on a stack of N D2-branes is the string field theory of the open strings of the A-model, namely a $U(N)$ -CS-theory.

The fundamental strings may, in turn, end on the D2-branes. Additional (coisotropic) branes in various dimensions might also be added [Kapustin, Orlov].

- The B-model

The B-model also contains fundamental strings. However, scattering amplitudes here only depend on the complex structure, and are completely independent w.r.t. the Kähler structure. Furthermore, the theory admits D(-1)-, D1-, D3-, and D5-branes, wrapped around holomorphic 0, 2, 4, and 6-submanifolds, respectively.

Mirror symmetry relates the amplitudes calculated on the two sides.

H QFT and the Jones polynomial

H.1 QFT and the Jones polynomials

Among the many useful applications of CS-theory as a 3D TQFT of Schwarz-type, is that it interprets the Jones polynomials (JPs).

JPs assign a Laurent polynomial in $t^{1/2}$ to any knot, $V(knot)$. Each JP can thus be completely determined by the unknot, $V(unknot) = 1$, and the Skein relations

$$\left(t^{1/2} - t^{-1/2}\right) V(L_o) = t^{-1} V(L_+) - t V(L_-), \quad (\text{H.1})$$

where L_{\pm}, L_o denote knots differing in between them only at a point, as shown in figure H.1.




Figure 90.

H.2 Schemes or algebraic stacks

A scheme is a mathematical structure that enlarges the notion of an algebraic variety, accounting for multiplicities, and allowing varieties over any commutative ring. Scheme theory was originally introduced by A. Grothendieck in 1960 as a key tool for solving open issues in algebraic geometry.

A scheme is a topological space, as well as a ringed space which is locally a spectrum of a commutative ring.

A morphisms of schemes can be used for defining algebraic surfaces. In many cases, the family of all varieties of a given type can itself be viewed as a variety of schemes.

H.3 Morse Theory and Classifying Spaces

The main idea of Morse Theory is to associate to a Morse functor

$$f : M \rightarrow \mathbf{R} \quad (\text{H.2})$$

on a closed Riemannian manifold, M , a category, \mathcal{C}_f , whose objects are the critical points of f , whereas the morphisms, $\overline{\mathcal{M}}(a, b)$, between two critical points, a and b , are piece-wise flow lines of the gradient flow of f connecting a and b . One can recover from the topological category \mathcal{C}_f the topology of M . As such, $\mathcal{M}(a, b)$ denotes the the moduli space of lines from a to b .

From local data, one cannot tell whether a given critical point is required by the topology or if, instead, it is removable. For the system in question to be nontrivial, we need to perform calculations that are sensitive of the presence of more than one critical point on the manifold in question. For this to be the case, we need to ensure that the potential energy in the problem, $V(\phi)$, has more than one minimum, specifically, one for each critical point, allowing for the possibility of tunnelling from one critical point to another.

Tunnelling can be calculated in the WKB-approximation, namely by means of instantons (using Witten's terminology), [444]. Tunnelling effects often remove spurious degeneracies arising

in perturbation theory, as is precisely the case here. From the instanton analysis, we know that the relevant instantons or tunnelling paths are the paths of steepest descent leading from one critical point to another. As such, they are therefore the solutions of the following equation

$$\frac{d\phi}{d\lambda} = \gamma^{ij} \frac{\partial h}{\partial \phi_j}, \quad (\text{H.3})$$

where λ denotes time in the supersymmetric nonlinear sigma-model, assuming ϕ is only a function of time.

The instanton calculation shows that the only relevant solutions of this equation are the ones connecting two critical points whose Morse indices differ by unity.

Instanton solutions or tunnelling paths in a given theory are extrema of this Lagrangian, written with a Euclidean metric, neglecting the fermionic content. The relevant action would therefore read

$$\begin{aligned} S &= \frac{1}{2} \int d\lambda \left(\gamma^{ij} \frac{d\phi^i}{d\lambda} \frac{d\phi^j}{d\lambda} + t^2 \gamma^{ij} \frac{\partial h}{\partial \phi^i} \frac{\partial h}{\partial \phi^j} \right) \\ &= \frac{1}{2} \int d\lambda \left| \frac{d\phi^i}{d\lambda} \pm t \gamma^{ij} \frac{\partial h}{\partial \phi^j} \right|^2 \mp t \int d\lambda \frac{dh}{d\lambda}. \end{aligned} \quad (\text{H.4})$$

The minimum action paths between any two critical points a and b are paths of steepest descent, and the action for such paths reduces to

$$S = t \left| h(B) - h(A) \right|. \quad (\text{H.5})$$

An example of a classifying space for the ∞ -cyclic group G is the circle as X . Moduli space is essentially a synonym for representing object and classifying space. Usually, the term classifying space is used in topology, whereas moduli space is used in algebraic geometry.

H.4 Donaldson invariants

Donaldson theory (DT), [445], is the study of the topology of 4-manifolds using moduli spaces of anti-self-dual instantons. Many of the theorems in DT can be proved by a much more convenient formalism, namely that of Seiberg-Witten (SW) Theory.

Even though this has been amply studied in the literature, there are still some open questions that remain to be addressed, phrased in terms of the following conjectures:

1. The *Witten conjecture*¹¹⁵: in Algebraic Geometry, it is related to the intersection numbers of stable classes on the moduli space of curves. In its formulation, it is a special case of a more general relation between integrable systems of Hamiltonian PDEs, and the geometry of certain 2D TFTs, also known as cohomological field theories¹¹⁶.
2. The *Atiyah-Floer conjecture*: connecting the instanton Floer homology with the Lagrangian intersection Floer homology¹¹⁷.

Witten showed that Donaldson invariants are correlation functions in a supersymmetric gauge theory, and that the infinite-dimensional Morse theory developed by Floer is part of a 2D QFT.

Mathematical gauge theory was later revolutionised when Seiberg and Witten deduced the long-distance behaviour of the supersymmetric gauge theory calculating the Donaldson invariants.

¹¹⁵The Witten conjecture was later extended by Virasoro.

¹¹⁶The latter were first studied by Kontsevich and Manin.

¹¹⁷The latter is, in turn, related to homological mirror symmetry.

Donaldson invariants are topological in nature, and are therefore scale-independent. Because of this, they found an alternative way of calculating them, this time by means of the so-called *Seiberg-Witten equations*.

The original motivation that led Witten to propose such conjecture was the existence of two different 2D models of quantum gravity that should be exhibiting the same partition function. These are: on the one hand, the intersection of number on the moduli stack of algebraic curves, and, on the other, it is the ln of of the T-function of the KdV-hierarchy, with T a translation operator such that

$$T(g)(x) = g(x + 1). \tag{H.6}$$

H.5 Seiberg-Witten theory

A Seiberg-Witten theory is an $\mathcal{N} = 2$ supersymmetric gauge theory whose low-energy effective action is exact in the massless limit. Such theory is also referred to as $\mathcal{N} = 2$ SYM theory, whose field content is a single $\mathcal{N} = 2$ vector supermultiplet, in complete analogy with the single vector gauge field (or connection) in ordinary YM theory. Its corresponding Lagrangian density in 2D reads

$$\mathcal{L}_{SYM} = \text{ImTr} \left(\frac{1}{4\pi} \int d^2\theta d^2\vartheta \mathcal{F}(\Psi) \right), \tag{H.7}$$

with θ, ϑ denoting the coordinates for the spinor directions of superspace, whereas \mathcal{F} is the prepotential. The minimal theory is defined by

$$\mathcal{F}(\Psi) = \frac{1}{2} \tau \Psi^2, \tag{H.8}$$

with τ the complex coupling constant.

H.6 From CS to Morse theory

H.6.1 Morse Theory

Morse Theory relates to the description of closed geodesics on smooth Riemannian manifolds. Since the first studies of R. Bott, the theory underwent several reformulations, which we will briefly overview in this section. The crucial feature of Morse Theory is its relation to the so-called critical-point theory.

The main idea is that of associating a Morse function

$$f : M \rightarrow \mathbf{R} \tag{H.9}$$

on a closed Riemannian manifold, M , a category, \mathcal{C}_f , whose objects, a, b , are the critical points of f , whereas the morphisms between two critical points of f are denoted by $\overline{\mathcal{M}}(a, b)$. The latter are piecewise flow lines of the gradient flow of f connecting a and b . One can recover the topology of M from the category \mathcal{C}_f . $\mathcal{M}(a, b)$ is the moduli space of flow-lines from a to b . This setup can be rephrased in terms of the space of loops, [A](#).

Morse Theories are classified in terms of:

1. Their dimensionality, which can either be finite or infinite.
2. The role of symmetry groups.
3. Their connection to higher or lower-dimensional TQFTs.

Its finite formulation, due to Bott, followed from approximating the description in terms of a finite set of polygons (i.e. piecewise geodesics). Furthermore, he restricted his studies on manifolds with Lie group acting on the manifold while preserving the Morse function, also known as *equivariant* Morse Theory.

On the other hand, Morse Theory in its infinite-dimensional version has many fascinating applications in lower-dimensional topology, to which we will come back momentarily.

H.6.2 Relation to Moduli Spaces of SQFTs

The most important feature of Morse Theory is its applicability to extended setups going beyond the work of Atiyah and Bott on Riemann surfaces.

Take a compact Lie group G , and M a smooth manifold. A G -connection on M gives a notion of parallel transport along paths, which can be considered as an element of G itself. For example, if $M \equiv S^2$, then parallel transport on a half-circle defines a map

$$S^1 \longrightarrow G, \tag{H.10}$$

relating the loop space of G and connections on the Riemann surfaces, the latter being the original setup investigated by Atiyah and Bott, namely the equivariant Morse Theory of the Yang-Mills (YM) functional

$$YM(A) = \frac{1}{2} \int_M |F_A|^2 d^D x. \tag{H.11}$$

To determine the critical points of (H.11), one needs to solve the associated Euler-Lagrange equation

$$d_A^* F_A = 0. \tag{H.12}$$

The minima of the energy functional (H.11) are flat connections with zero-curvature, which, from the Narasimhan-Seshadri Theorem corresponds to the space of stable bundles. Later developments in 4D by Atiyah, [170], Bott, [555], and Donaldson, [445], eventually lead to the study of instanton equations in 4D, further developed by Atiyah, Hitchin, and Singer, [98].

A great turnaround came when Floer started investigating 3D manifolds, [479], in which case the functional to be extremised is CS theory. The critical points of the functional are still flat connections on the given Riemannian manifold, but a crucial difference arises with respect to the 2D and 4D cases: the number of negative eigenvalues of the Hessian at the critical points is infinite-dimensional, implying traditional Morse Theory could not be applied straightforwardly. Importantly, Floer realised that one could define a *relative index* in between two critical points, by making use of a curve interpolating in between them. In such way, one is able to define chain complex, whose homology defines the Floer homology of the 3-manifold in question.

Furthermore, the latter can be related to the Donaldson invariants of a 4D manifold whose boundary is the 3D manifold whose Floer homology has just been defined. Indeed, Floer homology groups and Donaldson invariants follow some of the defining properties of a TQFT, such as the gluing property.

2 main lines of development in Morse Theory, [444]:

1. Witten was first motivated by supersymmetry models in lower-dimensional QFT.
2. The second line of development has been, instead, that of understanding that Morse functions decompose smooth manifolds into simple constituents.




Figure 91.

Mathematical constructions of TQFTs usually start with algebraic data which is then assigned to elementary bordisms between manifolds of fixed dimension. Invariants are obtained by decomposing arbitrary bordisms into elementary ones by means of Morse functions. The fact that this defines an invariant follows from comparing the results achieved from different Morse functions.

For the case of a 2D TFT, such algebraic data is a commutative Frobenius ring, first studied by Moore and Segal.

SQFTs from Morse Theory

We now briefly outline how Witten's approach shed light on the relation in between supersymmetric QFT (SQFT) and Morse Theory, from a mathematical perspective¹¹⁸.

Let us start by taking a smooth compact manifold M , and a smooth function

$$f : M \longrightarrow \mathbb{R}. \tag{H.13}$$

Then, introducing a deformation

$$0 \longrightarrow \Omega^1(X) \xrightarrow{d+t\epsilon(df)} \Omega^2(X) \xrightarrow{d+t\epsilon(df)} \Omega^3(X) \xrightarrow{d+t\epsilon(df)} \dots \tag{H.14}$$

of the ordinary de Rham complex, if f is a Morse function, and M is endowed with a Riemannian metric, then, in the $\lim_{t \rightarrow \infty}$, there is a close relation between the spectrum of the associated Laplace operator, Δ_t , and the critical points of f . From a physical point of view, these correspond to the different vacuum states of the quantum system. Quantum tunnelling in between them can be encoded as a differential on the \mathbb{Z} -graded abelian group generated by the vacuum states, or the critical points, namely where the grading is given by the Morse index. Smale was the first to realise this from a pure mathematical perspective, but Witten was the first to start the physical studies behind this, with the relevant system in question being that of supersymmetric quantum mechanics (SQM).

Once having a consistent understanding of SQM, one could lift the description by, either, increasing the number of spacetime dimensions, and/or the amount of supersymmetries at play, and subsequently studying its impact on the associated Morse Theory.

For the sake of completeness, we briefly outline a few examples of the statement we have just made.

1. Starting from SQM, doubling the amount of supersymmetry while preserving the dimensionality implies removing flows between the critical points of the associated Morse function. Upon raising the theory to 2D QFT (namely Landau-Ginzburg theory), one recovers nontrivial interpolating soliton solutions, with the latter being related to Picard-Lefschetz theory.

¹¹⁸See appendix E for descriptions of gauge theories in mathematical language.

2. For the case of 4D supersymmetric gauge theories, instead, one can study the tunnelling between vacua on a 3D manifold in terms of Floer homology.
3. In higher dimensions, with increasing amount of supersymmetry, namely 5D and 6D SCFTs, Witten proposed a new definition of the Jones polynomial and its categorification in terms of Khovanov homology. In this case, the critical point equations are special cases of the Kapustin-Witten equations introduced in the context of the Geometric Langlands Program (GLP), whereas the flow equations turn out being new equations in a 5D mathematical gauge theory.

H.6.3 The Cobordism Hypothesis

More generally to this has been the formulation of the so-called *cobordism hypothesis*. The Cobordism Hypothesis was first formulated by Baez and Dolan, and later proved by Hopkins and Lurie in 2D, and eventually in $D > 2$ from Lurie.

The Cobordism Hypothesis states that, given a symmetric monoidal (∞, n) -category, \mathcal{C} , which is fully dualisable and every k -morphism, for $1 \leq k \leq n - 1$, there is a bi-jection between the \mathcal{C} -valued symmetric monoidal functors of the cobordism category and the objects of \mathcal{C} .

The motivation for this is a statement we have already introduced in our treatment, namely that symmetric monoidal functors from the cobordism category correspond to TQFTs. In practice, the Cobordism Hypothesis is very similar in nature to the Eilenberg-Steenrod axioms for homology theories, with the latter essentially stating that, a homology theory is uniquely determined by its value from the point. In the cobordism case, this corresponds to saying that the bijection between \mathcal{C} -valued symmetric monoidal functors and the objects of \mathcal{C} is uniquely determined by its value for the point.

I 4D descendants of 6D $\mathcal{N} = (2, 0)$ SCFTs

I.1 Class \mathcal{S}

First, we briefly overview the properties of class \mathcal{S} theories, the calculation of the dimension of the Coulomb and Higgs branches, as well as the contribution from punctures to the 4D central charges, and how 3D SCFTs can be accommodated on nontrivial punctures. Then, we recall the basic motivations underlying the 2D/4D correspondence resulting from dimensional reduction of 6D $\mathcal{N} = (2, 0)$ SCFTs, needed in core sections of the present work. We end the section by recalling that the 3D theories resulting by twisted compactification on the non-invertible defects of 4D $\mathcal{N} = 4$ SYM are always of the ABJ-type, namely gauge theories with $U(M) \times U(N)$ gauge group, [332].

$\mathcal{N} = 2$ data

1. The moduli space: Coulomb, Higgs and mixed branches.
2. The charge lattice, Γ , with

$$\dim \Gamma \equiv 2r + f \tag{I.1}$$

where r denotes the rank of the gauge group and f the rank of the gauge flavour symmetry. BPS states are points in Γ .

3. An antisymmetric electromagnetic inner product on Γ

$$\circ : \Gamma \times \Gamma \longrightarrow \mathbb{C} \tag{I.2}$$

4. A central charge function

$$Z_u : \Gamma \longrightarrow \mathbb{C} \quad (\text{I.3})$$

where u denotes a generic point in the Coulomb branch.

Such data is conventionally encoded in a family of Riemann surfaces, $\{\Sigma_u\}$, known as *Seiberg-Witten curve*, which varies over the Coulomb branch, and whose homology lattice is identified with Γ . By definition, (I.3) is linear in the charges, and can therefore be expressed as a 1-form integral of the *Seiberg-Witten differential*, λ_u

$$Z_u(\gamma) \stackrel{\text{def.}}{=} \int_{\gamma} \lambda_u \quad , \quad \forall \gamma \in \Gamma \quad (\text{I.4})$$

Consequently, the electro-magnetic inner product (I.2) is naturally identified with the intersection of 1-cycles. Jointly, (Σ_u, λ_u) define the complete solution to the low-energy physics. On the Coulomb branch, this description is provided by a special Kähler geometry.

BPS quivers

The purpose of the present work requires restricting to *complete theories*. These include:

- Theories associated to triangulation surfaces.
- The quivers $E_n, \hat{E}_n, \hat{\hat{E}}_n$.
- The Derksen-Owen quivers, X_6, X_7 .

The natural relation between quivers and triangulations is manifest in terms of BPS state counting.

Assuming the theory of interest admits a set of $2r + f$ hypermultiplett states, $\{\gamma_i\}$, forms a positive integral basis for all particles, such as is the case for class \mathcal{S} theories, a BPS quiver is constructed by drawing a node $\forall \gamma_i$, and an arrow in between them, $\gamma_i \rightarrow \gamma_j$ if $\gamma_i \circ \gamma_j > 0$. If a site of the charge lattice

$$\gamma \equiv \sum_i n_i \gamma_i \in \Gamma \quad (\text{I.5})$$

is occupied by a BPS state, it is possible to determine its degeneracy and spin. The interactions of the BPS elementary particles are encoded in the 4-supercharge quiver QM of the given configuration, i.e. a 1D supersymmetric gauge theory with gauge and matter groups

$$G \stackrel{\text{def.}}{=} \prod_{\text{nodes}} U(n_i) \quad , \quad \text{Matter} \stackrel{\text{def.}}{=} \bigoplus_{\text{arrows}} B_{ij}^a \quad (\text{I.6})$$

where B_{ij}^a denotes the bifundamental field charged under G . To find the BPS states, we need to study the moduli space of supersymmetric ground states, \mathcal{M}_γ ,

$$\mathcal{M}_\gamma \stackrel{\text{def.}}{=} \left\{ R \equiv \left\{ B_{ij}^a : \mathbb{C}^{n_i} \rightarrow \mathbb{C}^{n_j} \right\} \mid \frac{\partial \mathcal{W}}{\partial B_{ij}^a} \equiv 0, R \text{ } \Pi \text{-stable} \right\} / \prod_i Gl(n_i, \mathbb{C}) \quad (\text{I.7})$$

Hence, \mathcal{M}_γ is the space of solutions to the F -term equations

$$\frac{\partial \mathcal{W}}{\partial B_{ij}^a} \equiv 0 \quad (\text{I.8})$$

subject to an additional stability constraint, also known as Π -stability, modulo the action of the complexified gauge group $Gl(n_i, \mathbb{C})$. Π -stability is the relevant condition for describing moduli space. In quiver representation theory, it means that, a representation R of a quiver Q is a complex vector space $\mathbb{C}^{n_i} \forall$ node i and a linear map $B_{ij}^a : \mathbb{C}^{n_i} \rightarrow \mathbb{C}^{n_j} \forall$ arrow a from i to j , corresponding to a choice of expectation values of matter fields in the corresponding quiver QM.

A subrepresentation S of R is a choice of vector subspaces $\mathbb{C}^{m_i} \subset \mathbb{C}^{n_i}$ for each node, and maps $b_{ij}^a : \mathbb{C}^{m_i} \rightarrow \mathbb{C}^{m_j}$ for each arrow, such that all diagrams look like the following

$$\begin{array}{ccc} \mathbb{C}^{n_i} & \xrightarrow{B_{ij}^a} & \mathbb{C}^{n_j} \\ \uparrow & & \uparrow \\ \mathbb{C}^{m_i} & \xrightarrow{b_{ij}^a} & \mathbb{C}^{m_j} \end{array}$$

Figure 92.

For a given charge associated to a representation R

$$\gamma_R \equiv \sum_j n_j \gamma_j \quad (\text{I.9})$$

the stabilisation condition corresponds to

$$\arg Z_u(S) < \arg Z_u(R) \quad (\text{I.10})$$

for S a subrepresentation of R . Upon identifying S with a parton state, Π -stability corresponds to forbidding decay channels of the kind $\gamma_R \rightarrow \gamma_S$, from which the definition (I.7) results.

The nodes of a BPS quiver are always Π -stable representations, due to the absence of destabilising subrepresentations. In terms of (I.9), this corresponds to setting $n_i \equiv \delta_{ij}$, meaning each node contributes with a multiplicity 1 hypermultiplet BPS state. States of higher spin or multiplicity do not turn into nodes in the quiver.

If \mathcal{M}_γ is nonempty, there is a BPS particle in the spectrum with charge γ . Since \mathcal{M}_γ has 4 supercharges, it is a Kähler manifold, and its cohomology forms representations of Lefschetz $SU(2)$. If \mathcal{M}_γ is a point, it defines a hypermultiplet. If, instead, $\mathcal{M}_\gamma \simeq \mathbb{P}^1$, it defines a vector multiplet.

BPS-wall crossing arises from the Π -stability condition if at some point u in \mathcal{M}_γ , $Z_u(S) \equiv Z_u(R)$, with S a subrepresentation of R .

I.2 Vacuum structure: Coulomb, Higgs and mixed branches

The full moduli space is unknown in most cases. For this reason, in the present work we will only be focussing on class \mathcal{S} -theories obtained by twisted compactification on a Riemann surface of 6D $\mathcal{N}=(2,0)$ on A_{N-1} , with generic moduli space

$$\mathcal{M} \stackrel{\text{def.}}{=} \bigcup_{\alpha} C_{\alpha} \times H_{\alpha} \quad (\text{I.11})$$

with C_{α}, H_{α} denoting the *Coulomb* and *Higgs branches* of the theory.

The former is defined by the Seiberg-Witten curve

$$x^N + \sum_{i=2}^N \phi_i(z) x^{N-i} = 0 \quad (\text{I.12})$$

where z is the coordinate of the base Riemann surface, and x is the one on the fiber of the cotangent bundle $K \equiv T^*C$. $\phi_i(z)$ is a section of the line bundle, $K^{\otimes i}$, namely it is the i^{th} differential on the Riemann surface C . In absence of punctures, its contribution to the dimension of the Coulomb branch reads

$$\dim(\phi_i) \equiv \dim H^0(C, K^{\otimes i}) = (2i-1)(g-1) \quad (\text{I.13})$$

The general expression in presence of regular punctures reads

$$\dim_{\mathbb{C}} C_{\alpha} \equiv (k-1)g + \sum_{s=1}^k (n_s^2 - 1)(g-1) + \sum_p \dim_{\mathbb{C}} C_{\alpha}(p) \quad (\text{I.14})$$

where k denotes the number of elements featuring in the partition $X \equiv [n_1, \dots, n_k]$, and

$$\dim_{\mathbb{C}} C_{\alpha}(p) \equiv \sum_{s=1}^k d(Y_s^D) = \sum_{s=1}^k \sum_{i=2}^k p_i \equiv \frac{1}{2} \sum_{s=1}^k \left(N^2 - \sum_{a=1}^{\ell} m_a^2 \right) \quad (\text{I.15})$$

where Y^D denotes the local contribution from a regular puncture.

The total dimension along the Higgs branch, instead, reads

$$\dim_{\mathbb{H}} H_{\alpha} \equiv k-1 + \sum_p \dim_{\mathbb{H}} H_{\alpha}(p) \quad (\text{I.16})$$

with

$$\dim_{\mathbb{H}} H_{\alpha}(p) \equiv d(Y') - d(Y) \quad (\text{I.17})$$




Figure 93. The vacuum structure of $\mathcal{N} = 2$ class \mathcal{S} theories, comprising a Coulomb (C), Higgs (H), and mixed branch.

Upon choosing a partition of N : $X \equiv [n_1, \dots, n_k]$, the maximally factorised SW curve can be expressed as follows

$$\prod_{s=1}^k \left(x^{n_s} + \sum_{i=2}^{n_s} \phi_{s,i}(z) x^{n_s-i} \right) = 0 \quad (\text{I.18})$$

I.3 Metrics

We now briefly overview the definition of the metric on the Riemann surface, following the work by Gaiotto, Moore, and Nietzke.

Take \mathcal{B} to be the Coulomb branch of a certain 4D $\mathcal{N} = 2$ supersymmetric gauge theory. At any point $u \in \mathcal{B}$, the gauge group is broken to a maximal torus $U(1)^r$. Denoting by $\hat{\Gamma}_u$ the charge lattice equipped with a pairing \langle, \rangle , with radical being the flavour charge lattice, $(\Gamma_{flav})_u$, the quotient charge sublattice of gauge charges can be defined as follows

$$\Gamma_u \stackrel{def.}{=} \hat{\Gamma}_u / (\Gamma_{flav})_u, \quad (\text{I.19})$$

with $\text{rank} \Gamma_u = 2r$.

The central charge homomorphism is

$$Z : \hat{\Gamma}_u \longrightarrow \mathbb{C}. \quad (\text{I.20})$$

Then, for every particle, one can define $Z_\gamma(u)$, with γ a local section of $\hat{\Gamma}$, as denoting the central charge of a particle of charge γ .

For a theory placed on $\mathbb{R}^3 \times S^1$, with R denoting the radius of the S^1 , for $R \ll 1$, the 4D theory is effectively 3D, and its moduli space is locally a product of hyperkähler manifolds

$$\mathcal{M}_{tot} = \text{HB}_{4D} \times \text{CB}_{3D} \quad (\text{I.21})$$

At generic nonsingular points of the 3D CB, the HB is absent, so one can just restrict to the 3D CB

$$\mathcal{M} \stackrel{def.}{=} \text{CB}_{3D}, \quad (\text{I.22})$$

such that one can define the map

$$\pi : \mathcal{M} \longrightarrow \mathcal{B}. \quad (\text{I.23})$$

That the metric on \mathcal{M} is hyperkähler follows from the theory being supersymmetric.

The main idea proposed by [130, 132, 132] is that of describing the metric by introducing holomorphic Darboux coordinates for \mathcal{M} (when viewing the latter as a holomorphic symplectic manifold).

Fixing $U \subset \mathcal{B}$ over which $\hat{\Gamma}$ can be trivialised, the holomorphic Darboux coordinates, $\chi_\gamma^{\vartheta, u_o}$, with $u_o \in U \subset \mathcal{B}$, are labelled by sections γ of $\hat{\Gamma}$ over U . These are functions on $\pi^{-1}(U) \times \mathbb{H}_\vartheta$, where

$$\mathbb{H}_\vartheta \stackrel{def.}{=} \left\{ \zeta : \vartheta - \frac{\pi}{2} < \arg \zeta < \vartheta + \frac{\pi}{2} \right\}. \quad (\text{I.24})$$

By construction, these coordinates, $\chi_\gamma^{\vartheta, u_o}$ experience a jump at specific values of ϑ , and the latter are the phases of the central charges of the BPS states in the vacuum labelled by u_o . Furthermore, these jumps are determined by the gauge charges of the BPS states.

such that the following properties are satisfied:

1. $\chi_{\gamma+\gamma'} = \chi_\gamma \chi_{\gamma'}$.
2. $\forall \xi \in \mathbb{C}^\times$, $\chi_\gamma(\cdot; \xi) \in \mathbb{C}^\times$, and is holomorphic.
3. $\{\chi_\gamma, \chi_{\gamma'}\} = \langle \gamma, \gamma' \rangle \chi_\gamma \chi_{\gamma'}$.
4. $\forall (u, \theta) \in \mathcal{M}$, $\chi_\gamma(u, \theta; \zeta)$ is holomorphic in ζ .
5. $\chi_\gamma(\cdot; \zeta) = \overline{\chi_{-\gamma}\left(\cdot; -\frac{1}{\zeta}\right)}$.
6. $\lim_{\zeta \rightarrow 0} \chi_\gamma(u, \theta; \zeta) \exp\left[-\frac{\pi R Z_\gamma(u)}{\zeta}\right]$ exists.
7. $\chi_\gamma^{\vartheta, u_o}$ features discontinuities at pairs (ϑ, u_o) for which there is some γ_{BPS} with

$$\arg(-Z_{BPS}(u_o)) = \vartheta. \quad (\text{I.25})$$

8. Fixing $\vartheta_o \in \mathbb{R}/2\pi\mathbb{Z}$, $u_o \in \mathcal{B}$, one can define

$$S_{\vartheta, u_o} \stackrel{def.}{=} \prod_{\gamma_{BPS}: \arg(-Z_{\gamma_{BPS}}(u_o)) = \vartheta_o} K_{\vartheta, u_o}^{\Omega(\gamma_{BPS}; u_o)}, \quad (\text{I.26})$$

with

$$K_{\vartheta, u_o} : \chi_\gamma \mapsto \chi_\gamma \left(1 - \sigma(\gamma_{BPS}) \chi_{\gamma_{BPS}}\right)^{\langle \gamma, \gamma_{BPS} \rangle}, \quad (\text{I.27})$$

$$\Rightarrow \left(\lim_{\vartheta \rightarrow \vartheta_o^+} \chi_\gamma^{\vartheta, u_o} \right) = S_{\vartheta_o, u_o} \left(\lim_{\vartheta \rightarrow \vartheta_o^-} \chi_\gamma^{\vartheta, u_o} \right). \quad (\text{I.28})$$

9. When $\zeta \in \mathbb{H}_\vartheta$, then

$$\chi_\gamma \sim \chi_\gamma^{\text{approx.}}. \quad (\text{I.29})$$

If we now consider a point $u \in \mathcal{B}$ and two different phases ϑ_\pm , such that $\vartheta_+ - \vartheta_- < \pi$, then, the two coordinate systems are related as follows

$$\chi_\gamma^{\vartheta_+, u} = S(\vartheta_-, \vartheta_+; u) \chi_\gamma^{\vartheta, u}, \quad (\text{I.30})$$

with

$$S(\vartheta_-, \vartheta_+; u) \stackrel{def.}{=} \prod_{\gamma_{BPS}: \vartheta_- < \arg(-Z_{\gamma_{BPS}}(u)) < \vartheta_+} K_{\vartheta, u_o}^{\Omega(\gamma_{BPS}; u)}, \quad (\text{I.31})$$

where the product decomposition is uniquely defined.

I.4 The Alday-Gaiotto-Tachikawa (AGT) correspondence: a brief overview

This can be generalised to arbitrary class \mathcal{S} -theories of $\mathfrak{su}(2)$ -type: the 4D $\mathcal{Z}_{S^1 \times S^3}$ associated to a Riemann surface of genus g and n -punctures, where we place the holonomy $\left(a_i, \frac{1}{a_i}\right) \in SU(2)_i$ for the flavour symmetry for the i^{th} puncture, corresponds to the 2D q -deformed T=Yang-Mills theory on the same Σ_g , where the punctures are described as holes with holonomy $\left(a_i, \frac{1}{a_i}\right) \in SU(2)$, thereby resulting in the following

$$\mathcal{Z}_{4D,g,n}(a_i) \equiv \frac{\prod_i K(a_i)}{K_o^{2g-2+n}} \mathcal{Z}_{2D,g,n,A=0}(a_i) \quad (\text{I.32})$$

where the prefactor can be reabsorbed under suitable redefinition of the handling operator. This correspondence was first found in [135].

Undeformed 2D YM

For a gauge theory on a cylinder, with holonomy

$$U \stackrel{\text{def.}}{=} \mathcal{P} \exp \int_0^L dx^1 A_1 \quad , \quad U \xrightarrow{G} g U g^{-1} \quad , \quad g \in G \quad (\text{I.33})$$




Figure 94.

the partition function behaves as

$$\mathcal{Z}_{A+A'}(U) \equiv e^{-Ac_2} \mathcal{Z}_{A'}(U) \quad (\text{I.34})$$

with c_2 the operator acting on $\chi_R(U)$ by $c_2(R)$. From this follows that $\mathcal{Z}_{A=0}(U)$ is enough to determine \mathcal{Z} for arbitrary A .

$$\mathcal{Z}_{A=0}(U) \equiv \alpha \delta(U) \quad (\text{I.35})$$

with $\delta(U)$ denoting the δ -function at the ideal of G and

$$\delta(U) \equiv \sum_R d_R \chi_R(U) \quad , \quad d_R \equiv \int_G dU \delta(U) \chi_R(U) \equiv \text{Tr} \mathbf{1} \equiv \dim R \quad (\text{I.36})$$

$$\mathcal{Z}_A(U) \equiv \alpha \sum_R e^{-Ac_2(R)} \dim R \chi_R(U) \quad (\text{I.37})$$

For a 2D surface with 2 boundary holonomies, U, V , instead, we have

$$Z_A(U, V) \equiv \sum_R e^{-Ac_2(R)} \chi_R(U) \chi_R(V) \quad (\text{I.38})$$

Any 2D surface can be cut into pieces, each one being a sphere with 3 holes. This way, we can recover the cylinder by gluing a disk to a hole of a 3-holes sphere. The partition function of the latter reads

$$Z_A(U, V, W) \equiv \frac{1}{\alpha} \sum_R \frac{e^{-Ac_2(R)}}{\dim R} \chi_R(U) \chi_R(V) \chi_R(W) \quad (\text{I.39})$$

Proceeding iteratively, the final result for an arbitrary number of holonomies becomes

$$Z_A(U_i) \equiv \alpha^{2-2g-n} \sum_R e^{-Ac_2(R)} \frac{\prod_i \chi_R(U_i)}{(\dim R)^{2g-2+n}} \quad (\text{I.40})$$

q-deformed 2D YM

The arguments outlined in the previous page are proper to the undeformed 2D YM theory. In order to build the correspondence with the 4D theory, as outlined in [135], the 2D theory needs to be deformed. This takes place as follows: quantising A into faces, f , to each edge e one assigns a dynamical variable $U_e \in G$. The total partition function of the system can therefore be expressed as follows

$$\mathcal{Z}_{tot} \equiv \int \prod_e dU_e \prod_f \mathcal{Z}_f \quad (\text{I.41})$$

with

$$\mathcal{Z}_f \equiv \sum_R e^{-A_f c_2(R)} \text{Tr}_R \prod_e U_e \quad (\text{I.42})$$

The deformation occurs upon requiring $U_e \in G_q$, with G_q being the *quantum group* obtained from the ordinary gauge group G upon making its matrix entries noncommutative. For the case of $SU(2)_q$

$$U_i^j \stackrel{\text{def.}}{=} \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}, \quad \bar{U}_i^j \stackrel{\text{def.}}{=} \begin{pmatrix} \delta & -q^{1/2}\gamma \\ -q^{1/2}\beta & \alpha \end{pmatrix} \quad (\text{I.43})$$

$$\Rightarrow \alpha\beta \equiv q^{1/2}\beta\alpha, \quad \alpha\delta - q^{1/2}\gamma\beta \equiv \delta\alpha - q^{-1/2}\gamma\beta \equiv 1 \quad (\text{I.44})$$

$$\Rightarrow \mathcal{D}^{i\bar{i}} U_i^j \bar{U}_i^{\bar{j}} \equiv D^{j\bar{j}} \neq \delta^{j\bar{j}} \quad (\text{I.45})$$

$$\mathcal{D}^{i\bar{i}} \stackrel{\text{def.}}{=} \begin{pmatrix} q^{-1/2} & 0 \\ 0 & q^{1/2} \end{pmatrix} \Rightarrow \delta^{i\bar{i}} \delta_{i\bar{i}} \equiv 2 \Rightarrow \boxed{D^{i\bar{i}} \delta_{i\bar{i}} \equiv q^{1/2} + q^{-1/2}} \quad (\text{I.46})$$

The boxed expression in (I.46) defines the *quantum dimension* of the 2D representation of $SU_q(2)$ and is basically the character, $\chi_{SU(2)}[q]$. For the $SU(N)$ case, (I.46) generalises as follows

$$\dim_q R \stackrel{def.}{=} \text{Tr}_q \text{diag} \left(q^{(N-1)/2}, q^{(N-3)/2}, \dots, q^{(1-N)/2} \right) \quad (\text{I.47})$$

Upon substituting this to the dimension of the representation in (I.40),

$$Z_{A,g}(U_i) \equiv \alpha^{2-2g-n} \sum_R e^{-Ac_2(R)} \frac{\prod_i \chi_R(U_i)}{(\dim_q R)^{2g-2+n}} \quad (\text{I.48})$$

The key point to notice is the fact that (I.48) can be calculated for a surface with arbitrary number of holes just by knowing \mathcal{Z}_{cyl} , $\mathcal{Z}_{3\text{-holes}}$ and suitable gluing. This statement can be readily understood by saying that correlation functions of a given theory can be determined just by knowing 2- and 3-point functions.

For example, a 4-punctured sphere is obtained by connecting 2 3-punctured spheres.



Figure 95.

4D class \mathcal{S} theories of $SU(2)$ -type

In a similar fashion, they can also be constructed from 3-punctured spheres and cylinders. In the 2D case, we saw how complex numbers are assigned to 2D surfaces. Once more, the main building blocks will be cylinders and punctured surface, with an $SU(2)$ flavour symmetry assigned to each puncture.



Figure 96. The 3-punctured sphere is equivalent to the $\mathcal{N} = 1$ chiral multiplet in the $(2, 2, 2)$ representation of $SU(2)_1 \times SU(2)_2 \times SU(2)_3$, constituting a half hypermultiplet.

The cylinder is associated to an $\mathcal{N}=2$ vector multiplet with gauge group $SU(2)$ whose complexified coupling is

$$\tau \stackrel{def.}{=} \frac{4\pi i}{g^2} + \frac{\theta}{2\pi} \quad , \quad g \in \mathbb{R} \quad , \quad \theta \sim \theta + 2\pi \quad (\text{I.49})$$

Given a collection of 3-punctured spheres, joining 2 of them with a cylinder, leads to a punctured torus. Parametrically, this corresponds to taking a cylinder with parameter τ and making the following identification

$$zw \equiv e^{2\pi i \tau} \equiv q \quad (\text{I.50})$$

with z, w indicating the positions of the 2 joined punctures. From the 4D gauge theory perspective, this corresponds to coupling 2 $SU(2)$ flavour symmetries to a dynamical $SU(2)$ gauge field with coupling constant τ . Once having joined the 2 punctures, their flavour symmetry groups are identified, therefore the hypermultiplet Q_{aiu} can be relabelled into triplets and singlets, $\{A_{Iu}\}, \{H_u\}$, defining an $\mathcal{N}=2$ $SU(2)$ gauge theory with enhanced $\mathcal{N}=4$ SUSY, corresponding to $\mathcal{N}=4$ SYM, namely a theory exhibiting S-duality.

S-duality states that, a theory with gauge group G and coupling constant τ is equivalent to the theory with dual gauge group G^\vee and coupling constant $-\frac{1}{\tau}$. For the case of $SU(2)$, $G \equiv G^\vee$.

For $\mathcal{N}=2$ Lagrangian theories, the partition function can be calculated performing a path integral on $S^1 \times S^3$. In addition to the gauge holonomies parameterising the BPS configurations on which integration is performed, there might also be additional flavour holonomies. As previously mentioned, the holonomy $g \in G$ around S^1 can be expressed as

$$g \equiv (z_1, \dots, z_r) \in U(1)^r \in G \quad (\text{I.51})$$

For an $\mathcal{N} = 2$ hypermultiplet consisting of $\mathcal{N} = 1$ chiral multiplets in a representation R of a symmetry G , the partition function is

$$\mathcal{Z}_R(q, g) \equiv \prod_{n \geq 0} \prod_w \frac{1}{11 - q^{n+1/2} z^w} \quad (\text{I.52})$$

with $w \equiv (w_1, \dots, w-r)$ the weights of R and

$$z^w \equiv \prod_i z_i^{w_i} \quad (\text{I.53})$$

A 3-punctured sphere, the partition function on $S^1 \times S^3$ with holonomies of $SU(2)_i \left(a_i, \frac{1}{a_i} \right) \in SU(2)_i$

$$\mathcal{Z}(a_1, a_2, a_3) \equiv \frac{K(a_1) K(a_2) K(a_3)}{K_o} \sum_{n \geq 0} \frac{\chi_n(a_1) \chi_n(a_2) \chi_n(a_3)}{\chi(q^{1/2})} \quad (\text{I.54})$$

where the RHS is equivalent to the amplitude of a q -deformed $SU(2)$ YM theory on a 3-punctured sphere.

I.5 3D SCFTs on punctures

Higgs branch of $T^\rho[\mathfrak{g}]$

Given a 4D $\mathcal{N}=4$ SYM with gauge group G on $\frac{1}{2}$ -space $\mathbb{R}^{2,1} \times \mathbb{R}^{\geq 0}$, obtained by dimensional reduction of a 6D $\mathcal{N}=(2,0)$ SCFT, admits $\frac{1}{2}$ -BPS boundary conditions for the adjoint fields specified by a homomorphism

$$\rho : \mathfrak{su}(2) \rightarrow \mathfrak{g}. \quad (\text{I.55})$$

The adjoint orbit for each element $e \in \mathfrak{g}$, $O_e^\mathfrak{g}$, is such that

$$O_e^\mathfrak{g} \stackrel{def.}{=} G_e \cdot e. \quad (\text{I.56})$$

The classification of such ρ s, up to conjugation, is equivalent to that of nilpotent elements in \mathfrak{g} . When $\mathfrak{g} \equiv \mathfrak{su}(N)$, a nilpotent orbit is specified by the size of the Jordan blocks

$$n \equiv n_1 + \dots + n_k \quad (\text{I.57})$$

or, equivalently, by the partition, $[n_i]$ of n . Similar considerations hold for $\mathfrak{g} \equiv \mathfrak{so}(n), \mathfrak{sp}(n)$.

The boundary conditions for $\mathcal{N}+4$ SYM are given by Neumann boundary conditions for $\Phi_{1,2,3}$ and Dirichlet BCs for $\phi_{4,5,6}$,

$$\Phi_{1,2,3}(s) \sim \frac{\rho(\tau_{1,2,3})}{s}, \quad \Phi_{4,5,6}(s) \Big|_{s=0} \equiv 0 \quad (\text{I.58})$$

Under S-duality of the 4D bulk, the gauge group $G \rightarrow G^\vee$, and at $s \equiv 0$ lives a 3D SCFT, $T^\rho[\mathfrak{g}]$, with G^\vee -flavour symmetry on the Coulomb branch, coupled to the restriction of the bulk gauge fields to the boundary.

For $\rho \equiv 0$, the Coulomb and Higgs branches of $T[\mathfrak{g}]$ are $\mathcal{N}_{\mathfrak{g}^\vee}, \mathcal{N}_{\mathfrak{g}}$, with the latter defining the nilpotent cone of \mathfrak{g} , namely the subset of \mathfrak{g} consisting of its nilpotent elements. Its dimension reads

$$\dim_{\mathbb{C}} \mathcal{N}_{\mathfrak{g}} \equiv \dim G - \text{rank } G \quad (\text{I.59})$$

Note that this is always an even number due to the fact that $\mathcal{N}_{\mathfrak{g}}$ is a hyperhähler cone.

More generally, turning on some Higgs vevs e , the low-energy limit of the theory at this point would feature singular moduli along the direction orthogonal to O_e , whereas those along the nilpotent orbit will be smooth. Hence, their contribution as free hypermultiplets to the total Higgs branch quaternionic dimension of $T^\rho[\mathfrak{g}]$ reads

$$\dim_{\mathbb{H}} H(T^\rho[\mathfrak{g}]) \equiv \frac{1}{2} (\dim G - \text{rank } G - \dim_{\mathbb{C}} O_e) \quad (\text{I.60})$$

For classical Lie algebras, $\mathfrak{g} = \mathfrak{su}(N), \mathfrak{so}(2N), \mathfrak{sp}(N), \mathfrak{so}(2\mathfrak{N} + 1)$, O_e is labelled by $[n_i]$ and its dimensions are known.

The Coulomb branch and the Spaltenstein map

$$\text{CB}(T^\rho[\mathfrak{g}]) \subset \text{CB}(T[\mathfrak{g}]) \quad (\text{I.61})$$

implying the LHS is the union of closures of nilpotent orbits in \mathfrak{g}^\vee . For \mathfrak{g} a classical Lie algebra, $T^\rho[\mathfrak{g}]$ can be constructed as a braneweb.

The Spaltenstein map leads to

$$d: \mathcal{N}_{\mathfrak{g}}/G \longrightarrow \mathcal{N}_{\mathfrak{g}^\vee}/G \quad (\text{I.62})$$

hence acting on the orbits as follows

$$d(O_p) \equiv O_q, \quad q \equiv p^\dagger \text{ for } \mathfrak{su}(N) \quad (\text{I.63})$$

Importantly,

$$\text{if } O \supseteq O' \Rightarrow d(O) \subseteq d(O') \text{ and } \text{CB}(T^\rho[\mathfrak{g}]) \subset \text{CB}(T^{\rho'}[\mathfrak{g}]) \quad (\text{I.64})$$

if $T^{\rho'}[\mathfrak{g}]$ can be Higgsed to $T^\rho[\mathfrak{g}]$. Rephrasing this in terms of the mixed branches of a given $T[\mathfrak{g}]$: given the maximum Coulomb branch $\mathcal{N}_{\mathfrak{g}^\vee}$ and a maximum Higgs branch $\mathcal{N}_{\mathfrak{g}}$, the mixed branches defining the total vacuum moduli space of $T[\mathfrak{g}]$ is

$$C \times H \equiv \left\{ (x, y) \in \mathcal{N}_{\mathfrak{g}_v} \times \mathcal{N}_{\mathfrak{g}} \mid d(O_x) \supseteq O_y \text{ and } d(O_y) \supseteq O_x \right\} \quad (\text{I.65})$$

such that, upon assigning a vev $e \in \mathcal{N}_{\mathfrak{g}}$ to the Higgs branch, the Coulomb branch is restricted to the closure of $d(O_e)$.

I.6 The Geometric Langlands Program (GLP)

The category of modules of a vertex operator algebra (VOA) is semisimple if and only if the VOA is rational. Any QFT has a trivial line operator. In a topological twist of a supersymmetric theory, it corresponds to an object $\mathbb{1}$ in the differential graded (dg) category. A dg-category is an additive category whose morphism spaces are dg vector spaces. Each space $\text{Hom}(L_i, L_j)$ has a cohomological \mathbb{Z} -grading and a nilpotent differential of degree 1, \mathcal{Q} , behaving as a derivation on a composition of morphisms. Equivalence relations are imposed by taking the \mathcal{Q} -cohomology.

A dg category, \mathcal{D} , can often be represented as a dg enhancement of the bounded derived category $D^b(\mathcal{C})$ of an abelian category \mathcal{C}

$$\mathcal{D} \simeq D^b(\mathcal{C}) \quad (\text{I.66})$$

If $\mathcal{D} \equiv D^b(\mathcal{C})$ is represented as a derived category, then the complex describing local operators at a junction of objects in it are

$$\text{Hom}_{\mathcal{D}}(L, L') = \text{Hom}_{\mathcal{C}}(L, L') \quad (\text{I.67})$$

The degree of the complex corresponds to U(1) R-charge of the local operator. R-charges manifesting in terms of higher-extension groups follows from having chosen to represent the intrinsic category of line operators \mathcal{D} as an enhancement of $D^b(\mathcal{C})$.

The category of lines in 3D TQFTs is expected to be a dg braided tensor category equipped with

1. fusion rules




Figure 97. Fusion in \mathcal{C}_{line} .

2. and braiding

Non-semisimplicity affects the structure of state spaces on surfaces Σ_g , which are closely related to the category of line operators. In a TQFT with semisimple dg category of line operators, \mathcal{D} , the torus Hilbert space is given by the Grothendieck group $K_0(\mathcal{D})$, with a basis labelled by simple objects. For higher genus, the Hilbert spaces are given by the fusion algebra of simple objects, with quantum dimensions provided by the Verlinde formula. When \mathcal{D} is not semisimple, state spaces are more complicated, in particular, for the torus, it is given by the Hochschild homology of \mathcal{D} , $HH_0(\mathcal{D})$.




Figure 98. Braiding in \mathcal{C}_{line} .

Admitting a semisimple category of line operators is a strong constraint to impose on a 3D TQFT; one such example being Chern-Simons theory with compact gauge group $\mathfrak{L}G\mathfrak{L}$ and level k . Indeed, CS is a TQFT of cohomological type, whose category of line operators \mathcal{D} is finite semisimple, with trivial dg structure. The simple objects defining it are Wilson lines labelled by irreps of G , $\{S_i\}$. There are only finitely many of them, due to an equivalence imposed by large gauge transformations. Furthermore, there are no local operators to define gauge-invariant junctions among different Wilson lines, namely, there are no local bulk operators apart from the identity, $\mathbb{1}$. On the other hand, topological twists of supersymmetric theories typically admit non-finite and non-semisimple categories of line operators.

3D $\mathcal{N}=4$ gauge theories admit 2 different topological twists:

1. a reduction of Witten's 4D $\mathcal{N}=2$ Donaldson twist (i.e. the A -twist)
2. an intrinsically 3D twist (the B -twist).

The two are related to the 2D A/B -twists under dimensional reduction on an S^1 . They also arise in 4D $\mathcal{N}=4$ SYM from the Kapustin-Witten A/B -twists. In such setups, semisimplicity is rare. Looking for a 3D TQFT that matches the structure of the CGP TQFT described above based on the non-semisimple category $U_q(\mathfrak{g})\text{-mod}$ at an even root of unity $q \equiv e^{i\pi/k}$. The theory is CS-like, containing a subset of line operators labelled by the same irreps of G at level k , matching the modules of $U_q(\mathfrak{g})$ surviving semisimplicity.

The starting point is the Gaiotto-Witten 3D $\mathcal{N}=4$ SCFTs, $T[G]$, defined as the S-duality interface of 4D $\mathcal{N}=4$ SYM. G is a compact simple Lie group depending on the complexified Lie algebra, \mathfrak{g} . The theory has $\tilde{G} \times \tilde{G}^V$ flavour symmetry, with \tilde{G}, \tilde{G}^V denoting the simply-connected forms of G and its Langlands dual, respectively. The two factors of the flavour group act on the Coulomb and Higgs branches of the moduli space of vacua of $T[G]$, which are Langlands dual nilpotent cones

$$\mathcal{M}_{Coul}[T[G]] \simeq \mathcal{N}^V \in \mathfrak{g}^V, \quad \mathcal{M}_{Higgs}[T[G]] \simeq \mathcal{N} \in \mathfrak{g} \quad (\text{I.68})$$

Requiring that $k \in \mathbb{Z} \simeq H^4(B\tilde{G})$, $k \geq h$, and $G \equiv \tilde{G}$ is simply connected, gauging the simply-connected \tilde{G} symmetry of $T[G]$ by introducing a 3D $\mathcal{N}=2$ gauge multiplet and a SUSY CS term at level K , the resulting theory

$$\mathcal{T}_{\tilde{G},k} \stackrel{def.}{=} T[G]/G_k \quad (\text{I.69})$$

which is still a 3D $\mathcal{N}=4$ SUSY theory, with Wilson line operators in the G -representation and G^V global symmetry. However, it is not topological due to the presence of superconformal matter in $T[G]$. To solve this issue, a topological twist is needed

$$\mathcal{T}_{\tilde{G},k}^A \sim \text{CS}[G_{k-h}] \otimes T[G]^A \quad (\text{I.70})$$

with resulting Hilbert space

$$\mathcal{H}_{\mathcal{T}_{\tilde{G},k}^A}(\Sigma) \sim \mathcal{H}_{G_{k-h}}(\Sigma) \otimes \mathcal{H}_{\mathcal{N}^V}(\Sigma) \quad (\text{I.71})$$

4D construction and 6D relations

The Gaiotto-Witten theories originate from 6D $\mathcal{N}=(2,0)$ SCFTs of ADE-type, \mathfrak{g} , on a product of a 3-manifold and a twisted cigar, $M \times D^2 \times_q S^1$, where $q \equiv e^{2\pi\Psi^V}$. The theory is topologically twisted along M and holomorphically topologically-twisted on $D^2 \times_q S^1$. At ∂D^2 , one places BCs labelled by a completely flat connection, \mathcal{A} , on M . Compactifying on the cigar circle and the noncontractible S^1 in various orders, the resulting theory is 4D YM on $M \times \mathbb{R}_+$ with different BCs.

As analysed by Kapustin and Witten within the context of the Geometric Langlands program, there is a prescription describing how to embed calculations in CS theory into GL-twisted 4D $\mathcal{N}=4$ SYM. A CS calculation on a 3-manifold \mathcal{M}_3 maps to a 4D gauge theory calculation $\mathcal{M}_3 \times \mathbb{R}^+$ with a specific BC deforming the standard Neumann BCs. The conventional notation for such generalised BCs is $B_{(p,q)}$. The analytically-continued CS level k is related to the coupling Ψ of the GL-twisted 4D theory as

$$k + h \equiv \Psi \stackrel{\text{def.}}{=} \frac{\theta}{2\pi} + \frac{4\pi i}{g_{YM}^2} \frac{t - t^{-1}}{t + t^{-1}} \quad (\text{I.72})$$

where $t \equiv \pm i$ corresponds to $\Psi \equiv \tau$.




Figure 99.




Figure 100.

When $G \equiv (P)SU(N)$, the configuration of figure 102 can be further lifted to branewebs in type-IIB string theory.




Figure 101. $\tilde{B}_{1,0}$ and $\tilde{B}_{0,1}$ denote Neumann and Dirichlet boundary conditions, respectively. They are related by the action of the S-duality wall.




Figure 102. Gauging the G -flavour symmetry of $T[G]$ at CS level k , corresponds to replacing one of the Dirichlet boundary conditions with a deformed Neumann BC $B_{1,k}$, obtained by placing an $T^{-k}S$ interface between ordinary Neumann and Dirichlet BCs.




Figure 103.

Physically, given an object L in \mathcal{C}_{ab} , one may perform a path-integral on a solid torus $D^2 \times S^1$ with line operator L inserted at its core to produce a state in the torus state space. This defines a map

$$\text{Ob}(\mathcal{C}_{ab}) \xrightarrow{\kappa} \mathcal{H}(T^2) \quad (\text{I.73})$$

with the following properties:

1. it is linear

$$\kappa(A \oplus B) \equiv \kappa(A) \oplus \kappa(B) \quad (\text{I.74})$$

- 2.

$$\kappa(A \otimes B) \equiv \kappa(A) \cdot \kappa(B) \quad (\text{I.75})$$

3. it factors through the Grothendieck ring $K_o(\mathcal{C}_{ab})$ because the states on the RHS don't see the difference between direct sums of linear operators $A \oplus B$ and nontrivial extensions (i.e. bound states)
4. images of κ belong to the cohomological degree-0 subspace of the state space $\mathcal{H}(T^2)$.

All this, together with the isomorphism $\mathcal{H}(T^2) \simeq HH_o(\mathcal{C})$, implies that κ induces the map

$$K_o(\mathcal{C}) \rightarrow HH_o(\mathcal{C}) \subset \mathcal{H}(T^2) \quad (\text{I.76})$$

Identity stalk and the flag manifold

$$D^b(\mathcal{C}_1^{ab}) \equiv \mathcal{C}_1 \simeq \text{Vect}^{\oplus 2} \oplus [\text{Coh}(T^*[2]\mathbb{P}^1)]^{\oplus k-1} \quad (\text{I.77})$$

where each of the two Vects is an abelian semisimple category with a single simple object, whereas, $[\text{Coh}(T^*[2]\mathbb{P}^1)]^{\oplus k-1}$, is a dg enrichment of the derived category. The direct sum signals that there are no morphisms between objects belonging to the two summands. The abelian and derived categories decompose as a direct sum of $K + 1$ blocks

$$\mathcal{C}_1^{ab} \simeq B_k^{ab+} \oplus B_k^{ab-} \oplus \bigoplus_{j=1}^{k+1} B_j^{ab} \quad (\text{I.78})$$

where $B_k^{ab\pm}$ denote the semisimple subcategories generated by the simple projectives S_k^\pm , whereas each B_j^{ab} consists of a subcategory generated by the pair (S_j^+, S_{k-j}^-) .

J Categories, knots, and entanglement

J.1 Higher-categorical structure of (linked minimal models) knots

This appendix contains additional background material underlying section C. In particular, the main focus is that of identifying fusion spaces associated with knots, [583]. A concrete calculation of knot multiplicity shows that the knot complement of a trefoil knot can store quantum information. Also spiral maps enabling to understand consistency relations for torus knots and spiral fusions of fluxes.

Immersion of A in M is defined by triples (A_+, A, \mathfrak{p}_A) . The projection $\mathfrak{p}_A : A \rightarrow M$ may or may not be an embedding. In particular, the Hilbert space induced by M on A might be smaller w.r.t. the one with which A is already equipped. Pictorially, this would appear as an overlap of A on M . At the level of the Hilbert space, the overlap corresponds to a double copy in the tensor product structure. The correct definition can be achieved by suitably lifting the refined state on local disks contained in $\tilde{\Gamma}(A_+)$.

The information convex set, $\Sigma(\Omega)$, of a 2-hole disk can be completely characterised. Any element belonging to it can be expressed as

$$\rho = \bigoplus_{a,b,c \in \mathcal{C}} p_{ab}^c \rho^{abc} \quad (\text{J.1})$$

with \mathcal{C} denoting the set of topological charges, $\{p_{ab}^c\}$ a probability distribution, and $\{\rho^{abc}\}$ a set of mutually orthogonal quantum states. For each triple (a, b, c) , $\{\rho^{abc}\}$ defines the state space of a finite-dimensional Hilbert space, whose dimension is given by the fusion multiplicities of the underlying anyon theory. (a, b) label the charges associated with the holes, and c the total charge of the disk.

The *information convex set* for an immersed region is defined as $\Sigma(\Omega) = \{\rho_\Omega\}$ such that

1. $\rho_\Omega \equiv \text{Tr}_{\Omega_+ \setminus \Omega} \rho_{\Omega_+}$.
2. ρ_{Ω_+} is consistent with $\sigma_\omega^p \forall \omega \in \tilde{\Gamma}(\Omega_+)$.

Elements of the information convex sets are closed under the merging operation as long as a certain mild condition is satisfied. The *merging theorem* states that immersed spaces, T_1, T_2 , with isomorphic ICSs, share the same number of elements. From this follows that, under the composition of density matrices belonging to each one of them, $\rho_{ABC} \in \Sigma(ABC)$, $\lambda_{BCD} \in \Sigma(BCD)$, the resulting merged state reads

$$\tau \stackrel{\text{def.}}{=} \rho \bowtie \lambda \quad , \quad \tau \in \Sigma(ABCD) \quad (\text{J.2})$$

For the case in which Ω is *sectorisable*, its ICS can be expressed in the following way

$$\Sigma(\Omega) \equiv \left\{ \sum_i p_i \rho_\Omega^i \mid \sum_i p_i = 1, p_i \geq 0 \right\} \quad (\text{J.3})$$

with $\{\rho_\Omega^i\}$ denoting the set of mutually orthogonal density matrices. The set of labels i forms the *set of superselection sectors*, \mathcal{C}_Ω , with $i = 1$ defining the *vacuum sector*. For any given sectorisable region Ω embedded in S^3 or B^3 , $\rho_\Omega^1 \equiv \sigma_\Omega$ is unique.

The *quantum dimension* can therefore be defined as follows

$$d_i \stackrel{\text{def.}}{=} \exp \left(\frac{S(\rho_\Omega^i) - S(\sigma_\Omega^*)}{2} \right) \quad (\text{J.4})$$

where σ_Ω^* denotes the vacuum state.

A set of *superselection sectors* can be assigned to any sectorisable region. Conventionally, we will be denoting them as $\{\mathcal{C}_I\}$, with the subscript I specifying the sector. Among the simplest choices one could make are: sphere shells, $\mathcal{C}_{\text{point}}$, solid tori, $\mathcal{C}_{\text{flux}}$, and torus shells, $\mathcal{C}_{\text{Hopf}}$. For the purpose of this work, the latter happens to be of particular importance. However, as we shall be arguing throughout, all 3 can be thought of as subcategories defining an n -category associated to an $(n + 1)$ -dimensional theory.

Its corresponding superselection sector type is the Hopf fibration, $\mathcal{C}_{\text{Hopf}} \equiv \{1, \eta, \zeta, \dots\}$. Superselection sectors are associated with different classes of loop excitations. Each one of them can be created by applying a suitable type of membrane operator. The description in terms of the information convex set is somehow dual to that of the excitations, since it focuses on their complement when embedded in an S^3 .

For a torus shell, $\Sigma(\mathbb{T})$ is a simplex with extreme points $\{\rho_\Sigma^\eta\}_{\eta \in \mathcal{C}_{\text{Hopf}}}$. The set of Hopf excitations admit a natural decomposition




Figure 104. A sphere shell, (on the LHS). On the RHS, the topological charges of the 2 connected components are mapped to a particle/antiparticle pair joined by a string.




Figure 105. A solid torus, \mathbb{T} , (on the LHS). On the RHS, a visualisation of a flux operator operator, η .




Figure 106. A torus shell, \mathbb{T} , (on the LHS). On the RHS, a visualisation of the Hopf link, resulting from the mutual intersection of η and η^v .

$$\mathcal{C}_{Hopf} \equiv \bigcup_{\mu \in \mathcal{C}_{flux}} \mathcal{C}_{Hopf}^{[\mu]} \quad (\text{J.5})$$

$\Sigma(\mathbb{T})$ comes equipped with nontrivial automorphisms, such as the one responsible for permuting the constituting loops in the Hopf link of its complement

$$v : \mathcal{C}_{Hopf} \longrightarrow \mathcal{C}_{Hopf} \quad , \quad v(\eta) \equiv \eta^v \quad (\text{J.6})$$

The set of shrinkable loops are not linked to any other by definition, and therefore provide a natural embedding for the fluxes

$$\mathcal{C}_{flux} \xrightarrow{\varphi} \mathcal{C}_{loop} \equiv \mathcal{C}_{Hopf}^{[1]} \subset \mathcal{C}_{Hopf} \quad (\text{J.7})$$

The embedding (J.5) implies the matching of the quantum dimensions

$$\sum_{a \in \mathcal{C}_{loop}} d_a^2 \equiv \sum_{\mu \in \mathcal{C}_{flux}} d_\mu^2 \quad (\text{J.8})$$

as well as the presence of a well defined map $\mathcal{C}_{loop} \rightarrow \mathcal{C}_{flux}$. Indeed, given a torus shell \mathbb{T} , and $T \subset \mathbb{T}$ a solid torus, an extreme point $\rho_{\mathbb{T}}^\eta \in \Sigma(\mathbb{T})$,

$$\text{Tr}_{\mathbb{T}/T} \rho_{\mathbb{T}}^\eta \equiv \rho_T^\mu \in \Sigma(T) \quad (\text{J.9})$$

Fusion rules and knot complements




Figure 107. On the LHS, a solid torus with 2 tori removed. On the RHS, the fusion of 2 loops in $\mathcal{C}_{Hopf}^{[mu]}$, from which the multiplicities $\{N_{ij}^h\}$ can be extracted.

The Hilbert space theorem states that,

$$\Sigma_I(\Omega) \simeq S(\mathbb{V}_I) \quad (\text{J.10})$$

where I labels the superselection sector and $S(\mathbb{V}_I)$ defines the state space of a finite-dimensional Hilbert space \mathbb{V}_I . From (J.10), it follows that the information convex set is completely characterised by the *fusion multiplicity*

$$N_I \equiv \dim \mathbb{V}_I \quad (\text{J.11})$$

For $N_I = 0$, $\Sigma_I(\Omega)$ is empty, whereas for $N_I = 1$ it contains a single element, which is an isolated fixed point. For $N_I = 2$ it is isomorphic to the Bloch ball and contains an infinite number of continuously parameterised extreme points. When $N_I > 1$, Ω can store quantum information.

the information convex sets of 3D regions with boundaries can be associated with a set of fusion spaces. The fusion of 2 loops, depicted on the RHS of figure 107, enables to derive the *fusion multiplicities*, $\{N_{ij}^h\}$,

$$h \times i \equiv \sum_j N_{ij}^h j \quad , \quad h, i, j \in \mathcal{C}_{Hopf}^{[\mu]} \quad (\text{J.12})$$

The *complement* of an unknot in B^3 is a shrinkable loop at a poin, with *shrinking rule*

$$\ell \equiv \sum_a N_\ell^a a \quad , \quad \ell \in \mathcal{C}_{loop} \quad , \quad a \in \mathcal{C}_{point} \quad (\text{J.13})$$




Figure 108. $B^3 \setminus T$ on the LHS, with its corresponding shrinking rule on the right.

The next essential ingredient that needs to be introduced is the *knot complement*, which plays an important role in their classification. For the purpose of our treatment, their key role resides in the fact that their information convex sets are characterised by nontrivial fusion data. For a given knot, K , we will denote its complement on the 3-sphere as

$$\Omega_K \stackrel{def.}{=} S^3/K \quad (\text{J.14})$$

with a thickened boundary $\partial\Omega_K$, i.e. a knotted embedded torus shell. It therefore follows that

$$\mathcal{C}_{\partial\Omega_K} \simeq \mathcal{C}_{Hopf} \quad (\text{J.15})$$

namely the 2 sets are identical up to permutations. The information convex set of the knot complement, $\Sigma(\Omega_K)$, is a convex hull of the subset $\Sigma_\zeta(\Omega_K)$, with $\zeta \in \mathcal{C}_{Hopf}$. In general, the set of superselection sectors that can exist on a knot K of an S^3 alone is a subset of \mathcal{C}_{Hopf} , implying that the knot excitations are a small number w.r.t. the Hopf link. However, they are more coherent, and therefore are able to store quantum information. The proof of this last statement was carried out in [583] and requires some additional elements we will now present.

J.2 Spiral maps, fusion multiplicities and holomorphic blocks

The *knot multiplicity*, $N_\zeta(K)$ is the dimension of the Hilbert space of $\Sigma_\zeta(\Omega_K)$

$$N_\zeta(K) \stackrel{def.}{=} \dim \mathbb{V}_\zeta(\Omega_K) \quad (\text{J.16})$$

For the unknot, (J.16) reduces to

$$N_\eta(\text{unknot}) \stackrel{def.}{=} \sum_{\mu \in \mathcal{C}_{flux}} \delta_{\eta, \varphi(\mu)} \quad , \quad \forall \eta \in \mathcal{C}_{Hopf} \quad (\text{J.17})$$

whereas for a general (p, q) -knot it reads as follows

$$N_\zeta^n(p, q) \stackrel{def.}{=} \dim \mathbb{V}_\zeta^n(T \setminus K_{p,q}) \quad , \quad \eta, \zeta \in \mathcal{C}_{Hopf} \quad (\text{J.18})$$




Figure 109. Example of $S^3 \setminus K(2, 3)$. With η, ζ denoting the unknotted and knotted boundaries, respectively.

with η, ζ labelling the unknotted (outer) and knot boundaries, respectively.

$\Sigma(\Omega)$ is completely determined by the *set of extreme points*, $\text{ext}(\Sigma(\Omega))$. In order to define the latter, some preliminary definitions are of order.




Figure 110. Ω is not sectorisable, but $\partial\Omega$ is, since it has 3 connected components, labelled $i = a, b, c$. Each one of them corresponds to a different superselection sector, collectively denoted $\mathcal{C}_{\partial\Omega}$.

For regions that are not sectorisable, such as the one depicted in figure 110, we can distinguish between Ω and its boundary defined as follows

$$\partial\Omega \stackrel{\text{def.}}{=} \mathcal{M}_{D-1} \times I \quad (\text{J.19})$$

where \mathcal{M}_{D-1} is a $(D-1)$ -dimensional closed manifold, and I an interval. $\Sigma(\partial\Omega)$ is a *simplex* since $\partial\Omega$ is sectorisable, namely it is possible to assign to it a finite set of superselection sectors labelled by $\mathcal{C}_{\partial\Omega}$. If $\partial\Omega$ contains multiple connected components, each component is sectorisable.

Under the action of partial tracing, every extreme point of $\Sigma(\Omega)$ reduces to one of $\Sigma(\partial\Omega)$.

Equipped with this machinery, it is now possible to define the *extreme point criterion*, stating that, for a given immersed Ω , $\rho_\Omega \in \Sigma(\Omega)$ is an extreme point if and only if

$$(S_\Omega + S_{\Omega \setminus \partial\Omega} - S_{\partial\Omega})_\rho \equiv 0 \quad (\text{J.20})$$

Furthermore, $\forall i \in \mathcal{C}_{\partial\Omega}$, corresponding to a nonempty $\Sigma_i(\Omega)$, the *quantum dimension* is defined as

$$\ln d_i \stackrel{def.}{=} S\left(\rho_\Omega^{i,e}\right) - S\left(\rho_\Omega^{1,e}\right) \quad (\text{J.21})$$

with $\rho_\Omega^{i,e}$ an extreme point of $\Sigma_i(\Omega)$, and $\rho_\Omega^{1,e}$ the vacuum reference state.

Once having identified the quantum dimensions associated to the different superselection sectors, we can calculate their *fusion multiplicities*, N_{ab}^c ,

$$d_a d_b = \sum_c N_{ab}^c d_c \quad (\text{J.22})$$

with (a, b, c) denoting the different topological charges involved in the fusion process.

J.3 The total quantum dimension

The *total* quantum dimensions can therefore be extracted as follows:

1. for points and fluxes

$$\mathcal{D} \stackrel{def.}{=} \sqrt{\sum_{a \in \mathcal{C}_{point}} d_a^2} \equiv \sqrt{\sum_{\mu \in \mathcal{C}_{flux}} d_\mu^2} \quad (\text{J.23})$$

2. for a knotted torus, instead, the definition can be relaxed, turning the equality into an upper bound on the topological entanglement entropy

$$\sum_{\zeta \in \mathcal{C}_{Hopf}} N_\zeta(K) d_\zeta \equiv \sum_{\substack{\mu, \nu \in \mathcal{C}_{flux} \\ t_p(\mu) \equiv t_q(\nu)}} \frac{d_\mu^2 d_\nu^2}{d_{t_p(\mu)}^2} \leq \mathcal{D}^4 \quad (\text{J.24})$$

For the case of K with arbitrary (p, q)

$$\sum_{\zeta \in \mathcal{C}_{Hopf}} N_\zeta^\eta(p, q) d_\zeta \equiv \sum_{\substack{\mu, \nu \in \mathcal{C}_{flux} \\ t_p(\mu) \equiv t_{(p,q)}(\nu)}} \frac{d_{\varphi^\vee} d_\eta}{d_{t_p(\mu)}^2} \leq \mathcal{D}^4 \quad (\text{J.25})$$

where

$$N_\zeta^\eta(p, q) \equiv \sum_{\mu \in \mathcal{C}_{flux}} N_{\zeta \varphi^\vee(\mu)}^\eta \quad (\text{J.26})$$

Importantly, the embedding operation

$$\varphi^\vee : \mathcal{C}_{flux} \hookrightarrow \mathcal{C}_{Hopf} \quad (\text{J.27})$$

really consists of 2 consecutive parts, namely

$$\varphi : \mathcal{C}_{flux} \hookrightarrow \mathcal{C}_{loop} \subset \mathcal{C}_{Hopf} \quad (\text{J.28})$$

and

$$\eta \rightarrow \eta^\vee \quad (\text{J.29})$$

from which follows that (J.26) is equivalent to a gauging procedure.

In a 4D TQFT, the entanglement entropy is given by the area-law plus a topological subleading correction

$$S_{EE} = \alpha \ell + \ln \mathcal{D} \quad (\text{J.30})$$

where α is a constant, ℓ is the area of the surface on which S_{EE} is being calculated, and \mathcal{D} is the quantum dimension. Following the Casini-Huerta-Myers prescription, the latter corresponds to the so-called boundary free energy of the 3D boundary to the 4D TQFT. It is a universal quantity, and its importance is crucial for the study of holographic embeddings of such theories.

For $\Sigma(\Omega_K)$, with corresponding maximum entropy state $\rho_{\Omega_K}^*$, the entropy w.r.t. the reference state can be extracted from the fusion multiplicities (J.25) and therefore reads

$$S(\rho_{\Omega_K}^*) - S(\sigma_{\Omega_K}) = \ln \left(\sum_{\zeta \in \mathcal{C}_{\text{Hopf}}} N_{\zeta}^{\eta}(p, q) d_{\zeta} \right) \quad (\text{J.31})$$

[analogy with Janus configurations]

K Sigma-models

K.1 The sigma-model

By definition, [86], a sigma-model is a field theory describing the field as a point particle confined to move on a fixed manifold. The latter could either be a Riemannian manifold, a Lie group, or a symmetric space.

When coupling it to a gauge field, we get the Landau-Ginzburg (LG)-model, [486].

The Lagrangian density for the sigma-model can be expressed in many different ways. One of the simplest expressions is the following

$$\mathcal{L} = \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n g_{ij}(\phi) \partial^{\mu} \phi_i \partial_{\mu} \phi_j, \quad (\text{K.1})$$

where $g_{ij}(\phi)$ denotes the metric tensor on the field space $\phi \in \Phi$, where Φ can be any Riemannian manifold.

In a more fully-geometric notation, it can be written as a fiber bundle with fibers Φ over a differentiable manifold M . Given a section

$$\phi: M \rightarrow \Phi, \quad (\text{K.2})$$

fix a point $x \in M$. Then, the pushforward at x is a map of tangent bundles

$$d_x \phi: T_x M \rightarrow T_{\phi(x)} \Phi, \quad (\text{K.3})$$

taking

$$\partial_{\mu} \mapsto \frac{\partial \phi^i}{\partial x^{\mu}} \partial_i. \quad (\text{K.4})$$

The sigma model action then reduces to

$$S = \frac{1}{2} \int_M d\phi \wedge \star d\phi. \quad (\text{K.5})$$

Among the most important interpretation of the classical sigma-model is that of non-interacting QM. Taking $\Phi \equiv \mathbf{C}$, (K.5) becomes

$$S = \frac{1}{2} \langle\langle \phi, \Delta \phi \rangle\rangle, \quad (\text{K.6})$$

with

$$\phi : M \rightarrow \mathbb{C} \quad (\text{K.7})$$

interpreted as a wavefunction, whose Laplacian corresponds to its associated kinetic energy. Hence, the classical sigma-model on \mathbf{C} can be interpreted as the QM of a free particle.

K.2 Riemann surfaces

The geodesic structure of a Riemannian manifold is described by the Hamilton-Jacobi equations. As previously mentioned, the M and Φ defining the sigma-model can both be taken to be Riemannian.

The cotangent bundle $T^*\Phi$ can always be locally trivialised, namely

$$T^*\Phi \Big|_U \simeq U \times \mathbf{R}^n. \quad (\text{K.8})$$

Given g_{ij} on Φ , the Hamiltonian function is defined as follows

$$H(q, p) \stackrel{\text{def.}}{=} \frac{1}{2} g^{ij}(q) p_i p_j, \quad (\text{K.9})$$

from which the Hamilton-Jacobi equations define the geodesic (or Hamiltonian) flow on Φ

$$\dot{q}^i = \frac{\partial H}{\partial p_i}, \quad \dot{p}_i = -\frac{\partial H}{\partial q^i}. \quad (\text{K.10})$$

The Hamiltonian description therefore enables us to interpret the sigma-model as the gluing of two energy functionals, namely the momenta in $T^*\Phi$ and T^*M .

K.3 Moduli spaces

In algebraic geometry, a moduli space is a geometric space (a scheme or an algebraic stack) whose points represent algebro-geometric objects of some kind, or their isomorphic classes. The main motivation for introducing moduli spaces is that of finding solutions of geometric problems.

A scheme is a mathematical structure that enlarges the notion of an algebraic variety, accounting for multiplicities, and allowing varieties defined on any commutative ring. Scheme theory was introduced by A. Grothendieck in 1960 for addressing deep problems in algebraic geometry.

A scheme is a topological space together with commutative rings for all of its open sets, which arises from gluing together spectra of commutative rings along their open subsets, i.e. ringed space which is locally a spectrum of a commutative ring.

Algebraic surfaces can be studied making use of morphisms of schemes. In many cases, the family of all varieties of a given type can itself be viewed as a variety of schemes, known as moduli space.

K.4 The nonlinear sigma-model

Considering Minkowski spacetime, and denoting with \mathcal{F} the space of fields, L the Lagrangian, and γ the variational 1-form, \mathcal{M} the space of classical solutions, and ω the local symplectic form. In its classical formulation, the Lagrangian encodes a classical Hamiltonian system once a specified time direction has been selected, thereby breaking Poincaré' invariance.

Upon taking a real-valued scalar field

$$\phi : M^n \rightarrow \mathbb{R}. \quad (\text{K.11})$$

The kinetic term in the Lagrangian would therefore read

$$\begin{aligned} L_{kin} &= \frac{1}{2} |d\phi|^2 \Big| d^n x \Big| \\ &= \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi \Big| d^n x \Big| \\ &= \frac{1}{2} \left[(\partial_0 \phi)^2 - \sum_{i=1}^{n-1} (\partial_i \phi)^2 \right] \Big| d^n x \Big|. \end{aligned} \quad (\text{K.12})$$

Adding a potential term of degree ≤ 2 of the kind

$$V : \mathbb{R} \rightarrow \mathbb{R} \quad (\text{K.13})$$

to (K.12), and assuming V is bounded below by choosing suitable coefficients of the polynomial, then one can eliminate the linear and constant terms by assuming the minimum of V is at the origin with $V = 0$ there. The total Lagrangian therefore becomes

$$L = \left(\frac{1}{2} |d\phi|^2 - \frac{m^2}{2} \phi^2 \right) \Big| d^n x \Big|. \quad (\text{K.14})$$

In non-free theories, the potential is no longer quadratic. The simplest example of an interacting theory is the case of the quartic potential. In either case, in a pure mathematical formulation, the non-linear σ -model is characterised by the following data:

1. A Riemannian manifold, X .
2. A potential energy function

$$V : X \rightarrow \mathbb{R}. \quad (\text{K.15})$$

In this formulation, the space of fields is the mapping space

$$\mathcal{F} = \left(\text{Map}(M^n, X) \right). \quad (\text{K.16})$$

The total Lagrangian therefore reads

$$L = \left[\frac{1}{2} |d\phi|^2 - \phi^* V \right] \Big| d^n x \Big|, \quad (\text{K.17})$$

from which the energy density reads

$$\mathcal{E}(\phi) = \left[\frac{1}{2} |\partial_0 \phi|^2 + \sum_{i=1}^{n-1} \frac{1}{2} |\partial_i \phi|^2 + V \right] \Big| d^{n-1} x \Big|, \quad (\text{K.18})$$

The space of solutions, namely the moduli space of vacua, \mathcal{M}_{vac} , is

$$\mathcal{M}_{vac} \subset \mathcal{F}\mathcal{E}_N, \quad (\text{K.19})$$

with $\mathcal{F}\mathcal{E}_N$ denoting the space of static fields of finite energy.

For the case of a real scalar field with potential V , one usually assumes that the minimum value is at $V = 0$, and therefore that the solutions lie at

$$\mathcal{M} \stackrel{\text{def.}}{=} V^{-1}(0). \quad (\text{K.20})$$

For example, if $X \equiv \mathbb{R}$, and $V = 0$, the resulting theory is that of a massless real scalar field. For this case, one finds that $\mathcal{M}_{vac} \simeq \mathbb{R}$. On the other hand, for the case of a scalar field with potential

$$V(\phi) = \frac{m^2}{2}\phi^2 \quad (\text{K.21})$$

\mathcal{M}_{vac} is a single point $\phi = 0$. For a quartic potential, instead, \mathcal{M}_{vac} is constituted by two points, i.e. the values of ϕ where V is minimised.

K.5 Principle bundles and connections

Formulating gauge theories in Lagrangian formulation that is suitable for a QFT, some preliminary differential geometry notations are needed.

The initial ingredients we start from are a manifold, M , and a Lie group, G . Then, a *principal G -bundle*, $P \rightarrow M$ is defined as a manifold P on which G acts freely on the right with quotient

$$P/G \simeq M, \quad (\text{K.22})$$

such that there exist local sections. If P', P are principal G -bundles over M , then an isomorphism

$$\varphi' : P' \longrightarrow P, \quad (\text{K.23})$$

is a smooth diffeomorphism which commutes with G , and induces the identity map on M . If $P \equiv P'$, such automorphism defines a gauge transformation of P .

For every M , there is a category of principal G -bundles and isomorphisms.

A *connection* on a principal G -bundle $\pi : P \rightarrow M$ is a G -invariant distribution in TP which is transverse to the vertical distribution $\ker d\pi$. A connection can therefore be expressed as a 1-form $A \in \Omega^1(P; \mathfrak{g})$ whose value at p is the projection

$$T_p P \longrightarrow V_p \simeq \mathfrak{g}, \quad (\text{K.24})$$

with kernel H_p .

Connections form a category, arranging in a set of equivalence classes under isomorphisms.

K.6 Mathematical Gauge theory

The prototypical example of a gauge theory is Maxwell's theory. In its quantised formulation, one needs to introduce Dirac's charge quantisation, replacing the group \mathbb{R} by the compact group $\mathbb{Z}/2\mathbb{Z}$. This statement can be generalised by replacing $\mathbb{Z}/2\mathbb{Z}$ with any compact Lie group G .

Hence, we are drawn to the conclusion that the Lagrangian formulation of the field theory in question is simply based on the following data:

1. A compact Lie group with Lie algebra, \mathfrak{g} .
2. A bi-invariant inner product on \mathfrak{g} , $\langle \cdot, \cdot \rangle$.

From this, the total Lagrangian for the gauge field reads

$$L = -\frac{1}{2} \langle F_A \wedge \star F_A \rangle \quad (\text{K.25})$$

from which the energy density reads

$$\mathcal{E}(A) = \left[\sum_{\mu < \nu} \frac{1}{2} |F_{\mu\nu}|^2 \right] |d^{n-1}x|, \quad (\text{K.26})$$

whose moduli space is simply

$$\mathcal{M}_{vac} = \text{pt}, \quad (\text{K.27})$$

where the calculation is understood to be performed on the space of equivalence classes of connections.

K.7 The gauged sigma-model

This is a very general bosonic field theory, combining pure gauge theory with the σ -model, both of which are greatly used in QFT. The data specifying the model in question are:

1. A Lie group, G , with Lie algebra, \mathfrak{g} .
2. A bi-invariant scalar product, \langle, \rangle , on \mathfrak{g} .
3. A Riemannian manifold, X , on which G acts by isometries.
4. A potential function

$$V : X \longrightarrow \mathbb{R}, \quad (\text{K.28})$$

invariant under G .

The space of fields, \mathcal{F} , consists of pairs

$$\mathcal{F} \stackrel{\text{def.}}{=} \{(A, \phi)\}, \quad (\text{K.29})$$

with A denoting a collection of principal G -bundles $P \rightarrow M$, and ϕ a section of the associated bundle over $P \times_G X \rightarrow M$. In most cases, it is convenient to view ϕ as an equivariant map

$$\phi : P \longrightarrow X. \quad (\text{K.30})$$

The space of fields can be thought of as a category, with an isomorphism

$$\varphi : P' \longrightarrow P \quad (\text{K.31})$$

of principal bundles, inducing an isomorphism of fields

$$(A', \phi') \longrightarrow (A, \phi). \quad (\text{K.32})$$

The global symmetry group of the theory is the subgroup of isometries of X commuting with the G -action, and preserving the potential, V .

The Lagrangian density, therefore reads

$$\mathcal{L} = -\frac{1}{2} |F_A|^2 + \frac{1}{2} |\mathcal{D}_A \phi|^2 - \phi^* V, \quad (\text{K.33})$$

where \mathcal{D}_A denotes the covariant derivative, coupling ϕ with A . The energy density of the pair (A, ϕ) , can readily be extracted from (K.33), and reads

$$\mathcal{E}(A, \phi) = \left[\sum_{\mu < \nu} \frac{1}{2} |F_{\mu\nu}|^2 + \sum_{\mu} \frac{1}{2} \left| (\partial_A)_\mu \phi \right|^2 + \phi^* V \right] |d^{D-1}x|, \quad (\text{K.34})$$

where D denotes the total number of spacetime dimensions.

Assuming V has a minimum at 0, one can look for solitons with 0-energy. By direct inspection of (K.34), it is possible to deduce that:

1. The first term implies that A is flat, hence, up to equivalence, it is the trivial connection with zero-curvature.
2. The second term tells us that ϕ has to be a constant for it to be a zero-energy solution.
3. The third term constraints the constant value of ϕ to lie in the $V^{-1}(0)$ set.

A trivial connection A has a group of automorphisms that are isomorphic to G , i.e. the group of global gauge transformations. For a vacuum solution, ϕ must be a constant function with values in $V^{-1}(0)$, and, therefore, $V^{-1}(0)$ inherits the same G -action from ϕ . Hence, the moduli space of vacua is a subquotient space of X

$$\boxed{\mathcal{M}_{vac} \stackrel{\text{def.}}{=} V^{-1}(0) / G}. \quad (\text{K.35})$$

For the case of certain supersymmetric field theories, X is Kähler or hyperkähler, and $V^{-1}(0)$ is the norm square of an appropriate moment map, in which case \mathcal{M}_{vac} reduces to a Kähler or hyperkähler quotient (with the latter actually being the setting where this had first been studied).

L (A)dS₃/ $T\bar{T}$ -deformed CFTs

As previously argued, the entropy of pure AdS is divergent, compatibly with the fact that bulk reconstruction enables to probe the IR bulk at arbitrary depth by evaluating S_{EE} on ever-increasing subregions of the conformal boundary.

Within the AdS/CFT correspondence, finiteness of bulk-reconstruction can be achieved by shifting the conformal boundary at a finite cutoff by deforming the CFT Lagrangian density by means of an irrelevant operator. In 2D, this is known as the $T\bar{T}$ -deformation, initially introduced by Zamolodchikov, [239]. The RG-flow equations driven by this operator interpolate between the UV and the IR.

For the 2D dual to an (A)dS₃, [240, 241], the action associated to the deformed theory reads

$$S_{T\bar{T}}^{AdS} = S_{CFT} + \lambda \int d^2x \sqrt{-\gamma} T\bar{T} \quad (\text{L.1})$$

$$S_{T\bar{T}}^{dS} = S_{CFT} + \lambda \int d^2x \sqrt{-\gamma} \left[T\bar{T} + \frac{1}{2\lambda^2} \right] \quad (\text{L.2})$$

respectively, with

$$T\bar{T} \stackrel{\text{def.}}{=} \frac{1}{8} (T^{ab}T_{ab} - (T_a^a)^2) \quad (\text{L.3})$$

The deformation of the CFT₂ Lagrangian induces a change in the partition functions, which, for AdS₃ reads

$$\delta \ln \mathcal{Z} = \ln \mathcal{Z}_{T\bar{T}} - \ln \mathcal{Z}_{CFT} = -\frac{1}{2} \int d^2x \sqrt{\gamma} \langle T^{ab} \rangle \delta \gamma_{ab} \quad (\text{L.4})$$

and *trace flow equations*, [239],

$$\langle T_a^a \rangle = -\frac{c}{24} \mathcal{R} - \frac{\lambda}{4} (\langle T^{ab} \rangle \langle T_{ab} \rangle - \langle T_a^a \rangle^2) \quad (\text{L.5})$$

where the 1st term defines the trace anomaly associated with a CFT₂.

The first application of these deformations for describing the gluing of bulk spacetimes from a holographic perspective, was carried out by H. Verlinde et al. in [240] for AdS₃ spacetimes, and further developed by Silverstein et al. [241] for local realisations of dS₃s in an asymptotically AdS₃ background, with the latter relying upon the dS_{D+1}/dS_D-correspondence (dS/dS for brevity). The fact that the latter defines a duality in between 2 dS theories follows from the fact that the deformed Lagrangian driving the RG-flow contains an additional term w.r.t. the AdS₃/CFT₂ case which is playing the role of a positive 2D cosmological constant. We will come back to this in greater detail when deriving the IR S_{EE} s in both setups. Before doing so, we will briefly outline the key features characterising the dS/dS-correspondence, motivating that the gluing of different spacetimes in the IR can be naturally accounted for in both correspondences alike.

According to the dS/dS correspondence, [244], a D+1-dimensional dS spacetime with curvature radius $1/H_{D+1}$ is equivalent to 2 interacting CFTs of D-dimensions with a cutoff H_{D+1} and coupled to gravity (cf. figure 111).




Figure 111. The conformal boundary is depicted in blue. The RT-surface is still a codimension-2 surface, in complete analogy with the AdS/CFT setup. For $D = 2$, it reduces to 2 antipodal points on an S^1 .

The analogy in between the 2 AdS/CFT and the dS/dS correspondences already shows at the level of the metric, which read¹¹⁹

$$ds_{D+1}^2 = dw^2 + \sin(h)^2 (w H) \left[-d\tau^2 + \frac{1}{H^2} \cosh^2(\tau H) d\Omega_{D-1}^2 \right] \quad (\text{L.6})$$

with the key difference being the change in the scale factor $\sin \rightarrow \sinh$, implying that the 2 match in the IR-region of the bulk, i.e. for small values of the arguments of the trigonometric/hyperbolic functions. From the CFT point of view this corresponds to the fact that the CFT on the dS/dS side has to be deformed by an irrelevant operator.

¹¹⁹sin is for dS and sinh for AdS.

L.1 $\mathcal{Z}_{S^2}^{T\bar{T}}$ and $S_{T\bar{T}}$ for AdS

For the case of a CFT₂, the S^2 -partition function can be determined upon integrating (L.4) over the metric defined by the line element

$$ds^2 = r^2(d\theta^2 + \sin^2\theta d\phi^2) \quad (\text{L.7})$$

The partition function changes under a change in r , [240, 241], hence, from (L.4)

$$\begin{aligned} \frac{\partial \ln \mathcal{Z}_{S^2}^{AdS, T\bar{T}}}{\partial r} &= -\frac{1}{r} \int d^2x \sqrt{\gamma} T_a^a = -\frac{1}{r} \int d\theta d\phi \sin\theta r^2 \frac{2}{\lambda} \left(1 - \sqrt{1 + \frac{c\lambda}{12r^2}}\right) \\ &\Rightarrow \frac{\partial \ln \mathcal{Z}_{S^2}^{AdS, T\bar{T}}}{\partial r} = -\frac{8\pi}{\lambda} \left(r - \sqrt{r^2 + \frac{c\lambda}{12}}\right) \end{aligned} \quad (\text{L.8})$$

$$\Rightarrow \ln \mathcal{Z}_{S^2}^{AdS, T\bar{T}} = -\frac{8\pi}{\lambda} \int dr \left(r - \sqrt{r^2 + \frac{c\lambda}{12}}\right) = \frac{8\pi}{\lambda} \left[-\frac{r^2}{2} + \frac{c\lambda}{12} \int dx \cosh^2 x\right]$$

where $r(x) \stackrel{\text{def.}}{=} \sqrt{\frac{c\lambda}{12}} \sinh x$; the partition function becomes

$$\ln \mathcal{Z}_{S^2}^{AdS, T\bar{T}} = \pi \frac{c}{3} \sinh^{-1} \left(\sqrt{\frac{12}{c\lambda}} r\right) - \frac{4\pi r^2}{\lambda} \left[1 - \sqrt{\frac{c\lambda}{12 r^2} + 1}\right] \quad (\text{L.9})$$

up to an overall integration constant. The entanglement entropy can be determined from (L.8) and (L.9), [240],

$$S_{T\bar{T}} = \left(1 - \frac{r}{2} \frac{\partial}{\partial r}\right) \ln \mathcal{Z}_{S^2}^{AdS, T\bar{T}} = \pi \frac{c}{3} \sinh^{-1} \left(\sqrt{\frac{12}{c\lambda}} r\right) \quad (\text{L.10})$$

For later purposes, it is important to highlight that, (L.9) and (L.10) differ by means of the following term

$$-\frac{r}{2} \frac{\partial}{\partial r} \ln \mathcal{Z}_{S^2}^{AdS, T\bar{T}} = \frac{4\pi r^2}{\lambda} \left[1 - \sqrt{\frac{c\lambda}{12 r^2} + 1}\right] \quad (\text{L.11})$$

and therefore, one could argue that, under a suitable choice of λ , (L.11) can be set to vanish. In such case, $S_{T\bar{T}}$ would be uniquely determined by $\ln \mathcal{Z}_{S^2}^{AdS, T\bar{T}}$. We will be coming back to the importance of this observation later on in section 16.2.1.

For $r \gg \sqrt{\lambda c}$,

$$\begin{aligned} \sinh^{-1} x &= \ln \left|x + \sqrt{1+x^2}\right| = \ln \left|x + x \sqrt{1 + \frac{1}{x^2}}\right| \sim \\ &\sim \ln |x| + \left(1 + \frac{1}{2x^2}\right) = \ln |x| + \frac{2x^2 + 1}{2x^2} \end{aligned}$$

and (L.10) therefore turns into

$$S_{T\bar{T}} = \frac{c}{3} \ln \left|\sqrt{\frac{12}{c\lambda}} r\right| + \frac{c}{3} \frac{2\left(\frac{12}{c\lambda} r^2 + 1\right)}{2\frac{12}{c\lambda} r^2} - \frac{c}{3} \frac{1}{\frac{24}{c\lambda} r^2} \quad (\text{L.12})$$

By further substituting $r(x) \stackrel{\text{def.}}{=} \sqrt{\frac{c\lambda}{12}} \sinh x$, (L.12) can be brought down to the more familiar expression

$$S_{T\bar{T}} = \frac{c}{3} \ln |\sinh x| + \frac{c^2 \lambda}{36} \coth^2 x - \frac{c}{6} \frac{1}{\sinh^2 x} \quad (\text{L.13})$$

where the 1^{st} term is the entanglement entropy for an interval of length r at finite temperature $\beta \sim \sqrt{12/c\lambda}$. Equation (L.13), to leading order in the coupling λ , therefore reads

$$S_{T\bar{T}} = \frac{c}{3} \ln \left| \sqrt{\frac{12}{c\lambda}} r \right| + \frac{c^2 \lambda}{72} \frac{1}{r^2} + \frac{c}{3} \quad (\text{L.14})$$

From (L.12) we can now see that in the UV-limit, i.e. as we take $\lim_{r \rightarrow 0} S_{T\bar{T}}$, the first two terms cancel each other out, thereby preventing the entanglement entropy from diverging. This can be checked explicitly through de l'Hopital's rule for the $\infty - \infty$ case.

L.2 $\mathcal{Z}_{S^2}^{T\bar{T}}$ and $S_{T\bar{T}}$ for dS

For dS₃/dS₂, the corresponding geodesic in the space of 2D theories is given by, [242],

$$\boxed{\frac{\delta \mathcal{L}}{\delta \lambda} = 2 T\bar{T} - \frac{1}{\lambda^2}} \quad (\text{L.15})$$

which differs from the AdS case by means of the 2^{nd} term, taking into account that the CFT₂ is now coupled to gravity. The corresponding change to the sphere partition function now reads

$$\frac{\partial}{\partial \lambda} \ln \mathcal{Z}_{S^2}^{dS, T\bar{T}} = -2 \int d^2 x \sqrt{\gamma} \langle T\bar{T} \rangle + \frac{1}{\lambda^2} \int d^2 x \sqrt{\gamma} = -8\pi r^2 \langle T\bar{T} \rangle + \frac{4\pi r^2}{\lambda^2} \quad (\text{L.16})$$

From the factorisation of the $T\bar{T}$ expectation value,

$$\langle T\bar{T} \rangle = -\frac{1}{4} \langle T_\theta^\theta \rangle^2 = -\frac{1}{4 \lambda^2} \left(\frac{\lambda c}{12r^2} - 2\sqrt{\frac{\lambda c}{12r^2} - 1} \right) \quad (\text{L.17})$$

$$\Rightarrow \frac{\partial}{\partial \lambda} \ln \mathcal{Z}_{S^2}^{dS, T\bar{T}} = \frac{2\pi r^2}{\lambda^2} \left(\frac{\lambda c}{12r^2} - 2\sqrt{\frac{\lambda c}{12r^2} - 1} \right) + \frac{4\pi r^2}{\lambda^2} \quad (\text{L.18})$$

which can be integrated w.r.t. λ resulting in

$$\ln \mathcal{Z}_{S^2}^{dS, T\bar{T}} = \pi \frac{c}{6} \ln \left| \frac{\lambda c}{12r^2} \right| - \frac{4\pi r^2}{\lambda} - 4\pi r^2 \int d\lambda \frac{1}{\lambda^2} \sqrt{\frac{\lambda c}{12r^2} - 1} \quad (\text{L.19})$$

The last term can be integrated by parts

$$\begin{aligned} -4\pi r^2 \int d\lambda \frac{1}{\lambda^2} \sqrt{\frac{\lambda c}{12r^2} - 1} &= -4r^2 \left(\frac{1}{\lambda} \sqrt{\frac{\lambda c}{12r^2} - 1} + \int d\lambda \frac{1}{2\lambda} \frac{1}{\sqrt{\frac{\lambda c}{12r^2} - 1}} \frac{c}{12r^2} \right) = \\ &= -4\pi r^2 \left(\frac{1}{\lambda} \sqrt{\frac{\lambda c}{12r^2} - 1} + \frac{c}{12r^2} \tan^{-1} \left(\frac{1}{\sqrt{\frac{\lambda c}{12r^2} - 1}} \right) \right) \end{aligned}$$

Gathering all the terms, the partition function thus reads

$$\ln \mathcal{Z}_{S^2}^{dS, T\bar{T}} = \pi \left[\frac{c}{3} \ln \left| \sqrt{\frac{\lambda c}{12} \frac{1}{r}} \right| - \frac{4r^2}{\lambda} \left(1 - \sqrt{\frac{\lambda c}{12r^2} - 1} \right) + \frac{c}{3} \tan^{-1} \left(\frac{1}{\sqrt{\frac{\lambda c}{12r^2} - 1}} \right) \right] \quad (\text{L.20})$$

from which the entanglement entropy thus follows, [242],

$$S_{T\bar{T}} = \left(1 - \frac{r}{2} \frac{\partial}{\partial r} \right) \ln \mathcal{Z}_{S^2}^{dS, T\bar{T}} = \pi \left[\frac{c}{6} + \frac{c}{3} \ln \left| \sqrt{\frac{\lambda c}{12} \frac{1}{r}} \right| + \frac{c}{3} \tan^{-1} \left(\frac{1}{\sqrt{\frac{\lambda c}{12r^2} - 1}} \right) \right] \quad (\text{L.21})$$

Notice that (L.21) and (L.20) differ by means of the following terms

$$-\frac{r}{2} \frac{\partial}{\partial r} \ln \mathcal{Z}_{S^2}^{dS, T\bar{T}} = -\frac{c}{6} + \frac{4r^2}{\lambda} \left(1 - \sqrt{\frac{\lambda c}{12r^2} - 1} \right) \quad (\text{L.22})$$

which, as also argued for the AdS case, (L.11), can be set to vanish under suitable choice of λ .

L.3 Comparing $S_{T\bar{T}}$ for AdS and dS

Comparing the expressions for the partition functions on the dS and AdS side, i.e. (L.20) and (L.9) respectively, we can rewrite the \sinh^{-1} term in the latter as a ln,

$$\sinh^{-1} \left(\sqrt{\frac{12}{c\lambda}} r \right) = \ln \left| \sqrt{\frac{12}{c\lambda}} r + \sqrt{\frac{12r^2}{c\lambda} + 1} \right| = \ln \left| \sqrt{\frac{12}{c\lambda}} r \left(1 + \sqrt{\frac{c\lambda}{12r^2} + 1} \right) \right|$$

which is reminiscent of the usual structure of S_{EE} for a finite-T CFT. The AdS partition function therefore becomes

$$\ln \mathcal{Z}_{S^2}^{AdS, T\bar{T}} = \pi \frac{c}{3} \ln \left| \sqrt{\frac{12}{c\lambda}} r \left(1 + \sqrt{\frac{c\lambda}{12r^2} + 1} \right) \right| + \frac{4\pi}{\lambda} \left[r \sqrt{r^2 + \frac{c\lambda}{12}} - r^2 \right] \quad (\text{L.23})$$

suggesting the factor of $\sqrt{\frac{12}{c\lambda}} = \frac{1}{\epsilon}$ can be interpreted as the cutoff at which the dS can be glued in the IR of the original AdS. By means of such regularisation, (L.20) can be rewritten in terms of ϵ as follows

$$\ln \mathcal{Z}_{S^2}^{dS, T\bar{T}} = \pi \left[\frac{c}{3} \ln \left| \sqrt{\frac{\lambda c}{12} \frac{1}{\epsilon}} \right| - \frac{4r^2}{\lambda} \left(1 - \frac{1}{\pi} \sqrt{\frac{\lambda c}{12r^2} - 1} \right) + \frac{c}{3} \tan^{-1} \left(\frac{1}{\sqrt{\frac{\lambda c}{12r^2} - 1}} \right) \right] \quad (\text{L.24})$$

Notice that only the ϵ -dependent term diverges one upon taking the UV-limit with $\lim_{r \rightarrow 0}$, i.e. when probing short-distance scales. From the redefinition (L.24), the entanglement entropy for the $T\bar{T}$ deformed dS-theory now reads, [242],

$$S_{T\bar{T}} = \left(1 - \frac{r}{2} \frac{\partial}{\partial r}\right) \ln \mathcal{Z}_{S^2}{}^{dS, T\bar{T}} = \pi \left[\frac{c}{6} + \frac{c}{3} \ln \left| \sqrt{\frac{\lambda c}{12} \frac{1}{\epsilon}} \right| + \frac{c}{3} \tan^{-1} \left(\frac{1}{\sqrt{\frac{\lambda c}{12 r^2} - 1}} \right) \right] \quad (\text{L.25})$$

In the work of Silverstein et al., [242], the 2^{nd} term in (L.25) is set to vanish by appropriately tuning the value of the CFT parameters and consequently of their corresponding bulk duals. For our purposes, though, this cutoff dependence plays a key role, and will therefore be kept explicit. As a further remark, notice that, for the case in which the dS well approximates the original AdS, the universal parts of the entanglement entropies on the AdS and dS side agree, since the \tan^{-1} term in (L.25) can be rewritten as \sinh^{-1} by making the argument imaginary

$$\tan^{-1} \left(\frac{1}{\sqrt{\frac{\lambda c}{12 r^2} - 1}} \right) = \sinh^{-1} \left(\sqrt{\frac{12}{c\lambda}} r \right) \quad (\text{L.26})$$

thereby proving that, under a suitable definition of ϵ , dS can arise from the RG-flow of a bulk AdS. We will be coming back to this when comparing localisation calculation with the actions for vacuum transitions.

L.4 Cutoff dependence of S_{EE}

The divergence of S_{EE} for pure AdS when evaluated at $z = 0$ follows from removing the UV-cutoff ϵ acting as a regulator, [288]. When deforming the CFT with an irrelevant operator, S_{EE} will still be dependent on the cutoff. However, its explicit dependence can be removed by appropriately tuning the parameters in the UV and IR of the theory, thereby leading to finite S_{EE} s.

L.5 Key points of $T\bar{T}$ -deformations

- They constitute an example of UV/IR-mixing, and therefore provide an interesting setup where to address unitarity issues similar to the ones arising within the context of the information loss paradox.
- dS can arise as the IR-limit of an RG-flow driven by an irrelevant operator starting from a UV-theory dual to AdS.
- Localisation techniques play a key role in evaluating the partition function for a given CFT. Given its key role in determining S_{EE} , the extremisation procedure rooted in the formalism is compatible with the method defining S_{TOT} by means of the FMP method. It is therefore reasonable to argue, as we shall see in section J, that these quantities must be proportional to each other.
- The cutoff radius ensures the finiteness of $S_{T\bar{T}}$ on both sides, explicitly featuring only universal terms.
- According to the dimensionality of the CFTs being involved, the behaviour of the universal terms is predicted by the g, c and a - *theorems*.

- The main advantage of having introduced $\lambda \neq 0$ is that, in the limit $r/\sqrt{c\lambda} \rightarrow 0$, $S_{EE} \rightarrow 0$ and does not diverge upon taking the UV-limit, unlike the usual CFT result. This proves that, removing the regulator, unitarity is still preserved. As we will see, removing the CFT-cutoff is equivalent to taking the limit of the vanishing cosmological constant in the $dS \rightarrow dS$ transition in absence of a black hole.

M Von Neumann algebras (VNAs)

M.1 Types of VNAs

A von Neumann algebra (VNA) is a $*$ -subalgebra M of $\mathcal{B}(\mathcal{H})$ that coincides with its bicommutant

$$M = M'' . \quad (\text{M.1})$$

A VNA $M \subset \mathcal{B}(\mathcal{H})$ is called a factor if $M \cap M' = \mathbb{C}\mathbf{1}_{\mathcal{B}(\mathcal{H})}$.

VN factors can be of different types:

1. I_n , or I_∞ , classified by their dimension.
2. II_1 or II_∞ , with

$$II_\infty \simeq II_1 \otimes \mathcal{B}(\mathcal{H}) . \quad (\text{M.2})$$

3. III_λ with $\lambda \in [0, 1]$.

For the case of type- II_1 , a *projection* in a VNA M is an element $p \in M$ such that

$$p^* = p = p^2 . \quad (\text{M.3})$$

The type of factor depends on the existence of small projections. Specifically, a linear functional τ on a VNA M is

- Positive, if $\tau(x^*x) \geq 0 \forall x \in M$.
- Faithful, if $\tau(x^*x) = 0 \Rightarrow x = 0$.
- Stable, if $\tau(1) = 1$.
- Tracial, if $\tau(xy) = \tau(yx) \forall x, y \in M$. A tracial state is simply referred to as trace.

M.2 Standard form of a II_1 factor

Let M be a factor with continuous and faithful trace τ . The pairing

$$\langle x, y \rangle = \tau(y^*x) \in \mathbb{C} \quad (\text{M.4})$$

defines an inner product on M . Let $L^2(M, \tau)$ be the associated completion of M . Then there is an embedding

$$\begin{aligned} M &\longrightarrow L^2(M) \\ x &\mapsto \hat{x} \\ 1 &\mapsto \Omega . \end{aligned} \quad (\text{M.5})$$

M is represented on a Hilbert space $L^2(M)$ by considering

$$\pi_\tau(x)\hat{y} = \hat{x}y \equiv x\hat{y} \quad (\text{M.6})$$

and extending it to a morphism

$$\pi_\tau : M \longrightarrow \mathcal{B}(L^2(M)). \quad (\text{M.7})$$

This representation is called the standard form of M , implying that

$$\hat{x} = x\hat{1} = x\hat{\Omega} = x\Omega, \quad (\text{M.8})$$

such that

$$\tau(x) = \langle x\Omega, \Omega \rangle. \quad (\text{M.9})$$

M.3 Representations and von Neumann dimensions

A representation of a factor M is a Hilbert space \mathcal{H} with a structure of an M -module.

There is a Theorem, stating that any representation \mathcal{H} of a II_1 factor M is equivalent to

$$p \left(L^2(M) \otimes \ell^2(\mathbb{N}) \right), \quad (\text{M.10})$$

where p is a projection

$$p = vv^* \in (M \times 1)' \quad (\text{M.11})$$

whose trace is the size of the representation \mathcal{H}

$$\dim_M \mathcal{H} \equiv \text{Tr}(vv^*) \in [0, \infty]. \quad (\text{M.12})$$

Properties of the VN dimension

If M is a type- II_1 factor, then

$$\dim_M \bigoplus_{j \in J} \mathcal{H}_j \equiv \sum_{j \in J} \dim_M \mathcal{H}_j. \quad (\text{M.13})$$

If $p \in M'$ is a projection, then

$$\dim_M L^2(M)_p = \tau(p). \quad (\text{M.14})$$

Representations are classified by their M -dimension, and it therefore follows that

$$\dim_M \mathcal{H} = \dim_M \mathcal{K} \iff \mathcal{H} \simeq \mathcal{K}. \quad (\text{M.15})$$

M.4 The Jones index

Let $N \subset M$ be type II_1 -factors. Then N is a representation on $L^2(M)$.

The *Jones index* of the subfactor $N \subset M$ is defined as follows

$$[M : N] \stackrel{\text{def.}}{=} \dim_N L^2(M). \quad (\text{M.16})$$

If N is represented on \mathcal{H} with $\dim_N \mathcal{H} < \infty$, then

$$[M : N] = \frac{\dim_N \mathcal{H}}{\dim_M \mathcal{H}}. \quad (\text{M.17})$$

N Other appendices to Part IV

N.1 Twist operators and Rényi entropies

The definition of entanglement entropy involves a path integral over a Riemann surface, where the boundary conditions dictated by the twist field operators¹²⁰, \mathcal{T} , encode the nontrivial topological features of the theory, [288]. This is of particular interest whenever dealing with theories with no 1:1 correspondence between the solutions to the e.o.m. and the original Lagrangian density.

A similar issue arises when performing coordinate transformations in 2D gravity theories, such as maximal extensions, ultimately leading to a rich web of dualities. The bulk-boundary correspondence of AdS/CFT, as well as the factorisation problem, provide further examples. Before delving into a more detailed analysis of concrete examples, we wish to highlight first the importance of one major quantity relating all these topics together, which in turn prompted the question of understanding its relation with non-invertible symmetries, namely the Rényi entropies.

For later convenience, we will briefly overview the main features of the arguments outlined by Cardy and Calabrese for proposing this quantity as the suitable candidate to study CFT₂s in all their richness. In doing so, we will also recapitulate how the emergence of the island becomes manifest from a pure CFT/BCFT calculation as the change in dominance of connected/disconnected channels within correlation functions of twist operators, stressing the importance of the fact that S_{EE} for a single interval in a BCFT shares the same behaviour as that of 2 intervals on a CFT.

4-point functions in a CFT

In this digression, we briefly outline the derivation of S_{EE} , following the techniques of [288]. In presence of conformal symmetry breaking, $\langle T_\mu^\mu \rangle \neq 0$, and the boundary-to-boundary propagator picks up a correction of the kind $\langle \mathcal{T}TT_\mu^\mu \rangle$. The map from the Riemann surface¹²¹ to the complex plane is defined as, [288],

$$z = \xi^{1/n} = \left(\frac{w-u}{w-v} \right)^{1/n} \Rightarrow \mathcal{R}_{n,1} \rightarrow \mathbb{C} \quad (\text{N.1})$$

and the trace of the energy-momentum tensor transforms accordingly as follows

$$T(w) = \left(\frac{dz}{dw} \right)^2 T(z) + \frac{c}{12} \{z, w\} \quad (\text{N.2})$$

with the 2nd term in N.2 denoting the *Schwarzian derivative*, from which its expectation value becomes

$$\langle T(w) \rangle_{\mathcal{R}_{n,1}} = \langle T(z) \rangle_{\mathbb{C}} + \frac{c(1-n^{-2})}{24} \frac{(v-u)^2}{(w-u)^2(w-v)^2} \frac{\langle \mathcal{T}_n(u,0) \tilde{\mathcal{T}}_n(v,0) T(w) \rangle_{\mathcal{L}^{(n),\mathbb{C}}}}{\langle \mathcal{T}_n(u,0) \tilde{\mathcal{T}}_n(v,0) \rangle} \quad (\text{N.3})$$

with twist operators $\mathcal{T}_n, \tilde{\mathcal{T}}_n$ being primary fields of conformal dimension $\Delta_n, \bar{\Delta}_n$ determined from the 2-point function

$$\langle \mathcal{T}_n(u,0) \tilde{\mathcal{T}}_n(v,0) \rangle_{\mathcal{L}^{(n),\mathbb{C}}} = |u-v|^{-2\Delta_n} \Rightarrow \Delta_n = \frac{c}{12} \left(n - \frac{1}{n} \right) . \quad (\text{N.4})$$

Following the work of Cardy and Calabrese, [288], the 4-point function of the twist operators characterising the Riemann surface on which a field theory lives, defines the partition function for 2 disjointed intervals on the FT side. The *nth-Rényi entropy* is defined as

¹²⁰ Twist field operators are located at the endpoints of the intervals w.r.t. which S_{EE} is being evaluated.

¹²¹... where the n -point functions of the twist operators are evaluated.

$$S_{EE}^{(n)} \stackrel{\text{def.}}{=} \text{Tr}(\rho_A)^n = \frac{\mathcal{Z}_n}{\mathcal{Z}^n} \quad , \quad (\text{N.5})$$

with

$$\begin{aligned} \mathcal{Z}_n &\stackrel{\text{def.}}{=} \langle \mathcal{T}(z_1) \bar{\mathcal{T}}(z_2) \bar{\mathcal{T}}(z_3) \mathcal{T}(z_4) \rangle_{\text{cyl}} = \\ &= \left(\frac{2\pi}{\beta} \right)^{\frac{c}{3} \left(n - \frac{1}{n} \right)} |w_1 w_2 w_3 w_4|^{\frac{c}{12} \left(n - \frac{1}{n} \right)} \langle \mathcal{T}(w_1) \bar{\mathcal{T}}(w_2) \bar{\mathcal{T}}(w_3) \mathcal{T}(w_4) \rangle_{\mathbb{C}} \end{aligned}$$

with the complex coordinates w_i being related to the spacetime coordinates (t, r) via the following change of variables

$$\begin{aligned} w_1 &= e^{-2\pi(r_I+t_I)/\beta} e^{i\pi/2} \quad , \quad w_2 = e^{2\pi(r_I-t_I)/\beta} e^{i\pi/2} \\ w_3 &= e^{-2\pi(r_R+t_R)/\beta} e^{-i\pi/2} \quad , \quad w_4 = e^{2\pi(r_R-t_R)/\beta} e^{-i\pi/2} \end{aligned}$$

Given a subregion A obtained by partial tracing over the density matrix of the total system, the Rényi entropies $S^{(n)}$, the entanglement entropy S_{EE} , the reduced density matrix ρ_A and the partition function \mathcal{Z} are related according to the following

$$\boxed{S_A^{EE} \stackrel{\text{def.}}{=} \lim_{n \rightarrow 1} S_A^{(n)} = -\lim_{n \rightarrow 1} \left[\frac{1}{1-n} \ln \text{Tr} \rho_A^n \right] = -\lim_{n \rightarrow 1} \frac{\partial}{\partial n} \text{Tr} \rho_A^n = -\lim_{n \rightarrow 1} \frac{\partial}{\partial n} \frac{\mathcal{Z}_n(A)}{\mathcal{Z}^n}} \quad (\text{N.6})$$

Key points:

- n is assumed to be analytically continued in order for the derivative to be defined in (N.6).
- The need to consider 1 interval rather than 2 for the case of a BCFT, suggests one should be looking for a more generic approach towards describing the behaviour of the system for $n \in \mathbb{R}$.
- The twist operators are playing the same role as the non-invertible defects of a QFT featuring global internal symmetries.

N.2 $S_{tot}^{\text{AdS} \rightarrow \text{AdSBH}}$

In this appendix we will briefly argue how to deal with transitions involving a black hole only on one side of the brane. At the level of the Lagrangian density, this amounts to turning on the mass deformation on, either, the interior or the exterior vacua. The procedure to obtain the total action turns out to be a combination of the cases outlined in sections 14 and 15, and we therefore quote the key steps, referring the reader to the aforementioned parts of this work for further details and intermediate steps.

The extrinsic curvatures on the 2 sides of the brane would now be

$$\left. \frac{\phi'}{L} \right|_{\pm} = \frac{\Lambda_+ - \Lambda_- \mp \frac{\kappa^2}{4}}{\kappa} \phi_b + \frac{\mathcal{B}}{\kappa}$$

Assuming the black hole lies in the inner spacetime¹²², the turning point can be determined from the potential

$$\phi_{1,2} = -\frac{\left(\Lambda_+ - \Lambda_- - \frac{\kappa^2}{4}\right) \mathcal{B}}{\kappa} \phi_o^2 \left[1 \pm \sqrt{1 - \frac{4\kappa^2 \left(\frac{\mathcal{B}^2}{\kappa^2} + \mathcal{C}\right)}{\left(\Lambda_+ - \Lambda_- - \frac{\kappa^2}{4}\right)^2 \mathcal{B}^2 \phi_o^2}} \right]$$

The bulk horizon contribution will only come from the AdSBH,

$$S_{hor} = \frac{\eta}{G} \frac{\mathcal{B}}{\Lambda_-} \left[1 - \sqrt{1 - \frac{4\mathcal{C}\Lambda_-}{\mathcal{B}^2}} \right]$$

whereas the brane action would now read

$$\begin{aligned} S_{brane} &= \frac{2\eta}{G} \int_{\phi_1}^{\phi_2} d\phi_b \left[\cosh^{-1} \left[\frac{\left(\Lambda_+ - \Lambda_- + \frac{\kappa^2}{4}\right) \phi_b + \mathcal{B}}{\kappa \sqrt{-\mathcal{C} - \mathcal{B}\phi_b - \Lambda_- \phi_b^2}} \right] - \cosh^{-1} \left[\frac{\left(\Lambda_+ - \Lambda_- - \frac{\kappa^2}{4}\right) \phi_b + \mathcal{B}}{\kappa \sqrt{-\mathcal{C} - \Lambda_+ \phi_b^2}} \right] \right] \\ &= \frac{2\eta}{G} \left[-\frac{\mathcal{B}}{2\Lambda_-} \ln \left| \frac{y_2}{y_1} \sqrt{\frac{1-y_1^2}{1-y_2^2}} \right| - \phi_2 \ln \left| \frac{A_2 y_2 \sqrt{1-y_4^2}}{A_4 y_4 \sqrt{1-y_2^2}} \right| + \phi_1 \ln \left| \frac{A_2 y_1 \sqrt{1-y_3^2}}{A_4 y_3 \sqrt{1-y_1^2}} \right| + \right. \\ &\quad \left. - \frac{1}{2} \sqrt{\frac{\mathcal{B}^2}{4\Lambda_-} - \mathcal{C}} \ln \left| \frac{1-y_2}{1+y_2} \frac{1+y_1}{1-y_1} \right| + \mu_+ \ln \left| \frac{1-y_4}{1+y_4} \frac{1+y_3}{1-y_3} \right| \right]. \end{aligned}$$

¹²²Labelled by the “-” sign.

N.3 No obstructions from the Generalised Second Law (GSL)

Wormhole traversability has been amply studied in the last 2 decades thanks to the development of the AdS/CFT correspondence, [203]. The birth of an inflating universe from an asymptotically flat background has been argued to violate wormhole traversability as a consequence of the *generalised 2nd law* (GSL), [308]. However, we would like to highlight that the results obtained in the present paper by means of the Hamiltonian formalism show that this does not hold in our case for several reasons that we will now outline. In light of the closing remarks in section 13, the total action for the transition can be recast in the form of the difference of 2 $S_{EE}^{T\bar{T}}$. Relying upon the holographic understanding of such kinds of theories, we can deduce that the transition is only involving a subregion of the entire spacetime, and is, therefore, compatible with the traversable wormhole picture first outlined by Gao, Jafferis and Wall (GAO) by means of double-trace deformations, [307]. In summary, in our setup, there is no obstruction to up-tunnelling due to the following features:

1. Transitions are local
2. The instanton is singular and the metric is degenerate
3. The transition amplitude is related to the Entanglement entropy, S_{EE} , rather than the Von Neumann entropy. This justifies the necessity for topology change to take place whenever describing vacuum transitions. Notice that, in the Hamiltonian formalism, transitions are only described on spatial slices and not on the entire manifold. For wormhole saddles of the kind expected to emerge in the path integral of quantum gravity, the entire spacetime should really be understood of as being multiply-connected.


As long as the manifold is simply connected, the action is related to a *monotonic* S_{EE} , and therefore behaves as the ordinary von Neumann entropy of a bipartite system. In any other case, though, such as for when describing up-tunnelling, the von Neumann entropy needs to be upgraded to S_{EE} , meaning spacetime is multiply-connected. The singular metric involved in up-tunnelling is unsuitable for applying the GSL, given that the latter takes into account the causal dependence in between different spacetimes at \mathcal{I}^\pm . But this is different from the local transitions we are dealing with. The Hamiltonian constraint defines the energy of the system, i.e. not of the entire spacetime, but only that of the local subregion involved in the transition.

Singular instantons

The present appendix is meant to be a short digression providing further insight into the singular instanton solution obtained by FGG. As also argued by the authors in [234], there are 2 possible ways in which the singularity might arise:

1. Either in the past of the spacetime, or
2. In the wormhole configuration mediating the tunnelling process (double analytic continuation).

The former can be avoided via the quantum tunnelling interpolating between 2 classical solutions, defined as *type-a* and *type-b* trajectories. The latter is unavoidable given the change in sign of the lapse function in the forbidden region, as already noticed in [210].



For the quantum process to take place, and effectively interpolate in between the 2 classical trajectories depicted in the figure above, the brane's mass, M , must be at least smaller w.r.t, M_S . i.e. the Schwarzschild mass. This ensures that the wall's trajectory intersects the horizon and the extrinsic curvature changes sign. In particular, [234] choose $M < M_D$. The reason for this is that, being M smaller than any other scale of the theory, it can be tuned to be arbitrarily small, ultimately enabling to circumvent the emergence from the initial singularity¹²³ as we will now briefly outline. Given a 2-surface, P , its past light-cone will inevitably intersect the singularity with at least one light ray. However, in the weak gravity limit, it is possible to create a bubble of suitable radius to be at the 1st turning point of the potential barrier, treating the nucleation process as a pure quantum fluctuation of a flat background.

N.4 Roots of a quartic

In this appendix we outline the procedure for determining the extremising value of $\bar{\rho}$ for an (A)dS₂ → (A)dS₂ in the thin-wall approximation, (13.56), outlined in section 13.2. Upon extremising B_{TOT} w.r.t. $\bar{\rho}$, we end up with a polynomial with all even powers of $\bar{\rho}$. In particular, redefining $x \stackrel{def.}{=} \bar{\rho}^2$, it can be brought down to the following form

$$x^4 + a_3x^3 + a_2x^2 + a_1x + a_0 = 0 \quad (\text{N.7})$$

By further redefining $y \stackrel{def.}{=} x + \frac{a_3}{4}$, (N.7) becomes

$$y^4 + py^2 + qy + r = 0 \quad (\text{N.8})$$

Adding and subtracting $y^2u + \frac{u^2}{4}$, (N.8) becomes

$$\left(y^2 + \frac{1}{2}u\right)^2 - \left[(u-p)x^2 - qx + \left(\frac{u^2}{4} - r\right)\right] = 0 \quad (\text{N.9})$$

¹²³[234] also argue in favour of this by means of the so-called trapped and anti-trapped surfaces, with the latter being those to which the Penrose Theorem applies.

After a suitable manipulation, it is possible to factorise it into the product of 2 quadratics subject to satisfying a cubic constraint equation, namely

$$q^2 = 4(u-p) \left(\frac{u^2}{4} - r \right) \quad \text{with} \quad \begin{cases} p \stackrel{\text{def.}}{=} a_2 - \frac{3}{8}a_3^2 \\ q \stackrel{\text{def.}}{=} a_1 - \frac{1}{2}a_2a_3 + \frac{1}{8}a_3^3 \\ r \stackrel{\text{def.}}{=} a_o - \frac{1}{4}a_1a_3 + \frac{1}{16}a_2a_3^2 - \frac{3}{256}a_3^4 \end{cases} \quad (\text{N.10})$$

The corresponding solution $u_o = t_o - \frac{p}{3}$, with

$$t_o \stackrel{\text{def.}}{=} \frac{2}{3}\sqrt{p^2 + 12r} \cos \left[\frac{1}{3} \cos^{-1} \left(\frac{2p^3 + 9(3q^2 - 8rp)}{18} \left(\frac{3}{p^2 + 12r} \right)^{3/2} \right) \right] \quad (\text{N.11})$$

Once having obtained such factorisation, the corresponding value of the turning point can be extracted. In the 4D case, extremisation leads to a relation which is equivalent to the junction conditions in BT, and given that the turning point can be analytically computed in such case, it also corresponds to the value of the bubble radius extremising B_{tot} . In this case, though, it is not so straightforward, therefore we need to calculate it explicitly from the extremising procedure. However, we anticipate that (13.56) prior to extremisation already shares the same structure as the total bounce for type-1 instantons as found by BT in 2D. The turning point reads

$$\bar{\rho}^2 = -\frac{2A_3 + A_5}{2} \left[1 \pm \sqrt{1 - \frac{4(A_3^2 + A_4 + A_5A_6)}{(2A_3 + A_5)^2}} \right] \quad (\text{N.12})$$

with

$$A_3 \stackrel{\text{def.}}{=} \frac{a_3}{4}, \quad A_4 \stackrel{\text{def.}}{=} \frac{1}{2} \left(t_o - \frac{p}{3} \right), \quad A_5 \stackrel{\text{def.}}{=} \sqrt{t_o - \frac{p}{3}}, \quad A_6 \stackrel{\text{def.}}{=} -\frac{1}{2} \sqrt{\frac{(t_o - \frac{p}{3})^2 - 4r}{t_o - \frac{4p}{3}}} \quad (\text{N.13})$$

O Geometric Invariant Theory (GIT)

Delving in the explicit description of the duality structure arising from class \mathcal{S} theories, to which Part V is devoted, requires some key preliminary algebro-geometric tools, specifically the definition of Coulomb and Higgs branches from geometric invariant theory, [140, 142–144, 346, 379, 598, 599, 614, 615].

This appendix recapitulates the essential tools required for performing GIT quotient constructions, highlighting its importance for realising 3D mirror symmetry. In particular, we focus on the calculation of the Hilbert series as specific examples of algebraic varieties of crucial interest in the study of categorical dualities for supersymmetric quiver gauge theories.

O.1 Equivariant cohomology

Equivariant cohomology, [386], also known as Borel cohomology, is a cohomology theory from algebraic topology which applies to topological spaces with a group action. It can be viewed as a generalisation of group cohomology. More explicitly, the equivariant cohomology ring of a space X with action of a topological group G is defined as the ordinary cohomology ring with coefficient ring Λ of the homotopy quotient $EG \times_G X$

$$H_G^\bullet(X; \Lambda) = H^\bullet(EG \times_G X; \Lambda). \quad (\text{O.1})$$




Figure 112. Key algebraic varieties and dualities between invariants.

The Kirwan map

The Kirwan map, first introduced in [138], states

$$H_G^\bullet(M) \rightarrow H^\bullet(M//_p G), \quad (\text{O.2})$$

where M is a Hamiltonian G -space, i.e. a symplectic manifold acted on by a Lie group G with a moment map

$$\mu: M \rightarrow \mathfrak{g}^*. \quad (\text{O.3})$$

$H^\bullet(M)$ is an equivariant cohomology ring of M , i.e. the cohomology ring of the homotopy quotient $EG \times_G M$ of M by G .

$$M//_p G = \mu^{-1}(p)/G \quad (\text{O.4})$$

is the symplectic quotient of M by G at a regular central value $p \in Z(\mathfrak{g}^*)$ of μ . It is defined as the map of equivariant cohomology induced by the inclusion

$$\mu^{-1}(p) \hookrightarrow M \quad (\text{O.5})$$

followed by the canonical isomorphism

$$H_G^\bullet(\mu^{-1}(p)) = H^\bullet(M//_p G). \quad (\text{O.6})$$

O.2 The GIT construction

Geometric invariant theory (GIT), [613] is a recipe for constructing quotients in algebraic geometry. An affine GIT quotient, or affine geometric invariant theory quotient of an affine scheme $X = \text{Spec} \mathcal{A}$ with an action by a group scheme G , is the affine scheme $\text{Spec}(\mathcal{A}^G)$, namely the prime spectrum of the ring of invariants of \mathcal{A} , and is denoted by $X//G$.

The input of the GIT construction consists of:

1. A complex reductive group, G .
2. A representation of G

$$X = \mathbf{A}^n. \quad (\text{O.7})$$

3. A character

$$\theta: G \rightarrow G. \quad (\text{O.8})$$

GIT takes this data and gives an open subset of X , namely the semistable locus

$$X_{\theta}^{ss}(G) \stackrel{\text{open}}{\subset} X, \quad (\text{O.9})$$

from which one can define the stack quotient as follows

$$X//_{\theta}(G) \stackrel{\text{def.}}{\equiv} [X_{\theta}^{ss}(G) / G]. \quad (\text{O.10})$$

As (O.10) suggests, a GIT quotient is therefore a categorical quotient of the locus of semistable points, i.e. the quotient of the semistable locus.

Examples of GIT are Grassmannians, whose definition goes as follows. Take G a complex (connected) reductive group. Let \mathcal{O} denote the formal power series ring $\mathbf{C}[[z]]$ and \mathcal{K} its fraction field $\mathbf{C}((z))$. The Grassmannian is defined by the group quotient

$$\text{Gr}_G \stackrel{\text{def.}}{=} G_{\mathcal{K}}/G_{\mathcal{O}}, \quad (\text{O.11})$$

where $G_{\mathcal{K}}$ and $G_{\mathcal{O}}$ denote the groups of \mathcal{K} and \mathcal{O} -valued points of G , respectively. More formally, Gr_G is defined as being the moduli space of pairs (\mathcal{P}, φ) of a G -bundle, \mathcal{P} , on the formal disk

$$D \stackrel{\text{def.}}{=} \text{Spec}(\mathcal{O}), \quad (\text{O.12})$$

and its trivialisation φ over the punctured disk

$$D^* \stackrel{\text{def.}}{=} \text{Spec}(\mathcal{K}). \quad (\text{O.13})$$

Similarly, for the case of a complex representation, \mathbf{N} , the moduli space is now parameterised by a triple

$$\mathcal{R} \stackrel{\text{def.}}{=} (\mathcal{P}, \varphi, s), \quad (\text{O.14})$$

where (\mathcal{P}, φ) is in Gr_G , and s is a section of an associated vector bundle $\mathcal{P}_N = \mathcal{P} \times_G \mathbf{N}$, such that it is sent to a regular section of the trivial bundle under φ .

For a given triple \mathcal{R} , there is a corresponding projection

$$\pi : \mathcal{R} \rightarrow \text{Gr}_G, \quad (\text{O.15})$$

where Gr_G is the affine Grassmannian, and G is a complex reductive group. There is a natural commutative ring object, \mathcal{A} , in the derived $G_{\mathcal{O}}$ -equivariant constructible category on Gr_G , $D_G(\text{Gr}_G)$, such that $H_{\bullet}^{G_{\mathcal{O}}}(\mathcal{R})$ can be constructed from $H_{\bullet}^{G_{\mathcal{O}}}(\text{Gr}_G, \mathcal{A})$ equipped with a commutative ring structure.

In presence of a collection of such commutative ring objects, $\{\mathcal{A}_i\}$, in $D_G(\text{Gr}_G)$, one can perform a *gluing construction* by defining a new commutative ring object as

$$\iota_{\Delta}(\boxtimes \mathcal{A}_i), \quad (\text{O.16})$$

where

$$\iota_{\Delta} : \text{Gr}_G \rightarrow \prod_i \text{Gr}_G \quad (\text{O.17})$$

is the diagonal embedding. One such example are star-shaped quivers, [377].

Its equivariant Borel-Moore homology, [623], can be defined by the $G_{\mathcal{O}}$ -equivariant Borel-Moore homology of

$$\mathcal{R}_{\leq \lambda} \equiv \mathcal{R} \cap \pi^{-1} \left(\bar{\text{Gr}}_G^\lambda \right) \quad (\text{O.18})$$

in such a way that an embedding

$$\mathcal{R}_\mu \hookrightarrow \mathcal{R}_\lambda \quad (\text{O.19})$$

will induce a map

$$H_\bullet^{G^\circ}(\mathcal{R}_{\leq \mu}) \rightarrow H_\bullet^{G^\circ}(\mathcal{R}_{\leq \lambda}) \quad (\text{O.20})$$

for $\mu \leq \lambda$.

More generally, the euivariant Borel-Moore homology can be expressed as a graded sum

$$H_\bullet^{G^\circ}(\mathcal{R}) \simeq \bigoplus_\gamma H_\bullet^{G^\circ}(\mathcal{R}^\gamma), \quad (\text{O.21})$$

with \mathcal{R}^γ the connected component corresponding to $\gamma \in \pi_1(G)$.

O.3 Moduli spaces: Coulomb and Higgs branches

The Coulomb branch is an affine algebraic variety whose coordinate ring is the equivariant Borel-Moore homology group, $H_\bullet^{G^\circ}(\mathcal{R})$ of a certain space \mathcal{R} , defining the variety of triples.

The Coulomb branch is defined as, [599],

$$\mathcal{M}_C \stackrel{\text{def.}}{=} \text{Spec } \mathcal{A}(G, \mathbf{N}) \equiv \text{Spec} \left(H_\bullet^{G^\circ}(\mathcal{R}), * \right). \quad (\text{O.22})$$

Similarly, one can define the quantised Coulomb branch as follows

$$\mathcal{M}_C \stackrel{\text{def.}}{=} \text{Spec } \mathcal{A}_\hbar(G, \mathbf{N}) \equiv \text{Spec} \left(H_\bullet^{G^\circ \times \mathbf{C}^\times}(\mathcal{R}), * \right), \quad (\text{O.23})$$

which, in terms of quantum homology, can also be re-expressed as follows,

$$\mathcal{M}_C \stackrel{\text{def.}}{=} \text{Spec} \left(QH_G^\bullet(G; HF^\bullet(X)) \right). \quad (\text{O.24})$$

On the other hand, the Higgs branch is defined as the spectrum of the categorical quotient, namely, [599],

$$\mathcal{M}_H \stackrel{\text{def.}}{=} \text{Spec} \left(QH^\bullet(X//G) \right) \simeq QH_{LG}^\bullet(X). \quad (\text{O.25})$$

O.4 The monopole formula

The monopole formula contains all the gauge-invariant chiral operators that acquire a non-zero expectation value¹²⁴ along the CB. It is particularly useful for investigating expected properties of Coulomb branches, and reads as follows

$$P_t^{G^\circ}(\mathcal{R}) = \sum_k t^{-k} \dim H_k^{G^\circ}(\mathcal{R}). \quad (\text{O.26})$$

As already remarked in [599], (O.26) is the same as the Hilbert series on the Coulomb branch of a 3D $\mathcal{N} = 4$ supersymmetric quiver gauge theory associated with the pair $(G_C, \mathbf{N} \oplus \mathbf{N}^*)$.

¹²⁴Their specific VEV depends on their respective dimensions and quantum numbers.

P Magnetic quivers

P.1 Magnetic Quivers: a unifying framework

This section outlines some further preliminary background tools needed in Part V for addressing the formulation of mirror symmetry from the perspective of geometric representation theory, [3]. Mostly inspired by [192] and further upcoming work by Teleman, [193], we argue how constructions as the ones addressed in [1], involving multiple gauging, can still be assigned a fiber functor, as in the Freed-Moore-Teleman symmetry TFT (SymTFT) setup, [102]; however, such fiber functor should not be associated to either of the absolute theories separated by the non-invertible defect, but, rather to a 3D theory whose partition function correctly accounts for the symplectic projection leading to either of the two absolute theories.

Our approach consists in a combination of the following:

1. Defining a fiber functor for different absolute theories connected by non-invertible defects, [1].
2. Relating mirror symmetry with the identification of a Drinfeld center, [193].
3. Recent advancements in the understanding of the Higgs branch (HB) structure of higher-dimensional quiver gauge theories with 8 supercharges by means of Coulomb branches (CBs) of magnetic quivers (MQs) associated to 3D $\mathcal{N} = 4$ gauge theories, [195–198, 222, 223].

In the current and next sections we mostly outline how the topics listed above combine together, and in the following sections we will provide a more detailed mathematical correspondence with the work of [192], highlighting how his constructions can effectively be generalised to higher dimensions and be naturally embedded in the setup of [102] precisely thanks to the understanding in terms of magnetic quivers of 3D $\mathcal{N} = 4$ theories addressed here.

One could see our proposal as further supporting the idea that higher-categorical symmetries probe representation theory structures.

In the present section, we review the correspondence between geometric and algebraic resolutions of framed Nakajima quiver varieties, [144, 344, 599], highlighting it as an interesting example of homological mirror symmetry. In particular, we emphasise the property the moment map and higher homologies need to satisfy to ensure agreement in between the calculation of the two Hilbert series. We conclude the section with a brief overview of Hasse diagram constructions via magnetic quivers for quiver gauge theories with 8 supercharges, highlighting an interesting 2-categorical structure arising when dealing with complete intersections.

Section P.1 is meant to provide a brief overview of some key elements we will be using throughout our treatment, emphasising the ones that are needed for building the connection with the work of [192, 193]¹²⁵, which will be the core focus of section 20. The first part of the present article is structured as follows:

- At first we review the correspondence between geometric and algebraic resolutions of framed Nakajima quiver varieties, [144, 344, 599], highlighting it as an interesting example of homological mirror symmetry¹²⁶. In particular, we emphasise the property the moment map and higher homologies need to satisfy to ensure agreement in between the two resolutions. Consequently, this is mapped to an equivalence in between the Hilbert series resulting from summing over characters.

¹²⁵Throughout the entire treatment, we will be assuming the basic knowledge of higher categories and ADE quivers. We refer the reader to the extended literature on both topics for detailed definitions and examples. Specific additional tools will be explained in due course when needed.

¹²⁶The main explanation for this statement will become manifest in Part III of our treatment.

- We then turn to a brief overview of Higgs and Coulomb branches¹²⁷ of quiver gauge theories as algebraic varieties, pointing out the role of magnetic quivers (MQs) for describing the HB of a quiver gauge theory with 8 supercharges and arbitrary flavour, [195–198, 222, 223]. In doing so, we highlight the importance of the role played by, both, the Hilbert series (HS) and highest weight generating function (HWG) in identifying different intersecting cones in the HB Hasse diagram, [195–198, 222, 223], suggesting a 2-categorical structure.

In section 20 we explain how gauging can be related to the poset ordering leading to the construction of Hasse diagrams, thanks to the unifying role of the moment map. We then explain how the identification of such moment map ensures the quiver gauge theory enjoys a generalised notion of homological mirror symmetry, with the latter corresponding to the presence of a Drinfeld center and a corresponding fiber functor for a 2-categorical structure, related to Rozansky-Witten theory, [343]. We conclude the section highlighting connections between the topics outlined in the present work and those of [192, 193], thereby opening the scene to the more mathematical treatment to which Part III is devoted.

As we shall see, the underlying motivation for building the connections explained in section 20 is the importance of identifying a unifying framework for realising mirror symmetry and its generalisations.

P.2 Intersecting cones

From the more mathematically-inclined overview of section E.3, we know how to obtain a configuration admitting homological mirror symmetry, and the conditions that must be satisfied for the Hilbert series to match on the GIT and derived representation scheme side. For the purpose of what we will be addressing in section 15, we will now overview a setup where homological mirror symmetry can be described in an interestingly generalised and controlled manner, suggesting this as a probe for investigating its underlying mathematical structure.




Figure 113. Moduli space with Coulomb branch (CB), \mathcal{M}_C , Higgs branch (HB), $\mathcal{M}_H \equiv \mathcal{M}_B \cup \mathcal{M}_M$, with its nontrivially intersecting mesonic and baryonic branch, denoted by \mathcal{M}_B and \mathcal{M}_M , respectively.

Higgs and Coulomb branches as algebraic varieties

The main focus of this section are quiver gauge theories with 8 supercharges, classical flavour and gauge groups, whose vacuum moduli space is related to the nilpotent orbits, \mathcal{O}_ρ , of classical

¹²⁷Simply denoted by HB and CB for convenience.

Lie algebras, \mathfrak{g} . For a given gauge group G , the nilpotent orbits of an algebra \mathfrak{g} are defined by equivalence classes of nilpotence conditions on representation matrices, [199]. From the theoretical physics point of view, the nilpotent conditions describe the way in which the scalar fields in the F -term equations vanish at the SUSY vacuum, and can be specified by a quiver gauge theory.

According to the Jacobson Morozov Theorem, these nilpotent orbits are in 1 to 1 correspondence with equivalence classes of embeddings of $\mathfrak{su}(2)$ into \mathfrak{g} . Each such embedding

$$\rho : \mathfrak{su}(2) \rightarrow \mathfrak{g} \quad (\text{P.1})$$

is a homomorphism, labelled, by, either, the partition of the representation of G , or by a characteristic, using Dynkin labels to specify the mapping of the roots and weights of \mathfrak{g} onto $\mathfrak{su}(2)$.

A *Slodowy slice*, $S_\rho \stackrel{\text{def.}}{=} \mathcal{O}_\rho^\perp$, is a space transverse to a nilpotent orbit, and therefore commuting with it, while living within the adjoint orbit of the ambient group G . These transverse spaces can be further restricted to their intersections with the closure of any enclosing nilpotent orbit $\overline{\mathcal{O}}_\sigma$, thereby leading to spaces labelled by pairs of nilpotent orbits, whose elements are *Slodowy interesections*

$$S_{\sigma,\rho} \stackrel{\text{def.}}{=} \overline{\mathcal{O}}_\sigma \cap S_\rho \quad (\text{P.2})$$

encoding, as particular examples, nilpotent orbits, Slodowy slices and Kraft-Procesi transitions¹²⁸.

(P.2) are the algebraic varieties of which a Hilbert series can be calculated¹²⁹, as we will explain later on. Prior to turning to that we outline some preliminary definitions that will be recursively used throughout the remainder of this work.

The *nilcone* is defined as

$$\overline{\mathcal{O}}_{max} \stackrel{\text{def.}}{=} \mathcal{N}. \quad (\text{P.3})$$

with dimension

$$|\mathcal{N}| \equiv |\mathfrak{g}| - \text{rank}[\mathfrak{g}]. \quad (\text{P.4})$$

Nilpotent orbits can be arranged as a *Hasse diagram* according to the inclusion relations of their closures:

$$\mathcal{N} \equiv \overline{\mathcal{O}}_{max} \supset \overline{\mathcal{O}}_{sub-reg} \supset \dots \overline{\mathcal{O}}_{min} \supset \overline{\mathcal{O}}_{trivial} \equiv \{0\}. \quad (\text{P.5})$$

From the definition of the Slodowy intersection, it follows that

$$S_{\mathcal{N},trivial} \equiv \overline{\mathcal{O}}_{max} \equiv \mathcal{N} \quad (\text{P.6})$$

By means of such terminology, the definition of the *Higgs* and *Coulomb branches* can be expressed as follows

$$\overline{\mathcal{O}}_{max} \stackrel{\text{def.}}{=} \text{HB}[\mathcal{M}_A(\rho, 0)] \quad , \quad S_{\mathcal{N},\rho^T} \stackrel{\text{def.}}{=} \text{CB}[\mathcal{M}_A(\rho, 0)] \quad , \quad \mathcal{M}_A(\rho, 0) \stackrel{\text{def.}}{=} \mathcal{L}_A(\rho^T) \quad (\text{P.7})$$

with $\mathcal{M}_A(\rho, 0)$ denoting a single-flavoured linear quiver.

¹²⁸The connection between the 3D boundary conditions on type-II brane systems in 4D $\mathcal{N} = 4$ CFTs and Slodowy intersections was highlighted in [123].

¹²⁹Hilbert series for Slodowy intersections can also be constructed by means of purely group theoretic methods, making use of localisation formulae related to the Hall Littlewood polynomials.

P.2.1 Magnetic quivers and Hasse diagrams

Having said this, we now briefly overview the prescription for constructing magnetic quivers and Hasse diagrams from quiver subtraction, [200]¹³⁰. The procedure can be summarised by the following steps:

1. Start from a certain quiver gauge theory with 8 supercharges. Focussing on SQCD theories, namely a 3-parameter family of theories labelled by (N_c, N_f, k) , denoting number of colours, flavours, and Chern-Simons level, respectively. As a framed Nakajima quiver variety, we assign to it an electric quiver of the kind 83. Its HB is an algebraic variety associated to a 5-brane web configuration.
2. Identify the number of maximal decompositions of the 5-brane web.
3. To each maximal decomposition associate the magnetic quiver (MQ) of a 3D $\mathcal{N} = 4$ gauge theory.
4. The HB of the SQCD theory is preserved under dimensional reduction, and is equivalent to the union of the CBs of the MQs identified in step 3.
5. Construct the Hasse diagram associated to the HB by implementing quiver subtraction¹³¹ on the MQs.

Most importantly, this procedure generalises the following relation:

$$\boxed{\text{HB}(T) \equiv \text{CB}(T^\vee)}, \tag{P.8}$$

with T a 3D $\mathcal{N} = 4$ quiver gauge theory, and T^\vee its mirror dual. (P.8) can also be re-expressed in terms of the electric and magnetic quivers

$$\boxed{\text{HB}(\text{EQ}) \equiv \text{CB}(\text{MQ})}. \tag{P.9}$$

The advantage of the procedure put forward by [195–198, 222, 223] is that, when T does not admit a unique T^\vee , its *generalised* mirror symmetric theory can still be recovered by specifying the MQs, or, equivalently, the generators of the cones featuring in the Hasse diagram, together with their intersections. For concreteness, we briefly overview some key examples of the procedure summarised above, encompassing the main features we will be needing to consider in section 15.

P.2.2 Rank-1 and rank-2 examples

The examples reproduced in this subsection are for 5D $\mathcal{N} = 1$ SQCD, with varying parameters (N_c, N_f, k) . Notice that, for $SU(2)$, $k = 0$, reason why it is conventionally omitted in the quiver representation. However, for $N_c > 2$ it might be $k \neq 0$, reason why in such case we specify this in the corresponding quiver.

For each case we draw brane intersections (NS5s-branes and (p,q)-branes are always present). When adding flavour, D5-branes also contribute to the intersection. On the right of each brane intersection, we show the Hasse diagram constructed by implementing quiver subtraction on the

¹³⁰Such terminology is explained in due course. We refer to the extensive literature, especially the one in the references, for more detailed explanations as well as several examples explicitly analysed.

¹³¹We are assuming familiarity with the notion of ADE quivers. We refer to the extended literature on the topic if needed.

magnetic quiver. Conventionally, the node at the bottom of the Hasse diagram denotes the original magnetic quiver, prior to implementing any quiver subtraction. At each step of the iteration, we draw a vertical line, ending at another node, which therefore denotes a reduced magnetic quiver. When a bifurcation takes place, it signals that there are multiple possible quiver subtractions that could be performed. This corresponds to the emergence of multiple cones in the Hasse diagram. In case no bifurcations are present, it means the cone is unique.

5D $SU(N_c)$, $N_f = 0$




Figure 114. In absence of flavour degrees of freedom, the 5-braneweb decomposition is unique (LHS) leading to a HB Hasse diagram characterised by a unique cone (RHS).

$$\text{HB}_\infty^{5\text{D}} \left(\begin{array}{c} 0 \\ \square \\ | \\ \circ \\ \text{SU}(N_c)_0 \end{array} \right) \equiv \mathbb{C}^3 / \mathbb{Z}_{N_c} \equiv \text{CB}^{3\text{D}} \left(\begin{array}{c} N_c \\ \text{---} \\ \circ \quad \circ \\ \text{---} \\ 1 \quad 1 \end{array} \right) \quad (\text{P.10})$$

5D $SU(2)$, $N_f = 2$

Figure 115. 5-braneweb with added flavour (LHS) and HB Hasse diagram with two cones.

$$\text{HB}_\infty^{5\text{D}} \left(\begin{array}{c} 2 \\ \square \\ | \\ \circ \\ \text{SU}(2) \end{array} \right) \equiv \text{CB}^{3\text{D}} \left(\begin{array}{c} 1 \\ \triangle \\ \circ \quad \circ \\ \text{---} \\ 1 \quad 1 \end{array} \right) \cup \text{CB}^{3\text{D}} \left(\begin{array}{c} N_c \\ \text{---} \\ \circ \quad \circ \\ \text{---} \\ 1 \quad 1 \end{array} \right) \quad (\text{P.11})$$



Figure 116. Adding flavour, there are two possible decompositions of the braneweb. Each decomposition is mapped to a different magnetic quiver. The union of their CBs is equivalent to the HB of the original 5D quiver gauge theory we started from. The intersection of the two cones is given by a single symplectic leaf.

5D $SU(2)$, $N_f = 4$




Figure 117. 5-braneweb and corresponding Hasse diagram for 5D $SU(2)$, $N_f = 4$.

$$\text{HB}_{\infty}^{5\text{D}} \left(\begin{array}{c} 4 \\ \square \\ \circ \\ SU(2) \end{array} \right) \equiv \text{CB}^{3\text{D}} \left(\begin{array}{c} 1 \quad 1 \\ \circ \quad \circ \\ \circ - \color{red}{\circ} - \color{blue}{\circ} - \circ \\ 1 \quad 2 \quad 2 \quad 1 \end{array} \right) \quad (\text{P.12})$$

5D $SU(3)_0$, $N_f = 6$




Figure 118. 5-braneweb for 5D $SU(3)_0$, $N_f = 6$. The cone is unique since $k = 0$.

$$\text{HB}_\infty^{5\text{D}} \left(\begin{array}{c} 6 \\ \square \\ \circ \\ \text{SU}(3)_0 \end{array} \right) \equiv \text{CB}^{3\text{D}} \left(\begin{array}{ccccc} & 1 & 2 & 1 & \\ & \circ & \circ & \circ & \\ \circ & \circ & \circ & \circ & \circ \\ 1 & 2 & 3 & 2 & 1 \end{array} \right) \quad (\text{P.13})$$

5D $\text{SU}(6)_2$ $\mathbf{N}_f = 8$




Figure 119. Adding a CS level $k \neq 0$ leads the coexistence of two maximal decompositions of the 5-braneweb. The corresponding Hasse diagram is therefore constituted by 2 intersecting cones. The intersection in this case goes along the vertical direction where the blue and red lines are parallel to each other.

$$\text{HB}_\infty^{5\text{D}} \left(\begin{array}{c} 8 \\ \square \\ \circ \\ \text{SU}(6)_2 \end{array} \right) \equiv \text{CB}^{3\text{D}} \left(\begin{array}{ccccccc} & & 1 & & & & \\ & 1 & \circ & 1 & & & \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \\ 1 & 2 & 2 & 2 & 1 & 2 & 1 \end{array} \right) \cup \text{CB}^{3\text{D}} \left(\begin{array}{ccccccc} & & & 1 & & & \\ & & & \circ & & & \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \\ 1 & 2 & 3 & 4 & 3 & 2 & 1 \end{array} \right) \quad (\text{P.14})$$

From the point of view of the HB of quiver gauge theories with 8 supercharges in 5, 4, and 3 dimensions share a similar structure. In particular, the ramification of the HB due to the presence of multiple cones is preserved when considering the derived category of BPS objects in the lower-dimensional theory.

The 2-categorical structure of complete intersections

As previously mentioned, there are quiver gauge theories with 8 supercharges that do not admit a mirror symmetric 3D $\mathcal{N} = 4$ theory description. The reason for this can be recast to the fact that its HB is not a single hyperkähler cone, but, rather, the union of two, with nontrivial intersection, such as the cases shown in figure 115 and 119.

For classical gauge groups $G \equiv U(N), Sp(N)$ or $USp(2N), SO(2N)$, the moduli space is a complete intersection, meaning it is an algebraic variety whose dimension, d , is given by the following expression

$$d \stackrel{def.}{=} g - r, \quad (\text{P.15})$$

where g, r respectively denote the generators and relations between them.

For the case in which a 3D $\mathcal{N} = 4$ gauge theory, T , with given gauge group and suitable choice of matter content, admits a mirror dual theory, T^\vee , the Hilbert series evaluated on the Coulomb branch (CB) of the former equals that of the Higgs branch (HB) of the latter,

$$\text{HS}[\text{CB}(T)] \equiv \text{HS}[\text{HB}(T^\vee)], \quad (\text{P.16})$$

hence

$$\text{HB}(\text{EQ}) \equiv \text{CB}(\text{MQ}). \quad (\text{P.17})$$

Perfect realisation of ordinary mirror symmetry in (P.16) follows from the HB and CB of the dual theory both being hyperkähler quotients. Recent developments in terms of magnetic quivers and Hasse diagrams highlighted the fact that setups where (P.16) effectively takes place involve Hasse diagrams featuring a single cone, such as in figure 117 and 118. This feature naturally emerges from the HS calculation in the sense that (P.18) would only account for the contribution from the generators of the unique cone involved in the construction of the diagram. On the other hand, for theories whose Hasse diagram is the union of two intersecting cones, (P.19) splits into terms associated to the generators of each individual cone, together with additional subtractive terms accounting for the nontrivial intersection in between them.

Extended calculation of HS for arbitrary gauge theories with 8 supercharges has been the main focus of several recent works, mostly [197, 198, 223]. As shown in [198], there are some cases in which one could trade a cone for an intersection, thereby indicating that the generators and the relations in between them encode the same information. This turns out to be of particular importance for our purposes, in particular in building the correspondence¹³² with the Drinfeld center¹³³.

Within the context of quiver gauge theories with 8 supercharges, the Hilbert series (HS) is a partition function counting chiral gauge-invariant operators, encoding the variety of vacua generated by such operators. The highest weight generating function (HWG), encodes the same information in a more succinct way, that is more useful to be dealing with for cases involving higher rank, [223].

The general expression reads

$$\text{HS}_{N_f}(t; x_1, \dots, x_k) \equiv \sum_j f_j(x_1, \dots, x_k) t^j, \quad (\text{P.18})$$

with t denoting a fugacity for the highest weight of the $SU(2)_R$ R -symmetry group providing the grading for the ring of functions, while $f_i(x_1, \dots, x_k)$ are sums of characters for irreducible representations of the global symmetry group. From this follows the equivalence with (E.28) and (E.29). In the notation of [223], this reads

$$\text{HS}_{N_f}(t; x_1, \dots, x_k) \equiv \int d\mu_G \text{HS} \left(\frac{\mathbb{C}[\mathcal{Q}, \tilde{\mathcal{Q}}]}{\sqrt{F\text{-terms}}} \right) \equiv \text{HS} \left(\frac{\mathbb{C}[M, \mathcal{Q}, \tilde{\mathcal{Q}}]}{\sqrt{\text{Rel.s}}} \right) \quad (\text{P.19})$$

¹³²A more detailed explanation of this will be provided in the next section, when comparing with [192].

¹³³As stated in the introduction, the focus of [3] was outlining the main proposal of connecting Drinfeld centers and magnetic quivers from a theoretical point of view, while reserving a more formal mathematical treatment of the same correspondence to followup work by the same author, [4, 5].

with M, \mathcal{Q} denoting mesons and baryons, respectively.

For the case of 4D $\mathcal{N} = 2$ SQCD with gauge group $G = SU(N_c)$ and N_f hypermultiplets in the fundamental representation of $SU(N_c)$, the calculation of P.19 requires:

1. solving the F -term equations, setting the derivatives of the superpotential to zero
2. identifying the gauge-invariant operators.

For $\mathcal{N} = 2$ $SU(2)$ $N_f = 2$, the HS is a sum of rational functions, [223],

$$\begin{aligned} \text{HS}(t; x_1, x_2) &\equiv \int d\mu_{SU(2)} F(t, z, x_1, x_2) \\ &\equiv \text{HS}(\mathbb{C}^2/\mathbb{Z}_2; t, x_1) + \text{HS}(\mathbb{C}^2/\mathbb{Z}_2; t, x_2) - 1 \end{aligned} \quad (\text{P.20})$$

with the first two terms corresponding to the two cones and the last term denoting the intersection in between them at the origin, as shown in figure 115. When expanding it in terms of powers of t , we get the Plethystic logarithm (PL),

$$\text{PL}(t; x_1, x_2) \equiv ([2; 0] + [0; 2])t^2 - ([2; 2] + 2[0; 0])t^4 + \dots \quad (\text{P.21})$$

with $[\cdot]$ denoting the characters of the corresponding representations of $SO(4)$. The character corresponding to a certain representation can be encoded in the corresponding Dynkin label. We can therefore choose a set of fugacities to keep track of them, enabling us to rewrite (P.20) and (P.21) as a highest weight generating function

$$\text{HWG}(t; \mu_1, \mu_2) \equiv \text{PE} \left[\left(\mu_1^2 + \mu_2^2 \right) t^2 - \mu_1^2 \mu_2^2 t^4 \right], \quad (\text{P.22})$$

with the first two terms corresponding to the generators of the two cones, and the last (subtractive) term to their intersection in the Hasse diagram.

Key points

The main message of this section can be summarised as follows:

- Quiver gauge theories with 8 supercharges are characterised by a structure fully specified by knowing the generators and intersections of the cones in their HB Hasse diagram. The latter is obtained by implementing quiver subtraction on MQs associated to maximal decompositions of the 5-brane webs associated to the electric quiver. As such, it admits a 2-categorical structure description.
- The moment map plays a key role in ensuring the identification of the underlying MQs. The flatness condition it is required to satisfy defines the HS, hence the HWG function, [344]. Combining the works of [195–198, 222, 223], this constitutes a sample realisation of homological mirror symmetry.
- In presence of more than one cone in the Hasse diagram, the identification of the MQs whose CBs' union equals the HB of the original theory, is a generalised statement of homological mirror symmetry.

We wish to emphasise that, while most of the above statements have been individually addressed in the literature, to the best of our knowledge, their combined application towards identifying the Drinfeld center in connection to CBs constitutes an original approach, to which Part V is devoted.

References

- [1] V. Pasquarella, *Categorical Symmetries and Fiber Functors from Multiple Condensing Homomorphisms from 6D $\mathcal{N} = (2, 0)$ SCFTs*, [arXiv:2305.18515 [hep-th]].
- [2] V. Pasquarella and F. Quevedo, *Vacuum transitions in two-dimensions and their holographic interpretation*, JHEP **05** (2023), 192 doi:10.1007/JHEP05(2023)192 [arXiv:2211.07664 [hep-th]].
- [3] V. Pasquarella, *Drinfeld Centers from Magnetic Quivers*, [arXiv:2306.12471 [hep-th]].
- [4] V. Pasquarella, *Moore-Tachikawa Varieties: Beyond Duality*, JHAP **3** (2023) no.4, 39-56 doi:10.22128/jhap.2023.741.1061 [arXiv:2310.01489 [hep-th]].
- [5] V. Pasquarella, *Factorisation Homology for Class \mathcal{S} Theories*, [arXiv:2312.06760 [hep-th]].
- [6] V. Pasquarella, *Particle Physics: a crash course for Mathematicians*, [arXiv:2404.08100 [hep-th]].
- [7] S. P. De Alwis, F. Muia, V. Pasquarella and F. Quevedo, *Quantum Transitions Between Minkowski and de Sitter Spacetimes*, Fortsch. Phys. **68** (2020) no.9, 2000069 doi:10.1002/prop.202000069
- [8] Green MB, Schwarz JH, Witten E. Superstring Theory: 25th Anniversary Edition. Cambridge University Press; 2012. Vol. 1
- [9] Green MB, Schwarz JH, Witten E. Superstring Theory: 25th Anniversary Edition. Cambridge University Press; 2012. Vol.2
- [10] Polchinski J. String Theory. Cambridge University Press; 1998. Vol.1
- [11] Polchinski J. String Theory. Cambridge University Press; 1998. Vol. 2
- [12] M. B. Green, *Supersymmetrical Dual String Theories and their Field Theory Limits: A Review*, Surveys High Energ. Phys. **3** (1984), 127-160 doi:10.1080/01422418308243456
- [13] J. H. Schwarz, *Superstring Theory*, Phys. Rept. **89** (1982), 223-322 doi:10.1016/0370-1573(82)90087-4
- [14] M. B. Green and J. H. Schwarz, *Anomaly Cancellation in Supersymmetric $D=10$ Gauge Theory and Superstring Theory*, Phys. Lett. B **149** (1984), 117-122 doi:10.1016/0370-2693(84)91565-X
- [15] D. J. Gross, J. A. Harvey, E. J. Martinec and R. Rohm, *The Heterotic String*, Phys. Rev. Lett. **54** (1985), 502-505 doi:10.1103/PhysRevLett.54.502
- [16] K. Wilson, Rev. Mod. Phys. 55, 583 (1983).
- [17] J. Polchinski, *What is string theory?*, [arXiv:hep-th/9411028 [hep-th]].
- [18] F. Quevedo, S. Krippendorff and O. Schlotterer, *Cambridge Lectures on Supersymmetry and Extra Dimensions*, [arXiv:1011.1491 [hep-th]].
- [19] B. C. Allanach, F. Quevedo and K. Suruliz, *Low-energy supersymmetry breaking from string flux compactifications: Benchmark scenarios*, JHEP **04** (2006), 040 doi:10.1088/1126-6708/2006/04/040 [arXiv:hep-ph/0512081 [hep-ph]].
- [20] W. Abdallah *et al.* [LHC Reinterpretation Forum], *Reinterpretation of LHC Results for New Physics: Status and Recommendations after Run 2*, SciPost Phys. **9** (2020) no.2, 022 doi:10.21468/SciPostPhys.9.2.022 [arXiv:2003.07868 [hep-ph]].

- [21] B. C. Allanach and H. Banks, *Hide and seek with the third family hypercharge model's Z' at the large hadron collider*, Eur. Phys. J. C **82** (2022) no.3, 279 doi:10.1140/epjc/s10052-022-10191-6 [arXiv:2111.06691 [hep-ph]].
- [22] *Charting the Fifth Force Landscape*, Phys. Rev. D **103** (2021) no.7, 075018 doi:10.1103/PhysRevD.103.075018 [arXiv:2009.12399 [hep-ph]].
- [23] B. Allanach, *Fits to measurements of rare heavy flavour decays*, [arXiv:2307.07532 [hep-ph]].
- [24] E. Hammou, Z. Kassabov, M. Madigan, M. L. Mangano, L. Mantani, J. Moore, M. M. Alvarado and M. Ubiali, *Hide and seek: how PDFs can conceal new physics*, JHEP **11** (2023), 090 doi:10.1007/JHEP11(2023)090 [arXiv:2307.10370 [hep-ph]].
- [25] Z. Kassabov, M. Madigan, L. Mantani, J. Moore, M. Morales Alvarado, J. Rojo and M. Ubiali, *The top quark legacy of the LHC Run II for PDF and SMEFT analyses*, JHEP **05** (2023), 205 doi:10.1007/JHEP05(2023)205 [arXiv:2303.06159 [hep-ph]].
- [26] S. Iranipour and M. Ubiali, *A new generation of simultaneous fits to LHC data using deep learning*, JHEP **05** (2022), 032 doi:10.1007/JHEP05(2022)032 [arXiv:2201.07240 [hep-ph]].
- [27] P. Nath, B. D. Nelson, H. Davoudiasl, B. Dutta, D. Feldman, Z. Liu, T. Han, P. Langacker, R. Mohapatra and J. Valle, *et al. The Hunt for New Physics at the Large Hadron Collider*, Nucl. Phys. B Proc. Suppl. **200-202** (2010), 185-417 doi:10.1016/j.nuclphysbps.2010.03.001 [arXiv:1001.2693 [hep-ph]].
- [28] S. Weinberg, *The Making of the standard model*, Eur. Phys. J. C **34** (2004), 5-13 doi:10.1140/epjc/s2004-01761-1 [arXiv:hep-ph/0401010 [hep-ph]].
- [29] L. McAllister and F. Quevedo, *Moduli Stabilization in String Theory*, [arXiv:2310.20559 [hep-th]].
- [30] M. Cicoli, M. Licheri, P. Piantadosi, F. Quevedo and P. Shukla, *Higher Derivative Corrections to String Inflation*, [arXiv:2309.11697 [hep-th]].
- [31] M. Cicoli, J. P. Conlon, A. Maharana, S. Parameswaran, F. Quevedo and I. Zavala, *String Cosmology: from the Early Universe to Today*, [arXiv:2303.04819 [hep-th]].
- [32] M. Cicoli, I. G. Etxebarria, F. Quevedo, A. Schachner, P. Shukla and R. Valandro, *The Standard Model quiver in de Sitter string compactifications*, JHEP **08** (2021), 109 doi:10.1007/JHEP08(2021)109 [arXiv:2106.11964 [hep-th]].
- [33] C. Crinò, F. Quevedo and R. Valandro, *On de Sitter String Vacua from Anti-D3-Branes in the Large Volume Scenario*, JHEP **03** (2021), 258 doi:10.1007/JHEP03(2021)258 [arXiv:2010.15903 [hep-th]].
- [34] S. AbdusSalam, S. Abel, M. Cicoli, F. Quevedo and P. Shukla, *A systematic approach to Kähler moduli stabilisation*, JHEP **08** (2020) no.08, 047 doi:10.1007/JHEP08(2020)047 [arXiv:2005.11329 [hep-th]].
- [35] M. Cicoli, K. Dutta, A. Maharana and F. Quevedo, *Moduli Vacuum Misalignment and Precise Predictions in String Inflation*, JCAP **08** (2016), 006 doi:10.1088/1475-7516/2016/08/006 [arXiv:1604.08512 [hep-th]].
- [36] M. Cicoli, F. Quevedo and R. Valandro, *De Sitter from T-branes*, JHEP **03** (2016), 141 doi:10.1007/JHEP03(2016)141 [arXiv:1512.04558 [hep-th]].
- [37] L. Aparicio, F. Quevedo and R. Valandro, *Moduli Stabilisation with Nilpotent Goldstino:*

- Vacuum Structure and SUSY Breaking*, JHEP **03** (2016), 036 doi:10.1007/JHEP03(2016)036 [arXiv:1511.08105 [hep-th]].
- [38] F. Quevedo, *Local String Models and Moduli Stabilisation*, Adv. Ser. Direct. High Energy Phys. **22** (2015), 219-243 doi:10.1142/9789814602686_0008
- [39] S. de Alwis, R. K. Gupta, F. Quevedo and R. Valandro, *On KKLT/CFT and LVS/CFT Dualities*, JHEP **07** (2015), 036 doi:10.1007/JHEP07(2015)036 [arXiv:1412.6999 [hep-th]].
- [40] S. S. AbdusSalam, C. P. Burgess and F. Quevedo, *MFV Reductions of MSSM Parameter Space*, JHEP **02** (2015), 073 doi:10.1007/JHEP02(2015)073 [arXiv:1411.1663 [hep-ph]].
- [41] F. Quevedo, *Local String Models and Moduli Stabilisation*, Mod. Phys. Lett. A **30** (2015) no.07, 1530004 doi:10.1142/S0217732315300049 [arXiv:1404.5151 [hep-th]].
- [42] S. de Alwis, R. Gupta, E. Hatefi and F. Quevedo, *Stability, Tunneling and Flux Changing de Sitter Transitions in the Large Volume String Scenario*, JHEP **11** (2013), 179 doi:10.1007/JHEP11(2013)179 [arXiv:1308.1222 [hep-th]].
- [43] C. P. Burgess, M. Cicoli and F. Quevedo, *String Inflation After Planck 2013*, JCAP **11** (2013), 003 doi:10.1088/1475-7516/2013/11/003 [arXiv:1306.3512 [hep-th]].
- [44] A. Font, F. Quevedo and S. Theisen, *A comment on continuous spin representations of the Poincare group and perturbative string theory*, Fortsch. Phys. **62** (2014), 975-980 doi:10.1002/prop.201400067 [arXiv:1302.4771 [hep-th]].
- [45] M. Cicoli, S. Krippendorff, C. Mayrhofer, F. Quevedo and R. Valandro, *D-Branes at del Pezzo Singularities: Global Embedding and Moduli Stabilisation*, JHEP **09** (2012), 019 doi:10.1007/JHEP09(2012)019 [arXiv:1206.5237 [hep-th]].
- [46] M. Cicoli, A. Maharana, F. Quevedo and C. P. Burgess, *De Sitter String Vacua from Dilaton-dependent Non-perturbative Effects*, JHEP **06** (2012), 011 doi:10.1007/JHEP06(2012)011 [arXiv:1203.1750 [hep-th]].
- [47] M. Cicoli, G. Tasinato, I. Zavala, C. P. Burgess and F. Quevedo, *Modulated Reheating and Large Non-Gaussianity in String Cosmology*, JCAP **05** (2012), 039 doi:10.1088/1475-7516/2012/05/039 [arXiv:1202.4580 [hep-th]].
- [48] S. Krippendorff, M. J. Dolan, A. Maharana and F. Quevedo, *D-branes at Toric Singularities: Model Building, Yukawa Couplings and Flavour Physics*, JHEP **06** (2010), 092 doi:10.1007/JHEP06(2010)092 [arXiv:1002.1790 [hep-th]].
- [49] R. Blumenhagen, J. P. Conlon, S. Krippendorff, S. Moster and F. Quevedo, *SUSY Breaking in Local String/F-Theory Models*, JHEP **09** (2009), 007 doi:10.1088/1126-6708/2009/09/007 [arXiv:0906.3297 [hep-th]].
- [50] S. S. AbdusSalam, B. C. Allanach, F. Quevedo, F. Feroz and M. Hobson, *Fitting the Phenomenological MSSM*, Phys. Rev. D **81** (2010), 095012 doi:10.1103/PhysRevD.81.095012 [arXiv:0904.2548 [hep-ph]].
- [51] S. Krippendorff and F. Quevedo, *Metastable SUSY Breaking, de Sitter Moduli Stabilisation and Kahler Moduli Inflation*, JHEP **11** (2009), 039 doi:10.1088/1126-6708/2009/11/039 [arXiv:0901.0683 [hep-th]].
- [52] J. P. Conlon, A. Maharana and F. Quevedo, *Towards Realistic String Vacua*, JHEP **05** (2009), 109 doi:10.1088/1126-6708/2009/05/109 [arXiv:0810.5660 [hep-th]].

- [53] M. Cicoli, C. P. Burgess and F. Quevedo, *Fibre Inflation: Observable Gravity Waves from IIB String Compactifications*, JCAP **03** (2009), 013 doi:10.1088/1475-7516/2009/03/013 [arXiv:0808.0691 [hep-th]].
- [54] J. P. Conlon, A. Maharana and F. Quevedo, *Wave Functions and Yukawa Couplings in Local String Compactifications*, JHEP **09** (2008), 104 doi:10.1088/1126-6708/2008/09/104 [arXiv:0807.0789 [hep-th]].
- [55] J. P. Conlon, R. Kallosh, A. D. Linde and F. Quevedo, *Volume Modulus Inflation and the Gravitino Mass Problem*, JCAP **09** (2008), 011 doi:10.1088/1475-7516/2008/09/011 [arXiv:0806.0809 [hep-th]].
- [56] M. Cicoli, J. P. Conlon and F. Quevedo, *General Analysis of LARGE Volume Scenarios with String Loop Moduli Stabilisation*, JHEP **10** (2008), 105 doi:10.1088/1126-6708/2008/10/105 [arXiv:0805.1029 [hep-th]].
- [57] C. P. Burgess, J. P. Conlon, L. Y. Hung, C. H. Kom, A. Maharana and F. Quevedo, *Continuous Global Symmetries and Hyperweak Interactions in String Compactifications*, JHEP **07** (2008), 073 doi:10.1088/1126-6708/2008/07/073 [arXiv:0805.4037 [hep-th]].
- [58] S. S. AbdusSalam, J. P. Conlon, F. Quevedo and K. Suruliz, *Scanning the Landscape of Flux Compactifications: Vacuum Structure and Soft Supersymmetry Breaking*, JHEP **12** (2007), 036 doi:10.1088/1126-6708/2007/12/036 [arXiv:0709.0221 [hep-th]].
- [59] M. Cicoli, J. P. Conlon and F. Quevedo, *Systematics of String Loop Corrections in Type IIB Calabi-Yau Flux Compactifications*, JHEP **01** (2008), 052 doi:10.1088/1126-6708/2008/01/052 [arXiv:0708.1873 [hep-th]].
- [60] J. P. Conlon, C. H. Kom, K. Suruliz, B. C. Allanach and F. Quevedo, *Sparticle Spectra and LHC Signatures for Large Volume String Compactifications*, JHEP **08** (2007), 061 doi:10.1088/1126-6708/2007/08/061 [arXiv:0704.3403 [hep-ph]].
- [61] D. Cremades, M. P. Garcia del Moral, F. Quevedo and K. Suruliz, *Moduli stabilisation and de Sitter string vacua from magnetised D7 branes*, JHEP **05** (2007), 100 doi:10.1088/1126-6708/2007/05/100 [arXiv:hep-th/0701154 [hep-th]].
- [62] J. P. Conlon, S. S. Abdussalam, F. Quevedo and K. Suruliz, *Soft SUSY Breaking Terms for Chiral Matter in IIB String Compactifications*, JHEP **01** (2007), 032 doi:10.1088/1126-6708/2007/01/032 [arXiv:hep-th/0610129 [hep-th]].
- [63] C. P. Burgess, P. G. Camara, S. P. de Alwis, S. B. Giddings, A. Maharana, F. Quevedo and K. Suruliz, *Warped Supersymmetry Breaking*, JHEP **04** (2008), 053 doi:10.1088/1126-6708/2008/04/053 [arXiv:hep-th/0610255 [hep-th]].
- [64] J. P. Conlon, D. Cremades and F. Quevedo, *Kahler potentials of chiral matter fields for Calabi-Yau string compactifications*, JHEP **01** (2007), 022 doi:10.1088/1126-6708/2007/01/022 [arXiv:hep-th/0609180 [hep-th]].
- [65] A. Avgoustidis, D. Cremades and F. Quevedo, *Wilson line inflation*, Gen. Rel. Grav. **39** (2007), 1203-1234 doi:10.1007/s10714-007-0454-y [arXiv:hep-th/0606031 [hep-th]].
- [66] J. P. Conlon and F. Quevedo, *Gaugino and Scalar Masses in the Landscape*, JHEP **06** (2006), 029 doi:10.1088/1126-6708/2006/06/029 [arXiv:hep-th/0605141 [hep-th]].
- [67] A. A. Belavin, A. M. Polyakov, A. S. Schwartz and Y. S. Tyupkin, *Pseudoparticle Solutions of the Yang-Mills Equations*, Phys. Lett. B **59** (1975), 85-87 doi:10.1016/0370-2693(75)90163-X

- [68] D. S. Freed and M. J. Hopkins, *Consistency of M-Theory on Non-Orientable Manifolds*, Quart. J. Math. Oxford Ser. **72** (2021) no.1-2, 603-671 doi:10.1093/qmath/haab007 [arXiv:1908.09916 [hep-th]].
- [69] D. S. Freed and M. J. Hopkins, *Invertible phases of matter with spatial symmetry*, Adv. Theor. Math. Phys. **24** (2020) no.7, 1773-1788 doi:10.4310/ATMP.2020.v24.n7.a3 [arXiv:1901.06419 [math-ph]].
- [70] D. S. Freed and C. Teleman, *Topological dualities in the Ising model*, Geom. Topol. **26** (2022), 1907-1984 doi:10.2140/gt.2022.26.1907 [arXiv:1806.00008 [math.AT]].
- [71] D. S. Freed, *Anomalies and Invertible Field Theories*, Proc. Symp. Pure Math. **88** (2014), 25-46 doi:10.1090/pspum/088/01462 [arXiv:1404.7224 [hep-th]].
- [72] D. S. Freed, *The cobordism hypothesis*, [arXiv:1210.5100 [math.AT]].
- [73] D. S. Freed, M. J. Hopkins and C. Teleman, *Consistent Orientation of Moduli Spaces*, doi:10.1093/acprof:oso/9780199534920.003.0019
- [74] D. S. Freed, M. J. Hopkins, J. Lurie and C. Teleman, *Topological Quantum Field Theories from Compact Lie Groups*, [arXiv:0905.0731 [math.AT]].
- [75] D. S. Freed, *Remarks on Chern-Simons Theory*, [arXiv:0808.2507 [math.AT]].
- [76] D. S. Freed, *Classical field theory and supersymmetry*,
- [77] D. S. Freed, D. R. Morrison and I. M. Singer, *Quantum field theory, supersymmetry, and enumerative geometry*,
- [78] D. S. Freed, M. J. Hopkins and C. Teleman, *Loop groups and twisted K-theory. II.*, [arXiv:math/0511232 [math.AT]].
- [79] D. S. Freed, M. J. Hopkins and C. Teleman, *Twisted K-theory and loop group representations. 1.*, [arXiv:math/0312155 [math.AT]].
- [80] D. S. Freed, *K theory in quantum field theory*, [arXiv:math-ph/0206031 [math-ph]].
- [81] D. S. Freed, *The Verlinde algebra is twisted equivariant K theory*, Turk. J. Math. **25** (2001), 159-167 [arXiv:math/0101038 [math.RT]].
- [82] D. S. Freed, *Dirac charge quantization and generalized differential cohomology*, [arXiv:hep-th/0011220 [hep-th]].
- [83] D. S. Freed and M. J. Hopkins, *On Ramond-Ramond fields and K theory*, JHEP **05** (2000), 044 doi:10.1088/1126-6708/2000/05/044 [arXiv:hep-th/0002027 [hep-th]].
- [84] D. S. Freed and E. Witten, *Anomalies in string theory with D-branes*, Asian J. Math. **3** (1999), 819 [arXiv:hep-th/9907189 [hep-th]].
- [85] P. Deligne, P. Etingof, D. S. Freed, L. C. Jeffrey, D. Kazhdan, J. W. Morgan, D. R. Morrison and E. Witten, *Quantum fields and strings: A course for mathematicians. Vol. 1, 2.*
- [86] D. S. Freed, *Five lectures on supersymmetry*, AMS, 1999, ISBN 978-0-8218-1953-1
- [87] D. Freed and K. Uhlenbeck, *Geometry and quantum field theory. Proceedings, Graduate Summer School on the Geometry and Topology of Manifolds and Quantum Field Theory, Park City, USA, June 22-July 20, 1991.*
- [88] D. S. Freed, *Higher algebraic structures and quantization*, Commun. Math. Phys. **159** (1994), 343-398 doi:10.1007/BF02102643 [arXiv:hep-th/9212115 [hep-th]].

- [89] D. S. Freed, *Extended structures in topological quantum field theory*, [arXiv:hep-th/9306045 [hep-th]].
- [90] D. S. Freed, *Characteristic numbers and generalized path integrals*,
- [91] D. S. Freed, *Locality and integration in topological field theory*, [arXiv:hep-th/9209048 [hep-th]].
- [92] D. S. Freed, *Classical Chern-Simons theory. Part 1*, Adv. Math. **113** (1995), 237-303 doi:10.1006/aima.1995.1039 [arXiv:hep-th/9206021 [hep-th]].
- [93] D. S. Freed, *Lectures on topological quantum field theory*, NATO Sci. Ser. C **409** (1993), 95-156
- [94] D. S. Freed and F. Quinn, *Chern-Simons theory with finite gauge group*, Commun. Math. Phys. **156** (1993), 435-472 doi:10.1007/BF02096860 [arXiv:hep-th/9111004 [hep-th]].
- [95] D. S. Freed and K. K. Uhlenbeck, *INSTANTONS AND FOUR - MANIFOLDS*
- [96] D. S. Freed, G. W. Moore and G. Segal, *The Uncertainty of Fluxes*, Commun. Math. Phys. **271** (2007), 247-274 doi:10.1007/s00220-006-0181-3 [arXiv:hep-th/0605198 [hep-th]].
- [97] G. Segal, *The definition of conformal field theory*,
- [98] N. J. Hitchin, G. B. Segal and R. S. Ward, *Integrable systems: Twistors, loop groups, and Riemann surfaces. Proceedings, Conference, Oxford, UK, September 1997*,
- [99] G. Segal, *Space from the point of view of loop groups*,
- [100] A. Pressley and G. Segal, *LOOP GROUPS*
- [101] G. B. Segal, *THE DEFINITION OF CONFORMAL FIELD THEORY*
- [102] D. S. Freed, G. W. Moore and C. Teleman, *Topological symmetry in quantum field theory*, [arXiv:2209.07471 [hep-th]].
- [103] D. S. Freed and C. Teleman, *Relative quantum field theory*, Commun. Math. Phys. **326** (2014), 459-476 doi:10.1007/s00220-013-1880-1 [arXiv:1212.1692 [hep-th]].
- [104] N. J. Proudfoot, *Hyperkahler analogues of Kahler quotients*, arXiv:math/0405233 [math.AG].
- [105] Weinberg S. *The Quantum Theory of Fields*. Cambridge University Press; 1995. Vol. 1
- [106] Weinberg S. *The Quantum Theory of Fields*. Cambridge University Press; 1996. Vol. 2
- [107] Weinberg S. *The Quantum Theory of Fields*. Cambridge University Press; 2000. Vol. 3
- [108] E. Witten, *Some comments on string dynamics*, [arXiv:hep-th/9507121 [hep-th]].
- [109] E. Witten, *Conformal Field Theory In Four And Six Dimensions*, [arXiv:0712.0157 [math.RT]].
- [110] E. Witten, *Geometric Langlands From Six Dimensions*, [arXiv:0905.2720 [hep-th]].
- [111] D. Ben-Zvi, *Theory X and Geometric Representation Theory*, talks at Mathematical Aspects of Six-Dimensional Quantum Field Theories IHES 2014, notes by Qiaochu Yuan.
- [112] G. Moore, *Applications of the six-dimensional (2,0) theories to Physical Mathematics*, Felix Klein lectures Bonn (2012).

- [113] J. J. Heckman, D. R. Morrison, T. Rudelius and C. Vafa, *Atomic Classification of 6D SCFTs*, Fortsch. Phys. **63** (2015), 468-530 doi:10.1002/prop.201500024 [arXiv:1502.05405 [hep-th]].
- [114] M. Del Zotto, J. J. Heckman, D. S. Park and T. Rudelius, *On the Defect Group of a 6D SCFT*, Lett. Math. Phys. **106** (2016) no.6, 765-786 doi:10.1007/s11005-016-0839-5 [arXiv:1503.04806 [hep-th]].
- [115] J. J. Heckman and T. Rudelius, *Top Down Approach to 6D SCFTs*, J. Phys. A **52** (2019) no.9, 093001 doi:10.1088/1751-8121/aafc81 [arXiv:1805.06467 [hep-th]].
- [116] P. C. Argyres, J. J. Heckman, K. Intriligator and M. Martone, *Snowmass White Paper on SCFTs*, [arXiv:2202.07683 [hep-th]].
- [117] J. J. Heckman, M. Hübner, E. Torres and H. Y. Zhang, *The Branes Behind Generalized Symmetry Operators*, Fortsch. Phys. **71** (2023) no.1, 2200180 doi:10.1002/prop.202200180 [arXiv:2209.03343 [hep-th]].
- [118] J. J. Heckman, M. Hubner, E. Torres, X. Yu and H. Y. Zhang, *Top down approach to topological duality defects*, Phys. Rev. D **108** (2023) no.4, 046015 doi:10.1103/PhysRevD.108.046015 [arXiv:2212.09743 [hep-th]].
- [119] F. Baume, J. J. Heckman, M. Hübner, E. Torres, A. P. Turner and X. Yu, *SymTrees and Multi-Sector QFTs*, Phys. Rev. D **109** (2024) no.10, 106013 doi:10.1103/PhysRevD.109.106013 [arXiv:2310.12980 [hep-th]].
- [120] M. Cvetič, J. J. Heckman, M. Hübner and E. Torres, *0-form, 1-form, and 2-group symmetries via cutting and gluing of orbifolds*, Phys. Rev. D **106** (2022) no.10, 106003 doi:10.1103/PhysRevD.106.106003 [arXiv:2203.10102 [hep-th]].
- [121] M. Cvetič, J. J. Heckman, M. Hübner and E. Torres, *Fluxbranes, generalized symmetries, and Verlinde's metastable monopole*, Phys. Rev. D **109** (2024) no.4, 046007 doi:10.1103/PhysRevD.109.046007 [arXiv:2305.09665 [hep-th]].
- [122] N. Seiberg and E. Witten, *Gauge dynamics and compactification to three-dimensions*, [arXiv:hep-th/9607163 [hep-th]].
- [123] D. Gaiotto and E. Witten, *S-Duality of Boundary Conditions In N=4 Super Yang-Mills Theory*, Adv. Theor. Math. Phys. **13** (2009) no.3, 721-896 doi:10.4310/ATMP.2009.v13.n3.a5 [arXiv:0807.3720 [hep-th]].
- [124] D. Gaiotto and J. Maldacena, *The Gravity duals of N=2 superconformal field theories*, JHEP **10** (2012), 189 doi:10.1007/JHEP10(2012)189 [arXiv:0904.4466 [hep-th]].
- [125] D. Gaiotto, A. Neitzke and Y. Tachikawa, *Argyres-Seiberg duality and the Higgs branch*, Commun. Math. Phys. **294** (2010), 389-410 doi:10.1007/s00220-009-0938-6 [arXiv:0810.4541 [hep-th]].
- [126] D. Gaiotto and E. Witten, *Supersymmetric Boundary Conditions in N=4 Super Yang-Mills Theory*, J. Statist. Phys. **135** (2009), 789-855 doi:10.1007/s10955-009-9687-3 [arXiv:0804.2902 [hep-th]].
- [127] D. Gaiotto and E. Witten, *Janus Configurations, Chern-Simons Couplings, And The theta-Angle in N=4 Super Yang-Mills Theory*, JHEP **06** (2010), 097 doi:10.1007/JHEP06(2010)097 [arXiv:0804.2907 [hep-th]].
- [128] D. S. Freed and A. Neitzke, *3d spectral networks and classical Chern-Simons theory*, [arXiv:2208.07420 [math.DG]].

- [129] A. Neitzke, *Hitchin Systems in $\mathcal{N} = 2$ Field Theory*, doi:10.1007/978-3-319-18769-3_3 [arXiv:1412.7120 [hep-th]].
- [130] D. Gaiotto, G. W. Moore and A. Neitzke, *Wall-Crossing in Coupled 2d-4d Systems*, JHEP **12** (2012), 082 doi:10.1007/JHEP12(2012)082 [arXiv:1103.2598 [hep-th]].
- [131] D. Gaiotto, G. W. Moore and A. Neitzke, *Four-dimensional wall-crossing via three-dimensional field theory*, Commun. Math. Phys. **299** (2010), 163-224 doi:10.1007/s00220-010-1071-2 [arXiv:0807.4723 [hep-th]].
- [132] D. Gaiotto, G. W. Moore and A. Neitzke, *Four-dimensional wall-crossing via three-dimensional field theory*, Commun. Math. Phys. **299** (2010), 163-224 doi:10.1007/s00220-010-1071-2 [arXiv:0807.4723 [hep-th]].
- [133] D. Gaiotto, A. Neitzke and Y. Tachikawa, *Argyres-Seiberg duality and the Higgs branch*, Commun. Math. Phys. **294** (2010), 389-410 doi:10.1007/s00220-009-0938-6 [arXiv:0810.4541 [hep-th]].
- [134] Y. Tachikawa, *On the 6d origin of discrete additional data of 4d gauge theories*, JHEP **05** (2014), 020 doi:10.1007/JHEP05(2014)020 [arXiv:1309.0697 [hep-th]].
- [135] L. F. Alday, D. Gaiotto and Y. Tachikawa, *Liouville Correlation Functions from Four-dimensional Gauge Theories*, Lett. Math. Phys. **91** (2010), 167-197 doi:10.1007/s11005-010-0369-5 [arXiv:0906.3219 [hep-th]].
- [136] G. W. Moore and Y. Tachikawa, *On 2d TQFTs whose values are holomorphic symplectic varieties*, Proc. Symp. Pure Math. **85** (2012), 191-208 doi:10.1090/pspum/085/1379 [arXiv:1106.5698 [hep-th]].
- [137] G. W. Moore and G. Segal, *D-branes and K-theory in 2D topological field theory*, [arXiv:hep-th/0609042 [hep-th]].
- [138] Kirwan, F. C. (1984). *Cohomology of Quotients in Symplectic and Algebraic Geometry*. (MN-31), Volume 31 (Vol. 104). Princeton University Press. <https://doi.org/10.2307/j.ctv10vm2m8>
- [139] H. Nakajima, *A mathematical definition of Coulomb branches of supersymmetric gauge theories and geometric Satake correspondences for Kac-Moody Lie algebras*, [arXiv:2201.08386 [math.RT]].
- [140] C. Teleman, *Coulomb branches for quaternionic representations*, [arXiv:2209.01088 [math.AT]].
- [141] A. Braverman, M. Finkelberg and H. Nakajima, *Towards a mathematical definition of Coulomb branches of 3-dimensional $\mathcal{N} = 4$ gauge theories, II*, Adv. Theor. Math. Phys. **22** (2018), 1071-1147 doi:10.4310/ATMP.2018.v22.n5.a1 [arXiv:1601.03586 [math.RT]].
- [142] C. Teleman, *The rôle of Coulomb branches in 2D gauge theory*, J. Eur. Math. Soc. **23** (2021) no.11, 3497-3520 doi:10.4171/jems/1071 [arXiv:1801.10124 [math.AG]].
- [143] C. Teleman. *The Quantization Conjecture Revisited.*, Annals of Mathematics 152, no. 1 (2000): 1–43. <https://doi.org/10.2307/2661378>.
- [144] A. Braverman, M. Finkelberg and H. Nakajima, *Coulomb branches of 3d $\mathcal{N} = 4$ quiver gauge theories and slices in the affine Grassmannian*, Adv. Theor. Math. Phys. **23** (2019), 75-166 doi:10.4310/ATMP.2019.v23.n1.a3 [arXiv:1604.03625 [math.RT]].

- [145] A. Braverman, M. Finkelberg and H. Nakajima, *Ring objects in the equivariant derived Satake category arising from Coulomb branches (with an appendix by Gus Lonergan)*, [arXiv:1706.02112 [math.RT]].
- [146] K. A. Intriligator and N. Seiberg, *Mirror symmetry in three-dimensional gauge theories*, Phys. Lett. B **387** (1996), 513–519 doi:10.1016/0370-2693(96)01088-X [arXiv:hep-th/9607207 [hep-th]].
- [147] M. Kontsevich, *Homological Algebra of Mirror Symmetry*, [arXiv:alg-geom/9411018 [math.AG]].
- [148] P. S. Aspinwall and D. R. Morrison, Topological field theory and rational curves, Comm. Math. Phys. 151 (1993), 245–262.
- [149] M. Bershadsky, S. Cecotti, H. Ooguri, and C. Vafa, Kodaira-Spencer theory of gravity and exact results for quantum string amplitudes, Comm. Math. Phys. 165 (1994), 311–427.
- [150] A. I. Bondal and M. M. Kapranov, Enhanced triangulated categories, Math. USSR-Sb. 70 (1991), 93–107.
- [151] P. Candelas, X. de la Ossa, P. S. Green, and L. Parkes, A pair of Calabi- Yau manifolds as an exactly soluble superconformal theory, in [Y], pp. 31–95.
- [152] B. Dubrovin, Integrable systems in topological field theory, Nuclear Phys. B 379 (1992), 627–685.
- [153] K. Fukaya, Morse homotopy, A[∞]-category and Floer homologies, MSRI pre-print No. 020–94 (1993).
- [154] M. Gerstenhaber and S. D. Schack, Algebraic cohomology and deformation theory, NATO ASI Ser. C (1988), 11–264.
- [155] M. Gromov, Pseudo holomorphic curves in symplectic manifolds, Invent. Math. 82 (1985), 307–347.
- [156] M. Kontsevich, Formal (non)-commutative symplectic geometry, The Gelfand Mathematical Seminars, 1990–1992 (L. Corwin, I. Gelfand, and J. Lepowsky, eds.), Birkhäuser, Basel and Boston, 1993, pp. 173–187.
- [157] M. Kontsevich, Enumeration of rational curves via torus actions, MPI preprint and hep-th/9405035 (1994).
- [158] M. Kontsevich and Yu. Manin, Gromov-Witten classes, quantum cohomology and enumerative geometry, Comm. Math. Phys. 164 (1994), 525–562.
- [159] D. R. Morrison, Mirror symmetry and rational curves on quintic threefolds: A guide for mathematicians, J. Amer. Math. Soc. 6 (1993), 223–247.
- [160] Y.-G. Oh, Floer Cohomology of Lagrangian intersections and pseudo-holomorphic discs I, II, Comm. Pure Appl. Math. 46:7 (1993), 949–1012.
- [161] Y. Ruan and G. Tian, Mathematical theory of quantum cohomology, Math. Res. Lett. 1 (1994), 269–278.
- [162] J. Stasheff, On the homotopy associativity of H-spaces I, II, Trans. Amer. Math. Soc. 108 (1963), 275–312.
- [163] J. L. Verdier, Categories dérivées. Quelques résultats (état 0), Séminaire de Geometrie Algébrique du Bois-Marie (SGA 42), Lecture Notes in Math., vol. 569, Springer-Verlag, Berlin and New York. 1977. nn. 262–311.

- [164] E. Witten, Topological sigma models. *Comm Math. Phys.* **118** (1988) 411–449
- [165] E. Witten, Two-dimensional gravity and intersection theory on moduli space, *Surveys Differential Geom.* **1** (1991). 243–310.
- [166] E. Witten, Mirror manifolds and topological field theory. in [Y], pp. 120–159.
- [167] E. Witten, Chern-Simons gauge theory as a string theory, preprint IASSNSHEP-92/45 and hep-th/9207094.
- [168] S. T. Yau, ed., *Essays on Mirror Manifolds*, International Press Co., Hong Kong, 1992.
- [169] L. Katzarkov, M. Kontsevich and T. Pantev, *Hodge theoretic aspects of mirror symmetry*, *Proc. Symp. Pure Math.* **78** (2008), 87-174 doi:10.1090/pspum/078/2483750 [arXiv:0806.0107 [math.AG]].
- [170] M. Atiyah and M. Marcolli, *Anyon Networks from Geometric Models of Matter*, *Quart. J. Math. Oxford Ser.* **72** (2021) no.1-2, 717-733 doi:10.1093/qmath/haab004
- [171] M. F. Atiyah and C. Zapata-Carratala, *K-Theory and the Jones Polynomial*, *Rev. Math. Phys.* **30** (2018) no.06, 1840002 doi:10.1142/S0129055X18400020
- [172] *Anyons in Geometric Models of Matter*, *JHEP* **07** (2017), 076 doi:10.1007/JHEP07(2017)076 [arXiv:1611.04047 [math-ph]].
- [173] M. F. Atiyah and G. W. Moore, *A Shifted View of Fundamental Physics*, [arXiv:1009.3176 [hep-th]].
- [174] M. Atiyah, *The Interaction between Geometry and Physics*, *Prog. Math.* **244** (2007), 1-15 doi:10.1007/0-8176-4467-9_1
- [175] M. Atiyah and G. Segal, *Twisted K-theory*, [arXiv:math/0407054 [math.KT]].
- [176] G. Segal, *Michael Atiyah's work in algebraic topology*, *Bull. Am. Math. Soc.* **58** (2021) no.4, 481-516 doi:10.1090/bull/1746
- [177] G. Segal, *Locality of Holomorphic Bundles, and Locality in Quantum Field Theory*, doi:10.1093/acprof:oso/9780199534920.003.0009
- [178] B. C. Allanach, B. Gripaios and J. Tooby-Smith, *Anomaly cancellation with an extra gauge boson*, *Phys. Rev. Lett.* **125** (2020) no.16, 161601 doi:10.1103/PhysRevLett.125.161601 [arXiv:2006.03588 [hep-th]].
- [179] B. C. Allanach, B. Gripaios and J. Tooby-Smith, *Geometric General Solution to the U(1) Anomaly Equations*, *JHEP* **05** (2020), 065 doi:10.1007/JHEP05(2020)065 [arXiv:1912.04804 [hep-th]].
- [180] J. Davighi, N. Lohitsiri and A. Debray, *Toric 2-group anomalies via cobordism*, *JHEP* **07** (2023), 019 doi:10.1007/JHEP07(2023)019 [arXiv:2302.12853 [hep-th]].
- [181] J. Davighi, B. Gripaios and N. Lohitsiri, *Anomalies of non-Abelian finite groups via cobordism*, *JHEP* **09** (2022), 147 doi:10.1007/JHEP09(2022)147 [arXiv:2207.10700 [hep-th]].
- [182] J. Davighi, B. Gripaios and O. Randal-Williams, *Differential cohomology and topological actions in physics*, [arXiv:2011.05768 [hep-th]].
- [183] J. Davighi, *Topological effects in particle physics phenomenology*, doi:10.17863/CAM.47560
- [184] B. Gripaios, O. Randal-Williams and J. Tooby-Smith, *Smooth generalized symmetries of quantum field theories*, [arXiv:2310.16090 [hep-th]].

- [185] B. Gripaios, O. Randal-Williams and J. Tooby-Smith, *Generalized symmetries of topological field theories*, JHEP **03** (2023), 087 doi:10.1007/JHEP03(2023)087 [arXiv:2209.13524 [hep-th]].
- [186] B. Gripaios, *Gauge anomalies of finite groups*, Phys. Rev. D **105** (2022) no.10, 105008 doi:10.1103/PhysRevD.105.105008 [arXiv:2201.11801 [hep-th]].
- [187] B. C. Allanach, B. Gripaios and J. Tooby-Smith, *Semisimple extensions of the Standard Model gauge algebra*, Phys. Rev. D **104** (2021) no.3, 035035 [erratum: Phys. Rev. D **106** (2022) no.1, 019901] doi:10.1103/PhysRevD.104.035035 [arXiv:2104.14555 [hep-th]].
- [188] J. Davighi and B. Gripaios, *Homological classification of topological terms in sigma models on homogeneous spaces*, JHEP **09** (2018), 155 [erratum: JHEP **11** (2018), 143] doi:10.1007/JHEP09(2018)155 [arXiv:1803.07585 [hep-th]].
- [189] B. Gripaios and O. Randal-Williams, *Topology of electroweak vacua*, Phys. Lett. B **782** (2018), 94-98 doi:10.1016/j.physletb.2018.05.013 [arXiv:1610.05623 [hep-th]].
- [190] B. Gripaios and D. Sutherland, *An operator basis for the Standard Model with an added scalar singlet*, JHEP **08** (2016), 103 doi:10.1007/JHEP08(2016)103 [arXiv:1604.07365 [hep-ph]].
- [191] D. Zagier, Note on the Landweber-Stong elliptic genus
- [192] C. Teleman, *Gauge theory and mirror symmetry*, [arXiv:1404.6305 [math-ph]].
- [193] D. Pomerleano and C. Teleman, to appear.
- [194] R. Melrose, Introduction to microlocal analysis, 2003.
- [195] A. Bourget, S. Cabrera, J. F. Grimminger, A. Hanany, M. Sperling, A. Zajac and Z. Zhong, *The Higgs mechanism — Hasse diagrams for symplectic singularities*, JHEP **01** (2020), 157 doi:10.1007/JHEP01(2020)157 [arXiv:1908.04245 [hep-th]].
- [196] A. Bourget, J. F. Grimminger, A. Hanany, M. Sperling and Z. Zhong, *Branes, Quivers, and the Affine Grassmannian*, Adv. Stud. Pure Math. **88** (2023), 331-435 doi:10.2969/aspm/08810331 [arXiv:2102.06190 [hep-th]].
- [197] A. Bourget, J. F. Grimminger, A. Hanany, R. Kalveks, M. Sperling and Z. Zhong, *A Tale of N Cones*, [arXiv:2303.16939 [hep-th]].
- [198] A. Bourget, S. Cabrera, J. F. Grimminger, A. Hanany and Z. Zhong, *Brane Webs and Magnetic Quivers for SQCD*, JHEP **03** (2020), 176 doi:10.1007/JHEP03(2020)176 [arXiv:1909.00667 [hep-th]].
- [199] A. Hanany and R. Kalveks, *Quiver Theories and Hilbert Series of Classical Slodowy Intersections*, Nucl. Phys. B **952** (2020), 114939 doi:10.1016/j.nuclphysb.2020.114939 [arXiv:1909.12793 [hep-th]].
- [200] S. Cabrera and A. Hanany, *Quiver Subtractions*, JHEP **09** (2018), 008 doi:10.1007/JHEP09(2018)008 [arXiv:1803.11205 [hep-th]].
- [201] D. Bigatti and L. Susskind, *TASI lectures on the holographic principle*, doi:10.1142/9789812799630_0012 [arXiv:hep-th/0002044 [hep-th]].
- [202] Weinberg S. Lectures on Quantum Mechanics. Cambridge University Press; 2012.
- [203] J. M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, Adv. Theor. Math. Phys. **2** (1998), 231-252 doi:10.1023/A:1026654312961 [arXiv:hep-th/9711200 [hep-th]].

- [204] D. Tong, *What is Quantum Field Theory and Why is it Incomplete?*,
<https://www.quantamagazine.org/what-is-quantum-field-theory-and-why-is-it-incomplete-20220810/>
- [205] B. C. Allanach, *Supersymmetry and Particle Physics*,
<https://www.youtube.com/watch?v=Yu1U1K3UpTk>
- [206] T. Azeyanagi, A. Karch, T. Takayanagi and E. G. Thompson, *Holographic calculation of boundary entropy*, JHEP **03** (2008), 054 doi:10.1088/1126-6708/2008/03/054 [arXiv:0712.1850 [hep-th]].
- [207] J. D. Brown and M. Henneaux, *Central Charges in the Canonical Realization of Asymptotic Symmetries: An Example from Three-Dimensional Gravity*, Commun. Math. Phys. **104** (1986), 207-226 doi:10.1007/BF01211590
- [208] S. R. Coleman and F. De Luccia, *Gravitational Effects on and of Vacuum Decay*, Phys. Rev. D **21** (1980), 3305 doi:10.1103/PhysRevD.21.3305
- [209] J. D. Brown and C. Teitelboim, *Neutralization of the Cosmological Constant by Membrane Creation*, Nucl. Phys. B **297** (1988), 787-836 doi:10.1016/0550-3213(88)90559-7
- [210] W. Fischler, D. Morgan and J. Polchinski, *Quantization of False Vacuum Bubbles: A Hamiltonian Treatment of Gravitational Tunneling*, Phys. Rev. D **42** (1990), 4042-4055 doi:10.1103/PhysRevD.42.4042
- [211] A. Almheiri, T. Hartman, J. Maldacena, E. Shaghoulian and A. Tajdini, *The entropy of Hawking radiation*, Rev. Mod. Phys. **93** (2021) no.3, 035002 doi:10.1103/RevModPhys.93.035002 [arXiv:2006.06872 [hep-th]].
- [212] S. W. Hawking, *Particle Creation by Black Holes*, Commun. Math. Phys. **43** (1975), 199-220 [erratum: Commun. Math. Phys. **46** (1976), 206] doi:10.1007/BF02345020
- [213] T. Banks, L. Susskind and M. E. Peskin, *Difficulties for the Evolution of Pure States Into Mixed States*, Nucl. Phys. B **244** (1984), 125-134 doi:10.1016/0550-3213(84)90184-6
- [214] S. B. Giddings, *The Black hole information paradox*, [arXiv:hep-th/9508151 [hep-th]].
- [215] E. Witten, *Gravity and the crossed product*, JHEP **10** (2022), 008 doi:10.1007/JHEP10(2022)008 [arXiv:2112.12828 [hep-th]].
- [216] V. Chandrasekaran, R. Longo, G. Penington and E. Witten, *An Algebra of Observables for de Sitter Space*, [arXiv:2206.10780 [hep-th]].
- [217] E. Witten, *Why Does Quantum Field Theory In Curved Spacetime Make Sense? And What Happens To The Algebra of Observables In The Thermodynamic Limit?*, [arXiv:2112.11614 [hep-th]].
- [218] P. Deligne and D.S. Freed, *Quantum Fields and Strings: A Course for Mathematicians*, Volume 1, American Mathematics Society, 1999.
- [219] D.S. Freed, *Classical Field Theory and Supersymmetry*, Lecture notes, 2001.
- [220] D. Ayala, J. Francis, *A factorisation homology primer*, arXiv:1903.10961 [math.AT].
- [221] Deligne, Pierre; Mumford, David. *The irreducibility of the space of curves of given genus*. Publications Mathématiques de l’IHÉS, Volume 36 (1969), pp. 75-109. http://www.numdam.org/item/PMIHES_1969_36_75_0/.

- [222] S. Cabrera, A. Hanany and M. Sperling, *Magnetic quivers, Higgs branches, and 6d $N=(1,0)$ theories*, JHEP **06** (2019), 071 [erratum: JHEP **07** (2019), 137] doi:10.1007/JHEP06(2019)071 [arXiv:1904.12293 [hep-th]].
- [223] G. Ferlito and A. Hanany, *A tale of two cones: the Higgs Branch of $Sp(n)$ theories with $2n$ flavours*, [arXiv:1609.06724 [hep-th]].
- [224] J. Maldacena, *Vacuum decay into Anti de Sitter space*, [arXiv:1012.0274 [hep-th]].
- [225] S. Ryu and T. Takayanagi, *Holographic derivation of entanglement entropy from AdS/CFT*, Phys. Rev. Lett. **96** (2006), 181602 doi:10.1103/PhysRevLett.96.181602 [arXiv:hep-th/0603001 [hep-th]].
- [226] V. E. Hubeny, M. Rangamani and T. Takayanagi, *A Covariant holographic entanglement entropy proposal*, JHEP **07** (2007), 062 doi:10.1088/1126-6708/2007/07/062 [arXiv:0705.0016 [hep-th]].
- [227] M. Van Raamsdonk, *Lectures on Gravity and Entanglement*, doi:10.1142/9789813149441_0005 [arXiv:1609.00026 [hep-th]].
- [228] M. Van Raamsdonk, *Spacetime from bits*, Science **370** (2020) no.6513, 198-202 doi:10.1126/science.aay9560
- [229] J. B. Hartle and S. W. Hawking, *Wave Function of the Universe*, Phys. Rev. D **28** (1983), 2960-2975 doi:10.1103/PhysRevD.28.2960
- [230] T. C. Bachlechner, *Inflation Expels Runaways*, JHEP **12** (2016), 155 doi:10.1007/JHEP12(2016)155 [arXiv:1608.07576 [hep-th]].
- [231] S. Céspedes, S. P. de Alwis, F. Muia, F. Quevedo, *Lorentzian vacuum transitions: open or closed?*, Phys. Rev. D **104**, 026013 (2021), [arXiv:2011.13936[hep-th]]
- [232] B. Banihashemi, T. Jacobson, A. Svesko and M. Visser, *The minus sign in the first law of de Sitter horizons*, [arXiv:2208.11706 [hep-th]].
- [233] K. Lee, E. J. Weinberg, *Decay of the True Vacuum in Curved Spacetime*, 10.113/PhysRev.D.36.1088
- [234] E. Farhi, A. H. Guth and J. Guven, *Is It Possible to Create a Universe in the Laboratory by Quantum Tunneling?*, Nucl. Phys. B **339** (1990), 417-490 doi:10.1016/0550-3213(90)90357-J
- [235] L. Susskind, *Three Impossible Theories*, [arXiv:2107.11688 [hep-th]].
- [236] M. Fujita, T. Takayanagi and E. Tonni, *Aspects of AdS/BCFT*, JHEP **11** (2011), 043 doi:10.1007/JHEP11(2011)043 [arXiv:1108.5152 [hep-th]].
- [237] I. Akal, Y. Kusuki, T. Takayanagi and Z. Wei, *Codimension two holography for wedges*, Phys. Rev. D **102** (2020) no.12, 126007 doi:10.1103/PhysRevD.102.126007 [arXiv:2007.06800 [hep-th]].
- [238] A. B. Zamolodchikov, *Renormalization Group and Perturbation Theory Near Fixed Points in Two-Dimensional Field Theory*, Sov. J. Nucl. Phys. **46** (1987), 1090
- [239] A. B. Zamolodchikov, *Expectation value of composite field T anti- T in two-dimensional quantum field theory*, [arXiv:hep-th/0401146 [hep-th]].
- [240] L. McGough, M. Mezei and H. Verlinde, *Moving the CFT into the bulk with $T\bar{T}$* , JHEP **04** (2018), 010 doi:10.1007/JHEP04(2018)010 [arXiv:1611.03470 [hep-th]].

- [241] V. Gorbenko, E. Silverstein and G. Torroba, *dS/dS and $T\bar{T}$* , JHEP **03** (2019), 085 doi:10.1007/JHEP03(2019)085 [arXiv:1811.07965 [hep-th]].
- [242] A. Lewkowycz, J. Liu, E. Silverstein and G. Torroba, *$T\bar{T}$ and EE, with implications for (A)dS subregion encodings*, JHEP **04** (2020), 152 doi:10.1007/JHEP04(2020)152 [arXiv:1909.13808 [hep-th]].
- [243] A. Strominger, *The dS / CFT correspondence*, JHEP **10** (2001), 034 doi:10.1088/1126-6708/2001/10/034 [arXiv:hep-th/0106113 [hep-th]].
- [244] M. Alishahiha, A. Karch, E. Silverstein and D. Tong, *The dS/dS correspondence*, AIP Conf. Proc. **743** (2004) no.1, 393-409 doi:10.1063/1.1848341 [arXiv:hep-th/0407125 [hep-th]].
- [245] A. Almheiri, R. Mahajan, J. Maldacena and Y. Zhao, *The Page curve of Hawking radiation from semiclassical geometry*, JHEP **03** (2020), 149 doi:10.1007/JHEP03(2020)149 [arXiv:1908.10996 [hep-th]].
- [246] G. Penington, *Entanglement Wedge Reconstruction and the Information Paradox*, JHEP **09** (2020), 002 doi:10.1007/JHEP09(2020)002 [arXiv:1905.08255 [hep-th]].
- [247] A. Almheiri, R. Mahajan and J. E. Santos, *Entanglement islands in higher dimensions*; SciPost Phys. **9** (2020) no.1, 001 doi:10.21468/SciPostPhys.9.1.001 [arXiv:1911.09666 [hep-th]].
- [248] S. Antonini, P. Simidzija, B. Swingle and M. Van Raamsdonk, *Cosmology as a holographic wormhole*, [arXiv:2206.14821 [hep-th]].
- [249] S. Antonini, P. Simidzija, B. Swingle and M. Van Raamsdonk, *Cosmology from the vacuum*, [arXiv:2203.11220 [hep-th]].
- [250] M. Van Raamsdonk, *Comments on wormholes, ensembles, and cosmology*, JHEP **12** (2021), 156 doi:10.1007/JHEP12(2021)156 [arXiv:2008.02259 [hep-th]].
- [251] M. Van Raamsdonk, *Cosmology from confinement?*, JHEP **03** (2022), 039 doi:10.1007/JHEP03(2022)039 [arXiv:2102.05057 [hep-th]].
- [252] T. Hartman, Y. Jiang and E. Shaghoulian, *Islands in cosmology*, JHEP **11** (2020), 111 doi:10.1007/JHEP11(2020)111 [arXiv:2008.01022 [hep-th]].
- [253] S. E. Aguilar-Gutierrez, A. Chatwin-Davies, T. Hertog, N. Pinzani-Fokeeva and B. Robinson, *Islands in Multiverse Models*, JHEP **11** (2021), 212 doi:10.1007/JHEP11(2021)212 [arXiv:2108.01278 [hep-th]].
- [254] H. Geng, S. Lüst, R. K. Mishra and D. Wakeham, *Holographic BCFTs and Communicating Black Holes*, jhep **08** (2021), 003 doi:10.1007/JHEP08(2021)003 [arXiv:2104.07039 [hep-th]].
- [255] H. Geng, Y. Nomura and H. Y. Sun, *Information paradox and its resolution in de Sitter holography*, Phys. Rev. D **103** (2021) no.12, 126004 doi:10.1103/PhysRevD.103.126004 [arXiv:2103.07477 [hep-th]].
- [256] Geng, H. , *Aspects of AdS2 quantum gravity and the Karch-Randall braneworld*, J. High Energ. Phys. **2022**, 24 (2022). [https://doi.org/10.1007/JHEP09\(2022\)024](https://doi.org/10.1007/JHEP09(2022)024), arXiv:2206.11277 [hep-th].
- [257] H. Geng, A. Karch, C. Perez-Pardavila, S. Raju, L. Randall, M. Riojas and S. Shashi, *Jackiw-Teitelboim Gravity from the Karch-Randall Braneworld*, Phys. Rev. Lett. **129** (2022) no.23, 231601 doi:10.1103/PhysRevLett.129.231601 [arXiv:2206.04695 [hep-th]].

- [258] K. Langhoff, C. Murdia and Y. Nomura, *Multiverse in an inverted island*, Phys. Rev. D **104** (2021) no.8, 086007 doi:10.1103/PhysRevD.104.086007 [arXiv:2106.05271 [hep-th]].
- [259] P. Betzios, E. Kiritsis and O. Papadoulaki, *Euclidean Wormholes and Holography*, JHEP **06** (2019), 042 doi:10.1007/JHEP06(2019)042 [arXiv:1903.05658 [hep-th]].
- [260] P. Betzios, E. Kiritsis and O. Papadoulaki, *Interacting systems and wormholes*, JHEP **02** (2022), 126 doi:10.1007/JHEP02(2022)126 [arXiv:2110.14655 [hep-th]].
- [261] R. Bousso and E. Wildenhain, *Islands in closed and open universes*, Phys. Rev. D **105** (2022) no.8, 086012 doi:10.1103/PhysRevD.105.086012 [arXiv:2202.05278 [hep-th]].
- [262] D. Anninos and D. A. Galante, *Constructing AdS_2 flow geometries*, JHEP **02** (2021), 045 doi:10.1007/JHEP02(2021)045 [arXiv:2011.01944 [hep-th]].
- [263] D. Anninos and E. Harris, *Interpolating geometries and the stretched dS_2 horizon*, [arXiv:2209.06144 [hep-th]].
- [264] Y. Chen, V. Gorbenko and J. Maldacena, *Bra-ket wormholes in gravitationally prepared states*, JHEP **02** (2021), 009 doi:10.1007/JHEP02(2021)009 [arXiv:2007.16091 [hep-th]].
- [265] B. Freivogel, V. E. Hubeny, A. Maloney, R. C. Myers, M. Rangamani and S. Shenker, *Inflation in AdS/CFT* , JHEP **03** (2006), 007 doi:10.1088/1126-6708/2006/03/007 [arXiv:hep-th/0510046 [hep-th]].
- [266] V. E. Hubeny, H. Maxfield, M. Rangamani and E. Tonni, *Holographic entanglement plateaux*, JHEP **08** (2013), 092 doi:10.1007/JHEP08(2013)092 [arXiv:1306.4004 [hep-th]].
- [267] Z. Fu and D. Marolf, *Bag-of-gold spacetimes, Euclidean wormholes, and inflation from domain walls in AdS/CFT* , JHEP **11** (2019), 040 doi:10.1007/JHEP11(2019)040 [arXiv:1909.02505 [hep-th]].
- [268] A. Almheiri and J. Polchinski, *Models of AdS_2 backreaction and holography*, JHEP **11** (2015), 014 doi:10.1007/JHEP11(2015)014 [arXiv:1402.6334 [hep-th]].
- [269] A. Karch and L. Randall, *Locally localized gravity*, JHEP **05** (2001), 008 doi:10.1088/1126-6708/2001/05/008 [arXiv:hep-th/0011156 [hep-th]].
- [270] L. Randall and R. Sundrum, *An Alternative to compactification*, Phys. Rev. Lett. **83** (1999), 4690-4693 doi:10.1103/PhysRevLett.83.4690 [arXiv:hep-th/9906064 [hep-th]].
- [271] R. Jackiw, *Lower Dimensional Gravity*, Nucl. Phys. B **252** (1985), 343-356 doi:10.1016/0550-3213(85)90448-1
- [272] C. Teitelboim, *Gravitation and Hamiltonian Structure in Two Space-Time Dimensions*, Phys. Lett. B **126** (1983), 41-45 doi:10.1016/0370-2693(83)90012-6
- [273] J. Maldacena, G. J. Turiaci and Z. Yang, *Two dimensional Nearly de Sitter gravity*, JHEP **01** (2021), 139 doi:10.1007/JHEP01(2021)139 [arXiv:1904.01911 [hep-th]].
- [274] P. Simidzija and M. Van Raamsdonk, *Holo-ween*, JHEP **12** (2020), 028 doi:10.1007/JHEP12(2020)028 [arXiv:2006.13943 [hep-th]].
- [275] M. Mirbabayi, *Uptunneling to de Sitter*, JHEP **09** (2020), 070 doi:10.1007/JHEP09(2020)070 [arXiv:2003.05460 [hep-th]].
- [276] L. Susskind, *De Sitter Space, Double-Scaled SYK, and the Separation of Scales in the Semiclassical Limit*, [arXiv:2209.09999 [hep-th]].

- [277] J. M. Schlenker and E. Witten, *No ensemble averaging below the black hole threshold*, JHEP **07** (2022), 143 doi:10.1007/JHEP07(2022)143 [arXiv:2202.01372 [hep-th]].
- [278] J. S. Schwinger, *On gauge invariance and vacuum polarization*, Phys. Rev. **82** (1951), 664-679 doi:10.1103/PhysRev.82.664
- [279] M. Banados, C. Teitelboim and J. Zanelli, *The Black hole in three-dimensional space-time*, Phys. Rev. Lett. **69** (1992), 1849-1851 doi:10.1103/PhysRevLett.69.1849 [arXiv:hep-th/9204099 [hep-th]].
- [280] D. Marolf and J. E. Santos, *Stability of the microcanonical ensemble in Euclidean Quantum Gravity*, [arXiv:2202.12360 [hep-th]].
- [281] D. Marolf and J. E. Santos, *The canonical ensemble reloaded: the complex-stability of Euclidean quantum gravity for black holes in a box*, JHEP **08** (2022), 215 doi:10.1007/JHEP08(2022)215 [arXiv:2202.11786 [hep-th]].
- [282] H. Geng, A. Karch, C. Perez-Pardavila, S. Raju, L. Randall, M. Riojas and S. Shashi, *Information Transfer with a Gravitating Bath*, SciPost Phys. **10** (2021) no.5, 103 doi:10.21468/SciPostPhys.10.5.103 [arXiv:2012.04671 [hep-th]].
- [283] H. Geng, A. Karch, C. Perez-Pardavila, S. Raju, L. Randall, M. Riojas and S. Shashi, *Inconsistency of islands in theories with long-range gravity*, JHEP **01** (2022), 182 doi:10.1007/JHEP01(2022)182 [arXiv:2107.03390 [hep-th]].
- [284] G. W. Gibbons and S. W. Hawking, *Action Integrals and Partition Functions in Quantum Gravity*, Phys. Rev. D **15** (1977), 2752-2756 doi:10.1103/PhysRevD.15.2752
- [285] C. Krishnan and E. di Napoli, *Can quantum de Sitter space have finite entropy?*, Class. Quant. Grav. **24** (2007), 3457-3463 doi:10.1088/0264-9381/24/13/019 [arXiv:hep-th/0602002 [hep-th]].
- [286] J. Chu, F. Deng and Y. Zhou, *Page curve from defect extremal surface and island in higher dimensions*, JHEP **10** (2021), 149 doi:10.1007/JHEP10(2021)149 [arXiv:2105.09106 [hep-th]].
- [287] F. Deng, J. Chu and Y. Zhou, *Defect extremal surface as the holographic counterpart of Island formula*, JHEP **03** (2021), 008 doi:10.1007/JHEP03(2021)008 [arXiv:2012.07612 [hep-th]].
- [288] P. Calabrese and J. L. Cardy, *Entanglement entropy and quantum field theory*, J. Stat. Mech. **0406** (2004), P06002 doi:10.1088/1742-5468/2004/06/P06002 [arXiv:hep-th/0405152 [hep-th]].
- [289] H. Casini, M. Huerta and R. C. Myers, *Towards a derivation of holographic entanglement entropy*, JHEP **05** (2011), 036 doi:10.1007/JHEP05(2011)036 [arXiv:1102.0440 [hep-th]].
- [290] M. V. Raamsdonk and C. Waddell, *Holographic and localization calculations of boundary F for $\mathcal{N} = 4$ SUSY Yang-Mills theory*, JHEP **02** (2021), 222 doi:10.1007/JHEP02(2021)222 [arXiv:2010.14520 [hep-th]].
- [291] M. Rozali, J. Sully, M. Van Raamsdonk, C. Waddell and D. Wakeham, *Information radiation in BCFT models of black holes*, JHEP **05** (2020), 004 doi:10.1007/JHEP05(2020)004 [arXiv:1910.12836 [hep-th]].
- [292] E. Witten, *Anti-de Sitter space and holography*, Adv. Theor. Math. Phys. **2** (1998), 253-291 doi:10.4310/ATMP.1998.v2.n2.a2 [arXiv:hep-th/9802150 [hep-th]].

- [293] C. Park, *Holographic RG flow triggered by a classically marginal operator*, Phys. Rev. D **105** (2022) no.4, 046004 doi:10.1103/PhysRevD.105.046004 [arXiv:2102.01829 [hep-th]].
- [294] M. J. Duff and J. T. Liu, *Complementarity of the Maldacena and Randall-Sundrum pictures*, Phys. Rev. Lett. **85** (2000), 2052-2055 doi:10.1088/0264-9381/18/16/310 [arXiv:hep-th/0003237 [hep-th]].
- [295] L. V. Iliesiu, J. Kruthoff, G. J. Turiaci and H. Verlinde, *JT gravity at finite cutoff*, SciPost Phys. **9** (2020), 023 doi:10.21468/SciPostPhys.9.2.023 [arXiv:2004.07242 [hep-th]].
- [296] M. J. Duff, J. T. Liu and H. Sati, *Complementarity of the Maldacena and Karch-Randall pictures*, Phys. Rev. D **69** (2004), 085012 doi:10.1103/PhysRevD.69.085012 [arXiv:hep-th/0207003 [hep-th]].
- [297] T. Anous, J. L. Karczmarek, E. Mintun, M. Van Raamsdonk and B. Way, *Areas and entropies in BFSS/gravity duality*, SciPost Phys. **8** (2020) no.4, 057 doi:10.21468/SciPostPhys.8.4.057 [arXiv:1911.11145 [hep-th]].
- [298] M. Van Raamsdonk and C. Waddell, *Finding $AdS^5 \times S^5$ in 2+1 dimensional SCFT physics*, JHEP **11** (2021), 145 doi:10.1007/JHEP11(2021)145 [arXiv:2109.04479 [hep-th]].
- [299] M. Alishahiha and F. Ardalan, *Central Charge for 2D Gravity on $AdS(2)$ and $AdS(2)/CFT(1)$ Correspondence*, JHEP **08** (2008), 079 doi:10.1088/1126-6708/2008/08/079 [arXiv:0805.1861 [hep-th]].
- [300] T. Hartman and A. Strominger, *Central Charge for $AdS(2)$ Quantum Gravity*, JHEP **04** (2009), 026 doi:10.1088/1126-6708/2009/04/026 [arXiv:0803.3621 [hep-th]].
- [301] E. K. Morvan, J. P. van der Schaar and M. R. Visser, *On the Euclidean Action of de Sitter Black Holes and Constrained Instantons*, [arXiv:2203.06155 [hep-th]].
- [302] S. Cooper, M. Rozali, B. Swingle, M. Van Raamsdonk, C. Waddell and D. Wakeham, *Black hole microstate cosmology*, JHEP **07** (2019), 065 doi:10.1007/JHEP07(2019)065 [arXiv:1810.10601 [hep-th]].
- [303] P. Pradhan, *Entropy Product Formula for Gravitational Instanton*, Adv. High Energy Phys. **2017** (2017), 7471640 doi:10.1155/2017/7471640 [arXiv:1609.07294 [gr-qc]].
- [304] A. Bondal, D. Orlov, *Derived categories of coherent sheaves*, arXiv:math/0206295 [math.AG].
- [305] D. Harlow and D. Jafferis, *The Factorization Problem in Jackiw-Teitelboim Gravity*, JHEP **02** (2020), 177 doi:10.1007/JHEP02(2020)177 [arXiv:1804.01081 [hep-th]].
- [306] O. J. C. Dias, P. Mitra and J. E. Santos, *New phases of $\mathcal{N} = 4$ SYM at finite chemical potential*, [arXiv:2207.07134 [hep-th]].
- [307] P. Gao, D. L. Jafferis and A. C. Wall, *Traversable Wormholes via a Double Trace Deformation* JHEP **12** (2017), 151 doi:10.1007/JHEP12(2017)151 [arXiv:1608.05687 [hep-th]].
- [308] A. C. Wall, *The Generalized Second Law implies a Quantum Singularity Theorem*, Class. Quant. Grav. **30** (2013), 165003 [erratum: Class. Quant. Grav. **30** (2013), 199501] doi:10.1088/0264-9381/30/19/199501 [arXiv:1010.5513 [gr-qc]].
- [309] O. Chacaltana, J. Distler and Y. Tachikawa, *Nilpotent orbits and codimension-two defects of 6d $N=(2,0)$ theories*, Int. J. Mod. Phys. A **28** (2013), 1340006 doi:10.1142/S0217751X1340006X [arXiv:1203.2930 [hep-th]].

- [310] Y. Tachikawa, *On the 6d origin of discrete additional data of 4d gauge theories*, JHEP **05** (2014), 020 doi:10.1007/JHEP05(2014)020 [arXiv:1309.0697 [hep-th]].
- [311] Milnor, J. W., & Moore, J. C. (1965). On the Structure of Hopf Algebras. Annals of Mathematics, 81(2), 211–264. <https://doi.org/10.2307/1970615>.
- [312] I. Bah, D. S. Freed, G. W. Moore, N. Nekrasov, S. S. Razamat and S. Schafer-Nameki, *A Panorama Of Physical Mathematics c. 2022*, [arXiv:2211.04467 [hep-th]].
- [313] G. W. Moore and N. Seiberg, *Classical and Quantum Conformal Field Theory*, Commun. Math. Phys. **123** (1989), 177 doi:10.1007/BF01238857.
- [314] G. W. Moore and N. Seiberg, *Naturality in Conformal Field Theory*, Nucl. Phys. B **313** (1989), 16-40 doi:10.1016/0550-3213(89)90511-7.
- [315] E. P. Verlinde, *Fusion Rules and Modular Transformations in 2D Conformal Field Theory*, Nucl. Phys. B **300** (1988), 360-376 doi:10.1016/0550-3213(88)90603-7.
- [316] D. Gaiotto, A. Kapustin, N. Seiberg and B. Willett, *Generalized Global Symmetries*, JHEP **02** (2015), 172 doi:10.1007/JHEP02(2015)172 [arXiv:1412.5148 [hep-th]].
- [317] L. Kong, T. Lan, X. G. Wen, Z. H. Zhang and H. Zheng, *Algebraic higher symmetry and categorical symmetry – a holographic and entanglement view of symmetry*, Phys. Rev. Res. **2** (2020) no.4, 043086 doi:10.1103/PhysRevResearch.2.043086 [arXiv:2005.14178 [cond-mat.str-el]].
- [318] L. Kong and Z. H. Zhang, *An invitation to topological orders and category theory*, [arXiv:2205.05565 [cond-mat.str-el]].
- [319] L. Kong and H. Zheng, *A mathematical theory of gapless edges of 2d topological orders. Part I*, JHEP **02** (2020), 150 doi:10.1007/JHEP02(2020)150 [arXiv:1905.04924 [cond-mat.str-el]].
- [320] L. Kong and H. Zheng, *A mathematical theory of gapless edges of 2d topological orders. Part II*, Nucl. Phys. B **966** (2021), 115384 doi:10.1016/j.nuclphysb.2021.115384 [arXiv:1912.01760 [cond-mat.str-el]].
- [321] L. Kong and H. Zheng, *Categorical computation*, Front. Phys. 18(2), 21302 (2023) [arXiv:2102.04814v2[quant-ph]].
- [322] D. Gaiotto and T. Johnson-Freyd, *Condensations in higher categories*, [arXiv:1905.09566 [math.CT]].
- [323] T. Johnson-Freyd and M. Yu, *Topological Orders in (4+1)-Dimensions*, SciPost Phys. **13** (2022) no.3, 068 doi:10.21468/SciPostPhys.13.3.068 [arXiv:2104.04534 [hep-th]].
- [324] T. Johnson-Freyd, *On the Classification of Topological Orders*, Commun. Math. Phys. **393** (2022) no.2, 989-1033 doi:10.1007/s00220-022-04380-3 [arXiv:2003.06663 [math.CT]].
- [325] M. Yu et al., *Anomalies for noninvertible surfaces and the 2D Deligne Theorem*, to appear.
- [326] V. Bashmakov, M. Del Zotto and A. Hasan, *On the 6d Origin of Non-invertible Symmetries in 4d*, [arXiv:2206.07073 [hep-th]].
- [327] V. Bashmakov, M. Del Zotto, A. Hasan and J. Kaidi, *Non-invertible Symmetries of Class S Theories*, [arXiv:2211.05138 [hep-th]].
- [328] L. Bhardwaj, S. Schafer-Nameki and A. Tiwari, *Unifying Constructions of Non-Invertible Symmetries*, [arXiv:2212.06159 [hep-th]].

- [329] L. Bhardwaj, L. E. Bottini, S. Schafer-Nameki and A. Tiwari, *Non-Invertible Higher-Categorical Symmetries*, [arXiv:2204.06564 [hep-th]].
- [330] J. Kaidi, K. Ohmori, Y. Zheng, *Kramers-Wannier like Duality Defects in (3+1)D Gauge Theories*, Phys. Rev. Lett. **128** (2022) 111601, [2111.01141].
- [331] J. Kaidi, K. Ohmori and Y. Zheng, *Kramers-Wannier-like Duality Defects in (3+1)D Gauge Theories*, Phys. Rev. Lett. **128** (2022) no.11, 111601 doi:10.1103/PhysRevLett.128.111601 [arXiv:2111.01141 [hep-th]].
- [332] J. Kaidi, G. Zafrir and Y. Zheng, *Non-invertible symmetries of $\mathcal{N} = 4$ SYM and twisted compactification*, JHEP **08** (2022), 053 doi:10.1007/JHEP08(2022)053 [arXiv:2205.01104 [hep-th]].
- [333] J. Kaidi, K. Ohmori and Y. Zheng, *Symmetry TFTs for Non-Invertible Defects*, [arXiv:2209.11062 [hep-th]].
- [334] Y. Choi, C. Cordova, P. S. Hsin, H. T. Lam and S. H. Shao, *Non-invertible Condensation, Duality, and Triality Defects in 3+1 Dimensions*, [arXiv:2204.09025 [hep-th]].
- [335] Y. Choi, C. Cordova, P. S. Hsin, H. T. Lam and S. H. Shao, *Noninvertible duality defects in 3+1 dimensions*, Phys. Rev. D **105** (2022) no.12, 125016 doi:10.1103/PhysRevD.105.125016 [arXiv:2111.01139 [hep-th]].
- [336] T. Bartsch, M. Bullimore, A. E. V. Ferrari and J. Pearson, *Non-invertible Symmetries and Higher Representation Theory II*, [arXiv:2212.07393 [hep-th]].
- [337] T. Johnson-Freyd, *Operators and higher categories in quantum field theory*, Lecture series.
- [338] L. Kong, *Anyon condensation and tensor categories*, Nucl. Phys. B **886** (2014), 436-482 doi:10.1016/j.nuclphysb.2014.07.003 [arXiv:1307.8244 [cond-mat.str-el]].
- [339] M. Yu, *Gauging Categorical Symmetries in 3d Topological Orders and Bulk Reconstruction*, [arXiv:2111.13697 [hep-th]].
- [340] W. Ji and X. G. Wen, *Non-invertible anomalies and mapping-class-group transformation of anomalous partition functions*, Phys. Rev. Research. **1** (2019), 033054 doi:10.1103/PhysRevResearch.1.033054 [arXiv:1905.13279 [cond-mat.str-el]].
- [341] J. C. Y. Teo, T. L. Hughes and E. Fradkin, *Theory of Twist Liquids: Gauging an Anyonic Symmetry*, Annals Phys. **360** (2015), 349-445 doi:10.1016/j.aop.2015.05.012 [arXiv:1503.06812 [cond-mat.str-el]].
- [342] M. Barkeshli, P. Bonderson, M. Cheng and Z. Wang, *Symmetry Fractionalization, Defects, and Gauging of Topological Phases*, Phys. Rev. B **100** (2019) no.11, 115147 doi:10.1103/PhysRevB.100.115147 [arXiv:1410.4540 [cond-mat.str-el]].
- [343] L. Rozansky and E. Witten, *HyperKahler geometry and invariants of three manifolds*, Selecta Math. **3** (1997), 401-458 doi:10.1007/s000290050016 [arXiv:hep-th/9612216 [hep-th]].
- [344] S. D'Alesio, *Derived representation schemes and Nakajima quiver varieties*, Selecta Math. **28** (2022) no.1, 20 doi:10.1007/s00029-021-00724-4
- [345] Y. Berest, G. Khachatryan, A. Ramadoss, *Derived representation schemes and cyclic homology*, Adv. Math. **245**, 625–689 (2013).
- [346] E. Gonzalez, C. Y. Mak and D. Pomerleano, *Coulomb branch algebras via symplectic cohomology*, [arXiv:2305.04387 [math.SG]].

- [347] A. Kapustin and L. Rozansky, *Three-dimensional topological field theory and symplectic algebraic geometry II*, Commun. Num. Theor. Phys. **4** (2010), 463-550
doi:10.4310/CNTP.2010.v4.n3.a1 [arXiv:0909.3643 [math.AG]].
- [348] A. Kapustin, L. Rozansky and N. Saulina, *Three-dimensional topological field theory and symplectic algebraic geometry I*, Nucl. Phys. B **816** (2009), 295-355
doi:10.1016/j.nuclphysb.2009.01.027 [arXiv:0810.5415 [hep-th]].
- [349] R. Bezrukavnikov, M. Finkelberg, and I. Mirković, *Equivariant homology and K-theory of affine Grassmannians and Toda lattices*, Compos.Math. **141** (2005) 746-768.
- [350] L. Bhardwaj, M. Bullimore, A. E. V. Ferrari and S. Schafer-Nameki, *Generalized Symmetries and Anomalies of 3d $N=4$ SCFTs*, [arXiv:2301.02249 [hep-th]].
- [351] D. Gaiotto and J. Kulp, *Orbifold groupoids*, JHEP **02** (2021), 132
doi:10.1007/JHEP02(2021)132 [arXiv:2008.05960 [hep-th]].
- [352] L. Kong, T. Lan, X. G. Wen, Z. H. Zhang and H. Zheng, *Algebraic higher symmetry and categorical symmetry – a holographic and entanglement view of symmetry*, Phys. Rev. Res. **2** (2020) no.4, 043086 doi:10.1103/PhysRevResearch.2.043086 [arXiv:2005.14178 [cond-mat.str-el]].
- [353] L. Kong and Z. H. Zhang, *An invitation to topological orders and category theory*, [arXiv:2205.05565 [cond-mat.str-el]].
- [354] L. Kong and H. Zheng, *A mathematical theory of gapless edges of 2d topological orders. Part I*, JHEP **02** (2020), 150 doi:10.1007/JHEP02(2020)150 [arXiv:1905.04924 [cond-mat.str-el]].
- [355] L. Kong and H. Zheng, *A mathematical theory of gapless edges of 2d topological orders. Part II*, Nucl. Phys. B **966** (2021), 115384 doi:10.1016/j.nuclphysb.2021.115384 [arXiv:1912.01760 [cond-mat.str-el]].
- [356] L. Kong and H. Zheng, *Categorical computation*, Front. Phys. **18**(2), 21302 (2023) [arXiv:2102.04814v2[quant-ph]].
- [357] D. Gaiotto and T. Johnson-Freyd, *Condensations in higher categories*, [arXiv:1905.09566 [math.CT]].
- [358] T. Johnson-Freyd and M. Yu, *Topological Orders in $(4+1)$ -Dimensions*, SciPost Phys. **13** (2022) no.3, 068 doi:10.21468/SciPostPhys.13.3.068 [arXiv:2104.04534 [hep-th]].
- [359] T. Johnson-Freyd, *On the Classification of Topological Orders*, Commun. Math. Phys. **393** (2022) no.2, 989-1033 doi:10.1007/s00220-022-04380-3 [arXiv:2003.06663 [math.CT]].
- [360] T. D. Décoppet and M. Yu, *Gauging noninvertible defects: a 2-categorical perspective*, Lett. Math. Phys. **113** (2023) no.2, 36 doi:10.1007/s11005-023-01655-1 [arXiv:2211.08436 [math.CT]].
- [361] T. Johnson-Freyd, *Operators and higher categories in quantum field theory*, Lecture series.
- [362] K. Hori and C. Vafa, *Mirror symmetry*, [arXiv:hep-th/0002222 [hep-th]].
- [363] A. Klemm, W. Lerche, P. Mayr, C. Vafa and N. P. Warner, *Selfdual strings and $N=2$ supersymmetric field theory*, Nucl. Phys. B **477** (1996), 746-766
doi:10.1016/0550-3213(96)00353-7 [arXiv:hep-th/9604034 [hep-th]].
- [364] C. Teleman, Western Hemisphere Colloquium on Geometry and Physics, online talk, 25h April 2023.

- [365] W. Crawley-Boevey, Geometry of the moment map for representations of quivers, *Compositio Math.* **126** (2001), no. 3, 257–293.
- [366] V. Bashmakov, M. Del Zotto and A. Hasan, *On the 6d Origin of Non-invertible Symmetries in 4d*, [arXiv:2206.07073 [hep-th]].
- [367] V. Bashmakov, M. Del Zotto, A. Hasan and J. Kaidi, *Non-invertible Symmetries of Class \mathcal{S} Theories*, [arXiv:2211.05138 [hep-th]].
- [368] J. Kaidi, K. Ohmori, Y. Zheng, *Kramers-Wannier like Duality Defects in (3+1)D Gauge Theories*, *Phys. Rev. Lett.* **128** (2022) 111601, [2111.01141].
- [369] J. Kaidi, K. Ohmori and Y. Zheng, *Kramers-Wannier-like Duality Defects in (3+1)D Gauge Theories*, *Phys. Rev. Lett.* **128** (2022) no.11, 111601 doi:10.1103/PhysRevLett.128.111601 [arXiv:2111.01141 [hep-th]].
- [370] Y. Choi, C. Cordova, P. S. Hsin, H. T. Lam and S. H. Shao, *Non-invertible Condensation, Duality, and Triality Defects in 3+1 Dimensions*, [arXiv:2204.09025 [hep-th]].
- [371] Y. Choi, C. Cordova, P. S. Hsin, H. T. Lam and S. H. Shao, *Noninvertible duality defects in 3+1 dimensions*, *Phys. Rev. D* **105** (2022) no.12, 125016 doi:10.1103/PhysRevD.105.125016 [arXiv:2111.01139 [hep-th]].
- [372] O. Aharony, N. Seiberg and Y. Tachikawa, *Reading between the lines of four-dimensional gauge theories*, *JHEP* **08** (2013), 115 doi:10.1007/JHEP08(2013)115 [arXiv:1305.0318 [hep-th]].
- [373] T. Johnson-Freyd, *Operators and higher categories in quantum field theory*, Lecture series.
- [374] L. Kong, *Anyon condensation and tensor categories*, *Nucl. Phys. B* **886** (2014), 436–482 doi:10.1016/j.nuclphysb.2014.07.003 [arXiv:1307.8244 [cond-mat.str-el]].
- [375] M. Yu, *Gauging Categorical Symmetries in 3d Topological Orders and Bulk Reconstruction*, [arXiv:2111.13697 [hep-th]].
- [376] W. Ji and X. G. Wen, *Non-invertible anomalies and mapping-class-group transformation of anomalous partition functions*, *Phys. Rev. Research.* **1** (2019), 033054 doi:10.1103/PhysRevResearch.1.033054 [arXiv:1905.13279 [cond-mat.str-el]].
- [377] F. Benini, Y. Tachikawa and D. Xie, *Mirrors of 3d Sicilian theories*, *JHEP* **09** (2010), 063 doi:10.1007/JHEP09(2010)063 [arXiv:1007.0992 [hep-th]].
- [378] O. Chacaltana, J. Distler and Y. Tachikawa, *Nilpotent orbits and codimension-two defects of 6d $N=(2,0)$ theories*, *Int. J. Mod. Phys. A* **28** (2013), 1340006 doi:10.1142/S0217751X1340006X [arXiv:1203.2930 [hep-th]].
- [379] D. Pomerleano, *Intrinsic mirror symmetry and categorical crepant resolutions*, arXiv:2103.01200 [math.SG].
- [380] Y. Berest, G. Khachatryan, A. Ramadoss, *Derived representation schemes and cyclic homology*, *Adv. Math.* **245**, 625–689 (2013).
- [381] R. Street, R. Walters, *Yoneda structures on 2-categories*, *Journal of Algebra*, **50**, 2, 350–379, 1978, 0021-8693, [https://doi.org/10.1016/0021-8693\(78\)90160-6](https://doi.org/10.1016/0021-8693(78)90160-6).
- [382] A. Kapustin and L. Rozansky, *Three-dimensional topological field theory and symplectic algebraic geometry II*, *Commun. Num. Theor. Phys.* **4** (2010), 463–550 doi:10.4310/CNTP.2010.v4.n3.a1 [arXiv:0909.3643 [math.AG]].

- [383] A. Kapustin, L. Rozansky and N. Saulina, *Three-dimensional topological field theory and symplectic algebraic geometry I*, Nucl. Phys. B **816** (2009), 295-355
doi:10.1016/j.nuclphysb.2009.01.027 [arXiv:0810.5415 [hep-th]].
- [384] John C. Baez and James Dolan, Higher-dimensional algebra and topological quantum field theory, J. Math. Phys. **36** (1995), no. 11, 6073–6105.
- [385] Perry, A. Hochschild cohomology and group actions. Math. Z. **297**, 1273–1292 (2021).
<https://doi.org/10.1007/s00209-020-02557-x>.
- [386] R. Bezrukavnikov, M. Finkelberg, and I. Mirković, *Equivariant homology and K-theory of affine Grassmannians and Toda lattices*, Compos.Math. **141** (2005) 746-768.
- [387] P. Tosteson, *Categorifications of rational Hilbert series and characters of FS^{op} modules*, Alg. Number Th. **16** (2022) 2433-2491, arXiv:2105.01711 [math.RT].
- [388] Radha Kessar, Markus Linckelmann, *On the Hilbert series of Hochschild cohomology of block algebras*, Journal of Algebra, Volume 371, 2012, Pages 457-461, ISSN 0021-8693,
<https://doi.org/10.1016/j.jalgebra.2012.07.020>.
- [389] A. Neitzke, *Hitchin Systems in $\mathcal{N} = 2$ Field Theory*, doi:10.1007/978-3-319-18769-3_3
[arXiv:1412.7120 [hep-th]].
- [390] J. Lurie, On the Classification of Topological Field Theories, Current Developments in Mathematics 2008 (2009), 129–280, arXiv:0905.0465 [math.CT].
- [391] N. Carqueville, F. Montiel Montoya, *Extending Landau-Ginzburg models to a point*, arXiv:1809.10965 [math.QA].
- [392] J. M. Maldacena, *Large N field theories, string theory and gravity*,
- [393] H. Lin, O. Lunin and J. M. Maldacena, *Bubbling AdS space and 1/2 BPS geometries*, JHEP **10** (2004), 025 doi:10.1088/1126-6708/2004/10/025 [arXiv:hep-th/0409174 [hep-th]].
- [394] J. M. Maldacena, *Quantum gravity as an ordinary gauge theory*,
- [395] G. T. Horowitz and J. M. Maldacena, *The Black hole final state*, JHEP **02** (2004), 008
doi:10.1088/1126-6708/2004/02/008 [arXiv:hep-th/0310281 [hep-th]].
- [396] J. M. Maldacena, *TASI 2003 lectures on AdS / CFT*, [arXiv:hep-th/0309246 [hep-th]].
- [397] S. Kachru, R. Kallosh, A. D. Linde, J. M. Maldacena, L. P. McAllister and S. P. Trivedi, *Towards inflation in string theory*, JCAP **10** (2003), 013 doi:10.1088/1475-7516/2003/10/013
[arXiv:hep-th/0308055 [hep-th]].
- [398] N. D. Lambert, H. Liu and J. M. Maldacena, *Closed strings from decaying D-branes*, JHEP **03** (2007), 014 doi:10.1088/1126-6708/2007/03/014 [arXiv:hep-th/0303139 [hep-th]].
- [399] L. F. Alday, D. Gaiotto and J. Maldacena, *Thermodynamic Bubble Ansatz*, JHEP **09** (2011), 032 doi:10.1007/JHEP09(2011)032 [arXiv:0911.4708 [hep-th]].
- [400] I. R. Klebanov and J. M. Maldacena, *Solving quantum field theories via curved spacetimes*, Phys. Today **62** (2009), 28-33 doi:10.1063/1.3074260
- [401] L. F. Alday and J. Maldacena, *Lectures on scattering amplitudes via AdS/CFT*, AIP Conf. Proc. **1031** (2008) no.1, 43-60 doi:10.1063/1.2972015
- [402] J. Maldacena, *Comments on emergent space-time*,

- [403] A. Dymarsky, S. S. Gubser, Z. Guralnik and J. M. Maldacena, *Calibrated surfaces and supersymmetric Wilson loops*, JHEP **09** (2006), 057 doi:10.1088/1126-6708/2006/09/057 [arXiv:hep-th/0604058 [hep-th]].
- [404] J. Maldacena, *Testing gauge/gravity duality on a quantum black hole*, Science **344** (2014), 806-807 doi:10.1126/science.1254597
- [405] A. Lewkowycz and J. Maldacena, *Exact results for the entanglement entropy and the energy radiated by a quark*, JHEP **05** (2014), 025 doi:10.1007/JHEP05(2014)025 [arXiv:1312.5682 [hep-th]].
- [406] T. Faulkner, A. Lewkowycz and J. Maldacena, *Quantum corrections to holographic entanglement entropy*, JHEP **11** (2013), 074 doi:10.1007/JHEP11(2013)074 [arXiv:1307.2892 [hep-th]].
- [407] J. Maldacena and L. Susskind, *Cool horizons for entangled black holes*, Fortsch. Phys. **61** (2013), 781-811 doi:10.1002/prop.201300020 [arXiv:1306.0533 [hep-th]].
- [408] A. Lewkowycz and J. Maldacena, *Generalized gravitational entropy*, JHEP **08** (2013), 090 doi:10.1007/JHEP08(2013)090 [arXiv:1304.4926 [hep-th]].
- [409] J. Maldacena, *The quantum spacetime*
- [410] D. Bigatti and L. Susskind, *The Holographic Principle*, NATO Sci. Ser. C **556** (2000), 179-226 doi:10.1007/978-94-011-4303-5_4
- [411] L. Susskind and N. Toumbas, *Wilson loops as precursors*, Phys. Rev. D **61** (2000), 044001 doi:10.1103/PhysRevD.61.044001 [arXiv:hep-th/9909013 [hep-th]].
- [412] J. Polchinski and L. Susskind, *Puzzles and paradoxes about holography*, [arXiv:hep-th/9902182 [hep-th]].
- [413] W. Fischler and L. Susskind, *Holography and cosmology*, [arXiv:hep-th/9806039 [hep-th]].
- [414] L. Susskind and E. Witten, *The Holographic bound in anti-de Sitter space*, [arXiv:hep-th/9805114 [hep-th]].
- [415] L. Susskind, *The Landscape and its Physics Foundations How String Theory Generates the Landscape*, Subnucl. Ser. **44** (2008), 161-252 doi:10.1142/9789812832467_0006
- [416] L. Susskind, G. Arcioni and S. de Haro, *Holographic Views of the World*, [arXiv:physics/0611143 [physics]].
- [417] L. Susskind, *The paradox of quantum black holes*, Nature Phys. **2** (2006) no.10, 665-677 doi:10.1038/nphys429
- [418] G. 't Hooft, L. Susskind, E. Witten, M. Fukugita, L. Randall, L. Smolin, J. Stachel, C. Rovelli, G. Ellis and S. Weinberg, *et al. A theory of everything?*, Nature **433** (2005), 257-259 doi:10.1038/433257a
- [419] L. Susskind and J. Lindesay, *An introduction to black holes, information and the string theory revolution: The holographic universe*
- [420] B. Freivogel and L. Susskind, *A Framework for the landscape*, Phys. Rev. D **70** (2004), 126007 doi:10.1103/PhysRevD.70.126007 [arXiv:hep-th/0408133 [hep-th]].
- [421] L. Dyson, J. Lindesay and L. Susskind, *Is there really a de Sitter/CFT duality?*, JHEP **08** (2002), 045 doi:10.1088/1126-6708/2002/08/045 [arXiv:hep-th/0202163 [hep-th]].

- [422] R. Bousso and L. Susskind, *The Multiverse Interpretation of Quantum Mechanics*, Phys. Rev. D **85** (2012), 045007 doi:10.1103/PhysRevD.85.045007 [arXiv:1105.3796 [hep-th]].
- [423] G. 't Hooft, *How studying black hole theory may help us to quantise gravity*, [arXiv:2211.10723 [gr-qc]].
- [424] G. 't Hooft, *Quantum mechanics, statistics, standard model and gravity*, Gen. Rel. Grav. **54** (2022) no.6, 56 doi:10.1007/s10714-022-02939-y
- [425] G. 't Hooft, *Quantum field theory with boundary conditions at the horizons*, Int. J. Mod. Phys. D **31** (2022) no.01, 2130008 doi:10.1142/S0218271821300081
- [426] N. Gaddam, N. Groenenboom and G. 't Hooft, *Quantum gravity on the black hole horizon*, JHEP **01** (2022), 023 doi:10.1007/JHEP01(2022)023 [arXiv:2012.02357 [hep-th]].
- [427] G. 't Hooft, *Black Hole Microstates*, doi:10.1142/9789811203961_0006
- [428] G. 't Hooft, *The quantum black hole as a theoretical lab, a pedagogical treatment of a new approach*, [arXiv:1902.10469 [gr-qc]].
- [429] O. Aharony, M. Berkooz, A. Karasik and T. Vaknin, *Supersymmetric field theories on $AdS_p \times S^q$* , JHEP **04** (2016), 066 doi:10.1007/JHEP04(2016)066 [arXiv:1512.04698 [hep-th]].
- [430] O. Aharony and M. Evtikhiev, *On four dimensional $N = 3$ superconformal theories*, JHEP **04** (2016), 040 doi:10.1007/JHEP04(2016)040 [arXiv:1512.03524 [hep-th]].
- [431] O. Aharony, M. Berkooz and S. J. Rey, *Rigid holography and six-dimensional $\mathcal{N} = (2, 0)$ theories on $AdS_5 \times S^1$* , JHEP **03** (2015), 121 doi:10.1007/JHEP03(2015)121 [arXiv:1501.02904 [hep-th]].
- [432] O. Aharony, S. S. Razamat, N. Seiberg and B. Willett, *3d dualities from 4d dualities*, JHEP **07** (2013), 149 doi:10.1007/JHEP07(2013)149 [arXiv:1305.3924 [hep-th]].
- [433] O. Aharony, D. Marolf and M. Rangamani, *Conformal field theories in anti-de Sitter space*, JHEP **02** (2011), 041 doi:10.1007/JHEP02(2011)041 [arXiv:1011.6144 [hep-th]].
- [434] O. Aharony and S. Kachru, *Stringy Instantons and Cascading Quivers*, JHEP **09** (2007), 060 doi:10.1088/1126-6708/2007/09/060 [arXiv:0707.3126 [hep-th]].
- [435] A. Castro, I. Coman, J. R. Fliss and C. Zukowski, *Coupling Fields to 3D Quantum Gravity via Chern-Simons Theory*, Phys. Rev. Lett. **131** (2023) no.17, 171602 doi:10.1103/PhysRevLett.131.171602 [arXiv:2304.02668 [hep-th]].
- [436] A. Castro, I. Coman, J. R. Fliss and C. Zukowski, *Keeping matter in the loop in dS_3 quantum gravity*, JHEP **07** (2023), 120 doi:10.1007/JHEP07(2023)120 [arXiv:2302.12281 [hep-th]].
- [437] A. Castro, *Probes in AdS_3 Quantum Gravity*, doi:10.1007/978-3-030-55777-5_36
- [438] A. Belin, A. Castro, C. A. Keller and B. Mühlmann, *The Holographic Landscape of Symmetric Product Orbifolds*, JHEP **01** (2020), 111 doi:10.1007/JHEP01(2020)111 [arXiv:1910.05342 [hep-th]].
- [439] A. Castro, V. Godet, F. Larsen and Y. Zeng, *Logarithmic Corrections to Black Hole Entropy: the Non-BPS Branch*, JHEP **05** (2018), 079 doi:10.1007/JHEP05(2018)079 [arXiv:1801.01926 [hep-th]].
- [440] A. Castro, D. M. Hofman and N. Iqbal, *Entanglement Entropy in Warped Conformal Field Theories*, JHEP **02** (2016), 033 doi:10.1007/JHEP02(2016)033 [arXiv:1511.00707 [hep-th]].

- [441] R. Bittleston and D. Skinner, *Twistors, the ASD Yang-Mills equations and 4d Chern-Simons theory*, JHEP **02** (2023), 227 doi:10.1007/JHEP02(2023)227 [arXiv:2011.04638 [hep-th]].
- [442] K. Roehrig and D. Skinner, *Ambitwistor strings and the scattering equations on $AdS_3 \times S^3$* , JHEP **02** (2022), 073 doi:10.1007/JHEP02(2022)073 [arXiv:2007.07234 [hep-th]].
- [443] T. Adamo, D. Skinner and J. Williams, *Twistor methods for AdS_5* , JHEP **08** (2016), 167 doi:10.1007/JHEP08(2016)167 [arXiv:1607.03763 [hep-th]].
- [444] E. Witten, *Supersymmetry and Morse theory*, J. Diff. Geom. **17** (1982) no.4, 661-692
- [445] Donaldson, Simon (1983), "An Application of Gauge Theory to Four Dimensional Topology", Journal of Differential Geometry, 18 (2): 279–315, MR 0710056.
- [446] D. Gaiotto, *Monster symmetry and Extremal CFTs*, JHEP **11** (2012), 149 doi:10.1007/JHEP11(2012)149 [arXiv:0801.0988 [hep-th]].
- [447] F. Denef, D. Gaiotto, A. Strominger, D. Van den Bleeken and X. Yin, *Black Hole Deconstruction*, JHEP **03** (2012), 071 doi:10.1007/JHEP03(2012)071 [arXiv:hep-th/0703252 [hep-th]].
- [448] D. Gaiotto, A. Strominger and X. Yin, *From $AdS(3)/CFT(2)$ to black holes/topological strings*, JHEP **09** (2007), 050 doi:10.1088/1126-6708/2007/09/050 [arXiv:hep-th/0602046 [hep-th]].
- [449] E. Hatefi and I. Y. Park, *More on closed string induced higher derivative interactions on D-branes*, Phys. Rev. D **85** (2012), 125039 doi:10.1103/PhysRevD.85.125039 [arXiv:1203.5553 [hep-th]].
- [450] L. Katzarkov, M. Kontsevich and T. Pantev, *Hodge theoretic aspects of mirror symmetry*, Proc. Symp. Pure Math. **78** (2008), 87-174 doi:10.1090/pspum/078/2483750 [arXiv:0806.0107 [math.AG]].
- [451] A. Grassi, Q. Hao and A. Neitzke, *Exponential Networks, WKB and Topological String*, SIGMA **19** (2023), 064 doi:10.3842/SIGMA.2023.064 [arXiv:2201.11594 [hep-th]].
- [452] A. Neitzke and A. Shehper, *Deformations of surface defect moduli spaces*, JHEP **10** (2021), 035 doi:10.1007/JHEP10(2021)035 [arXiv:2011.01970 [hep-th]].
- [453] C. Córdova and A. Neitzke, *Line Defects, Tropicalization, and Multi-Centered Quiver Quantum Mechanics*, JHEP **09** (2014), 099 doi:10.1007/JHEP09(2014)099 [arXiv:1308.6829 [hep-th]].
- [454] A. Neitzke, *Notes on a new construction of hyperkahler metrics*, [arXiv:1308.2198 [math.DG]].
- [455] A. Neitzke, *On a hyperholomorphic line bundle over the Coulomb branch*, [arXiv:1110.1619 [hep-th]].
- [456] A. Balasubramanian, J. Distler, R. Donagi and C. Perez-Pardavila, *The Hitchin Image in Type-D*, [arXiv:2310.05880 [hep-th]].
- [457] A. Balasubramanian, J. Distler and R. Donagi, *Families of Hitchin systems and $N = 2$ theories*, Adv. Theor. Math. Phys. **26** (2022) no.6, 1585-1667 doi:10.4310/ATMP.2022.v26.n6.a2 [arXiv:2008.01020 [hep-th]].
- [458] F. Beck, R. Donagi and K. Wendland, *Folding of Hitchin Systems and Crepant Resolutions*, Int. Math. Res. Not. **2022** (2022) no.11, 8370-8419 doi:10.1093/imrn/rnaa375 [arXiv:2004.04245 [math.AG]].

- [459] R. Donagi and D. R. Morrison, *Conformal field theories and compact curves in moduli spaces*, JHEP **05** (2018), 021 doi:10.1007/JHEP05(2018)021 [arXiv:1709.05355 [hep-th]].
- [460] T. Pantev and E. Sharpe, *Decomposition in Chern-Simons theories in three dimensions*, Int. J. Mod. Phys. A **37** (2022), 2250227 doi:10.1142/S0217751X2250227X [arXiv:2206.14824 [hep-th]].
- [461] T. Pantev and M. Wijnholt, *Hitchin's Equations and M-Theory Phenomenology*, J. Geom. Phys. **61** (2011), 1223-1247 doi:10.1016/j.geomphys.2011.02.014 [arXiv:0905.1968 [hep-th]].
- [462] R. Donagi and T. Pantev, *Langlands duality for Hitchin systems*, [arXiv:math/0604617 [math.AG]].
- [463] G. 't Hooft, *Quantum Foundations as a Guide for Refining Particle Theories*, [arXiv:2312.09396 [hep-th]].
- [464] G. 't Hooft, *The Standard Model and Beyond*, J. Phys. Conf. Ser. **1127** (2019) no.1, 012001 doi:10.1088/1742-6596/1127/1/012001
- [465] V. Braun, Y. H. He, B. A. Ovrut and T. Pantev, *The Exact MSSM spectrum from string theory*, JHEP **05** (2006), 043 doi:10.1088/1126-6708/2006/05/043 [arXiv:hep-th/0512177 [hep-th]].
- [466] V. Braun, Y. H. He, B. A. Ovrut and T. Pantev, *Vector bundle extensions, sheaf cohomology, and the heterotic standard model*, Adv. Theor. Math. Phys. **10** (2006) no.4, 525-589 doi:10.4310/ATMP.2006.v10.n4.a3 [arXiv:hep-th/0505041 [hep-th]].
- [467] S. Lüst, C. Vafa, M. Wiesner and K. Xu, *Holography and the KKLT scenario*, JHEP **10** (2022), 188 doi:10.1007/JHEP10(2022)188 [arXiv:2204.07171 [hep-th]].
- [468] H. Ooguri and C. Vafa, *On the Geometry of the String Landscape and the Swampland*, Nucl. Phys. B **766** (2007), 21-33 doi:10.1016/j.nuclphysb.2006.10.033 [arXiv:hep-th/0605264 [hep-th]].
- [469] M. R. Douglas and Z. Lu, *Finiteness of volume of moduli spaces*, [arXiv:hep-th/0509224 [hep-th]].
- [470] N. Arkani-Hamed, L. Motl, A. Nicolis and C. Vafa, *The String landscape, black holes and gravity as the weakest force*, JHEP **06** (2007), 060 doi:10.1088/1126-6708/2007/06/060 [arXiv:hep-th/0601001 [hep-th]].
- [471] X. G. Wen and E. Witten, *Electric and Magnetic Charges in Superstring Models*, Nucl. Phys. B **261** (1985), 651-677 doi:10.1016/0550-3213(85)90592-9
- [472] G. 't Hooft, *Magnetic Monopoles in Unified Gauge Theories*, Nucl. Phys. B **79** (1974), 276-284 doi:10.1016/0550-3213(74)90486-6
- [473] A. M. Polyakov, *Particle Spectrum in Quantum Field Theory*, JETP Lett. **20** (1974), 194-195 PRINT-74-1566 (LANDAU-INST).
- [474] M. C. Tan, *Vafa-Witten Theory: Invariants, Floer Homologies, Higgs Bundles, a Geometric Langlands Correspondence, and Categorification*, [arXiv:2305.15948 [hep-th]].
- [475] New invariants of 3- and 4-dimensional manifolds, The mathematical heritage of Hermann Weyl, 48., 285-299 M. Atiyah
- [476] L. Alvarez-Gaume and D. Z. Freedman, *Geometrical Structure and Ultraviolet Finiteness in the Supersymmetric Sigma Model*, Commun. Math. Phys. **80** (1981), 443 doi:10.1007/BF01208280

- [477] M. Abouzaid and C. Manolescu, *A sheaf-theoretic model for $SL(2, \mathbb{C})$ Floer homology*, J. Eur. Math. Soc. **22** (2020) no.11, 3641-3695 doi:10.4171/jems/994 [arXiv:1708.00289 [math.GT]].
- [478] Symmetries and stabilization for sheaves of vanishing cycles, arXiv preprint C. Brav, V. Bussi, D. Dupont, D. Joyce, B. Szendroi
- [479] Morse theory for Lagrangian intersections A. Floer J.Diff.Geom. 28 (1988) 513-47
- [480] An instanton-invariant for 3-manifolds A. Floer Commun.Math.Phys. 118 (1988) 215-40
- [481] Lectures on the Langlands program and conformal field theory, Frontiers in Number Theory, Physics, and Geometry II: On Conformal Field Theories, Discrete Groups and Renormalization,, 387-533 E. Frenkel
- [482] Surface operators and knot homologies, In New Trends in Mathematical Physics: Selected contributions of the XVth International Congress on Mathematical Physics,, (pp. 313-343) S. Gukov
- [483] The self-duality equations on a Riemann surface N.J. Hitchin Proc.Lond.Math.Soc. 3 (1987) 59-126
- [484] M. R. Douglas, *Gauge fields and D-branes*, J. Geom. Phys. **28** (1998), 255-262 doi:10.1016/S0393-0440(97)00024-7 [arXiv:hep-th/9604198 [hep-th]].
- [485] E. Witten, *Small instantons in string theory*, Nucl. Phys. B **460** (1996), 541-559 doi:10.1016/0550-3213(95)00625-7 [arXiv:hep-th/9511030 [hep-th]].
- [486] Vitaly Ginzburg, Lev Landau, On the Theory of Superconductivity, In: On Superconductivity and Superfluidity Springer (2009) (doi:1007/978-3-540-68008-64).
- [487] A note on quantum geometric Langlands duality, gauge theory, and quantization of the moduli space of flat connections, arXiv preprint A. Kapustin
- [488] Surface operators in four-dimensional topological gauge theory and Langlands duality, arXiv preprint A. Kapustin, K. Setter, K. Vyas
- [489] Electric-magnetic duality and the geometric Langlands program, arXiv preprint A. Kapustin, E. Witten
- [490] A quick view of Lagrangian Floer homology, Geometrical Themes Inspired by the N-body Problem,, 91-125 A. Pedroza
- [491] Topological quantum field theory and the geometric Langlands correspondence, California Institute of Technology K. Setter
- [492] C. Vafa and E. Witten, *A Strong coupling test of S duality*, Nucl. Phys. B **431** (1994), 3-77 doi:10.1016/0550-3213(94)90097-3 [arXiv:hep-th/9408074 [hep-th]].
- [493] H. Ooguri and T. Takayanagi, *Cobordism Conjecture in AdS*, [arXiv:2006.13953 [hep-th]].
- [494] D. Harlow and H. Ooguri, *Symmetries in quantum field theory and quantum gravity*, Commun. Math. Phys. **383** (2021) no.3, 1669-1804 doi:10.1007/s00220-021-04040-y [arXiv:1810.05338 [hep-th]].
- [495] N. Bao and H. Ooguri, *Distinguishability of black hole microstates*, Phys. Rev. D **96** (2017) no.6, 066017 doi:10.1103/PhysRevD.96.066017 [arXiv:1705.07943 [hep-th]].
- [496] J. Gomis, Z. Komargodski, H. Ooguri, N. Seiberg and Y. Wang, *Shortening Anomalies in Supersymmetric Theories*, JHEP **01** (2017), 067 doi:10.1007/JHEP01(2017)067 [arXiv:1611.03101 [hep-th]].

- [497] H. Ooguri and C. Vafa, *Non-supersymmetric AdS and the Swampland*, Adv. Theor. Math. Phys. **21** (2017), 1787-1801 doi:10.4310/ATMP.2017.v21.n7.a8 [arXiv:1610.01533 [hep-th]].
- [498] Y. Nakayama and H. Ooguri, *Bulk Local States and Crosscaps in Holographic CFT*, JHEP **10** (2016), 085 doi:10.1007/JHEP10(2016)085 [arXiv:1605.00334 [hep-th]].
- [499] N. Bao, S. Nezami, H. Ooguri, B. Stoica, J. Sully and M. Walter, *The Holographic Entropy Cone*, JHEP **09** (2015), 130 doi:10.1007/JHEP09(2015)130 [arXiv:1505.07839 [hep-th]].
- [500] H. Ooguri, *Lectures on topological string theory*, Lect. Notes Phys. **851** (2012), 233-263 doi:10.1007/978-3-642-25947-0_6
- [501] R. Dijkgraaf and C. Vafa, *Matrix models, topological strings, and supersymmetric gauge theories*, Nucl. Phys. B **644** (2002), 3-20 doi:10.1016/S0550-3213(02)00766-6 [arXiv:hep-th/0206255 [hep-th]].
- [502] R. Gopakumar and C. Vafa, *M theory and topological strings. 1.*, [arXiv:hep-th/9809187 [hep-th]].
- [503] R. Gopakumar and C. Vafa, *M theory and topological strings. 2.*, [arXiv:hep-th/9812127 [hep-th]].
- [504] H. Ooguri, A. Strominger and C. Vafa, *Black hole attractors and the topological string*, Phys. Rev. D **70** (2004), 106007 doi:10.1103/PhysRevD.70.106007 [arXiv:hep-th/0405146 [hep-th]].
- [505] M. Aganagic, H. Ooguri, C. Vafa and M. Yamazaki, *Wall Crossing and M-theory*, Publ. Res. Inst. Math. Sci. Kyoto **47** (2011), 569 [arXiv:0908.1194 [hep-th]].
- [506] R. Kitano, H. Ooguri and Y. Ookouchi, *Supersymmetry Breaking and Gauge Mediation*, Ann. Rev. Nucl. Part. Sci. **60** (2010), 491-511 doi:10.1146/annurev.nucl.012809.104540 [arXiv:1001.4535 [hep-th]].
- [507] H. Ooguri, *Geometry As Seen By String Theory*, [arXiv:0901.1881 [math.AG]].
- [508] H. Ooguri, *The Entropic Principle and the Landscape in SUSY Gauge Theories*, Subnucl. Ser. **45** (2009), 117-128 doi:10.1142/9789814293242_0005
- [509] T. Kawano, H. Ooguri and Y. Ookouchi, *Gauge Mediation in String Theory*, Phys. Lett. B **652** (2007), 40-42 doi:10.1016/j.physletb.2007.06.056 [arXiv:0704.1085 [hep-th]].
- [510] H. Ooguri, *Entropy of small black holes*, Prog. Theor. Phys. Suppl. **163** (2006), 355-357 doi:10.1143/PTPS.163.355
- [511] R. Dijkgraaf, R. Gopakumar, H. Ooguri and C. Vafa, *Baby universes in string theory*, Phys. Rev. D **73** (2006), 066002 doi:10.1103/PhysRevD.73.066002 [arXiv:hep-th/0504221 [hep-th]].
- [512] H. Ooguri, C. Vafa and E. P. Verlinde, *Hartle-Hawking wave-function for flux compactifications*, Lett. Math. Phys. **74** (2005), 311-342 doi:10.1007/s11005-005-0022-x [arXiv:hep-th/0502211 [hep-th]].
- [513] M. Aganagic, H. Ooguri, N. Saulina and C. Vafa, *Black holes, q-deformed 2d Yang-Mills, and non-perturbative topological strings*, Nucl. Phys. B **715** (2005), 304-348 doi:10.1016/j.nuclphysb.2005.02.035 [arXiv:hep-th/0411280 [hep-th]].
- [514] N. Nekrasov, H. Ooguri and C. Vafa, *S duality and topological strings*, JHEP **10** (2004), 009 doi:10.1088/1126-6708/2004/10/009 [arXiv:hep-th/0403167 [hep-th]].
- [515] A. Neitzke and C. Vafa, *Topological strings and their physical applications*, [arXiv:hep-th/0410178 [hep-th]].

- [516] P. Kraus, H. Ooguri and S. Shenker, *Inside the horizon with AdS / CFT*, Phys. Rev. D **67** (2003), 124022 doi:10.1103/PhysRevD.67.124022 [arXiv:hep-th/0212277 [hep-th]].
- [517] M. Aganagic, E. LePage and M. Rapcak, *Homological Link Invariants from Floer Theory*, [arXiv:2305.13480 [hep-th]].
- [518] M. Aganagic, *Homological Knot Invariants from Mirror Symmetry*, doi:10.4171/ICM2022/197 [arXiv:2207.14104 [math.GT]].
- [519] M. Aganagic, *Knot Categorification from Mirror Symmetry, Part II: Lagrangians*, [arXiv:2105.06039 [hep-th]].
- [520] M. Aganagic, *Knot Categorification from Mirror Symmetry, Part I: Coherent Sheaves*, [arXiv:2004.14518 [hep-th]].
- [521] M. Aganagic, K. Costello, J. McNamara and C. Vafa, *Topological Chern-Simons/Matter Theories*, [arXiv:1706.09977 [hep-th]].
- [522] M. Aganagic, E. Frenkel and A. Okounkov, *Quantum q -Langlands Correspondence*, Trans. Moscow Math. Soc. **79** (2018), 1-83 doi:10.1090/mosc/278 [arXiv:1701.03146 [hep-th]].
- [523] M. Aganagic, *String theory and math: Why this marriage may last. Mathematics and dualities of quantum physics*, Bull. Am. Math. Soc. **53** (2016) no.1, 93-115 doi:10.1090/bull/1517 [arXiv:1508.06642 [hep-th]].
- [524] M. Aganagic and S. Shakirov, *Gauge/Vortex duality and AGT*, doi:10.1007/978-3-319-18769-3_13 [arXiv:1412.7132 [hep-th]].
- [525] M. Aganagic, T. Ekhholm, L. Ng and C. Vafa, *Topological Strings, D-Model, and Knot Contact Homology*, Adv. Theor. Math. Phys. **18** (2014) no.4, 827-956 doi:10.4310/ATMP.2014.v18.n4.a3 [arXiv:1304.5778 [hep-th]].
- [526] I. García Etxebarria, *Branes and Non-Invertible Symmetries*, Fortsch. Phys. **70** (2022) no.11, 2200154 doi:10.1002/prop.202200154 [arXiv:2208.07508 [hep-th]].
- [527] M. Del Zotto and I. García Etxebarria, *Global structures from the infrared*, JHEP **11** (2023), 058 doi:10.1007/JHEP11(2023)058 [arXiv:2204.06495 [hep-th]].
- [528] F. Apruzzi, F. Bonetti, I. García Etxebarria, S. S. Hosseini and S. Schafer-Nameki, *Symmetry TFTs from String Theory*, Commun. Math. Phys. **402** (2023) no.1, 895-949 doi:10.1007/s00220-023-04737-2 [arXiv:2112.02092 [hep-th]].
- [529] M. Del Zotto, I. García Etxebarria and S. S. Hosseini, *Higher form symmetries of Argyres-Douglas theories*, JHEP **10** (2020), 056 doi:10.1007/JHEP10(2020)056 [arXiv:2007.15603 [hep-th]].
- [530] F. Albertini, M. Del Zotto, I. García Etxebarria and S. S. Hosseini, *Higher Form Symmetries and M-theory*, JHEP **12** (2020), 203 doi:10.1007/JHEP12(2020)203 [arXiv:2005.12831 [hep-th]].
- [531] I. García Etxebarria, B. Heidenreich and D. Regalado, *IIB flux non-commutativity and the global structure of field theories*, JHEP **10** (2019), 169 doi:10.1007/JHEP10(2019)169 [arXiv:1908.08027 [hep-th]].
- [532] I. García-Etxebarria and D. Regalado, *Exceptional $\mathcal{N} = 3$ theories*, JHEP **12** (2017), 042 doi:10.1007/JHEP12(2017)042 [arXiv:1611.05769 [hep-th]].
- [533] I. García Etxebarria and N. Iqbal, *A Goldstone theorem for continuous non-invertible symmetries*, JHEP **09** (2023), 145 doi:10.1007/JHEP09(2023)145 [arXiv:2211.09570 [hep-th]].

- [534] F. Apruzzi, F. Bonetti, D. S. W. Gould and S. Schafer-Nameki, *Aspects of Categorical Symmetries from Branes: SymTFTs and Generalized Charges*, [arXiv:2306.16405 [hep-th]].
- [535] S. Schafer-Nameki, *ICTP Lectures on (Non-)Invertible Generalized Symmetries*, [arXiv:2305.18296 [hep-th]].
- [536] L. Bhardwaj and S. Schafer-Nameki, *Generalized Charges, Part I: Invertible Symmetries and Higher Representations*, [arXiv:2304.02660 [hep-th]].
- [537] L. Bhardwaj, M. Bullimore, A. E. V. Ferrari and S. Schafer-Nameki, *Generalized Symmetries and Anomalies of 3d $N=4$ SCFTs*, [arXiv:2301.02249 [hep-th]].
- [538] L. Bhardwaj, L. E. Bottini, S. Schafer-Nameki and A. Tiwari, *Non-invertible symmetry webs*, SciPost Phys. **15** (2023) no.4, 160 doi:10.21468/SciPostPhys.15.4.160 [arXiv:2212.06842 [hep-th]].
- [539] L. Bhardwaj, S. Schafer-Nameki and A. Tiwari, *Unifying constructions of non-invertible symmetries*, SciPost Phys. **15** (2023) no.3, 122 doi:10.21468/SciPostPhys.15.3.122 [arXiv:2212.06159 [hep-th]].
- [540] M. van Beest, D. S. W. Gould, S. Schafer-Nameki and Y. N. Wang, *Symmetry TFTs for 3d QFTs from M-theory*, JHEP **02** (2023), 226 doi:10.1007/JHEP02(2023)226 [arXiv:2210.03703 [hep-th]].
- [541] L. Bhardwaj, S. Schafer-Nameki and J. Wu, *Universal Non-Invertible Symmetries*, Fortsch. Phys. **70** (2022) no.11, 2200143 doi:10.1002/prop.202200143 [arXiv:2208.05973 [hep-th]].
- [542] M. Hubner, D. R. Morrison, S. Schafer-Nameki and Y. N. Wang, *Generalized Symmetries in F-theory and the Topology of Elliptic Fibrations*, SciPost Phys. **13** (2022) no.2, 030 doi:10.21468/SciPostPhys.13.2.030 [arXiv:2203.10022 [hep-th]].
- [543] L. Bhardwaj, S. Giacomelli, M. Hübner and S. Schäfer-Nameki, *Relative defects in relative theories: Trapped higher-form symmetries and irregular punctures in class S*, SciPost Phys. **13** (2022) no.4, 101 doi:10.21468/SciPostPhys.13.4.101 [arXiv:2201.00018 [hep-th]].
- [544] C. Closset, S. Schäfer-Nameki and Y. N. Wang, *Coulomb and Higgs branches from canonical singularities. Part I. Hypersurfaces with smooth Calabi-Yau resolutions*, JHEP **04** (2022), 061 doi:10.1007/JHEP04(2022)061 [arXiv:2111.13564 [hep-th]].
- [545] F. Apruzzi, L. Bhardwaj, D. S. W. Gould and S. Schafer-Nameki, *2-Group symmetries and their classification in 6d*, SciPost Phys. **12** (2022) no.3, 098 doi:10.21468/SciPostPhys.12.3.098 [arXiv:2110.14647 [hep-th]].
- [546] C. Closset, S. Schafer-Nameki and Y. N. Wang, *Coulomb and Higgs Branches from Canonical Singularities: Part 0*, JHEP **02** (2021), 003 doi:10.1007/JHEP02(2021)003 [arXiv:2007.15600 [hep-th]].
- [547] F. Apruzzi, S. Schafer-Nameki, L. Bhardwaj and J. Oh, *The Global Form of Flavor Symmetries and 2-Group Symmetries in 5d SCFTs*, SciPost Phys. **13** (2022) no.2, 024 doi:10.21468/SciPostPhys.13.2.024 [arXiv:2105.08724 [hep-th]].
- [548] L. Bhardwaj, M. Hubner and S. Schafer-Nameki, *1-form Symmetries of 4d $N=2$ Class S Theories*, SciPost Phys. **11** (2021), 096 doi:10.21468/SciPostPhys.11.5.096 [arXiv:2102.01693 [hep-th]].
- [549] P. S. Aspinwall, T. Bridgeland, A. Craw, M. R. Douglas, A. Kapustin, G. W. Moore, M. Gross, G. Segal, B. Szendrői and P. M. H. Wilson, *Dirichlet branes and mirror symmetry*, AMS, 2009,

- [550] D. Ben-Zvi, A. Brochier and D. Jordan, *Integrating quantum groups over surfaces*, doi:10.1112/topo.12072 [arXiv:1501.04652 [math.QA]].
- [551] A. Brochier, D. Jordan and N. Snyder, *On dualizability of braided tensor categories*, Compos. Math. **157** (2021) no.3, 435-483 doi:10.1112/S0010437X20007630 [arXiv:1804.07538 [math.QA]].
- [552] S. Gunningham, D. Jordan and P. Safronov, *The finiteness conjecture for skein modules*, Invent. Math. **232** (2023) no.1, 301-363 doi:10.1007/s00222-022-01167-0 [arXiv:1908.05233 [math.QA]].
- [553] D. Jordan, *Langlands duality for skein modules of 3-manifolds*, [arXiv:2302.14734 [math.QA]].
- [554] D. Jordan, *Quantum character varieties*, [arXiv:2309.06543 [math.QA]].
- [555] The Yang-Mills equations over Riemann surfaces. Philos. Trans. Roy. Soc. London Ser. A / 308(1505):523-615 M.F. Atiyah, R. Bott
- [556] Factorization homology of topological manifolds D. Ayala, J. Francis J.Topol. 8 (2015) 1045-1084
- [557] D. Ayala and J. Francis, *The cobordism hypothesis*, [arXiv:1705.02240 [math.AT]].
- [558] D. Ayala, J. Francis and N. Rozenblyum, *Factorization homology I: Higher categories*, Adv. Math. **333** (2018), 1042-1177 [erratum: Adv. Math. **370** (2020), 107217] doi:10.1016/j.aim.2018.05.031 [arXiv:1504.04007 [math.AT]]
- [559] A. Y. Alekseev, H. Grosse and V. Schomerus, *Combinatorial quantization of the Hamiltonian Chern-Simons theory. 2.*, Commun. Math. Phys. **174** (1995), 561-604 doi:10.1007/BF02101528 [arXiv:hep-th/9408097 [hep-th]].
- [560] A. Y. Alekseev, *Integrability in the Hamiltonian Chern-Simons theory*, [arXiv:hep-th/9311074 [hep-th]].
- [561] M. Aganagic and S. Shakirov, *Knot Homology and Refined Chern-Simons Index*, Commun. Math. Phys. **333** (2015) no.1, 187-228 doi:10.1007/s00220-014-2197-4 [arXiv:1105.5117 [hep-th]].
- [562] J. C. Baez and J. Dolan, *Higher dimensional algebra and topological quantum field theory*, J. Math. Phys. **36** (1995), 6073-6105 doi:10.1063/1.531236 [arXiv:q-alg/9503002 [math.QA]].
- [563] Locally presentable and accessible categories, vol. 189 (Cambridge University Press,) J. Adámek, J. Rosický
- [564] J. C. Baez and J. Dolan, *Higher dimensional algebra and topological quantum field theory*, J. Math. Phys. **36** (1995), 6073-6105 doi:10.1063/1.531236 [arXiv:q-alg/9503002 [math.QA]].
- [565] Chiral algebras, vol. 51 of American Mathematical Society Colloquium Publications (American Mathematical Society, Providence, RI,) A. Beilinson, V. Drinfeld
- [566] Modular categories as representations of the 3dimensional bordism 2-category. ArXiv e-prints B. Bartlett, C.L. Douglas, C.J. Schommer-Pries, J. Vicary
- [567] Jr. Lectures on tensor categories and modular functors, vol. 21 of University Lecture Series (American Mathematical Society, Providence, RI,) B. Bakalov, A. Kirillov
- [568] Double affine Hecke algebras and generalized Jones polynomials Y. Berest, P. Samuelson Compos.Math. 152 (2016) 1333-1384

- [569] O. Aharony, A. Hanany and B. Kol, *Webs of (p,q) five-branes, five-dimensional field theories and grid diagrams*, JHEP **01** (1998), 002 doi:10.1088/1126-6708/1998/01/002 [arXiv:hep-th/9710116 [hep-th]].
- [570] A. Hanany and A. Zaffaroni, *Branes and six-dimensional supersymmetric theories*, Nucl. Phys. B **529** (1998), 180-206 doi:10.1016/S0550-3213(98)00355-1 [arXiv:hep-th/9712145 [hep-th]].
- [571] A. Hanany and J. Walcher, *On duality walls in string theory*, JHEP **06** (2003), 055 doi:10.1088/1126-6708/2003/06/055 [arXiv:hep-th/0301231 [hep-th]].
- [572] S. Franco and A. Hanany, *Geometric dualities in 4-D field theories and their 5-D interpretation*, JHEP **04** (2003), 043 doi:10.1088/1126-6708/2003/04/043 [arXiv:hep-th/0207006 [hep-th]].
- [573] A. Hanany, N. Prezas and J. Troost, *The Partition function of the two-dimensional black hole conformal field theory*, JHEP **04** (2002), 014 doi:10.1088/1126-6708/2002/04/014 [arXiv:hep-th/0202129 [hep-th]].
- [574] A. Hanany and B. Pioline, *(Anti-)instantons and the Atiyah-Hitchin manifold*, JHEP **07** (2000), 001 doi:10.1088/1126-6708/2000/07/001 [arXiv:hep-th/0005160 [hep-th]].
- [575] B. Feng, A. Hanany and Y. H. He, *Counting gauge invariants: The Plethystic program*, JHEP **03** (2007), 090 doi:10.1088/1126-6708/2007/03/090 [arXiv:hep-th/0701063 [hep-th]].
- [576] A. Hanany and D. Vegh, *Quivers, tilings, branes and rhombi*, JHEP **10** (2007), 029 doi:10.1088/1126-6708/2007/10/029 [arXiv:hep-th/0511063 [hep-th]].
- [577] S. Franco, A. Hanany, D. Martelli, J. Sparks, D. Vegh and B. Wecht, *Gauge theories from toric geometry and brane tilings*, JHEP **01** (2006), 128 doi:10.1088/1126-6708/2006/01/128 [arXiv:hep-th/0505211 [hep-th]].
- [578] S. Franco, A. Hanany, K. D. Kennaway, D. Vegh and B. Wecht, *Brane dimers and quiver gauge theories*, JHEP **01** (2006), 096 doi:10.1088/1126-6708/2006/01/096 [arXiv:hep-th/0504110 [hep-th]].
- [579] A. Hanany and K. D. Kennaway, *Dimer models and toric diagrams*, [arXiv:hep-th/0503149 [hep-th]].
- [580] S. Benvenuti, A. Hanany and N. Mekareeya, *The Hilbert Series of the One Instanton Moduli Space*, JHEP **06** (2010), 100 doi:10.1007/JHEP06(2010)100 [arXiv:1005.3026 [hep-th]].
- [581] A. Hanany and A. Zaffaroni, *The master space of supersymmetric gauge theories*, Adv. High Energy Phys. **2010** (2010), 427891 doi:10.1155/2010/427891
- [582] D. Forcella, A. Hanany and A. Zaffaroni, *Master Space, Hilbert Series and Seiberg Duality*, JHEP **07** (2009), 018 doi:10.1088/1126-6708/2009/07/018 [arXiv:0810.4519 [hep-th]].
- [583] J. L. Huang, J. McGreevy and B. Shi, *Knots and entanglement*, SciPost Phys. **14** (2023) no.6, 141 doi:10.21468/SciPostPhys.14.6.141 [arXiv:2112.08398 [hep-th]].
- [584] J. Cotler and A. Strominger, *Cosmic ER=EPR in dS/CFT*, [arXiv:2302.00632 [hep-th]].
- [585] I. R. Klebanov, D. Kutasov and A. Murugan, *Entanglement as a probe of confinement*, Nucl. Phys. B **796** (2008), 274-293 doi:10.1016/j.nuclphysb.2007.12.017 [arXiv:0709.2140 [hep-th]].
- [586] Z. Komargodski, K. Ohmori, K. Roumpedakis and S. Seifnashri, *Symmetries and strings of adjoint QCD₂*, JHEP **03** (2021), 103 doi:10.1007/JHEP03(2021)103 [arXiv:2008.07567 [hep-th]].

- [587] E. Witten, *Constraints on Supersymmetry Breaking*, Nucl. Phys. B **202** (1982), 253 doi:10.1016/0550-3213(82)90071-2
- [588] A. Schwimmer and N. Seiberg, “Comments on the N=2, N=3, N=4 Superconformal Algebras in Two-Dimensions,” Phys. Lett. B **184**, 191 (1987).
- [589] D. L. Jafferis, I. R. Klebanov, S. S. Pufu and B. R. Safdi, JHEP **06** (2011) 102, arXiv:1103.1181.
- [590] K. A. Intriligator and B. Wecht, Nucl. Phys. B **667**, 183 (2003) [arXiv:hep-th/0304128].
- [591] J. Penedones, K. Salehi Vaziri and Z. Sun, *Hilbert space of Quantum Field Theory in de Sitter spacetime*, [arXiv:2301.04146 [hep-th]].
- [592] H. P. Nilles, *Phenomenological aspects of supersymmetry*, [arXiv:hep-ph/9511313 [hep-ph]].
- [593] C. Csaki, *The Minimal supersymmetric standard model (MSSM)*, Mod. Phys. Lett. A **11** (1996), 599 doi:10.1142/S021773239600062X [arXiv:hep-ph/9606414 [hep-ph]].
- [594] P. Slavich, S. Heinemeyer, E. Bagnaschi, H. Bahl, M. Goodsell, H. E. Haber, T. Hahn, R. Harlander, W. Hollik and G. Lee, *et al. Higgs-mass predictions in the MSSM and beyond*, Eur. Phys. J. C **81** (2021) no.5, 450 doi:10.1140/epjc/s10052-021-09198-2 [arXiv:2012.15629 [hep-ph]].
- [595] W. Abdallah *et al.* [LHC Reinterpretation Forum], *Reinterpretation of LHC Results for New Physics: Status and Recommendations after Run 2*, SciPost Phys. **9** (2020) no.2, 022 doi:10.21468/SciPostPhys.9.2.022 [arXiv:2003.07868 [hep-ph]].
- [596] B. C. Allanach, F. Quevedo and K. Suruliz, *Low-energy supersymmetry breaking from string flux compactifications: Benchmark scenarios*, JHEP **04** (2006), 040 doi:10.1088/1126-6708/2006/04/040 [arXiv:hep-ph/0512081 [hep-ph]].
- [597] V. Pasquarella, to appear.
- [598] A. Braverman, M. Finkelberg and H. Nakajima, *Ring objects in the equivariant derived Satake category arising from Coulomb branches (with an appendix by Gus Lonergan)*, [arXiv:1706.02112 [math.RT]].
- [599] A. Braverman, M. Finkelberg and H. Nakajima, *Towards a mathematical definition of Coulomb branches of 3-dimensional $\mathcal{N} = 4$ gauge theories, II*, Adv. Theor. Math. Phys. **22** (2018), 1071-1147 doi:10.4310/ATMP.2018.v22.n5.a1 [arXiv:1601.03586 [math.RT]].
- [600] T. Dimofte and N. Garner, *Coulomb Branches of Star-Shaped Quivers*, JHEP **02** (2019), 004 doi:10.1007/JHEP02(2019)004 [arXiv:1808.05226 [hep-th]].
- [601] M. Bullimore, T. Dimofte and D. Gaiotto, *The Coulomb Branch of 3d $\mathcal{N} = 4$ Theories*, Commun. Math. Phys. **354** (2017) no.2, 671-751 doi:10.1007/s00220-017-2903-0 [arXiv:1503.04817 [hep-th]].
- [602] D. Jordan, *Quantum character varieties*, [arXiv:2309.06543 [math.QA]].
- [603] *Koszul duality between Higgs and Coulomb categories \mathcal{O}* , [arXiv:1611.06541 [math.RT]].
- [604] L. Gráf, B. Henning, X. Lu, T. Melia and H. Murayama, *Hilbert series, the Higgs mechanism, and HEFT*, JHEP **02** (2023), 064 doi:10.1007/JHEP02(2023)064 [arXiv:2211.06275 [hep-ph]].
- [605] M. P. Bento, J. P. Silva and A. Trautner, *The basis invariant flavor puzzle*, JHEP **01** (2024), 024 doi:10.1007/JHEP01(2024)024 [arXiv:2308.00019 [hep-ph]].

- [606] Anisha, S. Das Bakshi, J. Chakraborty and S. Prakash, *Hilbert Series and Plethystics: Paving the path towards 2HDM- and MLRSM-EFT*, JHEP **09** (2019), 035 doi:10.1007/JHEP09(2019)035 [arXiv:1905.11047 [hep-ph]].
- [607] T. Johnson-Freyd, *Operators and higher categories in quantum field theory*, Lecture series.
- [608] Conference Proceedings From Quiver Diagrams to Particle Physics Uranga, Angel M. Casacuberta, Carles Miró-Roig, Rosa Maria Verdera, Joan Xambó-Descamps, Sebastià European Congress of Mathematics 2001 Birkhäuser Basel Basel 978-3-0348-8266-8 10.1007/978-3-0348-8266-8 43
- [609] B. C. Allanach, B. Gripaios and J. Tooby-Smith, *Semisimple extensions of the Standard Model gauge algebra*, Phys. Rev. D **104** (2021) no.3, 035035 [erratum: Phys. Rev. D **106** (2022) no.1, 019901] doi:10.1103/PhysRevD.104.035035 [arXiv:2104.14555 [hep-th]].
- [610] A. Hanany and E. Witten, *Type IIB superstrings, BPS monopoles, and three-dimensional gauge dynamics*, Nucl. Phys. B **492** (1997), 152-190 doi:10.1016/S0550-3213(97)00157-0 [arXiv:hep-th/9611230 [hep-th]].
- [611] P. W. Higgs, *Broken symmetries, massless particles and gauge fields*, Phys. Lett. **12** (1964), 132-133 doi:10.1016/0031-9163(64)91136-9
- [612] P. W. Higgs, *Spontaneous Symmetry Breakdown without Massless Bosons*, Phys. Rev. **145** (1966), 1156-1163 doi:10.1103/PhysRev.145.1156
- [613] Deligne, Pierre; Mumford, David. The irreducibility of the space of curves of given genus. Publications Mathématiques de l’IHÉS, Volume 36 (1969), pp. 75-109. http://www.numdam.org/item/PMIHES_1969_36_75_0/.
- [614] H. Nakajima, *A mathematical definition of Coulomb branches of supersymmetric gauge theories and geometric Satake correspondences for Kac-Moody Lie algebras*, [arXiv:2201.08386 [math.RT]].
- [615] Y. Berest, G. Khachatryan, A. Ramadoss, *Derived representation schemes and cyclic homology*, Adv. Math. **245**, 625–689 (2013).
- [616] S. Cremonesi, A. Hanany, N. Mekareeya and A. Zaffaroni, *Coulomb branch Hilbert series and Three Dimensional Sicilian Theories*, JHEP **09** (2014), 185 doi:10.1007/JHEP09(2014)185 [arXiv:1403.2384 [hep-th]].
- [617] F. Apruzzi, I. Bah, F. Bonetti and S. Schafer-Nameki, *Noninvertible Symmetries from Holography and Branes*, Phys. Rev. Lett. **130** (2023) no.12, 121601 doi:10.1103/PhysRevLett.130.121601 [arXiv:2208.07373 [hep-th]].
- [618] B. Marsh, K. Rietsch, *The B-model connection and mirror symmetry for Grassmannians*, arXiv:1307.1085 [math.AG].
- [619] Radha Kessar, Markus Linckelmann, *On the Hilbert series of Hochschild cohomology of block algebras*, Journal of Algebra, Volume 371, 2012, Pages 457-461, ISSN 0021-8693, <https://doi.org/10.1016/j.jalgebra.2012.07.020>. bmodel,Ba,Frey:2018vpw,JP,Jansen:1994ym,BWB,stein,STY,PC1,PC2
- [620] D. D. Frey and T. Rudelius, *6D SCFTs and the classification of homomorphisms $\Gamma_{ADE} \rightarrow E_8$* , Adv. Theor. Math. Phys. **24** (2020) no.3, 709-756 doi:10.4310/ATMP.2020.v24.n3.a4 [arXiv:1811.04921 [hep-th]].
- [621] P. Bruillard, P. Gustafson, J. Yael Plavnik, E. Carson Rowell, *Dimension as a quantum statistic and the classification of metaplectic categories*, arXiv:1710.10284 [math.QA].

- [622] P. M. Vranas, *Domain wall fermions and applications*, Nucl. Phys. B Proc. Suppl. **94** (2001), 177-188 doi:10.1016/S0920-5632(01)00948-3 [arXiv:hep-lat/0011066 [hep-lat]].
- [623] B. Kostant, *Lie algebra cohomology and the generalized Borel-Weil theorem*, Ann. of Math. 74, 2, 1961, pp. 329-387.
- [624] A. Balibanu, *Steinberg slices and group-valued moment maps*, arXiv:2103.14704 [math.RT].
- [625] S.-T. Yau, *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation*, Communications on Pure and Applied Mathematics, 31 3 (1978) 339-411 [doi:10.1002/cpa.3160310304].
- [626] P. Candelas, X. de la Ossa, F. Rodriguez-Villegas, *Calabi-Yau Manifolds Over Finite Fields, I*, (arXiv:hep-th/0012233).
- [627] P. Candelas, X. de la Ossa, F. Rodriguez-Villegas, *Calabi-Yau Manifolds Over Finite Fields, II*, (arXiv:hep-th/0402133).