

Geometry of Invariant Almost Semi Kähler Submanifolds of Flag Manifolds

Neiton Pereira da Silva¹

Abstract

In this paper we discuss the geometry of homogeneous spaces which are almost Hermitian submanifolds of flag manifolds. We prove that such spaces are necessarily minimal submanifolds and in the case where these submanifolds are also flag manifolds, they are totally geodesic.

Mathematics Subject Classifications: 53C55; 53D15; 22F30 .

Keywords: Flag manifolds; almost Hermitian manifold; Almost semi Kähler manifolds.

1 Introduction

It is a well known fact that a Kähler submanifold is a minimal variety (see [13]). This result is also valid for others classes of almost Hermitian structures, [7]. In this paper we extend this result for invariant almost semi-Kähler submanifolds of flag manifolds.

An almost complex manifold is a manifold M equipped with an tensor field J of type $(1, 1)$, called an almost complex structure, such that $J^2 = -1$. Such a manifold is orientable and has even dimension, say $2m$. An elementary text on almost complex manifolds can be found in [9].

Recall the Nijenhuis tensor of J , defined by:

$$\frac{1}{2}N(X, Y) = [JX, JY] - J[JX, Y] - J[X, JY] - [X, Y], \quad X, Y \in \mathfrak{X}(M),$$

a well known theorem of Newlander and Nirenberg states that the almost complex manifold (M, J) is a complex manifold if and only if J is integrable, that is $N = 0$. An Almost Hermitian manifold (M, J, g) is an almost complex manifold (M, J) with a J -invariant Riemannian metric g , i.e.,

¹Neiton Pereira da Silva at UFU. Email: neiton@ufu.br

$g(X, Y) = g(JX, JY)$, for any $X, Y \in \mathfrak{X}(M)$. In the sequel we will abbreviate Almost Hermitian manifold by AH-manifold. The Kähler form of the almost Hermitian manifold (M, J, g) is the differential form $\Omega(X, Y) = g(X, JY)$, for any $X, Y \in \mathfrak{X}(M)$.

Following [11], associated to the Levi-Civita connection ∇ on M , we have the following formulas for the covariant derivative, the exterior differentiation and the codifferential of Ω :

$$(\nabla_X \Omega)(Y, Z) = g(Y, (\nabla_X J)Z), \quad 3d\Omega(X, Y, Z) = \sigma(\nabla_X \Omega)(Y, Z), \text{ and}$$

$$(\delta\Omega)(X) = \sum_{i=1}^m \{(\nabla_{E_i} \Omega)(E_i, X) + (\nabla_{JE_i} \Omega)(JE_i, X)\},$$

where σ denotes the cyclic sum over X, Y, Z and $\{E_1, \dots, E_m, JE_1, \dots, JE_m\}$ is a local orthonormal J-frame.

In [8], A. Gray and L. M. Hervella classify the AH-manifolds in 16 classes and show certain inclusion relations between them. In the next table we recall some of these classes, with the defining condition.

symbol	name	condition
K	Kähler	$\nabla J = 0$
AK	Almost Kähler	$d\Omega = 0$
NK	Nearly Kähler	$(\nabla_X J)X = 0$
QK	Quasi Kähler	$(\nabla_X J)Y + (\nabla_{JX} J)JY = 0$
ASK	Almost semi-Kähler	$\delta\Omega = 0$
SK	Semi-Kähler	$\delta\Omega = 0, N = 0$
		(here N is the Nijenhuis tensor)
H	Hermitian	$N = 0$

Certain types of almost Hermitian manifolds (e.g., H, AK) have been studied by several authors with the aim of generalizing the Kähler geometry.

A $2r$ -dimension submanifold M' of M is called holomorphic (or invariant or almost-complex) submanifold of M if for any $p \in M'$ the tangent space $T_p M'$ is J -invariant, i.e. $J(T_p M') = T_p M'$.

In [7], A. Gray obtained: *If M belongs to one of the classes QK, AK, NK, H, K, then any holomorphic submanifold M' of M belongs to the same class. Moreover, M' is necessarily minimal, except in the Hermitian case.*

Note that the classes ASK and SK are excluded from the above Gray's result. According [11], in general, it is not known if holomorphic submanifolds of almost semi-Kähler manifolds are minimal. However, if M' is a

holomorphic submanifold of an ASK-manifold with codimension 2, then M' is minimal (see [16]).

In this paper we show that Gray's result can be extended for the classes ASK and SK when $M = G/P$ is a flag manifold (i.e. M is a homogeneous space G/P where G is semi-simple Lie group and P is a parabolic subgroup) and G/P is endowed with a almost Hermitian structure (J, g) invariant by the action of G . More precisely, we prove that if $(G/P, J, g)$ is an invariant almost Hermitian flag manifold (which belong to the class ASK or SK). Then any invariant holomorphic submanifold M' of G/P belongs to the same class. Moreover, M' is necessarily minimal. See Theorem (7.4) and Corollary (7.5). We use the linear operators defined in [11] to study the minimality of M' .

In the case that an invariant holomorphic submanifold M' of G/P is also a flag manifold, we obtain an very interesting result which states that M' is a totally geodesic submanifold with respect to any invariant metric g on G/P , see Theorem 8.3. Here we used properties of Lie theory to obtain the result.

Others authors have studied (and have been studying) geometric properties of some classes of almost Hermitian structures on flag manifolds, see for example [12], [14] and [1]. The main idea here is that the definition conditions of the 16 classes can be parameterized in terms of roots of the Lie algebra \mathfrak{g} of G . Thus investigating geometric and others properties reduces to a study in terms of Lie theory.

In section 2, we recall the linear operators and others results used to obtain the main theorems of this paper. In section 3, we give the description of flag manifold as a even dimension real homogeneous space by means of Lie theory. In section 4, we set our notation for invariant almost Hermitian structures on flag manifolds recalling the notion of invariant almost complex structure and Riemannian metric on a flag manifold. In section 5, we compute explicitly the Riemannian connection of invariant Riemannian metric on the Weyl basis. In section 6, we show that the covariant derivative of the Kähler form is zero when computed in an invariant global orthonormal J -frame. As a consequence we obtain that all flag manifolds belong to the class ASK. In section 7, we study homogeneous holomorphic submanifolds of flag manifolds. One of the main results of this paper is Theorem 7.4. In section 8, we study the case where the homogeneous holomorphic submanifold is also a flag manifold and obtain the other main result, Theorem 8.3.

2 Almost Complex submanifolds

Now let (M, J, g) be an AH-manifold and M' a holomorphic submanifold. The almost complex structure induced on M' and the almost Hermitian Riemannian metric induced on M' will be also denoted by J and g , respectively. Note that $J(T_p M'^{\perp}) = T_p M'^{\perp}$, because if $Z \in T_p M'^{\perp}$ then $g(JZ, X) = -g(Z, JX) = 0$, for any $X \in T_p M'$, since $J(T_p M') = T_p M'$.

Recall the Gauss formula:

$$\nabla_X Y = \nabla'_X Y + \alpha(X, Y), \quad X, Y \in \mathfrak{X}(M')$$

where ∇' and α denote the Riemannian connection and the second fundamental form of M' , respectively.

An important type of holomorphic submanifold are the minimal one, in particular those totally geodesics. Next we recall these notions, see [15], p. 101, or [5] p.38, for details.

Definition 2.1. *Let (M', g, J) be an almost complex submanifold of (M, g, J) . The mean curvature of M' at $x \in M'$ is the normal vector*

$$H_x = \frac{1}{2r} \sum_{i=1}^r (\alpha(E_i, E_i) + \alpha(JE_i, JE_i))$$

where $\{E_1, \dots, E_r, JE_1, \dots, JE_r\}$ is a local orthonormal J -frame on TM' ,

- a) (M', g, J) is minimal if $H \equiv 0$;
- b) (M', g, J) is totally geodesic if $\alpha \equiv 0$.

Next, we denote by $\tilde{\mathfrak{X}}(M)$ the Lie algebra of vector fields tangent to M along M' . Thus,

$$\mathfrak{X}(M') \subset \tilde{\mathfrak{X}}(M) \subset \mathfrak{X}(M),$$

and

$$\tilde{\mathfrak{X}}(M) = \mathfrak{X}(M') \oplus \mathfrak{X}(M')^{\perp}. \quad (1)$$

Definition 2.2. *Let M' be a holomorphic submanifold of an AH-manifold (M, J, g) . The partial (or tangent) coderivative of Ω and the normal-coderivative of Ω , with respect to M' are the linear operators $\bar{\delta}\Omega$ and $\bar{\bar{\delta}}\Omega$, respectively, given by :*

$$\begin{aligned}
(\bar{\delta}\Omega)(X) &= \sum_{i=1}^r \{(\nabla_{E_i}\Omega)(E_i, X) + (\nabla_{JE_i}\Omega)(JE_i, X)\}, \\
(\bar{\bar{\delta}}\Omega)(X) &= \sum_{i=1}^{m-r} \{(\nabla_{F_i}\Omega)(F_i, X) + (\nabla_{JF_i}\Omega)(JF_i, X)\},
\end{aligned}$$

where $X \in \tilde{\mathfrak{X}}(M)$; $\{E_1, \dots, E_r, JE_1, \dots, JE_r\}$ is a local orthonormal J -frame on TM' , $\dim M' = 2r$ and $\{F_1, \dots, F_{m-r}, JF_1, \dots, JF_{m-r}\}$ is a local orthonormal J -frame on $(TM')^\perp$.

An easy computation shows that by using the Gauss formula,

$$\nabla_X J = \nabla'_X J + \alpha_X J, \quad X \in \mathfrak{X}(M') \quad (2)$$

where $(\alpha_X J)Y = \alpha(X, JY) - J\alpha(X, Y)$. Using equation (2), one proves that for any $Z \in \mathfrak{X}(M')^\perp$:

$$(\bar{\delta}\Omega)(Z) = 2rg(JH, Z) \quad (3)$$

where H is the mean curvature vector field of the submanifold M' (see [11] p. 63 or [7]). In particular, a almost complex submanifold M' of (M, J, g) is minimal if and only if $(\bar{\delta}\Omega) = 0$ on $\mathfrak{X}(M')^\perp$.

Proposition 2.3. [11] *Let M' be a holomorphic submanifold of an AH-manifold (M, J, g) . Then, for any $X \in \mathfrak{X}(M')$, one has:*

$$(\delta\Omega)(X) = (\delta'\Omega)(X) + (\bar{\bar{\delta}}\Omega)(X), \quad (4)$$

δ' denoting the codifferential operator on M' .

3 Flag manifolds

In this section we set up our notation and present the standard theory of partial (or generalized) flag manifolds associated with semisimple Lie algebras (see, for example, [1] or [4] for a similar description).

Let \mathfrak{g} be a finite-dimensional semisimple complex Lie algebra and G a Lie group with Lie algebra \mathfrak{g} . Consider a Cartan subalgebra \mathfrak{h} of \mathfrak{g} . We denote by R the system of roots of $(\mathfrak{g}, \mathfrak{h})$. A root $\alpha \in R$ is a linear functional on \mathfrak{g} . It determines uniquely an element $H_\alpha \in \mathfrak{h}$ by the Riesz representation

$\alpha(X) = B(X, H_\alpha)$, $X \in \mathfrak{g}$, with respect to the Killing form $B(\cdot, \cdot)$ of \mathfrak{g} . The Lie algebra \mathfrak{g} has the following decomposition

$$\mathfrak{g} = \mathfrak{h} \oplus \sum_{\alpha \in R} \mathfrak{g}_\alpha$$

where \mathfrak{g}_α is the one-dimensional root space corresponding to α . The eigenvectors $E_\alpha \in \mathfrak{g}_\alpha$ satisfy the following equation

$$[E_\alpha, E_{-\alpha}] = B(E_\alpha, E_{-\alpha}) H_\alpha. \quad (5)$$

We fix a system Σ of simple roots of R and denote by R^+ and R^- the corresponding set of positive and negative roots, respectively. Let $\Theta \subset \Sigma$ be a subset, define

$$R(\Theta) := \langle \Theta \rangle \cap R \quad \text{and} \quad R(\Theta)^\pm := \langle \Theta \rangle \cap R^\pm.$$

We denote by $R_\Theta := R \setminus R(\Theta)$ the complementary set of roots. In general, R_Θ is not a root system, see an example in [1].

The parabolic subalgebra \mathfrak{p}_Θ , associated to Θ , is defined by

$$\mathfrak{p}_\Theta := \mathfrak{h} \oplus \sum_{\alpha \in R^+} \mathfrak{g}_\alpha \oplus \sum_{\alpha \in R(\Theta)^-} \mathfrak{g}_\alpha.$$

Note that \mathfrak{p}_Θ contains the Borel subalgebra $\mathfrak{b}^+ = \mathfrak{h} \oplus \sum_{\alpha \in R^+} \mathfrak{g}_\alpha$.

The partial flag manifold determined by the choice of $\Theta \subset \Sigma$ is the homogeneous space $\mathbb{F}_\Theta = G/P_\Theta$, where P_Θ is the normalizer of \mathfrak{p}_Θ in G . In the special case $\Theta = \emptyset$, we obtain the maximal flag manifold $\mathbb{F} = G/B$ associated with \mathfrak{g} , where B is the normalizer of the Borel subalgebra.

Now we will see the construction of any flag manifold as the quotient U/K_Θ of a compact semisimple Lie group $U \subset G$ modulo the centralizer K_Θ of a torus in U . We fix once and for all a Weyl basis of \mathfrak{g} which amounts to giving $X_\alpha \in \mathfrak{g}_\alpha$, $H_\alpha \in \mathfrak{h}$ for $\alpha \in R$, with the standard properties:

$$B(X_\alpha, X_\beta) = \begin{cases} 1, & \alpha + \beta = 0, \\ 0, & \text{otherwise;} \end{cases} \quad [X_\alpha, X_\beta] = \begin{cases} H_\alpha \in \mathfrak{h}, & \alpha + \beta = 0, \\ n_{\alpha, \beta} X_{\alpha+\beta}, & \alpha + \beta \in R, \\ 0, & \text{otherwise.} \end{cases} \quad (6)$$

The real constants $n_{\alpha, \beta}$ are non zero if and only if $\alpha + \beta \in R$ and satisfy

$$\begin{cases} n_{\alpha, \beta} = -n_{-\alpha, -\beta} = -n_{\beta, \alpha} \\ n_{\alpha, \beta} = n_{\beta, \gamma} = n_{\gamma, \alpha}, \end{cases} \quad \text{if } \alpha + \beta + \gamma = 0.$$

Consider the following two-dimensional real spaces $\mathfrak{u}_\alpha = \text{span}_{\mathbb{R}}\{A_\alpha, iS_\alpha\}$, with $A_\alpha = X_\alpha - X_{-\alpha}$ and $S_\alpha = X_\alpha + X_{-\alpha}$, $\alpha \in R^+$. Then the real Lie algebra $\mathfrak{u} = i\mathfrak{h}_{\mathbb{R}} \oplus \sum \mathfrak{u}_\alpha$, $\alpha \in R^+$, is a compact real form of \mathfrak{g} , where $\mathfrak{h}_{\mathbb{R}}$ denotes the real vector space spanned by $\{H_\alpha : \alpha \in R\}$.

Let $U = \exp \mathfrak{u}$ be the compact real form of G corresponding to \mathfrak{u} . By the restriction of the action of G on \mathbb{F}_Θ , we can see that U acts transitively on \mathbb{F}_Θ , then $\mathbb{F}_\Theta = U/K_\Theta$, where $K_\Theta = P_\Theta \cap U$. The Lie algebra \mathfrak{k}_Θ of K_Θ is the set of fixed points of the conjugation $\tau: X_\alpha \mapsto -X_{-\alpha}$ of \mathfrak{g} restricted to \mathfrak{p}_Θ

$$\mathfrak{k}_\Theta = \mathfrak{u} \cap \mathfrak{p}_\Theta = i\mathfrak{h}_{\mathbb{R}} \oplus \sum_{\alpha \in R(\Theta)^+} \mathfrak{u}_\alpha.$$

The tangent space of $\mathbb{F}_\Theta = U/K_\Theta$ at the origin $o = eK_\Theta$ can be identified with the orthogonal complement (with respect to the Killing form) of \mathfrak{k}_Θ in \mathfrak{u}

$$T_o\mathbb{F}_\Theta = \mathfrak{m} = \sum_{\alpha \in R_\Theta^+} \mathfrak{u}_\alpha,$$

with $R_\Theta^+ = R_\Theta \cap R^+$. Thus we have $\mathfrak{u} = \mathfrak{k}_\Theta \oplus \mathfrak{m}$. The complexification of \mathfrak{m} is given by:

$$\mathfrak{m}^{\mathbb{C}} = \mathfrak{m} \otimes \mathbb{C} = \sum_{\alpha \in R_\Theta} \mathbb{C}X_\alpha.$$

4 Invariant almost Hermitian structures on flag manifolds

4.1 Invariant almost complex structures

In this section we describe almost complex structure on \mathbb{F}_Θ and Riemannian metric, both invariant by the action of the Lie group U , similar description can be found in [1], [14], or [12] for maximal flag manifolds.

An *almost complex structure* on $\mathbb{F}_\Theta = U/K_\Theta$ is a tensor field of type $(1, 1)$ that corresponds each $x \in \mathbb{F}_\Theta$ to a linear endomorphism $J_x: T_x\mathbb{F}_\Theta \rightarrow T_x\mathbb{F}_\Theta$ which satisfies $J_x^2 = -\text{Id}$. The almost complex structure on \mathbb{F}_Θ is invariant (or U -invariant) if

$$du_x \circ J_x = J_{ux} \circ du_x$$

for all $u \in U$. An *invariant almost complex structure* (*iacs* from now on) is determined by a linear endomorphism $J: \mathfrak{m} \rightarrow \mathfrak{m}$, which satisfies $J^2 = -\text{Id}$

and commutes with the adjoint action of K_Θ on \mathfrak{u} , that is,

$$\text{Ad}(k)J = J \text{Ad}(k), \quad \text{for all } k \in K_\Theta,$$

or, equivalently,

$$\text{ad}(L)J = J \text{ad}(L), \quad \text{for all } L \in \mathfrak{k}_\Theta.$$

We will use the same letter J to denote its extension to the complexification $\mathfrak{m}^\mathbb{C}$. Since $J^2 = -\text{Id}$, its eigenvalues are i and $-i$ and the corresponding eigenspaces are denoted by

$$T_o^{(1,0)}\mathbb{F}_\Theta = \{X \in T_o\mathbb{F}^\mathbb{C} : JX = iX\}$$

and

$$T_o^{(0,1)}\mathbb{F}_\Theta = \{X \in T_o\mathbb{F}^\mathbb{C} : JX = -iX\}.$$

Thus

$$\mathfrak{m}^\mathbb{C} = T_o\mathbb{F}_\Theta^\mathbb{C} = T_o^{(1,0)}\mathbb{F}_\Theta \oplus T_o^{(0,1)}\mathbb{F}_\Theta.$$

The eigenvectors with eigenvalue $+i$ (resp. $-i$) are called *of type (1,0)* (resp. *of type (0,1)*).

The invariance of J entails that $J(\mathfrak{g}_\alpha) = \mathfrak{g}_\alpha$ for all $\alpha \in R_\Theta$. The eigenvalues of J are $\pm i$ and the eigenvectors in $\mathfrak{m}^\mathbb{C}$ are X_α ; $\alpha \in R_\Theta$. Hence

$$J(X_\alpha) = i\varepsilon_\alpha X_\alpha$$

with $\varepsilon_\alpha = \pm 1$ satisfying $\varepsilon_{-\alpha} = -\varepsilon_\alpha$.

Thus an iacs on \mathbb{F}_Θ is completely prescribed by a set of signs $\{\varepsilon_\alpha\}_{\alpha \in R_\Theta}$ with $\varepsilon_{-\alpha} = -\varepsilon_\alpha$. Since \mathbb{F}_Θ is a homogeneous space of a complex Lie group it has a natural structure of a complex manifold. The associated integrable almost complex structure $J_\mathbb{C}$ is given by $\varepsilon_\alpha = 1$ if $\alpha > 0$. The conjugate structure $-J_\mathbb{C}$ is also integrable. A complete description of complex structures of flag manifolds can be found in [2].

4.2 Riemannian invariant metrics

Now we will describe Riemannian metrics $g(\cdot, \cdot)$ on \mathbb{F}_Θ which is J -invariant, i.e. $g(JX, JY) = g(X, Y)$ and also invariant by the action of U (or G). It is well known a Riemannian invariant metric on \mathbb{F}_Θ is completely determined by a real inner product $g(\cdot, \cdot)$ on $\mathfrak{m} = T_o\mathbb{F}_\Theta$ which is invariant by the adjoint

action of \mathfrak{k}_Θ . Besides that any real inner product $\text{ad}(\mathfrak{k}_\Theta)$ -invariant on \mathfrak{m} has the form

$$g(\cdot, \cdot) = - \sum_{\alpha \in R_\Theta^+} \lambda_\alpha B(\cdot, \cdot)|_{\mathfrak{u}_\alpha \times \mathfrak{u}_\alpha} \quad (7)$$

where the Killing form $B(\cdot, \cdot)$ is negative defined on $\mathfrak{m} \times \mathfrak{m}$ and $\lambda_\alpha > 0$.

Since \mathbb{F}_Θ is a reductive homogeneous space, i.e. $\text{ad}(\mathfrak{k}_\Theta)\mathfrak{m} \subset \mathfrak{m}$, we can decompose \mathfrak{m} into a sum of irreducible $\text{ad}(\mathfrak{k}_\Theta)$ submodules \mathfrak{m}_i of the module \mathfrak{m} :

$$\mathfrak{m} = \mathfrak{m}_1 \oplus \cdots \oplus \mathfrak{m}_s.$$

Thus a Riemannian invariant metric on \mathbb{F}_Θ is completely determined by a real inner product $g(\cdot, \cdot)$ on $\mathfrak{m} = T_o\mathbb{F}_\Theta$ which is invariant by the adjoint action of \mathfrak{k}_Θ . From Schur's Lemma any real inner product $\text{ad}(\mathfrak{k}_\Theta)$ -invariant on \mathfrak{m} is given by

$$g(\cdot, \cdot) = -\lambda_1 B(\cdot, \cdot)|_{\mathfrak{m}_1 \times \mathfrak{m}_1} - \cdots - \lambda_s B(\cdot, \cdot)|_{\mathfrak{m}_s \times \mathfrak{m}_s}. \quad (8)$$

Here $\lambda_i = \lambda_\alpha$ if $\mathfrak{u}_\alpha \subset \mathfrak{m}_i$, hence $\lambda_\alpha = \lambda_\beta$ if \mathfrak{u}_α and \mathfrak{u}_β belongs to the same irreducible $\text{ad}(\mathfrak{k}_\Theta)$ submodules \mathfrak{m}_i of the module \mathfrak{m} . We will call an inner product defined by equation (8) as a Riemannian invariant metric on \mathbb{F}_Θ . Any such inner product extends naturally to a symmetric bilinear form on the complexification $\mathfrak{m}^\mathbb{C}$ of \mathfrak{m} , we will denote this extension by the same notation $g(\cdot, \cdot)$.

Using the Weyl basis chosen it is easy to see that any iacs J is compatible with any invariant metric g on \mathbb{F}_Θ , i.e. $g(JX, JY) = g(X, Y)$, for any $X, Y \in \mathfrak{m}$. The pair (g, J) is called an *invariant almost Hermitian structure* on \mathbb{F}_Θ .

Remark 4.1. *By an easy computation, one see that an (invariant) J frame on \mathbb{F}_Θ is given by*

$$\left\{ A_\alpha / \sqrt{2\lambda_\alpha}, J(A_\alpha / \sqrt{2\lambda_\alpha}) = i\varepsilon_\alpha S_\alpha / \sqrt{2\lambda_\alpha}; \quad \alpha \in R_\Theta^+ \right\}.$$

5 The invariant Riemannian connection on \mathbb{F}_Θ

In this section we compute explicitly the invariant Levi-Civita connection on \mathbb{F}_Θ .

Recall the isomorphism $\mathfrak{m} \rightarrow T_o\mathbb{F}_\Theta$ which corresponds each $X \in \mathfrak{m}$ to the invariant vector field $X^* \in \mathfrak{X}(\mathbb{F}_\Theta)$:

$$X_o^* = \frac{d}{dt}(\exp(tX)) \cdot o|_{t=0}.$$

For each $X \in \mathfrak{m}$ the vector field X^* is a Killing vector field, since the Riemannian metric $g(\cdot, \cdot)$ is invariant. This is equivalent to

$$X^*g(Y, Z) = g([X^*, Y], Z) + g(Y, [X^*, Z]) \quad (9)$$

and

$$g(\nabla_Y X^*, Z) + g(\nabla_Z X^*, Y) = 0 \quad (10)$$

for all $Y, Z \in \mathfrak{X}(\mathbb{F}_\Theta)$, (see [15], pp.251).

Now recall that if $X, Y \in \mathfrak{m}$ then $[X^*, Y^*] = -[X, Y]^*$. Then using (9) and (10), it can be proved that the Riemannian connection is given by ([5], p. 183 or [3], p. 79) :

$$(\nabla_{X^*} Y^*)_o = -\frac{1}{2}[X, Y]_m + U(X, Y), \quad X, Y \in \mathfrak{m}$$

where $U: \mathfrak{m} \times \mathfrak{m} \rightarrow \mathfrak{m}$ is the symmetric bilinear map defined by

$$2g(U(X, Y), Z) = g([Z, X]_m, Y) + g(X, [Z, Y]_m). \quad (11)$$

We also denote $U(X, Y)$ its complexification $U: \mathfrak{m}^{\mathbb{C}} \times \mathfrak{m}^{\mathbb{C}} \rightarrow \mathfrak{m}^{\mathbb{C}}$. On the basis $\{X_\alpha\}$, $\alpha \in R_\Theta$, $U(X_\alpha, X_\beta) = \sum a_\gamma X_\gamma$ then

$$g(U(X_\alpha, X_\beta), X_{-\gamma}) = a_\gamma \lambda_\gamma.$$

Now $[X_\alpha, X_\beta]_m \neq 0 \iff \alpha + \beta \in R_\Theta = R \setminus R(\Theta)$ and in this case $[X_\alpha, X_\beta]_m = n_{\alpha, \beta} X_{\alpha + \beta}$.

Thus,

$$\begin{aligned} 2g(U(X_\alpha, X_\beta), X_{-\gamma}) &= g([X_{-\gamma}, X_\alpha]_m, X_\beta) + g(X_\alpha, [X_{-\gamma}, X_\beta]_m) \\ &= \begin{cases} n_{\alpha, \beta}(\lambda_\beta - \lambda_\alpha), & \text{if } \gamma = \alpha + \beta \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

Then

$$U(X_\alpha, X_\beta) = \begin{cases} \frac{n_{\alpha,\beta}}{2\lambda_{\alpha+\beta}}(\lambda_\beta - \lambda_\alpha)X_{\alpha+\beta}, & \text{if } \alpha + \beta \in R_\Theta \\ 0, & \text{otherwise.} \end{cases}$$

Thus we obtain the next result.

Proposition 5.1. *The Levi-Civita connection for invariant vector fields on \mathbb{F}_Θ is given by*

$$\begin{aligned} (\nabla_{X_\alpha^*} X_\beta^*)_o &= -\frac{1}{2}[X_\alpha, X_\beta]_m + U(X_\alpha, X_\beta) \\ &= \begin{cases} \frac{n_{\beta,\alpha}}{2\lambda_{\alpha+\beta}}(\lambda_{\alpha+\beta} + \lambda_\alpha - \lambda_\beta)X_{\alpha+\beta}, & \text{if } \alpha + \beta \in R_\Theta \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

For later using, we denote

$$r_{\alpha,\beta} = \frac{n_{\beta,\alpha}}{2\lambda_{\alpha+\beta}}(\lambda_{\alpha+\beta} + \lambda_\alpha - \lambda_\beta)$$

then

$$(\nabla_{X_\alpha^*} X_\beta^*)_o = r_{\alpha,\beta}X_{\alpha+\beta}. \quad (12)$$

A direct computation shows that

$$\begin{aligned} r_{\alpha,\beta} &= -r_{-\alpha,-\beta}; \\ r_{\alpha,\beta} + r_{\beta,\alpha} &= \frac{n_{\beta,\alpha}}{\lambda_{\alpha+\beta}}(\lambda_\alpha - \lambda_\beta). \end{aligned}$$

Lemma 5.2. *Let α, β be roots in R_Θ such that $\alpha + \beta$ and $\alpha - \beta$ are in R_Θ . Then*

- i) $\nabla_{S_\beta} A_\alpha = r_{\beta,\alpha}S_{\alpha+\beta} - r_{\beta,-\alpha}S_{\beta-\alpha}$
- ii) $\nabla_{S_\beta} S_\alpha = r_{\beta,\alpha}A_{\beta+\alpha} + r_{\beta,-\alpha}A_{\beta-\alpha}$
- iii) $\nabla_{A_\beta} A_\alpha = r_{\beta,\alpha}S_{\beta+\alpha} - r_{\beta,-\alpha}A_{\beta-\alpha}$

$$iv) \nabla_{A_\beta} S_\alpha = r_{\beta,\alpha} A_{\beta+\alpha} + r_{\beta,-\alpha} S_{\beta-\alpha}$$

$$\text{where } r_{\beta,\alpha} = \frac{n_{\alpha,\beta}}{2\lambda_{\alpha+\beta}} (\lambda_{\alpha+\beta} + \lambda_\beta - \lambda_\alpha).$$

Proof. All items follows by a direct computations, using $A_\alpha = X_\alpha - X_{-\alpha}$, $S_\alpha = X_\alpha + X_{-\alpha}$ and equation (12). \square

6 Almost semi Kähler Flag manifolds

In this section we show the covariant derivative of the Kähler form Ω has a special property, as an immediate consequence one obtains that \mathbb{F}_Θ always belong to class ASK for any invariant almost Hermitian structure (g, J) .

Consider the codifferential of Ω on a almost Hermitian Riemannian manifold (M, g, J) :

$$(\delta\Omega)(X) = \sum_{i=1}^m ((\nabla_{E_i}\Omega)(E_i, X) + (\nabla_{JE_i}\Omega)(JE_i, X)),$$

where $\{E_1, \dots, E_m, JE_1, \dots, JE_m\}$ is a local orthonormal J -frame. Recall that (M, g, J) is called Almost Semi Kähler (ASK) if $\delta\Omega = 0$.

In the case of flag manifolds, given a iacs J on \mathbb{F}_Θ , from the invariance of the vector fields, from remark 4.1, a global orthonormal J -frame at the origin is given by

$$\left\{ V_\alpha = A_\alpha / \sqrt{2\lambda_\alpha}, J(V_\alpha) = J(A_\alpha / \sqrt{2\lambda_\alpha}) = i\varepsilon_\alpha S_\alpha / \sqrt{2\lambda_\alpha}; \quad \alpha \in R_\Theta^+ \right\}. \quad (13)$$

Theorem 6.1. *For any invariant almost Hermitian structure on \mathbb{F}_Θ , one has*

$$(\nabla_{V_\beta}\Omega)(V_\beta, X) = (\nabla_{JV_\beta}\Omega)(JV_\beta, X) = 0$$

where $X \in \mathfrak{m}$.

Proof. Using the linearity of the invariant metric, the iacs and the Riemmanian connection, it is easy to see that if

$$X = \sum_{\alpha \in R_\Theta^+} a_\alpha V_\alpha + \sum_{\alpha \in R_\Theta^+} b_\alpha JV_\alpha$$

then

$$(\nabla_{V_\beta}\Omega)(V_\beta, X) = \sum_{\alpha \in R_\Theta^+} a_\alpha(\nabla_{V_\beta}\Omega)(V_\beta, V_\alpha) + \sum_{\alpha \in R_\Theta^+} b_\alpha(\nabla_{V_\beta}\Omega)(V_\beta, JV_\alpha)$$

$$(\nabla_{JV_\beta}\Omega)(JV_\beta, X) = \sum_{\alpha \in R_\Theta^+} a_\alpha(\nabla_{JV_\beta}\Omega)(JV_\beta, V_\alpha) + \sum_{\alpha \in R_\Theta^+} b_\alpha(\nabla_{JV_\beta}\Omega)(JV_\beta, JV_\alpha).$$

Now it is sufficient to see that each term in the four sum above is zero. In fact, since $(\nabla_X\Omega)(Y, Z) = g(Y, (\nabla_X J)Z)$, for $X, Y, Z \in \mathfrak{m}$, then

$$\begin{aligned} (\nabla_{V_\beta}\Omega)(V_\beta, V_\alpha) &= g(V_\beta, (\nabla_{V_\beta} J)V_\alpha) \\ &= g(V_\beta, \nabla_{V_\beta} JV_\alpha - J(\nabla_{V_\beta} V_\alpha)) \\ &= g\left(\frac{1}{\sqrt{2\lambda_\beta}}A_\beta, \frac{i}{2\sqrt{\lambda_\beta\lambda_\alpha}}\nabla_{A_\beta}S_\alpha - \frac{1}{2\sqrt{\lambda_\beta\lambda_\alpha}}J(\nabla_{A_\beta}A_\alpha)\right). \end{aligned}$$

From Lemma 5.2 the right side in last equation above is

$$\frac{1}{2\lambda_\beta\sqrt{2\lambda_\alpha}}\{(1 - \varepsilon_{\beta+\alpha})ir_{\beta,\alpha}g(A_\beta, A_{\beta+\alpha}) + (1 - \varepsilon_{\beta-\alpha})ir_{\beta,-\alpha}g(A_\beta, S_{\beta-\alpha})\} = 0,$$

because $g(A_\beta, A_{\beta+\alpha}) = 0 = g(A_\beta, S_{\beta-\alpha})$.

Analogously,

$$\begin{aligned} (\nabla_{V_\beta}\Omega)(V_\beta, JV_\alpha) &= \frac{i\varepsilon_\alpha}{2\lambda_\beta\sqrt{2\lambda_\alpha}}g(A_\beta, \nabla_{A_\beta}S_\alpha - J(\nabla_{A_\beta}S_\alpha)) \\ &= 0 \end{aligned}$$

since $g(A_\beta, \nabla_{A_\beta}S_\alpha) = g(A_\beta, J(\nabla_{A_\beta}S_\alpha)) = 0$. Using Lemma 5.2, by a similar computation, one obtains:

$$(\nabla_{JV_\beta}\Omega)(JV_\beta, V_\alpha) = 0 = (\nabla_{JV_\beta}\Omega)(JV_\beta, JV_\alpha).$$

□

Corollary 6.2. *Any invariant almost Hermitian structure on \mathbb{F}_Θ belongs to the almost semi Kähler class (ASK).*

7 Invariant almost Hermitian submanifolds of \mathbb{F}_Θ

Let $\mathbb{F}_\Theta = U/K_\Theta$ be a flag manifold. Consider $L \subset U$ a $2r$ -dimensional closed (and then compact) subgroup of U . Then $L \cap K_\Theta$ is a closed subgroup of L and the coset space $L/L \cap K_\Theta$ is a submanifold of \mathbb{F}_Θ such that the following diagram of principal bundles is commutative (see [10], p. 125)

$$\begin{array}{ccc} L & \xhookrightarrow{i} & U \\ \downarrow \pi_1 & & \downarrow \pi \\ L/L \cap K_\Theta & \xrightarrow{I} & U/K_\Theta. \end{array}$$

where π_1 and π are the natural mappings of L onto $L/L \cap K_\Theta$ and π of U onto U/K_Θ , respectively. The inclusion mapping of L into U is denoted by i and $I: L/L \cap K_\Theta \rightarrow U/K_\Theta$ denotes the mapping $l(L \cap U_\Theta) \mapsto i(l)K_\Theta$. We denote by \mathfrak{l} the Lie algebra of L . Since \mathfrak{l} is a subalgebra of the compact Lie algebra \mathfrak{u} , the Killing form $B(\cdot, \cdot)$ is negative definite on \mathfrak{l} . From the decomposition $\mathfrak{u} = \mathfrak{k}_\Theta \oplus \mathfrak{m}$ we obtain

$$\mathfrak{l} = \mathfrak{l} \cap \mathfrak{k}_\Theta \oplus \mathfrak{n}$$

where $\mathfrak{n} \subset \mathfrak{m}$ is the orthogonal complement of $\mathfrak{l} \cap \mathfrak{k}_\Theta$ in \mathfrak{l} with respect to $-B(\cdot, \cdot)$.

The U -invariance of $g(\cdot, \cdot)$ implies $g([X, Y], Z) = g(X, [Y, Z])$, for any X, Y, Z in \mathfrak{u} . In particular, for $X, Y \in \mathfrak{l} \cap \mathfrak{k}_\Theta$ and $Z \in \mathfrak{n}$, we have $0 = g([X, Y], Z) = g(X, [Y, Z])$, then $\text{ad}(\mathfrak{l} \cap \mathfrak{k}_\Theta)\mathfrak{n} \subset \mathfrak{n}$. The real tangent $T_o(L/L \cap K_\Theta)$ at the origin $o = L \cap K_\Theta$ is isomorphic to the real subspace \mathfrak{n} .

Lemma 7.1. *Let $(\mathbb{F}_\Theta = U/K_\Theta, J, g)$ be an invariant almost Hermitian flag manifold and $L/L \cap K_\Theta$ as above, then $J(\mathfrak{n}) = \mathfrak{n}$.*

Proof. Consider the following subset of the complementary roots

$$R_{\mathfrak{n}} = \left\{ \alpha \in R_\Theta^+ : g(X, V_\alpha) \neq 0, \quad \text{or} \quad g(X, JV_\alpha) \neq 0 \quad \text{for some} \quad X \in \mathfrak{n} \right\} \quad (14)$$

where the vectors V_α and JV_α were defined in (13).

Then the following subset of orthonormal J-frame $\{V_\alpha, J(V_\alpha); \alpha \in R_{\mathfrak{n}}\}$ span the tangent space \mathfrak{n} , because if $X \in \mathfrak{n}$,

$$X = \sum_{\alpha \in R_{\mathfrak{n}}} a_\alpha V_\alpha + \sum_{\alpha \in R_{\mathfrak{n}}} b_\alpha J V_\alpha$$

where $a_\alpha = g(X, V_\alpha)$ and $b_\alpha = g(X, J V_\alpha)$ are the real coefficients of this linear combination.

Thus,

$$JX = \sum_{\alpha \in R_{\mathfrak{n}}} a_\alpha J V_\alpha - \sum_{\alpha \in R_{\mathfrak{n}}} b_\alpha V_\alpha$$

with $\alpha \in R_{\mathfrak{n}}$. Now, for $\alpha \in R_\Theta^+$ and $L \in \mathfrak{k}_\Theta$,

$$g(V_\alpha, L) = 0 = g(J V_\alpha, L)$$

which is valid, in particular, for $\alpha \in R_{\mathfrak{n}} \subset R_\Theta^+$ and $L \in \mathfrak{l} \cap \mathfrak{k}_\Theta$. Then $g(JX, L) = 0$ for any $L \in \mathfrak{l} \cap \mathfrak{k}_\Theta$ and $J(\mathfrak{n}) \subset \mathfrak{n}$. Since $X = J(-JX)$, one has $\mathfrak{n} \subset J(\mathfrak{n})$. This proves that $J(\mathfrak{n}) = \mathfrak{n}$. \square

In the sequel we continue denoting by J and g their restriction to \mathfrak{n} .

Let $(\mathbb{F}_\Theta = U/K_\Theta, J, g)$ be an invariant almost Hermitian flag manifold, from the above Lemma, the restrictions of J and g to $\mathfrak{n} \subset \mathfrak{m}$ define an invariant almost Hermitian structure on $L/L \cap K_\Theta$ in a such way that $(L/L \cap K_\Theta, J, g)$ is an holomorphic submanifold of \mathbb{F}_Θ .

Now, since the vector fields are invariants, we have the following isomorphisms:

$$\begin{aligned} \mathfrak{X}(\mathbb{F}_\Theta) &= \mathfrak{m} = \tilde{\mathfrak{X}}(\mathbb{F}_\Theta), & \mathfrak{X}(L/L \cap K_\Theta) &= \mathfrak{n} \\ \mathfrak{X}(L/L \cap K_\Theta)^\perp &= \mathfrak{n}^\perp = \{X \in \mathfrak{m}; g(X, \mathfrak{n}) = 0\} \end{aligned}$$

and the equation (1) becomes

$$\mathfrak{m} = \mathfrak{n} \oplus \mathfrak{n}^\perp.$$

Lemma 7.2. *The orthogonal complement \mathfrak{n}^\perp admits a g -orthonormal basis $\{V_\alpha, J V_\alpha\}$, with $\alpha \in R_\Theta - R_{\mathfrak{n}}$.*

Proof. From the definition (14), if $\alpha \in R_\Theta - R_n$ then $g(X, V_\alpha) = g(X, JV_\alpha) = 0$, for all $X \in \mathfrak{n}$, hence $\{V_\alpha, JV_\alpha\} \subset \mathfrak{n}^\perp$. Note that any $Y \in \mathfrak{m}$ can be written as

$$\begin{aligned} Y &= Y_n + Y_{n^\perp} \\ &= \sum_{\alpha \in R_n} (a_\alpha V_\alpha + b_\alpha JV_\alpha) + \sum_{\alpha \in R_\Theta - R_n} (c_\alpha V_\alpha + d_\alpha JV_\alpha). \end{aligned}$$

By definition of R_n the g -orthogonal projection of Y on \mathfrak{n} is given by

$$Y_n = \sum_{\alpha \in R_n} (a_\alpha V_\alpha + b_\alpha JV_\alpha)$$

where $a_\alpha = g(Y, V_\alpha)$ and $b_\alpha = g(Y, JV_\alpha)$, then

$$Y_{n^\perp} = \sum_{\alpha \in R_\Theta - R_n} (c_\alpha V_\alpha + d_\alpha JV_\alpha).$$

Thus the subset of the J-frame $\{V_\alpha, JV_\alpha; \alpha \in R_\Theta - R_n\}$, spans \mathfrak{n}^\perp . Since we have an g -orthonormal subset, it is an invariant basis for \mathfrak{n}^\perp .

It remains to prove that $R_\Theta - R_n \neq \emptyset$. Above we proved that if $Y_{n^\perp} \neq 0$ then $R_\Theta - R_n \neq \emptyset$. Thus, if $R_\Theta = R_n$ then $\mathfrak{n}^\perp = 0$ and $L/L \cap K_\Theta$ has codimension 0. So $R_\Theta - R_n \neq \emptyset$. \square

Remark 7.3. *The subset $\{V_\alpha, JV_\alpha\}$, with $\alpha \in R_n$ of the invariant J-frame spans the space \mathfrak{n} but may not be a basis because it is possible that some of these vectors do not belong to \mathfrak{n} . We will see in next section that when $L/L \cap K_\Theta$ is also a flag manifold, this subset of the invariant J-frame becomes a basis of the tangent space \mathfrak{n} .*

Theorem 7.4. *Let $(\mathbb{F}_\Theta, J, g)$ be an invariant almost Hermitian flag manifold (which belong to the class ASK). Then any invariant holomorphic submanifold $L/L \cap K_\Theta$ of $(\mathbb{F}_\Theta, J, g)$ belongs to the class ASK. Moreover, $L/L \cap K_\Theta$ is necessarily minimal.*

Proof. Consider the invariant J-frame $\{V_\alpha, JV_\alpha\}$, $\alpha \in R_\Theta - R_n$, on \mathfrak{n}^\perp . For $X \in \mathfrak{n} \subset \mathfrak{m}$, the equation (4) sets:

$$(\delta\Omega)(X) = (\delta'\Omega)(X) + (\bar{\delta}\Omega)(X).$$

Now, by Corollary 6.2, $(\delta\Omega)(X) = 0$ and from Theorem 6.1 $(\bar{\delta}\Omega)(X) = 0$, computed on the invariant J-frame $\{V_\alpha, JV_\alpha\}$, $\alpha \in R_\Theta - R_n$. So $L/L \cap K_\Theta$ belongs to the class ASK.

To see that $L/L \cap K_\Theta$ is minimal, let $\{E_1, \dots, E_r, V_{\alpha_1}, \dots, V_{\alpha_s}, JE_1, \dots, JE_r, JV_{\alpha_1}, \dots, JV_{\alpha_s}\}$ be an adapted invariant orthonormal J-frame of \mathfrak{m} , such that $\{E_1, \dots, E_r, JE_1, \dots, JE_r\}$ is an orthonormal J-frame of \mathfrak{n} . For $X \in \mathfrak{m}$, we can write $X = X_n + X_{n^\perp}$ such that $X_n \in \mathfrak{n}$ and $X_{n^\perp} \in \mathfrak{n}^\perp$. Thus using Theorem 6.1, we have

$$\begin{aligned}
0 = (\delta\Omega)(X) &= \sum_{i=1}^r ((\nabla_{E_i}\Omega)(E_i, X) + (\nabla_{JE_i}\Omega)(JE_i, X)) \\
&+ \sum_{j=1}^s \underbrace{((\nabla_{V_{\alpha_j}}\Omega)(V_{\alpha_j}, X) + (\nabla_{JV_{\alpha_j}}\Omega)(JV_{\alpha_j}, X))}_{=0} \\
&= \sum_{i=1}^r ((\nabla_{E_i}\Omega)(E_i, X_n + X_{n^\perp}) + (\nabla_{JE_i}\Omega)(JE_i, X_n + X_{n^\perp})) \\
&= (\delta'\Omega)(X_n) + (\bar{\delta}\Omega)(X_{n^\perp})
\end{aligned}$$

From (3) we get that $(L/L \cap K_\Theta, J, g)$ is minimal since $(\delta'\Omega)(X_n) = 0$. \square

Corollary 7.5. *If $(\mathbb{F}_\Theta, J, g)$ belongs to the class SK. Then any invariant holomorphic submanifold $L/L \cap K_\Theta$ of \mathbb{F}_Θ belongs to the class SK. Moreover, $L/L \cap K_\Theta$ is necessarily minimal.*

8 Almost Hermitian flag submanifolds of \mathbb{F}_Θ

Let $\mathbb{F}_\Theta = G/P_\Theta$ be a flag manifold and L a Lie subgroup semisimple of G . Then $Q := L \cap P_\Theta$ is a parabolic subgroup of L .

Thus the flag manifold L/Q is a submanifold of \mathbb{F}_Θ such that the following diagram of principal bundles is commutative

$$\begin{array}{ccc}
L & \hookrightarrow & G \\
\downarrow Q & & \downarrow P_\Theta \\
L/Q & \hookrightarrow & G/P_\Theta
\end{array}$$

In terms of Lie theory we can obtain a submanifold L/Q of \mathbb{F}_Θ as following: let $\Theta' \subset \Sigma$ be a non empty subset of simple roots such that $\Theta' \not\subset \Theta$ and denote by $R' := \langle \Theta' \rangle \cap R$ the roots spanned by Θ' . Consider the following semisimple Lie subalgebra of \mathfrak{g} :

$$\mathfrak{l} = \mathfrak{h}' \oplus \sum_{\alpha \in R'} \mathfrak{g}_\alpha$$

where $\mathfrak{h}' = \sum_{\alpha \in R'} \mathbb{C}H_\alpha \subset \mathfrak{h}$ is a Cartan subalgebra of \mathfrak{l} such that R' is the root system associated to $(\mathfrak{l}, \mathfrak{h}')$. The following subalgebra

$$\mathfrak{q}_{\Theta'} = \mathfrak{l} \cap \mathfrak{p}_\Theta = \mathfrak{h}' \oplus \sum_{\alpha \in R'^+} \mathfrak{g}_\alpha \oplus \sum_{\alpha \in R'^- \cap R(\Theta)^-} \mathfrak{g}_\alpha$$

is a parabolic subalgebra in \mathfrak{l} because it contains $\mathfrak{b}' = \mathfrak{h}' \oplus \sum_{\alpha \in R'^+} \mathfrak{g}_\alpha$ which is a Borel subalgebra of \mathfrak{l} . Let L be the connected Lie group with Lie algebra \mathfrak{l} and denote by $\mathbb{F}_{\Theta'} = \{Ad_L(l)\mathfrak{q}_{\Theta'} : l \in L\}$ the set of subalgebras of \mathfrak{l} conjugate to $\mathfrak{q}_{\Theta'}$. Thus $\mathbb{F}_{\Theta'} = L/Q_{\Theta'}$, where $Q_{\Theta'} = \{l \in L; Ad_L(l)\mathfrak{q}_{\Theta'} = \mathfrak{q}_{\Theta'}\}$ is the normalizer of $\mathfrak{q}_{\Theta'}$ in L .

We can see $\mathbb{F}_{\Theta'}$ as a compact homogeneous space following a similar way used to obtain $\mathbb{F}_\Theta = U/K_\Theta$. Using the Weyl basis fixed previously, we see that

$$\mathfrak{v} = i\mathfrak{h}'_{\mathbb{R}} \oplus \sum_{\alpha \in R'^+} \mathfrak{u}_\alpha$$

is a compact real form of \mathfrak{l} , where $\mathfrak{h}'_{\mathbb{R}}$ denotes the real space vector spanned by $\{H_\alpha : \alpha \in R'\}$. Thus $V = \exp \mathfrak{v}$ is the compact real form of L corresponding to \mathfrak{v} . Restricting the action of L on $\mathbb{F}_{\Theta'}$ we see that

$$\mathbb{F}_{\Theta'} = V/K_{\Theta'}$$

where $K_{\Theta'} = V \cap Q_{\Theta'}$. The Lie algebra of $K_{\Theta'}$ is the fixed point of the conjugate $\tau: X_\alpha \mapsto -X_{-\alpha}$ of \mathfrak{l} restricted to $\mathfrak{q}_{\Theta'}$, given by

$$\mathfrak{k}_{\Theta'} = \mathfrak{v} \cap \mathfrak{q}_{\Theta'} = i\mathfrak{h}'_{\mathbb{R}} \oplus \sum_{\alpha \in R'^+ \cap R(\Theta)^+} \mathfrak{u}_\alpha.$$

Thus the real tangent space of $\mathbb{F}_{\Theta'}$ at the origin is identified with

$$T_o(V/K_{\Theta'}) = \mathfrak{n} = \sum_{\alpha \in R'^+ \setminus R(\Theta)^+} \mathfrak{u}_\alpha,$$

and its complexification is

$$\mathfrak{n}^{\mathbb{C}} = \mathfrak{n} \otimes \mathbb{C} = \sum_{\alpha \in R' \setminus R(\Theta)} \mathbb{C}X_{\alpha}.$$

Note that $R' \setminus R(\Theta) \neq \emptyset$, since $\Theta' \not\subset \Theta$. Of course $\mathfrak{n}^{\mathbb{C}}$ is a vector subspace of $\mathfrak{m}^{\mathbb{C}}$ and

$$\mathfrak{m}^{\mathbb{C}} - \mathfrak{n}^{\mathbb{C}} = \sum_{\alpha \in R_{\Theta} \setminus R'} \mathbb{C}X_{\alpha}.$$

Then $\mathbb{F}_{\Theta'}$ is a submanifold of \mathbb{F}_{Θ} of codimension $|R_{\Theta}| - |R' \cap R_{\Theta}| = |R| - |R(\Theta)| - |R' \cap R_{\Theta}|$.

From the previous section, it is clear that any invariant almost Hermitian structure (g, J) on \mathbb{F}_{Θ} can be restrict to $\mathbb{F}_{\Theta'}$ in a such way that $\mathbb{F}_{\Theta'}$ becomes an invariant holomorphic submanifold of \mathbb{F}_{Θ} . We will keep the same notation for g and J on $\mathbb{F}_{\Theta'}$.

Next, we denote by $Z_{\mathfrak{m}}$ (resp. $Z_{\mathfrak{n}}$) the component of $Z \in \mathfrak{g}$ in the subspace \mathfrak{m} (resp. \mathfrak{n}).

Proposition 8.1. *If $X, Y \in \mathfrak{n}$ then $[X, Y]_{\mathfrak{n}} = [X, Y]_{\mathfrak{m}}$.*

Proof. Since $[X, Y]_{\mathfrak{n}} = [X, Y]_{\mathfrak{m}} \iff [X, Y]_{\mathfrak{n}^{\mathbb{C}}} = [X, Y]_{\mathfrak{m}^{\mathbb{C}}}$ we just need to prove that $[X_{\alpha}, X_{\beta}]_{\mathfrak{n}^{\mathbb{C}}} = [X_{\alpha}, X_{\beta}]_{\mathfrak{m}^{\mathbb{C}}}$, for $\alpha, \beta \in R' \setminus R(\Theta)$.

If $\alpha, \beta \in R' \setminus R(\Theta)$ then $\alpha + \beta \in R$ or $\alpha + \beta \notin R$. If $\alpha + \beta \notin R$ then $[X_{\alpha}, X_{\beta}] = 0$, in particular $[X_{\alpha}, X_{\beta}]_{\mathfrak{n}^{\mathbb{C}}} = [X_{\alpha}, X_{\beta}]_{\mathfrak{m}^{\mathbb{C}}}$. If $\alpha + \beta \in R$ then $\alpha + \beta \in R'$, because R' is a root system. Now $\alpha + \beta \in R' \cap R(\Theta)$ or $\alpha + \beta \in R' \setminus R(\Theta)$. In the first case $[X_{\alpha}, X_{\beta}]_{\mathfrak{n}^{\mathbb{C}}} = 0 = [X_{\alpha}, X_{\beta}]_{\mathfrak{m}^{\mathbb{C}}}$. In the order case, $[X_{\alpha}, X_{\beta}]_{\mathfrak{n}^{\mathbb{C}}} = n_{\alpha, \beta} X_{\alpha + \beta} = [X_{\alpha}, X_{\beta}]_{\mathfrak{m}^{\mathbb{C}}}$. \square

Example 8.2. *Consider $\mathfrak{g} = \mathfrak{sl}(8, \mathbb{C})$. A Cartan subalgebra of $\mathfrak{sl}(8, \mathbb{C})$ has the form*

$$\mathfrak{h} = \left\{ \text{diag}(\varepsilon_1, \dots, \varepsilon_8) : \varepsilon_i \in \mathbb{C} \text{ and } \sum_{i=1}^8 \varepsilon_i = 0 \right\}.$$

The root system of $(\mathfrak{sl}(8, \mathbb{C}), \mathfrak{h})$ is given by linear maps $\alpha_{i,j} : \text{diag}(\varepsilon_1, \varepsilon_2, \dots, \varepsilon_8) \mapsto \varepsilon_i - \varepsilon_j$. The root system is

$$R = \{\pm\alpha_{i,j}, 1 \leq i < j \leq 8\}$$

and a choice of positive roots and simple roots associated are given by

$$R^+ = \{\alpha_{i,j}, 1 \leq i < j \leq 8\}, \quad \Sigma = \{\alpha_{i,i+1}, 1 \leq i \leq 8\}.$$

Now let $\Theta = \{\alpha_{1,2}, \alpha_{2,3}, \alpha_{5,6}\}$ and $\Theta' = \{\alpha_{1,2}, \alpha_{2,3}, \alpha_{3,4}, \alpha_{7,8}\}$ be subsets of Σ . Note that $\Theta' \not\subseteq \Theta$ and the Dynkin diagram of Θ' is not connected.

We have

$$R(\Theta) := \langle \Theta \rangle \cap R = \{\pm\alpha_{1,2}, \pm\alpha_{2,3}, \pm\alpha_{1,3}, \pm\alpha_{5,6}\}$$

$$\mathfrak{p}_\Theta := \mathfrak{b}^+ \oplus \mathfrak{g}_{\alpha_{2,1}} \oplus \mathfrak{g}_{\alpha_{3,2}} \oplus \mathfrak{g}_{\alpha_{3,1}} \oplus \mathfrak{g}_{\alpha_{6,5}},$$

where $\mathfrak{b}^+ = \mathfrak{h} \oplus \sum_{\alpha \in R^+} \mathfrak{g}_\alpha$ denotes the Borel subalgebra.

The real form compact of \mathfrak{p}_Θ is isomorphic to

$$\mathfrak{k}_\Theta = \mathfrak{su}(8) \cap \mathfrak{p}_\Theta = \mathfrak{s}(\mathfrak{u}(3) \times \mathfrak{u}(2) \times \mathfrak{u}(1)^3)$$

and $\mathbb{F}_\Theta = SU(8)/S(U(3) \times U(2) \times U(1) \times U(1) \times (1))$. Now

$$R' := \langle \Theta' \rangle \cap R = \{\pm\alpha_{1,2}, \pm\alpha_{2,3}, \pm\alpha_{1,3}, \pm\alpha_{3,4}, \pm\alpha_{1,4}, \pm\alpha_{2,4}, \pm\alpha_{7,8}\}.$$

A basis for a Cartan subalgebra associated to root system R' is given by the matrix of the form

$$H_{\alpha_{i,j}} = E_{ii} - E_{jj} \quad 1 \leq i \neq j \leq 4 \quad \text{and} \quad 7 \leq i \neq j \leq 8$$

where E_{rs} denotes the matrix 8×8 with 1 at position (r, s) and zeros everywhere else. Thus

$$\mathfrak{l} = \mathfrak{h}' \oplus \sum_{\alpha \in R'} \mathfrak{g}_\alpha = \mathfrak{sl}(4, \mathbb{C}) \times \mathfrak{sl}(2, \mathbb{C})$$

and the real compact form of \mathfrak{l} is

$$\mathfrak{v} = i\mathfrak{h}'_{\mathbb{R}} \oplus \sum_{\alpha \in R'^+} \mathfrak{u}_\alpha = \mathfrak{s}(\mathfrak{u}(4)) \times \mathfrak{s}(\mathfrak{u}(2)).$$

The following is a parabolic subalgebra of \mathfrak{l}

$$\mathfrak{q}_{\Theta'} = \mathfrak{l} \cap \mathfrak{p}_\Theta = \mathfrak{h}' \oplus \sum_{\alpha \in R'^+} \mathfrak{g}_\alpha \oplus \sum_{\alpha \in R'^- \cap R(\Theta)^-} \mathfrak{g}_\alpha = \mathfrak{h}' \oplus \sum_{\alpha \in R'^+} \mathfrak{g}_\alpha \oplus \mathfrak{g}_{\alpha_{2,1}} \oplus \mathfrak{g}_{\alpha_{3,2}} \oplus \mathfrak{g}_{\alpha_{3,1}}$$

its real compact form is given by

$$\mathfrak{k}_{\Theta'} = \mathfrak{v} \cap \mathfrak{q}_{\Theta'} = i\mathfrak{h}'_{\mathbb{R}} \oplus \sum_{\alpha \in R'^+ \cap R(\Theta)^+} \mathfrak{u}_\alpha = \mathfrak{s}(\mathfrak{u}(3) \times \mathfrak{u}(1)) \times \mathfrak{s}(\mathfrak{u}(1) \times \mathfrak{u}(1)).$$

Then $\mathbb{F}_{\Theta'} = S(U(4)) \times S(U(2))/S(U(3) \times U(1)) \times S(U(1) \times U(1))$. Thus we obtain the following commutative diagram

$$\begin{array}{ccc}
S(U(4)) \times S(U(2)) & \hookrightarrow & SU(8) \\
\downarrow & & \downarrow \\
\frac{S(U(4)) \times S(U(2))}{S(U(3) \times U(1)) \times S(U(1) \times U(1))} & \hookrightarrow & \frac{SU(8)}{S(U(3) \times U(2) \times U(1)^3)}.
\end{array}$$

Theorem 8.3. *If $(\mathbb{F}_\Theta, g, J)$ belongs to the class ASK or SK , then any holomorphic flag submanifold $\mathbb{F}_{\Theta'}$ of \mathbb{F}_Θ belongs to the same class. Moreover, $\mathbb{F}_{\Theta'}$ is a totally geodesic submanifold of \mathbb{F}_Θ with respect to any invariant metric g on \mathbb{F}_Θ .*

Proof. The first part follows immediately from Theorem 7.4 and Corollary 7.5.

Now, for $X, Y \in \mathfrak{n}$, from Gauss formula, the second fundamental form is given by:

$$\begin{aligned}
\alpha(X, Y) &= \nabla_X Y - \nabla'_X Y \\
&= -\frac{1}{2}[X, Y]_{\mathfrak{m}} + U(X, Y) + \frac{1}{2}[X, Y]_{\mathfrak{n}} - U'(X, Y).
\end{aligned}$$

where $\nabla'_X Y$ is the tangential component of $\alpha(X, Y)$ (hence the connection on $\mathbb{F}_{\Theta'}$) and $U': \mathfrak{n} \times \mathfrak{n} \rightarrow \mathfrak{n}$ is the symmetric bilinear map defined by

$$2g(U'(X, Y), Z) = g([Z, X]_{\mathfrak{n}}, Y) + g(X, [Z, Y]_{\mathfrak{n}}), \quad X, Y, Z \in \mathfrak{n}.$$

Using the Proposition 8.1, we see $U|_{\mathfrak{n} \times \mathfrak{n}} = U'$ and $\alpha(X, Y) = 0$.

□

References

- [1] Alves, L. A. ; Da Silva, N. P.: Invariant \mathcal{G}_1 structures on flag manifolds, *Geom. Dedicata*, Vol.213, 227-243, (2021).
- [2] Alekseevsky, D. V.; Arvanitoyeorgos, A.: Riemannian flag manifolds with homogeneous geodesics. *Trans. Amer. Math. Soc.* 359, 3769-3789 (2007)
- [3] Arvanitoyeorgos, A.: An Introduction to Lie Groups and the Geometry of Homogeneous Spaces, *STML 22 Amer. Math. Soc.*, (2003).

- [4] Arvanitoyeorgos, A.: Geometry of Flag Manifolds, International Journal of Geometric Methods in Modern Physics. Vol. 3, No. 05n06, 957-974 (2006).
- [5] Besse, A.L.: Einstein Manifolds. Springer-Verlag, (1987).
- [6] Burstall, Francis E. ; Calderbank, David M. J.: Submanifold geometry in generalized flag manifolds. Rend. Circ. Mat. Palermo (2) Suppl., 72, 13-41, (2004).
- [7] Gray, A.: Minimal varieties and almost Hermitian manifolds. Mich. Math. J., 12, 273-287, (1965).
- [8] Gray, A.; Hervella, L. M.: The sixteen classes of almost Hermitian manifolds and their linear invariants, Ann. Mat. Pura Appl. 123, 35-58 (1980)
- [9] Kobayashi, S.; Nomizu, K.: Foundations of Differential Geometry Vol. II. Wiley Classics Library, (1969).
- [10] Helgason S.: Differential geometry, Lie groups, and symmetric spaces. Providence, R.I.: American Mathematical Society, (2001).
- [11] Falcitelli, M.;Pastore, A. M.; Ianus, S.. Riemannian Submersions and Related Topics. [s.l.] World Scientific, (2004).
- [12] Negreiros, C. J. C.; San Martin, L. A. B.: Invariant almost Hermitian structures on flag manifolds, Advances in Mathematics, 178, 277-310 (2003)
- [13] de Rham, G.; On the area of complex manifolds, Notes, Institute for Advanced Princeton, (1957).
- [14] San Martin, L. A. B.; Silva, R. de C. J.: Invariant nearly-Kähler structures, Geometriae Dedicata, Vol. 121, 1, 143-154 (2006)
- [15] O’Neil, B.: Semi-Riemannian Geometry with Applications to Relativity, Academic Press, United Kingdom, (1983).
- [16] Watson, B. Almost Hermitian submersions. J. Diff. Geom., 11, 147-165, 1976.