

## REFLECTION POSITIVITY AND CHERN-SIMONS FUNCTIONAL INTEGRALS

JONATHAN WEITSMAN

ABSTRACT. We show that a mathematical version of the formal Chern-Simons functional integral of [26] for manifolds equipped with a reflection may be constructed in terms of a reflection positive functional, associated to the quadratic term in the Chern-Simons Lagrangian, on an algebra of functions on a Banach space  $\mathbf{A}$  of connections on the underlying 3-manifold. This construction yields a Hilbert space associated to a surface preserved by the reflection. A version of the cubic Bosonic interaction term in the Chern-Simons Lagrangian gives a self-adjoint operator on this Hilbert space, and by exponentiation, a unitary one parameter subgroup of operators. The vacuum expectation value of this one parameter subgroup is combined with an additional term associated to the ghost fields and their interaction, and an appropriate weak limit gives a partition function for the quantum field theory. This construction is nonperturbative. The theory is finite and does not require renormalization, as may be expected from perturbation theory [3]. It is natural to ask whether the resulting partition function is related to the manifold invariants of Witten and Reshetikhin-Turaev, or whether a more elaborate construction may be needed.

## CONTENTS

1. Introduction	2
1.1. Some Results from Constructive Quantum Field Theory	2
1.2. The Chern Simons functional integral	6
2. Manifolds with reflection	11
3. The configuration spaces $\mathcal{F}^B$ and $\mathcal{F}^F$	13
3.1. The one-particle configuration spaces $\mathcal{F}_1^B$ and $\mathcal{F}_1^F$	14
3.2. The Bose configuration space $\mathcal{F}_+^B$	14
3.3. The configuration space $\mathcal{F}^B(U)$ associated to $U \subset M$	15
3.4. The Fermionic configuration Space $\mathcal{F}^F$	16
4. Reflection Positivity and the Bosonic Hilbert Space	17
4.1. Reflection positivity for the one particle configuration space $\mathcal{F}_{1,+}^B$	18
4.2. Reflection positivity for $\mathcal{F}_+^B$	22
4.3. Is there a relation between $\mathcal{F}_+^B$ and $H^B(\Sigma)$ ?	27
5. The Fermion Hilbert space	28
5.1. The one particle configuration space $\mathcal{F}_1^F(U)$	29
5.2. Reflection positivity and $\mathcal{F}^F$	33
5.3. The Hilbert space $H(\Sigma)$	36
6. The interacting Theory	37
6.1. The Boson interaction term	39
6.2. The Fermion interaction term	40
6.3. The Partition function	41

Supported in part by a grant from the Simons Foundation (# 579801 to J.W.).  
June 19, 2024.

7.	Concluding remarks and speculations	42
7.1.	A finite dimensional expression that may be related to the Chern Simons path integral	43
7.2.	Relation to topological quantum field theory	44
7.3.	String Field Theory	44
	References	44
	Appendix A. Formulas for the Multiplication action	46

## 1. INTRODUCTION

Let  $M$  be a compact oriented 3-manifold, let  $G$  be a compact Lie group, and let  $k \in \mathbb{Z}$ . In [26] E. Witten showed that a formal functional integral

$$(1.1) \quad Z_k(M) = \int_{\mathcal{A}} dA e^{ik\text{CS}(A)}$$

of the exponential of the Chern-Simons functional over the space  $\mathcal{A}$  of connections on the trivial principal  $G$ -bundle on  $M$  should be interpreted as yielding a topological invariant of the three manifold  $M$ , and that integrals of certain functions in this formal functional integral should be interpreted as link invariants, which in the case  $M = S^3$  and  $G = SU(n)$  are the Jones polynomials [15] of links.

Invariants satisfying the properties expected from Witten's results were constructed by Reshetikhin and Turaev [22] and a vast literature has grown in this field which provides evidence that properties expected from the formal functional integral and its formal perturbation expansion are satisfied by these invariants.

The purpose of this paper is to give a mathematically well-defined construction of a version of the functional integral (1.1), for manifolds  $M$  equipped with a reflection, in terms of a reflection positive functional on an algebra of functions on an appropriate space of connections. As in [25], we do not show that this functional is bounded, and hence we cannot use measure theory. This is different from the case of Bosonic quantum field theories, which correspond to measures on configuration spaces.<sup>1</sup>

We begin with a brief review of some results in Constructive Quantum Field Theory which provide both analogies and contrasts to the construction in this paper. We refer the reader to [9] for an overview of this field as well as for complete proofs. We then sketch the methods required to deal with functional integrals of the type (1.1), where reflection positivity is used as a substitute for measure theory.

**1.1. Some Results from Constructive Quantum Field Theory.** Choose a representation of  $G$  giving a trace which produces a nondegenerate inner product on the Lie algebra  $\mathfrak{g}$ .<sup>2</sup> The Chern-Simons invariant appearing in (1.1) has the form

$$(1.2) \quad \text{CS}(A) = \frac{1}{4\pi} \text{tr} \int_M AdA + \frac{2}{3} A^3$$

<sup>1</sup>There is some resemblance to the case of purely Fermionic quantum field theories.

<sup>2</sup>For simplicity, we may consider the case where  $M$  is a rational homology sphere,  $G$  is a compact classical Lie group, and the inner product is given by negative of the trace in the defining representation.

It is therefore a sum of a quadratic polynomial and a higher degree polynomial in the connection form  $A$ . Such expressions occur commonly in Quantum Field Theory. One starting point in constructive quantum field theory is to consider a quadratic term as giving a Gaussian measure on some appropriate Banach space; the higher degree terms can be “cut off” to give functions on that space, and an appropriate limit of the integrals of these functions is the desired non-Gaussian measure.

Given a separable Hilbert space  $\mathbf{H}$ , there exists a Gaussian measure  $d\mu$  on a Banach space  $\mathbf{B} \supset \mathbf{H}$  characterized by the integrals of bounded functions  $\exp(ih(\cdot))$ , where  $h \in \mathbf{B}^* \subset \mathbf{H}^*$ ;<sup>3</sup> namely,

$$(1.3) \quad \int_{\mathbf{B}} d\mu(\phi) e^{ih(\phi)} = e^{-\frac{1}{2}\|h\|_{\mathbf{H}^*}^2};$$

See e.g. [12]. Equivalently, this measure may be characterized by its value on some unbounded, but integrable functions, of the form

$$p_{a_1, \dots, a_k} : \mathbf{B} \longrightarrow \mathbb{R}$$

where  $a_1, \dots, a_k \in \mathbf{B}^*$ , given by

$$p_{a_1, \dots, a_k}(\phi) = a_1(\phi) a_2(\phi) \dots a_k(\phi);$$

we then have, for  $k$  even,

$$(1.4) \quad \int_{\mathbf{B}} d\mu(\phi) p_{a_1, \dots, a_k}(\phi) = \frac{1}{2^{k/2} (k/2)!} \sum_{\sigma \in \Sigma_k} \langle a_{\sigma(1)}, a_{\sigma(2)} \rangle_{\mathbf{H}^*} \dots \langle a_{\sigma(k-1)}, a_{\sigma(k)} \rangle_{\mathbf{H}^*},$$

where  $\Sigma_k$  is the permutation group on  $k$  letters. (If  $k$  is odd, the integral is zero.)

This measure is an infinite dimensional analog of the finite dimensional Gaussian measure

$$dm(x) = e^{-\frac{1}{2}\|x\|^2} \frac{d^n x}{(2\pi)^{\frac{n}{2}}}$$

on a Euclidean space  $\mathbb{R}^n$ , where  $d^n x$  is Lebesgue measure; in this case the measure is defined on  $\mathbb{R}^n$  and satisfies the analog of the property (1.3). In infinite dimensions, neither the Lebesgue measure nor the function  $e^{-\frac{1}{2}\|x\|^2}$  has a reasonable analog, but the Gaussian measure has a good generalization, though only as a measure on a Banach space  $\mathbf{B}$  in which  $\mathbf{H}$  is a subspace of measure zero.

The difficulties of quantum field theory, involving the appearance of divergences in computations of expectations in such functional integrals, arise from the fact that the Banach space on which the measure lives is larger than the Hilbert space  $\mathbf{H}$ .

More precisely, let  $\mathbf{H}$  be some appropriate Sobolev space of functions on a compact Riemannian manifold  $M$  of dimension  $d$ . In quantum mechanics and quantum field theory, the standard choice is the Sobolev space  $\mathbf{H} = H_1(M)$ . Then the Banach space  $\mathbf{B}$  can be

---

<sup>3</sup>Here we denote by  $\mathbf{H}^*$  the Banach space dual of the Hilbert space  $\mathbf{H}$ . Of course, this is isomorphic to  $\mathbf{H}$ . In this paper,  $\mathbf{H}$  will be a Sobolev space of differential forms on some manifold  $M$ , of Sobolev class  $-s$ , and  $\mathbf{H}^*$  will be a similar Sobolev space of forms of class  $s$ .

taken to be the Sobolev space  $\mathbf{B} = H_{1-\frac{d}{2}-\epsilon}(M)$  for any  $\epsilon > 0$ . If  $d = 1$ , this can be improved and the space  $\mathbf{B}$  can be taken to be a space of continuous functions of Lipschitz class  $\frac{1}{2} - \epsilon$ .<sup>4</sup> If  $d > 1$ , however, the distributions in  $\mathbf{B}$  are not functions, and polynomial functions of the type  $\int_M \phi^k$  do not make sense as functions on  $\mathbf{B}$ .

In the case  $d = 1$ , where  $M = S^1$  with the standard metric, this difficulty does not arise, and there is no difficulty in defining the integral

$$(1.5) \quad Z_{QM}(\lambda) = \int_{\mathbf{B}} d\mu(\phi) e^{-\lambda \int_M P(\phi(x)) dx}$$

where  $P$  is any polynomial which is bounded below and  $\lambda > 0$ . The Feynman-Kac formula then relates the integral  $Z(\lambda)$  to Schrodinger operators on the real line. We have

$$(1.6) \quad Z_{QM}(\lambda) = C \operatorname{tr} e^{-\left(-\frac{1}{2} \frac{d^2}{dx^2} + \frac{1}{2} x^2 + \lambda P(x)\right)}$$

where the trace is taken over  $L_2(\mathbb{R})$  and where  $C$  is a constant independent of  $\lambda$ .<sup>5</sup> For now we note only that the Hilbert space on which the Schrodinger operator acts is not the Hilbert space  $\mathbf{H}$  used to define the path integral. The difference between these two spaces will be important below.

In the case of manifolds  $M$  of dimension  $d > 1$ , in order to overcome the problem in defining polynomial functions on the space  $\mathbf{B}$ , both formal quantum field theory and constructive quantum field theory proceed by two steps: Regularization and Renormalization.

For  $\phi \in \mathbf{B}$  and  $\epsilon > 0$ , let  $\phi_\epsilon \in C^\infty(M)$  denote the smooth function

$$\phi_\epsilon(x) = (e^{-\epsilon(\Delta+1)}\phi)(x)$$

where  $\Delta$  denotes the positive Laplacian on  $M$ .

We can use these smoothings of the distributions in  $\mathbf{B}$  to define polynomial functions on  $\mathbf{B}$ . To take the simplest example, the function  $F_\epsilon : \mathbf{B} \rightarrow \mathbb{R}$  given by the polynomial

$$(1.7) \quad F_\epsilon(\phi) = \int_M \phi_\epsilon(x)^3 dx$$

is a well defined function on  $\mathbf{B}$ . This is "regularization".

Now we wish to take  $\epsilon \rightarrow 0$ . The family of functions  $F_\epsilon$  are all square integrable with respect to the measure  $d\mu$ , but they do not have a limit in  $L_2(d\mu)$ . However, let

$$(1.8) \quad : F_\epsilon(\phi) := F_\epsilon(\phi) - 3 \int_M C_\epsilon(x) \phi_\epsilon(x) dx,$$

where  $C_\epsilon(x) = (e^{-\epsilon(\Delta+1)} \frac{1}{\Delta+1} e^{-\epsilon(\Delta+1)})(x, x)$ . This is "renormalization".

<sup>4</sup>The precise choice of the Banach space  $\mathbf{B}$  does not play much of a role in the construction. One aspect of this fact is that if we consider two choices of Banach space given by  $\mathbf{B} = H_{1-\frac{d}{2}-\epsilon}(M)$  and  $\mathbf{B}' = H_{1-\frac{d}{2}-\epsilon'}(M)$ , where  $\epsilon > \epsilon'$ , the open sets in  $\mathbf{B}'$  arising from the inclusion  $\mathbf{B}' \rightarrow \mathbf{B}$  give a very different topology on  $\mathbf{B}'$  than its native topology, but the same Borel sets, and therefore the same measure theory.

<sup>5</sup>The quadratic term in the Schrodinger operator arises from the constant term added to the Laplacian in the definition of the Sobolev space. This constant term has to be added to avoid problems with the constant mode in the Fourier expansion of  $\phi$ , known in the Physics literature as an infrared divergence. Such issues, though inessential, cause nontrivial difficulties we will have to surmount later in this paper.

Suppose now that  $d = 2$ . A computation using (1.4) shows that

$$(1.9) \quad \|\cdot\| : F_\epsilon(\phi) : - : F_{\epsilon'}(\phi) : \|_{L_2(\mathbf{B}, d\mu)}^2 \longrightarrow 0$$

as  $\epsilon, \epsilon' \longrightarrow 0$ .<sup>6</sup> Hence the family  $: F_\epsilon :$  converges in  $L_2(\mathbf{B}, d\mu)$  to a limit function  $: F :$ , and therefore the exponential

$$(1.10) \quad e^{i\lambda:F:}$$

is a bounded function on  $\mathbf{B}$ , hence integrable, for any  $\lambda \in \mathbb{R}$ . The resulting integral

$$(1.11) \quad \int_{\mathbf{B}} d\mu(\phi) e^{i\lambda:F(\phi):}$$

is the mathematical expression corresponding to the formal functional integral

$$(1.12) \quad \int d\phi e^{-\int_M (\frac{1}{2}|d\phi(x)|^2 + \frac{1}{2}\phi(x)^2 - i\lambda\phi(x)^3) dx}.$$

This formal expression, where the Lagrangian is the sum of a quadratic term and an imaginary cubic, may be viewed as a rough analog of the formal Chern-Simons functional integral (1.1).

Returning for the moment to (1.11), we see that as long as we are interested in integrals of functions of the form (1.10)—in Physics language, with imaginary coupling constant—the existence of the limit function  $: F :$ , which is equivalent to the convergence of the expectation of the square of a polynomial function (in Physics language, finiteness up to second order in perturbation theory), is sufficient to guarantee the existence of the functional integral *nonperturbatively*.<sup>7</sup>

Note that formally, the function  $: F :$  is a linear combination (1.8) of polynomials with infinite coefficients. This is an example of the infinite constants associated to renormalization in quantum field theory. However, mathematically, there is nothing mysterious about the limit in (1.9).<sup>8</sup>

**Remark 1.13.** We have chosen to consider the case of a cubic polynomial with pure imaginary coefficient due to the resemblance to the Chern-Simons functional. For purposes of the Feynman-Kac formula, it is necessary as in (1.6) to consider polynomials which are bounded below, such as quartics with a positive real coefficient. Take  $M = S^1 \times S^1$  in the flat metric. If we take

$$F_\epsilon(\phi) = \int_M \phi_\epsilon(x)^4 dx,$$

then the appropriate renormalization is

$$: F_\epsilon(\phi) := \int_M \phi_\epsilon(x)^4 dx - 6C_\epsilon \int_M \phi_\epsilon(x)^2 + 3C_\epsilon^2.$$

<sup>6</sup>This amounts to convergence of the value of the  $\Theta$  graph in second order perturbation theory. This in turn follows from the fact that the Green's function for the operator  $\Delta + 1$  on a compact two dimensional manifold has logarithmic singularities, and no worse, along the diagonal.

<sup>7</sup>In particular, one formally infinite adjustment (1.8) suffices to render all the terms in the formal perturbative series finite. It is perhaps possible to view the finiteness of the perturbation series for Chern-Simons gauge theory established by Axelrod-Singer [3] also as arising from the nonperturbative construction.

<sup>8</sup>If  $d > 2$ , the limit in (1.9) does not exist, and more sophisticated methods are required. See e.g. [11, 13, 8, 2].

(For  $S^1 \times S^1$  in the flat metric,  $C_\epsilon(x)$  is a constant we denote by  $C_\epsilon$ .) Again, the limit  $:F := \lim_{\epsilon \rightarrow 0} F_\epsilon$  exists by calculations using (1.4). However, the term  $-6C_\epsilon \int_M \phi_\epsilon^2$  in the definition of  $:F_\epsilon:$  is not positive, and consequently the limit function  $:F:$  is not bounded below. Thus integrability of the function  $e^{-:F:}$  requires more delicate estimates; see [9], Section 8.6. These results are based on the fact that estimates on the measure of the set where  $e^{-:F:}$  is large can be made in terms of estimates on integrals of polynomial functions, though more subtle than those of (1.9). There are also entirely different problems addressed in Constructive Quantum Field Theory to obtain the infinite volume limit and prove the Wightman axioms. None of these issues arise in our construction of Chern-Simons gauge theory.

**1.2. The Chern Simons functional integral.** We now outline the ideas required to construct a mathematical version of the Chern Simons functional integral.

Consider the Lagrangian (1.2)

$$\text{CS}(A) = \frac{1}{4\pi} \text{tr} \int_M AdA + \frac{2}{3} A^3,$$

which appears in the formal functional integral (1.1)

$$Z_k(M) = \int_{\mathcal{A}} dA e^{ik\text{CS}(A)}$$

multiplied by an imaginary factor  $ik$ . We would like to proceed by analogy to the formal expression (1.12)

$$\int d\phi e^{-\int_M (\frac{1}{2}|d\phi(x)|^2 + \frac{1}{2}\phi(x)^2 - i\lambda\phi(x)^3) dx}.$$

whose mathematically well-defined version is (1.11)

$$\int_{\mathbf{B}} d\mu(\phi) e^{i\lambda:F(\phi):}.$$

In (1.12), the quadratic term is real and the cubic is multiplied by an imaginary coupling.<sup>9</sup> The formal functional integral (1.1) can also be brought into this form by a formal change of variable. We therefore take as our starting point the formal functional integral

$$(1.14) \quad \int_{\mathcal{A}} dA e^{\text{tr} \int_M \frac{1}{2} AdA + i\lambda A^3}$$

where  $\lambda \in \mathbb{R}$ . The fact that perturbative Chern-Simons gauge theory has been a useful approach is one piece of evidence in favor of studying this type of functional integral.<sup>10</sup>

Consider then further the formal functional integrals (1.1) and (1.14). To begin with, the quadratic term appearing in both is degenerate, with the degeneracy corresponding

<sup>9</sup>This combination of a real, negative definite quadratic term and an imaginary cubic is necessary even in one dimension, to obtain convergent integrals over  $\mathbb{R}$  giving rise to Airy functions.

<sup>10</sup>A similar phenomenon, where the mathematically well-defined functional integrals give the objects of physical interest by a form of analytic continuation, is well-understood in constructive quantum field theory. Feynman's formal functional integrals take the form

$$Z_{\text{Formal}}(\lambda) = \int d\phi e^{i \int_M \phi \square \phi + \lambda P(\phi)},$$

to the gauge symmetry of the Chern-Simons function. The method of dealing with this degeneracy is standard: We choose a Riemannian metric on the three manifold  $M$ , and replace the space of all connections by the space of connections on  $M$  satisfying the gauge condition

$$(1.16) \quad d^*A = 0.$$

(We continue to denote this space by  $\mathcal{A}$ .) In order to avoid further complications, we restrict our attention to compact, connected manifolds with trivial real first homology group; i.e.

$$\begin{aligned} H^1(M, \mathbb{R}) &= H^2(M, \mathbb{R}) = 0 \\ H^0(M, \mathbb{R}) &= H^3(M, \mathbb{R}) = \mathbb{R}. \end{aligned}$$

In order to have any chance of the formal functional integral yielding topological invariants, the fact that the gauge choice (1.16) depends explicitly on the metric must be addressed by including additional ghost fields in the action. This modification of the formal functional integral (1.1) was addressed in [26, 3]. Following their methods, the formal functional integral we wish to address mathematically is

$$(1.17) \quad \int dAdc_0dc_2 e^{\text{tr} \int_M \frac{1}{2}(AdA - 2c_0dc_2) + i\lambda(A^3 - 6Ac_0c_2)}$$

where the connections  $A$  satisfy the gauge condition (1.16) and where the ghost field  $c_0$  is an anticommuting scalar with values in  $\mathfrak{g}$  lying in the orthogonal complement  $\Omega^0(M, \mathfrak{g})^\perp$  of the constants in  $\Omega^0(M, \mathfrak{g})$ <sup>11</sup> and the ghost field  $c_2$  is an anticommuting field with values in  $\Omega^2(M, \mathfrak{g})$  and satisfying the gauge condition  $d^*c_2 = 0$ .

Our aim then is to give precise meaning to all the elements appearing in (1.17). For now we note that the formal Berezin integral

$$(1.18) \quad \frac{1}{Z_{ghost}} \int dc_0dc_2 e^{-\text{tr} \int_M (c_0dc_2)}(\cdot)$$

(where  $Z_{ghost}$  is a formally infinite normalization factor) is a mnemonic device denoting the pairing of  $\bigwedge^*(\Omega^0(M, \mathfrak{g})^\perp)$  with  $\bigwedge^*(\Omega^2(M, \mathfrak{g}))$  induced by the inverse of the operator  $\star d : \Omega^0(M, \mathfrak{g})^\perp \rightarrow \Omega^2(M, \mathfrak{g})$  and causes complications, but virtually no difficulties, in our

---

where  $\square$  is the wave operator; whereas the non-Gaussian integrals of constructive quantum field theory correspond to formal expressions of the type

$$(1.15) \quad Z_{Euclidean}(\lambda) = \int d\phi e^{-\int_M \phi \Delta \phi + \lambda P(\phi)}.$$

Thus the objects of study in the formal Feynman integrals might be expected to be obtainable from formally related but well-defined objects in the non-Gaussian measures corresponding to (1.15) by analytic continuation. The issue of whether such an analytic continuation exists, and what properties it has, was extensively studied in constructive quantum field theory; we refer to [24, 9] for more information.

<sup>11</sup>The orthocomplement of the constants can be taken since the constant ghost modes decouple in the free part of the action (1.17).

construction. The main task of this paper is to interpret formal functional integrals of the type

$$(1.19) \quad \frac{1}{Z_{gauge}} \int_{\mathcal{A}} dA e^{\frac{1}{2} \text{tr} \int_M AdA}(\cdot)$$

(where  $Z_{gauge}$  is a formally infinite normalization factor) as giving a linear functional with useful positivity properties on a subalgebra of the algebra of functions on  $\mathcal{A}$ .

Now the quadratic term in the exponential in (1.19) is not positive, and therefore does not correspond in a naive way to a Gaussian measure. However, this quadratic term does possess a more subtle form of positivity, known in constructive quantum field theory as Reflection Positivity (or Osterwalder-Schrader Positivity) [19].

More explicitly, we will consider manifolds  $M$  equipped with a diffeomorphism  $R$ , which preserves, *but does not fix*,<sup>12</sup> a compact, connected, oriented submanifold  $\Sigma$  lying in  $M$ , and where  $M - \Sigma$  consists of two connected components  $M_+$  and  $M_- = RM_+$ . We require the action of  $R$  on  $\Sigma$  to lift to an action on  $H^1(\Sigma, \mathbb{R})$  which makes the bilinear form  $Q : H^1(\Sigma, \mathbb{R}) \otimes H^1(\Sigma, \mathbb{R}) \rightarrow \mathbb{R}$  given by

$$(1.20) \quad Q(a, b) = - \int_{\Sigma} R^* a \wedge b$$

*positive definite* on cohomology classes extending to  $M_-$ .<sup>13</sup> We then choose an  $R$ -invariant metric on  $M$  and study expressions given by formal expectations of functions on  $\mathcal{A}$  given by a product of linear functionals on  $\mathcal{A}$  in the formal integral (1.19), and given by

$$(1.21) \quad \frac{1}{Z_{gauge}} \int_{\mathcal{A}} dA e^{\frac{1}{2} \text{tr} \int_M AdA} \langle f, A \rangle \langle g, A \rangle = \langle f, Lg \rangle$$

where  $f, g \in \Omega^1(M, \mathfrak{g})$  satisfy  $d^* f = d^* g = 0$ ,  $L = (\star d) \Delta^{-1}$ ,  $\Delta$  is the Laplacian on  $\Omega^1(M, \mathfrak{g})$ , whose inverse exists due to the cohomological condition  $H^1(M, \mathbb{R}) = 0$ , and the inner product  $\langle \cdot, \cdot \rangle$  is the  $L^2$  inner product on one forms with values in  $\mathfrak{g}$  associated to the inner product on  $\mathfrak{g}$ .<sup>14</sup> (Then  $\frac{1}{2} \text{tr} \int_M AdA = -\frac{1}{2} \langle A, \star dA \rangle$  and formal application of the rule (1.4) gives (1.21).) The main theorem is then

**Theorem 1.** *Suppose  $f \in \Omega_c^1(M_+, \mathfrak{g})$  is a  $\mathfrak{g}$ -valued one form compactly supported on  $M_+$ , and satisfies  $d^* f = 0$ . Then*

$$(1.22) \quad \langle R^* f, Lf \rangle_{L_2} \geq 0$$

The positive semidefiniteness in (1.22) is the positive semidefiniteness of the expectations of certain quadratic polynomials in the formal functional integral associated to (1.19), and implies a similar positivity property for the expectations of *any* polynomial

<sup>12</sup>Reflection positivity in constructive quantum field theory generally considers reflections which *fix* a time-zero submanifold. In our case, the nontrivial action of  $R$  on  $\Sigma$  is crucial to the construction of a positive definite inner product.

<sup>13</sup>In fact, much less will suffice; we can consider any manifold formed by gluing two copies of a three manifold with boundary  $M_+$  by a diffeomorphism of the boundary having the appropriate action on cohomology. See Section 2.

<sup>14</sup>In the technical part of the paper, once we have interpreted formal functional integrals in terms of appropriate spaces of connections, the inner products appearing on the left hand side of (1.21) will be replaced by evaluation of distributions on elements of their dual spaces.



function arising from functions compactly supported on  $M_+$  (Theorem 2). Thus, the formal expectations motivated by (1.22) give a reflection positive linear functional on a space of functions on a Banach space  $\mathbf{A}$  of connections on  $M$ . In order to obtain a Hilbert space from this construction we must take a quotient by the those functions which are in the kernel of this reflection positive functional. We will show in Theorem 3 that the resulting Hilbert space is given by

$$(1.23) \quad H^B(\Sigma) = \text{Sym}^*(\Lambda)^-,$$

where  $\Lambda \subset H^1(\Sigma, \mathfrak{g})$  is the image of  $H^1(M_-, \mathfrak{g})$  in  $H^1(\Sigma, \mathfrak{g})$  and the symmetric algebra  $\text{Sym}^*(\Lambda)$  is completed in the norm arising from the bilinear form  $Q$  in (1.20) and the metric on  $\mathfrak{g}$ .<sup>15</sup>

Including the Fermions gives a full Hilbert space

$$H(\Sigma) = H^B(\Sigma) \otimes H^F(\Sigma)$$

where

$$H^F(\Sigma) = \bigwedge^* (H^0(\Sigma, \mathfrak{g}) \oplus H^2(\Sigma, \mathfrak{g}))$$

is a finite dimensional vector space. Thus the Hilbert space is a symmetric algebra in (a subspace of) the even homology of  $\Sigma$ , tensored with an alternating algebra on the odd cohomology of  $\Sigma$ . A subtlety is that the positive definite metric on  $H^F(\Sigma)$  is not the natural one from the point of view of topological quantum field theory; this is related to the fact that the Fermions are ghosts. We will show how to deal with this issue.

So far we have dealt with the quadratic term in the Chern-Simons functional, which is the only term occurring in abelian Chern-Simons gauge theory, and our Reflection positivity result falls naturally into what may be expected from such a theory. It remains to deal with the cubic term appearing in (1.17). (We continue to focus on the space  $\mathbf{A}$  of connections, and leave consideration of the ghosts for the technical part of the paper.) We wish to make sense of an expression of the form

$$(1.24) \quad e^{i\lambda \text{tr} \int_M A^3}.$$

Now unlike the case of cubic Bosonic quantum field theory, where such an expression can be interpreted as a function on the analog of the space of connections, our reflection positivity theorem only applies to polynomial functions<sup>16</sup>, so we have to interpret (1.24) in terms of the Hilbert space  $H^B(\Sigma)$ .

To do so, we write

$$\text{tr} \int_M A^3 = \Xi_B^+ + R^* \Xi_B^+$$

where

<sup>15</sup>In concrete terms, the identification in (1.23) is given by the map taking a one form  $f \in \Omega_c^1(M_+, \mathfrak{g})$  satisfying  $d^*f = 0$  to the cohomology class  $[Lf|_\Sigma] \in H^1(\Sigma, \mathfrak{g})$ . Since  $f$  is compactly supported on  $M_+$ , the one form  $Lf$  is closed on  $\Sigma$ , and extends to a closed form on  $M_-$ ; in other words,  $[Lf|_\Sigma] \in \Lambda$ .

<sup>16</sup>See Section 4.3 for a further discussion of this point.

$$(1.25) \quad \Xi_B^+ = \text{tr} \int_{M_+} A^3$$

is a polynomial<sup>17</sup> in the gauge field supported on  $M_+$ . So morally, we may expect that the interaction term corresponds to a selfadjoint multiplication operator on  $H^B(\Sigma)$ .

However, the polynomial  $\Xi_B^+$  is not quite an element of the configuration space, since it is not a limit of polynomials in terms of the form  $A(f)$  where  $f$  is a one form compactly supported on  $M_+$  and satisfying the condition  $d^*f = 0$ . One way of seeing that this difficulty is essential is to note that the expectation in the formal path integral of the product of  $\Xi_B^+$  with a cubic polynomial of the form  $A(Rf)A(Rg)A(Rh)$ , where  $f, g, h \in \Omega_c^1(M_+, \mathfrak{g})$  satisfy  $d^*f = d^*g = d^*h = 0$ , depends on the closed one forms  $Lf|_\Sigma, Lg|_\Sigma, Lh|_\Sigma$ , and not only on the cohomology classes (and Hilbert space images)  $[Lf|_\Sigma], [Lg|_\Sigma], [Lh|_\Sigma]$ ; see (6.3). So in order to define a Hilbert space element corresponding to  $\Xi_B^+$  we have to make some choices of lifts of cohomology classes to elements of the configuration space, and the Hilbert space element  $\xi_B^+$  corresponding to  $\Xi_B^+$  will depend on these choices.

Having made these choices, we obtain a cubic element  $\xi_B^+ \in H^B(\Sigma)$  of the Hilbert space, a corresponding densely defined symmetric operator  $\xi_B^+ \cdot + (\xi_B^+ \cdot)^*$ , a selfadjoint extension  $O_B = (\xi_B^+ \cdot + (\xi_B^+ \cdot)^*)^-$ , and a one parameter subgroup of bounded operators  $e^{i\lambda O_B}$  on  $H^B(\Sigma) \otimes \mathbb{C}$ . We may thus consider the "Bosonic partition function"

$$Z_B(\lambda) = \langle \Omega, e^{i\lambda O_B} \Omega \rangle$$

where  $\lambda \in \mathbb{R}$  and  $\Omega$  is the image of the polynomial 1 in  $H^B(\Sigma)$ .

The inclusion of the Fermion ghosts causes complications of various sorts, since, as we noted, the positive definite metric for the Fermion Hilbert space is not the correct one for topological field theory. However, this ghost Hilbert space is finite dimensional, so this can be managed at the cost of some complications. Again, the definition of the interaction term requires choices. We can then ask whether the combined partition function is independent of the choices we made in the definition of  $\xi_B^+$  and their equivalent for the Fermion interaction term.

It is also an open question whether this method of constructing a Chern-Simons interaction term is powerful enough to reproduce the manifold invariants of Witten and Reshetikhin-Turaev. In standard quantum field theory, it is necessary to take a time ordered exponential of interacting terms, not an ordinary exponential of the type we have studied. On the other hand, in a topological quantum field theory, where the Hamiltonian is zero and the effect of an interaction term depends only on topological data (see Appendix A for a manifestation of that), it is possible that this does not matter. See Questions 6.10, 6.11, and 6.12 for concrete statements.<sup>18</sup>

<sup>17</sup>The polynomial in (1.25) looks like a cubic, but in fact is linear in each of the components of the gauge field, and is an exterior product of those components, so its singularity properties are much better than that of a cubic of the type appearing in (1.10). One aspect of this is the vanishing of the Wick ordering correction. Since we cannot use this polynomial interaction, due to other reasons explained below, we do not attempt to study it further.

<sup>18</sup>A technical comment: The invariants of Witten and Reshetikhin-Turaev are defined for framed manifolds. Since we choose a metric on  $M$  once and for all, and normalize the quadratic functional integral associated to this metric to have value 1 on the polynomial 1, we are effectively normalizing the functional integral in a way that removes this framing dependence.

The remainder of the paper is structured as follows. In Section 2, we construct manifolds, which we require to be rational homology spheres, equipped with a reflection map adequate for an algebra of functions on the corresponding configuration spaces to carry a reflection positive functional. We also study cases where these manifolds satisfy further conditions which allow a more geometric construction of these reflection maps. In Section 3, we construct the configuration spaces for the gauge and ghost fields. There is a great deal of freedom in constructing such configuration spaces, and we make a convenient choice. We then proceed in Section 4 with the main result of the paper (Theorem 1 above) which shows that the quadratic term in the Chern-Simons function gives a reflection positive functional on the one particle configuration space of the gauge fields, and more generally a reflection positive functional on the polynomial algebra of fields (Theorem 2). We also compute the Hilbert space which this reflection positive functional associates with a two manifold given morally by the fixed set of the reflection. In Section 5, we study a similar construction for the Fermion, or ghost fields. The key result is Theorem 4, followed by Proposition 5.26. We then combine the results of Sections 4 and 5 and summarize them in Theorem 5. Finally in Section 6 we construct operators on these Hilbert spaces corresponding to the interaction terms, and in Theorem 6 obtain the partition function as a weak limit of vacuum expectations values of exponentials of these operators. We end with a discussion of possible relations to the manifold invariants of Witten and Reshetikhin-Turaev.

## 2. MANIFOLDS WITH REFLECTION

Let  $M_+$  be a smooth, compact, connected, oriented 3-manifold with boundary  $\Sigma$ , a smooth, compact, connected, oriented 2-manifold of genus  $g > 0$ . The orientation on  $\Sigma$  gives the vector space  $H^1(\Sigma, \mathbb{R})$  a symplectic structure  $\omega$  given by the intersection form. The image  $\Lambda_+$  of  $H^1(M_+, \mathbb{R})$  in  $H^1(\Sigma, \mathbb{R})$  is a Lagrangian subvariety.

Let  $S : \Sigma \rightarrow \Sigma$  be an orientation reversing diffeomorphism whose action  $S^*$  on  $H^1(\Sigma, \mathbb{R})$  satisfies<sup>19</sup>

$$(2.1) \quad \begin{aligned} (S^*)^2 &= 1 \\ \omega(S^*a, a) &> 0 \text{ for all nonzero } a \in \Lambda_+. \end{aligned}$$

Then  $\Lambda_{\mathbb{R}} = S^*\Lambda_+$  is also Lagrangian, and

$$(2.2) \quad \omega(S^*a, a) < 0 \text{ for all nonzero } a \in \Lambda_{\mathbb{R}}.$$

We can use  $S$  to construct a compact, connected, oriented 3-manifold  $M$  by gluing two copies of  $M_+$ , which we denote by  $M_+, M_-$ , by the diffeomorphism  $S$  of their common boundary. From now on we impose the assumption

$$H^1(M, \mathbb{R}) = 0.$$

Our construction of the functional integral will work for any rational homology sphere arising from a gluing satisfying the conditions (2.1). However, the construction will simplify considerably if the diffeomorphism  $S$  satisfies some finiteness conditions. One such

---

<sup>19</sup>Note that since the conditions (2.1) on  $S$  involve only the action  $S^*$  of  $S$  on cohomology, and since the representation of the mapping class group in  $Sp(2g, \mathbb{Z})$  given by the action on cohomology is a surjection onto  $Sp(2g, \mathbb{Z})$  (see [4]), any element of  $Sp(2g, \mathbb{Z})$  satisfying (2.1) arises from the action of some diffeomorphism.

condition is

**Condition A:**  $S$  is of finite order.

It simplifies further if  $S$  satisfies

**Condition B:**  $S^2 = \text{id}$ .

If Condition B is satisfied,  $M$  comes equipped with an orientation preserving diffeomorphism  $R$ , exchanging  $M_+$  and  $M_-$ , and whose action on  $\Sigma$  is given by  $S$ ; this is the situation considered in the introduction. To construct this orientation preserving diffeomorphism  $R : M \rightarrow M$ , write

$$M = (M_+ \amalg M_-) / \sim,$$

where  $m_+ \in M_+, m_- \in M_-$  satisfy  $m_+ \sim m_-$  if  $m_+ \in \partial M_+ = \Sigma, m_- \in \partial M_- = \Sigma$ , and  $m_- = Sm_+$  (as elements of  $\Sigma$ ). Then define  $R_+ : M_+ \rightarrow M_-$  by

$$R_+(m) = j(m)$$

and  $R_- : M_- \rightarrow M_+$  by

$$R_-(m) = j^{-1}(m)$$

where

$$(2.3) \quad j : M_+ \rightarrow M_-$$

is the identity diffeomorphism identifying the two copies  $M_+, M_-$  of  $M_+$ . Then  $R_+$  agrees with  $R_-$  on  $\Sigma$ , and gives a diffeomorphism of  $M$  preserving  $\Sigma$  and whose action on  $\Sigma$  is given by  $S$ . Choosing an invariant metric this diffeomorphism gives an involution of  $\Omega^*(M)$  preserving the de Rham operator and the Riemannian inner product, and exchanging the subspaces  $\Omega_c^*(M_+)$  and  $\Omega_c^*(M_-)$  of compactly supported differential forms.

We now choose a compact Lie group  $G$  and a representation of  $G$  whose associated trace gives a metric on  $\mathfrak{g}$ . This metric, combined with the Riemannian metric, gives a Hodge star operator and a metric on the differential forms with coefficients in  $\mathfrak{g}$ . The involution  $R^*$  then extends to the differential forms with coefficients in  $\mathfrak{g}$  and preserves the metric and Hodge star operator.

In the case of the weaker Condition A, we do not know how to construct such a diffeomorphism, but it is still possible to construct an involution exchanging forms compactly supported on  $M_+$  with those compactly supported on  $M_-$  with the properties needed to define a satisfactory form of reflection positivity. To do so, note that the identification  $j : M_+ \rightarrow M_-$  induces an isomorphism  $j^* : \Omega_c^*(M_-, \mathfrak{g}) \rightarrow \Omega_c^*(M_+, \mathfrak{g})$  on the compactly supported forms. To find a metric for which  $j^*$  preserves also the Hodge  $\star$  operator, choose an  $S$ -invariant metric  $g$  on  $\Sigma$ , and a metric on  $M_+$  taking the form  $g + (dt)^2$  on a collar neighborhood  $U = \Sigma \times [0, 1)$  of the boundary  $\Sigma$ . Gluing this metric to the same metric on  $M_-$  gives a metric on  $M$  which also has this collar form on a neighborhood of  $\Sigma \subset M$ . Then the map  $j^*$  satisfies

$$j^*(\star\xi) = \star j^*(\xi)$$

for any  $\xi \in \Omega_c^*(M_-, \mathfrak{g})$ . Since  $j$  is orientation preserving, the map  $j^*$  preserves the metric on compactly supported forms. We continue to denote the resulting map by

$$R^* : \Omega_c^*(M_{\pm}, \mathfrak{g}) \longrightarrow \Omega_c^*(M_{\mp}, \mathfrak{g}),$$

whether or not it arises from a diffeomorphism of  $M$ .

In the absence of Condition A, we can still obtain such a map as long as we restrict to forms compactly supported on the complement of a collar containing  $\Sigma$ . To do so, we construct a metric on  $M$  by again equipping  $M_+$  with a metric taking the collar form  $g + (dt)^2$  on a neighborhood of its boundary, and gluing  $M_+$  to a copy of  $M_-$  equipped with the same metric, using a collar  $\Sigma \times [-\delta, \delta]$  equipped with a metric given by  $g + (dt)^2$  near  $\Sigma \times \{-\delta\}$  and by  $S^*g + (dt)^2$  near  $\Sigma \times \{\delta\}$ . To perform the gluing, we use the identity on the boundary of  $M_-$  and the diffeomorphism  $S$  on the boundary of  $M_+$ . We therefore obtain a metric on  $M$  where the metrics on  $M_-$  and  $M_+$  agree on the complement of a  $\delta$  neighborhood  $W_\delta$  of  $\Sigma$ . Then the map  $j$  still preserves the metric, and therefore the Hodge  $\star$  operator, on the complement of  $W_\delta$ . By restricting  $j^* : \Omega_c^*(M_-, \mathfrak{g}) \longrightarrow \Omega_c^*(M_+, \mathfrak{g})$  to forms compactly supported away from this collar, we obtain a map

$$R^* : \Omega_c^*(M_{\pm}^\delta, \mathfrak{g}) \longrightarrow \Omega_c^*(M_{\mp}^\delta, \mathfrak{g}),$$

where  $M_{\pm}^\delta = M_{\pm} \cap (M - W_\delta)$ . The involution  $R^*$  preserves the Hodge  $\star$  operator, and therefore the metric on forms. It will then be necessary to take another limit, as  $\delta \longrightarrow 0$ , of the partition function and other quantum field theoretic objects.

In sum, we have chosen in all cases a Riemannian metric on  $M$ , along with an invariant metric on  $\mathfrak{g}$  arising from the trace in a representation. These endow the differential forms with coefficients in  $\mathfrak{g}$  with a Hodge star operator and a metric. We also have a reflection  $R^*$  which may or may not arise from the action of an involution on  $M$ , but which exchanges the differential forms on  $M$  compactly supported on  $M_+^\delta = M_+ \cap (M - W_\delta)$  (where in the case of Condition A, we may take  $\delta = 0$ ) with those compactly supported in its image in  $M_-$ , and preserves the metric and star operator on such forms.<sup>20</sup>

We will from now on consider mainly the case where Condition A is satisfied, and indicate the simplifications that occur in the case of Condition B, and the additional constructions needed in the absence of either condition.

### 3. THE CONFIGURATION SPACES $\mathcal{F}^B$ AND $\mathcal{F}^F$

In this section we construct the configuration spaces of our quantum field model. The configuration space  $\mathcal{F}^B$  for the gauge fields is the completion of the symmetric algebra on the one-particle configuration space  $\mathcal{F}_1^B$ . The space  $\mathcal{F}^B$  can also be viewed as a space of integrable functions on a space of connections  $\mathbf{A}$ , with respect to a Gaussian measure we construct on  $\mathbf{A}$ .

Similarly, we construct a one-particle configuration space  $\mathcal{F}_1^F$  corresponding to the ghost configuration space, and the completion  $\mathcal{F}^F$  of its alternating algebra is the configuration space for the ghost fields.

---

<sup>20</sup>We consider forms compactly supported on  $M_{\pm}$  as forms on  $M$  by extension by zero.

**3.1. The one-particle configuration spaces  $\mathcal{F}_1^B$  and  $\mathcal{F}_1^F$ .** Consider the subspace  $F_1^B = \ker(d^*) \subset \Omega^1(M, \mathfrak{g})$ , and complete it to a Hilbert space  $\mathcal{F}_1^B$  using the  $H_s$  Sobolev norm on  $\Omega^1(M, \mathfrak{g})$ , with  $s$  to be chosen later; we may take  $s = 4$ .<sup>21</sup> Since  $H^1(M, \mathbb{R}) = 0$ , the Laplacian  $\Delta = dd^* + d^*d$  on  $\Omega^1(M, \mathfrak{g})$  has an inverse  $\Delta^{-1}$  which extends to a bounded operator on  $\mathcal{F}_1^B$ . The operator

$$L = \star d \Delta^{-1}$$

is then also a bounded operator on  $\mathcal{F}_1^B$ . For any open set  $U \subset M$ , there are subspaces  $F_1^B(U) \subset F_1^B$  and  $\mathcal{F}_1^B(U) \subset \mathcal{F}_1^B$  consisting of forms compactly supported on  $U$ .

The odd variables, or Fermions, appearing in the action (1.17) are a scalar  $c_0 \in \Omega^0(M, \mathfrak{g})$  and a two form  $c_2 \in \Omega^2(M, \mathfrak{g})$ . Consider first the space of smooth forms  $\Omega^0(M, \mathfrak{g})$ . Inside this space we may consider the orthocomplement of the constants  $\mathfrak{g} \subset \Omega^0(M, \mathfrak{g})$  in the  $L^2$ -inner product; in other words the functions  $f \in \Omega^0(M, \mathfrak{g})$  with  $\int_M \star f = 0$ . We denote this subspace of  $\Omega^0(M, \mathfrak{g})$  by  $\Omega^0(M, \mathfrak{g})^\perp$ . Similarly we consider the space  $\ker(d_2^*) \subset \Omega^2(M, \mathfrak{g})$ . We then form the space  $\mathcal{F}_1^F$  by completing the space

$$F_1^F = \Omega^0(M, \mathfrak{g})^\perp \oplus \ker(d_2^*)$$

in the  $H_s$ <sup>22</sup> Sobolev norm given by the metric on  $M$  and the inner product on  $\mathfrak{g}$ .

On the space  $F_1^F$ , the operator  $\star d$  acts, interchanging the scalars and the two forms. Since we have taken the orthocomplement of the constants in  $\Omega^0$  and the kernel of  $d_2^*$  in  $\Omega^2$ , this operator has an inverse (as in [3]; see also [1])  $L = \star d \Delta^{-1}$ , which we continue to denote by  $L$ , a bounded operator on  $\mathcal{F}_1^F$ .

As in the Bose case, we may define a one particle configuration space corresponding to any open subset  $U \subset M$ . There is a slight subtlety due to the fact that in the construction of  $F_1^F$  we took the orthocomplement of the constants in the zero forms. So let  $F_1^F(U) = (a, b) \in \Omega^0(M, \mathfrak{g})^\perp \oplus \ker(d_2^*)$  with  $da$  and  $b$  compactly supported inside of  $U$ , and let  $\mathcal{F}_1^F(U)$  be the completion of  $F_1^F(U)$  in  $\mathcal{F}_1^F$ .

**3.2. The Bose configuration space  $\mathcal{F}^B$ .** We define the Bose configuration space by

$$\mathcal{F}^B = \text{Sym}^*(\mathcal{F}_1^B)^-,$$

the completion of the symmetric algebra  $\text{Sym}^*(\mathcal{F}_1^B)$  on  $\mathcal{F}_1^B$  in the norm induced on  $\text{Sym}^*(\mathcal{F}_1^B)$  by the Hilbert norm on  $\mathcal{F}_1^B$ . For future purposes we also define the dense subspace of polynomials  $\mathcal{F}_P^B = \text{Sym}^*(\mathcal{F}_1^B) \subset \mathcal{F}^B$ ; likewise the subspace  $\mathcal{F}_d^B = \text{Sym}^{\leq d}(\mathcal{F}_1^B) \subset \mathcal{F}^B$  of polynomials of degree  $\leq d$ .

**Remark 3.1.** The space  $\mathcal{F}^B$  has a description in terms of Gaussian integrals.

Let  $\mathbf{A}$  be the space of  $\mathfrak{g}$ -valued one forms  $A$  on  $M$  of Sobolev class  $-s - \frac{3}{2} - \epsilon$  for some  $\epsilon > 0$  and satisfying the condition  $d^*A = 0$ . On the space  $\mathbf{A}$  consider the the Gaussian measure  $d\mu$  with covariance  $\Delta^s$ ; again we may take  $s = 4$ .<sup>23</sup> In the terms used in the introduction, this measure can be obtained from the Hilbert space  $\mathbf{H}^* = \mathcal{F}_1^B$  by completing it to a Banach space  $\mathbf{B} = \mathbf{A}$ ; see [12].

This measure is characterized by integrals of exponentials, or alternatively, polynomial functions, on  $\mathbf{A}$  : Given  $f \in \mathbf{A}^* \subset \mathbf{H}^*$ ,

<sup>21</sup>The precise choice of  $s$  does not play much of a role in our construction, so long as it is chosen large enough. It is convenient for de Rham theory and the Hodge theorem to have forms with two continuous derivatives, which is the case if  $s > 7/2$ . We will need  $s > \frac{1}{4}$  for (4.17) and  $s$  to be an integer for Remark 3.5.

<sup>22</sup>We again choose the same  $s$  (i.e.  $s = 4$ ) for similar reasons to those in the Bose case.

<sup>23</sup>The formal path integral corresponds to the action  $S(A) = \langle A, \Delta^{-s} A \rangle_{L_2}$ .

$$(3.2) \quad \int_{\mathbf{A}} d\mu(A) e^{if(A)} = e^{-\frac{1}{2}\langle f, f \rangle_{\mathbf{H}^*}}$$

and given  $f_j \in \mathbf{A}^* \subset \mathbf{H}^*$ ,  $j = 1, \dots, m$ , the polynomial function  $F(A) = f_1(A) \dots f_m(A)$  is integrable, and for  $m$  even

$$(3.3) \quad \int_{\mathbf{A}} d\mu(A) f_1(A) \dots f_m(A) = \frac{1}{2^{m/2}(m/2)!} \sum_{\sigma \in \Sigma_m} \langle f_{\sigma(1)}, f_{\sigma(2)} \rangle_{\mathbf{H}^*} \dots \langle f_{\sigma(m-1)}, f_{\sigma(m)} \rangle_{\mathbf{H}^*}$$

where  $\Sigma_m$  is the permutation group on  $m$  letters. (For  $m$  odd, the integral is zero.)

Then  $\mathcal{F}_P^B$  is isometrically embedded in  $L^2(\mathbf{A}, d\mu)$ .<sup>24</sup>

In view of Remark 3.1, when convenient, we will sometimes consider elements of  $\mathcal{F}^B$  as functions on the space  $\mathbf{A}$ , interchangeably with their nature as elements of a completed symmetric algebra. So for example, if  $f_1, \dots, f_d \in F_1^B$ , the expression  $f_1(\cdot) \dots f_d(\cdot)$  is the function on  $\mathbf{A}$  associated to the element  $f_1 \otimes \dots \otimes f_d \in \mathcal{F}_P^B$ .

The following dense subsets of  $\mathcal{F}^B$  will be convenient to keep in mind.

*The ring of polynomials.* Let  $\mathcal{R} = \text{Sym}^*(F_1^B)$ .

Equivalently, we may use a different set of polynomials, which is also dense:

*The L-Wick ordered polynomials.* The  $L$ -Wick ordered polynomials  $\mathcal{O}$  are generated as a vector space by monomials of the form

$$(3.4) \quad : m_{f_1, \dots, f_m} := f_1(\cdot) \dots f_m(\cdot) :,$$

where  $f_1, \dots, f_m \in F_1^B$  as above. This is a polynomial of degree  $m$  with coefficients which can be extracted from the generating function

$$: e^{if(\cdot)} := e^{\frac{1}{2}\langle f, Lf \rangle_{L_2}} e^{if(\cdot)}.$$

Since any polynomial can be written as a linear combination of  $L$ -Wick ordered polynomials, these are also a dense subset of  $\mathcal{F}^B$ .<sup>25</sup>

For future reference we note that the definition of  $: m_{f_1, \dots, f_m} :$  can be extended also to the case where  $f_i \in \mathcal{F}_1^B$ .

**3.3. The configuration space  $\mathcal{F}^B(U)$  associated to  $U \subset M$ .** Inside  $\mathcal{F}^B$  we may consider the subspace  $\mathcal{R}(U) = \text{Sym}^*(F_1^B)(U)$ , and likewise the subspace  $\mathcal{O}(U)$  generated by elements of the form (3.4) where  $f_1, \dots, f_m$  are smooth and have compact support in  $U$ . The completion of any of these sets in  $\mathcal{F}^B$  is a Hilbert space we denote by  $\mathcal{F}^B(U)$ . This space may also be built up by completing the symmetric algebra of a Hilbert space, restricting to the case of one forms compactly supported in  $U$ : Letting  $\mathcal{F}_1^B(U) = \mathcal{F}_1^B \cap \mathcal{F}^B(U)$ , we have

$$\mathcal{F}^B(U) = (\text{Sym}^* \mathcal{F}_1^B(U))^-$$

and we define

$$\mathcal{F}_P^B(U) = \mathcal{F}_P^B \cap \mathcal{F}^B(U)$$

<sup>24</sup>This embedding uses Wick ordering. It is a bit simpler to build  $\mathcal{F}^B$  in the measure given by  $d\mu$  on the complexification of  $\mathbf{A}$ , as the completion of the space of real polynomials considered as functions on the complexified fields, since this avoids Wick ordering troubles.

<sup>25</sup>See further comments on  $L$ -Wick ordered polynomials in the next Section.

and

$$\mathcal{F}_d^B(U) = \mathcal{F}_d^B \cap \mathcal{F}^B(U)$$

**Remark 3.5.** If we choose  $s$  to be a nonnegative integer, the covariance in Remark 3.1 is a differential operator, and has zero correlations between disjoint open sets. Explicitly, if  $U, V$  are disjoint open sets, and  $P \in \mathcal{F}_P^B(U), Q \in \mathcal{F}_P^B(V)$ , then  $PQ \in \mathcal{F}_P^B$  and

$$\int d\mu P^2 Q^2 = \int d\mu P^2 \int d\mu Q^2;$$

alternatively,

$$\|PQ\|_2 = \|P\|_2 \|Q\|_2.$$

In other words,

**Lemma 3.6.** *Multiplication by  $P \in \mathcal{F}_P^B(U)$  is a bounded operator taking  $\mathcal{F}_P^B(V)$  to  $\mathcal{F}_P^B$ .*

**3.4. The Fermionic configuration Space  $\mathcal{F}^F$ .** Consider the alternating algebra  $\mathcal{F}_P^F = \bigwedge^* \mathcal{F}_1^F$ . The norm on  $\mathcal{F}_1^F$  makes  $\mathcal{F}_P^F$  into an inner product space, so it may be completed to a Hilbert space we call  $\mathcal{F}^F$ . Note that multiplication by an element of  $\mathcal{F}_1^F$  is bounded on  $\mathcal{F}^F$  and hence extends to a bounded operator on  $\mathcal{F}^F$ . (This is in contrast to the case of  $\mathcal{F}^B$ , where multiplication by an element of  $\mathcal{F}_1^B$  is not a bounded operator on  $\text{Sym}^*(\mathcal{F}_1^B)$ , and hence does not extend to an operator on  $\text{Sym}^*(\mathcal{F}_1^B)^\perp = \mathcal{F}^B$ .)

**Remark 3.7.** Alternatively, the inner product on  $\mathcal{F}^F$  can be considered as a functional on the larger alternating algebra  $\mathcal{U} = \mathcal{F}_L^F \otimes \mathcal{F}_R^F$  where  $\mathcal{F}_L^F, \mathcal{F}_R^F$  are two copies of  $\mathcal{F}^F$ ; the tensor product is the smallest alternating algebra containing both factors. Then any monomial  $m \in \mathcal{U}$  may be written as the product  $m = m_L m_R$  where  $m_L$  is a monomial in  $\mathcal{F}_L^F$  and  $m_R$  is a monomial in  $\mathcal{F}_R^F$ . We then define the functional  $\langle \cdot \rangle: \mathcal{U} \rightarrow \mathbb{R}$  by defining

$$(3.8) \quad \langle m_L^r m_R \rangle = \langle m_L, m_R \rangle$$

using the identification  $\mathcal{F}_L^F \simeq \mathcal{F}_R^F = \mathcal{F}^F$  and extending by linearity. (Here the notation  $m_L^r$  denotes the monomial  $(-1)^{\frac{|m_L|(|m_L|-1)}{2}} m_L$ , where  $|m_L|$  is the number of terms in  $m_L$ ; morally  $m_L^r$  is  $m_L$  written in reverse order.)<sup>26</sup>

In the language of Berezin integrals, the functional  $\langle \cdot \rangle$  is given by the Berezin integral

$$(3.9) \quad \langle p \rangle = \int dc_{0,L} dc_{0,R} dc_{2,L} dc_{2,R} e^{-\text{tr} \int_M {}^*c_{0,L} \Delta^{-s} c_{0,R} + {}^*c_{2,L} \Delta^{-s} c_{2,R}} p,$$

where  $p$  is any polynomial in the odd variables  $c_{0,L}, c_{0,R}, c_{2,L}, c_{2,R}$ , which can be thought of as an element of  $\mathcal{U}$ . These Berezin integrals are analogs of the Gaussian integrals of Remark 3.1.

As in the case of  $\mathcal{F}^B$ , we may consider the subalgebra  $\mathcal{F}^F(U) \subset \mathcal{F}^F$  corresponding to an open set  $U \subset M$ . Let  $\mathcal{F}_P^F(U) = \bigwedge^* \mathcal{F}_1^F(U) \subset \mathcal{F}^F$ , and let  $\mathcal{F}^F(U)$  be the completion of  $\mathcal{F}_P^F(U)$  in  $\mathcal{F}^F$ . Likewise we define  $\mathcal{F}_d^F(U)$  as the inhomogeneous polynomials of degree  $\leq d$ .

We now write<sup>27</sup>

<sup>26</sup>A technical point: There is no need to Wick order  $m_L$  and  $m_R$  on the left side of (3.8) since in the Berezin integral (3.9), any monomial of the form  $m_L$  or  $m_R$  is equal to its own Wick ordering.

<sup>27</sup>The tensor products below are tensor product of Hilbert spaces.



$$\mathcal{F} = \mathcal{F}^B \otimes \mathcal{F}^F$$

and, for any subset  $U \subset M$ ,

$$\mathcal{F}(U) = \mathcal{F}^B(U) \otimes \mathcal{F}^F(U)$$

as well as

$$\mathcal{F}_d(U) = \bigcup_{k+l=d} \mathcal{F}_k^B(U) \otimes \mathcal{F}_l^F(U)$$

and

$$\mathcal{F}_P(U) = \mathcal{F}_P^B(U) \otimes \mathcal{F}_P^F(U).$$

A special case is that where  $U = M_+^\delta$ ; we write  $\mathcal{F}_+^B = \mathcal{F}^B(M_+^\delta)$ ,  $\mathcal{F}_+^F = \mathcal{F}^F(M_+^\delta)$ , and  $\mathcal{F}_+ = \mathcal{F}(M_+^\delta)$  (where in the case of condition A or condition B being satisfied, we replace  $M_+^\delta$  by  $M_+$ ). We will write  $\mathcal{F}_{P,+}^B = \mathcal{F}_P^B \cap \mathcal{F}_+^B$ ,  $\mathcal{F}_{P,+}^F = \mathcal{F}_P^F \cap \mathcal{F}_+^F$ ,  $\mathcal{F}_{P,+} = \mathcal{F}_P \cap \mathcal{F}_+$ , and  $\mathcal{F}_{d,+}^B = \mathcal{F}_d^B \cap \mathcal{F}_+^B$ ,  $\mathcal{F}_{d,+}^F = \mathcal{F}_d^F \cap \mathcal{F}_+^F$ ,  $\mathcal{F}_{d,+} = \mathcal{F}_d \cap \mathcal{F}_+$ . Likewise we write  $\mathcal{R}_+$  and  $\mathcal{O}_+$ .

#### 4. REFLECTION POSITIVITY AND THE BOSONIC HILBERT SPACE

The formal functional integral

$$\int_{\mathcal{A}} e^{\frac{i}{2} \text{tr} \int_M \text{Ad}A}.$$

does not correspond in any known way to a measure on a Banach space of connections. However, it turns out that the expectations derived from application of the formal rule of the type (1.4) to polynomial functions in the formal functional integral (1.19)

$$(4.1) \quad \frac{1}{Z_{gauge}} \int_{\mathcal{A}} e^{\frac{1}{2} \text{tr} \int_M \text{Ad}A}(\cdot)$$

give rise to a reflection positive functional on the space of polynomials, in the following sense: Suppose that  $P, \tilde{P}$  are polynomials compactly supported in  $M_+$ .<sup>28</sup> Then application of the rules of the type (1.4) to the formal expression

$$q(P, \tilde{P}) = \frac{1}{Z_{gauge}} \int_{\mathcal{A}} e^{\frac{1}{2} \text{tr} \int_M \text{Ad}A} (R^* P) \tilde{P}$$

gives a positive semidefinite symmetric bilinear form on the space of polynomials. We may construct the Hilbert space  $H^B(\Sigma)$  by taking the quotient of the space of polynomials by the null elements, and completing in the norm induced by the bilinear form  $q$ . Along the way we prove that

$$H^B(\Sigma) = (\text{Sym}^* \Lambda)^-$$

where  $\Lambda \subset H^1(\Sigma, \mathfrak{g})$  is the image of  $H^1(M_-, \mathfrak{g})$  in  $H^1(\Sigma, \mathfrak{g})$ , and the closure is taken in the norm on the symmetric algebra  $\text{Sym}^* \Lambda$  arising from the norm on  $\Lambda$  given by the symmetric positive definite bilinear form

<sup>28</sup>In this informal discussion, we consider the case where Condition A is satisfied.

$$Q(a, b) = \text{tr} \int_{\Sigma} S^* a \wedge b,$$

for  $a, b \in \Lambda$ .

**4.1. Reflection positivity for the one particle configuration space  $\mathcal{F}_{1,+}^B$ .** The following theorem is the main idea in the construction of our reflection positive functional. Recall that  $\Delta$  is the Laplacian on differential forms, whose restriction to one-forms is invertible, and that  $L = \star d \Delta^{-1}$  is a compact operator. We now prove Theorem 1 from the introduction.

**Theorem 1.** *Let  $f \in \Omega^1(M, \mathfrak{g})$  be a smooth  $\mathfrak{g}$ -valued one form on  $M$  compactly supported in  $M_+$  (in the absence of Condition A,  $M_+^\delta$ ) and satisfying  $d^*f = 0$ . Then*

$$(4.2) \quad \langle R^* f, Lf \rangle_{L_2} \geq 0$$

*Proof:* We first consider the situation where  $M$  comes equipped with a reflection  $R$  (Condition B). We then have

$$(4.3) \quad \begin{aligned} \langle R^* f, Lf \rangle_{L_2} &= -\text{tr} \int_M R^* f \wedge \star Lf = \\ -\text{tr} \int_{M_-} R^* f \wedge \star Lf &= -\text{tr} \int_{M_-} dLR^* f \wedge Lf = \\ -\text{tr} \int_{M_-} LR^* f \wedge \star f &- \text{tr} \int_{-[\Sigma]} R^* Lf \wedge Lf. \end{aligned}$$

where we recall the orientation of  $\Sigma$  is that given by the orientation of  $M_+$ ; this gives rise to the negative sign in orientation of  $\Sigma$  in the final equality in (4.3). Since  $f$  is compactly supported on  $M_+$ ,

$$\text{tr} \int_{M_-} LR^* f \wedge \star f = 0.$$

But  $\star dLf = f$ , and therefore  $dLf$  is compactly supported on  $M_+$ . Hence  $Lf|_{\Sigma}$  is a closed  $\mathfrak{g}$ -valued one form on  $\Sigma$ , which extends to a closed one form  $Lf|_{M_-}$  on  $M_-$ ; in other words  $[Lf|_{\Sigma}] \in \Lambda = \Lambda_{\mathbb{R}} \otimes \mathfrak{g}$ . Then

$$\text{tr} \int_{\Sigma} R^* Lf \wedge Lf = \text{tr} \int_{\Sigma} R^* [Lf|_{\Sigma}] \wedge [Lf|_{\Sigma}],$$

where  $[Lf|_{\Sigma}]$  is the cohomology class of  $Lf|_{\Sigma}$  in  $H^1(\Sigma, \mathfrak{g})$ . But the bilinear form  $Q : H^1(\Sigma, \mathfrak{g}) \otimes H^1(\Sigma, \mathfrak{g}) \rightarrow \mathbb{R}$  given by

$$(4.4) \quad Q(a, b) = \text{tr} \int_{\Sigma} S^* a \wedge b$$

is symmetric and positive definite on  $\Lambda = \Lambda_{\mathbb{R}} \otimes \mathfrak{g}$  by (2.2).<sup>29</sup> Since  $R|_{\Sigma} = S$ ,

<sup>29</sup>Note that the trace differs from the positive inner product on the Lie algebra by a sign, so that the fact that the bilinear form in (2.2) is negative definite makes the form  $Q$  in (4.4) positive definite.

$$\mathrm{tr} \int_{\Sigma} R^*[Lf|_{\Sigma}] \wedge [Lf|_{\Sigma}] \geq 0.$$

In the more general case (Condition A), a slightly more elaborate argument is needed, but the essentials are the same. We begin with a Lemma.<sup>30</sup>

**Lemma 4.5.** *Let  $f \in \Omega_c^1(M_+, \mathfrak{g})$  and  $g \in \Omega_c^1(M_-, \mathfrak{g})$ . Then*

$$\mathrm{tr} \int_M f \wedge \star Lg = \mathrm{tr} \int_{\Sigma} Lf \wedge Lg.$$

*Proof.* We have

$$(4.6) \quad \begin{aligned} \mathrm{tr} \int_M f \wedge \star Lg &= \mathrm{tr} \int_{M_+} f \wedge \star Lg = \mathrm{tr} \int_{M_+} \star dLf \wedge \star Lg = \\ &= \mathrm{tr} \int_{M_+} Lf \wedge g + \mathrm{tr} \int_{\Sigma} Lf \wedge Lg. \end{aligned}$$

But  $g$  is compactly supported in  $M_-$ , so  $\mathrm{tr} \int_{M_+} Lf \wedge g = 0$ .  $\square$

Note that (in the notation of the above Lemma)  $dLf|_{\Sigma} = dLg|_{\Sigma} = 0$ , so that  $Lf$  and  $Lg$  are closed forms on  $\Sigma$ . We have shown that  $\mathrm{tr} \int_M f \wedge \star Lg = \mathrm{tr}([Lf|_{\Sigma}] \cdot [Lg|_{\Sigma}])$  where  $\cdot$  denotes the intersection form.

In fact, Lemma 4.5 applies to any splitting of  $M$  into two components by any submanifold  $\tilde{\Sigma} \subset M$ .

**Proposition 4.7.** *Let  $\tilde{\Sigma} \subset M$  be a compact connected oriented submanifold of  $M$  whose complement has two components  $M_1, M_2$ . Orient  $\tilde{\Sigma}$  as the boundary of  $M_1$ . Then if  $f_1 \in F_1^B(M_1)$ ,  $f_2 \in F_1^B(M_2)$ , we have*

$$\mathrm{tr} \int_M f_1 \wedge \star Lf_2 = \mathrm{tr} \int_{\tilde{\Sigma}} [Lf_1|_{\tilde{\Sigma}}] \wedge [Lf_2|_{\tilde{\Sigma}}].$$

Furthermore, if  $\Sigma_1, \Sigma_2$  are two isotopic compact submanifolds of  $M$ , and  $f$  is zero in the region between  $\Sigma_1$  and  $\Sigma_2$ , then  $[Lf|_{\Sigma_1}] = [Lf|_{\Sigma_2}]$  where the two cohomology groups are identified by the isotopy.<sup>31</sup>

We now continue studying the case where  $M$  is split by our original  $\Sigma$ .

**Lemma 4.8.** *Let  $f \in \Omega_c^1(M_+, \mathfrak{g})$ . Then*

$$[LR^*f|_{\Sigma}] = S^*[Lf|_{\Sigma}].$$

*Proof.* Denote by  $\phi_{\pm} \in \Omega^1(M_{\pm}, \mathfrak{g})$  the differential forms obtained by applying the maps  $j$  and  $j^{-1}$  exchanging  $M_+$  and  $M_-$  to the forms  $Lf|_{M_{\mp}}$ . Note that these are not compactly supported forms. Then we have

$$(4.9) \quad dLR^*f|_{M_{\pm}} = \star R^*f|_{M_{\pm}} = d\phi_{\pm}.$$

We wish to show that for any cohomology class  $\xi \in H^1(\Sigma, \mathbb{R})$

<sup>30</sup>Note that in the case where  $g = R^*f$ , Lemma 4.5 agrees with the result of Theorem 1.

<sup>31</sup>This is an indication that this theory has zero Hamiltonian.

$$\int_{\Sigma} ([ (LR^* f)|_{\Sigma} ] - S^*[Lf|_{\Sigma}]) \wedge \xi = 0.$$

Let  $\delta > 0$  be sufficiently small so that  $f$  is supported in  $M_+^{\delta}$ . Let  $\Sigma_{\pm} = \partial M_{\pm}^{\delta}$ . It will suffice to prove that there exists a basis  $\{\xi_i\}$  for  $H^1(\Sigma, \mathbb{R})$  so that for each  $i$ ,

$$\int_{\Sigma_+} ([ (LR^* f)|_{\Sigma_+} ] - [\phi_+|_{\Sigma_+}]) \wedge \xi_i = 0 \quad \text{or} \quad \int_{\Sigma_-} ([ (LR^* f)|_{\Sigma_-} ] - [\phi_-|_{\Sigma_-}]) \wedge \xi_i = 0.$$

Let  $i_{\pm} : \Sigma \rightarrow M_{\pm}$  denote the inclusion, and let  $\Lambda_{\pm} = i_{\pm}^*(H^1(M_{\pm}, \mathbb{R})) \subset H^1(\Sigma, \mathbb{R})$ . Since  $H^1(M, \mathbb{R}) = 0$ ,  $\Lambda_+ \cap \Lambda_- = \{0\}$  and  $\Lambda_+ \oplus \Lambda_- = H^1(\Sigma, \mathbb{R})$ . We claim that for any  $\xi \in \Lambda_{\pm}$ , there exists  $g \in \Omega_c^1(M_{\mp}, \mathbb{R})$  with  $d^*g = 0$  so that  $[Lg|_{\Sigma}] = \xi$ .<sup>32</sup> To see this, we take a form  $\tilde{\xi} \in \Omega^1(M, \mathbb{R})$  which is closed on  $M_{\pm}$  and on a neighborhood of  $\Sigma$ , and such that  $[\tilde{\xi}|_{\Sigma}] = \xi$ ; such a form exists by the definition of  $\Lambda_{\pm}$ . By the Hodge theorem, there exists  $\psi \in C^{\infty}(M)$  with  $d^*(\tilde{\xi} + d\psi) = 0$ . Let  $g = \star d(\tilde{\xi} + d\psi)$ . Then  $Lg = \tilde{\xi} + d\psi$ , so that  $[Lg|_{\Sigma}] = \xi$ . And  $g = \star d\tilde{\xi}$ , so that  $g$  is compactly supported on  $M_{\mp}$ .

Take a basis  $\{\xi_i^{\pm}\}$  for  $\Lambda_{\pm}$  and choose forms  $g_i^{\pm} \in \Omega_c^1(M_{\mp}, \mathbb{R})$  with  $d^*g_i^{\pm} = 0$  and  $[Lg_i^{\pm}|_{\Sigma}] = \xi_i^{\pm}$  for each  $i$ . Take  $\delta$  sufficiently small so that all the  $g_i^{\pm}$  (as well as  $f$ ) are supported in  $M_{\mp}^{\delta}$ . Consider first the case of the  $g_i^-$ . For each  $i$ , we wish to show that

$$\int_{\Sigma_-} (LR^* f - \phi_-) \wedge Lg_i^- = 0.$$

To see this, compute

$$\int_{\Sigma_-} (LR^* f - \phi_-) \wedge Lg_i^- = \int_{M_-^{\delta}} (dLR^* f - d\phi_-) \wedge Lg_i^- - (LR^* f - \phi_-) \wedge \star g_i^-.$$

But  $g_i^-|_{M_-} = 0$ , and  $(dLR^* f)|_{M_-} - d\phi_- = 0$  by (4.9).

A similar argument, substituting  $\Sigma_+$  for  $\Sigma_-$ , applies for the  $g_i^+$ .<sup>33</sup>

In the absence of Condition A, the same result applies to forms compactly supported on  $M_{\pm}^{\delta}$ . □

*End of Proof of Theorem 1:* The proof of Theorem 1 in the case of Condition A now follows from Lemma 4.5 and Lemma 4.8, along with condition (2.2). In the general case, where no finiteness condition is imposed on  $S$ , the same theorem holds restricting to forms compactly supported on the complement  $M_+^{\delta}$  of a collar neighborhood of  $\Sigma$ . □

We now form a Hilbert space using the bilinear form given by Theorem 1. Recall first that the space  $F_1^B(M_+) = \ker d^* \subset \Omega_c^1(M_+, \mathfrak{g})$  of smooth one-forms on  $M$  compactly supported on  $M_+$  is dense in  $\mathcal{F}_{1,+}^B$ . (In the absence of Condition A, replace  $M_+$  by  $M_+^{\delta}$ .) Let  $q : \mathcal{F}_{1,+}^B \otimes \mathcal{F}_{1,+}^B \rightarrow \mathbb{R}$  be the bilinear form given by

$$q(f, g) = \langle R^* f, Lg \rangle_{L_2}.$$

Then  $q$  is a positive semidefinite bilinear form on  $\mathcal{F}_{1,+}^B$ .

<sup>32</sup>We define the operator  $L$  on  $\mathbb{R}$ -valued one forms satisfying the gauge condition in the same way as for Lie algebra valued forms.

<sup>33</sup>In this case both of the relevant integrals vanish.

Let  $\mathcal{N}_1^B \subset \mathcal{F}_{1,+}^B$  denote the space of null elements of  $q$ . Then since  $q(f, f) \leq \|f\|_{\mathcal{F}_{1,+}^B}^2$ ,  $\mathcal{N}_1^B$  is a closed subspace of  $\mathcal{F}_{1,+}^B$ , and the quotient  $\mathcal{F}_{1,+}^B/\mathcal{N}_1^B$  can be completed in the norm given by  $q$  to give a Hilbert space  $H_1^B(\Sigma)$ . The computation in the proof of Theorem 1 shows there is a close relation between elements of  $\mathcal{F}_{1,+}^B$  and elements of  $\Lambda \subset H^1(\Sigma, \mathfrak{g})$ .

As in the proof of Theorem 1 we can construct a map  $\pi_\Sigma^B : F_1^B(M_+) \rightarrow H^1(\Sigma, \mathfrak{g})$  given by

$$\pi_\Sigma^B(f) = [Lf|_\Sigma].$$

**Proposition 4.10.** *The map  $\pi_\Sigma^B$  carries  $F_1^B(M_+)$  surjectively onto the image  $\Lambda \subset H^1(\Sigma, \mathfrak{g})$  of  $H^1(M_-, \mathfrak{g})$  in  $H^1(\Sigma, \mathfrak{g})$ , and carries the bilinear form  $q$  on  $F_1^B(M_+)$  to the positive definite bilinear form  $Q$  on  $\Lambda \subset H^1(\Sigma, \mathfrak{g})$ . (In the absence of Condition A, the same result holds with  $M_+$  replaced with  $M_+^\delta$ .)*

*Proof.* Let  $f, g$  be two smooth  $\mathfrak{g}$ -valued one forms on  $M$  compactly supported on  $M_+$ . By Lemma 4.5 and Lemma 4.8, we have  $q(f, g) = Q(R^*[Lf|_\Sigma], [Lg|_\Sigma])$ . Thus the map  $\pi_\Sigma^B : \ker(d_1^*) \subset \Omega^1(M_+, \mathfrak{g}) \rightarrow H^1(\Sigma, \mathfrak{g})$  given by  $\pi_\Sigma^B(f) = [Lf|_\Sigma]$  carries the bilinear form  $q$  (restricted to smooth forms) to the bilinear form  $Q$ .

To prove that the image of  $\pi_\Sigma^B$  lies in  $\Lambda$ , note that if  $f$  is a  $\mathfrak{g}$ -valued one form compactly supported on  $M_+$ , the one form  $Lf$  is closed on  $M_-$ . Thus the class  $[Lf|_\Sigma]$  extends to a class  $[Lf|_{M_-}]$ .

To prove surjectivity, we proceed as in the proof of Lemma 4.8. Suppose  $\xi$  is a closed  $\mathfrak{g}$ -valued one form on  $\Sigma$  extending to  $M_-$  as a closed form. Then  $\xi$  extends to a  $\mathfrak{g}$ -valued one form  $\tilde{\xi}$  on all of  $M$  whose restriction to  $M_-$  is closed; we may also take  $\tilde{\xi}$  closed in a neighborhood of  $\Sigma$ . By the Hodge Theorem and the vanishing of  $H^1(M, \mathbb{R})$ , there exists a  $\mathfrak{g}$ -valued function  $\phi$  such that  $d^*(\tilde{\xi} + d\phi) = 0$ . Then  $L \star d(\tilde{\xi} + d\phi) = \tilde{\xi} + d\phi$ , so that

$$[L(\star d(\tilde{\xi} + d\phi))|_\Sigma] = [\tilde{\xi}|_\Sigma + d(\phi|_\Sigma)] = [\xi].$$

Since  $\star d(\tilde{\xi} + d\phi) = \star d\tilde{\xi}$  is supported in  $M_+$ ,  $[\xi]$  is in the image of  $\pi_\Sigma$ . □

**Corollary 4.11.** *The map  $\pi_\Sigma^B$  extends to a map  $\pi_\Sigma^B : \mathcal{F}_{1,+}^B \rightarrow H^1(\Sigma, \mathfrak{g})$  and descends to an isomorphism  $\pi_\Sigma^B : \mathcal{F}_{1,+}^B/\mathcal{N}_1^B \rightarrow H_1^B(\Sigma) \simeq \Lambda$  carrying the positive definite form induced on  $\mathcal{F}_{1,+}^B/\mathcal{N}_1^B$  by  $q$  to the positive definite form  $Q$  on  $\Lambda$ .*

**Remark 4.12.** In view of Proposition 4.7, this construction can be performed for any splitting of  $M$  by a submanifold  $\tilde{\Sigma}$  isotopic to  $\Sigma$  into two components  $M_1, M_2$ . We obtain will a map  $\pi_{\tilde{\Sigma}}^B : \mathcal{F}_1^B(M_i) \rightarrow H^1(\tilde{\Sigma}, \mathfrak{g})$ ,  $i = 1, 2$ . Orient  $\tilde{\Sigma}$  as the boundary of  $M_1$ . The map  $\pi_{\tilde{\Sigma}}^B : \mathcal{F}_1^B(M_i) \rightarrow H^1(\tilde{\Sigma}, \mathfrak{g})$ ,  $i = 1, 2$ ; is an surjection onto the Lagrangian subspaces  $\Lambda$  and  $S^*\Lambda$ , respectively. We use the notation  $H_1^B(\tilde{\Sigma}) := \Lambda$  and continue to write  $\pi_{\tilde{\Sigma}}^B : \mathcal{F}_1^B(M_1) \rightarrow H_1^B(\tilde{\Sigma})$ , and similarly for the map  $\pi_{\tilde{\Sigma}}^B : \mathcal{F}_2^B(M_2) \rightarrow S^*\Lambda \simeq H_1^B(\tilde{\Sigma})$ .

**Proposition 4.13.** *If  $f_1 \in \mathcal{F}_1^B(M_1)$ ,  $f_2 \in \mathcal{F}_1^B(M_2)$ , then*

$$\int_M \star f_1 \wedge Lf_2 = \langle \pi_{\tilde{\Sigma}}^B(f_1), S^*\pi_{\tilde{\Sigma}}^B(f_2) \rangle_{H_1^B(\tilde{\Sigma})}.$$

*Furthermore, if  $\Sigma_1, \Sigma_2$  are two disjoint isotopic submanifolds of  $M$ , splitting  $M$  into three components,  $M_1, M_2$  and  $M_3 = \Sigma \times [0, 1]$ , with  $\partial M_3 = \Sigma_1 \cup \Sigma_2$ , then if  $f$  is compactly supported in  $M_1$  (or  $M_2$ ), then*

$$\pi_{\Sigma_1}(f) = \pi_{\Sigma_2}(f).$$

**4.2. Reflection positivity for  $\mathcal{F}_+^B$ .** We have now proved that the bilinear form  $q$  gives a reflection positive functional on  $\mathcal{F}_{1,+}^B$ . The form  $q(a, b)$  gives the expectation of the product of the linear functions on  $\mathbf{A}$  associated to the one forms  $a$  and  $b$  in the formal functional integral (4.1)

In this section we show that reflection positivity extends to  $\mathcal{R}_+$ , and hence to  $\mathcal{F}_{P,+}^B$ . To make a sensible definition corresponding to the formal functional integral (4.1), we define a functional  $\Phi : \mathcal{R} \rightarrow \mathbb{R}$ .

**Definition 4.14.** Let  $\Phi : \mathcal{R} \rightarrow \mathbb{R}$  be the linear functional which is zero on all monomials of the form  $f_1 \otimes \cdots \otimes f_n \in \mathcal{R}$  (where  $f_1, \dots, f_n \in F_1^B$ ) where  $n$  is odd, and whose value on a monomial  $f_1 \otimes \cdots \otimes f_n \in \mathcal{R}$  (where  $f_1, \dots, f_n \in F_1^B$ ) where  $n$  is even is given by

$$(4.15) \quad \Phi(f_1 \otimes \cdots \otimes f_n) = \frac{1}{2^{n/2}(n/2)!} \sum_{\sigma \in \Sigma_n} \langle f_{\sigma(1)}, Lf_{\sigma(2)} \rangle_{L_2} \cdots \langle f_{\sigma(n-1)}, Lf_{\sigma(n)} \rangle_{L_2}.$$

Here the inner product  $\langle \cdot, \cdot \rangle_{L_2}$  is the  $L_2$  inner product of  $\mathfrak{g}$ -valued one forms on  $M$ .

This functional corresponds to the formal functional integral

$$\frac{1}{Z_{gauge}} \int_{\mathcal{A}} e^{\frac{1}{2} \text{tr} \int_M AdA} f_1(A) \cdots f_n(A) dA$$

where  $Z_{gauge}$  is a formally infinite normalization constant.

**Remark 4.16.** Note also that we may scale the metric by a positive factor  $\alpha$  (corresponding to a scaling of the quadratic term in the free Lagrangian in (1.19)) to obtain a functional  $\Phi_\alpha$  obtained by scaling the right hand side of (4.15) by a factor of  $\alpha^{-\frac{n}{2}}$ .

Alternatively, we may use the fact that, given our choice of  $s = 4$ , the operator  $\Delta^{-s/2} L \Delta^{-s/2}$  is symmetric and Hilbert Schmidt,<sup>34</sup> as an operator on the  $L_2$  completion  $\overline{\mathcal{F}_1^B}$  of  $\mathcal{F}_1^B$ , and hence gives a symmetric element  $L_s \in \overline{\mathcal{F}_1^B} \otimes \overline{\mathcal{F}_1^B}$ , to write

$$(4.17) \quad \Phi(f_1 \otimes \cdots \otimes f_n) = \frac{1}{2^{n/2}(n/2)!} \langle L_s \otimes \cdots \otimes L_s, \Delta^{s/2} f_1 \otimes \cdots \otimes \Delta^{s/2} f_n \rangle_{\text{Sym}^* \overline{\mathcal{F}_1^B}}$$

Thus, given an inhomogeneous polynomial  $P = \sum_l P_l$  of degree  $d = 2k$ , where  $P_l$  is homogeneous of degree  $l$ , we may write

$$\Phi(P) = \left\langle \sum_{j=0}^k \frac{1}{2^j j!} L_s^{\otimes j}, \sum_l (\Delta^{s/2})^{\otimes l} P_l \right\rangle_{\text{Sym}^* \overline{\mathcal{F}_1^B}}$$

which shows that  $\Phi$  extends to a bounded linear functional on  $\mathcal{F}_d^B$  for any  $d$ : that is, for  $P \in \mathcal{F}_d^B$ ,

$$(4.18) \quad |\Phi(P)| \leq C_d \|P\|_{\mathcal{F}_d^B}$$

<sup>34</sup>For this property  $s > \frac{1}{4}$  would suffice.

for some constant  $C_d$  which depends on  $d$ . And  $\Phi$  gives a (possibly-unbounded) linear functional on  $\mathcal{F}_P^B$  (see Section 4.3).

We next define a bilinear form on  $\mathcal{R}_+$  which, since it extends the form  $q$  on linear polynomials, we continue to denote by  $q : \mathcal{R}_+ \otimes \mathcal{R}_+ \rightarrow \mathbb{R}$ , by

$$(4.19) \quad q(P, Q) = \Phi((R^*P)Q).$$

The bilinear form  $q$  then also extends to  $\mathcal{F}_{P,+}^B$  and is bounded on any  $\mathcal{F}_{d,+}^B$ .

We prove the following

**Theorem 2.** *Let  $P \in \mathcal{R}_+$ .*

*Then*

$$q(P, P) \geq 0.$$

Since  $\mathcal{R}_+ \cap \mathcal{F}_{d,+}^B$  is dense in  $\mathcal{F}_{d,+}^B$  for any  $d$ ,<sup>35</sup> this same result holds for  $\mathcal{F}_{d,+}^B$ . We denote the nullspace of the bilinear form  $q$  on  $\mathcal{F}_{d,+}^B$  by  $\mathcal{N}_d^B$ . The space  $\mathcal{N}_d^B$  is a closed subspace of  $\mathcal{F}_{d,+}^B$  by boundedness of  $\Phi$  on  $\mathcal{F}_{d,+}^B$ . The quotient  $\mathcal{F}_{d,+}^B / \mathcal{N}_d^B$  comes equipped with a bilinear form we also denote by  $q$  which is positive definite, and hence can be completed to a Hilbert space  $H_{\leq d}^B(\Sigma)$ . The resulting map  $\pi_{\Sigma}^B : \mathcal{F}_{d,+}^B \rightarrow H_{\leq d}^B(\Sigma)$  is bounded. Writing  $H_d^B(\Sigma) = H_{\leq d}^B(\Sigma) / H_{\leq d-1}^B(\Sigma)$ , taking direct sums and writing  $H_P^B(\Sigma) = \bigoplus_d H_d^B(\Sigma)$ , we obtain a map  $\pi_{\Sigma}^B : \mathcal{F}_{P,+}^B \rightarrow H_P^B(\Sigma)$ , which is bounded in any given degree. The space  $H_P^B(\Sigma)$  can in turn be completed to a Hilbert space  $H^B(\Sigma)$ . We will show below (see Proposition 4.26) that  $H_P^B(\Sigma) = \text{Sym}^* \Lambda \subset \text{Sym}^* H^1(\Sigma, \mathfrak{g})$ .

The proof of Theorem 2 requires the following Lemma, giving reflection invariance of the Wick ordering for forms compactly supported on  $M_+$  (or  $M_+^{\delta}$ , in the case where Condition A is absent).

**Lemma 4.20.** *Let  $f \in \Omega^1(M, \mathfrak{g})$  satisfy  $d^*f = 0$  and be compactly supported on  $M_+$ . Then*

$$\text{tr} \int_M f \wedge \star Lf = \text{tr} \int_M R^*f \wedge \star LR^*f.$$

*(In the absence of Condition A, replace  $M_+$  by  $M_+^{\delta}$ .)*

*Proof.* In the case of Condition B, this follows from the invariance of the metric under the involution  $R$ .<sup>36</sup>

In the case of Condition A, we may not have a globally defined involution  $R$ ; we have only an identification  $j$  of  $M_+$  with  $M_-$ . But write  $\psi = Lf$ , and define  $\phi \in \Omega^1(M_+, \mathfrak{g})$  by  $\phi = R^*((LR^*f)|_{M_-})$ ;<sup>37</sup> note that  $\phi$  is not a form with compact support. Then

<sup>35</sup>Recall that  $\mathcal{F}_{d,+}^B$  are the inhomogeneous polynomials of degree  $d$ .

<sup>36</sup>Note that for this invariance of the Wick ordering under reflection, it is crucial that the reflection  $R$  is orientation preserving, and hence preserves the propagator  $L$ . This is in contrast to the form of reflection positivity in the one particle space in [25], where the reflection is orientation reversing and reverses the sign of the propagator.

<sup>37</sup>This is a slight abuse of notation. What we mean by  $\phi = R^*((LR^*f)|_{M_-})$  is the form  $\phi \in \Omega^1(M_+, \mathfrak{g})$  (not of compact support) obtained from the form  $LR^*f|_{M_-} \in \Omega^1(M_-, \mathfrak{g})$  via the identification  $j$  of  $M_+$  with  $M_-$  in (2.3). We will only ever consider  $\phi$  restricted to the support of  $d\phi = \star f$  (which is also the support of  $d\psi$ ), so no problems arise.

$$(4.21) \quad d\psi|_{M_+} = \star f|_{M_+} = d\phi.$$

Thus

$$\operatorname{tr} \int_M \star f \wedge Lf = \operatorname{tr} \int_{M_+} \psi \wedge d\psi$$

and

$$\operatorname{tr} \int_M \star R^* f \wedge LR^* f = \operatorname{tr} \int_{M_+} \phi \wedge d\phi.$$

Then

$$(4.22) \quad \operatorname{tr} \int_{M_+} \psi \wedge d\psi - \phi \wedge d\phi = \operatorname{tr} \int_{M_+} (\psi - \phi) \wedge d\psi + \operatorname{tr} \int_{M_+} \phi \wedge d(\psi - \phi).$$

The second term on the right hand side of (4.22) vanishes by (4.21). The first term is given by

$$\operatorname{tr} \int_{M_+} (\psi - \phi) \wedge d\psi = \operatorname{tr} \int_{M_+} d(\psi - \phi) \wedge \psi - \operatorname{tr} \int_{\Sigma} [(LR^* f - R^* Lf)|_{\Sigma}] \wedge S^*[Lf|_{\Sigma}].$$

But  $d(\psi - \phi) = 0$  by equation (4.21), and  $[(LR^* f - R^* Lf)|_{\Sigma}] = 0$  by Lemma 4.8.

In the absence of Condition A, replace  $M_+$  by  $M_+^{\delta}$  and the argument goes through as before. □

We now return to the Proof of Theorem 2.

*Proof.* Since any polynomial in  $\mathcal{R}_+$  can be written as a linear combination of  $L$ -Wick ordered polynomials in  $\mathcal{O}_+$ , it suffices to prove reflection positivity for  $L$ -Wick ordered polynomials. By Lemma 4.20, for any  $L$ -Wick ordered monomial  $m_{f_1, \dots, f_n}$  where  $f_1, \dots, f_n \in F_1^B(M_+)$ ,

$$R^* : m_{f_1, \dots, f_n} := m_{R^* f_1, \dots, R^* f_n} :$$

Then Definition 4.14 shows that for any  $g_1, \dots, g_m \in F_1^B(M_+)$ , the expression

$$\Phi(R^* : m_{f_1, \dots, f_n} :: m_{g_1, \dots, g_m} :)$$

vanishes if  $m \neq n$  and is given by<sup>38</sup>

<sup>38</sup>Some comments about Wick ordering: This is most convenient to explain for the case of Hilbert Schmidt operators. Suppose  $L$  is a symmetric Hilbert Schmidt operator on a Hilbert space  $H$ . Then there exists  $v \in \operatorname{Sym}^2 H \subset H \otimes H$  with

$$(4.23) \quad \langle v, x \otimes y \rangle_{\operatorname{Sym}^2 H} = \langle x, Ly \rangle$$

for all  $x, y \in H$ .

Suppose  $P$  is a polynomial in  $\operatorname{Sym}^* H$  and define

$$\Phi(P) = \langle e^{iv} P, 1 \rangle_{\operatorname{Sym}^* H}$$



$$(4.25) \quad \Phi(R^* : m_{f_1, \dots, f_n} :: m_{g_1, \dots, g_m} :) = \langle R^* f_1 \otimes \dots \otimes R^* f_n, Lg_1 \otimes \dots \otimes Lg_n \rangle_{L_2}$$

if  $m = n$ ; here the inner product is the inner product on the symmetric algebra coming from  $L_2$  inner product on  $\mathfrak{g}$  valued forms. Therefore, the inner product  $q$  on polynomials in  $\mathcal{O}_+$  is the same as the inner product on  $\text{Sym}^* \mathcal{F}_{1,+}^B$  induced by the inner product given by  $q$  on  $\mathcal{F}_{1,+}^B$ . It is therefore positive semidefinite.

In the absence of Condition A, the same result holds, for forms compactly supported on the complement  $M_+^\delta$  of a collar neighborhood of  $\Sigma$ . □

In fact we have proved

**Proposition 4.26.** *The Hilbert space  $H_d^B(\Sigma)$  is given by*

$$H_d^B(\Sigma) \simeq \text{Sym}^d H_1^B(\Sigma).$$

*Proof.* The quadratic form associated to  $q$  restricted to  $\mathcal{F}_{1,+}^B$  defines a bounded, self adjoint, nonnegative operator on  $\mathcal{F}_{1,+}^B$ . We can therefore find an orthonormal basis for  $\mathcal{F}_{1,+}^B$  consisting of a finite number of smooth forms  $k_1, \dots, k_g$  forming a basis for the orthocomplement  $H_1^B(\Sigma)$  of the kernel  $\mathcal{N}_1^B$ , and orthogonal also for the bilinear form  $q$ , alongside an orthonormal basis of smooth forms  $\{l_i\}_{i=1}^\infty$  for  $\mathcal{N}_1^B$ . This gives us an orthonormal basis for each  $\text{Sym}^d(\mathcal{F}_{1,+}^B)$  consisting of the elements  $\{l_j \otimes p\}$ , where  $p = r_1 \otimes \dots \otimes r_{d-1}$  is any monomial of degree  $d-1$  in the  $k_i$ 's and  $l_i$ 's (i.e.  $r_i \in \{k_1, \dots, k_g, l_1, \dots\}$  for all  $i$ ), alongside the monomials  $k_{i_1} \otimes \dots \otimes k_{i_d}$ . This basis is not orthogonal for  $q$ . However, this basis for each  $\text{Sym}^{d'}(\mathcal{F}_{1,+}^B)$  yields a basis for  $\text{Sym}^{\leq d}(\mathcal{F}_{1,+}^B)$  given by the elements  $: m_{l_i, r_1, \dots, r_{d'}} :$ , where  $d' \leq d$ , and where again  $r_i \in \{k_1, \dots, k_g, l_1, \dots\}$  for all  $i$ , alongside the monomials  $: m_{k_{i_1} \dots k_{i_{d'}}} :$ , where  $d' \leq d$ . This basis is orthogonal for the bilinear form  $q_d = q|_{\text{Sym}^{\leq d}(\mathcal{F}_{1,+}^B)}$ . By (4.25), the bilinear form  $q_d$  has nullspace given by the span of the  $: m_{l_i, r_1, \dots, r_{d'}} :$ , and

where  $i_v$  is interior product in  $\text{Sym}^* H$ ; that is,

$$i_v(x \otimes y) = \langle v, x \otimes y \rangle_{\text{Sym}^2 H}.$$

Note that if  $L > 0$ , the functional  $\Phi$  has an alternative definition in terms of Gaussian integrals, as Gaussian measure with covariance  $L$ , where the Hilbert space  $\mathbf{H}^*$  of (1.3) is the completion of  $H$  in the norm given by  $\langle \cdot, L \cdot \rangle$ . The definition we have given for  $\Phi$  does not require positivity, although where positivity is not present,  $\Phi$  does not correspond to a measure, and is defined as it stands only for polynomials.

Write, for  $P \in \text{Sym}^* H$ ,

$$: P := e^{-i_v} P.$$

Then if  $P, Q \in \text{Sym}^* H$ ,

$$(4.24) \quad \Phi(: P :: Q :) = \langle P, \left( \sum_{k=0}^{\infty} \frac{1}{k!} \text{Sym}^k L \right) Q \rangle_{\text{Sym}^* H}.$$

If  $L$  is not Hilbert Schmidt, similar formulas hold as long as both sides of equation (4.24) are well defined. The Wick ordering can be defined by using equation (4.23) to replace the interior product  $i_v$  with an appropriate sequence of inner products. Equation (4.24) can be proved by cutting off  $L$  to obtain a sequence of finite rank operators, and then taking the limit on both sides. In our case an alternative argument uses transfer of regularity and the Hilbert Schmidt operator  $L_s$  of equation (4.17).

is nondegenerate, and equal to the bilinear form induced by  $q_1$  on  $\text{Sym}^{\leq d} H_1^B(\Sigma)$ , on the span of the  $: m_{k_{i_1} \dots k_{i_{d'}}} : .$  The result follows.  $\square$

In fact, since we have seen in Corollary 4.11 that there is an isometry  $\mathcal{F}_{1,+}^B / \mathcal{N}_1^B \simeq \Lambda$ , we have shown that there is an isometry

$$H^B(\Sigma) = (\text{Sym}^* \Lambda)^-,$$

identifying  $H^B(\Sigma)$  with the completion of the symmetric algebra  $\text{Sym}^* \Lambda \subset \text{Sym}^* H^1(\Sigma, \mathfrak{g})$ <sup>39</sup> in the norm induced on it by the norm on  $\Lambda$  arising from inner product coming from the orientation of  $\Sigma$ , the inner product on  $\mathfrak{g}$ , and the diffeomorphism  $S$  : that is the inner product

$$Q(a, b) = \text{tr} \int_{\Sigma} S^* a \wedge b$$

for  $a, b \in \Lambda$ . In terms of this identification, the map  $\pi_{\Sigma}^B$  is given on the dense subset of  $\mathcal{F}_{d,+}^B$  (for each  $d$ ) given by Wick ordered polynomials by writing<sup>40</sup>

$$(4.27) \quad \pi_{\Sigma}^B(: m_{f_1, \dots, f_{d'}} :) = [Lf_1|_{\Sigma}] \otimes \cdots \otimes [Lf_{d'}|_{\Sigma}]$$

for each  $d' \leq d$  and extending by linearity. The isomorphism induced by the map  $\pi_{\Sigma}^B$  then carries the norm on  $H_d^B(\Sigma)$  isometrically to the norm on  $\text{Sym}^d \Lambda$ . In summary we have proved

**Theorem 3.** *The map  $\pi_{\Sigma}^B$  given by (4.27) descends to an isomorphism*

$$H^B(\Sigma) \simeq (\text{Sym}^* \Lambda)^-$$

where  $(\text{Sym}^* \Lambda)^-$  is the completion of  $\text{Sym}^* \Lambda$  in the norm induced on  $\text{Sym}^* \Lambda$  by the positive definite bilinear form  $\text{Sym}^*(Q)$ .

The identification of  $H^B(\Sigma)$  with the completion of a symmetric algebra endows  $H^B(\Sigma)$  with a densely defined, self adjoint, nonnegative operator  $N_B$  on  $H^B(\Sigma)$  given by the polynomial degree, with domain  $\mathcal{D} = (\text{Sym}^* \Lambda) \subset H^B(\Sigma)$ .

**Remark 4.28.** In these terms the scaling of the quadratic term in the free Lagrangian by a factor of  $\alpha$  (equivalent to replacing  $\Phi$  by  $\Phi_{\alpha}$ ) is equivalent to replacing the metric  $\langle \cdot, \cdot \rangle$  on the symmetric algebra by the metric  $\langle \cdot, e^{-\log \alpha N_B} \cdot \rangle$ .

**Remark 4.29.** In view of Proposition 4.7, we can find a formula similar to (4.25) for any splitting of  $M$  by a compact connected submanifold  $\tilde{\Sigma}$  into two components  $M_1, M_2$ . Define  $\tilde{Q} : H^1(\tilde{\Sigma}, \mathfrak{g}) \otimes H^1(\tilde{\Sigma}, \mathfrak{g}) \rightarrow \mathbb{R}$  by

$$\tilde{Q}(a, b) = \text{tr} \int_{\tilde{\Sigma}} a \cup b.$$

Note that if  $\tilde{\Sigma} = \Sigma$ ,  $\tilde{Q}(a, b) = Q(S^* a, b)$ .<sup>41</sup>

<sup>39</sup>It is also possible to consider  $\text{Sym}^* \Lambda$  as an abelian subalgebra of the Weyl algebra on the vector space  $H^1(\Sigma, \mathfrak{g})$  equipped with the symplectic form  $\tilde{Q}$  given by the intersection form on  $\Sigma$  and defined in Remark 4.29 below; the subalgebra is abelian because  $\Lambda$  is Lagrangian in this symplectic form.

<sup>40</sup>Note that the map  $\pi_{\Sigma}^B$  is a filtered map, but not a graded map, of polynomials.

<sup>41</sup>The form  $\tilde{Q}$  is closely related to the Atiyah-Bott symplectic form on the space of connections on  $\Sigma$ .

**Proposition 4.30.** *Let  $\tilde{\Sigma} \subset M$  be a compact connected submanifold of  $M$  whose complement has two components  $M_1, M_2$ . Orient  $\tilde{\Sigma}$  as the boundary of  $M_1$ . Then if  $: m_{f_1, \dots, f_n} : \in \mathcal{O}(M_1)$  and  $: m_{g_1, \dots, g_n} : \in \mathcal{O}(M_2)$ , we have*

$$\Phi(: m_{f_1, \dots, f_n} :: m_{g_1, \dots, g_n} :) = \text{Sym}^* \tilde{Q}([Lf_1|_{\tilde{\Sigma}}] \otimes \cdots \otimes [Lf_n|_{\tilde{\Sigma}}], [Lg_1|_{\tilde{\Sigma}}] \otimes \cdots \otimes [Lg_n|_{\tilde{\Sigma}}])$$

We therefore define  $\hat{\pi}_{\tilde{\Sigma}}^B : \mathcal{F}_P^B(M_1) \oplus \mathcal{F}_P^B(M_2) \longrightarrow \text{Sym}^*(H^1(\tilde{\Sigma}, \mathfrak{g}))$  by taking

$$\hat{\pi}_{\tilde{\Sigma}}^B(: m_{f_1, \dots, f_n} :) = [Lf_1|_{\tilde{\Sigma}}] \otimes \cdots \otimes [Lf_n|_{\tilde{\Sigma}}]$$

and

$$\hat{\pi}_{\tilde{\Sigma}}^B(: m_{g_1, \dots, g_n} :) = [Lg_1|_{\tilde{\Sigma}}] \otimes \cdots \otimes [Lg_n|_{\tilde{\Sigma}}]$$

where  $: m_{f_1, \dots, f_n} : \in \mathcal{O}(M_1)$  and  $: m_{g_1, \dots, g_n} : \in \mathcal{O}(M_2)$ , and extending by linearity to  $\mathcal{F}_P^B(M_1) \oplus \mathcal{F}_P^B(M_2)$ . We then have the following

**Proposition 4.31.** *Let  $\tilde{\Sigma}$  be any submanifold of  $M$  isotopic to  $\Sigma$  as oriented manifolds, and splitting  $M$  into two components  $M_1, M_2$ . Assume the orientation of  $\tilde{\Sigma}$  agrees with that it inherits as the boundary of  $M_1$ . Then if  $P \in \mathcal{F}_P^B(M_1), Q \in \mathcal{F}_P^B(M_2)$ ,*

$$\Phi(PQ) = \langle (\hat{\pi}_{\tilde{\Sigma}}^B(P)), S^* \hat{\pi}_{\tilde{\Sigma}}^B(Q) \rangle_{H^B(\Sigma)}$$

where elements of  $\text{Sym}^* \Lambda \subset \text{Sym}^* H^1(\tilde{\Sigma}, \mathfrak{g})$  are identified with elements of  $H^B(\Sigma)$  using the isotopy.

**4.3. Is there a relation between  $\mathcal{F}_+^B$  and  $H^B(\Sigma)$ ?.** We consider the following question.

**Question:** Is there a bound

$$(4.32) \quad |\Phi(a)| \leq C \|a\|_{\mathcal{F}^B}$$

for all  $a \in \mathcal{R}_+$ ?

In quantum field theories with polynomial interactions in two dimensions, such a bound is immediate, but in our case, where the functional  $\Phi$  is not defined using the measure  $\mu$  of Remark 3.1, the existence of such a bound, especially since the operator  $L$  is not positive, is not so easy to see: A positive linear functional on a  $\mathbb{C}^*$  algebra is bounded, but I do not know of a comparable statement for a reflection positive functional.<sup>42</sup> The existence of such a bound (and the corresponding bound for Fermions) would simplify considerably the proof of the existence of the interacting theory: The elements  $e^{i\lambda \Xi_B^+}$  in  $\mathcal{F}_+ \otimes \mathbb{C}$  would immediately descend<sup>43</sup> to elements of  $H(\Sigma) \otimes \mathbb{C}$ , as would their counterparts for  $M_-$ ; in fact, this would hold for any  $M_+, M_-$  having common boundary  $\Sigma$ , without any need for a reflection. This would lead to the existence of partition functions in manifolds without reflection, and to strong bounds on those partition functions. It would be worthwhile looking into this as a guide to what kind of bounds, whether (4.32) or some weaker version, may be plausible. Thus there is some interplay between bounds of the form (4.32) and bounds on partition functions in topological quantum field theory. One possibility is that growth properties of manifold invariants which would preclude

<sup>42</sup>The functional  $\Phi$  can be extended to an algebra of bounded functions on  $\mathbf{A}$  by taking imaginary exponentials of linear functionals. For such functions arising from exponentiation of linear functions compactly supported on  $M_+$ ,  $\|(\overline{Rf})f\|_\infty = \|f\|_\infty^2$ , which resembles a  $\mathbb{C}^*$  condition. Another possible point of reference is Tomita-Takesaki theory.

<sup>43</sup>There is an additional point of concern about interpreting  $\Xi_B^+$  as an element of  $\mathcal{F}_+$ ; see Section 6.

such a bound may contradict the existence of certain Heegaard splittings, so there is a potential for topological consequences.

One way to approach this question would be to replace the measure  $d\mu$  on  $\mathbf{A}$  in Remark 3.1 with a measure  $d\mu_K$  with covariance  $L + KI$ , where  $K \in \mathbb{R}$  satisfies  $K > \|L\|$ . The covariance  $L + KI$  is positive, and the addition of  $L$  is inconsequential for the singularities of the fields, so it yields a measure on a Banach space of connections  $\mathbf{A}$  corresponding to the case  $s = 0$  (instead of  $s = 4$ ) in the construction of Remark 3.1; this space carries the White Noise measure on the space of connections. But the covariance  $L + KI$  is also reflection positive, since

$$(4.33) \quad \langle R^*f, (L + KI)f \rangle = \langle R^*f, Lf \rangle$$

for  $f$  compactly supported in  $M_+$ ; and (4.33) shows that the Hilbert space obtained from  $d\mu_K$  by taking fields compactly supported on  $M_+$  and then taking the quotient by the null vectors for the reflection positive functional is our same  $H^B(\Sigma)$ . Thus, replacing the functional  $\Phi$  by the expectation in the measure  $d\mu_K$ , the analog of bound (4.32) is satisfied, and we obtain a map from the configuration space of fields compactly supported on  $M_+$  to  $H^B(\Sigma)$ .

However, expectations in this new functional do not correspond in any way which I understand to Chern-Simons gauge theory; and I do not know how to take the limit  $K \rightarrow 0$ .

## 5. THE FERMION HILBERT SPACE

In this section we study the Fermion Hilbert space. Since the Fermions are ghosts, we may expect that positivity might not be straightforward, and there are in fact subtleties. The Fermion Hilbert space  $H^F(\Sigma)$  is finite dimensional, so those subtleties can be overcome.

We will see that as a vector space,  $H^F(\Sigma) = \bigwedge^*(\mathfrak{g} \oplus \mathfrak{g})$ , and therefore  $H^F(\Sigma)$  has the structure of an alternating algebra, which we can think of as the alternating algebra on the even cohomology  $H^0(\Sigma, \mathfrak{g}) \oplus H^2(\Sigma, \mathfrak{g})$ , just as the Bosonic Hilbert space was a symmetric algebra on (part of) the odd cohomology. This alternating algebra has a positive definite metric arising from the positive definite metric on  $\mathfrak{g} \oplus \mathfrak{g}$ . This positive definite form does arise from a reasonable form of reflection positivity on the linear part of the configuration space. But the required "twist" behaves badly under Wick ordering, so we do not get a reasonable form of reflection positivity for polynomials. On the other hand, the nondegenerate split linear bilinear form on  $\mathfrak{g} \oplus \mathfrak{g}$  does arise from reasonable formulas on the configuration space.<sup>44</sup>

Since  $H^F(\Sigma)$  is finite dimensional, the two inner products are related by a linear transformation; this gives a bounded operator on  $H^B(\Sigma) \otimes H^F(\Sigma)$ , allowing us to work with Hilbert spaces when needed in treating the interaction term.

---

<sup>44</sup>Note that it is conceivable that a subalgebra of the polynomials in the Fermi fields satisfies some form of reflection positivity even in the split linear metric. One example of such a phenomenon is given in [14]. I was not able to find such a subalgebra rich enough to contain the interaction term, and it is conceivable that due to the role of the ghosts in the interaction, no such subalgebra containing the interaction term can exist. It would be interesting to investigate the type of Spin-Statistics Theorems (see e.g. [24]) that apply in this situation.

5.1. **The one particle configuration space  $\mathcal{F}_1^F(U)$ .** Recall that the one particle configuration space  $\mathcal{F}_1^F(U)$  associated to an open set  $U \subset M$  was given by the completion of the space

$$F_1^F(U) = \Omega^0(U, \mathfrak{g})^\perp \oplus (\ker(d_2^*) \cap \Omega_c^2(U, \mathfrak{g}))$$

where the space  $\Omega^0(U, \mathfrak{g})^\perp$  is the space of functions on  $M$  which are constant outside a compact subset of  $U$ , and whose integral over  $M$  is zero. Our first theorem is the analog of Theorem 1 for Fermions.

Suppose  $\tilde{\Sigma}$  is a compact connected oriented submanifold of  $M$ , splitting  $M$  into two connected components  $U, V$ . Let  $f \in \Omega^0(U, \mathfrak{g})^\perp$  and  $g \in \ker(d_2^*) \cap \Omega_c^2(V, \mathfrak{g})$ . Then

$$(5.1) \quad \begin{aligned} f|_{M-U} &= c_f \\ Lg|_{M-V} &= c_g \end{aligned}$$

where  $c_f, c_g$  are constants.

**Theorem 4.** *Let  $\tilde{\Sigma}$  be a compact connected oriented two-dimensional submanifold of  $M$ , and suppose  $M - \tilde{\Sigma}$  consists of two components  $U, V$ . Let  $f \in \Omega^0(U, \mathfrak{g})^\perp$  and  $g \in \ker(d_2^*) \cap \Omega_c^2(V, \mathfrak{g})$ , with  $c_f, c_g$  as above.*

*Then*

$$(5.2) \quad \operatorname{tr} \int_M \star f \wedge Lg = -\operatorname{vol}(M) \operatorname{tr} c_f c_g.$$

*Proof.* We have

$$(5.3) \quad \begin{aligned} \operatorname{tr} \int_M \star f \wedge Lg &= \operatorname{tr} \int_U \star f c_g + \operatorname{tr} \int_V c_f \star Lg = \\ &= \operatorname{tr} c_g \left( - \int_V \star f \right) + \operatorname{tr} c_f \left( - \int_U \star Lg \right), \end{aligned}$$

using the fact that  $\int_M \star f = \int_M \star Lg = 0$ .

But

$$\operatorname{tr} c_g \left( - \int_V \star f \right) + \operatorname{tr} c_f \left( - \int_U \star Lg \right) = -\operatorname{vol}(M) \operatorname{tr} c_f c_g$$

as needed. □

We now define  $\pi_{\tilde{\Sigma}}^F : F_1^F(U) \longrightarrow \mathfrak{g} \oplus \mathfrak{g}$  by

$$\pi_{\tilde{\Sigma}}^F(f, g) = (c_f, c_g)$$

where as in (5.1)

$$(5.4) \quad \begin{aligned} f|_{M-U} &= c_f \\ Lg|_{M-U} &= c_g, \end{aligned}$$

and likewise  $\pi_{\tilde{\Sigma}}^F : F_1^F(V) \longrightarrow \mathfrak{g} \oplus \mathfrak{g}$  by

$$\pi_{\tilde{\Sigma}}^F(f, g) = (c_f, c_g)$$

where again

$$(5.5) \quad \begin{aligned} f|_{M-V} &= c_f \\ Lg|_{M-V} &= c_g. \end{aligned}$$

Then the map  $\pi_{\tilde{\Sigma}}^F$  extends to  $\mathcal{F}_1^F(U), \mathcal{F}_1^F(V)$ .<sup>45</sup> And if  $df$  and  $g$  are compactly supported in the connected components of  $U, V$  of  $M - \tilde{\Sigma}$  as above,

$$\mathrm{tr} \int_M \star f \wedge Lg = \langle \pi_{\tilde{\Sigma}}^F(f), \pi_{\tilde{\Sigma}}^F(g) \rangle_{\mathfrak{g} \oplus \mathfrak{g}}$$

where the inner product on  $\mathfrak{g} \oplus \mathfrak{g}$  is the split linear inner product<sup>46</sup>

$$(5.6) \quad \langle (a, b), (a', b') \rangle_{\mathfrak{g} \oplus \mathfrak{g}} = \mathrm{vol}(M)(\langle a, b' \rangle_{\mathfrak{g}} + \langle b, a' \rangle_{\mathfrak{g}}).$$

Note that, for any  $(f, g) \in F_1^F(M)$ ,

$$(5.7) \quad \mathrm{tr} \int_M \star g \wedge Lf = -\mathrm{tr} \int_M \star f \wedge Lg,$$

so that to obtain the split linear pairing on  $\mathfrak{g} \oplus \mathfrak{g}$ , the adjoint in  $\mathcal{F}_1^F(M)$  must include an additional negative sign on factors of  $c_2$ . In other words we write for  $(f, g) \in \mathcal{F}_1^F(M)$

$$(5.8) \quad f^c = f, g^c = -g,$$

or

$$(5.9) \quad (f, g)^c = (f, -g),$$

and then

$$(5.10) \quad \mathrm{tr} \int_M \star g^c \wedge Lf = \mathrm{tr} \int_M \star f^c \wedge Lg,$$

so that we that the following restatement of Theorem 4:

**Proposition 5.11.** *Let  $\tilde{\Sigma}$  be a compact connected oriented two-dimensional submanifold of  $M$ , and suppose  $M - \tilde{\Sigma}$  consists of two components  $U, V$ . Let  $f \in \Omega^0(U, \mathfrak{g})^\perp$  and  $g \in \ker(d_2^*) \cap \Omega_c^2(V, \mathfrak{g})$ . Then*

$$(5.12) \quad \mathrm{tr} \int_M \star g^c \wedge Lf = \mathrm{tr} \int_M \star f^c \wedge Lg = \langle \pi_{\tilde{\Sigma}}^F(f), \pi_{\tilde{\Sigma}}^F(g) \rangle_{\mathfrak{g} \oplus \mathfrak{g}}.$$

The split linear pairing on  $\mathfrak{g} \oplus \mathfrak{g}$  is not positive definite, but it is closely related to a positive definite pairing. Let  $J : \mathfrak{g} \oplus \mathfrak{g} \rightarrow \mathfrak{g} \oplus \mathfrak{g}$  be defined by

$$J(a, b) = (b, a).$$

Then the positive definite inner product on  $\mathfrak{g} \oplus \mathfrak{g}$  given by

<sup>45</sup>Where there is no danger of confusion, we denote  $(f, 0) \in F_1^F(M)$  simply by  $f$ , likewise  $(0, g) \in F_1^F(M)$  simply by  $g$ .

<sup>46</sup>Recall that the inner product on  $\mathfrak{g}$  is given by  $\langle a, b \rangle_{\mathfrak{g}} = -\mathrm{tr}(ab)$  for  $a, b \in \mathfrak{g}$ , which accounts for the sign difference between (5.2) and (5.6).

$$\langle (a, b), (a', b') \rangle_+ = \text{vol}(M)(\langle a, a' \rangle + \langle b, b' \rangle)$$

is related to the split linear pairing by

$$\langle \alpha, \beta \rangle = \langle \alpha, J\beta \rangle_+$$

where  $\alpha, \beta \in \mathfrak{g} \oplus \mathfrak{g}$ .

To lift the positive pairing to  $F_1^F(M)$ , we lift the map  $J$  to a map

$$\cdot^t : F_1^F(M) \longrightarrow F_1^F(M)$$

given by

$$(f, g)^t = (Lg, -\star df).$$

Note that for  $(f, g) \in F_1^F(M)$ ,

$$(5.13) \quad \text{tr} \int_M \star f \wedge Lg = \text{tr} \int_M \star f^t \wedge Lg^t$$

Then Theorem 4 can be equivalently stated as

**Proposition 5.14.** *Let  $U, V$  be the components of  $M - \tilde{\Sigma}$ . Let  $f \in \Omega^0(U, \mathfrak{g})^\perp$ ,  $\tilde{f} \in \Omega^0(V, \mathfrak{g})^\perp$  and  $g \in \ker(d_2^*) \cap \Omega_c^2(U, \mathfrak{g})$ ,  $\tilde{g} \in \ker(d_2^*) \cap \Omega_c^2(V, \mathfrak{g})$ .*

*Then*

$$\text{tr} \int_M \star \tilde{f}^t \wedge Lf = -\text{vol}(M) \text{tr} c_f c_{\tilde{f}}$$

*and*

$$\text{tr} \int_M \star \tilde{g}^t \wedge Lg = -\text{vol}(M) \text{tr} c_g c_{\tilde{g}}.$$

Alternatively, under the isomorphism

$$H^0(\tilde{\Sigma}, \mathfrak{g}) \oplus H^2(\tilde{\Sigma}, \mathfrak{g}) \simeq \mathfrak{g} \oplus \mathfrak{g},$$

the split linear form is (up to a factor of  $\text{vol}(M)$ ) the bilinear form on  $H^0(\tilde{\Sigma}, \mathfrak{g}) \oplus H^2(\tilde{\Sigma}, \mathfrak{g})$  arising from the intersection pairing of  $H^0(\tilde{\Sigma}, \mathfrak{g})$  with  $H^2(\tilde{\Sigma}, \mathfrak{g})$  and the inner product on  $\mathfrak{g}$ . And we can consider the map  $\pi_\Sigma^F$  as a map taking values in the one particle Fermionic space  $H^0(\tilde{\Sigma}, \mathfrak{g}) \oplus H^2(\tilde{\Sigma}, \mathfrak{g})$ . Note that this identification depends on an orientation of  $\Sigma$ , and hence the reflection  $R$  will introduce an additional sign in the two form  $c_2$ , which we have accounted for in the negative sign in the adjoint  $(\cdot)^c$  mentioned above.

Now consider the situation of Theorem 4 in the case where the splitting of  $M$  is given by  $\Sigma$ . In the absence of Condition B, there is a subtlety about the definition of  $R^* : \Omega^0(M_\pm, \mathfrak{g})^\perp \longrightarrow \Omega^0(M_\mp, \mathfrak{g})^\perp$ , since elements of  $\Omega^0(M_\pm, \mathfrak{g})^\perp$  are not compactly supported on  $M_\pm$ ; instead they are constant outside of  $M_\pm$ . However, if  $f \in \Omega^0(M_\pm, \mathfrak{g})^\perp$ ,  $f - c_f$  is compactly supported on  $M_\pm$ . We define the map  $R^* : \Omega^0(M_\pm, \mathfrak{g})^\perp \longrightarrow \Omega^0(M_\mp, \mathfrak{g})^\perp$  for  $f \in \Omega^0(M_\pm, \mathfrak{g})^\perp$  by

$$R^* f = R^*(f - c_f) + c_f.$$

Then  $R^*f \in \Omega^0(M_{\mp}, \mathfrak{g})^\perp$ , and

$$(5.15) \quad c_{R^*f} = c_f.$$

Also,

$$(5.16) \quad \star d(R^*f) = R^*(\star df).$$

In the absence of Condition A, we must restrict to forms constant outside of  $M_{\pm}^\delta$ , and then the same procedure goes through.

We then have the following result:

**Corollary 5.17.** *Let  $f \in \Omega^0(M_-^\delta, \mathfrak{g})^\perp$  and  $g \in \ker(d_2^*) \cap \Omega_c^2(M_+^\delta, \mathfrak{g})$ . Then*

$$(5.18) \quad \operatorname{tr} \int_M \star R^*f \wedge LR^*g = \operatorname{tr} \int_M \star f \wedge Lg.$$

In particular,

$$c_{R^*g} = c_g.$$

(Where Condition A holds, we may replace  $M_+^\delta$  by  $M_+$  and  $M_-^\delta$  by  $M_-$ .)

*Proof.* If Condition B is satisfied, this follows by  $R$ -invariance of the metric. Otherwise, still, by Theorem 4 and (5.15)-(5.16), it suffices to prove that

$$\operatorname{tr} \int_M \star f \wedge Lg = \operatorname{tr} \int_M \star f \wedge R^*LR^*g.$$

But

$$\operatorname{tr} \int_M \star f \wedge Lg = \operatorname{tr} \int_M dL f \wedge Lg = -\operatorname{tr} \int_M \star L f \wedge g$$

and, by a similar integration by parts,

$$\operatorname{tr} \int_M \star f \wedge R^*LR^*g = -\operatorname{tr} \int_M L f \wedge dR^*LR^*g = -\operatorname{tr} \int_M \star L f \wedge g.$$

□

A more subtle reasoning is needed for the next result, which is an analog of Lemma 4.20.

**Lemma 5.19.** *Let  $f \in \Omega^0(M_+, \mathfrak{g})^\perp$  and  $g \in \ker(d_2^*) \cap \Omega_c^2(M_+, \mathfrak{g})$ . Then*

$$(5.20) \quad \operatorname{tr} \int_M \star g \wedge Lf = \operatorname{tr} \int_M \star R^*g \wedge LR^*f.$$

(In the absence of Condition A, replace  $M_+$  by  $M_+^\delta$  and  $M_-$  by  $M_-^\delta$ .)

*Proof.* If Condition B is satisfied, this follows by  $R$ -invariance of the metric.

In the case of Condition A, we may not have a globally defined involution  $R$ , but we can argue as in the proof of Lemma 4.20.

First, we compute for the right hand side of (5.20)<sup>47</sup>

<sup>47</sup>By a slight abuse of notation, we denote by  $R^*LR^*f \in \Omega^2(M_+, \mathfrak{g})$  the pullback  $j^*((LR^*f)|_{M_-})$  of the form  $(LR^*f)|_{M_-} \in \Omega^2(M_-, \mathfrak{g})$  by the map  $j : M_+ \rightarrow M_-$  of (2.3). The form  $R^*LR^*f$  is not compactly supported, but its wedge product with the compactly supported form  $\star g$  may be integrated on  $M_+$ .



$$\mathrm{tr} \int_M \star R^* g \wedge LR^* f = \mathrm{tr} \int_{M_-} \star R^* g \wedge LR^* f = \mathrm{tr} \int_{M_+} \star g \wedge R^* LR^* f.$$

Then the difference between the right and left sides of (5.20) is given by

$$\mathrm{tr} \int_M \star g \wedge Lf - \mathrm{tr} \int_M \star R^* g \wedge LR^* f = \mathrm{tr} \int_{M_+} \star g \wedge (Lf - R^* LR^* f).$$

But

$$(5.21) \quad \begin{aligned} & \mathrm{tr} \int_{M_+} \star g \wedge (Lf - R^* LR^* f) = \mathrm{tr} \int_{M_+} dLg \wedge (Lf - R^* LR^* f) = \\ & -\mathrm{tr} \int_{M_+} Lg \wedge d(Lf - R^* LR^* f) + \mathrm{tr} \int_{\Sigma} Lg \wedge (Lf - R^* LR^* f). \end{aligned}$$

But, as forms on  $M_+$ ,

$$d(Lf - R^* LR^* f) = \star(f - R^* R^* f) = 0,$$

so that

$$\mathrm{tr} \int_{M_+} Lg \wedge d(Lf - R^* LR^* f) = 0.$$

On the other hand

$$Lg|_{\Sigma} = c_g$$

so

$$\mathrm{tr} \int_{\Sigma} Lg \wedge (Lf - R^* LR^* f) = \mathrm{tr} c_g \int_{\Sigma} (Lf - R^* LR^* f).$$

But

$$\int_{\Sigma} (Lf - R^* LR^* f) = \int_{M_+} (dLf - dR^* LR^* f) = \int_{M_+} (\star f - \star R^* R^* f) = 0.$$

In the absence of Condition A, the same result holds for forms compactly supported on the complement  $M_+^{\delta}$  of a collar neighborhood of  $\Sigma$ . □

**5.2. Reflection positivity and  $\mathcal{F}^F$ .** We consider the dual of the algebra  $F^F(M)$  as generated by fields  $c_0(f)$ ,  $c_2(g)$  where  $f \in \Omega^0(M, \mathfrak{g})^{\perp}$  and  $g \in \ker(d_2^*) \subset \Omega^2(M, \mathfrak{g})$ . On this algebra of fields we have the formal Berezin integral appearing in (1.17), which we can use to produce a functional on  $F^F(M)$ ; viz<sup>48</sup>

$$\Psi(f_1 \wedge \cdots \wedge f_m \wedge g_1 \wedge \cdots \wedge g_n) \sim \frac{1}{Z_{ghost}} \int dc_0 dc_2 e^{-\mathrm{tr} \int_M c_0 dc_2} c_0(f_1) \cdots c_0(f_m) c_2(g_1) \cdots c_2(g_n)$$

where  $Z_{ghost}$  is a formally infinite normalization constant.

<sup>48</sup>In this section, the wedge product  $\wedge$  denotes the product in the alternating algebra  $\mathcal{F}^F$  and its variants, not the wedge product of differential forms.

To define  $\Psi$  in concrete terms that avoid Berezin integrals, let  $f_i \in \Omega^0(M, \mathfrak{g})^\perp$ ,  $i = 1, \dots, m$  and  $g_j \in \ker(d_2^*) \subset \Omega^2(M, \mathfrak{g})$ ,  $j = 1, \dots, n$ , and define

$$\Psi(f_m \wedge \dots \wedge f_1 \wedge g_1 \wedge \dots \wedge g_n) = \frac{1}{m!} \det M,$$

where the  $m \times m$  matrix  $M$  is given by  $M = 0$  if  $m \neq n$  and by

$$M_{ij} = \langle f_i, Lg_j \rangle, i, j = 1, \dots, m$$

if  $m = n$ .

Alternatively, since the operator  $\Delta^{-s/2} L \Delta^{-s/2}$  is antisymmetric and Hilbert Schmidt on the  $L^2$  completion  $\overline{\mathcal{F}_1^F}$  of  $\mathcal{F}_1^F$  for our chosen  $s = 4$ ,<sup>49</sup> and hence gives an element  $L_s \in \wedge^2 \overline{\mathcal{F}_1^F}$ , we may write

(5.22)

$$\Psi(f_m \wedge \dots \wedge f_1 \wedge g_1 \wedge \dots \wedge g_n) = \left\langle \frac{1}{n!} \bigwedge^n L_s, \Delta^{s/2} f_m \wedge \dots \wedge \Delta^{s/2} f_1 \wedge \Delta^{s/2} g_1 \wedge \dots \wedge \Delta^{s/2} g_n \right\rangle_{\wedge^* \overline{\mathcal{F}_1^F}},$$

where the inner product is the inner product induced on the alternating algebra by the  $L_2$  inner product of  $\mathfrak{g}$ -valued forms; again this inner product is zero unless  $m = n$ . This shows, as in (4.18) that

$$|\Psi(P)| < C_d \|P\|_{\mathcal{F}_d^F}$$

for  $P \in \mathcal{F}_d^F$ . Thus  $\Psi$  extends to a bounded linear functional on  $\mathcal{F}_d^F$  for every  $d$ , and to a linear functional on  $\mathcal{F}_P^F$ .<sup>50</sup>

As in the Bosonic case, it will be convenient to work with  $L$ -Wick ordered polynomials.<sup>51</sup>

<sup>49</sup>For this property  $s > \frac{1}{4}$  would suffice.

<sup>50</sup>In fact,  $\Psi$  is a bounded linear functional, but we do not make any use of this.

<sup>51</sup>Some comments about Wick ordering: As in the Bosonic case this is most conveniently described first for Hilbert Schmidt operators.

Suppose  $L$  is an antisymmetric Hilbert Schmidt operator on a Hilbert space  $H$ . Then there exists  $v \in \wedge^2 H \subset H \otimes H$  with

$$\langle v, x \otimes y \rangle_{\wedge^2 H} = \langle x, Ly \rangle$$

for all  $x, y \in H$ .

Suppose  $P \in \wedge^* H$  is a polynomial and define

$$\Psi(P) = \langle e^{i_v} P, 1 \rangle_{\wedge^* H}$$

where  $i_v$  is interior product in  $\wedge^* H$ ; that is,

$$i_v(x \wedge y) = \langle v, x \wedge y \rangle_{\wedge^2 H}.$$

Write, for  $P \in \wedge^* H$ ,

$$: P := e^{-i_v} P.$$

Then if  $P, Q \in \wedge^* H$ ,

$$(5.23) \quad \Psi(: P :: Q :) = \langle P, \left( \sum_{k=0}^{\infty} \frac{1}{k!} \wedge^k L \right) Q \rangle_{\wedge^* H}.$$

A compact way of defining these is by taking another copy  $X$  of  $\mathcal{F}_1^F$  and defining

$$: e^{\langle x, y \rangle} := e^{-\frac{1}{2} \langle x, Lx \rangle} e^{\langle x, y \rangle}$$

for any  $x \in X$ ,  $y \in \mathcal{F}_1^F$  (see [7], page 22); as in the Bosonic case, the Wick ordered polynomials can be extracted from this formula.

As in the previous section, suppose  $\tilde{\Sigma}$  is a smooth compact connected submanifold of  $M$  splitting  $M$  into two components  $U, V$ . We now extend the definition of the map  $\pi_{\tilde{\Sigma}}^F$  to a map  $\pi_{\tilde{\Sigma}}^F : \mathcal{F}_P^F(U) \rightarrow \Lambda^*(\mathfrak{g} \oplus \mathfrak{g})$  by

$$\pi_{\tilde{\Sigma}}^F(: f_1 \wedge \cdots \wedge f_n \wedge g_1 \wedge \cdots \wedge g_m :) = \pi_{\tilde{\Sigma}}^F(f_1) \wedge \cdots \wedge \pi_{\tilde{\Sigma}}^F(f_n) \wedge \pi_{\tilde{\Sigma}}^F(g_1) \wedge \cdots \wedge \pi_{\tilde{\Sigma}}^F(g_m)$$

and extending by linearity, and similarly for forms compactly supported in  $V$ . This map is again bounded on each  $\mathcal{F}_d^F(U)$  and each  $\mathcal{F}_d^F(V)$ .

Given a monomial  $m$ , we extend the definition of the map  $(\cdot)^c$  in (5.8) by writing

$$(5.24) \quad m^c = (-1)^{s_1 + s_2} m,$$

where  $s_1 = \frac{|m|(|m|-1)}{2}$ ,  $|m|$  is the degree of  $m$ , and  $s_2$  is the number of factors of  $c_2$  in  $m$ . Intuitively, this monomial is obtained by reversing the order of the terms in  $m$  and attaching a  $-$  sign to each factor of  $c_2$ . Explicitly, if

$$m = f_1 \wedge \cdots \wedge f_k \wedge g_1 \wedge \cdots \wedge g_l,$$

where  $f_i, g_j$  are as above, then

$$m^c = -g_l \wedge \cdots \wedge -g_1 \wedge f_k \wedge \cdots \wedge f_1.$$

Note that for any monomial  $m$ ,

$$(5.25) \quad : m^c := : m :^c .$$

Likewise we extend the definition of the map  $\cdot^t$  to monomials by defining  $m^t$ , where  $m = f_1 \wedge \cdots \wedge f_n \wedge g_1 \wedge \cdots \wedge g_m$ , by

$$m^t = Lg_m \wedge \cdots \wedge Lg_1 \wedge (-\star df_n) \wedge \cdots \wedge (-\star df_1).$$

Extending by linearity, the definitions of  $\cdot^c$  and  $\cdot^t$  extend to all polynomials in  $\mathcal{F}_P^F(M)$ .

The following proposition is then an immediate consequence of the definition of  $\Psi$  and Theorem 4.

**Proposition 5.26.** *Let  $p \in \mathcal{F}_P^F(U)$  and  $q \in \mathcal{F}_P^F(V)$ . Then*

$$\Psi(: p^c : \wedge : q :) = \langle \pi_{\tilde{\Sigma}}^F(: p :), \pi_{\tilde{\Sigma}}^F(: q :) \rangle_{\Lambda^*(\mathfrak{g} \oplus \mathfrak{g})}$$

and therefore by (5.25)

$$\Psi(p^c \wedge q) = \langle \pi_{\tilde{\Sigma}}^F(p), \pi_{\tilde{\Sigma}}^F(q) \rangle_{\Lambda^*(\mathfrak{g} \oplus \mathfrak{g})}$$

---

As in the Bosonic case, if  $L$  is not Hilbert Schmidt, similar formulas hold as long as both sides of equation (5.23) are well defined; these can be proved by cutting off  $L$  to obtain a sequence of finite rank operators, and then taking the limit on both sides. In our case a transfer of regularity argument can be used, involving the operators  $L_s$  of equation (5.22).

where the inner product on  $\Lambda^*(\mathfrak{g} \oplus \mathfrak{g})$  is that given by the split linear form on  $\mathfrak{g} \oplus \mathfrak{g}$ .

In particular, if  $\tilde{\Sigma} = \Sigma, U = M_+, V = M_-$ ,

$$\Psi((R^*p)^c \wedge p) = \langle \pi_{\Sigma}^F(p), \pi_{\Sigma}^F(p) \rangle_{\Lambda^*(\mathfrak{g} \oplus \mathfrak{g})}.$$

Likewise we have

**Proposition 5.27.** *Let  $p \in \mathcal{F}_P^F(U)$  and  $q \in \mathcal{F}_P^F(V)$ . Then*

$$\Psi(\cdot : p^t : \wedge : q :) = \langle \pi_{\tilde{\Sigma}}^F(\cdot : p :), \pi_{\tilde{\Sigma}}^F(\cdot : q :) \rangle_+$$

where the metric  $\langle \cdot, \cdot \rangle_+$  on  $\Lambda^*(\mathfrak{g} \oplus \mathfrak{g})$  is the one given by the positive metric  $\langle \cdot, \cdot \rangle_+$  on  $\mathfrak{g} \oplus \mathfrak{g}$ .

Proposition 5.27, along with the reflection invariance of the Wick ordering (Corollary 5.17 and Lemma 5.19), hint at the possibility of a reflection positivity theorem for the fermions. However, the Wick ordering  $\cdot : \cdot :$  is not invariant under the transformation given by  $(\cdot)^t$ , and in fact the Fermions contain negative norm ("ghost") states.<sup>52</sup> We will thus have to handle the Boson-Fermion interaction term by different methods than those used for the Boson interaction term, which is a self adjoint operator.

In any event, we write  $H^F(\Sigma) = \Lambda^*(\mathfrak{g} \oplus \mathfrak{g})$ , with the positive definite metric  $\langle \cdot, \cdot \rangle_+$ .<sup>53</sup>

Likewise if  $\tilde{\Sigma}$  is any two dimensional submanifold of  $M$  isotopic to  $\Sigma$ , and  $U, V$  are the components of  $M - \tilde{\Sigma}$ , we write  $H^F(\tilde{\Sigma})$  for the Hilbert space associated to  $\tilde{\Sigma}$  (which is of course isomorphic to  $H^F(\Sigma)$ ) to make the relation to the geometry explicit.

Note that since  $H^F(\Sigma)$  is an alternating algebra on a finite dimensional vector space, it is equipped with a bounded, nonnegative, self-adjoint operator  $N_F : H^F(\Sigma) \rightarrow H^F(\Sigma)$  given by polynomial degree. Also, since  $H^F(\Sigma)$  is finite-dimensional, the two inner products on  $H^F(\Sigma) = \Lambda^*(\mathfrak{g} \oplus \mathfrak{g})$  are related by a (necessarily bounded) invertible linear operator  $\mathcal{J}_F = \Lambda^* J : H^F(\Sigma) \rightarrow H^F(\Sigma)$ .

**Remark 5.28.** As in the Bosonic case, scaling the quadratic term in the free Lagrangian by a factor of  $\alpha$  corresponds to replacing the functional  $\Psi$  by a functional  $\Psi_\alpha$  obtained by multiplying the right hand side of (5.22) by a factor of  $\alpha^{-n}$ . Equivalently, the inner products on  $H^F(\Sigma)$  can be modified by a factor of  $e^{-\log(\alpha)N_F}$ .

### 5.3. The Hilbert space $H(\Sigma)$ . We write

$$H(\Sigma) = H^B(\Sigma) \otimes H^F(\Sigma).$$

We also write

$$\pi_\Sigma = \pi_\Sigma^B \otimes \pi_\Sigma^F,$$

<sup>52</sup>For example, consider  $p = (Lg \otimes \xi) \wedge (g \otimes \xi)$  where  $g \in \ker(d_2^*) \subset \Omega_c^2(M_+^\delta, \mathbb{R})$  satisfies  $c_g = 0$  and  $\xi$  is a nonzero element of  $\mathfrak{g}$ . Then  $\Psi(Rp^t p) = -\|\xi\|_{\mathfrak{g}}^2 \|Lg\|_2^2$ .

<sup>53</sup>As for the gauge fields, where the symmetric algebra is a subalgebra of a Weyl algebra, this alternating algebra is embedded in a Clifford algebra on two copies of  $\mathfrak{g} \oplus \mathfrak{g}$  corresponding to ghost fields compactly supported on both sides of  $\Sigma$ .

reiterating that the map  $\pi_\Sigma$  is only defined on the dense subspace  $\mathcal{F}_{P,+} \subset \mathcal{F}_+$ . Similarly  $H(\tilde{\Sigma}), \pi_{\tilde{\Sigma}}$  for any compact connected smooth submanifold  $\tilde{\Sigma}$  isotopic to  $\Sigma$ .

The Hilbert space  $H(\Sigma)$  is equipped with a densely defined, nonnegative, self-adjoint operator  $N$  given by  $N = N_B + N_F$ , as well as with a bounded operator  $\mathcal{J} : H(\Sigma) \rightarrow H(\Sigma)$  given by  $\mathcal{J} = \text{id}_{H^B(\Sigma)} \otimes \mathcal{J}_F$ . And as before, a scaling of both quadratic terms in the free Lagrangian by a factor of  $\alpha$  is equivalent to a modification of the metric on the Hilbert space  $H(\Sigma)$  by the positive bounded self-adjoint operator  $e^{-\log(\alpha)N}$ .

We extend the action of the conjugation  $(\cdot)^c$  to  $\mathcal{F}_P(M)$  by defining its action on Bosonic fields to be trivial. Likewise define the map  $R^* : \mathcal{F}_P(M_+) \rightarrow \mathcal{F}_P(M_-)$  as the tensor product of the maps (both denoted  $R^*$ )  $R^* : \mathcal{F}_P^B(M_+) \rightarrow \mathcal{F}_P^B(M_-)$  and  $R^* : \mathcal{F}_P^F(M_+) \rightarrow \mathcal{F}_P^F(M_-)$ , with the obvious replacement of  $M_\pm$  with  $M_\pm^\delta$  if Condition A is not satisfied. Then the following Theorem summarizes the relation between the functionals  $\Phi$  and  $\Psi$  (which correspond to expectations in the formal functional integral arising from the quadratic part of the Chern Simons Lagrangian) and the Hilbert space  $H(\Sigma)$ .

**Theorem 5.** *Let  $p, q \in \mathcal{F}_{P,+}$ . Then*

$$(\Phi \otimes \Psi)((R^* p)^c q) = \langle \pi_\Sigma(p), \mathcal{J} \pi_\Sigma(q) \rangle.$$

## 6. THE INTERACTING THEORY

So far we have constructed a Hilbert space  $H(\Sigma) = H^B(\Sigma) \otimes H^F(\Sigma)$  with a positive definite inner product in which correlations of polynomial functions are as would be expected from the quadratic part of the formal Chern-Simons functional integral with action

$$S_F(A, c_0, c_2) = \frac{1}{2} \text{tr} \int_M AdA - 2c_0 dc_2.$$

We now wish to include the interaction term<sup>54</sup>

$$S_I(A, c_0, c_2) = \text{tr} \int_M A^3 - 6Ac_0c_2.$$

These interaction terms are local, and are morally of the form

$$S_I = \Xi_B^+ + R^* \Xi_B^+ + \Xi_{BF}^+ + R^*(\Xi_{BF}^+)^c,$$

where

$$(6.1) \quad \Xi_B^+(A) = \text{tr} \int_{M_+} A^3$$

and

$$(6.2) \quad \Xi_{BF}^+(A, c_0, c_2) = -6 \text{tr} \int_{M_+} Ac_0c_2$$

---

<sup>54</sup>We focus in this introduction on the case where Condition A holds.

are local interaction terms that have the appearance of elements of  $\mathcal{F}_{3,+}$ ,<sup>55</sup> and hence should yield polynomials of degree 3 in  $H(\Sigma)$ .

However, a closer study of the terms  $\Xi_B^+$  and  $\Xi_{BF}^+$  shows that in order to interpret them as elements of  $H(\Sigma)$ , some choices are necessary. This is due to the fact that elements of  $\mathcal{F}_{P,+}$  consist of local polynomials built from linear factors arising from differential forms compactly supported in<sup>56</sup>  $M_+$  and satisfying the gauge fixing conditions  $d^*(\cdot) = 0$ , while the integrals over  $M_+$  appearing in (6.1) and (6.2) are not of this form. To see this, choose an orthonormal basis  $\{e_\alpha\}$  for  $\mathfrak{g}$  and three smooth one forms  $f, g, h \in \Omega_c^1(M_+)$  satisfying the gauge condition. We would expect, morally,

$$(6.3) \quad \Phi(A(R^*f \otimes e_\alpha)A(R^*g \otimes e_\beta)A(R^*h \otimes e_\gamma)\Xi_B^+) = -f_{\alpha\beta\gamma} \int_{M_+} LR^*f \wedge LR^*g \wedge LR^*h.$$

where  $f_{\alpha\beta\gamma} = -\text{tr}(e_\alpha[e_\beta, e_\gamma])$ . Now the integral

$$\int_{M_+} LR^*f \wedge LR^*g \wedge LR^*h$$

does depend only on the boundary values  $Lf|_\Sigma, Lg|_\Sigma, Lh|_\Sigma \in Z^1(\Sigma)$ ; but not only on the cohomology classes  $[Lf|_\Sigma], [Lg|_\Sigma], [Lh|_\Sigma] \in H^1(\Sigma)$ . Similar problems arise for  $\Xi_{BF}^+$ . So it is not possible to interpret the interaction terms as elements of  $H(\Sigma)$  without making choices, specifically of an orthonormal basis  $\{b_i\}$  for  $\Lambda \subset H^1(\Sigma)$  and of one forms  $\psi_i$  with  $[L\psi_i|_\Sigma] = b_i$ , along with a similar choice for the Fermions. Once such a choice is made, equation (6.3) defines an element  $\xi_B^+ \in H_B(\Sigma)$  of degree 3. This gives a densely defined multiplication operator  $\xi_B^+ \cdot$ , and a symmetric operator  $\xi_B^+ \cdot + (\xi_B^+ \cdot)^*$ , which has a self adjoint extension  $O_B$  and gives rise to a one parameter subgroup  $e^{i\lambda O_B}$  on  $H_B(\Sigma) \otimes \mathbb{C}$ .

Similarly, the Fermion interaction term  $\Xi_{BF}^+$  gives rise, after choices, to an element  $\xi_{BF}^+ \in H(\Sigma)$  which is linear in the gauge field and quadratic in the Fermions. The interaction term does not correspond to the sum of this element and its adjoint in the positive metric on  $H(\Sigma)$ , but rather to the sum of this element and its adjoint in the indefinite inner product on  $H(\Sigma)$  arising from the split linear metric on  $\mathfrak{g} \oplus \mathfrak{g}$ . It therefore does not give a self adjoint operator on a Hilbert space, and does not give rise to a one parameter subgroup of unitary operators. However, we can consider the operator

$$O_{BF} = \xi_{BF}^+ \cdot + (\xi_{BF}^+ \cdot)^\dagger$$

where  $(\cdot)^\dagger = \mathcal{J}^{-1}(\cdot)^* \mathcal{J}$  denotes the adjoint in the indefinite inner product on  $H(\Sigma)$  arising from the split linear metric on  $\mathfrak{g} \oplus \mathfrak{g}$ . The operator is defined on the dense subset of  $H(\Sigma)$  consisting of elements of finite degree. Since  $\xi_{BF}$  is linear in the gauge field, and since the ghost Hilbert space is finite dimensional, so that the ghost field operators are bounded, we have a bound

$$O_{BF}^* O_{BF} \leq K_1 N_B + K_2$$

for some constants  $K_1, K_2$ . We can therefore exponentiate  $O_{BF}$  using the explicit power series to obtain a family

<sup>55</sup>Note that  $\Xi_B^+$  and  $\Xi_{BF}^+$  are Wick ordered polynomials.

<sup>56</sup>In the case of the zero form, with *derivative* compactly supported in  $M_+$

$$e^{-\epsilon N} e^{i\lambda O_{BF}}$$

for every  $\lambda \in \mathbb{R}$  and  $\epsilon > 0$  which are bounded for every  $\epsilon$ , and extend to all of  $H(\Sigma)$ .<sup>57</sup> We can then combine the operators  $e^{i\lambda O_B}$  and  $e^{-\epsilon N} e^{i\lambda O_{BF}}$ , and take a weak limit inspired by the Trotter product formula, to obtain what *morally* should be the partition function

$$\langle \Omega, \mathcal{J} e^{-\epsilon N + i\lambda(O_B + O_{BF})} \Omega \rangle,$$

where  $\Omega = \pi_\Sigma(1)$ . Since we have seen (see Remark 4.28 and Remark 5.28) that a factor of  $e^{-\epsilon N}$  amounts to a renormalization of the free Lagrangian, this is a reasonable definition of the partition function.<sup>58</sup>

In the remainder of this section, we focus on the case where Condition A is satisfied. If this is not the case, an additional weak limit has to be taken as  $\delta \rightarrow 0$ .

**6.1. The Boson interaction term.** We now define the Bosonic interaction term. Choose an orthonormal basis  $\{e_\alpha\}$  for  $\mathfrak{g}$  and write  $f_{\alpha\beta\gamma} = -\text{tr}(e_\alpha, [e_\beta, e_\gamma])$ . Choose a basis  $b_1, \dots, b_g$  for  $\Lambda \subset H^1(\Sigma)$ , and, using Proposition 4.10, choose one forms  $\phi_1, \dots, \phi_g \in F_1^B(M_+)$  such that

$$[L\phi_i|_\Sigma] = b_i.$$

We define the Bosonic interaction element

$$\xi_B^+ \in \text{Sym}^3(\Lambda) \subset H^B(\Sigma)$$

as the unique element of the finite dimensional vector space  $\text{Sym}^3(\Lambda)$  satisfying

$$\langle \xi_B^+, (b_i \otimes e_\alpha) \otimes (b_j \otimes e_\beta) \otimes (b_k \otimes e_\gamma) \rangle = -f_{\alpha\beta\gamma} \int_{M_+} LR^* \phi_i \wedge LR^* \phi_j \wedge LR^* \phi_k$$

for all  $\alpha, \beta, \gamma, i, j, k$ .

Then  $\xi_B^+$  gives a densely defined multiplication operator  $\xi_B^+ \cdot$  on  $H^B(\Sigma)$ , with domain  $\mathcal{D} = \text{Sym}^* \Lambda \subset H^B(\Sigma)$  given by the elements of finite degree. Since the domain of the adjoint  $(\xi_B^+ \cdot)^*$  contains  $\mathcal{D}$ , the sum

$$\xi_B^+ \cdot + (\xi_B^+ \cdot)^*$$

is a densely defined symmetric operator on the real Hilbert space  $H^B(\Sigma)$ , and hence has a (possibly non-unique) self-adjoint extension  $O_B$  densely defined on  $H^B(\Sigma) \otimes \mathbb{C}$ .<sup>59</sup> We may then exponentiate  $O_B$  to obtain a one parameter subgroup

$$e^{i\lambda O_B} : H^B(\Sigma) \otimes \mathbb{C} \rightarrow H^B(\Sigma) \otimes \mathbb{C}$$

of unitary operators.

By a slight abuse of notation we use the notation  $e^{i\lambda O_B}$  also for the one parameter subgroup of unitary operators acting as  $e^{i\lambda O_B} \otimes \text{id}$  on  $H^B(\Sigma) \otimes H^F(\Sigma) \otimes \mathbb{C} = H(\Sigma) \otimes \mathbb{C}$ .

<sup>57</sup>A similar method for handling Bose-Fermi interactions is in [20].

<sup>58</sup>Our actual construction is slightly different. See Section 6.3.

<sup>59</sup>This is because a densely defined symmetric operator on a real Hilbert space has equal deficiency indices as on operator on the complexified Hilbert space. See e.g. [18], p. 402, or [21] p. 143.

**Remark 6.4.** The fact that the interaction term  $\Xi_B^+$  is not naturally an element of  $H^B(\Sigma)$  raises the issue of whether a larger Hilbert space, big enough to contain  $\Xi_B^+$ , can be constructed from polynomials in the gauge fields built from  $\mathfrak{g}$ -valued one forms compactly supported on  $M_+$  but not satisfying the gauge conditions. Given  $f, g \in \Omega_c^1(M_+, \mathfrak{g})$ , we may compute, as in Lemma 4.5,

$$(6.5) \quad \text{tr} \int_M \star \pi(f) \wedge R^* L\pi(g)$$

where

$$\pi = d^* d\Delta^{-1}$$

is the projection onto one forms satisfying the gauge condition. A computation along the lines of that in Lemma 4.5 again reduces the integral in (6.5) to an integral over  $\Sigma$ , involving now three terms:<sup>60</sup>

$$(6.6) \quad \text{tr} \int_M \star \pi(f) \wedge R^* L\pi(g) = \text{tr} \int_\Sigma \star d\Delta^{-1} f \wedge \star d\Delta^{-1} R^* g + d^* \Delta^{-1} f \wedge d\Delta^{-1} R^* g + d\Delta^{-1} f \wedge d^* \Delta^{-1} R^* g$$

The first term pairs a one form (now not necessarily closed) arising from  $f$  with a similar one form arising from  $g$ , as in Lemma 4.5, and the two additional terms pair zero- and two-forms arising from  $f$  with similar forms arising from  $g$ . It would be interesting to see if any form of reflection positivity holds for the pairing given by (6.6), and to try to identify the resulting Hilbert space. The Hilbert space  $H^B(\Sigma)$  would then be a subspace of the larger Hilbert space hinted at by (6.6), and the element  $\xi_B^+ \in H^B(\Sigma)$  would be a projection of the element  $\Xi_B^+$  to this subspace. It is also conceivable that the effects of the terms corresponding to the pairing of zero-forms with two-forms cancel in some way with the contributions from the ghost fields, so that we effectively end up again with something like  $H^B(\Sigma)$ .

**6.2. The Fermion interaction term.** In order to define the Fermion interaction term, we must supplement the choices made in the previous section by choosing also a function  $f \in \Omega^0(M_+, \mathfrak{g})^\perp$  with  $c_f = 1$ . We define the Fermion interaction element as the unique element  $\xi_{BF}^+ \in \Lambda \otimes H^F(\Sigma) \subset H(\Sigma)$  satisfying

$$\langle \xi_{BF}^+, \mathcal{J}((b_i \otimes e_\alpha) \otimes (e_\beta \oplus 0) \wedge (0 \oplus e_\gamma)) \rangle = 6f_{\alpha\beta\gamma} \int_{M_+} LR^* \phi_i \wedge LR^* f$$

for all  $\alpha, \beta, \gamma, i$ , where we have identified the one particle subspace in  $H^F(\Sigma)$  with  $\mathfrak{g} \oplus \mathfrak{g}$ .

As before, the element  $\xi_{BF}^+$  gives a multiplication operator  $\xi_{BF}^+ \cdot$  densely defined on  $\mathcal{D} \otimes H^F(\Sigma)$ . A direct computation shows that the domain of the adjoint operator  $(\xi_{BF}^+ \cdot)^*$  contains  $\mathcal{D} \otimes H^F(\Sigma)$ .

Then the Fermion interaction term is given by the operator

$$O_{BF} = \xi_{BF}^+ \cdot + (\xi_{BF}^+ \cdot)^\dagger,$$

where the notation  $\cdot^\dagger$  denotes the adjoint in the bilinear form on  $H(\Sigma)$  given by the usual metric on  $H^B(\Sigma)$  and the split linear form on  $H^F(\Sigma)$ ; in terms of the Hilbert space metric, we have

$$(\xi_{BF}^+ \cdot)^\dagger = \mathcal{J}^{-1}(\xi_{BF}^+ \cdot)^* \mathcal{J}$$

where  $\mathcal{J}$  is the bounded linear operator on  $H(\Sigma)$  defined in Section 5.2 arising from the linear transformation on  $H^F(\Sigma)$  relating the split linear form to the metric. Since the

<sup>60</sup>In this equation the  $\star$  operator and the expression  $d^*$  refer to the Hodge star operator on  $M$ , not on  $\Sigma$ .



operator  $\mathcal{J}$  acts as the identity on  $H^B(\Sigma)$ , it preserves the domain  $\mathcal{D} \otimes H^F(\Sigma)$ , so that the operators  $(\xi_{BF}^+)^{\dagger}$  and  $O_{BF}$  are also defined on the dense domain  $\mathcal{D} \otimes H^F(\Sigma)$ .

Now the operator  $O_{BF}$  is not symmetric, so it cannot be extended to a self-adjoint operator and exponentiated to a one parameter subgroup of bounded operators. However,  $\xi_{BF}^+$  is linear in the gauge fields, so we have the inequality

$$(6.7) \quad O_{BF}^* O_{BF} \leq K_1 N_B + K_2$$

for some constants  $K_1, K_2 \geq 0$ .

We may therefore define, on  $\mathcal{D} \otimes H^F(\Sigma) \otimes \mathbb{C}$ , and for any  $\lambda \in \mathbb{C}$ , the operator

$$e^{-\epsilon N_B} e^{\lambda O_{BF}} := \sum_{k=0}^{\infty} \frac{1}{k!} \lambda^k e^{-\epsilon N_B} (O_{BF})^k$$

which, by (6.7), is bounded for every  $\epsilon > 0$ , and hence gives an operator on  $H(\Sigma) \otimes \mathbb{C}$ ; and we have the bound

$$(6.8) \quad \|e^{-\epsilon N_B/n} e^{i\lambda O_{BF}/n}\| \leq K \frac{1}{n}$$

for some constant  $K \geq 0$  (which depends on  $\epsilon$  and  $\lambda$  but not on  $n$ ) for any  $n > 0$ .

**6.3. The Partition function.** In this section we combine the results of Section 6.1 and Section 6.2 to define the partition function of our theory.

Let  $\lambda, \epsilon \in \mathbb{R}$  and, for any  $n > 0$ , let

$$(6.9) \quad Z_n(\epsilon, \lambda) = \langle \Omega, \mathcal{J}(e^{-\epsilon N/n} e^{i\lambda O_{BF}/n} e^{i\lambda O_B/n})^n \Omega \rangle$$

where  $\Omega$  is the identity element  $\Omega = 1 \in \text{Sym}^* \Lambda \otimes \bigwedge^*(\mathfrak{g} \oplus \mathfrak{g})$ .

By the estimate (6.8), the sequence  $Z_n$  is bounded, so that

**Theorem 6.** *The weak limit*

$$Z(\epsilon, \lambda) = \text{wk} \lim_{n \rightarrow \infty} Z_n(\epsilon, \lambda).$$

*exists.*

The function  $Z(\epsilon, \lambda)$  is the partition function of the quantum field theory.<sup>61</sup> Note that the theory is finite and does not require renormalization, as may be expected from perturbation theory [3].

In view of the Trotter product formula, and of the fact that inserting a factor of  $e^{-\epsilon N}$  in the inner product amounts to a scaling of the free Lagrangian (see Remark 4.28 and Remark 5.28), this seems like a reasonable definition.<sup>62</sup> But several questions come to mind.

<sup>61</sup>This method of taking limits through subsequences appears in constructive quantum field theory in Glimm-Jaffe's construction of the infinite volume limit in two dimensions in [10]. In that case, also, establishing uniqueness of the limit required further developments [9].

<sup>62</sup>It is also possible to contemplate a more elaborate construction, motivated by a time ordered exponential, obtained by cutting the manifold  $M_+$  and its mirror image into  $N$  slices, and viewing the corresponding interaction terms as giving a map between the Hilbert spaces corresponding to the boundaries of the slices. We would then compose the exponentials of these terms and take another weak limit as  $N \rightarrow \infty$ .

**Question 6.10.** *Is the weak limit unique?*

**Question 6.11.** *Assuming Conjecture 6.10, is this limit independent of choices?*<sup>63</sup>

**Question 6.12.** *Assuming Conjecture 6.10 and Conjecture 6.11, is this limit a topological invariant of  $M$ ? How is it related to the invariants of Witten, Reshetikhin, and Turaev?*

It may be that a positive answer to these questions can be expected if the coupling constant is quantized.

More generally, answering these questions may require ideas beyond reflection positivity. But reflection positivity does provide a context in which those questions can be formulated precisely.

**Remark 6.13.** Recalling the discussion in Section 4.3, we see that morally, the partition function should be a putative expression of the form

$$(6.14) \quad \langle \pi_\Sigma(e^{-i\lambda(\Xi_B^+ + \Xi_{BF}^+)}), \mathcal{J}\pi_\Sigma(e^{i\lambda(\Xi_B^+ + \Xi_{BF}^+)}) \rangle_{H(\Sigma) \otimes \mathbb{C}}.$$

Such an expression would be expected if we were to construct a topological field theory, which assigns an element of the Hilbert space associated to a two-manifold to any bounding three-manifold. And generalizations of (6.14) would then be available for any compact three manifold, not only for manifolds with reflection. However, as we noted in Section 4.3, and even leaving aside the fact that  $\Xi_B^+$  and  $\Xi_{BF}^+$  are not quite elements of  $\mathcal{F}_+$ , we do not know how to construct a map  $\mathcal{F}_+^B \rightarrow H^B(\Sigma)$ , and hence we do not know how to make sense of the expression  $\pi_\Sigma(e^{i\lambda(\Xi_B^+ + \Xi_{BF}^+)})$  unless we replace the operator  $L$  appearing in all our definitions of  $\Psi, \Phi$ , etc. by the operator  $KI + L$  where  $K > \|L\|$ . The expression (6.14) therefore makes sense only for  $K$  large, not for the geometrically relevant case  $K = 0$ .

The expression (6.9), while well defined for the geometrically relevant covariance  $L$ , is only available for manifolds with reflection.

This raises the issue of how an exponential of the form (6.9) is morally related to a putative partition function of the form (6.14). One way to look at  $\xi_B^+ + \xi_{BF}^+$  as an element of the Hilbert space of a three manifold with one boundary  $\Sigma$  and one "boundary" consisting of a point, to which topological quantum field theory would naturally assign the vacuum. The composition in topological quantum field theory of two such elements reduces essentially to multiplication, and so the exponentials in the morally correct (6.14) may be expected to be expressed in the mathematically well-defined (6.9).

One way to address these issues is via Questions 6.10-6.12.

## 7. CONCLUDING REMARKS AND SPECULATIONS

We conclude with some remarks and speculations about Chern-Simons functional integrals.

We begin with a general comment about the potential role of Chern-Simons functional integrals. First, and quite apart from topological quantum field theory, the question of providing a mathematical interpretation of functional integrals is of independent interest, and a longstanding problem in Mathematical Physics. Second, the construction we give indicates a possible close relationship between the topological quantum field theories associated with abelian Lie groups, which are related to the quadratic term in the Chern

<sup>63</sup>A speculation: The classical object corresponding to the sum of  $\xi_B^+$  and its adjoint vanishes morally due to the vanishing of the cup product  $\bigwedge^3 H^1(M) \rightarrow H^3(M)$ . Another aspect of this is the cancellation of the boundary terms in (6.3) between  $\Xi_B^+$  and its reflection. It might be that the quantization  $O_B$ , which is nonvanishing, may give rise to a partition function independent of choices as a type of anomalous quantum correction to the vanishing of the classical invariant.

Simons Lagrangian, and the topological quantum field theories associated to nonabelian groups. It would appear from the functional integral that nonabelian theories arise in some version of the abelian theory. This phenomenon may be interesting to study from a topological point of view. Finally, if functional integrals give an alternative construction of topological quantum field theories, and these were related to those constructed by topological methods, such a relation between topology and analysis could be seen as an infinite dimensional nonlinear analog of the de Rham theorem, or perhaps of the Atiyah-Singer index theorem.

**7.1. A finite dimensional expression that may be related to the Chern Simons path integral.** The Hilbert Space  $H^B(\Sigma)$  is the completion of a space of polynomials on a finite dimensional vector space  $\Lambda \subset H^1(\Sigma, \mathfrak{g})$ . The partition function and expectations of polynomials, which are morally given by integration on an infinite dimensional space of connections, should then also be expressible as path integrals over this finite dimensional space. The Fermion Hilbert space  $H^F(\Sigma)$  is finite dimensional, and the action is quadratic in the ghost variables, so the Fermi integral can be performed explicitly to add a determinant to this integral. One may speculate that since this is a topological quantum field theory, the path integral may be related to a finite dimensional integral. Where  $M$  is Heegaard split along  $\Sigma$ , we may consider an integral over the finite dimensional vector space  $\mathbb{R}^{2g} \otimes \mathfrak{g}$  roughly of the form

$$(7.1) \quad \int_{\Lambda} \left( \prod_{a,\alpha} dt_a^\alpha \right) e^{-t, (J^{-1} \otimes I)t} - i\lambda \sum_{\alpha,\beta,\gamma} \sum_{a,b,c} t_a^\alpha t_b^\beta t_c^\gamma f_{\alpha\beta\gamma} N_{abc} \det(K \otimes I + 6i\lambda \sum_{a,\alpha} t_a^\alpha B_a \otimes F_\alpha)$$

where the elements of the matrices  $J, K, B_a$ , and  $F_\alpha$  and the tensor  $N_{abc}$  are constants depending on the manifold  $M$  and the Heegaard splitting. For example, the constants  $N_{a,b,c}$  may be computed for elements  $a, b, c$  of the cohomology of  $\Sigma$  by extending closed forms representing them to forms  $A, B, C$  on  $M$  which are closed on  $M_-$  and computing the triple intersection  $\text{tr} \int_M \star dA \wedge \star dB \wedge \star dC$ , and then adding the result to a similar computation performed by exchanging  $M_+$  with  $M_-$  and replacing  $a, b, c$  with  $S^*a, S^*b, S^*c$ . And the bilinear form  $J$  arises from the intersection form on  $\Sigma$  and the diffeomorphism  $S$ . Then the positivity of the matrix  $J$  should imply that this integral is a matrix Airy function of the form studied in [16]. (The determinant appearing in the integrand is a polynomial in the  $t_a^\alpha$ , therefore integrable.)

Alternatively there may be as we mentioned a formulation in terms of an integral over a space of paths on  $\Lambda$ .<sup>64</sup>

The role of the group  $G$  is quite minimal in this finite dimensional path integral; in fact it appears that all that matters is the abelian path integral, which gives the matrix  $J$ , and some kind of lie algebra to make the alternating tensor  $N$  into a reasonable integrand.

In a more speculative vein, the perturbation of a quadratic integral on a symplectic manifold by a cubic term may hint at a deformation of the quantization of a  $\dim(G)$ -fold product of the Jacobian of a Riemann surface by a term determined by a bounding three manifold. This deformation involves the construction of a cubic term on  $H^1(\Sigma, \mathfrak{g})$  using

<sup>64</sup>The fact that the cup product map  $\bigwedge^3 H^1(M) \rightarrow H^3(M)$  vanishes may mean that an integral on the space of paths, with a path of connections on  $\Sigma$  ending at the trivial connection giving rise to a connection on  $M_+$ , is a more plausible construction. Note that the deformation of algebras of functions on a Poisson manifold referred to below also amounts to a functional integral over paths [5].

the Hodge star operator on a bounding three manifold and the structure constants of the Lie algebra  $\mathfrak{g}$ . This is reminiscent of the deformation quantization of [17].<sup>65</sup> In other contexts, the role of the Hodge star operator and of cubic deformations of algebras in finding manifold invariants beyond cohomology has been emphasized by D. Sullivan.

**7.2. Relation to topological quantum field theory.** The first aspect of a construction of topological quantum field theory would be to show that the limit partition function is a topological invariant. Beyond that, however, it would be necessary to construct the theory beyond manifolds with reflection. The most direct way to do this would be to assign an element of the Hilbert space  $H(\Sigma)$  to any manifold with boundary  $\Sigma$ . This in turn would arise from a map  $\mathcal{F}_+ \rightarrow H(\Sigma)$  as discussed in Remark 6.13.

Another question is the relation of our construction to the topological quantum field theory of [26, 22]. The Hilbert space we have associated to a two-manifold here is infinite-dimensional, whereas the constructions of [26, 22], arising from rational conformal field theory, give finite dimensional vector spaces. On the other hand, we have built our construction for generic values of the coupling  $\lambda$  lying on the real line whereas the constructions of [26, 22] require special values of the coupling lying in a different region of the complex plane. One analogy is to the representations of finite dimensional lie algebras, where for any choice of highest weight, one obtains an infinite dimensional Verma module; for dominant integral choices of this weight, this infinite dimensional module has a finite dimensional quotient. It is plausible that some similar phenomenon might appear here, where the generic quantum field theory we have constructed may be “analytically continued” (in analogy to the types of analytic continuation that give Minkowskian Quantum Field Theory in terms of Euclidean Quantum Field Theory) to values of the couplings where such a finite dimensional quotient may arise. In the context of knot theory, we may speculate that the representations of the Braid group constructed by Jones [15] may be related to quotients of sufficiently high tensor powers of the Burau representation.

**7.3. String Field Theory.** In a more speculative vein, proposals for String Field Theory have also been presented which involve a cubic functional integral not unlike that appearing in Chern-Simons gauge theory. We may also hope by analogy that “finiteness in second order perturbation theory implies existence of the nonperturbative theory in imaginary coupling,” as for the  $\phi^3$  theory described in Section 1.1. For superstring theory, the finiteness of second order perturbation theory has been established by d’Hoker and Phong [6]. If these superstring theories can be written in a satisfactory form as String Field Theory, this may hint at such a construction.

## REFERENCES

- [1] D. Adams, A note on the Faddeev-Popov determinant and Chern-Simons perturbation theory. *Letters in Mathematical Physics* 42, pages 205-214 (1997)
- [2] Aizenman, Michael; Duminil-Copin, Hugo. Marginal triviality of the scaling limits of critical 4D Ising and  $\phi_4^4$  models. *Ann. Math. (2)* 194, No. 1, 163-235 (2021)
- [3] Axelrod, Scott; Singer, I. M. Chern-Simons perturbation theory. *Proceedings of the XXth International Conference on Differential Geometric Methods in Theoretical Physics*, Vol. 1, 2 (New York, 1991), 3-45, World Sci. Publ., River Edge, NJ, 1992.

---

<sup>65</sup>Perhaps the tensors  $\epsilon_{ijk}$  and  $f_{\alpha\beta\gamma}$  combine to give a kind of even analog of a Poisson structure on  $\mathcal{A}$ . A similar appearance in Chern Simons gauge theory of an even analog of a symplectic form arising from the quadratic term in Chern-Simons gauge theory appears in [25].

- [4] H. Burkhardt, Grundzuge einer allgemeinen Systematik der hyperelliptischen Functionen I. Ordnung, *Math. Ann.* 35 (1889), no. 1-2, 198-296  
 –W. H. Meeks, III and J. Patrusky, Representing homology classes by embedded circles on a compact surface, *Illinois J. Math.* 22 (1978), no. 2, 262-269  
 –A. Putman, The symplectic representation of the mapping class group is surjective. <https://www3.nd.edu/~andyp/notes/SymplecticSurjective.pdf>
- [5] Alberto S. Cattaneo, Giovanni Felder. A path integral approach to the Kontsevich quantization formula. *Commun. Math. Phys.* 212 (2000) 591-611
- [6] D'Hoker, Eric; Phong, D. H. Lectures on two-loop superstrings. *Superstring theory*, 85-123, *Adv. Lect. Math. (ALM)*, 1, Int. Press, Somerville, MA, 2008.
- [7] J. Feldman, H. Knorrer, E. Trubowitz. *Fermionic Functional Integrals and the Renormalization Group*. American Mathematical Society, 2002.
- [8] R. Fernandez, J. Froehlich, A. D. Sokal, "Random Walks, Critical Phenomena, and Triviality in Quantum Field Theory". Springer (1992)
- [9] Glimm, James; Jaffe, Arthur. *Quantum physics. A functional integral point of view*. Second edition. Springer-Verlag, New York, 1987.
- [10] J. Glimm, A. Jaffe. The  $\lambda(\phi^4)_2$  Quantum Field Theory without Cutoffs. III. The Physical Vacuum. *Acta Math.* 125 (1970), 203-267
- [11] J. Glimm, A. Jaffe. Positivity of the  $\phi^4_3$  Hamiltonian. *Fortsch. Phys.* 21, 327-376 (1975)
- [12] Gross, Leonard. Abstract Wiener spaces. 1967 Proc. Fifth Berkeley Sympos. Math. Statist. and Probability (Berkeley, Calif., 1965/66), Vol. II: Contributions to Probability Theory, Part 1 pp. 31-42 Univ. California Press, Berkeley, Calif.
- [13] M. Hairer, Regularity structures and the dynamical  $\Phi^4_3$  model. arXiv:1508.05261
- [14] Arthur Jaffe and Fabio L. Pedrocchi. Reflection Positivity for Majoranas. *Ann. Henri Poincaré* 16 (2015), 189-203
- [15] Jones, Vaughan F. R. A polynomial invariant for knots via von Neumann algebras. *Bull. Amer. Math. Soc. (N.S.)* 12 (1985), no. 1, 103-111.
- [16] Kontsevich, Maxim. Intersection theory on the moduli space of curves and the matrix Airy function. *Comm. Math. Phys.* 147 (1992), no. 1, 1–23.
- [17] Kontsevich, Maxim. Deformation quantization of Poisson manifolds. *Lett. Math. Phys.* 66, No. 3, 157-216 (2003).
- [18] Lax, P. *Functional Analysis*. Wiley, 2002.
- [19] Osterwalder, Konrad; Schrader, Robert. Axioms for Euclidean Green's functions. *Comm. Math. Phys.* 31 (1973), no. 2, 83-112.  
 –Axioms for Euclidean Green's functions. II. *Commun. Math. Phys.* 42, 281-305 (1975).
- [20] Osterwalder, Konrad; Schrader, Robert. Euclidean Fermi fields and a Feynman-Kac formula for Boston-Fermion models. *Helv. Phys. Acta*, 46 (1973), 277-301
- [21] Reed, M., Simon, B. *Fourier Analysis, Self Adjointness*. Academic Press, 1975.
- [22] Reshetikhin, N.; Turaev, V. G. Invariants of 3-manifolds via link polynomials and quantum groups. *Invent. Math.* 103 (1991), no. 3, 547-597
- [23] Schwarz, A. S. The partition function of degenerate quadratic functional and Ray-Singer invariants. *Lett. Math. Phys.* 2 (1977/78), no. 3, 247-252.
- [24] Streater, R. F.; Wightman, A. S. *PCT, spin and statistics, and all that*. W. A. Benjamin, Inc., New York-Amsterdam 1964 viii+181 pp.
- [25] J. Weitsman, The Chern-Simons Functional Integral, Kauffman's Bracket Polynomial, and other link invariants. arXiv:2405.16775
- [26] Witten, Edward. Quantum field theory and the Jones polynomial. *Comm. Math. Phys.* 121 (1989), no. 3, 351-399




FIGURE 1. The manifold  $M$  cut into three open sets by two isotopic submanifolds  $\Sigma_1, \Sigma_2$

#### APPENDIX A. FORMULAS FOR THE MULTIPLICATION ACTION

In this section we briefly review a multiplication formula for the fields in the case where  $M$  is split into three components by submanifolds isotopic to  $\Sigma$ , which we will not use directly, but which we record to show the topological nature of the theory. Suppose  $\Sigma_1, \Sigma_2$  are two compact connected, disjoint submanifolds of  $M$ , each isotopic to  $\Sigma$ . Each of these splits  $M$  into two components, and the union  $\Sigma_1 \cup \Sigma_2$  splits  $M$  into three components  $U, V, W$  (see Figure 1).

We begin with formulas for the gauge fields. Using Proposition 4.31, we have

**Lemma A.1.** *Suppose  $p \in \mathcal{F}_P^B(U), q \in \mathcal{F}_P^B(V), r \in \mathcal{F}_P^B(W)$ . Then*

$$\Phi(pqr) = \langle \hat{\pi}_{\Sigma_1}^B(p), S^* \hat{\pi}_{\Sigma_1}^B(qr) \rangle_{H^B(\Sigma_1)} = \langle \hat{\pi}_{\Sigma_2}^B(pq), S^* \hat{\pi}_{\Sigma_2}^B(r) \rangle_{H^B(\Sigma_2)};$$

where elements of  $\text{Sym}^* \Lambda \subset \text{Sym}^* H^1(\Sigma_1, \mathfrak{g})$  and  $\text{Sym}^* \Lambda \subset \text{Sym}^* H^1(\Sigma_2, \mathfrak{g})$  are identified with elements of  $H^B(\Sigma)$  using the isotopy. Also,

$$\hat{\pi}_{\Sigma_1}^B(p) = \hat{\pi}_{\Sigma_2}^B(p).^{66}$$

Similarly for the ghosts, using Proposition 5.26, we have the following

<sup>66</sup>Note that as in Proposition 4.7, this indicates that the Hamiltonian in this theory is zero.

**Lemma A.2.** *Suppose  $p \in \mathcal{F}_P^F(U), q \in \mathcal{F}_P^F(V), r \in \mathcal{F}_P^F(W)$ . Then  $pq \in \mathcal{F}^F(U \cup V), qr \in \mathcal{F}^F(V \cup W)$ , and*

$$\Psi(pqr) = \langle \pi_{\Sigma_1}^F(p^c), \mathcal{J}_F \pi_{\Sigma_1}^F(qr) \rangle_{H^F(\Sigma_1)} = \langle \pi_{\Sigma_2}^F(q^c p^c), \mathcal{J}_F \pi_{\Sigma_2}^F(r) \rangle_{H^F(\Sigma_2)};$$

and

$$\pi_{\Sigma_1}^F(p) = \pi_{\Sigma_2}^F(p).$$

Combining Lemma A.2 with Lemma A.1, we then have the following formula for polynomials in the Bose and Fermi fields. Let  $\Sigma_1, \Sigma_2, U, V, W$  be as above. Assume that the orientation of  $\Sigma_1$  obtained from the isotopy agree with the orientation of the boundary of  $U$ . Write

$$\hat{\pi}_{\Sigma_i} = \hat{\pi}_{\Sigma_i}^B \otimes \pi_{\Sigma_i}^F, i = 1, 2.$$

Then<sup>67</sup>

**Proposition A.3.** *Suppose  $p \in \mathcal{F}_P(U), q \in \mathcal{F}_P(V), r \in \mathcal{F}_P(W)$ . Then*

$$(\Phi \otimes \Psi)(pqr) = \langle \hat{\pi}_{\Sigma_1}(p^c), \mathcal{J}(S^* \otimes \text{id}) \hat{\pi}_{\Sigma_1}(qr) \rangle_{H(\Sigma_1)} = \langle \hat{\pi}_{\Sigma_2}(q^c p^c), \mathcal{J}(S^* \otimes \text{id}) \hat{\pi}_{\Sigma_2}(r) \rangle_{H(\Sigma_2)},$$

where we extend the action of  $S^*$  on  $H^B(\Sigma)$  to the operator  $S^* \otimes \text{id}$  on  $H(\Sigma)$  by defining its action on Fermi fields to be the identity. Likewise we define the action of  $(\cdot)^c$  to  $\mathcal{F}_P(U)(U)$  by defining its action on Bose fields to be the identity. Furthermore

$$\hat{\pi}_{\Sigma_1}(p) = \hat{\pi}_{\Sigma_2}(p).$$

DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, MA 02115  
Email address: j.weitsman@neu.edu

---

<sup>67</sup>Note that the action of  $S^*$  on the even cohomology of  $\Sigma$  amounts to one negative sign on  $H^2$ , which is morally accounted for by the signs in the adjoint  $(\cdot)^c$ .