

# Study on relativistic transformations for thermodynamic quantities: Temperature of a moving system of quark-gluon plasma

A.S. Parvan<sup>a,b</sup>

<sup>a</sup>*Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna, Russia*

<sup>b</sup>*Department of Theoretical Physics, Horia Hulubei National Institute for R&D in Physics and Nuclear Engineering, Bucharest-Magurele, Romania*

---

## Abstract

Thermodynamics of systems moving with relativistic velocities is directly derived from the Lorentz transformation of four-momentum. It is shown that eight different perspectives on relativistic thermodynamics can be determined in the framework of the fundamental ensemble. However, only two sets of relativistic transformations for the physical thermodynamic quantities are consistent with the requirements of the special theory of relativity and the Hamiltonian and Lagrangian mechanics. These relativistic transformations are equivalent and correspond exactly to the Planck transformations for physical temperature and physical chemical potential. Other six sets of relativistic transformations of thermodynamic quantities are either unphysical due to the dependence of rest energy of the thermodynamic system on 3-velocity and 3-momentum or forbidden by the Hamiltonian and Lagrangian mechanics. It is found that the relativistic transformations for unphysical thermodynamic quantities corresponding to Planck's and Ott's transformations are unphysical and contradict the requirements of the special theory of relativity. The four classes of the obtained relativistic transformations of the physical thermodynamic quantities are compared by studying the thermodynamic properties of a relativistically moving system of the quark-gluon plasma (QGP).

*Keywords:* Relativistic thermodynamics, Lorentz transformations, Planck's formalism, Ott's formalism, Quark-gluon plasma

---

## 1. Introduction

The problem of the foundation of relativistic thermodynamics of a moving body remains unresolved [1, 2, 3]. There are two main views on relativistic thermodynamics: Planck's formalism [4, 5, 6] and Ott's one [7]. Relativistic transformations of temperature and other thermodynamic quantities in these two formalisms are mutually exclusive. In the Planck theory, the temperature  $T$  in the inertial reference frame which moves rectilinearly at velocity  $v$  relative to the thermodynamic system of temperature  $T_0$  at rest in another inertial reference frame is  $T = T_0/\gamma < T_0$ ; however, in the Ott formalism it is  $T = \gamma T_0 > T_0$ , where  $\gamma = (1 - v^2/c^2)^{-1/2}$ . This puzzle and other proposed formalisms led to a controversy in relativistic thermodynamics [8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35]. Therefore, a unique solution to this problem is a primary aim of relativistic thermodynamics of a moving body. This solution may be essential for constructing self-consistent relativistic statistical mechanics useful in high-energy physics, astrophysics, and cosmology.

Many works treated the Planck-Ott puzzle (see, e.g., the reviews [1, 2]). The main aim of this study is to solve this problem. For this purpose, we derive analytically four different classes of relativistic transformations for thermodynamic quantities considering the fundamental thermodynamic potential to

be either a relativistic Hamiltonian or a Lagrangian and an independent state variable to be either 3-momentum or 3-velocity. Each class is divided into two subclasses for physical and unphysical thermodynamic quantities assuming either the rest volume or relativistically contracted volume of a thermodynamic system to be an independent state variable. We have found that from eight sets of relativistic transformations only the relativistic transformations for physical thermodynamic quantities of class I corresponding to the Hamiltonian as a function of 3-momentum and of class III corresponding to the Lagrangian as a function of 3-velocity are consistent with the requirements of the special theory of relativity and the Hamiltonian and Lagrangian mechanics. These two sets of relativistic transformations obtained by the Hamiltonian and Lagrangian formalisms are equivalent and correspond to the Planck transformations for the physical temperature and physical chemical potential. For physical pressure, a new formula has been obtained. Relativistic transformations for physical and unphysical thermodynamic quantities of class II related to the Lagrangian as a function of 3-momentum and of class IV corresponding to the Hamiltonian as a function of 3-velocity are forbidden. Relativistic transformations for unphysical thermodynamic quantities of all classes are inconsistent with the requirements of the special theory of relativity.

The paper is organized as follows: In Section 2, we define the relativistic thermodynamic system. In Sections 3 and 4, we derive relativistic transformations for thermodynamic quantities. In Section 5, we study the properties of a moving system of the QGP. The conclusions are summarized in the final section.

---

*Email address:* parvan@theor.jinr.ru,  
parvan@theory.nipne.ro (A.S. Parvan)

## 2. Relativistic thermodynamic system: Fundamental ensemble

Let a relativistically moving thermodynamic system  $\mathcal{A}$  of mass  $M$  be given in two inertial reference frames  $K$  and  $K_0$ , which is at rest in the moving reference frame  $K_0$ . Then the 4-momentum of the thermodynamic system  $\mathcal{A}$  in the inertial reference frame  $K_0$  is  $P_0^\mu = (E_0/c, \mathbf{P}_0)$  with the 3-momentum  $\mathbf{P}_0 = 0$  and the total energy  $E_0 = Mc^2$ . Let the rest energy of the thermodynamic system  $E_0$  be a fundamental thermodynamic potential in the reference frame  $K_0$ , i.e.  $E_0 = E_0(S_0, V_0, N_0)$ , where  $S_0, V_0$  and  $N_0$  are the rest entropy, rest volume and rest number of particles, respectively. Then the differential of the thermodynamic potential  $E_0$  with respect to the thermodynamic state variables  $(S_0, V_0, N_0)$  can be written as [35]

$$dE_0 = T_0 dS_0 - p_0 dV_0 + \mu_0 dN_0 = \delta Q_0 - \delta W_0 + dE_{mat}^0, \quad (1)$$

$$T_0 \equiv \left( \frac{\partial E_0}{\partial S_0} \right)_{V_0, N_0}, \quad p_0 \equiv - \left( \frac{\partial E_0}{\partial V_0} \right)_{S_0, N_0}, \quad \mu_0 \equiv \left( \frac{\partial E_0}{\partial N_0} \right)_{S_0, V_0}. \quad (2)$$

Here  $T_0, p_0$  and  $\mu_0$  are respectively the temperature, pressure and chemical potential and  $\delta Q_0 = T_0 dS_0, \delta W_0 = p_0 dV_0$  and  $dE_{mat}^0 = \mu_0 dN_0$  are respectively the heat transfer, mechanical work due to a volume change and change in energy due to the exchange of matter for the thermodynamic system in the inertial reference frame  $K_0$ . Equation (1) represents the first law of thermodynamics in the inertial reference frame  $K_0$ , where the thermodynamic system is at rest.

The reference frame  $K_0$  and the thermodynamic system  $\mathcal{A}$  move rectilinearly relative to the fixed laboratory reference frame  $K$  with constant velocity  $\mathbf{v} = (v, 0, 0)$  along the positive directions of the  $x$  and  $x_0$  axes. The 4-momentum of the thermodynamic system  $\mathcal{A}$  in the inertial reference frame  $K$  is  $P^\mu = (E/c, \mathbf{P})$  with the 3-momentum  $\mathbf{P}$  and the total energy  $E$ . The Lorentz transformations for 4-momenta, when passing from the coordinate system  $K_0$  to the coordinate system  $K$ , have the following form [36]:

$$E = \gamma(E_0 + vP_x^0), \quad P_x = \gamma\left(P_x^0 + \frac{vE_0}{c^2}\right), \quad P_\perp = P_\perp^0, \quad (3)$$

$$\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (4)$$

where  $\gamma$  is the Lorentz factor. Substituting the value of the 3-momentum  $\mathbf{P}_0 = 0$  into Eq. (3), we obtain [36]

$$E = \gamma E_0, \quad \mathbf{P} = \frac{\mathbf{v}}{c^2} E, \quad (5)$$

where  $\mathbf{P} = (P_x, 0, 0)$ . The velocity  $\mathbf{v}$  of the frame  $K_0$  and the thermodynamic system  $\mathcal{A}$  can be written as [36]

$$\mathbf{v} = \frac{\mathbf{P}c^2}{E}. \quad (6)$$

The total energy  $E$ , which is the Hamiltonian, represents the fundamental thermodynamic potential of the thermodynamic

system  $\mathcal{A}$  in the reference frame  $K$ . The Lagrangian of the thermodynamic system  $\mathcal{A}$  in the reference frame  $K$  can be written as [36]:

$$L = \mathbf{v}\mathbf{P} - E = -\frac{E_0}{\gamma}. \quad (7)$$

The function,  $\bar{L} \equiv E - \mathbf{v}\mathbf{P} = -L$ , obtained from the fundamental thermodynamic potential  $E$  by the Legendre transform is a new thermodynamic potential which we will name the dual fundamental thermodynamic potential defining the dual fundamental ensemble. Nevertheless, in this paper, the negative Lagrangian,  $-L$ , and the usual names for fundamental thermodynamic potential and fundamental ensemble will be used.

The entropy  $S$  and the number of particles  $N$  are relativistic 4-invariants, and the volume  $V$  of the thermodynamic system undergoes the Lorentz contraction. Therefore, we have [1, 2]

$$S = S_0, \quad V = \frac{V_0}{\gamma}, \quad N = N_0. \quad (8)$$

For the relativistically moving thermodynamic system  $\mathcal{A}$ , four independent classes of thermodynamic quantities can be defined: I) the Hamiltonian is a function of 3-momentum, II) the Lagrangian is a function of 3-momentum, III) the Lagrangian is a function of 3-velocity, IV) the Hamiltonian is a function of 3-velocity. Let us consider these classes of functions.

## 3. Momentum as an independent state variable

### 3.1. Class I: Hamiltonian $E(\mathbf{P})$ as the fundamental thermodynamic potential

Let the 3-momentum  $\mathbf{P}$  be an independent state variable in the Hamilton function  $E$  in conformity with the requirements of the Hamiltonian mechanics.

#### 3.1.1. Independent physical state variables $(S_0, V_0, N_0, \mathbf{P})$

The rest energy of the thermodynamic system  $\mathcal{A}$  is a function of the state variables  $(S_0, V_0, N_0)$ , i.e.  $E_0 = E_0(S_0, V_0, N_0)$ . Then, using Eqs. (4)–(6), we obtain

$$\gamma(S_0, V_0, N_0, \mathbf{P}) = \sqrt{1 + \frac{\mathbf{P}^2 c^2}{E_0^2(S_0, V_0, N_0)}}. \quad (9)$$

Substituting Eq. (9) into Eq. (5), we obtain the Hamiltonian as

$$E(S_0, V_0, N_0, \mathbf{P}) = \gamma E_0 = \sqrt{\mathbf{P}^2 c^2 + E_0^2(S_0, V_0, N_0)}. \quad (10)$$

Substituting Eq. (10) into Eq. (6) and Eq. (9) into Eq. (8), we obtain the velocity and the volume of the system as

$$\mathbf{v}(S_0, V_0, N_0, \mathbf{P}) = \frac{\mathbf{P}c^2}{E} = \frac{\mathbf{P}c^2}{\sqrt{\mathbf{P}^2 c^2 + E_0^2(S_0, V_0, N_0)}}, \quad (11)$$

$$V(S_0, V_0, N_0, \mathbf{P}) = \frac{V_0}{\gamma} = \frac{V_0}{\sqrt{1 + \frac{\mathbf{P}^2 c^2}{E_0^2(S_0, V_0, N_0)}}}. \quad (12)$$

Let the relativistic Hamiltonian (10), which is the total energy of the thermodynamic system, be a fundamental thermodynamic potential in the inertial reference frame  $K$ . This function depends on the variables  $(S_0, V_0, N_0, \mathbf{P})$ . Thus, the variables  $(S_0, V_0, N_0, \mathbf{P})$  are independent state variables of the thermodynamic system. This is dictated by the fundamental thermodynamic potential  $E_0(S_0, V_0, N_0)$  in the inertial reference frame  $K_0$  and the Lorentz transformations (5), (10) considering momentum  $\mathbf{P}$  as an independent state variable. The total energy (10) depends on the rest volume  $V_0$  and does not depend on the contracted volume  $V$  because the rest energy  $E_0$  is a function of  $V_0$ . Therefore, the other thermodynamic quantities in the reference frame  $K$  are also dependent on  $V_0$  but not on  $V$  since the thermodynamic quantities in the fundamental ensemble are defined by the partial derivatives of the fundamental thermodynamic potential  $E$  with respect to the state variables. Thus, the rest volume  $V_0$  is an independent state variable and the volume  $V$  is a dependent variable. The infinitesimal physical changes in the rest entropy  $dS_0$ , rest volume  $dV_0$  and rest number of particles  $dN_0$  of the thermodynamic system  $\mathcal{A}$  in the inertial reference frame  $K_0$ , where  $\mathcal{A}$  is at rest, lead to direct physical changes in the internal energy  $E_0$  and total energy (10) of the system. Moreover, these changes are independent of momentum  $\mathbf{P}$  and velocity  $\mathbf{v}$  of  $\mathcal{A}$  in the reference frame  $K$  in accordance with the principles of the special theory of relativity. Therefore, for the independent rest volume  $V_0$ , the variables  $(S_0, V_0, N_0, \mathbf{P})$  are *independent physical state variables* and the thermodynamic quantities corresponding to them are *physical thermodynamic quantities*. It should be stressed that the state variables  $(S_0, V_0, N_0, \mathbf{P})$  can be changed independently from each other only by quasi-static thermodynamic processes which preserve the system  $\mathcal{A}$  in the thermodynamic equilibrium.

Taking the differential of the Hamilton function (10) with respect to the independent state variables  $(S_0, V_0, N_0, \mathbf{P})$  and using Eqs. (2), (5) and (6), we obtain

$$\begin{aligned} dE(S_0, V_0, N_0, \mathbf{P}) &= \frac{T_0}{\gamma} dS_0 - \frac{p_0}{\gamma} dV_0 + \frac{\mu_0}{\gamma} dN_0 + \mathbf{v}d\mathbf{P} \\ &= T_I dS_0 - p_I dV_0 + \mu_I dN_0 + \mathbf{v}_I d\mathbf{P}, \end{aligned} \quad (13)$$

where

$$\begin{aligned} \left( \frac{\partial E}{\partial S_0} \right)_{V_0 N_0 \mathbf{P}} &\equiv T_I, & \left( \frac{\partial E}{\partial V_0} \right)_{S_0 N_0 \mathbf{P}} &\equiv -p_I, \\ \left( \frac{\partial E}{\partial N_0} \right)_{S_0 V_0 \mathbf{P}} &\equiv \mu_I, & \left( \frac{\partial E}{\partial \mathbf{P}} \right)_{S_0 V_0 N_0} &\equiv \mathbf{v}_I = \mathbf{v}. \end{aligned} \quad (14)$$

Here  $T_I, p_I$  and  $\mu_I$  are respectively the physical temperature, physical pressure and physical chemical potential of class I for the thermodynamic system  $\mathcal{A}$  in the reference frame  $K$ ;  $\gamma$  and  $\mathbf{v}$  are calculated by Eqs. (9) and (11). Substituting Eq. (1) into (13), we have

$$\begin{aligned} dE(S_0, V_0, N_0, \mathbf{P}) &= \delta Q_I - \delta W_I + dE_{mat}^I + dE_{mov} \\ &= \frac{1}{\gamma} dE_0 + \mathbf{v}d\mathbf{P} = \frac{1}{\gamma} \delta Q_0 - \frac{1}{\gamma} \delta W_0 + \frac{1}{\gamma} dE_{mat}^0 + dE_{mov}, \end{aligned} \quad (15)$$

where  $\delta Q_I = T_I dS_0$ ,  $\delta W_I = p_I dV_0$ ,  $dE_{mat}^I = \mu_I dN_0$  and  $dE_{mov} = \mathbf{v}d\mathbf{P}$  are respectively the heat transfer, mechanical work due to a volume change, change in energy due to the exchange of matter and change in energy of motion of the thermodynamic system as a whole for the thermodynamic system in the laboratory reference frame  $K$ . Equation (15) represents the first law of thermodynamics in the laboratory reference frame  $K$  and the Lorentz transformation for the differential of energy  $dE_0$ .

Comparing Eqs. (13)–(15) with each other, we obtain the relativistic transformations for physical thermodynamic quantities of class I as

$$T_I = \frac{T_0}{\gamma}, \quad p_I = \frac{p_0}{\gamma}, \quad \mu_I = \frac{\mu_0}{\gamma}, \quad \mathbf{v}_I = \mathbf{v}, \quad (16)$$

$$\delta Q_I = \frac{1}{\gamma} \delta Q_0, \quad \delta W_I = \frac{1}{\gamma} \delta W_0, \quad dE_{mat}^I = \frac{1}{\gamma} dE_{mat}^0. \quad (17)$$

These are exactly the Planck transformations [1, 2, 4, 5, 6, 27, 35] for thermodynamic quantities with the exception of pressure. We have the invariance for the ratio  $p_I/T_I = p_0/T_0$ . Note that the formula for the relativistic transformation of physical pressure given in Eq. (16) was obtained for the first time.

### 3.1.2. Independent unphysical state variables $(S, V, N, \mathbf{P})$

Let the rest physical volume  $V_0$  be a dependent variable and the contracted volume  $V$  in the reference frame  $K$  be an independent state variable. Substituting the relations  $S_0 = S$  and  $N_0 = N$  from Eq. (8) into Eq. (12), we obtain

$$V = \frac{V_0}{\sqrt{1 + \frac{\mathbf{P}^2 c^2}{E_0^2(S, V_0, N)}}}. \quad (18)$$

Solving Eq. (18) with respect to  $V_0$ , we obtain the function  $V_0 = V_0(S, V, N, \mathbf{P})$ . Then, the fundamental thermodynamic potential  $E_0(S_0, V_0, N_0)$  in the reference frame  $K_0$  can be rewritten as

$$E_0(S, V, N, \mathbf{P}) = E_0(S, V_0(S, V, N, \mathbf{P}), N). \quad (19)$$

Substituting Eq. (19) into Eq. (10), we have

$$E(S, V, N, \mathbf{P}) = \sqrt{\mathbf{P}^2 c^2 + E_0^2(S, V_0(S, V, N, \mathbf{P}), N)}. \quad (20)$$

Substituting Eq. (19) into Eqs. (9) and (11), we obtain

$$\gamma(S, V, N, \mathbf{P}) = \sqrt{1 + \frac{\mathbf{P}^2 c^2}{E_0^2(S, V_0(S, V, N, \mathbf{P}), N)}}, \quad (21)$$

$$\mathbf{v}(S, V, N, \mathbf{P}) = \frac{\mathbf{P} c^2}{\sqrt{\mathbf{P}^2 c^2 + E_0^2(S, V_0(S, V, N, \mathbf{P}), N)}}. \quad (22)$$

Let the relativistic Hamiltonian (20) be a fundamental thermodynamic potential as a function of the independent state variables  $(S, V, N, \mathbf{P})$ . In this case, the internal rest energy  $E_0$  of the thermodynamic system  $\mathcal{A}$  in the inertial reference frame  $K_0$ , where  $\mathcal{A}$  is at rest, depend on the momentum  $\mathbf{P}$  of  $\mathcal{A}$  in the laboratory reference frame  $K$  (see Eq. (19)). In the special theory of relativity,  $E_0$  is independent of  $\mathbf{P}$  and  $\mathbf{v}$ . Thus, this is

an unphysical situation which violates the requirements of the special theory of relativity. For independent  $V$ , the rest volume  $V_0$ , rest energy  $E_0$ , rest temperature  $T_0$ , rest pressure  $p_0$  and the rest chemical potential  $\mu_0$  of the thermodynamic system  $\mathcal{A}$  in the inertial reference frame  $K_0$ , where  $\mathcal{A}$  is at rest, will change with momentum  $\mathbf{P}$  and velocity  $\mathbf{v}$  of the system at fixed values of  $S, V, N$  in the inertial reference frame  $K$ . This is not in accordance with the principles of the special theory of relativity. Moreover, it is difficult to imagine a physically possible situation in which the change of the contracted volume  $V$  leads to a change of the physical volume  $V_0$  of the thermodynamic system  $\mathcal{A}$  and produces the work  $\delta W$ . This violates the principle of causality. Only a change in the physical volume  $V_0$  of the thermodynamic system  $\mathcal{A}$  leads to a change in the contracted volume  $V$  but not vice versa. The volume  $V$  cannot be controlled thermodynamically as an independent quantity. Therefore, for the independent contracted volume  $V$ , the variables  $(S, V, N, \mathbf{P})$  are *independent unphysical state variables* and the thermodynamic quantities corresponding to them are *unphysical thermodynamic quantities*. The mechanical work  $\delta W$  due to a volume change is produced only by a change  $dV_0$  of the rest physical volume.

The differential of  $E$  with respect to the unphysical state variables  $(S, V, N, \mathbf{P})$  of the thermodynamic system  $\mathcal{A}$  in the reference frame  $K$  can be written as

$$dE(S, V, N, \mathbf{P}) = \tilde{T}_I dS - \tilde{p}_I dV + \tilde{\mu}_I dN + \tilde{\mathbf{v}}_I d\mathbf{P}, \quad (23)$$

$$\begin{aligned} \left(\frac{\partial E}{\partial S}\right)_{VNP} &\equiv \tilde{T}_I, & \left(\frac{\partial E}{\partial V}\right)_{SNP} &\equiv -\tilde{p}_I, \\ \left(\frac{\partial E}{\partial N}\right)_{SVP} &\equiv \tilde{\mu}_I, & \left(\frac{\partial E}{\partial \mathbf{P}}\right)_{SVN} &\equiv \tilde{\mathbf{v}}_I. \end{aligned} \quad (24)$$

Here  $\tilde{T}_I, \tilde{p}_I, \tilde{\mu}_I$  and  $\tilde{\mathbf{v}}_I$  are respectively the unphysical temperature, unphysical pressure, unphysical chemical potential and generalized velocity of class I for the thermodynamic system  $\mathcal{A}$  in the reference frame  $K$ .

Taking the differential of volume (18) with respect to the variables  $(S, V, N, \mathbf{P})$  and using the relations of independence of the state variables  $\partial V/\partial S = \partial V/\partial N = \partial V/\partial \mathbf{P} = 0$ , we obtain

$$\begin{aligned} dV_0(S, V, N, \mathbf{P}) &= -\alpha \frac{\mathbf{P}^2 c^2}{E^2} \frac{V_0 T_0}{E_0} dS - \alpha \frac{\mathbf{P}^2 c^2}{E^2} \frac{V_0 \mu_0}{E_0} dN \\ &+ \alpha \gamma dV + \frac{\alpha V_0}{\gamma E_0} \mathbf{v} d\mathbf{P}, \end{aligned} \quad (25)$$

where

$$\alpha \equiv \frac{1}{1 - \frac{\mathbf{P}^2 c^2}{E^2} \frac{V_0 p_0}{E_0}} = \frac{1}{1 - \frac{\mathbf{v}^2}{c^2} \frac{V_0 p_0}{E_0}}. \quad (26)$$

Substituting Eq. (20) into Eq. (24) and using the partial derivatives from Eq. (25), we obtain the relativistic transformations for the unphysical thermodynamic quantities (24) as

$$\tilde{T}_I = \frac{\alpha T_0}{\gamma}, \quad \tilde{p}_I = \alpha p_0, \quad \tilde{\mu}_I = \frac{\alpha \mu_0}{\gamma}, \quad \tilde{\mathbf{v}}_I = \alpha \left(1 - \frac{V_0 p_0}{E_0}\right) \mathbf{v}. \quad (27)$$

We have the invariance for the ratio  $\tilde{p}_I V / \tilde{T}_I = p_0 V_0 / T_0$ . It should be stressed that the relativistic transformations (27) for

the unphysical thermodynamic quantities (24) and the transformations (16) for the physical thermodynamic quantities (14) of class I are consistent because they correspond to the fundamental thermodynamic potential represented by the Hamilton function in which the 3-momentum is an independent state variable according to the requirements of the Hamiltonian mechanics. However, the relativistic transformations (27) are *unphysical* because  $V_0$  is a dependent variable and  $E_0, T_0, p_0$  and  $\mu_0$  depend on  $\mathbf{P}$  (see Eq. (19)) in contradiction with the requirements of the special theory of relativity.

Substituting Eq. (25) into Eq. (13) and using the relations  $dS_0 = dS$  and  $dN_0 = dN$ , we exactly obtain Eq. (23) and the relativistic transformations (27).

*Incomplete formula for the differential of volume.* Suppose that the differential of volume (8), (12) with respect to the state variables  $(S_0, V_0, N_0, \mathbf{P})$  is [2, 19] (cf. with Eq. (25))

$$dV = \frac{dV_0}{\gamma}, \quad (28)$$

i.e. the Lorentz factor (9) is constant and does not depend on the state variables,  $d\gamma(S_0, V_0, N_0, \mathbf{P}) = 0$ . In this case, it is easy to show that Eq. (28) is valid only for  $T_0 = p_0 = \mu_0 = 0$  and  $\mathbf{v} = 0$ . See also Eqs. (25) and (26). Note that Eq. (28) is not consistent with the formula for the Lorentz volume contraction (12). Substituting Eq. (28) into Eq. (13) and using the relations  $dS_0 = dS$  and  $dN_0 = dN$ , we obtain Eq. (23), where (cf. with Eqs. (27) and (16))

$$\tilde{T}_I = \frac{T_0}{\gamma}, \quad \tilde{p}_I = p_0, \quad \tilde{\mu}_I = \frac{\mu_0}{\gamma}, \quad \tilde{\mathbf{v}}_I = \mathbf{v}. \quad (29)$$

Relations (29) are exactly the Planck transformation formulae [1, 2, 4, 5, 6, 27, 35]. Thus, the Planck transformations (29) in the variables  $(S, V, N, \mathbf{P})$  derived from the incomplete formula for the differential of the volume (28) are valid only in the trivial case for  $T_0 = p_0 = \mu_0 = 0$  and  $\mathbf{v} = 0$ .

### 3.2. Class II: Lagrangian $L(\mathbf{P})$ as the fundamental thermodynamic potential

Let the 3-momentum  $\mathbf{P}$  be an independent state variable in the Lagrange function  $L$  in contradiction with the requirements of the Lagrangian mechanics.

#### 3.2.1. Independent physical state variables $(S_0, V_0, N_0, \mathbf{P})$

Let the rest volume  $V_0$  be an independent state variable. Substituting Eq. (9) into Eq. (7), we obtain the Lagrangian as

$$L(S_0, V_0, N_0, \mathbf{P}) = -\frac{E_0}{\gamma} = -\frac{E_0(S_0, V_0, N_0)}{\sqrt{1 + \frac{\mathbf{P}^2 c^2}{E_0^2(S_0, V_0, N_0)}}}. \quad (30)$$

Let the Lagrangian (30) be a fundamental thermodynamic potential as a function of the independent physical state variables  $(S_0, V_0, N_0, \mathbf{P})$ .

Taking the differential of the Lagrange function (30) with respect to the physical state variables  $(S_0, V_0, N_0, \mathbf{P})$  and using Eqs. (2), (5), (6) and (9), we obtain

$$dL(S_0, V_0, N_0, \mathbf{P}) = -\frac{\theta T_0}{\gamma} dS_0 + \frac{\theta p_0}{\gamma} dV_0 - \frac{\theta \mu_0}{\gamma} dN_0 + \frac{\mathbf{v}}{\gamma^2} d\mathbf{P} \\ = -T_{II} dS_0 + p_{II} dV_0 - \mu_{II} dN_0 + \mathbf{v}_{II} d\mathbf{P}, \quad (31)$$

$$\theta \equiv 1 + \frac{\mathbf{P}^2 c^2}{E^2} = 1 + \frac{\mathbf{v}^2}{c^2}, \quad (32)$$

where

$$\left( \frac{\partial L}{\partial S_0} \right)_{V_0, N_0, \mathbf{P}} \equiv -T_{II}, \quad \left( \frac{\partial L}{\partial V_0} \right)_{S_0, N_0, \mathbf{P}} \equiv p_{II}, \\ \left( \frac{\partial L}{\partial N_0} \right)_{S_0, V_0, \mathbf{P}} \equiv -\mu_{II}, \quad \left( \frac{\partial L}{\partial \mathbf{P}} \right)_{S_0, V_0, N_0} = \mathbf{v}_{II}. \quad (33)$$

Here,  $T_{II}$ ,  $p_{II}$  and  $\mu_{II}$  are respectively the physical temperature, physical pressure, physical chemical potential and generalized velocity of class II for the thermodynamic system  $\mathcal{A}$  in the reference frame  $K$ ;  $\gamma$ ,  $\mathbf{v}$  and  $V$  are calculated by Eqs. (9), (11) and (12), respectively.

Comparing Eqs. (31), (33) with each other, we obtain the relativistic transformations for physical thermodynamic quantities of class II as

$$T_{II} = \frac{\theta T_0}{\gamma}, \quad p_{II} = \frac{\theta p_0}{\gamma}, \quad \mu_{II} = \frac{\theta \mu_0}{\gamma}, \quad \mathbf{v}_{II} = \frac{\mathbf{v}}{\gamma^2}. \quad (34)$$

We have the invariance for the ratio  $p_{II}/T_{II} = p_0/T_0$ . Comparing Eq. (34) with Eq. (16), we obtain the relation between the physical thermodynamic quantities of classes I and II as

$$T_{II} = \theta T_I, \quad p_{II} = \theta p_I, \quad \mu_{II} = \theta \mu_I. \quad (35)$$

### 3.2.2. Independent unphysical state variables $(S, V, N, \mathbf{P})$

Let the rest physical volume  $V_0$  be a dependent variable and the contracted volume  $V$  in the reference frame  $K$  be an independent state variable. The Lagrangian (30) in the unphysical state variables  $(S, V, N, \mathbf{P})$  can be rewritten as (cf. with Eqs. (20) and (19))

$$L(S, V, N, \mathbf{P}) = -\frac{E_0(S, V_0(S, V, N, \mathbf{P}), N)}{\sqrt{1 + \frac{\mathbf{P}^2 c^2}{E_0^2(S, V_0(S, V, N, \mathbf{P}), N)}}. \quad (36)$$

The rest volume,  $V_0 = V_0(S, V, N, \mathbf{P})$ , is a solution of Eq. (18) and  $\gamma$  and  $\mathbf{v}$  are calculated by Eqs. (21) and (22), respectively. Let the Lagrangian (36) be a fundamental thermodynamic potential as a function of the independent unphysical state variables  $(S, V, N, \mathbf{P})$ .

The differential of  $L$  with respect to the state variables  $(S, V, N, \mathbf{P})$  of the thermodynamic system  $\mathcal{A}$  in the reference frame  $K$  can be written as

$$dL(S, V, N, \mathbf{P}) = -\tilde{T}_{II} dS + \tilde{p}_{II} dV - \tilde{\mu}_{II} dN + \tilde{\mathbf{v}}_{II} d\mathbf{P}, \quad (37)$$

where

$$\left( \frac{\partial L}{\partial S} \right)_{V, N, \mathbf{P}} \equiv -\tilde{T}_{II}, \quad \left( \frac{\partial L}{\partial V} \right)_{S, N, \mathbf{P}} \equiv \tilde{p}_{II}, \\ \left( \frac{\partial L}{\partial N} \right)_{S, V, \mathbf{P}} \equiv -\tilde{\mu}_{II}, \quad \left( \frac{\partial L}{\partial \mathbf{P}} \right)_{S, V, N} \equiv \tilde{\mathbf{v}}_{II}. \quad (38)$$

Here  $\tilde{T}_{II}$ ,  $\tilde{p}_{II}$ ,  $\tilde{\mu}_{II}$  and  $\tilde{\mathbf{v}}_{II}$  are respectively the unphysical temperature, unphysical pressure, unphysical chemical potential and the generalized velocity of class II for the thermodynamic system  $\mathcal{A}$  in the reference frame  $K$ . Substituting Eq. (36) into Eq. (38) and using partial derivatives from Eq. (25), we obtain the relativistic transformations for the unphysical thermodynamic quantities (38) of class II as

$$\tilde{T}_{II} = \frac{\alpha \theta T_0}{\gamma}, \quad \tilde{p}_{II} = \alpha \theta p_0, \quad \tilde{\mu}_{II} = \frac{\alpha \theta \mu_0}{\gamma}, \quad \tilde{\mathbf{v}}_{II} = \left(1 + \frac{V_0 p_0}{E_0}\right) \frac{\alpha \mathbf{v}}{\gamma^2}. \quad (39)$$

Here  $\gamma$ ,  $\mathbf{v}$  and  $\alpha$  are calculated by Eqs. (21), (22) and (26), respectively. The relativistic transformations (34) for the physical thermodynamic quantities (33) and the relativistic transformations (39) for the unphysical thermodynamic quantities (38) of class II are forbidden because they correspond to the fundamental thermodynamic potential represented by the Lagrange function in which the 3-momentum is an independent state variable in contradiction with the requirements of the Lagrangian mechanics. Moreover, the relativistic transformations (39) are *unphysical* because  $V_0$  is a dependent variable and  $E_0, T_0, p_0$  and  $\mu_0$  depend on  $\mathbf{P}$  (see Eq. (19)) in contradiction with the requirements of the special theory of relativity. We have the invariance for the ratio  $\tilde{p}_{II} V / \tilde{T}_{II} = p_0 V_0 / T_0$ . Comparing Eq. (39) with Eq. (27), we obtain the relation between the unphysical thermodynamic quantities of classes I and II as

$$\tilde{T}_{II} = \theta \tilde{T}_I, \quad \tilde{p}_{II} = \theta \tilde{p}_I, \quad \tilde{\mu}_{II} = \theta \tilde{\mu}_I. \quad (40)$$

Substituting Eq. (25) into Eq. (31) and using the relations  $dS_0 = dS$  and  $dN_0 = dN$ , we exactly obtain Eq. (37) and the relativistic transformations (39).

*Incomplete formula for the differential of volume.* Substituting Eq. (28) into Eq. (31) and using the relations  $dS_0 = dS$  and  $dN_0 = dN$ , we obtain

$$dL = -\frac{\theta T_0}{\gamma} dS + \theta p_0 dV - \frac{\theta \mu_0}{\gamma} dN + \frac{\mathbf{v}}{\gamma^2} d\mathbf{P}. \quad (41)$$

Comparing Eq. (41) with Eq. (37), we obtain (cf. with Eqs. (34) and (39))

$$\tilde{T}_{II} = \frac{\theta T_0}{\gamma}, \quad \tilde{p}_{II} = \theta p_0, \quad \tilde{\mu}_{II} = \frac{\theta \mu_0}{\gamma}, \quad \tilde{\mathbf{v}}_{II} = \frac{\mathbf{v}}{\gamma^2}. \quad (42)$$

Thus, the relativistic transformations (42) for the thermodynamic quantities (38) of class II in the variables  $(S, V, N, \mathbf{P})$  derived from the incomplete formula for the differential of the volume (28) are valid only in the trivial case for  $T_0 = p_0 = \mu_0 = 0$  and  $\mathbf{v} = 0$ .

## 4. Velocity as an independent state variable

### 4.1. Class III: Lagrangian $L(\mathbf{v})$ as the fundamental thermodynamic potential

Let the 3-velocity  $\mathbf{v}$  be an independent state variable in the Lagrange function  $L$  in conformity with the requirements of the Lagrangian mechanics.

#### 4.1.1. Independent physical state variables ( $S_0, V_0, N_0, \mathbf{v}$ )

Let the rest volume  $V_0$  be an independent state variable. The rest energy of the thermodynamic system  $\mathcal{A}$  is a function of the physical state variables ( $S_0, V_0, N_0$ ), i.e.  $E_0 = E_0(S_0, V_0, N_0)$ . The Lorentz factor  $\gamma(\mathbf{v})$  is calculated by Eq. (4). Substituting Eq. (4) into Eq. (7), we obtain the Lagrange function as

$$L(S_0, V_0, N_0, \mathbf{v}) = -\frac{E_0}{\gamma} = -E_0(S_0, V_0, N_0) \sqrt{1 - \frac{\mathbf{v}^2}{c^2}}. \quad (43)$$

Substituting Eqs. (4), (5) into Eq. (6) and Eq. (4) into Eq. (8), we have the 3-momentum and the volume of the system as

$$\mathbf{P}(S_0, V_0, N_0, \mathbf{v}) = \frac{\mathbf{v}}{c^2} E = \frac{\mathbf{v}}{c^2} \frac{E_0(S_0, V_0, N_0)}{\sqrt{1 - \frac{\mathbf{v}^2}{c^2}}}, \quad (44)$$

$$V(V_0, \mathbf{v}) = \frac{V_0}{\gamma} = V_0 \sqrt{1 - \frac{\mathbf{v}^2}{c^2}}. \quad (45)$$

Let the relativistic Lagrangian (43) be a fundamental thermodynamic potential. This function depends on the physical variables ( $S_0, V_0, N_0, \mathbf{v}$ ). Thus, the variables ( $S_0, V_0, N_0, \mathbf{v}$ ) are independent physical state variables of the thermodynamic system. This is dictated by the fundamental thermodynamic potential  $E_0(S_0, V_0, N_0)$  in the rest frame  $K_0$  and the Lagrange function (7) considering velocity  $\mathbf{v}$  as an independent state variable. The rest internal energy  $E_0$  is independent of  $\mathbf{v}$  in accordance with the requirements of the special theory of relativity. Therefore, the variables ( $S_0, V_0, N_0, \mathbf{v}$ ) and the thermodynamic quantities corresponding to them can be respectively considered as *independent physical state variables* and *physical thermodynamic quantities*.

Taking the differential of the Lagrange function (43) with respect to the physical state variables ( $S_0, V_0, N_0, \mathbf{v}$ ) and using Eqs. (2), (4) and (5), we obtain

$$\begin{aligned} dL(S_0, V_0, N_0, \mathbf{v}) &= -\frac{T_0}{\gamma} dS_0 + \frac{p_0}{\gamma} dV_0 - \frac{\mu_0}{\gamma} dN_0 + \mathbf{P} d\mathbf{v} \\ &= -T_{III} dS_0 + p_{III} dV_0 - \mu_{III} dN_0 + \mathbf{P}_{III} d\mathbf{v}, \end{aligned} \quad (46)$$

where

$$\begin{aligned} \left( \frac{\partial L}{\partial S_0} \right)_{V_0, N_0, \mathbf{v}} &\equiv -T_{III}, & \left( \frac{\partial L}{\partial V_0} \right)_{S_0, N_0, \mathbf{v}} &\equiv p_{III}, \\ \left( \frac{\partial L}{\partial N_0} \right)_{S_0, V_0, \mathbf{v}} &\equiv -\mu_{III}, & \left( \frac{\partial L}{\partial \mathbf{v}} \right)_{S_0, V_0, N_0} &\equiv \mathbf{P}_{III} = \mathbf{P}. \end{aligned} \quad (47)$$

Here  $T_{III}$ ,  $p_{III}$  and  $\mu_{III}$  are respectively the physical temperature, physical pressure and physical chemical potential of class III for the thermodynamic system  $\mathcal{A}$  in the reference frame  $K$ ;  $\gamma$  and  $\mathbf{P}$  are calculated by Eqs. (4) and (44).

Comparing Eq. (46) with Eq. (47), we obtain the relativistic transformations for physical thermodynamic quantities of class III as

$$T_{III} = \frac{T_0}{\gamma}, \quad p_{III} = \frac{p_0}{\gamma}, \quad \mu_{III} = \frac{\mu_0}{\gamma}, \quad \mathbf{P}_{III} = \mathbf{P}. \quad (48)$$

These are exactly the Planck transformations [1, 2, 4, 5, 6, 27, 35] for thermodynamic quantities with the exception of pressure. The transformation for the heat transfer can be written as  $\delta Q_{III} = \delta Q_0/\gamma$ . We have the invariance for the ratio  $p_{III}/T_{III} = p_0/T_0$ . Note that the dual thermodynamic quantities  $T_{III}$ ,  $p_{III}$  and  $\mu_{III}$  correspond to the dual fundamental thermodynamic potential  $\tilde{L}$  of the dual fundamental ensemble. Substituting Eq. (11) into the formula for the temperature from Eq. (48), we exactly obtain the temperature (16):  $T_I(S_0, V_0, N_0, \mathbf{P}) = T_{III}(S_0, V_0, N_0, \mathbf{v}(S_0, V_0, N_0, \mathbf{P}))$ . Similar relations are valid also for pressure and chemical potential. Thus, the relativistic transformations (48) and (16) for physical thermodynamic quantities of classes I and III are equivalent. This demonstrates that the Lagrangian and Hamiltonian mechanics give the same results for the relativistic transformations of physical thermodynamic quantities. Note that the formula for the relativistic transformation of physical pressure given in Eq. (48) was obtained for the first time.

#### 4.1.2. Independent unphysical state variables ( $S, V, N, \mathbf{v}$ )

Let the rest physical volume  $V_0$  be a dependent variable and the contracted volume  $V$  in the reference frame  $K$  be an independent state variable. Expressing  $V_0$  from Eq. (45), we get

$$V_0 = V_0(V, \mathbf{v}) = \frac{V}{\sqrt{1 - \frac{\mathbf{v}^2}{c^2}}}. \quad (49)$$

Then, the fundamental thermodynamic potential  $E_0(S_0, V_0, N_0)$  in the reference frame  $K_0$  can be rewritten as

$$E_0(S, V, N, \mathbf{v}) = E_0(S, V_0(V, \mathbf{v}), N). \quad (50)$$

Substituting Eq. (50) into Eq. (43), we obtain (cf. with Eqs. (20) and (36))

$$L(S, V, N, \mathbf{v}) = -E_0(S, V_0(V, \mathbf{v}), N) \sqrt{1 - \frac{\mathbf{v}^2}{c^2}}. \quad (51)$$

Substituting Eq. (50) into Eq. (44), we have the 3-momentum of the system as

$$\mathbf{P}(S, V, N, \mathbf{v}) = \frac{\mathbf{v}}{c^2} \frac{E_0(S, V_0(V, \mathbf{v}), N)}{\sqrt{1 - \frac{\mathbf{v}^2}{c^2}}}. \quad (52)$$

Let the relativistic Lagrangian (51) be a fundamental thermodynamic potential as a function of the independent state variables ( $S, V, N, \mathbf{v}$ ). In this case, the variables ( $S, V, N, \mathbf{v}$ ) and the thermodynamic quantities corresponding to them are *unphysical* because the fundamental thermodynamic potential  $E_0(S, V, N, \mathbf{v})$  in the inertial reference frame  $K_0$ , where  $\mathcal{A}$  is at rest, depends on the velocity  $\mathbf{v}$  of  $\mathcal{A}$  in the laboratory reference frame  $K$  in contradiction with the requirements of the special theory of relativity (see Eq. (50)).

The differential of  $L$  with respect to the unphysical state variables ( $S, V, N, \mathbf{v}$ ) of the thermodynamic system  $\mathcal{A}$  in the reference frame  $K$  can be written as

$$dL(S, V, N, \mathbf{v}) = -\tilde{T}_{III} dS + \tilde{p}_{III} dV - \tilde{\mu}_{III} dN + \tilde{\mathbf{P}}_{III} d\mathbf{v}, \quad (53)$$

where

$$\begin{aligned} \left(\frac{\partial L}{\partial S}\right)_{VN\mathbf{v}} &\equiv -\tilde{T}_{III}, & \left(\frac{\partial L}{\partial V}\right)_{S\mathbf{v}} &\equiv \tilde{p}_{III}, \\ \left(\frac{\partial L}{\partial N}\right)_{SV\mathbf{v}} &\equiv -\tilde{\mu}_{III}, & \left(\frac{\partial L}{\partial \mathbf{v}}\right)_{SVN} &\equiv \tilde{\mathbf{P}}_{III}. \end{aligned} \quad (54)$$

Here  $\tilde{T}_{III}$ ,  $\tilde{p}_{III}$ ,  $\tilde{\mu}_{III}$  and  $\tilde{\mathbf{P}}_{III}$  are respectively the unphysical temperature, unphysical pressure, unphysical chemical potential and generalized momentum of class III for the thermodynamic system  $\mathcal{A}$  in the reference frame  $K$ .

Taking the differential of the volume (49) with respect to the variables  $(V, \mathbf{v})$  and using the relations of independence of these state variables  $\partial V/\partial \mathbf{v} = 0$ , we obtain

$$dV_0(V, \mathbf{v}) = \gamma \left( dV + \frac{V_0}{E_0} \mathbf{P} d\mathbf{v} \right). \quad (55)$$

Substituting Eq. (51) into Eq. (54) and using the partial derivatives from Eq. (55), we obtain the relativistic transformation for the unphysical thermodynamic quantities (54) as (cf. with Eqs. (48) and (27))

$$\tilde{T}_{III} = \frac{T_0}{\gamma}, \quad \tilde{p}_{III} = p_0, \quad \tilde{\mu}_{III} = \frac{\mu_0}{\gamma}, \quad \tilde{\mathbf{P}}_{III} = \left( 1 + \frac{V_0 p_0}{E_0} \right) \mathbf{P}, \quad (56)$$

where  $\gamma$  and  $\mathbf{P}$  are calculated by Eqs. (4) and (52), respectively. These are exactly the Planck transformations [1, 2, 4, 5, 6, 27, 35] for the temperature, pressure and chemical potential. The relativistic transformations (48) for the physical thermodynamic quantities (47) and the Planck transformations (56) for the unphysical thermodynamic quantities (54) of class III are consistent because they correspond to the fundamental thermodynamic potential represented by the Lagrange function in which the 3-velocity is an independent state variable according to the requirements of the Lagrangian mechanics. However, the Planck transformations (56) are *unphysical* because  $V_0$  is a dependent variable and  $E_0, T_0, p_0$  and  $\mu_0$  are dependent on the velocity  $\mathbf{v}$  (see Eq. (50)) in contradiction with the requirements of the special theory of relativity. The transformation for the unphysical heat transfer can be written as  $\delta \tilde{Q}_{III} = \delta Q_0/\gamma$  [1, 2, 3]. We have also the invariance for the ratio  $\tilde{p}_{III} V/\tilde{T}_{III} = p_0 V_0/T_0$  [3, 17].

Substituting Eq. (55) into Eq. (46) and using the relations  $dS_0 = dS$  and  $dN_0 = dN$ , we exactly obtain Eq. (53) and the relativistic transformations (56).

*Incomplete formula for the differential of volume.* Suppose that the differential of volume (8), (45) with respect to the state variables  $(V_0, \mathbf{v})$  is given by Eq. (28) [2, 19], i.e. the Lorentz factor (4) is constant and does not depend on the velocity,  $\partial \gamma/\partial \mathbf{v} = 0$ . In this case, it is easy to show that Eq. (28) is valid only for  $\mathbf{P} = 0$ . See also Eq. (55). Note that Eq. (28) is not consistent with the formula for the Lorentz volume contraction (45). Substituting Eq. (28) into Eq. (46) and using the relations  $dS_0 = dS$  and  $dN_0 = dN$ , we obtain

$$dL = -\frac{T_0}{\gamma} dS + p_0 dV - \frac{\mu_0}{\gamma} dN + \mathbf{P} d\mathbf{v}. \quad (57)$$

Comparing Eq. (53) with Eq. (57), we exactly obtain the Planck transformation formulae [1, 2, 4, 5, 6, 27, 35] (cf. with Eqs. (29) and (56)):

$$\tilde{T}_{III} = \frac{T_0}{\gamma}, \quad \tilde{p}_{III} = p_0, \quad \tilde{\mu}_{III} = \frac{\mu_0}{\gamma}, \quad \tilde{\mathbf{P}}_{III} = \mathbf{P}. \quad (58)$$

Thus, the Planck transformations (58) in the variables  $(S, V, N, \mathbf{v})$  derived from the incomplete formula for the differential of the volume (28) are valid only for  $\mathbf{P} = 0$ . The transformations (58) are equivalent to the transformations (29).

#### 4.2. Class IV: Hamiltonian $E(\mathbf{v})$ as the fundamental thermodynamic potential

Let the 3-velocity  $\mathbf{v}$  be an independent state variable in the Hamilton function  $E$  in contradiction with the requirements of the Hamiltonian mechanics.

##### 4.2.1. Independent physical state variables $(S_0, V_0, N_0, \mathbf{v})$

Let the rest volume  $V_0$  be an independent state variable. Substituting Eq. (4) into Eq. (5), we obtain the Hamilton function as

$$E(S_0, V_0, N_0, \mathbf{v}) = \gamma E_0 = \frac{E_0(S_0, V_0, N_0)}{\sqrt{1 - \frac{v^2}{c^2}}}. \quad (59)$$

Let the Hamiltonian (59) be a fundamental thermodynamic potential as a function of the independent physical state variables  $(S_0, V_0, N_0, \mathbf{v})$ . The volume  $V$  is represented by Eq. (45).

Taking the differential of the Hamilton function (59) with respect to the physical state variables  $(S_0, V_0, N_0, \mathbf{v})$  and using Eqs. (2), (4) and (5), we obtain

$$\begin{aligned} dE(S_0, V_0, N_0, \mathbf{v}) &= \gamma T_0 dS_0 - \gamma p_0 dV_0 + \gamma \mu_0 dN_0 + \gamma^2 \mathbf{P} d\mathbf{v} \\ &= T_{IV} dS_0 - p_{IV} dV_0 + \mu_{IV} dN_0 + \mathbf{P}_{IV} d\mathbf{v}, \end{aligned} \quad (60)$$

where

$$\begin{aligned} \left(\frac{\partial E}{\partial S_0}\right)_{V_0 N_0 \mathbf{v}} &\equiv T_{IV}, & \left(\frac{\partial E}{\partial V_0}\right)_{S_0 N_0 \mathbf{v}} &\equiv -p_{IV}, \\ \left(\frac{\partial E}{\partial N_0}\right)_{S_0 V_0 \mathbf{v}} &\equiv \mu_{IV}, & \left(\frac{\partial E}{\partial \mathbf{v}}\right)_{S_0 V_0 N_0} &\equiv \mathbf{P}_{IV} = \gamma^2 \mathbf{P}. \end{aligned} \quad (61)$$

Here,  $T_{IV}$ ,  $p_{IV}$  and  $\mu_{IV}$  are respectively the physical temperature, physical pressure and physical chemical potential of class IV for the thermodynamic system  $\mathcal{A}$  in the reference frame  $K$ . Comparing Eq. (60) with Eq. (61), we obtain the relativistic transformation for the physical thermodynamic quantities (61) as

$$T_{IV} = \gamma T_0, \quad p_{IV} = \gamma p_0, \quad \mu_{IV} = \gamma \mu_0, \quad \mathbf{P}_{IV} = \gamma^2 \mathbf{P}, \quad (62)$$

where  $\gamma$  and  $\mathbf{P}$  are calculated by Eqs. (4) and (44). These are exactly the Ott transformations [1, 2, 7, 10, 35] for the thermodynamic quantities with the exception of pressure. We have the invariance for the ratio  $p_{IV}/T_{IV} = p_0/T_0$ . Comparing Eq. (62) with Eq. (48), we obtain the relation between the physical thermodynamic quantities of classes III and IV as

$$T_{IV} = \gamma^2 T_{III}, \quad p_{IV} = \gamma^2 p_{III}, \quad \mu_{IV} = \gamma^2 \mu_{III}. \quad (63)$$

#### 4.2.2. Independent unphysical state variables ( $S, V, N, \mathbf{v}$ )

Let the rest physical volume  $V_0$  be a dependent variable (49) and the contracted volume  $V$  in the reference frame  $K$  be an independent state variable. The Hamiltonian (59) in the unphysical state variables ( $S, V, N, \mathbf{v}$ ) can be rewritten as (cf. with Eqs. (20), (36), (51) and (50))

$$E(S, V, N, \mathbf{v}) = \frac{E_0(S, V_0(V, \mathbf{v}), N)}{\sqrt{1 - \frac{v^2}{c^2}}}. \quad (64)$$

Let the Hamiltonian (64) be a fundamental thermodynamic potential as a function of the independent unphysical state variables ( $S, V, N, \mathbf{v}$ ). The rest energy  $E_0$  is represented by Eq. (50).

The differential of  $E$  with respect to the unphysical state variables ( $S, V, N, \mathbf{v}$ ) of the thermodynamic system  $\mathcal{A}$  in the reference frame  $K$  can be written as

$$dE(S, V, N, \mathbf{v}) = \tilde{T}_{IV}dS - \tilde{p}_{IV}dV + \tilde{\mu}_{IV}dN + \tilde{\mathbf{P}}_{IV}d\mathbf{v}, \quad (65)$$

where

$$\begin{aligned} \left(\frac{\partial E}{\partial S}\right)_{VN\mathbf{v}} &\equiv \tilde{T}_{IV}, & \left(\frac{\partial E}{\partial V}\right)_{SN\mathbf{v}} &\equiv -\tilde{p}_{IV}, \\ \left(\frac{\partial E}{\partial N}\right)_{SV\mathbf{v}} &\equiv \tilde{\mu}_{IV}, & \left(\frac{\partial E}{\partial \mathbf{v}}\right)_{SVN} &\equiv \tilde{\mathbf{P}}_{IV}. \end{aligned} \quad (66)$$

Here  $\tilde{T}_{IV}$ ,  $\tilde{p}_{IV}$ ,  $\tilde{\mu}_{IV}$  and  $\tilde{\mathbf{P}}_{IV}$  are respectively the unphysical temperature, unphysical pressure, unphysical chemical potential and generalized momentum of class IV for the thermodynamic system  $\mathcal{A}$  in the reference frame  $K$ .

Substituting Eq. (64) into Eq. (66) and using the partial derivatives from Eq. (55), we obtain the relativistic transformation for the unphysical thermodynamic quantities (66) as

$$\tilde{T}_{IV} = \gamma T_0, \quad \tilde{p}_{IV} = \gamma^2 p_0, \quad \tilde{\mu}_{IV} = \gamma \mu_0, \quad \tilde{\mathbf{P}}_{IV} = \gamma^2 \left(1 - \frac{V_0 p_0}{E_0}\right) \mathbf{P}, \quad (67)$$

where  $\gamma$  and  $\mathbf{P}$  are calculated by Eqs. (4) and (52), respectively. These are exactly the Ott transformations [1, 2, 7, 10, 35] for the temperature, pressure and chemical potential. The relation for the pressure  $p = \gamma^2 p_0$  was first obtained by Sutcliffe in [10]. The relativistic transformation (62) for the physical quantities (61) and the Ott transformations (67) for the unphysical thermodynamic quantities (66) of class IV are forbidden because they correspond to the fundamental thermodynamic potential represented by the Hamilton function in which the 3-velocity is an independent state variable in contradiction with the requirements of the Hamiltonian mechanics. Moreover, the Ott transformations (67) (the Sutcliffe transformation for pressure) are *unphysical* because  $V_0$  is a dependent variable and  $E_0, T_0, p_0$  and  $\mu_0$  depend on the velocity  $\mathbf{v}$  (see Eq. (50)) in contradiction with the requirements of the special theory of relativity. The transformation for the unphysical heat transfer can be written as  $\delta \tilde{Q}_{IV} = \gamma \delta Q_0$  [1, 2, 3]. We have the invariance for the ratio  $\tilde{p}_{IV} V / \tilde{T}_{IV} = p_0 V_0 / T_0$  [3, 10]. Comparing Eq. (67) with Eq. (56), we obtain the relation between the unphysical thermodynamic quantities of classes III and IV as

$$\tilde{T}_{IV} = \gamma^2 \tilde{T}_{III}, \quad \tilde{p}_{IV} = \gamma^2 \tilde{p}_{III}, \quad \tilde{\mu}_{IV} = \gamma^2 \tilde{\mu}_{III}. \quad (68)$$

Substituting Eq. (55) into Eq. (60) and using the relations  $dS_0 = dS$  and  $dN_0 = dN$ , we exactly obtain Eq. (65) and the relativistic transformations (67).

*Incomplete formula for the differential of volume.* Substituting Eq. (28) into Eq. (60) and using the relations  $dS_0 = dS$  and  $dN_0 = dN$ , we obtain

$$dE = \gamma T_0 dS - \gamma^2 p_0 dV + \gamma \mu_0 dN + \gamma^2 \mathbf{P} d\mathbf{v}. \quad (69)$$

Comparing Eq. (69) with Eq. (65), we exactly obtain the Ott transformations [1, 2, 7, 10, 35] (cf. with Eq. (67)):

$$\tilde{T}_{IV} = \gamma T_0, \quad \tilde{p}_{IV} = \gamma^2 p_0, \quad \tilde{\mu}_{IV} = \gamma \mu_0, \quad \tilde{\mathbf{P}}_{IV} = \gamma^2 \mathbf{P}. \quad (70)$$

Thus, the Ott transformations (70) in the variables ( $S, V, N, \mathbf{v}$ ) derived from the incomplete formula for the differential of the volume (28) are valid only for  $\mathbf{P} = 0$ .

## 5. Applications: The moving system of quark-gluon plasma

Let us consider the ultrarelativistic ideal gas of quarks and gluons with  $N_f = 2 + 1$  flavors in the grand canonical ensemble at  $\mu_0 = 0$  (the Stefan–Boltzmann limit). Then the thermodynamic potential, energy, entropy and pressure of the thermodynamic system  $\mathcal{A}$  at rest in the inertial reference frame  $K_0$  can be written as [37]

$$\Omega_0 = -\frac{\pi^2 g V_0}{90} T_0^4, \quad E_0 = \frac{\pi^2 g V_0}{30} T_0^4 = -3\Omega_0, \quad (71)$$

$$S_0 = \frac{2\pi^2 g V_0}{45} T_0^3, \quad p_0 = \frac{\pi^2 g}{90} T_0^4 = -\frac{\Omega_0}{V_0} = \frac{E_0}{3V_0}, \quad (72)$$

where  $g$  is the effective degeneracy factor defined as [37, 38, 39]

$$\begin{aligned} g &= g_g + \frac{7}{8} g_q, \quad g_g = 2_{spin} \times (N_c^2 - 1), \\ g_q &= 2_{spin} \times 2_{q\bar{q}} \times N_c \times N_f. \end{aligned} \quad (73)$$

Here and throughout this paper we use the system of natural units  $\hbar = c = k_B = 1$ .

In the fundamental ensemble, the fundamental thermodynamic potential, temperature and pressure at  $\mu_0 = 0$  can be rewritten as

$$E_0 = \Omega_0 + T_0 S_0 + \mu_0 N_0 = \frac{3S_0}{4} \left(\frac{45S_0}{2\pi^2 g V_0}\right)^{1/3} = \frac{3S_0 T_0}{4}, \quad (74)$$

$$p_0 = \frac{S_0}{4V_0} \left(\frac{45S_0}{2\pi^2 g V_0}\right)^{1/3} = \frac{S_0 T_0}{4V_0}, \quad T_0 = \left(\frac{45S_0}{2\pi^2 g V_0}\right)^{1/3}. \quad (75)$$

Figure 1 represents the behaviour of the physical temperature and physical pressure in the laboratory reference frame  $K$  as functions of the 3-momentum  $\mathbf{P}$  of the system, which is at rest in the moving reference frame  $K_0$ , for the ultrarelativistic ideal gas of quarks and gluons with  $N_f = 2 + 1$  flavors in the volume  $V_0 = 1 \text{ fm}^3$  at  $S_0 = 189.995$  and  $\mu_0 = 0$ . In the reference frame  $K_0$ , the energy of the system  $E_0 = 56.998 \text{ GeV}$ , the temperature  $T_0 = 400 \text{ MeV}$  and the pressure  $p_0 = 18.999$






Figure 1: (Color online) The physical temperature  $T$  and physical pressure  $p$  in the laboratory reference frame  $K$  as functions of the 3-momentum  $\mathbf{P}$  of the system, which is at rest in the moving reference frame  $K_0$ , for the ultrarelativistic ideal gas of quarks and gluons with  $N_f = 2 + 1$  flavors in the volume  $V_0 = 1 \text{ fm}^3$  at  $S_0 = 189.995$  and  $\mu_0 = 0$ . The numbers  $I, II, III$  and  $IV$  denote four different classes of physical thermodynamic quantities defined in Eqs. (16), (34), (48) and (62), respectively.




Figure 2: (Color online) The equation of state for the ultrarelativistic ideal gas of quarks and gluons with  $N_f = 2 + 1$  flavors at  $S_0 = 189.995$ ,  $P_x = 100 \text{ GeV}$  and  $\mu_0 = 0$ . The symbol  $p_0$  denotes the rest pressure. For further details see the caption of Figure 1.

$\text{GeV}/\text{fm}^3$ . The curves  $I, II, III$  and  $IV$  are calculations for the ideal gas by Eqs. (16), (34), (48) and (62), respectively, using Eqs. (4), (9), (11) and (32). The Planck physical temperatures  $T_I$  and  $T_{III}$  are the same. They decrease with  $\mathbf{P}$  and tend to zero as  $\mathbf{P} \rightarrow \infty$ . The behaviour of the physical temperature  $T_{II}$  is

similar to the behavior of the temperatures  $T_I$  and  $T_{III}$ . The Ott temperature  $T_{IV}$  increases with  $\mathbf{P}$  and contrary to other temperatures tends to infinity as  $\mathbf{P} \rightarrow \infty$ . The behavior of the physical pressures  $p_I, p_{II}, p_{III}$  and  $p_{IV}$  is similar to the behavior of the physical temperatures  $T_I, T_{II}, T_{III}$  and  $T_{IV}$ .

Figure 2 represents the behaviour of the physical pressure in the laboratory reference frame  $K$  as a function of the rest volume  $V_0$  of the system, which is at rest in the moving reference frame  $K_0$ , for the ultrarelativistic ideal gas of quarks and gluons with  $N_f = 2 + 1$  flavors at entropy  $S_0 = 189.995$ , momentum  $P_x = 100 \text{ GeV}$  and chemical potential  $\mu_0 = 0$ . The curves  $I, II, III$  and  $IV$  are calculations for the ideal gas by Eqs. (16), (34), (48) and (62), respectively, using Eqs. (4), (9), (11) and (32). The curve  $p_0$  corresponds to the rest pressure (75). All the physical pressures  $p_I, p_{II}, p_{III}$  and  $p_{IV}$  decrease rapidly with  $V_0$ . The physical pressures  $p_I, p_{II}$  and  $p_{III}$  are less than the rest pressure  $p_0$ . However, the physical pressure  $p_{IV}$  is greater than the rest pressure  $p_0$ . In Figures 1 and 2, the Hamiltonian and Lagrangian mechanics allow only curves  $I$  and  $III$ , which are the same, and  $p_0$ .

## 6. Conclusions

In this work, we have derived eight different relativistic transformations for eight sets of thermodynamic quantities of the relativistically moving thermodynamic system. We divided these transformations into four classes considering the fundamental thermodynamic potential to be either the Hamilton or Lagrange function and the independent state variable to be either the 3-momentum or 3-velocity of the system. Each class is divided into two subclasses. The subclass of physical thermodynamic quantities is defined by the rest volume of the thermodynamic system as an independent state variable, and the subclass of unphysical thermodynamic quantities is defined by the contracted volume of the thermodynamic system in the laboratory reference frame as an independent state variable. The relativistic transformations for the thermodynamic quantities of classes  $I$  and  $III$  are consistent and allowed because they correspond to the fundamental thermodynamic potential represented either by the Hamilton function in which the 3-momentum is an independent state variable or by the Lagrange function in which the 3-velocity is an independent state variable according to the requirements of the Hamiltonian and Lagrangian mechanics. The relativistic transformations of classes  $II$  and  $IV$  are forbidden because they correspond to the fundamental thermodynamic potential represented either by the Lagrange function in which the 3-momentum is an independent state variable or by the Hamilton function in which the 3-velocity is an independent state variable in contradiction with the requirements of the Lagrangian and Hamiltonian mechanics. The four subclasses of the relativistic transformations for the unphysical thermodynamic quantities corresponding to the set of independent unphysical state variables are not consistent because the contracted volume is an independent state variable and the rest energy and other thermodynamic quantities of the thermodynamic system at rest depend on momentum and velocity of the system in the laboratory reference frame in disagreement with

the principles of the special theory of relativity and causality principle. Only four subclasses of relativistic transformations for physical thermodynamic quantities corresponding to the set of independent physical state variables are consistent because the rest volume is an independent state variable and the rest energy of the thermodynamic system at rest is independent of momentum and velocity of the system in the laboratory reference frame in agreement with the principles of the special theory of relativity.

We found that the relativistic transformations for the unphysical thermodynamic quantities of classes III and IV coincide with the Planck and Ott transformations for temperature, pressure and chemical potential. Thus, the Planck and Ott transformations corresponding to the set of independent unphysical state variables are unphysical. Moreover, these Ott transformations are forbidden by the Hamiltonian mechanics. We obtained that the relativistic transformations for the physical thermodynamic quantities of class IV coincide with the Ott transformations for temperature and chemical potential. These Ott transformations corresponding to the physical state variables are consistent but they are forbidden by the Hamiltonian mechanics. We found that the relativistic transformations for the physical thermodynamic quantities of classes I and III corresponding to the Hamiltonian and Lagrangian mechanics are equivalent and coincide with the Planck transformations for temperature and chemical potential. Thus, only these two equivalent transformations from eight ones are consistent and allowed by the Lagrangian and Hamiltonian mechanics. Other six transformations are either inconsistent or forbidden. The equivalence of the results obtained by the Lagrangian and Hamiltonian mechanics for the relativistic transformations of the physical thermodynamic quantities of classes I and III demonstrates that only these two relativistic transformations are valid. For the unphysical thermodynamic quantities of classes I and III, the Lagrangian and Hamiltonian mechanics provide the nonequivalent relativistic transformations. In this paper, the consistent formulae for the relativistic transformations of the physical pressures of classes I and III, which are equivalent, were obtained. It was shown that the formulae of Planck's and Sutcliffe's transformations of pressure are inconsistent.

We compared the relativistic transformations for physical thermodynamic quantities of classes I, II, III, and IV using them to describe the temperature and pressure of the relativistically moving system of the QGP.

We can apply the results obtained in this study to construct consistent relativistic statistical mechanics of a moving thermodynamic system. This statistical mechanics can help formulate local equilibrium statistical models for hadrons and partons to describe particle yields produced in heavy-ion and proton-proton collisions at high energies.

## Acknowledgments

This work was supported in part by the RSCF grant, N22-72-10028 and the Romanian Ministry of Research, Innovation and Digitalization, Project PN 23 21 01 01/2023.

## References

- [1] D. ter Haar, H. Wergeland, *Phys. Rep.* 1C (1971) 31.
- [2] C. Farias, V.A. Pinto, P.S. Moya, *Sci. Rep.* 7 (2017) 17657.
- [3] T.K. Nakamura, *Prog. Theor. Phys.* 128 (2012) 463.
- [4] K. von Mosengeil, *Annalen der Physik* 327, 5 (1907) 867-904.
- [5] M. Planck, *Annalen der Physik* 331, 6 (1908) 1-34; *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften*, June 13, 1907, pp. 542-570.
- [6] A. Einstein, *Jahrbuch der Radioaktivität und Elektronik* 4 (1907) 411-462.
- [7] H. Ott, *Z. Physik* 175 (1963) 70-104.
- [8] H. Arzeliès, *Nuovo Cimento B* 40 (1965) 333.
- [9] T.W.B. Kibble, *Nuovo Cimento B* 41 (1966) 72.
- [10] W.G. Sutcliffe, *Nuovo Cimento* 39 (1965) 683.
- [11] P.T. Landsberg, *Nature* 212 (1966) 571.
- [12] G. Cavalleri, G. Salgarelli, *Nuovo Cimento A* 62 (1969) 722.
- [13] R.G. Newburgh, *Nuovo Cimento B* 52 (1979) 219.
- [14] C.K. Yuen, *Am. J. Phys.* 38 (1970) 246.
- [15] H. Callen, G. Horwitz, *Am. J. Phys.* 39 (1971) 938.
- [16] H.-J. Treder, *Annalen der Physik* 489, 1 (1977) 23-29.
- [17] R.K. Pathria, *Proc. Phys. Soc.* 88 (1966) 791.
- [18] Ch. Fenech, J.P. Vigièr, *Phys. Lett. A* 215 (1996) 247.
- [19] N.G. van Kampen, *Phys. Rev.* 173 (1968) 295.
- [20] P.T. Landsberg, G.E.A. Matsas, *Phys. Lett. A* 223 (1996) 401.
- [21] J.J. Mareš, P. Hubík, J. Šesták, V. Špička, J. Křištofík, J. Stávek, *Physica E* 42 (2010) 484.
- [22] N. Papadatos, C. Anastopoulos, *Phys. Rev. D* 102 (2020) 085005.
- [23] G. Ares de Parga, B. López-Carrera, F. Angulo-Brown, *J. Phys. A: Math. Gen.* 38 (2005) 2821.
- [24] J.A. Heras, M.G. Osorno, *Eur. Phys. J. Plus* 137 (2022) 423.
- [25] J.J. Mareš, P. Hubík, V. Špička, *Fortschr. Phys.* 65 (2017) 1700018.
- [26] D. Mi, H.Y. Zhong, D.M. Tong, *Mod. Phys. Lett. A* 24 (2009) 73.
- [27] N. Agmon, *Foundations of Physics* 7 (1977) 331.
- [28] R. Balescu, *Physica* 40 (1968) 309.
- [29] J. Güémez, J.A. Mier, *Phys. Scr.* 98 (2023) 025001.
- [30] L. Gavassino, *Foundations of Physics* 50 (2020) 1554.
- [31] M. Przanowski, J. Tosiek, *Phys. Scr.* 84 (2011) 055008.
- [32] A. Montakhab, M. Ghodrati, M. Barati, *Phys. Rev. E* 79 (2009) 031124.
- [33] T.S. Biró, P. Ván, *Europhys. Lett.* 89 (2010) 30001.
- [34] X. Hao, S. Liu, L. Zhao, *Commun. Theor. Phys.* 75 (2023) 035601.
- [35] A.S. Parvan, *Annals of Physics* 401 (2019) 130.
- [36] L.D. Landau, E.M. Lifshitz, *The Classical Theory of Fields*, Vol. 2, third ed., Pergamon Press, 1971.
- [37] Kohsuke Yagi, Tetsuo Hatsuda, and Yasuo Miake, *Quark-Gluon plasma: from big bang to little bang*, Cambridge University Press, 2008.
- [38] G.E. Brown, H.A. Bethe, P.M. Pizzochero, *Phys. Lett. B* 263 (1991) 337.
- [39] A.S. Parvan, *Eur. Phys. J. A* 56 (2020) 192.