QUANTUM K-INVARIANTS VIA QUOT SCHEMES I

SHUBHAM SINHA AND MING ZHANG

ABSTRACT. We study the virtual Euler characteristics of sheaves over Quot schemes of curves, establishing that these invariants fit into a topological quantum field theory (TQFT) valued in $\mathbb{Z}[[q]]$. Utilizing Quot scheme compactifications alongside the TQFT framework, we derive presentations of the small quantum K-ring of the Grassmannian. Our approach offers a new method for finding explicit formulas for quantum K-invariants.

1. INTRODUCTION

1.1. **Overview.** Quantum K-theory was introduced by Givental [Giv] and Lee [Lee] as a K-theoretic generalization of Gromov–Witten theory. Let X be a projective manifold. Quantum K-invariants (or sometimes referred to as the K-theoretic Gromov–Witten invariants) are defined as Euler characteristics of K-theory classes on the Kontsevich's moduli spaces of stable maps to X. Using 2-pointed and 3-pointed, genus zero quantum K-invariants of X, one can define an associative product, called the quantum product, on the K-group of X. The resulting algebra is called the (small) quantum K-ring.

Let $\operatorname{Gr}(r, N)$ be the Grassmann variety of *r*-planes in \mathbb{C}^N . The small quantum *K*-ring of Grassmannians was first studied by Buch and Mihalcea in [BM]. Buch–Mihalcea's results rely on the analysis of the geometry of the Kontsevich moduli space $\overline{M}_{0,3}(\operatorname{Gr}(r, N), d)$, which compactifies the moduli space $\operatorname{Mor}_d(\mathbb{P}^1, \operatorname{Gr}(r, N))$ of degree-*d* morphisms from \mathbb{P}^1 to $\operatorname{Gr}(r, N)$. Note that the moduli space $\operatorname{Mor}_d(\mathbb{P}^1, \operatorname{Gr}(r, N))$ has a simpler compactification, the Grothendieck's Quot scheme [Gro]. In the early study of quantum cohomology of Grassmannians, many results have been obtained by using the Quot scheme compactification [Ber1, Ber2, BDW].

In this paper, we show that virtual Euler characteristics over Quot schemes of any genus g curve form a $\mathbb{Z}[[q]]$ -valued (1 + 1)-Topological Quantum Field Theory (TQFT). Additionally, we establish that the quantum K-ring QK(Gr(r, N)) can be fully reconstructed using the Quot schemes of \mathbb{P}^1 . In particular, our approach provides a new way to derive a ring presentation for QK(Gr(r, N)). We observe that fundamental properties of QK(Gr(r, N)), such as finiteness and S_3 -symmetry of the structure constants, are geometrically elucidated using this approach.

In part II [SZ], we establish explicit formulas for the Euler characteristics over Quot schemes of curves, similar to the Vafa–Intriligator formula for cohomological invariants. For the genus zero case, we employ torus localization for calculation, inspired by [OS] in K-theory (see also [MO1] for cohomology calculations). For higher-genus cases, we utilize the TQFT structure developed here. In this paper, some results critically depend on the vanishing results in Theorem 1.11 from [SZ]. 1.2. Quantum K-ring of the Grassmannian. Throughout this paper, we set X = Gr(r, N)and k = N - r. Let $P_{r,k}$ be the set of partitions contained in the $r \times k$ rectangular partition. The complement partition for $\lambda = (\lambda_1, \ldots, \lambda_r) \in P_{r,k}$ is given by $\lambda^* = (k - \lambda_r, \ldots, k - \lambda_1)$.

Let K(X) be the Grothendieck group of coherent sheaves (or, equivalently, vector bundles since X is smooth). For any partition $\lambda \in P_{r,k}$, let X_{λ} denote the Schubert cycle for a fixed complete flag of \mathbb{C}^N (see Section 2 for precise definitions). The K-theory classes of the structure sheaves of these Schubert cycles $\mathcal{O}_{\lambda} := [\mathcal{O}_{X_{\lambda}}]$ form a Z-linear basis for K(X). The structure constants $c_{\lambda,\mu}^{\nu}$ for the product

$$\mathcal{O}_{\lambda} \cdot \mathcal{O}_{\mu} = \sum_{\nu \in \mathcal{P}_{r,k}} c_{\lambda,\mu}^{\nu} \mathcal{O}_{\nu}$$

in K(X) are given by the K-theoretic Littlewood-Richardson rule in [Buc], which involves counting set-valued tableaux.

The K-group of Grassmannian admits another natural basis consisting of Schur bundles. Let $S \subset \mathcal{O}_X \otimes \mathbb{C}^N$ be the tautological subbundle over X. For any partition λ , we have a corresponding vector bundle $\mathbb{S}^{\lambda}(S)$, where \mathbb{S}^{λ} is the Schur functor associated to λ . For example, $\mathbb{S}^{(m)}(S) = \operatorname{Sym}^m S$ and $\mathbb{S}^{(1)^m}(S) = \wedge^m S$ where $(1)^m = (1, \ldots, 1)$. Let $A = (A^{\nu}_{\lambda})_{\lambda,\nu\in P_{r,k}}$ be the transition matrix between the Schur basis $\{\mathbb{S}^{\lambda}(S) : \lambda \in P_{r,k}\}$ and the Schubert basis $\{\mathcal{O}_{\nu} : \nu \in P_{r,k}\}$:

(1)
$$\mathcal{O}_{\lambda} = \sum_{\nu \in \mathbf{P}_{r,k}} \mathbf{A}_{\lambda}^{\nu} \, \mathbb{S}^{\nu}(S)$$

for any $\lambda \in \mathbf{P}_{r,k}$.

Next we recall the definition of Givental–Lee's quantum K-invariants over X. Let $\overline{M}_{0,n}(X,d)$ be the moduli stack of degree-d, genus zero, n-pointed stable maps into X. Let $\operatorname{ev}_i : \overline{M}_{0,n}(X,d) \to X$ denote the evaluation map at the *i*-th marking for $1 \leq i \leq n$. Then the quantum K-invariants are defined by

(2)
$$\langle \phi_1, \dots, \phi_n \rangle_{0,n,d} := \chi \left(\overline{M}_{0,n}(X,d), \prod_{i=1}^n \operatorname{ev}_i^*(\phi_i) \right),$$

where ϕ_1, \ldots, ϕ_n are K-theory classes on X. For partitions $\alpha, \beta, \gamma \in P_{r,k}$, we define

(3)
$$F_{\alpha,\beta} := \sum_{d \ge 0} q^d \langle \mathcal{O}_\alpha, \mathcal{O}_\beta \rangle_{0,2,d} \quad \text{and} \quad F_{\alpha,\beta,\gamma} := \sum_{d \ge 0} q^d \langle \mathcal{O}_\alpha, \mathcal{O}_\beta, \mathcal{O}_\gamma \rangle_{0,3,d},$$

where q is the Novikov variable. Here, $(F_{\alpha,\beta})_{\alpha,\beta\in P_{r,k}}$ is referred to as the quantized pairing matrix. Let $F^{\alpha,\beta}$ denote the entries of its inverse. The quantum K-ring of X, denoted by QK(X), is the $\mathbb{Z}[[q]]$ -module $K(X) \otimes_{\mathbb{Z}} \mathbb{Z}[[q]]$ equipped with the quantum K-product defined by

$$\mathcal{O}_{\lambda} \bullet \mathcal{O}_{\mu} = \sum_{\nu \in \mathcal{P}_{r,k}} N^{\nu}_{\lambda,\mu} \mathcal{O}_{\nu},$$

where the structure constants $N^{\nu}_{\lambda,\mu}$ are given by

(4)
$$N_{\lambda,\mu}^{\nu} := \sum_{\alpha \in \mathbf{P}_{r,k}} F^{\nu,\alpha} F_{\alpha,\lambda,\mu}$$

According to the general results in [Giv, Lee], the quantum K-product is commutative, associative, and a deformation of the tensor product. Furthermore, the structure constants $N^{\nu}_{\lambda,\mu}$ are polynomials in q (see [BM]), a property commonly known as the finiteness of the quantum K-ring.

1.3. Quot schemes. Fix a smooth projective curve C of genus g. Let $Quot_d(C, N, r)$ be the Quot scheme parametrizing short exact sequences

$$0 \to S \to \mathcal{O}_C^{\oplus N} \to Q \to 0$$

of coherent sheaves on C, where S has rank r and degree -d. When $C = \mathbb{P}^1$, the Quot scheme $\operatorname{Quot}_d(\mathbb{P}^1, N, r)$, is a smooth projective scheme and isomorphic to X when d = 0. Strømme [Str] studied geometric aspects of $\operatorname{Quot}_d(\mathbb{P}^1, N, r)$ using torus localization. In higher genus, $\operatorname{Quot}_d(C, N, r)$ is used to compactify the space $Mor_d(C, X)$ of degree-d morphisms from C to X and to calculate intersection numbers (see [BDW, Ber2, Ber1])). In [MO1], Marian and Oprea defined virtual fundamental classes for Quot schemes on curves of arbitrary genus and calculated virtual intersection numbers.

Let π_C and π be the projections from $Quot_d(C, N, r) \times C$ to C and $Quot_d(C, N, r)$, respectively. There exists a universal short exact sequence

$$0 \to \mathcal{S} \to \pi_C^* \mathcal{O}_C^{\oplus N} \to \mathcal{Q} \to 0.$$

of coherent sheaves over $\operatorname{Quot}_d(C, N, r) \times C$. For any $p \in C$, we denote by \mathcal{S}_p the restriction of the universal subbundle \mathcal{S} to $\operatorname{Quot}_d(C, N, r) \times \{p\}$.

Definition 1.1. For any partition $\lambda \in P_{r,k}$, we define the K-theory class

$$\widetilde{\mathcal{O}}_{\lambda} = \sum_{\nu \in \mathcal{P}_{r,k}} \mathcal{A}_{\lambda}^{\nu} \, \mathbb{S}^{\nu}(\mathcal{S}_p)$$

on $\operatorname{Quot}_d(C, N, r)$, where A^{ν}_{λ} are the coefficients of the transition matrix in (1) and $\mathbb{S}^{\nu}(\mathcal{S}_p)$ are the Schur functors applied to the vector bundle \mathcal{S}_p . Note that the point p is suppressed from the notation of $\widetilde{\mathcal{O}}_{\lambda}$. We define

$$\widetilde{\mathcal{O}}_{\nu}^* := \widetilde{\mathcal{O}}_{\nu^*} \cdot \det(\mathcal{S}_p),$$

where ν^* is the complement partition in the $r \times k$ rectangle.

Theorem 1.2. For any partitions $\lambda, \mu, \nu \in P_{r,k}$, the structure constants for the product in the quantum K-ring QK(Gr(r, N)) satisfy:

$$N_{\lambda,\mu}^{\nu} = \sum_{d \ge 0} q^d \chi(\operatorname{\mathsf{Quot}}_d(\mathbb{P}^1, N, r), \widetilde{\mathcal{O}}_{\lambda} \cdot \widetilde{\mathcal{O}}_{\mu} \cdot \widetilde{\mathcal{O}}_{\nu}^*).$$

The above theorem provides a more insightful description of the structure constants. For example, Theorem 1.2 immediately shows the S_3 -symmetry $N_{\lambda,\mu}^{\nu} = N_{\lambda,\nu^*}^{\mu^*}$. This symmetry was established for q = 0 (ordinary K-theory) in [Vak] and generally in [BM, Theorem 5.12], which is not evident from the original definition (4). Furthermore, the vanishing results of the Euler characteristics of Schur bundles $\mathbb{S}^{\lambda}(\mathcal{S}_p)$ over high-degree Quot schemes imply that $N_{\lambda\mu}^{\nu}$ is a polynomial in q of degree at most min $\{r, N - r\}$ (see Corollary 3.7).

1.4. Topological Quantum Field Theory. The Quot scheme $\operatorname{Quot}_d(C, N, r)$ may not be smooth when $C \neq \mathbb{P}^1$. A two-term perfect obstruction theory for $\operatorname{Quot}_d(C, N, r)$ was constructed in [MO1], yielding a virtual fundamental class via [LT, BF] and a virtual structure sheaf through [Lee]. When $d \gg 0$, the Quot scheme is an irreducible scheme of the expected dimension (see [BDW, PR]), and the virtual fundamental class and the virtual structure sheaf, denoted $\mathcal{O}_{Quot}^{\operatorname{vir}}$, equal the ordinary fundamental class and structure sheaf \mathcal{O}_{Quot} , respectively.

For any K-theory class ϕ on $\operatorname{Quot}_d(C, N, r)$, we define the virtual Euler characteristics by

$$\chi^{\mathrm{vir}}\big(\operatorname{\mathsf{Quot}}_d(C,N,r),\phi\big) := \chi\big(\operatorname{\mathsf{Quot}}_d(C,N,r),\mathcal{O}_{\operatorname{\mathsf{Quot}}}^{\mathrm{vir}}\cdot\phi\big).$$

Definition 1.3. For any tuples of partitions $\underline{\lambda} = (\lambda^1, \dots, \lambda^s)$ and $\underline{\nu} = (\nu^1, \dots, \nu^t)$ in $P_{r,k}$, we define the power series

$$N(g)_{\underline{\lambda}}^{\underline{\nu}} := \sum_{d \ge 0} q^d \chi^{\mathrm{vir}}(\mathrm{Quot}_d(C, N, r), \widetilde{\mathcal{O}}_{\lambda^1} \cdots \widetilde{\mathcal{O}}_{\lambda^s} \cdot \widetilde{\mathcal{O}}_{\nu^1}^* \cdots \widetilde{\mathcal{O}}_{\nu^t}^*),$$

where $\widetilde{\mathcal{O}}_{\lambda}$ and $\widetilde{\mathcal{O}}_{\lambda}^{*}$ are defined in Definition 1.1. Note that $N_{\lambda,\mu}^{\nu} = N(0)_{\lambda,\mu}^{\nu}$ by Theorem 1.2.

Next, we define a Topological Quantum Field Theory (TQFT) valued in the power series ring $\mathbf{R} = \mathbb{C}[[q]]$, as considered in [BP]. Let **2Cob** be the catergory where objects are disjoint union of circles and morphisms are diffeomorphism classes of cobordisms. The composition of morphisms is given by the concatenation of cobordisms.

Definition 1.4. A (1+1)-TQFT with coefficients in **R** is a symmetric monoidal functor

$F: \mathbf{2Cob} \to \mathbf{Mod}_{\mathbf{R}},$

where $\mathbf{Mod}_{\mathbf{R}}$ is the category of \mathbf{R} -modules, which is equivalent to:

- (i) Let M be an **R**-module associated to a circle, then F associates $M^{\otimes s}$ to the disjoint union of s circles.
- (ii) The trivial cobordism from a circle to a circle is associated to the identity map on M.
- (iii) The concatenation of two cobordisms correspond to the composition in $\mathbf{Mod}_{\mathbf{R}}$.

Consider the free $\mathbb{C}[[q]]$ -module $M = K(X) \otimes \mathbb{C}[[q]]$, with the basis $\{\mathcal{O}_{\lambda} : \lambda \in \mathbb{P}_{r,k}\}$. For a genus g cobordism from s circles to t circles, we define the morphism $M^{\otimes s} \to M^{\otimes t}$ using the matrix given by:

$$\mathcal{O}_{\lambda^1} \otimes \cdots \otimes \mathcal{O}_{\lambda^s} \to \sum_{\underline{\nu}} N(g)^{\underline{\nu}}_{\underline{\lambda}} \cdot \mathcal{O}_{\nu^1} \otimes \cdots \otimes \mathcal{O}_{\nu^t}.$$

Theorem 1.5. The invariants $N(g)_{\underline{\lambda}}^{\underline{\nu}}$ define a (1+1)-TQFT with values in the power series ring $\mathbb{C}[[q]]$. In particular, we prove the orthogonality relation

$$N(0)_{\lambda}^{\nu} = \sum_{d \ge 0} q^d \chi(\operatorname{Quot}_d(\mathbb{P}^1, N, r), \widetilde{\mathcal{O}}_{\lambda} \cdot \widetilde{\mathcal{O}}_{\nu}^*) = \delta_{\lambda, \nu},$$

and the gluing formula

$$\sum_{\mu} N(g_1)^{\underline{\mu}}_{\underline{\lambda}} \cdot N(g_2)^{\underline{\nu}}_{\underline{\mu}} = N(g+t-1)^{\underline{\nu}}_{\underline{\lambda}},$$

where $g = g_1 + g_2$ and the sum runs over all t-tuples, $\mu = (\mu^1, \dots, \mu^t)$, of partitions in $P_{r,k}$.



We shall see later that the above TQFT matches with the quantum-K TQFT for the Grassmannian. The novelty here is that all the invariants are expressed as (virtual) Euler characteristics over Quot schemes.

Remark 1.6. The cohomological Quot scheme invariants also exhibit a TQFT structure with \mathbb{C} -coefficients, called Grassmann/GL-Verlinde TQFT in [MO2], which differs from the standard TQFT coming from the quantum cohomology of the Grassmannian. Their proof relies on the fact that the cohomological invariants are enumerative in the large degree regime, which is not applicable in the K-theoretic setting.

1.5. Evaluation map. To present the next result, we will introduce notation that is basisindependent for defining invariants over the Quot scheme.

The Grassmannian $X = \operatorname{Gr}(r, N)$ has a GIT presentation $X = [M_{r \times N}^{ss}/\operatorname{GL}_r(\mathbb{C})]$, where $M_{r \times N}$ is the affine space of r by N complex matrices and $M_{r \times N}^{ss} \subset M_{r \times N}$ is the open subset of matrices of full rank. Let

$$\mathfrak{X} := [M_{r \times N} / \mathrm{GL}_r(\mathbb{C})]$$

be the quotient stack. Note that the K-theory of the stack $K(\mathfrak{X})$ is isomorphic to the ring $R(\operatorname{GL}_r(\mathbb{C}))$ of rational $\operatorname{GL}_r(\mathbb{C})$ -representations. Given a $\operatorname{GL}_r(\mathbb{C})$ -representation V, we define its associated K-theory classes on \mathfrak{X} as $M_{r\times N} \times_{\operatorname{GL}_r(\mathbb{C})} V \in K(\mathfrak{X})$, also denoted by V using the isomorphism $R(\operatorname{GL}_r(\mathbb{C})) \cong K(\mathfrak{X})$. An element $V \in K(\mathfrak{X})$ restricts to an element $V \in K(X)$ given by $M_{r\times N}^{ss} \times_{\operatorname{GL}_r(\mathbb{C})} V \in K(X)$.

In general, we can define K-theoretic Quot scheme invariants with insertions in $K(\mathfrak{X})$ as follows. We choose a point $p \in C$. Let

$$\widetilde{\operatorname{ev}}_p: \operatorname{\mathsf{Quot}}_d(C, N, r) \to \mathfrak{X}$$

be the 'stacky' evaluation morphism induced by the restriction $\mathcal{S}_p \to \mathcal{O}_p^{\oplus N}$.

Definition 1.7. For $V_1, \ldots, V_n \in K(\mathfrak{X})$, we define the K-theoretic Quot scheme invariant by

$$\langle \mathrm{V}_1, \ldots, \mathrm{V}_n \rangle_{g,d}^{\mathsf{Quot}} := \chi^{\mathrm{vir}} \left(\operatorname{\mathsf{Quot}}_d(C, N, r), \prod_{i=1}^n \widetilde{\mathrm{ev}}_p^*(\mathrm{V}_i) \right).$$

The above invariant is independent of the choices of p and the complex structure of the genus g curve C. We define the corresponding generating series by

$$\langle\!\langle \mathrm{V}_1,\ldots,\mathrm{V}_n
angle\!
angle_g^{\mathsf{Quot}} = \sum_{d\geq 0} q^d \langle \mathrm{V}_1,\ldots,\mathrm{V}_n
angle_{g,d}^{\mathsf{Quot}}$$

Let $S \in K(\mathfrak{X})$ be the element corresponding to the dual of the standard representation $(\mathbb{C}^r)^{\vee}$. Note that S restricts to the universal subbundle S over X = Gr(r, N). For $\lambda \in P_{r,k}$, we define the 'stacky' structure sheaf O_{λ} in $K(\mathfrak{X})$ by substituting \mathcal{S}_p with S in Definition 1.1. In the representation ring $R(GL_r(\mathbb{C})), O_{\lambda}$ is given by $\mathbb{G}^{\lambda}(\mathbb{C}^r)$, which is the Grothendieck functor applied to the standard representation [Len, §8] (see Section 2 for details). Note that the stacky pullback satisfies $\widetilde{ev}_p(S) = \mathcal{S}_p$. With this notation, the invariants in Definition 1.3 equal

$$N(g)_{\underline{\lambda}}^{\underline{\nu}} = \langle\!\langle \mathcal{O}_{\lambda^1}, \dots, \mathcal{O}_{\lambda^s}, \mathcal{O}_{\nu^1}^*, \dots, \mathcal{O}_{\nu^t}^* \rangle\!\rangle_g^{\mathsf{Quot}}$$

where $O_{\nu}^* = O_{\nu^*} \cdot \det(S) \in K(\mathfrak{X})$ is referred to as the quantized K-theoretic dual in this paper.

1.6. Quantum reduction map. Let QK(X) be the quantum K-ring of the Grassmannian X = Gr(r, N). To better understand the quantum K-ring, we consider the following:

Definition 1.8. The quantum reduction map $\kappa : R(\operatorname{GL}_r(\mathbb{C})) \otimes \mathbb{Z}[[q]] \to \operatorname{QK}(X)$ is defined to be the $\mathbb{Z}[[q]]$ -linear extension of the formula

(5)
$$\kappa(\mathbf{V}) = \sum_{\alpha \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\alpha}^* \rangle\!\rangle_0^{\mathsf{Quot}} \mathcal{O}_{\alpha},$$

where $V \in R(GL_r(\mathbb{C})) \cong K(\mathfrak{X})$.

Let $\mathcal{R}^{r,k}$ be the subspace of $R(\operatorname{GL}_r(\mathbb{C}))$ spanned by $\{O_\lambda\}_{\lambda\in P_{r,k}}$. The orthogonality relation in Theorem 1.5 implies that the map κ sends V to its restriction class $V \in K(X)$ for any representation V inside the 'rectangle box' $\mathcal{R}^{r,k}$. When V is outside of $\mathcal{R}^{r,k}$, the map κ 'reduces' it to an element that usually depends on q in the quantum K-ring. For example, Proposition 4.7 states that

$$\kappa(\det \mathbf{S}^{\vee}) = \sum_{d \ge 0} q^d \det S^{\vee} = \frac{\det S^{\vee}}{1-q},$$

where det S^{\vee} corresponds to the determinant of the standard $\operatorname{GL}_r(\mathbb{C})$ -representation.

Theorem 1.9. The quantum reduction map κ is a surjective ring homomorphism.

Since the product in the representation ring $R(\operatorname{GL}_r(\mathbb{C}))$ is well-understood, κ can be used to analyze and reconstruct the more intricate quantum K-ring. Specifically, a presentation of $\operatorname{QK}(X)$ can be formulated by identifying relations in the kernel of κ . To illustrate this point, we provide an independent proof of the relations in the quantum K-Whitney presentation for QK(X), proposed in the physics literature [GMSZ1] and proved mathematically in [GMSZ2, Theorem 1.1]: For any vector bundle $E \to X$ of rank e, we define the polynomial

$$\wedge_y E = 1 + yE + y^2 \wedge^2 E + \dots + y^e \wedge^e E.$$

Corollary 1.10. Let S and Q be the universal subbundle and quotient bundle over X = Gr(r, N). In QK(X),

(6)
$$\wedge_y S \bullet \wedge_y Q = (1+y)^N - \frac{q}{1-q} y^k (\wedge_y S - 1) \bullet \det Q.$$

The quantum K-Whitney presentation for QK(X) is given by the algebra generated by exterior powers of S and Q, modulo the relations given by (6). These relations are equivalent to those in the 'Coulomb branch' presentation in [GMSZ1] (see Corollary 4.9)

$$\wedge_y S \bullet \wedge_y Q = (1+y)^N - qy^k (\wedge_y S - 1) \cdot \det Q.$$

We re-establish (6) using the homomorphism property of κ from Theorem 1.9 and the vanishing results for the Euler characteristics over Quot scheme of \mathbb{P}^1 mentioned in the next subsection.

1.7. Vanishing of Euler characteristics. The proofs of Theorems 1.2, 1.5 and 1.9 rely on three main ingredients: the vanishing of Euler characteristics, degeneration formulas for the Quot schemes, and wall-crossing formulas. The latter two will be explained in the next two subsections and proved in Sections 5-7. The first is proved in [SZ] using equivariant torus localization (see also the first author's Ph.D. thesis [Sin1]). We recall the statement below.

Theorem 1.11 ([SZ]). Let $\lambda = (\lambda_1, \ldots, \lambda_r)$ be a nonempty partition. We have

$$\langle \mathbb{S}^{\lambda}(\mathbf{S}) \rangle_{0,d}^{\mathsf{Quot}} = \chi(\mathsf{Quot}_d(\mathbb{P}^1, N, r), \mathbb{S}^{\lambda}(\mathcal{S}_p)) = 0.$$

if one of the following two conditions holds

- (i) $d \ge \lambda_1 (N r);$
- (ii) $d \ge \lambda_1 2(N-r)$ with additional conditions $d > 0, r \ne N$, and $\lambda_r > 0$;

When d = 0, the Quot scheme is isomorphic to Gr(r, N) and the vanishing in part (i) agrees with the Borel–Weil–Bott theorem (c.f. [Kap]).

1.8. Wall-crossings formula. To compare the K-theoretic Quot scheme invariants with (Givental-Lee's) quantum K-invariants of the Grassmannian, we apply two sequences of wallcrossings. First, we apply the K-theoretic quasimap wall-crossing formula in [ZZ] to show that quantum K-invariants equal K-theoretic ($\epsilon = 0+$)-stable quasimap invariants. Second, we recall the light-to-heavy wall-crossing formula in [RZ2] which implies that K-theoretic Quot scheme invariants equal K-theoretic (0+)-stable graph quasimap invariants. The K-theoretic (0+)-stable graph quasimap invariants and K-theoretic ($\epsilon = 0+$)-stable quasimap invariants coincide when the number of markings is less than or equal to three.



More precisely, we prove the following (see Theorem 6.10 and Corollary 6.12):

Theorem 1.12. Let $V_1, V_2, V_3 \in \mathcal{R}^{r,k} \subset K(\mathfrak{X})$, where k = N - r > 0. Denote V_1, V_2, V_3 as their restrictions to K(X). Then

$$\langle \langle \mathbf{V}_1, \mathbf{V}_2, \mathbf{V}_3 \rangle \rangle_0^{\mathsf{Quot}} = \langle \langle V_1, V_2, V_3 \rangle \rangle_{0.3}$$

where the right-hand side is the generating series of the quantum K-invariants (2). The same holds for the 2-pointed and 1-pointed invariants.

Recall the definition of the correlation functions $F_{\lambda,\mu,\nu}$ and $F_{\lambda,\mu}$ of quantum K-invariants (see (3)). Theorem 1.12 shows that:

Corollary 1.13. For any partitions $\lambda, \mu, \nu \in P_{r,k}$, we have

$$F_{\lambda,\mu,\nu} = N(0)_{\lambda,\mu,\nu}$$
 and $F_{\lambda,\mu} = N(0)_{\lambda,\mu}$.

The quantum K-ring equipped with the quantized pairing $F_{\lambda,\mu}$ is a $\mathbb{Z}[[q]]$ -valued Frobenius algebra. The above corollary implies that the TQFT structure in Theorem 1.5 coincides with the one determined by the quantum K-Frobenius algebra QK(X).

Remark 1.14. The quantum K-invariants $\langle\!\langle \mathcal{O}_{\lambda}, \mathcal{O}_{\mu}, \mathcal{O}_{\nu}^{*} \rangle\!\rangle_{0,3}$ may not match the Euler characteristics $N_{\lambda,\mu}^{\nu} = \langle\!\langle \mathcal{O}_{\lambda}, \mathcal{O}_{\mu}, \mathcal{O}_{\nu}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}}$. The discrepancy is clarified in Remark 2.3, which shows that the quantized dual $\mathcal{O}_{\nu}^{*} \in K(\mathfrak{X})$ loses information when restricted to \mathcal{O}_{ν}^{*} over $X = \operatorname{Gr}(r, N)$.

1.9. **Degeneration formula.** In algebraic geometry, the degeneration formulas for Gromov–Witten invariants and Donaldson–Thomas theory were established in [Li1, Li2] and [MPT, LW], respectively. Subsequent work extended these formulas to K-theoretic Donaldson–Thomas invariants [Qu] and to K-theoretic quasimap invariants of Nakajima quiver varieties [Oko]. In Section 7, we prove the following two degeneration formulas for virtual K-theoretic invariants of the Quot scheme for curves of all genera:

Theorem 1.15. Let $F^{\alpha,\beta}$ be the inverse of the quantized pairing matrix define in (3).

(i) Let $V, W \in K(\mathfrak{X})$. Then

(7)
$$\langle\!\langle \mathbf{V}, \mathbf{W} \rangle\!\rangle_{g}^{\mathsf{Quot}} = \sum_{\alpha, \beta \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\alpha} \rangle\!\rangle_{g_{1}}^{\mathsf{Quot}} \cdot F^{\alpha, \beta} \cdot \langle\!\langle \mathbf{O}_{\beta}, \mathbf{W} \rangle\!\rangle_{g_{2}}^{\mathsf{Quot}},$$

where α and β range over all partitions in $P_{r,k}$, and $g_1 + g_2 = g$. (ii) For any $V \in K(\mathfrak{X})$ and g > 0:

(8)
$$\langle\!\langle \mathbf{V} \rangle\!\rangle_{g}^{\mathsf{Quot}} = \sum_{\alpha,\beta \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\alpha}, \mathbf{O}_{\beta} \rangle\!\rangle_{g-1}^{\mathsf{Quot}} \cdot F^{\alpha,\beta}$$

When $V, W \in \mathcal{R}^{r,2k}$ and g = 0, the degeneration formulas follow from the WDVV equation for quantum K-theory proved in [Giv] and [Lee]. Our proof of the above degeneration formulas leverages the good degeneration of Quot schemes developed in [LW] and is modeled on that in [Qu, §4].

Remark 1.16. Let $T \cong (\mathbb{C}^*)^N \subset \operatorname{GL}_N(\mathbb{C})$ be the maximal torus of diagonal matrices acting on $X = \operatorname{Gr}(r, N)$ and $\operatorname{Quot}_d(C, N, r)$. The vanishing results in Theorem 1.11, the wall-crossing result in Theorem 1.12, and the degeneration formulas in Theorem 1.15 all hold in the torusequivariant setting.

1.10. Future directions. In this paper, we explore the quantum K-theory of the Grassmannian via Quot schemes. The Quot scheme on \mathbb{P}^1 can be thought of as a quasimap space that parametrizes maps $\mathbb{P}^1 \to \mathfrak{X}$ to the quotient stack \mathfrak{X} such that the generic point of \mathbb{P}^1 is mapped to the Grassmannian $X \subset \mathfrak{X}$. In general, let G be a reductive group acting on an affine variety W and let θ be a character of G. Replacing X and \mathfrak{X} with the GIT quotient $W//_{\theta}G$ and the quotient stack [W/G] respectively, we define the quasimap space to $W//_{\theta}G$.

Under some mild conditions (see p.341 of [KOUY]), the quasimap space has a canonical perfect obstruction theory, as detailed in [CFKM, Theorem 7.2.2] and [KOUY, §10.7]. This allows us to define virtual cohomological invariants, useful for studying the quantum cohomology and Gromov–Witten–Ruan invariants of the GIT quotient.

This approach has been applied to target spaces such as isotropic Grassmannians and partial flag manifolds (see [CF1, CF3, CF2, Che, Sin2]). We hope to extend these results to K-theory, particularly to prove the degeneration formula for the K-theoretic quasimap invariants of a general GIT quotient and to use the quantum reduction map to derive presentations of its quantum K-ring.

1.11. Plan of the paper. In Section 2, we cover preliminaries K-theory, quantum K-theory, and virtual structure sheaves. In Section 3, we prove Theorem 1.2 and Theorem 1.5, assuming the vanishing Theorem 1.11, the wall-crossing result Corollary 1.13, and the degeneration formulas from Theorem 1.15. In Section 4, we introduce the quantum reduction map κ , prove Theorem 4.4, and use it to derive the Whitney presentation for QK(X). In Section 5 and 6, we prove the wall-crossing formulas used in Sections 3 and 7. Two degeneration formulas are proven in Section 7, which is crucially used in Section 3 to show the TQFT structure of the K-theoretic Quot scheme invariants.

1.12. Acknowledgment. Both authors are grateful to Leonardo C. Mihalcea for sharing insights on the K-theoretic divisor equation and his related conjecture with Buch [GMS⁺, Conjecture 4.3]. This paper, together with the adjoining [SZ], was inspired by our efforts to verify the K-theoretic divisor equation via Quot schemes. The first author thanks Alina Marian, Dragos Oprea, and Woonam Lim for valuable conversations about the Quot schemes of curve. The second author would like to thank Wei Gu and Du Pei for explaining the physics connected to this work, and thanks Reuven Hodges and Yongbin Ruan for helpful discussions.

S. SINHA AND M. ZHANG

2. Preliminaries

2.1. Algebraic representations of $\operatorname{GL}_r(\mathbb{C})$. Recall that a *partition* (with *r* nonnegative parts) is a weakly decreasing sequence of nonnegative integers $\lambda = (\lambda_1, \ldots, \lambda_r)$, where $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_r \geq 0$. Let

$$\mathbf{P}_{r,k} = \{\lambda = (\lambda_1, \dots, \lambda_r) \mid \lambda_1 \le k\}$$

be the set of partitions contained in the rectangle $(k)^r = (k, k, ..., k)$ with r rows and k columns.

Let \mathbb{S}^{λ} be the *Schur functor* associated to λ (see [FH, §6]) and let \mathbb{C}^{r} be the standard representation of $\operatorname{GL}_{r}(\mathbb{C})$. Then $\mathbb{S}^{\lambda}(\mathbb{C}^{r})$ is the unique irreducible (polynomial) representation of $\operatorname{GL}_{r}(\mathbb{C})$ of highest weight λ . For example, if $\lambda = (a)$, then $\mathbb{S}^{(a)}(\mathbb{C}^{r}) = \operatorname{Sym}^{a}(\mathbb{C}^{r})$; if $\lambda =$ $(1)^{b} := (1, \ldots, 1)$ with 1 repeated b times, then $\mathbb{S}^{(1)^{b}}(\mathbb{C}^{r}) = \wedge^{b}(\mathbb{C}^{r})$. Let $R^{0}(\operatorname{GL}_{r}(\mathbb{C}))$ denote the ring of polynomial representations of $\operatorname{GL}_{r}(\mathbb{C})$. Note that, for any $g \in \operatorname{GL}_{r}(\mathbb{C})$, the trace of g on $\mathbb{S}^{\lambda}(\mathbb{C}^{r})$ is given by

$$\chi_{\mathbb{S}^{\lambda}(\mathbb{C}^r)}(g) = s_{\lambda}(x_1, \dots, x_r),$$

where x_1, \ldots, x_k are the eigenvalues of g and s_{λ} is the *Schur polynomial*. This gives an isomorphism χ between $R^0(\operatorname{GL}_r(\mathbb{C}))$ and the ring of symmetric polynomials in r variables.

Let $R(\operatorname{GL}_r(\mathbb{C}))$ denote the ring of algebraic representations of $\operatorname{GL}_r(\mathbb{C})$. As a ring, $R(\operatorname{GL}_r(\mathbb{C}))$ is obtained from $R^0(\operatorname{GL}_r(\mathbb{C}))$ by inverting the determinant representation $\det(\mathbb{C}^r)$. The trace map χ extends to the isomorphism between $R(\operatorname{GL}_r(\mathbb{C}))$ and the ring of symmetric Laurent polynomials in the r variables x_1, \ldots, x_r .

Let $S := (\mathbb{C}^r)^{\vee}$ denote the dual of the standard representation. Taking the dual of representations gives an automorphism of the representation ring $R(\operatorname{GL}_r(\mathbb{C}))$. At the level of Laurent polynomials, it is given by mapping $x_i \to x_i^{-1}$. In particular, the representation ring $R(\operatorname{GL}_r(\mathbb{C}))$ is also generated by $\{1, S, \wedge^2 S, \ldots, \det(S), \det(S)^{\vee}\}$ over \mathbb{C} . Let \mathcal{R}^r denote subring $R(\operatorname{GL}_r(\mathbb{C}))$ generated by $\{1, S, \wedge^2 S, \ldots, \det(S)\}$, that is, \mathcal{R}^r corresponds to the subring consisting of symmetric polynomials in $x_1^{-1}, \ldots, x_r^{-1}$ inside the ring of symmetric Laurent polynomials.

Definition 2.1. Given a positive integer ℓ , let $\mathcal{R}^{r,\ell}$ be the linear subspace of \mathcal{R}^r spanned by polynomials in

$$\{S, \wedge^2 S, \ldots, det(S)\}$$

of degree at most ℓ . Note that each element $\wedge^i S$ has degree 1. The Jacobi-Trudi formula (of second type) implies that $\mathcal{R}^{r,\ell}$ is spanned by $\{\mathbb{S}^{\lambda}(S) \mid \lambda \in P_{r,\ell}\}$.

2.2. Basic notation in K-theory. Let X be a Deligne–Mumford (DM) stack with an action of an algebraic torus T. The group $K_0^T(X)$ (resp. $K_T^0(X)$) denotes the Grothendieck group of T-equivariant coherent sheaves (resp. T-equivariant locally free sheaves) on X. Given a (equivariant) coherent sheaf (or vector bundle) E, we denote by [E] or simply E its associated K-theory class. The group $K_0^T(X)$ has a $K_T^0(X)$ -module structure defined by the tensor product

$$K_T^0(X) \otimes K_0^T(X) \to K_0^T X, \quad ([E], [F]) \mapsto [E] \cdot [F] := [E \otimes_{\mathcal{O}_X} F].$$

We denote by $K_0(X)$ and $K^0(X)$ the non-equivariant K-groups of coherent sheaves and locally free sheaves, respectively. If X is nonsingular, the map $K^0(X) \to K_0(X)$ sending a vector bundle to its sheaf of sections is an isomorphism. In this case, we denote the non-equivariant K-group of X by K(X).

Let $pt = \operatorname{Spec} \mathbb{C}$. If we write $T = (\mathbb{C}^*)^N$, then

$$\Gamma := K_T^0(\mathrm{pt}) = K^0(BT) \cong \mathbb{Z}[\alpha_1^{\pm}, \dots, \alpha_N^{\pm}],$$

where α_i is the character of T given by the projection to the *i*-th factor. Given a character α of T, let \mathbb{C}_{α} denote the corresponding 1-dimensional representation of T.

Let F be a T-equivariant vector bundle on X. We define the K-theoretic Euler class of F by

$$\wedge_{-1}^{T}(F^{\vee}) := \sum_{i \ge 0} (-1)^{i} \wedge^{i} F^{\vee} \in K_{T}^{0}(X),$$

where F^{\vee} denotes the dual of F and $\wedge^i F^{\vee}$ denotes the *i*-th exterior power of F^{\vee} .

Given a *T*-equivariant morphism $f : X \to Y$, we have the pullback map $f^* : K_T^0(Y) \to K_T^0(X)$. If f is proper, we have the proper pushforward $f_* : K_0^T(X) \to K_0^T(Y)$ defined by

$$[F] \mapsto \sum_{n} (-1)^{n} [R^{n} f_{*} F]$$

Given a class $F \in K_0^T(X)$ on a proper DM stack X, we define its T-equivariant Euler characteristic as the proper pushforward to the point

$$\chi^T(X,F) := \sum_{i \ge 0} \operatorname{char}_T H^i(X,F) \in \Gamma.$$

where char_T denotes taking the character of a T-module. Furthermore, if X has a T-equivariant virtual struture sheaf $\mathcal{O}_X^{\text{vir}} \in K_0^T(X)$ (see Section 2.6), we define the T-equivariant virtual Euler characteristic by

$$\chi^{\operatorname{vir},T}(X,F) := \chi^T(X,\mathcal{O}_X^{\operatorname{vir}}\cdot F).$$

2.3. K-theory of the Grassmannian. We fix two positive integers r and N such that $N \ge r$. Throughout the paper, we set k = N - r and

$$X = \operatorname{Gr}(r, N) = \{ V \subset \mathbb{C}^N \mid \dim_{\mathbb{C}} V = r \}$$

to be the Grassmann variety of r-planes in \mathbb{C}^N .

Let $T \cong (\mathbb{C}^*)^N \subset \operatorname{GL}_N(\mathbb{C})$ be the maximal torus of diagonal matrices. Then T acts on X. There is a tautological subbundle $S \subset \mathcal{O}_X^{\oplus N} = X \times \mathbb{C}^N$ of rank r on X, and it has a natural T-linearization. We consider the following standard GIT presentation of X:

$$X = [M_{r \times N}^{ss} / \mathrm{GL}_r(\mathbb{C})],$$

where $M_{r \times N}$ is the affine space of r by N complex matrices and $M_{r \times N}^{ss} \subset M_{r \times N}$ is the open subset of matrices of full rank. Let

$$\mathfrak{X} = [M_{r \times N} / \mathrm{GL}_r(\mathbb{C})]$$

be the quotient stack. Let $\Gamma = K_T^0(\text{pt})$, then there is a homomorphism

(9)
$$R(\operatorname{GL}_r(\mathbb{C})) \to K^0_T(\mathfrak{X}), \quad \mathcal{V} \mapsto M_{r \times N} \times_{\operatorname{GL}_r(\mathbb{C})} \mathcal{V}.$$

Here the *T*-linearization on $M_{r\times N} \times_{\operatorname{GL}_r(\mathbb{C})} V$ is defined via $\lambda \cdot [m, v] = [m\lambda, v]$, where $\lambda \in T$, $m \in M_{r\times N}$, and $v \in V$. Since $K_T^0(\mathfrak{X})$ is a Γ -module, we can extend (9) to a surjective homomorphism $R(\operatorname{GL}_r(\mathbb{C})) \otimes \Gamma \to K_T^0(\mathfrak{X})$. We denote by $V \in K_T^0(X)$ the restriction of V to X, i.e.,

$$V = M_{r \times N}^{ss} \times_{\operatorname{GL}_r(\mathbb{C})} \operatorname{V}.$$

We abuse notation by referring to V as the K-theory class in $K^0_T(\mathfrak{X})$ associated with the $\operatorname{GL}_r(\mathbb{C})$ -module V, using Roman font to distinguish classes over the quotient stack \mathfrak{X} from those over the Grassmannian X.

Let \mathbb{C}^r denote the standard representation of $\operatorname{GL}_r(\mathbb{C})$, and let $S = (\mathbb{C}^r)^{\vee}$ be its dual. Let

(10)
$$\mathbf{S} := M_{r \times N} \times_{\mathrm{GL}_r(\mathbb{C})} (\mathbb{C}^r)^{\vee}$$

be the associated vector bundle to the representation S. Note that the restriction of S to X is the tautological subbundle S.

The K-theoretic Kirwan map is the composition of the homomorphism (9) with the restriction map $K_T^0(\mathfrak{X}) \to K_T^0(X)$. It induces the following linear isomorphism

$$\mathcal{R}^{r,k} \xrightarrow{\sim} K(X),$$

sending the representation $\mathbb{S}^{\lambda}(S)$ to the vector bundle $\mathbb{S}^{\lambda}(S)$. The collection $\{\mathbb{S}^{\lambda}(S)\}_{\lambda \in \mathcal{P}_{r,k}}$ is a \mathbb{Z} -basis of the K-group K(X) and a basis of the equivariant K-group $K_T^0(X)$ over $\Gamma = K_T^0(\operatorname{pt})$.

There is another \mathbb{Z} -basis of K(X) given by the *Schubert structure sheaves*. Recall that the *Schubert variety* for λ relative to the Borel subgroup of $\operatorname{GL}_r(\mathbb{C})$ stabilizing a complete flag $0 = F_0 \subset F_1 \subset \cdots \subset F_N = \mathbb{C}^N$ is defined by

$$X_{\lambda} = \{ V \in X \mid \dim(V \cap F_{k+i-\lambda_i}) \ge i, \quad \forall 1 \le i \le r \}.$$

The codimension of X_{λ} equals the weight $|\lambda| = \sum \lambda_i$ of λ . The K-theory class $\mathcal{O}_{\lambda} := [\mathcal{O}_{X_{\lambda}}]$ of the structure sheaf of X_{λ} is referred to as the Schubert structure sheaf. The set $\{\mathcal{O}_{\lambda}\}_{\lambda \in P_{r,k}}$ is a \mathbb{Z} -basis of K(X).

Recall that Schubert structure sheaves can be represented in terms Schur bundles in K(X)using a linear transformation given by

$$\mathcal{O}_{\lambda} = \sum_{\nu \in \mathcal{P}_{r,k}} \mathcal{A}_{\lambda}^{\nu} \, \mathbb{S}^{\nu}(S)$$

for any $\lambda \in P_{r,k}$. Below we use *stable Grothendieck polynomials* to describe the coefficient A_{λ}^{ν} . Furthermore we define the "stacky" structure sheaves $O_{\lambda} \in K(\mathfrak{X})$ such that the Kirwan

map (see [Buc, §8]) takes O_{λ} to the Schubert structure sheaf

$$\mathcal{O}_{\lambda} = \mathbb{G}^{\lambda}(S^{\vee}) \in K(X)$$

for $\lambda \in \mathbf{P}_{r,k}$.

2.4. Grothendieck functor. Let G_{λ} be the stable Grothendieck polynomial for the partition λ (see [Buc]). It has the following Weyl-type bialternant formula:

$$G_{\lambda}(x_1, \dots, x_r) = \frac{\det(x_j^{\lambda_i + r - i}(1 - x_j)^{i-1})_{1 \le i, j \le r}}{\det(x_j^{r-i})_{1 \le i, j \le r}}$$

By [Len, Theorem 2.2], we have the following expansion

(11)
$$G_{\lambda}(x_1,\ldots,x_r) = \sum_{\lambda \subseteq \mu \subseteq \hat{\lambda}} (-1)^{|\mu| - |\lambda|} g_{\lambda\mu} s_{\mu}(x_1,\ldots,x_r),$$

where $\hat{\lambda}$ is the unique maximal partition with r rows obtained from λ by adding at most i-1 boxes to its *i*th row for $2 \leq i \leq r$, and the coefficients $g_{\lambda,\mu}$ are nonnegative and count certain increasing skew tableaux.

According to [Len, §8], there is a well-defined virtual $\operatorname{GL}_r(\mathbb{C})$ -representation $\mathbb{G}^{\lambda}(\mathbb{C}^r)$ whose character satisfies

$$\chi_{\mathbb{G}^{\lambda}(V)}(g) = G_{\lambda}(1 - x_1^{-1}, \dots, 1 - x_r^{-1}),$$

where x_1, \ldots, x_r are the eigenvalues of g on V. Let λ be a partition with at most r parts, then

$$\mathbb{G}^{\lambda}(\mathbb{C}^{r}) = \sum_{\nu \in \mathcal{P}_{r,\lambda_{1}}} \mathcal{A}^{\nu}_{\lambda} \mathbb{S}^{\nu}(\mathcal{S}),$$

where $S = (\mathbb{C}^r)^{\vee}$ is the dual representation. The coefficients A^{ν}_{λ} can be calculated using (11) and expressing $s_{\mu}(1 - x_1^{-1}, \dots, 1 - x_r^{-1})$ as a linear combination of $s_{\nu}(x_1^{-1}, \dots, x_r^{-1})$.

The description of $g_{\lambda,\mu}$ in (11) and the fact that $s_{\lambda}(1 - x_1^{-1}, \ldots, 1 - x_r^{-1})$ can be written as a linear combination of $s_{\mu}(x_1^{-1}, \ldots, x_r^{-1})$ with $\mu \subseteq \lambda$ (see, e.g., [Mac, §3, Example 10]) implies that $\mathcal{R}^{r,k}$ in Definition 2.1 is spanned by $\{\mathbb{G}^{\lambda}(\mathbb{C}^r) : \lambda \in P_{r,k}\}$. This eludes to the fact that the Schubert structure sheaves $\{\mathcal{O}_{\lambda} : \lambda \in P_{r,k}\}$ give a linear basis for the K-theory of $\operatorname{Gr}(r, N)$.

Definition 2.2. For any partition λ (with r nonnegative parts), we define the stacky Schubert structure sheaf by

$$O_{\lambda} := \mathbb{G}^{\lambda}(S^{\vee}) \in K(\mathfrak{X}),$$

where S corresponds to the $\operatorname{GL}_r(\mathbb{C})$ representation $(\mathbb{C}^r)^{\vee}$ (see (10)).

Note that \mathcal{O}_{λ} equals zero in K(X) if λ is not contained in $(k)^r$, while O_{λ} is a non-zero element in $K(\mathfrak{X})$ for any partition λ with at most r parts.

Remark 2.3. Note that the quantized K-theoretic dual $O_{\lambda}^* = O_{\lambda}^* \cdot \det(S)$ restricts to the classical dual \mathcal{O}_{λ}^* for $\lambda \in P_{r,k}$. However, if we expand O_{λ}^* in terms of the stacky Schubert

structure sheaves, the expansion may involve O_{ν} with $\nu \notin P_{r,k}$ which restricts to zero in K(X). For example, when r = 2 and N = 4, we have

$$O_{(1)}^* = O_{(2,1)} \cdot \det S = O_{(2,1)} - O_{(2,2)} - O_{(3,1)} + O_{(3,2)}$$

while

$$\mathcal{O}_{(1)}^* = \mathcal{O}_{(1)}^*|_X = \mathcal{O}_{(2,1)} - \mathcal{O}_{(2,2)}$$

This shows that in general, the quantized K-theoretic duals may not live in K(X) and they contain more information than their classical counterparts.

2.5. Quantum *K*-theory. There are different variants of quantum *K*-invariants (see, for example, [Lee, RZ1, RZ2, Oko]). In this subsection, we recall the definition of genus-zero quantum *K*-invariants of Grassmannians in the sense of Givental–Lee [Lee, Giv, BM].

Let $X = \operatorname{Gr}(r, N)$ be a Grassmannian. For $d \in H_2(X) \cong \mathbb{Z}$, let $\overline{M}_{0,n}(X, d)$ be the moduli stack of degree-*d*, genus zero, *n*-pointed *stable maps* into *X*. This is a smooth proper DM stack equipped with *evaluation maps*

$$\operatorname{ev}_i : \overline{M}_{0,n}(X,d) \to X$$

for $1 \leq i \leq n$.

Definition 2.4. Let $E_1, \ldots, E_n \in K_T^0(X)$. We define the genus zero, n-pointed, T-equivariant quantum K-invariant with the insertions E_i 's by

$$\langle E_1, \dots, E_n \rangle_{0,n,d}^T := \chi^T \left(\overline{M}_{0,n}(X,d), \prod_{i=1}^n \operatorname{ev}_i^*(E_i) \right).$$

The corresponding correlation function is defined by

$$\langle\!\langle E_1,\ldots,E_n\rangle\!\rangle_{0,n}^T := \sum_{d\geq 0} q^d \langle E_1,\ldots,E_n\rangle_{0,n,d}^T,$$

where q is a formal variable called the Novikov variable. The non-equivariant quantum Kinvariant and its correlation function are denoted by $\langle E_1, \ldots, E_n \rangle_{0,n}$ and $\langle \langle E_1, \ldots, E_n \rangle \rangle_{0,n}$, respectively.

We assume k = N - r > 0. Let $\{\mathcal{O}_{\lambda}\}_{\lambda \in P_{r,k}}$ be a basis for K(X). We recall the definition of quantum K-ring [Giv, Lee, BM]. In general, we can choose any basis $\{\phi_{\lambda}\}_{\lambda \in P_{r,k}}$ of $K_T^0(X)$ and define T-equivariant quantum K-ring.

Recall the following generating series of the 3-pointed Givental-Lee's invariants

$$F_{\alpha,\beta,\gamma} := \langle\!\langle \mathcal{O}_{\alpha}, \mathcal{O}_{\beta}, \mathcal{O}_{\gamma} \rangle\!\rangle_{0,3} = \sum_{d \ge 0} q^d \langle \mathcal{O}_{\alpha}, \mathcal{O}_{\beta}, \mathcal{O}_{\gamma} \rangle_{0,3,d}$$

for partitions $\alpha, \beta, \gamma \in P_{r,k}$. Similarly we define $F_{\alpha,\beta} = \langle \langle \mathcal{O}_{\alpha}, \mathcal{O}_{\beta} \rangle \rangle_{0,2}$. Let $(F^{\alpha,\beta})$ be the matrix inverse to the quantized pairing matrix $(F_{\alpha,\beta})$.

Given three partitions $\lambda, \mu, \nu \in P_{r,k}$, we define the structure constant by

$$N_{\lambda,\mu}^{\nu} := \sum_{\alpha \in \mathbf{P}_{r,k}} F^{\nu,\alpha} F_{\alpha,\lambda,\mu}.$$

The WDVV equation in quantum K-theory (see [Giv]) shows that the quantum K-product defined by

$$\Phi_{\lambda} \bullet \Phi_{\mu} = \sum_{\nu} N^{\nu}_{\lambda,\mu} \Phi_{\nu}$$

is associative. The quantum K-ring of X, denoted by QK(X), is the $\mathbb{C}[[q]]$ -algebra $K(X) \otimes \mathbb{C}[[q]]$ equipped with the quantum K-product. If we take the non-equivariant limit, we obtain the quantum K-ring of X, denoted by QK(X).

Givental–Lee showed that the quantum K-ring QK(X) equipped with the quantized pairing $F_{\alpha,\beta}$ is a (formal) commutative *Frobenius algebra* with unit 1; see [Giv, Corollary 1] and [Lee, Proposition 12 (1)].

2.6. Perfect obstruction theories and virtual structure sheaves. Let $Quot_d(C, N, r)$ be the Quot scheme introduced in Section 1.3. Let S and Q be the universal subbundle and quotient sheaf over the universal curve $\pi : Quot_d(C, N, r) \times C \to Quot_d(C, N, r)$. Then

$$(R\pi_*\mathcal{H}om(\mathcal{S},\mathcal{Q}))$$

defines a perfect obstruction theory (POT) for $Quot_d(C, N, r)$ [MO1, Theorem 1].

In general, consider a projective morphism $\mathcal{C} \to B$ over a scheme B, where the fibers are prestable curves. Let $Q_{\mathcal{C}/B}$ denote the relative Quot scheme, parametrizing B-flat families of coherent quotients of rank-N trivial sheaves on the fibers. Define $Q' \subset Q_{\mathcal{C}/B}$ as the open locus corresponding to locally free subsheaves. Consider the universal sequence

$$0 \to \mathcal{S} \to \mathcal{O}_{\mathcal{C}_{\mathbf{Q}'}}^{\oplus N} \to \mathcal{Q} \to 0$$

over the universal curve $\pi : \mathcal{C}_{Q'} := Q' \times_B \mathcal{C} \to B$. According to the proof of [MOP, Theorem 2],

(12)
$$(R \mathcal{H}om_{\pi}(\mathcal{S}, \mathcal{Q}))^{\vee} = (R\pi_* \mathcal{H}om(\mathcal{S}, \mathcal{Q}))^{\vee}$$

defines a (2-term) perfect obstruction theory relative to $\nu : \mathbf{Q}' \to B$. If we consider the $T \cong (\mathbb{C}^*)^N$ -action on \mathbf{Q}' , induced by the action on $\mathcal{O}^{\oplus N}$, and the trivial *T*-action on *B*, the complex (12) defines a *T*-equivariant POT for the morphism $\nu_T : [\mathbf{Q}'/T] \to [B/T]$.

Note that the injection $\mathcal{S} \to \mathcal{O}_{\mathcal{C}_{\Omega'}}^{\oplus N}$ induces a map

$$\mathcal{C} \xrightarrow{f} \mathfrak{X}.$$

where $\mathfrak{X} = [M_{r \times N}/\mathrm{GL}_r(\mathbb{C})]$ is the quotient stack containing the Grassmannian X. Since the tangent complex $\mathbb{T}_{\mathfrak{X}}$ of \mathfrak{X} is quasi-isomorphic to $[\mathfrak{gl}_r(\mathbb{C}) \otimes \mathcal{O}_{M_{r \times N}} \to T_{M_{r \times N}}]$ (of amplitude in [-1,0]), we have

$$Lf^*\mathbb{T}_{\mathfrak{X}}\simeq [\mathcal{H}\mathrm{om}(\mathcal{S},\mathcal{S})
ightarrow (\mathcal{S}^{ee})^{\oplus N}]\simeq \mathcal{H}\mathrm{om}(\mathcal{S},\mathcal{Q}).$$

Hence, the POT (12) can also be written as

(13)
$$(R\pi_*Lf^*\mathbb{T}_{\mathfrak{X}})^{\vee},$$

which resembles that for stable maps (see, e.g. [Beh]).

By [Qu, Definition 2.2], the relative POT (12) (or, equivalently, (13)) induces virtual pullbacks

 $\nu^{!}: K_{0}(B) \rightarrow K_{0}(\mathbf{Q}') \text{ and } \nu^{!}_{T}: K_{0}^{T}(B) \rightarrow K_{0}^{T}(\mathbf{Q}').$

As an application, we define the (T-equivariant) virtual structure sheaf

$$\mathcal{O}_{\operatorname{\mathsf{Quot}}_d(C,N,r)}^{\operatorname{vir}} := \nu^! (\mathcal{O}_{\operatorname{Spec} \mathbb{C}}) \in K_0^T(\operatorname{\mathsf{Quot}}_d(C,N,r))$$

as the virtual pullback of $\mathcal{O}_{\operatorname{Spec}\mathbb{C}}$ along ν .

3. QUANTUM-K TQFT

In this section, we prove Theorem 1.2 and Theorem 1.5, assuming the wall-crossing result from Corollary 1.13 and the degeneration formulas from Theorem 1.15. We make critical use of a vanishing theorem for the Quot scheme from an adjoining paper [SZ]. Notably, this vanishing result, proven using Atiyah–Bott localization on the Quot scheme over \mathbb{P}^1 , is logically independent of the findings in this paper.

3.1. Orthogonality. Recall that $\mathcal{R}^{r,\ell}$ in Definition 2.1 is the subspace of the representation ring $R(\operatorname{GL}_r(\mathbb{C}))$ spanned by $\{\mathbb{S}^{\lambda}(S) \mid \lambda \in \mathbb{P}_{r,\ell}\}$. The classical Littlewood-Richardson rule implies the following very useful property: For any $V \in \mathcal{R}^{r,\ell}$ and $W \in \mathcal{R}^{r,m}$, the product

(14)
$$\mathbf{V} \cdot \mathbf{W} \in \mathcal{R}^{r,\ell+m}.$$

We use the the vanishing result in Theorem 1.11 to prove the following results.

Corollary 3.1. For any $V, W \in \mathcal{R}^{r,k} \cong K(X)$ and $d \ge m > 0$, we have $\langle \det(S)^m, V, W \rangle_{0,d}^{\mathsf{Quot}} = 0.$

Proof. Note that $det(S)^m = S^{(m)^r}(S)$, where $(m)^r = (m, \dots, m)$. Therefore,

$$\det(\mathbf{S})^m \cdot \mathbf{V} \cdot \mathbf{W} \in \mathcal{R}^{r,m+2k}$$

Since S has rank r, the Littlewood-Richardson rule/Pieri rule implies that det(S)·S^{λ}(S) = S^{ν}(S) where $\nu = (\nu_1, \ldots, \nu_r) = (\lambda_1 + 1, \ldots, \lambda_r + 1)$. Each term S^{ν}(S) in the Schur expansion of det(S)^m · V · W satisfies $\nu \in \mathcal{R}^{r,m+2k}$ and $\nu_r \ge m > 0$. For all such ν , part(ii) of Theorem 1.11 implies $\langle S^{\nu}(S) \rangle_{0,d}^{\mathsf{Quot}} = 0$ whenever $d \ge m$.

By setting m = 1 in the above corollary and using Theorem 1.12, we obtain an alternate proof of a conjecture of Buch–Mihalcea [GMS⁺, Conjecture 4.3] in the case of Grassmannians. This result may be thought of as a *divisor equation* for the K-theoretic Gromov-Witten invariants: For $V, W \in K_T^0(X)$ and d > 0, the quantum K-invariants satisfy

$$\langle \mathcal{O}_{(1)}, V, W \rangle_{0,3,d} = \langle V, W \rangle_{0,2,d}$$

For more general homogeneous spaces, the K-theoretic 'divisor equation' conjectuerd by Buch–Mihalcea plays crucial roles in studying quantum K-product. For example, it is used in the proof of the Chevalley formula for the equivariant quantum K-theory of cominuscule varieties [BCMP3, Theorem 3.9]. Gu–Mihalcea–Sharpe–Xu–Zhang–Zou [GMS⁺] also used it to

(partially) verify the conjectured complete set of relations for the equivariant quantum K-ring of the partial flag manifolds.

Proposition 3.2 (Orthogonality). We have

$$N(0)_{\lambda}^{\nu} = \langle\!\langle \mathcal{O}_{\lambda}, \mathcal{O}_{\nu}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} = \delta_{\lambda, \nu}$$

for $\nu, \lambda \in \mathbf{P}_{r,k}$.

Proof. The restriction of O_{ν}^* to Gr(r, N) is the classical K-theoretic dual $\mathcal{O}_{\nu}^* = (1 - \mathcal{O}_{(1)}) \cdot \mathcal{O}_{\nu^*}$ defined in [BM, §5.1] satisfying

(15)
$$\chi(X, \mathcal{O}_{\nu} \cdot \mathcal{O}_{\lambda}^*) = \delta_{\nu, \lambda}.$$

The degree zero part of the identity follows from (15) and the identity $(1 - \mathcal{O}_{(1)}) = \det(S)$. For d > 0, we have

$$\langle \mathcal{O}_{\lambda}, \mathcal{O}_{\nu}^{*} \rangle_{0,d}^{\mathsf{Quot}} = \langle \mathcal{O}_{\lambda}, \mathcal{O}_{\nu^{*}}, \det(\mathcal{S}) \rangle_{0,d}^{\mathsf{Quot}} = 0,$$

where the last equality follows from Corollary 3.1.

Remark 3.3. Note that O_{λ}^* may not correspond to a representation in $\mathcal{R}^{r,k}$. In fact, $\operatorname{Span}\{O_{\lambda}^* \mid \lambda \in P_{r,k}\} \subset K(\mathfrak{X})$ correspond to $\operatorname{Span}\{\mathbb{S}^{\lambda}(S) \mid \lambda \in P_{r,k+1} \text{ and } \lambda_r > 0\} \subset R(\operatorname{GL}_r(\mathbb{C}))$. Nevertheless, the above proposition shows that $\{O_{\lambda}^*\}$ are dual to $\{O_{\lambda}\}$ with respect to the pairing $\langle \langle -, - \rangle \rangle_0^{\operatorname{Quot}}$.

3.2. Structure constant. Corollary 1.13 states that $F_{\lambda,\mu,\nu} = N(0)_{\lambda,\mu,\nu}$ and $F_{\lambda,\mu} = N(0)_{\lambda,\mu}$ for any partitions $\lambda, \mu, \nu \in P_{r,k}$. Below we give a description of the inverse of the quantized pairing matrix $(F^{\alpha,\beta})$.

Corollary 3.4. We have

- (i) $F^{\lambda,\nu} = N(0)^{\lambda,\nu}$.
- (ii) $F^{\lambda,\nu}$ is a linear function in q.

Proof. It follows from the degeneration formula in Theorem 1.15 that

$$N(0)^{\lambda,\nu} = \langle\!\langle \mathcal{O}^*_{\lambda}, \mathcal{O}^*_{\nu} \rangle\!\rangle_0^{\mathsf{Quot}} = \sum_{\alpha,\beta \in \mathcal{P}_{r,k}} \langle\!\langle \mathcal{O}^*_{\lambda}, \mathcal{O}_{\alpha} \rangle\!\rangle_0^{\mathsf{Quot}} F^{\alpha,\beta} \langle\!\langle \mathcal{O}_{\beta}, \mathcal{O}^*_{\nu} \rangle\!\rangle_0^{\mathsf{Quot}} = F^{\lambda,\nu}.$$

Here we have used orthogonality property of Proposition 3.2 in the last equality. (ii) follows from the fact

$$\langle \mathcal{O}_{\nu}^{*},\mathcal{O}_{\lambda}^{*}\rangle_{0,d}^{\mathsf{Quot}} = \langle \mathcal{O}_{\nu^{*}},\mathcal{O}_{\lambda^{*}},\det(\mathcal{S})^{2}\rangle_{0,d}^{\mathsf{Quot}} = 0,$$

for any $d \ge 2$ by Corollary 3.1.

Remark 3.5. For a finite-dimensional Frobenius algebra, the formula in Corollary 3.4 (i) follows trivially from the orthogonality in Proposition 3.2. Here, in the infinite-dimensional setting, the proof relies on the degeneration formula. The above corollary also implies that $N(0)^{\lambda,\mu}$ is the inverse of the pairing matrix $N(0)_{\lambda,\mu}$.

As further applications of the quantized K-theoretic duals, we finish the proof of Theorem 1.2, i.e. show that the structure constant of the quantum K-product $N^{\nu}_{\lambda,\mu}$ equals $N(0)^{\nu}_{\lambda,\mu}$. *Proof of Theorem 1.2.* Recall that the structure constant is given by the sum

$$N_{\lambda,\mu}^{\nu} = \sum_{\alpha \in \mathcal{P}_{r,k}} F_{\lambda,\mu,\alpha} F^{\alpha,\nu}.$$

Using Corollary 1.13, we obtain

$$\begin{split} N_{\lambda,\mu}^{\nu} &= \sum_{\alpha \in \mathcal{P}_{r,k}} \langle\!\langle \mathcal{O}_{\lambda}, \mathcal{O}_{\mu}, \mathcal{O}_{\alpha} \rangle\!\rangle_{0}^{\mathsf{Quot}} F^{\alpha,\nu} \\ &= \sum_{\alpha,\beta \in \mathcal{P}_{r,k}} \langle\!\langle \mathcal{O}_{\lambda}, \mathcal{O}_{\mu}, \mathcal{O}_{\alpha} \rangle\!\rangle_{0}^{\mathsf{Quot}} F^{\alpha,\beta} \langle\!\langle \mathcal{O}_{\beta}, \mathcal{O}_{\nu}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} \\ &= \langle\!\langle \mathcal{O}_{\lambda}, \mathcal{O}_{\mu}, \mathcal{O}_{\nu}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} = N(0)_{\lambda,\mu}^{\nu}. \end{split}$$

Here we used the degeneration formula from Theorem 1.15 in the second-to-last equality.

The formula of the structure constant in the above corollary resembles that in a Frobenius algebra. It is not only more transparent and compact but also allows us to give intuitive proofs of some results of Buch–Mihalcea [BM].

Corollary 3.6 ([BM, Theorem 5.13]). The structure constants satisfy the S_3 -symmetry, i.e., $N^{\nu}_{\lambda,\mu} = N^{\mu^*}_{\lambda,\nu^*}$.

Proof. It follows from Theorem 1.2 and Definition 1.3 that

$$\langle\!\langle \mathcal{O}_{\lambda}, \mathcal{O}_{\mu}, \mathcal{O}_{\nu}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} = \langle\!\langle \mathcal{O}_{\lambda} \cdot \mathcal{O}_{\mu} \cdot \mathcal{O}_{\nu^{*}} \cdot \det(\mathcal{S}) \rangle\!\rangle_{0}^{\mathsf{Quot}}.$$

This easily implies the S_3 -symmetry.

Part (i) of Theorem 1.11 implies that the structure constants $N_{\lambda,\mu}^{\nu}$ are polynomials in the Novikov variable q of degree at most 2(N-r). In fact, by using the stronger vanishing result in part (ii) of Theorem 1.11, we can obtain the following optimal bounds of the degree of $N_{\lambda,\mu}^{\nu}$:

Corollary 3.7 ([BM, Corollary 5.8]). The structure constants $N_{\lambda,\mu}^{\nu}$ are polynomials in q of degree $\leq \min\{r, k\}$ with k = N - r.

Proof. Since $QK(Gr(r, N)) \cong QK(Gr(k, N))$, we only need to show that the degree of $N_{\lambda,\mu}^{\nu}$ is less than or equal to k. Using a similar argument to the proof of Corollary 3.1, each term $\mathbb{S}^{\nu}(S)$ in the Schur exapansion of $O_{\lambda} \cdot O_{\mu} \cdot O_{\nu^*} \cdot \det(S)$ is contained in $\mathcal{R}^{r,3k+1}$, and ν satisfies the condition $\nu_r > 0$. For all such ν , part(ii) of Theorem 1.11 implies $\langle \mathbb{S}^{\nu}(S) \rangle_{0,d}^{\mathsf{Quot}} = 0$ whenever $d \ge k+1$.

Remark 3.8. The property proved in the previous corollary is often referred to as finiteness of the quantum K-product. It was first proved for Grassmannians in [BM] and then generalized to cominuscule varieties [BCMP1, BCMP2] and flag manifolds [Kat, ACT1, ACT2]. As shown

in the proof of the above corollary, the finiteness is a consequence of the vanishing of certain K-theoretic Quot scheme invariants.

3.3. Proof of Theorem 1.5.

Lemma 3.9. For any $V, W \in K(\mathfrak{X})$ and $g_1 + g_2 = g$,

(16)
$$\langle\!\langle \mathbf{V}, \mathbf{W} \rangle\!\rangle_g^{\mathsf{Quot}} = \sum_{\rho \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\rho} \rangle\!\rangle_{g_1}^{\mathsf{Quot}} \cdot \langle\!\langle \mathbf{O}_{\rho}^*, \mathbf{W} \rangle\!\rangle_{g_2}^{\mathsf{Quot}}.$$

Proof. We repeatedly use the degeneration formula in Theorem 1.15.

$$\begin{split} \langle\!\langle \mathbf{V}, \mathbf{W} \rangle\!\rangle_{g}^{\mathsf{Quot}} &= \sum_{\rho, \beta \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\rho} \rangle\!\rangle_{g_{1}}^{\mathsf{Quot}} \cdot F^{\rho, \beta} \cdot \langle\!\langle \mathbf{O}_{\beta}, \mathbf{W} \rangle\!\rangle_{g_{2}}^{\mathsf{Quot}} \\ &= \sum_{\rho \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\rho} \rangle\!\rangle_{g_{1}}^{\mathsf{Quot}} \cdot \sum_{\alpha, \beta \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{O}_{\rho}^{*}, \mathbf{O}_{\alpha} \rangle\!\rangle_{0}^{\mathsf{Quot}} \cdot F^{\alpha, \beta} \cdot \langle\!\langle \mathbf{O}_{\beta}, \mathbf{W} \rangle\!\rangle_{g_{2}}^{\mathsf{Quot}} \\ &= \sum_{\rho \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\rho} \rangle\!\rangle_{g_{1}}^{\mathsf{Quot}} \cdot \langle\!\langle \mathbf{O}_{\rho}^{*}, \mathbf{W} \rangle\!\rangle_{g_{2}}^{\mathsf{Quot}}. \end{split}$$

Here we used the orthogonality relation in Proposition 3.2 for the second equality. \Box

To prove the gluing relation in Theorem 1.5, it is enough to show the following two gluing formulas:

• Let $\underline{\lambda} = (\lambda^1, \dots, \lambda^s)$ and $\underline{\nu} = (\nu^1, \dots, \nu^t)$ be tuples of partitions in $P_{r,k}$. Then for any $g_1 + g_2 = g$ and for any splitting of tuples $\underline{\lambda} = \underline{\lambda}_1 \cup \underline{\lambda}_2$ and $\underline{\nu} = \underline{\nu}_1 \cup \underline{\nu}_2$,

$$N(g)_{\underline{\lambda}}^{\underline{\nu}} = \sum_{\rho \in \mathcal{P}_{r,k}} N(g_1)_{\underline{\lambda_1},\rho}^{\underline{\nu_1}} \cdot N(g_2)_{\underline{\lambda_2}}^{\underline{\nu_2},\rho}.$$

Proof. Set $V = O_{\underline{\lambda_1}} \cdot O_{\underline{\nu_1}}^*$ and $W = O_{\underline{\lambda_2}} \cdot O_{\underline{\nu_2}}^*$ in (16).

• For any tuple of partitions $\underline{\lambda}$ and $\underline{\nu}$ and g > 0

$$N(g)_{\underline{\lambda}}^{\underline{\nu}} = \sum_{\rho \in \mathcal{P}_{r,k}} N(g-1)_{\underline{\lambda},\rho}^{\underline{\nu},\rho}$$

Proof. The second degeneration formula in Theorem 1.15 implies that

$$\begin{split} \langle\!\langle \mathbf{V} \rangle\!\rangle_{g}^{\mathsf{Quot}} &= \sum_{\alpha,\beta \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\alpha}, \mathbf{O}_{\beta} \rangle\!\rangle_{g-1}^{\mathsf{Quot}} \cdot F^{\alpha,\beta} \\ &= \sum_{\alpha,\beta \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\alpha}, \mathbf{O}_{\beta} \rangle\!\rangle_{g-1}^{\mathsf{Quot}} \cdot \langle\!\langle \mathbf{O}_{\beta}^{*}, \mathbf{O}_{\alpha}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} \\ &= \sum_{\alpha \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\alpha}, \mathbf{O}_{\alpha}^{*} \rangle\!\rangle_{g-1}^{\mathsf{Quot}}. \end{split}$$

Here we used Corollary 3.4 in the second equality, and (16) for the last equality. Set $V = O_{\underline{\lambda}} \cdot O_{\underline{\nu}}^*$ to obtain the required gluing formula.

S. SINHA AND M. ZHANG

4. QUANTUM REDUCTION MAP

In this section, we introduce the quatum reduction map κ and prove it is a ring homomorphism, as described in Theorem 1.9. We then explore some applications, including re-deriving the Whitney presentation of the quantum K-ring QK(X).

4.1. **Ring homomorphism property.** Recall that the *K*-theoretic Kirwan map is the following isomorphism

$$R(\operatorname{GL}_r(\mathbb{C})) \to K(\mathfrak{X}), \quad \mathcal{V} \mapsto M_{r \times N} \times_{\operatorname{GL}_r(\mathbb{C})} \mathcal{V},$$

sending a representation to its associated K-theory class on X. We define the following deformation of the K-theoretic Kirwan map:

Definition 4.1. The quantum reduction map $\kappa : R(\operatorname{GL}_r(\mathbb{C})) \otimes \mathbb{Z}[[q]] \to \operatorname{QK}(X)$ is defined to be the $\mathbb{Z}[[q]]$ -linear extension of the formula

(17)
$$\kappa(\mathbf{V}) = \sum_{\alpha \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\alpha}^* \rangle\!\rangle_0^{\mathsf{Quot}} \mathcal{O}_{\alpha},$$

where V, by abuse of notation, also denotes the K-theory class on \mathfrak{X} associated to $V \in R(\operatorname{GL}_r(\mathbb{C}))$.

Proposition 4.2. For any $V \in \mathcal{R}^{r,k}$, we have $\kappa(V) = V$ where

$$V = M_{r \times N}^{ss} \times_{\mathrm{GL}_r(\mathbb{C})} \mathbf{V} \in K(X).$$

Proof. Recall that $\mathcal{R}^{r,k}$ is spanned by $\{O_{\lambda} : \lambda \in P_{r,k}\}$. The statement follows from Proposition 3.2 and the fact that O_{λ} restricts to the class of structure sheaf \mathcal{O}_{λ} over X.

Lemma 4.3. Let $V \in R(GL_r(\mathbb{C})) \cong K(\mathfrak{X})$ be the associated K-theory class. We have

$$\langle \kappa(\mathrm{V}), \mathrm{W} \rangle _{0}^{\mathsf{Quot}} = \langle \! \langle \mathrm{V}, \mathrm{W}

 \rangle _{0}^{\mathsf{Quot}}$$

for any $W \in K(\mathfrak{X})$. Here we extend $\kappa(V)$ to a class in $\mathbb{Z}[[q]] \otimes K(\mathfrak{X})$ by replacing \mathcal{O}_{ν} by O_{ν} in (17).

Proof. It follows from the definition of κ (see (17)) and the identity (16) that

$$\langle\!\langle \kappa(\mathbf{V}), \mathbf{W} \rangle\!\rangle_0^{\mathsf{Quot}} = \sum_{\alpha \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\alpha}^* \rangle\!\rangle_0^{\mathsf{Quot}} \langle\!\langle \mathbf{O}_{\alpha}, \mathbf{W} \rangle\!\rangle_0^{\mathsf{Quot}} = \langle\!\langle \mathbf{V}, \mathbf{W} \rangle\!\rangle_0^{\mathsf{Quot}}$$

Theorem 4.4. The quantum reduction map κ is a surjective ring homomorphism.

Proof. It is clear from the definition that κ is surjective and additive. Let $V, W \in R(GL_r(\mathbb{C})) \cong K(\mathfrak{X})$, then by the definition of the quantum product, Theorem 1.2, and repeatedly using the

identity in (16), we obtain

$$\begin{split} \kappa(\mathbf{V}) \bullet \kappa(\mathbf{W}) &= \sum_{\alpha,\beta \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\alpha}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} \cdot \langle\!\langle \mathbf{W}, \mathbf{O}_{\beta}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} \mathcal{O}_{\alpha} \bullet \mathcal{O}_{\beta} \\ &= \sum_{\alpha,\nu \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\alpha}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} \cdot \langle\!\langle \mathbf{W}, \mathbf{O}_{\beta}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} \langle\!\langle \mathbf{O}_{\alpha}, \mathbf{O}_{\beta}, \mathbf{O}_{\nu}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} \mathcal{O}_{\nu} \\ &= \sum_{\alpha,\beta,\nu \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\alpha}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} \cdot \langle\!\langle \mathbf{W}, \mathbf{O}_{\alpha}, \mathbf{O}_{\nu}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} \mathcal{O}_{\nu} \\ &= \sum_{\nu \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{W}, \mathbf{O}_{\nu}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} \mathcal{O}_{\nu} \\ &= \kappa(\mathbf{V} \cdot \mathbf{W}). \end{split}$$

Remark 4.5. Let G be a complex reductive group and let Z be a smooth projective or convex quasi-projective polarized G-variety. In [GW], González and Woodward used K-theoretic affine gauged Gromov-Witten invariants to define the linearized quantum Kirwan map

$$D_{\alpha}\kappa_Z^G: \mathrm{QK}_G(Z) \to \mathrm{QK}(Z \not / G)$$

for any α in the G-equivariant quantum K-ring QK_G(Z). They showed that this map is a ring homomorphism preserving the (big) quantum K-products. We expect the non-equivariant limit of our reduction map κ to coincide with $D_0 \kappa_{M_r \times N}^{\mathrm{GL}_r(\mathbb{C})}$ in the case of Grassmannian and leave it for future investigation.

Remark 4.6. Consider the extended representation ring $R(\operatorname{GL}_r(\mathbb{C})) \otimes \mathbb{Z}[[q]]$, equipped with the pairing

(18)
$$V \otimes W \mapsto \langle\!\langle V, W \rangle\!\rangle_0^{\mathsf{Quot}} \quad V, W \in R(\mathrm{GL}_r(\mathbb{C})).$$

Note that this does not define a standard Frobenius algebra because the representation ring $R(\operatorname{GL}_r(\mathbb{C}))$ is infinite-dimensional and the pairing (18) is degenerate according to Lemma 4.3. It follows from Lemma 4.3 and Corollary 1.13 that the quantum reduction map κ preserves the pairings, i.e.,

$$\langle\!\langle \mathbf{V}, \mathbf{W} \rangle\!\rangle_0^{\mathsf{Quot}} = \langle\!\langle \kappa(\mathbf{V}), \kappa(\mathbf{W}) \rangle\!\rangle_{0,2}$$

for $V, W \in R(GL_r(\mathbb{C}))$.

4.2. Ring presentation for QK(X). The product structure in quantum K-ring QK(X) can be derived from the ring structure of the representation ring $R(GL_r(\mathbb{C}))$, i.e., Littlewood-Richardson rule, and the quantum reduction map κ defined in Section 4:

$$\mathcal{O}_{\lambda} \bullet \mathcal{O}_{\mu} = \kappa(\mathcal{O}_{\lambda}) \bullet \kappa(\mathcal{O}_{\mu}) = \sum_{\nu \in \mathcal{P}_{r,2k}} c_{\lambda,\mu}^{\nu} \kappa(\mathcal{O}_{\nu}),$$

where $c_{\lambda,\mu}^{\nu}$ are the K-theoretic Littlewood-Richardson coefficients. Using this strategy, we obtain a close formula for the structure constants of QK(Gr(2, N)) in [SZ].

We now use our technique to give an alternate proof of the quantum K-ring relations in [GMSZ2]. Let $0 \to S \to \mathcal{O}_X^{\oplus N} \to Q \to 0$ denote the universal sequence on $\operatorname{Gr}(r, N)$. Note that $\det(Q) = \det(S^{\vee})$. Using the classical orthogonality in (15) and the fact that $\chi(X, \mathcal{O}_{\nu}) = 1$ for all $\nu \in \mathbb{P}_{r,k}$, one can show that

(19)
$$\det(Q) = \sum_{\lambda \in \mathcal{P}_{r,k}} \mathcal{O}_{\lambda} \in K(X).$$

Proposition 4.7.

$$\kappa(\det(\mathbf{S}^{\vee})) = \frac{\det(Q)}{1-q}$$

Proof. For each $\lambda \in P_{r,k}$,

$$\langle\!\langle \det(\mathbf{S}^{\vee}), \mathbf{O}_{\lambda}^{*} \rangle\!\rangle_{0}^{\mathsf{Quot}} = \langle\!\langle \det(\mathbf{S}^{\vee}) \cdot \mathbf{O}_{\lambda^{*}} \cdot \det(\mathbf{S}) \rangle\!\rangle_{0}^{\mathsf{Quot}} = \langle\!\langle \mathbf{O}_{\lambda^{*}} \rangle\!\rangle_{0}^{\mathsf{Quot}} = \frac{1}{1-q}.$$

Then the proposition follows from (19) and the definition of κ .

Recall that $\mathcal{R}^{r,\ell}$ denotes the subspace of the representation ring $R(\operatorname{GL}_r(\mathbb{C}))$ spanned by $\{\mathbb{S}^{\lambda}(S) \mid \lambda \in P_{r,\ell}\}$. We will use the property (14) of the Littlewood-Richardson rule several times below.

Corollary 4.8. For $0 < i \le r$, the quantum K-product

$$\wedge^{i} S \bullet \det Q = (1 - q) \wedge^{i} S \cdot \det Q.$$

Proof. Using Proposition 4.7,

$$\wedge^{i} S \bullet \det Q = \kappa(\wedge^{i} S) \bullet (1-q)\kappa(\det(S^{\vee})) = (1-q)\kappa(\wedge^{i} S \cdot \det(S^{\vee})).$$

To show that $\kappa(\wedge^i \mathbf{S} \cdot \det(\mathbf{S}^{\vee})) = \wedge^i \mathbf{S} \cdot \det(\mathbf{S})^{-1}$, it suffices to prove that

$$\langle \wedge^i \mathbf{S}, \det(\mathbf{S}^{\vee}), \mathbf{O}_{\nu}^* \rangle_{0,d}^{\mathsf{Quot}} = \langle \wedge^i \mathbf{S}, \mathbf{O}_{\nu^*} \rangle_{0,d}^{\mathsf{Quot}} = 0.$$

for each $\nu \in P_{r,k}$, $d \ge 1$. Note that $O_{\nu^*} \in \mathcal{R}^{r,k}$, thus $\wedge^i S \cdot O_{\nu^*} \in \mathcal{R}^{r,k+1}$. Part (i) of Theorem 1.11 implies that $\langle V \rangle_{0,d}^{\mathsf{Quot}} = 0$ for any $V \in \mathcal{R}^{r,k+1}$ and $d \ge 1$, thus completing the proof.

We can express $\wedge^j Q$ in the Schur basis for K(X) by extracting the coefficient of y^j in

$$\wedge_y Q = \frac{\wedge_y(\mathcal{O}_X^{\oplus N})}{\wedge_y S} = \frac{(1+y)^N}{1+yS+y^2\wedge^2 S+\cdots y^r\wedge^r S}.$$

This coefficient is a polynomial in the exterior powers of S of degree at most j. Proposition 4.2 and (14) imply that there exists a unique class $\mathbf{H}_j \in \mathcal{R}^{r,j}$ such that $\kappa(\mathbf{H}_j) = \wedge^j Q$.

Corollary 4.9 ([GMSZ1]). In QK(X),

(20)
$$\wedge_y S \bullet \wedge_y Q = (1+y)^N - qy^k (\wedge_y S - 1) \cdot \det Q.$$

Proof. For j < k and any i,

$$\wedge^{i} S \bullet \wedge^{j} Q = \kappa(\wedge^{i} S \cdot H_{j}) = \wedge^{i} S \cdot \wedge^{j} Q.$$

Here the last equality is the consequence of Proposition 4.2 and observing that $\wedge^i \mathbf{S} \cdot \mathbf{H}_j \in \mathcal{R}^{r,j+1} \subset \mathcal{R}^{r,k}$. When j = k, Corollary 4.8 implies

$$\wedge_y S \bullet \det Q = \wedge_y S \cdot \det Q - q(\wedge_y S - 1) \cdot \det Q.$$

Combining the above two cases, we obtain the required identity (20).

Corollary 4.10 ([GMSZ2]). In QK(X), the 'quantum K-Whitney' relations are satisfied:

$$\wedge_y S \bullet \wedge_y Q = (1+y)^N - \frac{q}{1-q} y^k (\wedge_y S - 1) \bullet \det Q$$

Proof. Corollary 4.8 is equivalent to [GMSZ2, Theorem 1.3], and it states that $(\wedge_y S - 1) \bullet \det Q = (1 - q)(\wedge_y S - 1) \cdot \det Q$. The result then follows from Corollary 4.9.

4.3. Euler characteristic. Let $\chi^q : QK(X) \to \mathbb{Z}[[q]]$ denote the $\mathbb{Z}[[q]]$ -linear extension of the Euler characteristic that sends each Schubert structure sheaf \mathcal{O}_{λ} to $\chi(X, \mathcal{O}_{\lambda}) = 1$.

Proposition 4.11. Let $V \in R(GL_r(\mathbb{C})) \cong K(\mathfrak{X})$ be the associated K-theory class. We have

$$\langle\!\langle \mathbf{V} \rangle\!\rangle_0^{\mathsf{Quot}} = \frac{1}{1-q} \chi^q \left(X, \kappa(\mathbf{V}) \right)$$

Proof. It follows from Lemma 4.3 that $\langle\!\langle \mathbf{V} \rangle\!\rangle_0^{\mathsf{Quot}} = \langle\!\langle \kappa(V) \rangle\!\rangle_0^{\mathsf{Quot}}$. If $\lambda \in \mathbf{P}_{r,k}$, then

$$\langle\!\langle \mathcal{O}_{\lambda} \rangle\!\rangle_0^{\mathsf{Quot}} = \frac{1}{1-q} = \frac{1}{1-q} \chi(X, \mathcal{O}_{\lambda}).$$

We conclude the proof by noting that $\kappa(V)$ is a linear combination of $O_{\lambda}, \lambda \in P_{r,k}$ with coefficients in $\mathbb{Z}[[q]]$.

Remark 4.12. Consider the image of the quantized dual O_{λ}^* under the quantum reduction map $\kappa(O_{\lambda}^*) = \kappa(\det(S) \cdot O_{\lambda^*}) = \det(S) \bullet \mathcal{O}_{\lambda^*}$. It was observed in [BM, Theorem 5.15] that

$$\frac{\det(S)}{1-q} \bullet \mathcal{O}_{\nu^*}$$

is the $\mathbb{Z}[[q]]$ -linear dual to \mathcal{O}_{λ} with respect to χ^{q} . Indeed,

$$\frac{1}{1-q}\chi^q(\mathcal{O}_{\lambda} \bullet \det(S) \bullet \mathcal{O}_{\nu^*}) = \frac{1}{1-q}\chi^q\big(\kappa(\mathcal{O}_{\lambda} \cdot \mathcal{O}_{\nu}^*)\big) = \langle\!\langle \mathcal{O}_{\lambda} \cdot \mathcal{O}_{\nu}^* \rangle\!\rangle_0^{\mathsf{Quot}} = \delta_{\lambda,\nu}.$$

This can be viewed as a manifestation of the orthogonality relation in Proposition 3.2.

5. Quasimap wall-crossing

In this section, we prove that the quasimap invariants are independent of the stability parameter ϵ . The proof relies on the K-theoretic ϵ -wall-crossing formula in [ZZ, Theorem 1.3]. In particular, we show that the correction term vanishes by calculating a certain *I*-function.

5.1. K-theoretic ϵ -stable quasimap invariants.

Definition 5.1. An n-pointed, genus-zero quasimap of degree d to X = Gr(r, N) is a tuple

 $((C, \mathbf{p}), E, s)$

where

- (i) $(C, \mathbf{p}) = (C, p_1, \dots, p_n)$ is a prestable n-pointed curve, i.e., a connected, at most nodal, projective curve of genus-zero, together with n distinct and nonsingular markings on it.
- (ii) E is a vector bundle on C of rank r and degree d.
- (iii) $s \in H^0(E \otimes \mathcal{O}_C^{\oplus N})$ represents N sections which generically generate E.

A point $p \in C$ is called a *base point* if s does not span the fiber of E at p. For any $\epsilon \in \mathbb{Q}_{>0}$, let $Q_{0,n}^{\epsilon}(X,d)$ be the moduli stack of ϵ -stable quasimaps to X introduced in [Tod, CFKM]. It parametrizes tuples $((C, \mathbf{p}), E, s)$ of quasimaps satisfying the following

- (1) The base points of s are away from the nodes and the marked points.
- (2) $\det(E)^{\epsilon} \otimes \omega_C(\sum_{i=1}^n p_i)$ is ample,
- (3) for any point $p \in C$, the torsion subsheaf $\tau(Q)$ of $Q := \operatorname{coker}(E^{\vee} \xrightarrow{s^{\vee}} \mathcal{O}_{C}^{\oplus N})$ satisfies

$$\epsilon \cdot \operatorname{length} \tau(Q)_p \leq 1.$$

According to Theorem 1.1 and Lemma 2.15 of [Tod], the moduli stack $Q_{0,n}^{\epsilon}(X,d)$ is a smooth proper DM stack over \mathbb{C} . It is empty unless $-2 + n + \epsilon d > 0$. For a fixed degree d, the space $\mathbb{Q}_{>0}$ is divided into stability chambers by finitely many walls $\{1/d' \mid d \in \mathbb{Z}_{>0}, d' \leq d\}$. There are two extreme chambers $(2, \infty)$ and (0, 1/d], in which the stability conditions are denoted by $\epsilon = \infty$ and $\epsilon = 0+$, respectively. In these two extreme chambers, we recover the moduli spaces of stable maps and *stable quotients*, respectively, i.e., we have

$$\begin{aligned} Q_{0,n}^{\epsilon}(X,d) &\cong \overline{M}_{0,n}(X,d), \quad \epsilon > 2, \\ Q_{0,n}^{\epsilon}(X,d) &\cong Q_{0,n}^{\epsilon=0+}(X,d), \quad 0 < \epsilon \le 1/d \end{aligned}$$

Here $Q_{0,n}^{\epsilon=0+}(X,d)$ is the moduli stack of stable quotients introduced in [MOP].

Let $\operatorname{ev}_i : Q_{0,n}^{\epsilon}(X,d) \to X, 1 \leq i \leq n$ be the evaluation maps. The diagonal *T*-action on $\mathcal{O}_C^{\oplus N}$ induces a *T*-action on $Q_{0,n}^{\epsilon}(X,d)$.

Definition 5.2. We define the *T*-equivariant *K*-theoretic ϵ -stable quasimap invariant with insertions $E_1, \ldots, E_n \in K_T^0(X)$ by

$$\langle E_1,\ldots,E_n\rangle_{0,n,d}^{\epsilon,T} := \chi^T \left(Q_{0,n}^\epsilon(X,d),\prod_{i=1}^n \operatorname{ev}_i^*(E_i)\right).$$

5.2. ϵ -wall-crossing.

Proposition 5.3. Suppose $n \ge 3$ or n = 2 and d > 0. Then the *T*-equivariant *K*-theoretic ϵ -stable quasimap invariants are independent of ϵ , i.e.,

$$\langle E_1, \dots, E_n \rangle_{0,n,d}^{\epsilon,T} = \langle E_1, \dots, E_n \rangle_{0,n,d}^{\epsilon',T}$$

for any $\epsilon, \epsilon' \in \mathbb{Q}_{>0}$.

Proof. Let **t** be a generic element in $K_T^0(X)$. There is a natural S_n -action on $Q_{0,n}^{\epsilon}(X,d)$ defined by permuting the *n* marked points. Here S_n denotes the symmetric group on *n* letters. Consider the following (virtual) $(T \times S_n)$ -module

$$[\mathbf{t}, \dots, \mathbf{t}]_{0,n,d}^{\epsilon,T} := \sum_{j \ge 0} (-1)^j H^j \left(Q_{0,n}^{\epsilon}(X,d), \prod_{i=1}^n \mathrm{ev}_i^*(\mathbf{t}) \right).$$

Let $\epsilon = 1/d_0$ be a wall, where $d_0 \leq d$. Let $\epsilon_- < \epsilon_+$ be the stability conditions in the two adjacent chambers separated by ϵ_0 . Set $m = \lfloor d/d_0 \rfloor$. By the K-theoretic ϵ -wall-crossing formula [ZZ, Theorem 1.3],¹ we have

(21)
$$[\mathbf{t}, \dots, \mathbf{t}]_{0,n,d}^{\epsilon_{-},T} - [\mathbf{t}, \dots, \mathbf{t}]_{0,n,d}^{\epsilon_{+},T} = \sum_{j=1}^{m} [\mathbf{t}, \dots, \mathbf{t}, \mu_{d_0}(L), \dots, \mu_{d_0}(L)]_{0,n+j,d-jd_0}^{S_j,\epsilon_{+},T}$$

as virtual $(T \times S_n)$ -modules. Here

$$[\mathbf{t},\ldots,\mathbf{t},\mu_{d_0}(L),\ldots,\mu_{d_0}(L)]_{0,n+j,d-jd_0}^{S_j,\epsilon_+,T} := p_* \bigg(\prod_{i=1}^n \mathrm{ev}_i^*(\mathbf{t}) \cdot \prod_{a=1}^j \mathrm{ev}_{n+a}^*(\mu_{d_0}(\omega))|_{\omega=L_{n+a}}\bigg),$$

where $\mu_{d_0}(\omega)$ is a Laurent polynomial in ω defined by (22), L_{n+a} is the cotangent line bundle at the (n + a)-th marked point, and

$$p: [Q_{0,n+j}^{\epsilon_+}(X, d-jd_0)/(T \times S_j))] \to BT.$$

By Lemma 5.5, we have $\mu_d(\omega) = 0$, and therefore the right-hand side of (21) vanishes. Hence $[\mathbf{t}, \ldots, \mathbf{t}]_{0,n,d}^{\epsilon,T}$ is independent of ϵ .

Now set $\mathbf{t} = \sum_{i=1}^{n} \alpha_i E_i$ where α_i 's are formal variables. Then the coefficient of $\prod_{i=1}^{n} \alpha_i$ in the character of the virtual *T*-module $[\mathbf{t}, \ldots, \mathbf{t}]_{0,n,d}^{\epsilon,T}$ is given by

$$n! \cdot \langle E_1, \dots, E_n \rangle_{0,n,d}^{\epsilon,T} \in \Gamma.$$

This concludes the proof.

5.3. K-theoretic *I*-function and its truncations. To define the Laurent polynomial $\mu_d(\omega)$, we refer to the quasimap graph space $QG_{0,n}^{\epsilon=0+}(X,d)$ as introduced in [CFKM, §7.2], along with the \mathbb{C}^* -action on $QG_{0,n}^{\epsilon=0+}(X,d)$ induced from a \mathbb{C}^* -action on \mathbb{P}^1 . The relevant calculations of the fixed loci are detailed in [BCFK] and [RZ2, Appendix]. We will summarize these calculations to prove the vanishing of $\mu_d(\omega)$ as outlined in Lemma 5.5.

Definition 5.4. The $(\epsilon = 0+)$ -stable quasimap graph space, denoted $QG_{0,n}^{\epsilon=0+}(X,d)$, parameterizes tuples

$$((C, \mathbf{p}), E, s, \varphi)$$

where $\varphi : C \to \mathbb{P}^1$ is a degree one map, $((C, \mathbf{p}), E, s)$ is an n-pointed genus-zero quasimap of degree d to X, and it satisfies the following stability conditions:

 $^{^{1}}$ [ZZ, Theorem 1.3] is stated in the non-torus-equivariant setting, but it holds true in general and the proof is verbatim.

- (i) The base points of s are away from the nodes and the marked points.
- (ii) The line bundle

$$\det(E)^{\epsilon} \otimes \omega_C \big(\sum_{i=1}^n p_i\big) \otimes \varphi^*(\omega_{\mathbb{P}^1}^{-1} \otimes M),$$

is ample for every sufficiently small rational numbers $0 < \epsilon \ll 1$, where M is any ample line bundle on \mathbb{P}^1 .

Let $\pi : \mathcal{C} \to QG_{0,n}^{\epsilon=0+}(X,d)$ be the universal curve and let

$$0 \to \mathcal{S} \to \mathcal{O}_{\mathcal{C}}^{\oplus N} \to \mathcal{Q} \to 0$$

be the universal exact sequence over \mathcal{C} . Here $\mathcal{S} = \mathcal{E}^{\vee}$, where \mathcal{E} is the universal vector bundle. dle. The higher direct image $R^1\pi_*(\mathcal{H}om(\mathcal{S},\mathcal{Q}))$ vanishes because over each component of a geometric fiber C of π , the bundle $\mathcal{H}om(\mathcal{S},\mathcal{Q})|_C$ splits into a direct sum of line bundles of nonnegative degrees. Hence $\pi_*(\mathcal{H}om(\mathcal{S},\mathcal{Q}))$ is a vector bundle on $QG_{0,n}^{\epsilon=0+}(X,d)$. Let $\Phi: \mathcal{C} \to \mathbb{P}^1$ be the universal parametrization map and let $\mathcal{D}_i \subset \mathcal{C}$ be the divisor corresponding to the *i*-th marked point. According to the deformation theory of quasimaps (see, e.g., [RZ2, §6.2], [MOP, §3.2], [Tod, §2.4]), the moduli stack $QG_{0,n}^{\epsilon=0+}(X,d)$ is smooth, and its tangent bundle fits into the following exact sequence

$$\begin{aligned} 0 \to \pi_* \bigg(\Omega^{\vee}_{\pi} \bigg(-\sum_{i=1}^n \mathcal{D}_i \bigg) \bigg) \to \pi_* \left(\mathcal{H}om(\mathcal{S}, \mathcal{Q}) \right) \oplus \pi_* (\Phi^*(T_{\mathbb{P}^1})) \\ \to T_{QG_{0,n}^{\epsilon=0+}(X,d)} \to R^1 \pi_* \bigg(\Omega^{\vee}_{\pi} \bigg(-\sum_{i=1}^n \mathcal{D}_i \bigg) \bigg) \to 0. \end{aligned}$$

Let $T = (\mathbb{C}^*)^N$ be the diagonal torus acting on $QG_{0,n}^{\epsilon=0+}(X,d)$. Consider the \mathbb{C}^* -action on \mathbb{P}^1 defined by

$$t[\zeta_0,\zeta_1] = [t\zeta_0,\zeta_1], \quad t \in \mathbb{C}^*.$$

This induces a \mathbb{C}^* -action on $QG_{0,1}^{\epsilon=0+}(X,d)$ which commutes with the *T*-action. Let ω denote the character of this \mathbb{C}^* -action. We denote the unique marking by x_{\bullet} . Set 0 := [1 : 0] and $\infty := [0 : 1]$. Let $F_d \subset QG_{0,1}^{\epsilon=0+}(X,d)$ be the distinguished fixed-point component in $QG_{0,1}^{\epsilon=0+}(X,d)$ parametrizing \mathbb{C}^* -fixed quasimaps such that x_{\bullet} lies at ∞ and the only base point of s is at 0.

The graph space $QG_{0,1}^{\epsilon=0+}(X,d)$ and its distinguished \mathbb{C}^* -fixed-point component F_d are both smooth. The normal bundle $N := N_{F_d/QG_{0,1}^{\epsilon=0+}(X,d)}$ has two parts

- (i) a rank-1 part coming from deforming x_{\bullet} away from $\infty \in \mathbb{P}^1$, whose K-theoretic Euler class is $1 \omega^{-1}$;
- (ii) the moving part of the vector bundle $\pi_*(\mathcal{H}om(\mathcal{S}, \mathcal{Q}))$, which we denote by $N^{\mathrm{rel}} := N^{\mathrm{rel}}_{F_d/QG_{0,1}^{\epsilon=0+}(X,d)}$.

Let $\operatorname{ev}_{\bullet} : QG_{0,1}^{\epsilon=0+}(X,d) \to X \times \mathbb{P}^1$ be the evaluation map at the marking x_{\bullet} . Define $\widetilde{\operatorname{ev}}_{\bullet} := \pi_X \circ \operatorname{ev}_{\bullet}$, where $\pi_X : X \times \mathbb{P}^1 \to X$ is the projection. For d > 0, we define

$$\tilde{I}_d(\omega) := (1-\omega)(1-\omega^{-1})(\tilde{\operatorname{ev}}_{\bullet})_* \left(\frac{1}{\wedge_{-1}^{T\times\mathbb{C}^*}(N^{\vee})}\right) = (1-\omega)(\tilde{\operatorname{ev}}_{\bullet})_* \left(\frac{1}{\wedge_{-1}^{T\times\mathbb{C}^*}((N^{\operatorname{rel}})^{\vee})}\right).$$

This is a rational function in ω with coefficients in $K^0_T(X)$. We define

(22)
$$\mu_d(\omega) := \left[\tilde{I}_d(\omega)\right]_+$$

to be the Laurent polynomial part in the partial fraction decomposition of the rational function $\tilde{I}_d(\omega)$.

The distinguished \mathbb{C}^* -fixed-point component F_d is explicitly identified in [BCFK] (see also [RZ2]). The irreducible components of F_d are indexed by partitions $(d) := (d_1, d_2, \ldots, d_r)$ of d satisfying

$$\sum d_i = d$$
 and $0 \le d_1 \le d_2 \le \dots \le d_r$.

Let $\{r_i\}_{1 \le i \le j}$ be the jumping index of $\{d_i\}$, i.e., $0 \le d_1 = \cdots = d_{r_1} < d_{r_1+1} = \cdots = d_{r_2} < \cdots < d_{r_j+1} = \cdots = d_r$. The (restriction of the) evaluation map $\widetilde{\text{ev}}_{\bullet} : F_d \to X$ can be identified with the flag bundle map

$$\rho: \operatorname{Fl}(r_1, \ldots, r_j; S) \to X,$$

where $Fl(r_1, \ldots, r_i; S)$ denote the relative flag variety of type $\{r_i\}$. Let

$$0 \subset S_{r_1} \subset S_{r_2} \subset \cdots \subset S_{r_j} \subset S_{r_{j+1}} = \rho^* S$$

be the universal flag on $Fl(r_1, \ldots, r_j; S)$. Using the splitting principle, we write

$$\sum_{s=1}^{r_a - r_{a-1}} \mathcal{L}_{r_{a-1} + s} = (S_{r_a} / S_{r_{a-1}})^{\vee}.$$

Set $\bar{r}_a := r_a - r_{a-1}$ and $d_{ba} := d_{r_b} - d_{r_a}$ with $r_0 = 0$.

According to the computations detailed in [RZ2, Appendix A], which build on the analysis in [BCFK], the following explicit formula is derived:

(23)
$$\tilde{I}_d(\omega) = \sum_{(d)} \sum_{\sigma} (1-\omega) \cdot \sigma(I_{(d)})$$

where the permutation σ runs through all elements in $S_r/(S_{r_1} \times \cdots \times S_{r_{j+1}})$ and acts on the indices of \mathcal{L}_i 's in

$$\begin{split} I_{(d)} &:= \prod_{1 \le a < b \le j+1} \prod_{\substack{1 \le s \le \bar{r}_a \\ 1 \le t \le \bar{r}_b}} \prod_{c=1}^{d_{ba}-1} \mathcal{L}_{r_{b-1}+t}^{\vee} \mathcal{L}_{r_{a-1}+s} \omega^c \\ &\cdot \prod_{1 \le a < b \le j+1} \prod_{\substack{1 \le s \le \bar{r}_a \\ 1 \le t \le \bar{r}_b}} (-1)^{\bar{r}_a \bar{r}_b (d_{ba}-1)} \frac{1 - \mathcal{L}_{r_{b-1}+t}^{\vee} \mathcal{L}_{r_{a-1}+s} \omega^{d_{ba}}}{1 - \mathcal{L}_{r_{b-1}+t}^{\vee} \mathcal{L}_{r_{a-1}+s}} \\ &\cdot \frac{1}{\prod_{a=1}^{j+1} \prod_{s=1}^{\bar{r}_a} \prod_{b=1}^{d_{r_a}} \prod_{e=1}^{N} (1 - \mathcal{L}_{r_{a-1}+s}^{\vee} \alpha_e^{-1} \omega^b)}. \end{split}$$

The substitution of $\prod_{k=1}^{N} (1 - \mathcal{L}_{r_{a-1}+s}^{\vee} \alpha_k^{-1} q^b)$ in place of $(1 - \mathcal{L}_{r_{a-1}+s}^{\vee} q^b)^N$ in the non-torusequivariant *I*-function on [RZ2, p. 300] corresponds to taking into account the *T*-action on the trivial bundle \mathbb{C}^N in the Euler sequence on page 298 of [RZ2].

Lemma 5.5. The rational function $\tilde{I}_d(\omega)$ is regular at $\omega = 0$ and vanishes at $\omega = \infty$. As a consequence, we have $\mu_d(\omega) = [\tilde{I}_d(\omega)]_+ = 0$.

Proof. It is clear that $I_d(\omega)$ are regular at $\omega = 0$. To show that the rational function vanishes at $\omega = \infty$, we estimate the degrees of the numerator and denominator of the terms in (23).

We fix a partition (d) of d and a permutation σ . Since k > 0, the degree of the numerator of $(1 - \omega) \cdot \sigma(I_{(d)})$ in ω is bounded by

$$(24) \quad 1 + \sum_{1 \le a < b \le j+1} \frac{\bar{r}_a \bar{r}_b (d_{ba} - 1) d_{ba}}{2} + \sum_{1 \le a < b \le j+1} \bar{r}_a \bar{r}_b d_{ba} = 1 + \sum_{1 \le a < b \le j+1} \frac{\bar{r}_a \bar{r}_b (d_{ba} + 1) d_{ba}}{2}.$$

Note that the first summation is absent in the case r = 1. The degree of the denominators of (23) in ω is

(25)
$$\sum_{a=1}^{j+1} \frac{\bar{r}_a(d_{r_a}+1)d_{r_a}N}{2}$$

By using $\sum_{a < b} \bar{r}_a \leq r - 1$ and $0 < d_{ba} \leq d_{r_b}$, we obtain the following upper bound for the first term of the RHS of (24):

$$\sum_{1 \le a < b \le j+1} \frac{\bar{r}_a \bar{r}_b (d_{ba} + 1) d_{ba}}{2} \le \sum_{1 \le a < b \le j+1} \frac{\bar{r}_a \bar{r}_b (d_{r_b} + 1) d_{r_b}}{2} \le \frac{r-1}{2} \sum_{b=1}^{j+1} \bar{r}_b (d_{r_b} + 1) d_{r_b}.$$

It follows that the difference (25) - (24) is greater than or equal to

(26)
$$\frac{N - (r-1)}{2} \sum_{a=1}^{j+1} \bar{r}_a (d_{r_a} + 1) d_{r_a} - 1 > 0.$$

6. LIGHT-TO-HEAVY WALL CROSSING

In this section, we derive a light-to-heavy wall-crossing result for K-theoretic quasimap invariants in all genera using the tools from [RZ2, §6]. We then apply this result specifically to the genus-zero case to complete the proof of Theorem 1.12.

6.1. K-theoretic (0+)-stable graph quasimap invariants. We fix a smooth curve $(C, \mathbf{q}) = (C, q_1, \ldots, q_n)$ of genus g with n distinct markings. In this subsection, we consider quasimaps with one parametrized component isomorphic to C and their moduli space.

Definition 6.1. An *n*-pointed, genus g quasimap with one parametrized component of degree d to X = Gr(r, N) is a tuple

$$((C, \mathbf{p}), E, s, \varphi)$$

where

- (i) $((\widehat{C}, \mathbf{p}), E, s)$ is an n-pointed, genus g quasimap of degree d to X.
- (ii) $\varphi: \widehat{C} \to C$ is a morphism of degree one such that

(27)
$$\varphi(p_i) = q_i, \quad 1 \le i \le n.$$

By definition, \widehat{C} has an irreducible component \widehat{C}_0 such that $\varphi|_{\widehat{C}_0} : \widehat{C}_0 \xrightarrow{\sim} C$ is an isomorphism, and the rest of \widehat{C} is contracted by φ . We refer to \widehat{C}_0 as the parametrized component. An isomorphism between two quasimaps $((\widehat{C}, \mathbf{p}), E, s, \varphi)$ and $((\widehat{C}', \mathbf{p}'), E', s', \varphi')$ consists of an isomorphism $f : \widehat{C} \xrightarrow{\sim} \widehat{C}'$ of the underlying curves, along with an isomorphism $\sigma : E \to f^*E'$, such that

$$f(p_i) = p'_i, \ \varphi = \varphi' \circ f, \text{ and } \sigma(s) = f^*(s').$$

Definition 6.2. A quasimap $((\widehat{C}, \mathbf{p}), E, s, \varphi)$ with one parametrized component is (0+)-stable if the following two conditions hold:

- (i) The base points of s are away from the nodes and the markings.
- (ii) The line bundle

$$\det(E)^{\epsilon} \otimes \omega_{\widehat{C}} \Big(\sum_{i=1}^{n} p_i \Big) \otimes \varphi^*(\omega_C^{-1} \otimes M),$$

is ample for every sufficiently small rational numbers $0 < \epsilon \ll 1$, where M is any ample line bundle on C.

For a (0+)-stable tuple $((\widehat{C}, \mathbf{p}), E, s, \varphi)$, the source curve \widehat{C} is obtained from the parametrized component \widehat{C}_0 by adding to q_i , for all i, a (possibly empty) chain of rational curves with p_i on the extremal components, and (E, s) induces a rational map from \widehat{C} to X that is regular at the nodes and the markings p_i . We refer the reader to [RZ2, Fig. 2] for an illustration of the source curve \widehat{C} .

Let $Q_{C,n}^{0+}(X,d)$ denote the moduli stack of (0+)-stable quasimaps with one parametrized component as defined in Definition 6.2. It forms a closed substack of the moduli stack $\operatorname{Qmap}_{g,n}(X,d;C)$ defined in [CFKM, §7.2], which parametrizes the data from Definition 6.2 without the condition (27). According to [CFKM, Theorem 7.2.2], $\operatorname{Qmap}_{g,n}(X,d;C)$ is a proper DM stack, implying that $Q_{C,n}^{0+}(X,d)$ is also a proper DM stack.

Let \mathcal{S} and \mathcal{Q} be the universal subbundle and quotient sheaf on the universal curve $\pi : \mathcal{C} \to Q^{0+}_{C,n}(X,d)$. The complex $R \mathcal{H}om_{\pi}(\mathcal{S}, \mathcal{Q})$ defines a (*T*-equivariant) POT for $\nu : Q^{0+}_{C,n}(X,d) \to \mathcal{T}_{maps}(C,\mathbf{q})$, where $\mathcal{T}_{maps}(C,\mathbf{q})$ denotes the smooth Artin stack parametrizing the data $((\widehat{C},\mathbf{p}),\varphi))$ satisfying 27. Define

$$\mathcal{O}^{\mathrm{vir}}_{Q^{0+}_{C,n}(X,d)} := \nu^! (\mathcal{O}_{\mathcal{T}_{\mathrm{maps}}(C,\mathbf{q})}) \in K^T_0(Q^{0+}_{C,n}(X,d)).$$

Let $ev_i : Q_{C,n}^{0+}(X,d) \to X, 1 \le i \le n$ be the evaluation morphism.

Definition 6.3. Let $E_1, \ldots, E_n \in K_T^0(X)$. We define the *T*-equivariant *K*-theoretic (0+)stable graph quasimap invariant with the insertions E_i 's by

$$\langle E_1, \dots, E_n \rangle_{g,d}^{0+,T} := \chi^{\operatorname{vir},T} \left(Q_{C,n}^{0+}(X,d), \prod_{i=1}^n \operatorname{ev}_i^*(E_i) \right).$$

6.2. Light-to-heavy wall-crossing. In this subsection, we use the setup from [RZ2, §6.2] to derive a light-to-heavy wall-crossing formula that connects the K-theoretic Quot scheme invariants with the K-theoretic (0+)-stable graph quasimap invariants. Let m be an integer such that $1 \leq m \leq n$. We relabel the first m markings as x_1, \ldots, x_m and the last n - m markings as y_1, \ldots, y_{n-m} . Set $\mathbf{x} = (x_1, \ldots, x_m)$ and $\mathbf{y} = (y_1, \ldots, y_{n-m})$.

Definition 6.4. A quasimap $((\widehat{C}, \mathbf{x}; \mathbf{y}), E, s, \varphi)$ with one parametrized component to X is called (0+, 0+)-stable with (m, n-m)-weighted markings and degree d if :

- (i) The quasimap only has finitely many base points which are away from the nodes.
- (ii) x_1, \ldots, x_m are not base points of s.
- (iii) The \mathbb{Q} -line bundle

$$\det(E)^{\epsilon} \otimes \omega_{\widehat{C}} \left(\sum_{i=1}^{m} x_i + \epsilon' \sum_{j=1}^{n-m} y_j \right) \otimes \varphi^*(\omega_C^{-1} \otimes M)$$

is relatively ample for every sufficiently small rational numbers $0 < \epsilon, \epsilon' \ll 1$. Here M is any ample line bundle on C.

(iv) $\operatorname{rank}(E) = r \text{ and } \deg(E) = d.$

We refer to x_i as the *heavy* markings and y_i as the *light* markings. Note that each x_i is connected to the parametrized component $\widehat{C}_0 \cong C$ through a (possibly empty) chain of \mathbb{P}^1 's, and y_i is located on \widehat{C}_0 and corresponds to q_{m+i} . Let $Q_{C,m|n-m}^{0+,0+}(X,d)$ denote the moduli stack of (0+,0+)-stable quasimaps with one parametrized component to X with (m,n-m)-weighted markings. It follows from the definition that

(28)
$$Q_{C,0|n}^{0+,0+}(X,d) \cong \mathsf{Quot}_d(C,N,r), \quad Q_{C,n|0}^{0+,0+}(X,d) = Q_{C,n}^{0+}(X,d).$$

Similar to $Q_{C,n}^{0+}(X,d)$, the moduli stack $Q_{C,m|n-m}^{0+,0+}(X,d)$ is a proper DM stack and equipped with a (*T*-equivariant) virtual structure sheaf $\mathcal{O}_{Q_{C,m|n-m}^{0+,0+}(X,d)}^{\text{vir}}$. There are two types of evaluation morphisms:

$$\operatorname{ev}_{i}: Q^{0+,0+}_{C,m|n-m}(X,d) \to X, \quad i = 1, \dots, m,$$

 $\widetilde{\operatorname{ev}}_{j}: Q^{0+,0+}_{C,m|n-m}(X,d) \to \mathfrak{X} = [M_{r \times N}/\operatorname{GL}_{r}(\mathbb{C})], \quad j = 1, \dots, n-m.$

Let $V_1, \ldots, V_m, W_1, \ldots, W_{n-m}$ be (virtual) $\operatorname{GL}_r(\mathbb{C})$ -modules. We define their associated classes in $K_T^0(\mathfrak{X})$ by $M_{r \times N} \times_{\operatorname{GL}_r(\mathbb{C})} V_i$ and $M_{r \times N} \times_{\operatorname{GL}_r(\mathbb{C})} W_j$, which are still denoted by V_i and W_j by convention. Let \mathcal{V}_i and \mathcal{W}_j denote the restrictions of the above classes to X. **Definition 6.5.** We define the (m, n - m)-weighted invariant by

$$\langle \mathcal{V}_1, \cdots \mathcal{V}_m | \mathbf{W}_1, \dots, \mathbf{W}_{n-m} \rangle_{g,d,m|n-m}^{0+,T} := \chi^{\operatorname{vir},T} \left(Q_{C,m|n-m}^{0+,0+}(X,d), \prod_{i=1}^m \operatorname{ev}_i^*(\mathcal{V}_i) \cdot \prod_{j=1}^{n-m} \widetilde{\operatorname{ev}}_j^*(\mathbf{W}_j) \right).$$

The above invariant is a special case of the one in [RZ2, Definition 6.12], specifically when $e^* = 0$ and l = 0. Therefore, we can use the same proof to establish the following slight variation of [RZ2, Theorem 6.13]:

Proposition 6.6. Suppose that $W_1 \in \mathcal{R}^{r,k}$. We have

$$\langle \mathcal{V}_1, \cdots, \mathcal{V}_m | \mathbf{W}_1, \dots, \mathbf{W}_{n-m} \rangle_{g,d,m|n-m}^{0+,T} = \langle \mathcal{V}_1, \cdots, \mathcal{V}_m, \mathcal{W}_1 | \mathbf{W}_2 \dots, \mathbf{W}_{n-m} \rangle_{g,d,m+1|n-m-1}^{0+,T}$$

Proof. The master space $\widetilde{\mathcal{M}}_{C,m|n-m}$ introduced in [RZ2, §6] has commuting T- and \mathbb{C}^* -actions. By applying the K-theoretic localization formula for the \mathbb{C}^* -action on the master space, we obtain the T-equivariant version of [RZ2, (59)] that holds for T-equivariant Euler characteristics. We use the same argument as in the proof of [RZ2, Theorem 6.13] to reduce the problem to showing $\mu_d^{W_1}(\omega) = 0$ defined in (29) for any $W_1 \in \mathcal{R}^{r,k}$.

Recall the notations from Section 5.3. Consider $W = S^{\lambda}(S) \in K^{0}_{T}(\mathfrak{X})$ for an integer partition $\lambda \in P_{r,k}$. For d > 0, we define

$$I_d^{\mathrm{W}}(\omega) = (\widetilde{\mathrm{ev}}_{\bullet})_* \left(\frac{\mathbb{S}^{\lambda}(\mathcal{S}_0)}{\wedge_{-1}^{T \times \mathbb{C}^*}((N^{\mathrm{rel}})^{\vee})} \right),$$

where S_0 is the restriction of S to $F_d \times \{0\}$. Let

(29)
$$\mu_d^{\mathrm{W}}(\omega) = [I_d^{\mathrm{W}}(\omega)]_+$$

be the Laurent polynomial part in the partial fraction decomposition of $I_d^W(\omega)$. For a general $W \in \mathcal{R}^{r,k}$, $\mu_d^W(\omega)$ can be defined using linearity.

Moreover, the explicit localization calculation in [RZ2, Appendix A] shows that

$$I_d^{\mathrm{W}}(\omega) = \sum_{(d)} \sum_{\sigma} \sigma(\mathbb{S}^{\lambda}(\mathcal{K}_0^{(d)}) \cdot I_{(d)}),$$

where $I_{(d)}$ is defined in (23) and

$$\mathcal{K}_{0}^{(d)} = \sum_{a=1}^{j+1} \sum_{s=1}^{\bar{r}_{a}} \mathcal{L}_{r_{a-1}+s}^{\vee} \omega^{d_{r_{a}}}$$

Lemma 6.7. For any $W \in \mathcal{R}^{r,k}$, the rational function $I_d^W(\omega)$ is regular at $\omega = 0$ and vanishes at $\omega = \infty$. As a consequence, we have $\mu_d^W(\omega) = [I_d^W(\omega)]_+ = 0$.

Proof. Given $\lambda \in P_{r,k}$, we consider the $\operatorname{GL}_r(\mathbb{C})$ -representation $W = \mathbb{S}^{\lambda}((\mathbb{C}^r)^{\vee})$. Note that $\mathbb{S}^{\lambda}(\mathcal{K}_0^{(d)})$ with $\lambda \in P_{r,k}$ is a polynomial in ω whose degree is bounded above by kd. The proof

is similar to that of Lemma 5.5, except we have to add kd for the degree of $\mathbb{S}^{\lambda}(\mathcal{K}_{0}^{(d)})$ in place of 1 corresponding to the polynomial $(1 - \omega)$ in (23). In the last step, using the observation:

$$\sum_{a=1}^{j+1} \bar{r}_a (d_{r_a} + 1) d_{r_a} \ge 2 \sum_{a=1}^{j+1} d_{r_a} = 2d,$$

we establish the analogous bound from formula (26) as:

$$\frac{(N-(r-1))}{2}\sum_{a=1}^{j+1}\bar{r}_a(d_{r_a}+1)d_{r_a}-kd \ge (N-(r-1))d-kd=d>0.$$

Thus $I_d^W(\omega)$ vanishes at $\omega = \infty$.

By repeatedly applying Proposition 6.6 and using the isomorphisms (28), we obtain

Corollary 6.8. Suppose that $V_1, \ldots, V_n \in \mathcal{R}^{r,k}$. We have

$$\langle \mathbf{V}_1, \dots, \mathbf{V}_n \rangle_{g,d}^{\mathsf{Quot},T} = \langle \mathcal{V}_1, \dots, \mathcal{V}_n \rangle_{g,d}^{0+,T}.$$

6.3. Proof of Theorem 1.12. In this subsection, we take $C = \mathbb{P}^1$.

Lemma 6.9. There is a natural T-equivariant isomorphism

$$Q_{0,3}^{\epsilon=0+}(X,d) \cong Q_{\mathbb{P}^1,3}^{0+}(X,d)$$

Proof. There is a *T*-equivariant forgetful morphism

$$\mu: Q^{0+}_{\mathbb{P}^1,3}(X,d) \to Q^{\epsilon=0+}_{0,3}(X,d)$$

forgetting the parametrization φ . Let U be a scheme and let $(\pi : \mathcal{X} \to U, p_1, p_2, p_3; \mathcal{E}; s, \varphi)$ be a U-point of $Q_{0,3}^{\epsilon=0+}(X, d)$. Then the basepoint-free line bundle $\omega_{\pi}(p_1 + p_2 + p_3)$ induces an U-morphism $\varphi : \mathcal{X} \to U \times \mathbb{P}^1$ of degree 1. Hence we obtain an U-point of $Q_{0,3}^{\epsilon=0+}(X, d)$, and therefore a morphism

$$\nu: Q_{0,3}^{\epsilon=0+}(X,d) \to Q_{\mathbb{P}^1,3}^{0+}(X,d),$$

which is inverse to μ . This finishes the proof of the lemma.

Theorem 6.10. Let $V_1, V_2, V_3 \in \mathcal{R}^{r,k}$. Then we have

$$\langle \mathcal{V}_1, \mathcal{V}_2, \mathcal{V}_3 \rangle_{0,3,d}^T = \langle \mathbf{V}_1, \mathbf{V}_2, \mathbf{V}_3 \rangle_{0,d}^{\mathsf{Quot},T}.$$

Proof. By Proposition 5.3, we have

$$\langle \mathcal{V}_1, \mathcal{V}_2, \mathcal{V}_3 \rangle_{0,3,d}^{\epsilon = \infty, T} = \langle \mathcal{V}_1, \mathcal{V}_2, \mathcal{V}_3 \rangle_{0,3,d}^{\epsilon = 0+, T}$$

and the former is equal to the quantum K-invariant $\langle \mathcal{V}_1, \mathcal{V}_2, \mathcal{V}_3 \rangle_{0,3,d}^T$. It follows from Lemma 6.9 that

$$\langle \mathcal{V}_1, \mathcal{V}_2, \mathcal{V}_3 \rangle_{0,3,d}^{\epsilon=0+,T} = \langle \mathcal{V}_1, \mathcal{V}_2, \mathcal{V}_3 \rangle_{0,d}^{0+,T}$$

By Corollary 6.8, we have

$$\langle \mathcal{V}_1, \mathcal{V}_2, \mathcal{V}_3 \rangle_{0,d}^{0+,T} = \langle \mathbf{V}_1, \mathbf{V}_2, \mathbf{V}_3 \rangle_{0,d}^{\mathsf{Quot},T}$$

This concludes the proof.

32

Remark 6.11. It is well-known that 3-pointed, genus zero Gromov–Witten invariants of Grassmannians with Schubert class insertions coincide with the corresponding invariants defined using Quot schemes [Ber2]. A proof of this fact can be found in [CF2, §4], which utilizes the Kleiman's transversality theorem and a detailed description of the boundary of the Quot scheme. The wall-crossing strategy employed in this paper bypasses the analysis of the geometric properties of the moduli spaces.

Corollary 6.12. Let $V_1, V_2, V \in \mathbb{R}^{r,k}$. Then we have

$$\langle \mathcal{V}_1, \mathcal{V}_2 \rangle_{0,2,d}^T = \langle \mathbf{V}_1, \mathbf{V}_2 \rangle_{0,d}^{\mathsf{Quot},T}, \qquad \langle \mathcal{V} \rangle_{0,1,d}^T = \langle \mathbf{V} \rangle_{0,d}^{\mathsf{Quot},T}.$$

Proof. By Definition 1.7, we have

$$\begin{split} \langle \mathbf{V}_1, \mathbf{V}_2, \mathcal{O}_{\mathfrak{X}} \rangle_d^{\mathsf{Quot}, T} = \langle \mathbf{V}_1, \mathbf{V}_2 \rangle_d^{\mathsf{Quot}, T}, \\ \langle \mathbf{V}, \mathcal{O}_{\mathfrak{X}}, \mathcal{O}_{\mathfrak{X}} \rangle_d^{\mathsf{Quot}, T} = \langle \mathbf{V} \rangle_d^{\mathsf{Quot}, T}, \end{split}$$

where $\mathcal{O}_{\mathfrak{X}}$ is the trivial line bundle on \mathfrak{X} . Similar equalities hold for quantum K-invariants due to the string equation [Lee, (22)]. Hence Theorem 6.10 implies the corollary.

7. Degeneration formula

In this section, we adapt the constructions and arguments in [LW, Qu] to prove the degeneration formulas in Theorem 1.15 for K-theoretic Quot scheme invariants.

7.1. Expanded degenerations. We consider a simple degeneration of smooth projective curves. Let $\mathcal{C} \to B$ be a projective morphism from a smooth variety \mathcal{C} to a smooth pointed curve (B,0) such that the fibers outside 0 are smooth curves of genus g, and the fiber over 0 is a union of two smooth irreducible components C_{-} and C_{+} along the node x_{node} :

$$C_{\text{deg}} = C_{-} \coprod_{x_{\text{node}}} C_{+}.$$

Let g_- and g_+ be the genera of C_- and C_+ , respectively. Then $g_- + g_+ = g$. Let x_- and x_+ be the points on C_- and C_+ corresponding to the node x_{node} .

Expanded degenerations were introduced in [Li1, Li2]. We recall the setup in [LW]. Let 0 := [1 : 0] and $\infty := [0 : 1]$ be two points on \mathbb{P}^1 . There is a universal family $\mathfrak{C} \to \mathfrak{B}$ of expanded degenerations associated to the family $\mathcal{C} \to B$, whose singular fibers are expansions of C_{deg} obtained by inserting a chain of \mathbb{P}^1 's at the node x_{node} . More precisely, we have

$$C_{\deg}[n] = C_{-} \coprod_{x_{-}=0} \underbrace{\mathbb{P}^{1} \coprod_{\infty=0} \cdots \coprod_{\infty=0} \mathbb{P}^{1} \coprod_{\infty=x_{+}} C_{+},}_{n \text{ copies}}$$

where $n \ge 0$ and we identify x_- in C_- with 0 in the first copy of \mathbb{P}^1 , identify ∞ in the *i*-th \mathbb{P}^1 with 0 in the (i + 1)-th \mathbb{P}^1 , and identify ∞ in the last \mathbb{P}^1 with x_+ in C_+ . We refer to this

chain of \mathbb{P}^1 's as a *rational bridge* (of length n). The universal family fits into the following commutative diagram



where the first horizontal map is an isomorphism on smooth fibers and, on singular fibers, contracts the chains of \mathbb{P}^1 's to the node x_{node} in C_{deg} .

Remark 7.1. For detailed definitions of the stacks \mathfrak{E} and \mathfrak{B} , denoted as \mathfrak{X} and \mathfrak{C} respectively in [LW, §2.1, §2.2], we direct the reader to that source. Note that \mathfrak{B}_0 , the fiber of $\mathfrak{B} \to B$ over 0, is independent of \mathfrak{B} and is the same as \mathfrak{T}_0 in [ACFW].

For the relative pairs (C_{-}, x_{-}) and (C_{+}, x_{+}) , the universal families of expanded degenerations (or expanded relative pairs) are $(\mathfrak{C}_{-}, \mathfrak{x}_{-}) \to \mathcal{T}$ and $(\mathfrak{C}_{+}, \mathfrak{x}_{+}) \to \mathcal{T}$, respectively. Here, the base \mathcal{T} will be explained in Remark 7.2. The fibers of $(\mathfrak{C}_{\pm}, \mathfrak{x}_{\pm}) \to \mathcal{T}$ are expansions of (C_{\pm}, x_{\pm}) obtained by attaching chains of \mathbb{P}^{1} 's of various lengths to C_{\pm} at x_{\pm} :

$$(C_{-}[n], x_{-}[n]) := C_{-} \coprod_{x_{-}=0} \underbrace{\mathbb{P}^{1} \coprod_{n \text{ copies}} \cdots \coprod_{\infty=0} \mathbb{P}^{1}}_{n \text{ copies}},$$
$$(C_{+}[n], x_{+}[n]) := \underbrace{\mathbb{P}^{1} \coprod_{\infty=0} \cdots \coprod_{\infty=0} \mathbb{P}^{1} \coprod_{\infty=x_{+}} C_{+},}_{n \text{ copies}}$$

We refer to these chains of \mathbb{P}^1 's as *rational tails*, and the above expansions are sometimes referred to as *accordions*. We also need the family $(\mathfrak{x}_-, \mathfrak{C}_{\sim}, \mathfrak{x}_+) \to \mathcal{T}_{\sim}$ of nonrigid expanded degenerations associated to the node, whose fibers are chains of \mathbb{P}^1 's of length n for $n \geq 0$:

$$(x_{-}, \mathbb{P}^{1}[n]_{\sim}, D_{+}) := \underbrace{\mathbb{P}^{1} \coprod_{\infty = 0} \cdots \coprod_{\infty = 0} \mathbb{P}^{1}}_{n \text{ copies}}.$$

Remark 7.2. The stacks $\mathcal{T}, \mathcal{T}_{\sim}$ of expanded degenerations are independent of the choices of (C_{\pm}, x_{\pm}) and the same as those in [GV, ACFW]. According to [GV] (see also [ACFW, Proposition 4.2]), the stack \mathcal{T}_{\sim} can be identified with the moduli stack $\mathfrak{M}_{0,2}^{ss}$ of semistable, genus zero curves with two markings, where nodes only separate these markings.

Remark 7.3. Our notation \mathcal{T} for expanded relative pairs aligns with that in [Qu]; see Remark 4.3 therein for distinctions from the notation used in [LW].

7.2. Good degeneration of Quot schemes. We specify the definition of stable admissible quotients and their moduli space from [LW] to our setting. According to [LW, Definition 3.1], for a closed subscheme $D \subset W$ of a noetherian scheme, a coherent sheaf on W is said to be normal to D if $\operatorname{Tor}_{1}^{\mathcal{O}_{W}}(F, \mathcal{O}_{D}) = 0$. Let $W = C_{1} \coprod_{p} C_{2}$ be the union of two smooth curves intersecting at the node p. Let Z be a scheme and let \mathcal{F} be a Z-flat family of coherent sheaves

on $W \times Z$. Suppose for any $z \in Z$, the sheaf $\mathcal{F}_z = \mathcal{F} \otimes_{\mathcal{O}_Z} \mathbb{C}(z)$ is normal to p. After restricting to an affine neighborhood of p, the map (3.2) in [LW] is an isomorphism. According to [LW, Proposition 3.5], there is an open neighborhood $U \subset W \times Z$ of $p \times Z$ such that $\mathcal{F}|_U$ is flat over U. This implies that $\mathcal{F}|_U$ is torsion-free and, therefore, locally free at $p \times Z$.

As the family $\mathfrak{C} \to \mathfrak{B}$ is represented by a projective morphism, the relative Quot scheme of this family parametrizing rank N - r degree d quotients of the rank-N trivial sheaf is an algebraic stack projective over \mathfrak{B} . In [LW], the authors constructed an open substack $\mathfrak{Quot}_d(\mathfrak{C}/\mathfrak{B}, N, r)$ of the relative Quot scheme that parametrizes *stable* quotients. By [LW, Definition 4.2], the restriction of a quotient $\phi : \mathcal{O}_{\mathfrak{C}}^{\oplus N} \to \mathcal{Q}$ on any smooth fiber of $\mathfrak{C} \to \mathfrak{B}$ is automatically stable. Over the component $C_{\text{deg}}[n]$ of the singular fibers, the restriction of ϕ is said to be stable if its automorphism group, a subgroup of the automorphism group of $C_{\text{deg}}[n]$, is finite and it is *admissible* in the sense that \mathcal{Q} is normal to all the nodes in $C_{\text{deg}}[n]$. By [MOP, Lemma 1] and the discussion in the previous paragraph, the stability condition on ϕ introduced in [LW, Definition 4.2] is equivalent to that $\mathcal{Q}|_{C_{\text{deg}}[n]}$ is locally free at all the nodes and the restriction $\mathcal{Q}|_{\mathbb{P}^1}$ to any component of the rational bridges is of positive degree. Hence, in our setting of expanded degeneration of curves, we introduce the following equivalent definition:

Definition 7.4. A \mathfrak{B} -flat family of coherent quotients $\mathcal{O}^{\oplus N} \to \mathcal{Q}$ is stable if the restriction $\mathcal{Q}|_{C_{\text{deg}}[n]}$ is locally free at the nodes and has positive degrees over all components of the rational bridge in $C_{\text{deg}}[n]$.

Let $\mathfrak{Quot}_d(\mathfrak{C}/\mathfrak{B}, N, r)$ be the moduli space of flat families of stable quotients of rank N - rand degree d over the universal family $\mathfrak{C} \to \mathfrak{B}$ of expanded degenerations of curves. It is referred to as the *good degeneration* of Quot schemes in [LW]. We refer the reader to §4.1 therein for the definition of stable (admissible) quotients over general expanded degenerations.

Similarly, we define stable relative quotients on $(\mathfrak{C}_{\pm},\mathfrak{x}_{\pm}) \to \mathcal{T}$ and $(\mathfrak{x}_{-},\mathfrak{C}_{\sim},\mathfrak{x}_{+}) \to \mathcal{T}_{\sim}$ by requiring their restrictions to the fibers to be locally free at the nodes and the markings \mathfrak{x}_{\pm} and have positive degrees over all \mathbb{P}^{1} 's in the rational tails or chains. Let $\mathfrak{Quot}_{d}(\mathfrak{C}_{\pm}/\mathcal{T}, N, r)$ and $\mathfrak{Quot}_{d}(\mathfrak{C}_{\sim}/\mathcal{T}_{\sim}, N, r)$ denote the moduli spaces of flat families of stable relative quotients over the corresponding universal families of expanded degenerations. According to [LW, Theorem 4.14, 4.15], all these stacks, including $\mathfrak{Quot}_{d}(\mathfrak{C}/\mathfrak{B}, N, r)$ are separate, Deligne–Mumford stacks of finite type; the first three are proper, whereas $\mathfrak{Quot}_{d}(\mathfrak{C}/\mathfrak{B}, N, r)$ is proper over B. For simplicity, we set $\mathcal{M}^{d} := \mathfrak{Quot}_{d}(\mathfrak{C}/\mathfrak{B}, N, r), \mathcal{M}^{d}_{\pm} := \mathfrak{Quot}_{d}(\mathfrak{C}_{\pm}/\mathcal{T}, N, r)$, and $\mathcal{M}^{d}_{\sim} := \mathfrak{Quot}_{d}(\mathfrak{C}_{\sim}/\mathcal{T}_{\sim}, N, r)$.

Now we identify \mathcal{M}^d_{\pm} and \mathcal{M}^d_{\sim} with the quasimap moduli spaces introduced in Section 6. By Remark 7.2, we have $\mathcal{T}_{\sim} \cong \mathfrak{M}^{ss}_{0,2}$. Note that both \mathcal{M}^d_{\sim} and the moduli space $Q^{\epsilon=0+}_{0,2}(X,d)$ of $(\epsilon = 0+)$ -stable quasimaps, referred to as stable quotients in [MOP], are defined over \mathcal{T}_{\sim} with matching stability conditions (see Definition 7.4 and [MOP, §2]). Hence, we have the isomorphism over \mathcal{T}_{\sim} :

(30)
$$\mathcal{M}^d_{\sim} \cong Q^{\epsilon=0+}_{0,2}(X,d).$$

S. SINHA AND M. ZHANG

Let C be a smooth projective curve of genus g. Let $\mathfrak{M}_{g,1}(C,1)$ denote the stack of maps of degree 1 from 1-pointed prestable curves of genus g to C. In other words, $\mathfrak{M}_{g,1}(C,1)$ is the moduli stack of genus g, 1-pointed prestable curves with one parametrized component isomorphic to C. We choose a closed point $c \in C$. Let $\mathcal{T}_{maps}(C,c) \subset \mathfrak{M}_{g,1}(C,1)$ denote the locally closed substack parametrizing curves that are obtained from the parametrized component C by gluing at c a chain of rational curves, with the unique marking at the extremal component (c.f. [ACFW, Definition 3.1.2] in the case $C = \mathbb{P}^1$). Let $\mathcal{T}_{naive}(C,c)$ denote the stack of *naive expansions* of the pair (C,c) (see [ACFW, Definition 2.1.3]). Then by using the arguments in the proof of [ACFW, Lemma 3.1.3], one can show that

$$\mathcal{T}_{naive}(C,c) \cong \mathcal{T}_{maps}(C,c).$$

It follows from Lemma 3.2.2 and Proposition 3.2.3 in [ACFW] that

$$\mathcal{T} \cong \mathcal{T}_{\text{maps}}(C, c)$$

Let $Q_{C_{\pm},1}^{0+}(X,d)$ be the moduli stacks of (0+)-stable quasimaps with one parametrized component as defined in Definition 6.2. Consider the forgetful morphism $Q_{C_{\pm},1}^{0+}(X,d) \to \mathcal{T}_{maps}(C_{\pm},x_{\pm})$ forgetting the stable quotients. As a consequence of Definition 7.4, we have an isomorphism over \mathcal{T} :

$$\mathcal{M}^d_{\pm} \cong Q^{0+}_{C+,1}(X,d).$$

For later applications, we fix a point $y_{-} \in C_{-}$ (resp. $y_{+} \in C_{+}$) away from x_{-} (resp. x_{+}). Let $Q_{C_{\pm},1|1}^{0+,0+}(X,d_{\pm})$ be the moduli stacks of (0+,0+)-stable quasimap with one parametrized component and (1,1)-weighted markings, as defined in Definition 6.4. Here we assume the light marking is at y_{\pm} and the heavy marking is contracted to x_{\pm} via the parametrization map from source curves to C_{\pm} . Since the point y_{\pm} is fixed, adding a light point on the parametrized component at y_{\pm} does not change the moduli space. Hence, we have:

(31)
$$\mathcal{M}^d_{\pm} \cong Q^{0+,0+}_{C_+,1|1}(X,d).$$

7.3. Decomposition of the central fiber. Let $\mathfrak{C}_0, \mathfrak{B}_0$ and \mathcal{M}_0^d denote the fibers of $\mathfrak{C}, \mathfrak{B}$ and \mathcal{M}^d over $0 \in B$, respectively. In this section, we describe the structure of the central fiber \mathcal{M}_0^d . Set

$$\mathfrak{B}_0[n] := \mathcal{T} \times \underbrace{\mathcal{T}_{\sim} \times \cdots \times \mathcal{T}_{\sim}}_{n \text{ copies}} \times \mathcal{T}.$$

Then there is a natural map

$$\operatorname{gl}_n:\mathfrak{B}_0[n]\to\mathfrak{B}_0$$

induced by gluing the expanded relative pairs of (C_{\pm}, x_{\pm}) and non-rigid expanded degenerations associated to the node. By Remark 4.7 and the proof of Theorem 4.9 in [Qu] (see also [LW, Proposition 2.19] and [Lee, Lemma 3]), $\mathfrak{B}_0 \subset \mathfrak{B}$ is a complete intersection substack, $\mathfrak{B}_0[n]$ is the normalization of the codimension-*n* strata in \mathfrak{B}_0 , and we have

(32)
$$\sum_{n=0}^{\infty} (-1)^n (\mathrm{gl}_n)_* \mathcal{O}_{\mathfrak{B}_0[n]} = \mathcal{O}_{\mathfrak{B}_0}.$$

There are evaluation maps

$$\operatorname{ev}_{\pm} : \mathcal{M}^d_{\pm} \to X \quad \text{and} \quad (\operatorname{ev}^{\sim}_{-}, \operatorname{ev}^{\sim}_{+}) : \mathcal{M}^d_{\sim} \to X \times X$$

defined by the restrictions of the universal quotients to \mathfrak{x}_{\pm} . Given $d, n \geq 0$, we define

$$\Lambda_d^n = \Big\{ \underline{d} = (d_-, d_1, \dots, d_n, d_+) : d_- + d_+ + \sum_{i=1}^n d_i = d, d_\pm \ge 0, d_i > 0 \Big\}.$$

Given $\underline{d} \in \Lambda_d^n$, we set

$$\mathcal{M}_{\overline{0}}^{\underline{d}}[n] := \mathcal{M}_{-}^{d_{-}} \times_{X} \mathcal{M}_{\sim}^{d_{1}} \times_{X} \dots \mathcal{M}_{\sim}^{d_{n}} \times_{X} \mathcal{M}_{+}^{d_{+}}.$$

There is a gluing map $\iota_{\underline{d}} : \mathcal{M}_{0}^{\underline{d}}[n] \to \mathcal{M}_{0}^{d}$ of stable quotients, which fits into the following cartesian diagram

For $n \ge 0$, we think of $\coprod_{d \in \Lambda^n_d} \mathcal{M}^d_0[n]$ as the normalization of the codimension-*n* strata in \mathcal{M}^d_0 .

7.4. **Decomposition of the virtual structure sheaf.** Let $S_{\mathcal{M}_0^d}$ and $\mathcal{Q}_{\mathcal{M}_0^d}$ be the universal subbundle and quotient sheaf over the universal curve $\pi_d : \mathcal{M}_0^d \times_{\mathfrak{B}_0} \mathfrak{C}_0 \to \mathcal{M}_0^d$, whose pullbacks to $\mathcal{M}_0^{\underline{d}}[n]$ via $\iota_{\underline{d}}$ are denoted by $S_{\mathcal{M}_0^{\underline{d}}[n]}$ and $\mathcal{Q}_{\mathcal{M}_0^{\underline{d}}[n]}$, respectively. We define the relative POT of $\mathcal{M}_0^{\underline{d}}[n] \to \mathfrak{B}_0[n]$ by pulling back the relative POT $(R \operatorname{\mathcal{H}om}_{\pi_d}(\mathcal{S}_{\mathcal{M}_0^d}, \mathcal{Q}_{\mathcal{M}_0^d}))^{\vee}$ of $\mathcal{M}_0^d \to \mathfrak{B}_0$ via $\iota_{\underline{d}}$:

$$\left(R \operatorname{\mathcal{H}om}_{\pi_{\underline{d}}}(\mathcal{S}_{\mathcal{M}_{0}^{\underline{d}}[n]}, \mathcal{Q}_{\mathcal{M}_{0}^{\underline{d}}[n]})\right)^{\vee}$$

where $\pi_{\underline{d}} : \mathfrak{C}_0[n] \to \mathcal{M}_0^{\underline{d}}[n]$ is the universal curve. These two relative POTs induce virtual pullbacks for the two vertical maps in (33), and we define the virtual structure sheaves $\mathcal{O}_{\mathcal{M}_0^d}^{\mathrm{vir}}$ and $\mathcal{O}_{\mathfrak{M}_0^d}^{\mathrm{vir}}$ as the virtual pullbacks of $\mathcal{O}_{\mathfrak{B}_0}$ and $\mathcal{O}_{\mathfrak{B}_0[n]}$, respectively. According to [Qu, Theorem 2.6], virtual pullbacks commute with Gysin pullbacks. Hence, we have

(34)
$$(\operatorname{gl}_n)^! \mathcal{O}_{\mathcal{M}_0^d}^{\operatorname{vir}} = \mathcal{O}_{\mathcal{M}_0^d[n]}^{\operatorname{vir}}.$$

If we consider the standard $T = (\mathbb{C}^*)^N$ -action on \mathcal{M}_0^d and $\mathcal{M}_0^d[n]$, then the above equation also holds in the *T*-equivariant *K*-group.

Theorem 7.5. We have

$$\sum_{n\geq 0}\sum_{\underline{d}\in\Lambda_d^n}(-1)^n(\iota_{\underline{d}})_*\mathcal{O}_{\mathcal{M}_0^d[n]}^{\mathrm{vir}}=\mathcal{O}_{\mathcal{M}_0^d}^{\mathrm{vir}}$$

in $K_0^T(\mathcal{M}_0^d)$.

Proof. The theorem follows (32), (34), and the commutativity between virtual pullbacks and proper pushforwards [Qu, Proposition 2.4]. \Box

To split invariants over $\mathcal{M}_{\overline{0}}^{\underline{d}}[n]$, we first define the relative POTs for $\mathcal{M}_{\pm}^{d_{\pm}} \to \mathcal{T}$ and $\mathcal{M}_{\sim}^{d_{i}} \to \mathcal{T}_{\sim}$ using (13) (or equivalently (12)). Note that the (*T*-equivariant) isomorphisms (30) and (31) preserve these relative POTs. Let $\mathcal{O}_{\mathcal{M}_{\pm}^{d_{\pm}}}^{\text{vir}}$ and $\mathcal{O}_{\mathcal{M}_{\sim}^{d_{i}}}^{\text{vir}}$ be the virtual pullbacks of the structure sheaves of \mathcal{T} and \mathcal{T}_{\sim} , respectively. Consider the cartesian diagram:

Let $\mathfrak{C}'_0[n]$ denote the pullback of the univeral curve over $\mathcal{M}^{d_-}_- \times \mathcal{M}^{d_1}_\sim \times \cdots \mathcal{M}^{d_n}_\sim \times \mathcal{M}^{d_+}_+$ to $\mathcal{M}^{d}_0[n]$. Then we have a commutative diagram



where f and f' are the universal maps and p is the gluing map. By the stability condition of the quotients, $Lf^*\mathbb{T}_{\mathfrak{X}} = \mathcal{H}om(\mathcal{S}_{\mathcal{M}_0^d[n]}, \mathcal{Q}_{\mathcal{M}_0^d[n]})$ is locally free at the nodes of the universal curve. It follows from the arguments in the proof of the cutting-edges axiom in [Beh] that the relative POTs of $\mathcal{M}_0^d[n]$ and $\mathcal{M}_-^{d_-} \times \mathcal{M}_{\sim}^{d_1} \times \cdots \mathcal{M}^{d_n} \times \mathcal{M}_+^{d_+}$ are compatible with respect to $\Delta_X^{\times n}$ (see [BF]). Then by the functoriality of virtual pullbacks [Qu, Proposition 2.11], we have

(35)
$$\mathcal{O}_{\mathcal{M}_{0}^{d}[n]}^{\operatorname{vir}} = \left(\Delta_{X}^{\times n}\right)^{!} \left(\mathcal{O}_{\mathcal{M}_{-}^{d_{-}}}^{\operatorname{vir}} \boxtimes \mathcal{O}_{\mathcal{M}_{-}^{d_{1}}}^{\operatorname{vir}} \boxtimes \dots \mathcal{O}_{\mathcal{M}_{-}^{d_{n}}}^{\operatorname{vir}} \boxtimes \mathcal{O}_{\mathcal{M}_{+}^{d_{+}}}^{\operatorname{vir}}\right)$$

in $K_0^T(\mathcal{M}_0^{\underline{d}}[n])$.

7.5. Degeneration formula. We choose a point $y_{-} \in C_{-}$ (resp. $y_{+} \in C_{+}$) away from x_{-} (resp. x_{+}). After shrinking B if necessary, we extend these two markings to sections $\mathfrak{y}_{\pm}:\mathfrak{B}\to\mathfrak{C}$ of the expanded degeneration of curves.

Consider the 'stacky' evaluation map

$$\widetilde{\operatorname{ev}}_{y_{\pm}}:\mathcal{M}^{d_{\pm}}_{\pm} o \mathfrak{X}$$

induced by the restriction of the universal subbundles at y_{\pm} . By abuse of notation, we also use $\tilde{\mathrm{ev}}_{y_{\pm}}$ to denote the evaluation maps on $\mathcal{M}_{0}^{\underline{d}}[n]$ and $\mathcal{M}_{0}^{\underline{d}}$. For $\mathrm{V}, \mathrm{W} \in K_{T}^{0}(\mathfrak{X})$, we define

$$\langle\!\langle \mathbf{V}, \mathbf{W} \rangle\!\rangle_{\mathrm{deg}}^{T} := \sum_{d \ge 0} q^{d} \chi^{\mathrm{vir}, T} \left(\mathcal{M}_{0}^{d}, \widetilde{\mathrm{ev}}_{y_{-}}^{*}\left(\mathbf{V}\right) \cdot \widetilde{\mathrm{ev}}_{y_{+}}^{*}\left(\mathbf{W}\right) \right).$$

It follows from Theorem 7.5 that

$$\langle\!\langle \mathbf{V}, \mathbf{W} \rangle\!\rangle_{\mathrm{deg}}^{T} = \sum_{n \ge 0} (-1)^{n} \sum_{d \ge 0, \underline{d} \in \Lambda_{d}^{n}} q^{d} \chi^{\mathrm{vir}, T} \left(\mathcal{M}_{0}^{\underline{d}}[n], \widetilde{\mathrm{ev}}_{y_{-}}^{*}\left(\mathbf{V}\right) \cdot \widetilde{\mathrm{ev}}_{y_{+}}^{*}\left(\mathbf{W}\right) \right).$$

Define $F_{\alpha,\beta}^{\epsilon=0+} = g_{\alpha,\beta} + F'_{\alpha,\beta}$, where $g_{\alpha,\beta} := \chi^T(X, \mathcal{O}_{\alpha} \cdot \mathcal{O}_{\beta})$ and $F'_{\alpha,\beta} = \sum_{d>0} q^d \langle \mathcal{O}_{\alpha}, \mathcal{O}_{\beta} \rangle_{0,2,d}^{\epsilon=0+,T}$.

Using (35) and the isomorphisms (30), (31), we obtain

$$\sum_{d\geq 0,\underline{d}\in\Lambda_{d}^{n}}q^{d}\chi^{\operatorname{vir},T}\left(\mathcal{M}_{\overline{0}}^{\underline{d}}[n],\widetilde{\operatorname{ev}}_{y_{-}}^{*}(\mathbf{V})\cdot\widetilde{\operatorname{ev}}_{y_{+}}^{*}(\mathbf{W})\right) = \sum_{\alpha,\beta,a_{i},b_{i}\in\operatorname{P}_{r,k}}\langle\!\langle \mathbf{V},\mathcal{O}_{\alpha}\rangle\!\rangle_{C_{-},1|1}^{0+,T}$$
$$\cdot g^{\alpha,a_{1}}F_{a_{1},b_{1}}'g^{b_{1},a_{2}}\cdots F_{a_{m},b_{m}}'g^{b_{m},\beta}\cdot\langle\!\langle \mathcal{O}_{\beta},\mathbf{W}\rangle\!\rangle_{C_{+},1|1}^{0+,T}.$$

Here $g^{\alpha,\beta}$ denote the coefficients of the inverse of $(g_{\alpha,\beta})$. It was first observed in [Giv] (in quantum K-theory) that the combinatorics involved in the above equation can be simplified using the inverse matrix $F_{\epsilon=0+}^{\alpha,\beta}$, which equals to

$$g^{\alpha,\beta} + \sum_{m \ge 1, a_i, b_i \in \mathcal{P}_{r,k}} (-1)^m g^{\alpha,a_1} F'_{a_1,b_1} g^{b_1,a_2} \cdots F'_{a_m,b_m} g^{b_m,\beta}.$$

In sum, we obtain

$$\langle\!\langle \mathbf{V}, \mathbf{W} \rangle\!\rangle_{\mathrm{deg}}^{T} = \sum_{\alpha, \beta \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathcal{O}_{\alpha} \rangle\!\rangle_{C_{-},1|1}^{0+,T} \cdot F_{\epsilon=0+}^{\alpha,\beta} \cdot \langle\!\langle \mathcal{O}_{\beta}, \mathbf{W} \rangle\!\rangle_{C_{+},1|1}^{0+,T}$$

Hence by using the wall-crossing results from Proposition 6.6 and Proposition 5.3, we have

(36)
$$\langle\!\langle \mathbf{V}, \mathbf{W} \rangle\!\rangle_{\mathrm{deg}}^{T} = \sum_{\alpha, \beta \in \mathbf{P}_{r,k}} \langle\!\langle \mathbf{V}, \mathbf{O}_{\alpha} \rangle\!\rangle_{g_{-}}^{\mathsf{Quot}, T} \cdot F^{\alpha, \beta} \cdot \langle\!\langle \mathbf{O}_{\beta}, \mathbf{W} \rangle\!\rangle_{g_{+}}^{\mathsf{Quot}, T}$$

Let $q_{\mathfrak{B}} : \mathcal{M}^d \to \mathfrak{B}$ and $q_B : \mathcal{M}^d \to B$ be the good degeneration of Quot schemes. Let $i_b : b \hookrightarrow C$ be the regular embedding of a closed point. Consider the cartesian diagram

$$\begin{array}{ccc} \mathcal{M}_b^d & \longrightarrow \mathfrak{B}_b & \longrightarrow b \\ \downarrow^{\iota_b} & \downarrow & \downarrow^{i_b} \\ \mathcal{M}^d & \xrightarrow{q_{\mathfrak{B}}} \mathfrak{B} & \longrightarrow B \end{array}$$

We define the virtual structure sheaf of the good degeneration by

$$\mathcal{O}_{\mathcal{M}^d}^{\mathrm{vir}} := q^! (\mathcal{O}_{\mathfrak{B}}) \in K_0^T (\mathcal{M}^d).$$

Since the POT of \mathcal{M}_{b}^{d} is the pullback of the relative POT of q, we have

$$i_b^!(\mathcal{O}_{\mathcal{M}^d}^{\mathrm{vir}}) = \mathcal{O}_{\mathcal{M}_b^d}^{\mathrm{vir}}$$

Let $F \in K^0_T(\mathcal{M}^d)$. Then

$$\chi^{\operatorname{vir},T}\left(\mathcal{M}_{b}^{d},F|_{\mathcal{M}_{b}^{d}}\right) = \chi^{T}\left(\mathcal{M}^{d},F\cdot(\iota_{b})_{*}i_{b}^{!}(\mathcal{O}_{\mathcal{M}^{d}}^{\operatorname{vir}})\right) = \chi^{T}\left(B,i_{b}^{!}(q_{B})_{*}(F\cdot\mathcal{O}_{\mathcal{M}^{d}}^{\operatorname{vir}})\right)$$

is independent of b. In particular, if we fix a point $b \neq 0$ in B, then

(37)
$$\chi^{\operatorname{vir},T}\left(\mathcal{M}_{b}^{d},F|_{\mathcal{M}_{b}^{d}}\right) = \chi^{\operatorname{vir},T}\left(\mathcal{M}_{0}^{d},F|_{\mathcal{M}_{0}^{d}}\right).$$

Proof of Theorem 1.15 (i). Let $C = \mathfrak{C}_b$ be the fiber of $\mathfrak{C} \to B$ at $b \neq 0$. Then C is a smooth curve of genus g, and we have

$$\mathcal{M}_b^d \cong \mathsf{Quot}_d(C, N, r).$$

Set $F = \tilde{ev}_{\mathfrak{y}_{-}}^{*}(V) \cdot \tilde{ev}_{\mathfrak{y}_{+}}^{*}(W)$, where $\tilde{ev}_{\mathfrak{y}_{\pm}} : \mathcal{M}^{d} \to \mathfrak{X}$ denote the stacky evaluation maps at \mathfrak{y}_{\pm} . Then the degeneration formula (7) follows from the decomposition (36) and the deformation invariance of virtual Euler characteristics (37).

Remark 7.6. The degeneration formula (8) in Theorem 1.15 (ii) can be proven similarly by analyzing a simple degeneration $\mathcal{C}' \to \mathcal{B}$ over a smooth pointed curve $(\mathcal{B}, 0)$. Here, the fibers outside 0 are smooth curves of genus g, while the fiber at 0, C'_{deg} , is an irreducible curve with one node. The associated stack of expanded degenerations, $\mathfrak{C}' \to \mathfrak{B}$, has central fibers formed by inserting a chain of \mathbb{P}^1s (of various lengths) at the unique node of C'_{deg} (see [LW, §2.2]). The good degeneration of Quot schemes, $\mathfrak{Quot}_d(\mathfrak{C}'/\mathfrak{B}, N, r) \to \mathcal{B}$, parametrizes flat families of stable quotients of rank N - r and degree d over the universal family $\mathfrak{C}' \to \mathfrak{B}$.

Let \tilde{C} be the normalization of C'_{deg} , with x_{-} and x_{+} in \tilde{C} representing the preimages of the unique node in C'_{deg} . The central fiber of this degeneration of Quot schemes is 'virtually' a complete intersection substack, and its 'normalization' is

(38)
$$\coprod_{n,\underline{d}\in\tilde{\Lambda}^n_d} \tilde{\mathcal{M}}^{d_0}_0 \times_X \mathcal{M}^{d_1}_{\sim} \times_X \dots \mathcal{M}^{d_n}_{\sim} \times_X \left(\tilde{\mathcal{M}}^{d_0}_0 \right),$$

where $\tilde{\Lambda}_{d}^{n} = \{ \underline{d} = (d_{0}, d_{1}, \dots, d_{n}) : d_{0} + \sum_{i=1}^{n} d_{i} = d \text{ and } d_{i} > 0 \text{ for } 1 \leq i \leq n \},$ $\tilde{\mathcal{M}}_{0}^{d_{0}} \cong Q_{\tilde{C}, 1|2}^{0+, 0+}(X, d_{0}),$

and we put the further right $\tilde{\mathcal{M}}_{0}^{d_{0}}$ in parenthesis to indicate is is the same as the one on the left. Here, a source curve in $Q_{\tilde{C},2|2}^{0+,0+}(X,d_{0})$ is formed by attaching rational tails to \tilde{C} at x_{\pm} and placing heavy markings on the outermost components of these rational tails. The inclusion of one light marking on the parametrized component \tilde{C} does not alter the moduli space but allows us to define invariants with one insertion in $K_{T}^{0}(\mathfrak{X})$. Similar to the proof of Theorem 1.15 (i), we apply the deformation invariance of virtual Euler characteristics to relate $\langle\!\langle V \rangle\!\rangle_{g}^{Quot,T}$ to the alternating sum of virtual Euler characteristics over the components in (38). This sum can then be rewritten as the right side of the degeneration formula (8) using the wall-crossing formulas in Proposition 6.6 and Proposition 5.3.

References

- [ACFW] Dan Abramovich, Charles Cadman, Barbara Fantechi, and Jonathan Wise. Expanded degenerations and pairs. Comm. Algebra, 41(6):2346–2386, 2013.
- [ACT1] David Anderson, Linda Chen, and Hsian-Hua Tseng. On the finiteness of quantum K-theory of a homogeneous space. Int. Math. Res. Not. IMRN, (2):1313–1349, 2022.
- [ACT2] David Anderson, Linda Chen, and Hsian-Hua Tseng. On the finiteness of quantum K-theory of a homogeneous space. Int. Math. Res. Not. IMRN, (2):1313–1349, 2022.

- [BCFK] Aaron Bertram, Ionuţ Ciocan-Fontanine, and Bumsig Kim. Two proofs of a conjecture of Hori and Vafa. Duke Math. J., 126(1):101–136, 2005.
- [BCMP1] Anders S. Buch, Pierre-Emmanuel Chaput, Leonardo C. Mihalcea, and Nicolas Perrin. Finiteness of cominuscule quantum K-theory. Ann. Sci. Éc. Norm. Supér. (4), 46(3):477–494, 2013.
- [BCMP2] Anders S. Buch, Pierre-Emmanuel Chaput, Leonardo C. Mihalcea, and Nicolas Perrin. Rational connectedness implies finiteness of quantum K-theory. Asian J. Math., 20(1):117–122, 2016.
- [BCMP3] Anders S. Buch, Pierre-Emmanuel Chaput, Leonardo C. Mihalcea, and Nicolas Perrin. A Chevalley formula for the equivariant quantum K-theory of cominuscule varieties. Algebr. Geom., 5(5):568–595, 2018.
 - [BDW] Aaron Bertram, Georgios Daskalopoulos, and Richard Wentworth. Gromov invariants for holomorphic maps from Riemann surfaces to Grassmannians. J. Amer. Math. Soc., 9(2):529–571, 1996.
 - [Beh] K. Behrend. Gromov-Witten invariants in algebraic geometry. Invent. Math., 127(3):601-617, 1997.
 - [Ber1] Aaron Bertram. Towards a Schubert calculus for maps from a Riemann surface to a Grassmannian. Internat. J. Math., 5(6):811–825, 1994.
 - [Ber2] Aaron Bertram. Quantum Schubert calculus. Adv. Math., 128(2):289–305, 1997.
 - [BF] K. Behrend and B. Fantechi. The intrinsic normal cone. Invent. Math., 128(1):45– 88, 1997.
 - [BM] Anders S. Buch and Leonardo C. Mihalcea. Quantum K-theory of Grassmannians. Duke Math. J., 156(3):501–538, 2011.
 - [BP] Jim Bryan and Rahul Pandharipande. Curves in Calabi-Yau threefolds and topological quantum field theory. Duke Math. J., 126(2):369–396, 2005.
 - [Buc] Anders Skovsted Buch. A Littlewood-Richardson rule for the K-theory of Grassmannians. Acta Math., 189(1):37–78, 2002.
 - [CF1] Ionuţ Ciocan-Fontanine. Quantum cohomology of flag varieties. Internat. Math. Res. Notices, (6):263–277, 1995.
 - [CF2] Ionuţ Ciocan-Fontanine. On quantum cohomology rings of partial flag varieties. Duke Math. J., 98(3):485–524, 1999.
 - [CF3] Ionuţ Ciocan-Fontanine. The quantum cohomology ring of flag varieties. Trans. Amer. Math. Soc., 351(7):2695–2729, 1999.
- [CFKM] Ionuţ Ciocan-Fontanine, Bumsig Kim, and Davesh Maulik. Stable quasimaps to GIT quotients. J. Geom. Phys., 75:17–47, 2014.
 - [Che] Linda Chen. Quantum cohomology of flag manifolds. Adv. Math., 174(1):1–34, 2003.
 - [FH] William Fulton and Joe Harris. Representation theory, volume 129 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1991. A first course, Readings in Mathematics.

S. SINHA AND M. ZHANG

- [Giv] Alexander Givental. On the WDVV equation in quantum K-theory. volume 48, pages 295–304. 2000. Dedicated to William Fulton on the occasion of his 60th birthday.
- [GMS⁺] Wei Gu, Leonardo C. Mihalcea, Eric Sharpe, Weihong Xu, Hao Zhang, and Hao Zou. Quantum K Whitney relations for partial flag varieties. arXiv e-prints, page arXiv:2310.03826, October 2023.
- [GMSZ1] Wei Gu, Leonardo Mihalcea, Eric Sharpe, and Hao Zou. Quantum K theory of symplectic Grassmannians. J. Geom. Phys., 177:Paper No. 104548, 38, 2022.
- [GMSZ2] Wei Gu, Leonardo C. Mihalcea, Eric Sharpe, and Hao Zou. Quantum K theory of Grassmannians, Wilson line operators, and Schur bundles. arXiv e-prints, page arXiv:2208.01091, August 2022.
 - [Gro] Alexander Grothendieck. Techniques de construction et théorèmes d'existence en géométrie algébrique. IV. Les schémas de Hilbert. In Séminaire Bourbaki, Vol. 6, pages Exp. No. 221, 249–276. Soc. Math. France, Paris, 1995.
 - [GV] Tom Graber and Ravi Vakil. Relative virtual localization and vanishing of tautological classes on moduli spaces of curves. Duke Math. J., 130(1):1–37, 2005.
 - [GW] E. González and C. Woodward. Quantum Kirwan for quantum K-theory. In Facets of algebraic geometry. Vol. I, volume 472 of London Math. Soc. Lecture Note Ser., pages 265–332. Cambridge Univ. Press, Cambridge, 2022.
 - [Kap] M. M. Kapranov. Derived category of coherent sheaves on Grassmann manifolds. Izv. Akad. Nauk SSSR Ser. Mat., 48(1):192–202, 1984.
 - [Kat] Syu Kato. Loop structure on equivariant K-theory of semi-infinite flag manifolds. arXiv e-prints, page arXiv:1805.01718, May 2018.
- [KOUY] Bumsig Kim, Jeongseok Oh, Kazushi Ueda, and Yutaka Yoshida. Residue mirror symmetry for Grassmannians. In Schubert calculus and its applications in combinatorics and representation theory, volume 332 of Springer Proc. Math. Stat., pages 307–365. Springer, Singapore, [2020] ©2020.
 - [Lee] Y.-P. Lee. *Quantum K-theory. I. Foundations.* Duke Math. J., 121(3):389–424, 2004.
 - [Len] Cristian Lenart. Combinatorial aspects of the K-theory of Grassmannians. Ann. Comb., 4(1):67–82, 2000.
 - [Li1] Jun Li. Stable morphisms to singular schemes and relative stable morphisms. J. Differential Geom., 57(3):509–578, 2001.
 - [Li2] Jun Li. A degeneration formula of GW-invariants. J. Differential Geom., 60(2):199– 293, 2002.
 - [LT] Jun Li and Gang Tian. Virtual moduli cycles and Gromov-Witten invariants of algebraic varieties. J. Amer. Math. Soc., 11(1):119–174, 1998.
 - [LW] Jun Li and Baosen Wu. Good degeneration of Quot-schemes and coherent systems. Comm. Anal. Geom., 23(4):841–921, 2015.
 - [Mac] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Classic Texts in the Physical Sciences. The Clarendon Press, Oxford University Press, New York,

second edition, 2015. With contribution by A. V. Zelevinsky and a foreword by Richard Stanley, Reprint of the 2008 paperback edition [MR1354144].

- [MO1] Alina Marian and Dragos Oprea. Virtual intersections on the Quot scheme and Vafa-Intriligator formulas. Duke Math. J., 136(1):81–113, 2007.
- [MO2] Alina Marian and Dragos Oprea. GL Verlinde numbers and the Grassmann TQFT. Port. Math., 67(2):181–210, 2010.
- [MOP] Alina Marian, Dragos Oprea, and Rahul Pandharipande. The moduli space of stable quotients. Geom. Topol., 15(3):1651–1706, 2011.
- [MPT] D. Maulik, R. Pandharipande, and R. P. Thomas. Curves on K3 surfaces and modular forms. J. Topol., 3(4):937–996, 2010. With an appendix by A. Pixton.
- [Oko] Andrei Okounkov. Lectures on K-theoretic computations in enumerative geometry. In Geometry of moduli spaces and representation theory, volume 24 of IAS/Park City Math. Ser., pages 251–380. Amer. Math. Soc., Providence, RI, 2017.
- [OS] Dragos Oprea and Shubham Sinha. Euler characteristics of tautological bundles over Quot schemes of curves. Adv. Math., 418:Paper No. 108943, 45, 2023.
- [PR] Mihnea Popa and Mike Roth. Stable maps and Quot schemes. Invent. Math., 152(3):625–663, 2003.
- [Qu] Feng Qu. Virtual pullbacks in K-theory. Ann. Inst. Fourier (Grenoble), 68(4):1609– 1641, 2018.
- [RZ1] Yongbin Ruan and Ming Zhang. The level structure in quantum K-theory and mock theta functions. arXiv e-prints, page arXiv:1804.06552, Apr 2018.
- [RZ2] Yongbin Ruan and Ming Zhang. Verlinde/Grassmannian Correspondence and Rank 2 δ-Wall-Crossing. Peking Math. J., 2022.
- [Sin1] Shubham Sinha. Numerical Invariants of Quot Schemes of Curves. ProQuest LLC, Ann Arbor, MI, 2023. Thesis (Ph.D.)–University of California, San Diego.
- [Sin2] Shubham Sinha. The virtual intersection theory of isotropic quot schemes. J. Geom. Phys., 198:Paper No. 105128, 39, 2024.
- [Str] Stein Arild Strømme. On parametrized rational curves in Grassmann varieties. In Space curves (Rocca di Papa, 1985), volume 1266 of Lecture Notes in Math., pages 251–272. Springer, Berlin, 1987.
- [SZ] Shubham Sinha and Ming Zhang. Quantum K-invariants via quot schemes II.
- [Tod] Yukinobu Toda. Moduli spaces of stable quotients and wall-crossing phenomena. Compos. Math., 147(5):1479–1518, 2011.
- [Vak] Ravi Vakil. A geometric Littlewood-Richardson rule. Ann. of Math. (2), 164(2):371–421, 2006. Appendix A written with A. Knutson.
- [ZZ] Ming Zhang and Yang Zhou. *K-theoretic quasimap wall-crossing.* arXiv e-prints, page arXiv:2012.01401, December 2020.

S. SINHA AND M. ZHANG

INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS, TRIESTE *Email address:* ssinhal@ictp.it

Department of Mathematics, University of California, San Diego $\mathit{Email}\ address: \texttt{miz017@ucsd.edu}$