Improving Behrend's construction: Sets without arithmetic progressions in integers and over finite fields

Christian Elsholtz^{*}

Zach Hunter[†]

Laura Proske[‡]

Lisa Sauermann[§]

Abstract

We prove new lower bounds on the maximum size of subsets $A \subseteq \{1, \ldots, N\}$ or $A \subseteq \mathbb{F}_p^n$ not containing three-term arithmetic progressions. In the setting of $\{1, \ldots, N\}$, this is the first improvement upon a classical construction of Behrend from 1946 beyond lower-order factors (in particular, it is the first quasipolynomial improvement). In the setting of \mathbb{F}_p^n for a fixed prime p and large n, we prove a lower bound of $(cp)^n$ for some absolute constant c > 1/2 (for c = 1/2, such a bound can be obtained via classical constructions from the 1940s, but improving upon this has been a well-known open problem).

1 Introduction

The questions of estimating the maximum possible sizes of subsets of $\{1, \ldots, N\}$ and of \mathbb{F}_p^n without threeterm arithmetic progressions are among the most central problems in additive combinatorics. Let us denote the maximum possible size of such subsets by $r_3(N)$ and $r_3(\mathbb{F}_p^n)$, respectively. So, formally, $r_3(N)$ is the maximum possible size of a subset $A \subseteq \{1, \ldots, N\}$ such that there do not exist distinct $x, y, z \in A$ with x+z=2y, and similarly $r_3(\mathbb{F}_p^n)$ is the maximum possible size of a subset $A \subseteq \mathbb{F}_p^n$ such that there do not exist distinct $x, y, z \in A$ with x+z=2y. The problem of estimating $r_3(N)$ was raised by Erdős and Turán [18] in 1936, and has been intensively studied since then. It also has connections to problems in communication complexity, see [10, 23, 26]. The problem for \mathbb{F}_p^n has also been studied for several decades (see [28], see also [36, p. 142] for a slightly different but related setting).

In a breakthrough result in 2017, Ellenberg and Gijswijt [15] proved that for any prime $p \ge 3$, there is an upper bound of the form

$$r_3(\mathbb{F}_p^n) \le (c_p p)^n \tag{1.1}$$

for some constant $c_p < 1$ only depending on p (and their constant c_p converges to 0.841... for $p \to \infty$, see [3, Eq. (4.11)]). In the integer setting, in a more recent breakthrough Kelley and Meka [25] proved the upper bound

$$r_3(N) \le N \cdot \exp(-c(\log N)^{1/12})$$

for all $N \ge 3$, for some absolute constant c > 0. This drastically improved upon all the previous bounds, obtained over many decades by Roth [33], Heath-Brown [24], Szemerédi [40], Bourgain [6, 7], Sanders [37, 38] and Bloom–Sisask [4]. Afterwards, using a modification of their method, this bound was improved to

$$r_3(N) \le N \cdot \exp(-c(\log N)^{1/9})$$

by Bloom and Sisask [5].

^{*}Institute for Analysis and Number Theory, TU Graz, Austria. Email: elsholtz@math.tugraz.at. Supported by the joint FWF-ANR project Arithrand: FWF: I 4945-N and ANR-20-CE91-0006, and FWF grant DOC 183.

[†]Department of Mathematics, ETH Zürich, Switzerland. Email: zach.hunter@math.ethz.ch.

[‡]TU Graz, Austria. Email: proske@student.tugraz.at.

[§]Institute for Applied Mathematics, University of Bonn, Germany. Email: sauermann@iam.uni-bonn.de. Supported by the DFG Heisenberg Program.

These upper bounds for $r_3(N)$ match the shape of a classical lower bound for this problem due to Behrend [2] from 1946, which is of the form

$$r_3(N) > N \cdot 2^{-(2\sqrt{2} + o(1))} \sqrt{\log_2 N}.$$
(1.2)

Over the past almost eighty years, only the o(1)-term in this bound has been improved. In Behrend's original bound, this o(1)-term in the exponent encapsulated a factor of $(\log_2 N)^{-1/4}$, so the bound was of the form $r_3(N) \ge \Omega(N \cdot 2^{-2\sqrt{2}}\sqrt{\log_2 N} \cdot (\log_2 N)^{-1/4})$. In 2010, Elkin [13] improved this factor to $(\log_2 N)^{1/4}$ instead, and an alternative proof for the bound with this improved o(1)-term was found by Green and Wolf [22].

In this paper, we give the first improvement to Behrend's [2] classical lower bound beyond the o(1)-term in (1.2). As stated in the following theorem, we show that the constant factor $2\sqrt{2} \approx 2.828$ in the exponent can be improved to $2\sqrt{\log_2(24/7)} \approx 2.667$, proving that the classical bound (1.2) is not tight.

Theorem 1.1. We have

$$r_3(N) > N \cdot 2^{-(C+o(1))} \sqrt{\log_2 N}$$

with $C = 2\sqrt{\log_2(24/7)} < 2\sqrt{2}$.

Our proof of Theorem 1.1 is motivated by studying three-term progression free sets in \mathbb{F}_p^n , for a fixed relatively large prime p and large n. In this setting, one can adapt Behrend's construction [2] (as noted by Tao and Vu in their book on additive combinatorics [41, Exercise 10.1.3] and also observed by Alon, see [19, Lemma 17]) to show

$$r_3(\mathbb{F}_p^n) \ge \left(\frac{p+1}{2}\right)^{n-o(n)} \tag{1.3}$$

for any fixed prime p and large n. Alternatively, such a bound can also be shown via an adaptation of an earlier construction in the integer setting due to Salem and Spencer [35] from 1942 (see [1, Theorem 2.13]). The asymptotic notation o(n) in the bound (1.3) is for $n \to \infty$ with p fixed. The best quantitative bound for the o(n)-term in this statement is due to relatively recent work of the first author and Pach [17, Theorem 3.10], but beyond the o(n)-term, this bound has not been improved (except for specific small primes p, see the discussion below).

Comparing the upper and lower bounds for $r_3(\mathbb{F}_p^n)$ for a fixed (reasonably large) prime p and large n in (1.1) and (1.3), there is still a large gap. Both of these bounds are roughly of the form $(cp)^n$ with 0 < c < 1, but with different values of c. For the upper bound, the best known constant due to Ellenberg–Gijswijt [15] is $c \approx 0.85$ (when the fixed prime is large enough). For the lower bound the constant c = 1/2 from Behrend's construction [2] or alternatively the Salem–Spencer construction [35] has not been improved in more than eighty years despite a lot of attention, especially after the upper bound of Ellenberg–Gijswijt appeared (see e.g. the blog post [14] as well as the discussion thereafter). Here, we finally improve this constant in the lower bound to be strictly larger than 1/2.

Theorem 1.2. There is a constant c > 1/2 such that for every prime p and every sufficiently large positive integer n (sufficiently large in terms of p), we have $r_3(\mathbb{F}_p^n) \ge (cp)^n$.

Breaking the barrier of 1/2 in this result for $r_3(\mathbb{F}_p^n)$ relies on the same key insights as our lower bounds for $r_3(N)$ in Theorem 1.1 improving Behrend's construction. The problem for \mathbb{F}_p^n (the so-called finite field model) was popularized in the hope that improved upper bounds for $r_3(\mathbb{F}_p^n)$ would also lead to better upper bounds for $r_3(N)$ (see e.g. [21, 30, 43]). It turned out, however, that for \mathbb{F}_p^n much stronger bounds hold than in the integer setting (for fixed p, the Ellenberg–Gijswijt bound stated in (1.1) is of the form $r_3(\mathbb{F}_p^n) \leq (p^n)^{1-\gamma_p}$ for some $\gamma_p > 0$, but in the integer setting one cannot hope for a bound of the form $r_3(N) \leq N^{1-\gamma}$ for some fixed $\gamma > 0$, as shown by Salem–Spencer [35] and Behrend [2]). Nevertheless, this paper establishes a connection between lower bounds for $r_3(N)$ and for $r_3(\mathbb{F}_p^n)$.

Our proof of Theorem 1.2 shows that one can take any $c < \sqrt{7/24}$, for example c = 0.54. Even though this may not seem like a large improvement over 1/2, it is the first qualitative improvement over the constant 1/2 from the constructions of Salem–Spencer and Behrend from the 1940's. Both of these constructions lead to three-term progression free subsets of \mathbb{F}_p^n only consisting of vectors with all entries in $\{0, 1, \ldots, (p-1)/2\}$, i.e. they only use roughly half of the available elements in \mathbb{F}_p in each coordinate. The restriction of all entries

to $\{0, 1, \ldots, (p-1)/2\}$ is crucial in these constructions, as it ensures that there is no "wrap-around" over \mathbb{F}_p . However, such an approach cannot be used to obtain three-term progression free subset $A \subseteq \mathbb{F}_p^n$ of size $|A| > ((p+1)/2)^n$, so c = 1/2 is a significant barrier for this problem. In light of this, it may actually be considered a surprise that it is possible to obtain a constant c > 1/2.

For very small primes, better bounds were obtained with specific constructions depending on the particular prime. For example, in \mathbb{F}_3^n a lower bound of $2.2202^{n-o(n)}$ was recently obtained by Romera-Paredes et al. [32] using artifical intelligence building upon traditional methods from previous bound [9, 12, 42]. Naslund [29] informed us about forthcoming work with an approach related to Shannon capacity, proving a lower bound of $2.2208^{n-o(n)}$ for the maximum size of three-term progression free subsets of \mathbb{F}_3^n . In \mathbb{F}_5^n , the best known lower bounds is $(35^{1/3})^{n-o(n)}$ due to the first and third author, Pollak, Lipnik and Siebenhofer (see [16]), note that $35^{1/3} \approx 3.271$.

Our methods for improving the lower bounds for $r_3(N)$ and $r_3(\mathbb{F}_p^n)$ can be applied more generally for finding three-term progression free subsets in finite abelian groups. We recall that a three-term arithmetic progression in a finite abelian group G consists of three distinct elements $x, y, z \in G$ with x + z = y + y (see e.g. [11]), and let us denote the maximum possible size of a three-term progression free subset of G by $r_3(G)$. Brown and Buhler [8] proved in 1982 that $r_3(G) \leq o(|G|)$ for all finite abelian groups G (where the asymptotic notation o(|G|) is for growing group size |G|). Frankl, Graham and Rödl [20] gave a different proof of this statement in 1987, and further upper bounds were obtained by Meshulam [28] and Lev [27].

Of course, every finite abelian group G can be written in the form $\mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$, and the problem of estimating $r_3(\mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n})$ has been most intensively studied in the case $m_1 = \cdots = m_n$, i.e. in the setting of \mathbb{Z}_m^n . In breakthrough work, Croot-Lev-Pach [11] proved the upper bound $r_3(\mathbb{Z}_4^n) \leq 3.611^n$ by introducing a new polynomial method, which also led to the Ellenberg-Gijswijt bound for $r_3(\mathbb{F}_p^n)$ stated in (1.1). Using similar methods, Petrov and Pohoata [31] studied upper bounds for $r_3(\mathbb{Z}_8^n)$.

Generalizing Theorem 1.2, we show that for any integer $m \ge 2$ (not necessarily prime) and sufficiently large n, we have $r_3(\mathbb{Z}_m^n) \ge (cm)^n$ for some absolute constant c > 1/2. For odd m, the best previous lower bound for this problem has (as in the case of prime m) been of the form $((m + 1)/2)^{n-o(n)}$ (based on the Salem–Spencer construction [35] or the Behrend construction [2]). For even m, the best previous lower bound has been $((m + 2)/2)^{n-o(n)}$ due to the first author and Pach [17, Theorem 3.11].

In fact, we show the following more general lower bound for arbitrary abelian groups $\mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$.

Theorem 1.3. Consider integers m_1, \ldots, m_n (for some positive integer n), and let m be such that $2 \le m_i \le m$ for $i = 1, \ldots, n$. Then we have

$$r_3(\mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}) \ge \frac{(7/24)^{n/2}}{10^6 m^2 n^3} \cdot m_1 \cdots m_n.$$

Note that Theorem 1.3 immediately implies that for any fixed constant $c < \sqrt{7/24}$ and for every integer $m \ge 2$, we have $r_3(\mathbb{Z}_m^n) \ge (cm)^n$ if n is sufficiently large with respect to m and c. In particular, it implies Theorem 1.2.

It is plausible that by slight modifications of our method, one can obtain better numerical bounds in our results above. In particular, our construction relies on certain explicit two-dimensional building blocks of area close to 7/24, and an improved choice for these building blocks with larger area would automatically carry over to numerical improvements of the constant $C = 2\sqrt{\log_2(24/7)}$ in Theorem 1.1 and the constant 7/24 in Theorem 1.3. We see the main contribution of this paper as introducing this method, and using it to break the lower bounds for $r_3(N)$ and $r_3(\mathbb{F}_p^n)$ in (1.2) and (1.3) originating from the 1940's.

In the next section we give an overview of our proof approach and state some key propositions, from which we will deduce Theorems 1.1 to 1.3 in Section 3. To prove these propositions, we need to find suitable two-dimensional building blocks, and these will be constructed in Section 4.

Notation. As usual, for a point $x \in \mathbb{R}^n$, we denote the coordinates of x by x_1, \ldots, x_n . For real numbers $a, a' \in \mathbb{R}$, we write $a \equiv a' \mod 1$ if $a - a' \in \mathbb{Z}$. For two points $x \in \mathbb{R}^n$ and $y \in \mathbb{R}^n$, we write $x \equiv y \mod 1$ if we have $x_i \equiv y_i \mod 1$ for all $i = 1, \ldots, n$ (i.e. if we have $x - y \in \mathbb{Z}^n$). For a measurable subset $S \subseteq \mathbb{R}^n$, we denote its measure by $\mu(S)$.

Acknowledgements. We would like to thank Daniel Carter, Zachary Chase, Ben Green, Eric Naslund, and Benny Sudakov for interesting discussions and helpful comments.

2 Proof Overview

The first step in our constructions for large three-term progression free subsets of $\{1, \ldots, N\}$ or $\mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$ is to take a (randomized) embedding into a (high-dimensional) torus. This way, we can reduce both problems to finding a large subset of the torus without a three-term arithmetic progression such that the first and third term are reasonably far away from each other. Note that we cannot aim to find a large subset of the torus without any three-term arithmetic progressions, since it is well-known that any three-term progression free subset of the torus has measure zero (see for example [34]). So it is essential to only forbid three-term arithmetic progressions with far-away first and third term in our desired subset of the torus.

The following proposition states that there is indeed such a large-measure subset of a (high-dimensional) torus without three-term arithmetic progressions with far-away first and third term. Proving this proposition is the main difficulty of this paper.

Proposition 2.1. For any $0 < \delta < 1$ and any even positive integer n, there exists a measurable subset $S \subseteq [0,1)^n$ with measure $\mu(S) \ge 10^{-5} \delta^2 n^{-3} \cdot (7/24)^{n/2}$ such that for any $x, y, z \in S$ with $x + z \equiv 2y \mod 1$ we have $|x_i - z_i| < \delta$ for i = 1, ..., n.

The deductions of Theorems 1.1 to 1.3 from Proposition 2.1 will be discussed in Section 3.

We remark that it is easy to show a weaker version of Proposition 2.1 with a bound of $\mu(S) \ge \delta^2 \cdot (1/2)^{n+o(n)}$ for the measure of S. Indeed, there one can take $S \subseteq [0, 1/2)^n$ to be an appropriately chosen sphere shell of width on the order of δ^2 (notice that then for $x, y, z \in S$ we have $x + z \equiv 2y \mod 1$ if and only if x + z = 2y, since $S \subseteq [0, 1/2)^n$). From this weaker statement one can deduce (1.2) with Behrend's original constant $2\sqrt{2}$, this approach to proving Behrend's bound (with an improved o(1)-term) is due to Green–Wolf [22]. Obtaining the stronger bound on $\mu(S)$ in Proposition 2.1 (having a factor of $(7/24)^{n/2}$ instead of $(1/2)^n = (1/4)^{n/2}$), which leads to our improvement upon the constant $2\sqrt{2}$, is much more difficult. In particular, for n large in terms of δ , any subset $S \subseteq [0, 1)^n$ whose measure is as large as in Proposition 2.1 will have points $x, y, z \in S$ with $x + z \equiv 2y \mod 1$ but $x + z \neq 2y$, so the "modulo 1" cannot just be ignored like in the previous approach. In fact, there will always be $x, y, z \in S$ with $x + z \equiv 2y \mod 1$ and $x = z \neq y$. Therefore, in the conclusion of Proposition 2.1, it is crucial to bound the absolute values of the coordinates of x - z and not of x - y (which, as the step length of the "modulo 1" arithmetic progression x, y, z, would maybe be more natural to consider), since the analogous statement for x - y would be false.

In order to prove Proposition 2.1, we use subsets of a two-dimensional torus as building blocks. These twodimensional building blocks need to satisfy a somewhat technical condition, summarized in the following proposition.

Proposition 2.2. Let $0 < \varepsilon < 1$. Then there exists a measurable subset $T \subseteq [0,1)^2$ with measure $\mu(T) \ge 7/24 - \varepsilon$ and a measurable function $f: T \to [0,100/\varepsilon^2]$, such that for any $x, y, z \in T$ with $x + z \equiv 2y \mod 1$ we have

$$f(x) + f(z) \ge 2f(y) + (x_1 - z_1)^2 + (x_2 - z_2)^2.$$

To deduce Proposition 2.1 from Proposition 2.2, we define the set $S \subseteq [0, 1)^n$ in Proposition 2.1 to be a subset of the (n/2)-fold product set $T^{n/2} \subseteq [0, 1)^n$ consisting of those points $((x_1, x_2), (x_3, x_4), \ldots, (x_{n-1}, x_n)) \in$ $T^{n/2}$ where $f(x_1, x_2) + f(x_3, x_4) + \cdots + f(x_{n-1}, x_n)$ is contained in some small interval. In other words, the set $S \subseteq [0, 1)^n$ can be viewed as a "slice" of the product set $T^{n/2} \subseteq [0, 1)^n$ with respect to the function $f(x_1, x_2) + f(x_3, x_4) + \cdots + f(x_{n-1}, x_n)$. It is then not hard to show, using the conditions on the set T and the function f in Proposition 2.2, that such a set S satisfies the conditions in Proposition 2.1. The details of this proof are given at the end of this section.

In order to prove Proposition 2.2, we explicitly construct the desired subset $T \subseteq [0, 1)^2$ and the function f. The set T is a carefully chosen union of polygons in $[0, 1)^2$. The construction is somewhat ad-hoc and it seems plausible that the constant 7/24 in Propositions 2.2 and 2.1 (and consequently also the constants in Theorems 1.1 and 1.3) can be improved by finding a better construction of such a set T. However, the main novelty of this paper is our overall approach for proving Theorems 1.1 to 1.3, and showing that this approach can be used to beat the long-standing bounds on these problems.

The proof of Proposition 2.2 can be found in Section 4. We finish this section by showing that Proposition 2.2 indeed implies Proposition 2.1.

Proof of Proposition 2.1 assuming Proposition 2.2. For n = 2, Proposition 2.1 is trivially true (for example, we can take $S = [0, \delta)^2$). So we may assume that $n \ge 4$.

Define $\varepsilon = 1/n$ and let $T \subseteq [0,1)^2$ and the function f be as in Proposition 2.2. Note that then the (n/2)-fold product set $T^{n/2} \subseteq [0,1)^n$ has measure

$$\mu(T^{n/2}) = \mu(T)^{n/2} \ge (7/24 - \varepsilon)^{n/2} \ge \left(1 - \frac{24}{7}\varepsilon\right)^{n/2} \cdot \left(\frac{7}{24}\right)^{n/2} \ge 10^{-2} \cdot \left(\frac{7}{24}\right)^{n/2},$$

where we used that for all $a \in [0, 6/7]$ we have $1 - a \ge 10^{-(7/6)a}$ (indeed, for a = 0 we have $1 - 0 = 10^{-(7/6)\cdot 0}$, for a = 6/7 we have $1 - (6/7) = 1/7 > 1/10 = 10^{-(7/6)\cdot(6/7)}$, and the function $10^{-(7/6)a}$ is concave on the interval [0, 6/7]) and therefore (recalling that $\varepsilon = 1/n \le 1/4$)

$$\left(1 - \frac{24}{7}\varepsilon\right)^{n/2} \ge \left(10^{-(7/6)\cdot(24/7)\varepsilon}\right)^{n/2} = 10^{-2\varepsilon n} = 10^{-2}$$

On the set $T^{n/2}$, let us consider the function taking value $f(x_1, x_2) + f(x_3, x_4) + \cdots + f(x_{n-1}, x_n)$ for each $((x_1, x_2), (x_3, x_4), \dots, (x_{n-1}, x_n)) \in T^{n/2}$. This is a measurable function, with values in the interval $[0, 100n^3]$ (noting that $n \cdot 100/\varepsilon^2 = 100n^3$). Thus, for every integer $j = 0, \dots, \lfloor 100n^3 \cdot (2/\delta^2) \rfloor$, the set

$$S_j = \{ ((x_1, x_2), \dots, (x_{n-1}, x_n)) \in T^{n/2} \mid j \cdot \delta^2 / 2 \le f(x_1, x_2) + \dots + f(x_{n-1}, x_n) < (j+1) \cdot \delta^2 / 2 \}$$

is measurable, and the union of these sets S_j for $j = 0, ..., \lfloor 100n^3 \cdot (2/\delta^2) \rfloor$ is the entire product set $T^{n/2}$. Thus, there exists some $j \in \{0, ..., \lfloor 100n^3 \cdot (2/\delta^2) \rfloor\}$ such that

$$\mu(S_j) \ge \frac{\mu(T^{n/2})}{\lfloor 100n^3 \cdot (2/\delta^2) \rfloor + 1} \ge \frac{10^{-2} \cdot (7/24)^{n/2}}{4 \cdot 100n^3 \delta^{-2}} \ge 10^{-5} \delta^2 n^{-3} \cdot (7/24)^{n/2}.$$

It remains to show that for any $x, y, z \in S_j$ with $x + z \equiv 2y \mod 1$ we have $|x_i - z_i| < \delta$ for $i = 1, \ldots, n$. Indeed, for every $h = 1, \ldots, n/2$ we have $(x_{2h-1}, x_{2h}), (y_{2h-1}, y_{2h}), (z_{2h-1}, z_{2h}) \in T$ and $(x_{2h-1}, x_{2h}) + (z_{2h-1}, z_{2h}) \equiv 2(y_{2h-1}, y_{2h}) \mod 1$ and so by the condition in Proposition 2.2 we obtain

$$f(x_{2h-1}, x_{2h}) + f(z_{2h-1}, z_{2h}) \ge 2f(y_{2h-1}, y_{2h}) + (x_{2h-1} - z_{2h-1})^2 + (x_{2h} - z_{2ih})^2.$$

Summing this up for h = 1, ..., n/2, and recalling the definition of S_j , yields

$$2 \cdot (j+1) \cdot \delta^2/2 > f(x_1, x_2) + \dots + f(x_{n-1}, x_n) + f(z_1, z_2) + \dots + f(z_{n-1}, z_n)$$

$$\ge 2f(y_1, y_2) + \dots + 2f(y_{n-1}, y_n) + (x_1 - z_1)^2 + \dots + (x_n - z_n)^2$$

$$\ge 2 \cdot j \cdot \delta^2/2 + (x_1 - z_1)^2 + \dots + (x_n - z_n)^2,$$

and hence

$$(x_1 - z_1)^2 + \dots + (x_n - z_n)^2 < 2 \cdot \delta^2 / 2 = \delta^2$$

This implies that $|x_i - z_i| < \delta$ for i = 1, ..., n, as desired.

3 Deduction of main results

In this section we derive Theorems 1.1 to 1.3 from Proposition 2.1. We start with Theorem 1.3, and first prove the following version of it for the case of even n.

Proposition 3.1. Consider integers m_1, \ldots, m_n (for some positive integer n), and let m be such that $2 \le m_i \le m$ for $i = 1, \ldots, n$. If n is even, we have

$$r_3(\mathbb{Z}_{m_1}\times\cdots\times\mathbb{Z}_{m_n})\geq\frac{(7/24)^{n/2}}{10^5m^2n^3}\cdot m_1\cdots m_n.$$

As mentioned in the previous section, this statement is deduced from Proposition 2.1 by considering a (randomized) embedding of $\mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$ into the torus $[0,1)^n$. For a subset $S \subseteq [0,1)^n$ as in Proposition 2.1, one can then show that its pre-image is a three-term progression free subset. Using the lower bound on $\mu(S)$ in Proposition 2.1, this gives a lower bound for $r_3(\mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n})$.

Proof. Let $\delta = 1/m$, and let $S \subseteq [0,1)^n$ be a measurable subset as in Proposition 2.1. For any $a \in [0,1]^n$, let us now define a map $\varphi_a : \mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n} \to [0,1)^n$ as follows. Every point in $\mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$ can be represented (uniquely) by an n-tuple $(r_1, \ldots, r_n) \in \{0, 1, \ldots, m_1 - 1\} \times \cdots \times \{0, 1, \ldots, m_n - 1\}$. Let us now define $\varphi_a(r_1, \ldots, r_n) = (q_1, \ldots, q_n)$ for the unique point $q = (q_1, \ldots, q_n) \in [0, 1)^n$ with $(q_1, \ldots, q_n) \equiv$ $(a_1 + r_1/m_1, \ldots, a_n + r_n/m_n) \mod 1$.

Note that the map $\varphi_a : \mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n} \to [0,1)^n$ is injective for any $a \in [0,1]^n$. Furthermore, note that for any $a \in [0,1]^n$ and any $x, z \in \mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$ we either have $\varphi_a(x)_i = \varphi_a(z)_i$ or $|\varphi_a(x)_i - \varphi_a(z)_i| \ge 1/m_i \ge 1/m_i \ge 1/m = \delta$ (indeed, the different values occurring as the *i*-th coordinate of points in the image $\varphi_a(\mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n})$ form an arithmetic progression with step-width $1/m_i$). Hence for any $a \in [0,1]^n$ and any distinct $x, z \in \mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$, there exists a coordinate $i \in \{1, \ldots, n\}$ such that $|\varphi_a(x)_i - \varphi_a(z)_i| \ge \delta$. We claim that for any $a \in [0,1]^n$, the pre-image $\varphi_a^{-1}(S)$ of the set S from Proposition 2.1 is a threeterm progression free subset of $\mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$. Indeed, suppose that for some $a \in [0,1]^n$, there exist distinct $x, y, z \in \varphi_a^{-1}(S) \subseteq \mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$ with x + z = 2y. Then, as shown above, there must be a coordinate $i \in \{1, \ldots, n\}$ such that $|\varphi_a(x)_i - \varphi_a(z)_i| \ge \delta$. On the other hand, having x + z = 2y means that $x_i + z_i \equiv 2y_i \mod m_i$ for all $i = 1, \ldots, n$, where $(x_1, \ldots, x_n), (y_1, \ldots, y_n), (z_1, \ldots, z_n) \in \{0, 1, \ldots, m_1 - 1\} \times \cdots \times \{0, 1, \ldots, m_n - 1\}$ are the *n*-tuples corresponding to $x, y, z \in \mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$, respectively. Hence we obtain $(x_i/m_i) + (z_i/m_i) \equiv 2(y_i/m_i) \mod 1$ for $i = 1, \ldots, n$, and consequently

$$\varphi_a(x)_i + \varphi_a(z)_i \equiv (a_i + x_i/m_i) + (a_i + z_i/m_i) \equiv 2(a_i + y_i/m_i) \equiv 2\varphi_a(y)_i \mod 1$$

for i = 1, ..., n. This means that $\varphi_a(x) + \varphi_a(z) \equiv 2\varphi_a(y) \mod 1$. Since $\varphi_a(x), \varphi_a(y), \varphi_a(z) \in S$, the condition for the set S in Proposition 2.1 now implies that $|\varphi_a(x)_i - \varphi_a(z)_i| < \delta$ for all i = 1, ..., n. This is a contradiction, so the pre-image $\varphi_a^{-1}(S) \subseteq \mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$ is indeed three-term progression free for any $a \in [0, 1]^n$.

Finally, consider a uniformly random choice of $a \in [0, 1]^n$. For any $x \in \mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$ we have $x \in \varphi_a^{-1}(S)$ if and only if $\varphi_a(x) \in S$, and this happens with probability $\mu(S)$. Hence

$$\mathbb{E}[|\varphi_a^{-1}(S)|] = \sum_{x \in \mathbb{Z}_{m_1} \times \dots \times \mathbb{Z}_{m_n}} \mathbb{P}[x \in \varphi_a^{-1}(S)] = |\mathbb{Z}_{m_1} \times \dots \times \mathbb{Z}_{m_n}| \cdot \mu(S) \ge \frac{(7/24)^{n/2}}{10^5 m^2 n^3} \cdot m_1 \cdots m_n$$

recalling that $\mu(S) \ge 10^{-5} \delta^2 n^{-3} \cdot (7/24)^{n/2} = 10^{-5} (1/m)^2 n^{-3} \cdot (7/24)^{n/2}$. This implies that for some choice of $a \in [0, 1]^n$ we must have

$$|\varphi_a^{-1}(S)| \ge \frac{(7/24)^{n/2}}{10^5 m^2 n^3} \cdot m_1 \cdots m_n.$$

As shown above, the set $\varphi_a^{-1}(S) \subseteq \mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$ is three-term progression free, so this establishes the desired lower bound for $r_3(\mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n})$.

To deduce Theorem 1.3 from Proposition 3.1, we use the following simple fact.

Fact 3.2. For any finite abelian groups G and H, we have

$$r_3(G) \ge \frac{r_3(G \times H)}{|H|}.$$

Proof. Let $A \subseteq G \times H$ be a three-term progression free subset of $G \times H$ of size $|A| = r_3(G \times H)$. For any $h \in H$, let us define the map $\varphi_h : G \to G \times H$ by $\varphi_h(g) = (g, h)$ for all $g \in G$. For any $h \in H$, this map is injective and for any distinct $x, y, z \in G$, the images $\varphi_h(x), \varphi_h(y), \varphi_h(z) \in G \times H$ form a three-term arithmetic progression in $G \times H$ if and only if x, y, z form a three-term arithmetic progression in G. Thus, for any $h \in H$, the pre-image $\varphi_h^{-1}(A) \subseteq G$ is a three-term progression free subset of G and therefore has size $|\varphi_h^{-1}(A)| \leq r_3(G)$. On the other hand, by injectivity of φ_h we have $|\varphi_h^{-1}(A)| = |A \cap \varphi_h(G)|$. As the images $\varphi_h(G)$ for all $h \in H$ form a partition of $G \times H$, this implies

$$r_3(G \times H) = |A| = \sum_{h \in H} |A \cap \varphi_h(G)| = \sum_{h \in H} |\varphi_h^{-1}(A)| \le |H| \cdot r_3(G).$$

Rearranging yields the desired inequality.

Now, we are ready to prove Theorem 1.3

Proof of Theorem 1.3. Note that the inequality in the theorem statement holds trivially for $n \leq 2$ (since then the right-hand side is smaller than 1). So let us assume $n \geq 3$. If n is even, then the desired lower bound for $r_3(\mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n})$ follows immediately from Proposition 3.1. If n is odd, applying Proposition 3.1 with $m_{n+1} = m$ yields

$$r_3(\mathbb{Z}_{m_1} \times \dots \times \mathbb{Z}_{m_n} \times \mathbb{Z}_m) \ge \frac{(7/24)^{(n+1)/2}}{10^5 m^2 (n+1)^3} \cdot m_1 \cdots m_n \cdot m \ge \frac{(7/24)^{n/2}}{10^6 m^2 n^3} \cdot m_1 \cdots m_n \cdot m.$$

Applying Fact 3.2 to $G = \mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$ and $H = \mathbb{Z}_m$, we obtain

$$r_3(\mathbb{Z}_{m_1} \times \dots \times \mathbb{Z}_{m_n}) \ge \frac{r_3(\mathbb{Z}_{m_1} \times \dots \times \mathbb{Z}_{m_n} \times \mathbb{Z}_m)}{m} \ge \frac{(7/24)^{n/2}}{10^6 m^2 n^3} \cdot m_1 \cdots m_n.$$

Theorem 1.2 is an immediate consequence of Theorem 1.3.

Proof of Theorem 1.2. Let $1/2 < c < \sqrt{7/24}$. Then, as long as n is sufficiently large with respect to p, by Theorem 1.3 we have

$$r_3(\mathbb{F}_p^n) = r_3(\mathbb{Z}_p \times \dots \times \mathbb{Z}_p) \ge \frac{(7/24)^{n/2}}{10^6 p^2 n^3} \cdot p^n = \frac{(\sqrt{7/24} \cdot p)^n}{10^6 p^2 n^3} \ge (cp)^n.$$

Finally, it remains to derive Theorem 1.1.

I

Proof of Theorem 1.1. As we are proving an asymptotic statement, we may assume that N is sufficiently large. Let n be an even positive integer with

$$\frac{2 \cdot \sqrt{\log_2 N}}{\sqrt{\log_2(24/7)}} \le n \le \frac{2 \cdot \sqrt{\log_2 N}}{\sqrt{\log_2(24/7)}} + 2.$$
(3.1)

Let p_1, \ldots, p_n be the first *n* primes. Defining $p = \max\{p_1, \ldots, p_n\}$, we have $p \leq 100n \log_2 n$ (this follows from the prime number theorem, but there are also easier proofs for this weaker bound, see e.g. [39, p. 150] for a relatively short elementary proof). In particular, we can observe that $p_1 \cdots p_n \leq p^n \leq (100n \log_2 n)^n \leq (n^2/4)^n \leq (\log_2 N)^{\sqrt{\log_2 N}} < N$.

We claim that there are powers m_1, \ldots, m_n of the primes p_1, \ldots, p_m , respectively, with $N/p \leq m_1 \cdots m_n \leq N$ and $p_i \leq m_i < N^{1/n} \cdot p_i$ for $i = 1, \ldots, n$. Indeed, for each $i = 1, \ldots, n$, start by defining m'_i to be the unique power of p_i with $N^{1/n} \leq m'_i < N^{1/n} \cdot p_i$. Note that then we have $m'_1 \cdots m'_n \geq (N^{1/n})^n = N$, and $m'_i < N^{1/n} \cdot p_i$ for $i = 1, \ldots, n$. As long as $m'_1 \cdots m'_n > N$, let us decrease the product $m'_1 \cdots m'_n$ by choosing an index iwith $m'_i > p_i$ and dividing m'_i by p_i (again obtaining a power of p_i). In each step the product $m'_1 \cdots m'_n$ decreases by a factor of at most p. Thus, when the product $m'_1 \cdots m'_n$ stops being larger than N, it must attain a value between N/p and N. We can then define m_1, \ldots, m_n to be the values of m'_1, \ldots, m'_n at that point.

Note that m_1, \ldots, m_n are coprime (as they are powers of the distinct primes p_1, \ldots, p_n), and hence $\mathbb{Z}_{m_1 \cdots m_n} \cong \mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}$. Thus, for $m = \lfloor N^{1/n} \cdot p \rfloor$, we obtain

$$r_3(N) \ge r_3(m_1 \cdots m_n) \ge r_3(\mathbb{Z}_{m_1 \cdots m_n}) = r_3(\mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_n}) \ge \frac{(7/24)^{n/2}}{10^5 m^2 n^3} \cdot m_1 \cdots m_n$$

where at the last step we used Proposition 3.1 (and at the first two steps we used that every three-term progression free subset of $\mathbb{Z}_{m_1 \dots m_n}$ gives rise to a three-term progression free subset of $\{1, \dots, m_1 \dots m_n\}$ and hence in particular to a three-term progression free subset of $\{1, \dots, N\}$, recalling that $m_1 \dots m_n \leq N$). Rewriting the right-hand side yields

$$r_3(N) \ge \frac{(7/24)^{n/2}}{10^5 m^2 n^3} \cdot m_1 \cdots m_n \ge \frac{(7/24)^{n/2}}{10^5 (N^{1/n} p)^2 n^3} \cdot (N/p) = \frac{1}{10^5 p^3 n^3} \cdot N \cdot (7/24)^{n/2} \cdot N^{-2/n}$$

Recalling $p \leq 100n \log_2 n$, and recalling our choice of n, we can conclude that

$$\begin{split} r_{3}(N) &\geq \frac{1}{10^{5}(100n\log_{2}n)^{3}n^{3}} \cdot N \cdot 2^{-\log_{2}(24/7) \cdot n/2} \cdot 2^{-2(\log_{2}N)/n} \\ &\geq \frac{1}{10^{11}n^{6}(\log_{2}n)^{3}} \cdot N \cdot 2^{-\sqrt{\log_{2}(24/7)} \cdot \sqrt{\log_{2}N} - 2} \cdot 2^{-\sqrt{\log_{2}(24/7)} \cdot \sqrt{\log_{2}N}} \\ &\geq \frac{1}{10^{12}(2\sqrt{\log_{2}N})^{6}(\log_{2}\log_{2}N)^{3}} \cdot N \cdot 2^{-2\sqrt{\log_{2}(24/7)} \cdot \sqrt{\log_{2}N}} \\ &\geq \frac{N \cdot 2^{-2\sqrt{\log_{2}(24/7)} \cdot \sqrt{\log_{2}N}}}{10^{14}(\log_{2}N)^{3}(\log_{2}\log_{2}N)^{3}}. \end{split}$$

This shows that $r_3(N) \ge N \cdot 2^{-(2\sqrt{\log_2(24/7)} + o(1)) \cdot \sqrt{\log_2 N}}$, as desired.

We remark that it is also possible to deduce Theorem 1.1 directly from Proposition 2.1, eliminating the need to use estimates for the size of the *n*-th prime as in the proof above. Indeed, following an idea of Green and Wolf [22], given N, for any even positive integer n and any $a, b \in [0,1]^n$, one can consider the map $\varphi_{a,b} : \{1,\ldots,N\} \to [0,1)^n$ defined by $\varphi_{a,b}(x) \equiv a + xb \mod 1$ for all $x \in \{1,\ldots,N\}$. For a uniformly random choice of $b \in [0,1]^n$, the probability of having some $t \in \{1,\ldots,N\}$ such that $tb \equiv v \mod 1$ for a vector $v \in \mathbb{R}^n$ with $||v||_{\infty} \leq N^{-1/n}/4$ is at most $N \cdot (N^{-1/n}/2)^n \leq 1/2$ (indeed, for every $t \in \{1,\ldots,N\}$ the vector $tb \mod 1$ is uniformly distributed over $\mathbb{R}^n/\mathbb{Z}^n$ and so the probability of having $tb \equiv v \mod 1$ for a vector $v \in \mathbb{R}^n$ with $||v||_{\infty} \leq N^{-1/n}/4$ is $(N^{-1/n}/2)^n$). So we can fix some $b \in [0,1]^n$ such that $tb \neq v \mod 1$ for any $t \in \{1,\ldots,N\}$ and any vector $v \in \mathbb{R}^n$ with $||v||_{\infty} \leq N^{-1/n}/4$, yielding a measurable subset $S \subseteq [0,1)^n$ satisfying the conditions in the proposition. Now, for any $a \in [0,1]^n$ the pre-image $\varphi_{a,b}^{-1}(S) \subseteq \{1,\ldots,N\}$ is three-term progression free. Indeed, suppose there were $x, y, z \in \varphi_{a,b}^{-1}(S)$ with x > y > z and x + z = 2y. Then we would have $\varphi_{a,b}(x) + \varphi_{a,b}(z) \equiv (a + xb) + (a + zb) = 2a + (x + z)b = 2(a + yb) = 2\varphi_{a,b}(y) \mod 1$ and hence $|\varphi_{a,b}(x) - \varphi_{a,b}(z)|_{<} \delta$ for $i = 1, \ldots, n$ by the conditions on S (noting that $\varphi_{a,b}(x), \varphi_{a,b}(z), \varphi_{a,b}(z) \in S$), so $||\varphi_{a,b}(x) - \varphi_{a,b}(z)|_{<} < \delta = N^{-1/n}/4$. On the other hand, $(x - z)b = (a + xb) - (a + zb) \equiv \varphi_{a,b}(x) - \varphi_{a,b}(z)$ and 1, which yields a contradiction to our choice of b (as $x - z \in \{1, \ldots, N\}$). Thus, the pre-image $\varphi_{a,b}^{-1}(S) \subseteq \{1,\ldots, N\}$ is indeed three-term progression free for any $a \in [0,1]^n$. For uniformly random $a \in [0,1]^n$, the expected size of this pre-image is $\mathbb{E}[|\varphi_{a,b}^{-1}(S)|| = N \cdot \mu(S) \ge N \cdot 10^{-5}\delta^2 n^{-3} \cdot (7/24)^{n/2}$, since for any $x \in \{1,\ldots, N\}$ the vector $\varphi_{a,$

$$r_3(N) \ge N \cdot 10^{-5} \delta^2 n^{-3} \cdot (7/24)^{n/2} \ge \frac{1}{10^7 n^3} \cdot N \cdot (7/24)^{n/2} \cdot N^{-2/n}$$

for any choice of n. Optimizing n, we can again take n as in (3.1) and obtain

$$\begin{aligned} r_3(N) &\geq \frac{1}{10^7 (2\sqrt{\log_2 N})^3} \cdot N \cdot 2^{-\sqrt{\log_2 (24/7)} \cdot \sqrt{\log_2 N} - 2} \cdot 2^{-\sqrt{\log_2 (24/7)} \cdot \sqrt{\log_2 N}} \\ &\geq \frac{N \cdot 2^{-2\sqrt{\log_2 (24/7)} \cdot \sqrt{\log_2 N}}}{10^9 (\log_2 N)^{3/2}} \end{aligned}$$

if N is sufficiently large. This again shows $r_3(N) \ge N \cdot 2^{-(2\sqrt{\log_2(24/7)} + o(1)) \cdot \sqrt{\log_2 N}}$, with a slightly better bound for the o(1)-term compared to the previous proof.

4 Construction of the building blocks

In this section we prove Proposition 2.2, establishing the existence of our desired two-dimensional building blocks for our overall construction. Throughout this section we fix $0 < \varepsilon < 1$, as in the statement of Proposition 2.2.

We now define the desired subset $T \subseteq [0, 1)^2$ as follows.



Figure 4.1: The set T defined in Definition 4.1

Definition 4.1. Let us define

$$T_{1} = \left\{ (a,b) \in \left[\frac{1}{2},1\right) \times \left[0,1\right) \middle| \frac{2}{3} < a+b \le \frac{7}{6} \right\}$$

$$T_{2} = \left\{ (a,b) \in \left[\frac{1}{2},1\right) \times \left[0,\frac{1}{2}\right) \middle| \frac{7}{6} + \varepsilon \le a+b \le \frac{17}{12} \right\}$$

$$T_{3} = \left\{ (a,b) \in \left[0,\frac{1}{2}\right) \times \left[\frac{1}{2},1\right) \middle| \frac{7}{6} + \varepsilon \le a+b \le \frac{17}{12} \text{ and } 2a+b \ge \frac{3}{2} + \varepsilon \right\}$$

and $T = T_1 \cup T_2 \cup T_3 \subseteq [0,1)^2$.

For an illustration of these sets, see Figure 4.1. Note that the area $\mu(T)$ of the set T can be computed as follows. First, the area of T_1 is given by

$$\mu(T_1) = \left(\frac{1}{2}\right)^2 - \frac{1}{2} \cdot \left(\frac{1}{3}\right)^2 - \frac{1}{2} \cdot \left(\frac{1}{6}\right)^2 + \frac{1}{2} \cdot \left(\frac{1}{6}\right)^2 = \left(\frac{1}{2}\right)^2 - \frac{1}{2} \cdot \left(\frac{1}{3}\right)^2 = \frac{9-2}{36} = \frac{7}{36}.$$

Next, the area of T_2 is given by

$$\mu(T_2) = \frac{1}{2} \cdot \left(\frac{1}{3}\right)^2 - \frac{1}{2} \cdot \left(\frac{1}{12}\right)^2 - \mu\left(\{(a,b) \in [1/2,1) \times [0,1/2) \mid 7/6 \le a+b < 7/6 + \varepsilon\}\right)$$
$$\ge \frac{1}{2} \cdot \left(\frac{1}{3}\right)^2 - \frac{1}{2} \cdot \left(\frac{1}{12}\right)^2 - \frac{\varepsilon}{2} = \frac{16-1}{288} - \frac{\varepsilon}{2} = \frac{15}{288} - \frac{\varepsilon}{2}.$$

Finally, the area of T_3 is given by

$$\mu(T_3) = \mu\Big(\{(a,b) \in [1/4, 1/3) \times [1/2, 1) \mid 2a + b \ge 3/2 + \varepsilon\}\Big) \\ + \mu\Big(\{(a,b) \in [1/3, 1/2) \times [1/2, 1) \mid 7/6 + \varepsilon \le a + b \le 17/12\}\Big) \\ = \frac{1}{2} \cdot \Big(\frac{1}{12}\Big) \cdot \Big(\frac{1}{6}\Big) + \Big(\frac{1}{6}\Big)^2 - \frac{1}{2} \cdot \Big(\frac{1}{12}\Big)^2 + \frac{1}{2} \cdot \Big(\frac{1}{6}\Big)^2 \\ - \mu\Big(\{(a,b) \in [1/4, 1/3) \times [1/2, 1) \mid 3/2 \le 2a + b < 3/2 + \varepsilon\}\Big)$$

$$-\mu\Big(\{(a,b)\in[1/3,1/2)\times[1/2,1)\mid 7/6\le a+b\le 7/6+\varepsilon\}\Big)$$

$$\ge \frac{2+8-1+4}{288} - \frac{\varepsilon}{12} - \frac{\varepsilon}{6} = \frac{13}{288} - \frac{\varepsilon}{4}.$$

Thus, for the area of $T = T_1 \cup T_2 \cup T_3$ we obtain

$$\mu(T) = \mu(T_1) + \mu(T_2) + \mu(T_3) \ge \frac{7}{36} + \frac{15}{288} - \frac{\varepsilon}{2} + \frac{13}{288} - \frac{\varepsilon}{4} \ge \frac{56 + 15 + 13}{288} - \varepsilon = \frac{84}{288} - \varepsilon = \frac{7}{24} - \varepsilon.$$
(4.1)

It remains to show that there is a measurable function $f: T \to [0, 100/\varepsilon^2]$ such that the condition in Proposition 2.2 is satisfied. To define this function, we first need some more notation. Let us define the function $g: [0, 1) \to [0, 1/2)$ by setting

$$g(t) = \begin{cases} t^2 & \text{if } t \in [0, 1/2) \\ (t - 1/2)^2 & \text{if } t \in [1/2, 1) \end{cases}$$

for all $t \in [0, 1)$. Note that this function is measurable and satisfies $0 \le g(t) \le (1/2)^2 = 1/4$ for all $t \in [0, 1)$. Furthermore, for any $t, t' \in [0, 1)$ with $2t \equiv 2t' \mod 1$, we have g(t) = g(t'). Now, we can define the function $f: T \to [0, 100/\varepsilon^2]$ by

$$f(x) = \frac{24}{\varepsilon^2} \cdot (x_1 + x_2)^2 + 6 \cdot g(x_1)$$
(4.2)

for all $x \in T \subseteq [0,1)^2$, where we write the coordinates of x as $x = (x_1, x_2)$ as usual. Note that f is a measurable function and we indeed have $0 \le f(x) \le (24/\varepsilon^2) \cdot 2^2 + 6 \cdot 1/4 \le 100/\varepsilon^2$ for all $x \in T$. We need to show that for any $x, y, z \in T$ with $x + z \equiv 2y \mod 1$ we have

$$f(x) + f(z) \ge 2f(y) + (x_1 - z_1)^2 + (x_2 - z_2)^2.$$
(4.3)

In order to show this, we begin with some simple facts.

Fact 4.2. For any $s, t \in \mathbb{R}$, the following statements hold:

- (i) $s^2 + t^2 = 2 \cdot (s/2 + t/2)^2 + (s-t)^2/2$.
- (*ii*) $(s+t)^2 + t^2 \ge s^2/2$.
- (iii) If $t \ge 0$, then $(t+1/2)^2 \ge t^2 + 1/4$.

Proof. For (i), we observe

$$s^{2} + t^{2} = 2 \cdot \left(\frac{s+t}{2}\right)^{2} + 2 \cdot \left(\frac{s-t}{2}\right)^{2} = 2 \cdot \left(\frac{s}{2} + \frac{t}{2}\right)^{2} + \frac{(s-t)^{2}}{2}.$$

Plugging s + t and t into this equation, we obtain

$$(s+t)^2 + t^2 = 2 \cdot \left(\frac{s+t}{2} + \frac{t}{2}\right)^2 + \frac{s^2}{2} \ge \frac{s^2}{2},$$

showing (ii). For (iii), we simply note that

$$\left(t + \frac{1}{2}\right)^2 = t^2 + t + \frac{1}{4} \ge t^2 + \frac{1}{4}$$

if $t \geq 0$.

Fact 4.3. For the set $T \subseteq [0,1)^2$ in Definition 4.1, the following statements hold:

(i) For all $(a, b) \in T$, we have $2/3 < a + b \le 17/12$.

- (*ii*) For all $(a, b) \in T$, we have $g(a) \ge (a 1/2)^2$.
- (iii) For all $(a, b), (a', b') \in T$ with a + a' < 1, we have a + b + a' + b' > 11/6.

Proof. Note that (i) follows immediately from the definition of T.

To show (ii), first note that $g(a) = (a - 1/2)^2$ for all $a \in [1/2, 1)$. So we have $g(a) = (a - 1/2)^2$ for all $(a, b) \in T_1 \cup T_2$. It remains to consider $(a, b) \in T_3$, then $g(a) = a^2$. Furthermore, by $2a + b \ge 3/2 + \varepsilon > 3/2$ and b < 1, we have a > 1/4 and hence $g(a) = a^2 > a^2 - a + 1/4 = (a - 1/2)^2$.

It remains to show (iii), so consider $(a, b), (a', b') \in T$ with a+a' < 1. Then we must have a < 1/2 or a' < 1/2, so let us assume a' < 1/2 without loss of generality. This implies $(a', b') \in T_3$ and hence $a'+b' \ge 7/6+\varepsilon > 7/6$ by the definition of T_3 . Combining this with the lower bound in (i), we obtain

$$a + b + a' + b' > \frac{2}{3} + \frac{7}{6} = \frac{11}{6}.$$

We can now derive various consequences for points $x, y, z \in T$ such that $x + z \equiv 2y \mod 1$, with the aim of proving (4.3).

Lemma 4.4. Let $T \subseteq [0,1)^2$ be defined as in Definition 4.1. Consider points $x, y, z \in T$ such that $x + z \equiv 2y \mod 1$. Then we have either

$$y_1 + y_2 = \frac{x_1 + x_2}{2} + \frac{z_1 + z_2}{2}$$
$$x_1 + x_2 \quad z_1 + z_2$$

or

$$y_1 + y_2 = \frac{x_1 + x_2}{2} + \frac{z_1 + z_2}{2} - \frac{1}{2}$$

Proof. Let $y' = (y'_1, y'_2) \in [0, 1)^2$ be the mid-point of $x = (x_1, x_2)$ and $z = (z_1, z_2)$ (given by $y'_1 = (x_1 + z_1)/2$ and $y'_2 = (x_2 + z_2)/2$). Note that

$$(2y_1, 2y_2) = 2y \equiv x + z = (x_1, x_2) + (z_1, z_2) = (x_1 + z_1, x_2 + z_2) = (2y'_1, 2y'_2) \mod 1$$

meaning that $2y_1 \equiv 2y'_1 \mod 1$ and $2y_2 \equiv 2y'_2 \mod 1$. Hence $y_1 - y'_1 \in \{-1/2, 0, 1/2\}$ and $y_2 - y'_2 \in \{-1/2, 0, 1/2\}$. Since $(y_1, y_2) \notin [0, 1/2)^2$ (as $T \cap [0, 1/2)^2 = \emptyset$), we cannot have both $y_1 - y'_1 = -1/2$ and $y_2 - y'_2 = -1/2$. So we can conclude that $(y_1 + y_2) - (y'_1 + y'_2) \in \{-1/2, 0, 1/2, 1\}$.

The assertion of the lemma is that $(y_1 + y_2) - (y'_1 + y'_2) \in \{-1/2, 0\}$. Hence it suffices to prove $(y_1 + y_2) - (y'_1 + y'_2) < 1/2$. So let us assume for contradiction that $(y_1 + y_2) - (y'_1 + y'_2) \ge 1/2$.

Since $y_2 - y'_2 \le 1/2$, we must therefore have $y_1 \ge y'_1$. Similarly, we must have $y_2 \ge y'_2$, since $y_1 - y'_1 \le 1/2$. If $y'_1 < 1/2$ (i.e. if $x_1 + z_1 < 1$), by Fact 4.3(iii) we have

$$y_1' + y_2' = \frac{(x_1 + x_2) + (z_1 + z_2)}{2} > \frac{11/6}{2} = \frac{11}{12} = \frac{17}{12} - \frac{1}{2} \ge y_1 + y_2 - \frac{1}{2}$$

where the last inequality follows from Fact 4.3(i). This contradicts $(y_1 + y_2) - (y'_1 + y'_2) \ge 1/2$, so we must have $y'_1 \ge 1/2$.

Now, we obtain $1/2 \le y'_1 \le y_1 < 1$. Together with $y_1 - y'_1 \in \{-1/2, 0, 1/2\}$, this implies $y_1 = y'_1$. So from $(y_1 + y_2) - (y'_1 + y'_2) \ge 1/2$, we obtain $y_2 \ge y'_2 + 1/2 \ge 1/2$.

Thus, we have $y_1 = y'_1 \ge 1/2$ and $y_2 \ge 1/2$, meaning that $(y_1, y_2) \in [1/2, 1)^2$. By the definition of T, this implies $y_1 + y_2 \le 7/6$ and hence

$$y'_1 + y'_2 = \frac{(x_1 + x_2) + (z_1 + z_2)}{2} > \frac{2/3 + 2/3}{2} = \frac{2}{3} = \frac{7}{6} - \frac{1}{2} \ge y_1 + y_2 - \frac{1}{2},$$

where the first inequality follows from Fact 4.3(i). This is again a contradiction to $(y_1 + y_2) - (y'_1 + y'_2) \ge 1/2$.

Corollary 4.5. Let $T \subseteq [0,1)^2$ be defined as in Definition 4.1, and let $x, y, z \in T$ be points such that $x + z \equiv 2y \mod 1$. Then at least one of the following two statements holds

(a) $(x_1 + x_2)^2 + (z_1 + z_2)^2 \ge 2 \cdot (y_1 + y_2)^2 + \varepsilon^2/2.$ (b) $|x_1 + x_2 - z_1 - z_2| < \varepsilon$ and $(x_1 + x_2)^2 + (z_1 + z_2)^2 = 2 \cdot (y_1 + y_2)^2 + (x_1 + x_2 - z_1 - z_2)^2/2.$

Proof. By Lemma 4.4, either $y_1 + y_2 = (x_1 + x_2)/2 + (z_1 + z_2)/2$ or $y_1 + y_2 = (x_1 + x_2)/2 + (z_1 + z_2)/2 - 1/2$ must hold. In the latter case, we have

$$(x_1+x_2)^2 + (z_1+z_2)^2 \ge 2 \cdot \left(\frac{x_1+x_2}{2} + \frac{z_1+z_2}{2}\right)^2 = 2 \cdot (y_1+y_2+1/2)^2 \ge 2 \cdot (y_1+y_2)^2 + 1/2 \ge 2 \cdot (y_1+y_2)^2 + \varepsilon^2/2,$$

where the first inequality follows from Fact 4.2(i) and the second inequality from Fact 4.2(iii). So let us now assume that $y_1 + y_2 = (x_1 + x_2)/2 + (z_1 + z_2)/2$, then by Fact 4.2(i) we have

$$(x_1+x_2)^2 + (z_1+z_2)^2 = 2 \cdot \left(\frac{x_1+x_2}{2} + \frac{z_1+z_2}{2}\right)^2 + (x_1+x_2-z_1-z_2)^2/2 = 2 \cdot (y_1+y_2)^2 + (x_1+x_2-z_1-z_2)^2/2.$$

If $|x_1 + x_2 - z_1 - z_2| \ge \varepsilon$, this implies the inequality in (a). And if $|x_1 + x_2 - z_1 - z_2| < \varepsilon$, then (b) holds.

Recall that our goal is to prove (4.3) for any $x, y, z \in T$ with $x + z \equiv 2y \mod 1$. If (a) in Corollary 4.5 holds, then it is not hard to show inequality (4.3). The next lemma helps with showing the inequality in case (b) in Corollary 4.5 holds.

Lemma 4.6. Let $T \subseteq [0,1)^2$ be defined as in Definition 4.1, and let $x, z \in T$ be such that $|x_1+x_2-z_1-z_2| < \varepsilon$. Suppose that $x_1 \ge 1/2$ or $z_1 \ge 1/2$. Then we have $x_1 + z_1 \ge 1$.

Proof. Recall that for all points $(a, b) \in T_1$ we have $a + b \leq 7/6$, whereas for all points $(a, b) \in T_2 \cup T_3$ we have $a + b \geq 7/6 + \varepsilon$. Since $|x_1 + x_2 - z_1 - z_2| < \varepsilon$, this means that we either have $x, z \in T_1$ or $x, z \in T_2 \cup T_3$. If $x, z \in T_1$, then we have $x_1 \geq 1/2$ and $z_1 \geq 1/2$ and hence $x_1 + z_1 \geq 1$, as desired. So let us assume that $x, z \in T_2 \cup T_3$. If $x, z \in T_2$, we similarly have $x_1 \geq 1/2$ and $z_1 \geq 1/2$ and $z_1 \geq 1/2$ and hence $x_1 + z_1 \geq 1$. If $x, z \in T_3$, then we would have $x_1 < 1/2$ and $z_1 < 1/2$, which contradicts the assumption in the lemma.

So it only remains to consider the case that one of the points x and z is contained in the set T_2 , and the other point in the set T_3 . Let us assume without loss of generality that $x \in T_2$ and $z \in T_3$. Then we have

$$x_1 + z_1 = x_1 + 2z_1 + z_2 - (z_1 + z_2) \ge x_1 + 3/2 + \varepsilon - (z_1 + z_2) \ge x_1 + 3/2 + \varepsilon - (x_1 + x_2 + \varepsilon) = 3/2 - x_2 \ge 1,$$

where the first inequality follows from the definition of T_3 , the second inequality follows from the assumption $|x_1 + x_2 - z_1 - z_2| < \varepsilon$, and the third inequality follows from the definition of T_2 .

Corollary 4.7. Let $T \subseteq [0,1)^2$ be defined as in Definition 4.1, and let $x, y, z \in T$ be such that $x + z \equiv 2y \mod 1$ and $|x_1 + x_2 - z_1 - z_2| < \varepsilon$. Then we have

$$g(x_1) + g(z_1) \ge 2 \cdot g(y_1) + (x_1 - z_1)^2/2.$$

Proof. Since $x + z \equiv 2y \mod 1$, we have $x_1 + z_1 \equiv 2y_1 \mod 1$ and hence $2(x_1/2 + z_1/2) \equiv 2y_1 \mod 1$. So we have $g(y_1) = g(x_1/2 + z_1/2)$, and therefore the claimed inequality is equivalent to

$$g(x_1) + g(z_1) \ge 2 \cdot g(x_1/2 + z_1/2) + (x_1 - z_1)^2/2.$$

If $x_1 \ge 1/2$ or $z_1 \ge 1/2$, then by Lemma 4.6 we have $x_1/2 + z_1/2 \ge 1/2$ and hence $g(x_1/2 + z_1/2) = (x_1/2 + z_1/2 - 1/2)^2$. Thus, by Fact 4.3(ii) and Fact 4.2(i), in this case we obtain

$$g(x_1) + g(z_1) \ge (x_1 - 1/2)^2 + (z_1 - 1/2)^2 = 2 \cdot (x_1/2 + z_1/2 - 1/2)^2 + (x_1 - z_1)^2/2 = 2 \cdot g(x_1/2 + z_1/2) + (x_1 - z_1)^2/2,$$

as desired.

It remains to consider the case that $x_1 < 1/2$ and $z_1 < 1/2$, then we also have $x_1/2 + z_1/2 < 1/2$. Thus, observing that $g(x_1) = x_1^2$ and $g(z_1) = z_1^2$ and $g(x_1/2 + z_1/2) = (x_1/2 + z_1/2)^2$, by Fact 4.2(i) we obtain

$$g(x_1) + g(z_1) = x_1^2 + z_1^2 = 2 \cdot (x_1/2 + z_1/2)^2 + (x_1 - z_1)^2/2 = 2 \cdot g(x_1/2 + z_1/2) + (x_1 - z_1)^2/2.$$

Finally, we are ready to prove Proposition 2.2.

Proof of Proposition 2.2. Let $T \subseteq [0,1)^2$ be given as in Definition 4.1, and let $f: T \to [0,100/\varepsilon^2]$ be the measurable function defined in (4.2). In order to prove the proposition it remains to show the inequality

$$f(x) + f(z) \ge 2 \cdot f(y) + (x_1 - z_1)^2 + (x_2 - z_2)^2$$

for $x, y, z \in T \subseteq [0,1)^2$ with $x + z \equiv 2y \mod 1$. For any such $x, y, z \in T$, one of the two statements in Corollary 4.5 holds. If (a) holds, then we have

$$\begin{split} f(x) + f(z) &= \frac{24}{\varepsilon^2} \cdot (x_1 + x_2)^2 + 6 \cdot g(x_1) + \frac{24}{\varepsilon^2} \cdot (z_1 + z_2)^2 + 6 \cdot g(z_1) \\ &\geq \frac{24}{\varepsilon^2} \cdot \left((x_1 + x_2)^2 + (z_1 + z_2)^2 \right) \\ &\geq \frac{24}{\varepsilon^2} \cdot \left(2 \cdot (y_1 + y_2)^2 + \varepsilon^2 / 2 \right) \\ &= 2 \cdot \frac{24}{\varepsilon^2} \cdot (y_1 + y_2)^2 + 12 \\ &\geq 2 \cdot \frac{24}{\varepsilon^2} \cdot (y_1 + y_2)^2 + 2 \cdot 6 \cdot (1/4) + 1 + 1 \\ &\geq 2 \cdot \frac{24}{\varepsilon^2} \cdot (y_1 + y_2)^2 + 2 \cdot 6 \cdot g(y_1) + (x_1 - z_1)^2 + (x_2 - z_2)^2 \\ &= 2 \cdot f(y) + (x_1 - z_1)^2 + (x_2 - z_2)^2. \end{split}$$

If (b) in Corollary 4.5 holds, then in particular by Corollary 4.7 we have

$$g(x_1) + g(z_1) \ge 2 \cdot g(y_1) + (x_1 - z_1)^2/2.$$

Together with the equation in (b), this yields

$$\begin{split} f(x) + f(z) &= \frac{24}{\varepsilon^2} \cdot (x_1 + x_2)^2 + 6 \cdot g(x_1) + \frac{24}{\varepsilon^2} \cdot (z_1 + z_2)^2 + 6 \cdot g(z_1) \\ &= \frac{24}{\varepsilon^2} \cdot \left((x_1 + x_2)^2 + (z_1 + z_2)^2 \right) + 6 \cdot \left(g(x_1) + g(z_1) \right) \\ &\geq \frac{24}{\varepsilon^2} \cdot \left(2 \cdot (y_1 + y_2)^2 + (x_1 + x_2 - z_1 - z_2)^2 / 2 \right) + 6 \cdot \left(2 \cdot g(y_1) + (x_1 - z_1)^2 / 2 \right) \\ &= 2 \cdot \left(\frac{24}{\varepsilon^2} \cdot (y_1 + y_2)^2 + 6 \cdot g(y_1) \right) + \frac{12}{\varepsilon^2} \cdot (x_1 + x_2 - z_1 - z_2)^2 + 3 \cdot (x_1 - z_1)^2 \\ &\geq 2 \cdot f(y) + (x_1 - z_1)^2 + 2 \cdot (x_1 + x_2 - z_1 - z_2)^2 + 2 \cdot (x_1 - z_1)^2 \\ &\geq 2 \cdot f(y) + (x_1 - z_1)^2 + (x_2 - z_2)^2, \end{split}$$

where in the last step we used Fact 4.2(ii). In either case we proved the desired inequality, finishing the proof of Proposition 2.2. $\hfill \Box$

References

- N. Alon, A. Shpilka, and C. Umans, On sunflowers and matrix multiplication, Comput. Complexity 22 (2013), 219–243.
- F. A. Behrend, On sets of integers which contain no three terms in arithmetical progression, Proc. Nat. Acad. Sci. U.S.A. 32 (1946), 331–332.
- [3] J. Blasiak, T. Church, H. Cohn, J. A. Grochow, E. Naslund, W. F. Sawin, and C. Umans, *On cap sets and the group-theoretic approach to matrix multiplication*, Discrete Anal. 2017, Paper No. 3, 27pp.

- [4] T. F. Bloom and O. Sisask, Breaking the logarithmic barrier in Roth's theorem on arithmetic progressions, preprint, 2020, arXiv:2007.03528.
- [5] T. F. Bloom and O. Sisask, An improvement to the Kelley-Meka bounds on three-term arithmetic progressions, preprint, 2023, arXiv:2309.02353.
- [6] J. Bourgain, On triples in arithmetic progression, Geom. Funct. Anal. 9 (1999), 968–984.
- [7] J. Bourgain, Roth's theorem on progressions revisited, J. Anal. Math. 104 (2008), 155–192.
- [8] T. C. Brown and J. P. Buhler, A density version of a geometric Ramsey theorem, J. Combin. Theory Ser. A 32 (1982), 20–34.
- [9] A. R. Calderbank and P. C. Fishburn, Maximal three-independent subsets of {0,1,2}ⁿ, Des. Codes Cryptogr. 4 (1994), 203-211.
- [10] A. K. Chandra, M. L. Furst, and R. J. Lipton, *Multi-party protocols*, Proceedings of the Fifteenth Annual ACM Symposium on Theory of Computing (STOC '83), New York, NY, USA, 1983, 94–99.
- [11] E. Croot, V. F. Lev, and P. P. Pach, Progression-free sets in Zⁿ₄ are exponentially small, Ann. of Math. 185 (2017), 331–337.
- [12] Y. Edel, Extensions of generalized product caps, Des. Codes Cryptogr. **31** (2004), 5–14.
- [13] M. Elkin, An improved construction of progression-free sets, Israel J. Math. 184 (2011), 93–128.
- [14] J. S. Ellenberg, Sumsets and sumsets of subsets, blog post, 2016, https://quomodocumque.wordpress. com/2016/05/29/sumsets-and-sumsets-of-subsets/.
- [15] J. S. Ellenberg and D. Gijswijt, On large subsets of \mathbb{F}_q^n with no three-term arithmetic progression, Ann. of Math. **185** (2017), 339–343.
- [16] C. Elsholtz, G. F. Lipnik, and M. Siebenhofer, New constructions of large caps, Chapter 5 in Ph.D. Thesis of G. F. Lipnik, On Integer Partitions, Caps, Progression-Free Sets and q-Regular Sequences, Graz University of Technology, 2023.
- [17] C. Elsholtz and P. P. Pach, Caps and progression-free sets in \mathbb{Z}_m^n , Des. Codes Cryptogr. 88 (2020), 2133–2170.
- [18] P. Erdős and P. Turán, On Some Sequences of Integers, J. London Math. Soc. 11 (1936), 261–264.
- [19] J. Fox and H. T. Pham, Popular progression differences in vector spaces II, Discrete Anal. 2019, Paper No. 16, 39 pp.
- [20] P. Frankl, R. L. Graham, and V. Rödl, On subsets of abelian groups with no 3-term arithmetic progression, J. Combin. Theory Ser. A 45 (1987), 157–161.
- [21] B. Green, Finite field models in additive combinatorics, Surveys in combinatorics 2005, London Math. Soc. Lecture Note Ser., 327, Cambridge University Press, Cambridge, 2005, 1–27.
- [22] B. Green and J. Wolf, A note on Elkin's improvement of Behrend's construction, Additive number theory, Springer, New York, 2010, 141–144.
- [23] L. Hambardzumyan, T. Pitassi, S. Sherif, M. Shirley, A. Shraibman, An improved protocol for ExactlyN with more than 3 players, preprint, 2023, arXiv:2309.06554.
- [24] D. R. Heath-Brown, Integer sets containing no arithmetic progressions, J. London Math. Soc. 35 (1987), 385–394.
- [25] Z. Kelley and R. Meka, Strong bounds for 3-Progressions, 2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS), Santa Cruz, CA, USA, 2023, 933–973.

- [26] H. Lipmaa, Progression-free sets and sublinear pairing-based non-interactive zero-knowledge arguments, Theory of cryptography, Lecture Notes in Comput. Sci., 7194, Springer, Heidelberg, 2012, 169–189.
- [27] V. F. Lev, Progression-free sets in finite abelian groups, J. Number Theory 104 (2004), 162–169.
- [28] R. Meshulam, On subsets of finite abelian groups with no 3-term arithmetic progressions, J. Combin. Theory Ser. A 71 (1995), 168–172.
- [29] E. Naslund, Lower bounds for the Shannon Capacity of Hypergraphs, manuscript, 2024.
- [30] S. Peluse, *Finite field models in arithmetic combinatorics twenty years on*, preprint, 2023, arXiv:2312.08100.
- [31] F. Petrov, C. Pohoata, Improved bounds for progression-free sets in C_8^n , Israel J. Math. **236** (2020), 345–363.
- [32] B. Romera-Paredes, M. Barekatain, A. Novikov, M. Balog, M. P. Kumar, E. Dupont, F. J. R. Ruiz, J. S. Ellenberg, P. Wang, O. Fawzi, P. Kohli, and A. Fawzi, *Mathematical discoveries from program* search with large language models, Nature 625 (2024), 468–475.
- [33] K. F. Roth, On certain sets of integers, J. London Math. Soc. 28 (1953), 104–109.
- [34] S. Ruziewicz, Contribution à l'étude des ensembles des distances des points, Fund. Math. 7 (1925), 141–143.
- [35] R. Salem and D. C. Spencer, On sets of integers which contain no three terms in arithmetical progression, Proc. Nat. Acad. Sci. U.S.A. 28 (1942), 561–563.
- [36] R. Salem and D.C. Spencer, On sets which do not contain a given number of terms in arithmetical progression, Nieuw Arch. Wiskunde (2), 23 (1950), 133–143.
- [37] T. Sanders, On certain other sets of integers, J. Anal. Math. 116 (2012), 53–82.
- [38] T. Sanders, On Roth's theorem on progressions, Ann. of Math. 174 (2011), 619–636.
- [39] W. Sierpiński, *Elementary theory of numbers*, Translated from Polish by A. Hulanicki, Monogr. Mat., Tom 42, Państwowe Wydawnictwo Naukowe, Warsaw, 1964.
- [40] E. Szemerédi, Integer sets containing no arithmetic progressions, Acta Math. Hungar. 56 (1990), 155– 158.
- [41] T. Tao and V. H. Vu, Additive Combinatorics, Cambridge University Press, 2006.
- [42] F. Tyrrell, New lower bounds for cap sets, Discrete Anal. 2023, Paper No. 20, 18pp.
- [43] J. Wolf, Finite field models in arithmetic combinatorics ten years on, Finite Fields Appl. 32 (2015), 233–274.