The modulated soliton fields in the Goldstone boson model

Z.Shahbazi¹, S. Miraboutalebi¹ and F. Ahmadi^{2*}

¹Department of Physics, Islamic Azad University, North Tehran Branch, Tehran, 1651153311, Iran. ²Department of Physics, Shahid Rajaee Teacher Training University, Tehran 1678815811, Iran.

June 19, 2024

Abstract

It is well known that massless Goldstone bosons have not yet been observed. In the Goldstone boson model, after the spontaneous symmetry breaking under U(1), two coupled nonlinear equations are obtained for which we present the exact solitonic solutions. These solutions completely localize the energy density of the model and show the existence of two boson fields, one massive and the other massless. Also, it is seen that the solitonic waves are modulated and the massless wave rides on the massive wave. Finally, we calculate the charge density of the model, which again confirms the neutrality of these boson fields. Since massive bosons can be observed in the laboratory, these solitonic waves may be useful in tracking and detecting Goldstone bosons.

Keywords: Goldstone boson model; Spontaneous symmetry breaking; Exact solutions; Soliton boson fields.

1 Introduction

Symmetry plays an important role in our understanding of nature. Since the birth of science endeavor, symmetry structures in nature have attracted the minds of natural philosophers and physicists, looking for the origin of the laws of nature [1]. In some theories, spontaneous symmetry breaking (SSB) happens when the ground state does not respect the symmetry of the theory [2]. One such theory is that of the Goldstone boson model with a degenerate ground state. The symmetry of the model spontaneously breaks under continuous transformation $\phi(x,t) \rightarrow \phi(x,t)e^{i\alpha}$ and leads to the appearance of two spin-zero particles, a massless and a massive boson. The massless boson is known as the Nambu-Goldstone boson [3]. Since massless bosons have not any reasonable interaction strength, they cannot be observed easily, but the massless particles (pseudo-Nambu-Goldstone bosons) have many laboratory, astrophysical and cosmological implications [4, 5]. For example, they are considered as the generator of cosmological dark energy potential which causes present cosmic acceleration, so expansion of the universe can result from such particles [6, 7].

There are numerous reviews of the physics of Nambu-Goldstone bosons and their effective field theory [8]-[12]. The Goldstone boson model is the simplest model to undergo SSB which involves a complex scalar field with a self-interaction term described by the potential function $\mathcal{V}(\phi) = \mu^2 |\phi(x)|^2 + \lambda |\phi(x)|^4$. Regardless of the $\lambda |\phi(x)|^4$ term that ($\lambda > 0$), the govering equation of the model is the same as complex Klein-Gordon equation. In the Goldstone boson model after SSB, the complex field can be defined by two real fields which are derivatives around the ground state, so the Lagrangian of the model is defined according the fields that lead to two coupled nonlinear equations. The equations have already been treated by perturbation methods but suffer from problems such as inequality of the lagrangian with the initial lagrangian. The equivalence of Lagrangians holds only for exact

^{*}fahmadi@sru.ac.ir

solutions of the theory [13, 14]. In this work, the exact solutions of the coupled nonlinear equations are introduced by using tanh/sech method for which soliton solutions appear. Soliton solutions of the nonlinear partial differential equations (NPDEs) can illustrate many phenomena in physics, they are created as a result of a neutralization of the nonlinear and dispersive effects. The interesting feature of a soliton is that, after collision with another soliton it keeps its properties which is similar to that of a particles [15].

The tanh method is a powerful technique to obtain traveling wave solutions [16]-[28]. In other words, the method is appropriate to solve equations where dispersion, convection and reaction diffusion phenomena play an important role in such models [29]. For example the method is used to solve nonlinear Klein-Gordon equation [30]-[32], nonlinear fifth-order KDV equation [33] and nonlinear conduction and Burgers-Fisher equations [34]. The extended tanh method [35, 36] and the modified extended tanh-function method [37, 38] are the extended version of this approach.

In the Goldstone boson model to obtain exact solutions, at first we find the field equations in the form of two coupled (NPDEs), then by using the tanh method we convert the equations to nonlinear ordinary differential equations (NODEs). By solving the equations we get solitary solutions for which we choose the localized solutions. The soliton solutions are plotted and two modulated waves are observable in the figure for which the massless one rides on the massive one. Finally, the Hamiltonian density and charge density of the model are calculated. The figures of the energy density show that two localized fields exist in the model. It can also be seen that the charge density becomes zero, so the total charge and the electric dipole moment of the model become zero, which corresponds to the neutrality of the produced bosons. Since massive bosons can be observed in the laboratory, these solitonic waves may be useful in designing experiments to detect massless bosons. So, these studies could impact our understanding of dark matter, the problem of CP violation and the hierarchy problem [39].

This work organized as follows. In section two, the model is briefly presented and the field equations are derived. Also, the Hamiltonian density and charge-current density of the model are written. In section three, using the tanh/sech method, the soliton solutions are obtained and the figure of these modulated waves is plotted. In the first subsection of this section, the Hamiltonian density is determined which its figures show the localized energy density and two peaks close to each other. In the second subsection, the charge-current density of the model is calculated. Finally the conclusion are drawn in the last section. The relations and the model are written in natural units, namely, in $\hbar = c = m_e = \epsilon_0 = 1$ and the metric signature is (-, +, +, +).

2 The Model

The model is described by the Lagrangian density

$$\mathcal{L} = \partial^{\alpha} \phi^*(x^{\beta}) \partial_{\alpha} \phi(x^{\beta}) - \mu^2 |\phi(x^{\beta})|^2 - \lambda |\phi(x^{\beta})|^4 \,, \quad (\lambda > 0) \,, \tag{1}$$

where $\phi(x)$ is a classical complex scalar field, μ^2 and λ are arbitrary real parameters. It is obvious the Lagrangian density without interaction term $\lambda |\phi(x^{\beta})|^4$ is the same as the free complex Klein-Gordon field. The Lagrangian density is invariant under the following global transformations [3]

$$\phi(x^{\beta}) \longrightarrow \dot{\phi}(x^{\beta}) = \phi(x^{\beta})e^{i\gamma}, \qquad \phi^*(x^{\beta}) \longrightarrow \phi^*(x^{\beta}) = \phi^*(x)e^{-i\gamma}.$$
(2)

In order to find the ground state, minimum of the potential density must be obtained, so two different situations occur, depending on the sign of μ^2 . The potential density at $\phi = 0$ has a absolute minimum for $\mu^2 > 0$, it is clear that the (SSB) can not occur in this situation. For $\mu^2 < 0$, the potential density has a maximum at $\phi = 0$ and a set of absolute minimum at $\phi(x) = \phi_0 = (\frac{-\mu^2}{2\lambda})^{1/2} e^{i\theta}$, $0 \le \theta < 2\pi$. Since the Eq.(1) is invariance under the global phase transformation, the value of θ chosen is not important and it can be taken zero, therefore

$$\phi_0 = \left(\frac{-\mu^2}{2\lambda}\right)^{1/2} = \frac{1}{\sqrt{2}} \ \nu \,, \quad (\nu > 0) \,, \tag{3}$$

where $\nu = (\frac{-\mu^2}{\lambda})^{\frac{1}{2}}$ and ϕ_0 is real. It can be seen the (SSB) occurs by choosing one particular direction. The field $\phi(x^{\beta})$ around the minimum of the potential can be written as follows

$$\phi(x^{\beta}) = \frac{1}{\sqrt{2}} \left[\nu + \psi_1(x^{\beta}) + i\psi_2(x^{\beta}) \right], \qquad (4)$$

where $\psi_1(x^{\beta})$ and $\psi_2(x^{\beta})$ are real and measure the deviations of the field $\phi(x^{\beta})$ from the equilibrium ground state ϕ_0 . In turns of $\psi_1(x^{\beta})$ and $\psi_2(x^{\beta})$ the Lagrangian density become

$$\mathcal{L} = \frac{1}{2} \partial^{\alpha} \psi_{1}(x^{\beta}) \partial_{\alpha} \psi_{1}(x^{\beta}) - \lambda \nu^{2} \psi_{1}^{2}(x^{\beta}) + \frac{1}{2} \partial^{\alpha} \psi_{2}(x^{\beta}) \partial_{\alpha} \psi_{2}(x^{\beta}) - \lambda \nu \psi_{1}(x^{\beta}) \left[\psi_{1}^{2}(x^{\beta}) + \psi_{2}^{2}(x^{\beta}) \right] - \frac{1}{4} \lambda \left[\psi_{1}^{2}(x^{\beta}) + \psi_{2}^{2}(x^{\beta}) \right]^{2}.$$
(5)

According to the above Lagrangian, it is clear that the fields $\psi_1(x^\beta)$ and $\psi_2(x^\beta)$ are real Klein-Gordon fields and in the quantization process, both fields lead to neutral spin 0 particles that the ψ_1 boson with the mass $(\sqrt{2\lambda\nu^2})$ and the ψ_2 boson which has zero mass [3]. On the other hands, Eq.(1) and Eq.(5) are the same Lagrangian density expressed in terms of different variables. So, they are entirely equivalent and must lead to the same physical results. This equivalence only holds for exact solutions of the theory, but in perturbation theory, approximate solutions lead to a very different result [3]. In this work, we find the exact solutions for the above Lagrangian density and can get interesting information about the fields. Applying the Euler-Lagrange equation to the Lagrangian density (5), we get the following relations

$$\frac{1}{2}\Box\psi_1(x^\beta) + 2\lambda\nu^2\psi_1(x^\beta) + 3\lambda\nu\psi_1^2(x^\beta) + \lambda\nu\psi_2^2(x^\beta) + \lambda\psi_1(x^\beta)\left[\psi_1^2(x^\beta) + \psi_2^2(x^\beta)\right] = 0, \quad (6)$$

and

$$\frac{1}{2}\Box\psi_2(x^\beta) + 2\lambda\nu\psi_1(x^\beta)\psi_2(x^\beta) + \lambda\psi_2(x^\beta)\left[\psi_1^2(x^\beta) + \psi_2^2(x^\beta)\right] = 0,$$
(7)

since the above equations are coupled nonlinear equations, to get their exact solutions we use the tanh approach and obtain the soliton solutions in the next section. For energy density of the model, we calculate the following Hamiltonian density

$$\mathcal{H} = \frac{1}{2} \left[\dot{\psi_1}^2(x^\beta) + \dot{\psi_2}^2(x^\beta) + (\nabla \psi_1(x^\beta))^2 + (\nabla \psi_2(x^\beta))^2 \right] + \lambda \nu^2 \psi_1^2(x^\beta) + \lambda \nu \psi_1(x^\beta) \\ \times \left[\psi_1^2(x^\beta) + \psi_2^2(x^\beta) \right] + \frac{\lambda}{4} \left[\psi_1^2(x^\beta) + \psi_2^2(x^\beta) \right]^2 ,$$
(8)

and for charge-current density, using the definition (4), we have

$$J_{\alpha}(x^{\beta}) = q \left[-\nu \partial_{\alpha} \psi_2(x^{\beta}) - \partial_{\alpha} \psi_2(x^{\beta}) \psi_1(x^{\beta}) + \partial_{\alpha} \psi_1(x^{\beta}) \psi_2(x^{\beta}) \right],$$
(9)

which obviously satisfies the continuity equation, $\partial^{\alpha} J_{\alpha}(x^{\beta}) = 0$.

3 Soliton solutions

In this section, we are going to solve the coupled nonlinear Eqs.(6) and (7) in one dimension using the tanh approach. For this purpose, we first consider the following variable

$$\xi = g(x - wt), \tag{10}$$

where w is velocity of the traveling waves and g is wave number, that its inverse is proportional to width of the traveling waves. Then we replace ψ_1 and ψ_2 with

$$\psi_1(x,t) = V(\xi), \qquad \psi_2(x,t) = U(\xi),$$
(11)

where $V(\xi)$ and $U(\xi)$ will be the solitary solutions for the massive and massless fields respectively. Any static localized solution of the equations is a solitary wave which it can be transformed to a moving coordinate frame by a simple boost, so an advanced mode (x + wt) or a retarded mode (x - wt) is obtained and because of the nonlinearity, a simple linear combination of both of these modes does not provide a proper solution [40, 41]. By considering the above definitions, the Eqs.(6) and (7) become

$$\frac{\eta}{2}\frac{d^2V}{d\xi^2} + 2\lambda\nu^2 V + 3\lambda\nu V^2 + \lambda\nu U^2 + \lambda V[V^2 + U^2] = 0, \qquad (12)$$

and

$$\frac{\eta}{2}\frac{d^2U}{d\xi^2} + 2\lambda\nu VU + \lambda U[V^2 + U^2] = 0, \qquad (13)$$

where η is a constant defined as

$$\eta = g^2 \left(1 - w^2 \right) \,. \tag{14}$$

To display the solitary solutions as polynomials of sech function, the following series is supposed

$$V(\xi) = \mathbf{V}(Z) = \sum_{m=0}^{M} b_m Z^m , \qquad (15)$$

and

$$U(\xi) = \mathbf{U}(Z) = \sum_{n=0}^{N} a_n Z^n,$$
 (16)

where $Z = \operatorname{sech}(\xi)$. By putting the above series in Eqs.(12) and (13), we have

$$-\frac{\eta}{2} \left(\frac{d^2 \mathbf{V}(Z)}{dZ^2} \right) Z^4 - \eta \left(\frac{d \mathbf{V}(Z)}{dZ} \right) Z^3 + \eta \left(\frac{d^2 \mathbf{V}(Z)}{dZ^2} \right) Z^2 + \frac{\eta}{2} \left(\frac{d \mathbf{V}(Z)}{dZ} \right) Z + 2\lambda \nu^2 \mathbf{V}(Z)$$

+ $3\lambda \nu \mathbf{V}^2(Z) + \lambda \nu \mathbf{U}^2(Z) + \lambda \mathbf{V}(Z) \mathbf{U}^2(Z) + \lambda \mathbf{V}^3(Z) = 0,$ (17)

and

$$-\frac{\eta}{2} \left(\frac{d^2 \boldsymbol{U}(Z)}{dZ^2} \right) Z^4 - \eta \left(\frac{d \boldsymbol{U}(Z)}{dZ} \right) Z^3 + \eta \left(\frac{d^2 \boldsymbol{U}(Z)}{dZ^2} \right) Z^2 + \frac{\eta}{2} \left(\frac{d \boldsymbol{U}(Z)}{dZ} \right) Z + 2\lambda \nu \boldsymbol{V}(Z) \boldsymbol{U}(Z)$$

+ $\lambda \boldsymbol{U}(Z) \boldsymbol{V}^2(Z) + \lambda \boldsymbol{U}^3(Z) = 0.$ (18)

Now, considering Eqs.(17) and (18), parameters N and M in Eq.(15) and (16) can be found by balancing the highest-order linear term with the highest-order nonlinear term [16]. It turns out that N = M = 1, therefore we find

$$V(Z) = b_0 + b_1 Z \,, \tag{19}$$

and

$$U(Z) = a_0 + a_1 Z \,, \tag{20}$$

where b_0 , b_1 , a_0 , a_1 are four unknown parameters. Then, by inserting Eqs.(19) and (20) in the Eqs.(17) and (18), some equations of $Z = \operatorname{sech}(\xi)$ are obtained. In order to hold the equations for any Z, the coefficients for each power of Z should be zero. In this way, eight coupled equations of the unknown parameters of the theory are obtained. Solving these equations, an acceptable set of parameters can be found as follows

$$a_0 = 0, \quad a_1 = \pm \sqrt{2b_0^2 - b_1^2} \quad (2b_0^2 - b_1^2 > 0), \quad b_0 = b_0, \quad b_1 = b_1, \quad \lambda = \frac{\eta}{2b_0^2}, \quad \nu = -b_0,$$
(21)

where b_1 is free parameter but should satisfy the condition for a_1 in the above relation. Using above parameters, we get the soliton wave solutions of Eqs.(19) and (20) as follows

$$U(\xi) = \sqrt{2\nu^2 - b_1^2} \operatorname{sech}(\xi), \qquad V(\xi) = -\nu + b_1 \operatorname{sech}(\xi).$$
(22)

By substituting the above solutions in relation (4), $\phi(x^{\beta})$ is obtained as a normalizable wave function for each value of parameter b_1 . Therefore, without losing the generality of the problem, the parameter b_1 can be chosen such that the massless field is normalized. These solutions are plotted in Fig.(1), which shows both modulated solitons and the massless field rids on the massive one. It is notice that these fields must lead to the localized energy density, to be suitable for the model. So, we explore this subject in the next section.



Figure 1: The fields are plotted for $\nu = 1$ and $b_1 = 1.224$. The blue one is massless soliton and the green one is massive soliton.

3.1 The energy density of the model

For the energy density, using Eq.(8), we calculate the Hamiltonian density as follows

$$\mathcal{H}(\xi, w, g) = \frac{1}{2} \left[\left(\frac{\partial \psi_1(\xi)}{\partial \xi} \frac{\partial \xi}{\partial t} \right)^2 + \left(\frac{\partial \psi_2(\xi)}{\partial \xi} \frac{\partial \xi}{\partial t} \right)^2 + \left(\frac{\partial \psi_1(\xi)}{\partial \xi} \frac{\partial \xi}{\partial x} \right)^2 + \left(\frac{\partial \psi_2(\xi)}{\partial \xi} \frac{\partial \xi}{\partial x} \right)^2 \right] + \lambda \nu^2 \psi_1^2(\xi) \right]$$

$$(23)$$

By putting the obtained fields (22) in the above relationship, we have

$$\mathcal{H}(\xi, w, g) = \frac{g^2(w^2 + 1)}{2} \left[(b_1 \operatorname{sech}(\xi) \tanh(\xi))^2 + \left(\sqrt{2\nu^2 - b_1^2} \operatorname{sech}(\xi) \tanh(\xi)\right)^2 \right] + \frac{g^2(1 - w^2)}{2} \times \left[-\nu + b_1 \operatorname{sech}(\xi) \right]^2 + \frac{g^2(1 - w^2)}{2\nu} \times \left(-\nu + b_1 \operatorname{sech}(\xi) \right) \left[\left(-\nu + b_1 \operatorname{sech}(\xi) \right)^2 + \left(\sqrt{2\nu^2 - b_1^2} \operatorname{sech}(\xi)\right)^2 \right] + \frac{g^2(1 - w^2)}{8\nu^2} \left[\left(-\nu + b_1 \operatorname{sech}(\xi) \right)^2 + \left(\sqrt{2\nu^2 - b_1^2} \operatorname{sech}(\xi)\right)^2 \right]^2.$$
(24)

The Hamiltonian density is plotted in the Fig.(2) which is localized, and since the soliton waves are modulated in the Fig.(1), two peaks close to each other are observable in the Fig.(2) that shows the existence of two fields. Also, to make this clear, Fig.(3) is plotted for a specific g.

3.2 The Charge - Current density of the model

Using Eq.(9), we can find charge density of the model as

$$\rho(\xi, w, g) = q \left[-\nu \frac{\partial \psi_2(\xi)}{\partial \xi} - \frac{\partial \psi_2(\xi)}{\partial \xi} \psi_1(\xi) + \frac{\partial \psi_1(\xi)}{\partial \xi} \psi_2(\xi) \right] \times \left(\frac{\partial \xi}{\partial t} \right), \tag{25}$$



Figure 2: The energy density is plotted for $\nu = 1$, $b_1 = 1.224$ and w = 0.5. The figure shows two peaks that display the massive and the massless soliton fields.



Figure 3: The energy density is plotted for $\nu = 1$, $b_1 = 1.224$, w = 0.5 and g = 1.

by substituting the fields (22) in the above relation, we have

$$\rho(\xi) = -q\nu \left(\sqrt{2\nu^2 - b_1^2} \operatorname{sech}(\xi) tanh(\xi)\right) - q \left(\sqrt{2\nu^2 - b_1^2} \operatorname{sech}(\xi) tanh(\xi)\right) \left(-\nu + b_1 \operatorname{sech}(\xi)\right) \\
+ q \left(b_1 \operatorname{sech}(\xi) tanh(\xi)\right) \left(\sqrt{2\nu^2 - b_1^2} \operatorname{sech}(\xi)\right) = 0.$$
(26)

The above relationship shows that the charge density becomes zero, so the total charge and the electric dipole moment of the system are zero. These results are completely consistent with the neutrality of the produced bosons.

4 Conclusions

In this paper, we have studied the Goldstone boson model which is the simplest model to illustrate SSB under continuous transformation U(1), leading to the appearance of massless and massive spin-zero bosons. The massless bosons have never been observed, but could have many laboratory, astrophysical and cosmological implications. After SSB in the Goldstone boson model, two coupled nonlinear equations appear which have been treated by perturbation theory before, but faced with some problems which were discussed in the text. In this work, we obtained exact solutions for the coupled nonlinear equations by using tanh /sech method which is an effective technique to solve NPDEs. To apply the method we used a new transformation, namely we changed the coordinates by using a transformation which led us to solitonic solutions for which we chose the localized states. Referring to figure 1, we observed that the solitonic waves are modulated, with the massless solution riding on the massive one. Correspondingly, in the figures representing the energy density, two peaks could be seen that express the existence of two boson fields. It can also be seen that the charge-current density of the model is zero, which again confirms the neutrality of these boson fields. Since massive bosons can be observed in the laboratory, these solitonic waves may be useful in detecting Goldstone bosons and better understanding dark matter. Lastly, we note that the approach used in this paper can be extended to examine the model in a curved spacetime scenario.

5 Acknowledgment

The authors would like to thank the anonymous referee for her/his comments and suggestions to improve this paper. Also, the authors are grateful for the valuable comments and suggestions of Prof. Sepangi and Prof. Gousheh to improve the quality of the paper.

References

- T. Brauner, Spontaneous Symmetry Breaking and Nambu-Goldstone Bosons in Quantum Many-Body Systems, Symmetry 2: 609-657, (2010).
- [2] T.W.B. Kibble, History of electroweak symmetry breaking, Journal of Physics: Conference Series, 626, 012001, (2015).
- [3] F. Mandl, G. Shaw, Quantum Field Theory, 2nd Edition, John Wiley and Sons, Ltd, ISBN: 978-0-471-49683-0, (2010), 496 Pages.
- [4] G. Ballesteros, J. Redondo, A. Ringwald and C. Tamarit, Unifying inflation with the axion, dark matter, baryogenesis and the seesaw mechanism, Phys. Rev. Lett. 118, 7, 071802, (2017).
- [5] L. Di Luzio, M. Giannotti, E. Nardi and L. Visinelli, The landscape of QCD axion models, Phys. Rept. 870, (2020).
- [6] D. Adak, K. Dutta, Viable dark energy models using pseudo-Nambu-Goldstone bosons, Phys. Rev. D, 90, 043502, (2014).
- [7] J. E. Kim, Modeling small dark energy scale with quintessential pseudoscalar boson, Journal of the Korean Physical Society volume 64, 795-805, (2014).
- [8] H. Georgi, Effective Field Theory, Annu. Rev. Nucl. Part. Sci. 43, 209-252, (1993).
- [9] D. B. Kaplan, Effective Field Theories Lectures given at 7th Summer School in Nuclear Physics: Symmetries, Seattle, WA, USA, June 19-30, (1995).
- [10] A. V. Manohar, Effective Field Theories, in: H. Latal, W. Schweiger (Eds.), Perturbative and Nonperturbative Aspects of Quantum Field Theory: Proceedings of the 35. Internationale Universitatswochen fur Kern- und Teilchenphysik, Schladming, Austria, March 29, (1996), Vol. 479 of Lecture Notes in Physics, Springer, Berlin,Germany, (1997), pp. 311362.
- [11] A. Pich, Effective Field Theory, in: R. Gupta, A. Morel, E. de Rafael, F. David (Eds.), Probing the Standard Model of Particle Interactions, North Holland, Amsterdam, The Netherlands, (1999), pp. 9491049.

- [12] C. P. Burgess, Goldstone and pseudo-Goldstone bosons in nuclear, particle and condensed-matter physics, Phys. Rep. 330 (2000) 193261.
- [13] R. Alonso, M.B. Gavela, L. Merlo, S. Rigolin, J. Yepes, The effective chiral Lagrangian for a light dynamical Higgs particle, Physics Letters B, 722, 45, Pages 330-335, (2013).
- [14] K. Kanshin, The Goldstone Boson Higgs and the effective Lagrangian(s), Thesis: PhD U. Padua (main), Padua U. (2017).
- [15] A.M. Wazwaz, Partial Differential Equations and Solitary Waves Theory, Nonlinear Physical Science (2009) 457-449.
- [16] W. Malfliet, Solitary wave solutions of nonlinear wave equations, Am. J. Phys. 60, 650-654, (1992).
- [17] W. Malfliet, The tanh method: I. Exact solutions of nonlinear evolution and wave equations, Physica Scripta 54, 563-568, (1996).
- [18] W. Malfliet, The tanh method: II. Perturbation technique for conservative systems, Physica Scripta 54, 569-575, (1996).
- [19] A.H. Khater, W. Malfliet, D.K. Callebaut, E.S. Kamel, The tanh method, a simple transformation and exact analytical solutions for nonlinear reaction diffusion equations, Chaos Soliton. Fract. 14, 513-522, (2002).
- [20] E.J. Parkes, B.R. Duffy, An automated tanh-function method for finding solitary wave solutions to non-linear evolution equations, Comput. Phys. Commun. 98, 288-300, (1996).
- [21] E. Fan, Y.C. Hon, Generalized tanh method extended to special types of nonlinear equations, Z. Naturforsch 57a, 692-700, (2002).
- [22] K. Glasner, Nonlinear preconditioning for diffuse interfaces, J. Comput. Phys. 174, 695-711, (2001).
- [23] T. Kawahara, M. Tanaka, Interactions of traveling fronts: an exact solution of a nonlinear diffusion equation, Phys. Lett. 97A, (8), 311-314, (1983).
- [24] R. Conte, M. Mussette, Link between solitary waves and projective Riccati equations, J. Phys.: Math. Gen. 25, 5609-5623, (1992).
- [25] L. Huibin, W. Kelin, Exact solutions for two nonlinear equations: I, J. Phys. A: Math. Gen. 23, 3923-3928, (1990).
- [26] M. Wang, Exact solutions for a compound KdVBurgers equation, Phys. Lett. A, 213, 279-287, (1998).
- [27] W. Ma, Travelling wave solutions to a seventh order generalized KdV equation, Phys. Lett. A, 180, 221224, (1993).
- [28] A.M. Wazwaz, Partial Differential Equations: Methods and Applications, Balkema, The Netherlands, (2002).
- [29] A. J. M. Jawad, M. D. Petkovic, A. Biswas, Soliton solutions to a few coupled nonlinear wave equations by tanh method, IJST, 37A2: 109-115, (2013).
- [30] S. Sirendaoreji, Exact travelling wave solutions for four forms of nonlinear Klein-Gordon equations, Phys. Lett. A, 363, 440447, (2007).

- [31] A. M. Wazwaz, New traveling wave solutions to the Boussinesq and the Klein-Gordon equations, Commun. Nonlinear Sci Numer. Simu. 13, 889901, (2008).
- [32] A. M. Wazwaz, Compacton, solitons and periodic solutions for some forms of nonlinear Klein-Gordon equations, Chaos Solitons Fractals, 28, 1005-1013, (2006).
- [33] A. M. Wazwaz, The extended tanh method for new soliton solutions for many forms of the fifth-order KdV equations, Appl. Math. Comput. 184, 2, 1002-1014, (2007).
- [34] A. M.Wazwaz, The tanh method for generalized forms of nonlinear heat conduction and Burgers-Fisher equations, Appl. Math. Comput. 169, 321388, (2005).
- [35] A. M. Wazwaz, The extended tanh method for new compact and noncompact solutions for the KP-BBM and the ZK-BBM equations. Chaos, Solitons Fract. 38, 5, 15051516, (2008).
- [36] M. A. Abdou, The extended tanh method and its applications for solving nonlinear physical models, Appl. Math. Comput. 190, 988996, (2007).
- [37] A. A. Soliman, Extended Improved tanh-Function Method for Solving the Nonlinear Physical Problems, Acta Appl Math 104, 367383, (2008).
- [38] S. A. El-Wakil, M. A. Abdou, New exact travelling wave solutions using modified extended tanh-function method, Chaos Solitons Fractals, 31, 4, 840852, (2007).
- [39] D. F. J. Kimball, L. D. Duffy and, D. J. E. Marsh, Ultralight Bosonic Dark Matter Theory, (2023). In: D. F. J. Kimball, K. Van Bibber, (eds) The Search for Ultralight Bosonic Dark Matter. Springer, Cham. https://doi.org/10.1007/978-3-030-95852-7_2.
- [40] A. Jahangiri, S. Miraboutalebi, F. Ahmadi, A. A. Masoudi, The solitary solutions of nonlinear Klein-Gordon field with minimal length, phys. Lett. B, 818, 136351, (2021).
- [41] R. Rajaraman, Solitons and instantons, (1989).