

## LIMITS OF NODAL SURFACES AND APPLICATIONS

CIRO CILIBERTO AND CONCETTINA GALATI

ABSTRACT. Let  $\mathcal{X} \rightarrow \mathbb{D}$  be a flat family of projective complex 3-folds over a disc  $\mathbb{D}$  with smooth total space  $\mathcal{X}$  and smooth general fibre  $\mathcal{X}_t$ , and whose special fibre  $\mathcal{X}_0$  has double normal crossing singularities, in particular,  $\mathcal{X}_0 = A \cup B$ , with  $A, B$  smooth threefolds intersecting transversally along a smooth surface  $R = A \cap B$ . In this paper we first study the limit singularities of a  $\delta$ -nodal surface in the general fibre  $S_t \subset \mathcal{X}_t$ , when  $S_t$  tends to the central fibre in such a way its  $\delta$  nodes tend to distinct points in  $R$ . The result is that the limit surface  $S_0$  is in general the union  $S_0 = S_A \cup S_B$ , with  $S_A \subset A, S_B \subset B$  smooth surfaces, intersecting on  $R$  along a  $\delta$ -nodal curve  $C = S_A \cap R = S_B \cap B$ . Then we prove that, under suitable conditions, a surface  $S_0 = S_A \cup S_B$  as above indeed deforms to a  $\delta$ -nodal surface in the general fibre of  $\mathcal{X} \rightarrow \mathbb{D}$ . As applications we prove that there are regular irreducible components of the Severi variety of degree  $d$  surfaces with  $\delta$  nodes in  $\mathbb{P}^3$ , for every  $\delta \leq \binom{d-1}{2}$  and of the Severi variety of complete intersection  $\delta$ -nodal surfaces of type  $(d, h)$ , with  $d \geq h - 1$  in  $\mathbb{P}^4$ , for every  $\delta \leq \binom{d+3}{3} - \binom{d-h+1}{3} - 1$ .

## CONTENTS

1. Introduction	1
2. Limit singularity of a node of a surface in a threefold	3
2.1. The problem	3
2.2. Set up	3
2.3. Description of $\mathcal{S}_{ \mathcal{Y}_0}$ and of $\mathcal{S}_{ \mathcal{X}_0}$	5
2.4. Local deformation of a singularity of type $T_1$ to a node	7
3. Deformations of surfaces with $T_1$ singularities and nodes	7
3.1. Deformations and a smoothness criterion	8
3.2. Deformations of $T_1$ singularities	9
4. Applications	23
4.1. Severi varieties	23
4.2. The case of $\mathbb{P}^3$	24
4.3. Complete intersections in $\mathbb{P}^4$	25
References	26

## 1. INTRODUCTION

The main object of study in this article is Severi varieties of nodal surfaces on smooth, projective, complex threefolds. Severi varieties of nodal hypersurfaces on a smooth variety are a well know object of study in algebraic geometry, that goes back to well more than a century ago. Its importance is underlined by the relationships with other themes in the area. For example, the recent papers [7, 16] explore the relation of Severi varieties with the Hodge conjecture.

---

*Date:* 18/06/2024.

*2020 Mathematics Subject Classification.* 14B07, 14J17, 14C20.

*Key words and phrases.* Severi varieties of surfaces, deformations, nodes.

Our approach to the subject is via degenerations. Degenerations of smooth complex varieties to complex varieties with simple normal crossings is also a classical object of study. In particular it has been widely used by several authors for studying Severi varieties of nodal curves on surfaces. The method is powerful and enables one to obtain sharp results on the non-emptiness of some Severi varieties of curves (see, for instance, [3, 5, 6, 10], etc.).

One of the basic ideas in these papers is the well known and classical fact that the limit of a curve  $C_t$  with a node  $p_t$  on a smooth surface  $\mathcal{X}_t$ , when  $\mathcal{X}_t$  degenerates to a reducible surface  $\mathcal{X}_0 = A \cup B$ , with  $A$  and  $B$  smooth and meeting transversally along a smooth curve  $R = A \cap B$ , and  $p_t$  going to a point  $p_0 \in R$ , is a curve  $C_0 \subset \mathcal{X}_0$  with a tacnode in  $p_0$ , which appears scheme theoretically with multiplicity 2. This result is an easy consequence of the study of the versal deformation space of a tacnode, and its proof is in [2, 17]. This result has been proved also using limit linear systems techniques, see [9]. The present article intends to extend this result on the limit of a nodal curve to the case of nodal surfaces in threefolds, and we will take the point of view of [9]. In the sequel, a *node* of a surface will be an  $A_1$ -singularity.

Let  $\mathcal{X} \rightarrow \mathbb{D}$  be a flat family of projective complex 3-folds over a disc  $\mathbb{D}$  with smooth total space  $\mathcal{X}$  and smooth general fibre, and whose special fiber  $\mathcal{X}_0$  has double normal crossing singularities, in particular,  $\mathcal{X}_0 = A \cup B$ , with  $A, B$  smooth threefolds intersecting transversally along a smooth surface  $R = A \cap B$ .

First of all we will study in Section 2 the limit singularities of a  $\delta$ -nodal surface in the general fibre  $S_t \subset \mathcal{X}_t$ , when  $S_t$  tends to the central fibre in such a way that its  $\delta$  nodes tend to distinct points  $p_1, \dots, p_\delta$  in  $R$ . The result (see Theorem 2.2) is that the limit surface  $S_0$  is in general the union  $S_0 = S_A \cup S_B$ , with  $S_A \subset A$ ,  $S_B \subset B$  smooth surfaces, that cut out on  $R$  the same curve  $C$  having nodes at  $p_1, \dots, p_\delta$  and no further singularities. In this case we say that  $S_0$  presents a singularity of type  $T_1$  at every point  $p_i$ ,  $i = 1, \dots, \delta$ . The equations of a  $T_1$  singularity are given in (9). Finally in 2.4 we provide the local equation of (an example of) a local deformation of a singularity of type  $T_1$  to a node on the general fibre.

The central part of our paper is Section 3. First of all we prove in Lemma 3.3 that the only singularity of a surface  $S_t \subset \mathcal{X}_t$  to which a singularity of type  $T_1$  of a surface  $S_0 \subset \mathcal{X}_0$  may be deformed is a node. In §3.2.2 we describe the first order locally trivial deformations in  $\mathcal{X}_0$  of surfaces  $S_0 = S_A \cup S_B$  with  $T_1$  singularities on  $R$  and at most nodes elsewhere. In particular we find sufficient conditions for smoothness of the equisingular deformation locus of  $S_0$  in the relative Hilbert scheme of  $\mathcal{X}$ . If these conditions are verified, then the  $T_1$  singularities of  $S_0$  and its nodes can be smoothed independently inside  $\mathcal{X}_0$ . Next, in §3.2.3, we consider deformations of a surface  $S_0 \subset \mathcal{X}_0$ , with  $T_1$  singularities on  $R$  and at most nodes elsewhere, off the central fibre. We prove, in Theorem 3.18, that under suitable conditions, one can deform  $S_0$  off the central fibre  $\mathcal{X}_0$  to a surface  $S_t$  in the general fibre  $\mathcal{X}_t$ , with only nodes, that are the deformations of the nodes of  $S_0$  and of the  $T_1$  singularities of  $S_0$ , and that the space of this deformation is generically smooth of the expected dimension. Again, generic smoothness means that the nodes of the general surface  $S_t$  can be independently smoothed.

In Section 4 we give a couple of applications of our general result. Essentially we consider the following problem (see Problem 4.3). Let  $X$  be a smooth irreducible projective complex threefold. Let  $L$  be a line bundle on  $X$  such that the general surface in the linear system  $|L|$  is smooth and irreducible. Let  $V_\delta^{X, |L|}$  be the Severi variety of surfaces  $S$  in  $|L|$  which are reduced with only  $\delta$  nodes as singularities. The question we consider is: given  $X$  and  $L$  as above, which is the maximal value of  $\delta$  such that  $V_\delta^{X, |L|}$  has a generically smooth component of the expected codimension  $\delta$  in  $|L|$ ? We give contributions to this problem in two cases. The first one is for

$X = \mathbb{P}^3$  and  $L = \mathcal{O}_{\mathbb{P}^3}(d)$  (see Theorem 4.6), the second one is when  $X$  is a general hypersurface of degree  $h \geq 2$  in  $\mathbb{P}^4$  and  $L = \mathcal{O}_X(d)$  with  $d \geq h - 1$  (see Theorem 4.9).

To finish this introduction it is worth mentioning that the basic idea of a singularity of type  $T_1$  being a limit of a node, is already contained, although in a rather obscure form, in B. Segre's paper [18]. In this paper Segre considers, even more generally, the case of higher dimension. As a matter of fact, we believe that there should be no obstruction in extending our results in higher dimension too. However we did not dwell on this here, because we thought that the surface in threefold case already shows the complexity of the situation. We plan to come back on this in the future.

**Notation:** in what follows we use standard notation in algebraic geometry. In particular, we will denote by  $\sim$  the linear equivalence.

**Acknowledgments:** The Authors want to thank Th. Dedieu, B. Fantechi, R. Pardini, E. Sernesi, for useful discussions on the topics of this paper. The authors are member of GNSAGA of INdAM. In particular, the second author acknowledges funding from the GNSAGA of INdAM and the European Union - NextGenerationEU under the National Recovery and Resilience Plan (PNRR) - Mission 4 Education and research - Component 2 From research to business - Investment 1.1, Prin 2022 "Geometry of algebraic structures: moduli, invariants, deformations", DD N. 104, 2/2/2022, proposal code 2022BTA242 - CUP J53D23003720006.

## 2. LIMIT SINGULARITY OF A NODE OF A SURFACE IN A THREEFOLD

**2.1. The problem.** Let  $\mathcal{X} \rightarrow \mathbb{D}$  be a flat family of projective complex 3-folds over a disc  $\mathbb{D}$  with smooth total space  $\mathcal{X}$  and smooth general fiber  $\mathcal{X}_t$ , with  $t \in \mathbb{D} \setminus \{0\}$ , and whose special fiber  $\mathcal{X}_0$  has double normal crossing singularities, in particular,  $\mathcal{X}_0 = A \cup B$  has two smooth irreducible components  $A$  and  $B$ , intersecting transversally along a smooth surface  $R = A \cap B$ .

Let  $\mathcal{L}$  be a line bundle on  $\mathcal{X}$ . For each  $t \in \mathbb{D}$  we set  $\mathcal{L}_t = \mathcal{L}|_{\mathcal{X}_t}$ . We consider the following question. Roughly speaking, assume that for  $t \in \mathbb{D}$  general we have a surface  $S_t \in |\mathcal{L}_t|$  having a double point  $p_t$ . Assume that  $S_t$  tends to a surface  $S_0$  in  $\mathcal{X}_0$  with  $p_t$  tending to a point  $p_0 \in R$ . The question is: what is the singularity that  $S_0$  has at  $p_0$ . Let us make this setting more precise.

**2.2. Set up.** Let us fix  $p = p_0 \in R$ , which is a double point for the central fibre  $\mathcal{X}_0$  whereas  $\mathcal{X}$  is smooth at  $p$ . Hence there are no sections of  $\mathcal{X} \rightarrow \mathbb{D}$  passing through  $p$ . So let us consider a smooth bisection  $\gamma'$  of  $\mathcal{X} \rightarrow \mathbb{D}$  passing through  $p$ .

**Step 0.** Let us look at the following commutative diagram

$$\begin{array}{ccccc} \mathcal{Y} & \longrightarrow & \mathcal{X}' & \longrightarrow & \mathcal{X} \\ & \searrow & \downarrow & & \downarrow \\ & & \mathbb{D} & \xrightarrow{\nu_2} & \mathbb{D} \end{array}$$

where the rightmost square is cartesian and  $\nu_2 : u \in \mathbb{D} \rightarrow u^2 \in \mathbb{D}$ . Then  $\mathcal{X}'$  is singular along the counterimage of  $R$  (that by abuse of notation we still denote by  $R$ ), which is a locus of double points for  $\mathcal{X}'$ , with tangent cone a quadric cone of rank 3. The morphism  $\mathcal{Y} \rightarrow \mathcal{X}'$  is the desingularization of  $\mathcal{X}'$  obtained by blowing up  $\mathcal{X}'$  along  $R$ .

The induced morphism  $\pi : \mathcal{Y} \rightarrow \mathcal{X}$  is 2 : 1 outside the central fibre of  $\mathcal{Y}$ . In particular, for every  $t \neq 0$  there are exactly two fibres  $\mathcal{Y}_{u_1}$  and  $\mathcal{Y}_{u_2}$  of  $\mathcal{Y} \rightarrow \mathbb{D}$  isomorphic to the fibre  $\mathcal{X}_t$  of  $\mathcal{X} \rightarrow \mathbb{D}$  via  $\pi$ , where  $\{u_1, u_2\} = \nu_2^{-1}(t)$ . The family  $\mathcal{Y} \rightarrow \mathbb{D}$  has central fibre  $\mathcal{Y}_0 = A \cup \mathcal{E} \cup B$ , where, by abusing notation,  $A$  and  $B$

denote the proper transforms of  $A$  and  $B$  and  $\mathcal{E} \rightarrow R$  is a  $\mathbb{P}^1$ -bundle on  $R$ . The morphism  $\pi$  is totally ramified along  $A$  and  $B$  and it contracts  $\mathcal{E}$  to  $R$  in  $\mathcal{X}$ . In particular,  $A \cap \mathcal{E}$  and  $B \cap \mathcal{E}$  are two non-intersecting sections of  $\mathcal{E}$  both isomorphic to  $R$ . Denote by  $F$  the fibre of  $\mathcal{E} \rightarrow R$  over the point  $p \in R \subset \mathcal{X}_0$ . One has  $F \cong \mathbb{P}^1$ . Now the counterimage of  $\gamma'$  on  $\mathcal{Y}$  is the union of two sections of  $\mathcal{Y} \rightarrow \mathbb{D}$ , each intersecting  $\mathcal{Y}_0$  at a smooth point on  $F$ . We let  $\gamma$  be one of these two sections and  $q$  be the intersection point of  $\gamma$  and  $F$ .

Assume there exists an effective divisor  $\mathcal{S} \subset \mathcal{Y}$ , with  $\mathcal{S} \sim \pi^*(\mathcal{L})$ , having double points along  $\gamma$ . Let  $S$  be the image of  $\mathcal{S}$  in  $\mathcal{X}$  via the morphism  $\pi$ . Note that  $S$  has points of multiplicity 2 along the bisection  $\gamma'$ . For every  $t \neq 0$ , if  $\mathcal{Y}_{u_1}$  and  $\mathcal{Y}_{u_2}$ , with  $u_1^2 = u_2^2 = t$ , are the two fibres of  $\mathcal{Y} \rightarrow \mathbb{D}$  isomorphic to  $\mathcal{X}_t$  via  $\pi$ , we have

$$S_t = S \cap \mathcal{X}_t = S_{u_1} \cup S_{u_2},$$

where  $S_{u_i} = \pi(\mathcal{S}_{u_i})$  and  $\mathcal{S}_{u_i} = \mathcal{S} \cap \mathcal{Y}_{u_i}$ , for  $i = 1, 2$ . If  $t = 0$ , we have that

$$S \cap \mathcal{X}_0 = 2S_0 = 2(S_A \cup S_B),$$

where  $S_A = \pi(\mathcal{S} \cap A) \subset A$  and  $S_B = \pi(\mathcal{S} \cap B) \subset B$ .

We want to understand  $S|_{\mathcal{X}_0}$ . To do this, we will first understand  $\mathcal{S}|_{\mathcal{Y}_0}$ .

**Step 1.** Let  $\pi_1 : \mathcal{Y}^1 \rightarrow \mathcal{Y}$  be the blowing-up of  $\mathcal{Y}$  along  $\gamma$  with exceptional divisor  $\Gamma$ . We have a new family  $\mathcal{Y}^1 \rightarrow \mathbb{D}$  with general fibre the blow up of  $\mathcal{Y}_u \cong \mathcal{X}_{\nu_2(u)}$  at its intersection point with  $\gamma$  (that is also the point of multiplicity 2 of the surface  $\mathcal{S}_u$ ), and central fibre  $\mathcal{Y}_0^1 = A \cup \mathcal{E}' \cup B$ , where  $\mathcal{E}'$  is the blow-up of  $\mathcal{E}$  at  $q$ . Still denoting by  $F$  the proper transform of  $F$  in  $\mathcal{Y}^1$ , we have that the proper transform  $\mathcal{S}^1$  of  $\mathcal{S}$  in  $\mathcal{Y}^1$  satisfies

$$(1) \quad \mathcal{S}^1 \sim \pi_1^*(\mathcal{S}) - 2\Gamma.$$

We deduce that  $\mathcal{S}^1 \cdot F = -2$  and hence  $F \subset \mathcal{S}^1$ .

**Step 2.** Let now  $\pi_2 : \mathcal{Y}^2 \rightarrow \mathcal{Y}^1$  be the blow-up of  $\mathcal{Y}^1$  along  $F$  with new exceptional divisor  $\Theta$ . We have the new family  $\mathcal{Y}^2 \rightarrow \mathbb{D}$ , whose general fibre is the same as the general fibre of  $\mathcal{Y}^1 \rightarrow \mathbb{D}$ , and new central fibre  $\mathcal{Y}_0^2 = A' \cup \mathcal{E}'' \cup \Theta \cup B'$ , where  $A'$ ,  $\mathcal{E}''$  and  $B'$  are the blow-ups of  $A$ ,  $\mathcal{E}'$  and  $B$  at  $F \cap A$ ,  $F \subset \mathcal{E}'$  and  $B \cap F$  respectively. Notice that  $\Theta \rightarrow F$  is a  $\mathbb{P}^2$ -bundle on  $F$ , intersecting  $A'$  (resp.  $B'$ ) along a surface isomorphic to  $\mathbb{P}^2$ , which is a fibre of  $\Theta \rightarrow F$ , and at the same time is the exceptional divisor of the blow-up  $A' \rightarrow A$  at  $F \cap A$  (resp. of the blow-up  $B' \rightarrow B$  at  $F \cap B$ ). Moreover the surface  $E := \Theta \cap \mathcal{E}''$  has a  $\mathbb{P}^1$ -bundle structure  $E \rightarrow F$  and it is the exceptional divisor of  $\mathcal{E}''$ , arising from the blowing-up of  $F$  in  $\mathcal{E}'$ .

We claim that  $E \simeq \mathbb{F}_0$ . Indeed, since  $F \simeq \mathbb{P}^1$  and  $F$  is a fibre of  $\mathcal{E} \rightarrow R$ , we have that  $\mathcal{N}_{F|\mathcal{E}} \simeq \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}$ . This implies that  $\mathcal{N}_{F|\mathcal{E}'} = \mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$ , and hence  $E = \mathbb{P}(\mathcal{N}_{F|\mathcal{E}'}) = \mathbb{F}_0$ .

If  $\mathcal{S}^2$  is the proper transform of  $\mathcal{S}^1$  in  $\mathcal{Y}^2$ , by (1), we deduce that

$$(2) \quad \begin{aligned} \mathcal{S}^2|_{\Theta} &\sim \pi_2^*(\mathcal{S}^1)|_{\Theta} - m_F \Theta|_{\Theta} \\ &\sim -2f_{\Theta} + m_F(A' + B' + \mathcal{E}'')|_{\Theta} \\ &\sim -2f_{\Theta} + m_F(2f_{\Theta} + \mathcal{E}''|_{\Theta}) \\ &\sim (2m_F - 2)f_{\Theta} + m_F(\mathcal{E}''|_{\Theta}) \\ &\sim (2m_F - 2)f_{\Theta} + m_F E, \end{aligned}$$

where  $f_{\Theta}$  denotes the linear equivalence class of a fibre of  $\Theta \rightarrow F$  and  $m_F$  is the multiplicity of  $\mathcal{S}^1$  along  $F$ . Notice that  $\mathcal{S}^2|_{\Theta}$  must be an effective divisor because it is the restriction to  $\Theta$  of an effective divisor that does not contain  $\Theta$ . This implies the minimum value of  $m_F$  making  $\mathcal{S}^2|_{\Theta}$  effective is  $m_F = 1$ .

**2.3. Description of  $\mathcal{S}|_{\mathcal{Y}_0}$  and of  $\mathcal{S}|_{\mathcal{X}_0}$ .** We assume now  $m_F = 1$ . To better understand  $\mathcal{S}^2|_{\Theta} \sim E$ , we restrict  $\mathcal{S}^2|_{\Theta}$  to  $\mathcal{E}''$ . Let  $\sigma$  and  $f$ , with  $\sigma^2 = f^2 = 0$ , be the generators of the Picard group of  $E = \mathcal{E}'' \cap \Theta \cong \mathbb{F}_0$ . By restricting (2) to  $E$  one gets

$$(3) \quad \mathcal{S}^2|_E = \mathcal{S}^2|_{\Theta \cap \mathcal{E}''} \sim \mathcal{E}''|_E.$$

To compute  $\mathcal{E}''|_E$ , we use the obvious relation  $(A' + B' + \Theta + \mathcal{E}'')|_E = 0$ , which implies the following identity on  $E$

$$(4) \quad 2f + \Theta|_E + \mathcal{E}''|_E = 0.$$

Since  $E = \mathcal{E}'' \cap \Theta$ , then  $\Theta|_E$  is the class of  $\Theta^2 \cdot \mathcal{E}''$  which is clearly the class of the normal bundle  $\mathcal{N}_{E|\mathcal{E}''}$  of  $E$  in  $\mathcal{E}''$ . Similarly  $\mathcal{E}''|_E$  is the class of  $\Theta \cdot \mathcal{E}''^2 = c_1(\mathcal{N}_{E|\Theta})$ . Since  $E = \mathbb{P}(\mathcal{N}_{F|\mathcal{E}'})$ , denoting by  $\pi_E : E \rightarrow F$  the natural projection morphism, whose fiber is  $f$ , and by

$$e := \Theta|_E = \Theta^2 \cdot \mathcal{E}'' = c_1(\mathcal{N}_{E|\mathcal{E}''}),$$

we have that  $\mathcal{N}_{E|\mathcal{E}''} \subset \pi_E^*(\mathcal{N}_{F|\mathcal{E}'})$  is the tautological fibre bundle of  $E = \mathbb{P}(\mathcal{N}_{F|\mathcal{E}'})$ . So we get that

$$(5) \quad f \cdot e = -1$$

and

$$e^2 - c_1(\pi_E^*(\mathcal{N}_{F|\mathcal{E}'})) \cdot e + c_2(\pi_E^*(\mathcal{N}_{F|\mathcal{E}'})) = 0,$$

(see [12, p. 606]). Now

$$c_2(\pi_E^*(\mathcal{N}_{F|\mathcal{E}'})) = \pi_E^*(c_2(\mathcal{N}_{F|\mathcal{E}'})) = 0,$$

since  $\mathcal{N}_{F|\mathcal{E}'}$  is a vector bundle on  $F$  and  $\dim(F) = 1$ . So

$$e^2 - c_1(\pi_E^*(\mathcal{N}_{F|\mathcal{E}'})) \cdot e = e^2 - \pi_E^*(c_1(\mathcal{N}_{F|\mathcal{E}'})) \cdot e = e^2 - c_1(\mathcal{N}_{F|\mathcal{E}'})f \cdot e = e^2 + c_1(\mathcal{N}_{F|\mathcal{E}'}) = 0.$$

Thus

$$(6) \quad e^2 = -c_1(\mathcal{N}_{F|\mathcal{E}'}) = 2.$$

Set  $e = a\sigma + bf$ . By (5) and (6) one gets  $a = -1$  and

$$-2b = (-\sigma + bf)^2 = e^2 = 2, \quad \text{hence } b = -1.$$

Thus, we have

$$\Theta|_E = c_1(\mathcal{N}_{E|\mathcal{E}''}) = -\sigma - f,$$

hence by (4), we get

$$(7) \quad \mathcal{E}''|_E = c_1(\mathcal{N}_{E|\Theta}) = \sigma - f.$$

**Remark 2.1.** From (7) it follows that the divisor  $E$  (which does not move on  $\mathcal{E}''$  being there an exceptional divisor) does not move in  $\Theta$  either, since  $\mathcal{N}_{E|\Theta}$  is non-effective. Hence, by (2) and  $m_F = 1$ , we have  $\mathcal{S}^2|_E = E$ .

We are now also able to describe the divisor  $\mathcal{S}^2|_{A' \cap \mathcal{E}''} \cong \mathcal{S}^2|_{B' \cap \mathcal{E}''}$ . Indeed

$$\mathcal{S}^2|_{A' \cap \mathcal{E}''} \sim (\pi_2^* \pi_1^*(\mathcal{S}) - 2\pi_2^*(\Gamma) - \Theta)|_{A' \cap \mathcal{E}''} \sim \pi_2^* \pi_1^*(\mathcal{S})|_{A' \cap \mathcal{E}''} - \Theta \cap A' \cap \mathcal{E}'',$$

and, since  $\mathcal{S}^2|_{\Theta} = E$  by Remark 2.1, it contains the  $(-1)$ -curve  $\Theta \cap A' \cap \mathcal{E}'' = E \cap A'$  in its base locus with multiplicity 1. Thus  $\mathcal{S}^2|_{A' \cap \mathcal{E}''} = \mathcal{D}_A \cup (\Theta \cap A' \cap \mathcal{E}'')$ , where

$$\mathcal{D}_A \sim \pi_2^*(\pi_1^*(\mathcal{S}))|_{A' \cap \mathcal{E}''} - 2\Theta \cap A' \cap \mathcal{E}'',$$

and similarly for  $\mathcal{S}^2|_{B' \cap \mathcal{E}''}$ .

This analysis implies that:

**Theorem 2.2.** *Let  $\mathcal{S} \subset \mathcal{Y}$  be an effective Cartier divisor as in **Step 0**. Then the surface  $\mathcal{S}|_{\mathcal{Y}_0}$  is the union of three surfaces  $\mathcal{S}_A = \mathcal{S} \cap A$ ,  $\mathcal{S}_B = \mathcal{S} \cap B$  and  $\mathcal{S}_\mathcal{E} = \mathcal{S} \cap \mathcal{E}$ , where  $\mathcal{S}_A$  (resp.  $\mathcal{S}_B$ ) intersects  $A \cap \mathcal{E}$  (resp.  $B \cap \mathcal{E}$ ) along a curve which has a double point at the point  $F \cap A$  (resp.  $F \cap B$ ), these two curves are isomorphic, and  $\mathcal{S}_\mathcal{E}$  is a  $\mathbb{P}^1$ -bundle over any one of them.*

Accordingly,  $\mathcal{S}|_{\mathcal{X}_0} = 2S_0$ , with  $S_0 \in |\mathcal{L}_0|$  and  $S_0$  is the union of two surfaces  $S_A$ ,  $S_B$ , respectively isomorphic to  $\mathcal{S}_A, \mathcal{S}_B$ , intersecting along a curve in  $R$ , that has a double point at  $p$ .




FIGURE 1.

2.3.1. *Local equations of  $S_0$ .* We may assume that  $\mathcal{X}$  locally around  $p \in \mathcal{X}_0$  is embedded in  $\mathbb{A}^5$  with coordinates  $(x, y, z, u, t)$  with  $p$  corresponding to the origin. We may suppose that  $\mathcal{X}$  is defined by the equation  $xy = t$  and the map  $\mathcal{X} \rightarrow \mathbb{D}$  is given by  $(x, y, z, u, t) \mapsto t$ . So we will assume that  $A$  is defined by the equations  $x = t = 0$  and  $B$  by the equations  $y = t = 0$ , so that  $R$  is defined by  $x = y = t = 0$ .

The above analysis proves that the surfaces  $\mathcal{S}|_A$  and  $\mathcal{S}|_B$  belong to the restriction linear systems of  $\mathcal{L}$  to  $A$  and  $B$ , respectively, and moreover are tangent to  $R$  at the point  $p$ . Thus  $S_0 = S_A \cup S_B$  belongs to the linear system  $\mathcal{L}_0(2, p) \subset |\mathcal{L}_0|$  of surfaces with local equations at  $p$  given by

$$(8) \quad \begin{cases} (a_1x + b_1y) + f_2(x, y, z, u) = 0 \\ xy = 0, \end{cases}$$

with  $f_2(x, y, z, u)$  an analytic function with terms of degree at least 2.

**Definition 2.3.** Let  $S_0 = S_A \cup S_B$  be a surface that is the union of two irreducible components  $S_A, S_B$  intersecting along a curve  $C$ . Let  $p \in C$ . We will say that  $S$  has at  $p$  a *singularity of type  $T_1$*  if  $S_A$  and  $S_B$  are smooth at  $p$  and  $C$  has at  $p$  a node.

**Remark 2.4.** If in (8),  $a_1b_1 \neq 0$ , then  $S_0$  has a  $T_1$  singularity at the origin  $p$ , and, up to a linear change of coordinates, the local equations are given by

$$(9) \quad \begin{cases} x + y + f_2(x, y, z, u) = 0, \text{ with } f_2(0, 0, z, u) = 0 \text{ having a node at } \underline{0}, \\ xy = 0. \end{cases}$$

In the sequel we will also refer to  $\mathcal{L}_0(2, p)$  as the sublinear system of  $|\mathcal{L}_0|$  of surfaces with at least a  $T_1$  singularity at  $p$ .

**Remark 2.5.** We have that  $\mathcal{L}_0(2, p) \subset |\mathcal{L}_0|$  has dimension

$$\dim(\mathcal{L}_0(2, p)) \geq \dim |\mathcal{L}_0| - 3.$$

**2.4. Local deformation of a singularity of type  $T_1$  to a node.** In §2.3.1 we saw that a singularity of type  $T_1$  appears as a *generic* limit of a double point of a surface. In this section we will show that locally the converse happens, i.e., that locally a singularity of type  $T_1$  can be deformed to a node.

In local coordinates  $(x, y, z, u, t)$ , we consider as before the family of 3-folds  $\mathcal{X}_t : xy = t$ . We further consider the one parameter family of 3-folds in  $\mathbb{A}^4$  of local equation at  $\underline{0}$  given by

$$\mathcal{S}_\alpha : x - y - \alpha(t) - z^2 - u^2 = 0,$$

where  $\alpha(t)$  is a suitable function of  $t \in \mathbb{A}^1$  to be determined, such that  $\alpha(0) = 0$ . We will set  $S_t = \mathcal{S}_{\alpha(t)} \cap \mathcal{X}_t$  for any  $t \in \mathbb{A}^1$ . The surface  $S_0$  has a  $T_1$  singularity at  $\underline{0}$  and  $\mathcal{S}_\alpha$  is smooth. Our requirement on the function  $\alpha(t)$  is that for any  $t \neq 0$  there exists a singular point  $q(t) = (x(t), y(t), z(t), u(t)) \in S_t$ , i.e., such that

$$T_{q(t)}(\mathcal{S}_\alpha) = T_{q(t)}(\mathcal{X}_t).$$

This is equivalent to ask that there exists  $q(t) = (x(t), y(t), z(t), u(t)) \in \mathbb{A}^4$  satisfying

$$x(t) - y(t) - \alpha(t) - z(t)^2 - u(t)^2 = x(t)y(t) - t = 0$$

and

$$(x-x(t))-(y-y(t))-2z(t)(z-z(t))-2u(t)(u-u(t)) = c(t)\left(y(t)(x-x(t))+x(t)(y-y(t))\right),$$

for a non-zero  $c(t)$ . This implies

$$z(t) = u(t) = 0, x(t) = -y(t), \alpha(t) = 2x(t) \text{ and } t = -x(t)^2 = -\frac{\alpha(t)^2}{4}.$$

Thus, for every  $t \neq 0$  there exist exactly two divisors  $\mathcal{S}_{\alpha_i}$ , with  $i = 1, 2$  and  $\alpha_i(t)^2 = -4t$  so that

$$\mathcal{S}_{\alpha_i(t)} = \mathcal{S}_{\alpha_i} \cap \mathcal{Y}_t : \begin{cases} x = y + \alpha_i(t) + z^2 + u^2 \\ y(y + \alpha_i(t) + z^2 + u^2) = t \end{cases}$$

is a one-nodal surface, with tangent cone at  $q_i(t) = (\frac{\alpha_i(t)}{2}, -\frac{\alpha_i(t)}{2}, 0, 0)$  given by

$$TC_{q_i(t)}(\mathcal{S}_{\alpha_i(t)}) : x - y - \alpha_i(t) = 2\left(y + \frac{\alpha_i(t)}{2}\right)^2 - \alpha_i(t)z^2 - \alpha_i(t)u^2 = 0.$$

Notice that, for every  $i = 1, 2$ , we have that  $\alpha_i(t)$  is a well defined continuous function on  $\mathbb{D}_\epsilon^\circ = \mathbb{D}(\underline{0}, \epsilon) \setminus \{a + i0 \mid 0 < a < \epsilon\}$  (the disk cut along a radius), vanishing at 0 and holomorphic on  $\mathbb{D}_\epsilon^\circ \setminus \underline{0}$ . Each family  $\mathcal{S}_{\alpha_i} \rightarrow \mathbb{D}_\epsilon^\circ$ , for  $i = 1, 2$ , is not algebraic, while the complete intersection family of surfaces

$$(10) \quad \mathcal{D} : \begin{cases} (x - y - u^2 - z^2)^2 = -4t \\ xy = t. \end{cases}$$

is algebraic. As usual we set  $D_t = \mathcal{D} \cap \mathcal{X}_t$ . One has  $D_t = \mathcal{S}_{\alpha_1(t)} \cup \mathcal{S}_{\alpha_2(t)}$  for  $t \neq 0$  and non-reduced fibre  $D_0 = 2S_0$  for  $t = 0$ .

The locus  $x^2 + t = x + y = z = u = 0$ , whose general point is singular for  $D_t$ , is a bisection of  $\mathcal{X} \rightarrow \mathbb{A}^1$  passing through  $(\underline{0}, 0)$ .

### 3. DEFORMATIONS OF SURFACES WITH $T_1$ SINGULARITIES AND NODES

Throughout this section we will consider  $\mathcal{X} \rightarrow \mathbb{D}$  a family of projective complex 3-folds over a disc  $\mathbb{D}$  as in the previous section and we let  $\mathcal{H}^{\mathcal{X}|\mathbb{D}}$  be its relative Hilbert scheme, whose fiber over  $t \in \mathbb{D}$  is the Hilbert scheme of  $\mathcal{X}_t$  and it is denoted by  $\mathcal{H}^{\mathcal{X}_t}$ . Moreover we will consider  $S_0 = S_A \cup S_B \subset \mathcal{X}_0$ , with  $S_A \subset A$  and  $S_B \subset B$  an effective reduced Cartier divisor.

### 3.1. Deformations and a smoothness criterion.

**Definition 3.1.** Let  $S_0 = S_A \cup S_B \subset \mathcal{X}_0$ , with  $S_A \subset A$  and  $S_B \subset B$  be an effective reduced Cartier divisor and let

$$\mathcal{H}_{[S_0]}^{\mathcal{X}|\mathbb{D}} \rightarrow \mathbb{D}$$

be an irreducible component of the relative Hilbert scheme of  $\mathcal{X}$  containing  $[S_0]$ . A deformation of  $S_0$  in  $\mathcal{X}$  not in  $\mathcal{X}_0$  is the total space  $S \subset \mathcal{X}$  of an irreducible local  $r$ -multisection  $\gamma$  of  $\mathcal{H}_{[S_0]}^{\mathcal{X}|\mathbb{D}}$  passing through  $[S_0]$ . Equivalently, a deformation of  $S_0$  is an effective divisor

$$\begin{array}{ccc} S & \hookrightarrow & \mathcal{X} \\ & \searrow & \downarrow \\ & & \mathbb{D} \end{array}$$

dominating  $\mathbb{D}$ , whose central fibre is  $S \cap \mathcal{X}_0 = rS_0$ , i.e., the surface  $S_0$  counted with multiplicity  $r$ , and whose general fibre is a reduced surface with  $r$  irreducible components  $S \cap \mathcal{X}_t = S_t^1 \cup \dots \cup S_t^r$ , with  $[S_t^i] \in \mathcal{H}_{[S_0]}^{\mathcal{X}|\mathbb{D}}$ , for every  $i = 1, \dots, t$ . We will also say that every irreducible component  $S_t^i$  of  $S \cap \mathcal{X}_t$  is a deformation of  $S_0$  or that  $S_0$  is a limit of  $S_t^i$ . Let  $\mathcal{Y}$  be the smooth family of threefolds obtained from  $\mathcal{X} \rightarrow \mathbb{D}$  after a base change

$$\begin{array}{ccccc} \mathcal{Y} & \longrightarrow & \mathcal{X}' & \longrightarrow & \mathcal{X} \\ & \searrow & \downarrow & & \downarrow \\ & & \mathbb{D} & \xrightarrow{\nu_r} & \mathbb{D} \end{array}$$

of order  $r$  and after minimally desingularizing the total space of the obtained family.  $\mathcal{Y}$  has central fibre  $\mathcal{Y}_0 = A \cup \mathcal{E}_1 \cup \dots \cup \mathcal{E}_{r-1} \cup B$  with normal crossing singularities of multiplicity two, where every  $\mathcal{E}_i$  is a  $\mathbb{P}^1$ -bundle over  $\mathcal{E}_{i-1} \cap \mathcal{E}_i$ , intersecting  $\mathcal{E}_{i-1}$  and  $\mathcal{E}_{i+1}$ , with  $\mathcal{E}_0 = A$  and  $\mathcal{E}_r = B$ . We denote by  $\pi : \mathcal{Y} \rightarrow \mathcal{X}$  the induced morphism. Then the pullback divisor  $\pi^*(S) = \mathcal{S}^1 \cup \dots \cup \mathcal{S}^r$  has  $r$  irreducible distinct components  $\mathcal{S}^1, \dots, \mathcal{S}^r$ , where now every  $\mathcal{S}^i$  has irreducible general fibre and has central fibre given by  $\mathcal{S}_0^i = \mathcal{S}^i \cap \mathcal{X}_0 \cong S_0$ .

**Proposition 3.2.** *Let  $S_0 = S_A \cup S_B \subset \mathcal{X}_0$ , with  $S_A \subset A$  and  $S_B \subset B$ , be a reduced effective Cartier divisor as above. Let  $p$  be a point of the intersection curve  $C = S_A \cap S_B \subset R$  where  $S_A$  and  $S_B$  intersect transversally, i.e., such that  $S_A$  and  $S_B$  are smooth at  $p$  and  $T_p(S_A) \neq T_p(S_B)$ . Then for every deformation  $S \subset \mathcal{X}$  of  $S_0$  not in  $\mathcal{X}_0$ , we have that  $p$  is limit only of smooth points of the irreducible components of the general fibre of  $S$ , i.e., in a sufficiently small analytic neighborhood of  $p$  in  $\mathcal{X}$ , all irreducible components of the general fibre of  $S$  are smooth. In particular, if  $S_A$  and  $S_B$  intersect transversally along  $C$ , then  $S_0$  is limit only of smooth surfaces.*

*Proof.* Let  $S_0 = S_A \cup S_B \subset \mathcal{X}_0$  and  $p \in R = S_A \cap S_B$  as in the statement. Then, there exists an analytic coordinate system  $(x, y, z, u, t)$  of  $\mathcal{X}$  at  $p = \underline{0}$  and such that the local equation of  $S_0$  at  $p$  is given by  $xy = t = z + f_2(x, y, z, u) = 0$ , where  $f_2(x, y, z, u) \in (x, y, z, u)^2$ .

Assume that the assertion is not true. Let  $\pi : \mathcal{Y} \rightarrow \mathcal{X}$  be the morphism defined in Definition 3.1, from which we keep the notation. Then the chain of fibres  $F_p^1 \cup \dots \cup F_p^{r-1}$  of  $\pi^{-1}(S_0)$  contracted to  $p$  by  $\pi$ , intersects the singular locus of every divisor  $\mathcal{S}^i$ . In particular there exists an analytic  $s$ -multisection  $\gamma$  of  $\mathcal{X}$  (with  $s \geq r$ ) passing through  $p$ , whose general point is a singular point of an irreducible component of  $S \cap \mathcal{X}_t$ , with  $t$  general. Every analytic  $s$ -multisection of  $\mathcal{X} \rightarrow \mathbb{D}$  at  $p$  gives rise to  $s$  distinct continuous sections  $\gamma^1, \dots, \gamma^s$  over  $\mathbb{D}^\circ = \mathbb{D} \setminus \{a + i0 \mid 0 < a < 1\}$ , which are holomorphic on  $\mathbb{D}^\circ \setminus \underline{0}$ . If  $t$  varies in  $\mathbb{D}^\circ$ , then there exists a one-parameter analytic family of irreducible surfaces  $\mathcal{Z} \subset S$ , singular along  $\gamma^1$ , whose fibre  $\mathcal{Z}_t$  over  $t \neq 0$  is



an irreducible component of  $S \cap \mathcal{X}_t$  and whose fibre over 0 is  $\mathcal{Z}_0 = S_0$ . The equation of  $\mathcal{Z}_t$  in  $\mathbb{A}^4$  with coordinates  $(x, y, z, u)$  is given by

$$\begin{cases} p(x, y, z, u; t) = 0 \\ xy = t, \end{cases}$$

where  $p(x, y, z, u; t) = 0$  is an analytic function in  $(x, y, z, u)$ , whose coefficients are continuous functions in the variable  $t \in \mathbb{D}^\circ$  which are holomorphic on  $\mathbb{D}^\circ \setminus \underline{0}$ . If

$$\gamma^1(t) = (x(t), y(t), z(t), u(t)),$$

then, by the hypothesis that the general fibre of  $\mathcal{Z}$  is singular along  $\gamma$ , we have that

$$p(x, y, z, u; t) = c(t)(y(t)(x-x(t))+x(t)(y-y(t)))+g_2(x-x(t), y-y(t), z-z(t), u-u(t)),$$

where  $g_2(x-x(t), y-y(t), z-z(t), u-u(t)) \in (x-x(t), y-y(t), z-z(t), u-u(t))^2$ .

We moreover have that  $p(x, y, z, u; t)$  specializes to  $p(x, y, z, u; 0) = z + f_2(x, y, z, u) = z - z(0) + f_2(x-x(0), y-y(0), z-z(0), u-u(0))$  as  $t$  goes to 0. This is not possible. Thus every irreducible component of the general fibre of a deformation  $S \subset \mathcal{X}$  of  $S_0$  is smooth in a neighborhood of  $p$ .  $\square$

### 3.2. Deformations of $T_1$ singularities.

3.2.1. *Deformations not in  $\mathcal{X}_0$  of surfaces with  $T_1$  singularities.* In this section we prove that the only singularity of a surface in  $\mathcal{X}_t$ , with  $t \neq 0$ , to which a  $T_1$  singularity of a surface  $S_0 \subset \mathcal{X}_0$  may be deformed is a node.

**Lemma 3.3.** *Let  $S_0 = S_A \cup S_B \subset \mathcal{X}_0$  be a reduced effective Cartier divisor, with  $S_A \subset A$  and  $S_B \subset B$  as above. Let  $p$  be a point of the intersection curve  $C = S_A \cap S_B \subset R$  where  $S_0$  has a  $T_1$  singularity. Let  $S \subset \mathcal{X}$  be a deformation of  $S_0$  not in  $\mathcal{X}_0$ . Then there exists a sufficiently small analytic neighborhood of  $p$  in  $\mathcal{X}$  such that all irreducible components of the general fibre of  $S$  in that neighborhood are smooth or are 1-nodal.*

*Proof.* By Proposition 3.2, if  $S \subset \mathcal{X}$  is any deformation of  $S_0$  not in  $\mathcal{X}_0$ , then all irreducible components of the general fibre of  $S$  have, in a sufficiently small neighborhood of  $p$ , only isolated singularities. We want to prove that if the  $T_1$  singularity of  $S_0$  at  $p$  is limit of an isolated singularity, then this is a node. We argue as in the proof of Proposition 3.2.

Let  $\mathbb{D}_\epsilon = \mathbb{D}(\underline{0}, \epsilon) \subset \mathbb{A}^1$  be the open disc with center at the origin and radius  $\epsilon$  and let  $\mathbb{D}_\epsilon^\circ = \mathbb{D}(\underline{0}, \epsilon) \setminus \{a + i0 \mid 0 < a < \epsilon\}$ . We denote by  $t = a + ib$  the coordinate on  $\mathbb{D}_\epsilon^\circ$  and by  $(x, y, z, u)$  the coordinates in  $\mathbb{A}^4$ . In  $\mathbb{A}^4 \times \mathbb{D}_\epsilon^\circ$  we consider a one parameter family of 3-folds

$$\mathcal{S}_t : p(x, y, z, u; t) = 0, \quad t \in \mathbb{D}_\epsilon^\circ,$$

where  $p(x, y, z, u; t)$  is a polynomial in  $x, y, z, u$  whose coefficients are holomorphic functions on  $\mathbb{D}_\epsilon^\circ \setminus \underline{0}$ , continuous in  $\underline{0}$ , and the one parameter family of 3-folds

$$\mathcal{X}_t : xy = t, \quad t \in \mathbb{D}_\epsilon^\circ.$$

Assume that the surface

$$(11) \quad S_0 = \mathcal{S}_0 \cap \mathcal{X}_0 : \begin{cases} p(x, y, z, u; 0) = x + y + p_2(x, y, z, u) + o(3) = 0 \\ xy = 0 \end{cases}$$

has a  $T_1$  singularity at  $\underline{0}$ , where  $p_2(x, y, z, u)$  is the homogeneous part of degree 2 of  $p(x, y, z, u; 0)$ , where  $o(3)$  is the sum of terms of degree at least 3 in  $p(x, y, z, u; 0)$ , and where, by assumption,  $p_2(0, 0, z, u)$  has non-zero discriminant.

Assume that, for  $t \neq 0$ , there exists  $q(t) = (x(t), y(t), z(t), w(t); t) \in \mathcal{S}_t = \mathcal{S}_t \cap \mathcal{X}_t$  specializing to  $\underline{0}$ , as  $t$  goes to  $\underline{0}$  and such that  $\mathcal{S}_t$  has a singular point at  $q(t)$ . Thus,  $\mathcal{S}_t$  is smooth at  $q(t)$  since  $\mathcal{S}_0$  is smooth at  $q(0) = \underline{0}$  and we have that

$$T_{q(t)}(\mathcal{S}_t) = T_{q(t)}(\mathcal{X}_t).$$

In particular there exists a function  $c(t)$ , which is non-zero if  $t \neq 0$ , such that

$$\begin{aligned} y(t)(x - x(t)) + x(t)(y - y(t)) &= c(t) \frac{\partial p}{\partial x} \Big|_{q(t)} (x - x(t)) + c(t) \frac{\partial p}{\partial y} \Big|_{q(t)} (y - y(t)) \\ &+ c(t) \frac{\partial p}{\partial z} \Big|_{q(t)} (z - z(t)) + c(t) \frac{\partial p}{\partial u} \Big|_{q(t)} (u - u(t)), \end{aligned}$$

from which we deduce that

$$(12) \quad y(t) = c(t) \frac{\partial p}{\partial x} \Big|_{q(t)}, \quad x(t) = c(t) \frac{\partial p}{\partial y} \Big|_{q(t)},$$

$$(13) \quad \frac{\partial p}{\partial z} \Big|_{q(t)} = 0 \quad \text{and} \quad \frac{\partial p}{\partial u} \Big|_{q(t)} = 0.$$

As  $t$  goes to  $\underline{0}$ ,  $c(0) = 0$ , since  $x(t) \neq 0 \neq y(t)$  if  $t \neq 0$  but  $x(0) = y(0) = 0$  and  $\frac{\partial p}{\partial x} \Big|_{q(t)} \neq 0 \neq \frac{\partial p}{\partial y} \Big|_{q(t)}$  for any  $t$  in a neighborhood of  $\underline{0}$ . We now write down the local equations

$$(14) \quad \mathcal{X}_t : y(t)(x - x(t)) + x(t)(y - y(t)) + (x - x(t))(y - y(t)) = 0$$

of  $\mathcal{X}_t$  at  $q(t)$ , and the local equation

$$\begin{aligned} (15) \quad \mathcal{S}_t &: \frac{\partial p}{\partial x} \Big|_{q(t)} (x - x(t)) + \frac{\partial p}{\partial y} \Big|_{q(t)} (y - y(t)) \\ &+ \frac{\partial p}{\partial x \partial y} \Big|_{q(t)} (x - x(t))(y - y(t)) + \frac{\partial p}{\partial x \partial z} \Big|_{q(t)} (x - x(t))(z - z(t)) \\ &+ \frac{\partial p}{\partial x \partial u} \Big|_{q(t)} (x - x(t))(u - u(t)) + \frac{\partial p}{\partial y \partial z} \Big|_{q(t)} (y - y(t))(z - z(t)) \\ &+ \frac{\partial p}{\partial y \partial u} \Big|_{q(t)} (y - y(t))(u - u(t)) + \frac{\partial p}{\partial u \partial z} \Big|_{q(t)} (u - u(t))(z - z(t)) \\ &+ \frac{1}{2} \frac{\partial p}{\partial x^2} \Big|_{q(t)} (x - x(t))^2 + \frac{1}{2} \frac{\partial p}{\partial y^2} \Big|_{q(t)} (y - y(t))^2 + \frac{1}{2} \frac{\partial p}{\partial u^2} \Big|_{q(t)} (u - u(t))^2 \\ &+ \frac{1}{2} \frac{\partial p}{\partial z^2} \Big|_{q(t)} (z - z(t))^2 + o(3) = 0 \end{aligned}$$

of  $\mathcal{S}_t$  at  $q(t)$ , where  $o(3) \in (x - x(t), y - y(t), z - z(t), u - u(t))^3$ . By (14), one may write

$$(16) \quad x - x(t) = -\frac{x(t)(y - y(t))}{y - y(t) + y(t)}.$$

Let  $d$  be the maximum degree of  $x - x(t)$  in (15). By substituting in (15), by multiplying by  $y^d = (y - y(t) + y(t))^d$ , and by using that  $T_{q(t)}(\mathcal{S}_t) = T_{q(t)}(\mathcal{X}_t)$ , i.e.,  $y(t) \frac{\partial p}{\partial y} \Big|_{q(t)} = x(t) \frac{\partial p}{\partial x} \Big|_{q(t)}$  for any  $t \neq 0$ , we find that the local equation of  $\mathcal{S}_t = \mathcal{S}_t \cap \mathcal{X}_t$

is given by

$$\begin{aligned}
S_t &: \frac{\partial p}{\partial x}|_{q(t)} \left( -x(t)(y-y(t)) \right) \left( y-y(t)+y(t) \right)^{d-1} \\
&+ \left( \frac{\partial p}{\partial x \partial y}|_{q(t)}(y-y(t)) \right) \left( -x(t)(y-y(t)) \right) \left( y-y(t)+y(t) \right)^{d-1} \\
&+ \left( \frac{\partial p}{\partial x \partial z}|_{q(t)}(z-z(t)) \right) \left( -x(t)(y-y(t)) \right) \left( y-y(t)+y(t) \right)^{d-1} \\
&+ \left( \frac{\partial p}{\partial x \partial u}|_{q(t)}(u-u(t)) \right) \left( -x(t)(y-y(t)) \right) \left( y-y(t)+y(t) \right)^{d-1} \\
&+ \frac{1}{2} \frac{\partial p}{\partial x^2}|_{q(t)} \left( -x(t)(y-y(t)) \right)^2 \left( y-y(t)+y(t) \right)^{d-2} \\
&+ \frac{\partial p}{\partial y}|_{q(t)} \left( y-y(t) \right) \left( y-y(t)+y(t) \right)^d \\
&+ \left( \frac{\partial p}{\partial y \partial z}|_{q(t)}(z-z(t)) + \frac{\partial p}{\partial y \partial u}|_{q(t)}(u-u(t)) \right) \left( y-y(t) \right) \left( y-y(t)+y(t) \right)^d \\
&+ \left( \frac{1}{2} \frac{\partial p}{\partial y^2}|_{q(t)}(y-y(t))^2 + \frac{\partial p}{\partial u \partial z}|_{q(t)}(u-u(t))(z-z(t)) \right) \left( y-y(t)+y(t) \right)^d \\
&+ \left( \frac{1}{2} \frac{\partial p}{\partial u^2}|_{q(t)}(u-u(t))^2 + \frac{1}{2} \frac{\partial p}{\partial z^2}|_{q(t)}(z-z(t))^2 \right) \left( y-y(t)+y(t) \right)^d + o(3) \\
&= y(t)^{d-2} \left( \frac{\partial p}{\partial y}|_{q(t)}y(t) - \frac{\partial p}{\partial x \partial y}|_{q(t)}x(t)y(t) + \frac{1}{2} \frac{\partial p}{\partial x^2}|_{q(t)}x(t)^2 + \frac{1}{2} \frac{\partial p}{\partial y^2}|_{q(t)}y(t)^2 \right) \left( y-y(t) \right)^2 \\
&+ y(t)^{d-1} \left( y(t) \frac{\partial p}{\partial y \partial z}|_{q(t)} - x(t) \frac{\partial p}{\partial x \partial z}|_{q(t)} \right) \left( y-y(t) \right) \left( z-z(t) \right) \\
&+ y(t)^{d-1} \left( y(t) \frac{\partial p}{\partial y \partial u}|_{q(t)} - x(t) \frac{\partial p}{\partial x \partial u}|_{q(t)} \right) \left( y-y(t) \right) \left( u-u(t) \right) \\
&+ y(t)^d \left( \frac{1}{2} \frac{\partial p}{\partial z^2}|_{q(t)}(z-z(t))^2 + \frac{1}{2} \frac{\partial p}{\partial u^2}|_{q(t)}(u-u(t))^2 \right) \\
&+ y(t)^d \left( \frac{\partial p}{\partial u \partial z}|_{q(t)}(u-u(t))(z-z(t)) \right) + o(3) = 0.
\end{aligned}$$

Up to the irrelevant factor  $y(t)^{d-2}$ , the Hessian matrix at  $q(t)$  of the above polynomial is

$$(17) \quad H_{q(t)} = \begin{pmatrix} A_{11}(t) & A_{12}(t) & A_{13}(t) \\ A_{12}(t) & \frac{y(t)^2}{2} \frac{\partial p}{\partial z^2}|_{q(t)} & \frac{y(t)^2}{2} \frac{\partial p}{\partial z \partial u}|_{q(t)} \\ A_{13}(t) & \frac{y(t)^2}{2} \frac{\partial p}{\partial z \partial u}|_{q(t)} & \frac{y(t)^2}{2} \frac{\partial p}{\partial u^2}|_{q(t)} \end{pmatrix},$$

where

$$A_{11}(t) = \frac{\partial p}{\partial y}|_{q(t)}y(t) - \frac{\partial p}{\partial x \partial y}|_{q(t)}x(t)y(t) + \frac{1}{2} \frac{\partial p}{\partial x^2}|_{q(t)}x(t)^2 + \frac{1}{2} \frac{\partial p}{\partial y^2}|_{q(t)}y(t)^2,$$

$$A_{12}(t) = \frac{y(t)}{2} \left( y(t) \frac{\partial p}{\partial y \partial z}|_{q(t)} - x(t) \frac{\partial p}{\partial x \partial z}|_{q(t)} \right),$$

$$A_{13}(t) = \frac{y(t)}{2} \left( y(t) \frac{\partial p}{\partial y \partial u}|_{q(t)} - x(t) \frac{\partial p}{\partial x \partial u}|_{q(t)} \right).$$

Now  $S_t$  has a node at  $q(t)$  if and only if  $\det(H_{q(t)}) \neq 0$ . If we substitute the equalities (12) in  $H_{q(t)}$ , we see that this matrix has the first column divisible by  $c(t)$  and the second and third columns divisible by  $c(t)^2$ . Let  $B_{q(t)}$  be the matrix obtained by  $H_{q(t)}$  by dividing the first column by  $c(t)$  and the second and third columns by  $c(t)^2$ . We have that

$$(18) \quad B_{q(t)} = \begin{pmatrix} B_{11}(t) & B_{12}(t) & B_{13}(t) \\ B_{21}(t) & \frac{1}{2} \left( \frac{\partial p}{\partial x}|_{q(t)} \right)^2 \frac{\partial p}{\partial z^2}|_{q(t)} & \frac{1}{2} \left( \frac{\partial p}{\partial x}|_{q(t)} \right)^2 \frac{\partial p}{\partial z \partial u}|_{q(t)} \\ B_{31}(t) & \frac{1}{2} \left( \frac{\partial p}{\partial x}|_{q(t)} \right)^2 \frac{\partial p}{\partial z \partial u}|_{q(t)} & \frac{1}{2} \left( \frac{\partial p}{\partial x}|_{q(t)} \right)^2 \frac{\partial p}{\partial u^2}|_{q(t)} \end{pmatrix},$$

where

$$\begin{aligned}
B_{11}(t) &= \frac{\partial p}{\partial y}|_{q(t)} \frac{\partial p}{\partial x}|_{q(t)} + c(t) \left( -\frac{\partial p}{\partial x \partial y}|_{q(t)} \frac{\partial p}{\partial x}|_{q(t)} \frac{\partial p}{\partial y}|_{q(t)} \right. \\
&\quad \left. + \frac{1}{2} \frac{\partial p}{\partial x^2}|_{q(t)} \left( \frac{\partial p}{\partial y}|_{q(t)} \right)^2 + \frac{1}{2} \frac{\partial p}{\partial y^2}|_{q(t)} \left( \frac{\partial p}{\partial x}|_{q(t)} \right)^2 \right), \\
B_{21}(t) &= \frac{1}{2} \frac{\partial p}{\partial x}|_{q(t)} c(t) \left( \frac{\partial p}{\partial x}|_{q(t)} \frac{\partial p}{\partial y \partial z}|_{q(t)} - \frac{\partial p}{\partial y}|_{q(t)} \frac{\partial p}{\partial x \partial z}|_{q(t)} \right), \\
B_{31}(t) &= \frac{1}{2} \frac{\partial p}{\partial x}|_{q(t)} c(t) \left( \frac{\partial p}{\partial x}|_{q(t)} \frac{\partial p}{\partial y \partial u}|_{q(t)} - \frac{\partial p}{\partial y}|_{q(t)} \frac{\partial p}{\partial x \partial u}|_{q(t)} \right).
\end{aligned}$$

As  $t$  goes to 0,  $c(t)$  goes to 0 and the matrix  $B_{q(t)}$  specializes to the matrix

$$(19) \quad B_{\underline{0}} = \begin{pmatrix} \frac{\partial p}{\partial y}|_{\underline{0}} \frac{\partial p}{\partial x}|_{\underline{0}} & B_{12}(0) & B_{13}(0) \\ 0 & \frac{1}{2} \left( \frac{\partial p}{\partial x}|_{\underline{0}} \right)^2 \frac{\partial p}{\partial z^2}|_{\underline{0}} & \frac{1}{2} \left( \frac{\partial p}{\partial x}|_{\underline{0}} \right)^2 \frac{\partial p}{\partial z \partial u}|_{\underline{0}} \\ 0 & \frac{1}{2} \left( \frac{\partial p}{\partial x}|_{\underline{0}} \right)^2 \frac{\partial p}{\partial z \partial u}|_{\underline{0}} & \frac{1}{2} \left( \frac{\partial p}{\partial x}|_{\underline{0}} \right)^2 \frac{\partial p}{\partial u^2}|_{\underline{0}} \end{pmatrix}.$$

Using that  $\frac{\partial p}{\partial x}|_{\underline{0}} = 1 = \frac{\partial p}{\partial y}|_{\underline{0}}$ , we see that  $\det(B_{\underline{0}})$  coincides with the discriminant of the degree 2 homogeneous polynomial  $p_2(0, 0, z, u)$ , which is non zero by the hypothesis that  $S_0$  has a  $T_1$  singularity at  $\underline{0}$ . We finally deduce that  $\det(B_{q(t)}) \neq 0 \neq \det(H_{q(t)})$  and thus the surface  $S_t$  has a node at  $q(t)$  for  $t \neq 0$ .  $\square$

**3.2.2. Equisingular deformations of surfaces with  $T_1$  singularities.** We go on considering the setting we introduced at the beginning of Section 3. Assume that  $S_0 = S_A \cup S_B \subset \mathcal{X}_0$  is a surface with  $S_A$  and  $S_B$  smooth, intersecting transversally along  $R = A \cap B$ , except for  $\delta$  distinct points  $p_1, \dots, p_\delta \in S_A \cap S_B$ , where  $S_0$  has a singularity of type  $T_1$ . We recall the standard exact sequence

$$(20) \quad 0 \longrightarrow \Theta_{S_0} \longrightarrow \Theta_{\mathcal{X}|_{S_0}} \xrightarrow{\alpha} \mathcal{N}_{S_0|\mathcal{X}} \xrightarrow{\beta} T_{S_0}^1 \longrightarrow 0,$$

where  $\Theta_{S_0} = \mathfrak{hom}(\Omega_{S_0}^1, \mathcal{O}_{S_0})$  is the tangent sheaf of  $S_0$ ,  $\Theta_{\mathcal{X}|_{S_0}}$  is the tangent sheaf of  $\mathcal{X}$  restricted to  $S_0$ ,  $\mathcal{N}_{S_0|\mathcal{X}}$  is the normal bundle of  $S_0$  in  $\mathcal{X}$ , and  $T_{S_0}^1$  is the first cotangent sheaf of  $S_0$ . The latter is supported on the singular locus  $\text{Sing}(S_0) = S_A \cap S_B$ . The kernel  $\mathcal{N}'_{S_0|\mathcal{X}}$  of  $\beta$  is the so-called *equisingular normal sheaf to  $S_0$  in  $\mathcal{X}$* , whose global sections are the first order locally trivial deformations of  $S_0$  in  $\mathcal{X}$ .

In the sequel an equisingular (first order) deformation of  $S_0$  in  $\mathcal{X}$  will be a (first order) locally trivial deformation of  $S_0$  in  $\mathcal{X}$ .

We will denote by  $\mathcal{ES}_{[S_0]}^{\mathcal{X}_0} \subseteq \mathcal{H}^{\mathcal{X}_0} \subset \mathcal{H}^{\mathcal{X}|\mathbb{D}}$  the locally closed set of equisingular deformations of  $S_0$  in  $\mathcal{X}_0$ . Similarly, if  $p \in S_0$  is a point, we will denote by  $\mathcal{ES}_{[S_0], p}^{\mathcal{X}_0} \subseteq \mathcal{H}^{\mathcal{X}_0} \subset \mathcal{H}^{\mathcal{X}|\mathbb{D}}$  the locally closed set of deformations of  $S_0$  in  $\mathcal{X}_0$ , which are equisingular at  $p$ .

**Lemma 3.4.** *If  $\mathcal{ES}_{[S_0]}^{\mathcal{X}|\mathbb{D}} \subseteq \mathcal{H}^{\mathcal{X}|\mathbb{D}}$  is the locally closed set of equisingular deformations of  $S_0$  in  $\mathcal{X}$ , then  $\mathcal{ES}_{[S_0]}^{\mathcal{X}|\mathbb{D}}$  coincides with  $\mathcal{ES}_{[S_0]}^{\mathcal{X}_0} \subseteq \mathcal{H}^{\mathcal{X}_0}$ .*

*Proof.* This is a straightforward consequence of Lemma 3.3.  $\square$

If  $\mathbb{T}_\delta \subset \mathcal{H}^{\mathcal{X}_0}$  is the Zariski closure of the family of surfaces in  $\mathcal{X}_0$  with  $\delta$  singularities of type  $T_1$ , then every irreducible component of  $\mathcal{ES}_{[S_0]}^{\mathcal{X}_0}$  is a Zariski open set in an irreducible component of  $\mathbb{T}_\delta$ .

Consider the rational map

$$\varphi : \mathcal{H}^{\mathcal{X}_0} \dashrightarrow \mathcal{H}^R$$

where  $\mathcal{H}^R$  is the Hilbert scheme of  $R$ , and  $\varphi$  maps the general subscheme of  $\mathcal{X}_0$  to its intersection with  $R$ .

**Lemma 3.5.** *Let  $[S_0] \in \mathbb{T}_\delta$  be a point corresponding to a surface  $S_0 = S_A \cup S_B$  as above and suppose that  $[S_0]$  is a smooth point of the Hilbert scheme  $\mathcal{H}^{\mathcal{X}_0}$  so that there is a unique component  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$  of  $\mathcal{H}^{\mathcal{X}_0}$  containing  $S_0$ . Let  $C$  be the curve cut out by  $S_0$  on  $R$ . Assume that  $h^1(\mathcal{N}_{C|R} \otimes \mathcal{I}_{\{p_1, \dots, p_\delta\}|R}) = 0$ , where  $p_1, \dots, p_\delta$  are the nodes of  $C$ , which implies that  $\mathcal{H}^R$  is smooth at the point  $[C]$  and that the Severi variety of curves on  $R$  with  $\delta$  nodes is smooth at the point  $[C]$  of codimension  $\delta$  in the unique irreducible component  $\mathcal{H}_C^R$  of the Hilbert scheme  $\mathcal{H}^R$  containing  $[C]$ . Suppose moreover that the map*

$$(21) \quad \varphi|_{\mathcal{H}_{S_0}^{\mathcal{X}_0}} : \mathcal{H}_{S_0}^{\mathcal{X}_0} \dashrightarrow \mathcal{H}_C^R$$

is dominant.

Then there is an irreducible component  $\mathbb{T}$  of  $\mathbb{T}_\delta$  containing  $S_0$  that has codimension at most  $\delta$  in  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$ .

*Proof.* Let  $V \subset \mathcal{H}_C^R$  be the unique irreducible component of the locally closed set of curves on  $R$  with  $\delta$  nodes that contains the point  $[C]$ . Since  $[C]$  sits in the image of  $\varphi|_{\mathcal{H}_{S_0}^{\mathcal{X}_0}}$ ,  $V$  intersects the image of  $\varphi|_{\mathcal{H}_{S_0}^{\mathcal{X}_0}}$ . Since  $\varphi|_{\mathcal{H}_{S_0}^{\mathcal{X}_0}}$  is dominant, the intersection of  $V$  with the image of  $\varphi|_{\mathcal{H}_{S_0}^{\mathcal{X}_0}}$  is an open dense subset of  $V$ , hence there is an irreducible component  $\mathbb{T}$  of  $\mathbb{T}_\delta$  containing  $S_0$  such that the map

$$\varphi|_{\mathbb{T}} : \mathbb{T} \dashrightarrow V$$

is dominant. Let  $a$  be the dimension of the general fibre of  $\varphi|_{\mathbb{T}}$  and let  $b$  be the dimension of the general fibre of  $\varphi|_{\mathcal{H}_{S_0}^{\mathcal{X}_0}}$ . Of course  $a \geq b$ . We have

$$\dim(\mathbb{T}) = \dim(V) + a, \text{ and } \dim(\mathcal{H}_{S_0}^{\mathcal{X}_0}) = \dim(\mathcal{H}_C^R) + b$$

hence

$$\dim(\mathcal{H}_{S_0}^{\mathcal{X}_0}) - \dim(\mathbb{T}) = \dim(\mathcal{H}_C^R) - \dim(V) + b - a \leq \delta$$

and the assertion follows.  $\square$

**Remark 3.6.** Note that in the previous lemma, one has that  $\mathbb{T}$  has exactly codimension  $\delta$  in  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$  if and only if  $a = b$ .

**Lemma 3.7.** *Assume that  $S_0 = S_A \cup S_B \subset \mathcal{X}_0$  is a surface with  $S_A$  and  $S_B$  smooth, intersecting transversally along  $R = A \cap B$ , except for  $\delta$  points  $p_1, \dots, p_\delta \in S_A \cap S_B$ , where  $S_0$  has a singularity of type  $T_1$ . Then the equisingular first order infinitesimal deformations of  $S_0$  in  $\mathcal{X}$  coincide with the equisingular first order infinitesimal deformations of  $S_0$  in  $\mathcal{X}_0$ . More precisely, we have that*

$$(22) \quad H^0(S_0, \mathcal{N}'_{S_0|\mathcal{X}}) \subseteq H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_1, \dots, p_\delta\}|\mathcal{X}_0}),$$

where  $I_{\{p_1, \dots, p_\delta\}|\mathcal{X}_0}$  is the ideal sheaf of  $\{p_1, \dots, p_\delta\}$  in  $\mathcal{X}_0$ .

*Proof.* Let  $p = p_i$ , for  $i = 1, \dots, \delta$ , be a point where  $S_0$  has a  $T_1$  singularity. Consider the localized exact sequence

$$(23) \quad 0 \longrightarrow \mathcal{N}'_{S_0|\mathcal{X}, p} \longrightarrow \mathcal{N}_{S_0|\mathcal{X}, p} \longrightarrow T_{S_0, p}^1 \longrightarrow 0.$$

Let  $(x, y, z, u, t)$  be an analytic coordinate system of  $\mathcal{X}$  centered at  $p$  such that  $\mathcal{X}$  is given by  $xy = t$  and such that we have the following identifications:

- the local ring  $\mathcal{O}_{S_0, p} = \mathcal{O}_{\mathcal{X}, p} / \mathcal{I}_{S_0|\mathcal{X}, p}$  of  $S_0$  at  $p$  is identified with  $\mathbb{C}[x, y, z, u] / (h_1, h_2)$ , localized at the origin, where  $h_1(x, y, z, u) = x + y + h_{12}(x, y, z, u)$ ,  $h_{12}(x, y, z, u) \in (x, y, z, u)^2$  and  $h_{12}(0, 0, z, u) = 0$  having a node at  $\underline{0} = p$ , and  $h_2(x, y, z, u) = xy$ ;

- the  $\mathcal{O}_{S_0, p}$ -module  $\mathcal{N}_{S_0|\mathcal{X}, p}$  is identified with the free  $\mathcal{O}_{\mathcal{X}, p}$ -module  $\mathfrak{hom}_{\mathcal{O}_{\mathcal{X}, p}}(\mathcal{I}_{S_0|\mathcal{X}, p}, \mathcal{O}_{S_0, p})$ , generated by the morphisms  $h_1^*$  and  $h_2^*$ , defined by

$$h_i^*(s_1(x, y, z, u)h_1(x, y, z, u) + s_2(x, y, z, u)h_2(x, y, z, u)) = s_i(x, y, z, u), \text{ for } i = 1, 2$$

and, finally,

- the  $\mathcal{O}_{S_0, p}$ -module

$$\begin{aligned} (\Theta_{\mathcal{X}|S_0})_p &\simeq \Theta_{\mathcal{X}, p} \otimes \mathcal{O}_{S_0, p} \\ &\simeq \langle \partial/\partial x, \partial/\partial y, \partial/\partial z, \partial/\partial u, \partial/\partial t \rangle_{\mathcal{O}_{S_0, p}} / \langle \partial/\partial t - x\partial/\partial y - y\partial/\partial x \rangle \end{aligned}$$

is identified with the free  $\mathcal{O}_{\mathcal{X}, p}$ -module generated by the derivatives  $\partial/\partial x, \partial/\partial y, \partial/\partial z, \partial/\partial u$ .

With these identifications, the localization  $\alpha_p : (\Theta_{\mathcal{X}|S_0})_p \rightarrow \mathcal{N}_{S_0|\mathcal{X}, p}$  of the sheaf map  $\alpha$  from (20) is defined by

$$\begin{aligned} \alpha_p(\partial/\partial x) &= \left( s = s_1h_1 + s_2h_2 \mapsto \partial s/\partial x =_{\mathcal{O}_{S_0, p}} s_1\partial h_1/\partial x + s_2\partial h_2/\partial x \right) \\ &= (1 + \partial h_{12}/\partial x)h_1^* + yh_2^*, \\ \alpha_p(\partial/\partial y) &= (1 + \partial h_{12}/\partial y)h_1^* + xh_2^*, \\ \alpha_p(\partial/\partial z) &= (\partial h_{12}/\partial z)h_1^* \text{ and} \\ \alpha_p(\partial/\partial u) &= (\partial h_{12}/\partial u)h_1^*. \end{aligned}$$

By definition of  $\mathcal{N}'_{S_0|\mathcal{X}}$ , a local section  $s$  of  $\mathcal{N}'_{S_0|\mathcal{X}, p}$  is such that there exists a local section  $v$  of  $\Theta_{\mathcal{X}|S_0, p}$ , with

$$v = v_x(x, y, z, u)\partial/\partial x + v_y(x, y, z, u)\partial/\partial y + v_z(x, y, z, u)\partial/\partial z + v_u(x, y, z, u)\partial/\partial u,$$

such that  $s = \alpha_p(v)$ . Hence, locally at  $p$ , first order equisingular deformations of  $S_0$  in  $\mathcal{X}$  have equations

$$(24) \quad \begin{cases} x + y + h_{12}(x, y, z, u) & + \epsilon(v_x(1 + \partial h_{12}/\partial x) + v_y(1 + \partial h_{12}/\partial y) \\ & + v_z(\partial h_{12}/\partial z) + v_u(\partial h_{12}/\partial u)) = 0 \\ xy + \epsilon(yv_x + xv_y) & = 0. \end{cases}$$

The first equation above gives a first order infinitesimal deformation of the Cartier divisor cutting  $S_0$  on  $\mathcal{X}_0$ , while the second equation gives a first order infinitesimal deformation of  $\mathcal{X}_0$  in  $\mathcal{X}$ . More precisely, by the exact sequence

$$0 \longrightarrow \Theta_{\mathcal{X}_0} \longrightarrow \Theta_{\mathcal{X}|\mathcal{X}_0} \longrightarrow \mathcal{N}_{\mathcal{X}_0|\mathcal{X}} \cong \mathcal{O}_{\mathcal{X}_0} \longrightarrow T_{\mathcal{X}_0}^1 \cong \mathcal{O}_R \longrightarrow 0,$$

one sees that  $xy + \epsilon(yv_x + xv_y) = 0$  is the local equation at  $p$  of a first order equisingular deformation of  $\mathcal{X}_0$  in  $\mathcal{X}$ . But  $H^0(\mathcal{X}_0, \mathcal{N}'_{\mathcal{X}_0|\mathcal{X}}) = H^0(\mathcal{X}_0, \mathcal{I}_{R|\mathcal{X}_0}) = 0$ . It follows that the polynomial  $yv_x(x, y, z, u) + xv_y(x, y, z, u)$  in the second equation of (24) must be identically zero, proving the first assertion of the Lemma. In particular, by expanding  $v_x$  and  $v_y$  in Taylor series, we see that

$$v_x(\mathbb{Q}) = v_y(\mathbb{Q}) = 0.$$

Looking at the first equation of (24), we have that  $\frac{\partial h_{12}}{\partial z}(\mathbb{Q}) = \frac{\partial h_{12}}{\partial u}(\mathbb{Q}) = 0$  since  $h_{12}(x, y, z, u) \in (x, y, z, u)^2$ . This shows the inclusion (22).  $\square$

**Remark 3.8.** The argument in the proof of Lemma 3.7 proves more than stated. In fact it proves that if  $S_0$  is any surface in  $\mathcal{X}_0$  with  $\delta$  singularities of type  $T_1$  at  $p_1, \dots, p_\delta$  (and may be other singularities which we do not care about), the first order infinitesimal deformations of  $S_0$  in  $\mathcal{X}$  which are equisingular at  $p_1, \dots, p_\delta$  are a linear subspace of  $H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_1, \dots, p_\delta\}}|_{\mathcal{X}_0})$ .

**Corollary 3.9.** *Same hypotheses as in Lemma 3.5. Assume moreover that*

$$(25) \quad H^1(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_1, \dots, p_\delta\}|\mathcal{X}_0}) = 0$$

or, equivalently, that

$$(26) \quad H^1(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) = 0 \text{ and } h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_1, \dots, p_\delta\}|\mathcal{X}_0}) = h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - \delta$$

(assuring that the Hilbert scheme  $\mathcal{H}^{\mathcal{X}_0}$  is smooth at  $[S_0]$ ). Then the schemes  $\mathcal{E}\mathcal{S}_{[S_0]}^{\mathcal{X}_0}$  and  $\mathbb{T}_\delta$  are smooth at  $[S_0]$  of dimension  $h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_1, \dots, p_\delta\}|\mathcal{X}_0})$ , with tangent space  $T_{[S_0]}(\mathcal{E}\mathcal{S}_{[S_0]}^{\mathcal{X}_0}) \simeq T_{[S_0]}(\mathbb{T}_\delta) \simeq H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_1, \dots, p_\delta\}|\mathcal{X}_0}) \simeq H^0(S_0, \mathcal{N}'_{S_0|\mathcal{X}})$ .

*Proof.* Consider the exact sequence

$$(27) \quad 0 \longrightarrow \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_1, \dots, p_\delta\}|\mathcal{X}_0} \longrightarrow \mathcal{N}_{S_0|\mathcal{X}_0} \longrightarrow \mathcal{N}_{S_0|\mathcal{X}_0} \otimes \mathcal{O}_{\{p_1, \dots, p_\delta\}} \longrightarrow 0,$$

from which one deduces that  $h^1(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_1, \dots, p_\delta\}|\mathcal{X}_0}) = 0$  if and only if  $h^1(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) = 0$  and

$$h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_1, \dots, p_\delta\}|\mathcal{X}_0}) = h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - \delta.$$

Assume that  $h^1(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_1, \dots, p_\delta\}|\mathcal{X}_0}) = 0$ . Thus  $[S_0]$  is a smooth point of  $\mathcal{H}^{\mathcal{X}_0}$ . In particular there exists a unique component  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$  of  $\mathcal{H}^{\mathcal{X}_0}$  containing  $[S_0]$  and having dimension  $h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0})$  at  $[S_0]$ . Now, by Lemma 3.5, one has that

$$h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - \delta = \dim(\mathcal{H}_{S_0}^{\mathcal{X}_0}) - \delta \leq \dim_{[S_0]}(\mathcal{E}\mathcal{S}_{[S_0]}^{\mathcal{X}_0}) \leq \dim(T_{[S_0]}(\mathcal{E}\mathcal{S}_{[S_0]}^{\mathcal{X}_0})).$$

On the other hand, by (22), one has that

$$\dim(T_{[S_0]}(\mathcal{E}\mathcal{S}_{[S_0]}^{\mathcal{X}_0})) \leq h^0(S_0, \mathcal{N}'_{S_0|\mathcal{X}}) \leq h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_1, \dots, p_\delta\}|\mathcal{X}_0}) = h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - \delta.$$

The corollary follows.  $\square$

We note the following:

**Lemma 3.10.** *Let  $[S_0] \in \mathcal{H}^{\mathcal{X}|\mathbb{D}}$  be any point corresponding to a Cartier divisor  $S_0 = S_A \cup S_B \subset \mathcal{X}_0$ . Assume that  $H^1(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) = 0$ . Then the space of first order infinitesimal deformations of  $S_0$  in  $\mathcal{X}$  is given by*

$$H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}}) \simeq H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) \oplus H^0(S_0, \mathcal{O}_{S_0})$$

and

$$H^1(S_0, \mathcal{N}_{S_0|\mathcal{X}}) \simeq H^1(S_0, \mathcal{O}_{S_0})$$

is an obstruction space for  $\mathcal{O}_{\mathcal{H}^{\mathcal{X}|\mathbb{D}}, [S_0]}$ .

*Proof.* By the hypothesis we have  $\text{Ext}^1(\mathcal{O}_{S_0}, \mathcal{N}_{S_0|\mathcal{X}_0}) \simeq H^1(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) = 0$  and by the exact sequence

$$(28) \quad 0 \longrightarrow \mathcal{N}_{S_0|\mathcal{X}_0} \longrightarrow \mathcal{N}_{S_0|\mathcal{X}} \longrightarrow \mathcal{N}_{\mathcal{X}_0|\mathcal{X}}|_{S_0} = \mathcal{O}_{S_0} \longrightarrow 0,$$

we have that

$$\mathcal{N}_{S_0|\mathcal{X}} \simeq \mathcal{N}_{S_0|\mathcal{X}_0} \oplus \mathcal{O}_{S_0}.$$

The statement then follows by standard deformation theory.  $\square$

**Corollary 3.11.** *Let  $[S_0] \in \mathcal{H}^{\mathcal{X}|\mathbb{D}}$  be a point corresponding to a Cartier divisor  $S_0 = S_A \cup S_B \subset \mathcal{X}_0$ . Assume that  $[S_0]$  belongs to an irreducible component  $\mathcal{H}$  of  $\mathcal{H}^{\mathcal{X}|\mathbb{D}}$  that dominates  $\mathbb{D}$ . Suppose that  $H^1(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) = 0$ . Then  $[S_0]$  is a smooth point for  $\mathcal{H}^{\mathcal{X}|\mathbb{D}}$  and the  $\dim(\mathcal{H}) = \dim(\mathcal{H}_{[S_0]}^{\mathcal{X}_0}) + 1$ .*

*Proof.* One has  $\dim(\mathcal{H}) \geq \dim(\mathcal{H}_{[S_0]}^{\mathcal{X}_0}) + 1 = h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) + 1$ . On the other hand,  $\dim(\mathcal{H}) \leq h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}}) = h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) + 1$  by Lemma 3.10. The assertion follows.  $\square$

**Corollary 3.12.** *In the same setting as in Lemma 3.7 and same hypotheses as in Lemma 3.5, suppose that (25) (or equivalently (26)) holds, assuring that  $\mathbb{T}_\delta$  is smooth at  $[S_0]$  of dimension  $h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_1, \dots, p_\delta\}|\mathcal{X}_0}) = h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - \delta$ . Then for every positive integer  $r < \delta$ , the variety  $\mathbb{T}_r$  is non-empty and  $[S_0] \in \mathbb{T}_r$ . More precisely, in an analytic neighborhood of  $[S_0]$ ,  $\mathbb{T}_r$  consists of  $\binom{\delta}{r}$  smooth analytic branches each of dimension  $h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - r$  that intersect transversally at  $[S_0]$  along a smooth analytic branch of  $\mathbb{T}_\delta$ .*

*Informally speaking, this is as saying that the  $\delta$  singularities of type  $T_1$  of  $S_0$  can be independently smoothed inside  $\mathcal{X}_0$ .*

*Proof.* We first prove the assertion for  $r = 1$ .

Let  $\mathbb{T}$  be the closure of the subset of the Hilbert scheme  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$  consisting of all surfaces  $S'_0$  such that the intersection curve of  $S'_0$  with  $R = A \cap B$  is singular with at most nodes. Observe that  $[S_0] \in \mathbb{T}$ .

We claim that any irreducible component of  $\mathbb{T}$  that contains  $[S_0]$  has exactly codimension 1 in  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$ . Indeed, let  $\mathbb{T}'$  be such a component. Consider the dominant map  $\varphi|_{\mathcal{H}_{S_0}^{\mathcal{X}_0}}$  as in (21), which is defined at a general point of  $\mathbb{T}'$ .

By our hypotheses, the general element in  $\mathcal{H}_C^R$  is a smooth curve, hence  $\mathbb{T}'$  has codimension at least 1 in  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$ . Let  $\mathcal{T}$  be the image of the restriction of  $\varphi|_{\mathcal{H}_{S_0}^{\mathcal{X}_0}}$  to  $\mathbb{T}'$ . Then  $\mathcal{T}$  has codimension 1 in  $\mathcal{H}_C^R$ .

Let  $\alpha$  be the dimension of the general fibre of  $\varphi|_{\mathcal{H}_{S_0}^{\mathcal{X}_0}}$  and let  $\beta$  be the dimension of the general fibre of the restriction of  $\varphi|_{\mathcal{H}_{S_0}^{\mathcal{X}_0}}$  to  $\mathbb{T}'$ . One has  $\alpha \leq \beta$ . Then

$$\dim(\mathcal{H}_{S_0}^{\mathcal{X}_0}) = \dim(\mathcal{H}_C^R) + \alpha$$

and

$$\dim(\mathbb{T}') = \dim(\mathcal{T}) + \beta = \dim(\mathcal{H}_C^R) - 1 + \beta \geq \dim(\mathcal{H}_C^R) - 1 + \alpha = \dim(\mathcal{H}_{S_0}^{\mathcal{X}_0}) - 1.$$

Since  $\dim(\mathbb{T}') < \dim(\mathcal{H}_{S_0}^{\mathcal{X}_0})$  we have  $\dim(\mathbb{T}') = \dim(\mathcal{H}_{S_0}^{\mathcal{X}_0}) - 1$  and  $\alpha = \beta$ , as claimed.

Now, we consider a suitably small analytic open neighborhood  $U$  of  $[S_0]$  in  $\mathbb{T}$ . Every surface  $S'_0$  such that  $[S'_0] \in U$  has at most  $\delta$  singularities of type  $T_1$ .

Consider the variety  $I \subset U \times R$  consisting of all pairs  $([S'_0], q)$  with  $q$  a  $T_1$  singularity of  $S'_0$ . Let  $\pi_1 : I \rightarrow U$  and  $\pi_2 : I \rightarrow R$  be the two projections. The former one has finite fibres, implying the every irreducible component of  $I$  has dimension  $\dim(\mathcal{H}_{S_0}^{\mathcal{X}_0}) - 1$ . As for the latter, it is dominant, because we assume the hypotheses of Lemma 3.5. Moreover, if  $q \in R$  is a point, the fibre  $\pi_2^{-1}(q)$  is the locally closed set of surfaces  $S'_0$  in  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$  having a  $T_1$  singularity in  $q$ .

Let  $V_i$  be a sufficiently small analytic neighborhood of  $p_i$  in  $R$  for  $i = 1, \dots, \delta$ . By the above considerations,  $\pi_1(\pi_2^{-1}(V_i)) \subseteq U$  is an analytic open set that parametrizes deformations of  $S_0$  which are analytically equisingular at  $p_i$ . Hence, by Remark 3.8, the tangent space to  $\pi_1(\pi_2^{-1}(V_i))$  at  $[S_0]$  is contained in  $H^0(S, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{p_i|\mathcal{X}_0})$  and

$$\dim_{[S_0]}(\mathcal{H}_{S_0}^{\mathcal{X}_0}) - 1 = \dim_{[S_0]}(\pi_1(\pi_2^{-1}(V_i))) \leq h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{p_i|\mathcal{X}_0}).$$

If (25) holds, then  $h^1(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{p_i|\mathcal{X}_0}) = 0$  for all  $i = 1, \dots, \delta$ , and one has

$$h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{p_i|\mathcal{X}_0}) = h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - 1 = \dim_{[S_0]}(\mathcal{H}_{S_0}^{\mathcal{X}_0}) - 1.$$

Thus  $\pi_1(\pi_2^{-1}(V_i))$ , that is an open analytic subset of  $\mathcal{E}\mathcal{S}_{[S_0], p_i}^{\mathcal{X}_0}$ , is an analytic branch of  $U$  of dimension  $h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - 1$ , smooth at  $[S_0]$ .

Next we prove that the general element  $[S'_0]$  in  $\pi_1(\pi_2^{-1}(V_i))$  has a unique  $T_1$  singularity. We argue for the case  $i = 1$  and the proof is analogous in the other cases. Suppose this is not the case, and that  $S'_0$  has  $s$  singularities  $q_1, \dots, q_s$  of type  $T_1$  with  $s > 1$ . When  $S'_0$  specializes to  $S_0$ ,  $q_1, \dots, q_s$  specialize, say, to  $p_1, \dots, p_s$ . By



the same argument as above, the tangent space to  $\pi_1(\pi_2^{-1}(V_1))$  at  $[S'_0]$  is contained in  $H^0(S_0, \mathcal{N}_{S'_0|\mathcal{X}_0} \otimes I_{\{q_1, \dots, q_s\}|\mathcal{X}_0})$  and, under the hypothesis (25), one has

$$h^0(S'_0, \mathcal{N}_{S'_0|\mathcal{X}_0} \otimes I_{\{q_1, \dots, q_s\}|\mathcal{X}_0}) \leq h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_1, \dots, p_s\}|\mathcal{X}_0}) = h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - s < h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - 1$$

and this is a contradiction. This proves the assertion for  $r = 1$ .

Consider now the case  $\delta > r > 1$ . Fix  $p_{i_1}, \dots, p_{i_r}$  distinct points among  $p_1, \dots, p_\delta$ . The intersection

$$\mathfrak{T}_{i_1, \dots, i_r} := \bigcap_{j=1}^r \pi_1(\pi_2^{-1}(V_{i_j})),$$

that is an analytic open subset of  $\mathcal{ES}_{[S_0], p_{i_1}, \dots, p_{i_r}}^{\mathcal{X}_0}$ , is an analytic variety in  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$  parametrizing deformations of  $S_0$  that are analytic equisingular at the points  $p_{i_1}, \dots, p_{i_r}$ . With the same argument as above, one sees that, under the hypothesis (25),  $\mathfrak{T}_{i_1, \dots, i_r}$  is smooth of codimension  $r$  in  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$ , with tangent space at  $[S_0]$  given by  $H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\{p_{i_1}, \dots, p_{i_r}\}|\mathcal{X}_0})$ .

Moreover, again by the same argument as above, the general element  $S'_0$  in  $\mathfrak{T}_{i_1, \dots, i_r}$  has exactly  $r$  singularities of type  $T_1$  at points specializing to  $p_{i_1}, \dots, p_{i_r}$  when  $S'_0$  specializes to  $S_0$ . So  $\mathfrak{T}_{i_1, \dots, i_r}$  is a smooth analytic branch of  $\mathbb{T}_r$  containing  $[S_0]$ , and this ends the proof of the corollary.  $\square$

Let now  $\mathbb{T}_{\delta_A, \delta_B, \delta_R} \subseteq \mathcal{H}^{\mathcal{X}_0}$  be the Zariski closure of the family of surfaces  $S_0 = S_A \cup S_B$  in  $\mathcal{X}_0$  with  $\delta_A$  nodes on  $A$  and  $\delta_B$  nodes on  $B$  off  $R$  and  $\delta_R$  singularities of type  $T_1$  on  $R$ .

**Corollary 3.13.** *Let  $S_0 = S_A \cup S_B$  be a reduced effective Cartier divisor such that  $S_A$  and  $S_B$  have respectively  $\delta_A$  and  $\delta_B$  nodes  $p_{A,1}, \dots, p_{A,\delta_A}$  and  $p_{B,1}, \dots, p_{B,\delta_B}$  off  $R$ , are elsewhere smooth and intersect transversally along a curve  $C = S_A \cap S_B$ , except for  $\delta_R$  distinct points  $p_{R,1}, \dots, p_{R,\delta_R} \in C \subset R$  where  $S_0$  has singularities of type  $T_1$ . Let  $\mathcal{ES}_{[S_0]}^{\mathcal{X}_0}$  be the locally closed set of equisingular deformations of  $S_0$  in  $\mathcal{X}_0$ . Consider the ideal sheaf  $\mathcal{I}_{\mathfrak{Z}|\mathcal{X}_0}$  in  $\mathcal{X}_0$  of the 0-dimensional reduced scheme  $\mathfrak{Z}$  of length  $\delta = \delta_A + \delta_B + \delta_R$  given by*

$$\mathfrak{Z} = \sum_{i=1}^{\delta_A} p_{A,i} + \sum_{i=1}^{\delta_B} p_{B,i} + \sum_{i=1}^{\delta_R} p_{R,i}.$$

Thus

$$(29) \quad T_{[S_0]}(\mathcal{ES}_{[S_0]}^{\mathcal{X}_0}) \subseteq H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes \mathcal{I}_{\mathfrak{Z}|\mathcal{X}_0}).$$

If

$$(30) \quad h^1(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes \mathcal{I}_{\mathfrak{Z}|\mathcal{X}_0}) = h^1(C, \mathcal{N}_{C|R} \otimes \mathcal{I}_{\{p_{R,1}, \dots, p_{R,\delta_R}\}|R}) = 0$$

and the map

$$\varphi_{|\mathcal{H}_{S_0}^{\mathcal{X}_0}} : \mathcal{H}_{S_0}^{\mathcal{X}_0} \dashrightarrow \mathcal{H}_C^R$$

defined as in Lemma 3.5 is dominant, then the equality holds in (29), i.e., the locally closed set  $\mathcal{ES}_{[S_0]}^{\mathcal{X}_0}$  of locally trivial deformations of  $S_0$  in  $\mathcal{X}_0$  is smooth at  $[S_0]$  of dimension  $h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes \mathcal{I}_{\mathfrak{Z}|\mathcal{X}_0}) = h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - \delta$ . In particular there exists only one irreducible component  $\mathbb{T} \subset \mathbb{T}_{\delta_A, \delta_B, \delta_R}$  containing the point  $[S_0]$  (which is smooth at  $[S_0]$  and contains  $\mathcal{ES}_{[S_0]}^{\mathcal{X}_0}$  as a Zariski open set). Moreover, under these hypotheses, the singularities of  $S_0$  may be smoothed independently in  $\mathcal{X}_0$ . More precisely, for every  $\delta'_A \leq \delta_A$ ,  $\delta'_B \leq \delta_B$  and  $\delta'_R \leq \delta_R$  we have that  $\mathbb{T}_{\delta'_A, \delta'_B, \delta'_R}$  is non-empty and  $[S_0] \in \mathbb{T}_{\delta'_A, \delta'_B, \delta'_R}$ . In an analytic neighborhood of  $[S_0]$ ,  $\mathbb{T}_{\delta'_A, \delta'_B, \delta'_R}$  consists of

$$\begin{pmatrix} \delta_R \\ \delta'_R \end{pmatrix} \begin{pmatrix} \delta_A \\ \delta'_A \end{pmatrix} \begin{pmatrix} \delta_B \\ \delta'_B \end{pmatrix}$$

smooth analytic branches of dimension  $h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - \delta'$ , where  $\delta' = \delta'_A + \delta'_B + \delta'_R$ , that intersect transversally at  $[S_0]$  along a smooth analytic branch of  $\mathbb{T} \subset \mathbb{T}_{\delta_A, \delta_B, \delta_R}$ , corresponding to deformations of  $[S_0]$  preserving  $\delta'_R$  points of type  $T_1$  and  $\delta'_A$  nodes on  $A$  and  $\delta'_B$  nodes on  $B$ .

*Proof.* Let  $S_0$  be a surface as in the statement. The inclusion (29) follows from Lemma 3.7, Remark 3.8 and well known deformation theory of nodal surfaces (see [11, §2.3]). It can be proved by using (20). In particular, if one localizes (20) at a node  $p$  of  $S_0$ , then  $H^0(S_0, T_{S_0, p}^1) \cong \mathbb{C}$  can be identified with the tangent space to the versal deformation space of a node.

Now we want to prove that, under the hypotheses of the corollary, the locally closed set  $\mathcal{ES}_{[S_0]}^{\mathcal{X}_0}$  is smooth at  $[S_0]$  of codimension  $\delta$  in the Hilbert scheme  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$ .

Let  $C \subset R$  be the  $\delta_R$ -nodal curve cut out by  $S_0$  on  $R$ . By the hypothesis (30), one has that  $h^1(C, \mathcal{N}_{C|R} \otimes \mathcal{I}_{\{p_{R,1}, \dots, p_{R, \delta_R}\}}|R) = 0$ . This implies that  $[C]$  is a smooth point of the locally closed Severi variety  $\mathcal{V}_\delta$  of  $\delta$ -nodal curves in  $\mathcal{H}_C^R$ . Let  $V$  be the unique irreducible component of  $\mathcal{V}_\delta$  containing  $[C]$ . Now, as we saw in Lemma 3.5,  $\varphi_{|\mathcal{H}_{S_0}^{\mathcal{X}_0}}^{-1}(V)$  has at least one irreducible component of codimension at most  $\delta_R$  in  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$ . On the other hand, by Lemma 3.7 we have that

$$T_{[S_0]}(\varphi_{|\mathcal{H}_{S_0}^{\mathcal{X}_0}}^{-1}(V)) \subseteq H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes \mathcal{I}_{\{p_{R,1}, \dots, p_{R, \delta_R}\}}|\mathcal{X}_0)$$

and by (30) we have  $h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes \mathcal{I}_{\{p_{R,1}, \dots, p_{R, \delta_R}\}}|\mathcal{X}_0) = \dim(\mathcal{H}_{S_0}^{\mathcal{X}_0}) - \delta_R$ . Thus  $\varphi_{|\mathcal{H}_{S_0}^{\mathcal{X}_0}}^{-1}(V)$  is smooth at  $[S_0]$  of codimension  $\delta_R$ . We observe that  $\varphi_{|\mathcal{H}_{S_0}^{\mathcal{X}_0}}^{-1}(V)$  is an analytic open set of the variety  $\mathcal{ES}_{[S_0], p_{R,1}, \dots, p_{R, \delta_R}}^{\mathcal{X}_0}$  of deformations of  $S_0$  in  $\mathcal{X}_0$  that are locally trivial at every  $T_1$  singularity  $p_{R,i}$  and we just proved that

$$T_{[S_0]}(\mathcal{ES}_{[S_0], p_{R,1}, \dots, p_{R, \delta_R}}^{\mathcal{X}_0}) = T_{[S_0]}(\varphi_{|\mathcal{H}_{S_0}^{\mathcal{X}_0}}^{-1}(V)) = H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes \mathcal{I}_{\{p_{R,1}, \dots, p_{R, \delta_R}\}}|\mathcal{X}_0).$$

We moreover observe that the general element  $[S'_0]$  of  $\varphi_{|\mathcal{H}_{S_0}^{\mathcal{X}_0}}^{-1}(V)$  corresponds to a surface  $S'_0 = S'_A \cup S'_B$ , where  $S'_A$  and  $S'_B$  intersect transversally along a curve  $C'$  on  $R$ , except for  $\delta_R$  points  $p'_{R,1}, \dots, p'_{R, \delta_R} \in C'$ , which are singularities of type  $T_1$  of  $S'_0$ , and specialize to  $p_{R,1}, \dots, p_{R, \delta_R}$  as  $S'_0$  specializes to  $S_0$ .

We claim that  $S'_A$  and  $S'_B$  are smooth outside  $R$ . Indeed, since  $[S_0]$  belongs to  $\varphi_{|\mathcal{H}_{S_0}^{\mathcal{X}_0}}^{-1}(V)$ , the surface  $S'_0$  may have at most  $\delta'_A \leq \delta_A$  nodes  $p_{A,1}, \dots, p_{A, \delta'_A}$  on  $A$  and  $\delta'_B \leq \delta_B$  nodes  $p_{B,1}, \dots, p_{B, \delta'_B}$  on  $B$ , deformations of  $\delta'_A$  nodes of  $S_0$  on  $A$  and  $\delta'_B$  nodes of  $S_0$  on  $B$ . If this happens, denoting by  $\mathfrak{Z}'$  the scheme of singular points of  $S'_0$ , then  $T_{[S'_0]}(\mathcal{ES}_{[S_0], p_{R,1}, \dots, p_{R, \delta_R}}^{\mathcal{X}_0}) = T_{[S'_0]}(\varphi_{|\mathcal{H}_{S_0}^{\mathcal{X}_0}}^{-1}(V)) \subseteq H^0(S'_0, \mathcal{N}_{S'_0|\mathcal{X}_0} \otimes \mathcal{I}_{\mathfrak{Z}'|\mathcal{X}_0})$ . But, once again by (30) and by semicontinuity, one has that  $h^0(S'_0, \mathcal{N}_{S'_0|\mathcal{X}_0} \otimes \mathcal{I}_{\mathfrak{Z}'|\mathcal{X}_0}) = H^0(S'_0, \mathcal{N}_{S'_0|\mathcal{X}_0}) - \delta_R - \delta'_A - \delta'_B$ . It follows that  $\delta'_A = \delta'_B = 0$ , i.e.  $S'_A$  and  $S'_B$  are smooth off  $R$  and  $\varphi_{|\mathcal{H}_{S_0}^{\mathcal{X}_0}}^{-1}(V)$  is a locally closed set in one irreducible component  $\mathbb{T} \subset \mathbb{T}_{\delta_R}$ , smooth at  $[S_0]$ . We just proved that, under our hypotheses, one may deform  $S_0$  in  $\mathcal{X}_0$  by smoothing all nodes of  $S_0$  and by preserving all  $T_1$  singularities.

Let now  $\mathfrak{Z}_A$  and  $\mathfrak{Z}_B$  be respectively the scheme of nodes of  $S_0$  on  $A$  and  $B$ . Let  $\mathcal{ES}_{[S_0], \mathfrak{Z}_A, \mathfrak{Z}_B}^{\mathcal{X}_0}$  the scheme of deformations of  $S_0$  which are locally trivial at every node of  $S_0$ . By standard deformation theory of nodal surfaces, one has that, under the hypothesis  $h^1(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes \mathcal{I}_{\mathfrak{Z}_A \cup \mathfrak{Z}_B}|\mathcal{X}_0) = 0$  (that holds by (30)),  $\mathcal{ES}_{[S_0], \mathfrak{Z}_A, \mathfrak{Z}_B}^{\mathcal{X}_0}$  is smooth of codimension  $\delta_A + \delta_B$  in  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$  at  $[S_0]$  and moreover

$$T_{[S_0]}(\mathcal{ES}_{[S_0], \mathfrak{Z}_A, \mathfrak{Z}_B}^{\mathcal{X}_0}) = H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes \mathcal{I}_{\mathfrak{Z}_A \cup \mathfrak{Z}_B}|\mathcal{X}_0).$$

With a similar argument as above, one sees that the general element  $[S_0]$  of  $\mathcal{ES}_{[S_0],\mathfrak{z}_A,\mathfrak{z}_B}^{\mathcal{X}_0}$  corresponds to a surface  $\tilde{S}_0 = \tilde{S}_A \cup \tilde{S}_B$ , with  $\tilde{S}_A$  and  $\tilde{S}_B$  intersecting transversally along a smooth curve  $\tilde{C} \subset R$  and having, respectively,  $\delta_A$  and  $\delta_B$  nodes as singularities. In particular  $\mathcal{ES}_{[S_0],\mathfrak{z}_A,\mathfrak{z}_B}^{\mathcal{X}_0}$  is a locally closed set in an irreducible component  $\tilde{\mathbb{T}}$  of  $\mathbb{T}_{\delta_A,\delta_B,0}$ , of which  $[S_0]$  is a smooth point.

Now the equisingular deformation locus  $\mathcal{ES}_{[S_0]}^{\mathcal{X}_0}$  of  $S_0$  in  $\mathcal{X}_0$  is the intersection of the loci  $\mathcal{ES}_{[S_0],p_{R,1},\dots,p_{R,\delta_R}}^{\mathcal{X}_0}$  and  $\mathcal{ES}_{[S_0],\mathfrak{z}_A,\mathfrak{z}_B}^{\mathcal{X}_0}$ . Hence  $\mathcal{ES}_{[S_0]}^{\mathcal{X}_0}$  has codimension at most  $\delta$  in  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$ , because  $[S_0]$  is a smooth point of  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$ .

On the other hand, one has

$$\begin{aligned} T_{[S_0]}(\mathcal{ES}_{[S_0]}^{\mathcal{X}_0}) &= T_{[S_0]}(\mathcal{ES}_{[S_0],p_{R,1},\dots,p_{R,\delta_R}}^{\mathcal{X}_0}) \cap T_{[S_0]}(\mathcal{ES}_{[S_0],\mathfrak{z}_A,\mathfrak{z}_B}^{\mathcal{X}_0}) \\ &= H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes \mathcal{I}_{\{p_{R,1},\dots,p_{R,\delta_R}\}|\mathcal{X}_0}) \cap H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes \mathcal{I}_{\mathfrak{z}_A \cup \mathfrak{z}_B|\mathcal{X}_0}) \\ &= H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes \mathcal{I}_{\mathfrak{z}|\mathcal{X}_0}). \end{aligned}$$

By (30), we have

$$h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes \mathcal{I}_{\mathfrak{z}|\mathcal{X}_0}) = h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - \delta$$

this proves that  $\mathcal{ES}_{[S_0]}^{\mathcal{X}_0}$  is smooth at  $[S_0]$  of codimension exactly  $\delta$  in  $\mathcal{H}_{S_0}^{\mathcal{X}_0}$ , as wanted. This proves the first part of the corollary.

The second part is proved with analogous arguments as the ones used in the proof of Corollary 3.12.  $\square$

**Proposition 3.14.** *Let  $S_0 = S_A \cup S_B$  be a reduced effective Cartier divisor as in the statement of Corollary 3.13. Assume that  $[S_0]$  belongs to an irreducible component  $\mathcal{H}$  of  $\mathcal{H}^{\mathcal{X}|\mathbb{D}}$  that dominates  $\mathbb{D}$  and that (30) holds.*

*Let  $\mathcal{ES}_{[S_0],\mathfrak{z}_A,\mathfrak{z}_B}^{\mathcal{X}}$  be the locus in  $\mathcal{H}^{\mathcal{X}}$  of deformations of  $S_0$  which are equisingular at every node of  $S_0$ . Then  $\mathcal{ES}_{[S_0],\mathfrak{z}_A,\mathfrak{z}_B}^{\mathcal{X}}$  is generically smooth and it contains  $\mathcal{ES}_{[S_0],\mathfrak{z}_A,\mathfrak{z}_B}^{\mathcal{X}_0}$  as a subscheme of codimension 1.*

*In simple words,  $[S_0]$  can be deformed out of  $\mathcal{X}_0$  preserving the  $\delta_A + \delta_B$  nodes.*

*Proof.* From the hypotheses, it follows that  $H^1(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) = 0$ . Then, by Corollary 3.11,  $\mathcal{H}^{\mathcal{X}|\mathbb{D}}$  is smooth at  $[S_0]$  with dimension  $h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) + 1 = \dim(\mathcal{H}_{S_0}^{\mathcal{X}_0}) + 1$ . By standard deformation theory, there is an analytic neighborhood  $U$  of  $[S_0]$  in  $\mathcal{H}$  and a versal morphism

$$f : U \longrightarrow \prod_{p \in \mathfrak{z}_A + \mathfrak{z}_B} \Delta_p$$

where  $\Delta_p$  is the versal deformation space of a node, and therefore it has dimension 1. Let  $U' = U \cap \mathcal{H}_{S_0}^{\mathcal{X}_0}$ . Then  $f$  restricts to

$$g : U' \longrightarrow \prod_{p \in \mathfrak{z}_A + \mathfrak{z}_B} \Delta_p$$

The differential of  $g$  at  $[S_0]$  is

$$H^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) \rightarrow \prod_{p \in \mathfrak{z}_A + \mathfrak{z}_B} T_{S_0,p}^1 \cong \mathbb{C}^{\delta_A + \delta_B}$$

and this map is surjective by the hypotheses. Hence  $g$  has maximal rank at  $[S_0]$  and therefore also  $f$  is of maximal rank at  $[S_0]$ . Hence  $f^{-1}(0)$  and  $g^{-1}(0)$  are analytic subvarieties of  $U$  and  $U'$  respectively, smooth at  $[S_0]$  and of codimension  $\delta_A + \delta_B$  in  $U$  and  $U'$  respectively. By versality,  $f^{-1}(0)$  (resp.  $g^{-1}(0)$ ) coincides with  $\mathcal{ES}_{[S_0],\mathfrak{z}_A,\mathfrak{z}_B}^{\mathcal{X}}$  (resp.  $\mathcal{ES}_{[S_0],\mathfrak{z}_A,\mathfrak{z}_B}^{\mathcal{X}_0}$ ). The statement follows.  $\square$

**3.2.3. Global deformations of surfaces with  $T_1$  singularities to nodal surfaces.** In this section we will assume the following set up. We have the family  $\pi : \mathcal{X} \rightarrow \mathbb{D}$  as usual with its relative Hilbert scheme  $\mathcal{H}^{\mathcal{X}|\mathbb{D}}$ , whose fibre over  $t \in \mathbb{D}$  is the Hilbert scheme of  $\mathcal{H}^{\mathcal{X}_t}$  of  $\mathcal{X}_t$ .

Let  $\mathcal{V}_\delta^{\mathcal{X}|\mathbb{D}}$  be the Zariski closure in  $\mathcal{H}^{\mathcal{X}|\mathbb{D}}$  of the relative Severi variety  $\mathcal{W}_\delta^{\mathcal{X} \setminus \mathcal{X}_0|\mathbb{D} \setminus 0} \subset \mathcal{H}^{\mathcal{X} \setminus \mathcal{X}_0|\mathbb{D} \setminus 0}$  of  $\delta$ -nodal surfaces. We want to provide sufficient conditions for  $\mathcal{V}_\delta^{\mathcal{X}|\mathbb{D}}$  to be non-empty.

We will suppose that we have a line bundle  $\mathcal{L}$  on  $\mathcal{X}$  with the following properties:

- (1)  $h^0(\mathcal{X}_t, \mathcal{L}_t)$  is a constant  $r + 1$  in  $t$  and greater or equal than 4. In particular every surface  $S_0$  in  $|\mathcal{L}_0|$  belongs to an irreducible component  $\mathcal{H}$  of the relative Hilbert scheme  $\mathcal{H}^{\mathcal{X}|\mathbb{D}}$  that dominates  $\mathbb{D}$ ;
- (2)  $|\mathcal{L}_0|$  is base point free, so that we can assume that  $|\mathcal{L}_t|$  is base point free for all  $t \in \mathbb{D}$ ;
- (3) if  $p_t \in \mathcal{X}_t$  is a general point, then the general surface in  $|\mathcal{L}_t|$  with a singular point at  $p_t$  is singular only at finitely many points, for the general  $t \in \mathbb{D}$ .

In this setting we can consider the rank  $r$  projective bundle  $\bar{\pi} : \mathbb{P}(\pi_*(\mathcal{L})) \rightarrow \mathbb{D}$ . A point in  $\mathbb{P} := \mathbb{P}(\pi_*(\mathcal{L}))$  that maps to  $t \in \mathbb{D}$  is a non-zero section of  $H^0(\mathcal{X}_t, \mathcal{L}_t)$  up to a constant. In particular, if  $t \neq 0$ , a point in  $\mathbb{P}$  corresponds to a surface in  $|\mathcal{L}_t|$ . Consider the open Zariski subset  $\mathbb{P}' := \bar{\pi}^{-1}(\mathbb{D} \setminus \{0\})$ , which, by the above considerations, can be regarded as a subvariety of the relative Hilbert scheme of surfaces in  $\mathcal{X}$ . By a standard parameter count, one sees that there is a subscheme  $Z$  of pure codimension 1 in  $\mathbb{P}'$  whose points correspond to sections vanishing along singular surfaces. We will denote by  $\bar{Z}$  the closure of  $Z$  in  $\mathbb{P}$ , that has also codimension 1.

**Proposition 3.15.** *Set up as above with the following further condition: the subspace of sections of  $H^0(\mathcal{X}_0, \mathcal{L}_0)$  that vanish on  $R = A \cap B$ , with  $A$  and  $B$  the irreducible components of  $\mathcal{X}_0$ , has codimension strictly larger than 1 in  $H^0(\mathcal{X}_0, \mathcal{L}_0)$ .*

Let  $S_0 = S_A \cup S_B \subset \mathcal{X}_0$  be a surface corresponding to a section of  $\mathcal{L}_0$ . We suppose that:

- (a)  $S_A$  and  $S_B$  are smooth and intersect transversally along a curve  $C = S_A \cap S_B$ , except for a point  $p = p_1 \in C \subset R$  where  $S_0$  has a singularity of type  $T_1$  and the hypotheses of Lemma 3.5 hold for  $\delta = 1$ ;
- (b) the sublinear system  $\mathcal{L}_0(2, p)$  of  $|\mathcal{L}_0|$  of surfaces with at least a  $T_1$  singularity at  $p$  has codimension 3 in  $|\mathcal{L}_0|$ ;
- (c) (25) (for  $\delta = 1$ ) holds for  $S_0$ .

Then:

- (i)  $\mathbb{T}_1$  is smooth at  $[S_0]$  of codimension 1 in  $\mathcal{H}_0$ ;
- (ii)  $S_0$  can be deformed to a 1-nodal surface  $S_t \subset \mathcal{X}_t$ ;
- (iii) if  $\mathbb{T} \subset \mathbb{T}_1$  is the unique irreducible component containing  $[S_0]$ , then there exists a reduced, irreducible component  $\mathcal{V} \subset \mathcal{V}_1^{\mathcal{X}|\mathbb{D}}$  of dimension  $\dim(\mathcal{H}) - 1$  whose central fibre  $\mathcal{V}_0$  contains  $\mathbb{T}$  as an irreducible component.

Before giving the proof of the proposition, we make a preliminary lemma. For this we need some notation. Let  $I \subset |\mathcal{L}_0| \times R$ , with  $R = A \cap B$ , be the locally closed subset consisting of pairs  $(S_0, p)$  such that  $S_0$  cuts out on  $R$  a curve singular at  $p$ . We will consider the two projections  $\pi_1 : I \rightarrow |\mathcal{L}_0|$  and  $\pi_2 : I \rightarrow R$ . Note that if  $p \in R$ , then  $\pi_2^{-1}(p)$  can be identified with  $\mathcal{L}_0(2, p)$ .

**Lemma 3.16.** (i) *There is at most one irreducible component  $I'$  of  $I$  such that the restriction of  $\pi_2$  to  $I'$  is dominant to  $R$  via  $\pi_2$ .*

(ii) *If  $I'$  exists, and if its general element  $(S_0, p)$  is such that  $S_0$  cuts out on  $R$  a curve with finitely many singular points, then  $\mathcal{L}_0(2, p)$  has codimension 3 in  $|\mathcal{L}_0|$ . Moreover  $\dim(I') = \dim(|\mathcal{L}_0|) - 1$ , and its image in  $|\mathcal{L}_0|$  via  $\pi_1$  has codimension 1 in  $|\mathcal{L}_0|$ .*

(iii) If there is a pair  $(S_0, p)$  in  $I$  such that  $S_0$  cuts out on  $R$  a curve with finitely many singular points and if  $\mathcal{L}_0(2, p)$  has codimension 3 in  $|\mathcal{L}_0|$ , then  $(S_0, p)$  belongs to an irreducible component  $I'$  of  $I$  dominating  $R$  via  $\pi_2$ . For this component one has  $\dim(I') = \dim(|\mathcal{L}_0|) - 1$ , and its image in  $|\mathcal{L}_0|$  via  $\pi_1$  has codimension 1 in  $|\mathcal{L}_0|$ .

*Proof.* (i) Let  $I'$  be an irreducible component of  $I$  such that  $I'$  is dominant to  $R$  via  $\pi_2$ . If  $p \in R$  is a general point we know that  $\pi_2^{-1}(p)$  can be identified with  $\mathcal{L}_0(2, p)$ , and  $\mathcal{L}_0(2, p)$  is a projective space with dimension  $s$  independent on the general point  $p$ . This clearly implies that  $I'$  is unique.

(ii) Suppose the dominating component  $I'$  exists. With the same notation as above we have  $\dim(I') = \dim(R) + s = s + 2$ . On the other hand, by Remark 2.5,  $s \geq \dim(|\mathcal{L}_0|) - 3$ , hence  $\dim(I') \geq \dim(|\mathcal{L}_0|) - 1$ . By the hypotheses, the map  $\pi_1$ , restricted to  $I'$ , is generically finite onto the image and this image cannot be dense in  $|\mathcal{L}_0|$  by Bertini's theorem. Hence  $\dim(I') \leq \dim(|\mathcal{L}_0|) - 1$ , so the equality holds, and this implies that  $s = \dim(|\mathcal{L}_0|) - 3$ , as wanted.

(iii) Keep the same notation as above. The dimension of the fibre of  $\pi_2$  over a general point of  $R$  is  $r \geq \dim(|\mathcal{L}_0|) - 3 = \mathcal{L}_0(2, p) \geq 0$ . Moreover there is an open dense subset  $U$  of  $R$ , containing  $p$ , such that for all  $q \in U$  one has that  $\mathcal{L}_0(2, p)$  has dimension  $s = \dim(|\mathcal{L}_0|) - 3$ . Hence there is a component  $I'$  of  $I$  dominating  $R$  via  $\pi_2$ .  $\square$

We can now give the:

*Proof of Proposition 3.15.* We notice that by Corollary 3.11,  $[S_0]$  is a smooth point for  $\mathcal{H}^{\mathcal{X}|\mathbb{D}}$ . Part (i) follows by Corollary 3.9.

Let us prove part (ii). For this we go back to the notation introduced before the statement of Proposition 3.15. Consider then the intersection  $\bar{Z}_0$  of  $\bar{Z}$  with  $\bar{\pi}^{-1}(0) \cong |\mathcal{L}_0|$ , such that any of its irreducible components has codimension 1 in  $|\mathcal{L}_0|$ . By the hypotheses we made, if  $\bar{Z}'_0$  is any irreducible component of  $\bar{Z}_0$ , its general element does not contain  $R$ , hence it is a surface  $S'_0 \in |\mathcal{L}_0|$  that intersects  $R$  along a curve  $C'$ . By Proposition 3.2, the curve  $C'$  is singular.

**Claim 3.17.** *There is an irreducible component  $\bar{Z}'_0$  of  $\bar{Z}_0$ , such that for  $S'_0 \in \bar{Z}'_0$  general,  $S'_0$  intersects  $R$  in a curve  $C'$  that is singular at a general point  $p'$  of  $R$ . Moreover,  $S'_0$  is limit of reduced singular surfaces  $S_t \in |\mathcal{L}_t|$ .*

*Proof of the Claim 3.17.* This will be a consequence of the following fact that we are going to prove: given a general point  $p' \in R$ , there is some  $S'_0 \in \bar{Z}_0$  such that the curve  $C'$  cut out by  $S'_0$  on  $R$  is singular at  $p'$ . Indeed, given  $p' \in R$  general, take a smooth bisection  $\gamma'$  of  $\mathcal{X} \rightarrow \mathbb{D}$ , that passes through  $p'$ . As in §2.1, we can consider the family  $\mathcal{Y} \rightarrow \mathbb{D}$  obtained by desingularising the variety  $\mathcal{X}' \rightarrow \mathbb{D}$  gotten via 2-fold base change  $\nu_2 : \mathbb{D} \rightarrow \mathbb{D}$ . The variety  $\mathcal{Y} \rightarrow \mathbb{D}$  has a section  $\gamma$  that is mapped to  $\gamma'$  via the map  $\pi : \mathcal{Y} \rightarrow \mathcal{X}$ . We consider  $\pi^*(\mathcal{L})$ . Our assumption (3) implies that there are non-zero sections of  $\pi^*(\mathcal{L})$ , on  $\mathcal{Y} \setminus \mathcal{Y}_0$ , vanishing with multiplicity at least 2 along  $\gamma$ . The assertion is now a consequence of Theorem 2.2.  $\square$

By the hypothesis (b) and by Lemma 3.16(iii), the pair  $(S_0, p)$  belongs to the unique irreducible component  $I'$  of  $I$  dominating  $R$  via  $\pi_2$ , and  $I'$  has dimension equal to  $\dim(|\mathcal{L}_0|) - 1$ . Consider now the subset  $I'' \subseteq I$  of the pairs  $(S'_0, p')$  with  $S'_0 \in \bar{Z}'_0$ , where  $\bar{Z}'_0$  is as in Claim 3.17. We notice that  $I''$  also dominates  $R$  via  $\pi_2$ . So by Lemma 3.16(i),  $I''$  coincides with  $I'$ . This implies that  $\bar{Z}'_0 = \pi_1(I')$  hence  $S_0 \in \bar{Z}'_0$  and therefore the general surface  $S'_0 \in \bar{Z}'_0$  has a unique  $T_1$  singularity. By Lemma 3.3 the assertion (ii) follows.

To prove (iii) we remark first of all that (ii) implies that  $\mathcal{V}_1^{\mathcal{X}|\mathbb{D}}$  is non-empty and there is an irreducible component  $\mathcal{V}$  of  $\mathcal{V}_1^{\mathcal{X}|\mathbb{D}}$  that dominates  $\mathbb{D}$  and contains  $[S_0]$ . The general point in  $\mathcal{V}$  corresponds to a surface  $S_t$  with  $t \neq 0$ , with a unique node

at a general point  $p_t \in \mathcal{X}_t$ . Moreover, since  $[S_0]$  is a smooth point of  $\mathcal{H}$ , we have that  $[S_t]$  is a smooth point of  $\mathcal{H}$ , and by the hypothesis (c) and by semicontinuity, we have that  $h^1(S_t, N_{S_t|\mathcal{X}_t} \otimes I_{p_t}) = 0$ . This yields that  $\mathcal{V} \cap \mathcal{H}_t$  is smooth of dimension  $\dim(\mathcal{H}_t) - 1$ . Hence  $\mathcal{V}$  has dimension  $\dim(\mathcal{H}) - 1$ . To prove that  $\mathcal{V}$  is reduced, it suffices to prove that  $\mathcal{V}$  is smooth at  $[S_t]$ . To see this, consider the exact sequence

$$0 \rightarrow N'_{S_t|\mathcal{X}} \rightarrow N_{S_t|\mathcal{X}} \rightarrow T_{S_t}^1 \rightarrow 0$$

where  $T_{S_t}^1$  is supported on  $p_t$  with stalk  $\mathbb{C}$ , and  $H^0(S_t, N'_{S_t|\mathcal{X}})$  is the Zariski tangent space to  $\mathcal{V}$  at  $[S_t]$ . The map

$$H^0(S_t, N_{S_t|\mathcal{X}}) \rightarrow T_{S_t}^1 = \mathbb{C}$$

is surjective because  $S_t$  is smoothable inside  $\mathcal{H}$ , by the hypothesis (2) at the beginning of this section. Hence  $h^0(S_t, N'_{S_t|\mathcal{X}}) = h^0(S_t, N_{S_t|\mathcal{X}}) - 1 = \dim(\mathcal{H}) - 1$ , as wanted.  $\square$

We can now prove the main result of this section extending Proposition 3.15 to the case  $\delta > 1$ :

**Theorem 3.18.** *Set up as in Proposition 3.15. In particular we have the following condition: the subspace of sections of  $H^0(\mathcal{X}_0, \mathcal{L}_0)$  that vanish on  $R = A \cap B$  with  $A$  and  $B$  the irreducible components of  $\mathcal{X}_0$ , has codimension strictly larger than 1 in  $H^0(\mathcal{X}_0, \mathcal{L}_0)$ .*

Let  $S_0 = S_A \cup S_B \subset \mathcal{X}_0$  be a surface corresponding to a section of  $\mathcal{L}_0$ . We suppose that:

- (a)  $S_A$  and  $S_B$  have respectively  $\delta_A$  and  $\delta_B$  nodes  $p_{A,1}, \dots, p_{A,\delta_A}$  and  $p_{B,1}, \dots, p_{B,\delta_B}$  off  $R$ , are elsewhere smooth and intersect transversally along a curve  $C = S_A \cap S_B$ , except for  $\delta_R$  distinct points  $p_{R,1}, \dots, p_{R,\delta_R} \in C \subset R$  where  $S_0$  has singularities of type  $T_1$  and that the hypotheses of Lemma 3.5 hold;
- (b) the sublinear system  $\mathcal{L}_0(2, p_i)$  of  $|\mathcal{L}_0|$  of surfaces with at least a  $T_1$  singularity at  $p_i$  has codimension 3 in  $|\mathcal{L}_0|$ , for every  $1 \leq i \leq \delta$ ;
- (c) if  $\mathfrak{Z}$  is the 0-dimensional scheme of length  $\delta = \delta_A + \delta_B + \delta_R$  given by

$$\mathfrak{Z} = \sum_{i=1}^{\delta_A} p_{A,i} + \sum_{i=1}^{\delta_B} p_{B,i} + \sum_{i=1}^{\delta_R} p_{R,i}$$

then  $H^1(S_0, \mathcal{N}_{S_0|\mathcal{X}_0} \otimes I_{\mathfrak{Z}|\mathcal{X}_0}) = 0$  (where  $I_{\mathfrak{Z}|\mathcal{X}_0}$  is the ideal sheaf of the scheme  $\mathfrak{Z}$  in  $\mathcal{X}_0$ ).

Then:

- (i)  $S$  can be deformed to a  $\delta$ -nodal surface  $S_t \subset \mathcal{X}_t$ ;
- (ii) if  $\mathbb{T} \subseteq \mathbb{T}_{\delta_A, \delta_B, \delta_R}$  is the unique irreducible component containing  $[S]$ , then there exists an irreducible component  $\mathcal{V} \subset \mathcal{V}_\delta^{\mathcal{X}|\mathbb{D}}$  of dimension  $\dim(\mathcal{H}) - \delta$  whose central fibre  $\mathcal{V}_0$  contains  $\mathbb{T}$  as an irreducible component.

*Proof.* Again, by Corollary 3.11,  $[S_0]$  is a smooth point for  $\mathcal{H}^{\mathcal{X}|\mathbb{D}}$ . Moreover, by Corollary 3.13,  $\mathbb{T}_{\delta_A, \delta_B, \delta_R}$  is smooth at  $[S_0]$ .

We denote by  $\mathbb{T}$  the unique irreducible component of  $\mathbb{T}_{\delta_A, \delta_B, \delta_R}$  containing  $[S_0]$ , that is smooth at  $[S_0]$ .

Again by Corollary 3.13, in an analytic neighborhood of  $[S]$ ,  $\mathbb{T}$  consists of an analytic branch  $\mathcal{T}$  that is the transverse intersection of  $\delta$  smooth analytic branches of dimension  $h^0(S_0, \mathcal{N}_{S_0|\mathcal{X}_0}) - 1$ , each branch corresponding to the locus of deformations of  $S_0$  that are equisingular at a given point in  $\mathfrak{Z}$ , i.e., we have that

$$\mathcal{T} = \bigcap_{p \in \mathfrak{Z}} \mathcal{E}\mathcal{S}_{[S_0], p}^{\mathcal{X}_0}$$

The general element  $S'_0$  of  $\mathcal{ES}_{[S_0],p}^{\mathcal{X}_0}$  is a surface in  $\mathcal{X}_0$  that has a unique singularity analytically equivalent to the singularity of  $S_0$  at  $p$ , i.e., a node if  $p \in \mathfrak{Z}_A + \mathfrak{Z}_B$ , a  $T_1$  singularity otherwise.

By Proposition 3.14, for every  $p \in \mathfrak{Z}_A + \mathfrak{Z}_B$ ,  $\mathcal{ES}_{[S_0],p}^{\mathcal{X}_0}$  is contained in  $\mathcal{ES}_{[S_0],p}^{\mathcal{X}}$  as a subvariety of codimension 1, and  $\mathcal{ES}_{[S_0],p}^{\mathcal{X}}$  is an analytic branch of the Severi variety  $\mathcal{V}_1^{\mathcal{X}|\mathbb{D}}$ .

If  $p \in \mathfrak{Z}_R$ , then by the hypothesis (b), for general element  $S'_0$  of  $\mathcal{ES}_{[S_0],p}^{\mathcal{X}_0}$  the condition (b) of Proposition 3.15 holds. Then, by Proposition 3.15,  $\mathcal{ES}_{[S_0],p}^{\mathcal{X}_0}$  is contained, as a subvariety of codimension 1, in an analytic branch  $\mathcal{T}_p$  of  $\mathcal{V}_1^{\mathcal{X}|\mathbb{D}}$  having codimension 1 in  $\mathcal{H}$ , which is smooth at the general point corresponding to a 1-nodal surface.

Now, the intersection

$$\mathcal{T}' = \bigcap_{p \in \mathfrak{Z}_R} \mathcal{T}_p \cap \bigcap_{p \in \mathfrak{Z}_A + \mathfrak{Z}_B} \mathcal{ES}_{[S_0],p}^{\mathcal{X}}$$

has codimension at most  $\delta$  in  $\mathcal{H}$  and it contains the smooth analytic branch  $\mathcal{T}$  of  $\mathbb{T}$ , which has codimension  $\delta + 1$  in  $\mathcal{H}$ . The general element of  $\mathcal{T}'$  corresponds to a surface  $\tilde{S}$ , not contained in  $\mathcal{X}_0$ , with at least  $\delta$  singularities, precisely  $\delta_A$  (resp.  $\delta_B$ ) singularities in neighborhoods of the nodes  $p \in \mathfrak{Z}_A$  (resp.  $p \in \mathfrak{Z}_B$ ) and  $\delta_R$  singularities in neighborhoods of the  $T_1$  singularities  $p \in \mathfrak{Z}_R$ . Taking into account Lemma 3.3, we deduce that  $\tilde{S}$  has  $\delta$  nodes and no further singularities. This proves (i).

If  $[\tilde{S}] \subset \mathcal{X}_t$  and it has nodes at  $\tilde{q}_1, \dots, \tilde{q}_\delta$ , by semicontinuity, we have that  $H^0(\tilde{S}, \mathcal{N}_{\tilde{S}|\mathcal{X}_t} \otimes I_{\{\tilde{q}_1, \dots, \tilde{q}_\delta\}|\mathcal{X}_t})$  has dimension  $\dim(\mathcal{H}_t) - \delta$ . Thus  $\mathcal{T}'$  is an analytic branch containing the point  $[\tilde{S}]$  in an irreducible component  $\mathcal{V} \subset \mathcal{V}_\delta^{\mathcal{X}|\mathbb{D}}$  of dimension  $\dim(\mathcal{H}) - \delta$  whose central fibre  $\mathcal{V}_0$  contains  $\mathbb{T}$  as an irreducible component. This proves (ii).  $\square$

#### 4. APPLICATIONS

**4.1. Severi varieties.** Let  $X$  be a smooth irreducible projective complex threefold. Let  $L$  be a line bundle on  $X$  such that the general surface in the linear system  $|L|$  is smooth and irreducible. We denote by  $V_\delta^{X,|L|}$  the *Severi variety*, that is the locally closed subscheme in  $|L|$  parametrizing surfaces  $S$  in  $|L|$  which are reduced and with only  $\delta$  nodes as singularities. If  $[S] \in V_\delta^{X,|L|}$ , then the Zariski tangent space to  $V_\delta^{X,|L|}$  at  $[S]$  coincides with

$$T_{[S]}(V_\delta^{X,|L|}) \simeq H^0(S, \mathcal{O}_S(L) \otimes I_{N|S}),$$

where  $N$  is the reduced scheme of nodes of  $S$ . In particular,  $\dim(V_\delta^{X,|L|}) \leq h^0(S, \mathcal{O}_S(L) \otimes I_{N|S})$ . Moreover, by standard deformation theory,  $H^1(S, \mathcal{O}_S(L) \otimes I_{N|S})$  is an obstruction space for  $\mathcal{O}_{V_\delta^{X,|L|},[S]}$  and thus

$$h^0(S, \mathcal{O}_S(L) \otimes I_{N|S}) - h^1(S, \mathcal{O}_S(L) \otimes I_{N|S}) \leq \dim(V_\delta^{X,|L|}) \leq h^0(S, \mathcal{O}_S(L) \otimes I_{N|S}).$$

If  $h^1(S, \mathcal{O}_S(L) \otimes I_{N|S}) = 0$ , then  $V_\delta^{X,|L|}$  is smooth at  $[S]$  of dimension

$$h^0(S, \mathcal{O}_S(L) \otimes I_{N|S}) = \dim(|L|) - \delta.$$

In this case one says that  $[S]$  is a *regular point* of the  $\dim(V_\delta^{X,|L|})$ . An irreducible component  $V$  of  $\dim(V_\delta^{X,|L|})$  is said to be *regular* if it is regular at its general point.

**Remark 4.1.** Suppose  $V$  is a regular irreducible component of  $\dim(V_\delta^{X,|L|})$ . By standard deformation theory already used in Section 3, the nodes of the surface

corresponding to any smooth point in  $V$  can be independently smoothed. This implies that there are regular components of  $\dim(V_{\delta'}^{X,|L|})$  for any  $\delta' < \delta$ .

One can consider the following two questions.

**Problem 4.2.** *Given  $X$  and  $L$  as above, which is the maximal value of  $\delta$  such that the Severi variety  $V_{\delta}^{X,|L|}$  is non empty?*

**Problem 4.3.** *Given  $X$  and  $L$  as above, which is the maximal value of  $\delta$  such that the Severi variety  $V_{\delta}^{X,|L|}$  has a regular component?*

As for Problem 4.2, this is a classical and difficult question, for which there are several contributions, too many to be quoted here, probably the most efficient one is given by the Miyaoka's bound [15, Formulae (2) and (8)]. In particular the problem has been completely solved for  $X = \mathbb{P}^3$  and  $L = \mathcal{O}_{\mathbb{P}^3}(d)$  with  $d \leq 6$  (see, e.g., [14] and references therein). However, in this section we will not consider Problem 4.2 but we will give some contribution to Problem 4.3.

**Remark 4.4.** One could be tempted to believe that the maximal  $\delta$  for which the Severi variety is non-empty is bounded above by the dimension of  $|L|$ . This is not true. In fact there are classical examples, for  $X = \mathbb{P}^3$  and  $L = \mathcal{O}_{\mathbb{P}^3}(d)$  for suitable  $d$ , for which  $V_{\delta}^{X,|L|}$  is non-empty and  $\delta$  is greater than the dimension of  $|L|$  (cf. [1, [19]). In these cases every component of the Severi variety is not regular.

**Remark 4.5.** Referring to Problem 4.3, it is rather natural to conjecture that the  $\delta$  which answer the question should be bounded below by  $\delta_0 = \lfloor \frac{\dim(|L|)}{4} \rfloor$ . The reason for such a conjecture is the following: choose  $p_1, \dots, p_{\delta_0}$  general points on  $X$ . Since a double point imposes at most 4 conditions to  $|L|$ , certainly there are surfaces which are singular at every  $p_i$ . If the general such surface has only nodes at  $p_1, \dots, p_{\delta_0}$  and no other singularities then it belongs to a regular component of the Severi variety. However this heuristic argument is very difficult to be made rigorous in general.

**4.2. The case of  $\mathbb{P}^3$ .** In this section we give a contribution to Problem 4.3, in the case  $X = \mathbb{P}^3$  and  $L = \mathcal{O}_{\mathbb{P}^3}(d)$ . More precisely we will prove the following:

**Theorem 4.6.** *There is an irreducible, regular component of  $V_{\delta}^{\mathbb{P}^3, |\mathcal{O}_{\mathbb{P}^3}(d)|}$ , for any  $\delta \leq \binom{d-1}{2}$ .*

*Proof.* In view of Remark 4.1, it is sufficient to consider only the case  $\delta = \binom{d-1}{2}$ .

Let  $\mathcal{X}' = \mathbb{P}^3 \times \mathbb{D} \rightarrow \mathbb{D}$  be a trivial family. Let us consider  $\mathcal{X} \rightarrow \mathbb{D}$  the blow up of a point  $q$  in the central fibre  $\mathbb{P}^3$  over  $0 \in \mathbb{D}$ . Let  $\mathcal{X} \rightarrow \mathbb{D}$  be the new family. The fibre over  $t \in \mathbb{D} \setminus \{0\}$  of this family is  $\mathcal{X}_t \cong \mathbb{P}^3$ . The central fibre  $\mathcal{X}_0$  consists of two components  $A \cup B$ , where  $f : A \rightarrow \mathbb{P}^3$  is the blow-up of  $\mathbb{P}^3$  at  $q$ , whereas  $B \cong \mathbb{P}^3$  is the exceptional divisor in  $\mathcal{X}$ , and  $A \cap B = R \cong \mathbb{P}^2$  is the exceptional divisor in  $A$  and a plane in  $B$ .

On  $\mathcal{X}'$  there is a line bundle  $\mathcal{L}'$ , which is the pull-back via the first projection, of  $\mathcal{O}_{\mathbb{P}^3}(d)$ . We pull this back to  $\mathcal{X}$  and denote it  $\mathcal{L}$ . Now we consider on  $\mathcal{X}$  the line bundle  $\mathcal{L} \otimes \mathcal{O}_{\mathcal{X}}((1-d)B)$ . Its restriction to the general fibre  $\mathcal{X}_t$  is given by  $(\mathcal{L} \otimes \mathcal{O}_{\mathcal{X}}((1-d)B))|_{\mathcal{X}_t} \simeq \mathcal{O}_{\mathbb{P}^3}(d)$ . As for the restriction of  $\mathcal{L} \otimes \mathcal{O}_{\mathcal{X}}((1-d)B)$  to  $\mathcal{X}_0$ , we observe that  $(\mathcal{L} \otimes \mathcal{O}_{\mathcal{X}}((1-d)B))|_A \simeq \mathcal{O}_A(d) \otimes \mathcal{O}_A(-(d-1)R)$ , where  $\mathcal{O}_A(d) \simeq f^*(\mathcal{O}_{\mathbb{P}^3}(d))$ ; whereas  $(\mathcal{L} \otimes \mathcal{O}_{\mathcal{X}}((1-d)B))|_B \simeq \mathcal{O}_{\mathbb{P}^3}(d-1)$  and, finally, the restriction of  $\mathcal{L} \otimes \mathcal{O}_{\mathcal{X}}((1-d)B)$  to  $R$  is  $\mathcal{O}_{\mathbb{P}^2}(d-1)$ . One easily checks that the line bundle  $\mathcal{L} \otimes \mathcal{O}_{\mathcal{X}}((1-d)B)$  verifies the hypotheses (1), (2) and (3) at the beginning of Section 3.2.3.

We now consider on  $R$  a curve  $C$  which consists of the union of  $d-1$  general lines. Its has  $\delta = \binom{d-1}{2}$  nodes as singularities. By standard application of Bertini's



theorem, there exists a smooth surface  $S_B$  in  $B$  of degree  $d-1$  cutting out on  $R$  the curve  $C$ . Similarly there exists a smooth surface  $S_A \in |\mathcal{O}_A(d) \otimes \mathcal{O}_A(-(d-1)R)|$  restricting to  $C$  on  $R$ . To see this let  $(x, y, z)$  be an affine coordinates system on  $\mathbb{P}^3$  centered at  $q$ . If  $\phi_1(x, y, z) = 0$  is the equation of  $C$  in the plane at infinity, and  $\phi_2(x, y, z)$  is a general homogeneous polynomial of degree  $d$  in  $(x, y, z)$ , then the projective closure  $S_B$  of the degree  $d$  affine surface with equation  $\phi_1(x, y, z) + \phi_2(x, y, z) = 0$  has a point of multiplicity  $d-1$  at  $q$  and no other singularities, and its minimal resolution obtained by blowing up  $q$  is the required surface.

Now  $S_0 = S_A \cup S_B$  is a Cartier divisor in  $\mathcal{X}_0$  belonging to the linear system  $|\mathcal{L} \otimes \mathcal{O}_{\mathcal{X}}((1-d)B)|$ . Moreover  $S_0$  verifies all hypotheses of Theorem 3.18. In particular, if  $\mathfrak{Z}$  is the reduced scheme of the nodes of  $C$ , then  $\mathfrak{Z}$  imposes independent conditions to  $|\mathcal{L} \otimes \mathcal{O}_{\mathcal{X}}((1-d)B)|_R = |\mathcal{O}_{\mathbb{P}^2}(d-1)|$ , and therefore to  $|\mathcal{L} \otimes \mathcal{O}_{\mathcal{X}}((1-d)B)|$ , because the Severi varieties of nodal curves in the plane are well known to be regular. By applying Theorem 3.18, one may deform  $S_0$  to a surface  $S_t \subset \mathcal{X}_t$  with  $\delta$  nodes and no further singularities, which are deformations of the  $\delta$  singularities of type  $T_1$  of  $S_0$ . Finally the nodes of  $S_t$  impose independent conditions to surfaces of degree  $d$  on  $\mathcal{X}_t \simeq \mathbb{P}^3$ . Hence  $[S_t] \in V_{\delta}^{\mathbb{P}^3, |\mathcal{O}_{\mathbb{P}^3}(d)|}$  belongs to a regular component of the Severi variety.  $\square$

**Remark 4.7.** Taking into account Remark 4.5, we believe that the previous results is far from being sharp, not even asymptotically. Indeed we may expect that the bound on  $\delta$  for the existence of regular components of the Severi variety of nodal surfaces of degree  $d$  in  $\mathbb{P}^3$  could asymptotically go as  $\delta \sim \frac{d^3}{24}$ . Moreover our results could in principle be improved by imposing to  $S_0$  nodes off  $R$ , but we do not dwell on this here.

**Remark 4.8.** The known results about Problem 4.3 are very few. For example, in [13] one proves that, if  $V_{\delta}^{\mathbb{P}^3, |\mathcal{O}_{\mathbb{P}^3}(d)|}$  is non-empty, then every component of it is regular for  $d \leq 7$  and for  $d \geq 8$  and  $\delta \leq 4d - 5$ , and this last bound is sharp (the case  $d \leq 7$  was already proved in [8]). Nonemptiness results for  $d \leq 7$  are also well known (see, e.g., [14, p. 120]). In particular, our Theorem 4.6 is, at the best of our knowledge, new as soon as  $d \geq 8$ .

**4.3. Complete intersections in  $\mathbb{P}^4$ .** In this Section we want to provide a partial answer to Problem 4.3 in the case of complete intersections in  $\mathbb{P}^4$ .

Let  $X$  be a general hypersurface of degree  $h \geq 2$  in  $\mathbb{P}^4$ . We consider on  $X$  the linear system  $|\mathcal{O}_X(d)|$ . Our aim is to construct regular components of  $V_{\delta}^{X, |\mathcal{O}_X(d)|}$  with suitable  $\delta$ .

**Theorem 4.9.** *Let  $d \geq h - 1$  be an integer. There are regular components of  $V_{\delta}^{X, |\mathcal{O}_X(d)|}$  for*

$$\delta \leq \binom{d+3}{3} - \binom{d-h+1}{3} - 1.$$

*Proof.* As usual, to prove the theorem, it suffices to do the case

$$\delta = \binom{d+3}{3} - \binom{d-h+1}{3} - 1.$$

Let  $Y$  be a general hypersurface of  $\mathbb{P}^4$  of degree  $h-1$  and  $H$  be a general hyperplane, cutting  $Y$  along a surface  $R$ , which is a general surface of degree  $h-1$  in  $H \simeq \mathbb{P}^3$ . Let  $X$  be a general hypersurface of degree  $h$  and let us consider the pencil generated by  $X$  and  $Y \cup H$ . Specifically, if  $X$  has equation  $f = 0$ ,  $Y$  has equation  $g = 0$  and  $H$  has equation  $\ell = 0$ , we will consider the hypersurface  $\mathcal{X}''$  in  $\mathbb{P}^4 \times \mathbb{A}^1$ , with equation  $\{tf + g\ell = 0, \text{ with } t \in \mathbb{A}^1\}$ . Via the second projection  $\mathcal{X}'' \rightarrow \mathbb{A}^1$ , this becomes a flat family of 3-folds, with smooth general fibre  $\mathcal{X}_t''$ , corresponding to a general hypersurface of degree  $h$  in  $\mathbb{P}^4$ , and whose fibre over 0 is

$\mathcal{X}_0'' = Y \cup H \subset \mathbb{P}^4$ . We are interested to the singularities of  $\mathcal{X}''$  in a neighborhood of the central fibre, i.e., we are interested in what happens if  $t$  belongs to a disc  $\mathbb{D}$ , centered at the origin. Thus we consider the family

$$\mathcal{X}' = \{tf + g\ell = 0, \text{ with } t \in \mathbb{D}\} \rightarrow \mathbb{D}.$$

It is immediate to see that the singular locus of  $\mathcal{X}'$  coincides with the curve  $D : t = f = g = \ell = 0 \subset \mathcal{X}'_0$ , which is isomorphic to a smooth complete intersection curve of type  $(1, h-1, h)$  in  $\mathbb{P}^4$  cut out on  $R = Y \cap H$  by  $X$ . Moreover,  $\mathcal{X}'$  has double points along  $D$  with tangent cone a quadric of rank 4. We resolve these singularities by blowing-up  $\mathcal{X}'$  along  $D$ . One obtains a new family  $\tilde{\mathcal{X}} \rightarrow \mathbb{D}$  with the same general fibre as  $\mathcal{X}' \rightarrow \mathbb{D}$  and whose central fibre consists of three components  $\tilde{Y}$  and  $\tilde{H}$ , the blow-ups of  $Y$  and  $H$  along  $D$  and the exceptional divisor  $\tilde{\Theta}$  that is a  $\mathbb{P}^1 \times \mathbb{P}^1$  bundle over  $D$ . Now we can contract  $\tilde{\Theta}$  by contracting one of the two rulings of the  $\mathbb{P}^1 \times \mathbb{P}^1$  bundle. We choose to do this in the direction of  $Y$ . We obtain a new family of 3-folds  $\mathcal{X} \rightarrow \mathbb{D}$ , with  $\mathcal{X}$  smooth, with fiber  $\mathcal{X}_t = \mathcal{X}'_t$  over  $t \neq 0$ , and whose central fiber  $\mathcal{X}_0 = A \cup B$ , where now  $B = H \simeq \mathbb{P}^3$  and  $A = \text{Bl}_D(Y)$  is the blowing-up of  $Y$  along  $D$  and  $A$  and  $B$  intersect transversally along a surface isomorphic to  $R$ , which we still denote by  $R = A \cap B$ . The exceptional divisor  $\Theta$  in  $A = \text{Bl}_D(Y)$  is a  $\mathbb{P}^1$ -bundle on  $D \subset R$ , intersecting  $R$  along  $D$ . In particular  $\Theta \simeq \mathbb{P}(\mathcal{N}_{D|Y})$ .

Notice that one has a natural morphism  $\tilde{\mathcal{X}} \rightarrow \mathbb{P}^4$ . This factors through a morphism  $\phi : \mathcal{X} \rightarrow \mathbb{P}^4$ . The action of  $\phi$  on  $\mathcal{X}_0$  is as follows: it maps  $B$  isomorphically to  $H$  and it maps  $A$  to  $Y$  by contracting the exceptional divisor  $\Theta$ . Let us now set  $\mathcal{L}_d = \phi^*(\mathcal{O}_{\mathbb{P}^4}(d))$  and assume that  $d \geq h-1$ .

Recall that  $R$  is a general surface of degree  $h-1$  in  $\mathbb{P}^3$ , with  $h \geq 2$ . By [4],  $V_\delta^{R, |\mathcal{O}_R(d)|}$  is non-empty and contains a regular component  $V$  for

$$\delta = \dim(|\mathcal{O}_R(d)|) = \binom{d+3}{3} - \binom{d-h+1}{3} - 1.$$

So we can choose a general curve  $C$  in  $V$ , that is a complete intersection of type  $(h-1, d)$  on  $R$  with  $\delta$  nodes. Using Bertini's theorem, we can assume that there is a divisor  $S_0 \in \mathcal{L}_{d|\mathcal{X}_0}$  that cuts out  $C$  on  $R$  and  $S_0 = S_A \cup S_B$  (the notation is obvious), with  $S_A$  and  $S_B$  smooth.

Now  $S_0$  verifies all hypotheses of Theorem 3.18. In particular, if  $\mathfrak{Z}$  is the reduced scheme of nodes of  $C$ , then  $\mathfrak{Z}$  imposes independent conditions to  $\mathcal{L}_{d|\mathcal{X}_0}$ , because the component  $V$  of the Severi variety is regular. By applying Theorem 3.18, one may deform  $S_0$  to a surface  $S_t \subset \mathcal{X}_t$  with  $\delta$  nodes and no further singularities, which are deformations of the  $\delta$  singularities of type  $T_1$  of  $S_0$ . Finally the nodes of  $S_t$  impose independent conditions to surfaces in  $|\mathcal{O}_{X_t}(d)|$ . Hence  $[S_t] \in V_\delta^{\mathcal{X}_t, |\mathcal{O}_{X_t}(d)|}$  belongs to a regular component of the Severi variety, as wanted.  $\square$

## REFERENCES

- [1] A. Beauville: *Sur le nombre maximum de points doubles d'une surface dans  $\mathbb{P}^3$  ( $\mu(5) = 31$ )* Journées de Géométrie Algébrique d'Angers, Juillet 1979/Algebraic Geometry, Angers (1979), 207–215.
- [2] L. Caporaso, J. Harris: *Parameter spaces for curves on surfaces and enumeration of rational curves*, Compositio Math., **113**, (1998), 155–208.
- [3] X. Chen: *Rational curves on K3 surfaces*, J. Alg. Geom., **8**, (1999), 245–278.
- [4] L. Chiantini, C. Ciliberto: *On the Severi varieties of surfaces in  $\mathbb{P}^3$* , J. of Algebraic Geom., **8** (1999), 67–83.
- [5] C. Ciliberto, Th. Dedieu, C. Galati, A. Knutsen: *Severi varieties on blow-ups of the symmetric square of an elliptic curve*, Math. Nachr., **296** (2), (2023), 574–587.
- [6] C. Ciliberto, Th. Dedieu, C. Galati, A. Knutsen: *Nonemptiness of Severi varieties on Enriques surfaces*, Forum of Mathematics, Sigma, **11** (e52), (2023), 1–32.
- [7] V. Di Gennaro, D. Franco, *Intersection cohomology and Severi varieties*, in “The art of doing Algebraic Geometry”, Trends in Math., Springer Verlag, 2023.

- [8] A. Dimca: *On the syzygies and Hodge theory of nodal hypersurfaces*, Ann. Univer. Ferrara, **63** (1), (2017), 87–101.
- [9] C. Galati: *Degenerating curves and surfaces: first results*, Rend. Circ. Mat. Palermo, **58** (2009), 211–243.
- [10] C. Galati, A. Knutsen: *On the existence of curves with  $A_k$ -singularities on  $K3$  surfaces*, Math. Res. Lett., **21**, (2014), 1069–1109.
- [11] G.-M. Greuel, U. Karras: *Families of varieties with prescribed singularities*, Compositio Math., **69** (1), (1989), 83–110.
- [12] Ph. Griffiths, J. Harris: *Principles of Algebraic Geometry*, A. Wiley-Interscience Series of Texts, New York, 1978.
- [13] R. Kloosterman: *Nodal surfaces with obstructed deformations*, Geom. Dedicata, **190**, (2017), 143–150.
- [14] O. Labs: *Hypersurfaces with many singularities*, (2006), <https://openscience.ub.uni-mainz.de/bitstream/20.500.12030/1939/1/885.pdf>
- [15] Y. Miyaoka: *The Maximal Number of Quotient Singularities on Surfaces with Given Numerical Invariants*, Math. Ann., **268**, (1984), 159–171.
- [16] R. Thomas, *Nodes and the Hodge conjecture*, J. Alg. Geom., **14**, (2005), 177–185.
- [17] Z. Ran: *Families of plane curves and their limits: Enriques’ conjecture and beyond*, Annals of Math., **130** (1), (1989), 121–157.
- [18] B. Segre: *On limits of algebraic varieties, in particular of their intersections and tangential forms*, Proc. London Math. Soc., **47** (2), (1942), 351–403.
- [19] B. Segre: *Sul massimo numero di nodi delle superficie di dato ordine* Boll. Un. Mat. Ital. **2** (3) (1947), 204–212.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA “TOR VERGATA”, VIA DELLA RICERCA SCIENTIFICA, 00133 ROMA, ITALY.

*Email address:* `cilibert@mat.uniroma2.it`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DELLA CALABRIA, VIA P. BUCCI, CUBO 31B, 87036 ARCAVACATA DI RENDE (CS), ITALY.

*Email address:* `concettina.galati@unical.it`