

INTERPOLATED TIME-HÖLDER REGULARITY OF SOLUTIONS OF FULLY NONLINEAR PARABOLIC EQUATIONS

ALESSANDRO GOFFI

ABSTRACT. We show by the maximum principle parabolic interior Schauder estimates for a special class of fully nonlinear parabolic Isaacs equations, providing an Evans-Krylov result for the model equation $\min\{\inf_{\beta} L_{\beta}u, \sup_{\gamma} L_{\gamma}u\} - \partial_t u = 0$, where L_{β}, L_{γ} are linear operators with possibly variable Hölder coefficients. Along the way, we provide a short survey on the regularity theory for fully nonlinear parabolic equations of second order.

1. INTRODUCTION

In this note we provide a maximum principle approach to study interior maximal C^{α} smoothness (i.e. local $C^{2,\alpha}$ regularity with respect to the parabolic distance) of solutions to the following fully nonlinear parabolic model

$$(1) \quad F(D^2u) - \partial_t u = 0 \text{ in } Q_1 := B_1 \times (-1, 0] \subseteq \mathbb{R}^{n+1},$$

where we assume

$$(2) \quad \begin{cases} F(M) = \min\{F^{\cap}(M), F^{\cup}(M)\}, \quad \forall M \in \mathcal{S}_n, \\ F(0) = 0, \quad F^{\cap}, F^{\cup} \text{ are uniformly elliptic,} \\ F^{\cap} \text{ is concave in } M, \quad F^{\cup} \text{ is convex in } M. \end{cases}$$

The uniform ellipticity for an operator $G = G(M) : \text{Sym}_n \rightarrow \mathbb{R}$, Sym_n being the space of $n \times n$ symmetric matrices, reads as

$$\lambda \|N\| \leq G(M + N) - G(M) \leq \Lambda \|N\|, \quad \forall M \in \text{Sym}_n, \quad N \geq 0, \quad \lambda \leq \Lambda, \quad \|N\| = \sup_{|x|=1} |Nx|.$$

Date: June 27, 2024.

2020 Mathematics Subject Classification. 35B65.

Key words and phrases. Evans-Krylov theorem, Fully nonlinear parabolic equations, Isaacs equations.

The author is member of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). He was partially supported by the INdAM-GNAMPA project 2024, the King Abdullah University of Science and Technology (KAUST) project CRG2021-4674 “Mean-Field Games: models, theory and computational aspects” and by the project funded by the European Union – NextGenerationEU under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.1 - Call PRIN 2022 No. 104 of February 2, 2022 of Italian Ministry of University and Research; Project 2022W58BJ5 (subject area: PE - Physical Sciences and Engineering) “PDEs and optimal control methods in mean field games, population dynamics and multi-agent models” .

Note that the uniform ellipticity of F^\cap, F^\cup implies that F itself is uniformly elliptic. In this case we will call (1) uniformly parabolic. The result applies for instance to the evolution PDE

$$(3) \quad \min \left\{ \inf_{a \in \mathcal{A}} L_a u, \sup_{b \in \mathcal{B}} L_b u \right\} - \partial_t u = 0,$$

where \mathcal{A}, \mathcal{B} are arbitrary sets and L_a, L_b are linear operators of the form $L_k u = a_{ij}^k \partial_{ij} u + c_k$, $c_k = L_k 0 \in \mathbb{R}$, a_{ij}^k with constant coefficients. This class can be written in the Isaacs form $\inf_{\sigma \in \Sigma} \sup_{\theta \in \Theta} L_{\sigma\theta} u - \partial_t u = 0$, where the convex operators $(\sup_{\theta \in \Theta} L_{\sigma\theta} u)_{\sigma \in \Sigma}$ are all linear except for at most one σ . A prototype equation is the flow driven by the 3-operator

$$F_3(D^2u) - \partial_t u = \min\{L_1 u, \max\{L_2 u, L_3 u\}\} - \partial_t u,$$

where L_1, L_2, L_3 are linear uniformly elliptic operators with constant coefficients. This provides a time-dependent counterpart of a result by X. Cabré and L. Caffarelli [4] and it can be regarded as an Evans-Krylov type result for a (time-dependent) nonconvex/nonconcave equation. We recall that by the Evans-Krylov property we mean the boost of regularity

$$C^{2,1} \rightarrow C^{2+\alpha, 1+\alpha/2},$$

see Theorem 6.1 of [7] for the elliptic result.

The outline of the proof of the maximal parabolic C^α regularity under the assumption (2) is the following: one first proves that for the homogeneous equation $F(D^2u) - \partial_t u = 0$ we have

$$\partial_t u, Du \in C^{\alpha, \alpha/2}$$

for some small universal $\alpha \in (0, 1)$, by the Krylov-Safonov theory [22] (here $C_t^{\alpha/2}$ denotes the Hölder continuity in time, uniformly in space). Indeed, the first derivative (or its incremental quotient), both in time and space, satisfies a linear nondivergence form equation, since the equation is invariant in space-time. This step does not require any convexity or concavity assumptions on F , and it holds for continuous viscosity solutions of any uniformly parabolic equation, see Theorem 3.4. Then, for each time slice, which we freeze and do not display, we write

$$F(D^2u) = g$$

where $g = \partial_t u \in C^{\alpha, \alpha/2}$. This implies by the stationary (space) regularity of [4]

$$u \in C_x^{2, \alpha}$$

at each time layer. To reach parabolic $C^{2+\alpha, 1+\alpha/2}$ regularity of the solution to the parabolic equation (and in particular $\alpha/2$ time-regularity of second derivatives) we exploit a maximum principle method. More precisely, exploiting the information $D^2u \in C_x^\alpha$, we improve the time-regularity of the gradient showing that

$$Du \in C_t^{\frac{1+\alpha}{2}}$$

via the comparison principle. We then exploit an interpolation argument between $Du \in C_t^{(1+\alpha)/2}$ and $D^2u \in C_x^\alpha$ to conclude that $D^2u \in C_t^{\alpha/2}$, proving thus the claimed $C^{2+\alpha, 1+\alpha/2}$

regularity. Note that a direct interpolation with the sole information $Du \in C_t^{\alpha/2}$ available from the Krylov-Safonov theory provides a worse Hölder regularity exponent than the expected regularity in $C^{2+\alpha, 1+\alpha/2}$, which is however enough in general to reach classical smoothness.

In the case of the heat equation, a maximum principle approach was used to prove the time-Hölder regularity of the Hessian in [17] via an argument of A. Brandt [2].

Notably, our method provides a new proof of the parabolic $C^{2,\alpha}$ regularity for fully nonlinear, concave, uniformly parabolic equations using the stationary result and the maximum principle, and can be generalized to any parabolic equation that admits $C^{2,\alpha}$ interior elliptic estimates, see the end of Section 3. A comparison with other approaches to study the parabolic regularity of fully nonlinear equations is in Remark 3.7.

The study of these estimates are motivated by the challenging open question of finding some special structures of F guaranteeing classical smoothness of solutions when the parabolic dimension $n + 1 \geq 4$, cf. the introduction of [15] and Section 2 for references on the subject.

Following [4], we mention that these interior estimates could be used to prove an existence result for the Dirichlet problem by a refinement of the method of continuity based on weighted interior bounds. This approach was first introduced by J. H. Michael [27] for linear stationary PDEs and extended to linear parabolic equations in [31]. We do not pursue this direction here, but just recall that the application of the classical method of continuity requires estimates up to the boundary along with a smooth operator, namely F must be at least of class C^1 . Moreover, we emphasize that the process of smoothing fully nonlinear operators is known only for concave/convex equations, and hence one needs a different approach for other classes of non-smooth operators.

Plan of the paper. Section 2 provides a brief survey on the known literature of the $C^{2,\alpha}$ solvability of fully nonlinear parabolic equations. Section 3 is devoted to the application of the maximum principle approach to obtain $C^{2+\alpha, 1+\alpha/2}$ regularity for a time-dependent counterpart of an Isaacs model introduced by X. Cabré and L. Caffarelli.

Notation. We denote by $B_r(x)$ the ball of center x and radius r . When $x_0 = 0$ we simply write B_r .

$Q_r(x, t)$ is the parabolic cylinder $B_r(x) \times (t - r^2, t)$, and we write Q_r when $(x, t) = (0, 0)$. We denote, given an open set Ω , by $\partial_{\text{par}}(\Omega \times (a, b))$ the parabolic boundary of a set.

We denote by $|u|_{0;\Omega}$ the sup-norm of u , i.e. $\|u\|_{C(\Omega)}$ (both in the elliptic and the parabolic case).

Let $Q \subset \Omega \times (0, T)$ and $\alpha \in (0, 1)$. We denote by $d((x, t), (y, s)) = |x - y| + |t - s|^{\frac{1}{2}}$ the standard parabolic distance and

- $C^{\alpha,\alpha/2}(Q)$, $\alpha \in (0, 1]$, the space of those $u : Q \rightarrow \mathbb{R}$ such that

$$\|u\|_{C^{\alpha,\alpha/2}(Q)} := |u|_{0;Q} + [u]_{C^{\alpha,\alpha/2}(Q)} = |u|_{0;Q} + \sup_{\substack{(x,t),(y,s) \in Q, \\ (x,t) \neq (y,s)}} \frac{|u(x,t) - u(y,s)|}{d^\alpha((x,t), (y,s))};$$

- $C^{1+\alpha,(1+\alpha)/2}(Q)$ the space of functions u whose spatial gradient exists classically and equipped with the norm

$$\|u\|_{C^{1+\alpha,(1+\alpha)/2}(Q)} := |u|_{0;Q} + |Du|_{0;Q} + \sup_{\substack{(x,t),(y,s) \in Q, \\ (x,t) \neq (y,s)}} \frac{|u(x,t) - u(y,s)|}{d^{1+\alpha}((x,t), (y,s))}.$$

In particular, any $u \in C^{1+\alpha,(1+\alpha)/2}(Q)$ is such that each component of Du belongs to $C^{\alpha,\alpha/2}(Q)$ and u is Hölder continuous with exponent $(1+\alpha)/2$ in the time variable;

- $C^{2+\alpha,1+\alpha/2}(Q)$ the space of functions u such that

$$\|u\|_{C^{2+\alpha,1+\alpha/2}(Q)} := |u|_{0;Q} + \sum_{i=1}^n \|u_{x_i}\|_{C^{1+\alpha,(1+\alpha)/2}(Q)} + \|\partial_t u\|_{C^{\alpha,\alpha/2}(Q)}$$

This is equivalent to say that D^2u belongs to $C^{\alpha,\alpha/2}(Q)$ and $\partial_t u$ belongs to $C^{\alpha,\alpha/2}(Q)$: it is a consequence of Remark 8.8.7 in [20]. In this case we can consider the space $C^{2+\alpha,1+\alpha/2}(Q)$ equipped with the norm

$$\|u\|_{C^{2+\alpha,1+\alpha/2}(Q)} := |u|_{0;Q} + |Du|_{0;Q} + \|D^2u\|_{C^{\alpha,\alpha/2}(Q)} + \|\partial_t u\|_{C^{\alpha,\alpha/2}(Q)}.$$

For more properties on these spaces we refer to [20]. Furthermore, we denote by

$$\|u\|_{C^{2,1}(Q)} := \sum_{2i+j \leq 2} |\partial_t^i D_x^j u|_{0;Q}.$$

We will also use the equivalence (see e.g. p. 120 of [20]) between the Hölder seminorm $[u]_{C^{\alpha,\alpha/2}(Q_1)}$ previously defined and the seminorm

$$[u]'_{C^{\alpha,\beta}(Q_1)} = \sup_{\substack{x,y \in B_1, \\ x \neq y}} \frac{|u(x,t) - u(y,t)|}{|x-y|^\alpha} + \sup_{\substack{t,s \in (-1,0), \\ t \neq s}} \frac{|u(x,t) - u(x,s)|}{|t-s|^{\frac{\alpha}{2}}}.$$

2. A QUICK SURVEY ON THE (HIGHER) REGULARITY THEORY FOR FULLY NONLINEAR PARABOLIC EQUATIONS

In this section we survey on regularity results at the level of Hölder spaces for fully nonlinear parabolic equations, focusing on those of the form

$$(4) \quad F(D^2u) - \partial_t u = 0 \text{ in } B_1 \times (-1, 0] \subset \mathbb{R}^{n+1}.$$

We will in particular concentrate on structural conditions, possibly involving the dimension, guaranteeing classical solutions. The interested reader can find a complete updated account in the book [14] for the elliptic theory. We do not aim at discussing here $W_q^{2,1}$ regularity of solutions and we refer to [21] for more details and to the next section for the

main results on the low-regularity theory.

The simplest result for (4) says that solutions to uniformly parabolic equations in 3D (here we denote by D the dimension in space-time) are always classical without any other assumption on F , see [1, 15, 30] and the references therein. Its elliptic analogue was proved by L. Nirenberg (in the elliptic dimension $n = 2$) [14, Theorem 4.9], and its proof, based on the De Giorgi-Nash-Moser theory applied to the equation solved by second derivatives, does not seem to generalize to parabolic equations. It is also important to mention a result of S. Kruzhkov showing that the same result holds in the 2D case, cf. [18] and Section XIV.7 in [24].

Theorem 2.1. *Assume that F is uniformly elliptic and $n = 2$. Then, viscosity solutions to (4) (in dimension $n + 1 = 3$) are always classical and belong to $C_{\text{loc}}^{2+\alpha, 1+\alpha/2}$ for some small universal $\alpha \in (0, 1)$. It holds*

$$\|u\|_{C^{2+\alpha, 1+\alpha/2}(Q_{\frac{1}{2}})} \leq C\|u\|_{L^\infty(Q_1)}.$$

A second result holds under Cordes-type assumptions on the ellipticity in any space dimension, see Theorem 6.4 in [15] or [10].

Theorem 2.2. *Assume that F is uniformly elliptic and $\frac{\Lambda}{\lambda} \leq 1 + \delta$ for a small (explicit) constant δ depending only on n . Then viscosity solutions to (4) are always classical (in any dimension) and belong to $C_{\text{loc}}^{2+\alpha, 1+\alpha/2}$ for some small universal α . In addition, we have the regularity estimate*

$$\|u\|_{C^{2+\alpha, 1+\alpha/2}(Q_{\frac{1}{2}})} \leq C\|u\|_{L^\infty(Q_1)}.$$

We also mention that $C^{2+\alpha, 1+\alpha/2}$ estimates can be obtained without any concavity assumptions for short time horizons, see e.g. Chapter 8 of [26], [25] and the references therein. This corresponds to an elliptic result proved in [13] and saying that $C^{2,\alpha}$ and higher order estimates for an equation of the form $F(x, u, Du, D^2u) + \theta u = 0$ can be achieved if the zero-th order coefficient θ is large.

A counterpart of the result by Evans-Krylov [12, 21] provides parabolic $C^{2,\alpha}$ estimates under concavity assumptions on F , cf. [33], see also [24, 19, 21] for smooth operators.

Theorem 2.3. *Assume that F is uniformly elliptic and concave. Then viscosity solutions to (4) are always classical and belong to $C_{\text{loc}}^{2+\alpha, 1+\alpha/2}$ for some small universal α . It holds*

$$\|u\|_{C^{2+\alpha, 1+\alpha/2}(Q_{\frac{1}{2}})} \leq C\|u\|_{L^\infty(Q_1)}.$$

Concavity on the second derivatives was recently weakened to the requirement of convexity of the superlevel sets, cf. [15]. The latter also contains some $C^{1,1}$ estimates under concavity-type conditions at infinity implying, among others, $W_q^{2,1}$ estimates for non-homogeneous equations. A recent account on the treatment of fully nonlinear parabolic equations with relaxed concavity conditions can be found in [21].

We should mention that slightly before the Evans-Krylov theory, the paper [29] provided

an obstacle problem approach to reach parabolic $C^{2,\beta}$, $\beta \in (0, 1)$ small and universal, regularity for the model equation

$$\max\{\partial_t u - L_1 u, \partial_t u - L_2 u\} = 0$$

(here L_1, L_2 are uniformly elliptic linear operators) by reducing the problem to a variational inequality, an idea introduced by Brézis-Evans [3]. More references on the early regularity theory of fully nonlinear elliptic and parabolic equations can be found in [24, p. 382-384].

More recently, maximal C^α estimates were also achieved for flat solutions (i.e. with small L^∞ norm) in [34] without concavity assumptions on F , extending a result due to O. Savin.

We also emphasize that counterexamples to the smoothness of solutions when F is only uniformly parabolic can be found in the recent paper [30], which provides a time-dependent counterpart of the analysis in [28].

Different kind of results were studied in the paper [11], in which the author provided a partial regularity result, again without imposing any concavity condition on $D^2 u$, as initiated by Armstrong-Silvestre-Smart for the stationary problem, under the requirement that $F \in C^1$. It gives that a viscosity solution of a uniformly parabolic equation is $C^{2+\alpha, 1+\alpha/2}$ on the complement of a closed set having Hausdorff dimension at most $n + 1 - \varepsilon$, where ε depends on λ, Λ, n .

Finally, [8] studied the problem of the optimal parabolic regularity for the simple Bellman equation

$$\max\{L_1 u, L_2 u\} - \partial_t u = 0,$$

providing a counterexample to the $C^{2,1}$ regularity of its solution. It seems a difficult open question to find the optimal regularity of solutions to general Bellman equations, even with constant coefficients, assuming only F continuous and the uniform ellipticity. In this framework the best known elliptic and parabolic regularity in general dimension $n \geq 2$ is the Evans-Krylov $C^{2,\beta}$ regularity with β small. This problem was raised in [3, Remark 4.2], see also [14], and the paper [5] highlighted that for the elliptic PDE $\max\{L_1 u, L_2 u\} = 0$ in \mathbb{R}^2 , viscosity solutions are of class $C^{2,1}$, while it remains open the general case $n \geq 3$.

Summarizing, for a uniformly parabolic equation of the form (4) with F smooth the following results are known:

- If $u \in C_{x,t}^{2,1}$, then $u \in C^\infty$ by the linear theory, see [15, Proposition 5.3];
- $Du, \partial_t u \in C_{\text{loc}}^{\alpha, \alpha/2}$ always by the Krylov-Safonov theory, see Theorem 3.4;
- Solutions are always smooth for short times without concavity assumptions, see [25] and Chapter 8 of [26], or for arbitrary times when F is a smooth nonlinear strictly increasing function of the Laplacian, i.e. $F = f(\Delta u)$, see [32] and Section 8.5.2 of [26];
- In the parabolic dimensions $D = 2, 3$, solutions are always smooth, see Theorem 2.1 and [18];

- In any space dimension, if $\frac{\Lambda}{\lambda} \leq 1 + \delta(n)$, then the solutions are always smooth, see Theorem 2.2;
- If F is concave, then $u \in C^\infty$, see Theorem 2.3 and Corollary 8.7 in [15];
- $u \in C_{\text{loc}}^{2+\alpha, 1+\alpha/2}$ in the case of flat solutions, see [34];
- The assumption of concavity of F can be weakened to convexity of superlevel sets, see [15];
- Solutions of time-dependent Bellman equations cannot be $C^{2,1}$, see [8].

3. $C^{2+\alpha, 1+\alpha/2}$ REGULARITY FOR FULLY NONLINEAR PARABOLIC ISAACS EQUATIONS

Our main result is an Evans-Krylov theorem for

$$(5) \quad F(D^2u) - \partial_t u = 0 \text{ in } Q_1 = B_1 \times (-1, 0].$$

with

$$(6) \quad \begin{cases} F(M) = \min\{F^\cap(M), F^\cup(M)\}, \quad \forall M \in \mathcal{S}_n \\ F(0) = 0, \quad F^\cap, F^\cup \text{ are uniformly elliptic} \\ F^\cap \text{ is concave in } M, \quad F^\cup \text{ is convex in } M. \end{cases}$$

We recall that the Evans-Krylov theorem allows to pass from $C^{1,1}$ to $C^{2,\alpha}$ regularity in the stationary case, see e.g. [7, Chapter 6].

Theorem 3.1. *Let $u \in C(Q_1)$ be a viscosity solution to (5) with F satisfying (6). Then for some universal $\tilde{\alpha} \in (0, 1)$ depending on n, λ, Λ we have*

$$\|u\|_{C^{2+\tilde{\alpha}, 1+\tilde{\alpha}/2}(\bar{Q}_{\frac{1}{2}})} \leq C \|u\|_{L^\infty(Q_1)}.$$

where C is a constant depending on $\lambda, \Lambda, n, \tilde{\alpha}$.

Remark 3.2. Differently from the elliptic case, the condition $F(0) = 0$ cannot be in general dropped, as outlined in [21, Remark p. 257].

As a corollary, an argument by L. Caffarelli [6] provides the maximal C^α regularity for the non-homogeneous equation.

We denote by $\mathcal{M}_{\lambda, \Lambda}^\pm$ the Pucci's extremal operators with constants $0 < \lambda \leq \Lambda$ defined by

$$\begin{aligned} \mathcal{M}_{\lambda, \Lambda}^+(M) &= \sup_{\lambda I_n \leq A \leq \Lambda I_n} \text{Tr}(AM) = \Lambda \sum_{e_k > 0} e_k + \lambda \sum_{e_k < 0} e_k, \\ \mathcal{M}_{\lambda, \Lambda}^-(M) &= \sup_{\lambda I_n \leq A \leq \Lambda I_n} \text{Tr}(AM) = \lambda \sum_{e_k > 0} e_k + \Lambda \sum_{e_k < 0} e_k, \end{aligned}$$

where $e_k = e_k(M)$ are the eigenvalues of M . Moreover, $\underline{\mathcal{S}}(\lambda, \Lambda, f)$ stands for the space of continuous functions u in Ω that are subsolutions of the evolutive Pucci's maximal equation

$$\mathcal{M}_{\lambda, \Lambda}^+(D^2u) - \partial_t u = f(x, t) \text{ in the viscosity sense in } \Omega,$$

Similarly, $\overline{\mathcal{S}}(\lambda, \Lambda, f)$ will denote the space of continuous functions u in Ω that are supersolutions to the Pucci's minimal equation

$$\mathcal{M}_{\lambda, \Lambda}^-(D^2u) - \partial_t u = f(x, t) \text{ in the viscosity sense in } \Omega.$$

Solutions of fully nonlinear uniformly elliptic equations $F(D^2u) - \partial_t u = 0$ belong to the class $\mathcal{S}(\lambda, \Lambda, 0) = \overline{\mathcal{S}}(\lambda, \Lambda, 0) \cap \underline{\mathcal{S}}(\lambda, \Lambda, 0)$.

Before proving the result we recall some standard facts, see [22] and [16]:

Theorem 3.3 (Krylov-Safonov parabolic Hölder regularity). *Let $u \in C(Q_1)$ be a viscosity solution of $F(D^2u) - \partial_t u = 0$, F uniformly elliptic. Then, for some $\alpha > 0$ depending on n, λ, Λ we have that $u \in C^{\alpha, \alpha/2}(Q_{\frac{1}{2}})$ and the following regularity estimate holds*

$$\|u\|_{C^{\alpha, \alpha/2}(Q_{\frac{1}{2}})} \leq C \|u\|_{L^\infty(Q_1)},$$

where C is a constant depending on λ, Λ, n .

This Hölder property is the best one can get for equations with bounded and measurable coefficients in the parabolic dimension $n + 1 \geq 2$ by Theorem 1.1 (see also the discussion below) of [30]. The previous estimates are the cornerstone to deduce higher regularity of first-order: the next result holds for viscosity solutions of fully nonlinear parabolic homogeneous equations without any other assumption on F other than the uniform ellipticity, cf. Theorem 2.3 in [30] or Lemma 3 p. 257 in [21] (see also Corollary 5.7 in [7] for the elliptic case).

Theorem 3.4 (First-order space-time Hölder regularity). *Let $u \in C(Q_1)$ be a viscosity solution of $F(D^2u) - \partial_t u = 0$, F uniformly elliptic. Then, for some $\alpha > 0$ depending on n, λ, Λ we have that $\partial_t u, Du \in C^{\alpha, \alpha/2}(Q_{\frac{1}{2}})$ and*

$$\|\partial_t u\|_{C^{\alpha, \alpha/2}(Q_{\frac{1}{2}})} + \|Du\|_{C^{\alpha, \alpha/2}(Q_{\frac{1}{2}})} \leq C \|u\|_{L^\infty(Q_1)},$$

where C depends on λ, Λ, n .

We also have the following interior a priori $C^{2, \beta}$ estimate for the stationary non-homogeneous equation driven by operators satisfying (6), see Corollary 1.3 and Remark 1.4 of [4].

Theorem 3.5. *Let $u \in C(B_1)$ be a viscosity solution to $F(D^2u) = f(x)$ in B_1 with $f \in C^\beta(B_1)$ for some $0 < \beta < \bar{\beta}$, $\bar{\beta} \in (0, 1)$ being a universal constant, and F satisfying (6). Then, the following estimate holds*

$$\|u\|_{C^{2, \beta}(\overline{B}_{\frac{1}{2}})} \leq C (\|u\|_{L^\infty(B_1)} + \|f\|_{C^\beta(B_{\frac{3}{4}})}),$$

where C depends on $n, \lambda, \Lambda, \beta$.

Next we exploit the $C_x^{2, \beta}$ estimate and the evolution equation to improve the time-Hölder exponent of first-order derivatives from Theorem 3.4 via a comparison principle argument.

Proposition 3.6. *Suppose that there exists $\beta \in (0, 1)$ and a constant $C > 0$ such that*

$$|D^2u(x, \cdot) - D^2u(y, \cdot)| \leq C|x - y|^\beta, \quad x, y \in B_{\frac{1}{2}}.$$

Any $C^{2,1}$ solution to the uniformly parabolic equation (5) (without any other structural assumption on F) satisfies the estimate

$$|Du(\cdot, t) - Du(\cdot, s)| \leq C'|t - s|^{\frac{1+\beta}{2}} \text{ in } B_{\frac{1}{16}}.$$

where C' is a positive constant depending on $C, n, \beta, \lambda, \Lambda$.

Proof. First note that for $e \in \mathbb{R}^n$, $|e| = 1$,

$$u_h(x, t) := \frac{u(x + he, t) - u(x, t)}{h} \in \mathcal{S}(\lambda/n, \Lambda),$$

i.e. u_h is a solution to a nondivergence form equation in a smaller cylinder. This means (see e.g. [7, Proposition 5.5] or [33, Theorem 4.6]) that we have the validity of the following inequalities

$$(7) \quad \mathcal{M}_{\lambda, \Lambda}^+(D^2u_h) - \partial_t u_h \geq 0 \text{ in } Q_{\frac{1}{2}-h}$$

and

$$(8) \quad \mathcal{M}_{\lambda, \Lambda}^-(D^2u_h) - \partial_t u_h \leq 0 \text{ in } Q_{\frac{1}{2}-h}.$$

Moreover, the assumptions on the regularity of D^2u in space (for fixed times) imply that $u_h \in C_x^{1, \beta}$, $\beta \in (0, 1)$, uniformly with respect to h .

We give a proof using the maximum principle and the equivalence of the Hölder seminorms with those given by Taylor polynomials [20, Sections 3.3 and 8.5]. We have by Theorem 3.3.1 in [20]

$$|u_h(z', t) - u_h(z, t) - Du_h(z, t) \cdot (z' - z)| \leq C|z' - z|^{1+\beta} \text{ on } Q_{\frac{1}{2}-h}.$$

We apply the weighted Young's inequality to find for a positive constant \tilde{C}

$$C|z' - z|^{1+\beta} \leq \varepsilon + \tilde{C}\varepsilon^{-\frac{1-\beta}{1+\beta}}|z' - z|^2.$$

This implies the inequality

$$u_h(z', t) \leq u_h(z, t) + Du_h(z, t) \cdot (z' - z) + \varepsilon + \tilde{C}\varepsilon^{-\frac{1-\beta}{1+\beta}}|z' - z|^2.$$

We define

$$\Phi^+(z', t) = u_h(z, t) + Du_h(z, t) \cdot (z' - z) + \varepsilon + \tilde{C}\varepsilon^{-\frac{1-\beta}{1+\beta}}|z' - z|^2 + \nu(t - t).$$

One can check, after choosing $\nu \geq 2\tilde{C}n\Lambda\varepsilon^{-\frac{1-\beta}{1+\beta}}$, the validity of

$$\mathcal{M}_{\lambda, \Lambda}^+(D^2\Phi^+) - \partial_t \Phi^+ \leq 0,$$

so Φ^+ is a classical supersolution to the equation solved by u_h in $Q_{\frac{1}{4}}$, provided that $h < \frac{1}{8}$.

Recall that $u_h \in \mathcal{S}(\lambda/n, \Lambda)$ and thus solves a linear parabolic homogeneous equation in nondivergence form, see (7)-(8). Choosing ε small enough we have that $\Phi^+ \geq u_h$ on

$\partial B_{\frac{1}{4}} \times (-\frac{1}{16}, 0]$. By the maximum principle [9, Theorem 8.2] $\Phi^+ \geq u_h$ in the interior of the cylinder. Taking $z' = z$ and optimizing with respect to ε we conclude

$$u_h(z, t') \leq u_h(z, t) + \bar{C}_1(t' - t)^{\frac{1+\beta}{2}},$$

for a positive constant \bar{C}_1 . We can get a similar estimate from below by constructing a subsolution, say Φ^- , to the PDE

$$\mathcal{M}_{\lambda, \Lambda}^-(D^2\Phi^-) - \partial_t\Phi^- \geq 0$$

and applying the comparison principle. More precisely, we have

$$u_h(z', t) \geq u_h(z, t) + Du_h(z, t) \cdot (z' - z) - \varepsilon - \tilde{C}\varepsilon^{-\frac{1-\beta}{1+\beta}}|z' - z|^2.$$

We define

$$\Phi^-(z', t') = u_h(z, t) + Du_h(z, t) \cdot (z' - z) - \varepsilon - \tilde{C}\varepsilon^{-\frac{1-\beta}{1+\beta}}|z' - z|^2 - \nu(t' - t)$$

and choose $\nu \geq 2\tilde{C}n\Lambda\varepsilon^{-\frac{1-\beta}{1+\beta}}$ to conclude that $\mathcal{M}_{\lambda, \Lambda}^-(D^2\Phi^-) - \partial_t\Phi^- \geq 0$. By the maximum principle we have $\Phi^- \leq u_h$ in the interior of the cylinder, and proceeding as above we have for a positive constant \bar{C}_2 the estimate

$$u_h(z, t') \geq u_h(z, t) - \bar{C}_2(t' - t)^{\frac{1+\beta}{2}}.$$

□

We are now ready with the proof of the main result:

Proof of Theorem 3.1. Theorem 3.4 shows that viscosity solutions of (5) satisfy $\partial_t u \in C^{\tilde{\alpha}, \tilde{\alpha}/2}$ for some small $\tilde{\alpha} \in (0, 1)$ depending on λ, Λ and n . Recall that such a result does not depend on the structure of the operator and requires only its uniform ellipticity. Therefore, we can consider

$$F(D^2u) = \partial_t u \in C^{\tilde{\alpha}, \tilde{\alpha}/2}.$$

We freeze the time variable and apply Theorem 3.5 on each time slice to find

$$|D^2u(x, \cdot) - D^2u(y, \cdot)| \leq C|x - y|^{\tilde{\alpha}}.$$

Applying Proposition 3.6 with $\beta = \tilde{\alpha}$ and the space regularity of D^2u we conclude that

$$|Du(\cdot, t) - Du(\cdot, s)| \leq C|t - s|^{\frac{1+\tilde{\alpha}}{2}}.$$

Therefore, D^2u is time-Hölder continuous with exponent $\tilde{\alpha}/2$ after invoking the interpolation argument from Lemma 3.1 p. 80 of [23] applied to Du . We recall briefly that it says that if a function v is $C_t^{\gamma_1}$ and $Dv \in C_x^{\gamma_2}$, $\gamma_1, \gamma_2 \in (0, 1)$, then $Dv \in C_t^{\gamma_1\gamma_2/(1+\gamma_2)}$. This implies that $D^2u \in C^{\tilde{\alpha}, \tilde{\alpha}/2}$.

□

Remark 3.7. It is worth comparing the strategy outlined above with those already appeared in the literature of fully nonlinear parabolic equations. There are essentially two ways to achieve $C^{2+\alpha,1+\alpha/2}$ estimates for homogeneous fully nonlinear parabolic equations. The first one is to prove Evans-Krylov estimates directly for parabolic equations, following e.g. [19]. The second one is to consider (5) as $F(D^2u) = \partial_t u$, give an estimate of $\partial_t u$ and then consider the PDE for any fixed t as an elliptic one: this viewpoint however requires the development of a $C^{2,\alpha}$ theory for the non-homogeneous elliptic equation $F(D^2u) = f(x) \in C^\alpha$. The approach carried out in Chapter 12 of [21] follows this second idea, and exploits an interpolation argument between $D^2u(\cdot, t) \in C_x^\alpha$ and $\partial_t u(\cdot, t) \in C_x^\alpha$ for fixed times t (cf. Lemma 4 p. 259 in [21]), a different way with respect to our method. Related ideas were also employed in [1, 30] for parabolic equations in 3D.

It is worth pointing out that the foregoing approach applies to deduce interior $C^{2+\alpha,1+\alpha/2}$ bounds in the case of concave operators, with a different proof than Theorem 4.13 in [33] (in particular without using Lemma 4.16 therein). Here we provided a general strategy to deduce parabolic $C^{2+\alpha,1+\alpha/2}$ regularity through the maximum principle and the $C^{2,\alpha}$ regularity in the space variable only. We can generalize the previous argument providing an abstract result under the following hypothesis

(H) The stationary equation $F(D^2u) = f$, $f \in C^\alpha$, has interior $C^{2,\alpha}$ estimates for some $\alpha \in (0, 1)$ with constant c_e .

Theorem 3.8. *Let $u \in C(Q_1)$ be a viscosity solution of the equation $F(D^2u) - \partial_t u = 0$ in Q_1 . Assume that F is uniformly elliptic and that (H) holds. Then, $u \in C^{2+\alpha,1+\alpha/2}(Q_{\frac{1}{2}})$ and the following estimate holds*

$$\|u\|_{C^{2+\alpha,1+\alpha/2}(Q_{\frac{1}{2}})} \leq C$$

for a constant C depending on $n, \lambda, \Lambda, \alpha, c_e$.

We conclude by discussing the case of non-homogeneous equations. Schauder's estimates proved by L. Caffarelli [7, Section 8.1], see also [33], would imply $C^{2+\alpha,1+\alpha}$ estimates for the following equation

$$F(x, t, D^2u) - \partial_t u = f(x, t)$$

under the following assumptions on F :

- (i) For every $(x_0, t_0) \in Q_1$, the operator $F(x_0, t_0, \cdot)$ is the minimum of a concave and a convex operator (possibly depending on (x_0, t_0));
- (ii) $F(\cdot, M)$ and $f(\cdot)$ are Hölder continuous functions with respect to the parabolic distance.

This implies that $F(x, t, D^2u)$ can be an Isaacs operator of the form (3) with variable Hölder continuous coefficients with respect to the parabolic distance.

REFERENCES

- [1] B. Andrews. Fully nonlinear parabolic equations in two space variables. arXiv:0402235, 2004.
- [2] A. Brandt. Interior Schauder estimates for parabolic differential- (or difference-) equations via the maximum principle. *Israel J. Math.*, 7:254–262, 1969.

- [3] H. Brézis and L. C. Evans. A variational inequality approach to the Bellman-Dirichlet equation for two elliptic operators. *Arch. Rational Mech. Anal.*, 71(1):1–13, 1979.
- [4] X. Cabré and L. A. Caffarelli. Interior $C^{2,\alpha}$ regularity theory for a class of nonconvex fully nonlinear elliptic equations. *J. Math. Pures Appl. (9)*, 82(5):573–612, 2003.
- [5] L. Caffarelli, D. De Silva, and O. Savin. Two-phase anisotropic free boundary problems and applications to the Bellman equation in 2d. *Arch. Ration. Mech. Anal.*, 228(2):477–493, 2018.
- [6] L. A. Caffarelli. Interior a priori estimates for solutions of fully nonlinear equations. *Ann. of Math. (2)*, 130(1):189–213, 1989.
- [7] L. A. Caffarelli and X. Cabré. *Fully nonlinear elliptic equations*, volume 43 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1995.
- [8] L. A. Caffarelli and U. Stefanelli. A counterexample to $C^{2,1}$ regularity for parabolic fully nonlinear equations. *Comm. Partial Differential Equations*, 33(7-9):1216–1234, 2008.
- [9] M. G. Crandall, H. Ishii, and P.-L. Lions. User’s guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc. (N.S.)*, 27(1):1–67, 1992.
- [10] J. V. da Silva and M. S. Santos. Schauder and Calderón-Zygmund type estimates for fully nonlinear parabolic equations under “small ellipticity aperture” and applications. *Nonlinear Analysis*, 2023.
- [11] J.-P. Daniel. Quadratic expansions and partial regularity for fully nonlinear uniformly parabolic equations. *Calc. Var. Partial Differ. Equ.*, 54(1):183–216, 2015.
- [12] L. C. Evans. Classical solutions of fully nonlinear, convex, second-order elliptic equations. *Comm. Pure Appl. Math.*, 35(3):333–363, 1982.
- [13] L. C. Evans and P.-L. Lions. Fully nonlinear second order elliptic equations with large zeroth order coefficient. *Ann. Inst. Fourier*, 31(2):175–191, 1981.
- [14] X. Fernández-Real and X. Ros-Oton. *Regularity theory for elliptic PDE*, volume 28 of *Zurich Lectures in Advanced Mathematics*. EMS Press, Berlin, [2022] 2022.
- [15] A. Goffi. High-order estimates for fully nonlinear equations under weak concavity assumptions. *J. Math. Pures Appl. (9)*, 182:223–252, 2024.
- [16] C. Imbert and L. Silvestre. An introduction to fully nonlinear parabolic equations. In *An introduction to the Kähler-Ricci flow*, volume 2086 of *Lecture Notes in Math.*, pages 7–88. Springer, Cham, 2013.
- [17] B. F. Knerr. Parabolic interior Schauder estimates by the maximum principle. *Arch. Rational Mech. Anal.*, 75(1):51–58, 1980/81.
- [18] S. N. Kruzhkov. Nonlinear parabolic equations with two independent variables. *Trudy Moskov. Mat. Obsč.*, 16:329–346, 1967.
- [19] N. V. Krylov. *Nonlinear elliptic and parabolic equations of the second order. Transl. from the Russian by P. L. Buzdytsky*, volume 7 of *Math. Appl., Sov. Ser.* Kluwer Academic Publishers Group, Dordrecht, 1987.
- [20] N. V. Krylov. *Lectures on elliptic and parabolic equations in Hölder spaces*, volume 12 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1996.
- [21] N. V. Krylov. *Sobolev and viscosity solutions for fully nonlinear elliptic and parabolic equations*, volume 233 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2018.
- [22] N. V. Krylov and M. V. Safonov. A certain property of solutions of parabolic equations with measurable coefficients. *Math. USSR, Izv.*, 16:151–164, 1981.
- [23] O. A. Ladyzenskaja, V. A. Solonnikov, and N. N. Ural’ceva. *Linear and quasilinear equations of parabolic type*. Translations of Mathematical Monographs, Vol. 23. American Mathematical Society, Providence, R.I., 1968. Translated from the Russian by S. Smith.
- [24] G. M. Lieberman. *Second order parabolic differential equations*. World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
- [25] P.-L. Lions and P. E. Souganidis. Fully nonlinear second-order degenerate elliptic equations with large zeroth-order coefficients. *Indiana Univ. Math. J.*, 42(4):1525–1543, 1993.
- [26] A. Lunardi. *Analytic semigroups and optimal regularity in parabolic problems*. Mod. Birkhäuser Class. Basel: Birkhäuser, reprint of the 1995 hardback ed. edition, 2013.

- [27] J. H. Michael. A general theory for linear elliptic partial differential equations. *J. Differential Equations*, 23(1):1–29, 1977.
- [28] N. Nadirashvili, V. Tkachev, and S. Vlăduț. *Nonlinear elliptic equations and nonassociative algebras*, volume 200 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2014.
- [29] M. Schoenauer. Some regularizing effects for the Hamilton-Jacobi-Bellman equation for two parabolic operators. *Comm. Partial Differential Equations*, 6(8):929–949, 1981.
- [30] L. Silvestre. Singular solutions to parabolic equations in nondivergence form. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 23(2):993–1011, 2022.
- [31] J. van der Hoek. Existence via interior estimates for second order parabolic equations. In *Miniconference on nonlinear analysis (Canberra, 1984)*, volume 8 of *Proc. Centre Math. Anal. Austral. Nat. Univ.*, pages 221–232. Austral. Nat. Univ., Canberra, 1984.
- [32] W. von Wahl. On the equation $u' - f(\Delta u) = 0$. *Boll. Unione Mat. Ital., VII. Ser., A*, 1:437–441, 1987.
- [33] L. Wang. On the regularity theory of fully nonlinear parabolic equations. II. *Comm. Pure Appl. Math.*, 45(2):141–178, 1992.
- [34] Y. Wang. Small perturbation solutions for parabolic equations. *Indiana Univ. Math. J.*, 62(2):671–697, 2013.

DIPARTIMENTO DI MATEMATICA “TULLIO LEVI-CIVITA”, UNIVERSITÀ DEGLI STUDI DI PADOVA, VIA TRIESTE 63, 35121 PADOVA (ITALY)

Email address: `alessandro.goffi@unipd.it`