

Affordable mixed-integer Lagrangian methods: optimality conditions and convergence analysis

Alberto De Marchi

June 19, 2024

Abstract

Necessary optimality conditions in Lagrangian form and the augmented Lagrangian framework are extended to mixed-integer nonlinear optimization, without any convexity assumptions. Building upon a recently developed notion of local optimality for problems with polyhedral and integrality constraints, a characterization of local minimizers and critical points is given for problems including also nonlinear constraints. This approach lays the foundations for developing affordable sequential minimization algorithms with convergence guarantees to critical points from arbitrary initializations. A primal-dual perspective, a local saddle point property, and the dual relationships with the proximal point algorithm are also advanced in the presence of integer variables.

Keywords. Mixed-integer nonlinear programming, Necessary optimality conditions, Augmented Lagrangian framework, Lagrangian duality, Proximal point algorithm.

MSC 2020. [65K05](#), [90C06](#), [90C11](#), [90C30](#).

Contents

1	Introduction	2
1.1	Notation and Preliminaries	3
2	Optimality concepts	4
2.1	Stationarity concepts and first-order Lagrangian analysis	5
3	Augmented Lagrangian framework	8
3.1	Convergence analysis	9
4	Further characterizations	12
4.1	Lagrangian duality	12
4.2	Saddle points of the augmented Lagrangian	14
4.3	Relationship with Proximal Point Methods	15
5	Concluding Remarks	16
	References	16

University of the Bundeswehr Munich, Department of Aerospace Engineering, Institute of Applied Mathematics and Scientific Computing, 85577 Neubiberg, Germany. EMAIL alberto.demarchi@unibw.de, ORCID [0000-0002-3545-6898](https://orcid.org/0000-0002-3545-6898).

1 Introduction

Mixed-integer nonlinear programming (MINLP) offers a versatile template for capturing a variety of tasks and applications, but brings together “the combinatorial difficulty of optimizing over discrete variable sets with the challenges of handling nonlinear functions” [2]. Originating from the integer programming community, most approaches for MINLP rely on some sort of tree search for seeking globally optimal solutions, at least when some convexity is available. Our focus is on affordable techniques, also called heuristics, for addressing nonconvex MINLPs numerically. In particular, we are interested in iterative algorithms designed to converge in some sense to local solutions, not necessarily global minimizers, starting from arbitrary initial points [3, Chapter 6]. This allows us to handle large instances for a broad problem class, but requires defining a strong notion of local optimality, with the aim of striking a balance between global but expensive minima and local but affordable critical points. We seek as stationarity characterization that resembles, at least in spirit, the so called Karush-Kuhn-Tucker (or KKT) conditions in nonlinear programming, see e.g. [3, Chapter 3]. Although “in mixed-integer nonlinear programming, we do not know local optimality conditions comparable to the KKT conditions in continuous optimization” [12, Section 2], some advancements have been made based on an excess of multipliers and separation theorems [14]. In an attempt to upgrade our understanding, we study here a criticality concept for nonconvex MINLPs in simple Lagrangian terms. Building upon the optimality notions developed in [5], we dedicate this work to characterizing “local” minima with a Lagrangian perspective and then establishing convergence results for a class of augmented Lagrangian (AL) methods.

A mixed-integer linearization algorithm was proposed in [5] to address the minimization of a smooth function over a feasible set with mixed-integer linear structure, namely MINLP without nonlinear constraints. Nikitina et al. [15] recently developed a safeguarded AL scheme that exploits this affordable solver for tackling the AL subproblems. Although lacking a convergence analysis, they demonstrated the validity of their approach with extensive numerical tests. The present work provides solid theoretical foundations for the algorithmic design and numerical results obtained therein. Even beyond their AL scheme, we are motivated by the sequential (partially) unconstrained minimization framework [13], which includes (shifted) penalty [3] and barrier (or interior point) methods [10]. We discuss how this framework can be used to design other algorithms for MINLP, and in particular we indicate how similar arguments apply also to interior point approaches on the line of [10]. Methods based on sequential mixed-integer quadratic programming [12, 16] could benefit from these theoretical advances too. Other numerical approaches for MINLP, such as global methods or decomposition techniques [2, 19], could also exploit these principled heuristics to refine initial guesses, generate tighter bounds, and promote faster convergence.

Beyond numerical methods for MINLP, we enrich the theoretical framework and first-order analysis of mixed-integer optimization in Lagrangian terms, inspired by the celebrated KKT conditions in nonlinear programming. In the spirit of [14, 21, 17], we develop a theory of KKT-critical points, complemented by Lagrangian duality, saddle point properties, and relationships with the proximal points algorithm.

The problem template with nonconvex smooth objective and polyhedral, integrality, and nonlinear set-membership constraints reads

$$\text{minimize } f(x) \quad \text{over } x \in \mathcal{X} \quad \text{subject to } c(x) \in \mathcal{C} \quad (\text{P})$$

with $f: \mathcal{X} \rightarrow \mathbb{R}$ and $c: \mathcal{X} \rightarrow \mathbb{R}^m$ continuously differentiable functions, $\mathcal{C} \subset \mathbb{R}^m$ a nonempty closed convex set (projection-friendly in practice), and \mathcal{X} a nonempty closed set with mixed-

integer linear structure [5]. Set \mathcal{X} admits a description in the form

$$\mathcal{X} := \left\{ x \in \mathbb{R}^n \mid \begin{array}{l} A_e x = b_e, \ x_1 \leq x \leq x_u, \\ x_i \in \mathbb{Z} \ \forall i \in \mathcal{I} \end{array} \right\}$$

for some matrix A_e , vectors b_e , x_1 and x_u , and index set \mathcal{I} . In the following, we may refer to a partition of decision variables x into real-valued and integer-valued ones, respectively $\{x_i \mid i \notin \mathcal{I}\}$ and $\{x_i \mid i \in \mathcal{I}\}$. Furthermore, patterning [5, 15], we consider the following blanket assumptions.

Assumption 1.1. With regard to (P),

- (A1) $\{x \in \mathcal{X} \mid c(x) \in \mathcal{C}\}$ is nonempty and $\inf \{f(x) \mid x \in \mathcal{X}, c(x) \in \mathcal{C}\} \in \mathbb{R}$;
- (A2) functions f and c are continuously differentiable with locally Lipschitz derivatives;
- (A3) for all $i \in \mathcal{I}$ the set $\{a \in \mathbb{Z} \mid x \in \mathcal{X}, x_i = a\}$ is bounded.

The basic Assumption (A1) ensures that (P) is well-posed, namely that it is feasible and a solution exists. Differentiability of f and c in Assumption (A2) is intended with respect to real- and integer-valued variables, treating them all as real-valued ones to avoid exotic definitions or approximations, such as those in [12]. A practical situation that satisfies Assumption (A2) is when f and c depend linearly on the integer-valued variables, as supposed in [16]. Finally, Assumption (A3) guarantees that admissible values (with respect to \mathcal{X} alone) for the integer-valued decision variables lie in a bounded set. As it applies to integer-valued variables only, this boundedness requirement is reasonable and often satisfied in practice (trivially for binary variables). Following [5], we take advantage of Assumption (A3) to construct compact neighborhoods without explicitly localizing the integer-valued components.

Motivated by the numerical approach proposed in [15], we build a theoretical support for the optimality concepts and algorithms adopted there, and beyond, establishing convergence results under suitable assumptions. Our contributions can be summarized as follows:

- We derive and analyse necessary optimality conditions for (P) in Lagrangian form, comparable to the KKT system in continuous optimization—see Section 2.1.
- We prove the global convergence of a safeguarded augmented Lagrangian algorithm—see Algorithm 3.1 and Section 3.1. Providing a solid theoretical foundation for the numerical scheme investigated in [15], we generalize the affordable approach of [5] to sequential minimization schemes for MINLP.
- The Lagrangian system is further characterized in primal-dual terms, recovering generalized saddle-point properties and a close dual relationship with the proximal point algorithm—see Section 4.

1.1 Notation and Preliminaries

The set of natural, integer, and real numbers are denoted by \mathbb{N} , \mathbb{Z} , \mathbb{R} . The appearing spaces are equipped with the standard Euclidean inner product $\langle \cdot, \cdot \rangle$ and norm $\|\cdot\|$. Given a nonempty subset \mathcal{C} of \mathbb{R}^m , the *indicator* $\delta_{\mathcal{C}}: \mathbb{R}^m \rightarrow \mathbb{R} \cup \{\infty\}$, the *projection* $\text{proj}_{\mathcal{C}}: \mathbb{R}^m \rightarrow \mathcal{C}$, and the *distance* $\text{dist}_{\mathcal{C}}: \mathbb{R}^m \rightarrow \mathbb{R}$ are defined respectively by

$$\delta_{\mathcal{C}}(v) := \begin{cases} 0 & \text{if } v \in \mathcal{C}, \\ \infty & \text{otherwise,} \end{cases} \quad \text{proj}_{\mathcal{C}}(v) := \arg \min_{z \in \mathcal{C}} \|z - v\|, \quad \text{dist}_{\mathcal{C}}(v) := \min_{z \in \mathcal{C}} \|z - v\|.$$

The *normal cone* $\mathcal{N}_{\mathcal{C}}(z)$ of set $\mathcal{C} \subseteq \mathbb{R}^m$ at $z \in \mathcal{C}$ is given by

$$\mathcal{N}_{\mathcal{C}}(z) := \{v \in \mathbb{R}^m \mid \forall u \in \mathcal{C}: \langle v, u - z \rangle \leq 0\}.$$

For formal completeness, we define $\mathcal{N}_{\mathcal{C}}(z) := \emptyset$ if $z \notin \mathcal{C}$. We will make use of the following well known characterizations valid for a closed convex set $\mathcal{C} \subseteq \mathbb{R}^m$ [1]:

$$u \in \text{proj}_{\mathcal{C}}(z) \iff \forall w \in \mathcal{C}: \langle z - u, w - u \rangle \leq 0, \quad (1)$$

$$u \in \mathcal{N}_{\mathcal{C}}(z) \iff \forall \alpha > 0: z = \text{proj}_{\mathcal{C}}(z + \alpha u) \iff \exists \alpha > 0: z = \text{proj}_{\mathcal{C}}(z + \alpha u). \quad (2)$$

2 Optimality concepts

A point $\bar{x} \in \mathbb{R}^n$ is called *feasible* for (P) if $\bar{x} \in \mathcal{X}$ and $c(\bar{x}) \in \mathcal{C}$. It is clear how to define a *global* solution, or minimizer, x^* for (P),

$$x^* \in \mathcal{X}, \quad c(x^*) \in \mathcal{C}, \quad \forall x \in \mathcal{X}, c(x) \in \mathcal{C}: f(x^*) \leq f(x),$$

but what constitutes a suitable notion of *local* minimizer? Local notions, as opposed to global ones, depend on the concept of neighborhood and this, in turn, is very delicate in the mixed-integer context of (P). Following [5], we denote by $\|\cdot\|_{\text{LP}}$ an operator mapping x into a norm of the real-valued entries of x , that is, given the index set \mathcal{I} . Prominent examples are $\|v\|_{\text{LP}} := \max\{|v_i| \mid i \notin \mathcal{I}\}$ and $\|v\|_{\text{LP}} := \sum\{|v_i| \mid i \notin \mathcal{I}\}$, associated with ℓ_{∞} and ℓ_1 norms, respectively. Due to this partial localization, balls

$$\mathbb{B}_{\text{LP}}(x, \Delta) := \{w \in \mathbb{R}^n \mid \|w - x\|_{\text{LP}} \leq \Delta\}$$

induced by $\|\cdot\|_{\text{LP}}$ are *not* compact sets. Nevertheless, the intersection $\mathcal{X} \cap \mathbb{B}_{\text{LP}}(x, \Delta)$ is always a compact set, thanks to Assumption (A3), and thus represents a reasonable neighborhood of x —and a valid trust region stipulation—for any $x \in \mathcal{X}$ and $\Delta \geq 0$. Before proceeding, we should mention that adopting a polyhedral norm to define $\|\cdot\|_{\text{LP}}$ is favourable in practice, as the mixed-integer *linear* structure is not lost in the subproblems, but the theory applies with any norm. A local concept of solution for (P) can now be defined by means of these (partial) neighborhoods. Inspired by [5, Definition 2.1], local and global minimizers for (P) are characterized as follows.

Definition 2.1 (minimizers). A point $\bar{x} \in \mathbb{R}^n$ is called a *local minimizer* for (P) if it is feasible and there exists $\Delta > 0$ such that $f(\bar{x}) \leq f(x)$ for all feasible $x \in \mathbb{B}_{\text{LP}}(\bar{x}, \Delta)$. If the latter property additionally holds for all $\Delta > 0$, then \bar{x} is called a *global minimizer*.

For instances of (P) without integer-valued variables, namely $\mathcal{I} := \emptyset$, Definition 2.1 recovers the classical notion of local minima in nonlinear programming. Conversely, without real-valued variables, namely $\mathcal{I} := \{1, 2, \dots, n\}$, (P) is an integer program and Definition 2.1 effectively requires a global solution (since there is no actual localization in this case). Thus, we can observe that monitoring neighborhoods with $\|\cdot\|_{\text{LP}}$ leads to a stronger local optimality concept than a plain adaptation of continuous notions into the mixed-integer realm. Conversely, the combinatorial structure in (P) should be simple enough for practical purposes, e.g., mixed-integer linear.

Before delving into KKT-like optimality conditions for (P), let us recall some solution concepts for problems without nonlinear constraints. Following [5], consider the minimization of $\varphi: \mathcal{X} \rightarrow \mathbb{R}$ over \mathcal{X} as a basic template:

$$\text{minimize } \varphi(x) \quad \text{over } x \in \mathcal{X}. \quad (3)$$

A local notion of solutions for (3) is proposed in [5, Definition 2.2], inspired by [4, Definition 3.1] for the analogous minimization over a *convex* set. A first-order optimality measure associated to (3) (that is, to function φ and set \mathcal{X}) is defined in [5, Equation 4] and provides a metric $\Psi_{\varphi, \mathcal{X}}$ to monitor “optimality”: for all $x \in \mathcal{X}$ and $\Delta > 0$ it is given by

$$\Psi_{\varphi, \mathcal{X}}(x, \Delta) := \max_{w \in \mathcal{X} \cap \mathbb{B}_{\text{LP}}(x, \Delta)} \langle \nabla \varphi(x), x - w \rangle \geq 0. \quad (4)$$

Since $x, w \in \mathcal{X}$ in (4), $\Psi_{\varphi, \mathcal{X}}(\cdot, \Delta)$ is bounded from below by zero for all $\Delta > 0$. Then, a first-order optimality concept for the “unconstrained” problem (3) is defined as follows; cf. [5, Definition 2.4]..

Definition 2.2 (criticality). Given some $\varepsilon > 0$ and $\Delta > 0$, a point $\bar{x} \in \mathbb{R}^n$ is called ε - Δ -critical for (3) if $\bar{x} \in \mathcal{X}$ and $\Psi_{\varphi, \mathcal{X}}(\bar{x}, \Delta) \leq \varepsilon$. Given some $\varepsilon > 0$, a point $\bar{x} \in \mathbb{R}^n$ is called ε -critical for (3) if it is ε - Δ -critical for some $\Delta > 0$. A 0-critical point is simply called *critical*.

Definition 2.2 provides a valid concept to characterize candidate minimizers, necessary for optimality [5, Proposition 1], which is stronger than plain (M-)stationarity [8]. The criticality notion for “unconstrained”, or simply constrained, problems (3) will become important to characterize solutions to intermediate, auxiliary problems (referred to as subproblems). Moreover, defining an approximate counterpart of criticality allows us to consider inexact subproblem solutions, a strategy often (if not always) adopted in sequential minimization methods [3, 7, 9, 10, 20]. This is useful in accommodating iterative subsolvers with asymptotic convergence, and then in exploiting this property to reduce the overall computational effort.

2.1 Stationarity concepts and first-order Lagrangian analysis

What is a “critical point” for (P)? Treating the nonlinear constraints explicitly, let the Lagrangian function $\mathcal{L}: \mathcal{X} \times \mathbb{R}^m \rightarrow \mathbb{R}$ associated to (P) be defined, as usual, by

$$\mathcal{L}(x, y) := f(x) + \langle y, c(x) \rangle. \quad (5)$$

From the viewpoint of nonlinear programming, where stationarity of the Lagrangian plays a crucial role, we consider the following notion for KKT-like points of (P) based on Definition 2.2, put forward in [15, Definition 2.3]. Then, we are going to establish the (asymptotic) necessity of KKT-criticality for local optimality. Related concepts and results can be found in [6, 10, 7, 8].

Definition 2.3 (KKT-criticality). Given some $\Delta > 0$, a point $\bar{x} \in \mathbb{R}^n$ is called Δ -KKT-critical for (P) if $\bar{x} \in \mathcal{X}$ and there exists a multiplier $y \in \mathbb{R}^m$ such that

$$\Psi_{\mathcal{L}(\cdot, y), \mathcal{X}}(\bar{x}, \Delta) = 0 \quad \text{and} \quad y \in \mathcal{N}_{\mathcal{C}}(c(\bar{x})).$$

A point $\bar{x} \in \mathbb{R}^n$ is called *KKT-critical* for (P) if it is Δ -KKT-critical for some $\Delta > 0$.

KKT-criticality implicitly requires feasibility, since the normal cone $\mathcal{N}_{\mathcal{C}}(c(\bar{x}))$ must be nonempty. Moreover, by (4) the first condition can be rewritten as

$$\min_{x \in \mathcal{X} \cap \mathbb{B}_{\text{LP}}(\bar{x}, \Delta)} \langle \nabla f(\bar{x}) + c'(\bar{x})^\top y, x - \bar{x} \rangle = 0,$$

meaning that the Lagrangian function cannot be (locally) further minimized with respect to x while maintaining mixed-integer linear feasibility, in the sense of Definition 2.2, effectively replacing stationarity with criticality.

An asymptotic counterpart of [Definition 2.3](#) (sometimes also referred to as sequential or approximate) proves to be a key tool for convergence analysis; cf. [[3](#), [Definition 3.1](#)], [[7](#), [6](#)].

Definition 2.4 (AKKT-criticality). A point $\bar{x} \in \mathbb{R}^n$ is called *asymptotically KKT-critical* for (P) if $\bar{x} \in \mathcal{X}$ and there exist sequences $\{x^k\} \subset \mathbb{R}^n$, $\{y^k\} \subset \mathbb{R}^m$, $\{z^k\} \subseteq \mathcal{C}$, and $\{\Delta_k\} \subset \mathbb{R}_{++}$ such that $x^k \rightarrow \bar{x}$ and

$$\Psi_{\mathcal{L}(\cdot, y^k), \mathcal{X}}(x^k, \Delta_k) \rightarrow 0, \quad y^k \in \mathcal{N}_{\mathcal{C}}(z^k), \quad \text{and} \quad c(x^k) - z^k \rightarrow 0.$$

If a sequence $\{x^k\}$ has an accumulation point which is AKKT-critical, then finite termination can be attained with an approximate KKT-critical point, for any given tolerance $\varepsilon > 0$; see ε -KKT-criticality in [[15](#)].

A local minimizer for (P) is KKT-critical under validity of a suitable qualification condition. However, each local minimizer of (P) is always AKKT-critical, regardless of additional regularity. Related results can be found in [[3](#), [7](#), [6](#)].

Theorem 2.5. *Let $x^* \in \mathbb{R}^n$ be a local minimizer for (P). Then, x^* is AKKT-critical.*

Proof. By local optimality of x^* for (P) there exists $\delta > 0$ such that $f(x^*) \leq f(x)$ is valid for all feasible $x \in \mathbb{B}_{\text{LP}}(x^*, \delta)$; cf. [Definition 2.1](#). Consequently, x^* is the unique global minimizer of the localized problem

$$\begin{aligned} & \text{minimize} && f(x) + \|x - x^*\|^2 && (6) \\ & \text{over} && x \in \mathcal{X} \cap \mathbb{B}_{\text{LP}}(x^*, \delta) \\ & \text{subject to} && c(x) \in \mathcal{C}. \end{aligned}$$

Slightly deviating from the proof of [[7](#), [Proposition 2.5](#)], let us consider the penalized surrogate problem

$$\begin{aligned} & \text{minimize} && \pi_k(x) := f(x) + \|x - x^*\|^2 + \rho_k \text{dist}_{\mathcal{C}}^2(c(x)) && (7) \\ & \text{over} && x \in \mathcal{X} \cap \mathbb{B}_{\text{LP}}(x^*, \delta) \end{aligned}$$

where $k \in \mathbb{N}$ is arbitrary, $\rho_k > 0$, and the sequence $\{\rho_k\}_{k \in \mathbb{N}}$ satisfies $\rho_k \rightarrow \infty$ as $k \rightarrow \infty$.

Noting that the objective function of this optimization problem is lower semicontinuous while its feasible set is nonempty and compact (by feasibility of x^* , trust region stipulation, and [Assumption \(A3\)](#)), it possesses a global minimizer $x^k \in \mathcal{X}$ for each $k \in \mathbb{N}$, owing to Weierstrass' extreme value theorem. Without loss of generality, we assume $x^k \rightarrow \tilde{x}$ for some $\tilde{x} \in \mathcal{X} \cap \mathbb{B}_{\text{LP}}(x^*, \delta)$.

We now argue that $\tilde{x} = x^*$. To this end, we note that x^* is feasible to (7) with $c(x^*) \in \mathcal{C}$, which yields for each $k \in \mathbb{N}$ the (uniform, upper) estimate

$$f(x^k) + \|x^k - x^*\|^2 + \rho_k \text{dist}_{\mathcal{C}}^2(c(x^k)) \leq f(x^*). \quad (8)$$

Using $\rho_k \rightarrow \infty$, lower semicontinuity of f , finiteness of $f(x^*)$, closedness of \mathcal{C} , and the convergence $c(x^k) \rightarrow c(\tilde{x})$, taking the limit for $k \rightarrow \infty$ in (8) gives $c(\tilde{x}) \in \mathcal{C}$. Therefore, \tilde{x} is feasible for (P) and local optimality of x^* for (P) implies $f(x^*) \leq f(\tilde{x})$. Furthermore, exploiting (8) and the optimality of each $x^k \in \mathcal{X}$, we find

$$f(\tilde{x}) + \|\tilde{x} - x^*\|^2 \leq \liminf_{k \rightarrow \infty} \left\{ f(x^k) + \|x^k - x^*\|^2 + \rho_k \text{dist}_{\mathcal{C}}^2(c(x^k)) \right\} \leq f(x^*) \leq f(\tilde{x}).$$

Hence, $\tilde{x} = x^*$. Now we may assume without loss of generality that $\{x^k\}$ is taken from the interior of $\mathbb{B}_{\text{LP}}(x^*, \delta)$, as this is eventually the case, since $x^k \rightarrow x^*$. Thus, for each $k \in \mathbb{N}$, x^k globally minimizes π_k over \mathcal{X} , whose relevant criticality condition (necessary for optimality) reads, for some $\Delta_k > 0$,

$$\begin{aligned} 0 &= \Psi_{\pi_k, \mathcal{X}}(x^k, \Delta_k) \\ &= \max_{w \in \mathcal{X} \cap \mathbb{B}_{\text{LP}}(x^k, \Delta_k)} \langle \nabla f(x^k) + 2\rho_k c'(x^k)^\top [c(x^k) - \text{proj}_{\mathcal{C}}(c(x^k))] + 2(x^k - x^*), x^k - w \rangle \\ &= \max_{w \in \mathcal{X} \cap \mathbb{B}_{\text{LP}}(x^k, \Delta_k)} \langle \nabla_x \mathcal{L}(x^k, y^k) + 2(x^k - x^*), x^k - w \rangle \end{aligned}$$

where we set $y^k := 2\rho_k [c(x^k) - \text{proj}_{\mathcal{C}}(c(x^k))]$ for each $k \in \mathbb{N}$. Now, owing to continuous differentiability of \mathcal{L} and compactness of $\mathcal{X} \cap \mathbb{B}_{\text{LP}}(x^k, \Delta_k)$, by $x^k \rightarrow x^* \in \mathcal{X}$ we have

$$\begin{aligned} \lim_{k \rightarrow \infty} \Psi_{\mathcal{L}(\cdot, y^k), \mathcal{X}}(x^k, \Delta_k) &= \lim_{k \rightarrow \infty} \max_{w \in \mathcal{X} \cap \mathbb{B}_{\text{LP}}(x^k, \Delta_k)} \langle \nabla_x \mathcal{L}(x^k, y^k), x^k - w \rangle \\ &= \lim_{k \rightarrow \infty} \max_{w \in \mathcal{X} \cap \mathbb{B}_{\text{LP}}(x^k, \Delta_k)} \langle \nabla_x \mathcal{L}(x^k, y^k) + 2(x^k - x^*), x^k - w \rangle \\ &= \Psi_{\pi_k, \mathcal{X}}(x^k, \Delta_k) = 0. \end{aligned}$$

Thus, the conditions in [Definition 2.4](#) are a consequence of $x^k \rightarrow x^*$. Overall, this shows that any local minimizer x^* for [\(P\)](#) is AKKT-critical. \square

To link AKKT- and KKT-criticality, let us consider a constraint qualification (CQ) based on a geometric condition known as extended Robinson constraint qualification (ERCQ) [[21](#), [Definition 3.17](#)].

Definition 2.6 (Extended Robinson CQ). Let $x \in \mathbb{R}^n$ be arbitrary but fixed. We say that the *extended Robinson constraint qualification* (ERCQ) holds in x if

$$0 \in \text{int} \left(c(x) + c'(x)(\mathcal{X} - x) - \mathcal{C} \right).$$

Here, for brevity, operations involving set \mathcal{X} are meant to apply elementwise. Note that the condition defining ERCQ is the same as for the standard RCQ; the only difference is that, for the latter, x has to be feasible for [\(P\)](#), whereas ERCQ is defined for arbitrary points. Notice that when $\mathcal{X} = \mathbb{R}^n$ and $\mathcal{C} = \{0\}$, ERCQ reduces to the surjectivity of $c'(x)$, which is equivalent to LICQ in the Euclidean setting. In general, RCQ and ERCQ mean that for any sufficiently small perturbation ε of the constraints $c(x)$ there exists some feasible correction $d \in \mathcal{X} - x$ such that $c(x) + c'(x)d + \varepsilon \in \mathcal{C}$. This stability property is enough to bound the set of Lagrange multipliers and thus to guarantee that local minimizers are indeed KKT-critical.

Theorem 2.7. Let $x^* \in \mathbb{R}^n$ be AKKT-critical for [\(P\)](#). If ERCQ holds at x^* , then x^* is KKT-critical for [\(P\)](#).

Proof. Suppose that ERCQ holds at x^* and let $\{y^k\}$ be generated as in the proof of [Theorem 2.5](#). To fill the gap between AKKT- and KKT-criticality it suffices to show that the sequence $\{y^k\}$ remains bounded, so that it admits an accumulation point.

Assume, by contradiction, that $\{y^k\}$ is unbounded. Let us take a suitable subsequence $K \subseteq \mathbb{N}$ such that $\{\|y^k\|\}_{k \in K} \rightarrow \infty$ and $\{y^k / \|y^k\|\}_{k \in K} \rightarrow v \neq 0$ for some $v \in \mathcal{N}_{\mathcal{C}}(c(x^*))$ with $c'(x^*)^\top v = 0$. By ERCQ, we can take $t > 0$ small enough such that

$$0 = c(x^*) + c'(x^*)(\hat{x} - x^*) - w + tv$$

for some $w \in \mathcal{C}$ and $\hat{x} \in \mathcal{X}$. Thus, rearranging and using $c'(x^*)^\top v = 0$, we find that

$$\begin{aligned} 0 < \langle tv, v \rangle &= \langle w - c(x^*), v \rangle - \langle c'(x^*)(\hat{x} - x^*), v \rangle \\ &= \langle w - c(x^*), v \rangle - \langle \hat{x} - x^*, c'(x^*)^\top v \rangle = \langle w - c(x^*), v \rangle \leq 0. \end{aligned}$$

The last inequality follows from $v \in \mathcal{N}_{\mathcal{C}}(c(x^*))$, the characterizations (1)–(2), and the fact that $w \in \mathcal{C}$. The contradiction arising from $t, v \neq 0$ implies that $\{y^k\}$ remains bounded, concluding the proof by taking the limit at any convergent subsequence. \square

3 Augmented Lagrangian framework

Let us consider the MINLP (P) under Assumptions (A1)–(A3), which, under the lens of continuous optimization, can be seen as a nonlinear program with mixed-integer linear constraints, in the spirit of [5]. Since the restriction to \mathcal{X} is nonrelaxable but easy to satisfy, such constraint can be treated in a way essentially different from how nonlinear constraints are handled [3]. Motivated by [15], we study here an augmented Lagrangian (AL) method as an epitome for the class of sequential minimization schemes [13]. The AL framework has been broadly investigated and developed, giving rise to a variety of multifaceted ideas, of which we only scratch the surface here. The interested reader may refer to [3] for an overview, to [6, 8, 17, 21] for theoretical advances, and to [7, 20] for numerical aspects analogous to [15]. The main ingredient of AL methods is the AL function $\mathcal{L}_\mu: \mathcal{X} \times \mathbb{R}^m \rightarrow \mathbb{R}$, whose definition associated to (P) is

$$\mathcal{L}_\mu(x, y) := f(x) + \frac{1}{2\mu} \text{dist}_{\mathcal{C}}^2(c(x) + \mu y) - \frac{\mu}{2} \|y\|^2 \quad (9)$$

for some penalty parameter $\mu > 0$ and multiplier estimate $y \in \mathbb{R}^m$. This is a *partial* AL function in that it does not relax the simple constraint $x \in \mathcal{X}$, which is kept explicit in each subproblem. Notice that \mathcal{L} and \mathcal{L}_μ are smooth, with respect to both, primal and dual variables x and y , thanks to Assumption (A2) and convexity of \mathcal{C} . For later use, the partial derivatives of \mathcal{L}_μ read

$$\nabla_x \mathcal{L}_\mu(x, y) = \nabla f(x) + c'(x)^\top y_\mu(x, y), \quad \nabla_y \mathcal{L}_\mu(x, y) = c(x) - s_\mu(x, y) \quad (10)$$

where

$$s_\mu(x, y) := \text{proj}_{\mathcal{C}}(c(x) + \mu y), \quad y_\mu(x, y) := y + \frac{c(x) - s_\mu(x, y)}{\mu}. \quad (11)$$

Following the basic pattern of AL methods, Algorithm 3.1 proceeds by minimizing the AL function at each iteration, possibly inexactly and up to criticality, and updating the multiplier estimates and penalty parameters [3, Section 4.1]. Augmented Lagrangian subproblems require to

$$\text{minimize } \mathcal{L}_\mu(x, \hat{y}) \quad \text{over } x \in \mathcal{X} \quad (12)$$

given some $\mu > 0$ and $\hat{y} \in \mathbb{R}^m$. Feasibility of (12) follows from \mathcal{X} being nonempty, whereas well-posedness is due to (lower semi)continuity of $\mathcal{L}_\mu(\cdot, \hat{y})$ and is guaranteed if, e.g., \mathcal{X} is compact or f is bounded from below in \mathcal{X} . In fact, the existence of subproblem solutions is often just assumed [3, Assumption 6.1]. This issue could be circumvented by complementing the AL subproblems (12) with a localizing constraint, e.g., of trust region type [6, Remark 5.1]. Analogous in spirit to prox-boundedness [7], the following Assumption (A4) is weaker than typical coercivity or (level) boundedness assumptions but sufficient to yield well-posed subproblems.

Assumption 3.1. With regard to (P) and Algorithm 3.1,

- (A4) there exists $\bar{\mu} > 0$ such that for all $\mu \in (0, \bar{\mu}]$ and $\hat{y} \in \mathcal{Y}_s$ the function $\mathcal{L}_\mu(\cdot, \hat{y})$ is bounded from below over \mathcal{X} .

Although still unsatisfactory, demanding Assumption (A4) allows us to focus on the mixed-integer extension of generic AL methods to address MINLP.

Algorithm 3.1: Abstract safeguarded augmented Lagrangian method for (P)

Input: $\mu_0 \in (0, \bar{\mu}]$, $\varepsilon_0, \eta_0 > 0$, $\kappa_\mu, \theta_\mu \in (0, 1)$, $\mathcal{Y}_s \subseteq \mathbb{R}^m$ bounded

- 1 **for** $j = 0, 1, 2 \dots$ **do**
- 2 Select $\hat{y}^j \in \mathcal{Y}_s$
- 3 Find an ε_j -critical point x^j for $\mathcal{L}_{\mu_j}(\cdot, \hat{y}^j)$ over \mathcal{X}
- 4 Set $z^j \leftarrow \text{proj}_{\mathcal{C}}(c(x^j) + \mu_j \hat{y}^j)$, $v^j \leftarrow c(x^j) - z^j$, and $y^j \leftarrow \hat{y}^j + \mu_j^{-1} v^j$
- 5 **if** $j = 0$ **or** $\|v^j\| \leq \max\{\eta_j, \theta_\mu \|v^{j-1}\|\}$ **then**
- 6 set $\mu_{j+1} \leftarrow \mu_j$, **else** select $\mu_{j+1} \in (0, \kappa_\mu \mu_j)$
- 7 Select $\varepsilon_{j+1}, \eta_{j+1} \geq 0$ such that $\{\varepsilon_j\}, \{\eta_j\} \rightarrow 0$

The scheme outlined in Algorithm 3.1 is often referred to as *safeguarded* because the multiplier estimates \hat{y} are not allowed to grow too fast compared to the penalty parameter μ [3, 21, 20, 7]. In particular, it is required that $\|\mu_j \hat{y}^j\| \rightarrow 0$ as $\mu_j \rightarrow 0$, so that stronger global convergence properties can be attained. As a simple mechanism to ensure this property, multiplier estimates \hat{y} in Algorithm 3.1 are drawn from a bounded set $\mathcal{Y}_s \subseteq \mathbb{R}^m$. The dual safeguarding set \mathcal{Y}_s can be a generic hyperbox or can be tailored to the constraint set \mathcal{C} at hand [20]—see Section 4.1.

Subproblems (12) can be solved up to approximate criticality: given ε_j , at Step 3 we seek an ε_j -critical point $x^j \in \mathcal{X}$ for $\mathcal{L}_{\mu_j}(\cdot, \hat{y}^j)$, in the sense of Definition 2.2. For this task one can employ the mixed-integer linearization algorithm of [5], with guarantee of finite termination under Assumptions (A1)–(A4). Although the trust region radius Δ_j associated to the ε_j -criticality certificate does not need to be computed, it will be considered formally for the theoretical analysis. Given a (possibly inexact, first-order) solution x to (12), the dual update rule at Step 4 is designed toward the identity

$$\nabla_x \mathcal{L}_\mu(x, \hat{y}) = \nabla f(x) + c'(x)^\top y = \nabla_x \mathcal{L}(x, y), \quad (13)$$

as usual in AL methods. This allows to monitor the (outer) convergence with the (inner) subproblem tolerance; cf. Lemma 3.2 below.

Finally, Steps 5 to 7 are dedicated to monitoring primal feasibility (namely the conditions involving z^k in Definition 2.4) and updating the penalty parameter μ accordingly. Note that considering a sequence of primal tolerances $\{\eta_j\}$ allows to monitor primal convergence from a global perspective, slightly relaxing in fact other classical update rules [3, 6].

3.1 Convergence analysis

We begin our asymptotic analysis by collecting useful properties to characterize the iterations generated by Algorithm 3.1.

Lemma 3.2. *Let Assumptions (A1)–(A4) hold for (P) and consider the iterates of Algorithm 3.1. Then, for each $j \in \mathbb{N}$, Step 3 is well-posed and the iterates satisfy $x^j \in \mathcal{X}$, $z^j \in \mathcal{C}$, $y^j \in \mathcal{N}_{\mathcal{C}}(z^j)$, $\nabla_x \mathcal{L}_{\mu_j}(x^j, \hat{y}^j) = \nabla_x \mathcal{L}(x^j, y^j)$, and there exists some $\Delta_j > 0$ such that $\Psi_{\mathcal{L}(\cdot, y^j), \mathcal{X}}(x^j, \Delta_j) \leq \varepsilon_j$.*

Proof. Well-definedness of Algorithm 3.1 follows from the existence of solutions to the AL subproblems, which in turn is due to the standing Assumptions (A1)–(A4). In particular, the

feasible set \mathcal{X} is nonempty and closed, and the continuous real-valued cost function $\mathcal{L}_{\mu_j}(\cdot, \widehat{y}^j)$ is lower bound over \mathcal{X} , since $\mu_j \leq \bar{\mu}$, for all $j \in \mathbb{N}$.

Then, it is apparent that $x^j \in \mathcal{X}$ and $z^j \in \mathcal{C}$ for each $j \in \mathbb{N}$. Moreover, the assignments at [Step 4](#) gives that $z^j := \text{proj}_{\mathcal{C}}(c(x^j) + \mu_j \widehat{y}^j) = c(x^j) + \mu_j \widehat{y}^j - \mu_j y^j$, which is equivalent to $y^j \in \mathcal{N}_{\mathcal{C}}(z^j)$ by [\(2\)](#) and convexity of \mathcal{C} . By construction [\(13\)](#), the dual update rule readily yields $\nabla_x \mathcal{L}_{\mu_j}(x^j, \widehat{y}^j) = \nabla_x \mathcal{L}(x^j, y^j)$, and so the upper bound on the criticality measure and the existence of a suitable Δ_j follow from [Step 3](#). \square

We now turn to investigating properties of accumulation points, assuming their existence (which may follow from coercivity or level boundedness arguments). The following convergence results for [Algorithm 3.1](#) provides fundamental theoretical support for the numerical approach of [\[15\]](#). With [Theorem 3.3](#) we establish that feasible accumulation points of $\{x^j\}$ are AKKT-critical; see [\[7, Thm 3.3\]](#), [\[6, Thm 3.6\]](#) for analogous results.

Theorem 3.3. *Let Assumptions (A1)–(A4) hold. Consider a sequence $\{x^j\}$ generated by [Algorithm 3.1](#). Let x^* be an accumulation point of $\{x^j\}$ and $\{x^j\}_{j \in J}$ a subsequence such that $x^j \rightarrow_J x^*$. If x^* is feasible for (P), then x^* is AKKT-critical for (P).*

Proof. It is implicitly assumed [Algorithm 3.1](#) generates an infinite sequence of iterates $\{x^j\}$ with accumulation point x^* . Now we claim that the subsequences $\{x^j\}_{j \in J}$, $\{y^j\}_{j \in J}$, $\{z^j\}_{j \in J}$, $\{\Delta^j\}_{j \in J}$ satisfy the properties in [Definition 2.4](#), thus showing that x^* is AKKT-critical for (P). From [\(4\)](#) and [Lemma 3.2](#) we have that for all $j \in \mathbb{N}$

$$0 \leq \Psi_{\mathcal{L}(\cdot, y^j), \mathcal{X}}(x^j, \Delta_j) \leq \varepsilon_j$$

for some $\Delta_j > 0$. Hence, dual feasibility holds asymptotically owing to $\varepsilon_j \rightarrow 0$.

By assumption we have $x^j \rightarrow_J x^*$ with x^* feasible for (P), namely $x^* \in \mathcal{X}$ and $c(x^*) \in \mathcal{C}$. [Lemma 3.2](#) implies also that $y^j \in \mathcal{N}_{\mathcal{C}}(z^j)$ for each $j \in \mathbb{N}$. Finally, to demonstrate that $c(x^j) - z^j \rightarrow_J 0$ we consider two cases:

- If $\{\mu_j\}$ is bounded away from zero, the conditions at [Steps 5 and 6](#) of [Algorithm 3.1](#) and the construction of $\{\eta_j\}$ imply that $\|v^j\| := \|c(x^j) - z^j\| \rightarrow 0$, hence the assertion.
- If $\mu_j \rightarrow 0$, we exploit continuity of g , boundedness of $\{\widehat{y}^j\} \subseteq \mathcal{Y}_s$, feasibility of x^* , and closedness of \mathcal{C} . Combining these properties gives $c(x^j) + \mu_j \widehat{y}^j \rightarrow_J c(x^*) \in \mathcal{C}$ as $x^j \rightarrow_J x^*$. Therefore, $z^j \rightarrow_J c(x^*)$ as well, hence $c(x^j) - z^j \rightarrow_J 0$.

Overall, this proves that x^* is AKKT-critical for (P). \square

In contrast with global methods [\[3, Chapter 5\]](#), [\[8, Section 4.2\]](#), adopting affordable solvers for addressing [\(12\)](#) at [Step 3](#) impedes to guarantee that, in general, accumulation points are feasible or (globally) minimize an infeasibility measure. Thus, despite feasibility granted by Assumption (A1), [Algorithm 3.1](#) may not approach feasible points. In practice, however, for any fixed $\mu > 0$ and $\widehat{y} \in \mathbb{R}^m$, the AL subproblem [\(12\)](#) is equivalent to

$$\text{minimize} \quad \mu f(x) + \frac{1}{2} \text{dist}_{\mathcal{C}}^2(c(x) + \mu \widehat{y}) \quad \text{over} \quad x \in \mathcal{X}.$$

Hence, one can expect to find at least critical points of an infeasibility measure, as attested by the following result. Notice that this property requires mere boundedness of $\{\varepsilon_j\}$; cf. [\[3, Thm 6.3\]](#), [\[6, Proposition 3.7\]](#).

Theorem 3.4. *Let Assumptions (A1)–(A4) hold. Consider a sequence $\{x^j\}$ generated by Algorithm 3.1 with $\{\varepsilon_j\}$ merely bounded. Let x^* be an accumulation point of $\{x^j\}$ and $\{x^j\}_{j \in J}$ a subsequence such that $x^j \rightarrow_J x^*$. Then, x^* is a critical point for the feasibility problem*

$$\text{minimize } \mathcal{F}(x) := \frac{1}{2} \text{dist}_{\mathcal{C}}^2(c(x)) \quad \text{over } x \in \mathcal{X}.$$

Proof. It is implicitly assumed that Algorithm 3.1 generates an infinite sequence of iterates $\{x^j\}$ with accumulation point x^* . If $\{\mu_j\}$ is bounded away from zero, the conditions at Steps 5 and 6 of Algorithm 3.1 and the construction of $\{\eta_j\}$ imply that $\|v^j\| := \|c(x^j) - z^j\| \rightarrow 0$. By the upper bound $\|v^j\| \geq \text{dist}_{\mathcal{C}}(c(x^j))$ for each $j \in \mathbb{N}$, since $z^j \in \mathcal{C}$, taking the limit $j \rightarrow \infty$ yields $c(x^*) \in \mathcal{C}$ by continuity. Then, since $x^j \in \mathcal{X}$ for all $j \in \mathbb{N}$ and \mathcal{X} is closed, x^* is feasible for (P). Thus, x^* is a global minimizer of the feasibility problem and, by continuous differentiability of the objective function therein, x^* is critical for the feasibility problem.

Let us focus now on the case where $\{\mu_j\} \searrow 0$ and $x^* \in \mathcal{X}$ is infeasible for (P). First, we express what criticality entails for the feasibility problem above: a point $\bar{x} \in \mathbb{R}^n$ is critical if $\bar{x} \in \mathcal{X}$ and there exists some $\Delta > 0$ such that $\Psi_{\mathcal{F}, \mathcal{X}}(\bar{x}, \Delta) = 0$. Now, owing to (4) and Step 3, for all $j \in \mathbb{N}$ it is

$$\varepsilon_j \geq \Psi_{\mathcal{L}_{\mu_j}(\cdot, \hat{y}^j), \mathcal{X}}(x^j, \Delta_j) = \max_{w \in \mathcal{X} \cap \mathbb{B}_{\text{LP}}(x^j, \Delta_j)} \langle \nabla_x \mathcal{L}_{\mu_j}(x^j, \hat{y}^j), x^j - w \rangle \geq 0.$$

Multiplying by $\mu_j > 0$, by boundedness of $\{\varepsilon_j\}$ we have

$$0 \leq \max_{w \in \mathcal{X} \cap \mathbb{B}_{\text{LP}}(x^j, \Delta_j)} \langle \mu_j \nabla_x \mathcal{L}_{\mu_j}(x^j, \hat{y}^j), x^j - w \rangle \leq \mu_j \varepsilon_j \rightarrow 0.$$

Observing that $\mu_j \nabla_x \mathcal{L}_{\mu_j}(\cdot, \hat{y}^j)$ is locally Lipschitz continuous for all $\mu_j > 0$ by Assumption (A2), we have by [5, Lemma 3.5] and $x^j \rightarrow_J x^*$ that $\{\Delta_j\}_{j \in J}$ remains bounded away from zero. Furthermore, using $\{\mu_j\} \searrow 0$ yields

$$\mu_j \nabla_x \mathcal{L}_{\mu_j}(x^j, \hat{y}^j) \rightarrow_J c'(x^*)^\top [c(x^*) - \text{proj}_{\mathcal{C}}(c(x^*))] = \nabla \mathcal{F}(x^*)$$

by boundedness of $\{\hat{y}^j\}$ and $\{\nabla f(x^j)\}_{j \in J}$, the latter due to $x^j \rightarrow_J x^*$. Overall, taking the limit $j \rightarrow_J \infty$, we have that

$$\begin{aligned} 0 &= \lim_{j \rightarrow \infty} \max_{w \in \mathcal{X} \cap \mathbb{B}_{\text{LP}}(x^j, \Delta_j)} \langle \mu_j \nabla_x \mathcal{L}_{\mu_j}(x^j, \hat{y}^j), x^j - w \rangle \\ &= \max_{w \in \mathcal{X} \cap \mathbb{B}_{\text{LP}}(x^*, \Delta_*)} \langle \nabla \mathcal{F}(x^*), x^* - w \rangle = \Psi_{\mathcal{F}, \mathcal{X}}(x^*, \Delta_*) \end{aligned}$$

for some $\Delta_* > 0$, proving the result. \square

So far the focus has been on Algorithm 3.1, but how do these developments affect other numerical approaches for (P)? Being part of the AL framework, the scheme analysed in [11] can be naturally extended to handle MINLP. Its peculiarity is that, starting with a feasible point, convergence to feasible accumulation points can be guaranteed, thanks to a reset mechanism. Results similar to Theorems 3.3 and 3.4 can be readily obtained for this method too. Indeed, analogous findings seem to extend far beyond the penalty scheme considered in Section 3, possibly applying for a broad class of sequential minimization algorithms [13]. Although drawn in a different context, the arguments in [10, Section 4] give a valid proof pattern for interior point (or barrier) methods, among others.

For illustrative purposes, let us consider the special case of (P) with $\mathcal{C} := \mathbb{R}_+^m$. Introducing a barrier function $b: (0, \infty) \rightarrow \mathbb{R}$ to approximate the indicator $\delta_{\mathcal{C}}$, e.g., the classical log barrier

$b: t \mapsto -\log(t)$, and a barrier parameter $\beta > 0$ to control this approximation, one formulates a barrier subproblem—resembling (12)—of the form

$$\text{minimize } f(x) + \beta \sum_{i=1}^m b(c_i(x)) \quad \text{over } x \in \mathcal{X}.$$

Then, a sequence of subproblems is solved, possibly inexactly and up to criticality, with decreasing barrier parameters. Let us denote by x^j an ε_j -critical point for the barrier subproblem with parameter $\beta_j > 0$. Though with the drawback of requiring a strictly feasible point to start with (namely $x \in \mathcal{X}$, $c(x) < 0$), at every iteration it must be that $x^j \in \mathcal{X}$ and $c(x^j) < 0$, that is, this barrier scheme maintains (strict) feasibility. Moreover, echoing [Theorem 3.3](#), it is easy to show that, with $\beta_j, \varepsilon_j \rightarrow 0$, accumulation points of $\{x^j\}$ are AKKT-critical for (P); see [10, Thm 16].

4 Further characterizations

We now enrich the theoretical framework with results and interpretations beyond those motivated by [15], turning our attention to optimality conditions, Lagrangian duality, saddle point properties, and relationships with the classical proximal point algorithm.

For simplicity, we consider an optimization problem of the form (P) with $\mathcal{C} := \mathcal{K}$ a nonempty closed convex cone. Inspired by [21, Section 8.4], this assumption greatly simplifies the presentation thanks to the identity

$$\delta_{\mathcal{K}}^*(y) = \sup_{z \in \mathcal{K}} \langle z, y \rangle = \begin{cases} 0 & \text{if } y \in \mathcal{K}^\circ, \\ \infty & \text{otherwise} \end{cases} = \delta_{\mathcal{K}^\circ}(y), \quad (14)$$

which connects the indicator $\delta_{\mathcal{K}}: \mathbb{R}^m \rightarrow \mathbb{R} \cup \{\infty\}$ of a set $\mathcal{K} \subseteq \mathbb{R}^m$, the conjugate function $h^*: \mathbb{R}^m \rightarrow \mathbb{R} \cup \{\infty\}$ associated with a (proper and lower semicontinuous) function $h: \mathbb{R}^m \rightarrow \mathbb{R} \cup \{\infty\}$ [1, Definition 13.1], and the polar cone $\mathcal{K}^\circ \subseteq \mathbb{R}^m$ of a subset \mathcal{K} of \mathbb{R}^m [1, Definition 6.22], respectively

$$h^*(v) := \sup_{z \in \mathbb{R}^m} \{\langle z, v \rangle - h(z)\} \quad \text{and} \quad \mathcal{K}^\circ := \left\{ u \in \mathbb{R}^m \mid \sup_{v \in \mathcal{K}} \langle v, u \rangle \leq 0 \right\}.$$

4.1 Lagrangian duality

The necessary optimality conditions in [Definition 2.3](#) cannot be derived based on the Lagrangian function \mathcal{L} alone, but additional insights on the problem are needed to setup the complementarity system encapsulated in the expression $y \in \mathcal{N}_{\mathcal{K}}(c(x))$. Instead, a comprehensive first-order optimality analysis can be developed based on the *generalized* Lagrangian function, whose construction is briefly recalled following [17, 6, 8]. Introducing an auxiliary variable $s \in \mathbb{R}^m$, (P) can be rewritten as

$$\begin{aligned} & \text{minimize } f(x) && (\text{P}^S) \\ & \text{over } x \in \mathcal{X}, s \in \mathcal{K} \\ & \text{subject to } c(x) - s = 0, \end{aligned}$$

whose (classical) Lagrangian function, akin to (5), reads

$$\mathcal{L}^S(x, s, y) := f(x) + \langle y, c(x) - s \rangle.$$

Marginalization of \mathcal{L}^S with respect to s yields the generalized Lagrangian function $\ell: \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ associated to (P), given by

$$\ell(x, y) := \inf_{s \in \mathcal{K}} \mathcal{L}^S(x, s, y) = f(x) + \langle y, c(x) \rangle + \inf_s \{ \delta_{\mathcal{K}}(s) - \langle y, s \rangle \} = \mathcal{L}(x, y) - \delta_{\mathcal{K}}^*(y).$$

Then, observing the identity (14), the dual domain of ℓ , namely the set \mathcal{Y} of valid multipliers, is given by

$$\mathcal{Y} := \mathbb{R}^m \cap \text{dom } \delta_{\mathcal{K}}^* = \text{dom } \delta_{\mathcal{K}^\circ} = \mathcal{K}^\circ, \quad (15)$$

which corresponds to a nonempty closed convex cone in \mathbb{R}^m . Classical nonlinear programming is recovered by (neglecting integrality and) taking \mathcal{K} to be the standard constraint cone there: $\mathcal{K} := \{0\}$ and $\mathcal{K} := \mathbb{R}_-^m$ are associated respectively to $\mathcal{Y} := \mathbb{R}^m$ and $\mathcal{Y} := \mathbb{R}_+^m$. Then, with this insight about the dual domain, a sound yet simple stratagem for providing a safeguarding set to Algorithm 3.1 is to set $\mathcal{Y}_s := \mathcal{Y} \cap [-y_{\max}, y_{\max}]^m$ for some large $y_{\max} > 0$ [20, Section 3.1].

In contrast with the (classical) Lagrangian \mathcal{L} , the emergence of dual information from the generalized Lagrangian ℓ allows not only to obtain dual estimates tailored to \mathcal{K} , but also to express primal-dual first-order optimality conditions without direct access to (P). It is shown in [17], [8, Remark 3.5] that the generalized Lagrangian function ℓ is sufficient to write necessary optimality conditions for (P) when $\mathcal{X} = \mathbb{R}^n$ and \mathcal{K} is convex. These read

$$0 \in \partial_x \ell(x, y) \quad \text{and} \quad 0 \in \partial_y (-\ell)(x, y), \quad (16)$$

where the negative sign highlights the (generalized) saddle-point property of the primal-dual system. But how does (16) relate to Definition 2.3? Owing to the identity $\nabla_x \mathcal{L} = \nabla_x \ell$, the first criticality condition $\Psi_{\mathcal{L}(\cdot, y), \mathcal{X}}(x, \Delta) = 0$ in Definition 2.3 captures in fact an extension of $0 = \nabla_x \ell(x, y)$ to accommodate the mixed-integer linear constraint set \mathcal{X} . Inspired by the descent-ascent motive behind (16), the main definition we will use below is the following, with a character of primal-dual symmetry.

Definition 4.1 (Local saddle point). A pair $(x, y) \in \mathcal{X} \times \mathcal{Y}$ is called a *local saddle point* of $\mathcal{L}: \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ if

$$\Psi_{\mathcal{L}(\cdot, y), \mathcal{X}}(x, \Delta) = 0 \quad \text{and} \quad \Psi_{-\mathcal{L}(x, \cdot), \mathcal{Y}}(y, \Delta) = 0$$

for some $\Delta > 0$.

Theorem 4.2. Consider (P) and let $x \in \mathbb{R}^n$, $y \in \mathbb{R}^m$ be arbitrary but fixed. Then the following assertions are equivalent:

- (i) x is KKT-critical with multiplier y ;
- (ii) (x, y) is a local saddle point of \mathcal{L} .

Proof. Since both KKT-critical and local saddle points demand that $x \in \mathcal{X}$ and $\Psi_{\mathcal{L}(\cdot, y), \mathcal{X}}(x, \Delta) = 0$ holds for some $\Delta > 0$, it remains to consider the second part of Definitions 2.3 and 4.1, namely the equivalence of $y \in \mathcal{N}_{\mathcal{K}}(c(x))$ and $\Psi_{-\mathcal{L}(x, \cdot), \mathcal{Y}}(y, \Delta) = 0$. We proceed by deriving a sequence of identities. Observing that

$$0 = \Psi_{-\mathcal{L}(x, \cdot), \mathcal{Y}}(y, \Delta) = \max_{w \in \mathcal{Y} \cap \mathbb{B}_{\text{LP}}(y, \Delta)} \langle -\nabla_y \mathcal{L}(x, y), y - w \rangle \geq 0$$

can be rewritten with a universal quantifier as

$$\forall w \in \mathcal{Y} \cap \mathbb{B}_{\text{LP}}(y, \Delta): \langle -\nabla_y \mathcal{L}(x, y), y - w \rangle = \langle y + \nabla_y \mathcal{L}(x, y) - y, w - y \rangle \leq 0,$$

the characterization (1) of projections onto convex sets yields

$$y = \text{proj}_{\mathcal{Y} \cap \mathbb{B}_{\text{LP}}(y, \Delta)} (y + \nabla_y \mathcal{L}(x, y)).$$

Since all variables in y are real-valued and the ball $\mathbb{B}_{\text{LP}}(y, \Delta)$ is compact convex and centered at $y \in \mathcal{Y}$, the previous identity is equivalent to $y = \text{proj}_{\mathcal{Y}} (y + \nabla_y \mathcal{L}(x, y))$ for all $\Delta > 0$. Using the property (2) of normal cones and the partial derivative of \mathcal{L} in (5), we obtain $\nabla_y \mathcal{L}(x, y) = c(x) \in \mathcal{N}_{\mathcal{Y}}(y)$. Exploiting now the definition of \mathcal{Y} (15), the polar-conjugacy relation (14) implies that $c(x) \in \partial \delta_{\mathcal{K}^\circ}(y) = \partial \delta_{\mathcal{K}}^*(y)$. Finally, owing to [18, Proposition 11.3], this is equivalent to $y \in \partial \delta_{\mathcal{K}}(c(x)) = \mathcal{N}_{\mathcal{K}}(c(x))$, which also implies the inclusion $c(x) \in \mathcal{K}$, concluding the proof. \square

4.2 Saddle points of the augmented Lagrangian

Inspired by the primal-dual characterization of KKT-critical points in Section 4.1, here we show that KKT-criticality for (P) is also associated to a local saddle point property of the *augmented* Lagrangian function. This trait, recently re-investigated by Rockafellar [17] for a broad problem class, allows to interpret the update rule at Step 4 as a dual gradient ascent step for the augmented Lagrangian, thus making Algorithm 3.1 a primal descent, dual ascent method; see also [21, Section 8.1].

We begin with some preliminary observations.

Lemma 4.3. *Consider (P) and let $x \in \mathcal{X}$, $y \in \mathbb{R}^m$, and $\Delta, \mu > 0$ be arbitrary but fixed. Then the following assertions are equivalent:*

- (i) $y \in \mathcal{N}_{\mathcal{K}}(c(x))$;
- (ii) $\nabla_y \mathcal{L}_\mu(x, y) = 0$;
- (iii) $\Psi_{-\mathcal{L}_\mu(x, \cdot), \mathcal{Y}}(y, \Delta) = 0$.

In particular, these conditions imply the inclusions $c(x) \in \mathcal{K}$ and $y \in \mathcal{Y}$.

Proof. Owing to (10), condition (ii) can be rewritten as $c(x) = \text{proj}_{\mathcal{K}}(c(x) + \mu y)$ and, since $\mu > 0$, property (2) implies the equivalence of (i) and (ii). Now, patterning the proof of Theorem 4.2, we obtain that (iii) is equivalent to $\nabla_y \mathcal{L}_\mu(x, y) \in \mathcal{N}_{\mathcal{Y}}(y)$. Then, the implication (ii) \implies (iii) is clear, and it remains to focus on the converse one.

Let us consider now the maximization of $\mathcal{L}_\mu(x, \cdot)$ over \mathbb{R}^m , that is, dropping the restriction to \mathcal{Y} —as well as the trust region in (4). Then, any (unconstrained) solution $\tilde{y} \in \mathbb{R}^m$ necessarily satisfies $\nabla_y \mathcal{L}_\mu(x, \tilde{y}) = 0$, which is equivalent to $\tilde{y} \in \mathcal{N}_{\mathcal{K}}(c(x))$ by combining (10)–(11) and (2). Furthermore, owing to convexity of \mathcal{K} and [18, Proposition 11.3], this inclusion coincides with $c(x) \in \mathcal{N}_{\mathcal{K}^\circ}(\tilde{y})$, meaning in particular that $\tilde{y} \in \mathcal{K}^\circ = \mathcal{Y}$ by (15). Thus, since the unconstrained optimum \tilde{y} satisfies in fact the restriction to \mathcal{Y} , it is optimal for the constrained problem too. Indeed, by convexity of \mathcal{Y} , \tilde{y} remains optimal also considering a trust region $\mathbb{B}_{\text{LP}}(\tilde{y}, \Delta)$, for any $\Delta > 0$, thus showing that (iii) \implies (ii).

Finally, the inclusions follow respectively from the normal cone $\mathcal{N}_{\mathcal{K}}(c(x))$ being nonempty in (i) and from the restriction $y \in \mathcal{Y}$ in (4) for (iii). \square

The following is the main result of this section.

Theorem 4.4. Consider (P) and let $x \in \mathbb{R}^n$, $y \in \mathbb{R}^m$ be arbitrary but fixed. Then the following assertions are equivalent:

- (i) x is KKT-critical with multiplier y ;
- (ii) (x, y) is a local saddle point of \mathcal{L}_μ for some $\mu > 0$;
- (iii) (x, y) is a local saddle point of \mathcal{L}_μ for all $\mu > 0$.

Proof. We prove the equivalence via a loop of implications. Note that (iii) \implies (ii) is straightforward.

(ii) \implies (i) Let (x, y) be a local saddle point of \mathcal{L}_μ for some $\mu > 0$. Then Lemma 4.3 implies that $c(x) \in \mathcal{K}$ and $y \in \mathcal{N}_{\mathcal{K}}(c(x))$. Therefore, by combining with (10)–(11) and properties (1)–(2), we obtain the identity

$$\nabla_x \mathcal{L}_\mu(x, y) = \nabla f(x) + c'(x)y = \nabla_x \mathcal{L}(x, y). \quad (17)$$

Therefore, since $\Psi_{\mathcal{L}_\mu(\cdot, y), \mathcal{X}}(x, \Delta) = 0$ holds for some $\Delta > 0$, it must be also $\Psi_{\mathcal{L}(\cdot, y), \mathcal{X}}(x, \Delta) = 0$. Thus, x is KKT-critical for (P) with multiplier y .

(i) \implies (iii) Let $\mu > 0$ be arbitrary but fixed and x a KKT-critical point with multiplier y . Then, $c(x) \in \mathcal{K}$ and $y \in \mathcal{N}_{\mathcal{K}}(c(x))$ hold owing to KKT-criticality. Hence, on the one hand, Lemma 4.3 implies that the second equality in Definition 4.1 is satisfied. On the other hand, this furnishes again (17), and thus KKT-criticality of (x, y) yields $\Psi_{\mathcal{L}_\mu(\cdot, y), \mathcal{X}}(x, \Delta) = \Psi_{\mathcal{L}(\cdot, y), \mathcal{X}}(x, \Delta) = 0$. With $\mu > 0$ being arbitrary, this shows that (x, y) is a local saddle point of \mathcal{L}_μ for all $\mu > 0$. \square

4.3 Relationship with Proximal Point Methods

Connections of augmented Lagrangian methods with duality and the proximal point algorithm (PPA) have been recently discussed in Hilbert spaces [21, Section 8.4] and explored in the broad setting of generalized nonlinear programming [17]. We turn now to exploring these properties in the context of MINLP. Considering (P), the associated Lagrangian function (5), and the dual domain \mathcal{Y} (15), we define for all $y \in \mathcal{Y}$

$$Q(y) := \inf_{x \in \mathcal{X}} \mathcal{L}(x, y) = \inf_{x \in \mathcal{X}} \{f(x) + \langle y, c(x) \rangle\}$$

so that the natural “dual” problem of (P) is given by

$$\text{maximize } Q(y) \quad \text{over } y \in \mathcal{Y}.$$

Note that Q is a concave function since it is an infimum of affine functions. Then, by convexity of \mathcal{Y} , the above is a concave maximization problem, that is, a convex minimization problem. The PPA consists in applying the recursion

$$y^{j+1} := \text{prox}_{-\nu_j Q}(y^j)$$

with parameter $\nu_j > 0$, where the central ingredient is the *proximal mapping* (or proximity operator) associated to the problem, given by

$$\text{prox}_{-\nu Q}(w) := \arg \min_{y \in \mathcal{Y}} \left\{ -Q(y) + \frac{1}{2\nu} \|y - w\|^2 \right\}$$

for any $\nu > 0$ [1, Chapter 24], [17, Section 2]. Note that the function occurring inside the arg min is strongly convex, hence it admits a unique minimizer, and thus the proximal mapping is well-defined and single-valued. We will demonstrate that this iterative procedure is strongly related to the augmented Lagrangian method, whose basic iteration with parameter $\mu_j > 0$ reads

$$x^{j+1} \in \arg \min_{x \in \mathcal{X}} \mathcal{L}_{\mu_j}(x, y^j), \quad z^{j+1} := \text{proj}_{\mathcal{K}}(c(x^{j+1}) + \mu_j y^j), \quad y^{j+1} := y^j + \frac{c(x^{j+1}) - z^{j+1}}{\mu_j},$$

where $z^{j+1} \in \mathcal{K}$ and $y^{j+1} \in \mathcal{N}_{\mathcal{K}}(z^{j+1})$ hold by construction; see Lemma 3.2.

The main result in this section is the following.

Theorem 4.5. *Consider (P) and let $w \in \mathbb{R}^m$, $\mu > 0$ be arbitrary but fixed. Let \bar{x} be a critical point for $\mathcal{L}_{\mu}(\cdot, w)$ over \mathcal{X} . Define the associated $\bar{s} := \text{proj}_{\mathcal{K}}(c(\bar{x}) + \mu w)$ and $\bar{y} := w + [c(\bar{x}) - \bar{s}]/\mu$. Then $\bar{y} = \text{prox}_{\mu Q}(w) \in \mathcal{Y}$ and $\bar{x} \in \mathcal{X}$ is a critical point for the infimum defining $Q(\bar{y})$, namely for $\mathcal{L}(\cdot, \bar{y})$ over \mathcal{X} .*

Proof. We prove the claim by showing that (\bar{x}, \bar{y}) is a local saddle point of the function

$$h: \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}, \quad h(x, y) := \mathcal{L}(x, y) - \frac{\mu}{2} \|y - w\|^2$$

which brings together the dual function Q with the quadratic proximal term. To verify this saddle property, note that the definition of \bar{x} and \bar{y} implies by (10)–(11) that

$$\nabla_x \mathcal{L}_{\mu}(\bar{x}, w) = \nabla f(\bar{x}) + c'(\bar{x})^\top \bar{y} = \nabla_x \mathcal{L}(\bar{x}, \bar{y}) = \nabla_x h(\bar{x}, \bar{y}).$$

Then, by Definition 2.2, there exists some $\Delta > 0$ such that

$$0 = \Psi_{\mathcal{L}_{\mu}(\cdot, w), \mathcal{X}}(\bar{x}, \Delta) = \Psi_{\mathcal{L}(\cdot, \bar{y}), \mathcal{X}}(\bar{x}, \Delta) = \Psi_{h(\cdot, \bar{y}), \mathcal{X}}(\bar{x}, \Delta),$$

hence \bar{x} is a critical point for $h(\cdot, \bar{y})$ over \mathcal{X} . On the other hand, $h(\bar{x}, \cdot)$ is a strictly concave quadratic function of the form

$$h(\bar{x}, \cdot): y \mapsto f(\bar{x}) + \langle y, c(\bar{x}) \rangle - \frac{\mu}{2} \|y - w\|^2 = -\frac{\mu}{2} \left\| y - w + \frac{c(\bar{x})}{\mu} \right\|^2 + c_{\circ},$$

where $c_{\circ} \in \mathbb{R}$ is a constant independent of y . Therefore, the unique maximizer \tilde{y} of $h(\bar{x}, \cdot)$ over the convex set \mathcal{Y} is determined by the necessary optimality condition $\nabla_y h(\bar{x}, \tilde{y}) \in \mathcal{N}_{\mathcal{Y}}(\tilde{y})$. Using the definition of h , (10)–(11), (15), and the identity (14), this can be rewritten as $c(\bar{x}) + \mu(w - \tilde{y}) \in \mathcal{N}_{\mathcal{K}^{\circ}}(\tilde{y}) = \partial \delta_{\mathcal{K}}^*(\tilde{y})$. Then, by convexity of \mathcal{K} and [18, Proposition 11.3], this is equivalent to $\tilde{y} \in \mathcal{N}_{\mathcal{K}}(c(\bar{x}) + \mu(w - \tilde{y}))$. Finally, the definition of \bar{s} and characterization (2) yield the identity

$$\bar{s} := \text{proj}_{\mathcal{K}}(c(\bar{x}) + \mu w) = c(\bar{x}) + \mu(w - \tilde{y}),$$

showing that the unique maximizer \tilde{y} coincides in fact with \bar{y} , concluding the proof. \square

5 Concluding Remarks

The results in this paper offer solid theoretical foundations for employing continuous optimization techniques to address mixed-integer nonlinear programming, at least as principled heuristics. It remains an open question how to overcome Assumption (A3). When localizing both real- and integer-valued variables, enough freedom should be left for the latter, but not necessarily for the former. In particular, one should prevent that some integers become effectively fixed, leading to weaker optimality conditions. Another topic for future research concerns the possibility to waive Assumption (A4) without interfering with the convergence guarantees of available methods.

References

- [1] Heinz H. Bauschke and Patrick L. Combettes. *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*. Springer, Cham, CH, 2017.
- [2] Pietro Belotti, Christian Kirches, Sven Leyffer, Jeff Linderoth, James Luedtke, and Ashutosh Mahajan. Mixed-integer nonlinear optimization. *Acta Numerica*, 22:1–131, 2013.
- [3] Ernesto G. Birgin and José M. Martínez. *Practical Augmented Lagrangian Methods for Constrained Optimization*. Society for Industrial and Applied Mathematics, Philadelphia, PA, US, 2014.
- [4] Richard H. Byrd, Nicholas I. M. Gould, Jorge Nocedal, and Richard A. Waltz. On the convergence of successive linear-quadratic programming algorithms. *SIAM Journal on Optimization*, 16(2):471–489, 2005.
- [5] Alberto De Marchi. Mixed-integer linearity in nonlinear optimization: a trust region approach. *arXiv:2310.17285*, 2023.
- [6] Alberto De Marchi. Implicit augmented Lagrangian and generalized optimization. *Journal of Applied and Numerical Optimization*, 6(2):291–320, 2024.
- [7] Alberto De Marchi, Xiaoxi Jia, Christian Kanzow, and Patrick Mehlitz. Constrained composite optimization and augmented Lagrangian methods. *Mathematical Programming*, 201(1):863–896, 2023.
- [8] Alberto De Marchi and Patrick Mehlitz. Local properties and augmented Lagrangians in fully nonconvex composite optimization. *Journal of Nonsmooth Analysis and Optimization*, 2024.
- [9] Alberto De Marchi and Andreas Themelis. Proximal gradient algorithms under local Lipschitz gradient continuity. *Journal of Optimization Theory and Applications*, 194(3):771–794, 2022.
- [10] Alberto De Marchi and Andreas Themelis. An interior proximal gradient method for nonconvex optimization. *Open Journal of Mathematical Optimization*, 2024.
- [11] Brecht Evens, Puya Latafat, Andreas Themelis, Johan Suykens, and Panagiotis Patrinos. Neural network training as an optimal control problem: An augmented lagrangian approach. In *2021 60th IEEE Conference on Decision and Control (CDC)*, pages 5136–5143, 2021.
- [12] Oliver Exler, Thomas Lehmann, and Klaus Schittkowski. A comparative study of SQP-type algorithms for nonlinear and nonconvex mixed-integer optimization. *Mathematical Programming Computation*, 4(4):383–412, 2012.
- [13] Anthony V. Fiacco and Garth P. McCormick. *Nonlinear Programming: Sequential Unconstrained Minimization Techniques*. Wiley, New York, NY, US, 1968.
- [14] Johannes Jahn and Martin Knossalla. Lagrange theory of discrete-continuous nonlinear optimization. *Journal of Nonlinear and Variational Analysis*, 2(3):317–342, 2018.
- [15] Viktoriya Nikitina, Alberto De Marchi, and Matthias Gerdt. Hybrid optimal control with mixed-integer Lagrangian methods. *arXiv:2403.06842*, 2024.

- [16] Rien Quirynen and Stefano Di Cairano. Sequential quadratic programming algorithm for real-time mixed-integer nonlinear MPC. In *2021 60th IEEE Conference on Decision and Control (CDC)*, pages 993–999, Austin, TX, US, 2021.
- [17] R. Tyrrell Rockafellar. Convergence of augmented lagrangian methods in extensions beyond nonlinear programming. *Mathematical Programming*, 199(1):375–420, 2023.
- [18] R. Tyrrell Rockafellar and Roger J. B. Wets. *Variational Analysis*, volume 317. Springer, Heidelberg, DE, 1998.
- [19] Sebastian Sager, Michael Jung, and Christian Kirches. Combinatorial integral approximation. *Mathematical Methods of Operations Research*, 73(3):363–380, 2011.
- [20] Pantelis Sopasakis, Emil Fresk, and Panagiotis Patrinos. OpEn: Code generation for embedded nonconvex optimization. *IFAC-PapersOnLine*, 53(2):6548–6554, 2020.
- [21] Daniel Steck. *Lagrange Multiplier Methods for Constrained Optimization and Variational Problems in Banach Spaces*. PhD thesis, University of Würzburg, Würzburg, DE, 2018.