

Slow rates of approximation of U-statistics and V-statistics by quadratic forms of Gaussians

Kevin Han Huang and Peter Orbanz

Gatsby Unit, University College London

Abstract. We construct examples of degree-two U- and V-statistics of n i.i.d. heavy-tailed random vectors in $\mathbb{R}^{d(n)}$, whose ν -th moments exist for $\nu > 2$, and provide tight bounds on the error of approximating both statistics by a quadratic form of Gaussians. In the case $\nu = 3$, the error of approximation is $\Theta(n^{-1/12})$. The proof adapts a result of Huang, Austern and Orbanz [12] to U- and V-statistics. The lower bound for U-statistics is a simple example of the concept of variance domination used in [12].

1 Introduction

We consider distributional approximations of degree-two U- and V-statistics that are of the form

$$u_n(x_1, \dots, x_n) := \frac{\sum_{1 \leq i \neq j \leq n} k_u(x_i, x_j)}{n(n-1)} \quad \text{and} \quad v_n(x_1, \dots, x_n) := \frac{\sum_{1 \leq i, j \leq n} k_v(x_i, x_j)}{n^2}.$$

Here, $x_1, \dots, x_n \in \mathbb{R}^{d(n)}$ are elements of a high-dimensional Euclidean space, i.e. the dimension $d(n)$ may depend on n , and $k_u : \mathbb{R}^{d(n)} \times \mathbb{R}^{d(n)} \rightarrow \mathbb{R}$ and $k_v : \mathbb{R}^{d(n)} \times \mathbb{R}^{d(n)} \rightarrow \mathbb{R}$ are symmetric functions. These statistics arise frequently in the estimation of two-way interactions. Applications include gene-set testing [6], change-point detection [21] and kernel-based tests in machine learning [10].

Classically, with $d(n) = d$ fixed, it is well-known that the distributions of both statistics admit either a Gaussian limit or a sum-of-chi-squares limit (up to reweighting and centering of the sum), depending on a degeneracy condition [17]. Recently, the high-dimensional case, where $d(n) \rightarrow \infty$ as $n \rightarrow \infty$, has gained attention due to its relevance in machine learning applications. Recent work investigates (variants of) U-statistics that admit a Gaussian limit despite degeneracy [6, 7, 13, 18, 22], or a sum-of-chi-squares limit despite non-degeneracy [3, 11]. In all cases, the error of Gaussian approximation for both U- and V-statistics is known to be $O(n^{-1/2})$ [5], and is not improvable [2]. The behavior of non-Gaussian approximation is more nuanced: The known upper bounds depend on the number of non-zero eigenvalues of the Hilbert-Schmidt operator associated with the kernel u of the U-statistic, ranging from $O(n^{-1})$ for five non-zero eigenvalues [9], $O(n^{-1/12})$ for one non-zero eigenvalue [23] (and control of an 18/5th moment), to $O(n^{-1/14})$ without eigenvalue assumptions [11]. This note constructs a set of i.i.d. random vectors $X = (X_i)_{i \leq n}$ in $\mathbb{R}^{d(n)}$, whose ν -th moments exist for $\nu \in (2, 3]$, and functions k_u and k_v , such that the following error bounds hold for the approximation by a quadratic form of Gaussians.

Theorem 1. Fix $\nu \in (2, 3]$. Let χ_1^2 be a chi-squared random variable with 1 degree of freedom, $\xi \sim \mathcal{N}(0, 1)$ be independent of χ_1^2 and $\overline{\chi_1^2} = \chi_1^2 - 1$. There exist some constants $c, C > 0$, $N \in \mathbb{N}$ and a sequence $(\sigma_n)_{n \in \mathbb{N}}$ with $\sigma_n \rightarrow 0$, where X , k_u and k_v depend on σ_n , such that for all $n > N$ and $d(n) \in \mathbb{N}$,

$$cn^{-\frac{\nu-2}{4\nu}} \leq \sup_{t \in \mathbb{R}} \left| \mathbb{P}(\sqrt{n(n-1)} u_n(X) \leq t) - \mathbb{P}(\sigma_n \xi + \overline{\chi_1^2} \leq t) \right| \leq Cn^{-\frac{\nu-2}{4\nu}},$$

$$cn^{-\frac{\nu-2}{4\nu}} \leq \sup_{t \in \mathbb{R}} \left| \mathbb{P}(n v_n(X) \leq t) - \mathbb{P}(\sigma_n \xi + \chi_1^2 \leq t) \right| \leq Cn^{-\frac{\nu-2}{4\nu}}.$$

Remark 1. (i) When $\nu = 3$, Theorem 1 says that the approximation error is $\Theta(n^{-\frac{1}{12}})$. (ii) In the construction, we choose σ_n to decay as $\Theta(n^{-\frac{\nu-2}{2\nu}})$. The approximation becomes a chi-squared approximation in the limit $n \rightarrow \infty$, but at a very slow rate. (iii) Since ξ can be obtained as the limiting distribution of a partial sum of weighted and centred chi-squared variables, Theorem 1 can be read as a result on the approximation of $u_n(X)$ and $v_n(X)$ by infinite sums of weighted and shifted chi-squares. For further discussions on when a Gaussian component may emerge from an infinite sum of weighted chi-squares, see [3].

Theorem 1 is an instance of the broader theme of Gaussian universality: Consider a set of i.i.d. random vectors $Y = (Y_i)_{i \leq n}$ in \mathbb{R}^b and the collection of Gaussian vectors

$$Z = (Z_1, \dots, Z_n) \quad \text{where} \quad Z_i \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(\mathbb{E}[Y_1], \text{Var}[Y_1]).$$

The term Gaussian universality refers to the fact that, for a large class of functions $f : (\mathbb{R}^b)^n \rightarrow \mathbb{R}$, $f(Y)$ is close to $f(Z)$ in distribution as $n \rightarrow \infty$. This phenomenon can be observed even when the law of Y_1 is allowed to vary in n , and has found applications in probability, statistics and machine learning [1, 8, 14, 15, 19, 20]. Indeed, a crucial intermediate step in Theorem 1 is to write $u_n(X) = \tilde{u}_n(Y)$ and $v_n(X) = \tilde{v}_n(Y)$ for some intermediate functions \tilde{u}_n and \tilde{v}_n , and approximate them by $\tilde{u}_n(Z)$ and $\tilde{v}_n(Z)$ respectively. The lower bound of Theorem 1 adapts a degree- m polynomial $p_m^*(X)$ constructed in our recent work [12]—inspired by Senatov [16]—which has a slow universality approximation error on the order $\Omega(n^{-\frac{\nu-2}{2\nu m}})$. The upper bound of Theorem 1 improves upon our generic universality result in [12] by using an argument specific to $p_m^*(X)$, instead of applying Lindeberg’s technique. To extend these results to u_n and v_n , our proof also exploits a technique called variance domination in [12], which is summarised in Proposition 2 in Section 3.

An open question from [12] was whether the slow $n^{-1/12}$ universality approximation error holds also for degree-two U-statistics and V-statistics, which see more real-life applications. Yanushkevichiene [23] conjectures that the $n^{-1/12}$ rate is unimprovable for degree-two U-statistics in view of the construction by Senatov [16]. Theorem 1 answers this question in the affirmative. An implication is that, without additional structural assumptions on the data distribution or the kernel functions k_u and k_v , the slow $n^{-1/12}$ rate of quadratic-form-of-Gaussian approximation for U-statistics and V-statistics is not improvable. In Section 2, we provide the explicit constructions of k_u , k_v and X . All proofs are included in Section 4.

2 Construction of k_u , k_v and X

We first recall the lower bound construction from [12] in the case $m = 2$. The construction involves a polynomial $p_n^* : (\mathbb{R}^2)^n \rightarrow \mathbb{R}$ given by

$$p_n^*(y_1, \dots, y_n) := \frac{1}{\sqrt{n}} \sum_{i=1}^n y_{i1} + \left(\frac{1}{\sqrt{n}} \sum_{i=1}^n y_{i2} \right)^2 \quad \text{for } y_i = (y_{i1}, y_{i2}) \in \mathbb{R}^2.$$

The collection of \mathbb{R}^2 random vectors $Y = (Y_1, \dots, Y_n)$ are generated as follows. For a fixed $\sigma_0 > 0$ and $\nu \in (2, 3]$, write $\sigma_n = \min\{\sigma_0 n^{-(\nu-2)/2\nu}, 1\}$. Note that this is the sequence $\sigma_n \rightarrow 0$ in Theorem 1. Let U_{σ_n} be the discrete random variable supported at three points with

$$U_{\sigma_n} = \begin{cases} -6^{-1/2} \sigma_n^{-2/(\nu-2)} & \text{with probability } 2\sigma_n^{2\nu/(\nu-2)}, \\ 0 & \text{with probability } 1 - 3\sigma_n^{2\nu/(\nu-2)}, \\ 2 \times 6^{-1/2} \sigma_n^{-2/(\nu-2)} & \text{with probability } \sigma_n^{2\nu/(\nu-2)}. \end{cases}$$

U_{σ_n} is constructed such that it becomes increasingly heavy-tailed as $\sigma_n \rightarrow 0$. Now let $U_{1;\sigma_n}, \dots, U_{n;\sigma_n}$ be i.i.d. copies of U_{σ_n} . The i.i.d. random vectors $Y = (Y_i)_{i \leq n}$ are generated by

$$Y_{i;\sigma_n} := \left(\frac{1}{\sqrt{2}} U_{i;\sigma_n} + \frac{\sigma_n}{\sqrt{2}} \xi_{i1}, \xi_{i2} \right) \quad \text{where } \xi_{11}, \xi_{12}, \dots, \xi_{n1}, \xi_{n2} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1).$$

To adapt $p_n^*(Y)$ to our setup, we first observe that p_n^* can be rewritten as a V-statistic:

$$p_n^*(y_1, \dots, y_n) = \frac{1}{n} \sum_{i,j=1}^n \left(\frac{y_{i1}}{2\sqrt{n}} + \frac{y_{j1}}{2\sqrt{n}} + y_{i2} y_{j2} \right) = n \tilde{v}_n(y_1, \dots, y_n),$$

where we have defined, for $y_1, \dots, y_n \in \mathbb{R}^2$ and $a_1, a_2, b_1, b_2 \in \mathbb{R}$,

$$\tilde{v}_n(y_1, \dots, y_n) := \frac{1}{n^2} \sum_{i,j=1}^n \tilde{k}_v(y_i, y_j), \quad \tilde{k}_v((a_1, a_2), (b_1, b_2)) := \frac{a_1}{2\sqrt{n}} + \frac{b_1}{2\sqrt{n}} + a_2 b_2.$$

Moreover, since we have no restrictions on how $d(n)$ depends on n , there are non-unique choices of a function $\phi_{d(n)} : \mathbb{R}^{d(n)} \rightarrow \mathbb{R}$ and a probability measure $\mu_{d(n)}$ on $\mathbb{R}^{d(n)}$ such that

$$X_1 \sim \mu_{d(n)} \quad \Leftrightarrow \quad \phi_{d(n)}(X_1) \stackrel{d}{=} Y_{1;\sigma_n}.$$

Our construction of the V-statistic is thus given by taking $X_i \stackrel{\text{i.i.d.}}{\sim} \mu_{d(n)}$ and $k_v(x_1, x_2) := \tilde{k}_v(\phi_{d(n)}(x_1), \phi_{d(n)}(x_2))$, which gives

$$v_n(X) = \frac{1}{n^2} \sum_{1 \leq i, j \leq n} k_v(X_i, X_j) \stackrel{d}{=} \tilde{v}_n(Y) = \frac{1}{n} p_n^*(Y),$$

where $\stackrel{d}{=}$ denotes equality in distribution. This makes the lower bound from [12] immediately applicable. For the U-statistics construction, we observe that

$$\begin{aligned} p_n^*(y_1, \dots, y_n) &= \frac{1}{\sqrt{n(n-1)}} \sum_{i \neq j}^n \left(\frac{y_{i1}}{2\sqrt{n-1}} + \frac{y_{j1}}{2\sqrt{n-1}} + \frac{\sqrt{n-1}}{\sqrt{n}} y_{i2} y_{j2} \right) + \frac{1}{n} \sum_{i=1}^n y_{i2}^2 \\ &= \sqrt{n(n-1)} \tilde{u}_n(y_1, \dots, y_n) + R_n(y_1, \dots, y_n), \end{aligned}$$

where we have defined, for $y_1, \dots, y_n \in \mathbb{R}^2$ and $a_1, a_2, b_1, b_2 \in \mathbb{R}$,

$$\begin{aligned} \tilde{u}_n(y_1, \dots, y_n) &:= \frac{1}{n(n-1)} \sum_{i \neq j}^n \tilde{k}_u(y_i, y_j), \quad R_n(y_1, \dots, y_n) := \frac{1}{n} \sum_{i=1}^n y_{i2}^2, \\ \tilde{k}_u((a_1, a_2), (b_1, b_2)) &:= \frac{a_1}{2\sqrt{n-1}} + \frac{b_1}{2\sqrt{n-1}} + \frac{\sqrt{n-1}}{\sqrt{n}} a_2 b_2. \end{aligned}$$

Therefore, our construction of the U-statistic is to take $k_u(x_1, x_2) := \tilde{k}_u(\phi_{d(n)}(x_1), \phi_{d(n)}(x_2))$, which gives

$$u_n(X) = \frac{1}{n(n-1)} \sum_{1 \leq i \neq j \leq n} k_u(X_i, X_j) \stackrel{d}{=} \tilde{u}_n(Y) = p_n^*(Y) - R_n(Y).$$

The main technical task is thus to show that $p_n^*(Y)$ approximates a chi-squared distribution and that $R_n(Y)$ has negligible effect other than centering the chi-squared distribution.

3 Variance domination

One main technique used in the proofs is the idea of variance domination: When computing the limit of a sum of two dependent, real-valued random variables $X' + Y'$, it suffices to ignore Y' in the limit provided that the variance of Y' is negligible compared to that of X' . The same idea is used in [12] for extending their universality results to non-polynomial functions. The next result summarises the technique.

Proposition 2. Let X' and Y' be two \mathbb{R} -valued, possibly dependent random variables with $\mathbb{E}[Y'] = 0$. Then for every $t \in \mathbb{R}$,

$$\begin{aligned} & |\mathbb{P}(X' + Y' \leq t) - \mathbb{P}(X' \leq t)| \\ & \leq \inf_{\epsilon > 0} \left(\max \{ \mathbb{P}(X' \in (t - \epsilon, t]), \mathbb{P}(X' \in (t, t + \epsilon]) \} + \frac{\text{Var}[Y']}{\epsilon^2} \right). \end{aligned}$$

If we further have $\sigma_{X'}^2 := \text{Var}[X'] > 0$, then

$$\begin{aligned} & |\mathbb{P}(\sigma_{X'}^{-1}(X' + Y') \leq t) - \mathbb{P}(\sigma_{X'}^{-1}X' \leq t)| \\ & \leq \inf_{\epsilon > 0} \left(\max \{ \mathbb{P}(\sigma_{X'}^{-1}X' \in (t - \epsilon, t]), \mathbb{P}(\sigma_{X'}^{-1}X' \in (t, t + \epsilon]) \} + \frac{\text{Var}[Y']}{\text{Var}[X']\epsilon^2} \right). \end{aligned}$$

Remark 2. Several adaptations are useful: (i) To swap the roles of $X' + Y'$ and X' , one may replace X' and Y' above by $X' + Y'$ and $-Y'$ respectively; (ii) To replace $\sigma_{X'}$ by another normalisation, e.g. $\sqrt{\text{Var}[X' + Y']}$, one may rescale t and ϵ simultaneously.

Proposition 2 formalises the variance domination effect: If $\text{Var}[Y']/\text{Var}[X'] = o(1)$, by choosing $\epsilon = (\text{Var}[Y']/\text{Var}[X'])^{1/3}$, Proposition 2 implies that the c.d.f. difference $|\mathbb{P}(\sigma_{X'}^{-1}(X' + Y') \leq t) - \mathbb{P}(\sigma_{X'}^{-1}X' \leq t)| = o(1)$. Meanwhile, to get a finite-sample control, one needs an anti-concentration bound on $\sigma_{X'}^{-1}X'$. By a triangle inequality, it also suffices if $\sigma_{X'}^{-1}X'$ is well-approximated in distribution by some random variable Z' with an anti-concentration bound. For polynomials of a random vector following a log-concave probability measure, a celebrated anti-concentration result is available due to [4]. This in particular applies to polynomials of Gaussian random vectors.

Fact 3 (Carbery-Wright inequality, Theorem 8 of [4]). Let $q_m(\mathbf{x})$ be a degree- m polynomial of $\mathbf{x} \in \mathbb{R}^d$ taking values in \mathbb{R} , and η be an \mathbb{R}^d -valued random vector following a log-concave probability measure. Then there exists a constant C independent of q_m , d , m or η such that, for every $\epsilon > 0$,

$$\mathbb{P}(|q_m(\eta)| \leq \epsilon) \leq Cm \epsilon^{1/m} (\mathbb{E}[|q_m(\eta)|^2])^{-1/2m}.$$

This immediately implies the following corollary of Proposition 2:

Corollary 4. Let X' , Y' and $\sigma_{X'} > 0$ be defined as in Proposition 2, and $q_m(\eta)$ be given as in Fact 3. Suppose $\text{Var}[q_m(\eta)] = \text{Var}[\sigma_{X'}^{-1}X'] = 1$. Then there is an absolute constant $C > 0$ such that for every $t \in \mathbb{R}$,

$$\begin{aligned} & \sup_{t \in \mathbb{R}} |\mathbb{P}(X' + Y' \leq t) - \mathbb{P}(X' \leq t)| \\ & \leq Cm \left(\frac{\text{Var}[Y']}{\text{Var}[X']} \right)^{\frac{1}{2m+1}} + 2 \sup_{t \in \mathbb{R}} |\mathbb{P}(\sigma_{X'}^{-1}X' \leq t) - \mathbb{P}(q_m(\eta) \leq t)|. \end{aligned}$$

Proof of Corollary 4. The result follows from combining the second statement of Proposition 2 and Fact 3, noting that $\mathbb{E}[|q_m(\eta)|^2] \geq \text{Var}[q_m(\eta)] = 1$, choosing $\epsilon = (\text{Var}[Y']/\text{Var}[X'])^{m/(2m+1)}$ and taking a supremum over $t \in \mathbb{R}$. \square

Remark 3. The generality of Proposition 2 comes at a cost: It does not provide the tightest control on the c.d.f. difference in general, as the variance ratio term comes from the Markov's inequality. When one has more information about the tail behaviour of $Y'/\sqrt{\text{Var}[X']}$, the bound can usually be improved.

4 Proofs

Throughout this section, we denote the collection of Gaussian vectors

$$Z = (Z_1, \dots, Z_n) \quad \text{where} \quad Z_i \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(\mathbb{E}[Y_{1;\sigma_n}], \text{Var}[Y_{1;\sigma_n}]) = \mathcal{N}\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma_n^2 & 0 \\ 0 & 1 \end{pmatrix}\right).$$

We state three intermediate results used in the proof of Theorem 1. The first two results concerns the Gaussian universality approximations of $\tilde{v}_n(Y)$ and $\tilde{u}_n(Y)$. We improve the upper bound of [12], $n^{-\frac{\nu-2}{4\nu+2}}$, by using an argument specific to the construction instead of the generic Lindeberg's technique used in [12]. Note that the σ_0 -dependence below arises because Y and Z are implicitly parameterised by σ_0 .

Lemma 5. Fix $\nu \in (2, 3]$. Then there exist some absolute constants $c, C, \sigma_0 > 0$ and $N \in \mathbb{N}$, such that for all $n \geq N$,

$$cn^{-\frac{\nu-2}{4\nu}} \leq \sup_{t \in \mathbb{R}} |\mathbb{P}(n\tilde{v}_n(Y) \leq t) - \mathbb{P}(n\tilde{v}_n(Z) \leq t)| \leq Cn^{-\frac{\nu-2}{4\nu}}.$$

Lemma 6. Fix $\nu \in (2, 3]$ and let σ_0 be given as in Lemma 5. Then there exist some absolute constants $c', C' > 0$ and $N' \in \mathbb{N}$, such that for all $n \geq N'$,

$$c'n^{-\frac{\nu-2}{4\nu}} \leq \sup_{t \in \mathbb{R}} \left| \mathbb{P}\left(\sqrt{n(n-1)}\tilde{u}_n(Y) \leq t\right) - \mathbb{P}\left(\sqrt{n(n-1)}\tilde{u}_n(Z) \leq t\right) \right| \leq C'n^{-\frac{\nu-2}{4\nu}}.$$

Now observe that the universality approximations can be expressed as

$$\begin{aligned} n\tilde{v}_n(Z) &= \frac{1}{\sqrt{n}} \sum_{i=1}^n Z_{i1} + \frac{1}{n} \sum_{i,j=1}^n Z_{i2}Z_{j2} \stackrel{d}{=} \sigma_n\xi + \chi_1^2, \\ \sqrt{n(n-1)}\tilde{u}_n(Z) &= \frac{1}{\sqrt{n}} \sum_{i=1}^n Z_{i1} + \frac{1}{n} \sum_{i \neq j} Z_{i2}Z_{j2} \stackrel{d}{=} \sigma_n\xi + \frac{1}{n} \sum_{i \neq j} Z_{i2}Z_{j2}. \end{aligned}$$

The next result allows the quadratic part of $\sqrt{n(n-1)}\tilde{u}_n(Z)$ to be approximated by the centred chi-squared variable $\overline{\chi}_1^2$ plus a small Gaussian component $\sigma_n\xi$.

Lemma 7. There exists some absolute constant $C'' > 0$ such that for all $n \in \mathbb{N}$,

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P}\left(\sqrt{n(n-1)}\tilde{u}_n(Z) \leq t\right) - \mathbb{P}\left(\sigma_n\xi + \overline{\chi}_1^2 \leq t\right) \right| \leq C''n^{-1/5}.$$

4.1. Proof of Theorem 1

Recall that by construction, $u_n(X) \stackrel{d}{=} \tilde{u}_n(Y)$, $v_n(X) \stackrel{d}{=} \tilde{v}_n(Y)$ and $n\tilde{v}_n(Z) \stackrel{d}{=} \sigma_n\xi + \chi_1^2$. The V-statistic bound then follows directly from Lemma 5: There exist some constants $\sigma_0, c_1, C_1 > 0$ and $N_1 \in \mathbb{N}$ such that for all $n \geq N_1$,

$$c_1n^{-\frac{\nu-2}{4\nu}} \leq \sup_{t \in \mathbb{R}} \left| \mathbb{P}(nv_n(X) \leq t) - \mathbb{P}(\sigma_n\xi + \chi_1^2 \leq t) \right| \leq C_1n^{-\frac{\nu-2}{4\nu}}.$$

By using the triangle inequality to combine the U-statistic bounds from Lemma 6 and Lemma 7, there is another absolute constant $C_2 > 0$ such that

$$\begin{aligned} \sup_{t \in \mathbb{R}} \left| \mathbb{P}\left(\sqrt{n(n-1)}u_n(X) \leq t\right) - \mathbb{P}\left(\sigma_n\xi + \overline{\chi}_1^2 \leq t\right) \right| &\leq C_1n^{-\frac{\nu-2}{4\nu}} + C_2n^{-\frac{1}{5}}, \\ \sup_{t \in \mathbb{R}} \left| \mathbb{P}\left(\sqrt{n(n-1)}u_n(X) \leq t\right) - \mathbb{P}\left(\sigma_n\xi + \overline{\chi}_1^2 \leq t\right) \right| &\geq c_1n^{-\frac{\nu-2}{4\nu}} - C_2n^{-\frac{1}{5}}. \end{aligned}$$

Since $\frac{\nu-2}{4\nu} \leq \frac{1}{8} < \frac{1}{5}$ for $\nu \in (2, 3]$, the $n^{-1/5}$ term can be ignored when n is large. In particular, there exist $c \in (0, c_1]$, $C > C_1$ and an integer $N \geq N_1$ such that for all $n \geq N$,

$$cn^{-\frac{\nu-2}{4\nu}} \leq \sup_{t \in \mathbb{R}} \left| \mathbb{P}(nv_n(X) \leq t) - \mathbb{P}(\sigma_n\xi + \chi_1^2 \leq t) \right| \leq Cn^{-\frac{\nu-2}{4\nu}}.$$

□

4.2. Proof of Proposition 2

The proof relies on the following result:

Lemma 8 (Lemma 25 of [12]). *Let X' and Y' be two real-valued random variables. For any $a, b \in \mathbb{R}$ and $\epsilon > 0$, we have*

$$\begin{aligned}\mathbb{P}(a \leq X' + Y' \leq b) &\leq \mathbb{P}(a - \epsilon \leq X' \leq b + \epsilon) + \mathbb{P}(|Y'| \geq \epsilon), \\ \mathbb{P}(a \leq X' + Y' \leq b) &\geq \mathbb{P}(a + \epsilon \leq X' \leq b - \epsilon) - \mathbb{P}(|Y'| \geq \epsilon).\end{aligned}$$

Proof of Proposition 2. By Lemma 8 followed by the Markov's inequality, we get that for every $t \in \mathbb{R}$ and $\epsilon > 0$,

$$\begin{aligned}\mathbb{P}(X' + Y' \leq t) &\leq \mathbb{P}(X' \leq t + \epsilon) + \mathbb{P}(|Y'| \geq \epsilon) \leq \mathbb{P}(X' \leq t + \epsilon) + \frac{\text{Var}[Y']}{\epsilon^2}, \\ \mathbb{P}(X' + Y' \leq t) &\geq \mathbb{P}(X' \leq t - \epsilon) - \frac{\text{Var}[Y']}{\epsilon^2}.\end{aligned}$$

Subtracting $\mathbb{P}(X' \leq t)$ from both sides, we get that

$$|\mathbb{P}(X' + Y' \leq t) - \mathbb{P}(X' \leq t)| \leq \max \{ \mathbb{P}(X' \in (t - \epsilon, t]), \mathbb{P}(X' \in (t, t + \epsilon]) \} + \frac{\text{Var}[Y']}{\epsilon^2},$$

and taking an infimum over $\epsilon > 0$ gives the first bound. The second bound follows by rescaling t and ϵ at the same time by $\sigma'_X > 0$. \square

4.3. Proof of Lemma 5

The lower bound follows directly from Theorem 2 of [12] by noting that $n\tilde{\nu}_n = p_n^*$, so it suffices to prove the upper bound. Denote

$$S_n := \frac{1}{\sqrt{n}} \sum_{i=1}^n \left(\frac{1}{\sqrt{2}} U_{i;\sigma_n} + \frac{\sigma_n}{\sqrt{2}} \xi_{i1} \right)$$

such that, for $\xi \sim \mathcal{N}(0, 1)$ independent of all other variables,

$$n\tilde{\nu}_n(Y) \stackrel{d}{=} S_n + \chi_1^2 \quad \text{and} \quad n\tilde{\nu}_n(Z) \stackrel{d}{=} \sigma_n \xi + \chi_1^2.$$

By Lemmas 19 and 20 of [12], under the choice of σ_0 and \tilde{N} in Theorem 2 of [12], we have that for all $n \geq \tilde{N}$,

$$\begin{aligned}\sup_{x \in \mathbb{R}} \left| \mathbb{P}(S_n < x) - \mathbb{P}(\sigma_n \xi < x) - \frac{A}{n^{1/2} \sigma_n^{\nu/(\nu-2)}} \left(1 - \frac{x^2}{\sigma_n^2} \right) e^{-x^2/(2\sigma_n^2)} \right| &\leq \frac{2}{n \sigma^{2\nu/(\nu-2)}}, \\ |\mathbb{P}(S_n < x) - \mathbb{P}(\sigma_n \xi < x)| &\leq B \left(\frac{1}{n^{1/2} \sigma_n^{\nu/(\nu-2)}} e^{-x^2/16\sigma_n^2} + \frac{1}{n^{3/2} x^4 \sigma_n^{(8-\nu)/(\nu-2)}} \right)\end{aligned}$$

for some absolute constants $A, B > 0$. By splitting an integral and using these bounds,

$$\begin{aligned}
& |\mathbb{P}(n \tilde{v}_n(Y) < t) - \mathbb{P}(n \tilde{v}_n(Z) < t)| \\
&= \left| \int_{-\infty}^{\infty} (\mathbb{P}(S_n < t - y^2) - \mathbb{P}(\sigma_n \xi < t - y^2)) \frac{e^{-y^2/2}}{\sqrt{2\pi}} dy \right| \\
&\leq \int_{|t-y^2| < \sigma_n} |\mathbb{P}(S_n < t - y^2) - \mathbb{P}(\sigma_n \xi < t - y^2)| \frac{e^{-y^2/2}}{\sqrt{2\pi}} dy \\
&\quad + \int_{|t-y^2| \geq \sigma_n} |\mathbb{P}(S_n < t - y^2) - \mathbb{P}(\sigma_n \xi < t - y^2)| \frac{e^{-y^2/2}}{\sqrt{2\pi}} dy \\
&\leq \frac{A}{\sqrt{2\pi} n^{1/2} \sigma_n^{\nu/(\nu-2)}} \int_{|t-y^2| < \sigma_n} \left| 1 - \frac{(t-y^2)^2}{\sigma_n^2} \right| e^{-\frac{(t-y^2)^2}{2\sigma_n^2} - \frac{y^2}{2}} dy + \frac{4\sqrt{\sigma_n}}{n\sigma_n^{2\nu/(\nu-2)}} \\
&\quad + \frac{B}{\sqrt{2\pi} n^{1/2} \sigma_n^{\nu/(\nu-2)}} \int_{|t-y^2| \geq \sigma_n} e^{-\frac{(t-y^2)^2}{16\sigma_n^2} - \frac{y^2}{2}} dy \\
&\quad + \frac{B}{\sqrt{2\pi} n^{3/2} \sigma_n^{(8-\nu)/(\nu-2)}} \int_{|t-y^2| \geq \sigma_n} \frac{1}{(t-y^2)^4} e^{-\frac{y^2}{2}} dy \\
&\leq \frac{2A\sqrt{\sigma_n}}{\sqrt{2\pi} n^{1/2} \sigma_n^{\nu/(\nu-2)}} + \frac{4\sqrt{\sigma_n}}{n\sigma_n^{2\nu/(\nu-2)}} + \frac{B\sqrt{\sigma_n}}{\sqrt{2\pi} n^{1/2} \sigma_n^{\nu/(\nu-2)}} \int_{|\sigma_n^{-1}t-y^2| \geq 1} e^{-\frac{(\sigma_n^{-1}t-y^2)^2}{16}} dy \\
&\quad + \frac{B\sqrt{\sigma_n}}{\sqrt{2\pi} n^{3/2} \sigma_n^{(8-\nu)/(\nu-2)} \sigma_n^4} \int_{|\sigma_n^{-1}t-y^2| \geq 1} \frac{1}{(\sigma_n^{-1}t-y^2)^4} dy.
\end{aligned}$$

In the last line, we have used a change-of-variable $y \mapsto \sqrt{\sigma_n}y$ and noted that $e^{-y^2/2} \leq 1$. Now fix $N \geq \tilde{N}$ such that $\sigma_n = \sigma_0 n^{-(\nu-2)/(2\nu)}$ for all $n \geq N$, which only depends on the absolute constant $\sigma_0 > 0$; in this case,

$$\frac{\sqrt{\sigma_n}}{n^{1/2} \sigma_n^{\nu/(\nu-2)}} = \sigma_0^{\frac{1}{2} - \frac{\nu}{\nu-2}} n^{-\frac{\nu-2}{4\nu}}, \quad \frac{\sqrt{\sigma_n}}{n^{3/2} \sigma_n^{(8-\nu)/(\nu-2)} \sigma_n^4} = \sigma_0^{\frac{1}{2} - \frac{3\nu}{\nu-2}} n^{-\frac{\nu-2}{4\nu}}.$$

Then there exists some constant A' that depends only on σ_0 such that, for all $n > N$,

$$\begin{aligned}
& |\mathbb{P}(n \tilde{v}_n(Y) < t) - \mathbb{P}(n \tilde{v}_n(Z) < t)| \\
&\leq A' n^{-\frac{\nu-2}{4\nu}} \left(1 + \int_{|\sigma_n^{-1}t-y^2| \geq 1} e^{-\frac{(\sigma_n^{-1}t-y^2)^2}{16}} dy + \int_{|\sigma_n^{-1}t-y^2| \geq 1} \frac{1}{(\sigma_n^{-1}t-y^2)^4} dy \right) \\
&=: A' n^{-\frac{\nu-2}{4\nu}} (1 + I_1(\sigma_n^{-1}t) + I_2(\sigma_n^{-1}t)),
\end{aligned}$$

where we write $I_1(\tau) := \int_{|\tau-y^2| \geq 1} e^{-(\tau-y^2)^2/16} dy$ and $I_2(\tau) := \int_{|\tau-y^2| \geq 1} (\tau-y^2)^{-4} dy$. To handle $I_1(\tau)$, we split the integral further and use a change-of-variable with $z = y^2$ to obtain

$$\begin{aligned}
I_1(\tau) &\leq \int e^{-(\tau-y^2)^2/16} dy = \int_{y^2 < 1} e^{-(\tau-y^2)^2/16} dy + 2 \int_{y^2 \geq 1, y \geq 0} e^{-(\tau-y^2)^2/16} dy \\
&\leq 2 + 2 \int_{z \geq 1} e^{-(\tau-z)^2/16} \frac{1}{2\sqrt{z}} dz \\
&\leq 2 + \int_{z \geq 1} e^{-(\tau-z)^2/16} dz \leq 2 + \sqrt{2\pi} \times 8 \int \frac{e^{-(\tau-z)^2/16}}{\sqrt{2\pi} \times 8} dz = 2 + 4\sqrt{\pi}.
\end{aligned}$$

A similar strategy applied to $I_2(\tau)$ gives

$$\begin{aligned}
I_2(\tau) &= \int_{|\tau-y^2| \geq 1, y^2 < 1} \frac{1}{(\tau-y^2)^4} dy + 2 \int_{|\tau-y^2| \geq 1, y^2 \geq 1, y \geq 0} \frac{1}{(\tau-y^2)^4} dy \\
&\leq 2 + 2 \int_{|\tau-z| \geq 1, z \geq 1} \frac{1}{(\tau-z)^4} \frac{1}{2\sqrt{z}} dz \\
&\leq 2 + \int_{|\tau-z| \geq 1} \frac{1}{(\tau-z)^4} dz = 2 + \int_{|z'| \geq 1} \frac{1}{(z')^4} d(z') = \frac{8}{3}.
\end{aligned}$$

Substituting these two bounds back and noting that the resulted bounds do not depend on t , we get that there exist some constants $C > 0$ and $N \in \mathbb{N}$ that depend only on the absolute constant $\sigma_0 > 0$ such that for all $n \geq N$,

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P}(n \tilde{v}_n(Y) < t) - \mathbb{P}(n \tilde{v}_n(Z) < t) \right| \leq C n^{-\frac{\nu-2}{4\nu}}.$$

□

4.4. Proof of Lemma 6

Notice that

$$\sqrt{n(n-1)} \tilde{u}_n = p_n^* - R_n = n \tilde{v}_n - R_n,$$

and we already have the universality approximation bound for $n \tilde{v}_n$ from Lemma 5. It suffices to apply variance domination to approximate $\sqrt{n(n-1)} \tilde{u}_n$ by $n \tilde{v}_n$, which requires us to compute the relevant variances. Write $Z_i = (Z_{i1}, Z_{i2})$, where Z_{i1} is the Gaussian component that match the first two moments of $U_{i;\sigma_n} + 2^{-1/2} \sigma_n \xi_{i1}$ and Z_{i2} is the Gaussian component that matches $\xi_{i2} \sim \mathcal{N}(0, 1)$ in distribution. Then by independence,

$$\begin{aligned} \sigma_*^2 &:= \text{Var}[p_n^*(Y)] = \text{Var}[p_n^*(Z)] = \text{Var}\left[\frac{1}{\sqrt{n}} \sum_{i=1}^n Z_{i1}\right] + \text{Var}\left[\left(\frac{1}{\sqrt{n}} \sum_{i=1}^n Z_{i2}\right)^2\right] \\ &\geq \text{Var}\left[\left(\frac{1}{\sqrt{n}} \sum_{i=1}^n Z_{i2}\right)^2\right] = \text{Var}[Z_{12}^2] = 2. \end{aligned}$$

By independence again,

$$\text{Var}[R_n(Y)] = \text{Var}[R_n(Z)] = \text{Var}\left[\frac{1}{n} \sum_{i=1}^n Z_{i2}^2\right] = \frac{\text{Var}[Z_{12}^2]}{n} = \frac{2}{n},$$

and also note that

$$\mathbb{E}[R_n(Y)] = \mathbb{E}[R_n(Z)] = 1.$$

We now prove the upper bound by combining the variance domination result in Corollary 4 and the upper bound of Lemma 5. Let σ_0 be given as in Lemma 5. Since

$$\sqrt{n(n-1)} \tilde{u}_n(Y) = (p_n^*(Y) - 1) - (R_n - \mathbb{E}[R_n(Y)]),$$

there are some absolute constants $\tilde{C}' > 0$ and $N \in \mathbb{N}$ such that for all $n \geq N$ and every $t \in \mathbb{R}$,

$$\begin{aligned} & \left| \mathbb{P}(\sqrt{n(n-1)} \tilde{u}_n(Y) \leq t) - \mathbb{P}(p_n^*(Y) - 1 \leq t) \right| \\ & \leq \tilde{C}' \left(\frac{\text{Var}[R_n(Y)]}{\text{Var}[p_n^*(Y)]} \right)^{\frac{1}{5}} + 2 \sup_{\tau \in \mathbb{R}} \left| \mathbb{P}(\sigma_*^{-1}(p_n^*(Y) - 1) \leq \tau) - \mathbb{P}(\sigma_*^{-1}(p_n^*(Z) - 1) \leq \tau) \right| \\ & \leq \tilde{C}' n^{-\frac{1}{5}} + 2C n^{-\frac{\nu-2}{4\nu}} \leq \tilde{C}'_* n^{-\frac{\nu-2}{4\nu}} \end{aligned}$$

for some absolute constants $C, \tilde{C}'_* > 0$; in the last line, we have used that $\frac{\nu-2}{4\nu} \leq \frac{1}{8} < \frac{1}{5}$ for $\nu \in (2, 3]$. Similarly we have

$$\left| \mathbb{P}(\sqrt{n(n-1)} \tilde{u}_n(Z) \leq t) - \mathbb{P}((p_n^*(Z) - 1) \leq t) \right| \leq \tilde{C}'_* n^{-\frac{\nu-2}{4\nu}}.$$

Combining both bounds with the upper bound of Lemma 5 by a triangle inequality, we get the desired upper bound that for some absolute constant $C' > 0$ and all $n \geq N$,

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P}(\sqrt{n(n-1)} \tilde{u}_n(Y) \leq t) - \mathbb{P}(\sqrt{n(n-1)} \tilde{u}_n(Z) \leq t) \right| \leq C' n^{-\frac{\nu-2}{4\nu}}.$$

For the lower bound, we apply Lemma 21 of [12]: There exist some constants $c_* > 0$ and $\tilde{N} \in \mathbb{N}$ depending only on the fixed constant σ_0 —which is chosen to be the one used in Theorem 2 of [12] and also Lemma 5—such that for any $n \geq \tilde{N}$,

$$\mathbb{P}(p_n^*(Z) < -2\sigma_n) - \mathbb{P}(p_n^*(Y) < -2\sigma_n) \geq c_* n^{-\frac{\nu-2}{4\nu}}.$$

To exploit this lower bound, we shall apply Lemma 8 directly: For any $\epsilon > 0$,

$$\begin{aligned} & \mathbb{P}(\sqrt{n(n-1)} \tilde{u}_n(Z) < -2\sigma_n - \sigma_* \epsilon - 1) - \mathbb{P}(\sqrt{n(n-1)} \tilde{u}_n(Y) < -2\sigma_n - \sigma_* \epsilon - 1) \\ &= \mathbb{P}(\sqrt{n(n-1)} \tilde{u}_n(Z) < -2\sigma_n - \sigma_* \epsilon - 1) - \mathbb{P}(p_n^*(Z) - 1 < -2\sigma_n - 2\sigma_* \epsilon - 1) \\ &\quad - \mathbb{P}(-2\sigma_n - 2\sigma_* \epsilon \leq p_n^*(Z) < -2\sigma_n) \\ &\quad + \mathbb{P}(p_n^*(Z) < -2\sigma_n) - \mathbb{P}(p_n^*(Y) < -2\sigma_n) \\ &\quad + \mathbb{P}(p_n^*(Y) - 1 < -2\sigma_n - 1) - \mathbb{P}(\sqrt{n(n-1)} \tilde{u}_n(Y) < -2\sigma_n - \sigma_* \epsilon - 1) \\ &\geq -2\mathbb{P}(|R_n(Y) - 1| \geq \sigma_* \epsilon) - \mathbb{P}(|p_n^*(Z) + 2\sigma_n + \sigma_* \epsilon| \leq \sigma_* \epsilon) + c_* n^{-\frac{\nu-2}{4\nu}} \\ &\stackrel{(a)}{\geq} -\frac{2\text{Var}[R_n(Y)]}{\sigma_*^2 \epsilon^2} - \frac{\tilde{c}'_*(\sigma_* \epsilon)^{1/2}}{(\mathbb{E}[|p_n^*(Z) + 2\sigma_n + \sigma_* \epsilon|^2])^{1/4}} + c_* n^{-\frac{\nu-2}{4\nu}} \\ &\stackrel{(b)}{\geq} -\frac{2}{\epsilon^2 n} - \tilde{c}'_* \epsilon^{1/2} + c_* n^{-\frac{\nu-2}{4\nu}} \end{aligned}$$

for some absolute constant $\tilde{c}'_* > 0$. In (a), we have used the Markov's inequality and the Carbery-Wright inequality (Fact 3); in (b), we have plugged in the bounds on $\text{Var}[R_n(Y)]$ and σ_* , and noted that $\mathbb{E}[|p_n^*(Z) + 2\sigma_n + \sigma_* \epsilon|^2] \geq \text{Var}[p_n^*(Z)] = \sigma_*^2$. Taking $\epsilon = n^{-2/5}$ gives

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P}(\sqrt{n(n-1)} \tilde{u}_n(Y) \leq t) - \mathbb{P}(\sqrt{n(n-1)} \tilde{u}_n(Z) \leq t) \right| \geq c_* n^{-\frac{\nu-2}{4\nu}} - \left(\frac{4}{5} + \tilde{c}'_*\right) n^{-\frac{1}{5}},$$

Since $\frac{\nu-2}{4\nu} \leq \frac{1}{8} < \frac{1}{5}$, there are some absolute constants $c' > 0$ and $N' \geq \tilde{N}$ such that the desired lower bound holds for all $n \geq N'$. \square

4.5. Proof of Lemma 7

Recall from the proof of Lemma 6 that $\mathbb{E}[R_n(Z)] = 1$, $\text{Var}[R_n(Z)] = \frac{2}{n}$ and that $\text{Var}[n\tilde{v}_n(Z)] = \text{Var}[p_n^*(Z)] \geq 2$. Since

$$\sqrt{n(n-1)} \tilde{u}_n(Z) = n\tilde{v}_n(Z) - \frac{1}{n} \sum_{i=1}^n Z_{i2}^2 = (n\tilde{v}_n(Z) - 1) - \left(\frac{1}{n} \sum_{i=1}^n Z_{i2}^2 - 1\right),$$

we can again apply variance domination (Corollary 4) to obtain that

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P}(\sqrt{n(n-1)} \tilde{u}_n(Z) \leq t) - \mathbb{P}(n\tilde{v}_n(Z) - 1 \leq t) \right| \leq C \left(\frac{\text{Var}[R_n(Z)]}{\text{Var}[n\tilde{v}_n(Z) - 1]} \right)^{\frac{1}{5}} = C n^{-\frac{1}{5}}$$

for some absolute constant $C > 0$. Noting that $n\tilde{v}_n(Z) - 1 \stackrel{d}{=} \sigma_n \xi + \overline{\chi_1^2}$ finishes the proof. \square

Acknowledgements

This work has been supported by the Gatsby Charitable Foundation.

References

- [1] M. Bayati, M. Lelarge, and A. Montanari. Universality in polytope phase transitions and message passing algorithms. *Ann. Appl. Probab.*, 25(2):753 – 822, 2015.
- [2] V. Bentkus, F. Gotze, and R. Zitikis. Lower estimates of the convergence rate for u-statistics. *Ann. Probab.*, 22(4):1707–1714, 1994.
- [3] B. B. Bhattacharya, S. Das, S. Mukherjee, and S. Mukherjee. Asymptotic distribution of random quadratic forms. *arXiv:2203.02850*, 2022.
- [4] A. Carbery and J. Wright. Distributional and l^q norm inequalities for polynomials over convex bodies in \mathbb{R}^n . *Math. Res. Lett.*, 8(3):233–248, 2001.
- [5] L. H. Chen, L. Goldstein, and Q.-M. Shao. *Normal approximation by Stein’s method*, volume 2. Springer, 2011.
- [6] S. X. Chen and Y.-L. Qin. A two-sample test for high-dimensional data with applications to gene-set testing. *Ann. Statist.*, 38(2):808 – 835, 2010.
- [7] H. Gao, R. Wang, and X. Shao. Dimension-agnostic change point detection. *arXiv:2303.10808*, 2023.
- [8] F. Gerace, F. Krzakala, B. Loureiro, L. Stephan, and L. Zdeborová. Gaussian universality of perceptrons with random labels. *Phys. Rev. E*, 109(3):034305, 2024.
- [9] F. Götze and A. Y. Zaitsev. Explicit rates of approximation in the clt for quadratic forms. *Ann. Probab.*, 42(1): 354–397, 2014.
- [10] A. Gretton, K. M. Borgwardt, M. J. Rasch, B. Schölkopf, and A. Smola. A kernel two-sample test. *J. Mach. Learn. Res.*, 13(1):723–773, 2012.
- [11] K. H. Huang, X. Liu, A. Duncan, and A. Gandy. A high-dimensional convergence theorem for U-statistics with applications to kernel-based testing. In *The Thirty Sixth Annual Conference on Learning Theory*, pages 3827–3918. PMLR, 2023.
- [12] K. H. Huang, M. Austern, and P. Orbanz. Gaussian universality for approximately polynomial functions of high-dimensional data. *arXiv:2403.10711*, 2024.
- [13] I. Kim and A. Ramdas. Dimension-agnostic inference using cross U-statistics. *Bernoulli*, 30(1):683 – 711, 2024.
- [14] A. Montanari and B. N. Saeed. Universality of empirical risk minimization. In *Conference on Learning Theory*, pages 4310–4312. PMLR, 2022.
- [15] L. Pesce, F. Krzakala, B. Loureiro, and L. Stephan. Are Gaussian data all you need? The extents and limits of universality in high-dimensional generalized linear estimation. In *International Conference on Machine Learning*, pages 27680–27708. PMLR, 2023.
- [16] V. V. Senatov. *Normal Approximation*. De Gruyter, 1998.
- [17] R. J. Serfling. *Approximation theorems of mathematical statistics*, volume 162. John Wiley & Sons, 1980.
- [18] S. Shekhar, I. Kim, and A. Ramdas. A permutation-free kernel two-sample test. *Advances in Neural Information Processing Systems*, 35:18168–18180, 2022.
- [19] T. Tao and V. Vu. Random matrices: universality of local eigenvalue statistics. *Acta Math.*, 206:127–204, 2011.
- [20] T. Tao and V. Vu. Random matrices: universality of local spectral statistics of non-hermitian matrices. *Ann. Probab.*, 43:782–874, 2015.
- [21] R. Wang, C. Zhu, S. Volgushev, and X. Shao. Inference for change points in high-dimensional data via self-normalization. *Ann. Statist.*, 50(2):781–806, 2022.
- [22] J. Yan and X. Zhang. Kernel two-sample tests in high dimension: interplay between moment discrepancy and dimension-and-sample orders. *Biometrika*, 110(2):411–430, 2022.
- [23] O. Yanushkevichiene. On bounds in limit theorems for some U-statistics. *Theory Probab. Appl.*, 56(4):660–673, 2012.