

Small maximal clusters are very unlikely in critical random graphs

Umberto De Ambrogio*

June 19, 2024

Abstract

We describe a probabilistic methodology, based on random walk estimates, to obtain *exponential* upper bounds for the probability of observing unusually *small* maximal components in two classical (near-)critical random graph models. More specifically, we analyse the near-critical Erdős-Rényi model $\mathbb{G}(n, p)$ and the random graph $\mathbb{G}(n, d, p)$ obtained by performing near-critical p -bond percolation on a simple random d -regular graph and show that, for each one of these models, the probability that the size of a largest component is smaller than $n^{2/3}/A$ is at most of order $\exp(-A^{3/2})$. The exponent $3/2$ is known to be *optimal* for the near-critical $\mathbb{G}(n, p)$ random graph, whereas for the near-critical $\mathbb{G}(n, d, p)$ model the best known upper bound for the above probability was of order $A^{-3/5}$. As a secondary result we show, by means of an *optimized* version of the martingale method of Nachmias and Peres, that the above probability of observing an unusually small maximal component is at most of order $\exp(-A^{3/5})$ in other two critical models, namely a random intersection graph and the quantum random graph; this stretched-exponential bounds also improve upon the known (polynomial) bounds available for these other two critical models.

1 Introduction

In recent years, the problem of deriving exponential bounds for the probability of observing unusually *large* maximal components in critical random graphs has gained considerable interest; see e.g. [12, 13, 22, 30, 31] as well as Section 7.1.1 in the PhD thesis of Dhara [16].

In this work we consider the complementary and less investigated problem of determining *exponential* upper bounds for the probability of observing unusually *small* maximal components in critical random graphs. In other words, we are interested in providing exponential (upper) bounds for probabilities of the type

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G})| < k), \quad (1)$$

where $\mathcal{C}_{\max}(\mathbb{G})$ is a maximal cluster of some (critical) random graph \mathbb{G} and $k \in \mathbb{N}$ (for us, $k = n^{2/3}/A$, with $n^{2/3}$ being the correct order of $|\mathcal{C}_{\max}(\mathbb{G})|$ in the regime we are interested in).

To the best of our knowledge, the only other work we are aware of that investigates this problem is that of Pittel [30] in the context of the (near-critical) Erdős-Rényi model $\mathbb{G}(n, p)$. Before looking at Pittel's result, let us recall the definition of the $\mathbb{G}(n, p)$ model.

This random graph, which is obtained by performing p -bond percolation on the complete graph on n vertices (i.e. it is constructed by independently retaining each edge with probability p and deleting it with probability $1 - p$), is known to undergo a phase transition as np passes 1. Specifically, letting $p = p(n) := \mu/n$, if $\mu < 1$ then $|\mathcal{C}_{\max}(\mathbb{G}(n, p))|$ is of order $\log n$; if $\mu = 1$ (the *critical case*), then $|\mathcal{C}_{\max}(\mathbb{G}(n, p))|$ is of order $n^{2/3}$; and if $\mu > 1$, then $|\mathcal{C}_{\max}(\mathbb{G}(n, p))|$ is of order n . See e.g. the monographs [7], [20] or [24] for more details.

Actually we know more: if $\mu = \mu(n) = 1 + \lambda n^{-1/3}$ then $|\mathcal{C}_{\max}|$ is still of order $n^{2/3}$ [20]; this is the so-called *critical window*. Pittel [30] considered this regime and he showed (among many other things) that, in the near-critical Erdős-Rényi random graph $\mathbb{G}(n, p)$ with $p = p(n) = (1 + \lambda n^{-1/3})/n$ and $\lambda \in \mathbb{R}$, as $A \rightarrow \infty$

$$\lim_{n \rightarrow \infty} \mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, p))| < n^{2/3}/A) = C(\lambda) \exp\left(-c_1 \lambda^2 A^{1/2} - c_2 \lambda A - c_2 A^{3/2}\right), \quad (2)$$

*Ludwig Maximilians Universität München, Department of Mathematical Sciences. umbidea@gmail.com

where c_1, c_2 and c_3 are positive constants (independent of λ) and $C(\lambda)$ is a positive constant which depends on λ . We refer the reader to Corollary 2 in Pittel's paper [30] for the definition (and numerical estimates) of the constants appearing in (2).

To the best of our knowledge, with the exception of (2), all other upper bounds available in the literature for the probability of observing *small* maximal clusters in (near-)critical random graphs are of the type $A^{-\gamma}$, where $\gamma \in (0, 1)$.

For instance, Nachmias and Peres [27] used martingale arguments to analyse the component structure in the random graph obtained by performing p -bond percolation on a random d -regular graph. This random graph, denoted by $\mathbb{G}(n, d, p)$, is the random graph on n vertices obtained by first drawing uniformly at random a d -regular simple graph on $[n]$ and then performing independent p -bond percolation on it.

Alon, Benjamini and Stacey [3] showed that $\mathbb{G}(n, d, \mu/(d-1))$ has a phase transition similar to the one observed in the $\mathbb{G}(n, p)$ model. Indeed, $\mathbb{G}(n, d, \mu/(d-1))$ undergoes a phase transition as μ passes 1: the size of the largest component $|\mathcal{C}_{\max}(\mathbb{G}(n, d, \mu/(d-1)))|$ is of order $\log n$ when $\mu < 1$, and of order n when $\mu > 1$.

Nachmias and Peres [27] analysed (using probabilistic arguments) the critical window of this model and showed that, when $p = p(n, d) = (1 + \lambda n^{-1/3})/(d-1)$ with $\lambda \in \mathbb{R}$ and $d \geq 3$ both fixed, there exists a positive constant $c = c(\lambda, d)$ which depends on λ, d such that, for all large enough $A > 0$ and n ,

$$\mathbb{P}\left(|\mathcal{C}_{\max}(\mathbb{G}_{n,d,p})| < n^{2/3}/A\right) \leq cA^{-1/2}. \quad (3)$$

More recently, Joos and Perarnau [25] showed that $|\mathcal{C}_{\max}(\mathbb{G}(n, d, p))|$ is of order $n^{2/3}$ for all $d = d(n) \in \{3, \dots, n-1\}$ when $p = (1 + \lambda n^{-1/3})/(d-1)$ (with $\lambda \in \mathbb{R}$ fixed), extending the result of Nachmias and Peres [27] and confirming a prediction of the two authors on a question of Benjamini. In particular, Joos and Perarnau [25] showed that (in the above setting)

$$\mathbb{P}\left(n^{2/3}/A < |\mathcal{C}_{\max}(\mathbb{G}(n, p))| < An^{2/3}\right) \geq 1 - cA^{-1/2}.$$

We remark that the (probabilistic) methodology used by Nachmias and Peres [27] was introduced by the same authors in [28], with the purpose of giving simple proofs that $|\mathcal{C}_{\max}(\mathbb{G}(n, p))|$ is of order $n^{2/3}$ when $p = (1 + \lambda n^{-1/2})/n$. They showed (among other things) that when $p = 1/n$ then

$$\mathbb{P}\left(|\mathcal{C}_{\max}(\mathbb{G}(n, p))| < n^{2/3}/A\right) \leq cA^{-3/5} \quad (4)$$

for some constant $c > 0$ and for all sufficiently large A and n . (In [28] the authors also considered the probability of observing large maximal clusters and p of the form $p = (1 + \lambda n^{-1/3})/n$ with $\lambda \in \mathbb{R}$, see Section 1.2 below.)

The martingale method of Nachmias and Peres [27, 28], which lead to the bounds displayed in (3) and (4), has been successfully employed by other authors to analyse the (near-)critical behaviour of several other random graphs, leading to polynomial (upper) bounds of the above.

For instance, Hatami and Molloy [19] adapted such a methodology to determine the critical window for a random graph on a given degree sequence, whereas Dembo, Levit and Vadlamani [15] used the method developed by Nachmias and Peres to study the near-critical behaviour of the *quantum* random graph (see Section 3 for a brief introduction to this model). Furthermore, the same technique was also used by the author and Pachon [11] to analyse (a specific instance of) the critical Norros-Reittu model [29], whose near-critical behaviour was first established by van der Hofstad [21].

1.1 Main result

Pittel's proof of (2) heavily relies on combinatorial arguments and seems difficult to be adapted to other models. On the other hand, the probabilistic argument of Nachmias and Peres [27, 28], although very general and robust, does not seem sufficient to retrieve the optimal exponent $3/2$ which appears in the leading term of (2). Indeed, although we can boost (by means of few modifications) the original martingale argument [27, 28] to obtain upper bounds of the type $\exp(-A^{3/5})$ in several critical models, it seems that the method can't be pushed to achieve upper bounds of the (correct) order $\exp(-A^{3/2})$; we refer the reader to Section 3 for a detailed explanation concerning why we believe this to be the case.

The goal of this work is to introduce a robust probabilistic methodology, simply based on random walk estimates, to derive *sharp* exponential upper bounds for the probability of observing unusually *small* maximal clusters in (near-)critical random graphs. We do so by analysing the near-critical models $\mathbb{G}(n, p)$ and $\mathbb{G}(n, d, p)$ and show that $|\mathcal{C}_{\max}(\mathbb{G})| \geq n^{2/3}/A$ with probability at least of order $1 - e^{-A^{3/2}}$ when either $\mathbb{G} = \mathbb{G}(n, p)$ or $\mathbb{G} = \mathbb{G}(n, d, p)$. Here is our main result.

Theorem 1.1. *Let $p = (1 + \lambda n^{-1/3})/n$ with $\lambda \in \mathbb{R}$ fixed. There exist constants $A_0 = A_0(\lambda) > 0$ and $c > 0$ such that, for any $A_0 \leq A = o(n^{2/3})$ and for all large enough n , we have*

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, p))| < n^{2/3}/A) \leq \exp(-cA^{3/2}). \quad (5)$$

Let $p = (1 + \lambda n^{-1/3})/(d - 1)$ with $\lambda \in \mathbb{R}, d \geq 3$ both fixed. There exist constants $A_0 = A_0(\lambda, d) > 0$ and $c = c(d) > 0$ such that, for any $A_0 \leq A = O(n^{1/6})$ and for all large enough n , we have

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, d, p))| < n^{2/3}/A) \leq \exp(-cA^{3/2}). \quad (6)$$

Remark 1. We emphasize that (5) is not new (see (2) above), but our proof is purely probabilistic; on the other hand, (6) is new and most likely is of the right order; moreover, it substantially improves upon the known bound displayed in (3). We remark that Theorem 1.1 only considers the near-critical models $\mathbb{G}(n, p)$ and $\mathbb{G}(n, d, p)$, but we believe that the argument used to establish (5) and (6) is robust. One could try to adapt the methodology introduced in this work to prove similar tail bounds in other (near-)critical random graphs. We refer the reader to Section 3 for a detailed discussion (and related results) along these lines.

Remark 2. We emphasize that, in Theorem 1.1, A is allowed to depend on n . The condition $A = O(n^{1/6})$, which we require in our upper bound for the $\mathbb{G}(n, d, p)$ random graph could be changed to $A = o(n^{2/3})$ at the cost of adding a term of order $\exp(-A^{1/2}n^{1/6})$ to (6). Moreover, the condition $A = o(n^{2/3})$ which we require in our upper bound for the $\mathbb{G}(n, p)$ model could be relaxed to $A \leq \varepsilon n^{2/3}$ for a sufficiently small constant $\varepsilon > 0$. However, when $A \asymp n^{2/3}$, then $n^{2/3}/A \asymp 1$ and so we would be bounding from above the probability that the size of a maximal cluster is at most some *constant*, which is not very interesting. Therefore we can say that our upper bound in (5) concerns *all* the relevant values of A . On the other hand, it could be possible to weaken the requirement $A = O(n^{1/6})$ which we made for the $\mathbb{G}(n, d, p)$ random graph at the expense of longer computations, but we refrained to do so. Indeed, we believe the bounds in (5) and (6) to be more interesting for ‘small’ (possibly dependent of n) values of A , being $n^{2/3}$ the correct order for the size of a largest cluster in both (near-)critical models. We also remark that the parameter λ , which we consider as fixed, could be allowed to depend on n at the expense of more careful computations. In this case, one would need to keep track of the dependence between A and λ . Furthermore, if needed, one could give bounds on the ‘smallest values’ of A_0 for which (5) and (6) hold; similarly, one might try to give bounds on the smallest n for which (5) and (6) are verified (as it is done e.g. in [28]). However, we refrained to do so in order to keep the computations shorter.

1.2 Related work

There are several works focusing on the problem of deriving sharp tail bounds for the probability of observing unusually *large* clusters.

Concerning the Erdős-Rényi random graph $\mathbb{G}(n, p)$, Pittel [30] also showed that, in the near-critical regime $p = (1 + \lambda n^{-1/3})/n$ with $\lambda \in \mathbb{R}$, then

$$\lim_{n \rightarrow \infty} A^{3/2} \exp\left(\frac{A^3}{8} - \frac{\lambda A^2}{2} + \frac{\lambda^2 A}{2}\right) \mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, p))| > An^{2/3}) \quad (7)$$

converges, as $A \rightarrow \infty$, to a specific constant, which is stated to be $(2\pi)^{-1/2}$ but should be $(8/9\pi)^{1/2}$, as remarked by Roberts [31].

Along the same lines, van der Hofstad, Kleim and Van Leeuwen [22] proved similar results in the context of inhomogeneous random graphs.

Both Pittel [30] and Roberts [31] relied on a combinatorial formula for the expected number of components with k vertices and $k + \ell$ edges, which is specific to the $\mathbb{G}_{n,p}$ model and appears difficult to adapt to other random graphs.

A first, purely probabilistic approach to establish an exponential upper bound for the probability displayed in (7) was introduced by Nachmias and Peres [28], who used martingale arguments to show that for all large enough A and n , when $p = 1/n$ then

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, p))| > An^{2/3}) \leq \frac{4}{A} \exp(-A^2(A-4)/32). \quad (8)$$

They also considered the near-critical model where $p = (1 + \lambda n^{-1/3})/n$ for fixed $\lambda \in \mathbb{R}$ and established a similar bound.

The same martingale-based method was subsequently used by the same authors in [27], where they analysed the (near-critical) $\mathbb{G}(n, d, p)$ model.

Amongst other results (some of which were discussed earlier), Nachmias and Peres [27] proved that, in the near-critical regime where $p = (1 + \lambda n^{-1/3})/(d-1)$ and $\lambda \in \mathbb{R}, d \geq 3$ are both *fixed*, there are positive constants $c = c(\lambda, d)$ and $C = C(\lambda, d)$ which depend on both λ and d such that, for any $A > 0$ and for all large enough n ,

$$\mathbb{P}\left(|\mathcal{C}_{\max}(\mathbb{G}(n, d, p))| > An^{2/3}\right) \leq \frac{C}{A} \exp(-cA^3). \quad (9)$$

With the purpose of introducing a *probabilistic* methodology capable of recovering the correct asymptotic of Pittel displayed in (7), more recently the author and Roberts [12] introduced a new methodology, based on (probabilistic) tools such as generalised ballot theorems [1] and strong embedding with Brownian motion [8], to derive *matching* upper and lower bounds for the probability of observing unusually *large* maximal clusters in critical random graphs.

In particular, for the near-critical $\mathbb{G}(n, p)$ model with $p = (1 + \lambda n^{-1/3})/n$ and $\lambda \in \mathbb{R}$ (possibly dependent on n), the authors showed in [12] that there exist constants $A_0 > 0, n_0 \in \mathbb{N}$ such that, if $A_0 \leq A = A(n) = o(n^{1/30})$ and $\lambda = \lambda(n)$ satisfies $|\lambda| \leq A/3$, then for all $n \geq n_0$

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, p))| > An^{2/3}) \asymp \frac{c_2}{A^{3/2}} \exp\left(-\frac{A^3}{8} + \frac{\lambda A^2}{2} - \frac{\lambda^2 A}{2}\right) \quad (10)$$

(where we use the symbol \asymp to mean that the expression on the left-hand side is bounded from above and below by a constant times the expression on the right-hand side), thus improving upon the Nachmias and Peres bound given in (8) and retrieving the asymptotic result of Pittel displayed in (7).

Subsequently, the methodology which lead to (10) was adapted by the same authors to study the near-critical $\mathbb{G}(n, d, p)$ random graph where $p = (1 + \lambda n^{-1/3})/(d-1)$ with $\lambda \in \mathbb{R}$ (possibly dependent on n) and $d \geq 3$ *fixed*. For this model, it was shown in [13] that there exist constants $A_0 > 0, n_0 \in \mathbb{N}$ such that, if $A_0 \leq A = A(n) = o(n^{1/30})$ and $\lambda = \lambda(n)$ satisfies $|\lambda| \leq A(1 - 2/d)[3(d-1)]^{-1}$, then for all $n \geq n_0$

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, d, p))| > An^{2/3}) \asymp \frac{c_2}{A^{3/2}} \exp\left(-\frac{A^3(d-1)(d-2)}{8d^2} + \frac{\lambda A^2(d-1)}{2d} - \frac{\lambda^2 A(d-1)}{2(d-2)}\right),$$

thus considerably improving upon the bound displayed in (9).

Remark 3. Observe that, in the particular case where $\lambda = 0$ (that is, in the purely critical regime), the constant $c(d) := (d-1)(d-2)/d^2$ which multiplies the leading term $A^3/8$ in the exponential above tends to 1 as $d \rightarrow \infty$. We suspect that this phenomenon is true more generally. Specifically, let \mathbb{G} be a random graph that, when considered at criticality of its parameter(s), exhibits maximal components containing $\asymp n^{2/3}$ vertices. We believe that, at criticality, there exist constants $c_1, c_2 > 0$ with $c_1 < c_2$ such that, for all large enough n and (not ‘too large’) $A = A(n)$,

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G})| > An^{2/3}) \asymp A^{-3/2} e^{-C^* \frac{A^3}{8}},$$

where $C^* = 1$ if $\mathbb{G} = \mathbb{G}(n, 1/n)$, otherwise C^* is a constant which depends on some parameter κ related to the random graph \mathbb{G} under consideration satisfying $C^*(\kappa) \rightarrow 1$ as $\kappa \rightarrow \infty$. Moreover, we further believe that (at criticality) there are constants $C, c > 0$ such that, for all large enough n and (not ‘too large’) $A = A(n)$,

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G})| < n^{2/3}/A) \leq e^{-cA^{3/2}}.$$

1.3 Some useful tools

Here we recall some results which we will use later on. We start by recalling a Chernoff bound for the Binomial distribution. The following version comes from [24].

Lemma 1.1 ([24, Theorem 2.1]). *Let $B_{N,P}$ be a binomial random variable of parameters N and p . Then for every $x \geq 0$ we have*

$$\mathbb{P}(B_{N,P} \geq NP + x) \leq \exp\left(-\frac{x^2}{2(NP + x/3)}\right).$$

Next we recall Doob's inequality and the Optional Stopping Theorem; proofs of both statements can be found in any advanced probability textbook.

Theorem 1.2. *Let $(M_t)_{t \in \mathbb{N}_0}$ be a positive submartingale. Then, for every $x > 0$, we have*

$$\mathbb{P}\left(\max_{t \in \{0\} \cup [n]} M_t \geq x\right) \leq \mathbb{E}[M_n]/x.$$

Theorem 1.3. *Let $(M_t)_{t \in \mathbb{N}_0}$ be a submartingale (resp. martingale, supermartingale) with respect to a filtration $(\mathcal{F}_t)_{t \in \mathbb{N}_0}$. Let σ_1 and σ_2 be bounded stopping times with respect to $(\mathcal{F}_t)_{t \in \mathbb{N}_0}$ such that $\sigma_1 \leq \sigma_2$. Then the random variables M_{σ_1} and M_{σ_2} are integrable and $\mathbb{E}[M_{\sigma_2} | \mathcal{F}_{\sigma_1}] \geq M_{\sigma_1}$ (resp. $\mathbb{E}[M_{\sigma_2} | \mathcal{F}_{\sigma_1}] = M_{\sigma_1}$, $\mathbb{E}[M_{\sigma_2} | \mathcal{F}_{\sigma_1}] \leq M_{\sigma_1}$).*

1.4 Notation and structure of the paper

Here we introduce the notation used throughout the article and describe the structure of the rest of the paper.

Notation. We let $\mathbb{N} = \{1, 2, \dots\}$ and define $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$. For $n \in \mathbb{N}$ we set $[n] := \{1, 2, \dots, n\}$. Given two sequences of real numbers $(x_n)_{n \geq 1}$ and $(y_n)_{n \geq 1}$ we write: (1) $x_n = O(y_n)$ if there exist $N \in \mathbb{N}$ and $C \in [0, \infty)$ such that $|x_n| \leq C|y_n|$ for all $n \geq N$; (2) either $x_n = o(y_n)$ or $x_n \ll y_n$ if $x_n/y_n \rightarrow 0$ as $n \rightarrow \infty$; and (3) either $x_n = \Theta(y_n)$ or $x_n \asymp y_n$ if $x_n = O(y_n)$ and $y_n = O(x_n)$. We often write c, C to denote constants in $(0, \infty)$, and use these letters many times in a single proof even though their actual values may change from line to line.

When talking about random variables, the abbreviation iid means *independent and identically distributed*. Moreover, given random variables X and Y , we write \leq_{sd} if $\mathbb{P}(X \geq z) \leq \mathbb{P}(Y \geq z)$ for every $z \in \mathbb{R}$, and say that X is stochastically dominated by Y . We write $\text{Bin}(N, q)$ to denote the binomial distribution of parameters $N \in \mathbb{N}_0, q \in [0, 1]$ and $\text{Unif}([0, 1])$ to denote the uniform distribution on $[0, 1]$. Given a distribution function F , we use the notation $X =_d F$ to mean that X has the distribution F .

Let $\mathbb{G} = (V, E)$ be any (undirected, possibly random) multigraph. Given two vertices $u, v \in V$, we write $u \sim v$ if $\{u, v\} \in E$ and say that vertices u and v are neighbours. We often write uv as shorthand for the edge $\{u, v\}$. We write $u \leftrightarrow v$ if there exists a path of occupied edges connecting vertices u and v , where we adopt the convention that $v \leftrightarrow v$ for every $v \in V$. We denote by $\mathcal{C}(v) := \{u \in V : u \leftrightarrow v\}$ the component containing vertex $v \in V$. We define a largest component $\mathcal{C}_{\max}(\mathbb{G})$ of \mathbb{G} to be some cluster $\mathcal{C}(v)$ for which $|\mathcal{C}(v)|$ is maximal, so that $|\mathcal{C}_{\max}(\mathbb{G})| = \max_{v \in V} |\mathcal{C}(v)|$.

Structure of the paper. The rest of the paper is organised as follows. In Section 2 we prove Theorem 1.1, whereas in Section 3 we describe how to modify the martingale argument of [27, 28] with the purpose of obtaining stretched exponential estimates in several critical models.

2 Proof of Theorem 1.1

Here the goal is to show that $|\mathcal{C}_{\max}(\mathbb{G})| \geq n^{2/3}/A$ with probability at least of order $1 - e^{-A^{3/2}}$ when either $\mathbb{G} = \mathbb{G}(n, p)$ and $p = (1 + \lambda n^{-1/3})/n$ or $\mathbb{G} = \mathbb{G}(n, d, p)$ and $p = (1 + \lambda n^{-1/3})/(d - 1)$ (with $\lambda \in \mathbb{R}, d \geq 3$ fixed).

We begin by providing an heuristic derivation of these estimates and we do so in terms of the $\mathbb{G}(n, p)$ random graph; the proof for the $\mathbb{G}(n, d, p)$ model follows the same line of thoughts.

2.1 Heuristic derivation of the bounds stated in Theorem 1.1

In order to study $|\mathcal{C}_{\max}(\mathbb{G}(n, p))|$ we use (as it is standard, see e.g. [20]) an *exploration process*, which is an algorithmic procedure to sequentially reveal the connected components of $\mathbb{G}(n, p)$ and that reduces the study of component sizes to the analysis of the trajectory of a (self-interacting) discrete-time, non-negative stochastic process.

Roughly speaking, an exploration process (in this setting) works as follows. We start by specifying an ordering of the vertices. Then, at time $t = 0$, one of the nodes, say u , is declared *active*, whereas all the remaining nodes are *unseen* and there are no *explored* vertices. At time $t = 1$, we reveal the *unseen* neighbours of u , which are declared active. The step terminates by declaring u explored. Then we iterate the procedure: at time $t = 2$, we select one of the active nodes (if any), say v , and reveal its unseen neighbours, which are declared active, whereas v itself becomes explored. If at the start of some step t the set of active vertices is *empty*, then we pick the first (with respect to the ordering fixed at the very beginning) *unseen* vertex (if any), say w , reveal its unseen neighbours, which are declared active and change the status of w to explored. The procedure terminates when *all* the vertices are in status explored.

We notice that the exploration of a connected component, started at some time t , continues as long as there are active vertices, say until time $t + k$, so that the size of the underlying cluster equals k , the number of steps of the procedure (starting from time t) for which the set of active vertices is non-empty.

Denote by Y_t the number of *active* vertices at the end of step t in the exploration process. Following the above description, we use such a procedure to bound (from above) the probability that $\mathcal{C}_{\max}(\mathbb{G}(n, p))$ contains less than $n^{2/3}/A$ vertices by the probability that the positive excursions of Y_t never last for more than $n^{2/3}/A$ steps.

More precisely, defining t_i to be the first time $t > t_{i-1}$ at which $Y_t = 0$ (prior to the end of the procedure) and denoting by \mathcal{C}_i the i -th explored cluster, whose exploration starts at time t_i so that $|\mathcal{C}_i| = t_i - t_{i-1}$, we can bound from above the probability that $\mathcal{C}_{\max}(\mathbb{G}(n, p))$ contains less than $n^{2/3}/A$ vertices by the probability that each excursion length $t_i - t_{i-1}$ is at most $n^{2/3}/A$:

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, p))| < n^{2/3}/A) = \mathbb{P}(t_i - t_{i-1} < T \forall i \in [N]), \quad (11)$$

where we denote by N the (random) number of clusters in $\mathbb{G}(n, p)$ (i.e. the number of excursions needed to reveal all the components in the graph).

Let us start by making a simple, yet useful observation. On the event where $t_i - t_{i-1} < n^{2/3}/A$ for each i , we have

$$n = \sum_{i=1}^N |\mathcal{C}_i| = \sum_{i=1}^N (t_i - t_{i-1}) < N \frac{n^{2/3}}{A},$$

whence $N > An^{1/3} =: L$. Therefore, setting $T := n^{2/3}/A$ for ease of notation, we can write

$$\mathbb{P}(t_i - t_{i-1} < T \forall i \in [N]) \leq \mathbb{P}(t_i - t_{i-1} < T \forall L/2 \leq i \leq L)$$

and

$$\mathbb{P}(t_i - t_{i-1} < T \forall L/2 \leq i \leq L) = \prod_{i=L/2}^L \left(1 - \mathbb{P}(t_i - t_{i-1} \geq T | t_j - t_{j-1} < T \forall L/2 \leq j \leq i-1)\right). \quad (12)$$

We claim that, for *each* $L/2 \leq i \leq L$,

$$\mathbb{P}(t_i - t_{i-1} \geq T | t_j - t_{j-1} < T \forall L/2 \leq j \leq i-1) \gtrsim T^{-1/2}. \quad (13)$$

Observe that, if (13) were true, then (recalling the value of T and using the classical inequality $1 + x \leq e^x$) the expression on the right-hand side of (12) would satisfy

$$\begin{aligned} & \prod_{i=L/2}^L \left(1 - \mathbb{P}(t_i - t_{i-1} \geq T | t_j - t_{j-1} < T \forall L/2 \leq j \leq i-1)\right) \\ & \lesssim \left(1 - T^{-1/2}\right)^{L/2} \leq \exp(-L/(2T^{1/2})) \lesssim \exp(-A^{3/2}), \end{aligned}$$

which is the desired bound. Hence we are left with the problem of showing (13). To this end, recall the description of the exploration process given earlier. It is clear that, if at time $t - 1$ there is at least one active node then, denoting by η_t the number of *unseen* vertices which become active at step t , we have $Y_t = Y_{t-1} + \eta_t - 1$: indeed, the number of active vertices at time t equals the number of active nodes at time $t - 1$, plus the newly discovered (necessarily unseen) neighbours of the vertex under investigation at step t , minus one (since the status of the vertex under exam passes from active to explored). Consequently, if the number of active vertices stays positive from a time s until a time $s + t$, then we can write $Y_{s+t} = 1 + \sum_{k=1}^t (\eta_{s+k} - 1)$. Moreover, conditional on the history of the exploration process until time $s + k - 1$, it is clear from the model definition and the description of the exploration process that the random variable η_{s+k} has the $\text{Bin}(n - s - k + 1 - Y_{s+k-1}, p)$ distribution (where $n - s - k + 1 - Y_{s+k-1}$ is the number of unseen vertices at the start of step t).

Keeping these considerations in mind and ignoring the term Y_{s+k-1} from the first parameter of the binomial distribution (this approximation is justified by the fact that the number of active vertices is never ‘too large’), we write

$$\mathbb{P}(t_i - t_{i-1} \geq T | t_j - t_{j-1} < T \ \forall L/2 \leq j \leq i - 1) \approx \mathbb{P}\left(\sum_{k=1}^t (\text{Bin}_k(n - t_{i-1} - k, p) - 1) > 0 \ \forall t \leq T\right).$$

Let us *assume* that $t_L \lesssim LT^{1/2}$; then, for each $L/2 \leq i \leq L$, the exploration of the i -th cluster \mathcal{C}_i starts before time $LT^{1/2}$, which implies that, when starting the exploration of \mathcal{C}_i , the number of explored vertices is $\lesssim LT^{1/2}$ (for each $L/2 \leq i \leq L$). Under this assumption, after noticing that $k \leq T = n^{2/3}/A \ll A^{1/2}n^{2/3} = LT^{1/2}$ (whence $LT^{1/2} + k \leq 2LT^{1/2}$ for all large enough n), we can couple the $\text{Bin}_k(n - t_{i-1} - k, p)$ random variables appearing within the above probability with independent $\text{Bin}_k(n - 2LT^{1/2}, p)$ random variables in such a way that

$$\text{Bin}_k(n - t_{i-1} - k, p) \geq \text{Bin}_k(n - 2LT^{1/2}, p) \text{ for each } k \in [T].$$

Hence we can bound

$$\mathbb{P}\left(\sum_{k=1}^t (\text{Bin}_k(n - t_{i-1} - k, p) - 1) > 0 \ \forall t \leq T\right) \gtrsim \mathbb{P}\left(\sum_{k=1}^t (\text{Bin}_k(n - 2LT^{1/2}, p) - 1) > 0 \ \forall t \leq T\right). \quad (14)$$

Now the process considered in the last probability is a random walk with a *small* negative drift. Such a small drift does not affect the behaviour of the process, which from a practical point of view can be regarded as a \mathbb{Z} -valued, mean-zero random walk. More specifically, we can approximate the probability in (14) by $\mathbb{P}(S_t > 0 \ \forall t \leq T)$, where S_t a random walk with iid $\text{Bin}(n, p) - 1$ increments. It is well-known that a mean-zero random walk with iid increments having a finite second moment stays above zero for T steps with probability $\asymp T^{-1/2}$, thus establishing the claim (13).

2.2 Proof of Theorem 1.1: the $\mathbb{G}(n, p)$ model

We start by providing a formal description of the exploration process introduced in Section 2.1. Fix an ordering of the n vertices with v first. Let us denote by \mathcal{A}_t , \mathcal{U}_t and \mathcal{E}_t the (random) sets of active, unseen and explored vertices at the end of step $t \in \mathbb{N}_0$, respectively. Then, for any given $t \in \mathbb{N}_0$, we can partition the vertex set as $[n] = \mathcal{A}_t \cup \mathcal{U}_t \cup \mathcal{E}_t$ (a disjoint union), so that in particular $\mathcal{U}_t = [n] \setminus (\mathcal{A}_t \cup \mathcal{E}_t)$ at each step t .

Algorithm 1. At time $t = 0$, vertex v is declared active whereas all other vertices are declared unseen, so that $\mathcal{A}_0 = \{v\}$, $\mathcal{U}_0 = [n] \setminus \{v\}$ and $\mathcal{E}_0 = \emptyset$. For every $t \in \mathbb{N}$, the algorithm proceeds as follows.

- (a) If $|\mathcal{A}_{t-1}| \geq 1$, we let u_t be the first active vertex (here and in what follows, the term *first* refers to the ordering that we have fixed at the beginning of the procedure).
- (b) If $|\mathcal{A}_{t-1}| = 0$ and $|\mathcal{U}_{t-1}| \geq 1$, we let u_t be the first unseen vertex.
- (c) If $|\mathcal{A}_{t-1}| = 0 = |\mathcal{U}_{t-1}|$ (so that $\mathcal{E}_{t-1} = [n]$), we halt the procedure.

Denote by \mathcal{D}_t the (random) set of *unseen* neighbours of u_t , i.e. we set $\mathcal{D}_t := \{x \in \mathcal{U}_{t-1} \setminus \{u_t\} : u_t \sim x\}$ and note that $\mathcal{U}_{t-1} \setminus \{u_t\} = \mathcal{U}_{t-1}$ if $\mathcal{A}_{t-1} \neq \emptyset$, since $u_t \in \mathcal{A}_{t-1}$ in this case. Then we update $\mathcal{U}_t := \mathcal{U}_{t-1} \setminus (\mathcal{D}_t \cup \{u_t\})$, $\mathcal{A}_t := (\mathcal{A}_{t-1} \setminus \{u_t\}) \cup \mathcal{D}_t$ and $\mathcal{E}_t := \mathcal{E}_{t-1} \cup \{u_t\}$.

Remark 4. Note that, since in the procedure **Algorithm 1** we explore *one vertex* at each step, we have $\mathcal{A}_t \cup \mathcal{U}_t \neq \emptyset$ for every $t \leq n - 1$ and $\mathcal{A}_n \cup \mathcal{U}_n = \emptyset$ (as $\mathcal{E}_n = [n]$). Thus the algorithm runs for n steps.

Denoting by η_t the (random) number of unseen vertices that we add to the set of active nodes at time t , since at the end of each step i in which $|\mathcal{A}_{i-1}| \geq 1$ we remove the (active) vertex u_i from \mathcal{A}_{i-1} (after having revealed its unseen neighbours), we have the recursion

- $|\mathcal{A}_t| = |\mathcal{A}_{t-1}| + \eta_t - 1$, if $|\mathcal{A}_{t-1}| > 0$;
- $|\mathcal{A}_t| = \eta_t$, if $|\mathcal{A}_{t-1}| = 0$.

We let $t_0 := 0$ and recursively we define t_i to be the first time after t_{i-1} at which the number of active vertices hits zero; that is,

$$t_i := \min\{t \geq t_{i-1} + 1 : Y_t = 0\}, \text{ for } i \in [N], \quad (15)$$

where we denote by N the (random) number of excursions (i.e. the number of clusters). Setting $T := \lceil n^{2/3}/A \rceil$, we write

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, p))| < n^{2/3}/A) \leq \mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, p))| < T) = \mathbb{P}(t_i - t_{i-1} < T \forall i \in [N]).$$

As remarked in Section 2.1, if $t_i - t_{i-1} < T$ for all $i \in [N]$ then $n = \sum_{i=1}^N (t_i - t_{i-1}) < NT$ and hence $N > n/T$. Setting $L := \lfloor n/(2T) \rfloor$, we see that $N > 2L$. Hence we can bound

$$\mathbb{P}(t_i - t_{i-1} < T \forall i \in [N]) \leq \mathbb{P}(t_i - t_{i-1} < T \forall L + 1 \leq i \leq 2L). \quad (16)$$

Let us introduce the following (good) event:

$$\mathcal{G}_j = \mathcal{G}_j(B) := \{t_j \leq BLT^{1/2}\},$$

where $L \leq j \leq 2L$ and B is some large positive constant (independent of A, n, λ). The event \mathcal{G}_j tells that we start exploring the j -th cluster before time $BLT^{1/2}$, which also means that when starting exploring the j -th cluster, the number of explored vertices is at most $BLT^{1/2}$ (recall that at each step of the exploration exactly *one* vertex becomes explored). A good control on the t_j 's is needed when bounding from *below* the probability that $t_i - t_{i-1}$ is larger than T , which is the same as the probability that the number of active vertices stays above zero for T steps starting from time t_{i-1} . If the number of explored vertices were 'too large' when starting the exploration of \mathcal{C}_i , then the latter probability would be 'too small', because the negative drift of the increments $\eta_{t_{i-1}+k} - 1$ would be 'too large' (in absolute value) to keep the process above zero for T steps.

Going back to (16), we continue by writing

$$\mathbb{P}(t_i - t_{i-1} < T \forall L + 1 \leq i \leq 2L) \leq \mathbb{P}(\{t_i - t_{i-1} < T \forall L + 1 \leq i \leq 2L\} \cap \mathcal{G}_{2L}) + \mathbb{P}(\mathcal{G}_{2L}^c).$$

The next lemma controls the probability of the (bad) event \mathcal{G}_{2L}^c .

Lemma 2.1. *There exist positive constants $B_0, A_0 = A_0(\lambda) > 0$ such that, for any $A \geq A_0, B \geq B_0$ and for all large enough n , we have*

$$\mathbb{P}(\mathcal{G}_{2L}^c) \leq \exp(-cA^{3/2}),$$

for some positive constant $c = c(B) > 0$.

Therefore, in what follows, we can focus on the probability

$$\mathbb{P}(\{t_i - t_{i-1} < T \forall L + 1 \leq i \leq 2L\} \cap \mathcal{G}_{2L}). \quad (17)$$

Let us define $\mathcal{F}_j := \mathcal{F}_{t_j}$ for $j \in [N]$, the σ -algebra consisting of all the information collected by the exploration process by time t_j , i.e. by the time at which we start the exploration of the j -th component.

With the next result we show that, on the event \mathcal{G}_{i-1} , we have $t_i - t_{i-1} \geq T$ with probability $\gtrsim T^{-1/2}$ for each $L + 1 \leq i \leq 2L$.

Lemma 2.2. *There exist positive constants $B_0, A_0 = A_0(\lambda) > 0$ such that, for any $A \geq A_0, B \geq B_0$ and for all large enough n , on the event \mathcal{G}_{i-1} we have*

$$\mathbb{P}(t_i - t_{i-1} \geq T | \mathcal{F}_{i-1}) \geq \frac{c_0}{T^{1/2}}$$

for each $L + 1 \leq i \leq 2L$, for some constant $c_0 = c_0(B) > 0$.

It is now very easy to show that the expression in (17) is at most $\exp(-cA^{3/2})$ (for some $c > 0$). Indeed, using the tower property of conditional expectation, together with the fact that $\mathcal{G}_j \subset \mathcal{G}_{j-1}$, we can write

$$\begin{aligned} & \mathbb{P}(\{t_i - t_{i-1} < T \ \forall L + 1 \leq i \leq 2L\} \cap \mathcal{G}_{2L}) \\ & \leq \mathbb{E} \left[\mathbb{1}_{\{t_i - t_{i-1} < T \ \forall L + 1 \leq i \leq 2L - 1\} \cap \mathcal{G}_{2L-1}} (1 - \mathbb{P}(t_{2L} - t_{2L-1} \geq T | \mathcal{F}_{2L-1})) \right]. \end{aligned} \quad (18)$$

Thanks to Lemma 2.2, the expression on the right-hand side of (18) is at most

$$(1 - c_0 T^{-1/2}) \mathbb{P}(\{t_i - t_{i-1} < T \ \forall L + 1 \leq i \leq 2L - 1\} \cap \mathcal{G}_{2L-1}).$$

Iterating and using the classical inequality $1 + x \leq e^x$ (which is valid for all $x \in \mathbb{R}$), we conclude that for all large enough A and n

$$\mathbb{P}(\{t_i - t_{i-1} < T \ \forall L + 1 \leq i \leq 2L\} \cap \mathcal{G}_{2L}) \leq (1 - c_0 T^{-1/2})^L \leq \exp(-c_0 L T^{-1/2}). \quad (19)$$

Recalling the definition of L and T , it is easy to see that $L T^{-1/2} \geq A^{3/2}/4$ for all large enough n (the constant 4 is clearly not best possible but sufficient for our purposes) and hence, putting all pieces together, we conclude that there is a constant $c > 0$ such that, for all large enough n ,

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, p))| < n^{2/3}/A) \leq \exp(-cA^{3/2}).$$

This concludes the proof of Theorem 1.1 for the $\mathbb{G}(n, p)$ random graph. What is left to do is to prove Lemmas 2.1 and 2.2; this is done in the following section.

2.2.1 Proofs of Lemmas 2.1 and 2.2

In this section we establish the two auxiliary facts which were stated, without proof, while proving the main theorem for the $\mathbb{G}(n, p)$ random graph.

Proof of Lemma 2.1. Recall that $\mathcal{G}_{2L} := \{t_{2L} \leq BLT^{1/2}\}$. To bound from above the probability of the complementary event, we proceed as follows. Let $S_0 := 1$ and define recursively $S_t := S_{t-1} + \eta_t - 1$, where the random variable η_i , conditional on the history of the exploration process until time $i - 1$ (denoted here by \mathcal{H}_{i-1}), has the $\text{Bin}(n - (i - 1) - Y_{i-1}, p)$ distribution.

The \mathbb{Z} -valued random process $(S_t)_{t \in \mathbb{N}_0}$ is closely related to the process of active vertices $(Y_t)_{t \in \mathbb{N}_0}$; in particular, they coincide up to time t_1 (the time at which we completes the exploration of the first cluster). However, at time $t_1 + 1$, we have $Y_{t_1+1} = \eta_{t_1+1}$ whereas $S_{t_1+1} = \eta_{t_1+1} - 1$. The two processes Y_{t_1+t} and S_{t_1+t} have the same increments for $t_1 + 1 < t \leq t_2$, but then again $Y_{t_2+1} = \eta_{t_2+1}$ whereas $S_{t_2+1} = \eta_{t_2+1} - 2$, and so on

Then $(-\min_{j \in [t]} S_j) + 1$ denotes the number of disjoint clusters that have been fully explored by time $t \in [n]$ (see e.g. Chapter 5.2.4 in [20]). When $Y_t = 0$, we have fully explored a cluster, and this occurs precisely when $S_t = \min_{j \in [t-1]} S_j - 1$, i.e., the process $(S_t)_{t \in [n]}$ reaches a new all-time minimum.

Now observe that, if $t_{2L} > BLT^{1/2}$, then the number of components that are fully explored up to time $\lfloor BLT^{1/2} \rfloor$ is smaller than $2L$ (exactly because t_{2L} is the time at which we start exploring the $(2L)$ -th cluster). Therefore we can bound

$$\mathbb{P}(\mathcal{G}_{2L}^c) \leq \mathbb{P}\left(-\min_{j \leq \lfloor BLT^{1/2} \rfloor} S_j < 2L - 1\right) = \mathbb{P}\left(\max_{j \leq \lfloor BLT^{1/2} \rfloor} (-S_j) < 2L - 1\right) \leq \mathbb{P}(S_{\lfloor BLT^{1/2} \rfloor} > -2L). \quad (20)$$

Now, by definition of S_i , we have

$$S_{\lfloor BLT^{1/2} \rfloor} = 1 + \sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} (\eta_i - 1) \text{ and } (\eta_i | \mathcal{H}_{i-1}) =_d \text{Bin}(n - (i - 1) - Y_{i-1}, p) \leq_{sd} \text{Bin}(n - (i - 1), p).$$

By coupling the η_i with independent $\text{Bin}(n - (i - 1), p)$ random variables δ_i in such a way that $\eta_i \leq \delta_i$ for each i , we arrive at

$$\mathbb{P}(S_{\lfloor BLT^{1/2} \rfloor} > -2L) \leq \mathbb{P}\left(\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} (\delta_i - 1) > -2L\right).$$

Now let δ'_i be independent random variables with the $\text{Bin}(i - 1, p)$ distribution, also independent of the δ_i . Then, setting $\xi_i := \delta_i + \delta'_i \stackrel{d}{=} \text{Bin}(n, p)$ for each i , we see that

$$\mathbb{P}\left(\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} (\delta_i - 1) > -2L\right) = \mathbb{P}\left(\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} (\xi_i - 1) > \sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \delta'_i - 2L\right). \quad (21)$$

Since

$$\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \delta'_i \stackrel{d}{=} \text{Bin}\left(\frac{M}{2}, p\right) \text{ where } M := (\lfloor BLT^{1/2} \rfloor - 1)(\lfloor BLT^{1/2} \rfloor - 2) \sim B^2 L^2 T,$$

we can use Lemma 1.1 to bound

$$\mathbb{P}\left(\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \delta'_i \leq (Mp)/2 - L\right) \leq \exp\left(-c \frac{L^2}{Mp}\right) \leq \exp\left(-\frac{c}{Tp}\right)$$

where $c = c(B) > 0$ is a constant which depends on B . Since $TP = O(A^{-1}n^{-1/3})$,

$$\exp\left(-\frac{c}{Tp}\right) \leq \exp(-cAn^{1/3}).$$

Therefore, going back to (21) and setting

$$\ell := \lfloor BLT^{1/2} \rfloor(1 - np) + (Mp)/2 - 3L,$$

we can bound

$$\begin{aligned} \mathbb{P}\left(\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} (\xi_i - 1) > \sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \delta'_i - 2L\right) \\ \leq \mathbb{P}\left(\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} (\xi_i - 1) > (Mp)/2 - 3L\right) + \exp(-cAn^{1/3}) \\ = \mathbb{P}\left(\text{Bin}(n\lfloor BLT^{1/2} \rfloor, p) > np\lfloor BLT^{1/2} \rfloor + \ell\right) + \exp(-cAn^{1/3}). \end{aligned} \quad (22)$$

Now note that, by taking a large enough A (in particular, $A > B^2\lambda^2$ suffices) we obtain

$$\ell \geq (Mp)/2 - \left(3L + \frac{|\lambda|\lfloor BLT^{1/2} \rfloor}{n^{1/3}}\right) \geq (Mp)/2 - 4L.$$

Therefore the last probability in (22) is at most

$$\mathbb{P}\left(\text{Bin}(n\lfloor BLT^{1/2} \rfloor, p) > np\lfloor BLT^{1/2} \rfloor + (Mp)/2 - 4L\right). \quad (23)$$

A simple computation shows that, for all large enough A and n ,

$$(Mp)/2 - 4L \geq \frac{B^2 L^2 T}{2n} - 5L = L\left(\frac{B^2 LT}{2n} - 5\right) \geq L(B^2/2 - 5 - o(1)) \geq L(B^2/2 - 6),$$

which is positive provided that B is large enough. Hence, using once again Lemma 1.1, we can bound from above the probability in (23) by

$$\mathbb{P}\left(\text{Bin}(n\lfloor BLT^{1/2} \rfloor, p) > np\lfloor BLT^{1/2} \rfloor + L(B^2/2 - 6)\right) \leq \exp\left(-c \frac{L}{T^{1/2}}\right) \leq \exp(-cA^{3/2}) \quad (24)$$

for some constant $c = c(B) > 0$. Since $An^{1/3} \gg A^{3/2}$ (because of our assumption $A \ll n^{2/3}$), we see that the term $\exp(-cAn^{1/3})$ in (22) is much smaller than $\exp(-cA^{3/2})$ which appears on the right-hand side of (24), whence the desired result follows.

Proof of Lemma 2.2. Here the goal is to show that there is a constant $c_0 = c_0(B) > 0$ such that, for each $L + 1 \leq i \leq 2L$, on the event \mathcal{G}_{i-1} it holds that

$$\mathbb{P}(t_i - t_{i-1} \geq T | \mathcal{F}_{i-1}) \geq \frac{c_0}{T^{1/2}}$$

for all large enough n , where we recall that $\mathcal{F}_{i-1} = \mathcal{F}_{t_{i-1}}$ is the σ -algebra consisting of all the information gathered by the exploration process until time t_{i-1} (i.e. until the moment at which the procedure finishes revealing the $(i-1)$ -th cluster). Then, setting $Y_t^i := Y_{t_i+t}$ and $\eta_t^i := \eta_{t_i+t}$ to simplify notation, we can write

$$\mathbb{P}(t_i - t_{i-1} \geq T | \mathcal{F}_{i-1}) = \mathbb{P}(Y_t^i > 0 \forall t < T | \mathcal{F}_{i-1}) \geq \mathbb{P}\left(1 + \sum_{k=1}^t (\eta_k^i - 1) > 0 \forall t \leq T | \mathcal{F}_{i-1}\right), \quad (25)$$

where we recall that, setting $\mathcal{F}_{i-1}(k-1) := \mathcal{F}_{t_{i-1}+k-1}$, we have

$$(\eta_k^i | \mathcal{F}_{i-1}(k-1)) =_d \text{Bin}_k(n - t_{i-1} - k + 1 - Y_{k-1}^i, p). \quad (26)$$

Before proceeding with the actual proof, let us make a few observations and give some hints on how we intend to proceed.

Note that the η_k^i are not independent, which makes our analysis non-trivial. The plan is to replace these random variables with a sequence of *independent* variables which are easier to analyse. To this end we first observe that, on \mathcal{G}_{i-1} , we have $t_{i-1} < BLT^{1/2}$; moreover, recalling that $L \sim An^{1/3}$, $T \sim n^{2/3}/A$ and $A \ll n^{2/3}$ by assumption, we have $k \leq t < T \ll LT^{1/2} \ll n$. If we can show that Y_{k-1}^i is at most $LT^{1/2}$ too, then we would obtain $t_{i-1} + k + Y_{k-1}^i \leq 3BLT^{1/2}$ and we could couple the $\text{Bin}_k(n - t_{i-1} - k + 1 - Y_{k-1}^i, p)$ random variables with independent $\text{Bin}_k(n - 3BLT^{1/2}, p)$ random variables in such a way that

$$\text{Bin}_k(n - t_{i-1} - k + 1 - Y_{k-1}^i, p) \geq \text{Bin}_k(n - 3BLT^{1/2}, p),$$

thus removing the two sources of randomness from the first parameter of the (conditional) distribution in (26). At this stage, we would have a random walk $\sum_{k=1}^t [\text{Bin}_k(n - 3BLT^{1/2}, p) - 1]$ with independent and identically distributed increments, having a *small* negative drift; the drift is so small that it does not impact much the behaviour of the process, which can be regarded as a random walk with iid, mean zero increments having finite second moment. Putting all pieces together, we would bound from below the expression in (25) by the probability that a mean-zero \mathbb{Z} -valued random walk with the latter properties stays above zero for T steps, which is known to be $\asymp T^{-1/2}$ (as desired).

We now make rigorous the above simple strategy, and start by replacing the η_k^i with iid random variables having the $\text{Bin}(n - 3L\lceil BT^{1/2} \rceil, p)$ law.

The next result is basically just a rephrasing of Lemma 4.1 in [12] adapted to our current need; we include a proof for completeness.

Lemma 2.3. *There exists a sequence $(\delta_k)_{k \leq T}$ of independent random variables, with $\delta_k =_d \text{Bin}(n - 3L\lceil BT^{1/2} \rceil, p)$ for each k , such that on the event \mathcal{G}_{i-1} it holds that*

$$\mathbb{P}\left(1 + \sum_{k=1}^t (\eta_k^i - 1) > 0 \forall t \leq T | \mathcal{F}_{i-1}\right) \geq \mathbb{P}\left(1 + \sum_{k=1}^t (\delta_k - 1) > 0 \forall t \leq T\right) - \exp(-cLT^{1/2})$$

for all large enough n , for some constant $c = c(B) > 0$.

Proof. Recall that u_k is the vertex that is explored at step t in the exploration of $\mathbb{G}(n, p)$. Denoting by X_k^v the indicator that vertex u_k is a neighbour of $v \in [n]$, we note that the random variable η_k^0 can be expressed as

$$\eta_k^i = \sum_{v \in [n] \setminus (\mathcal{A}_{k-1}^i \cup \mathcal{E}_{k-1}^i)} X_k^v,$$

with $\mathcal{E}_j^i := \mathcal{E}_{t_{i-1}+j}$ being the set of explored vertices at time $t_{i-1}+j$. Let us denote by $\mathcal{R}_j^i := \mathcal{A}_j^i \cup \mathcal{E}_j^i$ the set of vertices which are either active or explored at time j (we use \mathcal{R} for ‘revealed’). For convenience, we also give a name to the set of *unseen* vertices which become active at step j of the exploration of \mathcal{C}_{i-1}

(i.e. at time $t_{i-1} + j$); denote such a set by $\mathcal{A}_j^{i,*}$, so that then $\mathcal{R}_{k-1}^i = \bigcup_{j=0}^{k-1} \mathcal{A}_j^{i,*}$ (with $\mathcal{A}_0^{i,*} = \{u_{t_{i-1}+1}\}$, the vertex from which we start the exploration of \mathcal{C}_i).

For each $1 \leq k \leq T$, if $|\mathcal{R}_{k-1}^i| < 3L\lceil BT^{1/2} \rceil$ (which occurs if $Y_{k-1}^i < BLT^{1/2}$, because $|\mathcal{R}_{k-1}^i| = Y_{k-1}^i + t_{i-1} + k - 1 < Y_{k-1}^i + 2BLT^{1/2}$ on the event \mathcal{G}_{i-1}) we let $\mathcal{B}_{k-1}^{i,*}$ be any subset of the vertices $[n]$ such that

$$\mathcal{A}_{k-1}^{i,*} \subset \mathcal{B}_{k-1}^{i,*}; \quad \mathcal{B}_{k-1}^{i,*} \cap \mathcal{R}_{k-2}^i = \emptyset; \quad |\mathcal{B}_{k-1}^{i,*} \cup \mathcal{R}_{k-2}^i| = 3L\lceil BT^{1/2} \rceil \ll n.$$

If $|\mathcal{R}_{k-1}^i| \geq 3L\lceil BT^{1/2} \rceil$ then we simply let $\mathcal{B}_{k-1}^{i,*} = \mathcal{A}_{k-1}^{i,*}$. Next we define

$$h_k := |\mathcal{B}_{k-1}^{i,*} \cup \mathcal{R}_{k-2}^i| - 3L\lceil BT^{1/2} \rceil \geq 0.$$

Take a (doubly-infinite) sequence $(I_j^k : j, k \in \mathbb{N})$ of independent Bernoulli random variables of parameter p , also independent of everything else. Then define

$$\delta_k := \sum_{v \in [n] \setminus (\mathcal{B}_{k-1}^{i,*} \cup \mathcal{R}_{k-2}^i)} X_v^v + \sum_{i=1}^{h_k} I_j^k.$$

Note that $|[n] \setminus (\mathcal{B}_{k-1}^{i,*} \cup \mathcal{R}_{k-2}^i)| = n - 3L\lceil BT^{1/2} \rceil$ by construction and hence

$$|[n] \setminus (\mathcal{B}_{k-1}^{i,*} \cup \mathcal{R}_{k-2}^i)| + h_k = n - 3L\lceil BT^{1/2} \rceil.$$

Moreover, the random variables δ_k are independent because $\{X_v^k : v \in (R_{k-2}^i)^c\}$ are independent and independent of $\{X_v^j : v \in (R_{j-2}^i)^c\}$ for any $j \neq k$; they are also independent of the indicators associated to the edges revealed up to time t_{i-1} , whence independent of \mathcal{F}_{i-1} .

We also observe that if $|\mathcal{R}_{k-1}^i| < 3L\lceil BT^{1/2} \rceil$, then $|\mathcal{B}_{k-1}^{i,*} \cup \mathcal{R}_{k-2}^i| = 3L\lceil BT^{1/2} \rceil$ and so $h_k = 0$. Since we also have $\mathcal{A}_{k-1}^{i,*} \subset \mathcal{B}_{k-1}^{i,*}$ by construction, we see that if $|\mathcal{R}_{k-1}^i| < 3L\lceil BT^{1/2} \rceil$ then $\eta_k^i \geq \delta_k$. Thus, recalling that t_i denotes the first time at which the set of active vertices becomes empty for the first time after t_{i-1} , we see that the probability which appears in the statement of the lemma is at least

$$\mathbb{P}\left(1 + \sum_{k=1}^t (\delta_k - 1) > 0 \forall t \leq T, Y_k^i < BLT^{1/2} \forall k < (t_i - t_{i-1}) \wedge T | \mathcal{F}_{i-1}\right), \quad (27)$$

where the last inequality follows from the fact that, on the event $\{Y_k^i < BLT^{1/2} \forall k < T\}$, we have $|\mathcal{R}_{k-1}^i| < 3L\lceil BT^{1/2} \rceil$ for each k as discussed earlier, and so $\eta_k^i \geq \delta_k$ for $k < T$ (and $(t_i - t_{i-1}) \wedge T = T$ on the event $\{1 + \sum_{k=1}^t (\eta_k^i - 1) > 0 \forall t \leq T - 1\}$). To conclude, we observe that the probability on the right-hand side of (27) is at least

$$\mathbb{P}\left(1 + \sum_{k=1}^t (\delta_k - 1) > 0 \forall t \leq T\right) - \mathbb{P}\left(\exists k < (t_i - t_{i-1}) \wedge T : Y_k^i \geq BLT^{1/2} | \mathcal{F}_{i-1}\right),$$

where we have used the above mentioned independence for the last inequality. There remains to bound from above the second probability on the right-hand side of the last expression, which is easily done. Indeed, using a union bound together with the fact that $Y_k^i = 1 + \sum_{j=1}^k (\eta_j^i - 1)$ for $k \leq t_i - t_{j-1}$, we can write

$$\mathbb{P}\left(\exists k < (t_i - t_{i-1}) \wedge T : Y_k^i \geq BLT^{1/2} | \mathcal{F}_{i-1}\right) \leq \sum_{k=1}^{T-1} \mathbb{P}\left(1 + \sum_{j=1}^k (\eta_j^i - 1) \geq BLT^{1/2} | \mathcal{F}_{i-1}\right).$$

By coupling the η_k^i with independent $\text{Bin}(n, p)$ random variables ξ_k in such a way that $\eta_k^i \leq \xi_k$ and using Lemma 1.1, we see that the probability on the right-hand side of the last expression is at most

$$\sum_{k=1}^{T-1} \exp\left(-c \frac{B^2 i^2 T}{knp + BiT^{1/2}}\right) \leq \exp\left(-cLT^{1/2}[1 - \log(T)/(cLT^{1/2})]\right) \leq \exp\left(-cLT^{1/2}\right),$$

where we have used that $\log(T) \leq T \ll LT^{1/2}$. \square

In order to bound from below the probability that the random walk formed by the δ_k stays above zero for T steps (see Lemma 2.3), we use the following result, which is taken from [12].

Lemma 2.4. [Lemma 4.6 in [12]] Take $n \in \mathbb{N}$, $h_n \geq 0$, $a_n \in (-1, \infty)$ satisfying $na_n \in \mathbb{Z}$, $b_n \in (-1, n-1)$ and $t_n \in \mathbb{N}$. Suppose that $M_t = 1 + \sum_{i=1}^t (W_i - 1)$ where the W_i are independent $\text{Bin}(n(1+a_n), (1+b_n)/n)$ random variables. Let $\mu_n = (1+a_n)(1+b_n)$. Then

$$\begin{aligned} & \mathbb{P}(M_t > 0 \quad \forall t \in [t_n], M_{t_n} \in [h_n, 2h_n]) \\ & \geq (\mu_n \wedge 1)^{2h_n} \mu_n^{t_n-1} e^{(1-\mu_n)t_n} \mathbb{P}(\hat{M}_t > 0 \quad \forall t \in [t_n], \hat{M}_{t_n} \in [h_n, 2h_n]) - \frac{t_n}{n} (1+a_n)(1+b_n)^2 \end{aligned}$$

where $\hat{M}_t = 1 + \sum_{i=1}^t (\hat{W}_i - 1)$, and $(\hat{W}_i)_{i=1}^{t_n}$ is a sequence of independent Poisson random variables with mean one.

We apply this lemma with

$$a_n := -3L \lceil BT^{1/2} \rceil / n \text{ and } b_n := \lambda / n^{1/3}.$$

Note that clearly $na_n \in \mathbb{Z}$ and, since $LT^{1/2}/n \sim A^{1/2}/n^{1/3} \ll 1$ (as we assumed that $A \ll n^{2/3}$), we also have $a_n > -1$ for all sufficiently large n . Moreover, it is clear that $b_n \in (-1, n-1)$ provided n is large enough.

Now, setting $\mu_n := (1+a_n)(1+b_n)$ as in the statement of Lemma 2.4, it is not difficult to see that μ_n is negative provided $A \geq A_0$ for a large enough $A_0 = A_0(\lambda)$ which depends on λ and moreover

$$|1 - \mu_n| = O\left(\frac{LT^{1/2}}{n}\right) \asymp \frac{A^{1/2}}{n^{1/3}} \ll 1.$$

Hence, as $n \rightarrow \infty$, we can write

$$\log(\mu_n) = \log(1 - (1 - \mu_n)) = -(1 - \mu_n) + O((1 - \mu_n)^2). \quad (28)$$

Now using the above Taylor expansion we see that, for all large enough n ,

$$(\mu_n \wedge 1)^{2 \lceil T^{1/2} \rceil} \mu_n^{T-1} e^{(1-\mu_n)T} \geq c \mu_n^{2 \lceil T^{1/2} \rceil} \exp(-cT(1-\mu_n)^2) \geq c \mu_n^{2 \lceil T^{1/2} \rceil},$$

where the last inequality follows from the fact that $T(1-\mu_n)^2 = O(1)$. Moreover, by expressing $\mu_n^{2 \lceil T^{1/2} \rceil} = e^{2 \lceil T^{1/2} \rceil \log(\mu_n)}$ and using once more the expansion in (28), we conclude that

$$(\mu_n \wedge 1)^{2 \lceil T^{1/2} \rceil} \mu_n^{T-1} e^{(1-\mu_n)T} \geq c$$

for some positive constant $c = c(B) > 0$ which depends solely on B . But then we can bound, using Lemma 2.4 (note that for us $t_n = T$ and $h_n = \lceil T^{1/2} \rceil$)

$$\begin{aligned} & \mathbb{P}\left(\sum_{k=1}^t (\delta_k - 1) > 0 \quad \forall t \leq T\right) \\ & \geq \mathbb{P}\left(\sum_{k=1}^t (\delta_k - 1) > 0 \quad \forall t \leq T, \sum_{k=1}^T (\delta_k - 1) \in [\lceil T^{1/2} \rceil, 2 \lceil T^{1/2} \rceil]\right) \\ & \geq c(B) \mathbb{P}(M_t > 0 \quad \forall t \leq T, M_T \in [\lceil T^{1/2} \rceil, 2 \lceil T^{1/2} \rceil]) - \frac{T}{2n} \end{aligned} \quad (29)$$

for all large enough n , where $M_t = 1 + \sum_{i=1}^t (W_i - 1)$ and $(W_i)_{i=1}^T$ is a sequence of independent Poisson random variables with mean one. To finish the proof, we use the following generalised ballot theorem:

Theorem 2.1 (Addario-Berry and Reed [1]). Suppose X is a random variable satisfying $\mathbb{E}[X] = 0$, $\text{Var}(X) > 0$, $\mathbb{E}[X^{2+\alpha}] < \infty$ for some $\alpha > 0$, and X is a lattice random variable with period d (meaning that dX is an integer random variable and d is the smallest positive real number for which this holds). Then given independent random variables X_1, X_2, \dots distributed as X with associated partial sums $S_t = \sum_{i=1}^t X_i$, for all j such that $0 \leq j = O(\sqrt{n})$ and such that j is a multiple of $1/d$ we have

$$\mathbb{P}(S_t > 0 \quad \forall t \in [n], S_n = j) = \Theta\left(\frac{j+1}{n^{3/2}}\right). \quad (30)$$

Going back to (29) and using (30) we see that

$$\mathbb{P}(M_t > 0 \forall t \leq T, M_T \in [[T^{1/2}], 2[T^{1/2}]]) \geq c \sum_{j=[T^{1/2}] }^{2[T^{1/2}]} \frac{j}{T^{3/2}} \geq cT^{-1/2}. \quad (31)$$

Finally, observe that the term $T/2n$ on the right-hand side of (29) is $O((An^{1/2})^{-1})$, which is much smaller than $A^{1/2}/n^{1/3} \asymp T^{-1/2}$.

Therefore we conclude that there exists a constant $c_0 = c_0(B) > 0$ which depends solely on B such that, for all large enough A, n ,

$$\mathbb{P}\left(\sum_{k=1}^t (\delta_k - 1) > 0 \forall t \leq T\right) \geq c_0 T^{-1/2},$$

for each $L + 1 \leq i \leq 2L$. Combining this last estimate together with (25) and Lemma 2.3 we obtain the desired result.

2.3 Proof of Theorem 1.1: the $\mathbb{G}(n, d, p)$ model

We start by describing a method for exploring the components of the graph $\mathbb{G}(n, d, p)$, which we recall is the random graph obtained by performing bond percolation with parameter p on a realisation $\mathbb{G}(n, d)$ of a d -regular graph sampled uniformly at random from the set of all d -regular (simple) graphs on $[n]$. The exploration process we use here is taken verbatim from [13] (see also [20] and [27]).

In order to describe such procedure, we first recall the *configuration model*; this is an algorithm, which is due to Bollobás [6], that gives us a way of choosing a graph $\mathbb{G}(n, d)$ uniformly at random from the set of all d -regular graphs on n vertices, provided that dn is even.

The configuration model. Start with dn stubs, labelled (v, i) for $v \in [n]$ and $i \in [d]$. Choose a stub (V_0, I_0) in some way (the manner of choosing may be deterministic or random) and pair it uniformly at random with another stub (W_0, J_0) . Say that these two stubs are *matched* and put $\{V_0, W_0\} \in E$. Then at each subsequent step $k \in \{1, \dots, nd/2 - 1\}$, choose a stub (V_k, I_k) in some way from the set of unmatched stubs, and pair it uniformly at random with another unmatched stub (W_k, J_k) . Say that these two stubs are matched and put $\{V_k, W_k\} \in E$.

At the end of this process, the resulting object $G = ([n], E)$ is uniformly chosen amongst all d -regular *multigraphs* on $[n]$, i.e. it may have multiple edges or self-loops. However, with probability converging to $\exp((1 - d^2)/4)$ it is a simple graph, and conditioning on this event, it is uniformly chosen amongst all d -regular (simple) graphs on $[n]$.

The exploration process we employ will use the configuration model to generate components of $\mathbb{G}'(n, d, p)$, the p -percolated version of a uniformly random d -regular multigraph $\mathbb{G}'(n, d)$. When we talk about whether an edge of $\mathbb{G}'(n, d)$ is *retained*, we mean whether it is present in $\mathbb{G}'(n, d, p)$.

During our exploration process, each stub (or half-edge) of $\mathbb{G}'(n, d)$ is either *active*, *unseen* or *explored*, and its status changes during the course of the process. We write $\mathcal{A}_t, \mathcal{U}_t$ and \mathcal{E}_t for the sets of active, unseen and explored stubs at the end of the t -th step of the exploration process, respectively.

Given a stub h of $\mathbb{G}(n, d)$, we denote by $v(h)$ the vertex incident to h (in other words, if $h = (u, i)$ for some i then $v(h) = u$) and we write $\mathcal{S}(h)$ for the set of *all* stubs incident to $v(h)$ in $\mathbb{G}'(n, d)$ (that is, $\mathcal{S}(h) = \{(v(h), i) : i \in [d]\}$; note in particular that $h \in \mathcal{S}(h)$).

Algorithm 2. Let V_n be a vertex selected uniformly at random from $[n]$. At step $t = 0$ we declare *active* all stubs incident to V_n , while all the other $d(n - 1)$ stubs are declared *unseen*. Therefore we have that $|\mathcal{A}_0| = d$, $|\mathcal{U}_0| = d(n - 1)$ and $|\mathcal{E}_0| = 0$. For every $t \geq 1$, we proceed as follows.

- (a) If $|\mathcal{A}_{t-1}| \geq 1$, we choose (in an arbitrary way) one of the active stubs, say e_t , and we pair it with a stub h_t picked uniformly at random from $[dn] \setminus (\mathcal{E}_{t-1} \cup \{e_t\})$, i.e. from the set of all unexplored stubs after having removed e_t .
 - (a.1) If $h_t \in \mathcal{U}_{t-1}$ **and** the edge $e_t h_t$ is retained in the percolation (the latter event occurs with probability p , independently of everything else), then all the unseen stubs in $\mathcal{S}(h_t) \setminus \{h_t\}$ are declared active, while e_t and h_t are declared explored. Formally we update

- * $\mathcal{A}_t := (\mathcal{A}_{t-1} \setminus \{e_t\}) \cup (\mathcal{U}_{t-1} \cap \mathcal{S}(h_t) \setminus \{h_t\});$
 - * $\mathcal{U}_t := \mathcal{U}_{t-1} \setminus \mathcal{S}(h_t);$
 - * $\mathcal{E}_t := \mathcal{E}_{t-1} \cup \{e_t, h_t\}.$
- (a.2) If $h_t \in \mathcal{U}_{t-1}$ **but** the edge $e_t h_t$ is not retained in the percolation, then we simply declare e_t and h_t explored while the status of all other stubs remain unchanged. Formally we update
- * $\mathcal{A}_t := \mathcal{A}_{t-1} \setminus \{e_t\};$
 - * $\mathcal{U}_t := \mathcal{U}_{t-1} \setminus \{h_t\};$
 - * $\mathcal{E}_t := \mathcal{E}_{t-1} \cup \{e_t, h_t\}.$
- (a.3) If $h_t \in \mathcal{A}_{t-1}$, then we simply declare e_t and h_t explored while the status of all other stubs remain unchanged. Formally we update
- * $\mathcal{A}_t := \mathcal{A}_{t-1} \setminus \{e_t, h_t\};$
 - * $\mathcal{U}_t := \mathcal{U}_{t-1};$
 - * $\mathcal{E}_t := \mathcal{E}_{t-1} \cup \{e_t, h_t\}.$
- (b) If $|\mathcal{A}_{t-1}| = 0$ **and** $|\mathcal{U}_{t-1}| \geq 1$, we pick (in an arbitrary way) an unseen stub e_t , we declare active *all* the unseen stubs in $\mathcal{S}(e_t)$ (thus e_t at least is declared active), so that the number of active stubs is non-zero, and then we proceed as in step (a).
- (c) Finally, if $|\mathcal{A}_{t-1}| = 0$ **and** $|\mathcal{U}_{t-1}| = 0$, then all the stubs have been paired and we terminate the procedure.

Remark 5. Note that, since in the above procedure we explore *two half-edges* at each step, we have $\mathcal{A}_t \cup \mathcal{U}_t \neq \emptyset$ for $t \leq (dn/2) - 1$ (recall that dn is even) and $\mathcal{A}_{dn/2} \cup \mathcal{U}_{dn/2} = \emptyset$ (as $\mathcal{E}_{dn/2} = \{(u, i) : u \in [n], i \in [d]\}$, the set of all stubs). Thus the algorithm runs for $dn/2$ steps.

For $t \geq 1$ we define the event $R_t := \{e_t h_t \in \mathbb{G}(n, d, p)\}$ that the edge $e_t h_t$ revealed during the t -th step of the exploration process is retained in the percolation.

Observe that, if $|\mathcal{A}_{t-1}| \geq 1$, denoting by η_t the number of unseen half-edges that become active at step t , we can write

$$\eta_t = |\mathcal{A}_t| - |\mathcal{A}_{t-1}| = \mathbb{1}_{\{h_t \in \mathcal{U}_{t-1}\}} \mathbb{1}_{R_t} |\mathcal{S}(h_t) \cap \mathcal{U}_{t-1} \setminus \{h_t\}| - \mathbb{1}_{\{h_t \in \mathcal{A}_{t-1}\}} - 1. \quad (32)$$

In words, assuming $|\mathcal{A}_{t-1}| \geq 1$, the number of active stubs at the end of step t decreases by two if h_t is an active stub; it decreases by one if h_t is unseen and the edge $e_t h_t$ is not retained in the percolation, or if h_t is the unique unseen stub incident to $v(h_t)$ and the edge $e_t h_t$ is retained in the percolation; and it increases by $m - 2 \in \{0, 1, \dots, d - 2\}$ if $v(h_t)$ has $m \in \{2, \dots, d\}$ unseen stubs at the end of step $t - 1$ and the edge $e_t h_t$ is retained in the percolation.

Moreover, note that if $|\mathcal{A}_{t-1}| = 0$, then

$$\eta_t = |\mathcal{S}(e_t) \cap \mathcal{U}_{t-1} \setminus \{e_t\}| + \mathbb{1}_{\{h_t \in \mathcal{U}_{t-1} \setminus \mathcal{S}(e_t)\}} \mathbb{1}_{R_t} |\mathcal{S}(h_t) \cap \mathcal{U}_{t-1} \setminus \{h_t\}| - \mathbb{1}_{\{h_t \in \mathcal{S}(e_t)\}}. \quad (33)$$

Set $Y_t := |\mathcal{A}_t|$ for $t \in [dn/2] \cup \{0\}$. We define $t_0 := 0$ and recursively we set t_i to be the first time after t_{i-1} at which the number of active vertices hits zero; that is,

$$t_i := \min\{t \geq t_{i-1} + 1 : Y_t = 0\}, \quad (34)$$

for $i \geq 1$ (prior to the end of the procedure). The next result, which corresponds to Lemma 10 in [27], gives us a way to obtain an upper bound for the probability that $\mathcal{C}_{\max}(\mathbb{G}(n, d, p))$ contains less than $n^{2/3}/A$ vertices in terms of the (random) lengths of the positive excursions of the process Y_t . We refer the reader to [10] for an *intuitive* explanation.

Lemma 2.5 (Lemma 10 in [27]). *For $i \in \mathbb{N}$, denote by \mathcal{C}_i the i -th explored component in $\mathbb{G}(n, d, p)$. Then $t_i - t_{i-1} \leq (d - 1)|\mathcal{C}_i|$ for every i (until the end of the exploration process).*

We now proceed in the exact same way as we did for the $\mathbb{G}(n, p)$ model. Setting $T' := \lceil n^{2/3}/A \rceil$ and using the above lemma we bound

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, d, p))| < n^{2/3}/A) \leq \mathbb{P}(t_i - t_{i-1} < (d - 1)T' \forall i \in [N]),$$

where we denote by N the number of positive excursions of the process Y_t until the end of **Algorithm 2**. Define $T := (d-1)T$. If $t_i - t_{i-1} < T$ for all $i \in [N]$ then $(dn)/2 = \sum_{i=1}^N (t_i - t_{i-1}) < NT$ and hence $N > (dn)/(2T)$. Setting $L := \lfloor (dn)/(4T) \rfloor$, we see that $N > 2L$. Hence we can bound

$$\mathbb{P}(t_i - t_{i-1} < T \forall i \in [N]) \leq \mathbb{P}(t_i - t_{i-1} < T \forall L+1 \leq i \leq 2L).$$

As we did when analysing the $\mathbb{G}(n, d, p)$ model, we define the (good) event

$$\mathcal{G}_j := \{t_j \leq BLT^{1/2}\},$$

where $L \leq j \leq 2L$ and B is some large positive constant (independent of A, n, λ but possibly dependent on d). The events \mathcal{G}_j are needed to bound from above the number of explored half-edges while revealing the cluster of a vertex in $\mathbb{G}'(n, d, p)$; such an upper bound on the number of explored stubs is needed in order to derive a meaningful lower bound for the probability that the number of active half-edges stays above zero for the required number of steps.

We continue by writing

$$\mathbb{P}(t_i - t_{i-1} < T \forall L+1 \leq i \leq 2L) \leq \mathbb{P}(\{t_i - t_{i-1} < T \forall L+1 \leq i \leq 2L\} \cap \mathcal{G}_{2L}) + \mathbb{P}(\mathcal{G}_{2L}^c).$$

The next lemma controls the probability of the (bad) event \mathcal{G}_{2L}^c .

Lemma 2.6. *There exist positive constants $B_0 = B_0(d), A_0 = A_0(\lambda, d) > 0$ such that, for any $A \geq A_0, B \geq B_0$ and for all large enough n , we have*

$$\mathbb{P}(\mathcal{G}_{2L}^c) \leq \exp(-cA^{3/2}),$$

for some positive constant $c = c(B) > 0$.

Therefore in what follows we can focus on the probability

$$\mathbb{P}(\{t_i - t_{i-1} < T \forall L+1 \leq i \leq 2L\} \cap \mathcal{G}_{2L}) \quad (35)$$

Let us define $\mathcal{F}_j := \mathcal{F}_{t_j}$ for $j \in [N]$, the σ -algebra consisting of all the information collected by the exploration process by time t_j , i.e. by the time at which we start the exploration of the j -th component. With the next result we show that, on the event \mathcal{G}_{i-1} , we have $t_i - t_{i-1} \geq T$ with probability $\gtrsim T^{-1/2}$.

Lemma 2.7. *There exist positive constants $B_0 = B_0(d), A_0 = A_0(\lambda, d) > 0$ such that, for any $A \geq A_0, B \geq B_0$ and for all large enough n , on the event \mathcal{G}_{i-1} we have*

$$\mathbb{P}(t_i - t_{i-1} \geq T | \mathcal{F}_{i-1}) \geq \frac{c_0}{T^{1/2}}$$

for each $L+1 \leq i \leq 2L$, for some constant $c_0 = c_0(B, d) > 0$.

It is now very easy to show that the expression in (35) is at most $\exp(-cA^{3/2})$ (for some $c > 0$). Indeed, proceeding as we did for the $\mathbb{G}(n, p)$ model, we can write

$$\begin{aligned} & \mathbb{P}(\{t_i - t_{i-1} < T \forall L+1 \leq i \leq 2L\} \cap \mathcal{G}_{2L}) \\ & \leq \mathbb{E} \left[\mathbb{1}_{\{t_i - t_{i-1} < T \forall L+1 \leq i \leq 2L-1\} \cap \mathcal{G}_{2L-1}} (1 - \mathbb{P}(t_{2L} - t_{2L-1} \geq T | \mathcal{F}_{2L-1})) \right]. \end{aligned} \quad (36)$$

Thanks to Lemma 2.7 we know that the expression on the right-hand side of (36) is at most

$$(1 - c_0 T^{-1/2}) \mathbb{P}(\{t_i - t_{i-1} < T \forall L+1 \leq i \leq 2L-1\} \cap \mathcal{G}_{2L-1}).$$

Iterating and using the classical inequality $1 + x \leq e^x$ (which is valid for all $x \in \mathbb{R}$), we conclude that

$$\mathbb{P}(\{t_i - t_{i-1} < T \forall L+1 \leq i \leq 2L\} \cap \mathcal{G}_{2L}) \leq (1 - c_0 T^{-1/2})^L \leq \exp(-(c_0 L) T^{-1/2}). \quad (37)$$

Recalling the definition of L and T , it is easy to see that $LT^{-1/2} \geq cA^{3/2}$ for all large enough n , for some constant $c = c(d) > 0$; hence, putting all pieces together, we conclude that there is a constant $c > 0$ such that, for all large enough n ,

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}'(n, d, p))| < n^{2/3}/A) \leq \exp(-cA^{3/2}). \quad (38)$$

Finally, writing \mathbb{S}_n for the event that the multigraph $\mathbb{G}'(n, d)$ underlying $\mathbb{G}'(n, d, p)$ is simple, we have

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}'(n, d, p))| < n^{2/3}/A | \mathbb{S}_n) \leq \frac{\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}'(n, d, p))| < n^{2/3}/A)}{\mathbb{P}(\mathbb{S}_n)}.$$

Since $\mathbb{P}(\mathbb{S}_n) \sim e^{(1-d^2)/4}$ as $n \rightarrow \infty$, using (38) we arrive at

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, d, p))| < n^{2/3}/A) = \mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}'(n, d, p))| < n^{2/3}/A | \mathbb{S}_n) \leq \exp(-cA^{3/2})$$

for some (new) constant $c = c(d) > 0$. This concludes the proof of Theorem 1.1. What is left to do is giving proofs of Lemmas 2.6 and 2.7; this is done in the following section.

2.3.1 Proofs of Lemmas 2.6 and 2.7

In this section we establish the two auxiliary facts which were stated, without proof, while proving the main theorem for the $\mathbb{G}(n, d, p)$ model.

Proof of Lemma 2.6. Recall that $\mathcal{G}_{2L} := \{t_{2L} \leq BLT^{1/2}\}$. To bound from above the probability of the complementary event, we employ the same strategy used for the $\mathbb{G}(n, p)$ model. Let $S_0 := d$ and define recursively $S_i := S_{i-1} + \hat{\eta}_i - 1$, where the random variable $\hat{\eta}_i$ is defined (for $i \leq (dn)/2$) by

$$\hat{\eta}_i := |\mathcal{S}(e_i) \cap \mathcal{U}_{i-1} \setminus \{e_i\}| + \mathbb{1}_{\{h_i \in \mathcal{U}_{i-1} \setminus \mathcal{S}(e_i)\}} \mathbb{1}_{R_i} |\mathcal{S}(h_i) \cap \mathcal{U}_{i-1} \setminus \{h_i\}| - \mathbb{1}_{\{h_i \in \mathcal{A}_{i-1} \cup \mathcal{S}(e_i)\}} - 1. \quad (39)$$

Recalling (32) and (33) we immediately see that $\hat{\eta}_i = \eta_i$ when either $1 \leq i \leq t_1$ or $i \in (t_{j-1} + 1, t_j]$ for some $1 < j \leq N$, whereas $\hat{\eta}_{t_{j-1}+1} = \eta_{t_{j-1}+1} - 1$ for $1 < j \leq N$. Indeed, if $1 \leq i \leq t_1$, then e_i is taken from the set of active half-edge, so that $\mathcal{S}(e_i) \cap \mathcal{U}_{i-1} \setminus \{e_i\} = \mathcal{S}(e_i) \cap \mathcal{U}_{i-1} = \emptyset$ (whence $|\mathcal{S}(e_i) \cap \mathcal{U}_{i-1} \setminus \{e_i\}| = 0$) and also $\mathcal{A}_{i-1} \cup \mathcal{S}(e_i) = \mathcal{A}_{i-1}, \mathcal{U}_{i-1} \setminus \mathcal{S}(e_i) = \mathcal{U}_{i-1}$; thus $\hat{\eta}_i = \eta_i$. The other cases are treated similarly.

Then $(-\min_{j \in [t]} S_j) + 1$ denotes the number of disjoint clusters that have been fully explored by time $t \in [n]$. When $Y_t = 0$, we have fully explored a cluster, and this occurs precisely when $S_t = \min_{j \in [t-1]} S_j - 1$, i.e., the process $(S_t)_{t \in [n]}$ reaches a new all-time minimum.

Proceeding in the exact same way as we did in the proof of Lemma 2.1, we arrive at

$$\mathbb{P}(\mathcal{G}_{2L}^c) \leq \mathbb{P}\left(\max_{j \leq \lfloor BLT^{1/2} \rfloor} (-S_j) < 2L\right). \quad (40)$$

To continue, we separate $\hat{\eta}_t$ into its constituent by introducing

$$X_{i,1} := \mathbb{1}_{\{h_i \in \mathcal{U}_{i-1} \setminus \mathcal{S}(e_i)\}} \mathbb{1}_{R_i} |\mathcal{S}(h_i) \cap \mathcal{U}_{i-1} \setminus \{h_i\}| - 1$$

and

$$X_{i,2} := |\mathcal{S}(e_i) \cap \mathcal{U}_{i-1} \setminus \{e_i\}| - \mathbb{1}_{\{h_i \in \mathcal{A}_{i-1} \cup \mathcal{S}(e_i)\}}.$$

Then clearly $\hat{\eta}_i = X_{i,1} + X_{i,2}$ and hence we can write

$$S_j = d + \sum_{i=1}^j \hat{\eta}_i = d + \sum_{i=1}^j X_{i,1} + \sum_{i=1}^j X_{i,2}.$$

Observe that, on the event within the probability on the right-hand side of (40), we have

$$\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} |\mathcal{S}(e_i) \cap \mathcal{U}_{i-1} \setminus \{e_i\}| \leq d \sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \mathbb{1}_{\{|\mathcal{A}_{i-1}|=0\}} = d \max_{j \in \lfloor BLT^{1/2} \rfloor} (-S_j) < 2dL,$$

where the first inequality follows from the observation that $|\mathcal{S}(e_i) \cap \mathcal{U}_{i-1} \setminus \{e_i\}| > 0$ if, and only if, $\mathcal{A}_{i-1} = \emptyset$ (otherwise e_i would be selected from the set of active stubs and hence we would have $\mathcal{S}(e_i) \cap \mathcal{U}_{i-1} = \emptyset$), in which case it is at most d .

Therefore, the probability on the right-hand side of (40) equals

$$\mathbb{P}\left(d + \sum_{i=1}^j X_{i,1} > -2L - \sum_{i=1}^j X_{i,2} \quad \forall j \leq \lfloor BLT^{1/2} \rfloor, \sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} |\mathcal{S}(e_i) \cap \mathcal{U}_{i-1} \setminus \{e_i\}| < 2dL\right). \quad (41)$$

Since $X_{i,2} \leq |\mathcal{S}(e_i) \cap \mathcal{U}_{i-1} \setminus \{e_i\}| \in \mathbb{N}_0$, on the (second) event in (41) we have

$$\sum_{i=1}^j X_{i,2} \leq \sum_{i=1}^j |\mathcal{S}(e_i) \cap \mathcal{U}_{i-1} \setminus \{e_i\}| \leq \sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} |\mathcal{S}(e_i) \cap \mathcal{U}_{i-1} \setminus \{e_i\}| < 2dL \text{ for each } j \leq \lfloor BLT^{1/2} \rfloor.$$

Whence the probability in (41) is at most

$$\mathbb{P}\left(d + \sum_{i=1}^j X_{i,1} > -4dL \forall j \leq \lfloor BLT^{1/2} \rfloor\right) \leq \mathbb{P}\left(d + \sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} X_{i,1} > -4dL\right).$$

In order to bound the probability on the right-hand side of the last inequality, we follow [13].

Denote by $\mathcal{V}_k^{(d)}$ the set of vertices having d unseen stubs at step k of the exploration process; we call these vertices *fresh*. We also introduce the event $F_k := \{v(h_k) \in \mathcal{V}_{k-1}^{(d)}\}$ that $v(h_k)$ is fresh. Then, defining

$$\eta'_i := \mathbb{1}_{R_i}(d-2) + \mathbb{1}_{R_i} \mathbb{1}_{F_i} - 1,$$

we have $X_{i,1} \leq \eta'_i$. Indeed, the inequality is actually an identity whenever $v(h_i)$ has $m \in \{d-1, d\}$ unseen stubs attached to it and R_i occurs, whereas $X_{i,1} < \eta'_i$ in all the other cases.

Then we can bound

$$\mathbb{P}\left(d + \sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} X_{i,1} > -4dL\right) \leq \mathbb{P}\left(d + \sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \eta'_i > -4dL\right). \quad (42)$$

In order to bound from above the last probability we substitute (as in [13]) the dependent indicators $\mathbb{1}_{F_i}$ that appear in the definition of η'_i with independent $\{0, 1\}$ -valued random variables. To this end notice that, conditional on everything that occurred up to the end of step $i-1$ in the exploration process, vertex $v(h_i)$ is fresh with probability

$$\frac{d|\mathcal{V}_{i-1}^{(d)}|}{dn - 2(i-1) - 1}.$$

Thus, in order to substitute the $\mathbb{1}_{F_i}$ with (larger) independent indicator random variables, we need an upper bound for the number of fresh vertices that we expect to observe at each step $i \leq \lfloor BLT^{1/2} \rfloor$ of the exploration process.

The next result, which is basically the same as Lemma 3.1 in [13], states that it is very unlikely to have more than $n - 1 - i + i^2/2n$ fresh vertices at the i -th step of the exploration process.

Lemma 2.8. *Let $a_n(i) := n - 1 - i + i^2/2n$. Then, for every $m \geq 1$ and all large enough n , we have that*

$$\mathbb{P}\left(\exists i \leq \lfloor BLT^{1/2} \rfloor : |\mathcal{V}_i^{(d)}| > a_n(i) + m\right) \leq BLT^{1/2} \exp\left(-c \frac{mn^{1/2}}{LT^{1/2}}\right)$$

where $c = c(d)$ is some finite constant that depends only on d .

Arguing as in [13] we observe that, if $|\mathcal{V}_i^{(d)}| \leq a_n(i) + m$ for all $i \leq \lfloor BLT^{1/2} \rfloor$, then we can write

$$\eta'_i = \mathbb{1}_{R_i}(d-2) + \mathbb{1}_{R_i} \mathbb{1}_{F_i} \mathbb{1}_{\{|\mathcal{V}_{i-1}^{(d)}| \leq a_n(i-1) + m\}} - 1 =: \eta''_i, \text{ for } i \leq \lfloor BLT^{1/2} \rfloor. \quad (43)$$

Then we can bound from above the probability on the right-hand side of (42) by

$$\begin{aligned} \mathbb{P}\left(d + \sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \eta'_i > -4dL, |\mathcal{V}_i^{(d)}| \leq a_n(i) + m \forall i \leq \lfloor BLT^{1/2} \rfloor\right) &+ BLT^{1/2} \exp\left(-c \frac{mn^{1/2}}{LT^{1/2}}\right) \\ &\leq \mathbb{P}\left(d + \sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \eta''_i > -4dL\right) + BLT^{1/2} \exp\left(-c \frac{mn^{1/2}}{LT^{1/2}}\right). \end{aligned}$$

Conditional on everything that has occurred up to the end of step $i - 1$ in the exploration process, the random variable $\mathbb{1}_{R_i} \mathbb{1}_{F_i} \mathbb{1}_{\{|\mathcal{V}_{i-1}^{(d)}| \leq a_n(i-1)+m\}}$ which appears in (43) equals 1 with probability

$$p \mathbb{1}_{\{|\mathcal{V}_{i-1}^{(d)}| \leq a_n(i-1)+m\}} \frac{d \left| \mathcal{V}_{i-1}^{(d)} \right|}{dn - 2(i-1) - 1} \leq p \frac{d(a_n(i-1) + m)}{dn - 2(i-1) - 1}.$$

Therefore, arguing precisely as in [13], if $(U_i)_{i \geq 1}$ is an iid sequence of $U([0, 1])$ random variables (also independent from all other random quantities involved), then

$$\mu_i := \mathbb{1}_{R_i}(d - 2) + \mathbb{1}_{R_i} \mathbb{1}_{\{U_i \leq \frac{d(a_n(i-1)+m)}{dn-2(i-1)-1}\}} - 1 \quad (44)$$

defines a sequence of *independent* random variables that should be larger than the η_i'' . The next result can be proved in the exact same way as Proposition 3.3 in [13].

Lemma 2.9. *Let $(U_i)_i$ be a sequence of iid random variables, also independent from all other random variables involved, with $U_1 =_d U([0, 1])$. For each $i \leq \lfloor BLT^{1/2} \rfloor$ let μ_i be as in (44) above. Then*

$$\mathbb{P}\left(d + \sum_{i=1}^t \eta_i'' > -4dL\right) \leq \mathbb{P}\left(d + \sum_{i=1}^t \mu_i > -4dL\right).$$

In order to provide an upper bound for the above quantity we would like to turn the independent (but not identically distributed) μ_i into iid random variables ξ_i . To this end, keeping in mind the definition of μ_i given in (44), following [13] we define

$$\mu'_i := \mathbb{1}_{R_i} \mathbb{1}_{\{U_i > \frac{d(a_n(i-1)+m)}{dn-2(i-1)-1}\}} \quad (45)$$

and set, for $i \leq \lfloor BLT^{1/2} \rfloor$,

$$\xi_i := \mu_i + \mu'_i = \mathbb{1}_{R_i}(d - 2) + \mathbb{1}_{R_i} - 1 = \mathbb{1}_{R_i}(d - 1) - 1. \quad (46)$$

By adding the (random) sums $\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \mu'_i$ to the $\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \mu_i$ we can rewrite the probability which appears in Lemma 2.9 as

$$\mathbb{P}\left(d + \sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \xi_i > \sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \mu'_i - 4dL\right). \quad (47)$$

In order to control the (random) sum $\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \mu'_i$ we use Lemma 3.4 in [13]

Lemma 2.10. *Define*

$$q(t) = q_{n,d}(t) := p \left(1 - \frac{2}{d}\right) \frac{t(t-1)}{2n}, \quad t \leq \lfloor BLT^{1/2} \rfloor, t \in \mathbb{N}. \quad (48)$$

Then, for all large enough n , if $m = O(n^{1/2})$ we have

$$\mathbb{P}\left(\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \mu'_i \leq q(\lfloor BLT^{1/2} \rfloor) - h\right) \leq C e^{-\frac{hn^{1/2}}{\lfloor BLT^{1/2} \rfloor}}, \quad (49)$$

where $C = C(B, d)$ is a constant that depends on d, B .

(We remark that the factor T which multiplies the exponential term in Lemma 3.4 of [13] should not be there.) Thanks to the last lemma applied with $h := L$ and recalling the definition of ξ_i given in (46) above, if n is large enough we can easily upper bound the probability in (47) by

$$\begin{aligned} \mathbb{P}\left(\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \xi_i > q(\lfloor BLT^{1/2} \rfloor) - L - 4dL - d\right) &+ \mathbb{P}\left(\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \mu'_i \leq q(\lfloor BLT^{1/2} \rfloor) - L\right) \\ &\leq \mathbb{P}\left(\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \mathbb{1}_{R_i} > \lfloor BLT^{1/2} \rfloor p + y\right) + e^{-c \frac{n^{1/2}}{T^{1/2}}}, \end{aligned}$$

where we set

$$y := \left(q(\lfloor BLT^{1/2} \rfloor) - 5dL \right) (d-1)^{-1} + \lfloor BLT^{1/2} \rfloor ((d-1)^{-1} - p).$$

Recalling that $L \sim An^{1/3}$, $T \sim n^{2/3}/A$ and $p = (1 + \lambda n^{-1/3})/(d-1)$ we obtain (for all large enough n)

$$q(\lfloor BLT^{1/2} \rfloor) - 5dL \geq L \left(\frac{B^2 LT(d-2)}{4nd(d-1)} - 5d \right) \geq L \left(\frac{B^2(d-2)}{5d(d-1)} - 5d \right).$$

Moreover,

$$\lfloor BLT^{1/2} \rfloor ((d-1)^{-1} - p) \geq -|\lambda| BLT^{1/2} (d-1)^{-1} n^{-1/3} \geq -|\lambda| BL (d-1)^{-1} A^{-1/2}$$

so that (if n is large enough)

$$y \geq (d-1)^{-1} L \left(\frac{B^2(d-2)}{5d(d-1)} - 5d - |\lambda| BA^{-1/2} \right) \geq c'L$$

for some constant $c' = c'(B, d) > 0$, provided $B = B(d)$ is sufficiently large and $A \geq A_0$ for some large enough $A_0 = A_0(\lambda, d) > 0$. Hence, since $\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \mathbb{1}_{R_i}$ has the $\text{Bin}(\lfloor BLT^{1/2} \rfloor, p)$ law, using Lemma 1.1 we arrive at

$$\begin{aligned} \mathbb{P} \left(\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \mathbb{1}_{R_i} > \lfloor BLT^{1/2} \rfloor p + y \right) &\leq \mathbb{P} \left(\sum_{i=1}^{\lfloor BLT^{1/2} \rfloor} \mathbb{1}_{R_i} > \lfloor BLT^{1/2} \rfloor p + c'L \right) \\ &\leq \exp \left(-c \frac{L}{T^{1/2}} \right) \leq \exp(-cA^{3/2}), \end{aligned}$$

for some constant $c = c(d) > 0$. Putting all pieces together, we have shown that, if $m = O(n^{1/2})$, then for all large enough A, n we have

$$\mathbb{P}(\mathcal{G}_{2L}^c) \leq \exp(-cA^{3/2}) + \exp\left(-c \frac{n^{1/2}}{T^{1/2}}\right) + BLT^{1/2} \exp\left(-c \frac{mn^{1/2}}{LT^{1/2}}\right).$$

Since, by assumption, $A = O(n^{1/6})$, taking $m = L$ (which is indeed $O(n^{1/2})$) we see that the second and third terms on the right-hand side of the last expression are both at most $\exp(-\Theta(A^{3/2}))$, thus completing the proof of the lemma.

Proof of Lemma 2.7. Here the goal is to show that there is a constant $c_0 = c_0(B, d) > 0$ such that, for each $L + 1 \leq i \leq 2L$, on the event \mathcal{G}_{i-1} it holds that

$$\mathbb{P}(t_i - t_{i-1} \geq T | \mathcal{F}_{i-1}) \geq \frac{c_0}{T^{1/2}}$$

for all large enough n , where we recall that $\mathcal{F}_{i-1} = \mathcal{F}_{t_{i-1}}$ is the σ -algebra consisting of all the information gathered by the exploration process until time t_{i-1} (i.e. until the moment at which we start revealing the $(i-1)$ -th cluster). Then, setting $Y_t^i := Y_{t_i+t}$ and $\eta_t^i := \eta_{t_i+t}$ to simplify notation, we can write

$$\mathbb{P}(t_i - t_{i-1} \geq T | \mathcal{F}_{i-1}) = \mathbb{P}(Y_t^i > 0 \forall t \leq T-1 | \mathcal{F}_{i-1}) \geq \mathbb{P}\left(d_i + \sum_{k=1}^t \eta_k^i > 0 \forall t \leq T | \mathcal{F}_{i-1}\right), \quad (50)$$

where $d_i \geq 1$ represents the number of unseen stubs attached to the node from which we start exploring (at time $t_{i-1} + 1$) the i -th cluster. Recall that, if $Y_{k-1}^i \geq 1$, then

$$\eta_k^i = \mathbb{1}_{\{h_k^i \in \mathcal{U}_{k-1}^i\}} \mathbb{1}_{R_k^i} | \mathcal{S}(h_k^i) \cap \mathcal{U}_{k-1}^i \setminus \{h_k^i\} | - \mathbb{1}_{\{h_k^i \in \mathcal{A}_{k-1}^i\}} - 1, \quad (51)$$

where denotes the half-edge selected at time $t_{i-1} + k$ whereas $\mathcal{U}_{k-1}^i, \mathcal{A}_{k-1}^i$ are the set of unseen stubs and set of active half-edges, respectively. We further denote by $\mathcal{V}_{k,i}^{(d)}$ the set of (fresh) vertices with d unseen stubs at time $t_{i-1} + k$ and $F_k^i := \{v(h_k^i) \in \mathcal{V}_{k-1,i}^{(d)}\}$, the event that $v(h_k^i)$ is fresh.

Since we want to bound the probability in (50) from below, we need to approximate the η_k^i with *smaller* random variables, sufficiently close to the former but easier to be dealt with. To this end, following [13], we define

$$\delta_k^i := \mathbb{1}_{R_k^i} \mathbb{1}_{F_k^i} (d-1) - \mathbb{1}_{\{h_k^i \in \mathcal{A}_{k-1}^i\}} - 1 \quad (52)$$

and we observe that, on the event where $d_i + \sum_{k=1}^t \eta_k^i > 0$ for all $t \leq T$, we have

$$\eta_k^i \geq \delta_k^i. \quad (53)$$

Indeed, if at step $t_{i-1} + k$ we pick h_k^i from the set of active vertices, then $\eta_k^i = \delta_k^i$; similarly, if h_k^i is incident to a fresh vertex (whence in particular h_k^i is unseen), then again $\eta_k^i = \delta_k^i$. In all other cases (i.e. when h_k^i is unseen but the vertex to which is attached has less than d unseen half-edges incident to it), then $\eta_k^i \geq \delta_k^i$.

The reason why the δ_k^i are a good approximation of the η_k^i is because, during the first phase of the exploration process, most vertices are fresh (and hence for most steps $i \leq T$ we expect that $\eta_k^i = \delta_k^i$).

Then, using (53), we bound from below the probability in (50) by

$$\mathbb{P}\left(d_i + \sum_{k=1}^t \delta_k^i > 0 \forall t \leq T \mid \mathcal{F}_{i-1}\right). \quad (54)$$

To establish the required lower bound for this probability, we proceed exactly as in Section 4.2 of [13]. Actually, in our case the proof is even simpler, because we are *not* conditioning on the event \mathbb{S}_n that the multigraph $\mathbb{G}'(n, d)$ underlying $\mathbb{G}'(n, d, p)$ is simple (removing the conditioning on \mathbb{S}_n was the content of Section 4.2.1 in [13]). We will not give all the details here, but nevertheless, for the sake of clarity, we will carefully explain the required adaptations needed from [13] to the current setting.

The idea is to replace the δ_k^i with the simpler

$$\chi_k^i := \mathbb{1}_{R_k^i} \mathbb{1}_{F_k^i} (d-1) - 1 \quad (55)$$

and then to show that, for any $\gamma \in (0, 1/2)$, we can find a constant c_0 (which depends on B, d) such that, for all sufficiently large A, n , on \mathcal{G}_{i-1} we can bound

$$\mathbb{P}\left(d_i + \sum_{k=1}^t \chi_k^i \geq t^\gamma \forall t \leq T \mid \mathcal{F}_{i-1}\right) \geq c_0' T^{-1/2} \quad (56)$$

for some constant $c_0' = c_0'(B, d) > 0$. Subsequently we show that for $\gamma \in (0, 1/2)$ (and for all large enough A, n), on \mathcal{G}_{i-1} we have

$$\mathbb{P}\left(\exists t \leq (t_i - t_{i-1}) \wedge T : \sum_{k=1}^t \mathbb{1}_{\{h_k^i \in \mathcal{A}_{k-1}^i\}} \geq t^\gamma \mid \mathcal{F}_{i-1}\right) \ll T^{-1/2}. \quad (57)$$

Combining (56) and (57) we obtain the desired lower bound on (54); this is the content of Lemma 2.11 below. (As in [13], we note that the choice of γ is not important above; one may choose, for example, $\gamma = 1/4$ in both (56) and (57). We retain the general γ in the proofs since this is no extra work.)

Lemma 2.11. *Let χ_k^i be as in (55). Suppose that (56) and (57) hold. Then, there exists a constant $c_0 = c_0(B, d) > 0$ such that, for all sufficiently large A, n , on \mathcal{G}_{i-1} we have*

$$\mathbb{P}\left(d_i + \sum_{k=1}^t \eta_k^i > 0 \forall t \leq T \mid \mathcal{F}_{i-1}\right) \geq c_0 T^{-1/2},$$

for each $L+1 \leq i \leq 2L$.

Proof. We follow the argument given in Section 4.2.6 in [13]. Note that

$$\mathbb{P}\left(d_i + \sum_{k=1}^t \delta_k^i > 0 \forall t \leq T \mid \mathcal{F}_{i-1}\right) = \mathbb{P}\left(d_i + \sum_{k=1}^t \delta_k^i > 0 \forall t \leq (t_i - t_{i-1}) \wedge T \mid \mathcal{F}_{i-1}\right) \quad (58)$$

because, if $d_i + \sum_{k=1}^t \delta_k^i > 0$ for all $t \leq T$, then (thanks to (53)) also $d_i + \sum_{k=1}^t \eta_k^i > 0$ for all $t \leq T$ and hence we must have that $t_i - t_{i-1} > T$ (so that $(t_i - t_{i-1}) \wedge T = T$).

Now the probability on the right-hand side of (58) is clearly at least

$$\mathbb{P}\left(d_i + \sum_{k=1}^t \chi_k^i > t^\gamma \forall t \leq (t_i - t_{i-1}) \wedge T, \sum_{k=1}^t \mathbb{1}_{\{h_k^i \in \mathcal{A}_{k-1}^i\}} < t^\gamma \forall t \leq (t_i - t_{i-1}) \wedge T \mid \mathcal{F}_{i-1}\right),$$

and the last probability is at least the difference between (56) and (57), which is at least (for all large enough n) $c_0 T^{-1/2}$ with $c_0 := c_0'/2$, as desired. \square

There remains to establish (56) and (57). To this end, we continue with the following Lemma, whose proof follows the exact same steps as Proposition 4.11 in [13].

Lemma 2.12. *Define $\Delta_k^i := \mathbb{1}_{R_k^i} \mathbb{1}_{\{U_k \leq 1 - (4BLT^{1/2})/n\}}$ for $k \leq T$, where $(U_k)_{k \in \mathbb{N}}$ is a sequence of iid $\text{Unif}([0, 1])$ random variables, independent of everything else. Then*

$$\mathbb{P}\left(\sum_{k=1}^t \chi_k^i \geq t^\gamma \quad \forall t \leq T \mid \mathcal{F}_{i-1}\right) \geq \mathbb{P}\left(\sum_{i=1}^t \Delta_k^i \geq t^\gamma \quad \forall t \leq T\right).$$

The intuition here is that, since $|\mathcal{V}_{k,i}^{(d)}| \geq n - 1 - 2(t_{i-1} + k)$ for each $k \geq 1$ (as at most two vertices can be removed from the set of fresh nodes at each step), the (conditional) probability (on the event \mathcal{G}_{i-1}) that h_k^i is fresh when $k \leq T$ equals

$$\frac{d|\mathcal{V}_{i,k-1}^{(d)}|}{dn - 2(t_{i-1} - k - 1) - 1} \geq \frac{d(n - 2t_{i-1} - 2k)}{dn} = 1 - \frac{2(t_{i-1} + k)}{n} \geq 1 - \frac{4BLT^{1/2}}{n},$$

provided that n is large enough. Thus (with $\mathcal{F}_{i-1}(k-1) := \mathcal{F}_{t_{i-1}+k-1}$) for all large enough n we obtain

$$\mathbb{P}(F_k^i \mid \mathcal{F}_{i-1}(k-1)) \geq 1 - \frac{4BLT^{1/2}}{n} = \mathbb{P}(U_k \leq 1 - (4BLT^{1/2})/n).$$

which then (recalling the definitions of χ_k^i and Δ_k^i) shows that χ_k^i stochastically dominates Δ_k^i for each k .

We do not give the actual proof of Lemma 2.12 because, as we said earlier, it can be established in the exact same way as the proof of Proposition 4.11 in [13]. (We remark that the $1 - (4BLT^{1/2})/n$ in the definition of Δ_k^i given here translates into $1 - T'/n$ in Proposition 4.11 of [13].)

The next step is the following lemma, which is almost the same as Lemma 4.13 in [13].

Lemma 2.13. *Let Δ_k^i be as in Lemma 2.12. Let $(D_k)_{k \leq T}$ be a sequence of iid random variables with $\mathbb{P}(D_k = d - 2) = 1/(d - 1) = 1 - \mathbb{P}(D_k = -1)$. There exists constants $A_0 = A_0(\lambda, d) > 0$ and $c = c(d) > 0$ such that, for any $A \geq A_0, \varepsilon \in (0, 1)$ and for all large enough n , we have*

$$\mathbb{P}\left(\sum_{k=1}^t \Delta_k^i \geq t^\gamma \quad \forall t \leq T\right) \geq c \mathbb{P}\left(\sum_{k=1}^t D_k \geq t^\gamma \quad \forall t \leq T, \sum_{k=1}^T D_k \in [\varepsilon T^{1/2}, T^{1/2}/\varepsilon]\right).$$

Proof. This is basically Lemma 4.13 in [13] (whence we won't provide all the details), the proof of which is based on an exponential change of measure. We set

$$\frac{d\hat{\mathbb{P}}}{d\mathbb{P}} \Big|_{\mathcal{F}_t} := \frac{\exp\left(\nu \sum_{k=1}^t \Delta_k^i\right)}{\mathbb{E}\left[\exp\left(\nu \sum_{k=1}^t \Delta_k^i\right)\right]}, \quad t \in \mathbb{N}_0$$

where ν chosen in such a way to obtain the desired distribution for the Δ_k^i . In particular, in our case we set

$$\nu := \frac{1}{d-1} \log\left(1 - p\left(1 - \frac{4BLT^{1/2}}{n}\right)\right) - \frac{1}{d-1} \log\left((d-2)p\left(1 - \frac{4BLT^{1/2}}{n}\right)\right). \quad (59)$$

(The term $4BLT^{1/2}$ here corresponds to the term T' in Lemma 4.13 of [13].) With this choice of ν , the sequence $(\Delta_k^i)_{k \in [T]}$ is iid with

$$\hat{\mathbb{P}}(\Delta_k^i = d - 2) = \frac{1}{d-1} = 1 - \hat{\mathbb{P}}(\Delta_k^i = -1), \quad (60)$$

which is the required distribution. Proceeding in the exact same way as in the proof of Lemma 4.13 in [13], we arrive at

$$\mathbb{P}\left(\sum_{k=1}^t \Delta_k^i \geq t^\gamma \quad \forall t \leq T\right) \geq e^{-|\nu|T^{1/2}\varepsilon^{-1}} \mathbb{E}[e^{\nu \Delta_1^i}]^T \hat{\mathbb{P}}\left(\sum_{k=1}^t \Delta_k^i \geq t^\gamma \quad \forall t \leq T, \sum_{k=1}^T \Delta_k^i \in [\varepsilon T^{1/2}, T^{1/2}/\varepsilon]\right).$$

Thanks to (60), we only need to check that

$$e^{-|\nu|T^{1/2}\varepsilon^{-1}} \mathbb{E}[e^{\nu \Delta_1^i}]^T \geq c \quad (61)$$

for some constant $c = c(d) > 0$. To this end, we start by noticing that (by definition of ν)

$$\mathbb{E}[e^{\nu\Delta_1^i}] = \exp\left(O\left(\frac{\lambda^2}{n^{2/3}} + \frac{\lambda BLT^{1/2}}{n^{4/3}} + \frac{(BLT^{1/2})^2}{n^2}\right)\right);$$

whence, recalling that $T \sim n^{2/3}/A$ and $L \sim An^{1/3}$, we see that

$$T\left(\frac{\lambda^2}{n^{2/3}} + \frac{\lambda BLT^{1/2}}{n^{4/3}} + \frac{(BLT^{1/2})^2}{n^2}\right) \asymp \frac{\lambda^2}{A} + \frac{\lambda}{A^{1/2}} + 1.$$

Therefore $\mathbb{E}[e^{\nu\Delta_1^i}]^T \geq c$. Moreover,

$$|\nu|T^{1/2} \leq C\left(1 + \frac{|\lambda|}{A^{1/2}} + \frac{A^{1/2}}{n^{1/3}}\right)$$

and so (since $A \ll n^{2/3}$) we see that $e^{-|\nu|T^{1/2}\varepsilon^{-1}} \geq c > 0$ too, as required. \square

Finally, using Lemma 4.14 in [13] (with T in place of T'), we obtain

$$\mathbb{P}\left(\sum_{k=1}^t D_k \geq t^\gamma \forall t \leq T, \sum_{k=1}^T D_k \in [\varepsilon T^{1/2}, T^{1/2}/\varepsilon]\right) \geq cT^{-1/2}. \quad (62)$$

Combining Lemmas 2.12 and 2.13 together with (62) we conclude that indeed (56) holds. There remains to establish (57); this is the content of the next lemma, which is established following the steps carried out in Section 4.2.5 of [13].

Lemma 2.14. *On the event \mathcal{G}_{i-1} , for all large enough n we have*

$$\mathbb{P}\left(\exists t \leq (t_i - t_{i-1}) \wedge T : \sum_{i=1}^t \mathbb{1}_{\{h_k^i \in \mathcal{A}_{k-1}^i\}} \geq t^\gamma \mid \mathcal{F}_{i-1}\right) \ll T^{-1/2}. \quad (63)$$

Proof. As we said prior to the statement of the lemma, the proof follows the same steps carried out in Section 4.2.5 of [13]. The strategy there was to use a union bound to show that, for small values of t (i.e. during the first phase of the exploration process), it is very unlikely that h_k^i is selected from the set of active stubs. On the other hand, for large values of t , say $t \geq b$ with $b = b(A, n)$ to be selected, we show that it is unlikely to observe a large time at which the sum of indicators in (63) is larger than b^γ (provided b is large enough).

Keeping the above explanation in mind, we start by observing that, since $|\mathcal{A}_k^i| \leq dk$ (which follows after noticing that, at each step of the exploration process, at most d stubs become active), we have

$$\mathbb{P}\left(\exists t \leq (t_i - t_{i-1}) \wedge b : h_k^i \in \mathcal{A}_{k-1}^i \mid \mathcal{F}_{i-1}\right) \leq \sum_{k=1}^b \frac{d(k-1)}{dn - 2(t_{i-1} + k - 1) - 1} \leq \frac{2b^2}{n}$$

for all large enough n , where for the last inequality we have used that, on \mathcal{G}_{i-1} , we have $t_{i-1} + k \leq BLT^{1/2} + T \leq 2BLT^{1/2}$ and $BLT^{1/2} \asymp A^{1/2}n^{2/3} \ll n$ (as $A \ll n^{2/3}$).

Next we consider the case of *large* t , i.e. we take $t > b$. In this case, we set

$$\mathcal{B} := \{|\mathcal{A}_k^i| \leq \ell \forall t \leq (t_i - t_{i-1}) \wedge T\}, \ell \in \mathbb{N},$$

and observe that

$$\begin{aligned} \mathbb{P}\left(\sum_{i=1}^{(t_i - t_{i-1}) \wedge T} \mathbb{1}_{\{h_k^i \in \mathcal{A}_{k-1}^i\}} \geq b^\gamma \mid \mathcal{F}_{i-1}\right) \\ \leq \mathbb{P}\left(\left\{\sum_{i=1}^{(t_i - t_{i-1}) \wedge T} \mathbb{1}_{\{h_k^i \in \mathcal{A}_{k-1}^i\}} \geq b^\gamma\right\} \cap \mathcal{B} \mid \mathcal{F}_{i-1}\right) + \mathbb{P}(\mathcal{B}^c \mid \mathcal{F}_{i-1}). \end{aligned} \quad (64)$$

Let us ignore for the moment the second probability on the right-hand side of (64). Arguing exactly as in Section 4.2.5 of [13], it is not difficult to see that

$$\mathbb{P}\left(\left\{\sum_{i=1}^{(t_i - t_{i-1}) \wedge T} \mathbb{1}_{\{h_k^i \in \mathcal{A}_{k-1}^i\}} \geq b^\gamma\right\} \cap \mathcal{B} \mid \mathcal{F}_{i-1}\right) \leq \exp(-b^\gamma + cT\ell n^{-1})$$

for some constant $c > 0$. But then combining the last estimates we can bound from above the probability which appears in the statement of the lemma by

$$\begin{aligned} & \mathbb{P}(\exists t \leq (t_i - t_{i-1}) \wedge b : h_k^i \in \mathcal{A}_{k-1}^i \mid \mathcal{F}_{i-1}) + \mathbb{P}(\exists b < t \leq (t_i - t_{i-1}) \wedge T : \sum_{i=1}^t \mathbb{1}_{\{h_k^i \in \mathcal{A}_{k-1}^i\}} \geq t^\gamma \mid \mathcal{F}_{i-1}) \\ & \leq \frac{2b^2}{n} + \mathbb{P}\left(\sum_{i=1}^{(t_i - t_{i-1}) \wedge T} \mathbb{1}_{\{h_k^i \in \mathcal{A}_{k-1}^i\}} \geq b^\gamma \mid \mathcal{F}_{i-1}\right) \\ & \leq \frac{2b^2}{n} + \exp(-b^\gamma + cT\ell n^{-1}) + \mathbb{P}(\mathcal{B}^c \mid \mathcal{F}_{i-1}). \end{aligned} \quad (65)$$

There remains to bound the (conditional) probability of \mathcal{B}^c , given \mathcal{F}_{i-1} . This is easily done (here we follow closely Lemma 4.6 in [13]). Note that

$$\mathbb{P}(\mathcal{B}^c \mid \mathcal{F}_{i-1}) \leq \mathbb{P}(\exists t \leq (t_i - t_{i-1}) \wedge T : d_i + \sum_{k=1}^t D_k^i > \ell), \quad (66)$$

where $D_k^i := \mathbb{1}_{R_k^i}(d-1) - 1$ (the upper bound is justified by the fact that, for $t \leq t_i - t_{i-1}$, we can write $|\mathcal{A}_t^i| = d_i + \sum_{k=1}^t \eta_k^i$ and each η_k^i is at most D_k^i ; moreover, the R_k^i do not depend on \mathcal{F}_{i-1} , as they only pertain to the percolation process on $\mathbb{G}(n, d)$). Recalling that R_k^i occurs with probability $p = (1 + \lambda n^{-1/3})/(d-1)$ (being the event that the edge revealed at step k is retained), we see that $\sum_{k=1}^t D_k^i$ is a sub-martingale for $\lambda > 0$ (an assumption which can be made without loss of generality); an application of lemma 1.2 (i.e. Doob's inequality) as in Lemma 4.6 of [13] yields that the probability on the right-hand side of (66) is at most

$$C \exp\left(-\frac{\ell^2}{p(d-1)T} + \frac{\ell}{2}(1 - 1/[p(d-1)])\right) \leq C \exp\left(-\frac{\ell^2}{p(d-1)T} + C' \frac{\ell|\lambda|}{n^{1/3}}\right) \quad (67)$$

for positive constants $C, C' > 0$. Setting for instance $b := \lceil A^{1/5} n^{1/3} \rceil$ and $\ell := n^{1/3+\gamma/3} A$ we see that $T\ell n^{-1} \ll b^\gamma$ as well as $\ell^2/T \gg \ell/n^{1/3}$. Thus, the exponential term in (65) is at most e^{-cb^γ} , whereas the expression on the right-hand side of (67) is at most $e^{-c\ell^2/T}$. Both these terms are, for all large enough n , at most $b^2/n \asymp A^{2/5}/n^{1/3}$, which in turn is $\ll A^{1/2}/n^{1/3} \asymp T^{-1/2}$, concluding the proof of the lemma. \square

3 Boosting the martingale method of Nachmias and Peres

The proof of Theorem 1.1 rely on random walk estimates which are easily applicable provided that the (conditional) distribution of the number of unseen vertices/stubs becoming active at any step of the underlying exploration process is 'simple enough'. When the above-mentioned conditional law is more complicated, an analysis like the one we provided in Section 2 becomes more involved. To this end, the goal of this section is to show how to optimize the (very robust) martingale argument of Nachmias and Peres [27, 28] in order to obtain *stretched*-exponential upper bounds for the probability of observing an unusually small maximal cluster in critical random graphs. With our adjustments, we can show that the above probability is at most $\exp(-A^{3/5})$ in other two (critical) models (which are introduced below), thus considerably improving upon the existing polynomial bounds available in the literature for such random graphs; this is the content of Theorem 3.1 below. (We remark that the same methodology used to establish the bounds of Theorem 3.1 applies to the models treated in Theorem 1.1; however, since we already derived *better* bounds for those random graphs in Section 2, we won't treat them again here.)

For the random graphs treated in Theorem 3.1 we are content with the stretched bounds given here, mostly because we believe that the optimised version of the martingale argument (described below) is interesting on its own and provides a good alternative to the more precise methodology of Section 2. However, we believe that by adapting the same argument used to analyse the near-critical $\mathbb{G}(n, p)$ and $\mathbb{G}(n, d, p)$ models, it should be possible to show that the probability of observing an unusually small maximal components in these other (critical) models is also at most of order $\exp(-A^{3/2})$. However, such an adaptation requires some work, since the (conditional) distributions of the random variables η_i are now more complicated: they are still binomials, but now also the success probabilities are random

(and consequently the random walk estimates used to treat the models of Theorem 1.1 requires some work to be implemented).

Before stating Theorem 3.1, we first need to introduce the critical random graphs considered there.

Given $n, k \in \mathbb{N}$, let $V := \{v_1, \dots, v_n\}$ and $W := \{w_1, \dots, w_k\}$. The random *bipartite* graph $\mathbb{B}(n, k, p)$ is obtained by performing p -bond percolation on the complete bipartite graph with bipartition (V, W) where each node v_i is linked to each w_j (for $i \in [n], j \in [k]$). The random intersection graph $\mathbb{G}(n, k, p)$ model is constructed from $\mathbb{B}(n, k, p)$ in the following way. The vertex set is V and each pair of distinct vertices $\{v_i, v_j\}$ is present as an edge if, and only if, v_i and v_j share *at least one* common neighbour in $\mathbb{B}(n, k, p)$. We call V the set of *vertices*, whereas we call W the set of *attributes* (or features or auxiliary vertices). This random graph model was analysed by Behrisch [2] for the case $k = k(n) = n^\alpha$ and $p^2k = c/n$, with $\alpha, c \in (0, 1) \cup (1, \infty)$. As shown by Stark [33], the vertex degree distribution (i.e. the distribution of the degree of a vertex selected uniformly at random) is highly dependent on the value of α . However, as shown by Deijfen and Kets [14], the clustering of the graph is controllable only when $\alpha = 1$. Indeed, when $\alpha \in (0, 1)$, the clustering coefficient converges to 1 as $n \rightarrow \infty$, while it converges to 0 (as $n \rightarrow \infty$) when $\alpha > 1$. On the other hand, in the regime $\alpha = 1$, the clustering coefficient converges to β (as $n \rightarrow \infty$) and so, by specifying the value of β , one can control the clustering of the graph. This latter regime where $k = \Theta(n)$ was subsequently investigated by Lageras and Lindholm [26]. Specifically, in [26] the authors considered the $\mathbb{G}_{n,k,p}$ random graph with $k = \lfloor \beta n \rfloor$ and $p = \gamma/n$, where $\gamma, \beta > 0$ are model parameters and proved that the $\mathbb{G}_{n,k,p}$ model undergoes a phase transition as $\beta\gamma^2$ passes 1. Indeed, setting $\mu = \beta\gamma^2$, they proved that if $\mu < 1$ (sub-critical case) then with probability tending to one there is no component in $\mathbb{G}_{n,k,p}$ with more than $O(\log(n))$ vertices, while if $\mu > 1$ (super-critical case) then, with probability tending to one, there exists a unique giant component of size $n\delta$ (with $\delta \in (0, 1)$ constant) and the size of the second largest component is at most of order $\log(n)$. It was then shown in [9, 10] that, in the (critical) case $\mu = 1$, there is a constant $c = c(\gamma, \beta) > 0$ depending on γ, β such that, for all large enough A and n ,

$$\mathbb{P}\left(|\mathcal{C}_{\max}(\mathbb{G}_{n,k,p})| > An^{2/3}\right) \leq cA^{-3/2} \text{ and } \mathbb{P}\left(|\mathcal{C}_{\max}(\mathbb{G}_{n,k,p})| < n^{2/3}/A\right) \leq cA^{-1/2}. \quad (68)$$

The second model we consider in this section is the so-called (critical) *quantum random graph*. Here we only provide an informal definition of such a model and refer the interested reader to [15] and Section 3.3 below for the rigorous definition (see also [10]).

We have n circles of length $\beta > 0$, punctured with *independent* Poisson point processes of intensity $\lambda > 0$. Because of these (Poisson) processes of holes, circles can be written as *finite disjoint* unions of connected intervals. The edges of the graph link intervals on distinct circles and they are generated as follows. Given any two (distinct) circles $u \neq v$, we run another *independent* Poisson point process of intensity $1/n$ on a (third) circle of length $\beta > 0$. If such a point process jumps at a time which is contained in both an interval of u and an interval of v (recall the decomposition mentioned earlier), then these two intervals are considered directly connected. The resulting random graph, denoted by $\mathbb{G}(n, \beta, \lambda)$, is called *quantum random graph*. As explained in [15], the notion of component of a point $x = (u, t)$ (where u is an element of $[n]$ while t is a point of the circle) in this random graph ensemble is well-defined and the *size* of a component is the number of *intervals* it contains. The critical parameter (curve) for the quantum random graph in the (β, λ) -parameter space was identified by Ioffe and Levit [23] by comparison with a critical branching process whose offspring distribution is the so-called *cut-gamma* law $\Gamma_\theta(2, 1)$, which is the distribution of $J := (J_1 + J_2) \wedge \theta$ for independent random variables J_1, J_2 with the $\text{Exp}(1)$ law. Setting $F(\lambda\beta) := 2(1 - e^{-\lambda\beta}) - \lambda\beta e^{-\lambda\beta}$, criticality occurs when

$$F(\beta, \lambda) := \lambda^{-1}F(\lambda\beta) = 1,$$

with $\lambda^{-1}F(\lambda\beta)$ corresponding to the expected length of an interval I in the quantum random graph. (It was shown in [23] that, if $F(\beta, \lambda) > 1$, then a giant component of order $\Theta(n)$ emerges, whereas when $F(\beta, \lambda) < 1$ all components are typically of order $O(\log(n))$; see [15] and [23]). Dembo, Levit and Vadlamani [15] analysed the (near-)critical behaviour of this model and showed that, when $F(\beta, \lambda) = 1$, then $|\mathcal{C}_{\max}(\mathbb{G}(n, \beta, \lambda))|$ is of order $n^{2/3}$. Amongst many other things, they showed that there is a constant $c = c(\beta) > 0$ depending on β such that, for all sufficiently large A and n ,

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, \beta, \lambda))| > An^{2/3}) \leq cA^{-3/2} \text{ and } \mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, \beta, \lambda))| < n^{2/3}/A) \leq cA^{-3/5}. \quad (69)$$

By means of the optimised martingale version we can give much better tail bounds (compared to (68) and (69)) for the probability that $|\mathcal{C}_{\max}(\mathbb{G})| < n^{2/3}/A$ when either $\mathbb{G} = \mathbb{G}(n, k, p)$ or $\mathbb{G} = \mathbb{G}(n, \beta, \lambda)$.

Theorem 3.1. *Let $p = \gamma/n$ and $k = \lfloor \beta n \rfloor$, with $\gamma, \beta > 0$ such that $\beta\gamma^2 = 1$. There exist constants $A_0 = A_0(\gamma, \beta) > 0$ and $c = c(\beta, \lambda) > 0$ such that, for any $A_0 \leq A = O(n^{5/12})$ and for all large enough n , we have*

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, k, p))| < n^{2/3}/A) \leq \exp(-cA^{3/5}). \quad (70)$$

Let $F(\beta, \lambda) = 1$. There exist constants $A_0 = A_0(\lambda, \beta) > 0$ and $c = c(\lambda, \beta) > 0$ such that, for any $A_0 \leq A = O(n^{5/12})$ and for all large enough n , we have

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}(n, \beta, \lambda))| < n^{2/3}/A) \leq \exp(-cA^{3/5}). \quad (71)$$

Remark 6. The requirement $A = O(n^{5/12})$ can be weakened to $A = o(n^{2/3})$, at the expenses of adding a term like $\exp(-cn^{1/3}/A^{1/5})$ to both bounds appearing in (70) and (71), which considerably improve upon (68) and (69).

Remark 7. As we pointed out at the start of this section, the same method used to establish Theorem 3.1 can also be used to prove similar bounds for the critical random graphs $\mathbb{G}(n, p)$ and $\mathbb{G}(n, d, p)$; however, we have already established stronger estimates in Section 2 and hence there is no reason to use the optimised martingale argument to obtain bounds of the type $\exp(-cA^{3/5})$ for these two models.

We continue by providing a detailed, but informal explanation on how to optimize the martingale argument introduced in [27, 28] to obtain the stretched exponential upper bounds displayed in Theorem 3.1; we do so in terms of the critical $\mathbb{G}(n, p)$ model, but the same strategy works for the critical random graphs $\mathbb{G}(n, k, p)$, $\mathbb{G}(n, \beta, \lambda)$ (and $\mathbb{G}(n, d, p)$).

3.1 Heuristic derivation of the stretched exponential bounds

Here we show how the original argument of [28] can be boosted to show that $|\mathcal{C}_{\max}(\mathbb{G}(n, 1/n))| \geq n^{2/3}/A$ with probability at least of order $1 - \exp(-A^{3/5})$; we provide the explanation in terms of the $\mathbb{G}(n, 1/n)$ model for simplicity of exposition, but the same argument applies to the other models considered in this work.

Recall the exploration process for the $\mathbb{G}(n, p)$ model, with Y_t denoting the number of active vertices at time $t \in \mathbb{N}$ in the exploration process of Section 2 for the $\mathbb{G}(n, p)$ model. As usual, $t_0 := 0$ and denote by t_i the first time $t > t_{i-1}$ at which $Y_t = 0$, for $i \in \mathbb{N}$ (prior to the end of the procedure). Then

$$\mathbb{P}(\mathcal{C}_{\max}(\mathbb{G}(n, p))) \leq \mathbb{P}(t_i - t_{i-1} < n^{2/3}/A \ \forall i). \quad (72)$$

Following Nachmias and Peres [28], the basic idea is then to show that it is (very) likely for the process Y_t to go above some large enough level $\ell = \ell(n, A)$ before some time $\mathbb{N} \ni T = T(\ell) \ll n$ and then to remain positive for at least $n^{2/3}/A$ steps.

Hence we bound from above the probability on the right-hand side of (72) by

$$\mathbb{P}(t_i - t_{i-1} < n^{2/3}/A \ \forall i, \exists t < T : Y_t \geq \ell) + \mathbb{P}(Y_t < \ell \ \forall t < T). \quad (73)$$

At this stage we introduce the necessary changes to the original argument of [28] in order to show that both terms in (73) are at most $\lesssim \exp(-cA^{3/5})$.

Before starting the actual (heuristic) derivation, we make a general observation concerning the choice of ℓ and T . The first probability appearing in (73) is at most the probability that the process Y_t , after having reached level ℓ (and considered from the the first time it does so), reaches zero in less than $n^{2/3}/A$ steps (otherwise we would find a positive excursion of Y_t lasting for *more* than $n^{2/3}/A$ steps). If we want this probability to be small, the level ℓ has to be chosen large; intuitively, thinking of Y_t as a \mathbb{Z} -valued, mean-zero random walk (this approximation is justified later), we need ℓ to be much bigger than the square root of the time, whence we require that $\ell \gg n^{1/3}/A^{1/2}$. This forces T to be large, otherwise in the second probability in (73) we would be asking the number of active vertices Y_t to remain below a large value for a short time interval, a likely event. In particular, we need T to be much bigger than ℓ^2 , i.e. we require $T \gg \ell^2$. But, as we will see below, the increments of the process Y_t have a (negative) drift of order $-T/n$, whence a very large value of T would cause Y_t to reach zero too quickly, thus making the first probability in (73) large. All in all, it becomes evident that a trade-off between ℓ and T is needed in order to make *both* terms in (73) small at the same time.

Let us now illustrate the simple argument which we use to obtain stretched exponential bounds for both expressions in (73). Let $1 \ll h, m \in \mathbb{N}$ (possibly dependent on A, n) and $\delta > 0$ (independent of A, n). To bound the second term in (73), i.e. the probability that the number of active vertices stays

below ℓ for T steps, we set $\ell := h\delta$ and split the (discrete) interval $[T] := \{1, \dots, T\}$ into m (smaller) disjoint, connected sub-intervals of length h^2 and we show that, in each one of these m smaller intervals, the process has a *constant, positive* probability (which depends on δ) to go above the level $h\delta$ in h^2 steps, provided $\delta < 1$ is sufficiently small. This is true because the random process Y_t behaves like a \mathbb{Z} -valued random walk with iid increments having mean zero and finite second moment, and such a random walk has a constant positive probability (independent of h) to be above $h\delta$ after h^2 steps. Denoting by $c = c(\delta) < 1$ the positive constant bounding from above the probability that Y_t stays below $h\delta$ for h^2 steps and iterating the argument m times, we obtain that Y_t stays below $h\delta$ for $T = mh^2$ consecutive steps with probability at most $c^m = \exp(-m \log(1/c))$.

To obtain an exponential (upper) bound for the first probability in (73) instead, we proceed as follows. Let τ be the minimum between the first time $t \in \mathbb{N}$ at which the number of active vertices Y_t goes above $h\delta$ and $T = mh^2$. Denote by η_t the number of *unseen* vertices which are added to the set of active nodes at step t . We start by noticing that the process $Y_{\tau+s}$ (started at time τ) must reach zero in *less* than $n^{2/3}/A$ steps, otherwise we would discover a positive excursion consisting of more than $n^{2/3}/A$ steps. Hence there is a first time $t < n^{2/3}/A$ at which $Y_{\tau+t} = 0$. Observe that at time $\tau + i$ we add $\eta_{\tau+i}$ vertices to the set of active nodes and remove one vertex from such a set (as long as the set of active nodes remains non-empty); thus, since the number of active vertices stays positive from time τ until time $\tau + t$, we can write $Y_{\tau+s} = Y_\tau + \sum_{i=1}^s (\eta_{\tau+i} - 1)$ for $s \leq t$. But we know that $Y_\tau \geq h\delta$ (and $Y_{\tau+t} = 0$), whence we arrive at $\sum_{i=1}^t (\eta_{\tau+i} - 1) \leq -h\delta$. Conditional on the history of the exploration process until step $\tau + i - 1$, the random variable $\eta_{\tau+i}$ has the $\text{Bin}(n - \tau - i + 1 - Y_{\tau+i-1}, p)$ distribution (with $n - \tau - i + 1 - Y_{\tau+i-1}$ being the number of *unseen* vertices available at time $\tau + i$). Since the number of active vertices is never ‘too large’, to continue our derivation we approximate $\text{Bin}(n - \tau - i - 1 - Y_{\tau+i-1}, p) \approx \text{Bin}(n - \tau - i, p)$. Then, roughly, $\sum_{i=1}^t (\eta_{\tau+i} - 1) \leq -h\delta$ occurs if, and only if, $\sum_{i=1}^t (\text{Bin}_i(n - \tau - i, p) - 1) \leq -h\delta$ happens. Since $\tau \leq mh^2$ and $t \leq n^{2/3}/A$, the latter event ‘implies’ that $\sum_{i=1}^t (\text{Bin}_i(n - mh^2 - n^{2/3}/A, p) - 1) \leq -h\delta$ occurs. Moreover, using the fact that binomial random variables concentrate around their mean (whence in particular $\sum_{i=1}^t \text{Bin}_i(T + n^{2/3}/A, p) \approx (tmh^2)/n + t/(n^{1/3}A)$, where we used that $p = 1/n$ and $t < n^{2/3}/A$), we conclude that the probability of $\sum_{i=1}^t (\text{Bin}_i(n - mh^2 - n^{2/3}/A, p) - 1) \leq -h\delta$ happening at some $t < n^{2/3}/A$ is (very) roughly the same as the probability of $\sum_{i=1}^t (\text{Bin}_i(n, p) - 1) \leq -h\delta + (tmh^2)/n + t/(n^{1/3}A)$ occurring at some $t < n^{2/3}/A$. Now notice that, since $t < n^{2/3}/A$, we can bound

$$(tmh^2)/n \leq (mh^2)/(n^{1/3}A) \text{ and } t/(n^{1/3}A) \leq n^{1/3}/A^2$$

and hence the last probability is at most the probability of seeing a time $t < n^{2/3}/A$ at which $\sum_{i=1}^t (\text{Bin}_i(n, p) - 1) \leq -h\delta + (mh^2)/(n^{1/3}A) + n^{1/3}/A^2$, which is small only if both $(mh^2)/(n^{1/3}A)$ and $n^{1/3}/A^2$ are sufficiently smaller than $h\delta$, say smaller than $(h\delta)/3$. In this case then, we would be bounding from above the probability that a mean-zero random walk (with supposedly iid, well-behaved) $\text{Bin}_i(n, p) - 1$ increments goes below $\asymp -h$ in less than $n^{2/3}/A$ steps. An application of Doob’s submartingale inequality yields that this occurs with probability at most $\lesssim \exp(-(h^2A)/n^{2/3})$. Combining this estimate with our earlier bound of order e^{-m} we see that the sum of the two terms in (73) is at most of order

$$\exp(-(h^2A)/n^{2/3}) + \exp(-m). \quad (74)$$

The first exponential is small only if $h \gg n^{1/3}/A^{1/2}$ (in which case the earlier constraint $n^{1/3}/(2A^2) \leq (h\delta)/3$ is clearly satisfied provided A is large enough). Recalling the other constraint we imposed earlier, namely that $(mh^2)/(An^{1/3}) \leq h/3$, or equivalently $h \leq (An^{1/3})/(3m)$, we immediately see the necessity of having

$$n^{1/3}/A^{1/2} \ll h \leq (An^{1/3})/(3m),$$

which leads to $m \ll A^{3/2}$, thus excluding right away the optimal bound of order $\exp(-A^{3/2})$. The best choice seems to be something like $m = A, h = n^{1/3}$, in which case the expression in (74) would be $\lesssim \exp(-A)$; however, as we will see later on, a further technical requirement forbids this choice. Indeed, we will also need $(mh^3)/n$ to be at most of (small) constant order, and taking m, h as above would violate the latter requirement. Hence, including this extra constraint, the optimal choice seems to be $h = n^{1/3}/A^{1/5}, m = A^{3/5}$, thus leading to the bound $\lesssim \exp(-A^{3/5})$ (under the assumption $A = O(n^{5/12})$).

3.2 The $\mathbb{G}(n, k, p)$ model

Recall that we are interested in the case where $k = \lfloor \beta n \rfloor$ and $p = \gamma/n$, for some $\beta\gamma > 0$ which are model parameters. Moreover, throughout we assume that $\gamma^2\beta = 1$.

As we did in the previous sections, we start by describing an exploration process to reveal the connected components of the random intersection graph $\mathbb{G}_{n,k,p}$.

Fix an ordering of the n vertices in V with v first. Let us denote by \mathcal{A}_t , \mathcal{U}_t and \mathcal{E}_t the (random) sets of active, unseen and explored *vertices* at the end of step $t \in \mathbb{N}_0$, respectively. Then, for every $t \in \mathbb{N}_0$, we have $V = \mathcal{A}_t \cup \mathcal{U}_t \cup \mathcal{E}_t$ (a disjoint union), so that in particular $\mathcal{U}_t = V \setminus (\mathcal{A}_t \cup \mathcal{E}_t)$ at each step t . Moreover, we denote by \mathcal{D}_t the (random) sets of discovered *attributes* by the end of step $t \in \mathbb{N}_0$, so that $W \setminus \mathcal{D}_t$ is the set of fresh attributes at the end of step t .

Algorithm 3. At time $t = 0$, vertex v is declared active whereas all other vertices are declared unseen; moreover, all the attributes are fresh. Therefore $\mathcal{A}_0 = \{v\}$, $\mathcal{E}_0 = \emptyset$ and $\mathcal{D}_0 = \emptyset$. For every $t \in \mathbb{N}$, the algorithm proceeds as follows.

- (a) If $|\mathcal{A}_{t-1}| \geq 1$, we let u_t be the first active node (here and in what follows, the term *first* refers to the ordering that we have fixed at the very beginning of the procedure).
- (b) If $|\mathcal{A}_{t-1}| = 0$ and $|\mathcal{U}_{t-1}| \geq 1$, we let u_t be the first unseen vertex.
- (c) if $|\mathcal{A}_{t-1}| = 0 = |\mathcal{U}_{t-1}|$ (so that $\mathcal{E}_{t-1} = V$), we halt the procedure.

Let us denote by \mathcal{N}_t the (random) set of (fresh) attributes that are linked to u_t ; formally,

$$\mathcal{N}_t := \{w \in W \setminus \mathcal{D}_{t-1} : w \sim u_t\}. \quad (75)$$

Moreover, we denote by \mathcal{R}_t the (random) set of unseen neighbours of u_t that are linked to *at least one* of the attributes in \mathcal{N}_t ; formally, we set $\mathcal{R}_t := \{u \in \mathcal{U}_{t-1} \setminus \{u_t\} : u \sim w \text{ for some } w \in \mathcal{N}_t\}$ (and note that $\mathcal{U}_{t-1} \setminus \{u_t\} = \mathcal{U}_{t-1}$ if $\mathcal{A}_{t-1} \neq \emptyset$). Then we update $\mathcal{U}_t := \mathcal{U}_{t-1} \setminus (\mathcal{R}_t \cup \{u_t\})$, $\mathcal{A}_t := (\mathcal{A}_{t-1} \setminus \{u_t\}) \cup \mathcal{R}_t$, $\mathcal{E}_t := \mathcal{E}_{t-1} \cup \{u_t\}$ and $\mathcal{D}_t := \mathcal{D}_{t-1} \cup \mathcal{N}_t$.

Remark 8. Since in the procedure **Algorithm 3** we explore *one vertex* at each step, we have $\mathcal{A}_t \cup \mathcal{U}_t \neq \emptyset$ for every $t \leq n-1$ and $\mathcal{A}_n \cup \mathcal{U}_n = \emptyset$ (as $\mathcal{E}_n = V$). Thus the algorithm runs for n steps.

Denoting by η_t the (random) number of unseen vertices that we add to the set of active nodes at time t , since at the end of each step i in which $|\mathcal{A}_{i-1}| \geq 1$ we remove the (active) vertex u_i from \mathcal{A}_{i-1} (after having revealed its unseen neighbours), we have the recursion

- $|\mathcal{A}_t| = |\mathcal{A}_{t-1}| + \eta_t - 1$ if $|\mathcal{A}_{t-1}| > 0$;
- $|\mathcal{A}_t| = \eta_t$ if $|\mathcal{A}_{t-1}| = 0$.

Let us denote by Y_t and U_t the number of active and unseen vertices at the end of step $t \in [n] \cup \{0\}$, respectively, so that

$$Y_t = |\mathcal{A}_t| \text{ and } U_t = |\mathcal{U}_t| = n - t - Y_t.$$

Denote by \mathcal{F}_t the σ -algebra generated by all the information collected by the exploration process until the end of step t . Then, conditional on \mathcal{F}_{t-1} and \mathcal{N}_t , the random variable η_t depends on the past until time $t-1$ only through Y_{t-1} and, in particular, setting $N_t := |\mathcal{N}_t|$ we have

$$\eta_t =_d \text{Bin}(U_{t-1} - \mathbb{1}_{\{Y_{t-1}=0\}}, 1 - (1-p)^{N_t}) \text{ for } t \in [n]. \quad (76)$$

We remark that each random variable N_t satisfies $N_t \leq_{sd} \text{Bin}(k, p)$; this is immediate from the description of **Algorithm 3** given above.

Define $t_0 := 0$ and $t_i := \min\{t \geq t_{i-1} + 1 : Y_t = 0\}$, for $i \geq 1$ (as long as we do not enter Step (c) in **Algorithm 3**). Then, denoting by \mathcal{C}_i the i -th component revealed during the exploration process, we have $|\mathcal{C}_i| = t_i - t_{i-1}$ for all i and so, setting $T := \lceil n^{2/3}/A \rceil$, we can write

$$\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}_{n,k,p})| < n^{2/3}/A) = \mathbb{P}(t_i - t_{i-1} < n^{2/3}/A \forall i) \leq \mathbb{P}(t_i - t_{i-1} < T \forall i). \quad (77)$$

Following the strategy described in Section 3.1, we let $1 \ll h, m \in \mathbb{N}$ (possibly dependent on A, n) and denote by $\delta \in (0, 1)$ a constant that we specify later. Then we bound

$$\begin{aligned} \mathbb{P}(t_i - t_{i-1} < T \forall i) &\leq \mathbb{P}(t_i - t_{i-1} < T \forall i, \exists t \in [mh^2] : Y_t \geq h\delta) \\ &\quad + \mathbb{P}(Y_t < h\delta \forall t \in [mh^2]) \end{aligned} \quad (78)$$

and control these two terms separately. We start by bounding the first term on the right-hand side of (78) with of Proposition 3.1 below and subsequently, with Proposition 3.2, we control the second probability on the right-hand side of (78).

Proposition 3.1. *Suppose that $(mh^2)/n = O(1)$ and $T^2/(hn) \leq \delta/2$. Then, for all large enough n , we have*

$$\mathbb{P}(t_i - t_{i-1} < T \forall i, \exists t \in [mh^2] : Y_t \geq h\delta) \leq e^{-ch} \vee e^{-(ch^2)/T},$$

for some constant $c = c(\gamma, \beta, \delta)$ which depends on γ, β and δ .

Proof. Let us denote by τ' the first time $t \geq 1$ at which $Y_t \geq h\delta$ and set $\tau := \tau' \wedge mh^2$. Note that, on the event $\{\exists t \in [mh^2] : Y_t \geq h\delta\}$, we have $Y_\tau \geq h\delta$; moreover, if $t_i - t_{i-1} < T$ for every i , then necessarily there exists a (first) time $t < T$ such that $Y_{\tau+s} > 0$ for all $s \leq t-1$ and $Y_{\tau+t} = 0$ (otherwise we would find an excursion lasting for more than T steps, which is impossible on the event $\{t_i - t_{i-1} < T \forall i\}$). Thus we can express $Y_{\tau+t}$ as $Y_{\tau+t} = Y_\tau + \sum_{i=1}^t (\eta_{\tau+i} - 1)$. Hence, since on the event $\{Y_\tau \geq h\delta\}$ we have $-Y_\tau \leq -h\delta$, we can bound

$$\begin{aligned} \mathbb{P}(t_i - t_{i-1} < T \forall i, \exists t \in [mh^2] : Y_t \geq h\delta) &\leq \mathbb{P}(\exists t < T : Y_{\tau+s} > 0 \forall s \leq t-1, Y_{\tau+t} = 0, Y_\tau \geq h\delta) \\ &\leq \mathbb{P}(\exists t < T : Y_{\tau+s} > 0 \forall s \leq t-1, \sum_{i=1}^t (\eta_{\tau+i} - 1) \leq -h\delta). \end{aligned} \quad (79)$$

In order to control (from above) the number of active vertices which we observe in the time window $[\tau, \tau + T] \cap \mathbb{N}$ (the need for a uniform upper bound on Y_t will become clear in a moment), we define, for $t < T$, the events

$$\mathcal{H}_1(t) := \{Y_{\tau+s} > 0 \forall s \leq t-1, \sum_{i=1}^t (\eta_{\tau+i} - 1) \leq -h\delta\} \text{ and } \mathcal{H}_2(t) := \{Y_{\tau+z} < h \forall z \leq t-1\}.$$

Then the probability on the right-hand side of (79) is at most

$$\mathbb{P}\left(\bigcup_{t < T} \mathcal{H}_1(t) \cap \mathcal{H}_2(t)\right) + \mathbb{P}\left(\bigcup_{t < T} \{Y_{\tau+s} > 0 \forall s \leq t-1\} \cap \mathcal{H}_2^c(t)\right). \quad (80)$$

With the next lemma we show that the second probability on the right-hand side of (80) is small.

Lemma 3.1. *Suppose that $(mh^2)/n = O(1)$. Then, for all large enough n , we have*

$$\mathbb{P}\left(\bigcup_{t < T} \{Y_{\tau+s} > 0 \forall s \leq t-1\} \cap \mathcal{H}_2^c(t)\right) \leq e^{-ch} \vee e^{-(ch^2)/T},$$

for some constant $c = c(\gamma, \beta) > 0$ which depends on γ, β .

Proof. We can bound from above the probability which appears in the statement of the lemma by

$$\mathbb{P}\left(\exists t < T : \sum_{i=1}^t (X_{\tau+i} - E_{\tau+i}) \geq h-1 - \sum_{i=1}^t E_{\tau+i}\right), \quad (81)$$

where we set $X_{\tau+i} := \eta_{\tau+i} - 1$ and $E_{\tau+i} := \mathbb{E}[X_{\tau+i} | \mathcal{F}_{\tau+i-1}]$. Now recall from (76) that, conditional on $\mathcal{F}_{\tau+i-1}$ and τ , the random variable $\eta_{\tau+i}$ is distributed as

$$\text{Bin}(n - \tau - (i-1) - Y_{\tau+i-1} - \mathbb{1}_{\{Y_{\tau+i-1}=0\}}, 1 - (1-p)^{N_{\tau+i}}). \quad (82)$$

Therefore, using the classical bound $(1-x)^\ell \geq 1 - \ell p$ (which is valid for all $x > -1, \ell \in \mathbb{N}$), we see that

$$1 - (1-p)^{N_{\tau+i}} \leq N_{\tau+i} p \quad (83)$$

and so, as $N_t \leq_{sd} \text{Bin}(k, p)$ for each t and $nkp^2 \leq \gamma^2\beta = 1$, we obtain that

$$E_{\tau+i} \leq np\mathbb{E}[N_{\tau+i}] - 1 \leq nkp^2 - 1 \leq 0. \quad (84)$$

Thus the probability on the right-hand side of (81) is, for any given $r \in (0, 1)$, at most

$$\mathbb{P}(\max_{t < T} e^r \sum_{i=1}^t (X_{\tau+i} - E_{\tau+i}) \geq e^{r(h-1)}). \quad (85)$$

In a moment we will need a lower bound on $|\mathcal{D}_{\tau+i-1}|$, the number of discovered attributes by the end of time $\tau + i - 1$. To this end, we define

$$X'_{\tau+i} := \mathbb{1}_{\{|\mathcal{D}_{\tau+i-1}| \leq \omega\}} X_{\tau+i} \text{ and } E'_{\tau+i} := \mathbb{1}_{\{|\mathcal{D}_{\tau+i-1}| \leq \omega\}} E_{\tau+i} \quad (86)$$

where $\omega \geq 1$ has to be specified. Now observe that, as $\mathcal{D}_t \subset \mathcal{D}_{t+1}$ at all times t (since more and more attributes become discovered as the exploration proceeds), we obtain

$$\mathbb{P}(\exists t < mh^2 + T : |\mathcal{D}_t| > \omega) \leq \mathbb{P}(|\mathcal{D}_{mh^2+T}| > \omega) \leq e^{-r\omega} \mathbb{E}[e^{r\text{Bin}(k,p)}]^{mh^2+T}.$$

Taking $\omega := 2pk(mh^2 + T)$ and $0 < r < 1$, it is immediate to show that the expression on the right-hand side of the last inequality is at most $\exp\{-rpk(mh^2 + T)(1 - r)\}$ whence, setting e.g. $r = 1/2$, we arrive at

$$\mathbb{P}(\exists t < mh^2 + T : |\mathcal{D}_t| > 2pk(mh^2 + T)) \leq e^{-\frac{pk(mh^2+T)}{4}} \leq e^{-c'(mh^2+T)} \quad (87)$$

for some constant $c' = c'(\gamma, \beta) > 0$ depending on γ, β . Thus, going back to (85) and setting

$$\mathcal{W} := \{|\mathcal{D}_t| \leq 2pk(mh^2 + T) \ \forall t < mh^2 + T\}, \quad (88)$$

noticing that (for each $i \in [T]$) we have $X'_{\tau+i} = X_{\tau+i}$ and $E'_{\tau+i} = E_{\tau+i}$ on \mathcal{W} , we can bound (thanks to (87))

$$\begin{aligned} \mathbb{P}(\max_{t < T} e^r \sum_{i=1}^t (X_{\tau+i} - E_{\tau+i}) \geq e^{r(h-1)}) &\leq \mathbb{P}(\{\max_{t < T} e^r \sum_{i=1}^t (X_{\tau+i} - E'_{\tau+i}) \geq e^{r(h-1)}\} \cap \mathcal{W}) + \mathbb{P}(\mathcal{W}^c) \\ &\leq \mathbb{P}(\max_{t < T} e^r \sum_{i=1}^t (X'_{\tau+i} - E'_{\tau+i}) \geq e^{r(h-1)}) + e^{-c'(mh^2+T)}. \end{aligned}$$

Since the process defined by $\exp\{r \sum_{i=1}^t (X'_{\tau+i} - E'_{\tau+i})\}$ is a positive submartingale (relative to $\mathcal{F}_{\tau+t}$), we can use Doob's inequality to conclude that

$$\mathbb{P}(\max_{t < T} e^r \sum_{i=1}^t (X'_{\tau+i} - E'_{\tau+i}) \geq e^{r(h-1)}) \leq e^{-r(h-1)} \mathbb{E}[e^{r \sum_{i=1}^{T-1} (X'_{\tau+i} - E'_{\tau+i})}]. \quad (89)$$

Using (82) together with the chain of inequalities $1 - x \leq e^{-x} \leq 1 - x + x^2/2$ (which are valid for all $x \geq 0$) we see that

$$1 - (1 - p)^{N_{\tau+i}} \geq pN_{\tau+i} - \frac{1}{2}(pN_{\tau+i})^2.$$

Moreover, since on the event $\{|\mathcal{D}_{\tau+i-1}| \leq 2pk(mh^2 + T)\}$ we have $\text{Bin}(k - 2pk(mh^2 + T), p) \leq_{sd} N_{\tau+i}$, we obtain (on such event)

$$p\mathbb{E}[N_{\tau+i} | \mathcal{F}_{\tau+i-1}] \geq p^2(k - 2pk(mh^2 + T))$$

and

$$p^2\mathbb{E}[N_{\tau+i}^2 | \mathcal{F}_{\tau+i-1}] \leq p^2\mathbb{E}[\text{Bin}^2(k, p)] \leq p^2(kp + (kp)^2) = O_{\gamma, \beta}(1/n^2),$$

so that

$$p\mathbb{E}[N_{\tau+i} | \mathcal{F}_{\tau+i-1}] - \frac{1}{2}p^2\mathbb{E}[N_{\tau+i}^2 | \mathcal{F}_{\tau+i-1}] \geq p^2k - O(p^3k(mh^2 + T)).$$

Thus, recalling that $\gamma^2\beta = 1$ (and since $p^3k \leq (\gamma^2\beta)/n^2$), we can conclude that, on $\{|\mathcal{D}_{\tau+i-1}| \leq 2pk(mh^2 + T)\}$,

$$\begin{aligned} \mathbb{E}[\eta_{\tau+i} | \mathcal{F}_{\tau+i-1}] &\geq (n - \tau - (i - 1) - Y_{\tau+i-1} - \mathbb{1}_{\{Y_{\tau+i-1}=0\}})(p^2k - O(p^3k(mh^2 + T))) \\ &\geq 1 - \frac{\tau + i - 1 + Y_{\tau+i-1} + \mathbb{1}_{\{Y_{\tau+i-1}=0\}}}{n} - O_{\gamma, \beta}((mh^2 + T)/n^2). \end{aligned} \quad (90)$$

Whence, recalling the definition of $E'_{\tau+i}$, we obtain

$$E'_{\tau+i} \geq -\mathbb{1}_{\{|\mathcal{D}_{\tau+i-1}| \leq 2pk(mh^2+T)\}} \left(\frac{\tau+i-1 + Y_{\tau+i-1} + \mathbb{1}_{\{Y_{\tau+i-1}=0\}}}{n} + O_{\gamma,\beta}((mh^2+T)/n^2) \right).$$

A repeated application of the inequality $e^x \leq 1+x+x^2$ (which is valid for all $x \in [0,1]$) together the fact that $e^x \geq 1+x$ for all $x \in \mathbb{R}$ yields

$$\mathbb{E}[e^{rX'_{\tau+i}} | \mathcal{F}_{\tau+i-1}] \leq \exp \left(-r\mathbb{1}_{\{|\mathcal{D}_{\tau+i-1}| \leq 2pk(mh^2+T)\}} \frac{\tau+i-1 + Y_{\tau+i-1} + \mathbb{1}_{\{Y_{\tau+i-1}\}}}{n} + O_{\gamma,\beta}(r^2) \right). \quad (91)$$

Thus we arrive at

$$\mathbb{E}[e^{r(X'_{\tau+i} - E'_{\tau+i})} | \mathcal{F}_{\tau+i-1}] = e^{-rE'_{\tau+i}} \mathbb{E}[e^{rX'_{\tau+i}} | \mathcal{F}_{\tau+i-1}] \leq e^{O_{\gamma,\beta}(r^2) + O_{\gamma,\beta}(r(mh^2+T)/n^2)}.$$

It follows that

$$\mathbb{E}[e^{r \sum_{i=1}^{T-1} (X'_{\tau+i} - E'_{\tau+i})}] \leq e^{O_{\gamma,\beta}(r^2) + O_{\gamma,\beta}(r(mh^2+T)/n^2)} \mathbb{E}[e^{r \sum_{i=1}^{T-1} (X'_{\tau+i} - E'_{\tau+i})}]$$

and iterating we arrive at

$$\mathbb{E}[e^{r \sum_{i=1}^{T-1} (X'_{\tau+i} - E'_{\tau+i})}] \leq e^{O_{\gamma,\beta}(r^2T) + O_{\gamma,\beta}(T(mh^2+T)/n^2)}.$$

Note that, as $T/n \ll 1$ and $(mh^2)/n = O(1)$ by assumption, we obtain

$$O_{\gamma,\beta}(rT(mh^2+T)/n^2) = O_{\gamma,\beta}(r(mh^2+T)/n) = O_{\gamma,\beta}(r).$$

Hence, when n is large enough, we can bound from above the expression on the right-hand side of (89) by

$$c \exp(-rh + O_{\gamma,\beta}(r^2T)),$$

for some constant $c = c(\gamma, \beta) > 0$. Now if $T/h \leq C$ for some constant $C > 0$, we can find a small enough $r = r(C, \gamma, \beta) > 0$ (which does not depend on n) such that $O_{\gamma,\beta}((rT)/h) \leq 1/2$ and hence we obtain

$$c \exp\{-rh + O_{\gamma,\beta}(r^2T)\} \leq ce^{-(rh)/2}.$$

On the other hand, if $T/h \gg 1$, then we can find a large enough constant $C = C(\gamma, \beta)$ such that, taking $r = h/(CT) \ll 1$,

$$c \exp(-rh + O_{\gamma,\beta}(r^2T)) \leq ce^{-h^2/(C'T)}$$

for some other constant $C' > 0$ (depending on γ, β). All in all (making use of (87)), we conclude that there is a large enough constant $c = c(\gamma, \beta) > 0$ which depends on γ, β such that, for all large enough n ,

$$\mathbb{P}(\exists t < T : \sum_{i=1}^t (X_{\tau+i} - E_{\tau+i}) \geq h-1) \leq e^{-ch} \vee e^{-(ch^2)/T} + e^{-c'(mh^2+T)}.$$

Since $mh^2 \gg h^2/T$, the last exponential term is of smaller order and the desired conclusion follows. \square

Going back to (80) we see that, in order to complete the proof of Proposition 3.1, we still need to bound (from above) the probability

$$\mathbb{P}\left(\bigcup_{t < T} \mathcal{H}_1(t) \cap \mathcal{H}_2(t)\right) \leq \mathbb{P}\left(\bigcup_{t < T} \left\{ \sum_{i=1}^t (\eta_{\tau+i} - 1) \leq -h\delta \right\} \cap \mathcal{H}_2(t)\right), \quad (92)$$

which we do next.

Lemma 3.2. *Suppose that $(mh^2)/n = O(1)$ and $T^2/(hn) \leq \delta/2$. Then, for all large enough n , we have*

$$\mathbb{P}\left(\bigcup_{t < T} \left\{ \sum_{i=1}^t (\eta_{\tau+i} - 1) \leq -h\delta \right\} \cap \mathcal{H}_2(t)\right) \leq e^{-ch} \vee e^{-(ch^2)/T},$$

for some constant $c = c(\gamma, \beta, \delta) > 0$ which depends on γ, β and δ .

Proof. On the event $\mathcal{H}_2(t)$ we can write

$$\eta_{\tau+i} - 1 = \mathbb{1}_{\{Y_{\tau+i-1} < h\}}(\eta_{\tau+i} - 1) =: X_{\tau+i}$$

and hence, given any $r \in (0, 1)$, the probability which appears in the statement of the lemma is at most

$$\mathbb{P}\left(\bigcup_{t < T} \left\{ \sum_{i=1}^t X_{\tau+i} \leq -h\delta \right\}\right) = \mathbb{P}\left(\max_{t < T} e^{-r \sum_{i=1}^t X_{\tau+i}} \geq e^{rh\delta}\right).$$

As before, we need a lower bound on the number of discovered attributes at the end of step $\tau + i - 1$ to control the (conditional) first moment of the $X_{\tau+i}$ given $\mathcal{F}_{\tau+i-1}$. Defining \mathcal{W} as in (88) and $X'_{\tau+i}$ as in (86), we bound from above the expression on the right-hand side of the last inequality by

$$\mathbb{P}\left(\max_{t < T} e^{-r \sum_{i=1}^t X'_{\tau+i}} \geq e^{rh\delta}\right) + \mathbb{P}(\mathcal{W}^c).$$

It follows from (84) that (as $r > 0$) the process $e^{-r \sum_{i=1}^t X'_{\tau+i}}$ is a submartingale (relative to $\mathcal{F}_{\tau+t}$), whence we can use Doob's inequality to conclude that

$$\mathbb{P}\left(\max_{t < T} e^{-r \sum_{i=1}^t X'_{\tau+i}} \geq e^{rh\delta}\right) \leq e^{-rh\delta} \mathbb{E}\left[e^{-r \sum_{i=1}^{T-1} X'_{\tau+i}}\right]. \quad (93)$$

Since $\tau \leq mh^2$ (by definition) and recalling (90), by similar computations as those that led to (91) we obtain, setting $\mathcal{C}_i := \{Y_{\tau+i-1} < h, |\mathcal{D}_{\tau+i-1}| < 2pk(mh^2 + T)\}$,

$$\begin{aligned} \mathbb{E}\left[e^{-r X'_{\tau+i}} | \mathcal{F}_{\tau+i-1}\right] &= e^{r \mathbb{1}_{\mathcal{C}_i}} \mathbb{E}\left[e^{-r \mathbb{1}_{\mathcal{C}_i} \eta_{\tau+i}} | \mathcal{F}_{\tau+i-1}\right] \\ &\leq \exp\left(r \frac{mh^2 + i + h}{n} + r O_{\gamma, \beta}(r(mh^2 + T)/n^2) + O_{\gamma, \beta}(r^2)\right). \end{aligned}$$

Whence we arrive at

$$\mathbb{E}\left[e^{-r \sum_{i=1}^{T-1} X'_{\tau+i}}\right] \leq \exp\left(r \frac{mh^2 + T - 1 + h}{n} + O_{\gamma, \beta}\left(\frac{r(mh^2 + T)}{n^2}\right) + O_{\gamma, \beta}(r^2)\right) \mathbb{E}\left[e^{-r \sum_{i=1}^{T-2} X_{\tau+i}}\right]$$

and iterating we obtain

$$\mathbb{E}\left[e^{-r \sum_{i=1}^{T-1} X'_{\tau+i}}\right] \leq \exp\left(r \frac{mh^2}{n} + r \frac{T^2}{n} + \frac{hT}{n} + O_{\gamma, \beta}(rT(mh^2 + T)/n^2) + O_{\gamma, \beta}(r^2 T)\right).$$

Since $T/n \ll 1$, $(mh^2)/n = O(1)$ and $T^2/(hn) \leq \delta/2$ we see that, for all large enough n , the expression on the right-hand side of (93) is at most

$$c \exp\left\{- (rh\delta)/2 + O_{\beta}(r^2 T)\right\},$$

for some constant $c = c(\gamma, \beta) > 0$. Now if $T/h \leq C$ for some constant $C > 0$, we can find a small enough $r = r(C, \gamma, \beta, \delta) > 0$ (which does not depend on n) such that $O_{\gamma, \beta}((rT)/h) \leq 1/4$ and hence we obtain

$$c \exp\left\{- (rh\delta)/2 + O_{\gamma, \beta}(r^2 T)\right\} \leq ce^{-(rh\delta)/4}.$$

On the other hand, if $T/h \gg 1$, then we can find a large enough constant $C = C(\gamma, \beta)$ such that, taking $r = (h\delta)/(CT) \ll 1$,

$$c \exp\left\{- (rh\delta)/2 + O_{\gamma, \beta}(r^2 T)\right\} \leq c \exp\left\{-c'(h\delta)^2/T\right\}$$

for some other constant $c' = c'(\gamma, \beta) > 0$ (depending on γ, β). All in all, we conclude that there is a large enough constant $c = c(\gamma, \beta, \delta) > 0$ which depends on λ, β and δ such that, for all large enough n ,

$$e^{-r(h-1)} \mathbb{E}\left[e^{-r \sum_{i=1}^{T-1} X_{\tau+i}}\right] \leq e^{-ch} \vee e^{-(ch^2)/T} + \mathbb{P}(\mathcal{W}^c);$$

using (87) the desired conclusion follows. \square

The proof of the proposition is completed combining (79), (80) and the last two lemmas. \square

Next we control the second probability on the right-hand side of (78).

Proposition 3.2. *Suppose that $(mh^3)/n = O(1)$. Then there is $\delta_0 = \delta_0(\gamma, \beta) > 0$ such that, if $0 < \delta \leq \delta_0$, then for all large enough n we have*

$$\mathbb{P}(Y_t < h\delta \ \forall t \in [mh^2]) \leq \exp(-mc_0)$$

for some constant $c_0 = c_0(\delta) > 0$.

Proof. Define $I_i := ((i-1)h^2, ih^2] \cap \mathbb{N}$ for $i \in [m]$, so that $[mh^2] = \cup_{i=1}^m I_i$ (a disjoint union), and let τ^* be the first time $t \in \mathbb{N}$ at which $|\mathcal{D}_t| > 2pkmh^2$. Noticing that $\tau^* > mh^2$ implies $\tau^* > ih^2$ for every $i \in [m]$, we can bound from above the probability that $Y_t < h\delta$ for all $t \in [mh^2]$ by

$$\mathbb{E} \left[\mathbb{1}_{\cap_{i \in [m-1]} \{Y_t < h\delta \ \forall t \in I_i, \tau^* > ih^2\}} \mathbb{P}(Y_t < h\delta \ \forall t \in I_m, \tau^* > mh^2 | \mathcal{F}_{(m-1)h^2}) \right] + \mathbb{P}(\tau^* \leq mh^2). \quad (94)$$

The last probability in (94) can be bounded as in the previous proposition; in particular, it is at most $\exp(-cmh^2)$ for some constant $c = c(\gamma, \beta) > 0$. We claim that the (conditional) probability which appears within the last expectation is at most $4\delta^2$, that is

$$\mathbb{P}(Y_t < h\delta \ \forall t \in I_m, \tau^* > mh^2 | \mathcal{F}_{(m-1)h^2}) \leq 4\delta^2; \quad (95)$$

we establish (95) along the lines of [28]. To this end, define $M_m(t) := Y_{(m-1)h^2+t}$ and $\mathcal{G}_m(t) := \mathcal{F}_{(m-1)h^2+t}$ for $t \in \{0\} \cup [h^2]$; then the probability in (95) equals

$$\mathbb{P}(M_m(t) < h\delta \ \forall t \in [h^2], \tau^* > mh^2 | \mathcal{G}_m(0)).$$

Define τ'_m to be the first time $t \geq 1$ such that $M_m(t) \geq h\delta$ and set $\tau_m := \tau'_m \wedge h^2$. Note that, if $\tau^* > mh^2$, then $|\mathcal{D}_{(m-1)h^2+t}| \leq 2pkmh^2$ for $t \in [h^2]$. Thus, denoting by τ_m^* the first time $t \in [h^2]$ at which $|\mathcal{D}_{(m-1)h^2+t}| > 2pkmh^2$, we see that $\tau^* > mh^2$ implies $\tau_m^* > h^2$. Moreover, if $M_m(t) < h\delta$ for all $t \in [h^2]$ and $\tau^* > mh^2$, then $\tau_m \wedge \tau_m^* = \tau_m = h^2$. Therefore, using (the conditional version of) Markov's inequality, we obtain

$$\begin{aligned} \mathbb{P}(M_m(t) < h\delta \ \forall t \in [h^2], \tau^* > mh^2 | \mathcal{G}_m(0)) &\leq \mathbb{P}(M_m(t) < h\delta \ \forall t \in [h^2], \tau_m^* > h^2 | \mathcal{G}_m(0)) \\ &\leq \mathbb{P}(\tau_m \wedge \tau_m^* = h^2 | \mathcal{G}_m(0)) \\ &\leq \frac{\mathbb{E}[\tau_m \wedge \tau_m^* | \mathcal{G}_m(0)]}{h^2}. \end{aligned} \quad (96)$$

We want to use Theorem 1.3 to bound from above the (conditional) expected value of $\tau_m \wedge \tau_m^*$ given $\mathcal{G}_m(0)$. In particular, if we manage to show that

$$\mathbb{E}[\tau_m \wedge \tau_m^* | \mathcal{G}_m(0)] \leq 4(h\delta)^2, \quad (97)$$

then the ratio in (96) is at most $4\delta^2$ and we are done. Hence we can focus on showing (97). To this end, we first give bounds on the first and second moment of the η_i .

Let $t \leq \tau_m \wedge \tau_m^*$. Recalling (76) and using the tower property together with the bound $(1-x)^\ell \geq 1 - \ell x$ (which is valid for every $x > -1/\ell \in \mathbb{N}$) we immediately obtain

$$\mathbb{E}[\eta_{(m-1)h^2+t}^2 | \mathcal{F}_{(m-1)h^2+t-1}] \leq 2 + \gamma;$$

similarly, $\mathbb{E}[\eta_{(m-1)h^2+t} | \mathcal{F}_{(m-1)h^2+t-1}] \leq 1$. Next, observe that, if $t \leq \tau_m \wedge \tau_m^*$, since $(m-1)h^2 + t - 1 \leq mh^2$

$$\mathbb{E}(\eta_{(m-1)h^2+t} | N_{(m-1)h^2+t}, \mathcal{F}_{(m-1)h^2+t-1}) \geq (n - mh^2 - h\delta)(pN_{(m-1)h^2+t} - (pN_{(m-1)h^2+t})^2/2), \quad (98)$$

where the last inequality follows from the usual bound $e^{-x} \leq 1 - x + x^2/2$ (valid for all $x \geq 0$). Since each N_i is stochastically dominated by the $\text{Bin}(k, p)$ distribution, we obtain (recalling the definition of p, k)

$$p^2 \mathbb{E}[N_{(m-1)h^2+t}^2 | \mathcal{F}_{(m-1)h^2+t-1}] \leq p^2(kp + (kp)^2) = O_{\gamma, \beta}(1/n^2). \quad (99)$$

Moreover, if $t \leq \tau_m \wedge \tau_m^*$ we have

$$p \mathbb{E}[N_{(m-1)h^2+t} | \mathcal{F}_{(m-1)h^2+t-1}] = p^2(k - |\mathcal{D}_{(m-1)h^2+t-1}|) \geq p^2(k - 2pkmh^2). \quad (100)$$

Therefore taking conditional expectation $\mathbb{E}[\cdot|\mathcal{F}_{(m-1)h^2+t-1}]$ on both sides of (98) we obtain, using (99) and (100),

$$\mathbb{E}[\eta_{(m-1)h^2+t}|\mathcal{F}_{(m-1)h^2+t-1}] \geq 1 - \frac{cmh^2}{n} \quad (101)$$

for all large enough n , where $c = c(\gamma, \beta) > 0$ is constant. Furthermore (as $t \leq \tau_m \wedge \tau_m^*$)

$$\mathbb{E}[\eta_t^2|\mathcal{F}_{t-1}] \geq 2 - \frac{cmh^2}{n}; \quad (102)$$

the same bounds in (101) and (102) hold when $Y_{(m-1)h^2+t-1} = 0$. Using above estimates in (101) and (102) we see that, if $0 \leq t \leq \tau_m \wedge \tau_m^*$ and $Y_{(m-1)h^2+t-1} \geq 1$, then

$$\mathbb{E}[M_m^2(t)|\mathcal{G}_m(t-1)] \geq M_m^2(t-1) + 2(\eta_{(m-1)h^2+t} - 1) + (\eta_{(m-1)h^2+t} - 1)^2 \geq M_m^2(t-1) + 1 - \frac{c\delta mh^3}{n}.$$

The same bound is also satisfied when $Y_{(m-1)h^2+t-1} = 0$. By taking a small enough δ we see that the process defined by

$$M_m^2(t \wedge \tau_m \wedge \tau_m^*) - (t \wedge \tau_m \wedge \tau_m^*)/2, \quad t \in \{0\} \cup [h^2]$$

is a submartingale. Applying Theorem 1.3 with the (bounded) stopping times $\sigma_1 = 0$ and $\sigma_2 = \tau_m \wedge \tau_m^*$ we obtain

$$\mathbb{E}[M_m^2(\tau_m \wedge \tau_m^*) - (\tau_m \wedge \tau_m^*)/2 | \mathcal{G}_m(0)] \geq M_m^2(0),$$

from which it follows that

$$\mathbb{E}[\tau_m \wedge \tau_m^* | \mathcal{G}_m(0)] \leq 2\mathbb{E}[M_m^2(\tau_m \wedge \tau_m^*) | \mathcal{G}_m(0)].$$

There remains to bound the expected value on the right-hand side of the last inequality. Using once again the estimates on the moments of the η_i given earlier and the definition of $\mathcal{G}_m(0)$ we obtain

$$\mathbb{E}[M_m^2(\tau_m \wedge \tau_m^*) | \mathcal{G}_m(0)] \leq 2(h\delta)^2$$

for all large enough n . It follows that $\mathbb{E}[\tau_m \wedge \tau_m^* | \mathcal{F}_{(m-1)h^2}] \leq 4(h\delta)^2$, from which we obtain (97). Therefore the probability on the left-hand side of (96) is at most $4\delta^2$, establishing (95). Substituting the bound of (95) into (94) and iterating m times we obtain the desired inequality. \square

Let $m := \lceil A^{3/5} \rceil$ and $h := \lceil n^{1/3}/A^{1/5} \rceil$. Combining (77), (78) together with Propositions 3.1 and 3.2 (and recalling that, by assumption, $A = O(n^{5/12})$) we see that $\mathbb{P}(|\mathcal{C}_{\max}(\mathbb{G}_{n,k,p})| < n^{2/3}/A) \leq \exp(-cA^{3/5})$, as desired.

3.3 The $\mathbb{G}(n, \beta, \lambda)$ model

In this last section we consider the (critical) *quantum* version of the Erdős-Rényi random graph, which we defined non-rigorously in the introductory section.

We refer the reader to [15] and references therein for explanations on such terminology, moving on instead to the precise description of the model, as it is given in [15] (the description given here is taken verbatim from [10]).

Let \mathbb{S}_β denote the circle of length $\beta > 0$ and set $\mathbb{G}_{n,\beta} := [n] \times \mathbb{S}_\beta$, so that to each element $v \in [n]$ we associate the copy $\mathbb{S}_\beta^v = \{v\} \times \mathbb{S}_\beta$ of \mathbb{S}_β . In this way, a point in $\mathbb{G}_{n,\beta}$ has *two* coordinates: its site $v \in [n]$ and the location on the circle $t \in \mathbb{S}_\beta$. Then we create, within each \mathbb{S}_β^v , finitely many holes, according to independent Poisson point processes $(\mathcal{P}_v : v \in [n])$ of intensity $\lambda > 0$. As a result, each punctured circle $\mathbb{S}_\beta^v \setminus \mathcal{P}_v$ can be expressed as the (disjoint) union of $k_v \in \mathbb{N}$ connected intervals, namely

$$\mathbb{S}_\beta^v \setminus \mathcal{P}_v = \bigcup_{j=1}^{k_v} I_j^v, \quad (103)$$

where $|\mathcal{P}_v|$ equals k_v (unless $|\mathcal{P}_v| = 0$, in which case $k_v = 1$ and the circle \mathbb{S}_β^v remains intact).

We add edges in the following (random) manner. To each unordered pair $\{u, v\}$ of (distinct) vertices $u, v \in [n]$, we associate a further circle $\mathbb{S}_\beta^{u,v}$ of length β and a Poisson point process $\mathcal{L}_{u,v} = \mathcal{L}_{v,u}$ on $\mathbb{S}_\beta^{u,v}$ with intensity $1/n$. The processes $\mathcal{L}_{u,v}$ are assumed to be independent for different (u, v) and also independent of $(\mathcal{P}_z : z \in [n])$.

Then, two intervals I_j^u, I_ℓ^v ($u \neq v$) of the decomposition (103) are considered to be directly connected if there exists some $t \in \mathcal{L}_{u,v}$ such that $(u, t) \in I_j^u$ and $(v, t) \in I_\ell^v$. In simple words, two intervals I_j^u and I_ℓ^v (with $u \neq v$) of the decomposition (103) are linked by an edge if they both include a time t at which the Poisson point process $\mathcal{L}_{u,v}$ (on $\mathbb{S}_\beta^{u,v}$) jumps.

The resulting random graph, which is obtained after connecting intervals in the way we have just described, is denoted by $\mathbb{G}_{n,\beta,\lambda}$.

As remarked in [15], setting $\mathcal{P} := \cup_{u \in [n]} \mathcal{P}_u$ (a *finite* collection of points), the decomposition $\mathbb{G}(n, \beta, \lambda) = \mathcal{C}_1 \cup \dots \cup \mathcal{C}_N$ into maximal components is well-defined and, moreover, each point $x = (v, t) \in \mathbb{S}_\beta^v$ is almost surely not in \mathcal{P} . Hence, the notion of component $\mathcal{C}(x)$ of x is also well-defined and throughout the size of a component is the number of *intervals* it contains. The following equivalent description of the $\mathbb{G}_{n,\beta,\lambda}$ model will be convenient for our purposes [15]. Let $\lambda > 0$ and assign to each vertex $v \in [n]$ a circle of length $\theta := \lambda\beta > 0$ with an independent rate one Poisson process \mathcal{P}_v of holes on it. The links between each pair of punctured circles are created by means of iid Poisson processes of intensity $(\lambda n)^{-1}$ (which are also independent of the rate 1 Poisson processes of holes). The resulting random graph, denoted by $\mathbb{G}_{n,\theta}$, has the same law as $\mathbb{G}_{n,\beta,\lambda}$ (see [15]).

As we did for the binomial model and the random graph obtained by bond percolation on a random d -regular graph, in order to establish the upper bound stated in Theorem 1.1 for the $\mathbb{G}_{n,\theta} =_d \mathbb{G}_{n,\beta,\lambda}$ model, the idea is to use an algorithm to sequentially construct an instance of this random graph, interval by interval.

In the following description, which is taken from [15] the vertices (which we recall are circles) have first been labelled with numbers $1, \dots, n$ and, at the end of each time step $t \in \mathbb{N}$ of the algorithm, exactly one interval becomes *explored*.

Algorithm 4. At time $t = 0$, we fix the vertex $w_0 = 1$ and choose a point s_0 uniformly at random on $\mathbb{S}_\theta^{w_0}$. The point (w_0, s_0) is declared *active*, whereas the remaining space is declared *neutral*. Hence, denoting by \mathcal{A}_t the set of active points at the end of step $t \in \mathbb{N}_0$, we have $|\mathcal{A}_0| = 1$. For every $t \in \mathbb{N}$, the algorithm works as follows.

- (a) If $|\mathcal{A}_{t-1}| \geq 1$, we choose an active point (w_t, s_t) whose vertex has the smallest index among all active points. In case of a tie (which may occur since each \mathbb{S}_θ^v could contain several active points), we choose the active point which chronologically appeared earlier than the others on the same vertex.
- (b) If $|\mathcal{A}_{t-1}| = 0$ and there exists at least one neutral circle, we choose w_t to be the neutral vertex with the smallest index and select s_t uniformly at random on $\mathbb{S}_\theta^{w_t}$. Then we declare the point (w_t, s_t) active.
- (c) If $|\mathcal{A}_{t-1}| = 0$ and there is no neutral circle left, we choose w_t to be the vertex of smallest index among the vertices having some neutral part and select s_t uniformly at random on the neutral part of $\mathbb{S}_\theta^{w_t}$. Then we declare the point (w_t, s_t) active.
- (d) If $|\mathcal{A}_{t-1}| = 0$ and there is no neutral part available on any circle, then we terminate the procedure.

The algorithm proceeds as follows. Using iid $\text{Exp}(1)$ random variables J_-^t, J_+^t and writing s_t^1, s_t^2 for the points on the neutral space of $\mathbb{S}_\theta^{w_t}$ around s_t , we extract out of the *maximal* neutral interval $\{w_t\} \times (s_t^1, s_t^2)$ around the (active) point (w_t, s_t) the sub-interval $I_t := \{w_t\} \times \tilde{I}_t$, where

$$\tilde{I}_t := (s_t^1 \vee (s_t - J_-^t), s_t^2 \wedge (s_t + J_+^t)).$$

If $\mathbb{S}_\theta^{w_t}$ is neutral (apart from active points), we take $s_t^1 = -\infty = -s_t^2$ (so that $\tilde{I}_t = (s_t - J_-^t, s_t + J_+^t)$) and $\tilde{I}_t = \mathbb{S}_\theta$ (the whole circle of length θ) if $J_-^t + J_+^t \geq \theta$. We then remove from the list of active points \mathcal{A}_{t-1} *all* those points which got encompassed by the interval I_t , including the active point currently under investigation (w_t, s_t) . The links in the graph connected to all such points, other than (w_t, s_t) , are considered to be *surplus edges*.

The connections of I_t are then constructed in the following manner.

- (i) For $i \neq w_t$, we regard \tilde{I}_t as a subset of $\mathbb{S}_\theta^{w_t, i}$ and sample the process of links $\mathcal{L}_{w_t, i}$ for times r restricted to \tilde{I}_t . That is, we run the process $(\mathcal{L}_{w_t, i}(r))_{r \in \tilde{I}_t}$ and, denoting by $j_1^i, \dots, j_{L_i}^i \in \tilde{I}_t$ the jumps of this process (if any), then we create a link between each pair of points $(w_t, j_\ell^i), (i, j_\ell^i)$, $\ell \in [L_i]$.

- (ii) Subsequently, we erase all links between I_t and points on already explored intervals, and record each (i, j_ℓ^i) ($\ell \in [L_i], i \neq w_t$) on the neutral space as an active point, labelled with the time (order) of its registration (see part (a) of the algorithm, where such ordering is used).

After examining all the connections from I_t , we declare such interval *explored* and increase t by one. (We notice that the procedure halts after finitely many steps, because the number of intervals is finite.)

Denote by ζ_t the *new* links that are formed during step t in **Algorithm 4**. Moreover, we let Spl_t denote the number of surplus edges found by the end of the first t steps in the procedure. Define

$$\eta_t := \zeta_t - (\text{Spl}_t - \text{Spl}_{t-1}) \quad (104)$$

and note that, since at those times t where $|\mathcal{A}_{t-1}| \geq 1$ we add to the list of active points the ζ_t new links found during step t but we remove those (active) points which are encompassed by I_t (including (w_t, s_t)), we have the recursion

- $|\mathcal{A}_t| = |\mathcal{A}_{t-1}| + \eta_t - 1$, if $|\mathcal{A}_{t-1}| \geq 1$;
- $|\mathcal{A}_t| = \eta_t$, if $|\mathcal{A}_{t-1}| = 0$.

To bound (from above) the probability that $\mathcal{C}_{\max}(\mathbb{G}(n, \beta, \lambda))$ contains less than $n^{2/3}/A$ nodes, we follow the strategy adopted in [15]. Specifically, we consider a more *restrictive* exploration process, which leads to shorter (positive) excursions compared to those of $|\mathcal{A}_t|$. The advantage of such a new procedure is that it yields a simpler (conditional) distribution for the random variable η_i .

Reduced exploration process. This procedure, after creating the first active point on each node $u \in [n]$, it *voids* all space on that vertex apart from the relevant interval around that point, thus sequentially producing components with no more intervals than does the original exploration process. Specifically, at each step $t \in \mathbb{N}$ of this procedure, either I_t is originated from a point that was active at the end of step $t-1$ or, if $|\mathcal{A}_{t-1}| = 0$, it is formed from a point (w_t, s_t) with s_t selected uniformly at random on the completely neutral circle $\mathbb{S}_\theta^{w_t}$, if such a circle exists; otherwise, we halt the procedure. Moreover, this new (restrictive) exploration process keeps *at most* one link from I_t to any (as of yet) never visited (in particular, neutral) circle \mathbb{S}_θ^i , erasing all other connections which are being formed during step t by the original exploration process, namely **Algorithm 4**.

Remark 9. Since in the above procedure we void all space on a node after creating the first active point on it, we see that the algorithm runs for n steps.

It is important to notice that such restrictive exploration process *does not* have surplus links and, moreover, its number of active points $|\mathcal{A}_t^*|$ also satisfies $|\mathcal{A}_0^*| = 1$ and a recursion of the type:

- $|\mathcal{A}_t^*| = |\mathcal{A}_{t-1}^*| + \eta_t^* - 1$, if $|\mathcal{A}_{t-1}^*| \geq 1$;
- $|\mathcal{A}_t^*| = \eta_t^*$, if $|\mathcal{A}_{t-1}^*| = 0$,

with η_t^* representing the number of new active points created at time t .

Let us denote by Y_t and U_t the number of active points and neutral vertices at the end of step $t \in [n] \cup \{0\}$ in the reduced exploration process, respectively, so that

$$Y_t = |\mathcal{A}_t^*| \text{ and } U_t = n - t - Y_t.$$

Denote by \mathcal{F}_t the σ -algebra generated by the information collected by the exploration process until the end of step t . Then, conditional on \mathcal{F}_{t-1} and J_t , the random variable η_t^* depends on the past until time $t-1$ only through Y_{t-1} and

$$\eta_t^* =_d \text{Bin}(U_{t-1} - \mathbb{1}_{\{Y_{t-1}=0\}}, 1 - e^{-J_t/(\lambda n)}), \quad (105)$$

where we recall that the J_i are iid $\Gamma_\theta(2, 1)$ -distributed random variables.

As before, the component sizes are given by $t_i^* - t_{i-1}^*$, where we set $t_0^* := 0$ and let $t_i^* := \min\{t \geq t_{i-1}^* + 1 : Y_t = 0\}$ for $i \geq 1$ (as long as there is at least one neutral circle).

Using the reduced exploration process we can bound

$$\mathbb{P}(t_i - t_{i-1} < n^{2/3}/A \ \forall i) \leq \mathbb{P}(t_i^* - t_{i-1}^* < n^{2/3}/A \ \forall i). \quad (106)$$

Then, arguing as we did for the random graphs $\mathbb{G}_{n,p}$ and $\mathbb{G}_{n,d,p}$, we take $1 \ll h = h(n) \in \mathbb{N}$, $m \in \mathbb{N}$ (possibly dependent on n), and denote by $\delta \in (0, 1)$ a constant that we specify later. Then, setting $T := \lceil n^{2/3}/A \rceil$, we bound from above the probability on the right-hand side of (106) by

$$\mathbb{P}(t_i^* - t_{i-1}^* < T \ \forall i, \exists t \in [mh^2] : Y_t \geq h\delta) + \mathbb{P}(Y_t < h\delta \ \forall t \in [mh^2]). \quad (107)$$

The two terms which appear in the last display are bounded (from above) by means of the following two propositions.

Proposition 3.3. *Suppose that $(mh^2)/n = O(1)$ and $T^2/(hn) \leq \delta/2$. Then, for all large enough n , we have*

$$\mathbb{P}(t_i^* - t_{i-1}^* < T \ \forall i, \exists t \in [mh^2] : Y_t \geq h\delta) \leq e^{-ch} \vee e^{-(ch^2)/T},$$

for some constant $c = c(\lambda, \beta, \delta) > 0$ which depends on λ, β and δ .

Proof. The proof follows the exact same steps carried out in the proof of Proposition 3.1 and therefore we omit the details; we just compute the relevant expectations needed to adapt the argument.

Recall from (105) that, conditional on $\mathcal{F}_{\tau+i-1}$, the random variable $\eta_{\tau+i}^*$ is distributed as

$$\text{Bin}(n - \tau - (i - 1) - Y_{\tau+i-1} - \mathbb{1}_{\{Y_{\tau+i-1}=0\}}, 1 - e^{-J_{\tau+i}/(\lambda n)}). \quad (108)$$

Therefore, using the classical bounds $e^x \geq 1 + x$ (which is valid for all $x \in \mathbb{R}$) we see that

$$1 - e^{-J_{\tau+i}/(\lambda n)} \leq \frac{J_{\tau+i}}{\lambda n}$$

and so we obtain

$$E_{\tau+i} \leq n\mathbb{E}[J_{\tau+i}]/(\lambda n) - 1 \leq 0. \quad (109)$$

Using (108) together with the bound $e^{-x} \leq 1 - x + x^2/2$ (which is valid for all $x \geq 0$) we obtain

$$\mathbb{E}[\eta_{\tau+i}^* | \mathcal{F}_{\tau+i-1}] \geq 1 - \frac{\tau + i - 1 + Y_{\tau+i-1} + \mathbb{1}_{\{Y_{\tau+i-1}=0\}}}{n} - O_{\lambda, \beta}(n^{-1}), \quad (110)$$

whence

$$E_{\tau+i} \geq -\frac{\tau + i - 1 + Y_{\tau+i-1} + \mathbb{1}_{\{Y_{\tau+i-1}=0\}}}{n} - O_{\lambda, \beta}(n^{-1}).$$

Moreover, a repeated application of the inequality $e^x \leq 1 + x + x^2$ (which is valid for all $x \in [0, 1]$) together with the fact that $e^x \geq 1 + x$ for all $x \in \mathbb{R}$ yields

$$\mathbb{E}[e^{rX_{\tau+i}} | \mathcal{F}_{\tau+i-1}] \leq \exp\left(-r \frac{\tau + i - 1 + Y_{\tau+i-1} + \mathbb{1}_{\{Y_{\tau+i-1}=0\}}}{n} + O_{\lambda, \beta}(r^2)\right). \quad (111)$$

Thus we obtain

$$\mathbb{E}[e^{r(X_{\tau+i} - E_{\tau+i})} | \mathcal{F}_{\tau+i-1}] = e^{-rE_{\tau+i}} \mathbb{E}[e^{rX_{\tau+i}} | \mathcal{F}_{\tau+i-1}] \leq e^{O_{\lambda, \beta}(r^2) + O_{\lambda, \beta}(1/n)}.$$

Since $\tau \leq mh^2$ (by definition) and recalling (110), by similar computations as the ones that led to (111) we obtain

$$\mathbb{E}[e^{-rX_{\tau+i}} | \mathcal{F}_{\tau+i-1}] \leq \exp\left(r \frac{mh^2 + i + h}{n} + O_{\lambda, \beta}(r/n) + O_{\lambda, \beta}(r^2)\right).$$

The result follows arguing as in Proposition 3.1. \square

Proposition 3.4. *Suppose that $(mh^3)/n = O(1)$. Then there is $\delta_0 = \delta_0(\lambda, \beta) > 0$ such that, if $0 < \delta \leq \delta_0$, then for all large enough n we have*

$$\mathbb{P}(Y_t < h\delta \ \forall t \in [mh^2]) \leq \exp(-mc_0)$$

for some constant $c_0 = c_0(\delta) > 0$.

Proof. The proof follows the same steps carried out in the proof of Proposition 3.2 and hence we omit most of the details.

First of all note that, recalling (105) and using the tower property together with the bound $e^x \geq 1+x$ (which is valid for every x) we immediately obtain

$$\begin{aligned} \mathbb{E}[(\eta_t^*)^2 | \mathcal{F}_{t-1}] &= \mathbb{E}_{\mathcal{F}_{t-1}}[\mathbb{E}((\eta_t^*)^2 | \mathcal{F}_{t-1}, J_t)] \leq \mathbb{E}_{\mathcal{F}_{t-1}}[\mathbb{E}(\text{Bin}^2(n, 1 - e^{-J_t/(\lambda n)}) | J_t)] \\ &\leq \mathbb{E}_{\mathcal{F}_{t-1}}[n(1 - e^{-J_t/(\lambda n)})] + \mathbb{E}_{\mathcal{F}_{t-1}}[(n(1 - e^{-J_t/(\lambda n)}))^2] \\ &\leq C \end{aligned}$$

where $C = C(\lambda, \beta) := 1 + \lambda^{-2} \mathbb{E}[J_1^2]$ depends on λ, β ; similarly, $\mathbb{E}[\eta_t^* | \mathcal{F}_{t-1}] \leq 1$ (at all times t). Next, suppose that $1 \leq Y_{t-1} < h\delta$. Then we can bound, using (110),

$$\mathbb{E}[\eta_t^* | Y_{t-1}] \geq 1 - \frac{2mh^2}{n} - O_{\lambda, \beta}(1/n), \quad (112)$$

for all large enough n . Moreover, using the inequalities $e^x \geq 1+x$ and $e^{-x} \leq 1-x+x^2/2$ (which are valid for all $x \in \mathbb{R}$ and $x \geq 0$, respectively) we see that (as $1 \leq Y_{t-1} < h\delta$ and $t \leq mh^2$)

$$\begin{aligned} \mathbb{E}[(\eta_t^*)^2 | \mathcal{F}_{t-1}] &= \mathbb{E}_{\mathcal{F}_{t-1}}[\mathbb{E}[\text{Bin}^2(n - (t-1) - Y_{t-1}, 1 - e^{-J_t/(\lambda n)}) | J_t]] \\ &\geq (n - mh^2 - h\delta) \left(\frac{1}{n} - O_{\lambda, \beta}(1/n^2) \right) + (n - mh^2 - h\delta)^2 \mathbb{E}[(1 - e^{-J_t/(\lambda n)})^2]. \end{aligned} \quad (113)$$

Now since

$$\mathbb{E}[(1 - e^{-J_t/(\lambda n)})^2] \geq \mathbb{E}[1 - e^{-J_t/(\lambda n)}]^2 \geq n^{-2} \left(1 - \frac{\mathbb{E}[J_t^2]}{2\lambda^2 n} \right)^2 \geq n^{-2} (1 - O_{\lambda, \beta}(1/n)),$$

we see that the last term on the right-hand side of (113) is at least

$$1 - \frac{mh^2 + h\delta}{n} - O_{\lambda, \beta}(1/n),$$

provided n is sufficiently large. Since the second-last term on the right-hand side of (113) satisfies

$$(n - mh^2 - h\delta) \left(\frac{1}{n} - O_{\lambda, \beta}(1/n) \right) \geq 1 - \frac{mh^2 + h\delta}{n} - O_{\lambda, \beta}(1/n)$$

and $h\delta \leq mh^2$, we conclude that

$$\mathbb{E}[(\eta_t^*)^2 | Y_{t-1}] \geq 2 - 4\frac{mh^2}{n} - O_{\lambda, \beta}(1/n) \geq 2 - \frac{cmh^2}{n} \quad (114)$$

for some constant $c = c(\lambda, \beta) > 0$; the same bounds in (112) and (114) hold when $Y_{t-1} = 0$. The desired result then follows arguing as in Proposition 3.2 \square

The required upper bound on the probability that $|\mathcal{C}_{\max}(\mathbb{G}(n, \beta, \lambda))| < n^{2/3}/A$ can be obtained combining Propositions 3.3 and 3.4, using the same values for h, m employed when analysing the $\mathbb{G}(n, k, p)$ model.

Acknowledgements

The author thanks Matthew Roberts for a useful discussion on this problem when the former was a PhD student at the University of Bath.

References

- [1] L. Addario-Berry and B.A. Reed. Ballot theorems, old and new. In *Horizons of Combinatorics*, pages 9–35. Springer, 2008.

- [2] M. Behrisch. Component evolution in random intersection graphs. In *The Electronical Journal of Combinatorics*, 14(1), 2007.
- [3] N. Alon, I. Benjamini and A. Stacey. Percolation on finite graphs and isoperimetric inequalities. In *The Annals of Probability*, 32: 1727–1745, 2004.
- [4] S. Bhamidi, R. van der Hofstad and J.S.H. van Leeuwaarden. Novel scaling limits for critical inhomogeneous random graphs. *The Annals of Probability*, 40(6):2299–2361, 2012.
- [5] S. Bhamidi, R. van der Hofstad, and J.S.H. van Leeuwaarden. Scaling limits for critical inhomogeneous random graphs with finite third moments. *Electronic Journal of Probability*, 15:1682–1702, 2010.
- [6] B. Bollobás. A probabilistic proof of an asymptotic formula for the number of labelled regular graphs. *European Journal of Combinatorics*, 1(4):311–316, 1980.
- [7] B. Bollobás. *Random graphs*, volume 73 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 2001.
- [8] S. Chatterjee. A new approach to strong embeddings. *Probability Theory and Related Fields*, 152(1-2):231–264, 2012.
- [9] U. De Ambroggio. An elementary approach to component sizes in critical random graphs. *Journal of Applied Probability*, 59(4):1228–1242, 2022.
- [10] U. De Ambroggio. A simple path to component sizes in critical random graphs. *SIAM Journal of Discrete Mathematics*, 38 (2):1492–1525, 2024.
- [11] U. De Ambroggio and A. Pachon. Upper bounds for the largest components in critical inhomogeneous random graphs. *ALEA, Lat. Am. J. Probab. Math. Stat.*, 20:1315–1358, 2023.
- [12] U. De Ambroggio and M.I. Roberts. Unusually large components in near-critical Erdős-Rényi graphs via ballot theorems. *Combinatorics, Probability & Computing*, 31(5):840–869, 2022.
- [13] U. De Ambroggio and M.I. Roberts. The probability of unusually large components for critical percolation on random d -regular graphs. *Electronic Journal of Probability*, 28:1-55, 2023.
- [14] M. Deijfen and M. Kets. Random intersection graphs with tunable degree distribution and clustering. *Probability in the Engineering and Informational Sciences*, 23(4):661–674, 2009.
- [15] A. Dembo, A. Levit and S. Vadlamani. Component sizes for large quantum Erdős-Rényi graph near criticality. *The Annals of Probability*, 47(2):1185–1219, 2019.
- [16] S. Dhara. Critical Percolation on Random Networks with Prescribed Degrees, PhD Thesis (2018).
- [17] S. Dhara, R. van der Hofstad, J.S.H. van Leeuwaarden and S. Sen. Critical window for the configuration model: finite third moment degrees. *Electronic Journal of Probability*, 22(16):1–33, 2017.
- [18] A. Goerdts. The giant component threshold for random regular graphs with edge faults. *Theoretical Computer Science*, 259:307–321, 2001.
- [19] H. Hatami and M. Molloy. The scaling window for a random graph with a given degree sequence. *Random Structures & Algorithms*, 41: 99-123, 2012.
- [20] R van der Hofstad. *Random graphs and complex networks*, volume 1. Cambridge University Press, 2016.
- [21] R. van der Hofstad. Critical behavior in inhomogeneous random graphs. *Random Struct. Alg.*, 42(4): 480-508 (2013).
- [22] R. van der Hofstad, S. Kliem and J.S.H. van Leeuwaarden. Cluster tails for critical power-law inhomogeneous random graphs. *Journal of Statistical Physics*, 171(1):38–95, 2018.

- [23] D. Ioffe and A. Levit. Long range order and giant components of quantum random graphs. *Markov Processes and Related Fields*, 13(3):469–492, 2018.
- [24] S. Janson, T. Luczak and A. Rucinski. *Random graphs*, volume 45. John Wiley & Sons, 2011.
- [25] F. Joos and G. Perarnau. Critical percolation on random regular graphs. *Proc. Amer. Math. Soc.*, 146:3321–3332, 2018.
- [26] A.N. Lageras and M. Lindholm A note on the component structure in random intersection graphs with tunable clustering. *The Electronic Journal of Combinatorics*, 15(1), 2008.
- [27] A. Nachmias and Y. Peres. Critical percolation on random regular graphs. *Random Structures & Algorithms*, 36(2):111–148, 2010.
- [28] A. Nachmias and Y. Peres. The critical random graph, with martingales. *Israel Journal of Mathematics*, 176:29–41, 2010.
- [29] I. Norros and H. Reittu. On a conditionally Poissonian graph process. *Advances in Applied Probability*, 38(1):59-75 (2006).
- [30] B. Pittel. On the largest component of the random graph at a nearcritical stage. *Journal of Combinatorial Theory, Series B*, 82(2):237–269, 2001.
- [31] M.I. Roberts. The probability of unusually large components in the near-critical Erdős-Rényi graph. *Advances in Applied Probability*, 50(1):245–271, 2017.
- [32] M.I. Roberts and B. Şengül. Exceptional times of the critical dynamical Erdős-Rényi random graph. *The Annals of Applied Probability*, 28(4):2275–2308, 2018.
- [33] D. Stark. The vertex degree distribution of random intersection graphs. *Random Structures & Algorithms*, 24(3):249–258., 2004.