

On the structure of the complement $\overline{\text{fib}}$ of the set fib of fibbinary numbers in the set of positive natural numbers

A.J. Macfarlane ¹

Centre for Mathematical Sciences, D.A.M.T.P.
Wilberforce Road, Cambridge CB3 0WA, UK

Abstract

The set fib of fibbinary numbers is defined via a bijection between the set \mathbb{N} of natural numbers and fib. Since the elements of the set fib do not exhaust \mathbb{N} , the structure of the complement $\overline{\text{fib}}$ of fib in \mathbb{N} , is of interest. An explicit expression $\overline{\text{fib}} = \bigcup_{k \geq 1} \Phi_k$ is obtained in terms of certain well-defined sets Φ_k , $k \geq 1$. The key to its proof lies in first considering the odd numbers involved in this statement: a general treatment, with full justification, of the binary representations of the odd numbers is developed, and exploited in showing the expression quoted for $\overline{\text{fib}}$ to be correct. The main results of the article can also be viewed as providing partitions of the set of natural numbers, and also of its subset of odd numbers, that follow from the introduction of the set fib, and of its subset of odd integers.

Keywords: Fibonacci number, Zeckendorf representation, fibbinary number, complement, binary representation

Mathematical Subject Classification: 11A63, 11A67

1 Introduction

A recent paper [1], describes the bijection \mathcal{Z} between the set \mathbb{N} of positive natural numbers and the set fib of fibbinary numbers, see [1, 2], and [A003714](#) in [3], which underlies the definition of the positive integers $\text{fib}(n)$, $n \geq 1$. The bijection [1, Eqs. (1),(29)] in question

$$\begin{aligned} \mathcal{Z} : \mathbb{N} &\rightarrow \text{fib} \\ n &\mapsto \text{fib}(n), \end{aligned} \quad (1)$$

depends on the use of Zeckendorf's theorem [4–6], [7, p. 125] and the Zeckendorf representation $z(n)$ of $n \in \mathbb{N}$. The paper [1] contains more information and detail on these matters than section two here provides.

The elements of the set fib do not exhaust \mathbb{N} , and the purpose of this paper is to describe the complement $\overline{\text{fib}}$ in \mathbb{N} of fib

$$\overline{\text{fib}} = \mathbb{N} - \text{fib}. \quad (2)$$

In particular a set theoretic expression for $\overline{\text{fib}}$ is offered and proved to be correct. In terms of sets Φ_k , $k \geq 1$, defined in section three, this formula reads as

$$\overline{\text{fib}} = \bigcup_{k=1}^{\infty} \Phi_k. \quad (3)$$

To approach (3), introduce the set odfib , [A022341](#) in [3], [1, 2], which is the subset of fib which contains all of its odd elements, and the sets Ψ_k , $k \geq 1$, consisting of the all of the odd elements of Φ_k . The major part of the effort in completing the proof of (3) resides in proving correct the expression

$$\overline{\text{odfib}} = \bigcup_{k=1}^{\infty} \Psi_k. \quad (4)$$

¹*e-mail:* a.j.macfarlane@damtp.cam.ac.uk

for the complement $\overline{\text{odfib}}$

$$\overline{\text{odfib}} = \mathbb{N}_{\text{odd}} - \text{odfib}. \quad (5)$$

of fib in the set of positive odd numbers \mathbb{N}_{odd} , [A005408](#) in [3]. After this proof is completed, that of (2) follows easily.

The results (2) and (3) can also be recast in the form

$$\mathbb{N} = \text{fib} + \bigcup_{k=1}^{\infty} \Phi_k, \quad (6)$$

that is, as a partition of the natural numbers. Similarly (4) and (5) provide a partition of the odd numbers.

Section two summarises well-known material about the sets fib and odfib and the bijection \mathcal{Z} , as needed for later sections. Section three defines the sets Φ_k , $k \geq 0$, adopting the convenient notation $\text{fib} = \Phi_0$, and introduces tables that give valuable insight into the structure of these sets. The elements of the sets Ψ_k , $k \geq 0$, with $\Psi_0 = \text{odfib}$, can be read off the results quoted for the sets Φ_k or off the associated tables. Tables 1 and 2 relate to fib and Φ_1 . The latter is typical of the tables that can be drawn up for Φ_k for any $k \geq 1$. Explicit expressions are given for $\Phi_k(n)$, $\Psi_k(n)$, $k \geq 1$, in terms of $\Phi_0(n) = \text{fib}(n)$, $\Psi_0(n) = \text{odfib}(n)$.

In section four, the first odd number array, ONA1 is presented as Table 3. Its first or $k = 0$ column contains the integers of odfib. Its k -th column, $k \geq 1$, contains the integers of Ψ_k . It will be seen that all odd integers are present each one exactly once in the array ONA1. To prove this the vital step is to construct the second odd number array, ONA2. This is obtained from the array ONA1 by replacing each of its elements by its unique binary representation, BR, written in an appropriate way. Proof of (4), given in detail, exploits the information so presented explicitly in ONA2.

The display in ONA2 of the BRs of its odd number entries is of significance in its own right. It can be seen to provide a BR interpretation of the formula (38) for $\Psi_k(n)$, and hence also for the formula (37) for $\Phi_k(n)$. See section five.

2 The Zeckendorf representation and the fibbinary numbers

2.1 The Zeckendorf representation

Let F denote the set of Fibonacci numbers

$$\begin{aligned} F &= \{F_n, \quad n \geq 2\} \\ &= \{1, 2, 3, 5, 8, 13, 21, 34, 55, \dots\}. \end{aligned} \quad (7)$$

Zeckendorf's theorem [4–7] states that every positive integer can be expressed uniquely as a sum of non-consecutive Fibonacci numbers $F_n \in F$. This leads to the Zeckendorf representation – notation $z(n)$ and abbreviation ZR – of the integers $n \in \mathbb{N}$, $n \geq 1$. This can be written uniquely in the form

$$z(n) = \sum_{i=2}^r a_i F_i, \quad (8)$$

where

$$\begin{aligned} a_i &\in \{0, 1\}, \quad a_r = 1, \\ a_i a_{i+1} &= 0, \quad 2 \leq i < r, \end{aligned} \quad (9)$$

and r is the largest integer such that $F_r \leq n$. The entry a_2 is referred to as the least significant bit, LSB, and (8) as the Zeckendorf condition, ZC.

For $z(n)$, given by (8), write also

$$z(n) = (a_r, a_{r-1}, \dots, a_2)_F = (n)_F, \quad (10)$$

where, within $(n)_F$, n denotes the string

$$a_r a_{r-1} \dots a_2, \quad (11)$$

of ones and zeros obtained from (2). Define also the length $l(n)$ of the ZR of n , that is, the total number of ones and zeros in the string (11).

If the integers $n \in \mathbb{N}$ are listed in ZR form, it is seen that they separate into Fibonacci subsets \mathcal{F}_k of cardinality F_k such that $l(n) = k$ for all $n \in \mathcal{F}_k$.

2.2 The fibbinary numbers

The definition of $z(n)$ of (8) motivates introduction of the set

$$\text{fib} = \{\text{fib}(n), \quad n = 1, 2, \dots\}, \quad (12)$$

[A003714](#) in [3], [1, 2], of fibbinary numbers, wherein the positive integer $\text{fib}(n)$ is defined by

$$\text{fib}(n) = \sum_{i=2}^r a_i 2^{i-2}, \quad (13)$$

and use is made of the same set of coefficients a_i , $2 \leq i \leq r$, as are used in (8). Eq. (13) is the binary representation, BR, of $\text{fib}(n)$. As for the integers $n \in \mathbb{N}$ in ZR form (8), the $\text{fib}(n)$ separate into Fibonacci subsets \mathcal{F}_k of cardinality F_k with $l(\text{fib}(n)) = l(n)$. Accordingly the listing of the elements of fib is presented as

$$\begin{aligned} \text{fib} = & \{\text{fib}(n), \quad n = 1, 2, \dots\} \\ = & \{1; 2; 4, 5; 8, 9, 10; 16, 17, 18, 20, 21 = \text{fib}(12); 32, 33, 34, 36, 37, 40, 41, 42 = \text{fib}(20); \\ & 64, 65, 66, 68, 69, 72, 73, 74, 80, 81, 82, 84, 85 = \text{fib}(33); \\ & 128, \dots, 170 = \text{fib}(54); 256, \dots, 341 = \text{fib}(88); \dots\}, \end{aligned} \quad (14)$$

using semi-colons to indicate separation of the entries into Fibonacci subsets. The largest element of the \mathcal{F}_k , $k = 1, 2, \dots$, subset of fib is $\text{fib}(F_{k+2} - 1)$, as noted explicitly in (14) for $k = 5, \dots, 9$.

Writing (13) as

$$\text{fib}(n) = (a_r, a_{r-1} \dots a_2)_2 = (n)_2, \quad (15)$$

in analogy with (11), see the same string (10) used for n in $(n)_F$ and $(n)_2$ but with distinct meanings. Respect of the Zeckendorf condition is a strict requirement in both uses.

It is clear that the map (1) from \mathbb{N} to fib defines a bijection which may be expressed now more explicitly as

$$\begin{aligned} \mathcal{Z} : \mathbb{N} & \rightarrow \text{fib} \\ n = (n)_F & \mapsto \text{fib}(n) = (n)_2. \end{aligned} \quad (16)$$

2.3 The set fib and its subset odfib

Define next the important subset odfib of fib of odd fibbinary numbers, [A022341](#) in [3], [1, 2]. This subset contains all the odd elements of fib each one exactly once:

$$\begin{aligned} \text{odfib} = & \{\text{odfib}(n), \quad n = 0, 1, \dots\} \\ = & \{1, 5; 9; 17, 21; 33, 37, 41; 65, 69, 73, 81, 85; \\ & 129, 133, 137, 145, 149, 161, 165, 169; \\ & 257, \dots, 341; 513, \dots\}. \end{aligned} \quad (17)$$

The even fibbinary numbers of evfib , [A022342](#) in [3], are not used here. The elements of the subsets (17) are given in terms of those of fib by

$$\begin{aligned} \text{odfib}(n) & = 4\text{fib}(n) + 1 = (n01)_2, \quad n = 0, 1, \dots, \\ \text{evfib}(n) & = 2\text{fib}(n) = (n0)_2, \quad n = 1, 2, \dots, \end{aligned} \quad (18)$$

where $\text{fib}(0) = 0$ is used so that (18) reads correctly for $n = 0$. Note also that (18) gives

$$l(\text{odfib}(n)) = l(\text{fib}(n)) + 2 = l(n) + 2. \quad (19)$$

3 The sets Φ_k , $k \geq 1$, and their subsets Ψ_k , $k \geq 1$.

The definition of all sets Φ_k , $k \geq 0$ of interest here depends upon

- a) their initial element α_k , and
- b) the following key property:

If $j \in \Phi_k$, $k \geq 0$, then so also does the even integer $2j$, and the odd integer $4j + 1$, but $4j - 1 \notin \Phi_k$.

Given α_k , the elements of Φ_k can be obtained by applying the key property. For the set $\Phi_0 = \text{fib}$, with $\Phi_0(1) = \alpha_0 = 1$, (14) gives a listing of its elements.

To construct the sets Φ_k , $k \geq 1$ define their initial elements

$$\alpha_k = 4k - 1, \quad (20)$$

odd integers not present in fib , and apply property b). This gives rise to the following:

$$\begin{aligned} \Phi_k &= \{\Phi_k(n), k \geq 1, n \geq 1, \Phi_k(1) = \alpha_k.\}, \\ \Phi_1 &= \{3; 6; 12, 13; 24, 25, 26; 48, 49, 50, 52, 53 = \Phi_3(12); \\ &\quad 96, 97, 98, 100, 101, 104, 105, 106 = \Phi_3(20); \\ &\quad 192, 193, 194, 196, 197, 200, 201, 202, 208, 209, 210, 212, 213 = \Phi_3(33); \dots \}, \end{aligned} \quad (21)$$

$$\begin{aligned} \Phi_2 &= \{7; 14; 28, 29; 56, 57, 58; 112, 113, 114, 116, 117; \\ &\quad 224, 225, 226, 228, 229, 232, 233, 234; 448, \dots, 469; \dots \}, \end{aligned} \quad (22)$$

$$\begin{aligned} \Phi_3 &= \{11; 22; 44, 45; 88, 89, 90; 176, 177, 178, 180, 181; \\ &\quad 352, 353, 354, 356, 357, 360, 361, 362; 706, \dots, 725; \dots \}, \end{aligned} \quad (23)$$

$$\begin{aligned} \Phi_4 &= \{15; 30; 60, 61; 120, 121, 122; 240, 241, 242, 244, 245; \\ &\quad 480, 481, 482, 484, 485, 488, 489, 490; 960, \dots, 981; \dots \}, \end{aligned} \quad (24)$$

$$\Phi_5 = \{19; 38; 76, 77; 152, 153, 154; 304, 305, 306, 308, 309; 608, \dots, 618; 1216, \dots, 1237; \dots \}, \quad (25)$$

$$\Phi_6 = \{23; 46, 47; 92, 93; 184, 185, 186; 368, \dots, 373; 736, \dots, 746; 1472, \dots, 1493; \dots \}. \quad (26)$$

The sets Ψ_k , $k \geq 0$ are defined to be the sets of all odd elements of Φ_k , $k \geq 0$. For $k = 0$, $\Psi_0 = \text{odfib}$, with elements shown in (17). The elements of Ψ_k , $k \geq 1$, may be read off (21 – 26).

$$\begin{aligned} \Psi_k &= \{\Psi_k(n), k \geq 1, n \geq 0, \\ \Psi_1 &= \{3; 13; 25; 49, 53; 97, 101, 105; 193, 197, 201, 209, 213; \dots \}, \end{aligned} \quad (27)$$

$$\Psi_2 = \{7; 29; 57; 113, 117; 225, 229, 233; 449, 453, 457, 465, 469; \dots \}, \quad (28)$$

$$\Psi_3 = \{11; 45; 89; 177, 181; 353, 357, 361; 705, 709, 713, 721, 725; \dots \}, \quad (29)$$

$$\Psi_4 = \{15; 61; 121; 241, 245; 481, 485, 489; 961, 965, 969, 979, 981; \dots \}, \quad (30)$$

$$\Psi_5 = \{19; 77; 153; 305, 309; 605, 609, 613; 1217, 1221, 1225, 1233, 1237; \dots \}, \quad (31)$$

$$\Psi_6 = \{23; 93; 185; 369, 373; 737, 741, 747; 1473, 1477, 1481, 1489, 1493; \dots \}. \quad (32)$$

As in [1], the structures of sets such as the sets Φ_k , $k \geq 0$, are seen clearly by presenting their elements in table form. The rules governing the construction of such tables are fully specified in section four of [1]. Table 1, copied from [1, Table 2], gives the fibbinary table for $\text{fib} = \Phi_0$.

1												
2												
4								5				
8					9			10				
16			17		18			20			21	
32		33	34		36		37	40		41	42	
0	1	2	3	4	5	6	7	8	9	10	11	12

Table 1: The fibbinary table.

Table 2 gives the corresponding table for Φ_1 . This is typical of the tables for the Φ_k , $k \geq 1$.

3												
6												
12								13				
24					25			26				
48		49			50			52			53	
96		97	98		100		101	104		105	106	
0	1	2	3	4	5	6	7	8	9	10	11	12

Table 2: The table for the set Φ_1 .

Various features of such tables are of use in section four. First, for all $k \geq 0$, at the head of all columns of each table all the odd integers of Ψ_k are found. Second, below each such odd integer, call it j , in the same column, its even multiples $2^r j$, $r \geq 1$, are found. Thus, if the BR of j is written $(j)_2$, then these multiples have BR $(j0^r)_2$. This is important: once mastery of the BRs of the odd numbers has been achieved, the same for the even numbers, and hence all numbers, follows easily. Third, the elements of the Fibonacci subsets \mathcal{F}_i , $i = 1, \dots, 6$, of the set displayed are aligned along the rows $i = 1, \dots, 6$, of the table.

Next, note that the structure of the set $\Phi_0 = \text{fib}$ can be expressed in the form

$$\Phi_0 = \sum_{s=0}^{\infty} \alpha_0(s), \quad \alpha_0(1) = 1,$$

$$\alpha_0(s) = \text{odfib}(s) \{2^r, r = 0, 1, \dots, \}. \quad (33)$$

Similarly, for $k \geq 1$,

$$\Phi_k = \sum_{s=0}^{\infty} \alpha_k(s), \quad \alpha_k(1) = 4k - 1,$$

$$\alpha_k(s) = \Psi_k(s) \{2^r, r = 0, 1, \dots, \}. \quad (34)$$

Alongside the result (18)

$$\Psi_0(n) = \text{odfib}(n) = 4\text{fib}(n) + 1 = 4\Phi_0(n) + 1 \quad (35)$$

it can be checked that

$$\Psi_k(n) = 4\Phi_k(n) + 1, \quad k \geq 1, \quad n \geq 1, \quad (36)$$

is correct. Although $\alpha_k = 4k - 1$ holds, by definition, (20), (36) cannot be extended to the case $n = 0$, since $\Phi_k(0)$, $k \geq 1$, is not defined. Setting $\Phi_k(0) = k - \frac{1}{2}$ makes no sense, so that $\Phi_k(0)$, $k \geq 1$, is left undefined. This minor awkwardness is caused by the use of accepted (and natural) definitions of fib and odfib.

An explicit expression for $\Phi_k(n)$, $k \geq 1$, is easy to derive. It reads as

$$\Phi_k(n) = \Phi_0(n) + (2k - 1)2^{l(n)}, \quad (37)$$

where $l(j)$ is the length of j , or the total number of ones and zeros in the ZR of j . The result $\Phi_0(n) = \text{fib}(n)$ does not conform to (37). Also for $k \geq 1$, $n \geq 0$,

$$\Psi_k(n) = \Psi_0(n) + (2k - 1)2^{l(\Psi_0(n))}, \quad (38)$$

and, again, $\Psi_0(n) = \text{odfib}(n)$, $n \geq 0$, does not conform to (38). The result (38) is very important, in full agreement with the picture of the BRs of the odd numbers developed in section four, and commented on in section five. To show that (38) is consistent with (37), use (18) and (19). See also the examples (43) and (44) in section five.

4 The odd numbers arrays ONA1 and ONA

Using the definitions, (27 – 32), of the sets $\Psi_k, k \geq 0$, define the first odd number array ONA1, Table 3.

Ψ_0	Ψ_1	Ψ_2	Ψ_3	Ψ_4	Ψ_5	Ψ_6
1	3	7	11	15	19	23
5	13	29	45	61	77	93
9	25	57	89	121	153	185
17	49	113	177	241	305	369
21	53	117	181	245	309	373
33	97	225	353	481	609	737
37	101	229	357	485	613	741
41	105	233	361	489	617	745
65	193	449	705	961	1217	1473
69	197	453	709	965	1221	1477
73	201	457	713	969	1225	1481
81	209	465	721	977	1233	1489
85	213	469	725	981	1237	1493

Table 3: The first odd number array, ONA1.

The columns of ONA1 contain the integers of the sets $\Psi_k, k \geq 0$. The rows of ONA1 are numbered $n = 0, 1, \dots$, with entries $\Psi_0(0) = \text{odfib}(0) = 1$ and $\Psi_k(0) = \alpha_k = 4k - 1, k \geq 1$, in the top row.

The second odd number array ONA2 is obtained by replacing each element of ONA1 by its unique binary representation, BR, giving rise to Table 4 for ONA2.

Ψ_0	Ψ_1	Ψ_2	Ψ_3	Ψ_4	Ψ_5	Ψ_6
0:1	1:1	11:1	101:1	111:1	1001:1	1011:1
0:101	1:101	11:101	101:101	111:101	1001:101	1011:101
0:1001	1:1001	11:1001	101:1001	111:1001	1001:1001	1011:1001
0:10001	1:10001	11:10001	101:10001	111:10001	1001:10001	1011:10001
0:10101	1:10101	11:10101	101:10101	111:10101	1001:10101	1011:10101
0:10 ⁴ 1	1:10 ⁴ 1	11:10 ⁴ 1	101:10 ⁴ 1	111:10 ⁴ 1	1001:10 ⁴ 1	1011:10 ⁴ 1

Table 4: The second odd number array, ONA2
, showing the binary representations of the elements of ONA1.

It is sometimes convenient to refer to the portion of an array which lies to the right of the double lines of the arrays as an ONA array proper.

The elements of $\Psi_0 = \text{odfib}$ in the first column of ONA2 respect the Zeckendorf condition, ZC, of (9). By the definitions of fib and odfib, all the elements of fib which do so are therefore present in Φ_0 . The elements of the sets Ψ_k , $k \geq 1$, all violate the ZC, and hence belong to $\overline{\text{odfib}}$. The ONA arrays proper in Table 3 and 4 consist entirely of odd numbers which violate the ZC. It is to be proved that all integers of \mathbb{N}_3 which do so are present.

Since all elements of ONA2 proper violate the ZC, their BRs must contain at least one pair 11 of consecutive entries equal to one. In ONA2, the rightmost 11 pair of any BR is written as 1 : 1, with the colon separating that BR into two parts. To the left of the colon, find the principal part PP of the BR, with its rightmost digit equal to one, so that, viewed itself as the BR of some number, the PP represents an odd number. To the right of the colon, find the odfib part OP. The placing of the colon ensures that the OP part of any BR respects the ZC, and represents an element of odfib. Next, note two observations regarding ONA2 proper. First, the PP of the $\Psi_k(n)$, $n \geq 0$, are independent of n for each $k \geq 0$, with these PP equal to 0 for $\Psi_0(n)$, and equal to $(2k - 1)$ for $\Psi_k(n)$, $k \geq 1$. Second, the OP of the $\Psi_k(n)$, $k \geq 0$, are independent of k , with OP equal to $\text{odfib}(n)$.

A colon has been inserted to the left of the odfib entries of the $k = 0$ columns. Their BRs therefore have $\text{PP} = 0$.

The stage is now set to approach the proof of (4). All the elements of Table 4 are odd, by construction of the sets defining its columns. Since BRs are unique no odd number appears twice in Table 4. Put otherwise, with entries written in (PP, OP) form, any two distinct entries differ either in PP or OP parts or both. Also, by construction, the PP part of any entry is either 0 for the odfib elements of the $k = 0$ columns, or else $(2k - 1)$ in columns $k \geq 1$, so that all possible odd numbers are found for the PP parts of all the entries of columns $k \geq 1$. Thus, with the elements of Table 4 written in (PP, OP) form, each part ranges over all the allowed possibilities. No other odd number can arise by the method of construction followed here. It is therefore claimed that all odd numbers are present in Table 4. Those of odfib occur in the $k = 0$ column, the rest occupy the other columns, so that the truth of (4) follows. To complete the justification of this statement, suppose it is alleged that some odd number ν is missing from Table 4. Either ν respects the ZC, in which case $\nu \in \text{fib}$ and is indeed present in the first column, or else $\nu \in \overline{\text{odfib}}$, in which case its unique BR must contain at least one 11 pair. Find the rightmost such pair, and identify the PP and OP parts of ν . But all possible BRs of the form (PP, OP) have been identified. So ν must coincide with exactly one of them, and the allegation is refuted.

It is now claimed that the truth of (3) follows from that of (4). In the sets $\Phi_0 = \text{fib}$, (14), and Φ_k , $k \geq 1$, (27 - 32), it is clear from Tables such as Tables 1 and 2 that all the even elements of such sets are found by reading down the columns of the tables headed by the odd entries. It follows that all the even multiples of all the odd numbers are accounted for, that is, all the even numbers, and hence all $n \in \mathbb{N}$. Further they, all the integers $n \in \mathbb{N}$, separate into the elements of fib, which respect the ZC, and the rest which do not, and hence belong to $\overline{\text{fib}}$. Thus (3), and hence (2), follow.

5 A few comments

In Table 4 it is shown that the odd numbers $\Psi_k(n)$, $k \geq 1$, $n \geq 0$, can be expressed uniquely in the form

$$\begin{aligned} (\text{PP}, \text{OP}) &= ((2k - 1), \text{odfib}(n)), \quad k \geq 1, \\ &= (0, \text{odfib}(n)), \quad k = 0. \end{aligned} \tag{39}$$

This picture is in full agreement with (38). The second term of (38), interpreted in binary, corresponds to $(2k - 1)$ in binary followed by a number of zeros equal to the length of $\text{odfib}(n)$. This means binary addition to this of the first term $\text{odfib}(n)$ is trivial, producing results of the form seen in the entries of Table 4.

Some examples, first for Ψ sets and (38):

$$\begin{aligned} 57 = \Psi_2(2) &= (11 : 1001)_2 = (1001)_2 + (11)_2 2^4 \\ &= 9 + 3.16 = \text{odfib}(2) + (2.2 - 1)2^{l(\text{odfib}(2))}. \end{aligned} \quad (40)$$

$$\begin{aligned} 481 = \Psi_4(5) &= (111 : 100001)_2 = (100001)_2 + (111)_2 2^6 \\ &= 33 + 7.64 = \text{odfib}(5) + (2.4 - 1)2^{l(\text{odfib}(5))}. \end{aligned} \quad (41)$$

Examples for elements of Φ sets and (37):

$$\begin{aligned} 114 = \Phi_2(10) &= (11 : 10010)_2 = (10010)_2 + (11)_2 2^5 \\ &= 18 + 3.32 = \text{fib}(10) + (2.2 - 1)2^{l(\text{fib}(10))}. \end{aligned} \quad (42)$$

$$\begin{aligned} 485 = \Phi_4(17) &= (111 : 100101)_2 = (100101)_2 + (111)_2 2^6 \\ &= 37 + 7.64 = \text{fib}(17) + (2.4 - 1)2^{l(17)} \end{aligned} \quad (43)$$

Note that (43) can also be viewed in the context of (38):

$$\begin{aligned} 485 = \Psi_4(6) &= 37 + 448 \\ &= \text{odfib}(6) + (2.4 - 1)2^{l(\text{odfib}(6))}, \end{aligned} \quad (44)$$

with $\text{fib}(17) = 37 = \text{odfib}(6)$.

References

- [1] A. J. Macfarlane, *On the fibbinary numbers and the Wythoff array*, arXiv:2405.18128.
- [2] L. Lindros, A. Silk and H. Wang, *Odd fibbinary numbers and the golden ratio*, *Fibonacci Quarterly* **52** 61–65 (2014).
- [3] N. J. A. Sloane, *The On-line Encyclopedia of Integer Sequences*, <http://oeis.org>
- [4] E. Zeckendorf, *Représentations des nombres naturels par une somme de nombres de Fibonacci*, *Bul. Soc. Roy. Sci. Liège*, **41** 179–182 (1972), with the proof of 1939.
- [5] C. G. Lekkerkerker, *Voorstelling van natuurlijke getallen door een som van getallen van Fibonacci*, *Simon Steven* **29** 190–195 (1952).
- [6] N. J. A. Sloane, *Classic sequences in the OEIS*, oeis.org/classic.html.
- [7] J-P. Allouche and J. Shallit, *Automatic sequences: theory, applications, generalisations*, Cambridge University Press, 2003.