# Spin-independent interactions of Dirac Fermionic Dark Matter in the composite Higgs models

M. G. Belyakova, R. Nevzorov

I. E. Tamm Department of Theoretical Physics, Lebedev Physical Institute, Leninsky prospect 53, 119991 Moscow, Russia

#### Abstract

According to recent measurements, dark matter magnetic dipole moment is strongly constrained. In the composite Higgs models the magnetic dipole moment of the Dirac dark matter fermion and its mass can be suppressed by the approximate U(1)symmetry. We consider  $E_6$  inspired composite Higgs model ( $E_6$ CHM) with U(1)symmetry violating operators, which give rise to dark matter's mass and coupling constant to Higgs boson. The dependence of the spin-independent dark matternucleon scattering cross section on the  $E_6$ CHM parameters is explored. We argue that there are regions of the parameter space which are still safe from all current constraints and may lead to spectacular LHC signatures.

### 1 Introduction

Numerous astrophysical and cosmological observations confirm the presence of a relatively large non-luminous (dark) component in the matter content of the Universe. The explanation of these observations requires the existence of galaxy halos composed of dark matter (DM) objects. Most likely the corresponding non-luminous component is constituted of nonrelativistic particles which interact gravitationally and possibly by weak interactions. Candidates for DM state occur naturally in a variety of extensions of the standard model (SM) and may possess diverse properties. All such states can contribute to DM content of the Universe. It is especially interesting to explore the nature of DM candidates within well motivated extensions of the SM that permit to almost stabilize the hierarchy between Grand Unification (or Planck) and electroweak (EW) scales. Here we focus on the composite Higgs models. They are based on the ideas which were proposed in the 70's [1] and 80's [2]. These extensions of the SM contain two sectors [3]. The weakly-coupled sector involves elementary states with the quantum numbers of the SM fermions and SM gauge bosons. The second, strongly interacting sector lead to a set of resonances. In particular, it is expected that the breakdown of an approximate global symmetry near the scale f in this sector gives rise to pseudo-Nambu-Goldstone bosons (pNGBs). The present experimental data indicate that the compositeness scale f should be substantially larger than 1 TeV. Four of the pNGBs form Higgs doublet H. The vacuum expectation value (VEV)  $v \simeq 246 \,\text{GeV}$  of H breaks the EW symmetry inducing the masses of all SM particles.

In principle, the strongly interacting sector in the composite Higgs models may also result in the neutral Dirac fermion  $\chi$  that can contribute to the DM in our Universe. Such fermion tends to have a magnetic dipole moment  $\mu_{\chi}$  that within these extensions of the SM is of the order of  $\mu_{\chi} \sim e/f$  where e is the electron charge. The limits on  $\mu_{\chi}$  set by DM direct detection experiments imply that  $f \gtrsim 10^4 \text{ TeV} [4]-[5]^1$ . In this case an enormous degree of tuning, which is about  $\xi \simeq v^2/f^2$ , is required to get a Higgs boson with mass 125 GeV. However in some composite Higgs models the interaction in the Lagrangian

$$\mathcal{L}_{MDM} = \frac{\mu_{\chi}}{2} \bar{\chi}_R \, \sigma^{\mu\nu} \chi_L F_{\mu\nu} + h.c. \tag{1}$$

can be suppressed by the approximate global symmetry. This symmetry may also ensure that  $\chi$  is the lightest composite state.

In this note we investigate the interaction of the Dirac DM fermion with nucleons in the framework of the  $E_6$  inspired composite Higgs model ( $E_6$ CHM) [7]– [10]. Within this model the composite sector is invariant under the transformations of

<sup>&</sup>lt;sup>1</sup>Other bounds are considerably weaker [6].

 $SU(6) \times U(1)_L \times U(1)_B$  symmetry where  $U(1)_L$  and  $U(1)_B$  are associated with the conservations of the lepton and baryon numbers respectively. The breakdown of approximate SU(6) symmetry near the scale f down to SU(5), which involves the SM  $SU(3)_C \times SU(2)_W \times U(1)_Y$  gauge group, gives rise to the composite Higgs doublet H. In the limit when the lightest neutral exotic fermion has a mass which is considerably smaller then 1 TeV the E<sub>6</sub>CHM possesses an additional approximate  $U(1)_E$  symmetry that allows to suppress the interaction (1) in the Lagrangian. We examine the dependence of the spin-independent part of the dark matter-nucleon scattering cross section on the parameters of the model in this case and consider its possible LHC signatures.

The layout of this paper is as follows. In the next section we briefly review the composite Higgs models and  $E_6$ CHM. In section 3 we discuss the generation of masses of exotic fermions and dark matter within the  $E_6$ CHM. The spin-independent interactions of the lightest neutral exotic fermion with nucleons are explored in section 4. Section 5 concludes the paper.

#### 2 Composite Higgs models and E<sub>6</sub>CHM

In the minimal composite Higgs model (MCHM) [11] the invariance of the Lagrangian of the strongly interacting sector with respect to  $SO(5) \times U(1)_X$  symmetry transformations is imposed. Near the scale f this approximate global symmetry is broken down to  $SO(4) \times U(1)'_X \cong SU(2)_W \times SU(2)_R \times U(1)'_X$ , which includes the  $SU(2)_W \times U(1)_Y$  gauge group as a subgroup. The corresponding breakdown leads to four pNGBs that compose the Higgs doublet H. In this case  $SU(2)_{cust} \subset SU(2)_W \times SU(2)_R$  [12] protects the Peskin– Takeuchi  $\hat{T}$  parameter [13] against the contributions induced by new states. The contributions of new resonances to the electroweak observables within the composite Higgs models were analysed in Refs. [14]– [22]. The implications of these extensions of the SM were also studied for Higgs physics [17]– [18], [23]– [26], gauge coupling unification [27]– [28], dark matter [15], [24], [28]– [29] and collider phenomenology [16]– [17], [19], [26], [30]– [32]. Different extensions of the MCHM were considered in Refs. [15], [23]– [24], [28]– [29], [33].

At low energies the SM fields are superpositions of the corresponding elementary states and their composite partners. According to this partial compositeness framework [34,35] the couplings of the SM particles to the composite Higgs and other composite states are proportional to its fraction of compositeness. Since all SM fermions except the top quark are relatively light their fractions of compositeness tend to be rather small resulting in the partial suppression of flavour–changing processes in the composite Higgs models [34]. Within these models the constraints caused by the non–diagonal flavour transitions in the quark and lepton sectors were analysed in Refs. [20]– [22], [30], [36]– [37] and [31], [37]– [39], respectively. In general the corresponding constraints require the compositeness scale f to be larger than 10 TeV [20]– [21], [30], [36], [38]. Nevertheless in the composite Higgs models with additional global flavour symmetries this bound can be substantially weakened [19]– [20], [30]– [31], [37], [40].

The particle content, the global and gauge symmetries of the E<sub>6</sub>CHM may originate from  $E_6 \times G_0$  Supersymmetric (SUSY) Grand Unified Theory (GUT). Fields belonging to the weakly-coupled sector participate in the  $E_6$  interactions only whereas strongly interacting sector comprises multiplets which are charged under both the  $G_0$  and  $E_6$ gauge groups. Near the GUT scale  $M_X \sim 10^{16} \text{ GeV } G_0$  and  $E_6$  can be broken to their subgroups G and  $SU(3)_C \times SU(2)_W \times U(1)_Y$  so that SU(6) remains an approximate global symmetry of the strongly coupled sector. The Lagrangian of the E<sub>6</sub>CHM may also possess  $U(1)_L$  and  $U(1)_B$  global symmetries if different multiplets of the elementary leptons and quarks stem from different fundamental 27-dimensional representations of  $E_6$ . All other components of the corresponding 27-plets gain masses of the order of  $M_X$ . Such splitting of the fundamental representations of  $E_6$  can take place within a six-dimensional orbifold GUT model with N = 1 supersymmetry (SUSY) [7,8]. In this case the breakdown of SUSY can occur somewhat below the GUT scale  $M_X^2$ . The process of the generation of baryon asymmetry within the E<sub>6</sub>CHM was explored in [8,9].

Approximate  $U(1)_B$  and  $U(1)_L$  symmetries permit to get the appropriate suppression of the proton decay rate in the E<sub>6</sub>CHM. Hereafter we assume that  $U(1)_B$  is almost exact. The approximate  $U(1)_L$  symmetry forbids operators that give rise to unacceptably large masses of the left-handed neutrino in the composite Higgs models. Tiny Majorana masses of the left-handed neutrino can be still induced if in the weakly-coupled sector  $U(1)_L$  is broken down to  $Z_2^L = (-1)^L$ , where L is a lepton number.

Below the scale f the approximate SU(6) symmetry is broken down to SU(5) giving rise to eleven pNGB states. The SU(5) symmetry does not contain  $SU(2)_{cust}$  subgroup. As a consequence the electroweak precision measurements set stringent lower limit  $f \gtrsim 5 - 6$  TeV in the E<sub>6</sub>CHM [7]. The generators of the SU(5) subgroup of SU(6) and eleven generators from the coset SU(6)/SU(5) are denoted here by  $T^a$  and  $T^{\hat{a}}$  respectively, where  $\text{Tr}\left(T^aT^b\right) = \frac{1}{2}\delta_{ab}$ ,  $\text{Tr}\left(T^{\hat{a}}T^{\hat{b}}\right) = \frac{1}{2}\delta_{\hat{a}\hat{b}}$  and  $\text{Tr}\left(T^aT^{\hat{b}}\right) = 0$ . For the eleven pNGB states it is convenient to use the following non–linear representation [7]

$$\Omega^{T} = \Omega_{0}^{T} \Sigma^{T} = e^{i \frac{\phi_{0}}{\sqrt{15}f}} \left( C\phi_{1} \quad C\phi_{2} \quad C\phi_{3} \quad C\phi_{4} \quad C\phi_{5} \quad \cos\frac{\tilde{\phi}}{\sqrt{2}f} + \sqrt{\frac{3}{10}}C\phi_{0} \right),$$

$$C = \frac{i}{\tilde{\phi}} \sin\frac{\tilde{\phi}}{\sqrt{2}f}, \qquad \tilde{\phi} = \sqrt{\frac{3}{10}\phi_{0}^{2} + |\phi_{1}|^{2} + |\phi_{2}|^{2} + |\phi_{3}|^{2} + |\phi_{4}|^{2} + |\phi_{5}|^{2}},$$
(2)

<sup>&</sup>lt;sup>2</sup>The phenomenological aspects of the  $E_6$  inspired models with low-scale SUSY breaking were examined in [41].

where a 6-component unit vector  $\Omega$  is a fundamental representation of SU(6) given by

$$\Omega^{T} = \Omega_{0}^{T} \Sigma^{T}, \qquad \Omega_{0}^{T} = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \qquad \Sigma = e^{i\Pi/f}, \qquad \Pi = \Pi^{\hat{a}} T^{\hat{a}}.$$

In Eq. (2) fields  $\phi_1, \phi_2, \phi_3, \phi_4$  and  $\phi_5$  are complex whereas  $\phi_0$  is a real field. Because  $\phi_0$  and  $\tilde{\phi}$  are invariant under the preserved SU(5) vector  $\Omega$  transforms as  $\mathbf{5} + \mathbf{1}$  under the transformation of the unbroken SU(5), where a 5-component vector is formed by  $\mathbf{5} \sim (\phi_1 \phi_2 \phi_3 \phi_4 \phi_5)$ . The SU(5) singlet state  $\mathbf{1} = \phi_0$  is a real SM singlet field. The first two components of  $\mathbf{5}$ , i.e.  $(\phi_1 \phi_2)$ , transform as an  $SU(2)_W$  doublet and therefore they are associated with the SM-like Higgs doublet H. Three other components of  $\mathbf{5}$ , i.e.  $(\phi_3 \phi_4 \phi_5)$ , form an  $SU(3)_C$  triplet T. Since Higgs doublet has B = L = 0, all components of  $\Omega$  do not carry any lepton and/or baryon numbers.

In the leading approximation the Lagrangian describing the interactions of the pNGBs can be written as

$$\mathcal{L}_{pNGB} = \frac{f^2}{2} \left| \mathcal{D}_{\mu} \Omega \right|^2.$$
(3)

The effective potential  $V(H, T, \phi_0)$ , that vanishes in the exact SU(6) symmetry limit, is induced by the interactions of the elementary fermions and gauge bosons with their composite partners, which break global SU(6) symmetry. The analysis performed within the composite Higgs models, which are similar to the E<sub>6</sub>CHM, indicates that there is a substantial region of the parameter space where the EW symmetry is broken to  $U(1)_{em}$ , corresponding to electromagnetism, while the  $SU(3)_C$  symmetry is preserved [15], [28]. It was also shown that in such models the parameters may be tuned so that 125 GeV Higgs state can be obtained [28]. The  $SU(3)_C$  triplet scalar T is much heavier than the SM–like Higgs scalar in this case.

#### 3 Exotic fermions and dark matter in the $E_6$ CHM

Since t-quark is considerably heavier than other SM fermions, the compositeness fraction of the right-handed top quark  $t^c$  should be of the order of unity. In fact, the E<sub>6</sub>CHM implies that  $t^c$  is entirely composite. This can happen if the weakly-coupled sector includes the following set of fermion multiplets [7]

$$(q_i, d_i^c, \ell_i, e_i^c) + u_{\alpha}^c + \bar{q} + \bar{d}^c + \bar{\ell} + \bar{e}^c,$$
 (4)

where  $\alpha = 1, 2$  and i = 1, 2, 3. In Eq. (4)  $e_i^c$ ,  $u_{\alpha}^c$ ,  $d_i^c$  represent the right-handed charged leptons, up- and down-type quarks,  $\ell_i$  and  $q_i$  are associated with the left-handed lepton and quark doublets, whereas extra exotic fermions  $\bar{e}^c$ ,  $\bar{\ell}$ ,  $\bar{q}$  and  $\bar{d}^c$  have exactly opposite  $SU(3)_C \times SU(2)_W \times U(1)_Y$  quantum numbers to the right-handed charged leptons, lefthanded lepton doublets, left-handed quark doublets and right-handed down-type quarks, respectively. The set of multiplets (4) is chosen so that the weakly-coupled sector contains all SM fermions except  $t^c$  and anomaly cancellation takes place. It is expected that the dynamics of strongly interacting sector results in the  $\mathbf{10} + \overline{\mathbf{5}}$  fermion multiplets of SU(5) below the compositeness scale f. Most components of these SU(5) multiplets get combined with  $e^c$ ,  $\bar{\ell}$ ,  $\bar{q}$  and  $\bar{d}^c$  leading to vector-like fermions with masses of the order of f. However the components of the 10-plet associated with the right-handed top quark  $t^c$ survive to the EW scale. The particle content of the weakly-coupled sector (4) leads to the approximate unification of the SM gauge couplings near the scale  $M_X \sim 10^{15} - 10^{16}$  GeV.

The composite  $\mathbf{10} + \overline{\mathbf{5}}$  multiplets of SU(5) can stem from one  $\mathbf{15}$ -plet and two  $\overline{\mathbf{6}}$ plets ( $\overline{\mathbf{6}}_1$  and  $\overline{\mathbf{6}}_2$ ) of SU(6) that have the following decomposition in terms of SU(5)representations:  $\mathbf{15} = \mathbf{10} \oplus \mathbf{5}$  and  $\overline{\mathbf{6}} = \overline{\mathbf{5}} \oplus \mathbf{1}$ . The components of  $\mathbf{15}$ ,  $\overline{\mathbf{6}}_1$  and  $\overline{\mathbf{6}}_2$  decompose under  $SU(3)_C \times SU(2)_W \times U(1)_Y \times U(1)_B$  as follows:

$$\mathbf{5} \rightarrow Q = \left(3, 2, \frac{1}{6}, -\frac{1}{3}\right), \\
 t^{c} = \left(\overline{3}, 1, -\frac{2}{3}, -\frac{1}{3}\right), \\
 t^{c} = \left(1, 1, 1, -\frac{1}{3}, -\frac{1}{3}\right), \\
 D = \left(3, 1, -\frac{1}{3}, -\frac{1}{3}\right), \\
 \overline{L} = \left(1, 2, \frac{1}{2}, -\frac{1}{3}\right); \\
 \overline{L} = \left(1, 2, \frac{1}{2}, -\frac{1}{3}\right); \\
 \overline{N}_{2} = \left(1, 1, 0, -\frac{1}{3}\right).$$

$$\overline{B}_{1} \rightarrow D_{1}^{c} = \left(\overline{3}, 1, \frac{1}{3}, \frac{1}{3}\right), \\
 L_{1} = \left(1, 2, -\frac{1}{2}, \frac{1}{3}\right), \\
 L_{1} = \left(1, 2, -\frac{1}{2}, \frac{1}{3}\right), \\
 N_{1} = \left(1, 2, -\frac{1}{2}, -\frac{1}{3}\right), \\
 L_{2} = \left(1, 2, -\frac{1}{2}, -\frac{1}{3}\right), \\
 \overline{N}_{2} = \left(1, 1, 0, -\frac{1}{3}\right).$$

$$(5)$$

The first and second quantities in brackets are the  $SU(3)_C$  and  $SU(2)_W$  representations, whereas the third and fourth quantities are  $U(1)_Y$  and  $U(1)_B$  charges.

The right-handed top quark  $t^c$  is contained in **15**-plet of SU(6). Then the baryon number conservation requires all components of this multiplet to carry the same baryon number B = -1/3. The masses of D and  $\overline{L}$  components of **15**-plet are induced through interaction

$$\mathcal{L}_{SU(6)}^d = h_N f(\mathbf{15} \times \overline{\mathbf{6}}_{\mathbf{1}} \times \Omega^{\dagger}) + h.c. \,. \tag{6}$$

The dimensionless coupling  $h_N$  is expected to be of the order of unity. The corresponding operator is allowed only if all components of  $\overline{6}_1$  have baryon number B = 1/3. The non-zero mass of  $N_1$  and  $\overline{N}_2$  can be induced if the baryon number of  $\overline{6}_2$  is either 1/3 or (-1/3). Here we assume that  $D_2^c$ ,  $L_2$  and  $N_2$  carry baryon number B = -1/3. Then the corresponding mass term is generated after the breakdown of SU(6) symmetry through interaction

$$\mathcal{L}^{n}_{SU(6)} = g_N f(\overline{\mathbf{6}}_1 \times \Omega)(\overline{\mathbf{6}}_2 \times \Omega) + h.c., \qquad (7)$$

where  $g_N$  is a dimensionless coupling. As a result the full set of mass terms associated with exotic states in the E<sub>6</sub>CHM can be written as follows

$$\mathcal{L}_{mass} = \mu_q \bar{q} Q + \mu_e \bar{e}^c E^c + \mu_D D_1^c D + \mu_L \overline{L} L_1 + \mu_d \bar{d}^c D_2^c + \mu_l \bar{\ell} L_2 + \mu_N \overline{N}_2 N_1 + h.c. , \quad (8)$$

where  $\mu_N \simeq g_N f$ ,  $\mu_D \simeq \mu_L \simeq h_N f$  and  $\mu_q \sim \mu_e \sim \mu_d \sim \mu_l \sim f$ . All exotic fermions mentioned above do not carry any lepton number.

Eq. (8) indicates that in the limit  $g_N = 0$  the E<sub>6</sub>CHM Lagrangian is invariant under the transformations of  $U(1)_E$  global symmetry defined as

$$\overline{\mathbf{6}}_{\mathbf{2}} \longrightarrow e^{i\beta}\overline{\mathbf{6}}_{\mathbf{2}}, \qquad \overline{d}^{c} \longrightarrow e^{-i\beta}\overline{d}^{c}, \qquad \overline{\ell} \longrightarrow e^{-i\beta}\overline{\ell}.$$
(9)

The approximate  $U(1)_E$  symmetry ensures that the lightest exotic fermion  $\chi$  is mostly superposition of  $N_1$  and  $N_2$ . The baryon number conservation does not permit such lightest exotic state to decay. This can be understood in terms of the discrete  $Z_3$  symmetry which is known as baryon triality [15], [42]. The corresponding symmetry transformations are given by

$$\Psi \longrightarrow e^{2\pi i B_3/3} \Psi, \qquad B_3 = (3B - n_C)_{\text{mod } 3}, \tag{10}$$

where B is the baryon number of the multiplet  $\Psi$  and  $n_C$  is the number of colour indices  $(n_C = 1 \text{ for the colour triplet and } n_C = -1 \text{ for antitriplet})$ . Since the baryon number is preserved, the E<sub>6</sub>CHM Lagrangian is also invariant under the symmetry transformations (10). All SM bosons and SM fermions have  $B_3 = 0$  while exotic fermions and the scalar colour triplet T carry either  $B_3 = 2$  or  $B_3 = 1$ . As a consequence the lightest exotic fermion  $\chi$  with non-zero  $B_3$  charge cannot decay into SM particles and should therefore be stable.

Thus  $\chi$  may account for all or some of the observed cold dark matter density. Because the lightest exotic fermion carries baryon number, some part of the baryon asymmetry of our Universe is stored in the dark matter sector. Moreover, in the E<sub>6</sub>CHM the relic abundance of the lightest exotic fermions might be induced by the same mechanism that results in the baryon asymmetry. In particular, this can happen if the mass of the lightest exotic fermion  $m_{\chi}$  is quite close to half the mass of the SM singlet pNGB state  $\phi_0$ , i.e.  $m_A/2$ , so that  $\chi \bar{\chi}$  annihilation is efficient enough. The ratio of the baryon charges  $B_{\chi}$ and  $B_N$  accumulated by  $\chi$  and nucleons can be estimated as

$$\frac{B_{\chi}}{B_N} \simeq \frac{1}{3} \left(\frac{\rho_{\chi}}{\rho_N}\right) \left(\frac{m_N}{m_{\chi}}\right) \,, \tag{11}$$

where  $\rho_N$  and  $\rho_{\chi}$  are contributions of nucleons and lightest exotic fermions to the total energy density, whereas  $m_N$  and  $m_{\chi}$  are the masses of nucleons and  $\chi$ , respectively. For  $\rho_{\chi} \lesssim 5\rho_N$  this ratio can be bigger than 0.01 only if  $m_{\chi} \lesssim 200$  GeV.

## 4 Spin-independent interactions of dark matter with nucleons in the $E_6$ CHM

In the composite Higgs models with  $f \gg v$  the spin-independent interactions of the Dirac fermionic Dark Matter with nucleons are mediated by the *t*-channel exchange of the 125 GeV Higgs scalar, Z-boson and photon. However the contributions of diagrams involving virtual photon or virtual Higgs state may be strongly suppressed by the approximate  $U(1)_E$  global symmetry. To simplify our analysis here we assume that the magnetic dipole moment of the lightest exotic fermion  $\mu_{\chi}$  is much smaller than the present experimental limit  $\sim 10^{-8} \,\text{GeV}^{-1}$  and the interaction (1) can be ignored. At the same time we allow the  $U(1)_E$  symmetry violating operators that give rise to  $m_{\chi}$  and the coupling of  $\chi$  to the 125 GeV Higgs boson.

The  $SU(2)_W$  doublets  $\bar{\ell}$ ,  $\bar{L}$ ,  $L_1$  and  $L_2$  contain electrically neutral components  $\bar{\nu}_{\ell}$ ,  $\bar{\nu}_L$ ,  $\nu_1$ and  $\nu_2$ . These components mix with  $N_1$  and  $\bar{N}_2$  after the breakdown of the EW symmetry. When  $U(1)_E$  symmetry is preserved to a very good approximation the conservation of the baryon number allows only a few terms in the E<sub>6</sub>CHM Lagrangian that give rise to such mixing, i.e.

$$\mathcal{L}_{mix} = h_N(\overline{L}H^c)N_1 + \tilde{h}_N(\overline{\ell}H^c)\overline{N}_2 + h.c.$$
(12)

In this limit the left-handed and right-handed components of the lightest exotic fermion  $\chi$  are given by

$$\chi_L \simeq N_1 \cos \theta_1 - \nu_1 \sin \theta_1 , \qquad \chi_R \simeq N_2 \cos \theta_2 - \bar{\nu}_2 \sin \theta_2 , \qquad (13)$$

where  $\tan \theta_1 \simeq \frac{v}{\sqrt{2}f}$  and  $\tan \theta_2 \simeq \frac{\tilde{h}_N v}{\sqrt{2}\mu_\ell}$ . The mixing angles  $\theta_1$  and  $\theta_2$  determine the strength of the interaction of  $\chi$  with the Z-boson. The corresponding part of the Lagrangian can be written as

$$\mathcal{L}_{Z\chi} = \overline{\chi} (a_V^{\chi} \gamma^{\mu} + a_{PV}^{\chi} \gamma^{\mu} \gamma^5) \chi Z_{\mu} ,$$

$$a_V^{\chi} = \frac{\overline{g}}{4} (\sin^2 \theta_1 - \sin^2 \theta_2) , \qquad a_{PV}^{\chi} = \frac{\overline{g}}{4} (\sin^2 \theta_1 + \sin^2 \theta_2) .$$
(14)

In Eq. (14)  $\bar{g} = \sqrt{g^2 + g'^2}$  whereas g and g' are  $SU(2)_W$  and  $U(1)_Y$  gauge couplings. Since  $v \ll f$  the mixing angles  $\theta_1$  and  $\theta_2$  as well as the couplings  $a_V^{\chi}$  and  $a_{PV}^{\chi}$  are always small. Therefore it is convenient to use the following parameterisation

$$a_{V}^{\chi} = \frac{\bar{g}v^{2}}{8f^{2}}c_{V}^{\chi}, \qquad c_{V}^{\chi} \simeq 1 - \left(\frac{\tilde{h}_{N}f}{\mu_{\ell}}\right)^{2}, \qquad (15)$$
$$a_{PV}^{\chi} = \frac{\bar{g}v^{2}}{8f^{2}}c_{PV}^{\chi}, \qquad c_{PV}^{\chi} \simeq 1 + \left(\frac{\tilde{h}_{N}f}{\mu_{\ell}}\right)^{2}.$$

When  $\mu_{\ell} \gtrsim \tilde{h}_N f$ , the dimensionless coefficients  $c_V^{\chi}$  and  $c_{PV}^{\chi}$  are of the order of unity.

Using the Lagrangian describing the interactions of the Z-boson with quarks

$$\mathcal{L}_{Zq} = \sum_{q} \frac{\bar{g}}{2} \bar{q} (a_{V}^{q} \gamma^{\mu} + a_{PV}^{q} \gamma^{\mu} \gamma^{5}) q Z_{\mu} = \frac{\bar{g}}{2} J_{NC}^{\mu} Z_{\mu} ,$$

$$a_{V}^{q} = T_{3q} - 2s_{W}^{2} Q_{q} , \qquad a_{PV}^{q} = T_{3q} ,$$
(16)

one can compute in the leading approximation the hadronic matrix elements

$$\langle N'|J_{NC}^{\mu}|N\rangle = \overline{\Psi}_{N}'(a_{V}^{N}\gamma^{\mu} + a_{PV}^{N}\gamma^{\mu}\gamma^{5})\Psi_{N},$$
  
$$a_{V}^{p} \simeq \frac{1}{2} - 2s_{W}^{2}, \qquad a_{PV}^{p} \simeq -a_{V}^{n} \simeq -a_{PV}^{n} \simeq \frac{1}{2}$$
(17)

In Eq. (16)–(17) N = p, n and  $s_W \simeq g'/\bar{g}$  while  $T_{3q}$  and  $Q_q$  are the third component of isospin and the electric charge of the quark q respectively.

The interaction of the  $125 \,\text{GeV}$  Higgs boson h with the lightest exotic fermion is determined by the following term in the Lagrangian

$$\mathcal{L}_{H\chi} = \frac{\varepsilon_H}{f} H^{\dagger} H(\overline{N}_2 N_1) + h.c. \,. \tag{18}$$

The  $U(1)_E$  symmetry forbids operator (18) so that in this limit the dimensionless parameter  $\varepsilon_H$  vanishes. In the case of approximate  $U(1)_E$  symmetry the lightest exotic fermion  $\chi$  gains non-zero mass  $m_{\chi} \ll f$  while  $\varepsilon_H$  is expected to be small, i.e.  $\varepsilon_H \ll 1$ . As a consequence the breakdown of the EW symmetry leads to the small coupling of the 125 GeV Higgs state h to  $\chi$ 

$$\mathcal{L}_{h\chi\chi} = g_{h\chi\chi}(\overline{\chi}\chi)h, \qquad \qquad g_{h\chi\chi} \simeq \varepsilon_H \frac{v}{f}.$$
(19)

The strength of the interaction of nucleons with the Higgs state h is defined by the coupling

$$g_{hNN} = a_S^N \frac{m_N}{v} \,, \tag{20}$$

where

$$a_S^N = \sum_{q=u,d,s} f_{Tq}^N + \frac{2}{27} \sum_{Q=c,b,t} f_{TQ}^N , \qquad (21)$$

$$\langle N|m_q \bar{q}q|N\rangle = m_N f_{Tq}^N, \qquad \qquad f_{TQ}^N = 1 - \sum_{q=u,d,s} f_{Tq}^N$$

The coefficients  $f_{Tq}^N$  are related to the  $\pi$ -nucleon  $\sigma$  term and the spin content of the nucleon. In our analysis we fix  $f_{Tq}^p \simeq f_{Tq}^n \simeq f_{Tq}$ , i.e.  $a_S^p \simeq a_S^n \simeq a_S$ , as well as  $f_{Tu} \simeq 0.0153$ ,  $f_{Td} \simeq 0.0191$  and  $f_{Ts} \simeq 0.0447$  which are the default values used in micrOMEGAs [43].

In the leading approximation the spin–independent  $\chi - N$  scattering cross section averaged over initial spins and summed over final spins takes the form

$$\sigma = \frac{m_r^2}{\pi} \left| \frac{g_{h\chi\chi} a_S m_N}{v m_h^2} - \frac{\bar{g} a_V^{\chi} \langle a_V \rangle}{2M_Z^2} \right|^2, \qquad m_r = \frac{m_{\chi} m_N}{m_{\chi} + m_N}. \tag{22}$$



Figure 1: The dependence of the spin-independent  $\chi - N$  scattering cross section  $\sigma$  on the compositeness scale f for  $\varepsilon_H = 0$  (Left) and  $\varepsilon_H = -0.1$  (Right). The solid and dotted lines correspond to  $c_V^{\chi} = 1$  and  $c_V^{\chi} = 0.5$  respectively. The highest, second lowest and lowest dashed lines represent the experimental limits on  $\sigma$  for  $m_{\chi} = 1$  TeV,  $m_{\chi} = 500$  GeV and  $m_{\chi} = 200$  GeV.

In Eq. (22)  $M_Z$  and  $m_h$  are the masses of the Z-boson and Higgs scalar, i.e.  $M_Z \simeq 91.2 \text{ GeV}$  and  $m_h \simeq 125 \text{ GeV}$ , while  $\langle a_V \rangle$  is given by

$$\langle a_V \rangle = \frac{1}{A} \left( Z a_V^p + (A - Z) a_V^n \right),$$

where A and Z are the nucleon number and charge of the target nucleus. Focussing on the case of xenon target nucleus we set  $A \approx 130$  and Z = 54.

The results of our analysis are summarised in Fig. 1. In this figure the results of theoretical calculations of the spin-independent  $\chi - N$  scattering cross section in the E<sub>6</sub>CHM are compared with the corresponding experimental limits which are 50 yb, 140 yb and 300 yb for the dark matter masses 200 GeV, 500 GeV and 1 TeV respectively [44,45]. Since the E<sub>6</sub>CHM does not possess the  $SU(2)_{cust}$  symmetry we focus on the part of the parameter space that corresponds to  $f \gtrsim 5$  TeV as it was discussed in Section 2. To avoid too large fine-tuning associated with the stabilisation of the EW scale we restrict our consideration to  $f \lesssim 10$  TeV. Here we also assume that  $m_{\chi} \gg m_N$  so that  $\sigma$  does not change much when  $m_{\chi}$  varies. As a consequence the spin-independent  $\chi - N$  scattering cross section (22) mostly depends on f,  $c_V^{\chi}$  and  $\varepsilon_H$ . As follows from Fig. 1  $\sigma$  diminishes with increasing f. Moreover the spin-independent  $\chi$ -nucleon scattering cross section also

reduces when the absolute value of  $c_V^{\chi}$  decreases. Therefore the scenarios with  $f \gtrsim 5 \text{ TeV}$ and  $m_{\chi} \lesssim 1 \text{ TeV}$  have not been ruled out. The small and negative values of  $\varepsilon_H$  results in further reduction of  $\sigma$ . Because of this for  $\varepsilon_H \simeq -0.1$  and  $f \simeq 5 \text{ TeV}$  one can still find the phenomenologically viable scenarios even if  $m_{\chi} \lesssim 500 \text{ GeV}$ . In the near future the experiments XENONnT [46] and LUX–ZEPLIN (LZ) [47], may set even more stringent constraints on the spin-independent  $\chi - N$  scattering cross section.

#### 5 Conclusions

Recent measurements set stringent constraints on the magnetic dipole moment  $\mu_{\chi}$  of the Dirac DM fermions. Such fermions may compose dark matter density in the composite Higgs models. However the experimental limits on  $\mu_{\chi}$  imply that in general the compositeness scale f in these models has to be larger than  $10^4$  TeV which makes the stabilization of the EW scale rather problematic. In this article we argued that in the composite Higgs models the approximate U(1) symmetry can result in the relatively light Dirac DM fermion with suppressed magnetic dipole moment and small coupling to the 125 GeV Higgs state h. In particular, we considered the  $E_6$  inspired composite Higgs model ( $E_6$ CHM) in which the strongly interacting sector possesses the approximate SU(6) symmetry and the right-handed top quark is composite. The particle content of the  $E_6$ CHM involves a set of exotic fermions which contains two SM singlet states  $N_1$  and  $\overline{N}_2$  with baryon numbers 1/3 and (-1/3). These states form Dirac fermion  $\chi$ . If the E<sub>6</sub>CHM Lagrangian is invariant under the transformations of global  $U(1)_E$  symmetry then the mass of  $\chi$ , its magnetic dipole moment and coupling to h vanish. In the case of the approximate  $U(1)_E$ symmetry  $\chi$  tends to be the lightest exotic fermion in the spectrum. The baryon number conservation ensures that the Dirac fermion  $\chi$  is stable.

The SU(6) symmetry breaking near the scale f down to its SU(5) subgroup, which contains the SM gauge group, leads to the pNGB states in the E<sub>6</sub>CHM that form the SM Higgs doublet H, the scalar colour triplet T and the SM singlet  $\phi_0$ . The lower bound on the compositeness scale f in this model caused by the electroweak precision measurements is about 5 TeV. Because of this the coupling of the lightest exotic fermion  $\chi$  to Z boson is quite suppressed. Neglecting  $\mu_{\chi}$  we explored the spin-independent interactions of  $\chi$  with nucleons which are dominated by t-channel exchange of the Z boson and the SM Higgs scalar in this case. It was demonstrated that the present experimental limits haven't ruled out the scenarios with  $f \leq 10$  TeV and  $m_{\chi} \leq 1$  TeV.

When  $f \simeq 5 - 6$  TeV the scalar colour triplet T may be lighter than 2 TeV. If such states do exist at so low scales, they can be accessed at the LHC. In collider experiments these scalars can only be created in pairs. At the LHC the pairs of  $T\overline{T}$  are mostly produced through the gluon fusion. After being produced T sequentially decays into b-quark and  $\overline{\chi}$  giving rise to some enhancement of the cross section of

$$pp \to b\bar{b} + E_T^{\text{miss}} + X \,,$$

$$\tag{23}$$

where  $E_T^{\text{miss}}$  is the missing energy and transverse momentum associated with  $\chi$  and  $\overline{\chi}$  in the final state whereas X can include extra charged leptons and/or jets that may come from the decays of intermediate states. Similar enhancement of the cross section of the process (23) can be caused by the presence of relatively light superpartner of the *b*-quark. If the mass of the DM fermion is smaller than 800 GeV then ATLAS experiment excluded the bottom squarks with masses below 1200 – 1270 GeV at 95% confidence level [48]. Limits from CMS are comparable [49]. At the same time no limit can be placed for the DM fermion mass above 800 GeV. The experimental bound mentioned here are fully applicable to the case of the scalar colour triplet T. Its discovery at the LHC may permit to distinguish the E<sub>6</sub>CHM from other extensions of the SM.

### Acknowledgements

The authors are grateful to M. I. Vysotsky for valuable comments and remarks.

### References

- H. Terazawa, K. Akama, Y. Chikashige, Phys. Rev. D 15 (1977) 480; H. Terazawa, Phys. Rev. D 22 (1980) 184.
- [2] S. Dimopoulos, J. Preskill, Nucl. Phys. B 199 (1982) 206; D. B. Kaplan, H. Georgi, Phys. Lett. B 136 (1984) 183; D. B. Kaplan, H. Georgi, S. Dimopoulos, Phys. Lett. B 136 (1984) 187; H. Georgi, D. B. Kaplan, P. Galison, Phys. Lett. B 143 (1984) 152; T. Banks, Nucl. Phys. B 243 (1984) 125; H. Georgi, D. B. Kaplan, Phys. Lett. B 145 (1984) 216; M. J. Dugan, H. Georgi, D. B. Kaplan, Nucl. Phys. B 254 (1985) 299; H. Georgi, Nucl. Phys. B 266 (1986) 274.
- [3] B. Bellazzini, C. Csáki, J. Serra, Eur. Phys. J. C 74 (2014) 5, 2766 [arXiv:1401.2457 [hep-ph]].
- [4] B. Ali *et al.* [PICO], Phys. Rev. D 106 (2022) no.4, 042004 [arXiv:2204.10340 [astro-ph.CO]].
- [5] X. Ning *et al.* [PandaX], Nature **618** (2023) 7963, 47; J. Buch, J. Fan, J. S. C.Leung, Phys. Rev. D **101** (2020) 063026 [arXiv:1910.06356 [hep-ph]].

- [6] T. Hambye, X. J. Xu, JHEP **11** (2021) 156 [arXiv:2106.01403 [hep-ph]].
- [7] R. Nevzorov, A. W. Thomas, Phys. Rev. D 92 (2015) 075007 [arXiv:1507.02101 [hep-ph]].
- [8] R. Nevzorov, Universe 8 (2022) 33.
- [9] R. Nevzorov and A. W. Thomas, Phys. Lett. B 774 (2017) 123 [arXiv:1706.02856 [hep-ph]]. R. Nevzorov and A. W. Thomas, EPJ Web Conf. 191 (2018) 02004 [arXiv:1808.01982 [hep-ph]]; R. Nevzorov and A. W. Thomas, Phys. Part. Nucl. 51 (2020) no.4, 709 [arXiv:2001.09843 [hep-ph]].
- [10] R. Nevzorov, A. W. Thomas, EPJ Web Conf. **125** (2016) 02021 [arXiv:1608.00320 [hep-ph]];
   R. Nevzorov, A. W. Thomas, J. Phys. G **44** (2017) 075003 [arXiv:1605.07313 [hep-ph]].
- [11] K. Agashe, R. Contino and A. Pomarol, Nucl. Phys. B 719 (2005) 165 [hepph/0412089].
- [12] P. Sikivie, L. Susskind, M. B. Voloshin and V. I. Zakharov, Nucl. Phys. B 173 (1980) 189.
- [13] M. E. Peskin and T. Takeuchi, Phys. Rev. D 46 (1992) 381.
- [14] K. Agashe, R. Contino, Nucl. Phys. B 742 (2006) 59 [hep-ph/0510164]; K. Agashe, R. Contino, L. Da Rold, A. Pomarol, Phys. Lett. B 641 (2006) 62 [hep-ph/0605341];
  G. F. Giudice, C. Grojean, A. Pomarol, R. Rattazzi, JHEP 0706 (2007) 045 [hep-ph/0703164]; R. Barbieri, B. Bellazzini, V. S. Rychkov, A. Varagnolo, Phys. Rev. D 76 (2007) 115008 [arXiv:0706.0432 [hep-ph]]; P. Lodone, JHEP 0812 (2008) 029 [arXiv:0806.1472 [hep-ph]]; M. Gillioz, Phys. Rev. D 80 (2009) 055003 [arXiv:0806.3450 [hep-ph]]; C. Anastasiou, E. Furlan, J. Santiago, Phys. Rev. D 79 (2009) 075003 [arXiv:0901.2117 [hep-ph]]; G. Panico, A. Wulzer, JHEP 1109 (2011) 135 [arXiv:1106.2719 [hep-ph]]; S. De Curtis, M. Redi, A. Tesi, JHEP 1204 (2012) 042 [arXiv:1110.1613 [hep-ph]]; D. Marzocca, M. Serone, J. Shu, JHEP 1208 (2012) 013 [arXiv:1205.0770 [hep-ph]]; A. Orgogozo, S. Rychkov, JHEP 1306 (2013) 014 [arXiv:1211.5543 [hep-ph]]; D. Pappadopulo, A. Thamm, R. Torre, JHEP 1307 (2013) 058 [arXiv:1303.3062 [hep-ph]]; C. Grojean, O. Matsedonskyi, G. Panico, JHEP 1310 (2013) 160 [arXiv:1306.4655 [hep-ph]].
- [15] M. Frigerio, J. Serra, A. Varagnolo, JHEP **1106** (2011) 029 [arXiv:1103.2997 [hepph]].

- [16] M. Carena, E. Ponton, J. Santiago and C. E. M. Wagner, Nucl. Phys. B 759 (2006) 202 [hep-ph/0607106]; A. Pomarol, J. Serra, Phys. Rev. D 78 (2008) 074026 [arXiv:0806.3247 [hep-ph]]; D. Pappadopulo, A. Thamm and R. Torre, JHEP 1307 (2013) 058 [arXiv:1303.3062 [hep-ph]].
- [17] B. Bellazzini, C. Csaki, J. Hubisz, J. Serra, J. Terning, JHEP **1211** (2012) 003 [arXiv:1205.4032 [hep-ph]].
- [18] M. Gillioz, R. Grober, C. Grojean, M. Muhlleitner, E. Salvioni, JHEP **1210** (2012) 004 [arXiv:1206.7120 [hep-ph]]; A. Azatov, J. Galloway, Int. J. Mod. Phys. A **28** (2013) 1330004 [arXiv:1212.1380]; A. Falkowski, F. Riva and A. Urbano, JHEP **1311** (2013) 111 [arXiv:1303.1812 [hep-ph]]; A. Azatov, R. Contino, A. Di Iura, J. Galloway, Phys. Rev. D **88** (2013) 7, 075019 [arXiv:1308.2676 [hep-ph]]; M. Gillioz, R. Gröber, A. Kapuvari, M. Mühlleitner, JHEP **1403** (2014) 037 [arXiv:1311.4453 [hep-ph]].
- [19] R. Barbieri, G. Isidori, D. Pappadopulo, JHEP 0902 (2009) 029 [arXiv:0811.2888
   [hep-ph]]; O. Matsedonskyi, JHEP 1502 (2015) 154 [arXiv:1411.4638 [hep-ph]].
- [20] R. Barbieri, D. Buttazzo, F. Sala, D. M. Straub, A. Tesi, JHEP **1305** (2013) 069 [arXiv:1211.5085 [hep-ph]].
- [21] C. Csaki, A. Falkowski, A. Weiler, JHEP 0809 (2008) 008 [arXiv:0804.1954 [hep-ph]];
   K. Agashe, A. Azatov, L. Zhu, Phys. Rev. D 79 (2009) 056006 [arXiv:0810.1016 [hep-ph]].
- [22] N. Vignaroli, Phys. Rev. D 86 (2012) 115011 [arXiv:1204.0478 [hep-ph]].
- [23] B. Gripaios, A. Pomarol, F. Riva, J. Serra, JHEP 0904 (2009) 070 [arXiv:0902.1483 [hep-ph]]; J. Mrazek, A. Pomarol, R. Rattazzi, M. Redi, J. Serra, A. Wulzer, Nucl. Phys. B 853 (2011) 1 [arXiv:1105.5403 [hep-ph]]; M. Redi, A. Tesi, JHEP 1210 (2012) 166 [arXiv:1205.0232 [hep-ph]]; E. Bertuzzo, T. S. Ray, H. de Sandes, C. A. Savoy, JHEP 1305 (2013) 153 [arXiv:1206.2623 [hep-ph]]; M. Montull, F. Riva, JHEP 1211 (2012) 018 [arXiv:1207.1716 [hep-ph]]; M. Chala, JHEP 1301 (2013) 122 [arXiv:1210.6208 [hep-ph]].
- [24] M. Frigerio, A. Pomarol, F. Riva, A. Urbano, JHEP **1207** (2012) 015 [arXiv:1204.2808 [hep-ph]].
- [25] R. Contino, C. Grojean, M. Moretti, F. Piccinini, R. Rattazzi, JHEP 1005 (2010)
   089 [arXiv:1002.1011 [hep-ph]]; I. Low, A. Vichi, Phys. Rev. D 84 (2011) 045019

[arXiv:1010.2753 [hep-ph]]; R. Contino, D. Marzocca, D. Pappadopulo, R. Rattazzi, JHEP **1110** (2011) 081 [arXiv:1109.1570 [hep-ph]]; A. Azatov, J. Galloway, Phys. Rev. D 85 (2012) 055013 [arXiv:1110.5646 [hep-ph]]; R. Contino, M. Ghezzi, M. Moretti, G. Panico, F. Piccinini, A. Wulzer, JHEP 1208 (2012) 154 [arXiv:1205.5444 [hep-ph]]; R. Contino, M. Ghezzi, C. Grojean, M. Muhlleitner, M. Spira, JHEP 1307 (2013) 035 [arXiv:1303.3876 [hep-ph]]; C. Delaunay, C. Grojean, G. Perez, JHEP **1309** (2013) 090 [arXiv:1303.5701 [hep-ph]]; A. Banfi, A. Martin, V. Sanz, JHEP 1408 (2014) 053 [arXiv:1308.4771 [hepph]]; M. Montull, F. Riva, E. Salvioni, R. Torre, Phys. Rev. D 88 (2013) 095006 [arXiv:1308.0559 [hep-ph]]; R. Contino, C. Grojean, D. Pappadopulo, R. Rattazzi, A. Thamm, JHEP **1402** (2014) 006 [arXiv:1309.7038 [hep-ph]]; T. Flacke, J. H. Kim, S. J. Lee, S. H. Lim, JHEP 1405 (2014) 123 [arXiv:1312.5316 [hep-ph]]; C. Grojean, E. Salvioni, M. Schlaffer, A. Weiler, JHEP 1405 (2014) 022 [arXiv:1312.3317 [hepph]]; M. Carena, L. Da Rold, E. Pontón, JHEP 1406 (2014) 159 [arXiv:1402.2987 [hep-ph]]; A. Carmona, F. Goertz, JHEP **1505** (2015) 002 [arXiv:1410.8555 [hep-ph]]; G. Buchalla, O. Cata, C. Krause, Nucl. Phys. B 894 (2015) 602 [arXiv:1412.6356 [hep-ph]].

- [26] A. Pomarol, F. Riva, JHEP **1208** (2012) 135 [arXiv:1205.6434 [hep-ph]]; O. Matsedonskyi, G. Panico, A. Wulzer, JHEP **1301** (2013) 164 [arXiv:1204.6333 [hep-ph]].
- [27] K. Agashe, A. Delgado, R. Sundrum, Annals Phys. **304** (2003) 145 [hep-ph/0212028];
   T. Gherghetta, Phys. Rev. D **71** (2005) 065001 [hep-ph/0411090].
- [28] J. Barnard, T. Gherghetta, T. S. Ray, A. Spray, JHEP **1501** (2015) 067 [arXiv:1409.7391 [hep-ph]].
- [29] M. Asano and R. Kitano, JHEP **1409** (2014) 171 [arXiv:1406.6374 [hep-ph]]; M. Chala, G. Nardini and I. Sobolev, Phys. Rev. D **94** (2016) 055006 [arXiv:1605.08663 [hep-ph]].
- [30] M. Redi, A. Weiler, JHEP **1111** (2011) 108 [arXiv:1106.6357 [hep-ph]].
- [31] M. Redi, JHEP **1309** (2013) 060 [arXiv:1306.1525 [hep-ph]].
- [32] K. Agashe, A. Belyaev, T. Krupovnickas, G. Perez, J. Virzi, Phys. Rev. D 77 (2008) 015003 [hep-ph/0612015]; B. Lillie, L. Randall and L. T. Wang, JHEP 0709 (2007) 074 [hep-ph/0701166]; K. Agashe, H. Davoudiasl, S. Gopalakrishna, T. Han, G. Y. Huang, G. Perez, Z. G. Si, A. Soni, Phys. Rev. D 76 (2007) 115015 [arXiv:0709.0007 [hep-ph]]; M. Carena, A. D. Medina, B. Panes, N. R. Shah,

C. E. M. Wagner, Phys. Rev. D 77 (2008) 076003 [arXiv:0712.0095 [hep-ph]]; R. Contino, G. Servant, JHEP 0806 (2008) 026 [arXiv:0801.1679 [hep-ph]]; K. Agashe, S. Gopalakrishna, T. Han, G. Y. Huang, A. Soni, Phys. Rev. D 80 (2009) 075007 [arXiv:0810.1497 [hep-ph]]; J. Mrazek, A. Wulzer, Phys. Rev. D 81 (2010) 075006 [arXiv:0909.3977 [hep-ph]]; K. Agashe, A. Azatov, T. Han, Y. Li, Z. G. Si, L. Zhu, Phys. Rev. D 81 (2010) 096002 [arXiv:0911.0059 [hep-ph]]; G. Dissertori, E. Furlan, F. Moortgat, P. Nef, JHEP 1009 (2010) 019 [arXiv:1005.4414 [hep-ph]]; N. Vignaroli, Phys. Rev. D 86 (2012) 075017 [arXiv:1207.0830 [hep-ph]]; G. Cacciapaglia, A. Deandrea, L. Panizzi, S. Perries, V. Sordini, JHEP 1303 (2013) 004 [arXiv:1211.4034] [hep-ph]]; A. De Simone, O. Matsedonskyi, R. Rattazzi, A. Wulzer, JHEP 1304 (2013) 004 [arXiv:1211.5663 [hep-ph]]; J. Li, D. Liu, J. Shu, JHEP 1311 (2013) 047 [arXiv:1306.5841 [hep-ph]]; M. Redi, V. Sanz, M. de Vries, A. Weiler, JHEP 1308 (2013) 008 [arXiv:1305.3818 [hep-ph]]; C. Delaunay, T. Flacke, J. Gonzalez-Fraile, S. J. Lee, G. Panico, G. Perez, JHEP 1402 (2014) 055 [arXiv:1311.2072 [hep-ph]]; O. Matsedonskyi, F. Riva, T. Vantalon, JHEP 1404 (2014) 059 [arXiv:1401.3740] [hep-ph]]; H. C. Cheng, J. Gu, JHEP **1410** (2014) 002 [arXiv:1406.6689 [hep-ph]]; B. Gripaios, T. Müller, M. A. Parker, D. Sutherland, JHEP 1408 (2014) 171 [arXiv:1406.5957 [hep-ph]]; A. Azatov, G. Panico, G. Perez, Y. Soreq, JHEP 1412 (2014) 082 [arXiv:1408.4525 [hep-ph]]; M. Backović, T. Flacke, J. H. Kim, S. J. Lee, JHEP **1504** (2015) 082 [arXiv:1410.8131 [hep-ph]]; S. Kanemura, K. Kaneta, N. Machida, T. Shindou, Phys. Rev. D 91 (2015) 11, 115016 [arXiv:1410.8413 [hepph]; M. Chala, J. Juknevich, G. Perez and J. Santiago, JHEP 1501 (2015) 092 [arXiv:1411.1771 [hep-ph]]; A. Thamm, R. Torre, A. Wulzer, arXiv:1502.01701 [hepph]; A. Azatov, D. Chowdhury, D. Ghosh, T. S. Ray, arXiv:1505.01506 [hep-ph]; J. Serra, arXiv:1506.05110 [hep-ph].

[33] J. Barnard, T. Gherghetta, T. S. Ray, JHEP 1402 (2014) 002 [arXiv:1311.6562 [hep-ph]]; G. Ferretti, D. Karateev, JHEP 1403 (2014) 077 [arXiv:1312.5330 [hep-ph]]; G. Cacciapaglia, F. Sannino, JHEP 1404 (2014) 111 [arXiv:1402.0233 [hep-ph]]; A. Hietanen, R. Lewis, C. Pica, F. Sannino, JHEP 1407 (2014) 116 [arXiv:1404.2794 [hep-lat]]; G. Ferretti, JHEP 1406 (2014) 142 [arXiv:1404.7137 [hep-ph]]; A. Parolini, Phys. Rev. D 90 (2014) 11, 115026 [arXiv:1405.4875 [hep-ph]]; M. Geller, O. Telem, Phys. Rev. Lett. 114 (2015) 191801 [arXiv:1411.2974 [hep-ph]]; B. Gripaios, M. Nardecchia, S. A. Renner, JHEP 1505 (2015) 006 [arXiv:1412.1791 [hep-ph]]; M. Low, A. Tesi, L. T. Wang, Phys. Rev. D 91 (2015) 094506 [arXiv:1501.07890 [hep-ph]]; M. Golterman, Y. Shamir, Phys. Rev. D 91 (2015) 094506 [arXiv:1502.00390 [hep-ph]]; J. Erdmenger, N. Evans, W. Porod, K. S. Rigatos, Phys. Rev. Lett. 126 (2021) 071602 [arXiv:2009.10737 [hep-ph]]; J. Erdmenger, N. Evans, W. Porod,

K. S. Rigatos, JHEP **02** (2021) 058 [arXiv:2010.10279 [hep-ph]].

- [34] R. Contino, T. Kramer, M. Son, R. Sundrum, JHEP 0705 (2007) 074 [hepph/0612180].
- [35] D. B. Kaplan, Nucl. Phys. B **365** (1991) 259.
- [36] M. Blanke, A. J. Buras, B. Duling, S. Gori, A. Weiler, JHEP 0903 (2009) 001
   [arXiv:0809.1073 [hep-ph]]; O. Gedalia, G. Isidori, G. Perez, Phys. Lett. B 682 (2009)
   200 [arXiv:0905.3264 [hep-ph]].
- [37] R. Barbieri, D. Buttazzo, F. Sala, D. M. Straub, JHEP **1207** (2012) 181 [arXiv:1203.4218 [hep-ph]].
- [38] K. Agashe, A. E. Blechman, F. Petriello, Phys. Rev. D 74 (2006) 053011 [hep-ph/0606021]; C. Csaki, Y. Grossman, P. Tanedo, Y. Tsai, Phys. Rev. D 83 (2011) 073002 [arXiv:1004.2037 [hep-ph]].
- [39] C. Csaki, C. Delaunay, C. Grojean, Y. Grossman, JHEP 0810 (2008) 055
   [arXiv:0806.0356 [hep-ph]]; F. del Aguila, A. Carmona, J. Santiago, JHEP 1008
   (2010) 127 [arXiv:1001.5151 [hep-ph]].
- [40] G. Cacciapaglia, C. Csaki, J. Galloway, G. Marandella, J. Terning, A. Weiler, JHEP 0804 (2008) 006 [arXiv:0709.1714 [hep-ph]]; M. Redi, Eur. Phys. J. C 72 (2012) 2030 [arXiv:1203.4220 [hep-ph]]; M. König, M. Neubert, D. M. Straub, Eur. Phys. J. C 74 (2014) 7, 2945 [arXiv:1403.2756 [hep-ph]].
- [41] Y. Daikoku, D. Suematsu, Phys. Rev. D 62 (2000) 095006; T. Hambye, E. Ma, M. Raidal, U. Sarkar, Phys. Lett. B 512 (2001) 373; S. F. King, S. Moretti, R. Nevzorov, Phys. Rev. D 73 (2006) 035009; S. F. King, S. Moretti, R. Nevzorov, Phys. Lett. B 634 (2006) 278; S. F. King, S. Moretti, R. Nevzorov, arXiv:hep-ph/0601269; S. F. King, S. Moretti, R. Nevzorov, AIP Conf. Proc. 881 (2007) 138; S. F. King, S. Moretti, R. Nevzorov, Phys. Lett. B 650 (2007) 57; R. Howl, S. F. King, JHEP 0801 (2008) 030; P. Athron, S. F. King, D. J. Miller, S. Moretti, R. Nevzorov, arXiv:0810.0617 [hep-ph]; S. Hesselbach, D. J. Miller, G. Moortgat-Pick, R. Nevzorov and M. Trusov, Phys. Lett. B 662 (2008) 199; P. Athron, J. P. Hall, R. Howl, S. F. King, D. J. Miller, S. Moretti and R. Nevzorov, Nucl. Phys. B Proc. Suppl. 200-202 (2010), 120-129; J. P. Hall, S. F. King, R. Nevzorov, S. Pakvasa and M. Sher, PoS QFTHEP2010 (2010) 069 [arXiv:1012.5365 [hep-ph]]; P. Athron, D. Stockinger, A. Voigt, Phys. Rev. D 86 (2012) 095012; R. Nevzorov, Phys. Rev. D 87 (2013) 015029; P. Athron, M. Binjonaid, S. F. King, Phys. Rev. D 87

(2013) 115023; D. J. Miller, A. P. Morais, P. N. Pandita, Phys. Rev. D 87 (2013) 015007; M. Sperling, D. Stöckinger, A. Voigt, JHEP 1307 (2013) 132; R. Nevzorov, Phys. Rev. D 89 (2014) 5, 055010; R. Nevzorov, S. Pakvasa, Phys. Lett. B 728 (2014) 210; M. Sperling, D. Stöckinger, A. Voigt, JHEP 1401, 068 (2014); P. Athron, M. Mühlleitner, R. Nevzorov, A. G. Williams, JHEP 1501 (2015) 153; P. Athron, D. Harries, R. Nevzorov and A. G. Williams, Phys. Lett. B 760 (2016) 19 [arXiv:1512.07040 [hep-ph]]; P. Athron, D. Harries, R. Nevzorov and A. G. Williams, JHEP 1612 (2016) 128 [arXiv:1610.03374 [hep-ph]]; P. Athron, A. W. Thomas, S. J. Underwood and M. J. White, Phys. Rev. D 95 (2017) 035023 [arXiv:1611.05966 [hep-ph]]; R. Nevzorov, Phys. Lett. B 779 (2018), 223-229 [arXiv:1710.11533 [hepph]]; R. Nevzorov, Int. J. Mod. Phys. A 33 (2018) no.31, 1844007 [arXiv:1805.08260 [hep-ph]]; S. F. King, S. Moretti and R. Nevzorov, Symmetry 12 (2020) no.4, 557 [arXiv:2002.02788 [hep-ph]]; R. Nevzorov, Phys. Atom. Nucl. 83 (2020) no.2, 338-350; R. Nevzorov, Symmetry 14 (2022) no.10, 2090 [arXiv:2209.00505 [hep-ph]]; R. B. Nevzorov, Usp. Fiz. Nauk 193 (2023) no.6, 577-613; R. Nevzorov, Universe 9 (2023) no.3, 137 [arXiv:2304.04629 [hep-ph]].

- [42] K. Agashe, G. Servant, Phys. Rev. Lett. 93 (2004) 231805 [hep-ph/0403143];
   K. Agashe, G. Servant, JCAP 0502 (2005) 002 [hep-ph/0411254].
- [43] G. Bélanger, F. Boudjema, A. Pukhov, A. Semenov, Comput. Phys. Commun. 185 (2014) 960 [arXiv:1305.0237 [hep-ph]].
- [44] J. Aalbers *et al.* [LZ], Phys. Rev. Lett. **131** (2023) no.4, 041002 [arXiv:2207.03764 [hep-ex]].
- [45] E. Aprile et al. [XENON], Phys. Rev. Lett. 131 (2023) no.4, 041003 [arXiv:2303.14729 [hep-ex]].
- [46] E. Aprile et al. [XENON], JCAP 11 (2020), 031 [arXiv:2007.08796 [physics.ins-det]].
- [47] D. S. Akerib *et al.* [LUX-ZEPLIN], Phys. Rev. D **101** (2020) 052002 [arXiv:1802.06039 [astro-ph.IM]].
- [48] G. Aad *et al.* [ATLAS], JHEP **05** (2021), 093 [arXiv:2101.12527 [hep-ex]].
- [49] A. M. Sirunyan et al. [CMS], Eur. Phys. J. C 80 (2020) 3 [arXiv:1909.03460 [hep-ex]].