

MIRROR PARTNER FOR A KLEIN QUARTIC POLYNOMIAL

ALEXEY BASALAEV

ABSTRACT. The results of A. Chiodo, Y. Ruan and M. Krawitz associate the mirror partner Calabi–Yau variety X to a Landau–Ginzburg orbifold (f, G) if f is an invertible polynomial satisfying Calabi–Yau condition and the group G is a diagonal symmetry group of f . In this paper we investigate the Landau–Ginzburg orbifolds with a Klein quartic polynomial $f = x_1^3x_2 + x_2^3x_3 + x_3^3x_1$ and G being all possible subgroups of $\mathrm{GL}(3, \mathbb{C})$, preserving the polynomial f and also the pairing in its Jacobian algebra. In particular, G is not necessarily abelian or diagonal. The zero–set of polynomial f , called Klein quartic, is a genus 3 smooth compact Riemann surface. We show that its mirror Landau–Ginzburg orbifold is (f, G) with G being a $\mathbb{Z}/2\mathbb{Z}$ –extension of a Klein four–group.

1. INTRODUCTION

Mirror symmetry, in one of its formulations, relates a complex variety X to a so–called *Landau–Ginzburg orbifold* (f, G) ([IV90, V89, W93]). In the latter f defines a quasihomogeneous isolated singularity and G is its symmetry group. In particular $f \in \mathbb{C}[x_1, \dots, x_N]$ is a polynomial, whose partial derivatives all vanish simultaneously if and only if $x_1 = \dots = x_N = 0$ and there is a set of natural numbers d_f, d_1, \dots, d_N , s.t. $f(\lambda^{d_1}x_1, \dots, \lambda^{d_N}x_N) = \lambda^{d_f}f(x_1, \dots, x_N)$ for any $\lambda \in \mathbb{C}^*$.

The symmetry group G is a subgroup of $\mathrm{GL}_f := \{g \in \mathrm{GL}(\mathbb{C}^N) \mid f(g \cdot x) = f(x)\}$. The diagonal operator $j_f = \mathrm{diag}(\exp(2\pi\sqrt{-1}\frac{d_1}{d_f}), \dots, \exp(2\pi\sqrt{-1}\frac{d_N}{d_f}))$ belongs to GL_f , showing that GL_f is never trivial. The variety X is assumed to be either quasismooth or a smooth orbifold.

Both X and (f, G) have some essential \mathbb{C} –algebras associated to them. These are the cohomology ring $H^*(X)$ if X is smooth or orbifold cohomology ring $H_{orb}^*(X)$ otherwise (cf. [AGV08]). Note that $H_{orb}^*(X) = H^*(X)$ if X is smooth. To a pair (f, G) one associates the Hochschild cohomology ring $\mathrm{HH}^*(f, G)$ of the category of G –equivariant matrix factorizations $\mathrm{MF}_G(f)$. The variety X is said to be mirror to the Landau–Ginzburg orbifold (f, G) if there is a \mathbb{C} –algebra homomorphism $H_{orb}^*(X) \cong \mathrm{HH}^*(f, G)$.

Date: July 18, 2024.

1.1. Landau-Ginzburg – Calabi-Yau mirror symmetry. Large class of mirror pairs was discovered by Chiodo, Ruan and Krawitz in [CR11, K09]. In particular, let (f, G) be such that

- (1) f is the so-called invertible polynomial. Namely f is a sum of exactly N monomials;
- (2) Calabi–Yau condition holds: $d_1 + \dots + d_N = d_f$;
- (3) G acts diagonally on \mathbb{C}^N and is a subgroup of $\mathrm{SL}(N, \mathbb{C})$.

Then the Berglund–Hübsch–Henningson dual Landau-Ginzburg orbifold (\tilde{f}, \tilde{G}) (see [BH93, BH95, Kru94]) can be used to construct the mirror variety by

$$X_{\tilde{f}, \tilde{G}} := \left\{ (x_1, \dots, x_N) \in \mathbb{P}(d_1, \dots, d_N) \mid \tilde{f}(x_1, \dots, x_N) = 0 \right\} / (\tilde{G} / \langle j_{\tilde{f}} \rangle).$$

Theorem 14 of [CR11] together with Theorem 2.4 of [K09] give that there is a \mathbb{C} -vector space isomorphism $\tau : H_{orb}^*(X_{\tilde{f}, \tilde{G}}) \cong \mathrm{HH}^*(f, G)$. This suggests calling $X_{\tilde{f}, \tilde{G}}$ a mirror pair of (f, G) . Up to now τ is proved to be an algebra homomorphism only for the certain examples (cf. [MR11, MS12, BT3]). The major complication here lies in computing the algebra structure of $H_{orb}^*(X_{\tilde{f}, \tilde{G}})$ while the algebra structure of $\mathrm{HH}^*(f, G)$ was computed explicitly in [BT2, BTW16, BTW17, HLL20, BI22] using the technique developed by D.Shklyarov in [Sh20].

Let's comment on the conditions above. Condition 1 restricts us to work with the invertible polynomials only, for which the construction of Berglund–Hübsch–Henningson can be applied. Condition 2 has the name *Calabi–Yau condition* because it guarantees that the first Chern class of $X_{\tilde{f}, \tilde{G}}$ vanishes. It follows from Condition 3 that $\langle j_{\tilde{f}} \rangle \subseteq \tilde{G}$ — this is the special property of the dual group construction.

1.2. Admissible symmetry groups. The algebra $H_{orb}^*(X_{\tilde{f}, \tilde{G}})$ is endowed with a pairing for any f . However one can introduce an essential pairing on $\mathrm{HH}^*(f, G)$ only if f satisfies the Calabi–Yau condition (cf. [PV12]). Generally in mirror symmetry one considers the groups G , s.t. $G \subseteq \mathrm{SL}(N, \mathbb{C})$. These groups are distinguished by the fact that they preserve volume form of \mathbb{C}^N and therefore the pairing of both algebras.

1.3. Mirror symmetry for curves. More general mirror symmetry results were established in the context toric degenerations and homological algebra. In particular, Ruddat in [R17] and Gross-Katzarkov-Ruddat in [GKR17] investigated the mirrors of the varieties of general type from the point of view of toric degenerations via the relations on the Hodge numbers.

Efimov and Seidel [E12, S11] established an equivalence between the (suitably defined) triangulated categories of a genus $g \geq 2$ surface S_g and Landau–Ginzburg orbifold

(f_g, G) with $f_g = x_1^{2g+1} + x_2^{2g+1} + x_3^{2g+1} + x_1x_2x_3$. Here the group G is generated by the diagonal operator $\text{diag}(\zeta, \zeta, \zeta^{2g-1})$, s.t. $\zeta = \exp(\frac{2\pi\sqrt{-1}}{2g+1})$. Without giving the details of this mirror theorem we quote the following consequence: there is a \mathbb{C} -algebra isomorphism $H^*(S_g) \cong \text{HH}^*(f_g, G)$ (cf. [E12, S11, Sh20]).

1.4. Klein quartic. Consider the polynomial $f = x_1^3x_2 + x_2^3x_3 + x_3^3x_1$. We have $d_1 = d_2 = d_3 = 1$ and $d_f = 4$. It is invertible with the Berglund–Hübsch dual $\tilde{f} = f$. Take $J := \langle j_f \rangle$. Then the variety $X_{\tilde{f}, J}$ is a smooth genus 3 curve S_3 . Its symmetry group is isomorphic to $\text{PSL}(2, 7)$ — simple nonabelian of order 168 (cf. [K79, E98]).

Denote $\text{SL}_f := \text{GL}_f \cap \text{SL}(N, \mathbb{C})$. For any $G \subseteq \text{SL}_f$ denote by \widehat{G} the central extension of G by $\mathbb{Z}/2\mathbb{Z}$ obtained by adding the $-\text{id} \in \text{GL}_f$ to G .

Theorem 1. *Let $f = x_1^3x_2 + x_2^3x_3 + x_3^3x_1$ be a Klein quartic polynomial and S_3 stand for the smooth genus 3 Riemann surface. Then*

- (a) $\dim \text{HH}^*(f, G) > \dim H^*(S_3)$ for any $G \subseteq \text{SL}_f$.
- (b) $\dim \text{HH}^*(f, \widehat{G}) = \dim H^*(S_3)$ for $\widehat{G} \subseteq \widehat{\text{SL}}_f$ if and only if $G \cong V_4$ is conjugate to a Klein four-group.
- (c) *There is a Frobenius algebra isomorphism $\text{HH}^*(f, \widehat{V}_4) \cong H^*(S_3)$.*

Proof. Proof of the theorem occupies Section 4 along with some preparational propositions given in the preceding sections. □

This theorem suggests that the mirror partner of $X_{\tilde{f}, J}$ is (f, \widehat{V}_4) .

Very important consequence of this theorem is the following

Corollary 2. *In mirror symmetry it's not enough to consider only the symmetry groups acting diagonally even for invertible polynomials.*

This is important to stress that the theorem above considers not only diagonal symmetry groups. It also covers some symmetry groups G that are not abelian. Such symmetry groups are not widely considered from the point of view of Landau–Ginzburg orbifolds. The following list of citations is pretty much the full list of publications assuming nonabelian orbifolds [M, BI21, BI22, BI24, WWP, CJMPW23, EGZ20, EGZ18]. And even in this list (except [BI24]) one considers only the semidirect product groups $G^d \rtimes S$ with G^d acting diagonally and S being a subgroup of a symmetric group. To our knowledge, no investigation of a simple group was made before except [BI24].

1.5. Acknowledgements. The work of Alexey Basalaev was supported by the Theoretical Physics and Mathematics Advancement Foundation "BASIS".

2. NOTATIONS AND TERMINOLOGIES

For a non-negative integer N and a polynomial $f = f(x_1, \dots, x_N) \in \mathbb{C}[x_1, \dots, x_N]$, the *Jacobian algebra* $\text{Jac}(f)$ of f is a \mathbb{C} -algebra defined as

$$\text{Jac}(f) = \mathbb{C}[x_1, \dots, x_N] \Big/ \left(\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_N} \right). \quad (1)$$

The polynomial f is said to define an isolated singularity if $\text{Jac}(f)$ is a finite-dimensional. Then set $\mu_f := \dim_{\mathbb{C}} \text{Jac}(f)$ and call it the *Milnor number* of f . For $N = 0$ set $\text{Jac}(f) := \mathbb{C}$ with $\mu_f = 1$.

Let the *Hessian* of f be defined as the following determinant:

$$\text{hess}(f) := \det \left(\frac{\partial^2 f}{\partial x_i \partial x_j} \right)_{i,j=1,\dots,N}. \quad (2)$$

Its class is nonzero in $\text{Jac}(f)$ giving the \mathbb{C} -bilinear nondegenerate pairing η_f called *residue pairing* (see [GH94, Chapter 5], [AGV85, Section 5.11]). The value $\eta_f([u], [v])$ is taken as the projection of the product $[u][v]$ to the \mathbb{C} -span of $[\text{hess}(f)]$. In particular, we fix

$$\eta_f([1], [\text{hess}(f)]) = 1.$$

Remark 3. One notes immediately that under the coordinate-wise action of $g \in \text{GL}_f$ we have $g \cdot \text{hess}(f) = (\det g)^2 \text{hess}(f)$. This shows that the groups $G \subseteq \{g \in \text{GL}_f \mid \det g = \pm 1\}$ preserve the pairing η_f . We will comment more on this in Proposition 8.

The \mathbb{C} -algebra (A, \circ) with a non-degenerate bilinear form η_A is called a *Frobenius algebra* if $\eta_A(u, v \circ w) = \eta_A(u \circ v, w)$ for all $u, v, w \in A$. The pair $(\text{Jac}(f), \eta_f)$ is an example of a Frobenius algebra.

2.1. Symmetries. Given a quasihomogeneous polynomial $f = f(x_1, \dots, x_N)$ consider the *maximal group of linear symmetries* of f defined by

$$\text{GL}_f := \{g \in \text{GL}(N, \mathbb{C}) \mid f(g \cdot \mathbf{x}) = f(\mathbf{x})\}.$$

Remark 4. This group contains the so-called maximal group of *diagonal* symmetries of f defined by $G_f^d := \{(\lambda_1, \dots, \lambda_N) \in (\mathbb{C}^*)^N \mid f(\lambda_1 x_1, \dots, \lambda_N x_N) = f(x_1, \dots, x_N)\}$.

For each $g \in \text{GL}_f$, denote by $\text{Fix}(g)$ the *fixed locus* of g

$$\text{Fix}(g) := \{(x_1, \dots, x_N) \in \mathbb{C}^N \mid g \cdot (x_1, \dots, x_N) = (x_1, \dots, x_N)\}. \quad (3)$$

This is an eigenvalue 1 subspace of g and therefore a linear subspace of \mathbb{C}^N . By $N_g := \dim_{\mathbb{C}} \text{Fix}(g)$ denote its dimension and by $f^g := f|_{\text{Fix}(g)}$ the restriction of f to the fixed

locus of g . For $g \in G_f^d$ this linear subspace is furthermore a span of a collection of standard basis vectors.

Proposition 5. *For any $g \in \mathrm{GL}_f$ with $N_g > 0$ there is a choice of coordinates on $\mathrm{Fix}(g)$ linear in x_i , such that the polynomial f^g defines a quasihomogeneous singularity.*

Proof. Let $\tilde{x}_1, \dots, \tilde{x}_{N_g}, \tilde{x}_{N_g+1}, \dots, \tilde{x}_N$ be the coordinates of \mathbb{C}^N , s.t. $\tilde{x}_1, \dots, \tilde{x}_{N_g}$ are dual to the eigenspace 1 basis vectors of g . In this coordinates we have $f^g = f|_{\tilde{x}_{N_g+1}=\dots=\tilde{x}_N=0}$. The proof follows now by the same argument as in Proposition 5 of [ET13]. \square

Denote also

$$\mathrm{SL}_f := \mathrm{GL}_f \cap \mathrm{SL}(N, \mathbb{C}).$$

This group will be important later on because it preserves the volume form of \mathbb{C}^N .

3. PHASE SPACE

This section is devoted to the definition of $\mathrm{HH}^*(f, G)$ and also some computational aspects of it.

3.1. The total space. For each $g \in \mathrm{GL}_f$ fix a generator of a one-dimensional vector space $\Lambda(g) := \bigwedge^{N-N_g}(\mathbb{C}^N/\mathrm{Fix}(g))$. Denote it by ξ_g .

Define $\mathcal{B}_{tot}(f)$ to be the \mathbb{C} -vector spaces of dimension $\sum_{g \in \mathrm{GL}_f} \dim \mathrm{Jac}(f^g)$

$$\mathcal{B}_{tot}(f) := \bigoplus_{g \in \mathrm{GL}_f} \mathrm{Jac}(f^g)\xi_g, \quad (4)$$

Each direct summand $\mathrm{Jac}(f^g)\xi_g$ will be called the g -th sector and denoted by \mathcal{B}'_g . Any $h \in \mathrm{GL}_f$ induces a map

$$h: \mathrm{Fix}(g) \rightarrow \mathrm{Fix}(hgh^{-1}).$$

and hence

$$h: \Lambda(g) \rightarrow \Lambda(hgh^{-1}).$$

Since we have fixed the generators ξ_\bullet , the latter map provides a constant $\rho_{h,g} \in \mathbb{C}^*$ such that

$$h(\xi_g) = \rho_{h,g}\xi_{hgh^{-1}}.$$

Then $\rho_{h_2, h_1 g h_1^{-1}} \rho_{h_1, g} = \rho_{h_2 h_1, g}$. Define the action of GL_f on \mathcal{B}_{tot} by

$$h^*([p(\mathbf{x})]\xi_g) = \rho_{h,g}[p(h^{-1} \cdot \mathbf{x})]\xi_{hgh^{-1}}. \quad (5)$$

For a finite $G \subseteq \mathrm{GL}_f$ put

$$\mathcal{B}_{tot, G} := \bigoplus_{g \in G} \mathcal{B}'_g = \bigoplus_{g \in G} \mathrm{Jac}(f^g)\xi_g \subseteq \mathcal{B}_{tot}$$

and define as a vector space

$$\mathrm{HH}^*(f, G) := (\mathcal{B}_{tot, G})^G. \quad (6)$$

Namely, the linear span of the \mathcal{B}_{tot} vectors that are invariant with respect to the action of all elements of G .

The latter direct sum and the group action can be intertwined as follows. Let \mathcal{C}^G stand for the set of representatives of the conjugacy classes of G and $Z(g)$ for the centralizers of $g \in G$. Denote $\mathcal{B}_g := (\mathcal{B}'_g)^{Z(g)}$. Then we have

$$\mathrm{HH}^*(f, G) \cong \bigoplus_{g \in \mathcal{C}^G} \mathcal{B}_g = \bigoplus_{g \in \mathcal{C}^G} (\mathcal{B}'_g)^{Z(g)}. \quad (7)$$

cf. [BI22, Proposition 42] for the proof.

It's much more complicated to define the algebra structure of $\mathrm{HH}^*(f, G)$ and we don't do it here because it will not be needed later on. We will only need that the product \circ of $\mathrm{HH}^*(f, G)$ respects the G -grading: $\circ : \mathcal{B}_g \otimes \mathcal{B}_h \rightarrow \mathcal{B}_{gh}$, its unit is $1\xi_{\mathrm{id}} \in \mathcal{B}_{\mathrm{id}}$ and $\circ : \mathcal{B}_{\mathrm{id}} \otimes \mathcal{B}_{\mathrm{id}} \rightarrow \mathcal{B}_{\mathrm{id}}$ coincides with the product of $(\mathrm{Jac}(f))^G$.

3.2. Computational statements. The following proposition follows immediately from the definition but will be very important later on.

Proposition 6. *Let f define an isolated singularity. Then we have*

- (i): *Let $g \in \mathrm{GL}_f$ then $g^*(\xi_{g^k}) = \det g \cdot \xi_{g^k}$ for any k , s.t. $g^k \neq \mathrm{id}$.*
- (ii): *Let $G \subseteq \mathrm{GL}_f$ be abelian. Then $\mathrm{HH}^*(f, G) = \bigoplus_{g \in G} (\mathrm{Jac}(f^g)\xi_g)^G$.*

Proof. This is immediate. □

Proposition 7. *Let $G_1, G_2 \subseteq \mathrm{GL}_f$ be conjugate. Then $\mathrm{HH}^*(f, G_1) \cong \mathrm{HH}^*(f, G_2)$.*

Proof. Let $G_2 = \tilde{g}G_1\tilde{g}^{-1}$ for some $\tilde{g} \in \mathrm{GL}(\mathbb{C}^N)$. Consider the decomposition of Eq.(7) for both groups. We show that

$$\Psi : [\phi(x)]\xi_h \mapsto [\phi(g \cdot x)]\xi_{\tilde{g}h\tilde{g}^{-1}}, \quad \forall h \in G_1$$

establishes the isomorphism $\mathrm{HH}^*(f, G_1) \xrightarrow{\sim} \mathrm{HH}^*(f, G_2)$.

Obviously $\mathcal{C}^{G_2} = \{\tilde{g}r\tilde{g}^{-1} \mid r \in \mathcal{C}^{G_1}\}$ and $Z(\tilde{g}h\tilde{g}^{-1}) = \tilde{g}Z(h)\tilde{g}^{-1}$. Note that we have the isomorphism $\mathrm{Fix}(h) \xrightarrow{\sim} \mathrm{Fix}(\tilde{g}h\tilde{g}^{-1})$. This gives $\mathcal{B}'_h = \mathcal{B}'_{\tilde{g}h\tilde{g}^{-1}}$.

In order to finish the proof it remains to show that the invariants under the action of $Z(h)$ and $Z(\tilde{g}h\tilde{g}^{-1})$ agree. Let $r = \tilde{g}s\tilde{g}^{-1} \in \tilde{g}Z(h)\tilde{g}^{-1}$. Then we have

$$r^*\xi_{\tilde{g}h\tilde{g}^{-1}} = \rho \cdot \xi_{\tilde{g}s\tilde{g}^{-1}} \quad \text{and} \quad s^*\xi_{\tilde{g}h\tilde{g}^{-1}} = \rho' \cdot \xi_s$$

for some $\rho, \rho' \in \mathbb{C}^*$. The following computation shows that $\rho = \rho'$.

$$r^* \xi_{\tilde{g}h\tilde{g}^{-1}} = (\tilde{g}s\tilde{g}^{-1})^* \xi_{\tilde{g}h\tilde{g}^{-1}} = s^* \xi_h$$

where the last equality follows from the fact that s acts on $\Lambda(h)$ by the same multiple as $\tilde{g}s\tilde{g}^{-1}$ acts on $\Lambda(\tilde{g}h\tilde{g}^{-1})$. \square

The following proposition shows that the pairing restricts to the invariants of $G \subseteq \widehat{\text{SL}}_f$.

Proposition 8. *Let $G \subset \text{GL}_f$ be s.t. $\det g = \pm 1$ for any $g \in G$. Consider the polynomials $\phi_1(\mathbf{x}), \phi_2(\mathbf{x}) \in \mathbb{C}[\mathbf{x}]$, s.t. $\eta_f([\phi_1], [\phi_2]) \neq 0$.*

Then if $[\phi_1] \neq 0$ in $(\text{Jac}(f))^G$, then $[\phi_2] \neq 0$ in $(\text{Jac}(f))^G$ as well.

Proof. Any $g \in \text{GL}_f$ acts on the hessian polynomial by $g(\text{hess}(f)) = (\det g)^2 \text{hess}(f)$. Let $\eta_f([\phi_1], [\phi_2]) = \alpha \in \mathbb{C}^*$. The following equality holds in $\mathbb{C}[\mathbf{x}]$ for some polynomials p_1, \dots, p_N

$$\phi_1(\mathbf{x}) \cdot \phi_2(\mathbf{x}) - \alpha \text{hess}(f) = \sum_{k=1}^N p_k(\mathbf{x}) \frac{\partial f}{\partial x_k}.$$

The Jacobian ideal of f is preserved under the G -action. Hence under the assumption of the proposition we have

$$[\phi_1(\mathbf{x})] \cdot [\phi_2(\mathbf{x})] = \alpha [\text{hess}(f)] \neq 0 \in (\text{Jac}(f))^G$$

what completes the proof. \square

3.3. Jac(f, G) for a Klein quartic. The following propositions are specific for f being the Klein quartic polynomial. We consider in details the cases of $G \subseteq \text{GL}_f$ with $\det g = 1$ and $(\det g)^2 = 1$.

3.3.1. *Consider $g \in G \setminus \{\text{id}\}$ with $\det g = 1$. The fixed locus of g can be either 0 or 1-dimensional. If $\text{Fix}(g) = 0$, any element $h \in Z(g)$ acts on ξ_g by multiplication by $\det h$. In particular, ξ_g is G -invariant if $G \subset \text{SL}(3, \mathbb{C})$ and is not G -invariant if $-\text{id} \in G$.*

If $\text{Fix}(g) \cong \mathbb{C}$, we have $f^g = \tilde{x}^4$ where \tilde{x} is a coordinate of $\text{Fix}(g)$. This gives $\mathcal{B}'_g \cong \mathbb{C}\langle [1], [\tilde{x}], [\tilde{x}^2] \rangle \xi_g$. Let $h \in Z(g)$ act on \tilde{x} by $\rho \in \mathbb{C}^*$. Namely $h(\tilde{x}) = \rho \tilde{x}$. Then we have

$$h^*([\tilde{x}^k] \xi_g) = \frac{\det h}{\rho^{k+1}} \cdot [\tilde{x}^k] \xi_g.$$

In order to compute the direct sum contribution of g to $\text{HH}^*(f, G)$ it remains to check the multiples ρ of all the elements of $Z(g)$.

3.3.2. *Consider $g \in G$ with $\det g = -1$.* The fixed locus of g can be either 0, 1–dimensional or 2–dimensional. The first two cases are treated exactly in the same way as above and we concentrate on the last one. Let \tilde{x}_1, \tilde{x}_2 be the coordinates on $\text{Fix}(g)$. Then f^g is quasihomogeneous with weights $(1/4, 1/4)$. After some rescaling of the variables f^g can either be $\tilde{x}_1^4 + \tilde{x}_2^4$, $\tilde{x}_1^3\tilde{x}_2 + \tilde{x}_2^4$ or $\tilde{x}_1^3\tilde{x}_2 + \tilde{x}_2^3\tilde{x}_1$.

We have $\mathcal{B}'_g \cong \mathbb{C}^9$ generated by the classes $[\tilde{x}_1^{a_1}\tilde{x}_2^{a_2}]$ for some pairs (a_1, a_2) . Since $\det g = -1$, we have $g^*(\xi_g) = -\xi_g$ and $g^*([\tilde{x}_1^{a_1}\tilde{x}_2^{a_2}]\xi_g) = -[\tilde{x}_1^{a_1}\tilde{x}_2^{a_2}]\xi_g$.

3.3.3. *Consider $g = \text{id}$.* While computing the vector space $\text{HH}^*(f, G)$ one has to find the G –invariants of $\text{Jac}(f)\xi_{\text{id}}$. We have $h^*(\xi_{\text{id}}) = \xi_{\text{id}}$ and it remains to compute the G –invariants of $\text{Jac}(f)$ itself. This goal can be achieved by symmetrization procedure. Namely, if $\{\phi_k(\mathbf{x})\}_{k=1}^\mu$ are the monomials whose classes generate $\text{Jac}(f)$, then the classes of $\{\sum_{g \in G} \phi_k(g \cdot \mathbf{x})\}_{k=1}^\mu$ generate $(\text{Jac}(f))^G$. However, some of these classes might be zero or linearly dependant.

The basis monomials of ϕ_\bullet can be taken to be $x_1^{a_1}x_2^{a_2}x_3^{a_3}$, s.t. $\sum_k a_k \leq 6$ and $0 \leq a_k \leq 2$. Also $[\text{hess}(f)] = 756[(x_1x_2x_3)^2]$. Assign the grading $\sum_k a_k$ to each of these monomials. The vector space $\text{Jac}(f)$ decomposes into the direct sum of the graded pieces. The dimensions of these pieces can be composed into the following vector

$$1, 3, 6, 7, 6, 3, 1.$$

This vector is symmetric by its middle point — the 4–th component because the pairing η_f is non–degenerate.

The grading introduced is preserved under the coordinate–wise action of $g \in \text{GL}_f$. Similarly $(\text{Jac}(f))^G$ decomposes into the direct sum of the graded pieces and the dimensions of these pieces can be written by a vector. This vector will be symmetric by its middle point due to Proposition 8.

4. ADMISSIBLE SYMMETRY GROUPS

This section gathers the computations of $\text{HH}^*(f, G)$ for a Klein quartic polynomial f and all admissible symmetry groups $G \subset \text{GL}_f$. Namely, the groups G , s.t. $\det g = \pm 1$ for any $g \in G$.

4.1. **The subgroups.** The group $\text{SL}(f)$ is generated by the following $\text{SL}(3, \mathbb{C})$ elements

$$R = \frac{\sqrt{-1}}{\sqrt{7}} \begin{pmatrix} \zeta - \zeta^6 & \zeta^2 - \zeta^5 & \zeta^4 - \zeta^3 \\ \zeta^2 - \zeta^5 & \zeta^4 - \zeta^3 & \zeta - \zeta^6 \\ \zeta^4 - \zeta^3 & \zeta - \zeta^6 & \zeta^2 - \zeta^5 \end{pmatrix}, \quad T = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad S = \begin{pmatrix} \zeta^4 & 0 & 0 \\ 0 & \zeta^2 & 0 \\ 0 & 0 & \zeta \end{pmatrix}.$$

where $\zeta = \exp(2\pi\sqrt{-1}/7)$. This was observed already by F.Klein [K79]. The following list gives the classification of the subgroups of $\mathrm{SL}(f)$ up to conjugation

- (a) 8 conjugate elementary abelian groups of order 7,
- (b) 28 conjugate cyclic groups of order 3,
- (c) 21 conjugate cyclic groups of order 4,
- (d) 21 conjugate cyclic groups of order 2,
- (e) two classes of 7 conjugate dihedral abelian Klein 4-groups of order 4,
- (f) 28 dihedral nonabelian groups of order 6,
- (g) 21 dihedral nonabelian groups of order 8,
- (h) 8 nonabelian groups of order 21,
- (i) two classes of 7 nonabelian conjugates of the symmetric group of degree 4,
- (j) two classes of 7 nonabelian conjugates of the alternating group of degree 4.

For any group G from the list above we also consider its extension by $\mathbb{Z}/2\mathbb{Z}$ obtained by adding an element $-\mathrm{id} \in \mathrm{GL}(3, \mathbb{C})$. Denote such group by \widehat{G} . Obviously, if g_1, \dots, g_r are all the representatives of the conjugacy classes of G , then $\pm g_1, \dots, \pm g_r$ are the representatives of the conjugacy classes of \widehat{G} .

In the following computations we make use of different statements of Section 3 without quoting them each time. In particular, we make use of Eq. (7) to find $\mathrm{HH}^*(f, G)$ and Section 3.2,3.3 to compute the G -invariants. In particular, for any G beneath we find the basis of $\mathrm{HH}^*(f, G)$.

For the $\mathbb{Z}/2\mathbb{Z}$ extended groups \widehat{G} only the computations of \widehat{V}_4 and $\widehat{\mathrm{SL}}_f$ are given explicitly. The computations of all the other groups are derived easily by using the statements of Section 3.

4.2. The group SL_f . Let $G := \langle R, T, S \rangle$ be the full group SL_f . It is well-known to have 6 conjugacy classes of orders 1, 24, 24, 21, 42, 56. We have

$$\mathcal{C}^G = \{\mathrm{id}, S, S^3, R, RS^3, T\}.$$

The centralizers are $Z(S) = Z(S^3) = \langle S \rangle$, $Z(T) = \langle T \rangle$, $Z(R) = \langle TRSR S^5, TRS^3RS^6 \rangle$ and $Z(RS^3) = \langle RS^3 \rangle$.

The elements S and S^3 don't have 1 as an eigenvalue, while all other do. The multiples of the centralizers are ± 1 for R , and all equal to 1 for T and RS^3 . This gives

$$\mathcal{B}_S = \mathbb{C}\langle \xi_S \rangle, \quad \mathcal{B}_{S^3} = \mathbb{C}\langle \xi_{S^3} \rangle, \quad \mathcal{B}_R = \mathbb{C}\langle [\tilde{x}]\xi_R \rangle \quad (8)$$

$$\mathcal{B}_{RS^3} = \mathbb{C}\langle \xi_{RS^3}, [\tilde{x}]\xi_{RS^3}, [\tilde{x}^2]\xi_{RS^3} \rangle, \quad \mathcal{B}_T = \mathbb{C}\langle \xi_T, [\tilde{x}]\xi_T, [\tilde{x}^2]\xi_T \rangle \quad (9)$$

The identity sector \mathcal{B}_{id} is spanned by

$$1\xi_{\text{id}} \quad \text{and} \quad [5x_2^2x_3^2x_1^2 - x_3x_1^5 - x_1x_2^5 - x_2x_3^5]\xi_{\text{id}}.$$

Both classes are G -invariant and non-zero in $\text{Jac}(f)$.

$\text{HH}^*(f, G)$ has $2 + 9 = 11$ basis elements in total.

4.3. The group (a). Let G be generated by S . This is an order 7 group and $\text{Fix}(g) = 0$ for any nonidentical element. This gives $\mathcal{B}_g = \mathbb{C} \cdot \xi_g$ for any such g .

In the identity sector the space \mathcal{B}_{id} is spanned by the elements

$$\xi_{\text{id}}, \quad [x_1x_2x_3]\xi_{\text{id}}, \quad [(x_1x_2x_3)^2]\xi_{\text{id}}.$$

$\text{HH}^*(f, G)$ has $3 + 6 = 9$ basis elements in total.

4.4. The group (b). Let G be generated by T . We have $\mathcal{C}^G = \{\text{id}, T, T^2\}$ and 1 is an eigenvalue of T, T^2 . This gives

$$\mathcal{B}_g = \mathbb{C}\langle \xi_g, [(x_1 + x_2 + x_3)]\xi_g, [(x_1 + x_2 + x_3)^2]\xi_g \rangle, \quad g = T, T^2.$$

The identity sector basis is given by 11 elements $[\phi(x)]\xi_{\text{id}}$ with $\phi(x)$ taken from the list below

$$\begin{aligned} &1, \quad x_1 + x_2 + x_3, \quad x_1^2 + x_2^2 + x_3^2, x_2x_3 + x_1x_2 + x_1x_3, \quad x_3x_1^2 + x_2^2x_1 + x_2x_3^2, x_1x_2x_3, \\ &x_2x_1^2 + x_3^2x_1 + x_2^2x_3, \quad x_1x_2x_3(x_1 + x_2 + x_3), \quad x_1^2x_2^2 + x_1^2x_3^2 + x_2^2x_3^2, \\ &x_1x_2x_3(x_2x_3 + x_1x_2 + x_1x_3), \quad x_1^2x_2^2x_3^2. \end{aligned}$$

The corresponding dimension vector reads

$$1, 1, 2, 3, 2, 1, 1.$$

$\text{HH}^*(f, G)$ has $11 + 2 \cdot 3 = 17$ basis elements in total.

4.5. The group (c). Let G be generated by $g := RSR S^5$. The generator has 1 as an eigenvalue, $\mathcal{C}^G = G$ and

$$\mathcal{B}_{g^k} \cong \mathbb{C}\langle [1]\xi_{g^k}, [\tilde{x}]\xi_{g^k}, [\tilde{x}^2]\xi_{g^k} \rangle, \quad k = 1, 2, 3.$$

The identity sector basis is given by 9 elements $[\phi(x)]\xi_{\text{id}}$ with the dimension vector

$$1, 1, 2, 1, 2, 1, 1.$$

The polynomials $\phi(x)$ can be taken to be the symmetrizations of

$$1, \quad x_1, \quad x_3^2, x_2x_3, \quad x_2x_3^2, \quad x_1^2x_2^2, x_1^2x_2x_3, \quad x_1(x_2x_3)^2, \quad (x_1x_2x_3)^2.$$

$\mathrm{HH}^*(f, G)$ has $9 + 3 \cdot 3 = 18$ basis elements in total.

4.6. The group (d). Let G be generated by $g = RT$. The generator has 1 as an eigenvalue. We have $\mathcal{C}^G = G$ and $\mathcal{B}_g \cong \mathbb{C}^3$.

The identity sector basis is given by 15 elements $[\phi(x)]\xi_{\mathrm{id}}$ with the dimension vector

$$1, 1, 4, 3, 4, 1, 1.$$

The polynomials $\phi(x)$ can be taken to be the symmetrizations of

$$1, x_1, x_3^2, x_2x_3, x_2^2, x_1x_3, x_1x_2x_3, x_1x_3^2, x_2x_3^2, x_1^2x_2^2, x_1^2x_2x_3, x_1^2x_3^2, x_1x_2^2x_3, x_1(x_2x_3)^2, (x_1x_2x_3)^2.$$

$\mathrm{HH}^*(f, G)$ has $15 + 3 = 18$ basis elements in total.

4.7. The group (e) - Klein 4-group. Let $G := \{\mathrm{id}, RS^2RS, SRS^6, S^2RS^3RS\}$. It's not hard to see that G is indeed homomorphic to a Klein four-group. All nontrivial group elements have eigenvalues $(1, -1, -1)$.

We have $\mathcal{C}^G = G$ and the centralizers being all G . For the nonidentical elements g , the multiples of the centralizers are ± 1 and therefore

$$\mathcal{B}_g = \mathbb{C}\langle[\tilde{x}]\rangle\xi_g \quad \forall g \in G \setminus \{\mathrm{id}\}.$$

The identity sector basis is given by 9 elements $[\phi(x)]\xi_{\mathrm{id}}$ with the dimension vector

$$1, 0, 3, 1, 3, 0, 1.$$

The polynomials $\phi(x)$ can be taken to be the symmetrizations of

$$1, x_3^2, x_2x_3, x_2^2, x_2x_3^2, x_1^2x_2^2, x_1^2x_2x_3, x_1^2x_3^2, (x_1x_2x_3)^2.$$

$\mathrm{HH}^*(f, G)$ has $9 + 3 = 12$ basis elements in total.

4.8. The group (f) - D_6 . Let G be generated by $a = T$ and $b = R$.

We have $\mathcal{C}^G = \{\mathrm{id}, a, b\}$. All these elements have 1 as an eigenvalue. The corresponding centralizers are $Z(a) = \langle a \rangle$, $Z(b) = \langle b \rangle$.

The multiples of the centralizers are all equal to 1 in both cases. Due to this we have

$$\mathcal{B}_a = \mathbb{C}\langle\xi_a, [\tilde{x}]\xi_a, [\tilde{x}^2]\xi_a\rangle, \quad \mathcal{B}_b = \mathbb{C}\langle\xi_b, [\tilde{x}]\xi_b, [\tilde{x}^2]\xi_b\rangle.$$

The identity sector basis is given by 7 elements $[\phi(x)]\xi_{\mathrm{id}}$ with the dimension vector

$$1, 0, 2, 1, 2, 0, 1.$$

The polynomials $\phi(x)$ can be taken to be the symmetrizations of

$$1, x_3^2, x_2x_3, x_1x_2x_3, x_1^2x_2^2, x_1^2x_2x_3, (x_1x_2x_3)^2.$$

$\text{HH}^*(f, G)$ has $7 + 2 \cdot 3 = 13$ basis elements in total.

4.9. **The group (g) - D_8 .** Let G be generated by $a = RS^3$ and $b = RS^2RS$.

We have $\mathcal{C}^G = \{\text{id}, a^2, b, ab, a\}$. All these elements have 1 as an eigenvalue. The corresponding centralizers are $Z(a^2) = G$, $Z(b) = \langle a^2, b \rangle$, $Z(ab) = \langle a^2, ab \rangle$, $Z(a) = \langle a \rangle$.

The multiples of the centralizers are ± 1 for a^2 , b and ab . The multiples of the centralizers for a are $\pm\sqrt{-1}, 1$. Due to this we have

$$\mathcal{B}_{a^2} = \mathbb{C}\langle[\tilde{x}]\xi_{a^2}\rangle, \mathcal{B}_b = \mathbb{C}\langle[\tilde{x}]\xi_b\rangle, \mathcal{B}_{ab} = \mathbb{C}\langle[\tilde{x}]\xi_{ab}\rangle, \mathcal{B}_a = 0.$$

The identity sector basis is given by 6 elements $[\phi(x)]\xi_{\text{id}}$ with the dimension vector

$$1, 0, 2, 0, 2, 0, 1.$$

The polynomials $\phi(x)$ can be taken to be the symmetrizations of

$$1, \quad x_3^2, x_2x_3, \quad x_1^2x_2^2, x_1^2x_2x_3, \quad (x_1x_2x_3)^2.$$

$\text{HH}^*(f, G)$ has $6 + 3 = 9$ basis elements in total.

4.10. **The group (h).** Let G be generated by T and S .

We have $\mathcal{C}^G = \{\text{id}, T, T^2, S, S^2\}$. Both S and S^2 do not have 1 as an eigenvalue and therefore give one-dimensional sectors. The elements T and T^2 both have 1 as an eigenvalue and the centralizers in both cases are $\langle T \rangle \subset G$. We have

$$\mathcal{B}_S = \mathbb{C} \cdot \xi_S, \mathcal{B}_{S^2} = \mathbb{C} \cdot \xi_{S^2}, \mathcal{B}_T = \mathbb{C}\langle\xi_T, [\tilde{x}]\xi_T, [\tilde{x}^2]\xi_T\rangle, \mathcal{B}_{T^2} = \mathbb{C}\langle\xi_{T^2}, [\tilde{x}]\xi_{T^2}, [\tilde{x}^2]\xi_{T^2}\rangle$$

The identity sector basis is given by 3 elements $[\phi(x)]\xi_{\text{id}}$ with the dimension vector

$$1, 0, 0, 1, 0, 0, 1.$$

The polynomials $\phi(x)$ can be taken to be

$$1, x_1x_2x_3, (x_1x_2x_3)^2.$$

$\text{HH}^*(f, G)$ has $3 + 8 = 11$ basis elements in total.

4.11. **The group (i).** Consider

$$c_2 := TS^5RS^6, \quad c_3 := TS^4, \quad c_4 := TRS^2RS^3, \quad v_4 := T^2RS^6RS^4.$$

These elements have orders 2, 3, 4, 2 respectively. The first three represent 2-cycle, 3-cycle and 4-cycle in S_4 embedded in SL_f as a group $G := \langle c_2, c_4 \rangle$. The element v_4 above is the Klein 4-group element of G .

We have $\mathcal{C}^G = \{\text{id}, c_2, c_3, c_4, v_4\}$. All the elements listed have 1 as an eigenvalue. The centralizers are $Z(c_2) = \{\text{id}, RSR^2, RS^4RS^4, RS^5RS^6\}$ giving the multiples ± 1 ; $Z(c_3) = \{\text{id}, TS^4, T^2S^5\}$ giving the multiples 1; $Z(c_4) = \{\text{id}, TRSR^5, TRS^2RS^3, RS^6RS\}$ giving the multiples 1; $Z(v_4) = \{\text{id}, RS^4RS^3, R, T^2RS^2RS^6, T^2RS^3RS, T^2RS^5RS^2, T^2RS^6RS^4, TRSR^5\}$ giving the multiples ± 1 . We have

$$\mathcal{B}_{c_3} = \mathbb{C}\langle \xi_{c_3}, [\tilde{x}_1]\xi_{c_3}, [\tilde{x}_1^2]\xi_{c_3} \rangle, \quad \mathcal{B}_{c_4} = \mathbb{C}\langle \xi_{c_4}, [\tilde{x}_1]\xi_{c_4}, [\tilde{x}_1^2]\xi_{c_4} \rangle, \quad \mathcal{B}_{c_2} = \mathbb{C}\langle [\tilde{x}_1]\xi_{c_2} \rangle, \quad \mathcal{B}_{v_4} = \mathbb{C}\langle [\tilde{x}_1]\xi_{v_4} \rangle.$$

The identity sector basis is given by 4 elements $[\phi(x)]\xi_{\text{id}}$ with the dimension vector

$$1, 0, 1, 0, 1, 0, 1.$$

The polynomials $\phi(x)$ can be taken to be

$$1, x_3^2, (x_1x_2)^2, (x_1x_2x_3)^2.$$

$\text{HH}^*(f, G)$ has $4 + 8 = 12$ basis elements in total.

4.12. The group (j). Consider the subgroup $A_4 \subset S_4$ for S_4 embedded in SL_f as in the case above. Let the notation be as above.

We have $\mathcal{C}^G = \{\text{id}, c_3, v_4\}$. All the elements listed have 1 as an eigenvalue. The centralizers are $Z(v_4) = \{\text{id}, T^2RS^3RS, T^2RS^5RS^2, T^2RS^6RS^4\}$ giving the multiples ± 1 ; $Z(c_3) = \{\text{id}, TS^4, T^2S^5\}$ giving the multiple 1 exclusively. We have

$$\mathcal{B}_{c_4} = \mathbb{C}\langle \xi_{c_4}, [\tilde{x}_1]\xi_{c_4}, [\tilde{x}_1^2]\xi_{c_4} \rangle, \quad \mathcal{B}_{v_4} = \mathbb{C}\langle [\tilde{x}_1]\xi_{v_4} \rangle.$$

The identity sector basis is given by 10 elements $[\phi(x)]\xi_{\text{id}}$ with the dimension vector

$$1, 0, 3, 2, 3, 0, 1.$$

The polynomials $\phi(x)$ can be taken to be

$$1, x_3^2, x_2x_3, x_2^2, x_1x_3^2, x_2x_3^2, (x_1x_2)^2, x_1^2x_2x_3, (x_1x_3)^2, (x_1x_2x_3)^2.$$

$\text{HH}^*(f, G)$ has $10 + 4 = 14$ basis elements in total.

4.13. $\mathbb{Z}/2\mathbb{Z}$ -extension of the Klein 4-group. In this section denote $\widehat{G} := \{\pm g \mid g \in G\}$ with $g \in G := \{\text{id}, RS^2RS, SRS^6, S^2RS^3RS\}$.

Then for any $g \in G$ the eigenvalues of g are $(1, -1, -1)$ and those of $-g$ are $(-1, 1, 1)$. We get $\text{Fix}(-g) \cong \mathbb{C}^2$, however the sector $\mathcal{B}_{-g} = 0$ as explained in Section 3.3.2. Similarly $\mathcal{B}_g = (\mathbb{C}\langle [\widehat{x}] \rangle \xi_g)^{\widehat{G}} = 0$ because $[\widehat{x}]\xi_g$ is not invariant under the action of $-\text{id}$.

We conclude that

$$\text{HH}^*(f, \widehat{G}) = (\text{Jac}(f)\xi_{\text{id}})^{\widehat{G}} \cong (\text{Jac}(f))^{\widehat{G}} \cdot \xi_{\text{id}}.$$

By using the computations done for the group G we have

$$\mathrm{HH}^*(f, \widehat{G}) \cong \mathbb{C}\langle [1], [x_3^2], [x_2x_3], [x_2^2], [x_1^2x_2^2], [x_1^2x_2x_3], [x_1^2x_3^2], [(x_1x_2x_3)^2] \rangle \xi_{\mathrm{id}}.$$

It is 8–dimensional with the product restricted from $\mathrm{Jac}(f)$. The only non–zero products that do not involve the element $1\xi_{\mathrm{id}}$ are

$$\begin{aligned} [x_2^2]\xi_{\mathrm{id}} \circ [x_1^2x_3^2]\xi_{\mathrm{id}} &= [(x_1x_2x_3)^2]\xi_{\mathrm{id}}, & [x_3^2]\xi_{\mathrm{id}} \circ [x_1^2x_2^2]\xi_{\mathrm{id}} &= [(x_1x_2x_3)^2]\xi_{\mathrm{id}} \\ [x_2x_3]\xi_{\mathrm{id}} \circ [x_1^2x_2x_3]\xi_{\mathrm{id}} &= [(x_1x_2x_3)^2]\xi_{\mathrm{id}}. \end{aligned}$$

4.14. $\mathbb{Z}/2\mathbb{Z}$ –**extension of SL_f** . In this section denote $\widehat{G} := \{\pm g \mid g \in G\}$ with $g \in \mathrm{SL}_f$.

$$\mathcal{C}^{\widehat{G}} = \{\pm \mathrm{id}, \pm S, \pm S^3, \pm R, \pm RS^3, \pm T\}.$$

We have $\mathcal{B}_{\pm S} = \mathcal{B}_{\pm S^3} = \mathcal{B}_{-\mathrm{id}} = 0$ because the fixed locus of these elements is 0 and the respective generator ξ_g is not preserved by the action of $-\mathrm{id}$.

The fixed loci of R and $-R$ are 1 and 2–dimensional respectively.

$$\mathcal{B}_R \cong (\mathbb{C}\langle [\tilde{x}]\xi_R \rangle)^{(-\mathrm{id})} = 0$$

and $\mathcal{B}_{-R} = 0$ as explained in Section 3.3.2.

For $g \in \{\pm RS^3, \pm T\}$ the fixed locus is 1–dimensional giving

$$\mathcal{B}_g \cong (\mathbb{C}\langle \xi_g, [\tilde{x}]\xi_g, [\tilde{x}^2]\xi_g \rangle)^{(-\mathrm{id})} \cong \mathbb{C}\langle \xi_{Rg}, [\tilde{x}^2]\xi_g \rangle. \quad (10)$$

The identity sector is spanned by

$$1\xi_{\mathrm{id}} \quad \text{and} \quad [5x_2^2x_3^2x_1^2 - x_3x_1^5 - x_1x_2^5 - x_2x_3^5]\xi_{\mathrm{id}}.$$

We conclude that $\mathrm{HH}^*(f, \widehat{G})$ is of dimension $2 \cdot 4 + 2 = 10$.

REFERENCES

- [AGV08] D. Abramovich, T. Graber, A. Vistoli, *Gromov–Witten theory of Deligne–Mumford stacks*, Amer. J. Math. 130 (2008), no. 5, 1337–1398.
- [AGV85] V. Arnold, A. Gusein-Zade, A. Varchenko, *Singularities of Differentiable Maps*, vol I Monographs in Mathematics, 82. Birkhäuser Boston, Inc., Boston, MA, 1985
- [BT2] A. Basalaev, A. Takahashi, *Hochschild cohomology and orbifold Jacobian algebras associated to invertible polynomials*, Journal of noncommutative geometry, Vol. 14. No. 3. pp. 861–877 (2020).
- [BT3] A. Basalaev, A. Takahashi, *Mirror Symmetry for a Cusp Polynomial Landau–Ginzburg Orbifold*, International Mathematics Research Notices, Vol. 2022, No. 19, pp. 14865–14922
- [BTW16] A. Basalaev, A. Takahashi, E. Werner, *Orbifold Jacobian algebras for invertible polynomials*, Journal of Singularities, Vol. 26 (2023), 92–127.
- [BTW17] A. Basalaev, A. Takahashi, E. Werner, *Orbifold Jacobian algebras for exceptional unimodal singularities*, Arnold Math J. 3, pp. 483–498 (2017).

- [BI21] A. Basalaev, A. Ionov, *Mirror map for Fermat polynomials with a nonabelian group of symmetries*, Theoretical and Mathematical Physics, 209(2), 1491–1506., 2021.
- [BI22] A. Basalaev, A. Ionov, *Hochschild cohomology of Fermat type polynomials with non-abelian symmetries*, Journal of Geometry and Physics, 174 (2022), 104450.
- [BI24] A. Basalaev, A. Ionov, *Hodge Diamonds of the Landau–Ginzburg Orbifolds*, SIGMA 20 (2024), 024.
- [BH95] P. Berglund, M. Henningson, *Landau–Ginzburg orbifolds, mirror symmetry and the elliptic genus*. Nuclear Phys. B 433, pp. 311–32 (1995).
- [BH93] P. Berglund, T. Hübsch, *A generalized construction of mirror manifolds*. Nuclear Phys. B 393, pp. 377–91 (1993).
- [CJMPW23] A. Clawson, D. Johnson, D. Morais, N. Priddis, C. B. White, *Mirror Map for Landau–Ginzburg models with nonabelian groups*. Journal of Geometry and Physics, Volume 199, 2024, 105161.
- [CR11] A. Chiodo, Y. Ruan. *LG/CY correspondence: the state space isomorphism*. Adv. Math. 227, no. 6 pp. 2157–88 (2011).
- [EGZ18] W. Ebeling, S. Gusein-Zade, *A version of the Berglund–Hübsch–Henningson duality with non-abelian groups*, International Mathematical Research Notices Vol. 2021, Issue 16, August 2021, Pages 12305–12329
- [EGZ20] W. Ebeling, S. Gusein-Zade, *Dual Invertible Polynomials with Permutation Symmetries and the Orbifold Euler Characteristic*. Symmetry, Integrability and Geometry: Methods and Applications, 16, 1–15 (2020).
- [ET13] W. Ebeling, A. Takahashi, *Variance of the exponents of orbifold Landau–Ginzburg models*, MATH RES LETT, 20 (2013), no.01, 51–65.
- [E12] A. Efimov, *Homological mirror symmetry for curves of higher genus*, Adv. Math. 230 (2012), no. 2, 493–530.
- [E98] N. Elkies, *The Klein quartic in number theory*. In The Eightfold Way: The Beauty of Klein’s Quartic Curve, ed. Sylvio Levi, 51–102. Mathematical Sciences Research Institute publications, 35. Cambridge: Cambridge University Press.
- [GKR17] M. Gross, L. Katzarkov, H. Ruddat, *Towards mirror symmetry for varieties of general type*, Adv. Math. 308 (2017) 208–275
- [GH94] P. Griffiths, J. Harris. *Principles of algebraic geometry*. John Wiley and Sons, 1994.
- [HLL20] W. He, Si Li, Y. Li. *G-twisted braces and orbifold Landau–Ginzburg models*. Communications in Mathematical Physics 373.1 (2020): 175-217.
- [IV90] K. A. Intriligator, C. Vafa, *Landau–Ginzburg Orbifolds*. Nucl. Phys., B339:95–120 (1990).
- [K79] F. Klein, *Ueber die Transformationen siebenter Ordnung der elliptischen Funktionen*, Math. Annalen 14 (1879), 428–471.
- [K09] M. Krawitz, *FJRW rings and Landau–Ginzburg Mirror Symmetry*, (2009) PhD thesis, The University of Michigan.
- [Kre94] M. Kreuzer, *The mirror map for invertible LG models*, PHYS LETT B **328** (1994), no.3-4, 312–318.
- [KS] M. Kreuzer, H. Skarke *On the Classification of Quasihomogeneous Functions*, Communications in mathematical physics 150 (1992): 137–147.

- [MR11] T. Milanov, Y. Ruan, *Gromov-Witten theory of elliptic orbifold \mathbb{P}^1 and quasi-modular forms* arXiv preprint 1106.2321 (2011).
- [MS12] T. Milanov, Y. Shen, *Global mirror symmetry for invertible simple elliptic singularities*. arXiv preprint 1210.6862 (2012).
- [M] D. Mukai, *Nonabelian Landau-Ginzburg orbifolds and Calabi-Yau/Landau-Ginzburg correspondence*, (2017) arXiv preprint: 1704.04889.
- [PV12] A. Polishchuk, A. Vaintrob, *Chern characters and Hirzebruch–Riemann–Roch formula for matrix factorizations*. Duke Mathematical Journal 161.10 (2012): 1863.
- [R17] H. Ruddat, *Perverse curves and mirror symmetry*. J. ALGEBRAIC GEOMETRY 26 (2017), 17–42
- [Sh20] D. Shklyarov, *On Hochschild invariants of Landau–Ginzburg orbifolds*, Advances in Theoretical and Mathematical Physics, Vol. 24, pp. 189–258 (2020).
- [S11] P. Seidel, *Homological mirror symmetry for the genus two curve*. J. Algebraic Geom. 20 (2011), no. 4, 727–769
- [V89] C. Vafa, *String vacua and orbifoldized LG models*. Modern Physics Letters A 4.12 pp. 1169–1185 (1989).
- [W93] E. Witten, *Phases of $N=2$ theories in two dimensions*. Nuclear Physics B. Aug 16;403(1-2):159–222 (1993).
- [WWP] J. Ward, M. Williams, N. Priddis *Mirror Symmetry for Nonabelian Landau-Ginzburg Models*, SIGMA. Symmetry, Integrability and Geometry: Methods and Applications. (2020), 16, 059.

A. BASALAEV:

FACULTY OF MATHEMATICS, NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS,
 USACHEVA STR., 6, 119048 MOSCOW, RUSSIAN FEDERATION, AND
 SKOLKOVO INSTITUTE OF SCIENCE AND TECHNOLOGY, NOBELYA STR., 3, 121205 MOSCOW, RUSSIAN
 FEDERATION

Email address: a.basalaev@skoltech.ru