

C^1 -robust homoclinic tangencies

Dongchen Li

Department of Mathematics, Imperial College London

June 19, 2024

Abstract. The aim of this paper is twofold. First, motivated by the nearly-affine blender system found in [LT24], we introduce standard blenders and their variations, and prove their fundamental properties on the generation of C^1 -robust tangencies. Next, as an application, we show that unfolding a homoclinic tangency to a hyperbolic periodic point can produce uncountably many C^1 -robust homoclinic tangencies, provided that either this point is involved in a coindex-1 heterodimensional cycle, or the central dynamics near it are not essentially two-dimensional.

Keywords. blender, heterodimensional cycle, homoclinic tangency, nonhyperbolic dynamics.

AMS subject classification. 37C05, 37C29, 37C75, 37D25.

Contents

1	Introduction	2
1.1	Robustness of tangencies: from C^2 to C^1	3
1.2	Creating C^1 -robust homoclinic tangencies from heterodimensional cycles	4
1.3	Strengthening the Newhouse theorem	5
1.4	C^1 -Robust homoclinic tangencies in parameter families	6
2	Standard blenders	7
2.1	Partially hyperbolic Markov partitions	7
2.2	Robust heterodimensional non-transverse intersections	9
3	Separated standard blenders	11
3.1	Base-Center structure	11
3.2	Creation of a robust homoclinic tangency. Proof of Theorem 2	12
4	Arrayed standard blenders	15
4.1	Base-Center-Gap structure	15
4.2	Reproducing lemma	17
4.3	Robust presence of uncountably many homoclinic tangencies. Proof of Theorem 3	20
5	Nearly-affine blender system (NABS)	21
5.1	NABS near heterodimensional cycles. Proof of Theorem 4	23
5.2	Robust homoclinic tangencies in parameter families. Proof of Corollary 6	26
5.3	Standard blenders arisen form an NABS. Proof of Theorem 5.3	26
	Acknowledgments	32
A	NABS obtained in [LT24]	32

arXiv:2406.12500v1 [math.DS] 18 Jun 2024

A.1 Proof of Theorem 5.6	32
A.2 NABS in the saddle-focus and double-focus cases	33
References	35

1 Introduction

A *homoclinic tangency* refers to a non-transverse intersection between the stable and unstable manifolds of a saddle¹ O of a diffeomorphism f . Since the Kupka-Smale Theorem asserts that, C^r -generically, invariant manifolds of saddles intersect transversely, the homoclinic tangency to O cannot be robust. However, this situation changes when O is replaced by a non-trivial hyperbolic basic (i.e., compact, transitive and locally maximal) set.

Definition 1.1 (Robust homoclinic tangencies). Let a C^r diffeomorphism f have a non-trivial hyperbolic basic set Λ . We say that Λ has a C^r -robust homoclinic tangency if there exists a C^r -neighborhood \mathcal{U} of f such that for every $g \in \mathcal{U}$ the continuation Λ_g of Λ has an orbit of non-transverse intersection in $W^u(\Lambda_g) \cap W^s(\Lambda_g)$.

The first example of robust homoclinic tangencies was given by Newhouse [New70], where he showed the existence of C^2 -robust homoclinic tangencies for surface diffeomorphisms. Later, he further proved [New79] that C^2 -robust homoclinic tangencies are, in fact, abundant:

every C^r ($r = 2, \dots, \infty, \omega$) surface diffeomorphism which has a saddle with a homoclinic tangency is accumulated in the C^r topology by diffeomorphisms which have hyperbolic basic sets exhibiting C^2 -robust homoclinic tangencies.

A generalization to higher dimensions with the same regularity was done in [GTS93b, PV94, Rom95]. Those open sets of homoclinic tangencies carry many important dynamical phenomena: coexistence of infinitely many sinks [New74, New79], super-exponential growth of periodic orbits [Kal00], and abundance of strange attractors [MV93], among others.

It is natural to ask whether the robust homoclinic tangencies in the Newhouse construction are C^1 -robust. A key ingredient in Newhouse’s proof is the stable intersection between two Cantor sets obtained from the stable and unstable laminations of two horseshoes with large *thickness*. Roughly speaking, large thickness implies a non-empty intersection between the Cantor sets, and each intersection point corresponds to a heteroclinic tangency between the two horseshoes. The C^2 -robust tangencies follow from the continuity dependence of thickness on C^2 maps diffeomorphisms. However, Ures showed [Ure95] that the thickness loses continuous dependence on diffeomorphisms of class C^1 , and Moreira further built in [Mor11] a general theory that no horseshoes of C^1 surface diffeomorphisms can have C^1 -robust tangencies. Since all higher dimensional generalization of the Newhouse theorem more or less rely on the original two dimensional construction, the above-mentioned results strongly indicate that Newhouse’s thick horseshoe construction cannot produce C^1 -robust homoclinic tangencies. In this paper, we prove

Theorem 1. *Every C^r ($r = 1, \dots, \infty, \omega$) diffeomorphism having a homoclinic tangency to a saddle with $d_{\text{eff}} > 2$ can be approximated in the C^r topology by a diffeomorphism having a hyperbolic basic set which exhibits uncountably many C^1 -robust homoclinic tangencies.*

¹ By saddles we mean hyperbolic periodic points with multipliers both inside and outside the unit circle. The multipliers are the eigenvalues of the linearization matrix of $f^{\text{per}(O)}$ at O .

The condition $d_{\text{eff}} > 2$ will be made precise in Definition 1.3, which means that the central dynamics near the saddle are not essentially two-dimensional. This theorem follows directly from a more collaborative version – Theorem 5 in Section 1.3.

1.1 Robustness of tangencies: from C^2 to C^1

In contrast with the essentially two-dimensional Newhouse construction, there are different constructions in dimension three or higher which lead to C^1 -robust homoclinic tangencies. Examples with specific global constructions were presented in [Sim72, Asa08]. A more general local approach was proposed by Bonatti and Díaz in [BD12] using the so-called *blender-horseshoes*, which are a special type of hyperbolic basic sets and serve a role analogous to the thick horseshoes in the Newhouse construction. They proved that a homoclinic tangency to the blender-horseshoe can be made C^1 -robust by an arbitrarily C^r -small perturbation.

Inspired by the work [BD12], we introduce the notions of *separated standard blenders* and *arrayed standard blenders*, which are variations of the *standard blenders* defined in [LT24]; precise definitions are given in Sections 3 and 4, respectively. For brevity, we will also refer to them as separated and arrayed blenders. They are both given as the locally maximal invariant sets induced from partially hyperbolic Markov partitions (see Definition 2.1). In particular, separated blenders are a generalization of blender-horseshoes. Let \mathcal{M} be a manifold of dimension at least three and denote by $\text{Diff}^r(\mathcal{M})$ the space of C^r diffeomorphisms of \mathcal{M} .

Theorem 2. *Let $f \in \text{Diff}^r(\mathcal{M})$, $r = 1, \dots, \infty, \omega$, have a separated standard blender Λ with a homoclinic tangency. Then, f can be approximated in the C^r topology by a diffeomorphism g such that Λ_g exhibits a C^1 -robust homoclinic tangency.*

This result is proved in Section 3.2. The arrayed blenders carry a more delicate structure and can produce a large number of robust tangencies.

Theorem 3. *Let $f \in \text{Diff}^r(\mathcal{M})$, $r = 1, \dots, \infty, \omega$, have an arrayed standard blender Λ with a homoclinic tangency. Then, f can be approximated in the C^r topology by a diffeomorphism g such that Λ_g exhibits uncountably many C^1 -robust homoclinic tangencies.*

We prove Theorem 3 in Section 4.3. Standard blenders are hyperbolic sets having a partially hyperbolic structure with a one-dimensional center. We note that the structure of separated blenders have the potential to be extended to the case where the central dimension is larger than one, and hence be used to create robust tangencies of a *higher corank*, that is, the tangent spaces of the stable and unstable manifolds of the blender at the tangency point share a common subspace of dimension larger than one, see Remark 4.5. For recent progresses on high corank tangencies, see [BR17, Asa22, Min24]. Although arrayed blenders seem to be a more powerful version of the separated ones, the construction of arrayed blenders is different, as it relies on the one-dimensional center in an essential way, see Remark 4.5.

It is worth mentioning that arrayed blenders can be embedded into parameter families such that they become the parablenders introduced by Berger [Ber16], which are key to build an open region, the *Berger domain*, in the space of finite-parameter families of diffeomorphisms where generic families display infinitely many sinks for an open set of parameter values. See [BCP22] for more recent progress in this direction. In a forthcoming paper [LT], we use arrayed blenders to show that Berger domains can be found near every homoclinic tangency of the sink-producing type [Tur96, GST08].

1.2 Creating C^1 -robust homoclinic tangencies from heterodimensional cycles

Standard blenders and blender-horseshoes are both obtained near coindex-1 heterodimensional cycles, but the former require only C^r -small perturbations, while the latter essentially need C^1 -small perturbations which are large in the C^r topology. Let us recall some definitions. The index of a transitive hyperbolic set, denoted by $\text{ind}(\cdot)$, refers to the rank of its unstable bundle.

Definition 1.2 (Heterodimensional cycles). We say that a diffeomorphism has a *heterodimensional cycle* involving two saddles O_1 and O_2 if $\text{ind}(O_1) \neq \text{ind}(O_2)$, and $W^u(O_1) \cap W^s(O_2) \neq \emptyset$ and $W^u(O_2) \cap W^s(O_1) \neq \emptyset$. The difference of the indices is called the *coindex* of the cycle.

We enumerate the involved saddles in the order that $\text{ind}(O_1) + 1 = \text{ind}(O_2)$. The *center-stable* and *center-unstable* multipliers of O_i ($i = 1, 2$) are those nearest to the unit circle from inside and, respectively, from outside. In general position, each O_i has either one simple and real center-stable multiplier or two, forming a conjugate complex (nonreal) pair. Similar for the center-unstable multiplier. Denote by λ the center-stable multiplier of O_1 and by γ the center-unstable multiplier of O_2 . Then, a heterodimensional cycle falls into one of the three different cases:

- *saddle*: both λ and γ are real;
- *saddle-focus*: one of λ and γ is real and the other is not; and
- *double-focus*: both λ and γ are not real.

Recall that standard blenders carry a partially hyperbolic structure with a one-dimensional center. They are further called *center-stable* (*cs*) or *center-unstable* (*cu*), depending on whether the central dynamics are contraction or expansion; see Definition 2.10. The standard blenders near a heterodimensional cycle are obtained as invariant sets of the first return map along the cycle. In the saddle-focus and double-focus cases, there is flexibility (due to the rotations brought by complex central multipliers) for realizing both contracting and expanding central dynamics. However, in the saddle case, this contraction/expansion is relatively rigid and is governed by a derivative of the transition map along the transverse heteroclinic intersection of the cycle, which is denoted as α in [LT24, Equation (2.4)]. See Section 5.1 for a brief explanation, and [LT24, Section 2.3] for more details. Recall that two transitive hyperbolic sets are *homoclinically related* if they have the same index, and the unstable manifold of each set intersects the stable manifold of the other set transversely. To facilitate the presentation, we call a transitive hyperbolic set C^1 -wild if it exhibits uncountably many C^1 -robust homoclinic tangencies, and use $\overset{\text{h.r.}}{\sim}$ to denote the homoclinic relation.

Theorem 4. *Let $f \in \text{Diff}^r(\mathcal{M})$, $r = 1, \dots, \infty, \omega$, have a coindex-1 heterodimensional cycle involving two saddles O_1 and O_2 . Then, f can be approximated in the C^r topology by a diffeomorphism g which has at least one standard blender, being simultaneously separated and arrayed. More specifically,*

- *in the saddle case,*
 - *when $|\alpha| \leq 1$, there is one such blender $\Lambda_1 \overset{\text{h.r.}}{\sim} O_{1,g}$,*
 - *when $|\alpha| \geq 1$, there is one such blender $\Lambda_2 \overset{\text{h.r.}}{\sim} O_{2,g}$, and*
- *in other cases, there are two such blenders Λ_1 and Λ_2 with $\Lambda_1 \overset{\text{h.r.}}{\sim} O_{1,g}$ and $\Lambda_2 \overset{\text{h.r.}}{\sim} O_{2,g}$, where*
 - *in saddle-focus case,*
 - * Λ_1 *is C^1 -wild if γ is nonreal,*
 - * Λ_2 *is C^1 -wild if λ is nonreal, and*
 - *in the double-focus case, g can be chosen such that either Λ_1 or Λ_2 is C^1 -wild.*

Moreover, for $i = 1, 2$, if we additionally assume that O_i has a homoclinic tangency, then the blender Λ_i is always C^1 -wild. In all cases, Λ_1 is center-stable and Λ_2 is center-unstable.

Note that, in the saddle-focus case, both Λ_1 and Λ_2 are C^1 -wild if either O_1 has a homoclinic tangency and λ is nonreal, or O_2 has a homoclinic tangency and γ is nonreal; in the double-focus case, this happens if one of O_1 and O_2 has a homoclinic tangency.

The above two theorems follow from a more detailed Theorem 5.5 in Section 5.1, which considers the coexistence of robust homoclinic tangencies and robust heterodimensional dynamics. Its proof is based on the analysis of the *nearly-affine blender system* (see Definition 5.1), discovered near coindex-1 heterodimensional cycles in [LT24]. These systems serve as a source for generating all types of standard blenders.

1.3 Strengthening the Newhouse theorem

In light of the result in [LLST22], Theorem 4 strengthens the classical Newhouse theorem (and its higher dimensional generalization) that every homoclinic tangency is accumulated by C^2 -robust ones in the C^r ($r > 1$) topology to C^1 -robustness for all homoclinic tangencies associated with saddles of *effective dimension larger than two*, which we define as follows. Let us denote the multipliers of a saddle O as

$$|\lambda_{d_s}| \leq \dots \leq |\lambda_1| < 1 < |\gamma_1| \leq \dots \leq |\gamma_{d_u}|.$$

In general position, there is only one real or one pair of conjugate complex center-stable multipliers, namely, λ_1 or λ_1 and $\lambda_2 = \lambda_1^*$. The same for center-unstable multipliers. Let us denote by d_{cs} and d_{cu} the numbers of center-stable and -unstable multipliers. If O has a generic homoclinic tangency, then, according to [GTS93a, BC15], the orbits of O and the tangency lie in a center invariant manifold \mathcal{M}^c of dimension $(d_{cs} + d_{cu})$, which is tangent to the eigenspace corresponding to the central multipliers. We are particularly interested in the dynamics restricted to \mathcal{M}^c , as everything born from the homoclinic bifurcation must lie in \mathcal{M}^c , see [Tur96].

Definition 1.3 (Effective dimension). We say that a saddle has *effective dimension larger than two*, denoted by $d_{\text{eff}} > 2$, if $\dim \mathcal{M}^c > 2$ and the dynamics restricted to it are neither area-contracting nor area-expanding. More specifically, $d_{\text{eff}} > 2$ if

1. $\lambda_1 = \lambda_2^*$ is not real, γ_1 is real, and $|\lambda_1 \gamma_1| > 1$, or
2. λ_1 is real, $\gamma_1 = \gamma_2^*$ is not real, and $|\lambda_1 \gamma_1| < 1$, or
3. $\lambda_1 = \lambda_2^*$ and $\gamma_1 = \gamma_2^*$ are not real, and $|\lambda_1 \gamma_1| \neq 1$.

This definition is the same as the effective dimension introduced in [Tur96], denoted by d_{eff}^b , with $d_{\text{eff}} = d_{\text{eff}}^b - 1$. The difference is that d_{eff}^b is associated with the bifurcation of a homoclinic tangency, rather than the saddle itself. It was established in [GST08] that a homoclinic bifurcation of dimension d_{eff}^b can produce periodic points which have d_{eff}^b multipliers on the unit circle, and hence lead to the coexistence of saddles of $d_{\text{eff}}^b + 1$ different indices. For our purpose, the condition $d_{\text{eff}} > 2$ (or $d_{\text{eff}}^b > 1$) ensures that unfolding a homoclinic tangency yields two saddles of different indices, which allows for the potential creation of a heterodimensional cycle involving these saddles. It is indeed the case by the following

Theorem ([LLST22, Theorem 1]). *Let $f \in \text{Diff}^r(\mathcal{M})$, $r = 1, \dots, \infty, \omega$, have a homoclinic tangency to a saddle O with $d_{\text{eff}} > 2$. Then, f can be approximated in the C^r topology by a diffeomorphism which has a coindex-1 heterodimensional cycle involving the continuation of O and another saddle Q satisfying $\text{ind}(Q) = \text{ind}(O) + 1$ if $|\lambda_1 \gamma_1| > 1$, and $\text{ind}(Q) = \text{ind}(O) - 1$ if $|\lambda_1 \gamma_1| < 1$.*

Here the assumption $|\lambda_1 \gamma_1| > 1$ means that O_1 belongs either to case 1 of Definition 1.3, or to case 3 with $|\lambda_1 \gamma_1| > 1$. Similarly for the assumption $|\lambda_1 \gamma_1| < 1$.

By the (multidimensional) Newhouse theorem, up to an arbitrarily C^r -small perturbation, one can assume that f has a hyperbolic basic set having a C^2 -robust homoclinic tangency. Moreover, this set is homoclinically related to O [GTS93b, GLM]. Hence, after applying the perturbation involved in the above theorem, one can, by the lambda lemma, recover the homoclinic tangency to the continuation of O , by an additional arbitrarily C^r -small perturbation. After that, invoking the saddle-focus case of Theorem 4 immediately gives

Theorem 5. *Let $f \in \text{Diff}^r(\mathcal{M})$, $r = 1, \dots, \infty, \omega$, have a homoclinic tangency to a saddle O with $d_{\text{eff}} > 2$. Then, f can be approximated in the C^r topology by a diffeomorphism g which has a standard cs-blender and a standard cu-blender, Λ_1 and Λ_2 with $\text{ind}(\Lambda_1) + 1 = \text{ind}(\Lambda_2)$, each of which is simultaneously separated and arrayed, and exhibits uncountably many C^1 -robust homoclinic tangencies. Moreover, $O_g \stackrel{\text{h.r.}}{\sim} \Lambda_1$ if $|\lambda_1 \gamma_1| > 1$, and $O_g \stackrel{\text{h.r.}}{\sim} \Lambda_2$ if $|\lambda_1 \gamma_1| < 1$.*

Remark 1.4. The condition $d_{\text{eff}} > 2$ is *necessary* for creating heterodimensional cycles from the bifurcation of a single² generic homoclinic tangency, and, hence, necessary for creating C^1 -robust homoclinic tangencies through the mechanism of heterodimensional bifurcations. To see this, note that any heterodimensional cycle born from the homoclinic bifurcation of O is also a cycle for $f|_{\mathcal{M}^c}$. So, the involved hyperbolic periodic points, when viewed as those of $f|_{\mathcal{M}^c}$, must also be saddles of different indices, which is impossible when $d_{\text{eff}} \leq 2$.

The robust presence of uncountably many homoclinic tangencies also appears in the two-dimensional Newhouse construction (though only C^2 -robust). This was not proved in his original paper, but follows from the main result in [HKY93]. For a detailed explanation see [BB23, Corollary 2.17].

1.4 C^1 -Robust homoclinic tangencies in parameter families

It is well-known that the C^2 -robust homoclinic tangencies guaranteed by the Newhouse theorem can be found within a generic unfolding of a homoclinic tangency. There are similar results regarding the C^1 -robust homoclinic tangencies of the blenders in Theorem 5.

We consider a generic homoclinic tangency to the saddle O with $d_{\text{eff}} > 2$, where the genericity conditions are given in [LLST22, Section 2.3] and can be fulfilled by an arbitrarily C^r -small perturbation. Let us take two points $P_1 \in W_{loc}^u(O)$ and $P_2 \in W_{loc}^s(O)$ from the orbit of tangency with $P_2 = f^n(P_1)$, and define a transition map T as the restriction of f^n to a small neighborhood of P_1 . Then, we can introduce a splitting parameter μ as the functional measuring the distance between $T(W_{loc}^u(O))$ and $W_{loc}^s(O)$ near the point P . We also use the arguments of the central multipliers λ and γ as other parameters, denoted by ω_1 and ω_2 .

Let $f \in \text{Diff}^r(\mathcal{M})$, $r = 3, \dots, \infty, \omega$, have a generic homoclinic tangency to a saddle O with $d_{\text{eff}} > 2$. Any family $\{f_\varepsilon\}$ with at least two parameters and $f_0 = f$ is called a *proper unfolding* if,

- when $\lambda_1 \gamma_1 > 1$, $\frac{\partial(\mu, \omega_1)}{\partial \varepsilon}$ has rank 2; and
- when $\lambda_1 \gamma_1 < 1$, $\frac{\partial(\mu, \omega_2)}{\partial \varepsilon}$ has rank 2.

Corollary 6. *For any proper unfolding $\{f_\varepsilon\}$, there exists a sequence $\varepsilon_i \rightarrow 0$ such that f_{ε_i} has a hyperbolic basic set exhibiting uncountably many C^1 -robust homoclinic tangencies. More specifically, this set is the blender Λ_2 of Theorem 5 when $\lambda_1 \gamma_1 > 1$, and is Λ_1 when $\lambda_1 \gamma_1 < 1$.*

² Heterodimensional cycles can also be created in case 1 of Definition 1.3 if O has two orbits of homoclinic tangency which intersect the same strong-stable/unstable leaf, see [LT20].

We prove this corollary in Section 5.2. It is an immediate consequence of a parametric result of [LLST22] (which requires f to be at least C^3) and the proof of Theorems 2 and 3. Note that the obtained blender that exhibits C^1 -robust homoclinic tangencies has an index different from O , so it cannot be homoclinically related to O . However, as shown in Theorem 5.5, this blender and O form C^1 -robust heterodimensional dynamics.

In the next three sections, we define standard blenders, separated standard blenders, and arrayed standard blenders, respectively, which are characterized by Markov partitions, and prove their fundamental properties. Finally, in Section 5, we use the results in [LT24] to prove Theorem 4.

2 Standard blenders

The notion of blenders was introduced by Bonatti and Díaz for constructing robust non-hyperbolic transitive diffeomorphisms [BD96]. A blender is a hyperbolic basic (i.e., compact, transitive and locally maximal) set with the property that the projection of its stable or unstable lamination to certain central subspace has a non-empty interior. As a consequence, the lamination acts as if it has one more topological dimension and hence can produce robust non-transverse intersections with other submanifolds.

Blenders have been used to obtain important dynamical phenomena in a wide range of settings. For a list of these applications we refer the readers to [Li24] and the references therein. Due to different purposes, blenders have emerged in various forms during the last decades, see, e.g. [NP12, BKR14, BBD16]. In particular, the blenders arisen from heterodimensional cycles are characterized by Markov partitions of a finite (and possibly large) number of elements, which motivated the notion of standard blenders given in [LT24, Appendix]. In this section, we first introduce partially hyperbolic Markov partitions and reformulate the definition of standard blenders. After that, we prove the fundamental property of standard blenders that they can produce robust non-transverse heterodimensional intersections, see Proposition 2.11.

2.1 Partially hyperbolic Markov partitions

Let $\{\Pi_i\}$ be a finite collection of pairwise disjoint closed subsets of a manifold \mathcal{M} of dimension $d \geq 3$, and $g : U \rightarrow \mathcal{M}$ be any diffeomorphism of some neighborhood U of $\bigcup \Pi_i$ such that $g(\Pi_i) \subset U$. Up to dividing Π_i into subsets, we can assume that the intersection $\Pi_i \cap g(\Pi_j)$ has at most one connected component. We call the pair $(\{\Pi_i\}, g)$ a *Markov partition* if for each pair (i, j) it holds that $g^n(\Pi_i) \cap \Pi_j \neq \emptyset$ for some $n \geq 0$. Denote by $\text{int}(\cdot)$ the interior of a set.

Definition 2.1 (cs-Markov partitions). A Markov partition $(\{\Pi_i\}, g)$ is called *center-stable (cs) partially hyperbolic* if

- (Horizontal domains) there exist charts (U, φ_i) , convex closed subsets $\mathbb{X}_i \subset \mathbb{R}$, $\mathbb{Y}_i \subset \mathbb{R}^{d_u}$, $\mathbb{Z}_i \subset \mathbb{R}^{d_{ss}}$ with $1 + d_u + d_{ss} = d$, and functions $\psi_i : \mathbb{X}_i \times \mathbb{R}^{d_u} \times \mathbb{Z}_i \rightarrow \text{int}(\mathbb{Y}_i)$ such that

$$\varphi_i(\Pi_i) = \{(x, \psi_i(x, y, z), z) : (x, y, z) \in \mathbb{X}_i \times \mathbb{Y}_i \times \mathbb{Z}_i\}; \quad (1)$$

- (Vertical images) for every pair (i, j) with $g(\Pi_i) \cap \Pi_j \neq \emptyset$, there exists a function $g_{ij}^\times : \mathbb{X}_i \times \mathbb{Y}_j \times \mathbb{Z}_i \rightarrow \text{int}(\mathbb{X}_j \times \mathbb{Y}_i \times \mathbb{Z}_j)$ which is a contraction, i.e., $\left\| \frac{\partial g_{ij}^\times}{\partial(x, y, z)} \right\| < 1$ for some suitable norm, such

that one has $(\bar{x}, \bar{y}, \bar{z}) = \varphi_j \circ g \circ \varphi_i(x, y, z)$ if and only if

$$(\bar{x}, \bar{y}, \bar{z}) = g_{ij}^\times(x, \bar{y}, \bar{z}); \quad (2)$$

- (hyperbolicity with a partially hyperbolic structure) with denoting by (dx, dy, dz) the vectors in the tangent spaces, there exist on $\varphi(\Pi_i)$ the cone fields

$$\begin{aligned} \mathcal{C}^u &= \{(dx, dy, dz) : |dx| + \|dz\| < K_i^u |dy|\}, \\ \mathcal{C}^s &= \{(dx, dy, dz) : \|dy\| < K_i^s (|dx| + \|dz\|)\}, \\ \mathcal{C}^{ss} &= \{(dx, dy, dz) : |dx| + \|dy\| < K_i^{ss} \|dz\|\}, \end{aligned} \quad (3)$$

for some $K_i \in (0, 1)$ satisfying that, for any $P \in \Pi_i$ and $\bar{P} = g(P) \in \Pi_j$,

- \mathcal{C}^u is forward-invariant: \mathcal{C}_P^u is mapped strictly inside $\mathcal{C}_{\bar{P}}^u$ by the differential $D(\varphi_j \circ g \circ \varphi_i^{-1})$,
- \mathcal{C}^s and \mathcal{C}^{ss} are backward-invariant: $\mathcal{C}_{\bar{P}}^{s/ss}$ is mapped strictly inside $\mathcal{C}_P^{s/ss}$ by $D(\varphi_i \circ g^{-1} \circ \varphi_j^{-1})$,
- vectors in \mathcal{C}^u are uniformly expanded by $D(\varphi_j \circ g \circ \varphi_i^{-1})$ and vectors in $\mathcal{C}^{s/ss}$ are uniformly contracted by $D(\varphi_i \circ g^{-1} \circ \varphi_j^{-1})$.

Remark 2.2. The existence of the cone fields in (3) implies a hyperbolic splitting of the tangent bundle $T\Pi_i$ with a partially hyperbolic structure: $T\Pi_i = E^{cs} \oplus E^u \oplus E^{ss}$. The stronger contraction in z -directions implies that the first component of g_{ij}^\times , say g_{ij1}^\times , has its first partial derivative with respect to x bounded away from zero. Thus, $\partial g_{ij1}^\times / \partial x$ is either negative or positive for all $(\bar{y}, \bar{z}) \in \mathbb{Y}_j \times \mathbb{Z}_i$.

Let us make some simple observations from the above definition. To simplify the notations, we will always identify Π_i with $\varphi_i(\Pi_i)$ throughout Sections 2–4, unless otherwise stated.

By (1) and (2), each preimage $H_{ij} := \Pi_i \cap g^{-1}(\Pi_j)$ is a *horizontal strips* in the sense that it has full (x, z) -size and small y -size in Π_i . That is, H_{ij} has a skew-product structure as in (1), but with some function ψ_{ij} satisfying $|\psi_{ij}(x, y, z)| < |\psi_i(x, y, z)|$. Similarly, each image $V_{ji} := \Pi_i \cap g(\Pi_j)$ is a *vertical strip* in the sense that $g(\Pi_j)$ has full y -size and small (x, z) -size in $\mathbb{X}_i \times \mathbb{Y}_i \times \mathbb{Z}_i$ by (2), and hence V_{ji} has full y -size and small (x, z) -size in Π_i by (1).

Example 2.3. A simple example of a cs-Markov partition can be generated by a three-dimensional horseshoe, as illustrated in Figure 2.1. The map g contracts the cube in the (x, z) -directions, expand it in the y -direction, and then bend it such that its top crosses the cube along the y -direction. This results in a cs-Markov partition $\{\Pi_1, \Pi_2\}$, and the above-mentioned horizontal and vertical strips are in fact substrips inside Π_i and $g(\Pi_i)$.




Figure 2.1: A three-dimensional Horseshoe

Convention. For a Markov partition $(\{\Pi_i\}_{i=1}^N, g)$, iterations of g is defined inductively by $g^{n+1}(\Pi_i) = \bigcup_{j=1}^N g(g^n(\Pi_i) \cap \Pi_j)$.

Denote $\Pi = \bigcup \text{int}(\Pi_i)$. The Markov partition induces a zero-dimensional hyperbolic basic set $\Lambda := \bigcap_{n \in \mathbb{Z}} g^n(\Pi)$ consisting of points whose orbits never leave Π . To see this, note first that, since hyperbolicity follows directly from the existence of the cone fields, it suffices to show that this invariant set is non-empty. By definition, the map g_{ij}^\times takes $\mathbb{X}_i \times \mathbb{Y}_j \times \mathbb{Z}_i$ into $\text{int}(\mathbb{X}_j \times \mathbb{Y}_i \times \mathbb{Z}_j)$ and is a contraction. It then follows from a fixed-point lemma on products of metric spaces (see e.g. [Shi67, Theorem 5.8]) that the set of points whose orbits by g lie entirely in Π , i.e., the set Λ , is in one-to-one correspondence to the set $\{1, \dots, \text{card}\{\Pi_i\}\}^{\mathbb{Z}}$, or a subset of it when there exist i and j such that $g(\Pi_i) \cap \Pi_j = \emptyset$. More specifically, each point $P \in \Lambda$ corresponds to a coding $(i_n)_n \in \{1, \dots, \text{card}\{\Pi_i\}\}^{\mathbb{Z}}$ such that its orbit $\{P_n := g^n(P)\}$ satisfies $P_n \in \Pi_{i_n}$. In particular, Λ is non-empty. By construction, the stable manifold $W^s(\Lambda)$ consists of points whose forward iterates never leave Π , while the unstable manifold $W^u(\Lambda)$ consists of points whose backward iterates never leave Π .

Definition 2.4 (cu-Markov partitions). We say that the $(\{\Pi_i\}, g)$ is a *center-unstable partially hyperbolic (cu)* Markov partition if its inverse $(\{g(\Pi_i)\}, g^{-1})$ is a cs-Markov partition.

2.2 Robust heterodimensional non-transverse intersections

The complexity of the dynamics arisen from a cs-Markov partition, and, in particular, our results, are tightly related to the behavior of the backward iteration of curves tangent to the cone field \mathcal{C}^{ss} , which we all *ss-discs* defined below. We are interested the “recurrence” of such discs that the preimage of every ss-disc contains a new ss-disc.

Let $\partial_x \Pi_i$ be the x -boundary of Π_i , namely, the set $\partial \mathbb{X}_i \times \mathbb{Y}_i \times \mathbb{Z}_i$. We also have the y -boundary $\partial_y \Pi_i$ and z -boundary $\partial_z \Pi_i$. Similarly, we define the x, y, z -boundaries of horizontal strips H of Π_i . For a vertical strip of the form $V = g(H) \cap \Pi_i$ for some horizontal strips $H \subset \Pi_j$, we have $\partial_\rho V = g(\partial_\rho H) \cap \Pi_i$ for $\rho = x, y, z$.

Definition 2.5 (ss-discs and u-discs). Let \mathcal{C}^{ss} and \mathcal{C}^u be the cone fields in (3) associated to the cs-Markov partition $(\{\Pi_i\}, g)$. An *ss-disc* S of Π_i is a submanifold tangent to the cone field \mathcal{C}^{ss} such that it lies in $\Pi_i \setminus (\partial_x \Pi_i \cup \partial_y \Pi_i)$ and is given by some smooth function $(x, y) = s(z)$ defined for all $z \in \mathbb{Z}_i$. A *u-disc* S^u of Π_i is a submanifold tangent to the cone field \mathcal{C}^u such that it lies in $\Pi_i \setminus (\partial_x \Pi_i \cup \partial_z \Pi_i)$ and is given by some smooth function $(x, z) = s^u(y)$ defined for all $y \in \mathbb{Y}_i$.

Lemma 2.6. *Let S be an ss-disc of Π_i and V be a vertical strip of the form $V = g(H)$ for some horizontal strip $H \subset \Pi_j$. If S crosses V , i.e., $S \cap V \subset (V \setminus \partial_x V)$, then $g^{-1}(S \cap V)$ is an ss-disc of Π_j .*

Proof. Since V has full y -size, S intersects all faces of $\partial_z V = g(\partial_z H) \cap \Pi_i$. By construction, we have $\partial_z H \subset \partial_z \Pi_j$. It follows that $g^{-1}(S \cap V)$ is a smooth submanifold intersecting all faces of $\partial_z \Pi_j$. It is further an ss-disc of Π_j due to the invariance of the cone field \mathcal{C}^{ss} . \square

Definition 2.7 (Base of Π_i). We associate to each i an open interval $\mathbb{X}_i^B \subset \mathbb{X}_i$ and define

- *base:* $\Pi_i^B = \Pi_i \cap (\mathbb{X}_i^B \times \mathbb{Y}_i \times \mathbb{Z}_i)$, $H^B = H \cap \Pi_i^B$, $V^B = g(\tilde{H}^B) \cap \Pi_i^B$,

where H and V are any horizontal and vertical strips of Π_i , with $V = g(\tilde{H}) \cap \Pi_i$ for some \tilde{H} of Π_j . We say that an ss-disc S *crosses* the base Π_i^B if $S \cap \Pi_i \subset \Pi_i^B$; similarly for all horizontal strips H and vertical strips V .

Definition 2.8 (Covering property). A cs-Markov partition $(\{\Pi_i\}, g)$ has the *covering property* if

(A1) for all Π_i , every ss-disc S crossing the base Π_i^B crosses the base V^B of some vertical strip.

This property is easy to achieve. Since V has full y -size, whether S intersects V essentially depends on the x -coordinates of its points. When the cone constants in (3) are small, one can find, for some reference point z_i^* , an interval I such that an ss-disc crosses V whenever $s_x(z^*) \in I$, where s_x is the x -component of its defining function. The covering property is essentially the existence of sufficiently many vertical strips whose corresponding intervals I cover \mathbb{X}_i^B .

Example 2.9. Continue with Example 2.3, we define the bases by darker colors as illustrated in Figure 2.2. The covering property is satisfied since every ss-disc crossing Π_i^B must cross the one of the two intersections $\Pi_i^B \cap g(\Pi_1^B)$ and $\Pi_i^B \cap g(\Pi_2^B)$, each of which is the base of a vertical strip.




Figure 2.2: Π_i with bases

Definition 2.10 (Standard blenders). Given a cs-Markov partition $(\{\Pi_i\}, g)$ that satisfies the covering property, the locally maximal invariant set $\Lambda := \bigcap_{n \in \mathbb{Z}} g^n(\bigcup \text{int}(\Pi_i))$ is called a *standard center-stable (cs) blender*. It is called a *standard center-unstable (cu) blender* if the Markov partition is center-unstable.

As discussed after Definition 2.1, a standard blender is a zero-dimensional hyperbolic basic set, restricted to which g is conjugate to a full shift of $\text{card}\{\Pi_i\}$ symbols. For cu-Markov partitions, one has, instead of (3), forward-invariant cone fields $\mathcal{C}^u, \mathcal{C}^{uu}$ and backward-invariant cone field \mathcal{C}^s , which lead to the notions of uu- and s-discs, completely parallel to the ss- and u-discs in Definition 2.5.

Proposition 2.11 (Robust non-transverse intersections). *The unstable manifold of a standard cs-blender intersects every ss-disc which crosses some Π_i^B ; and the stable manifold of a standard cu-blender intersects every uu-disc which crosses some Π_i^B .*

This is the fundamental property of blenders as introduced in the discovery paper [BD96] of blenders.

Proof. We only prove the proposition for cs-blenders, as the arguments for cu-blenders are parallel. Let Λ be the blender induced from a Markov partition $(\{\Pi_i\}, g)$. By the covering property, any ss-disc S crossing Π_i^B crosses some V^B . In particular, it crosses V , and hence, by Lemma 2.6, the preimage $S_1 := g^{-1}(S \cap V)$ is an ss-disc of some Π_{i_1} . By construction, S_1 crosses $\Pi_{i_1}^B$. Repeating this process yields a coding $\underline{i} = (i_n)$ and a sequence of ss-discs S_n crossing $\Pi_{i_n}^B$. The images $\hat{S}_n := g^n(S_n)$ are

nested compact sets in S , so one can find a point $P \in \bigcap \hat{S}_n$ such that its backward iterates never leave $\bigcup \Pi_i$. Thus, $P \in W^u(\Lambda)$ by definition of $W^u(\Lambda)$, and, in particular, it belongs to the unstable leaf $\ell^u := \bigcup_{n \geq 0} g^n(\Pi_{i_n}^B)$. (See the discussion in the end of Section 2.1.) \square

Remark 2.12. It follows from the proof that if an ss-disc crosses some V^B , then it must intersect an unstable leaf belongs to V^B .

3 Separated standard blenders

It is established in [LT24, Theorem D] (and also in [Li24, Corollary A] as an improved version) that a heterodimensional cycle produces an infinite sequence of standard blenders, instead of just finitely many. This sequence is called a nearly-affine blender system and is explained in Section 5. An important observation is that an appropriate superposition of finitely many standard blenders from this system can yield more powerful ones, including the separated blenders which we discuss in this section.

Separated blenders have the fundamental property that homoclinic tangencies associated with them can be made C^1 -robust, see Theorem 2. In this regard, they work as the blender-horseshoes in [BD12], while in a more relaxed setting.

3.1 Base-Center structure

We introduce an additional structure for a cs-Markov partition $(\{\Pi_i\}, g)$. Recall that we have identified Π_i with $\mathbb{X}_i \times \mathbb{Y}_i \times \mathbb{Z}_i$.

Definition 3.1 (Base-Center structure of Π_i). One can associate to each i two open intervals $\mathbb{X}_i^C \subset \mathbb{X}_i^B \subset \mathbb{X}_i$ and define the following structure:

- *base*: $\Pi_i^B = \Pi_i \cap (\mathbb{X}_i^B \times \mathbb{Y}_i \times \mathbb{Z}_i)$, $H^B = H \cap \Pi_i^B$, $V^B = g(\tilde{H}^B) \cap \Pi_i^B$,
- *center*: $\Pi_i^C = \Pi_i \cap (\mathbb{X}_i^C \times \mathbb{Y}_i \times \mathbb{Z}_i)$, $H^C = H \cap \Pi_i^C$, $V^C = g(\tilde{H}^C) \cap \Pi_i^B$,

where H and V are any horizontal and vertical strips of Π_i , with $V = g(\tilde{H}) \cap \Pi_i$ for some \tilde{H} of Π_j . We say that an ss-disc S *crosses* $\Pi_i^{B/C}$ if $S \cap \Pi_i \subset \Pi_i^{B/C}$; similarly for the strips.

We say that two sets are *separated* if there is no ss-disc intersecting both of them.

Definition 3.2 (Base-Center covering property). A cs-Markov partition $(\{\Pi_i\}, g)$ has the *Base-Center covering property* if, in addition to (A1), the cubes Π_i can be structured such that, with $S \in \Pi_i^B$ and $V \in \Pi_i$ denoting ss-discs and vertical strips, the following are satisfied for every i :

- (A2) if $S \cap \partial_x V_1^B \neq \emptyset$, then S crosses some center V_2^C which is separated from V_1^C , and
- (A3) if $S \cap \Pi_i^C \neq \emptyset$, then S crosses some center V^C .

Property (A2) says that the vertical strips are ‘aligned’, while property (A3) requires sufficient vertical strips whose centers cover Π_i^C . Figure 3.1 provides an illustration of properties (A1)–(A3), where (A2) for the strips V_1 and V_2 are drawn explicitly with cone field \mathcal{C}^{ss} (in red).




Figure 3.1: The fulfilment of properties (A1)–(A3), where the red cylinders represent the areas covered by the strong-stable cone field \mathcal{C}^{ss}

Remark 3.3. Denote

$$\mathcal{W} = \{\ell^u \subset W^u(\Lambda) : g^{-n}(\ell^u) \text{ belongs to some } \Pi_{i_n}^C \text{ for every } n \geq 0\}.$$

Condition (A3) implies that any ss-disc crossing Π_i^C intersects some unstable leaf $\ell^u \in \mathcal{W}$. To see this, one just needs to repeat the proof of Proposition 2.11 with replacing Π_{i_n} by $\Pi_{i_n}^C$.

Remark 3.4. One can replace Π^B by the interior $\text{int } \Pi_i$ for less notions, but then the condition (A2) becomes stronger. For example..

Definition 3.5 (Separated standard blenders). A standard blender is called *separated* if its inducing Markov partition satisfies the aligned covering property.

3.2 Creation of a robust homoclinic tangency. Proof of Theorem 2

This result was established in [BD12, Theorem 4.9] for blender-horseshoes, and one can check that they are separated cu-blenders induced from a cu-Markov partition with 2 elements. Here we prove it for all separated blenders. The proof is based on the notion of *folding manifolds* introduced in [BD12], which is defined below with modifications to fit the general setting of the present paper. We will see that Theorem 2 is a direct consequence of the fact *every folding manifold has a tangency with $W^u(\Lambda)$* . To proof of this fact contains two steps. First, we show (Lemma 3.8) that every folding manifold, if has no tangency with $W^u(\Lambda)$, then must contain another folding manifold in its preimage. Next, we show (Proposition 3.9) that such property implies the existence of a tangency.

Let Λ be a separated cs-blender with the inducing Markov partition $(\{\Pi_i\}, g)$, and let \mathcal{W} be the collection of unstable leaves defined in Remark 3.3.

Definition 3.6 (Folding manifolds). A submanifold $\mathcal{S} \subset \Pi_i^B$ of dimension $(d_{ss} + 1)$ is called a *folding manifold* of the blender Λ if there exists a smooth family $\{S_t\}_{t \in [0,1]}$ of ss-discs crossing Π_i^B such that

- $\mathcal{S} = \bigcup_{t \in [0,1]} S_t$, and
- S_0 and S_1 intersect a leaf $\ell^u \in \mathcal{W}$.

See Figure 3.2 for an illustration. In comparison with the definition in [BD12], the folding manifolds there are required to intersect, instead of any leaf in $W^u(\Lambda)$, one of the local unstable manifolds of two prescribed saddles in Λ , and to satisfy that $S_t \cap \ell^u = \emptyset$ for $t \in (0, 1)$.




Figure 3.2: A folding manifold

For any u -disc S^u with a defining function $(x, z) = (s_1^u(y), s_2^u(y))$, consider the family $\{S_t^u\}$ of u -discs with each S_t^u given by $(x, z) = (s_1^u(y) + t, s_2^u(y))$. Take any interval $I \subset \mathbb{R}$ such that $\bigcup_{t \in I} S_t^u$ covers Π_i . We denote

$$\mathcal{L}(S^u) = \Pi_i \cap \bigcup_{t \in I} S_t^u. \quad (4)$$

Lemma 3.7. *Any folding manifold \mathcal{S} has some point $P \in \mathcal{S}$ such that $T_P \mathcal{S} \cap \mathcal{C}^u \neq \emptyset$.*

Proof. Let ℓ^u be the leaf defining the folding manifold, and $\mathcal{L} := \mathcal{L}(\ell^u)$ be defined as in (4). Denote by $P_0 = (x_0, y_0, z_0)$ and $P_1 = (x_1, y_1, z_1)$ the intersections points of $S_0 \cap \ell^u$ and $S_1 \cap \ell^u$, respectively. Take the smooth curve $\gamma \subset \mathcal{S} \cap \mathcal{L}$ connecting P_1 and P_2 such that $\gamma(t) = (\gamma_x(t), \gamma_y(t), \gamma_z(t)) \in S_t$ with $\gamma(0) = P_0$ and $\gamma(1) = P_1$.

Let the defining function of ℓ^u be $(x, z) = (s_1^u(y), s_2^u(y))$. By construction we have $\gamma_z(t) = s_2^u(\gamma_y(t))$. Since ℓ^u is tangent to \mathcal{C}^u , we have

$$\|\gamma'_z(t)\| = \left\| \frac{ds_2^u(\gamma_y(t))}{dy} \cdot \gamma'_y(t) \right\| < K^u \|\gamma'_y(t)\|,$$

where K^u is the cone constant in (3). It follows that, if $\gamma'_x(t)$ vanishes at some $t' \in (0, 1)$, then $T_{\gamma(t')} \mathcal{S} \cap \mathcal{C}^u \neq \emptyset$. Now suppose that $T_{\gamma(t)} \mathcal{S} \cap \mathcal{C}^u = \emptyset$ for all t . In particular, $\gamma'_x(t)$ never vanishes, and hence $|\gamma_x(t)|$ is strictly increasing. Since the tangent spaces of γ are transverse to \mathcal{C}^u everywhere, for any given variation in y -coordinates, say $\|y_1 - y_2\|$, the corresponding variation Δx_γ in the x -coordinate of γ must be strictly larger than that of any curve tangent to \mathcal{C}^u . However, by construction, we have $\Delta x_\gamma = |x_1 - x_0|$, which is equal to the x -variation of any smooth curve in ℓ^u connecting P_1 and P_2 . \square

Lemma 3.8. *If \mathcal{S} is a folding manifold of Λ , then either \mathcal{S} has a non-transverse intersection with $W^u(\Lambda)$, or the preimage $g^{-1}(\mathcal{S})$ contains a folding manifold.*

Proof. Let V_0 be the vertical strip containing the leaf ℓ^u of the folding manifold.

(Case 1): If every ss-disc in \mathcal{S} crosses V_0 , then $g^{-1}(\mathcal{S} \cap V_0)$ is a folding manifold. Indeed, by (A1) this preimage lies in some Π_j^B and consists of ss-discs, and the leaf $g^{-1}(\ell^u)$ belongs to \mathcal{W} by definition. The projection of this case to the (x, y) -space is sketched in the left picture of Figure 3.3.

(Case 2): Now suppose that V_0 is not crossed by all ss-discs of \mathcal{S} . In particular, there exists S_{t_1} intersecting $\partial_x V_0^B$ for some $t_1 \in (0, 1)$. By (A2) there is some vertical strip V_1 such that S_t crosses V_1^C . It follows from Remark 3.3 that S_t intersects a leaf $\ell_1^u \in \mathcal{W}$. If ℓ_1^u is tangent to \mathcal{S} , then we are done.




Figure 3.3: Two cases of the position of a folding manifold

Claim. *If ℓ_1^u is transverse to \mathcal{S} , then ℓ_1^u intersects S_{t_2} for some $t_2 \in (0, 1) \setminus \{t_1\}$.*

Let us prove the claim. Let ℓ^u be the leaf in the definition of the folding manifold. We attach to each point of ℓ^u an ss-disc such that they form a continuous family including S_0 and S_1 . Call its union \mathcal{H} . Next, consider the cu-surface $\mathcal{L} := \mathcal{L}(\ell_1^u)$ as in (4). By dimensional reason, ℓ_1^u divides \mathcal{L} into two connected components C_L and C_R , the left and right ones in the obvious sense. The curve $\mathcal{H} \cap \mathcal{L}$ must lie entirely in either C_L or C_R , for otherwise there exists an ss-disc intersecting both $\ell^u \subset V_0^C$ and $\ell_1^u \subset V_1^C$, contradicting (A2). In particular, we have that $S_0 \cap \mathcal{L}$ and $S_1 \cap \mathcal{L}$ lie in the same component. The claim then follows from the transversality assumption as it says that, for a small neighborhood I of t_1 , the intersection $(\bigcup_{t \in I} S_t) \cap \mathcal{L}$ has points in both components.

By construction, $\mathcal{S}' := \bigcup_{t \in [t_1, t_2]} S_t$ is a folding manifold. See the right picture of Figure 3.3 for an illustration. If every ss-disc of \mathcal{S}' crosses the base of V_1 , then we go to case 1 and finish the proof. Otherwise, we repeat the above procedure and obviously we must arrive at case 1 after finite repetitions. \square

Proposition 3.9. *If \mathcal{S} is a folding manifold of a separated standard cs-blender Λ , then \mathcal{S} has a non-transverse intersection with $W^u(\Lambda)$.*

The proof of this proposition is the same as that of [BD12, Proposition 4.4], where one replaces [BD12, Remark 4.3] and [BD12, Lemma 4.5] by Lemmas 3.7 and 3.8, respectively. For completeness, we reprove it here.

Sketch of proof. The idea is similar to that of the proof of Proposition 2.11, with a folding manifold \mathcal{S} in place of an ss-disc. Suppose in Lemma 3.8 there are no tangencies found in every backward iterate of the folding manifold \mathcal{S} . Then, we obtain a nested sequence $\{\hat{\mathcal{S}}_n \subset \mathcal{S}\}_{n \geq 0}$ with $\mathcal{S}_0 = \mathcal{S}$ such that $\mathcal{S}_n := g^{-n}(\hat{\mathcal{S}}_n)$ is a folding manifold for each n , and $\mathcal{S}' := \bigcap \hat{\mathcal{S}}_n \subset W^u(Q)$ for some point $Q \in \Lambda$ by the connectedness of \mathcal{S}' .

Lemma 3.7 implies that there exists for each n a point $P_n \in \mathcal{S}_n$ such that $T_{P_n} \mathcal{S}_n$ contains a non-zero vector v_n which also lies in \mathcal{C}^u . Denote $\hat{P}_n = g^n(P) \in \hat{\mathcal{S}}_n$ and $\hat{v}_n = Dg^n(v_n) \in T_{\hat{P}_n} \hat{\mathcal{S}}_n$. Passing to subsequences if necessary, one has

$$\hat{P}_n \rightarrow \hat{P} \in \mathcal{S}' \quad \text{and} \quad \hat{v}_n \rightarrow \hat{v} \in T_{\hat{P}} \mathcal{S}'$$

for some point \hat{P} and vector \hat{v} . Let us denote by \mathcal{C}_P^u the cone of cone field \mathcal{C}^u at point P . By the uniform hyperbolicity, the size of the iterated cone $Dg^n(\mathcal{C}_P^u)$ goes to zero exponentially as $n \rightarrow +\infty$. Since one has the hyperbolic splitting $T_P \Pi_i = E_P^{ss} \oplus E_P^{cs} \oplus E_P^u$, it follows that $Dg^n(\mathcal{C}_{P_n}^u)$, and hence \hat{v}_n ,

converge to the unstable space $E_{\hat{P}_n}^u$ as $n \rightarrow +\infty$. Thus, by continuity of the splitting one has $\hat{v} \in E_{\hat{P}}^u$, which is just $T_{\hat{P}}W^u(Q)$. \square

Proof of Theorem 2. We prove only for a separated cs-blender Λ , and the proof for a cu-blender is the same. By construction, for any local unstable leaf $\ell^u \in \Pi_i$, the union of its iterates $\bigcup g^n(\ell^u)$ is dense in $W^u(\Lambda)$. Let us further take ℓ^u from \mathcal{W} (defined in Remark 3.3). By an arbitrarily C^r -small perturbation, we can find a piece W^s of $W^s(\Lambda)$ tangent to ℓ^u . With an additional small perturbation if necessary, we make the tangency quadratic. Then, since W^s is foliated by strong-stale leaves which particularly are ss-disc, unfolding the tangency will create inside W^s a folding manifold attached to ℓ^u . The theorem now follows from Proposition 3.9 and the fact that being a folding manifold is a C^1 -robust property. \square

4 Arrayed standard blenders

We are now in the position to introduce arrayed blenders, which are used to produce uncountably many robust homoclinic tangencies, see Theorem 3. Roughly speaking, a k -arrayed blender is a proper combination of k standard blenders such that the central direction is covered k times in an aligned fashion.

4.1 Base-Center-Gap structure

Recall that we say two sets A and B are separated if no ss-disc intersecting both of them. We now denote this by $A \parallel B$, and write $A \nparallel B$ if there is some ss-disc connecting them. By Remark 2.2 the first component of the map $g|_H$ for a horizontal strip H is strictly monotone in x . We say that $g|_H$ is orientation-preserving if this component is increasing in x , and orientation-reversing if it is decreasing in x .

Definition 4.1 (Base-Center-Gap (BCG) structure of Π_i). For each i consider two open intervals \mathbb{X}_i^C and \mathbb{X}_i^B satisfying $\overline{\mathbb{X}_i^C} \subset \mathbb{X}_i^B \subset \mathbb{X}_i$, and two points x_i^L, x_i^R in the left and, respectively, right (by the usual orientation on \mathbb{R}) components of $\mathbb{X}_i^B \setminus \overline{\mathbb{X}_i^C}$, and denote

$$\mathring{\Pi}_i^L := \Pi_i \cap (\{x_i^L\} \times \mathbb{Y}_i \times \mathbb{Z}_i) \quad \text{and} \quad \mathring{\Pi}_i^R := \Pi_i \cap (\{x_i^R\} \times \mathbb{Y}_i \times \mathbb{Z}_i).$$

We define

- *base*: $\Pi_i^B = \Pi_i \cap (\mathbb{X}_i^B \times \mathbb{Y}_i \times \mathbb{Z}_i)$,
- *center*: $\Pi_i^C = \Pi_i \cap (\mathbb{X}_i^C \times \mathbb{Y}_i \times \mathbb{Z}_i)$,
- *left gap*: Π_i^L , the union of all u-discs S^u satisfying $S^u \nparallel \mathring{\Pi}_i^L$,
- *right gap*: Π_i^R , the union of all u-discs S^u satisfying $S^u \nparallel \mathring{\Pi}_i^R$.

For any horizontal strip H and vertical strip V with $V = g(\tilde{H}) \cap \Pi_i$ for some \tilde{H} of Π_j , we denote

$$H^{B/C/L/R} = H \cap \Pi_i^{B/C/L/R}, \quad \mathring{H}^{L/R} = H \cap \mathring{\Pi}_i^{L/R}, \quad V^{B/C} = g(\tilde{H}^{B/C}) \cap \Pi_i^B,$$

The gaps for vertical strips are defined as follows: if $g|_H$ is orientation-preserving, then

$$\mathring{V}^{L/R} = g(\mathring{H}^{L/R}) \cap \Pi_i^B, \quad V^{L/R} = g(H^{L/R}) \cap \Pi_i^B;$$

if $g|_H$ is orientation-reversing, then

$$\mathring{V}^{L/R} = g(\mathring{H}^{R/L}) \cap \Pi_i^B, \quad V^{L/R} = g(H^{R/L}) \cap \Pi_i^B;$$

We say that an ss-disc S *crosses* $\Pi_i^{B/C/L/R}$ if $S \cap \Pi_i \subset \Pi_i^{B/C/L/R}$; similarly for the strips.

See Figure 4.1 for an illustration, where the size of the gaps (in gray) are determined by the size of the cone fields.




Figure 4.1: The Base-Center-Gap structure projected to \mathbb{X}_i

For a finite collection \mathcal{V} of different vertical strips in Π_i , its base, center and gaps, denote by $\mathcal{V}^B, \mathcal{V}^C, \mathcal{V}^L, \mathcal{V}^R$, are defined as the collections of the corresponding parts of the vertical strips in it.

Definition 4.2 (Arrays of vertical strips). A finite collection \mathcal{V} of k different vertical strips in Π_i is called a *vertical k -array* of Π_i if there exists some ss-disc crossing every $V \in \mathcal{V}$; such ss-discs are said to *cross the array*, and the same for crossing of base, center, gaps of \mathcal{V} .

Let $\partial_x^{L/R} \Pi_i$ be the left/right x -boundary in the obvious sense (see the discussion above Definition 2.5), and the same for $\partial_x^{L/R} H$. Let $V \subset \Pi_i$ be of the form $V = g(H) \cap \Pi_i$ for some $H \subset \Pi_j$. If $g|_H$ is orientation-preserving, then

$$\partial_x^L V = g(\partial_x^L H) \cap \Pi_i, \quad \partial_x^R V = g(\partial_x^R H) \cap \Pi_i,$$

and, if $g|_H$ is orientation-reversing in x , then

$$\partial_x^L V = g(\partial_x^R H) \cap \Pi_i, \quad \partial_x^R V = g(\partial_x^L H) \cap \Pi_i.$$

Definition 4.3 (Base-Center-Gap (BCG) covering property for arrays). A cs-Markov partition $(\{\Pi_i\}, g)$ has the *k -arrayed BCG covering property* if Π_i can be structured such that, with $S \subset \Pi_i^B, V \subset \Pi_i, \mathcal{V} \subset \Pi_i$ denoting its ss-discs, vertical strips and, respectively, vertical k -arrays, the following are satisfied:

- (B1) any S crossing Π_i^B crosses at least one vertical array \mathcal{V}^B of bases,
- (B2) if $S \cap \partial_x^R V^B \neq \emptyset$ for some $V \in \mathcal{V}_1$, then S crosses some \mathcal{V}_2^C such that $\mathcal{V}_2^B \parallel \mathcal{V}_1^L$,
- (B2') if $S \cap \partial_x^L V^B \neq \emptyset$ for some $V \in \mathcal{V}_1$, then S crosses some \mathcal{V}_2^C such that $\mathcal{V}_2^B \parallel \mathcal{V}_1^R$,
- (B3) if $S \cap \partial_x^L \Pi_i^C \neq \emptyset$, then S crosses some \mathcal{V}^C such that $\mathcal{V}^B \parallel \Pi_i^L$,
- (B3') if $S \cap \partial_x^R \Pi_i^C \neq \emptyset$, then S crosses some \mathcal{V}^C such that $\mathcal{V}^B \parallel \Pi_i^R$, and
- (B4) $\Pi_i^{L/R} \parallel \Pi_i^C$ for any Π_i and $V^{L/R} \parallel V^C, V^{L/R} \parallel \partial_x^{L/R} V^B$ for any V .

We say that a cu-Markov partition satisfies the BCG covering property if its inverse satisfies it.

Property (B1) is just the covering property (A1), but stated for arrays. Figure 4.2 provides an illustration of the fulfilment of other properties, except for the symmetric ones: (B3') and $\Pi_i^R \parallel \Pi_i^C$

in (B4), where, in particular, properties (B2), (B2'), (B3) are shown explicitly with the cone field \mathcal{C}^{ss} (in red).




Figure 4.2: The fulfilment of properties (B2)–(B4), where the red cylinders represent the areas covered by the strong-stable cone field \mathcal{C}^{ss}

Definition 4.4 (k -arrayed standard blenders). Given a partially hyperbolic Markov partition that satisfies the k -arrayed BCG covering property, its locally maximal set is called a k -arrayed standard blender. The blender is further called center-stable (cs) for cs-Markov partitions and center-unstable (cu) for cu-Markov partitions.

Remark 4.5. Comparing Definitions 3.2 and 4.3, one sees that an arrayed blender is not necessarily a separated one, and vice versa. The reason is that the BCG covering property enables us to distinguish the left and right (to define the folding manifolds in Section 4.2), based on the fact that the central dynamics in \mathbb{X} is one dimensional. In contrast, the BC covering property essentially provides the notions of inside and outside of Π_i^C so that one can define folding manifolds which are large enough (in the central direction) – it crosses the boundary $\partial\mathbb{X}_i^C$ and hence have parts both inside and outside Π_i^C . Since $\partial\Pi_i^C$ is essentially determined by $\partial\mathbb{X}_i^C$, the BC covering property can be generalized to the case of d_c -dimensional ($d_c > 1$) center by distinguish the inside and outside of $\partial\mathbb{X}_i^C = S^{d_c}$, the d_c -dimensional sphere. Hence, we believe that separated blenders can be constructed for Markov partitions with higher center dimension to create robust tangencies of higher corank, that is, tangencies at which the two tangent spaces share a common subspace of dimension larger than 1.

4.2 Reproducing lemma

Let $\mathcal{S} \subset \Pi_i^B$ be a submanifold of dimension $(d_{ss} + 1)$ which is the union of a smooth family $\{S_t\}_{t \in [0,1]}$ of ss-discs crossing Π_i^B .

Definition 4.6 (Prefolding manifolds). The submanifold \mathcal{S} is called a *right/left-prefolding manifold* if

- S_0 and S_1 intersect a leaf $\ell^u \subset (W^u(\Lambda) \cap \Pi_i^{L/R})$, and
- $S_t \cap \Pi_i^C \neq \emptyset$ for some $t \in (0, 1)$.

Definition 4.7 (Folding manifolds). The submanifold \mathcal{S} is called a *right/left-folding manifold related to a vertical strip V* if

- S_0 and S_1 intersect a leaf $\ell^u \subset (W^u(\Lambda) \cap V^{L/R})$,

- $S_t \cap \ell^u = \emptyset$ for all $t \in (0, 1)$, and
- $S_t \cap V^C \neq \emptyset$ for some $t \in (0, 1)$;

it is further called *exact* if every ss-disc of \mathcal{S} crosses V^B .

See Figure 4.3 for an illustration.




Figure 4.3: A right-prefolding manifold (\mathcal{S}_1), a right-folding manifold (\mathcal{S}_2) related to V_2 , and an exact right-folding manifold (\mathcal{S}_3) related to V_1

Definition 4.8 (Folding arrays). The submanifold \mathcal{S} is called a *right/left-folding array related to a vertical k -array $\mathcal{V} = \{V_i\}$* if, for $i = 1, \dots, k$, there exist unstable leaves ℓ_i^u and $t_i, t'_i \in (0, 1]$ with $t_i < t'_i$ such that $\mathcal{S}_i := \bigcup_{t \in [t_i, t'_i]} S_t$ are right-/left-folding manifolds related to V_i . It is further called *exact* if all \mathcal{S}_i are exact.

Lemma 4.9 (Reproducing lemma). *Let the k -arrayed BCG covering property be satisfied. If \mathcal{S} is an exact folding array related to $\mathcal{V} = \{V_i\}_{i=1}^k$, then for each i the preimage $g^{-1}(\mathcal{S} \cap V_i)$ contains an exact folding array in some Π_j .*

The novelty in this result is the nonstop reproducing process, in contrast with Lemma 3.8 where the process terminates upon detecting a tangency (since the arguments there for generating a new folding manifold is based on the absence of tangencies). This property is crucial for obtaining uncountably many tangencies in Proposition 4.14.

Since for each i the intersection $\mathcal{S} \cap V_i$ contains an exact folding manifold related to V_i , Lemma 4.9 is an immediate consequence of the following three lemmas.

Lemma 4.10. *The preimage by g of an exact folding manifold is a prefolding manifold.*

Lemma 4.11. *Every prefolding manifold contains a folding array.*

Lemma 4.12. *Every folding array contains an exact folding array.*

Proof of Lemma 4.10. Let \mathcal{S} be an exact folding manifold related to V . It follows from definitions of the BCG structure and folding manifolds that one has $g^{-1}(\mathcal{S} \cap V^{L/R/C}) \subset \Pi_j^{L/R/C}$ for some j if $g^{-1}|_V$ is orientation preserving or $g^{-1}(\mathcal{S} \cap V^{L/R/C}) \subset \Pi_j^{R/L/C}$ if $g^{-1}|_V$ is orientation reversing. Since the cone field \mathcal{C}^{ss} is backward-invariant, in either case the preimage is a prefolding manifold. \square

Proof of Lemma 4.11. Let $\mathcal{S} = \bigcup_{t \in [0,1]_{S_t}}$ be a right-prefolding manifold of Π_i . Since some S_t crosses Π_i^C by definition, there exists S_{t_0} such that $S_{t_0} \cap \partial_x^L \Pi_i^C \neq \emptyset$. Let \mathcal{V} be the vertical k -array given by (B3). Take any $V_j \in \mathcal{V}$. Since S_{t_0} crosses \mathcal{V}^C , it particularly crosses V_j^C . We prove the lemma in several steps.

Step 1: there exists $t_1 \in (0, t_0)$ such that $S_{t_1} \cap \mathring{V}_j^L \neq \emptyset$ (see Definition 4.3).

By the continuity of S_t on t and the fact that ss-discs are defined for all z values (see Definition 2.5), there must exist some $t_1 \in (0, t_0)$ such that $S_{t_1} \cap \mathring{V}_j^L \neq \emptyset$ or $S_{t_1} \cap \mathring{V}_j^R \neq \emptyset$. We prove below that the latter case implies the existence of some $t' \in (0, t_0)$ with $S_{t'} \cap \mathring{V}_j^L \neq \emptyset$. Denote by ℓ^u the unstable leaf in the definition of a prefolding manifold. Let us attach to each point of ℓ^u an ss-disc, including S_0 and S_1 , to form a su-surface \mathcal{H} of codimension one which divides Π_i into two connected components, the left one C_L and the right one C_R in the obvious sense. So, we have $\Pi_i \setminus \mathcal{H} = C_L \cup C_R$. Similarly, we take a family of ss-discs including S_{t_0} such that every disc intersects V_j^C . It forms another codimension-one surface \mathcal{H}_0 , which lies in C_R by (B3) and further divides C_R into two components C_{R_1} and C_{R_2} , labeled from left to right. Property (B4) implies $V_j^L \subset \Pi_i \setminus C_{R_2}$ and $V_j^R \subset C_{R_2}$. Since (B3) also implies $V_j^L \subset C_R$, one has $V_j^L \subset C_{R_1}$. The desired statement then follows immediately from the continuity since before intersecting \mathring{V}_j^R the family $\{S_t\}$ must go through C_{R_1} , and hence intersect \mathring{V}_j^L .

Step 2: S_{t_1} intersects an unstable leaf $\ell_1^u \in W^u(\Lambda) \cap V_j^L$.

Property (B1) implies that every ss-disc crossing Π_i^B intersects $W^u(\Lambda)$ (see Proposition 2.11). In particular, there exists some unstable leaf $\ell_1^u \subset W^u(\Lambda)$ intersecting S_{t_1} , which lies in V_j^B by Remark 2.12. We in fact have $\ell_1^u \subset V_j^L$. To see this, let $V_j = g(H_j)$ for some horizontal strip H_j . Assume that $g|_{H_j}$ is orientation preserving in x , so $\mathring{V}_j^L = g(\mathring{H}_j^L)$. By the invariance of the unstable foliation and the cone field \mathcal{C}^{ss} , we have that the ss-disc $(g|_{H_j})^{-1}(S_{t_1})$ intersects both \mathring{H}_j^L and the unstable leaf $(g|_{H_j})^{-1}(\ell_1^u)$. Hence, $\ell_1^u \subset V_j^L$ by the definition of gaps. Obviously, the result also holds when $g|_H$ is orientation reserving in x .

Step 3: there exists $t'_1 \in [t_1, t_0)$ such that $S_{t'_1}$ intersects ℓ_1^u and $S_t \cap \ell_1^u = \emptyset$ for $t \in (t'_1, t_0]$.

Such t'_1 does not exist only if S_t intersects ℓ_1^u for every $t \in [t_1, t_0]$. This is impossible since S_{t_0} belongs to \mathcal{H}_0 , the right boundary of C_{R_1} , while $\ell_1^u \subset \overline{V}_j^L \subsetneq C_{R_1}$ by (B3) and (B4).

Step 4: there exists $t_2 \in (t_0, 1)$ such that S_{t_2} intersects ℓ_1^u and $S_t \cap \ell_1^u = \emptyset$ for $t \in [t_0, t_2)$.

Since S_{t_0} and S_{t_1} belong to the right, and, respectively, left boundaries of C_{R_1} while $\ell_1^u \subset \overline{V}_j^L \subsetneq C_{R_1}$, the desired value t_2 follows from the continuity of $\{S_t\}$.

Step 5: completion of the proof.

By construction, $\mathcal{S}_j = \bigcup_{t \in [t'_1, t_2]} S_t$ is a right-folding manifold of V_j . Repeat the above arguments for all k vertical strips in \mathcal{V} , we find k right-folding manifolds \mathcal{S}_j and their union is a right-folding array related to \mathcal{V} . The proof for the case where \mathcal{S} is a left-prefolding manifold is the same, with using (B3') instead of (B3). \square

Remark 4.13. It can be seen from the proof that, with using (B2) instead of (B3), if \mathcal{S} is a right-folding manifold \mathcal{S} related to some vertical strip V (instead of a right-prefolding manifold) such that $\mathcal{S} \cap \partial_x^R V \neq \emptyset$, then \mathcal{S} contains a folding array given by (B2). Similarly, for left-folding manifolds, using (B2').

Proof of Lemma 4.12. Let $\mathcal{S} = \bigcup_{t \in [0,1]} S_t$ be a right-folding array of a vertical k -array \mathcal{V}_0 . If \mathcal{S} is not

exact, then by definition there exists some $V_0 \in \mathcal{V}_0$ such that the right-folding manifold \mathcal{S}_0 associated with V_0 is not exact. So, \mathcal{S}_0 contains some ss-disc S_{t_0} such that S_{t_0} intersects $\partial_x^L V_0^B$ or $\partial_x^R V_0^B$.

We claim that $S_{t_0} \cap \partial_x^R V_0^B \neq \emptyset$. To prove the claim, first note that the gap V_0^L divides V_0 into two connected components C_L and C_R , the left and right ones, such that $\partial_x^L V_0^B \subset C_L$. It is evident that if there exists an ss-disc intersecting both $\partial_x^L V_0^B$ and C_R , then one can construct another ss-disc intersecting $\partial_x^L V_0^B$ and V^L . On the other hand, if $\ell^u \subset V_0^L$ is the unstable leaf associated with \mathcal{S}_0 as in Definition 4.7, then $\ell^u \cap S_{t_0} = \emptyset$, and hence S_{t_0} intersects $(V^L \cup C_R)$. The claim then follows, since $S_{t_0} \cap \partial_x^L V_0^B \neq \emptyset$ would contradict $\partial_x^L V_0^B \parallel V^L$ given by (B4).

Thus, by (B2), there exists a vertical k -array \mathcal{V}_1 such that S_{t_0} crosses \mathcal{V}_1^C . One then obtains a right-folding array of \mathcal{V}_1 by the same arguments in the proof of Lemma 4.11, with $\mathcal{S}_0, V_0, \mathcal{V}_1$, (B2) in place of $\mathcal{S}, \Pi_i, \mathcal{V}$, (B3), see Remark 4.13. If this folding array is not exact, then we repeat this procedure. Since there are only finitely many vertical arrays and the new folding array obtained at each step is strictly smaller than the previous one, we will obtain an exact folding array after finite steps. The proof for the case where \mathcal{S} is a left-folding array is the same, with (B2') in place of (B2). \square

4.3 Robust presence of uncountably many homoclinic tangencies. Proof of Theorem 3

Let Λ be a k -arrayed cs-blender induced from a cs-Markov partition $(\{\Pi_i\}, g)$. Let us denote by \mathcal{V}_{ij} the vertical arrays of Π_i involved in the k -arrayed BCG covering property in Definition 4.3, where $j = 1, \dots, j^*$ for some integer $j^* > 1$ depending on i . We label the corresponding vertical strips by $V_{ijs} \in \mathcal{V}_{ij}$ with $s = 1, \dots, k$. We call the connected components of $W^u(\Lambda) \cap \Pi_i$ the local unstable leaves of $W^u(\Lambda)$ (in Π_i). Denote by $\Sigma_+^k := \{1, \dots, k\}^{\mathbb{N}_0}$ the set of one-sided sequences of k symbols.

Proposition 4.14. *For any exact folding array \mathcal{S} of Λ , each $\underline{s} = (s_n)_{n \geq 0} \in \Sigma_+^k$ corresponds to a sequence $(i_n, j_n, s_n)_{n \geq 0}$ such that the local unstable leaf defined by*

$$\ell^u(\underline{s}) = \left(\bigcap_{n \geq 0} g^n(V_{i_n j_n s_n}) \right) \cap \Pi_{i_0} \quad (5)$$

intersects \mathcal{S} non-transversely. Moreover, this correspondence is injective.

Proof. Suppose \mathcal{S} is related to $\mathcal{V}_{i_0 j_0}$ in Π_{i_0} . By the exactness of \mathcal{S} , for any $s_0 \in \{1, \dots, k\}$, the intersection $\mathcal{S} \cap V_{i_0 j_0 s_0}$ contains an exact folding manifold related to $V_{i_0 j_0 s_0}$. The reproducing lemma (Lemma 4.9) implies that the preimage $g^{-1}(\mathcal{S} \cap V_{i_0 j_0 s_0})$ contains an exact folding array related to an array $\mathcal{V}_{i_1 j_1}$ in some Π_{i_1} , where i_1 and j_1 depend on the choice of s_0 . For any $s_1 \in \{1, \dots, k\}$, the intersection of the new exact folding array with $V_{i_1 j_1 s_1}$ contains an exact folding manifold. Invoking the reproducing lemma on $V_{i_1 j_1 s_1}$, we find another exact folding array related to some array $\mathcal{V}_{i_2 j_2}$. Repeating this procedure infinitely many times, we obtain a sequence $\underline{s} = (s_n)_{n \geq 0} \in \Sigma_+^k$. By construction, it corresponds to a sequence $(i_n, j_n, s_n)_{n \geq 0}$ with i_n and j_n depending on s_{n-1} for $n > 0$ such that, with $\mathcal{S}_0(\underline{s}) := \mathcal{S}$, each of the preimages $\mathcal{S}_{n+1}(\underline{s}) := g^{-1}(\mathcal{S}_n(\underline{s}) \cap V_{i_n j_n s_n})$ contains an exact folding manifold. Since Lemma 3.7 obviously holds for exact folding manifolds, one can argue as in the proof of Proposition 3.9 to conclude that $\mathcal{S} \cap W^u(\Lambda)$ has a non-transverse intersection point in $\bigcap_{n \geq 0} \hat{\mathcal{S}}_n(\underline{s})$, where $\hat{\mathcal{S}}_n(\underline{s}) := g^n(\mathcal{S}_n(\underline{s}))$. By construction, this intersection point belongs to the leaf defined by (5), and for different sequences the corresponding leaves are different. \square

Proof of Theorem 3. By the symmetry of the problem, we only prove for a k -arrayed cs-blender Λ .

The goal is to create an exact folding array in $W^s(\Lambda)$ and apply Proposition 4.14. By Lemmas 4.11 and 4.12, it suffices to find a prefolding manifold. The robustness in the theorem is automatic since being a prefolding manifold or an (exact) folding array is a C^1 -robust property.

For that, consider any Π_i . By (B1) (and Proposition 2.11), there exists a local unstable leaf $\ell^u \subset W^u(\Lambda) \cap \Pi_i^L$. By construction this leaf must contain a point in Λ . Indeed, ℓ^u intersects all local stable leaves in Π_i and all intersection points belong to the locally maximal set Λ . Since periodic points are dense in Λ , we can take some periodic point P such that $W_{loc}^u(P) \subset \Pi_i^L$. Similarly, we find a periodic point P' with $W_{loc}^u(P') \subset \Pi_i^R$. Since $W^u(P)$ is dense in $W^u(\Lambda)$, by an arbitrarily small perturbation, we can create a quadratic tangency between a piece W^s of $W^s(\Lambda)$ and $W_{loc}^u(P)$. There are now two cases.

The first case is when W^s at the tangency point is pointing to the right in the x -direction, see the upper picture in Figure 4.4. In this case, we unfold the tangency by moving W^s to the right in the x -direction and denote by μ this displacement. So the tangency exists at $\mu = 0$, and there are two transverse intersection points of W^s with $W_{loc}^u(P)$ at $\mu > 0$. Note that W^s is foliated by strong-stable leaves (which in particular are ss-discs), so these intersection points belong to two strong-stable leaves in W^s , denoted by ℓ_0^{ss} and ℓ_1^{ss} . Let τ be the period of P . Due to the contraction in x , for any fixed integer $n > 0$, there exists $\mu_1 > 0$ such that $g^{-n\tau}(W^s)$ intersects the right boundary $\partial_x^R \Pi_i$ at $\mu = \mu_1$. By continuity, one can find $\mu_2 \in (0, \mu_1)$ such that $g^{-n\tau}(W^s)$ intersects Π_i^C but still lies in Π_i^B at $\mu = \mu_2$. Now, the part of $g^{-n\tau}(W^s)$ bounded by $g^{-n\tau}(\ell_0^{ss})$ and $g^{-n\tau}(\ell_1^{ss})$ is a right-prefolding manifold, as desired. Moreover, μ_2 , and hence the size of the perturbation, tend to zero as $n \rightarrow \infty$.

In the second case, W^s at the tangency point is pointing to the left, see the lower picture in Figure 4.4. We now only perturb the tangency to a quadratic one and do not unfold it. Let us write it as a union of strong-stable leaves, $W^s = \bigcup_{t \in [0,1]} \ell_t^{ss}$, where the tangency point belongs to $\ell_{t_0}^{ss}$ for some $t_0 \in (0, 1)$. Due to the contraction in x , the preimage $g^{-n\tau}(W^s)$ for a large n must intersect $W_{loc}^u(P')$ in at least two leaves $g^{-n\tau}(\ell_{t_1}^{ss})$ and $g^{-n\tau}(\ell_{t_2}^{ss})$ satisfying $t_1 \in (0, t_0)$ and $t_2 \in (t_0, 1)$. Evidently, the part of $g^{-n\tau}(W^s)$ bounded by these two leaves, i.e., $\bigcup_{t \in [t_1, t_2]} g^{-n\tau}(\ell_t^{ss})$, is a left-prefolding manifold. \square




Figure 4.4: Creating prefolding manifolds in the two cases in the proof of Theorem 3, projected to the (x, y) -space

5 Nearly-affine blender system (NABS)

In this section, we prove Theorem 4. We begin with introducing nearly-affine blender systems, which generalizing the renormalization results obtained in [LT24].

Denote $d = \dim \mathcal{M}$. Let $\{\Pi_\delta\}_{\delta>0}$ be a family of embedded d -dimensional unit cubes, that is, there is a family of charts $\{(U_\delta, \psi_\delta)\}_{\delta>0}$ with $\Pi_\delta \subset U_\delta$ and $\varphi_\delta(\Pi_\delta) = [-1, 1]^d \subset \mathbb{R}^d$. For each fixed δ , we associate with Π_δ a family $\mathcal{F}_\delta := \{F_{i,\delta}\}_{i \in \mathbb{N}}$ of diffeomorphisms where the intersections of domains of $F_{i,\delta}$ with Π_δ are pairwise disjoint and so are the intersections of ranges of $F_{i,\delta}$ with Π_δ .

Definition 5.1 (Nearly-affine blender system). The family $\{\Pi_\delta, \mathcal{F}_\delta\}$ is called a *center-stable nearly-affine blender system (cs-NABS)* if the charts $\{(U_\delta, \varphi_\delta)\}$ can be chosen such that, for any $F_{i,\delta} \in \mathcal{F}_\delta$ and for any point

$$(x, y, z) \in \psi_\delta(\Pi_\delta) = [-1, 1] \times [-1, 1]^{d_u} \times [-1, 1]^{d_{ss}}, \quad 1 + d_u + d_{ss} = d,$$

one has that $(\bar{x}, \bar{y}, \bar{z}) = \varphi_\delta \circ F_{i,\delta} \circ \varphi_\delta^{-1}(x, y, z)$ if and only if

$$\begin{aligned} \bar{x} &= A_i(\delta)x + B_i(\delta) + \psi_{1,i}(x, \bar{y}, z; \delta) =: \hat{\psi}_{1,i}(x, \bar{y}, z; \delta), \\ \bar{y} &= \psi_{2,i}(x, \bar{y}, z; \delta), \\ \bar{z} &= \psi_{3,i}(x, \bar{y}, z; \delta), \end{aligned} \tag{6}$$

where the coefficients A_i, B_i and functions ψ satisfy that

- (S1) $|A_i(\delta)| - \alpha = o(1)_{\delta \rightarrow 0} + o(1)_{\delta \cdot i \rightarrow \infty}$ for some constant $\alpha \in (0, 1)$. Here it means that given any $\varepsilon > 0$, there exists $\delta_0 > 0$ satisfying that, for any $\delta \in (0, \delta_0)$, one can find $N(\delta) \in \mathbb{N}$ such that $|A_i(\delta)| - \alpha < \varepsilon$ for all $i > N(\delta)$,
- (S2) for every fixed δ , the set $\{B_i(\delta)\}_{i \in \mathbb{N}}$ is dense in an interval I , containing 0 and independent of δ and i , and
- (S3) the maps $\psi_{j,i}$ ($j = 1, 2, 3$) are defined on $\{(x, \bar{y}, z) \in \Pi\}$ and $\|\psi_{j,i}\|_{C^1} = o(1)_{\delta \rightarrow 0} + o(1)_{\delta \cdot i \rightarrow \infty}$.

The family $\{\Pi_\delta, \mathcal{F}_\delta\}$ is called a *center-unstable nearly-affine blender system (cu-NABS)* if (S1) is replaced by

$$(S1') \quad |A_i(\delta)| - \alpha' = o(1)_{\delta \rightarrow 0} + o(1)_{\delta \cdot i \rightarrow \infty} \text{ for some constant } \alpha' \in (1, \infty).$$

It is called a *full nearly-affine blender system* if it is the union of a cs-NABS and a cu-NABS.

Let us show that a cs-NABS defines infinitely many cs-Markov partitions, which are of the same form as the one in Example 2.3. For brevity, we identify $F_{i,\delta}$ with $\varphi_\delta \circ F_{i,\delta} \circ \varphi_\delta^{-1}$, so we can take $\Pi_{i,\delta} = [-1, 1]^d =: \Pi$. We fix a δ and consider any finite subset \mathcal{I} of

$$\hat{\mathcal{I}} = \{i \in \mathbb{N} : \sup_{(x, \bar{y}, z) \in \Pi} |\hat{\phi}_{1,i}(x, \bar{y}, z)| < 1\}. \tag{7}$$

Denote by Π_i the intersection of Π with the domain of $F_{i,\delta}$ and define $g_\delta : \bigcup_{i \in \mathcal{I}} \Pi_i \rightarrow \Pi$ by $g_\delta(x, y, z) = F_{i,\delta}(x, y, z)$ if $(x, y, z) \in \Pi_i$. Up to shrinking Π and rescaling it to the unit size, we can assume that g_δ is a diffeomorphism on a small neighborhood of $\bigcup_{i \in \mathcal{I}} \Pi_i$.

Claim 5.2. *For any fixed δ , the pair $(\{\Pi_i\}_{i \in \mathcal{I}}, g_\delta)$ is a cs-Markov partition, and we call it a cs-Markov partition of $\{\Pi_\delta, \mathcal{F}_\delta\}$ with the specification (δ, \mathcal{I}) .*

One sees that Example 2.3 illustrates the case where $\text{card } \mathcal{I} = 2$.

Proof. To verify the the first item of Definition 2.1, one sees that by taking Π is immediate since By (6) one has

$$\Pi_i = \{(x, y, z) : x \in [-1, 1], y \in \psi_{2,i}(x, [-1, 1]^{d_u}, z; \delta), z \in [-1, 1]^{d_{ss}}\}, \tag{8}$$

and property (S3) guarantees that $\psi_{2,i}$ takes values in $(-1, 1)$ for all small δ and large i . The first item of Definition 2.1 is then satisfied by taking $\mathbb{X}_i \times \mathbb{Y}_i \times \mathbb{Z}_i = [-1, 1]^d$ for all i . The second item is immediate since the right-hand of (6), considered as a map $(x, \bar{y}, z) \mapsto (\bar{x}, y, \bar{z})$, is a contraction by (S1) and (S3), and takes Π into itself by (S3) and the choice of \mathcal{I} . The cone fields in (3) can be obtained by direct computations with formula (6) and the cone coefficients satisfy

$$K := \max\{K_i^u, K_i^s, K_i^{ss}\} = o(1)_{\delta \rightarrow 0} + o(1)_{\delta \cdot i \rightarrow \infty}, \quad (9)$$

see [LT24, Lemma 3.1]. The claim is proven. \square

When the NABS $\{\Pi_\delta, \mathcal{F}_\delta\}$ is center-unstable, we can also define cu-Markov partitions from the same procedure as above. To see this, we use the Implicit Function Theorem to find x as a function of (\bar{x}, \bar{y}, z) from the first equation of (6):

$$x = \frac{1}{A_i(\delta)} \bar{x} - B_i(\delta) + \tilde{\psi}_{1,i}(\bar{x}, \bar{y}, z),$$

for some $\tilde{\psi}_{1,i}$ with $\|\tilde{\psi}_{1,i}\|_{C^1} = o(1)_{\delta \rightarrow 0} + o(1)_{\delta \cdot i \rightarrow \infty}$. Then, one has that $(x, y, z) = \varphi_\delta \circ F_{i,\delta}^{-1} \circ \varphi_\delta^{-1}(\bar{x}, \bar{y}, \bar{z})$ if and only if the above equation and the last two of (6) are satisfied. Clearly, this defines a cs-NABS $\{\Pi_\delta, \mathcal{F}'_\delta\}$ with $\mathcal{F}'_\delta = \{F_{i,\delta}^{-1}\}$, and cs-Markov partitions of this NABS are cu-Markov partition of the original one.

Theorem 5.3. *Let $\{\Pi_\delta, \mathcal{F}_\delta\}$ be a nearly-affine blender system. Up to replacing \mathcal{F}_δ by some iteration $\mathcal{F}_\delta^n := \{F_{i,\delta}^n\}$, for any integer $k \geq 1$, there exists a specification (δ, \mathcal{I}) such that the corresponding partially hyperbolic Markov partition induces a standard blender which is simultaneously separated and k -arrayed. It is center-stable/-unstable if $\{\Pi_\delta, \mathcal{F}_\delta\}$ is center-stable/-unstable.*

We postpone the proof of this theorem to Section 5.3, after we prove Theorem 4.

5.1 NABS near heterodimensional cycles. Proof of Theorem 4

In this section, we prove Theorem 5.5 below, which implies Theorem 4, and strengthens the main finding of [LT24] that any coindex-1 heterodimensional cycle can generate

Definition 5.4 (C^1 -robust heterodimensional dynamics). We say that a diffeomorphism f has C^1 -robust heterodimensional cycles involving two hyperbolic basic sets Λ and Σ if there exists a C^1 neighborhood \mathcal{U} of f such that $W^u(\Lambda_g) \cap W^s(\Sigma_g) \neq \emptyset$ and $W^u(\Sigma_g) \cap W^s(\Lambda_g) \neq \emptyset$ for every $g \in \mathcal{U}$.

Let $f \in \text{Diff}^r(\mathcal{M})$, $r = 1, \dots, \infty, \omega$, have a coindex-1 heterodimensional cycle involving two saddles O_1 and O_2 with $\text{ind}(O_1) + 1 = \text{ind}(O_2)$. Recall that we denote by λ the center-stable multiplier of O_1 and by γ the center-unstable multiplier of O_2 .

For two transitive hyperbolic sets Λ_1 and Λ_2 with different indices, by saying Λ_1 form a C^1 -cycle with Λ_2 , we mean they form C^1 -robust heterodimensional dynamics. Recall that, by saying a transitive hyperbolic Λ is C^1 -wild, we mean it exhibits uncountably many C^1 -robust homoclinic tangencies, and $\overset{\text{h.r.}}{\sim}$ denotes the homoclinic relation between two hyperbolic sets.

Theorem 5.5. *Given any integer $k > 1$, f can be approximated in the C^r topology by a diffeomorphism g , which has up to two standard blenders, a center-stable one Λ_1 and a center-unstable one Λ_2 , each of which is simultaneously separated and k -arrayed with $\Lambda_i \overset{\text{h.r.}}{\sim} O_{i,g}$ ($i = 1, 2$). More specifically,*

- in the saddle case,
 - when $|\alpha| \leq 1$, Λ_1 exists, forming a C^1 -cycle with a non-trivial hyperbolic basic set $\Lambda_2 \stackrel{\text{h.r.}}{\simeq} O_{2,g}$, and
 - when $|\alpha| \geq 1$, Λ_2 exists, forming a C^1 -cycle with a non-trivial hyperbolic basic set $\Lambda_1 \stackrel{\text{h.r.}}{\simeq} O_{1,g}$;
- in the saddle-focus case, both Λ_1 and Λ_2 exist, forming a C^1 -cycle, and,
 - Λ_1 is C^1 -wild if γ is nonreal, and
 - Λ_2 is C^1 -wild if λ is nonreal;
- in the double-focus case, both Λ_1 and Λ_2 exist, forming a C^1 -cycle, and either Λ_1 or Λ_2 is C^1 -wild.

In all cases, Λ_i is always C^1 -wild if O_i originally has a homoclinic tangency.

To prove the theorem, we now summarize the renormalization results in [LT24], which indicate that each heterodimensional cycle is associated with an NABS. Take a *fragile* heteroclinic orbit \mathcal{T}^0 from the non-transverse intersection $W^u(O_1) \cap W^s(O_2)$ and a *robust* heteroclinic orbit \mathcal{T}^1 from the transverse intersection $W^u(O_2) \cap W^s(O_1)$. We consider the closed invariant set

$$\Gamma := \mathcal{O}(O_1) \cup \mathcal{O}(O_2) \cup \mathcal{T}^0 \cup \mathcal{T}^1, \quad (10)$$

where $\mathcal{O}(\cdot)$ denotes the orbit of a point. In [LT24], a heterodimensional cycle involving O_1 and O_2 is defined as the set Γ . To facilitate the presentation, we will do so in the rest of the section.

The key result in [LT24] regarding blenders is the renormalization of the first return maps for a non-degenerate cycle Γ of a C^r ($r = 2, \dots, \infty, \omega$) diffeomorphism, where the non-degeneracy conditions can be fulfilled by an arbitrarily C^r -small perturbation. Each first return map is a composition of four maps: two local maps near the orbits of O_1 and O_2 which are the restrictions of the period maps, and two transition maps along the heteroclinic orbits \mathcal{T}^0 and \mathcal{T}^1 , connecting neighborhoods of two points in $\mathcal{T}^{0,1}$ with one near O_1 and the other near O_2 . By considering different iteration time near the two saddle orbits, we obtain a family of first return maps which after renormalization form an NABS. Whether this NABS is center-stable or center-unstable is determined by the contraction/expansion in the central dynamics. As discussed above Theorem 4, the central dynamics for saddle cycles are tightly related to the quantity α , which in fact appears as certain first partial derivative of the transition map \mathcal{T}^1 . In particular, $|\alpha| \neq 1$ for non-degenerate saddle cycles and the resulting NABS satisfies $A_i \rightarrow \alpha$ in (6). We refer the readers to [LT24, Sections 2.1–2.3] for a detailed description of the first return maps and non-degeneracy conditions.

Theorem 5.6 ([LT24]). *Let $f \in \text{Diff}^r(\mathcal{M})$, $r = 2, \dots, \infty, \omega$, have a non-degenerate heterodimensional cycle Γ of coindex 1, and let U be any neighborhood of Γ . Then, up to an arbitrarily C^r -small perturbation which does not destroy Γ when it is of saddle-focus or double-focus type, there exist a family $\{\Pi_\delta\}_\delta$ of embedded cubes in U and sequences $\{n_i(\delta)\}_{i,\delta}$ of positive integers which are first return times³ of Π_δ such that the family $\{\Pi_\delta, \mathcal{F}_\delta\}$ with $\mathcal{F}_\delta = \{F_{i,\delta} := f^{n_i(\delta)}|_{f^{-n_i(\delta)}(\Pi_\delta)}\}$ is a nearly-affine blender system. This system is*

- either center-stable or center-unstable if Γ is of saddle type, depending on whether $|\alpha| < 1$ or $|\alpha| > 1$, and
- full if Γ is of saddle-focus or double-focus type.

Any standard cs-blender Λ_1 and cu-blender Λ_2 arisen from the NABS satisfy, up to a further arbitrarily C^r -small perturbation, that

³ Namely, $f^n(f^{-n_i(\delta)}(\Pi_\delta) \cap \Pi_\delta) \cap \Pi_\delta = \emptyset$ for $0 < n < n_i(\delta)$.

- in the saddle case,
 - if $|\alpha| < 1$, either $\Lambda_1 \stackrel{\text{h.r.}}{\sim} O_1$ or it forms a C^1 -cycle with a non-trivial hyperbolic basic set $\Lambda_2' \stackrel{\text{h.r.}}{\sim} O_2$,
 - if $|\alpha| > 1$, either $\Lambda_2 \stackrel{\text{h.r.}}{\sim} O_2$ or it forms a C^1 -cycle with a non-trivial hyperbolic basic set $\Lambda_1' \stackrel{\text{h.r.}}{\sim} O_1$, and,

moreover, if either O_1 or O_2 is contained in a non-trivial hyperbolic set, then, in each of the above two cases, the two results hold simultaneously; and
- in the saddle-focus and double-focus cases, Λ_1 and Λ_2 form a C^1 -cycle with $\Lambda_1 \stackrel{\text{h.r.}}{\sim} O_1$ and $\Lambda_2 \stackrel{\text{h.r.}}{\sim} O_2$, where the former homoclinic relation can be obtained without destroying the original heterodimensional cycle if γ is nonreal, and the same for the latter if λ is nonreal.

This theorem follows from several results in [LT24] and is not stated explicitly there. We detail its references in the [Appendix](#). We also need the following result:

Lemma 5.7 ([BDL]). *Let O_1 and O_2 have a heterodimensional cycle of coindex-1, and let at least one of the central multipliers λ and γ be nonreal. Then, there exists an arbitrarily C^r -small ($r = 1, \dots, \infty, \omega$) perturbation which yields a quadratic homoclinic tangency*

- to the continuation of O_1 if γ is not real; and
- to the continuation of O_2 if λ is not real.

Note that, even if both λ and γ are not real, only one of the above two statements can be guaranteed. For completeness, let us sketch the main idea of the proof of this lemma. We only discuss the first statement, as the second one can be achieved by considering f^{-1} . By the definition of a heterodimensional cycle, we can find two pieces $W^u \subset W^u(O_1)$ and $W^s \subset W^s(O_1)$ near O_2 , intersecting $W_{loc}^s(O_2)$ and $W_{loc}^u(O_2)$, respectively. Since γ is complex, the forward iterates of W^u are rotating in the central direction, and, in particular, the rotation angles form a dense in 2π when the argument of γ , say ω , is irrational. When the cycle is in general position, one can find an iterate of W^u such that it has a tangent vector with arbitrarily small angle to some tangent vector of W^s . Thus, a tangency can be created by unfolding the cycle and changing the argument of λ , say ω , at the same time. Indeed, it is proven in [BDL] that, if the cycle is non-degenerate, then the family $\{f_{\mu,\omega}\}$, where μ denotes the distance between W^u and $W_{loc}^s(O_2)$ and $f_{0,\omega^*} = f$, has a sequence $\{(\mu_i, \omega_i)\} \rightarrow (0, \omega^*)$ of parameter values for each of which the continuation of O_1 has a homoclinic tangency.

Proof of Theorem 5.5. We prove for the three cases separately. For brevity, by a perturbation we always mean an arbitrarily C^r -small one, and we use the same notations for the continuations of hyperbolic sets after a perturbation.

(*Saddle-focus case.*) The existence of the desired blenders, their homoclinic relations to O_1 and O_2 , and the fact that they form a C^1 -cycle follow directly from Theorem 5.6. We thus only prove the part regarding the existence of robust homoclinic tangencies. By considering f^{-1} , it suffices to prove that Λ_1 is C^1 -wild if either

- (1) O_1 has a homoclinic tangency, or
- (2) $\gamma = |\gamma|e^{i\omega}$ with $\omega \in (0, \pi)$.

Let us start with case (1). By the multidimensional Newhouse theorem, we first, while keep the heterodimensional cycle, unfold this tangency to obtain a hyperbolic basic set Λ_0 which has a C^2 -robust homoclinic tangency and, by [GTS93b, GLM], is homoclinically related to O_1 . Next, we

use a further perturbation to make the heterodimensional cycle non-degenerate, and then apply Theorems 5.6 and 5.3. After that, we obtain a separated and k -arrayed blender Λ which is homoclinically related to O_1 , and hence also homoclinically related to Λ_0 . By the lambda lemma, the leaves of the stable/unstable invariant laminations of Λ accumulate on those of Λ_0 . So, one can perturb f again to create a homoclinic tangency to Λ , while keeping Λ as an separated and k -arrayed blender (since this is a C^1 -open property). Finally, we invoke Theorem 3 to conclude the proof for the case where O_1 has a homoclinic tangency.

We proceed to consider case (2), where the center-unstable multiplier γ of O_2 is nonreal. By Theorems 5.6 and 5.3, we first perturb f to obtain a standard blender which is both separated and k -arrayed, and is homoclinically related to O_1 . Since this perturbation does not destroy the heterodimensional cycle, we apply Lemma 5.7 to create a homoclinic tangency to O_1 by a further perturbation. The proof is then completed with an application of the lambda lemma and Theorem 3.

(*Double-focus case.*) The proof is the same as for the saddle-focus case.

(*Saddle case.*) The proof in this case is almost identical to case (1) of the saddle-focus case, except that the C^1 -cycle involving Λ_1 and Λ_2' is not automatic. To obtain this, we need to use the ‘moreover’ part of the saddle case in Theorem 5.6, namely, we need to show that O_1 or O_2 is contained in non-trivial hyperbolic set. But this is immediate, because, after the first step of case (1), either O_1 or O_2 (depending on which one has the tangency) is homoclinically related to the hyperbolic basic set Λ_0 . \square

5.2 Robust homoclinic tangencies in parameter families. Proof of Corollary 6

Let us assume $|\lambda_1\gamma_1| > 1$, and the proof of the other case is the same by considering f^{-1} . Denote the argument of λ_1 by ω and its value at f by ω^* . It also suffices to assume $\varepsilon = (\mu, \omega)$. It is established in [LLST22, Theorem 5.1] that there exists a sequence $(\mu_i, \omega_i) \rightarrow (0, \omega^*)$ such that, for every sufficiently large i , the map f_{μ_i, ω_i} has a heterodimensional cycle involving O and Q , a periodic point with $\text{ind}(Q) = \text{ind}(O) + 1$; and, moreover, this cycle at $(\mu, \omega) = (\mu_i, \omega_i)$ unfolds generically when μ varies from μ_i . Let us take $(\mu, \omega) = (\mu_i, \omega_i)$ and adjust the parameter values as follows, where all changes can be arbitrarily small.

1. change μ and ω so that, with the cycle being kept, a standard blender Λ_2 appears and is homoclinically related to Q , by [LLST22, Lemma 6.2 and Theorem 6.1],
2. change μ and ω so that a quadratic homoclinic tangency to Q is created, by Lemma 5.7 (and the discussion followed),
3. change μ to create a quadratic tangency to Λ_2 , by the lambda-lemma, and
4. change μ to create C^1 -robust homoclinic tangencies, as in the proof of Theorem 3.

Denote those new parameter values as (μ'_i, ω'_i) . Repeating the above procedure for all sufficiently large i , we find the desired sequence of parameter values $\{(\mu'_i, \omega'_i)\}$.

5.3 Standard blenders arisen form an NABS. Proof of Theorem 5.3

We will only consider the cs-NABS, and the proof for the center-unstable case is completely parallel. Let us identify $F_{i,\delta}$ with $\varphi_\delta \circ F_{i,\delta} \circ \varphi_\delta^{-1}$. So, $F_{i,\delta}$ is a map defined on Π_i given by (8) and satisfy the cross-form relation (6). We prove Theorem 5.3 in several steps. Roughly speaking, we first find a specification set \mathcal{I} such that the BCG covering property in Definition 4.3 is satisfied, and

hence the cs-Markov partition $(\{\Pi_i\}_{i \in \mathcal{I}}, g_\delta)$ induces a k -arrayed blender. Then, we show that by ‘duplicating’ this set \mathcal{I} the obtained arrayed blender can also be a separated one.

5.3.1 Preliminaries. Let us define the BCG structure for Π_i in Definition 4.1 (which also includes the BC structure in Definition 3.2). By (8), the sets $\mathbb{X}_i, \mathbb{Y}_i, \mathbb{Z}_i$ in Definition 2.1 are the same for all i in our case, and, after dropping the subscript, they are $\mathbb{X} = [-1, 1], \mathbb{Y} = [-1, 1]^{d_u}, \mathbb{Z} = [-1, 1]^{d_{ss}}$. Denote $\Pi = [-1, 1]^d$. We take real numbers x_B, x_C, x_G with $0 < x_C < x_G < x_B < 1$ to be determined, and define for all i

$$\mathbb{X}^B = (-x_B, x_B), \quad \mathbb{X}^C = (-x_C, x_C), \quad x^L = -x_G, \quad x^R = x_G,$$

and

$$\begin{aligned} \Pi^B &= (-x_B, x_B) \times \mathbb{Y} \times \mathbb{Z}, & \Pi^C &= (-x_C, x_C) \times \mathbb{Y} \times \mathbb{Z}, \\ \mathring{\Pi}^L &= \{-x_G\} \times \mathbb{Y} \times \mathbb{Z}, & \mathring{\Pi}^R &= \{x_G\} \times \mathbb{Y} \times \mathbb{Z}, \\ \Pi_i^B &= \Pi_i \cap \Pi^B, & \Pi_i^C &= \Pi_i \cap \Pi^C, & \mathring{\Pi}_i^L &= \Pi_i \cap \mathring{\Pi}^L, & \mathring{\Pi}_i^R &= \Pi_i \cap \mathring{\Pi}^R. \end{aligned} \quad (11)$$

Consider the map g defined on all Π_i by $g(x, y, z) = F_{i, \delta}(x, y, z)$ if $(x, y, z) \in \Pi_i$. By (6), we have

$$\begin{aligned} V_i &:= g(\Pi_i) = \{(\hat{\psi}_{1,i}(x, y, z), y, \psi_{3,i}(x, y, z)) : (x, y, z) \in \mathbb{X} \times \mathbb{Y} \times \mathbb{Z}\}, \\ V_i^{B/C} &:= g(\Pi_i^{B/C}) = \{(\hat{\psi}_{1,i}(x, y, z), y, \psi_{3,i}(x, y, z)) : (x, y, z) \in \mathbb{X}^{B/C} \times \mathbb{Y} \times \mathbb{Z}\}, \\ \mathring{V}_i^{L/R} &:= g(\mathring{\Pi}_i^{L/R}) = \{(\hat{\psi}_{1,i}(x, y, z), y, \psi_{3,i}(x, y, z)) : (x, y, z) \in \{x^{L/R}\} \times \mathbb{Y} \times \mathbb{Z}\}. \end{aligned} \quad (12)$$

Note that the vertical strips which constitute the vertical arrays in the BCG covering property (Definition 4.3) are those inside Π_j . Hence, in our case, they are of the form $V_{ij} = V_i \cap \Pi_j$. Similarly, we have $V_{ij}^{B/C/L/R} = V_i^{B/C/L/R} \cap \Pi_j$. Combining this fact with the above BCG structure of Π_i , we obtain

Lemma 5.8. *The hyperbolic basic set Λ induced from the cs-Markov partition $(\{\Pi_i\}_{i \in \mathcal{I}}, g_\delta)$ is a k -arrayed blender if the BCG covering property is satisfied with Π and $\mathcal{V} \subset \{V_i : i \in \mathcal{I}\}$, in place of Π_i and $\mathcal{V} \subset \{V_{ij} : i, j \in \mathcal{I}\}$.*

Recall Definition 2.5 that an ss-disc S is of the form $(x, y) = (s_1(z), s_2(z))$ for $z \in \mathbb{Z}$ and a u-disc S^u is of the form $(x, z) = (s_1^u(y), s_2^u(y))$ for $y \in \mathbb{Y}$. Let Δ_S^{ss} and $\Delta_{S^u}^u$ be the maximal deviation of the x -coordinate of an ss-disc S and a u-disc S^u , i.e.,

$$\Delta_S^{ss} = \max_{z_1, z_2 \in [-1, 1]^{d_{ss}}} |s_1(z_1) - s_1(z_2)| \quad \text{and} \quad \Delta_{S^u}^u = \max_{y_1, y_2 \in [-1, 1]^{d_{ss}}} |s_1^u(y_1) - s_1^u(y_2)|.$$

Denote by Δ_i^{ss} the supremum of Δ_S^{ss} over all ss-discs S crossing Π_i , and by Δ_i^u the supremum of $\Delta_{S^u}^u$ over all u-discs S^u crossing Π_i . One sees from (9) that $\max\{\Delta_i^{ss}, \Delta_i^u\} = o(1)_\delta + o(1)_{\delta \cdot i \rightarrow \infty}$. (Recall the meaning of this notation in (S1).)

By (6), the image $\hat{\psi}_{1,i}(\mathbb{X}^{B/C} \times \mathbb{Y} \times \mathbb{Z})$ contains the interval $(B_i - A_i x_B + \|\psi_{1,i}\|, B_i + A_i x_B - \|\psi_{1,i}\|)$. By the form of V_i^B in (12), one can easily check⁴ that any ss-disc which lies entirely in

$$(B_i - |A_i| x_B + \|\psi_{1,i}\|, B_i + |A_i| x_B - \|\psi_{1,i}\|) \times \mathbb{Y} \times \mathbb{Z} \quad (13)$$

crosses V_i^B in the sense of Definition 2.7. By (S1), (S3) and (9), we have

$$\Delta_i := \max\{|A_i| - \alpha, \|\psi_{1,i}\|, \Delta_i^{ss}, \Delta_i^u\} = o(1)_\delta + o(1)_{\delta \cdot i \rightarrow \infty}. \quad (14)$$

⁴ An elaborated computation can be found in the proof of [LT24, Lemma 4.10].

Lemma 5.9. *If an ss-disc crosses Π^{B} and intersects $\mathbb{X}(V_i^{\text{B}}) \times \mathbb{Y} \times \mathbb{Z}$, then it crosses V_i^{B} , where $\mathbb{X}(V_i^{\text{B}}) = (B_i - \alpha x_{\text{B}} + 3\Delta_i, B_i + \alpha x_{\text{B}} - 3\Delta_i)$.*

Proof. By Definition 2.7 of crossing, the ss-disc lies entirely in $(-x_{\text{B}}, x_{\text{B}}) \times \mathbb{Y} \times \mathbb{Z}$. By the choice of Δ , if it intersects the cube in the statement of the lemma, then it must lie entirely in the cube given by (13). \square

5.3.2 Finding 1-arrayed standard blenders. Let us now find a specification set \mathcal{I} such that $(\{\Pi_i\}_{i \in \mathcal{I}}, g_\delta)$ satisfies 1-arrayed BCG covering property as in Definition 4.3, i.e., every vertical array has only one vertical strip. After that, we show that the k -arrayed covering property can be satisfied by repeating the same argument k times. By Lemma 5.8, we verify the properties (B1)–(B4) with Π and $\mathcal{V} \subset \{V_i : i \in \mathcal{I}\}$ in Lemmas 5.10–5.14 below.

Take any $x_{\text{B}} \in (0, 1)$ such that $[-x_{\text{B}}, x_{\text{B}}]$ is contained in the interval in which $\{B_i\}$ is dense, given by (S2). Let $N > 0$ be an integer to be determined. We divide the interval $[-x_{\text{B}}, x_{\text{B}}]$ equally into N subintervals and denote the endpoints by x_n with $n = 0, \dots, N$, $x_0 = -x_{\text{B}}$ and $x_N = x_{\text{B}}$. By 14, we can take a sequence $\{\delta_s\} \rightarrow 0$ and a family $\{\mathcal{I}_s\}_{s \in \mathbb{N}}$ of sets $\mathcal{I}_s = \{i_1(s), \dots, i_N(s)\}$ such that

$$\Delta_{i_n(s)} \rightarrow 0 \quad \text{and} \quad B_{i_n(s)} - x_n \rightarrow 0 \quad \text{as} \quad s \rightarrow \infty.$$

We further require the family of pairs $\{(\delta_s, \mathcal{I}_s)\}$ is chosen such that $\Delta_{i_n(s)} \rightarrow 0$ as $s \rightarrow \infty$. Let $\lfloor \cdot \rfloor$ denote the largest integer smaller than a given number.

Lemma 5.10. *Given any integer N with $N \geq \lfloor 1/\alpha \rfloor + 2$, the specification sets \mathcal{I}_s satisfy property (B1) for all sufficiently large s .*

Proof. Note that, if the union of the associated intervals $\mathbb{X}(V_i^{\text{B}})$ with $i \in \mathcal{I}$ covers $[-x_{\text{B}}, x_{\text{B}}]$, then every ss-disc crossing Π^{B} must intersect $\mathbb{X}(V_i^{\text{B}}) \times \mathbb{Y} \times \mathbb{Z}$ for some i , and hence cross V_i^{B} by Lemma 5.9, which gives (B1).

Taking

$$\tilde{\Delta}_s = \max_{n=1, \dots, N} \{\Delta_{i_n(s)}, |B_{i_n(s)} - x_n|\}, \quad (15)$$

one sees that $\mathbb{X}(V_{i_n(s)}^{\text{B}})$ contains the subinterval

$$\tilde{\mathbb{X}}(V_{i_n(s)}^{\text{B}}) := (x_n - \alpha x_{\text{B}} + 4\tilde{\Delta}_s, x_n + \alpha x_{\text{B}} - 4\tilde{\Delta}_s).$$

Hence, $[-x_{\text{B}}, x_{\text{B}}]$ is covered by $\bigcup_n \mathbb{X}(V_{i_n(s)}^{\text{B}})$ if it is covered by $\bigcup_n \tilde{\mathbb{X}}(V_{i_n(s)}^{\text{B}})$. But the latter is obvious for all sufficiently large s , since

$$\sum_n \left| \tilde{\mathbb{X}}(V_{i_n(s)}^{\text{B}}) \cap [-x_{\text{B}}, x_{\text{B}}] \right| > (2\alpha x_{\text{B}} - 8\tilde{\Delta}_s) \cdot (\alpha^{-1} + 1) = 2x_{\text{B}} + 2\alpha x_{\text{B}} + o(1)_{s \rightarrow \infty},$$

where $|\cdot|$ denotes the length of an interval. It then follows from the discussion at the beginning that (B3) holds for all \mathcal{I}_s with sufficiently large s . \square

Lemma 5.11. *The specification sets \mathcal{I}_s satisfy property (B4) for all sufficiently large s .*

Proof. By the choice of Δ_i in (14), Definition 4.1 and the BCG structure in (11), we find that

$$\Pi^{\text{L}} \subset (-x_{\text{G}} - 3\Delta_i, -x_{\text{G}} + 3\Delta_i) \times \mathbb{Y} \times \mathbb{Z} \quad \text{and} \quad \Pi^{\text{R}} \subset (x_{\text{G}} - 3\Delta_i, x_{\text{G}} + 3\Delta_i) \times \mathbb{Y} \times \mathbb{Z}, \quad (16)$$

where we use three Δ_i to bound $\|\psi_{1,i}\|$, the x -deviations of the ss-discs and of the u-discs, respectively. Recall the separation notation \parallel is defined before Definition 4.3. It is immediate from (11) that $\Pi_i^{L/R} \parallel \Pi_i^C$ if $x_G - x_C > 4\Delta_i$, which is automatic for all sufficiently large i .

Let us check $V_i^R \parallel V^C$. Recall that $|A_i - \alpha| < \Delta_i$. We find from (6) and (12) that

$$\begin{aligned} V_i^R &\subset ((\alpha + \Delta_i)(x_G - 3\Delta_i) + B_i - \Delta_i, (\alpha + \Delta_i)(x_G + 3\Delta_i) + B_i + \Delta_i) \times \mathbb{Y} \times \mathbb{Z}, \\ V_i^C &\subset (-(\alpha + \Delta_i)x_C + B_i - \Delta_i, (\alpha + \Delta_i)x_C + B_i + \Delta_i) \times \mathbb{Y} \times \mathbb{Z}. \end{aligned}$$

Thus, $V_i^R \parallel V^C$ if

$$((\alpha + \Delta_i)(x_G - 3\Delta_i) + B_i - \Delta_i) - ((\alpha + \Delta_i)x_C + B_i + \Delta_i) > \Delta_i,$$

where the Δ_i on the right-hand side is used to bound the x -deviation of the ss-discs. This inequality is satisfied when $x_G - x_C > 9\Delta_i\alpha^{-1}$. By a similar discussion, the same condition can be derived from $V_i^L \parallel V^C$ and $V_i^{L/R} \parallel \partial_x^{L/R}V^B$. The lemma follows immediately since by construction $i \in \mathcal{I}_s$ tend to infinity as $s \rightarrow \infty$. \square

Lemma 5.12. *Properties (B3) and (B3') are satisfied by $\mathcal{I}_s = \{x_0, \dots, x_N\}$ for all sufficiently large s if*

- N is odd and $-x_C = x_n$ for some $0 < n < (N - 1)/2$, and
- $x_G > x_C + \alpha x_B$.

Proof. Let $\tilde{\Delta}_s$ be given by (15). By (6) and (12),

$$V_{i_n(s)}^C \supset (x_n - \alpha x_C + 3\tilde{\Delta}_s, x_n + \alpha x_C - 3\tilde{\Delta}_s) \times \mathbb{Y} \times \mathbb{Z}, \quad (17)$$

where the term $3\tilde{\Delta}_s$ comes from $|\alpha - A_{i_n(s)}|$, $|x_C - B_{i_n(s)}|$, and $\|\psi_{1,i_n(s)}\|$. Since every ss-disc intersecting $\partial_x^L \Pi^C = \{-x_C\} \times \mathbb{Y} \times \mathbb{Z}$ has its x -deviation bounded by $2\tilde{\Delta}_s$ from $x = -x_C = x_n$, it must cross the above cube if $2\alpha x_C - 6\tilde{\Delta}_s > 2\tilde{\Delta}_s$, which is automatic for all large s . This proves the first part of (B3).

To check $V_{i_n(s)}^B \parallel \Pi^L$, we obtain from (6) and (12) that

$$V_{i_n(s)}^B \subset (x_n - \alpha x_B - 3\tilde{\Delta}_s, x_n + \alpha x_B + 3\tilde{\Delta}_s) \times \mathbb{Y} \times \mathbb{Z}. \quad (18)$$

Since Π^L belongs to the first cube in (16) and $x_n = -x_C$, it follows that $V_{i_n(s)}^B \parallel \Pi^L$ if $-x_C - \alpha x_B + x_G > 7\tilde{\Delta}_s$, which is fulfilled by all large s due to the second condition in the lemma. Since N is odd, the discussion for (B3') is the same by symmetry. \square

Recall that the points x_n divide $[-x_B, x_B]$ into N subintervals of the same length.

Lemma 5.13. *Properties (B2) and (B2') are satisfied by $\mathcal{I}_s = \{x_0, \dots, x_N\}$ for all sufficiently large s if*

$$\alpha(x_B - x_C) < x_{n+1} - x_n < \alpha(x_B + x_C). \quad (19)$$

More specifically, for $n = 0, \dots, N - 1$, any ss-disc intersecting $\partial_x^R V_{i_n(s)}^B$ crosses $V_{i_{n+1}(s)}^C$, and $V_{i_{n+1}(s)}^B \parallel V_{i_n(s)}^L$; and, for $n = 1, \dots, N$, any ss-disc intersecting $\partial_x^L V_{i_{n+1}(s)}^B$ crosses $V_{i_n(s)}^C$, and $V_{i_n(s)}^B \parallel V_{i_{n+1}(s)}^R$.

Proof. Again by (6) and (12), we find that $\partial_x^R V_{i_n(s)}^B$ lies between the following two hyperplanes:

$$\{x_n + \alpha x_B - 3\tilde{\Delta}_s\} \times \mathbb{Y} \times \mathbb{Z} \quad \text{and} \quad \{x_n + \alpha x_B + 3\tilde{\Delta}_s\} \times \mathbb{Y} \times \mathbb{Z},$$

while, by (17),

$$V_{i_{n+1}(s)}^C \supset (x_{n+1} - \alpha x_C + 3\tilde{\Delta}_s, x_{n+1} + \alpha x_C - 3\tilde{\Delta}_s) \times \mathbb{Y} \times \mathbb{Z}.$$

Thus, an ss-disc which intersects $\partial_x^R V_{i_n(s)}^B$ will cross $V_{i_{n+1}(s)}^C$ if

$$\begin{aligned} x_n + \alpha x_B - 3\tilde{\Delta}_s - (x_{n+1} - \alpha x_C + 3\tilde{\Delta}_s) &> \tilde{\Delta}_s, \\ -(x_n + \alpha x_B + 3\tilde{\Delta}_s) + x_{n+1} + \alpha x_C - 3\tilde{\Delta}_s &> \tilde{\Delta}_s, \end{aligned}$$

or

$$x_{n+1} - x_n < \alpha(x_B + x_C) - 7\tilde{\Delta}_s \quad \text{and} \quad x_{n+1} - x_n > \alpha(x_B - x_C) + 7\tilde{\Delta}_s,$$

which holds for all sufficiently large s by (19). This proves the first part of (B2).

To check the second part, we find from (18)

$$V_{i_{n+1}(s)}^B \subset (x_{n+1} - \alpha x_B - 3\tilde{\Delta}_s, x_{n+1} + \alpha x_B + 3\tilde{\Delta}_s) \times \mathbb{Y} \times \mathbb{Z},$$

and from (6) and (12)

$$V_{i_n(s)}^L \subset (-\alpha x_G + x_n - 7\tilde{\Delta}_s, -\alpha x_G + x_n + 7\tilde{\Delta}_s) \times \mathbb{Y} \times \mathbb{Z}.$$

Obviously, we have $V_{i_{n+1}(s)}^B \parallel V_{i_n(s)}^L$ if

$$x_{n+1} - \alpha x_B - 3\tilde{\Delta}_s - (-\alpha x_G + x_n + 6\tilde{\Delta}_s) > \tilde{\Delta}_s,$$

which holds for all sufficiently large s by (19). The proof of (B2') is completely parallel due to the symmetry. \square

Lemma 5.14. *If $\alpha < 1/20$, then x_C, x_G, N can be chosen such that the conditions in Lemmas 5.10, 5.12 and 5.13 are satisfied.*

Proof. Since $x_{n+1} - x_n = 2x_B/N$, inequality (19) can be rewritten as

$$\frac{2x_B}{\alpha(x_B + x_C)} < N < \frac{2x_B}{\alpha(x_B - x_C)}. \quad (20)$$

We claim that, for every N satisfying this inequality and has the form $N = 8m + 1$ with $m \geq 1$, one can define the desired x_C, x_G .

Let us prove the claim. Take $x_C = -x_{m+1}$, so $x_C/x_B = (m - 1/2)/(4m + 1/2) \in (1/9, 1/4)$, and hence

$$\frac{1}{9}x_B < x_C < \frac{1}{4}x_B. \quad (21)$$

This along with (20) implies

$$\frac{8}{5\alpha} < N < \frac{9}{4\alpha}. \quad (22)$$

The condition of Lemma 5.10 is satisfied since $8/(5\alpha) - (\lfloor 1/\alpha \rfloor + 2) > 32 - 22 > 0$. The first condition of Lemma 5.12 is automatic by construction, and the second one is equivalent to require $x_C + \alpha x_B < 1$, which is also satisfied since $x_C + \alpha x_B < x_B/4 + x_B/20 < 1$. So, x_C can be taken as any number satisfying the second condition of Lemma 5.12. The claim is proven.

It remains to show that there indeed exist integers of the form $N = 8m + 1$ satisfying (20). But

this is obvious since

$$\frac{2x_B}{\alpha(x_B - x_C)} - \frac{2x_B}{\alpha(x_B + x_C)} > \frac{9}{4\alpha} - \frac{9}{5\alpha} > 9.$$

□

5.3.3 Finding k -arrayed standard blenders. Lemmas 5.10–5.14 imply that, when $\alpha < 1/20$, there exists a family $\{\mathcal{I}_s\}$ of specification sets defined above Lemma 5.10 such that for every sufficiently large s , the cs-Markov partition $(\{\Pi_i\}_{i \in \mathcal{I}_s}, g_\delta)$ induces a 1-arrayed blender. To obtain a k -arrayed blender for any $k > 1$, we find k families $\{\mathcal{I}_s^j\}_{s \in \mathbb{N}, 1 \leq j \leq k}$ via Lemmas 5.10–5.14 by taking $\tilde{\Delta}_s = \max_j \{\tilde{\Delta}_s^j\}$, where $\tilde{\Delta}_s^j$ are defined as (15) but now with the superscript. Then, for every small δ and large s , the cs-Markov partition $(\{\Pi_i\}_{i \in \bigcup_j \mathcal{I}_s^j}, g_\delta)$ induces a k -arrayed blender. To remove the assumption $\alpha < 1/20$, one simply considers the iteration g^n for some n such that the new contraction coefficient $\alpha^n < 1/20$.

5.3.4 Building a separated blender: completion of the proof. Recall that the BC structure (see Definition 3.2) is defined as in (11). Denote by \mathcal{I} the specification set of the found k -arrayed blender. In what follows, we show that by adding two more copies of \mathcal{I} with some shifts in the x -direction, the resulting bigger specification set yields a standard blender which is simultaneously k -arrayed and separated.

Recall the numbers x_i ($i = 0, \dots, N$) defined above Lemma 5.10 and denote $D = x_1 - x_0$. We take

$$x'_i = x_i + \frac{1}{3}D, \quad x'_{-1} = x_0 - \frac{2}{3}D, \quad x''_i = x_i + \frac{2}{3}D, \quad x''_{-1} = x_0 - \frac{1}{3}D.$$

Let \mathcal{I}' and \mathcal{I}'' be the specification sets found in the same way as \mathcal{I} , but with $\{x'_i\}$ and $\{x''_i\}$ instead of $\{x_i\}$ in Lemmas 5.10–5.14. Evidently, the standard blender of $\hat{\mathcal{I}} = \mathcal{I} \cup \mathcal{I}' \cup \mathcal{I}''$ is still a k -arrayed one, where the extra points x'_{-1} and x''_{-1} are for ensuring that property (B2') is satisfied by each of the array set given by \mathcal{I}' and \mathcal{I}'' .

We now prove that the blender of $\hat{\mathcal{I}}$ is also a separated one. Property (A1) is immediate from (B1). To verify (A2), it suffices to show that, instead of $\mathcal{V}_2^B \parallel \mathcal{V}_1^L$ in (B2), the stronger property $\mathcal{V}_2^C \parallel \mathcal{V}_1^C$ holds, and similarly for (B2'). Due to the symmetry, we only prove $\mathcal{V}_2^C \parallel \mathcal{V}_1^C$. One just needs to replace $V_{i_{n+1}(s)}^B$ and $V_{i_n(s)}^L$ in the second part of the proof of Lemma 5.13 by $V_{i_{n+1}(s)}^C$ and $V_{i_n(s)}^C$. This yields a condition that can be satisfied for all large s if $x_{n+1} - x_n > 2\alpha x_C$, which, by the choice of x_C in Lemma 5.14 and the fact that $x_{n+1} - x_n = 2x_B/N$, is equivalent to

$$\alpha < \frac{4m + \frac{1}{2}}{(8m + 1)(m - \frac{1}{2})}.$$

This can be satisfied for any fixed m by consider iterations of the NABS, as explained in Section 5.3.3. Thus, (A2) is satisfied.

We proceed to check a stronger version of (A3) that every ss-disc intersecting Π_i^B will cross some V^C . This is just a repetition of the proof of Lemma 5.10, with requiring that $[-x_B, x_B]$ is covered by $\mathbb{X}(V)_i^C$, instead of $\mathbb{X}(V)_i^B$, with $i \in \hat{\mathcal{I}}$. It follows that the desired property holds if $\text{card } \hat{\mathcal{I}} = 3N \geq \lfloor x_B/\alpha x_C \rfloor + 2$, which is automatic by (21) and (22). Thus, the k -arrayed blender of $\hat{\mathcal{I}}$ is also separated, and the proof of Theorem 5.3 is completed.

Acknowledgment

I am grateful to Dmitry Turaev for important conversations during the preparation of this paper, especially on the formulation of the arrayed standard blenders. I also thank Masayuki Asaoka, Xiaolong Li, Dmitrii Mints and Zhiyuan Zhang for useful discussions.

A NABS obtained in [LT24]

A.1 Proof of Theorem 5.6

We now collect several results in [LT24] which together imply Theorem 5.6. In some cases, we will also refer to [Li24] where the corresponding improved results are obtained. The statements there hold for all standard blenders arisen from the NABS near heterodimensional cycles, which in particular include the separated and arrayed ones introduced in this paper.

In [LT24, Section 2.5], saddle heterodimensional cycles are further divided into two subcases, type I and type II, according to whether the transition map along \mathcal{T}^0 satisfies an orientation-related condition or not. The existence of the NABS for type-I cycles follows from the renormalization formulas (3.26) and (4.21) of [LT24], with replacing (X, Y, Z) by $(\delta x, \delta y, \delta z)$ in these two formulas. The part on the homoclinic relation and C^1 -cycle follows from [LT24, Theorem 2], while the additional ‘moreover’ part concerning the case where one of the saddles is contained in a non-trivial hyperbolic set is by [LT24, Corollary 3] (see also [LT24, Remark 2.2 and Theorem 4] which imply the corollary). The NABS part for type-II cycles is due to [LT24, Theorem 5] that saddle cycles of type II are accumulated by those of type I, and other parts are from [LT24, Corollary 2]. The proofs are based on the analysis of first return maps on a family $\{U_\delta\}$ of shrinking neighborhoods of some point $P \in \Gamma$, where δ denotes the size. A single renormalization map $\varphi : U \rightarrow \mathbb{R}^d$ for some $U \supset U_\delta$ is considered, that is, $\varphi_\delta = \varphi|_{U_\delta}$. In this case, one has $\{n_i(\delta_2)\} \subset \{n_i(\delta_1)\}$ for any $\delta_2 < \delta_1$.

The NABS part of Theorem 5.6 for saddle-focus and double-focus cycles follows from formulas (6.18) and, respectively, (6.44) of [LT24], with replacing (X, Y, Z) and, respectively, (U, V, W) by $(\delta x, \delta y, \delta z)$. A more concise presentation of these two formulas can be found in [Li24, Sections 2.2 and 2.3]. The convergence requirements in (S1) and (S1’) are described in detail by Lemma A.1 below.

The part on the homoclinic relation and C^1 -cycle is given by [LT24, Theorem 7]. In these two cases, the proofs are still based on a family of shrinking neighborhoods but now the involved renormalization maps depend on the size of the neighborhoods.

The first perturbation in Theorem 5.6 is to achieve certain arithmetic properties on the central multipliers, which guarantee the existence of the nearly-affine blender systems. These properties are fully classified in [Li24, Corollary A]. The exception is for saddle cycles of type II, where the cycle is additionally unfolded to create those of type I. The second perturbation for homoclinic relations and C^1 -cycles are done by unfolding the heterodimensional cycle, and this can be done within generic finite-parameter families, where a complete discussion is given in by [LT24, Theorem 2] for the saddle case and [Li24, Theorem C] for other cases.

A.2 NABS in the saddle-focus and double-focus cases

Theorem 5.6 in these two cases establishes the coexistence of a cs-NABS and a cu-NABS, and they carry additional properties not used in this paper. First, in contrast to the saddle case, where A_i in (6) converge to the quantity α , we have

Lemma A.1 (Full central contraction/expansion). *In the saddle-focus and double-focus cases, it holds that for any $\alpha \in \mathbb{R} \setminus \{0, \pm 1, \pm \infty\}$, there exists a sequence $\{i_n\}_{n \in \mathbb{N}}$ such that $A_{i_n} \rightarrow \alpha$ as $n \rightarrow \infty$ and $\{B_{i_n}\}$ is dense near 0.*

Recall that the map (6) are obtained as the first return map along the heterodimensional cycle, and they are of the form $F_i := F_{k_i, m_i}$, where k_i and m_i are the number of iterations spent near O_1 and O_2 , respectively. The result in the saddle-focus case follows from the proof of [LT24, Proposition 6.2], or, more specifically, the last displayed equation in the proof. For convenience we present it here with a concise form.

$$A_{k_i, m_i} = \frac{(B_{k_i, m_i} - c) \sin(k\omega + \eta_1)}{\sin(k\omega + \eta_2)},$$

where $A_{k_i, m_i}, (B_{k_i, m_i} - c)$ are the quantities A_i and B_i in (6), ω is the argument of the complex center-stable multiplier of O_1 , and $c, \eta_{1,2}$ are constants with $\eta_1 \neq \eta_2 + n\pi$. The arithmetic condition of [LT24, Theorem 7] ensures that $\{k\omega/2\pi, B_{k,m}\}_{k,m \in \mathbb{N}}$ is dense in \mathbb{R}^2 . This implies that, given any α , one can choose (k_i, m_i) such that $\{(B_{k_i, m_i} - c)\}$ is dense near a small neighborhood of 0, and, in the same time, $A_{k_i, m_i} \rightarrow \alpha$, proving the lemma in the case where the center-stable multipliers of O_1 are nonreal. When O_2 has nonreal center-unstable multipliers, one just considers f^{-1} . This lemma for the double-focus case follows from a reduction to the saddle-focus one in the proof of [LT24, Proposition 6.7].

Lemma A.2 (Overlapping). *The cs- and cu-NABSs, which constitute the full NABS in the saddle-focus and double-focus cases, share the same family $\{(U_\delta, \varphi_\delta)\}$ of charts in Definition 5.1.*

This lemma follows directly from [LT24, Propositions 6.2 and 6.7]. An immediate consequence due to Proposition 2.11 is that, for any pair of standard cs- and cu-blender arisen from such full NABS, denoted as Λ^{cs} and Λ^{cu} , one has that $W^u(\Lambda^{cs})$ intersects $W^s(\Lambda^{cu})$ C^1 -robustly.

References

- [Asa08] Asaoka, M. Hyperbolic sets exhibiting C^1 -persistent homoclinic tangency for higher dimensions. *Proc. Amer. Math. Soc.*, 136(2):677–686, 2008. [Cited on page 3.]
- [Asa22] Asaoka, M. Stable intersection of cantor sets in higher dimension and robust homoclinic tangency of the largest codimension. *Trans. Amer. Math. Soc.*, 375:873–908, 2022. [Cited on page 3.]
- [BB23] Berger, P. and Biebler, S. Emergence of wandering stable components. *J. Amer. Math. Soc.*, 36:397–482, 2023. [Cited on page 6.]
- [BBD16] Bochi, J., Bonatti, C., and Díaz, L. J. Robust criterion for the existence of nonhyperbolic ergodic measures. *Comm. Math. Phys.*, 344(3):751–795, 2016. [Cited on page 7.]
- [BC15] Bonatti, C. and Crovisier, S. Center manifolds for partially hyperbolic sets without strong unstable connections. *J. Inst. Math. Jussieu*, 2015. [Cited on page 5.]

- [BCP22] Berger, P., Crovisier, S., and Pujals, E. Germ-typicality of the coexistence of infinitely many sinks. *Adv. Math.*, 407:108528, 2022. [Cited on page 3.]
- [BD96] Bonatti, C. and Díaz, L. J. Persistent nonhyperbolic transitive diffeomorphisms. *Ann. of Math. (2)*, 143(2):357–396, 1996. [Cited on pages 7 and 10.]
- [BD12] Bonatti, C. and Díaz, L. J. Abundance of C^1 -robust homoclinic tangencies. *Trans. Amer. Math. Soc.*, 364(10):5111–5148, 2012. [Cited on pages 3, 11, 12, and 14.]
- [BDL] Barrientos, P. G., Díaz, L. J., and Li, D. Non-dominated cycles of codimension one. *in preparation*. [Cited on page 25.]
- [Ber16] Berger, P. Generic family with robustly infinitely many sinks. *Invent. Math.*, 205(1):121–172, 2016. [Cited on page 3.]
- [BKR14] Barrientos, P. G., Ki, Y., and Raibekas, A. Symbolic blender-horseshoes and applications. *Nonlinearity*, 27(12):2805–2839, 2014. [Cited on page 7.]
- [BR17] Barrientos, P. G. and Raibekas, A. Robust tangencies of large codimension. *Nonlinearity*, 30(12):4369, 2017. [Cited on page 3.]
- [GLM] Gourmelon, N., Li, D., and Mints, D. Stabilization of homoclinic tangencies. *in preparation*. [Cited on pages 6 and 25.]
- [GST08] Gonchenko, S. V., Shilnikov, L. P., and Turaev, D. V. On dynamical properties of multi-dimensional diffeomorphisms from Newhouse regions. I. *Nonlinearity*, 21(5):923–972, 2008. [Cited on pages 3 and 5.]
- [GTS93a] Gonchenko, S. V., Turaev, D. V., and Shilnikov, L. P. Dynamical phenomena in multi-dimensional systems with a structurally unstable homoclinic Poincaré curve. *Dokl. Akad. Nauk*, 330(2):144–147, 1993. [Cited on page 5.]
- [GTS93b] Gonchenko, S. V., Turaev, D. V., and Shilnikov, L. P. On the existence of newhouse regions near systems with non-rough poincaré homoclinic curve (multidimensional case). *Russian Acad. Sci. Dokl. Math.*, 47(2):268–273, 1993. [Cited on pages 2, 6, and 25.]
- [HKY93] Hunt, B. R., Kan, I., and Yorke, J. A. When cantor sets intersect thickly. *Trans. Amer. Math. Soc.*, 339:869–888, 1993. [Cited on page 6.]
- [Kal00] Kaloshin, V. Generic diffeomorphisms with superexponential growth of number of periodic orbits. *Comm Math Phys*, 211:253–271, 2000. [Cited on page 2.]
- [Li24] Li, D. Blender-producing mechanisms and a dichotomy for local dynamics for heterodimensional cycles. *preprint arXiv:2403.15818v1*, 2024. [Cited on pages 7, 11, and 32.]
- [LLST22] Li, D., Li, X., Shinohara, K., and Turaev, D. V. Robust heterodimensional cycles in two-parameter unfolding of homoclinic tangencies. *preprint arXiv:2203.14075v1*, 2022. [Cited on pages 5, 6, 7, and 26.]
- [LT] Li, D. and Turaev, D. V. Typicality of robust coexistence of infinitely many sinks for families of diffeomorphisms. *in preparation*. [Cited on page 3.]
- [LT20] Li, D. and Turaev, D. Persistent heterodimensional cycles in periodic perturbations of Lorenz-like attractors. *Nonlinearity*, 33(3):971–1015, 2020. [Cited on page 6.]

- [LT24] Li, D. and Turaev, D. V. Persistence of heterodimensional cycles. *Invent. Math.*, 236:1413–1504, 2024. [Cited on pages 1, 3, 4, 5, 7, 11, 21, 23, 24, 25, 27, 32, and 33.]
- [Min24] Mints, D. High order homoclinic tangencies of corank 2. *preprint arXiv:2404.08989*, 2024. [Cited on page 3.]
- [Mor11] Moreira, C. G. There are no C^1 -stable intersections of regular cantor sets. *Acta Math.*, 206(2):311–323, 2011. [Cited on page 2.]
- [MV93] Mora, L. and Viana, M. Abundance of strange attractors. *Acta Math.*, 171:1–71, 1993. [Cited on page 2.]
- [New70] Newhouse, S. E. Nondensity of axiom A (a) on \mathbb{S}^2 . *Global analysis*, 1:191–202, 1970. [Cited on page 2.]
- [New74] Newhouse, S. E. Diffeomorphisms with infinitely many sinks. *Topology*, 13:9–18, 1974. [Cited on page 2.]
- [New79] Newhouse, S. E. The abundance of wild hyperbolic sets and nonsmooth stable sets for diffeomorphisms. *Publ. Math. IHES*, (50):101–151, 1979. [Cited on page 2.]
- [NP12] Nassiri, M. and Pujals, E. R. Robust transitivity in Hamiltonian dynamics. *Ann. Sci. Éc. Norm. Supér. (4)*, 45(2):191–239, 2012. [Cited on page 7.]
- [PV94] Palis, J. and Viana, M. High dimension diffeomorphisms displaying infinitely many periodic attractors. *Ann. of Math. (2)*, 140(1):207–250, 1994. [Cited on page 2.]
- [Rom95] Romero, N. Persistence of homoclinic tangencies in higher dimensions. *Ergodic Theory and Dynamical Systems*, 15:735–757, 1995. [Cited on page 2.]
- [Shi67] Shilnikov, L. P. On a poincaré-birkhoff problem. *Math. USSR Sbornik*, 3:91–102, 1967. [Cited on page 9.]
- [Sim72] Simon, C. P. Instability in $\text{Diff}^r(T^3)$ and the nongenericity of rational zeta functions. *Trans. Amer. Math. Soc.*, 174:217–242, 1972. [Cited on page 3.]
- [Tur96] Turaev, D. On dimension of non-local bifurcational problems. *Internat. J. Bifur. Chaos Appl. Sci. Engrg.*, 6(5):919–948, 1996. [Cited on pages 3 and 5.]
- [Ure95] Ures, R. Abundance of hyperbolicity in the C^1 topology. *Ann. Sci. École Norm. Sup. (4)*, 28(6):747–760, 1995. [Cited on page 2.]